

# GEOMETRIC INEQUALITIES FOR QUASI-EINSTEIN MANIFOLDS

RAFAEL DIÓGENES, JACIANE GONÇALVES AND ERNANI RIBEIRO JR

ABSTRACT. In this article, we investigate some geometric inequalities for quasi-Einstein manifolds. We use the generalized Reilly's formulas by Qiu-Xia and Li-Xia to establish new boundary estimates and an isoperimetric type inequality for compact quasi-Einstein manifolds with boundary. Boundary estimates in terms of the first eigenvalue of the Jacobi operator and the Hawking mass are also established. In particular, we present a Heintze-Karcher type inequality for a compact domain on a quasi-Einstein manifold.

## 1. INTRODUCTION

A complete Riemannian manifold  $(M^n, g)$ ,  $n \geq 2$ , will be called *m-quasi-Einstein manifold*, or simply *quasi-Einstein manifold*, if there exists a smooth potential function  $f$  on  $M^n$  such that

$$(1.1) \quad Ric_f^m = Ric + \nabla^2 f - \frac{1}{m} df \otimes df = \lambda g,$$

for some constants  $\lambda$  and  $m > 0$  (cf. [12, 21]). Here,  $\nabla^2 f$  stands for the Hessian of  $f$  and  $Ric$  is the Ricci tensor of  $g$ . In this case,  $f$  is a quasi-Einstein potential. We say that a quasi-Einstein manifold is *trivial* if its potential function  $f$  is constant, otherwise, we say that it is *nontrivial*.

The  $m$ -Bakry-Émery Ricci tensor  $Ric_f^m$  is a natural object in geometric analysis and it is directly related to the study of smooth metric measure spaces and diffusion operators by Bakry and Émery [5]. Observe that  $\infty$ -quasi-Einstein manifold is precisely a gradient Ricci soliton. Moreover, when  $m = 1$ , we consider in addition that  $\Delta e^{-f} + \lambda e^{-f} = 0$  in order to obtain the *static spaces*, which have special interest due to the connection with general relativity. As discussed by Besse [12, pg. 267], a quasi-Einstein manifold is a base of a warped product Einstein metric. Besides, quasi-Einstein manifolds have attracted interest in physics due to their relation with the geometry of a degenerate Killing horizon and horizon limit (see, e.g., [3, 4, 56]). Notice that, for  $m < \infty$ , we may consider the function  $u = e^{-\frac{f}{m}}$  on  $M^n$ , which in turn implies that  $\nabla u = -\frac{u}{m} \nabla f$  and hence, (1.1) becomes

$$(1.2) \quad \nabla^2 u = \frac{u}{m} (Ric - \lambda g).$$

In [32, 33], He, Petersen and Wylie started a fruitful study of quasi-Einstein manifolds with boundary. In this context, a Riemannian manifold  $(M^n, g)$  with (nonempty) boundary  $\partial M$  is an *m-quasi-Einstein manifold* if there exists a smooth potential function  $u$  on  $M^n$

---

*Date:* June 6, 2024.

*2020 Mathematics Subject Classification.* Primary 53C20, 53C25; Secondary 53C65.

*Key words and phrases.* quasi-Einstein manifolds; isoperimetric inequality; boundary estimates; Heintze-Karcher inequality.

R. Diógenes was partially supported by CNPq/Brazil [Grant: 310680/2021-2].

J. Gonçalves was partially supported by CAPES/Brazil - Finance Code 001.

E. Ribeiro was partially supported by CNPq/Brazil [309663/2021-0 & 403344/2021-2], CAPES/Brazil and FUNCAP/Brazil [ITR-0214-00116.01.00/23].

satisfying the system

$$(1.3) \quad \begin{cases} \nabla^2 u = \frac{u}{m}(Ric - \lambda g) & \text{in } M, \\ u > 0 & \text{on } \text{int}(M), \\ u = 0 & \text{on } \partial M. \end{cases}$$

According to [32, Theorem 4.1], nontrivial compact  $m$ -quasi-Einstein manifolds with (non-empty) boundary  $\partial M$  have necessarily  $\lambda > 0$ . Nontrivial examples of compact and noncompact quasi-Einstein manifolds  $(M^n, g, u, \lambda)$  can be found in, e.g., [12, 13, 14, 19, 20, 21, 22, 28, 32, 43, 50, 51, 55]. For what follows, it is important to recall some of them.

(1) *Compact with boundary:*

- (i) the standard hemisphere  $(\mathbb{S}_+^n, dr^2 + \sin^2(r)g_{\mathbb{S}^{n-1}}, u = \cos(r), \lambda > 0)$ , where  $r$  is a height function with  $r \leq \frac{\pi}{2}$ .
- (ii)  $(I \times \mathbb{S}^{n-1}, dt^2 + \frac{n-2}{\lambda}g_{\mathbb{S}^{n-1}}, u(t, x) = \sin(ct), \lambda > 0)$ , where  $c > 0$  is a constant;
- (iii)  $(\mathbb{S}_+^{p+1} \times \mathbb{S}^q, dr^2 + \sin^2 r g_{\mathbb{S}^p} + \frac{q-1}{p+m}g_{\mathbb{S}^q}, u(x, y) = \cos(r(x)), \lambda = p + m)$ , where  $r$  is a height function on  $\mathbb{S}_+^{p+1}$  with  $r \leq \frac{\pi}{2}$  and  $q > 1$ .

(2) *Noncompact:*

- (a)  $([0, \infty) \times F, dt^2 + g_F, u(t, x) = Ct, \lambda = 0)$ ;
- (b) the hyperbolic space  $(\mathbb{H}^n, dt^2 + \sqrt{-k} \sinh^2(\sqrt{-k}t)g_{\mathbb{S}^{n-1}}, u = C \cosh(\sqrt{-k}t), \lambda < 0)$ ;
- (c)  $([0, \infty) \times N, dt^2 + \sqrt{-k} \cosh^2(\sqrt{-k}t)g_{\mathbb{S}^{n-1}}, u(t, x) = C \sinh(\sqrt{-k}t), \lambda < 0)$ ;
- (d)  $(\mathbb{R} \times F, dt^2 + e^{2\sqrt{-k}t}g_F, u(t, x) = Ce^{2\sqrt{-k}t}, \lambda < 0)$ ,

where  $F$  is Ricci flat,  $N$  is an Einstein metric with negative Ricci curvature,  $C$  is an arbitrary positive constant and  $k = \frac{\lambda}{m+n-1}$ .

He, Petersen and Wylie [32] proved that a nontrivial quasi-Einstein manifold with constant Ricci curvature must be isometric to either Example (i), or (a), or (b), or (c), or (d). For the case of compact with boundary, Costa, Ribeiro and Zhou [24] proved recently that a 3-dimensional simply connected compact quasi-Einstein manifold with boundary and constant scalar curvature must be isometric to either Example (i), or (ii). For dimension  $n = 4$ , they proved that a 4-dimensional simply connected compact quasi-Einstein manifold with boundary and constant scalar curvature is isometric to either Example (i), or (ii), or (iii). Despite of that, the classification of compact quasi-Einstein manifolds with boundary and constant scalar curvature still open for dimension  $n \geq 5$ .

Geometric inequalities are fundamental objects of study in geometry. Such inequalities are useful in proving novel obstruction results and put away possible new examples of a special kind of manifolds (or metric). Among the geometric inequalities that motivate the present work, we may mention the isoperimetric and Heintze-Karcher inequalities. The isoperimetric inequality is one of the oldest and most famous inequality in geometry. In  $\mathbb{R}^n$ , the isoperimetric inequality asserts that if  $M \subset \mathbb{R}^n$  is a compact domain with smooth boundary  $\partial M$ , then

$$(1.4) \quad |\partial M| \geq \frac{|\partial \mathbb{B}_1^n|}{Vol(\mathbb{B}_1^n)^{\frac{n-1}{n}}} Vol(M)^{\frac{n-1}{n}},$$

where  $|\partial M|$  denotes the  $(n-1)$ -dimensional volume of  $\partial M$  and  $Vol(M)$  is the volume of  $M$ . In special cases, the  $n$ -dimensional isoperimetric inequality is equivalent to the Sobolev inequality on  $\mathbb{R}^n$  (see [44] and [52]). While the Heintze-Karcher inequality asserts that, for a bounded domain  $\Omega$  in  $\mathbb{R}^n$  with smooth and strictly mean-convex boundary  $\partial\Omega$ , namely,  $H > 0$  on  $\partial\Omega$ ,

$$(1.5) \quad \int_{\partial\Omega} \frac{1}{H} dS \geq \frac{n+1}{n} |\Omega|,$$

where  $H$  is the mean curvature of the  $\partial\Omega$ . This inequality was first outlined by Heintze and Karcher [34] in 1978 and in the form (1.5) by Ros [53] in 1987.

In the last years, there has been a lot of progress on the boundary and volume estimates for special manifolds (or metrics), e.g., static spaces,  $V$ -static spaces, Einstein-type manifolds and critical metrics. For instance, motivated by the classical isoperimetric inequality and a result due Shen [54] and Boucher, Gibbons and Horowitz [17], it was established some boundary and volume estimates for critical metrics of the volume functional in [6, 7, 8, 10, 25, 30, 45, 57] and static spaces in [1, 15, 16, 17, 23, 35, 39, 48], etc. In the same spirit, Diógenes, Gadelha and Ribeiro [27] showed that a compact  $m$ -quasi-Einstein manifold with connected boundary and constant scalar curvature must satisfy

$$(1.6) \quad |\partial M| \leq n \sqrt{\frac{\lambda}{m+n-1}} \text{Vol}(M).$$

In our first result, we shall establish a new estimate for the area of the boundary of a compact  $m$ -quasi-Einstein manifold. In particular, no assumption on the scalar curvature is assumed. More precisely, we have the following result.

**Theorem 1.** *Let  $(M^n, g, u, \lambda)$ ,  $n \geq 3$ , be a compact, oriented  $m$ -quasi-Einstein manifold with connected boundary and  $m > 1$ . Then we have*

$$(1.7) \quad |\partial M| \leq \frac{n\lambda}{\sqrt{(\lambda_1(m+n-1) + \lambda)(m+n-1)}} \text{Vol}(M),$$

where  $\lambda_1$  is the first non-zero eigenvalue of the Laplacian operator. Moreover, if equality holds in (1.7), then  $(M^n, g)$  is isometric, up to scaling, to the standard hemisphere  $\mathbb{S}_+^n$ .

**Remark 1.** *Notice that the estimate (1.7) improves (1.6). Indeed, we have*

$$\frac{n\lambda}{\sqrt{(\lambda_1(m+n-1) + \lambda)(m+n-1)}} < \frac{n\lambda}{\sqrt{\lambda(m+n-1)}} = n \sqrt{\frac{\lambda}{m+n-1}}.$$

In the sequel, we investigate an isoperimetric type inequality for compact  $m$ -quasi-Einstein manifolds with connected boundary and constant scalar curvature. To do so, we first establish a lower bound estimate (depending on the volume of  $M^n$ ) for the area of the boundary. To be precise, we have the following result.

**Theorem 2.** *Let  $(M^n, g, u, \lambda)$ ,  $n \geq 3$ , be a compact, oriented  $m$ -quasi-Einstein manifold with connected boundary, constant scalar curvature  $R$  and  $m > 1$ . Then we have:*

$$(1.8) \quad |\partial M| \geq \frac{(m-1)\alpha}{(m-2)\alpha + \lambda m} \sqrt{\frac{2\lambda mn + \alpha(n(m-1) + 4)}{2n(m+1)m}} \text{Vol}(M),$$

where  $\alpha = n\lambda - R$  is a positive constant. Moreover, if equality holds in (1.8), then  $M^n$  is isometric, up to scaling, to the standard hemisphere  $\mathbb{S}_+^n$ .

A relevant observation is that the proof of Theorem 2 is inspired by the work of Baltazar, Diógenes and Ribeiro [6] combined with a generalized Reilly's formula obtained by Qiu and Xia [48]. As a consequence of Theorem 2, we obtain the following isoperimetric type inequality for compact  $m$ -quasi-Einstein manifolds with connected boundary and constant scalar curvature.

**Corollary 1.** *Let  $(M^n, g, u, \lambda)$ ,  $n \geq 3$ , be a compact, oriented  $m$ -quasi-Einstein manifold with connected boundary, constant scalar curvature  $R$  and  $m > 1$ . Then we have:*

$$(1.9) \quad |\partial M| \geq \Lambda(\alpha, m, n, u)^{\frac{1}{n}} \text{Vol}(M)^{\frac{n-1}{n}},$$

where

$$\Lambda(\alpha, m, n, u) = \frac{\alpha}{m|\nabla u|_{\partial M}} \left( \frac{(m-1)\alpha}{(m-2)\alpha + \lambda m} \sqrt{\frac{2\lambda mn + \alpha(n(m-1) + 4)}{2n(m+1)m}} \right)^{n-1} \int_M u dV_g$$

and  $\alpha = n\lambda - R$ . Moreover, if equality holds in (1.9), then  $M^n$  is isometric, up to scaling, to the standard hemisphere  $\mathbb{S}_+^n$ .

**Remark 2.** It would be interesting to see if the constant  $\Lambda(\alpha, m, n, u)$  in (1.9) can be improved to only depend on the dimension  $n$ ,  $m$  and the volume of the unit ball. We highlight that  $|\nabla u|_{\partial M}$  is a positive constant along  $\partial M$  (see Propositions 2.2 and 2.3 in [32]). Also, it follows from [32, Corollary 4.3] that if a nontrivial compact quasi-Einstein manifold with (nonempty) boundary has constant scalar curvature  $R$ , then  $R < n\lambda$ . Whence,  $\alpha > 0$ .

It is known that the boundary  $\partial M$  of a quasi-Einstein manifold is totally geodesic with the induced metric (see Section 2). Therefore, it is natural to seek for an estimate for the area of the boundary  $\partial M$  in terms of the eigenvalue of Jacobi operator of  $\partial M$ . We recall that given an arbitrary function  $\varphi \in C^\infty(\partial M)$ , the Jacobi operator (or stability operator)  $J$  acting in  $\varphi$  is given by

$$J(\varphi) = \Delta_{\partial M} \varphi + (Ric(\nu, \nu) + |\text{III}|^2)\varphi,$$

where  $\Delta_{\partial M}$  stands for the Laplacian operator on  $\partial M$ ,  $Ric(\nu, \nu)$  is the Ricci curvature of  $M$  in the direction of the outward unit normal vector field  $\nu$  and  $\text{III}$  is the second fundamental form of  $\partial M$ . Besides, let  $\beta_1$  be the first eigenvalue of the Jacobi operator  $J$ , i.e.,  $J(\varphi) = -\beta_1\varphi$ . Thus,  $\beta_1$  is given by

$$(1.10) \quad \beta_1 = \inf_{\varphi \neq 0} \frac{-\int_{\partial M} \varphi J(\varphi) dS_g}{\int_{\partial M} \varphi^2 dS_g}.$$

Specifically, the *index* of  $\partial M$  is the number of (counted with multiplicity) negative eigenvalues of  $J$ . In [8, Theorem 1.10], Barros and Silva proved an estimate for the area of the boundary of a triple static space involving the first eigenvalue of the Jacobi operator, provided that the boundary  $\partial M$  is Einstein and  $\inf R^{\partial M} > 0$ . A similar result was obtained by Costa et al. [23] for a static perfect fluid space-time under the same conditions. In the following, we shall establish a boundary estimate for a compact quasi-Einstein manifold with boundary in terms of the first eigenvalue of the Jacobi operator  $\beta_1$ .

**Theorem 3.** *Let  $(M^n, g, u, \lambda)$ ,  $n \geq 3$ , be a compact  $m$ -quasi-Einstein manifold with connected Einstein boundary and  $m > 1$ . Then we have:*

$$(1.11) \quad |\partial M| \leq \left( \frac{(n-1)(n-2)(m+n-1)}{2(m+n-1)\beta_1 + n(n-1)\lambda} \right)^{\frac{n-1}{2}} \omega_{n-1},$$

where  $\omega_{n-1}$  denotes the volume of the round unit sphere  $\mathbb{S}^{n-1}$ . Moreover, if equality holds in (1.11), then  $\partial M$  is isometric to the round sphere  $\mathbb{S}^{n-1}$ .

One new feature in Theorem 3 is that no scalar curvature condition is assumed. Furthermore, as we will see in Remark 5, the estimate obtained in Theorem 3 improves the estimate established in [27, Theorem 1].

In [27], Diógenes, Gadelha and Ribeiro proved a boundary estimate for compact quasi-Einstein manifolds with boundary in terms of the Brown-York mass. Here, as a consequence of the arguments in the proof of Theorem 3, we shall obtain boundary estimates for 3-dimensional compact quasi-Einstein manifolds involving the Hawking mass. The Hawking

mass<sup>1</sup> of a 2-surface  $\Sigma$  is given by

$$(1.12) \quad \mathfrak{m}_H(\Sigma) = \frac{|\Sigma|^{\frac{1}{2}}}{(16\pi)^{\frac{3}{2}}} \left( 8\pi\chi(\Sigma) - \int_{\Sigma} H^2 dS_g - \frac{2}{3}\Lambda|\Sigma| \right),$$

where  $\Lambda = \inf_M R$  and  $\chi(\Sigma)$  is the Euler characteristic of  $\Sigma$ ; see [9, 31]. The problem of finding explicit lower bounds on Hawking mass is very intriguing; see, e.g., [40, 46, 47]. More precisely, we have the following result.

**Corollary 2.** *Let  $(M^3, g, u, \lambda)$  be a 3-dimensional compact  $m$ -quasi-Einstein manifold with connected boundary and  $m > 1$ . Then the following assertions hold:*

(i)

$$(1.13) \quad \mathfrak{m}_H(\partial M) + \frac{4\lambda}{(m+2)} \left( \frac{|\partial M|}{16\pi} \right)^{\frac{3}{2}} \leq \frac{1}{2} \sqrt{\frac{(m+2)}{\lambda}}.$$

Moreover, if equality holds in (1.13), then  $\partial M$  is isometric to the round sphere  $\mathbb{S}^2$ .

(ii)

$$(1.14) \quad \mathfrak{m}_H(\partial M) \geq 4 \left( \beta_1 + \frac{2\lambda}{m+2} \right) \left( \frac{|\partial M|}{16\pi} \right)^{\frac{3}{2}} \geq 0.$$

**Remark 3.** *Note that (1.14) can be seen as a Penrose inequality for the boundary (see [36]).*

We now turn our attention to the Heintze-Karcher inequality. Recently, there has been a significant progress concerning the Heintze-Karcher type inequalities. For instance, Brendle [18] obtained Heintze-Karcher type inequalities on substatic warped product spaces. By using an elliptic method, Li and Xia [41, 42] established Heintze-Karcher type inequalities for bounded domains in general substatic Riemannian manifolds. Among other results, they proved that letting  $\Omega \subset M$  be a bounded domain with connected strictly mean-convex boundary  $\partial\Omega$  such that  $f|_{\Omega} > 0$  on an  $n$ -dimensional substatic Riemannian manifold  $(M^n, g, f)$ , then the following Heintze-Karcher type inequality holds:

$$(1.15) \quad \int_{\partial\Omega} \frac{f}{H} dS \geq \frac{n}{n-1} \int_{\Omega} f d\Omega.$$

Moreover, if equality holds in (1.15), then  $\partial\Omega$  is umbilical (see [42, Theorem 1.3]). Furthermore, a rigidity statement for the equality case of (1.15) was established by Borghini, Fogagnolo and Pinamonti in [16]; see also related results in [37].

In our next result, as an application of the generalized Reilly's formula obtained by Li and Xia [42], we shall establish an Heintze-Karcher type inequality for a compact domain on a Riemannian manifold satisfying (1.2).

**Theorem 4.** *Let  $(M^n, g, u, \lambda)$ ,  $n \geq 3$ , be an  $n$ -dimensional (compact or noncompact)  $m$ -quasi-Einstein manifold with  $m > 1$  and  $\text{Ric} \geq \frac{(n-1)}{m+n-1}\lambda$ . Consider  $\Omega \subset \text{int}(M)$  be a compact domain with connected strictly mean-convex boundary  $\partial\Omega$ . Then we have*

$$(1.16) \quad \int_{\partial\Omega} \frac{u}{H} dS \geq \frac{n}{n-1} \int_{\Omega} u d\Omega.$$

Moreover, if equality holds in (1.16), then  $\Omega$  is a geodesic ball and  $(M^n, g)$  is isometric, up to scaling, to either Example (i), or (a), or (b), or (c), or (d).

---

<sup>1</sup>The Hawking mass is often utilized as a lower bound for the Bartnik quasi-local mass (see [46, 47]).

**Remark 4.** *As we shall see later, a quasi-Einstein manifold satisfying the Ricci bound condition assumed in Theorem 4 must be a substatic Riemannian manifold. Consequently, the Heintze-Karcher type inequality may follow from Li and Xia [42]. An advantage here is that the rigidity statement for the equality case of (1.16) is established. We also observe that such a Ricci bound condition implies that  $R \geq \frac{n(n-1)}{m+n-1}\lambda$ , which holds in general for compact quasi-Einstein manifolds; see [32, Remark 5.1] and [21, Proposition 3.6]. Anyway, it should be interesting to obtain an Heintze-Karcher type inequality for compact domain on quasi-Einstein manifolds by removing the Ricci bound condition.*

The rest of this paper is organized as follows. In Section 2, we review some basic facts on  $m$ -quasi-Einstein manifolds. Moreover, we present a couple of key lemmas that will be used in the proofs of the main results. Section 3 collects the proofs of Theorems 1, 2, 3 and 4 and Corollaries 1 and 2.

## 2. BACKGROUND

In this section, we review some basic facts and a couple of lemmas that will be useful in the proof of the main results. We start by recalling that the fundamental equation of an  $m$ -quasi-Einstein manifold  $(M^n, g, u, \lambda)$  (possibly) with boundary  $\partial M$  is given by

$$(2.1) \quad \nabla^2 u = \frac{u}{m}(\text{Ric} - \lambda g),$$

where  $u > 0$  in the interior of  $M^n$  and  $u = 0$  on  $\partial M$ . In particular, taking the trace of (2.1) we arrive at

$$(2.2) \quad \Delta u = \frac{u}{m}(R - \lambda n).$$

Plugging this fact into (2.1) yields

$$(2.3) \quad u\overset{\circ}{\text{Ric}} = m\overset{\circ}{\nabla}^2 u,$$

where  $\overset{\circ}{T} = T - \frac{\text{tr}T}{n}g$  stands for the traceless part of  $T$ .

Since  $u > 0$  in the interior of  $M^n$  and  $u = 0$  on the boundary  $\partial M$ , one deduces that  $\nu = -\frac{\nabla u}{|\nabla u|}$  is the outward unit normal vector. Besides, it follows from [32, Propositions 2.2 and 2.3] that  $|\nabla u| \neq 0$  is constant along  $\partial M$ . Thereby, we set an orthonormal frame given by  $\{e_1, \dots, e_{n-1}, e_n = \nu\}$ . The second fundamental form at  $\partial M$  satisfies

$$(2.4) \quad \mathbb{I}l_{ij} = \langle \nabla_{e_i} \nu, e_j \rangle = -\frac{1}{|\nabla u|} \nabla_i \nabla_j u = 0,$$

for any  $1 \leq i, j \leq n-1$ . Consequently,  $\partial M$  is totally geodesic. By Gauss equation

$$R_{ijkl}^{\partial M} = R_{ijkl} - h_{il}h_{jk} + h_{ik}h_{jl},$$

we then infer

$$(2.5) \quad R_{ijkl}^{\partial M} = R_{ijkl}.$$

Taking the trace in (2.5), we have

$$(2.6) \quad R_{ik}^{\partial M} = R_{ik} - R_{inkn}$$

and

$$(2.7) \quad R^{\partial M} = R - 2R_{nn}.$$

We now collect some well-known equations for quasi-Einstein manifolds (cf. [21, 26, 38]).

**Proposition 1.** *Let  $(M^n, g, u, \lambda)$ ,  $n \geq 3$ , be an  $n$ -dimensional quasi-Einstein manifold. Then we have*

$$(2.8) \quad \frac{1}{2}u\nabla R = -(m-1)\text{Ric}(\nabla u) - (R - (n-1)\lambda)\nabla u;$$

$$(2.9) \quad u\Delta u + (m-1)|\nabla u|^2 + \lambda u^2 = \mu,$$

where  $\mu$  is a constant;

$$(2.10) \quad \frac{1}{2}\Delta R + \frac{m+2}{2u}\langle \nabla u, \nabla R \rangle = -\frac{m-1}{m} \left| \text{Ric} - \frac{R}{n}g \right|^2 - \frac{(n+m-1)}{nm}(R-n\lambda) \left( R - \frac{n(n-1)}{n+m-1}\lambda \right).$$

As a consequence of Proposition 1, we obtain the following lemma.

**Lemma 1.** *Let  $(M^n, g, u, \lambda)$ ,  $n \geq 3$ , be an  $n$ -dimensional quasi-Einstein manifold with boundary  $\partial M$  and  $m > 1$ . Then, at  $\partial M$ , we have:*

$$(m+1)R = (m-1)R^{\partial M} + 2(n-1)\lambda.$$

*Proof.* Since  $u$  vanishes on the boundary  $\partial M$ , we may use (2.8) to infer

$$\text{Ric}(\nabla u, \nabla u) = -\frac{(R - (n-1)\lambda)}{m-1}|\nabla u|^2$$

on  $\partial M$ . Consequently,

$$R_{nn} = -\frac{(R - (n-1)\lambda)}{m-1}.$$

Hence, by using (2.7),

$$\frac{R - R^{\partial M}}{2} = -\frac{(R - (n-1)\lambda)}{m-1}$$

and the result follows.  $\square$

It is known from [32, Remark 5.1] that the scalar curvature  $R$  of a nontrivial compact quasi-Einstein manifold  $M^n$  with boundary  $\partial M$  must satisfy

$$(2.11) \quad R \geq \frac{n(n-1)}{m+n-1}\lambda.$$

In particular, it follows from Eq. (2.10) that if equality occurs in (2.11), then  $M^n$  is necessarily Einstein. As observed in [21, Proposition 3.6], (2.11) also holds in the case of compact without boundary.

As mentioned in the Introduction, the Jacobi operator  $J$  acting in  $\varphi$  is given by

$$J(\varphi) = \Delta_{\partial M} \varphi + (R_{nn} + |\mathbb{I}\mathbb{I}|^2)\varphi,$$

for all  $\varphi \in C^\infty(\partial M)$ . Besides, we consider  $\beta_1$  to be the first eigenvalue of the Jacobi operator  $J$ , i.e.,

$$(2.12) \quad \beta_1 = \inf_{\varphi \neq 0} \frac{-\int_{\partial M} \varphi J(\varphi) dS_g}{\int_{\partial M} \varphi^2 dS_g}.$$

With aid of this notation, we get the following proposition.

**Proposition 2.** *Let  $(M^n, g, u, \lambda)$ ,  $n \geq 3$ , be a compact  $n$ -dimensional quasi-Einstein manifold with boundary  $\partial M$  and  $m > 1$ . Then we have:*

$$\beta_1 \geq -\frac{(n-1)}{m+n-1}\lambda.$$

Moreover, equality holds if and only if  $R = \frac{n(n-1)}{m+n-1}\lambda$  on  $\partial M$  and the eigenfunction associated to  $\beta_1$  is constant.

*Proof.* Firstly, on integrating by parts, we have

$$\begin{aligned} -\int_{\partial M} \varphi J(\varphi) dS_g &= -\int_{\partial M} \varphi \Delta_{\partial M} \varphi dS_g - \int_M R_{nn} \varphi^2 dS_g \\ (2.13) \qquad \qquad \qquad &= \int_{\partial M} |\nabla_{\partial M} \varphi|^2 dS_g - \int_M R_{nn} \varphi^2 dS_g. \end{aligned}$$

On the other hand, by using (2.8) on  $\partial M$  and (2.11), one obtains that

$$\begin{aligned} R_{nn} &= \frac{(n-1)\lambda - R}{m-1} \\ &\leq \frac{1}{m-1} \left( (n-1)\lambda - \frac{n(n-1)}{m+n-1}\lambda \right) \\ (2.14) \qquad \qquad \qquad &= \frac{(n-1)}{m+n-1}\lambda. \end{aligned}$$

Plugging this into (2.13) yields

$$-\int_{\partial M} \varphi J(\varphi) dS_g \geq \int_{\partial M} |\nabla_{\partial M} \varphi|^2 dS_g - \frac{(n-1)}{m+n-1}\lambda \int_{\partial M} \varphi^2 dS_g$$

for all  $\varphi \in C^\infty(\partial M)$ . In particular, choosing  $\varphi$  such that  $J(\varphi) = -\beta_1 \varphi$ , one sees that

$$\begin{aligned} \beta_1 &\geq \frac{\int_{\partial M} |\nabla_{\partial M} \varphi|^2 dS_g}{\int_{\partial M} \varphi^2 dS_g} - \frac{(n-1)}{m+n-1}\lambda \\ (2.15) \qquad \qquad \qquad &\geq -\frac{(n-1)}{m+n-1}\lambda. \end{aligned}$$

The case of equality follows directly from (2.14) and (2.15). So, the proof is completed.  $\square$

**Remark 5.** *It is easy to check from Proposition 2 that*

$$(2.16) \qquad \qquad \qquad \frac{2}{n-1}\beta_1 + \frac{n}{m+n-1}\lambda \geq \frac{(n-2)}{m+n-1}\lambda.$$

*This implies that our estimate obtained in Theorem 3 improves the estimate established in [27, Theorem 1].*

By assuming that the scalar curvature of the boundary  $R^{\partial M}$  of a quasi-Einstein manifold is constant, we also obtain the following result.

**Proposition 3.** *Let  $(M^n, g, u, \lambda)$ ,  $n \geq 3$ , be an  $n$ -dimensional quasi-Einstein manifold with boundary  $\partial M$  and  $m > 1$ . Suppose that the scalar curvature of  $\partial M$  is constant. Then*

$$\beta_1 = \frac{R - (n-1)\lambda}{m-1}.$$



*Proof.* Choosing  $\varphi \equiv 1$  in (2.12), we obtain

$$\beta_1 |\partial M| \leq - \int_{\partial M} R_{nn} dS_g$$

and by using (2.8) on  $\partial M$ , one deduces that

$$\beta_1 |\partial M| \leq \int_{\partial M} \left( \frac{R - (n-1)\lambda}{m-1} \right) dS_g.$$

By Lemma 1, we have that  $R$  is constant on  $\partial M$  and hence,

$$\beta_1 \leq \frac{R - (n-1)\lambda}{m-1}.$$

Next, we derive the reverse inequality. It follows from (2.13) and (2.8) that

$$- \int_{\partial M} \varphi J(\varphi) dS_g = \int_{\partial M} |\nabla_{\partial M} \varphi|^2 dS_g + \frac{R - (n-1)\lambda}{m-1} \int_{\partial M} \varphi^2 dS_g.$$

Now, taking  $\varphi$  such that  $J(\varphi) = -\beta_1 \varphi$ , we infer

$$\begin{aligned} \beta_1 &= \frac{\int_{\partial M} |\nabla_{\partial M} \varphi|^2 dS_g}{\int_{\partial M} \varphi^2 dS_g} + \frac{R - (n-1)\lambda}{m-1} \\ &\geq \frac{R - (n-1)\lambda}{m-1}. \end{aligned}$$

Thus, we obtain the asserted equality.  $\square$

In order to proceed, we recall a very important result so called *generalized Reilly's formula* that was established by Qiu and Xia [48].

**Proposition 4** ([48]). *Let  $(M^n, g)$  be an  $n$ -dimensional compact Riemannian manifold with boundary  $\partial M$ . Given two functions  $f$  and  $u$  on  $M$  and  $k \in \mathbb{R}$ , we have the following identity:*

$$\begin{aligned} &\int_M f ((\Delta u + knu)^2 - |\nabla^2 u + kug|^2) dV_g = (n-1)k \int_M (\Delta f + nkf) u^2 dV_g \\ &+ \int_M (\nabla^2 f - (\Delta f)g - 2(n-1)kfg + fRic) (\nabla u, \nabla u) dV_g \\ &+ \int_{\partial M} f \left[ 2 \left( \frac{\partial u}{\partial \nu} \right) \Delta_{\partial M} u + H \left( \frac{\partial u}{\partial \nu} \right)^2 + \mathbb{III}(\nabla_{\partial M} u, \nabla_{\partial M} u) + 2(n-1)k \left( \frac{\partial u}{\partial \nu} \right) u \right] dS_g \\ &+ \int_{\partial M} \frac{\partial f}{\partial \nu} (|\nabla_{\partial M} u|^2 - (n-1)ku^2) dS_g, \end{aligned}$$

where  $\mathbb{III}$  and  $H = \text{tr}(\mathbb{III})$  stand for the second fundamental form and the mean curvature of  $\partial M$ , respectively.

Notice that the classical Reilly's formula is obtained by considering  $f = 1$  and  $k = 0$  in the above expression. We refer to [39, Proposition 1] for an alternative proof of Proposition 4. It is known that the classical Reilly's formula is particularly efficient for manifolds with nonnegative Ricci curvature. Interestingly, the formula obtained in Proposition 4 is also useful for manifolds allow negative curvature. In [39], Kwong and Miao have used Proposition 4 to prove a functional inequality on the boundary of static manifolds. Similarly, Araújo, Freitas and Santos [2] used such a proposition to establish an integral inequality for the boundary of a bounded domain in a quasi-Einstein manifold. While Diógenes, Pinheiro

and Ribeiro [29] employed the generalized Reilly's formula to prove new sharp integral estimates for critical metrics of the volume functional on compact manifolds with boundary.

Now, we are going to use Proposition 4 to establish a key lemma that will be applied in the proof of Theorem 2.

**Lemma 2.** *Let  $(M^n, g, u, \lambda)$ ,  $n \geq 3$ , be a compact oriented  $m$ -quasi-Einstein manifold with connected boundary and constant scalar curvature. Then we have*

$$\frac{m+1}{m} \int_M u Ric(\nabla u, \nabla u) dV_g = - \int_M u |\nabla^2 u|^2 dV_g + \frac{(n-2)(n\lambda - R) + n\lambda}{mn} \int_M u |\nabla u|^2 dV_g.$$

*Proof.* We start by using (2.2) to infer that  $u$  must satisfy

$$\begin{cases} \Delta u + n\beta u = 0 & \text{in } M, \\ u = 0 & \text{on } \partial M, \end{cases}$$

where  $\beta = \frac{(n\lambda - R)}{mn}$ . Hence, choosing  $u = f$  and  $\kappa = \beta$ , it follows from Proposition 4 that

$$- \int_M u \left| \nabla^2 u - \frac{\Delta u}{n} g \right|^2 dV_g = \int_M \left( \nabla^2 u - (\Delta u)g + \frac{2(n-1)}{n} (\Delta u)g + u Ric \right) (\nabla u, \nabla u) dV_g,$$

where we also have used that  $u = 0$  on  $\partial M$ . Therefore, by (1.2), one obtains that

$$\begin{aligned} - \int_M u |\nabla^2 u|^2 dV_g &= \int_M \left( \frac{u}{m} (Ric - \lambda g) + \frac{(n-2)}{n} (\Delta u)g + u Ric \right) (\nabla u, \nabla u) dV_g \\ &= \frac{m+1}{m} \int_M u Ric(\nabla u, \nabla u) dV_g - \frac{\lambda}{m} \int_M u |\nabla u|^2 dV_g \\ &\quad + \frac{(n-2)}{n} \int_M (\Delta u) |\nabla u|^2 dV_g \\ &= \frac{m+1}{m} \int_M u Ric(\nabla u, \nabla u) dV_g - \frac{\lambda}{m} \int_M u |\nabla u|^2 dV_g \\ &\quad + \frac{(n-2)(R - n\lambda)}{mn} \int_M u |\nabla u|^2 dV_g, \end{aligned}$$

where in the last equality we have used again (2.2). So, the proof is completed.  $\square$

To conclude this section, we recall another *generalized Reilly's formula* obtained subsequently by Li and Xia in [42, Theorem 1.1].

**Proposition 5** ([42]). *Let  $(M^n, g)$  be an  $n$ -dimensional smooth Riemannian manifold and  $\Omega \subset M$  be a bounded domain with smooth boundary  $\partial\Omega$ . Let  $V \in C^\infty(\overline{\Omega})$  be a smooth function such that  $\frac{\nabla^2 V}{V}$  is continuous up to  $\partial\Omega$ . Then for any  $f \in C^\infty(\overline{\Omega})$ , the following integral identity holds:*

$$\begin{aligned} &\int_\Omega V \left( \left( \Delta f - \frac{\Delta V}{V} f \right)^2 - \left| \nabla^2 f - \frac{\nabla^2 V}{V} f \right|^2 \right) d\Omega \\ &= \int_{\partial\Omega} \left( V \mathbb{I}(\nabla_{\partial\Omega} f, \nabla_{\partial\Omega} f) + 2V \frac{\partial f}{\partial \nu} \Delta_{\partial\Omega} f + V H \left( \frac{\partial f}{\partial \nu} \right)^2 + \frac{\partial V}{\partial \nu} |\nabla_{\partial\Omega} f|^2 + 2f \nabla^2 V(\nabla_{\partial\Omega} f, \nu) \right) dS_g \\ &\quad + \int_{\partial\Omega} \left( -2f \frac{\partial f}{\partial \nu} \left( \Delta_{\partial\Omega} V + H \frac{\partial V}{\partial \nu} \right) - f^2 \frac{\nabla^2 V - \Delta V g}{V}(\nabla V, \nu) \right) dS_g \\ &\quad + \int_\Omega (\Delta V g - \nabla^2 V + V Ric) \left( \nabla f - \frac{\nabla V}{V} f, \nabla f - \frac{\nabla V}{V} f \right) d\Omega. \end{aligned}$$

Here,  $\nu$  is the outward unit normal vector,  $\text{III} e H = \text{tr}(\text{III})$  are the second fundamental form and the mean curvature of  $\partial\Omega$ , respectively.

Notice that the classical Reilly's formula is obtained by considering  $V = 1$ .

### 3. PROOF OF THE MAIN RESULTS

In this section, we shall present the proofs of Theorems 1, 2, 3 and 4 and Corollaries 1 and 2.

#### 3.1. Proof of Theorem 1.

*Proof.* To begin with, upon integrating (2.9) over  $M$ , we use (2.2) to infer

$$\begin{aligned}
\mu \text{Vol}(M) &= \int_M \frac{u^2}{m} (R - n\lambda) dV_g + (m-1) \int_M |\nabla u|^2 dV_g + \lambda \int_M u^2 dV_g \\
&\geq -\frac{n\lambda}{m+n-1} \int_M u^2 dV_g + (m-1) \int_M |\nabla u|^2 dV_g + \lambda \int_M u^2 dV_g \\
(3.1) \quad &= \frac{m-1}{m+n-1} \lambda \int_M u^2 dV_g + (m-1) \int_M |\nabla u|^2 dV_g,
\end{aligned}$$

where in the second line we have used the estimate (2.11). Next, from the Rayleigh-quotient characterization of the first nonzero eigenvalue of the Laplacian  $\lambda_1$ , one sees that

$$\int_M |\nabla u|^2 dV_g \geq \lambda_1 \int_M u^2 dV_g,$$

where

$$\lambda_1 = \inf_{H_0^{1,2}(M), u \neq 0} \frac{\int_M |\nabla u|^2 dV_g}{\int_M u^2 dV_g}.$$

Plugging this fact into (3.1) yields

$$\mu \text{Vol}(M) \geq \frac{m-1}{m+n-1} (\lambda_1(m+n-1) + \lambda) \int_M u^2 dV_g.$$

By Holder's inequality, one obtains that

$$(3.2) \quad \mu \text{Vol}(M)^2 \geq \frac{(m-1)}{(m+n-1)} (\lambda_1(m+n-1) + \lambda) \left( \int_M u dV_g \right)^2,$$

On the other hand, combining (2.2) and (2.11), we deduce

$$(3.3) \quad \Delta u \geq -\frac{n\lambda}{m+n-1} u,$$

and by Stokes' theorem, one sees that

$$\int_M u dV_g \geq \frac{m+n-1}{n\lambda} \int_M (-\Delta u) dV_g = \frac{m+n-1}{n\lambda} |\nabla u|_{|\partial M}^2 |\partial M|.$$

Substituting this into (3.2), we arrive at

$$\mu \text{Vol}(M)^2 \geq (m-1) \frac{(\lambda_1(m+n-1) + \lambda)(m+n-1)}{n^2 \lambda^2} |\nabla u|_{|\partial M}^2 |\partial M|^2.$$

Next, since  $u|_{\partial M} = 0$ , by evaluating (2.9) on  $\partial M$ , one sees that  $\mu = (m-1) |\nabla u|_{|\partial M}^2$ . Therefore, rearranging terms, we have

$$(3.4) \quad |\partial M| \leq \frac{n\lambda}{\sqrt{(\lambda_1(m+n-1) + \lambda)(m+n-1)}} \text{Vol}(M),$$

which proves (1.7). Moreover, if equality holds in (3.4), then

$$R = \frac{n(n-1)}{m+n-1} \lambda$$

Hence, it follows from (2.10) that  $|\mathring{Ric}| = 0$  on  $M$ , i.e.,  $M^n$  is Einstein and in this case, it suffices to apply Proposition 3.1 of [32] to conclude that  $M^n$  is isometric to the standard hemisphere  $\mathbb{S}_+^n$ . This finishes the proof of the theorem.  $\square$

### 3.2. Proof of Theorem 2.

*Proof.* Initially, by the classical Böchner's formula, we have

$$u\Delta|\nabla u|^2 = 2u\text{Ric}(\nabla u, \nabla u) + 2u\langle \nabla(\Delta u), \nabla u \rangle + 2u|\nabla^2 u|^2.$$

Upon integrating this expression over  $M$ , we use (2.2) and the fact that  $M$  has constant scalar curvature in order to infer

$$(3.5) \quad \begin{aligned} \int_M u\Delta|\nabla u|^2 dV_g &= 2 \int_M u\text{Ric}(\nabla u, \nabla u) dV_g - \frac{2\alpha}{m} \int_M u|\nabla u|^2 dV_g + 2 \int_M u|\nabla^2 u|^2 dV_g \\ &= 2 \int_M u\text{Ric}(\nabla u, \nabla u) dV_g - \frac{2\alpha}{m} \int_M u|\nabla u|^2 dV_g + 2 \int_M u|\mathring{\nabla}^2 u|^2 dV_g \\ &\quad + \frac{2}{n} \int_M u(\Delta u)^2 dV_g, \end{aligned}$$

where  $\mathring{\nabla}^2 u = \nabla^2 u - \frac{\Delta u}{n}g$  and  $\alpha = n\lambda - R$ .

On the other hand, by using the Stokes' theorem and the fact that  $u = 0$  on  $\partial M$ , one sees that

$$\begin{aligned} \int_M u\Delta|\nabla u|^2 dV_g &= \int_M |\nabla u|^2 \Delta u dV_g \\ &\quad + \int_{\partial M} \left( u \left\langle \nabla|\nabla u|^2, -\frac{\nabla u}{|\nabla u|} \right\rangle - |\nabla u|^2 \left\langle \nabla u, -\frac{\nabla u}{|\nabla u|} \right\rangle \right) dS_g \\ &= -\frac{\alpha}{m} \int_M u|\nabla u|^2 dV_g + |\nabla u|_{\partial M}^3 |\partial M|, \end{aligned}$$

which compared with (3.5) and using once more (2.2) yields

$$\begin{aligned} |\nabla u|_{\partial M}^3 |\partial M| &= 2 \int_M u\text{Ric}(\nabla u, \nabla u) dV_g - \frac{\alpha}{m} \int_M u|\nabla u|^2 dV_g \\ &\quad + 2 \int_M u|\mathring{\nabla}^2 u|^2 dV_g + \frac{2}{n} \int_M u(\Delta u)^2 dV_g \\ &= 2 \int_M u\text{Ric}(\nabla u, \nabla u) dV_g - \frac{\alpha}{m} \int_M u|\nabla u|^2 dV_g \\ &\quad + 2 \int_M u|\mathring{\nabla}^2 u|^2 dV_g + \frac{2\alpha^2}{nm^2} \int_M u^3 dV_g. \end{aligned}$$

Now, we may invoke Lemma 2 to infer

$$(3.6) \quad \begin{aligned} |\nabla u|_{\partial M}^3 |\partial M| &= \frac{2}{m+1} \int_M u|\mathring{\nabla}^2 u|^2 dV_g + \frac{2\alpha^2}{m^2 n} \int_M u^3 dV_g \\ &\quad + \frac{2mn\lambda + \alpha(n(m-1) - 4m)}{n(m+1)m} \int_M u|\nabla u|^2 dV_g. \end{aligned}$$

At the same time, by (2.2), Stokes' theorem and the fact that  $u|_{\partial M} = 0$ , we have

$$\frac{\alpha}{m} \int_M u^3 dV_g = - \int_M u^2 (\Delta u) dV_g = \int_M \langle \nabla u^2, \nabla u \rangle dV_g = 2 \int_M u |\nabla u|^2 dV_g.$$

Substituting this into (3.6) gives

$$|\nabla u|_{\partial M}^3 |\partial M| = \frac{2}{m+1} \int_M u |\mathring{\nabla}^2 u|^2 dV_g + \frac{(2nm\lambda + \alpha(n(m-1) + 4))\alpha}{2nm^2(m+1)} \int_M u^3 dV_g,$$

so that,

$$(3.7) \quad |\nabla u|_{\partial M}^3 |\partial M| \geq \frac{(2nm\lambda + \alpha(n(m-1) + 4))\alpha}{2nm^2(m+1)} \int_M u^3 dV_g.$$

Proceeding, upon integrating (2.9) over  $M^n$ , we use once more the Stokes' theorem and (2.2) to infer

$$\begin{aligned} \mu Vol(M) &= \int_M u \Delta u dV_g + (m-1) \int_M |\nabla u|^2 dV_g + \lambda \int_M u^2 dV_g \\ &= -(m-2) \int_M u \Delta u dV_g + \lambda \int_M u^2 dV_g \\ (3.8) \quad &= \left( \frac{(m-2)\alpha + \lambda m}{m} \right) \int_M u^2 dV_g. \end{aligned}$$

Besides, by Holder's inequality and (2.2), one sees that

$$\begin{aligned} \int_M u^2 dV_g &= \int_M u^{\frac{3}{2}} u^{\frac{1}{2}} dV_g \\ &\leq \left( \int_M u^3 dV_g \right)^{\frac{1}{2}} \left( \int_M u dV_g \right)^{\frac{1}{2}} \\ &= \left( \int_M u^3 dV_g \right)^{\frac{1}{2}} \left( \frac{m}{\alpha} \int_M (-\Delta u) dV_g \right)^{\frac{1}{2}} \\ (3.9) \quad &= \left( \int_M u^3 dV_g \right)^{\frac{1}{2}} \left( \frac{m |\nabla u|_{\partial M} |\partial M|}{\alpha} \right)^{\frac{1}{2}}, \end{aligned}$$

where we have used that  $\alpha = n\lambda - R$  is positive, which follows from the fact that  $M^n$  has constant scalar curvature (see Remark 2, or [32, Corollary 4.3]). Next, we already know from the proof of Theorem 1 that  $\mu = (m-1) |\nabla u|_{\partial M}^2$  and hence, plugging (3.9) into (3.8), one concludes that

$$(3.10) \quad Vol(M) \leq \left( \frac{(m-2)\alpha + \lambda m}{(m-1)\sqrt{m\alpha} |\nabla u|_{\partial M}^{\frac{3}{2}}} \right) |\partial M|^{\frac{1}{2}} \left( \int_M u^3 dV_g \right)^{\frac{1}{2}}$$

Whence, it suffices to invoke (3.7) to achieve

$$(3.11) \quad Vol(M) \leq \left( \frac{(m-2)\alpha + \lambda m}{(m-1)\alpha} \right) \sqrt{\frac{2n(m+1)m}{2\lambda mn + \alpha[n(m-1) + 4]}} |\partial M|$$

which proves the stated inequality.

Finally, if equality holds in (3.11), then (3.7) also becomes an equality. Of which, it follows that  $|\mathring{\nabla}^2 u|^2 = 0$  and by using (2.3), one sees that  $M^n$  is an Einstein manifold. Thereby, we are in position to apply Proposition 3.1 of [32] to conclude that  $M^n$  is isometric, up to scaling, to the standard hemisphere  $\mathbb{S}_+^n$ . So, the proof is completed.  $\square$

### 3.3. Proof of Corollary 1.

*Proof.* To begin with, one observes that

$$\begin{aligned} |\partial M|^n &= |\partial M| |\partial M|^{n-1}. \\ &= \frac{\alpha}{m |\nabla u|_{|\partial M}} \left( \int_M u dV_g \right) |\partial M|^{n-1}, \end{aligned}$$

where we have used (2.2) and the Stokes' theorem. Consequently, by Theorem 2, one sees that

$$(3.12) \quad |\partial M| \geq \Lambda(\alpha, m, n, u)^{\frac{1}{n}} \text{Vol}(M)^{\frac{n-1}{n}},$$

where

$$\Lambda(\alpha, m, n, u) = \frac{\alpha}{m |\nabla u|_{|\partial M}} \left( \frac{(m-1)\alpha}{(m-2)\alpha + \lambda m} \sqrt{\frac{2\lambda mn + (\alpha(n(m-1) + 4))}{2n(m+1)m}} \right)^{n-1} \int_M u dV_g,$$

as asserted. Furthermore, the case of equality follows directly from Theorem 2. Thus, we finish the proof of the corollary.  $\square$

### 3.4. Proof of Theorem 3.

*Proof.* We start by claiming that  $R^{\partial M} > 0$ . Indeed, it follows from Lemma 1 and (2.11) that

$$\begin{aligned} (m-1)R^{\partial M} &= (m+1)R - 2(n-1)\lambda \\ &\geq \frac{(m+1)n(n-1)}{m+n-1} \lambda - 2(n-1)\lambda \\ &= \frac{(m-1)(n-1)(n-2)}{m+n-1} \lambda. \end{aligned}$$

Since  $m > 1$  and  $M^n$  is compact, we then obtain

$$(3.13) \quad R^{\partial M} \geq \frac{(n-1)(n-2)}{m+n-1} \lambda > 0,$$

as claimed.

From now on, we adapt the arguments by Barros-Silva [8]. Firstly, since  $\partial M$  is totally geodesic, we obtain from (1.10) that

$$\beta_1 \int_{\partial M} \varphi^2 dS_g \leq - \int_{\partial M} \varphi J(\varphi) dS_g = \int_{\partial M} |\nabla^{\partial M} \varphi|^2 dS_g - \int_{\partial M} R_{nn} \varphi^2 dS_g$$

for any  $\varphi \in C^\infty(\partial M)$ . Thereby, choosing  $\varphi \equiv 1$ , one sees that

$$(3.14) \quad \begin{aligned} \beta_1 |\partial M| &\leq - \int_{\partial M} R_{nn} dS_g = -\frac{1}{2} \int_{\partial M} (R - R^{\partial M}) dS_g \\ &\leq \frac{1}{2} \int_{\partial M} R^{\partial M} dS_g - \frac{n(n-1)}{2(m+n-1)} \lambda |\partial M| \end{aligned}$$

where we have used (2.7) and (2.11).

On the other hand, since  $\partial M$  is Einstein and  $R^{\partial M} > 0$ , we may write

$$\text{Ric}^{\partial M} = \frac{R^{\partial M}}{n-1} = (n-2)\varepsilon,$$

where  $\varepsilon = \frac{R^{\partial M}}{(n-1)(n-2)}$ . Whence, it follows from Bonnet-Myer's theorem that  $\text{diam}_{g_{\partial M}}(\partial M) \leq \frac{\pi}{\sqrt{\varepsilon}}$ . Furthermore, by Bishop-Gromov's theorem, we have

$$(3.15) \quad \text{Vol} \left( B_{\frac{\pi}{\sqrt{\varepsilon}}}^{\partial M} \right) (p) \leq \text{Vol} (\mathbb{S}_{g_\varepsilon}^{n-1}) = \varepsilon^{-\frac{(n-1)}{2}} \omega_{n-1},$$

for any point  $p \in \partial M$ , where  $g_\varepsilon = \frac{1}{\varepsilon} g_{\mathbb{S}^{n-1}}$  and  $\omega_{n-1}$  denotes the volume of the standard unit sphere  $\mathbb{S}^{n-1}$ . These facts together yield

$$|\partial M| \leq \text{Vol} \left( B_{\frac{\pi}{\sqrt{\varepsilon}}}^{\partial M} \right) \leq \varepsilon^{-\frac{(n-1)}{2}} \omega_{n-1}.$$

so that

$$(3.16) \quad R^{\partial M} \leq (n-1)(n-2) (\omega_{n-1})^{\frac{2}{n-1}} |\partial M|^{-\frac{2}{n-1}}.$$

This substituted into (3.14) gives

$$2\beta_1 \leq (n-1)(n-2) (\omega_{n-1})^{\frac{2}{n-1}} |\partial M|^{-\frac{2}{n-1}} - \frac{n(n-1)}{m+n-1} \lambda.$$

Consequently,

$$(3.17) \quad \left( \frac{2}{n-1} \beta_1 + \frac{n}{m+n-1} \lambda \right) |\partial M|^{\frac{2}{n-1}} \leq (n-2) (\omega_{n-1})^{\frac{2}{n-1}},$$

Now, it suffices to use Remark 5 to conclude that  $\frac{2}{n-1} \beta_1 + \frac{n}{m+n-1} \lambda > 0$  and hence, the stated inequality follows.

Finally, if equality holds in (3.17), then (3.15) also becomes an equality. Thus, it follows from the equality case for the Bishop-Gromov's theorem that the boundary is isometric to a round sphere  $\mathbb{S}^{n-1}$ . This finishes the proof of the theorem.  $\square$

### 3.5. Proof of Corollary 2.

*Proof.* Firstly, we invoke the Gauss-Bonnet formula to deduce

$$\frac{1}{2} \int_{\partial M} R^{\partial M} dS_g = \int_{\partial M} K^{\partial M} dS_g = 2\pi \chi(\partial M).$$

So, it suffices to use (3.13) to conclude that  $\chi(\partial M) > 0$  and  $\partial M$  is a 2-sphere. Next, since  $\partial M$  is Einstein, we may combine (3.16) and (3.13) in order to infer

$$(3.18) \quad |\partial M| \leq \frac{m+2}{\lambda} 4\pi.$$

On the other hand, by (2.11),  $\chi(\partial M) = 2$  and the fact that  $\partial M$  is totally geodesic, we see that

$$(3.19) \quad \begin{aligned} \mathfrak{m}_H(\partial M) &= \frac{|\partial M|^{\frac{1}{2}}}{(16\pi)^{\frac{3}{2}}} \left( 16\pi - \frac{2}{3} \Lambda |\partial M| \right) \\ &\leq \frac{|\partial M|^{\frac{1}{2}}}{(16\pi)^{\frac{1}{2}}} - \frac{4}{m+2} \lambda \left( \frac{|\partial M|}{16\pi} \right)^{\frac{3}{2}} \\ &\leq \frac{1}{2} \sqrt{\frac{(m+2)}{\lambda}} - \frac{4}{m+2} \lambda \left( \frac{|\partial M|}{16\pi} \right)^{\frac{3}{2}}, \end{aligned}$$

where in the last inequality we have used (3.18). This therefore gives the first stated inequality. Moreover, if equality holds in (3.19), it suffices to invoke the equality case of Theorem 3.

We now deal with the second inequality. It follows from the Hawking mass definition and the fact that  $\chi(\partial M) = 2$  that

$$(3.20) \quad \mathfrak{m}_H(\partial M) = \frac{4|\partial M|^{\frac{1}{2}}}{(16\pi)^{\frac{3}{2}}} \left( 4\pi - \frac{\Lambda}{6}|\partial M| \right).$$

At the same time, choosing  $\varphi \equiv 1$  in (1.10), we then use (2.7) and the Gauss-Bonnet formula in order to deduce

$$\begin{aligned} \beta_1|\partial M| &\leq - \int_{\partial M} R_{nn} dS_g \\ &= \int_{\partial M} K^{\partial M} dS_g - \frac{1}{2} \int_{\partial M} R dS_g \\ &= 4\pi - \frac{1}{2} \int_{\partial M} R dS_g \\ &\leq 4\pi - \frac{\Lambda}{2}|\partial M|. \end{aligned}$$

This substituted into (3.20) yields

$$\mathfrak{m}_H(\partial M) \geq 4 \left( \beta_1 + \frac{\Lambda}{3} \right) \left( \frac{|\partial M|}{16\pi} \right)^{\frac{3}{2}}.$$

Now, we use (2.11) to infer  $\Lambda \geq \frac{6}{m+2}\lambda$ . Consequently,

$$\mathfrak{m}_H(\partial M) \geq 4 \left( \beta_1 + \frac{2\lambda}{m+2} \right) \left( \frac{|\partial M|}{16\pi} \right)^{\frac{3}{2}} \geq 0,$$

where we have used Proposition 2 to conclude that  $\beta_1 + \frac{2\lambda}{m+2} \geq 0$ . Thus, the proof is finished.  $\square$

### 3.6. Proof of Theorem 4.

*Proof.* Initially, we consider the auxiliar tensor

$$P = Ric - \frac{(n-1)\lambda - R}{m-1}g.$$

With aid of this notation, we see from (2.1) and (2.2) that

$$\begin{aligned} (\Delta u)g - \nabla^2 u + uRic &= \frac{u}{m} [(R - (n-1)\lambda)g + (m-1)Ric] \\ &= \frac{(m-1)}{m} uP. \end{aligned}$$

In particular, since  $m > 1$ ,  $M^n$  is a substatic manifold if and only if  $P \geq 0$ .

On the other hand, it is not hard to check that the Ricci bound assumption implies

$$\begin{aligned} P &\geq \left( \frac{(n-1)\lambda}{m+n-1} - \frac{((n-1)\lambda - R)}{m-1} \right) g \\ &= \frac{R(m+n-1) - n(n-1)\lambda}{(m-1)(m+n-1)} g. \end{aligned}$$

By tracing the assumption on Ricci, one sees that  $R(m+n-1) \geq n(n-1)\lambda$ . Consequently,  $P \geq 0$  and hence,  $M^n$  is a substatic Riemannian manifold. Thus, the Heintze–Karcher type inequality follows from Theorem 1.3 in [42].



At the same time, we are going to present here another proof of the asserted inequality in order to discuss the equality case. To do so, we divide the proof into two parts and adapt the arguments by Li-Xia [42]. We start by considering the following boundary value problem

$$(3.21) \quad \begin{cases} \Delta f - \frac{\Delta u}{u} f = 1 & \text{in } \Omega, \\ f = 0 & \text{on } \partial\Omega. \end{cases}$$

Notice that by [42, Lemma 2.5], the first eigenvalue  $\lambda_1(\Delta - q, \Omega) > 0$ , where  $q := \frac{\Delta u}{u}$ . Consequently, by the standard elliptic PDE theory (see [11]), (3.21) admits a unique smooth solution  $f \in C^\infty(\bar{\Omega})$ . Thereby, using Proposition 5 for such a solution and choosing  $V = u$ , we see that

$$(3.22) \quad \begin{aligned} \int_{\Omega} u d\Omega &= \int_{\Omega} u \left| \nabla^2 f - \frac{\nabla^2 u}{u} f \right|^2 d\Omega + \int_{\partial\Omega} u H \left( \frac{\partial f}{\partial \nu} \right)^2 dS_g \\ &+ \int_{\Omega} (\Delta u g - \nabla^2 u + u Ric) \left( \nabla f - \frac{\nabla u}{u} f, \nabla f - \frac{\nabla u}{u} f \right) d\Omega. \end{aligned}$$

By using the fact that

$$(3.23) \quad \left| \nabla^2 f - \frac{\nabla^2 u}{u} f \right|^2 \geq \frac{1}{n} \left| \Delta f - \frac{\Delta u}{u} f \right|^2 = \frac{1}{n},$$

we then obtain

$$(3.24) \quad \begin{aligned} \frac{n-1}{n} \int_{\Omega} u d\Omega &\geq \int_{\partial\Omega} u H \left( \frac{\partial f}{\partial \nu} \right)^2 dS_g \\ &+ \int_{\Omega} (\Delta u g - \nabla^2 u + u Ric) \left( \nabla f - \frac{\nabla u}{u} f, \nabla f - \frac{\nabla u}{u} f \right) d\Omega. \end{aligned}$$

On the other hand, our assumption on Ricci curvature guarantees that  $R \geq \frac{n(n-1)}{m+n-1} \lambda$ . This jointly with (2.2) and the fact that  $u > 0$  in  $\Omega$  yields

$$(3.25) \quad \Delta u \geq -\frac{n\lambda}{m+n-1} u$$

in  $\Omega$ , which combined with (1.2) gives

$$\Delta u g - \nabla^2 u + u Ric \geq -\frac{n\lambda}{m+n-1} u g + \frac{m-1}{m} u Ric + \frac{\lambda}{m} u g.$$

Plugging this into (3.24), we see that

$$(3.26) \quad \begin{aligned} \frac{n-1}{n} \int_{\Omega} u d\Omega &\geq \int_{\partial\Omega} u H \left( \frac{\partial f}{\partial \nu} \right)^2 dS_g \\ &+ \int_M \frac{(m-1)}{m} u \left( Ric - \frac{(n-1)}{m+n-1} \lambda \right) \left( \nabla f - \frac{\nabla u}{u} f, \nabla f - \frac{\nabla u}{u} f \right) d\Omega \\ &\geq \int_{\partial\Omega} u H \left( \frac{\partial f}{\partial \nu} \right)^2 dS_g, \end{aligned}$$

where in the last line we used again the assumption  $Ric \geq \frac{n-1}{m+n-1} \lambda$  and  $m > 1$ .

Next, it follows from (3.21) that  $u\Delta f - (\Delta u)f = u$  in  $\Omega$ . Thus, by Green's identity, one obtains that

$$\begin{aligned}\int_{\Omega} u d\Omega &= \int_{\Omega} u\Delta f d\Omega - \int_{\Omega} (\Delta u)f d\Omega \\ &= \int_{\partial\Omega} u \frac{\partial f}{\partial\nu} dS_g - \int_{\partial\Omega} f \frac{\partial u}{\partial\nu} dS_g \\ &= \int_{\partial\Omega} u \frac{\partial f}{\partial\nu} dS_g,\end{aligned}$$

where in the last line we used that  $f|_{\partial\Omega} = 0$ . By Hölder's inequality and (3.26), we have

$$\begin{aligned}\left(\int_{\Omega} u d\Omega\right)^2 &= \left(\int_{\partial\Omega} u \frac{\partial f}{\partial\nu} dS_g\right)^2 \leq \int_{\partial\Omega} u H \left(\frac{\partial f}{\partial\nu}\right)^2 dS_g \int_{\partial\Omega} \frac{u}{H} dS_g \\ &\leq \frac{n-1}{n} \int_{\Omega} u d\Omega \int_{\partial\Omega} \frac{u}{H} dS_g,\end{aligned}$$

so that

$$n \int_{\Omega} u d\Omega \leq (n-1) \int_{\partial\Omega} \frac{u}{H} dS_g.$$

which proves the asserted inequality.

Now, we deal with the equality case. Indeed, if equality holds, one deduces from (3.23) and (3.25) that

$$(3.27) \quad \nabla^2 f - \frac{\nabla^2 u}{u} f = \frac{1}{n} g$$

and

$$R = \frac{n(n-1)}{m+n-1} \lambda$$

in  $\Omega$ . Since  $u$  and  $g$  are real analytic in harmonic coordinates (cf. Proposition 2.4 in [32]), we conclude that

$$R = \frac{n(n-1)}{m+n-1} \lambda$$

in  $M$ . This into (2.10) therefore implies that  $(M^n, g)$  is an Einstein manifold.

We now have two cases to be analyzed, namely,  $\lambda = 0$  and  $\lambda \neq 0$ . Firstly, if  $\lambda = 0$ , it suffices to invoke Proposition 3.1 of [32] to conclude that  $M^n$  is isometric to  $[0, \infty) \times F$ , where  $F$  is Ricci flat. In particular, we have  $\nabla^2 u = \frac{u}{m}(Ric - \lambda g) = 0$  in  $M$ , which compared with (3.27), gives  $\nabla^2 f = \frac{1}{n}g$  in  $\Omega$ . Taking into account that  $f|_{\partial\Omega} = 0$ , it suffices to apply Lemma 3 of [49] to conclude that  $\Omega$  is a geodesic ball.

On the other hand, if  $\lambda \neq 0$ , then we may use again Proposition 3.1 of [32] to infer that  $M^n$  is isometric to either  $\mathbb{S}_+^n$ , or  $\mathbb{H}^n$ , or  $[0, \infty) \times N$ , or  $\mathbb{R} \times F$ . In particular, one sees that

$$\frac{\nabla^2 u}{u} = \frac{1}{m}(Ric_g - \lambda g) = -\frac{\lambda}{m+n-1}g$$

and by (3.27), we have

$$\nabla^2 f = -\frac{\lambda}{m+n-1}fg + \frac{1}{n}g.$$

By considering  $F := f - K$ , where  $K = \frac{m+n-1}{n\lambda}$ , it is easy to check that

$$\nabla^2 F = -\frac{\lambda}{m+n-1}Fg$$

in  $\Omega$ . Moreover,  $F|_{\partial\Omega}$  is constant. Therefore, it suffices to apply Theorem B of [49] to conclude that  $\Omega$  is a geodesic ball of constant sectional curvature  $\frac{\lambda}{m+n-1}$ . So, the proof is completed.  $\square$

**Acknowledgement.** *The authors want to thank B. Leandro for his interest in this work and for pointing out the preprint [40]. Moreover, they want to thank A. Barros and N. Pinheiro for helpful conversations about this subject.*

#### REFERENCES

- [1] Ambrozio, L.: *On static three-manifolds with positive scalar curvature*. J. Diff. Geom. 7 (2017) 1–45.
- [2] Araújo, M., Freitas, A. and Santos, M.: *About an integral inequality and rigidity of  $m$ -quasi-Einstein manifolds*. Lett. Math. Phys. 113 (2023) 115.
- [3] Bahuaud, E., Gunasekaran, S., Kunduri, K. and Woolgar, E.: *Static near-horizon geometries and rigidity of quasi-Einstein manifolds*. Lett. Math. Phys. 112 (2022) 116.
- [4] Bahuaud, E., Gunasekaran, S., Kunduri, K. and Woolgar, E.: *Rigidity of quasi-Einstein metrics: The incompressible case*. Lett. Math. Phys. 114 (2024) 8.
- [5] Bakry, D. and Émery, M.: *Diffusions Hypercontractives*. Séminaire de probabilités, XIX, 1983/84. Lecture Notes in Math., vol. 1123, Springer, Berlin, (1985) 177–206.
- [6] Baltazar, H., Diógenes, R. and Ribeiro Jr., E.: *Isoperimetric inequality and Weitzenböck type formula for critical metrics of the volume*. Israel J. Math. 234 (2019) 309–329.
- [7] Barbosa, E., Lima, L. and Freitas, A.: *The generalized Pohozaev-Schoen identity and some geometric applications*. Commun. Anal. Geom. 28 (2020) 223–242.
- [8] Barros, A. and da Silva, A.: *Rigidity for critical metrics of the volume functional*. Math. Nachr. 291 (2019) 709–719.
- [9] Barros, A., Batista, R., and Cruz, T.: *Hawking mass and local rigidity of minimal surfaces in three-manifolds*. Commun. Anal. Geom. 25 (2017) 1–23.
- [10] Batista, R., Diógenes, R., Ranieri, M. and Ribeiro Jr., E.: *Critical metrics of the volume functional on compact three-manifolds with smooth boundary*. J. Geom. Analysis. 27 (2017) 1530–1547.
- [11] Berestycki, H., Nirenberg, L. and Varadhan, S.: *The principal eigenvalue and maximum principle for second-order elliptic operators in general domains*. Commun. Pure Appl. Math. 47 (1994) 47–92.
- [12] Besse, A. L.: *Einstein Manifolds*. Springer-Verlag, Berlin (1987).
- [13] Böhm, C.: *Inhomogeneous Einstein metrics on low-dimensional spheres and other low-dimensional spaces*. Invent. Math. 134 (1) (1998) 145–176.
- [14] Böhm, C.: *Non-compact cohomogeneity one Einstein manifolds*. Bull. Soc. Math. France. 127 (1999) 135–177.
- [15] Borghini, S. and Mazzieri, L.: *On the mass of static metrics with positive cosmological constant: II*. Commun. Math. Phys. 377 (2020) 2079–2158.
- [16] Borghini, S., Fogagnolo, M. and Pinamonti, A.: *The equality case in the substatic Heintze-Karcher inequality*. arXiv:2307.04253 [math.DG] (2023).
- [17] Boucher, W., Gibbons, G. and Horowitz, G.: *Uniqueness theorem for anti-de Sitter spacetime*. Phys. Rev. D. (3) 30 (1984) 2447–2451.
- [18] Brendle, S.: *Constant mean curvature surfaces in warped product manifolds*. Publ. Math. Inst. Hautes Études Sci., 117 (2013) 247–269.
- [19] Case, J.: *The nonexistence of quasi-Einstein metrics*. Pacific J. Math. 248 (2010), no.2, 277–284.
- [20] Case, J.: *Smooth metric measure spaces, quasi-Einstein metrics, and tractors*. Cent. Eur. J. Math. 10 (2012), no.5, 1733–1762.
- [21] Case, J., Shu, Y. and Wei, G.: *Rigidity of quasi-Einstein Metrics*. Diff. Geom. Appl. 29 (2011), no.1, 93–100.
- [22] Catino, G., Mantegazza, C., Mazzieri, L. and Rimoldi, M.: *Locally conformmally flat quasi-Einstein manifolds*. J. Reine Angew. Math. (Crelle’s Journal), 675 (2013), 181–189.
- [23] Costa, J., Diógenes, R., Pinheiro, N. and Ribeiro Jr, E.: *Geometry of static perfect space time*. Class. Quantum Grav. 40 (2023) 205012.
- [24] Costa, J., Ribeiro Jr, E. and Zhou, D.: *Rigidity of compact quasi-Einstein manifolds with boundary*. arXiv:2401.16929 [math.DG] (2024).
- [25] Corvino, J.: Eichmair, M. and Miao, P.: *Deformation of scalar curvature and volume*. Math. Ann. 357 (2013) 551–584.
- [26] Diógenes, R. and Gadelha, T.: *Compact quasi-Einstein manifolds with boundary*. Math. Nachr. 295 (2022) 1690–1708.
- [27] Diógenes, R., Gadelha, T. and Ribeiro Jr, E.: *Geometry of compact quasi-Einstein manifolds*. Manuscripta Math. 169 (2022) 167–183.

- [28] Diógenes, R., Gadelha, T. and Ribeiro Jr, E.: *Remarks on quasi-Einstein manifolds with boundary*. Proc. Amer. Math. Soc. 150, (2022) 351–363.
- [29] Diógenes, R., Pinheiro, N. and Ribeiro Jr, E.: *Integral and boundary estimates for critical metrics of the volume functional*. arXiv:2207.12344 [math.DG] (2022), to appear in Israel J. Math.
- [30] Fang, Y. and Yuan, W.: *Brown-York mass and positive scalar curvature II: Besse’s conjecture and related problems*. Ann. Glob. Anal. Geom. 56 (2019) 1–15.
- [31] Hawking, S.W.: *Gravitational radiation in an expanding universe*. J. Math. Phys. 9 (1968) 598–604.
- [32] He, C., Petersen, P. and Wylie, W.: *On the classification of warped product Einstein metrics*. Comm. Anal. Geom. 20 (2012), no.2, 71–311.
- [33] He, C., Petersen, P. and Wylie, W.: *Warped product Einstein metrics over spaces with constant scalar curvature*. Asian J. Math. 18 (2014), no. 1, 159–189.
- [34] Heintze, E. and Karcher, H.: *A general comparison theorem with applications to volume estimates for submanifolds*. Ann. Sci. École Norm. Sup. (4) 11.4 (1978) 51–470.
- [35] Hijazi, O., Montiel, S. and Raulot, S.: *Uniqueness of the de Sitter spacetime among static vacua with positive cosmological constant*. Ann. Glob. Anal. Geom. 47 (2014) 167–178.
- [36] Huiskens, G. and Ilmanen, T.: *The inverse mean curvature flow and the Riemannian Penrose inequality*. J. Diff. Geom. 59 (2001), no. 3, 353–437.
- [37] Jia, X., Wang, G., Xia, C. and Zhang, X.: *Heintze-karcher inequality and capillary hypersurfaces in a wedge*. arXiv:2209.13839, to appear in Ann. Sc. Norm. Super. Pisa Cl. Sci.
- [38] Kim, D.-S. and Kim, Y. H.: *Compact Einstein warped product spaces with nonpositive scalar curvature*. Proc. Amer. Math. Soc. 131 (2003) 2573–2576.
- [39] Kwong, K. and Miao, P.: *A functional inequality on the boundary of static manifolds*. Asian J. Math. (4) 21 (2017) 687–696.
- [40] Leandro, B. and Sabo, G.: *Sharp lower bound for the charged Hawking mass in the electrostatic space*. To appear in Math. Research Lett. (2024).
- [41] Li, J. and Xia, C.: *An integral formula for affine connections*. J. Geom. Anal. 27 (2017) 2539–2556.
- [42] Li, J. and Xia, C.: *An integral formula and its applications on sub-static manifolds*. J. Diff. Geom. 113(3) (2019) 493–518.
- [43] Lü, H., Page, D. and Pope, C.: *New inhomogeneous Einstein metrics on sphere bundles over Einstein-Kähler manifolds*. Phys. Lett. B 593 (1-4) (2004) 218–226.
- [44] Maz’ya, M.: *Lectures on isoperimetric and isocapacitary Inequalities in the theory of Sobolev spaces*. Contemp. Math. 338 (2003) 307–340.
- [45] Miao, P. and Tam, L.-F.: *On the volume functional of compact manifolds with boundary with constant scalar curvature*. Calc. Var. PDE. 36 (2009) 141–171.
- [46] Miao, P., Wang, Y and Xie, N.: *On Hawking mass and Bartnik mass of CMC surfaces*. Math. Research Lett. 27 (2020) 855–885.
- [47] Mondino, A. and Templeton-Browne, A.: *Some rigidity results for the Hawking mass and a lower bound for the Bartnik capacity*. J. Lond. Math. Soc. (2) 106 (2022) 1844–1896.
- [48] Qiu, G. and Xia, C.: *A generalization of Reilly’s formula and its applications to a new Heintze-Karcher type inequality*. Int. Math. Res. Not. IMRN (17) (2015) 7608–7619.
- [49] Reilly, R. C.: *Geometric applications of the solvability of Neumann problems on a Riemannian manifold*. Archive for Rational Mech. Anal. (1) 75 (1980) 23–29.
- [50] Ribeiro Jr, E. and Tenenblat, K.: *Noncompact quasi-Einstein manifolds conformal to a Euclidean space*. Math. Nachr. 294 (2021) 132–144.
- [51] Rimoldi, M.: *A remark on Einstein warped products*. Pacific J. Math. 252 (2011) 207–218.
- [52] Ros, A.: *The isoperimetric problem*. Global theory of minimal surfaces. MSRI. 2 (2001) 175–209.
- [53] Ros, A.: *Compact hypersurfaces with constant higher order mean curvatures*. Rev. Mat. Iberoamericana 3. 3-4 (1987) 447–453.
- [54] Shen, Y.: *A note on Fischer-Marsden’s conjecture*. Proc. Amer. Math. Soc. 125 (1997) 901–905.
- [55] Wang, L.: *On noncompact  $\tau$ -quasi-Einstein metrics*. Pacific J. Math. 254 (2011) 449–464.
- [56] Wylie, W.: *Rigidity of compact static near-horizon geometries with negative cosmological constant*. Lett. Math. Phys. (2023) 113:29.
- [57] Yuan, W.: *Volume comparison with respect to scalar curvature*. Analysis & PDE, 16 (2023) 1–34.

(R. Diógenes) UNILAB, INSTITUTO DE CIÊNCIAS EXATAS E DA NATUREZA, RUA JOSÉ FRANCO DE OLIVEIRA, s/n, 62790-970, REDENÇÃO / CE, BRAZIL.  
*Email address:* rafaeldiogenes@unilab.edu.br

(J. Gonçalves) UNIVERSIDADE FEDERAL DO CEARÁ - UFC, DEPARTAMENTO DE MATEMÁTICA, CAMPUS DO PICI, AV. HUMBERTO MONTE, BLOCO 914, 60455-760, FORTALEZA / CE, BRAZIL  
*Email address:* mjaciane05@alu.ufc.br

(E. Ribeiro Jr) UNIVERSIDADE FEDERAL DO CEARÁ - UFC, DEPARTAMENTO DE MATEMÁTICA, CAMPUS  
DO PICI, AV. HUMBERTO MONTE, BLOCO 914, 60455-760, FORTALEZA / CE, BRAZIL  
*Email address:* [ernani@mat.ufc.br](mailto:ernani@mat.ufc.br)