

# Dissipative realization of Kondo models

Martino Stefanini,<sup>1,\*</sup> Yi-Fan Qu,<sup>2</sup> Tilman Esslinger,<sup>3</sup> Sarang Gopalakrishnan,<sup>4</sup> Eugene Demler,<sup>2</sup> and Jamir Marino<sup>1</sup>

<sup>1</sup>*Institut für Physik, Johannes Gutenberg-Universität Mainz, D-55099 Mainz, Deutschland*

<sup>2</sup>*Institute for Theoretical Physics, ETH Zürich, 8093 Zurich, Switzerland*

<sup>3</sup>*Institute for Quantum Electronics & Quantum Center, ETH Zurich, 8093 Zurich, Switzerland*

<sup>4</sup>*Department of Electrical and Computer Engineering, Princeton University, Princeton, NJ 08544*

(Dated: June 7, 2024)

We propose a dissipative implementation of a variety of Kondo models by means of strong two-body losses localized on a few impurity sites of a fermionic lattice—a setup which is suited to experiments with ultracold atomic gases. We study in detail the simplest scenario of just one dissipated site, showing that it is effectively described by the Anderson impurity model with infinite repulsion, perturbed by a small residual dissipation. We compute a number of signatures of the Kondo effect in transport across the impurity, finding a competition between the Kondo resonance and the residual dissipation. Our dissipative setup can be generalized to two or more sites subject to losses—realizing an impurity with spin 1 or higher—and more reservoirs, opening up the possibility of simulating several kinds of Kondo-like models with ultracold atoms.

*Introduction*— The Kondo effect [1] is one of the simplest and most iconic models in the physics of strongly correlated systems. It emerges when an interacting impurity can exchange particles with a gapless fermionic bath. The hybridization of the impurity levels with the bath’s states causes the emergence of a very narrow many-body resonance (the Kondo, or Abrikosov-Suhl resonance) pinned at the chemical potential of the reservoir, whose properties dominate the low-energy physics and lead to a number of fascinating phenomena, such as universal behavior of various thermodynamics quantities [1–3], and an almost perfect conductance through the impurity [4–7]. The Kondo effect first arose as the the origin of the anomalous behavior of metals doped with magnetic impurities [1–3, 8–16], but since its inception it has shaped our understanding of a variety of phenomena in strongly correlated systems: besides fueling the development of renormalization group ideas [17–20], it forms the building block of the physics of heavy fermion compounds [1, 3, 21, 22], it has been instrumental in describing transport in quantum dots [23–25] and to the understanding of the Mott transition within dynamical mean field theory [26]. In addition to the well-understood physics of the original Kondo model [27–30], generalized Kondo models in which the impurity spin is higher than 1/2 and interacts with fermions of different flavors [1, 31–34] have been considered. Such generalizations are important for realistic models of magnetic impurities in metals [16, 31, 35, 36], but they have been recognized to show remarkable properties, from non-monotonic conductance [37–39] to non-Fermi liquid ground states [32–34, 40, 41], and are still actively researched.

With the advent of quantum simulation with ultracold atomic gases, new possibilities have arisen for the study of Kondo physics in regimes that would be otherwise inaccessible to traditional solid state setups, such as the exploration of real-time dynamics. In this Letter, we propose and discuss a dissipative route to the implemen-

tation of both standard and more exotic Kondo models. We show how a strong, localized two-body loss [42–46] within a noninteracting fermionic gas can provide the correlations necessary to induce the Kondo effect. The advantages of using dissipation with respect to unitary dynamics [47, 48] are that it is simple to localize and suited to a transport setup [49–57], while the irreversible nature of losses drives the system to the desired Kondo state. We remark *en passant* that dissipation can also be used to introduce entirely new features, such as engineering of non-Hermitian versions of the Kondo model [58–60] and measurement-induced crossovers in continuously monitored quantum dots [59, 61].

Our main finding is the characterization of the Kondo effect induced by a strong two-body loss localized on a single site connected to two reservoirs of noninteracting fermions. We argue that for an infinitely strong dissipation this system realizes the well-known Anderson impurity model (AIM) with infinite repulsion, and we derive the dissipative corrections for finite dissipation rate. We analyze the typical signatures of the Kondo effect (Kondo resonance in the spectral function, enhanced differential conductance at zero bias and suppressed decay of magnetization) both in dynamics and in the local steady state, and characterize the competition between the Kondo effect and the residual losses.

We conclude by describing how to realize higher-spin Kondo models by distributing the dissipation on more than one site, provided they are strongly coupled among themselves. Since there is no *a priori* limit on the number of reservoirs coupled to the impurity sites, our proposal would pave the way to the experimental study of generalized higher-spin, many-flavor Kondo models in ultracold atoms.

*Model*— We consider the system depicted in the leftmost part of Fig. 1, composed of two reservoirs (or leads) of noninteracting, spinful fermions that are connected to a central region (the “dot”) whose sites are




FIG. 1. Using strong, localized two-body losses to realize the Anderson impurity model with infinite repulsion. Left panel: sketch of the simplest dissipative setup considered in the main text. The two tubes represent reservoirs of noninteracting, spinful fermions (possibly at different chemical potentials) that are allowed to tunnel to the central site, which is subjected to strong two-body losses. Inset: sketch of the dark states of the isolated impurity subjected to the two-body loss. The hatched area represents the dissipated states. Central panel: impurity spectral function in the local steady state, showing the smearing of the Kondo resonance as the effective dissipation is increased. The parameters are  $\varepsilon_d = -2\Gamma_T$  and  $\Gamma_T = 10^{-2}W$  (we take the half-bandwidth  $W$  as our energy unit), for unbiased reservoirs at  $\mu = 0$  and zero initial temperature. Inset: zero-bias conductance across the impurity, normalized to  $g_0 = 2/h$  ( $h$  being Planck's constant). Lighter colors correspond to larger  $|\varepsilon_d|/\Gamma_T \in \{0.5, 1, 1.5, 2, 2.5, 3\}$ . Right panel: decay rate of the impurity magnetization, showing a crossover from the Kondo regime  $\Gamma_{\text{decay}} \sim T_K$  to the incoherent regime  $\Gamma_{\text{decay}} \sim \kappa_\sigma$ . In these plots  $\Gamma_T = 10^{-2}W$ .

subjected to a two body loss—whenever two opposite-spin fermions occupy one of the dissipative sites, they can both be lost from the system with a certain rate  $\gamma$ . We are going to comment on the many-site and many-leads scenario at the end of the paper. In the rest of this Letter we will focus on the simplest case of a single dot site hosting a single orbital, and coupled to two reservoirs. In the conditions common to experiments with ultracold atoms [42–44, 50] losses are Markovian, and the dynamics of the system's density matrix  $\rho(t)$  can be described by a Lindblad master equation  $\dot{\rho}(t) = -i[H, \rho(t)] + \gamma(L\rho(t)L^\dagger - \{L^\dagger L, \rho(t)\}/2)$  with jump operator  $L = d_\downarrow d_\uparrow$  (where  $d_\sigma$  annihilates a fermion with spin  $\sigma$  on the dot, with  $\sigma \in \{\uparrow, \downarrow\} = \{+, -\}$ ). The Hamiltonian has the generic form  $H = H_{\text{dot}} + H_{\text{leads}} + H_{\text{tun}}$ , where the first term describes the local energy level(s) of the dot region, the second term describes the reservoirs and the third accounts for the hybridization (i.e. single-particle hopping) between the dot and the leads. We are going to assume that no interactions are present in any portion of the system, so that the only source of correlations are the two-body losses.

We are interested in the regime of strong dissipation in which the decay rate  $\gamma$  is the largest energy scale. We will consider both the dynamics of the system after turning on the dissipation, and the properties of the local steady state that forms around the dot at long times. Since the dissipation removes pairs of opposite-spin fermions, it is clear that in the limit  $\gamma \rightarrow +\infty$  the system remains confined to a subspace with no double occupancies on the dot site(s). In the case of a single site, this subspace is spanned by three dark states, namely states which are

simultaneously eigenstates of  $H_{\text{dot}}$  and annihilated by the jump  $L$ : the empty dot  $|0\rangle$  and the singly occupied dot  $|\sigma\rangle$ . The dynamics within the dark subspace is generated by the exchange of particles with the leads, according to the Hamiltonian:

$$H_{\text{eff}} = \varepsilon_d \sum_{\sigma} X_{\sigma\sigma} + \sum_{p\sigma} \varepsilon_{p\alpha} c_{p\sigma\alpha}^\dagger c_{p\sigma\alpha} + \sum_{p\sigma} (V_{p\alpha} X_{\sigma 0} c_{p\sigma\alpha} + \text{H.c.}), \quad (1)$$

where we introduced the dot energy  $\varepsilon_d$ ,  $c_{p\sigma\alpha}$  annihilates a fermion in lead  $\alpha \in \{R, L\}$  with momentum  $p$  and single-particle energy  $\varepsilon_{p\alpha} = \varepsilon_p - \mu_\alpha$ , possibly biased by a chemical potential  $\mu_\alpha$ . The tunneling between the dot and the leads is governed by the amplitudes  $V_{p\alpha}$ . For most purposes, the leads are fully described by the level width function [62]  $\Gamma_\alpha(\omega) \equiv \sum_p |V_{p\alpha}|^2 \delta(\omega - \varepsilon_p) \equiv \Gamma_\alpha \xi(\omega)$ , which we parametrize in terms of the energy scale  $\Gamma_\alpha$ , which contributes to the width  $\Gamma_T \equiv \sum_\alpha \Gamma_\alpha$  of the dot levels in the absence of dissipation, and the shape function  $\xi(\omega)$  that sets the bandwidth  $W \gg |\varepsilon_d|, \Gamma_\alpha$ . We are going to assume symmetric leads  $\Gamma_\alpha = \Gamma$  and a flat shape function  $\xi(\omega) = \theta(W - |\omega|)$  ( $\theta$  being the Heaviside function). The operators  $X_{\sigma\sigma} \equiv |\sigma\rangle\langle\sigma|$ ,  $X_{\sigma 0} = X_{0\sigma}^\dagger \equiv |\sigma\rangle\langle 0|$  and  $X_{00} \equiv |0\rangle\langle 0|$  are the known as Hubbard operators [3]. The Hamiltonian (1) is the well-known Anderson impurity model (AIM) with infinite repulsion [1, 3, 14, 31], a strongly correlated model whose properties have been extensively studied. In particular, the model is known to show the Kondo effect—namely, its low-energy physics is dominated by a narrow many-body resonance pinned at the chemical potential of the leads.

It is important to understand to what extent the Kondo phenomenology associated with the AIM survives once we move away from the ideal  $\gamma \rightarrow +\infty$  limit. As long as  $\gamma$  is still the largest energy scale of the system we can derive corrections to the dynamics in the constrained subspace by means of an adiabatic elimination [63] or, equivalently, via a dissipative generalization of the Schrieffer-Wolff transformation [64], the details of which are reported in [65]. Physically, in presence of a strong dissipation  $\gamma \gg \Gamma$  any double occupancy on the dot site is rapidly removed from the system before it can be replenished from the leads during a transient that lasts a few  $\gamma^{-1}$ . The dynamics at later times is effectively projected on the subspace of no double occupancies [66], in which it is effectively described by the master equation  $\dot{\rho}(t) = -i[H_{\text{eff}}, \rho(t)] + L_{\text{eff}}\rho(t)L_{\text{eff}}^\dagger - \{L_{\text{eff}}^\dagger L_{\text{eff}}, \rho(t)\}/2$ , where the Hamiltonian (1) governs the coherent part, while there is a residual dissipation with jump operator  $L_{\text{eff}} \equiv 2\gamma^{-1/2} \sum_{p\sigma\alpha} \sigma V_{p\alpha} X_{0\bar{\sigma}} c_{p\sigma\alpha}$ . The effective dissipation accounts for the virtual processes in which a fermion with spin  $\sigma$  hops from a lead to the dot while the latter is already occupied by a fermion of opposite spin  $\bar{\sigma}$ , causing both fermions to be lost from the system due to the dissipation.

*Signatures of Kondo*— For a strictly infinite dissipation, the dynamics is governed by the effective AIM Hamiltonian (1), which guarantees the presence of Kondo physics—such as the Kondo resonance in the impurity spectral function and the maximal differential conductance at zero bias. Therefore, a first important task is to assess to what extent a finite but large value of  $\gamma$  alters the well-known Kondo features. A second interesting question is how these features emerge from the uncorrelated system at  $\gamma = 0$  once the dissipation is increased.

During the years, a large number of different numerical techniques have been employed to tackle the dynamics of the AIM and of the Kondo model for isolated systems—to mention only some of the most recent, we can list methods based on the time-dependent variational principle [67, 68], on the influence functional [69] and on matrix product states [70]. In the present work we need to deal with a dissipative dynamics, and we employ two different numerical approaches. One is based on quantum trajectories [71] and on a variational Ansatz for the state of the system along the trajectories [72–74]. The detailed description of this method and its main results is the subject of the companion paper [75]. The advantage of this method is that it can be applied for any value of  $\gamma$ , so that we can observe the full crossover towards the Kondo phenomenology as the dissipation is increased. The main observables accessible to the method are currents and single-time averages and correlation functions.

In the present Letter, we take an alternative approach, and work directly with the effective model (1) and the residual dissipation  $L_{\text{eff}}$  to observe the effects of a large but finite  $\gamma$ . In this regime, we use the slave boson rep-

resentation [3, 76, 77] and we apply the widely used non-crossing approximation (NCA) [1, 31, 78–82] to derive Kadanoff-Baym equations for the Keldysh Green’s functions of the auxiliary particles [65]. To the same leading perturbative order as the NCA self-energy, it suffices to treat the effective dissipation at the mean-field level. The crucial feature of this extended NCA is that the resulting approximation is conserving [78–84]—namely, the approximate dynamics respects the relevant conservation laws—a necessary feature for real-time simulations.

Before describing the main results, we wish to remark that our dissipative model leads to an effective AIM physics only in the sense that a local steady state forms around the dot, as it is common with dissipative impurities [85–97]. The true stationary states in our model are fully ferromagnetic Dicke states [65, 98] that extend to the leads. When starting from an initial spin-balanced configuration (before losses are turned on), such fully polarized states can not be reached due to spin conservation. A related yet possible scenario is the formation of a finite size ferromagnetic bubble around the impurity site. The larger the bubble, the stronger the suppression of losses. On the other hand, larger bubbles are energetically unfavorable and their formation requires a less likely statistical fluctuation. In the NCA analysis the possibility of ferromagnetic bubbles is not included, because it requires introducing spin symmetry breaking. We will discuss conditions for observing ferromagnetic bubbles in the companion paper [75].

We have computed the real-time dynamics of the effective model after a quench of the tunneling  $\Gamma$ , and we have analyzed its properties in the local steady state at late times. We first discuss the stationary properties, which are clearer to understand. The main signature of Kondo behavior is the presence of the Kondo resonance in the impurity spectral function  $A_\sigma(\omega) \equiv -\text{Im} G_{d\sigma}^R(\omega)/\pi$ , where  $G_{d\sigma}^R(\omega)$  is the Fourier transform of the retarded impurity Green’s function  $G_{d\sigma}^R(t-t') \equiv -i\theta(t-t') \langle \{d_\sigma(t), d_\sigma^\dagger(t')\} \rangle$  [65], computed in the local steady state [99]. We show it in the central panel of Fig. 1 for an impurity tuned to the Kondo regime  $\varepsilon_d = -2\Gamma_T$  in presence of unbiased reservoirs. We observe the typical two-peak shape that one finds in the AIM [1, 3, 31]. There is a broad peak centered close to the single-particle energy  $\varepsilon_d$ , whose width is set by the original tunneling rate  $\Gamma$ . This peak reflects the rapid exchange of charge between the dot and the leads. The second peak is the Kondo resonance: a much narrower feature, pinned at the reservoirs’ chemical potential  $\mu_{R,L} = 0$ , which signals the presence of a long-lived spin degree of freedom. In the  $\gamma \rightarrow \infty$  limit, the width of the peak is set by the Kondo temperature  $T_K \equiv [2\Gamma_T(\mu - \varepsilon_d)]^{1/2} \exp[-\pi(\mu - \varepsilon_d)/\Gamma_T]$  [1]. In the full model there would be also an extremely broad peak at the same energy  $\omega \approx \varepsilon_d$ , with width of order  $\gamma$  and a highly reduced height, corresponding to the doubly-occupied dot. The effect of a finite dissipation





as simply the Dicke states [98, 106, 107] associated with  $H_{\text{dot}}$  [65]. The resulting dark subspace is organized in multiplets of increasing spin  $S = 0, \dots, \ell_d/2$  and particle numbers  $n = 2S$ , with higher spins possessing lower energy (although there are  $\binom{\ell_d}{2S}$  different multiplets with the same spin  $0 < S < \ell_d/2$  but different energies, where  $(\cdot)$  is the binomial coefficient). We notice that this procedure is completely general, in the sense that it does not rely on any particular structure of the dot region, except for spin rotational invariance. In this regard, our treatment is similar to that of the Hubbard Hamiltonian with infinite repulsion [108–110].

In the setup with two sites there is an extra complication with respect to the single-site case: the slowest-decaying eigenstate of the Lindbladian has a decay rate  $\gamma_1$  which is nonmonotonic in  $\gamma$ , with a maximum  $\gamma_1^*$  at  $\gamma = 8J$  followed by a slow decrease  $\gamma_1 \sim J^2/\gamma$ . In contrast, all other bright states have decay rates of order of  $\gamma$ . So, to keep the occupation of all bright states suppressed by dissipation, we need to have a large hopping  $J \sim \gamma$ , and hence a large dot energy  $\varepsilon_d \sim \gamma$  to maintain the spin 1 states at energy lower than the spin 1/2 ones,  $2\varepsilon_d < \varepsilon_d - J$ . The nonmonotonic behavior of the smallest decay rate, and the consequent need for fine tuning, is likely to be generic for higher  $\ell_d$  [65].

If we introduce back the coupling to the leads, the latter will mediate transitions between multiplets with neighboring values of  $S$ , and we obtain a model that belongs to the family of the ionic models [1, 31, 36], that have been extensively studied in the context of magnetic impurities in metals. These models can be mapped on higher-spin Kondo models by the usual Schrieffer-Wolff transformation [1, 3, 15]. In the case of two sites, the lowest-energy multiplet would act like a spin 1 impurity. On the other hand, there is no constraint on the number of possible leads, thus opening to the possibility of studying over-screened realizations of these Kondo-like models, that are known to possess non-Fermi liquid ground states [1, 32, 33, 38, 39].

*Acknowledgments*— We acknowledge useful discussions with D. Sels, A. Gomez Salvador, R. Andrei, M. Kiselev. M.S. and J.M. have been supported by the DFG through the grant HADEQUAM-MA7003/3-1. Y.F. and E.D. acknowledge support by the SNSF project 200021.212899, the NCCR SPIN of the Swiss NSF., the Swiss State Secretariat for Education, Research and Innovation (contract number UeM019-1). E.D. also acknowledges support from the ARO grant number W911NF-20-1-0163. T.E., Y.Q., and E.D. acknowledge funding by the ETH grant. J.M. acknowledges the Pauli Center for hospitality. We gratefully acknowledge the computing time granted through the project “DysQCorr” on the Mogon II supercomputer of the Johannes Gutenberg University Mainz ([hpc.uni-mainz.de](http://hpc.uni-mainz.de)), which is a member of the AHRP (Alliance for High Performance Computing in Rhineland Palatinate, [www.ahrp.info](http://www.ahrp.info)),

and the Gauss Alliance e.V.

---

\* [mstefa@uni-mainz.de](mailto:mstefa@uni-mainz.de)

- [1] A. C. Hewson, *The Kondo Problem to Heavy Fermions*, Cambridge Studies in Magnetism (Cambridge University Press, 1993).
- [2] J. Kondo, Theory of Dilute Magnetic Alloys, in *Solid State Physics*, Vol. 23, edited by F. Seitz, D. Turnbull, and H. Ehrenreich (Academic Press, 1970) pp. 183–281.
- [3] P. Coleman, *Introduction to Many-Body Physics* (Cambridge University Press, 2015).
- [4] T. K. Ng and P. A. Lee, On-Site Coulomb Repulsion and Resonant Tunneling, *Phys. Rev. Lett.* **61**, 1768 (1988).
- [5] L. I. Glazman and M. E. Raikh, Resonant kondo transparency of a barrier with quasilocal impurity states, *JETP Letters* **47**, 378 (1988).
- [6] N. S. Wingreen and Y. Meir, Anderson model out of equilibrium: Noncrossing-approximation approach to transport through a quantum dot, *Phys. Rev. B* **49**, 11040 (1994).
- [7] W. G. van der Wiel, S. D. Franceschi, T. Fujisawa, J. M. Elzerman, S. Tarucha, and L. P. Kouwenhoven, The Kondo Effect in the Unitary Limit, *Science* **289**, 2105 (2000).
- [8] J. Kondo, Resistance Minimum in Dilute Magnetic Alloys, *Progress of Theoretical Physics* **32**, 37 (1964).
- [9] W. de Haas, J. de Boer, and G. van den Berg, The electrical resistance of gold, copper and lead at low temperatures, *Physica* **1**, 1115 (1934).
- [10] A. M. Clogston, B. T. Matthias, M. Peter, H. J. Williams, E. Corenzwit, and R. C. Sherwood, Local magnetic moment associated with an iron atom dissolved in various transition metal alloys, *Phys. Rev.* **125**, 541 (1962).
- [11] A. A. Abrikosov, Electron scattering on magnetic impurities in metals and anomalous resistivity effects, *Physics Physique Fizika* **2**, 5 (1965).
- [12] H. Suhl, Formation of Local Magnetic Moments in Metals, *Phys. Rev. Lett.* **19**, 442 (1967).
- [13] A. Blandin and J. Friedel, Propriétés magnétiques des alliages dilués. Interactions magnétiques et antiferromagnétisme dans les alliages du type métal noble-métal de transition, *J. Phys. Radium* **20**, 160 (1959).
- [14] P. W. Anderson, Localized magnetic states in metals, *Phys. Rev.* **124**, 41 (1961).
- [15] J. R. Schrieffer and P. A. Wolff, Relation between the Anderson and Kondo Hamiltonians, *Phys. Rev.* **149**, 491 (1966).
- [16] B. Coqblin and J. R. Schrieffer, Exchange interaction in alloys with cerium impurities, *Phys. Rev.* **185**, 847 (1969).
- [17] P. W. Anderson and G. Yuval, Exact Results in the Kondo Problem: Equivalence to a Classical One-Dimensional Coulomb Gas, *Phys. Rev. Lett.* **23**, 89 (1969).
- [18] P. W. Anderson, A poor man’s derivation of scaling laws for the Kondo problem, *Journal of Physics C: Solid State Physics* **3**, 2436 (1970).
- [19] M. Fowler and A. Zawadowski, Scaling and the renormalization group in the Kondo effect, *Solid State Com-*

- munications **9**, 471 (1971).
- [20] K. G. Wilson, The renormalization group: Critical phenomena and the Kondo problem, *Rev. Mod. Phys.* **47**, 773 (1975).
  - [21] F. Steglich, J. Aarts, C. D. Bredl, W. Lieke, D. Meschede, W. Franz, and H. Schäfer, Superconductivity in the Presence of Strong Pauli Paramagnetism:  $\text{CeCu}_2\text{Si}_2$ , *Phys. Rev. Lett.* **43**, 1892 (1979).
  - [22] G. R. Stewart, Heavy-fermion systems, *Rev. Mod. Phys.* **56**, 755 (1984).
  - [23] M. Pustilnik and L. Glazman, Kondo effect in quantum dots, *Journal of Physics: Condensed Matter* **16**, R513 (2004).
  - [24] D. Goldhaber-Gordon, H. Shtrikman, D. Mahalu, D. Abusch-Magder, U. Meirav, and M. A. Kastner, Kondo effect in a single-electron transistor, *Nature* **391**, 156 (1998).
  - [25] S. M. Cronenwett, T. H. Oosterkamp, and L. P. Kouwenhoven, A Tunable Kondo Effect in Quantum Dots, *Science* **281**, 540 (1998).
  - [26] A. Georges, G. Kotliar, W. Krauth, and M. J. Rozenberg, Dynamical mean-field theory of strongly correlated fermion systems and the limit of infinite dimensions, *Rev. Mod. Phys.* **68**, 13 (1996).
  - [27] P. Nozières, A “Fermi-liquid” description of the Kondo problem at low temperatures, *Journal of Low Temperature Physics* **17**, 31 (1974).
  - [28] N. Andrei, Diagonalization of the Kondo Hamiltonian, *Phys. Rev. Lett.* **45**, 379 (1980).
  - [29] N. Andrei, K. Furuya, and J. H. Lowenstein, Solution of the Kondo problem, *Rev. Mod. Phys.* **55**, 331 (1983).
  - [30] P. Vignani, Exact solution of sd exchange model at  $T=0$ , *Soviet Journal of Experimental and Theoretical Physics Letters* **31**, 364 (1980).
  - [31] N. E. Bickers, Review of techniques in the large- $N$  expansion for dilute magnetic alloys, *Rev. Mod. Phys.* **59**, 845 (1987).
  - [32] A. O. Gogolin, A. A. Nersisyan, and A. M. Tsvelik, *Bosonization and Strongly Correlated Systems* (Cambridge University Press, Cambridge, 1998).
  - [33] T. Giamarchi, *Quantum Physics in One Dimension*, 1st ed. (Clarendon Press, Oxford, 2003).
  - [34] D. L. Cox and A. Zawadowski, Exotic Kondo effects in metals: Magnetic ions in a crystalline electric field and tunnelling centres, *Advances in Physics* **47**, 599 (1998).
  - [35] P. Nozières and A. Blandin, Kondo effect in real metals, *J. Phys. France* **41**, 193 (1980).
  - [36] L. Hirst, Theory of the coupling between conduction electrons and moments of 3d and 4f ions in metals, *Advances in Physics* **27**, 231 (1978).
  - [37] R. M. Potok, I. G. Rau, H. Shtrikman, Y. Oreg, and D. Goldhaber-Gordon, Observation of the two-channel Kondo effect, *Nature* **446**, 167 (2007).
  - [38] D. B. Karki, C. Mora, J. von Delft, and M. N. Kiselev, Two-color Fermi-liquid theory for transport through a multilevel Kondo impurity, *Phys. Rev. B* **97**, 195403 (2018).
  - [39] D. B. Karki, A. I. Pavlov, and M. N. Kiselev, Multistage Kondo effect in a multiterminal geometry: A modular quantum interferometer, *Phys. Rev. B* **105**, L041410 (2022).
  - [40] A. W. W. Ludwig and I. Affleck, Exact, asymptotic, three-dimensional, space- and time-dependent, Green’s functions in the multichannel Kondo effect, *Phys. Rev. Lett.* **67**, 3160 (1991).
  - [41] P. D. Sacramento and P. Schlottmann, Thermodynamics of the n-channel Kondo model for general n and impurity spin S in a magnetic field, *Journal of Physics: Condensed Matter* **3**, 9687 (1991).
  - [42] B. Yan, S. A. Moses, B. Gadway, J. P. Covey, K. R. A. Hazzard, A. M. Rey, D. S. Jin, and J. Ye, Observation of dipolar spin-exchange interactions with lattice-confined polar molecules, *Nature* **501**, 521 (2013).
  - [43] B. Zhu, B. Gadway, M. Foss-Feig, J. Schachenmayer, M. L. Wall, K. R. A. Hazzard, B. Yan, S. A. Moses, J. P. Covey, D. S. Jin, J. Ye, M. Holland, and A. M. Rey, Suppressing the loss of ultracold molecules via the continuous quantum Zeno effect, *Phys. Rev. Lett.* **112**, 070404 (2014).
  - [44] K. Sponselee, L. Freystatzky, B. Abeln, M. Diem, B. Hundt, A. Kochanek, T. Ponath, B. Santra, L. Mathey, K. Sengstock, and C. Becker, Dynamics of ultracold quantum gases in the dissipative Fermi-Hubbard model, *Quantum Sci. Technol.* **4**, 014002 (2018).
  - [45] T. Tomita, S. Nakajima, Y. Takasu, and Y. Takahashi, Dissipative Bose-Hubbard system with intrinsic two-body loss, *Phys. Rev. A* **99**, 031601 (2019).
  - [46] N. Syassen, D. M. Bauer, M. Lettner, T. Volz, D. Dietze, J. J. García-Ripoll, J. I. Cirac, G. Rempe, and S. Dürr, Strong Dissipation Inhibits Losses and Induces Correlations in Cold Molecular Gases, *Science* **320**, 1329 (2008).
  - [47] L. Riegger, N. Darkwah Oppong, M. Höfer, D. R. Fernandes, I. Bloch, and S. Fölling, Localized magnetic moments with tunable spin exchange in a gas of ultracold fermions, *Phys. Rev. Lett.* **120**, 143601 (2018).
  - [48] M. Kanász-Nagy, Y. Ashida, T. Shi, C. u. u. u. P. m. c. Moca, T. N. Ikeda, S. Fölling, J. I. Cirac, G. Zaránd, and E. A. Demler, Exploring the anisotropic Kondo model in and out of equilibrium with alkaline-earth atoms, *Phys. Rev. B* **97**, 155156 (2018).
  - [49] S. Krinner, T. Esslinger, and J.-P. Brantut, Two-terminal transport measurements with cold atoms, *Journal of Physics: Condensed Matter* **29**, 343003 (2017).
  - [50] L. Corman, P. Fabritius, S. Häusler, J. Mohan, L. H. Dogra, D. Husmann, M. Lebrat, and T. Esslinger, Quantized conductance through a dissipative atomic point contact, *Phys. Rev. A* **100**, 053605 (2019).
  - [51] M. Lebrat, S. Häusler, P. Fabritius, D. Husmann, L. Corman, and T. Esslinger, Quantized conductance through a spin-selective atomic point contact, *Phys. Rev. Lett.* **123**, 193605 (2019).
  - [52] M.-Z. Huang, J. Mohan, A.-M. Visuri, P. Fabritius, M. Talebi, S. Wili, S. Uchino, T. Giamarchi, and T. Esslinger, Superfluid signatures in a dissipative quantum point contact, *Phys. Rev. Lett.* **130**, 200404 (2023).
  - [53] M. Lebrat, P. Grišins, D. Husmann, S. Häusler, L. Corman, T. Giamarchi, J.-P. Brantut, and T. Esslinger, Band and correlated insulators of cold fermions in a mesoscopic lattice, *Phys. Rev. X* **8**, 011053 (2018).
  - [54] R. Labouvie, B. Santra, S. Heun, S. Wimberger, and H. Ott, Negative differential conductivity in an interacting quantum gas, *Phys. Rev. Lett.* **115**, 050601 (2015).
  - [55] R. Labouvie, B. Santra, S. Heun, and H. Ott, Bistability in a driven-dissipative superfluid, *Phys. Rev. Lett.* **116**, 235302 (2016).

- [56] M. Will, J. Marino, H. Ott, and M. Fleischhauer, Controlling superfluid flows using dissipative impurities, *SciPost Phys.* **14**, 064 (2023).
- [57] G. Barontini, R. Labouvie, F. Stubenrauch, A. Vogler, V. Guarrera, and H. Ott, Controlling the Dynamics of an Open Many-Body Quantum System with Localized Dissipation, *Phys. Rev. Lett.* **110**, 035302 (2013).
- [58] M. Nakagawa, N. Kawakami, and M. Ueda, Non-Hermitian Kondo Effect in Ultracold Alkaline-Earth Atoms, *Phys. Rev. Lett.* **121**, 203001 (2018).
- [59] M. Hasegawa, M. Nakagawa, and K. Saito, Kondo effect in a quantum dot under continuous quantum measurement (2021), [arXiv:2111.07771 \[cond-mat.mes-hall\]](#).
- [60] P. Kattel, A. Zhakenov, P. R. Pasnoori, P. Azaria, and N. Andrei, Dissipation driven phase transition in the non-Hermitian Kondo model (2024), [arXiv:2402.09510 \[cond-mat.str-el\]](#).
- [61] M. Vanhoecke and M. Schirò, Kondo-Zeno crossover in the dynamics of a monitored quantum dot (2024), [arXiv:2405.17348 \[cond-mat.str-el\]](#).
- [62] H. Haug and A.-P. Jauho, *Quantum Kinetics in Transport and Optics of Semiconductors*, 2nd ed., Springer Series in Solid-State Sciences (Springer Berlin, Heidelberg, 2007).
- [63] J. J. García-Ripoll, S. Dürr, N. Syassen, D. M. Bauer, M. Lettner, G. Rempe, and J. I. Cirac, Dissipation-induced hard-core boson gas in an optical lattice, *New Journal of Physics* **11**, 013053 (2009).
- [64] E. M. Kessler, Generalized Schrieffer-Wolff formalism for dissipative systems, *Phys. Rev. A* **86**, 012126 (2012).
- [65] See the Supplemental Material at [\[url will be inserted by publisher\]](#) for technical details.
- [66] This is not strictly true, of course: there is a residual double occupancy of order  $\langle d_{\uparrow}^{\dagger} d_{\uparrow} d_{\downarrow}^{\dagger} d_{\downarrow} \rangle \sim \gamma^{-1}$ , that is responsible for the effective loss that is suppressed as  $1/\gamma$ —an instance of the well-known Zeno effect [43].
- [67] Y. Ashida, T. Shi, M. C. Bañuls, J. I. Cirac, and E. Demler, Solving quantum impurity problems in and out of equilibrium with the variational approach, *Phys. Rev. Lett.* **121**, 026805 (2018).
- [68] Y. Ashida, T. Shi, M. C. Bañuls, J. I. Cirac, and E. Demler, Variational principle for quantum impurity systems in and out of equilibrium: Application to Kondo problems, *Phys. Rev. B* **98**, 024103 (2018).
- [69] J. Thoenniss, M. Sonner, A. Lerose, and D. A. Abanin, Efficient method for quantum impurity problems out of equilibrium, *Phys. Rev. B* **107**, L201115 (2023).
- [70] M. M. Wauters, C.-M. Chung, L. Maffi, and M. Burrello, Simulations of the dynamics of quantum impurity problems with matrix product states, *Phys. Rev. B* **109**, 115101 (2024).
- [71] A. J. Daley, Quantum trajectories and open many-body quantum systems, *Advances in Physics* **63**, 77 (2014).
- [72] S. Bravyi and D. Gosset, Complexity of quantum impurity problems, *Communications in Mathematical Physics* **356**, 451 (2017).
- [73] I. Snyman and S. Florens, Efficient impurity-bath trial states from superposed Slater determinants, *Phys. Rev. B* **104**, 195136 (2021).
- [74] W. Zhang and T. Shi (2024), in preparation.
- [75] Y.-F. Qu, M. Stefanini, T. Esslinger, S. Gopalakrishnan, J. Marino, and E. Demler, in preparation (2024).
- [76] P. Coleman, New approach to the mixed-valence problem, *Phys. Rev. B* **29**, 3035 (1984).
- [77] S. E. Barnes, New method for the Anderson model, *Journal of Physics F: Metal Physics* **6**, 1375 (1976).
- [78] M. H. Hettler, J. Kroha, and S. Hershfield, Nonequilibrium dynamics of the Anderson impurity model, *Phys. Rev. B* **58**, 5649 (1998).
- [79] N. S. Wingreen and Y. Meir, Anderson model out of equilibrium: Noncrossing-approximation approach to transport through a quantum dot, *Phys. Rev. B* **49**, 11040 (1994).
- [80] N. Sivan and N. S. Wingreen, Single-impurity Anderson model out of equilibrium, *Phys. Rev. B* **54**, 11622 (1996).
- [81] D. C. Langreth and P. Nordlander, Derivation of a master equation for charge-transfer processes in atom-surface collisions, *Phys. Rev. B* **43**, 2541 (1991).
- [82] H. Shao, D. C. Langreth, and P. Nordlander, Many-body theory for charge transfer in atom-surface collisions, *Phys. Rev. B* **49**, 13929 (1994).
- [83] G. Baym, Self-consistent approximations in many-body systems, *Phys. Rev.* **127**, 1391 (1962).
- [84] M. Fabrizio, *A Course in Quantum Many-Body Theory*, 1st ed., Graduate Texts in Physics (Springer Cham, 2022).
- [85] P. E. Dolgirev, J. Marino, D. Sels, and E. Demler, Non-Gaussian correlations imprinted by local dephasing in fermionic wires, *Phys. Rev. B* **102**, 100301(R) (2020).
- [86] V. Alba and F. Carollo, Noninteracting fermionic systems with localized losses: Exact results in the hydrodynamic limit, *Phys. Rev. B* **105**, 054303 (2022).
- [87] F. Tonielli, R. Fazio, S. Diehl, and J. Marino, Orthogonality catastrophe in dissipative quantum many-body systems, *Physical Review Letters* **122**, 040604 (2019).
- [88] F. Tonielli, N. Chakraborty, F. Grusdt, and J. Marino, Ramsey interferometry of non-Hermitian quantum impurities, *Physical Review Research* **2**, 032003 (2020).
- [89] H. Fröml, C. Muckel, C. Kollath, A. Chiocchetta, and S. Diehl, Ultracold quantum wires with localized losses: Many-body quantum Zeno effect, *Phys. Rev. B* **101**, 144301 (2020).
- [90] T. Müller, M. Gievers, H. Fröml, S. Diehl, and A. Chiocchetta, Shape effects of localized losses in quantum wires: Dissipative resonances and nonequilibrium universality, *Phys. Rev. B* **104**, 155431 (2021).
- [91] H. Fröml, A. Chiocchetta, C. Kollath, and S. Diehl, Fluctuation-induced quantum Zeno effect, *Phys. Rev. Lett.* **122**, 040402 (2019).
- [92] A. P. Chaudhari, S. P. Kelly, R. J. Valencia-Tortora, and J. Marino, Zeno crossovers in the entanglement speed of spin chains with noisy impurities, *Journal of Statistical Mechanics: Theory and Experiment* **2022**, 103101 (2022).
- [93] U. Javed, J. Marino, V. Oganesyan, and M. Kolodrubetz, Counting edge modes via dynamics of boundary spin impurities, *Physical Review B* **108**, L140301 (2023).
- [94] U. Javed, R. J. Valencia-Tortora, J. Marino, V. Oganesyan, and M. Kolodrubetz, Zeno physics of the Ising chain with symmetry-breaking boundary dephasing, [arXiv preprint arXiv:2404.04255](#) (2024).
- [95] B. Doyon and N. Andrei, Universal aspects of nonequilibrium currents in a quantum dot, *Phys. Rev. B* **73**, 245326 (2006).
- [96] M. Schiro and O. Scarlatella, Quantum impurity models

- coupled to Markovian and non-Markovian baths, *The Journal of Chemical Physics* **151**, 044102 (2019).
- [97] M. Vanhovecke and M. Schirò, Diagrammatic Monte Carlo for dissipative quantum impurity models, *Phys. Rev. B* **109**, 125125 (2024).
  - [98] M. Foss-Feig, A. J. Daley, J. K. Thompson, and A. M. Rey, Steady-state many-body entanglement of hot reactive fermions, *Phys. Rev. Lett.* **109**, 230501 (2012).
  - [99] See [65] for the full time-dependent generalization of  $A_\sigma(\omega)$ , displayed in the Figure.
  - [100] A.-M. Visuri, T. Giamarchi, and C. Kollath, Symmetry-protected transport through a lattice with a local particle loss, *Phys. Rev. Lett.* **129**, 056802 (2022).
  - [101] S. Uchino, Comparative study for two-terminal transport through a lossy one-dimensional quantum wire, *Phys. Rev. A* **106**, 053320 (2022).
  - [102] A.-M. Visuri, J. Mohan, S. Uchino, M.-Z. Huang, T. Esslinger, and T. Giamarchi, DC transport in a dissipative superconducting quantum point contact, *Phys. Rev. Res.* **5**, 033095 (2023).
  - [103] In principle, one could describe the particle losses as caused by the coupling to a third, empty reservoir [101].
- This point of view might be sometimes useful for interpreting the results.
- [104] Y. Meir and N. S. Wingreen, Landauer formula for the current through an interacting electron region, *Phys. Rev. Lett.* **68**, 2512 (1992).
  - [105] Although it is known that the NCA tends to overestimate the conductance by a few percent [6].
  - [106] R. H. Dicke, Coherence in spontaneous radiation processes, *Phys. Rev.* **93**, 99 (1954).
  - [107] F. T. Arecchi, E. Courtens, R. Gilmore, and H. Thomas, Atomic coherent states in quantum optics, *Phys. Rev. A* **6**, 2211 (1972).
  - [108] E. H. Lieb and F. Y. Wu, Absence of Mott Transition in an Exact Solution of the Short-Range, One-Band Model in One Dimension, *Phys. Rev. Lett.* **20**, 1445 (1968).
  - [109] B. Douçot and X. G. Wen, Instability of the Nagaoka state with more than one hole, *Phys. Rev. B* **40**, 2719 (1989).
  - [110] A. Mielke, The one-dimensional Hubbard model for large or infinite  $U$ , *Journal of Statistical Physics* **62**, 509 (1991).



# Supplemental Material — Dissipative realization of Kondo models

Martino Stefanini,<sup>1</sup> Yi-Fan Qu,<sup>2</sup> Tilman Esslinger,<sup>3</sup> Sarang Gopalakrishnan,<sup>4</sup> Eugene Demler,<sup>2</sup> and Jamir Marino<sup>1</sup>

<sup>1</sup>*Institut für Physik, Johannes Gutenberg-Universität Mainz, D-55099 Mainz, Deutschland*

<sup>2</sup>*Institute for Theoretical Physics, ETH Zurich, 8093 Zurich, Switzerland*

<sup>3</sup>*Institute for Quantum Electronics & Quantum Center, ETH Zurich, 8093 Zurich, Switzerland*

<sup>4</sup>*Department of Electrical and Computer Engineering, Princeton University, Princeton, NJ 08544*

(Dated: June 7, 2024)

## DERIVATION OF THE EFFECTIVE DYNAMICS

In this Section we derive the effective master equation governing the dynamics of the single-site dissipative dot at large dissipation.

We consider the strongly dissipative limit in which the loss rate is the largest energy scale  $\gamma \gg W, |\varepsilon_d|, \Gamma$ . In other words, we will take the purely dissipative dynamics as the zeroth-order solution and expand in the Hamiltonian contribution to the Lindblad equation. This approach is slightly different from the usual Schrieffer-Wolff mapping of the Anderson impurity model to the Kondo model [1, 2], which treats the uncoupled dot-leads system as the unperturbed dynamics and expands in  $H_{\text{tun}}$ , i.e. in  $\Gamma$ . In the present model, this would amount to  $\gamma \sim W \sim |\varepsilon_d| \gg \Gamma$ . Although it is possible to adopt this approach also in the dissipative case, it yields considerably more complicated expressions for the effective dynamics, while adding very little to the physical description of the system.

The starting point of our analysis is the purely dissipative dot site, decoupled from the rest of the system. The master equation becomes

$$\frac{d}{dt}\rho = \mathcal{L}_0\rho, \quad (1)$$

where

$$\mathcal{L}_0\rho \equiv \gamma(L\rho L^\dagger - \frac{1}{2}\{L^\dagger L, \rho\}) \quad (2)$$

We label the states of the dissipative site as  $|\alpha\rangle$ , where  $\alpha \in \{0, \uparrow, \downarrow, d = \uparrow\downarrow\}$ , and vectorize the density matrix as [3]  $\rho = \sum_{\alpha\beta} \rho_{\alpha\beta} |\alpha\rangle\langle\beta| \rightarrow |\rho\rangle = \sum_{\alpha\beta} \rho_{\alpha\beta} |\alpha\beta\rangle$ . Then, it is easy to see that in the 16-dimensional  $|\alpha\beta\rangle$  basis the Liouville superoperator  $\mathcal{L}_0$  is almost diagonal:

$$\begin{aligned} \mathcal{L}_0|\alpha\beta\rangle &= 0 \quad \text{for all } \alpha, \beta \in \{0, \uparrow, \downarrow\}, \\ \mathcal{L}_0|\alpha d\rangle &= -\frac{\gamma}{2}|\alpha d\rangle \quad \text{for all } \alpha \in \{0, \uparrow, \downarrow\}, \\ \mathcal{L}_0|d\alpha\rangle &= -\frac{\gamma}{2}|d\alpha\rangle \quad \text{for all } \alpha \in \{0, \uparrow, \downarrow\}, \\ \mathcal{L}_0|dd\rangle &= \gamma|00\rangle \end{aligned}$$

The only coupled states are  $|00\rangle$  and  $|dd\rangle$ , and the diagonalization of the corresponding  $2 \times 2$  matrix yields the remaining two eigenvalues of  $\mathcal{L}_0$ :

$$\left. \begin{aligned} \mathcal{L}_0|\phi_0^R\rangle &= 0, \quad |\phi_0^R\rangle = |00\rangle \\ (\phi_0^L|\mathcal{L}_0 &= 0, \quad (\phi_0^L| = \langle 00| + \langle dd| \end{aligned} \right\} \quad \lambda_0 = 0 \quad (3a)$$

$$\left. \begin{aligned} \mathcal{L}_0|\phi_d^R\rangle &= -\gamma|\phi_d^R\rangle, \quad |\phi_d^R\rangle = -|00\rangle + |dd\rangle \\ (\phi_d^L|\mathcal{L}_0 &= -\gamma(\phi_d^L|, \quad (\phi_d^L| = \langle dd| \end{aligned} \right\} \quad \lambda_2 = -\gamma \quad (3b)$$

As usual, we are normalizing the states as  $(\phi_a^L|\phi_b^R) = \delta_{ab}$ . Summing up, there are nine stationary (dark) states with eigenvalue  $\lambda_0 = 0$ , six states with eigenvalue  $\lambda_1 = -\frac{\gamma}{2}$ , and a non-degenerate state with  $\lambda_2 = -\gamma$ . For later use, we report here the superprojectors on the three eigenspaces:

$$\mathcal{P}_0 \equiv |00\rangle\langle dd| + \sum_{\alpha, \beta < d} |\alpha\beta\rangle\langle\alpha\beta|, \quad (4a)$$

$$\mathcal{P}_1 \equiv \sum_{\alpha < d} (|\alpha d\rangle\langle\alpha d| + |d\alpha\rangle\langle d\alpha|), \quad (4b)$$

$$\mathcal{P}_2 \equiv |dd\rangle\langle dd| - |00\rangle\langle dd| \quad (4c)$$

The  $\mathcal{P}_0$  projector defines the slow subspace in which the dynamics will be confined at large dissipation. It is useful to express the above superprojectors in terms of operators in the ordinary Hilbert space:

$$\mathcal{P}_0\rho = X_{0d}\rho X_{d0} + X_{<}\rho X_{<}, \quad (5a)$$

$$\mathcal{P}_1\rho = X_{<}\rho X_d + X_d\rho X_{<}, \quad (5b)$$

$$\mathcal{P}_2\rho = X_d\rho X_d - X_{0d}\rho X_{d0}, \quad (5c)$$

where we have introduced the ordinary projectors (Hubbard operators [1, 2, 4])

$$X_{\alpha\beta} \equiv |\alpha\rangle\langle\beta|, \quad (6a)$$

$$X_\alpha \equiv X_{\alpha\alpha} = |\alpha\rangle\langle\alpha|, \quad (6b)$$

$$X_{<} \equiv \sum_{\alpha < d} X_\alpha \quad (6c)$$

We turn to consider the full Lindblad master equation

$$\frac{d}{dt}\rho = (\mathcal{L}_0 + \mathcal{L}_H)\rho, \quad (7)$$

where  $\mathcal{L}_H = -i[H, \cdot]$  is the Hamiltonian part, which we will treat as a perturbation. Following [5] and [3], the effective dynamics in the slow subspace  $\mathcal{P}_0\rho = \rho$  in the  $\gamma \rightarrow +\infty$  limit is generated by the effective Lindblad superoperator  $\mathcal{L}_{\text{eff}} = \mathcal{L}_1 + \mathcal{L}_2 + \mathcal{O}(\gamma^{-2})$ , where

$$\mathcal{L}_1 = \mathcal{P}_0\mathcal{L}_H\mathcal{P}_0, \quad (8a)$$

$$\mathcal{L}_2 = -\mathcal{P}_0\mathcal{L}_H(\mathcal{Q}_0\mathcal{L}_0\mathcal{Q}_0)^{-1}\mathcal{L}_H\mathcal{P}_0, \quad (8b)$$

and  $\mathcal{Q}_0 = \mathbb{1} - \mathcal{P}_0$  is the complementary projector to  $\mathcal{P}_0$ . The first-order term is just the Hamiltonian part projected on the stationary subspace of  $\mathcal{L}_0$ ,  $\mathcal{L}_1\rho = \mathcal{P}_0\mathcal{L}_H\mathcal{P}_0\rho$ . We classify the Hamiltonian terms according to the commutativity of the corresponding superoperator  $-i[H, \cdot]$  with  $\mathcal{P}_0$ . The dot Hamiltonian  $H_{\text{dot}}$  and the leads' one  $H_{\text{leads}}$  both commute (the latter since  $\mathcal{P}_0$  acts as the identity on the leads states). The tunneling term  $H_{\text{tun}}$  is the only one which does not commute, since it connects the slow subspace  $\mathcal{P}_0\rho = \rho$  with the other subspaces. The decomposition  $d_\sigma = X_{0\sigma} + \sigma X_{\bar{\sigma}d}$  induces the further subdivision  $H_{\text{tun}} = H_{\text{tun}}^0 + H_{\text{tun}}^1$ , with

$$\begin{aligned} H_{\text{tun}}^0 &= \sum_{\sigma} (X_{\sigma 0}\Psi_{\sigma} + \Psi_{\sigma}^{\dagger}X_{0\sigma}), \\ H_{\text{tun}}^1 &= \sum_{\sigma} \sigma (X_{d\bar{\sigma}}\Psi_{\sigma} + \Psi_{\sigma}^{\dagger}X_{\bar{\sigma}d}), \end{aligned} \quad (9)$$

where  $\Psi_{\sigma} \equiv \sum_{p\alpha} V_{p\alpha}c_{p\sigma\alpha}$ . Notice that the projectors  $X_{\sigma 0}$ ,  $X_{d\sigma}$  and their conjugates are of fermionic nature in the sense that they anticommute with  $\Psi_{\sigma}$ ,  $\Psi_{\sigma}^{\dagger}$ , while all other projectors are bosonic (i.e. commuting with the  $\Psi$ s). The first term  $H_{\text{tun}}^0$  commutes with  $\mathcal{P}_0$ , while the second is annihilated by it. So, if we call  $H_0 = H_{\text{dot}} + H_{\text{leads}} + H_{\text{tun}}^0$  the part of the Hamiltonian that commutes with  $\mathcal{P}_0$ , we have

$$\mathcal{L}_1\rho = \mathcal{P}_0(\mathcal{L}_{H_0} + \mathcal{L}_{H_{\text{tun}}^1})\mathcal{P}_0\rho = \mathcal{L}_{H_0}\mathcal{P}_0^2\rho + \mathcal{P}_0\mathcal{L}_{H_{\text{tun}}^1}\mathcal{P}_0\rho = \mathcal{L}_{H_0}\rho \quad (10)$$

where we used  $\rho = \mathcal{P}_0\rho$ . The Hamiltonian  $H_0$  is the Anderson impurity model with infinite dissipation quoted in Eq. (1) in the main text.

The second-order term  $\mathcal{L}_2$  yields the effective dissipation. Since  $\mathcal{Q}_0\mathcal{L}_0\mathcal{Q}_0 = -\gamma/2\mathcal{P}_1 - \gamma\mathcal{P}_2$ , we have  $(\mathcal{Q}_0\mathcal{L}_0\mathcal{Q}_0)^{-1} = -2/\gamma\mathcal{P}_1 - 1/\gamma\mathcal{P}_2$ . We evaluate Eq. (8b) starting from its rightmost terms:

$$(\mathcal{Q}_0\mathcal{L}_0\mathcal{Q}_0)^{-1}\mathcal{L}_H\mathcal{P}_0\rho = \left(-\frac{2}{\gamma}\mathcal{P}_1 - \frac{1}{\gamma}\mathcal{P}_2\right)(\mathcal{L}_{H_0} + \mathcal{L}_{H_{\text{tun}}^1})\rho = \frac{2}{\gamma}\mathcal{P}_1\mathcal{L}_{H_{\text{tun}}^1}\rho. \quad (11)$$

In the expression above, the terms proportional to  $\mathcal{L}_{H_0}\rho$  vanish because of the commutativity of  $\mathcal{L}_{H_0}$  with both projectors  $\mathcal{P}_{1,2}$ , which entails  $\mathcal{P}_{1,2}\mathcal{L}_{H_0}\mathcal{P}_0 = \mathcal{P}_{1,2}\mathcal{P}_0\mathcal{L}_{H_0} = 0$ . The term proportional to  $\mathcal{P}_2\mathcal{L}_{H_{\text{tun}}^1}\rho$  also vanishes because if  $\rho$  belongs to the slow subspace  $\mathcal{P}_0$ ,  $\mathcal{L}_{H_{\text{tun}}^1}\rho$  yields terms belonging to the  $\mathcal{P}_1$  subspace, but not to the  $\mathcal{P}_2$  one. Then

we expand

$$\begin{aligned}
\frac{2}{\gamma} \mathcal{P}_1 \mathcal{L}_{H_{\text{tun}}^1} \rho &= \frac{2i}{\gamma} (X_{<} H_{\text{tun}}^1 \rho X_d - X_{<} \rho H_{\text{tun}}^1 X_d + X_d H_{\text{tun}}^1 \rho X_{<} - X_d \rho H_{\text{tun}}^1 X_{<}) \\
&= \frac{2i}{\gamma} \sum_{\sigma} \sigma [\Psi_{\sigma}^{\dagger} X_{\bar{\sigma}d} \rho X_d - X_{<} \rho \Psi_{\sigma}^{\dagger} X_{\bar{\sigma}d} + X_{d\bar{\sigma}} \Psi_{\sigma} \rho X_{<} - X_d \rho X_{d\bar{\sigma}} \Psi_{\sigma}] \\
&= \frac{2i}{\gamma} \sum_{\sigma} \sigma [-X_{<} \rho \Psi_{\sigma}^{\dagger} X_{\bar{\sigma}d} + X_{d\bar{\sigma}} \Psi_{\sigma} \rho X_{<}] ,
\end{aligned}$$

where in the last equality we used  $X_{\alpha d} \rho X_{d\beta} = 0$  for  $\rho$  in the slow subspace. Then,

$$\begin{aligned}
\mathcal{L}_2 &= -\mathcal{P}_0 (\mathcal{L}_{H_0} + \mathcal{L}_{H_{\text{tun}}^1}) \mathcal{P}_1 \mathcal{L}_{H_{\text{tun}}^1} \rho = -\frac{2}{\gamma} \mathcal{L}_{H_0} \mathcal{P}_0 \mathcal{P}_1 \mathcal{L}_{H_{\text{tun}}^1} \rho - \frac{2}{\gamma} \mathcal{P}_0 \mathcal{L}_{H_{\text{tun}}^1} \mathcal{P}_1 \mathcal{L}_{H_{\text{tun}}^1} \rho = -\frac{2}{\gamma} \mathcal{P}_0 \mathcal{L}_{H_{\text{tun}}^1} \mathcal{P}_1 \mathcal{L}_{H_{\text{tun}}^1} \rho \\
&= -\frac{2}{\gamma} \mathcal{P}_0 \sum_{\sigma\tau} \sigma\tau (-2X_{d\bar{\tau}} \Psi_{\tau} \rho \Psi_{\sigma}^{\dagger} X_{\bar{\sigma}d} + X_{<} \rho \Psi_{\sigma}^{\dagger} X_{\bar{\sigma}\bar{\tau}} \Psi_{\tau} + \Psi_{\sigma}^{\dagger} X_{\bar{\sigma}\bar{\tau}} \Psi_{\tau} \rho X_{<}) \\
&= \frac{4}{\gamma} \sum_{\sigma\tau} \sigma\tau (X_{d\bar{\tau}} \Psi_{\tau} \rho \Psi_{\sigma}^{\dagger} X_{\bar{\sigma}d} - \frac{1}{2} \rho \Psi_{\sigma}^{\dagger} X_{\bar{\sigma}\bar{\tau}} \Psi_{\tau} - \frac{1}{2} \Psi_{\sigma}^{\dagger} X_{\bar{\sigma}\bar{\tau}} \Psi_{\tau} \rho) ,
\end{aligned}$$

where we recognize the dissipative part of the effective Lindblad superoperator,

$$\mathcal{L}_{\text{eff}} \rho = L_{\text{eff}} \rho L_{\text{eff}}^{\dagger} - \frac{1}{2} \{L_{\text{eff}}^{\dagger} L_{\text{eff}}, \rho\} \quad (12)$$

with

$$L_{\text{eff}} \equiv \frac{2}{\gamma^{1/2}} \sum_{\sigma} \sigma X_{0\bar{\sigma}} \Psi_{\sigma} = \frac{2}{\gamma^{1/2}} \sum_{p\alpha\sigma} \sigma V_{p\alpha} X_{0\bar{\sigma}} c_{p\sigma\alpha} , \quad (13)$$

which is the one quoted in the main text.

## DETAILS ON THE NON-CROSSING APPROXIMATION

In this Section, we provide analytical details of our implementation of the non-crossing approximation and on our real-time calculations.

The Anderson impurity model with infinite repulsion that represents the effective unitary dynamics of our dissipative model is not suitable to the usual perturbative techniques based on Green's functions, since the Hubbard operators  $X_{\alpha\beta}$  do not obey a standard fermionic or bosonic algebra and so their correlation functions cannot be extracted through Wick's theorem. So, we convert the effective Hamiltonian to a form that is suitable to standard perturbative treatments. This is achieved through the slave boson mapping of the Hubbard operators to a boson  $b$  and a spinful auxiliary fermion  $f_{\sigma}$  [1, 6, 7]:

$$\begin{aligned}
X_{\sigma 0} &= |\sigma\rangle\langle 0| = d_{\sigma}^{\dagger} (\mathbb{1} - n_{\bar{\sigma}}) = f_{\sigma}^{\dagger} b \\
X_{0\sigma} &= |0\rangle\langle \sigma| = (\mathbb{1} - n_{\bar{\sigma}}) d_{\sigma} = b^{\dagger} f_{\sigma} \\
X_{00} &= |0\rangle\langle 0| = (\mathbb{1} - n_{\uparrow})(\mathbb{1} - n_{\downarrow}) = b^{\dagger} b
\end{aligned} \quad (14)$$

with the physical subspace defined by  $Q \equiv b^{\dagger} b + \sum_{\sigma} f_{\sigma}^{\dagger} f_{\sigma} = 1$ . The physical meaning of the mapping is that the auxiliary fermions create single occupancies  $|\sigma\rangle = f_{\sigma}^{\dagger} |\text{vac}\rangle$ , while the bosons create the empty site  $|0\rangle = b^{\dagger} |\text{vac}\rangle$ , where  $|\text{vac}\rangle$  is the vacuum in the extended Hilbert space. The effective Hamiltonian becomes

$$H_{\text{eff}} = \sum_{\sigma} \varepsilon_{\sigma} f_{\sigma}^{\dagger} f_{\sigma} + \sum_{p\sigma} \varepsilon_p c_{p\sigma\alpha}^{\dagger} c_{p\sigma\alpha} + \sum_{p\sigma} V_p (b f_{\sigma}^{\dagger} c_{p\sigma\alpha} + \text{H.c.}) , \quad (15)$$

and the effective jump operator reads

$$L_{\text{eff}} = \frac{2}{\gamma^{1/2}} b^{\dagger} \sum_{p\sigma} \sigma V_p f_{\bar{\sigma}} c_{p\sigma\alpha} , \quad (16)$$




FIG. 1. Upper row: diagrammatic representation of the Luttinger-Ward functional truncated to the second order in the dot-leads hopping. The thick, aquamarine lines are the Green's functions of the auxiliary fermions, the dotted red lines are those of the auxiliary bosons and the dashed gray lines are the local ones of the leads. The lower rows of the figures depict the noninteracting fermionic and bosonic self-energies derived from  $\Phi$ . In each self-energy, the first diagram is the usual NCA self-energy coming from the effective AIM Hamiltonian, while the second term is the mean-field contribution from the effective dissipation.

corresponding to the Lindbladian dissipative action [8]:

$$\begin{aligned}
 S_d^{\text{eff}} = & -i \frac{4\eta}{\gamma} \int dt \sum_{\substack{p, p' \\ \sigma, \sigma'}} \sigma \sigma' V_p V_{p'} \left[ b_-(t) \bar{c}_{p\sigma-}(t) \bar{f}_{\bar{\sigma}-}(t) \bar{b}_+(t) f_{\bar{\sigma}'+}(t) c_{p'\sigma'+}(t) \right. \\
 & - \frac{1}{2} b_+(t) \bar{c}_{p\sigma+}(t) \bar{f}_{\bar{\sigma}+}(t) \bar{b}_+(t-0) f_{\bar{\sigma}'+}(t-0) c_{p'\sigma'+}(t-0) \\
 & \left. - \frac{1}{2} b_-(t) \bar{c}_{p\sigma-}(t) \bar{f}_{\bar{\sigma}-}(t) \bar{b}_-(t+0) f_{\bar{\sigma}'-}(t+0) c_{p'\sigma'-}(t+0) \right], \quad (17)
 \end{aligned}$$

where  $t \pm 0$  indicates an infinitesimal shift of the times that is necessary to correctly calculate tadpole contributions to the self-energies.

The non-crossing approximation (NCA) amounts to a self-consistent perturbation theory (i.e. a skeleton expansion of the self-energy) in the impurity-lead hopping  $V_p$  truncated at the leading order [4, 7, 9–13], namely the second. The residual dissipation needs to be included at the same order. Since the dissipative vertex Eq. (17) is already of order  $V_p^2$ , we only need to include it at the mean-field (i.e. tadpole) level. The resulting approximation to the self-energy is then conserving [9, 11–13]. The Luttinger-Ward functional [14] truncated to the second order in the impurity-bath hopping is shown in Fig. 1, along with the relevant self-energies.

The only nontrivial aspect of the slave boson mapping is that the constraint  $Q = 1$  has to be taken into account exactly. This can be done in a standard fashion [4, 9–13] by endowing the auxiliary particles with a fictitious chemical potential that is taken to infinity to extract the physical observables. Following Refs. [12, 13], we implement this constraint by discarding all terms that feature one or more lesser Green's function (bosonic or fermionic) in the retarded self-energy, and keeping at most one lesser Green's function in the expression of the lesser self-energy.

We are going to derive the equations describing the time evolution of the system prepared in a factorized state  $\rho_0 = \chi_d \otimes \rho_l$  between the impurity  $\chi_d$  and the leads  $\rho_l$ . For our purposes, the initial state of the impurity is completely characterized by the initial occupancy of the dot site  $n_\sigma^0 = \langle d_\sigma^\dagger d_\sigma \rangle_\chi$ , which coincides with the auxiliary fermion occupancy  $\langle f_\sigma^\dagger f_\sigma \rangle = \langle d_\sigma^\dagger d_\sigma \rangle$  (and determines the initial boson occupancy  $\langle b^\dagger b \rangle_\chi = 1 - \sum_\sigma n_\sigma^0$  through the constraint). The leads are assumed to be prepared in their own equilibrium states  $\rho_l = \bigotimes_\alpha \exp(-\beta H_\alpha)/Z_\alpha$ , where  $H_\alpha = \sum_{p\sigma\alpha} (\varepsilon_p - \mu_\alpha) c_{p\sigma\alpha}^\dagger c_{p\sigma\alpha}$  and  $Z_\alpha = \text{Tr} \exp(-\beta H_\alpha)$ . The subsequent evolution can be thought as a quench in



which the dot sites and the leads are initially disconnected (i.e.  $V_p = 0$ ) and suddenly put into contact at time  $t = 0$  by turning on the dot-leads hopping  $V_p$ . Hence, we will consider a time-dependent  $V_p(t) = \theta(t)V_p$ , which is necessary to set the correct limits in some integrals over time. The time dependence of the tunneling terms, i.e. the perturbation, makes all self energies  $\Sigma_{f\sigma}(t, t')$ ,  $\Pi(t, t')$  vanish whenever either of their time arguments is negative. Thus, we will always consider  $t \geq 0$ ,  $t' \geq 0$  in the following equations, and we will consequently omit the  $\theta$  functions in front of the self-energies.

The contour-ordered NCA self-energies associated to the unitary part of the dynamics of the auxiliary particles  $f_\sigma$  and  $b$  are given by [4, 7, 9–13]

$$\begin{aligned}\Sigma_{f\sigma}(t, t') &= i\Delta_\sigma(t, t')B(t, t') \quad \text{for auxiliary fermions,} \\ \Pi(t, t') &= -i \sum_\sigma \Delta_\sigma(t', t)G_\sigma(t, t') \quad \text{for bosons}\end{aligned}\tag{18}$$

where  $G_\sigma(t, t') \equiv -i \langle \mathcal{T} f_\sigma(t) f_\sigma^\dagger(t') \rangle$  and  $B(t, t') \equiv -i \langle \mathcal{T} b(t) b^\dagger(t') \rangle$  are the contour-ordered Green's functions for the auxiliary fermions and the bosons, respectively ( $\mathcal{T}$  is the contour-ordering symbol). The function  $\Delta_\sigma(t, t')$  is the local Green's function of the leads, namely  $\Delta_\sigma(t, t') \equiv \sum_{p, \alpha, p', \alpha'} V_p(t) g_{p\sigma\alpha, p'\sigma\alpha'}(t, t') V_p(t')$ , where  $g_{p\sigma\alpha, p'\sigma\alpha'}(t, t') \equiv -i \langle \mathcal{T} c_{p\sigma\alpha}(t) c_{p'\sigma\alpha'}^\dagger(t') \rangle$ . Applying Langreth's rules [15] and the projection onto  $Q = 1$  we obtain the various components of the self-energies:

$$\begin{aligned}\Sigma_{f\sigma}^\geq(t, t') &= i\Delta_\sigma^\geq(t - t')B^\geq(t, t') , \\ \Sigma_{f\sigma}^{R,A}(t, t') &= i\Delta_\sigma^\geq(t - t')B^{R,A}(t, t')\end{aligned}\tag{19}$$

and

$$\begin{aligned}\Pi^\geq(t, t') &= -i \sum_\sigma \Delta_\sigma^\leq(t' - t)G_\sigma^\geq(t, t') , \\ \Pi^{R,A}(t, t') &= -i \sum_\sigma \Delta_\sigma^\leq(t' - t)G_\sigma^{R,A}(t, t') .\end{aligned}\tag{20}$$

The projection on the physical subspace has two effects on the self-energies. The first is that some contributions are discarded, while the second is that the local Green's function  $\Delta_\sigma(t, t')$  has to be substituted by its *unperturbed* version  $\Delta_\sigma(t, t')$ . This occurs because any correction to  $\Delta_\sigma(t, t')$  must contain at least one lesser Green's function of the fermions or the bosons [12, 13], contrary to the projection rule. This substitution is not an approximation, and it does not imply that the leads are not affected by the impurity. In fact, we will show in the section devoted to transport that the knowledge of the Green's functions of the auxiliary particles is sufficient to compute  $g_{p\sigma\alpha, p'\sigma\alpha'}(t, t')$ .

Since the unperturbed leads are assumed to be in thermodynamic equilibrium, we have [remembering  $V_p(t) = \theta(t)V_p$ ]  $\Delta_\sigma(t, t') \equiv \theta(t)\theta(t')\Delta_\sigma(t-t')$  and we can express  $\Delta_\sigma(t-t')$  through its Fourier transform  $\Delta_\sigma(\omega) = \int dt e^{i\omega^+(t-t')} \Delta_\sigma(t-t')$  (where  $\omega^\pm \equiv \omega \pm i0$  denotes an infinitesimal shift along the imaginary axis):  $\Delta_\sigma(\omega) = \sum_\alpha \Delta_{\sigma\alpha}(\omega)$  where

$$\Delta_{\sigma\alpha}^{R,A}(\omega) = \int \frac{d\varepsilon}{2\pi} \frac{\Gamma_\alpha(\varepsilon)}{\omega^\pm - \varepsilon} = \mathcal{P} \int \frac{d\varepsilon}{2\pi} \frac{\Gamma_\alpha(\varepsilon)}{\omega - \varepsilon} \mp \frac{i}{2} \Gamma_\alpha(\omega)\tag{21a}$$

$$\Delta_{\sigma\alpha}^<(\omega) = i\Gamma_\alpha(\omega)F_\alpha(\omega)\tag{21b}$$

$$\Delta_{\sigma\alpha}^>(\omega) = -i\Gamma_\alpha(\omega)[1 - F_\alpha(\omega)]\tag{21c}$$

We have introduced the level width function  $\Gamma_\alpha(\omega) \equiv 2\pi \sum_p V_{p\alpha}^2 \delta(\omega - \varepsilon_p)$  and the Fermi distribution  $F_\alpha(\omega) \equiv \{\exp[\beta(\omega - \mu_\alpha)] + 1\}^{-1}$  of the lead  $\alpha$ , with inverse temperature  $\beta$  [16]. Notice that we keep the label  $\sigma$  for the sake of generality, although we never consider the presence of magnetic fields or initially spin-unbalanced leads that could lead to an explicit spin dependence of  $\Delta$ . Also, we temporarily keep the label  $\alpha$  on  $\Gamma_\alpha$ , although we will always consider two identical baths with  $\Gamma_L = \Gamma_R$ .

In our calculations, we assume zero temperature  $\beta \rightarrow \infty$  and we take a flat density of states  $\Gamma(\omega) = \Gamma\xi(\omega)$ , with  $\xi(\omega) = \theta(W - |\omega|)$ , for which

$$\begin{aligned}\Delta_{\sigma\alpha}^<(t) &= -\frac{\Gamma}{2\pi} \frac{e^{-i\mu_\alpha t} - e^{iWt}}{t} \\ \Delta_{\sigma\alpha}^>(t) &= \frac{\Gamma}{2\pi} \frac{e^{-iWt} - e^{-i\mu_\alpha t}}{t} .\end{aligned}\tag{22}$$

The dissipative contributions to the retarded self-energies are

$$\begin{aligned}\Sigma_{f\sigma}^{\text{diss},R}(t,t') &= -i\frac{2}{\gamma}B^>(t,t)\Delta_{\sigma}^<(t,t)\delta(t-t') \\ &\stackrel{\text{proj.}}{\rightarrow} -i\frac{2}{\gamma}[B^>(t,t) - B^<(t,t)]\Delta_{\sigma}^<(t,t)\delta(t-t') = -\frac{2}{\gamma}\Delta_{\sigma}^<(t,t)\delta(t-t') \equiv -\frac{i}{2}\kappa_{\sigma}\delta(t-t') \\ \Pi^{\text{diss},R}(t,t') &= -\frac{2i}{\gamma}\sum_{\sigma}\Delta_{\sigma}^<(t,t)G_{\sigma}^<(t,t)\delta(t-t') \stackrel{\text{proj.}}{\rightarrow} 0\end{aligned}\quad (23)$$

where  $\stackrel{\text{proj.}}{\rightarrow}$  stands for the projection procedure of discarding all lesser fermionic or bosonic Green's functions, mentioned in the previous paragraphs. In the above Equations we introduced the effective dissipation rate

$$\kappa_{\sigma} \equiv -i\frac{4}{\gamma}\Delta_{\sigma}^<(t,t) = \sum_{\alpha}\frac{2\Gamma_{\alpha}}{\pi\gamma}\int_{-\infty}^{\infty}d\omega\xi(\omega)F_{\alpha}(\omega), \quad (24)$$

where  $\xi(\omega)$  is the cutoff function defined by  $\Gamma_{\alpha}(\omega) = \Gamma_{\alpha}\xi(\omega)$ . For a flat level width function  $\xi(\omega) = \theta(W - |\omega|)$  we have

$$\kappa_{\sigma} = 2\sum_{\alpha}\frac{\Gamma_{\alpha}}{\pi\gamma}(\mu_{\alpha} + W), \quad (25)$$

while for a general shape  $\xi(\omega)$  we can estimate  $\kappa \sim \Gamma/\gamma \cdot W$ . Thus, the effective loss rate depends on the full band-shape of the leads, and grows with the bandwidth.

The lesser self-energies read

$$\begin{aligned}\Sigma_{f\sigma}^{\text{diss},<}(t,t') &= 0 \\ \Pi^{\text{diss},<}(t,t') &= \frac{4i}{\gamma}\Delta_{\sigma}^<(t,t)G_{\sigma}^<(t,t)\delta(t-t') \stackrel{\text{proj.}}{\rightarrow} \frac{4i}{\gamma}\sum_{\sigma}\Delta_{\sigma}^<(t,t)G_{\sigma}^<(t,t)\delta(t-t') = -\sum_{\sigma}\kappa_{\sigma}G_{\sigma}^<(t,t)\delta(t-t')\end{aligned}\quad (26)$$

As anticipated, the leading order self-energies from the effective interaction are mean-field-like corrections, because they are local in time.

*Dyson equations* Using the self-energies written above, we can write down the Dyson equations determining the dynamics of the retarded Green's functions

$$\begin{aligned}(i\partial_t - \varepsilon_{\sigma})G_{\sigma}^R(t,t') &= \delta(t-t') - \frac{i}{2}\kappa_{\sigma}G_{\sigma}^R(t,t') + i\int_{t'}^t d\bar{t}\Delta_{\sigma}^>(t-\bar{t})B^R(t,\bar{t})G_{\sigma}^R(\bar{t},t'), \\ (-i\partial_{t'} - \varepsilon_{\sigma})G_{\sigma}^R(t,t') &= \delta(t-t') - \frac{i}{2}\kappa_{\sigma}G_{\sigma}^R(t,t') + i\int_{t'}^t d\bar{t}G_{\sigma}^R(t,\bar{t})\Delta_{\sigma}^>(\bar{t}-t')B^R(\bar{t},t')\end{aligned}\quad (27)$$

for the auxiliary fermions, and

$$\begin{aligned}i\partial_t B^R(t,t') &= \delta(t-t') - i\sum_{\sigma}\int_{t'}^t d\bar{t}\Delta_{\sigma}^<(\bar{t}-t)G_{\sigma}^R(t,\bar{t})B^R(\bar{t},t'), \\ -i\partial_{t'} B^R(t,t') &= \delta(t-t') - i\sum_{\sigma}\int_{t'}^t d\bar{t}B^R(t,\bar{t})\Delta_{\sigma}^<(t'-\bar{t})G_{\sigma}^R(\bar{t},t')\end{aligned}\quad (28)$$

for the bosons. The advanced Green's functions can be obtained as  $G^A(t,t') = [G^R(t',t)]^*$ , with  $G = G_{\sigma}$  or  $B$ . The boundary conditions for the Dyson equations are  $G_{\sigma}^R(t+0,t) = B^R(t+0,t) = -i$ , which simply enforces the (anti-)commutation relations. A peculiar behavior of the Dyson equations in the NCA is that they only involve retarded functions, which greatly simplifies their numerical solution. In particular, the retarded Green's function that are obtained are time-translational invariant:  $G_{\sigma}^R(t,t') \equiv G_{\sigma}^R(t-t')$ ,  $B^R(t,t') \equiv B^R(t-t')$ . For  $\kappa = 0$  we obtain the same solution as in equilibrium. If this behavior might seem unphysical, let us recall that it regards the dynamics of two unphysical, auxiliary particles  $f_{\sigma}$  and  $b$  of which the physical fermion  $d_{\sigma} = b^{\dagger}f_{\sigma}$  [cf. Eqs. (14)] is “composed”. Indeed, the object that we aim to compute is the physical Green's function for the  $d_{\sigma}$  operators, which in the language of the slave boson representation is a four-point vertex. For this object, the time evolution of the retarded and lesser components are intertwined.

*Kinetic equations* The lesser components of the auxiliary fermion Green's functions are determined by

$$\begin{aligned} (i\partial_t - \varepsilon_\sigma)G_\sigma^<(t, t') &= -\frac{i}{2}\kappa_\sigma G_\sigma^<(t, t') + i[(\Delta_\sigma^> B^R) * G_\sigma^<](t, t') + i[\Delta_\sigma^< B^< * G_\sigma^A](t, t') , \\ (-i\partial_{t'} - \varepsilon_\sigma)G_\sigma^<(t, t') &= +\frac{i}{2}\kappa_\sigma G_\sigma^<(t, t') + [G_\sigma^R * i\Delta_\sigma^< B^<](t, t') + [G_\sigma^< * i\Delta_\sigma^> B^A](t, t') \end{aligned} \quad (29)$$

the above Equations have been written in a compact notation that treats the Green's functions as matrices in the two time indices. We have introduced two different “products” for the Green's functions: a direct product  $(AB)(t, t') = A(t, t')B(t, t')$  and the time convolution (i.e. the matrix product)  $(A * B)(t, t') \equiv \int d\bar{t} A(t, \bar{t})B(\bar{t}, t')$ . The time evolution of the diagonal component reads

$$\begin{aligned} \text{id}_t G_\sigma^<(t, t) &= [i\partial_t G_\sigma^<(t, t') + i\partial_{t'} G_\sigma^<(t, t')]|_{t'=t} = \\ &= -i\kappa_\sigma G_\sigma^<(t, t) + 2 \text{Re} \left\{ i \int_0^t d\bar{t} \left[ \Delta_\sigma^<(t - \bar{t}) B^<(t, \bar{t}) G_\sigma^A(\bar{t}, t) + \Delta_\sigma^>(t - \bar{t}) B^R(t, \bar{t}) G_\sigma^<(\bar{t}, t) \right] \right\} \end{aligned} \quad (30)$$

The boundary conditions are simply the initial dot populations:  $G_\sigma(0, 0) = n_{d\sigma}(0)$ .

For the bosons we similarly obtain

$$\begin{aligned} i\partial_t B^<(t, t') &= -\sum_\sigma \kappa_\sigma G_\sigma^<(t, t) B^A(t - t') \\ &\quad - i \sum_\sigma [(\Delta_\sigma^<)^T G_\sigma^R * B^<](t, t') - i \sum_\sigma [(\Delta_\sigma^>)^T G_\sigma^< * B^A](t, t') , \\ -i\partial_{t'} B^<(t, t') &= -\sum_\sigma \kappa_\sigma G_\sigma^<(t, t) B^R(t - t') \\ &\quad - [B^R * i \sum_\sigma (\Delta_\sigma^>)^T G_\sigma^<](t, t') - [B^< * i \sum_\sigma (\Delta_\sigma^<)^T G_\sigma^A](t, t') \end{aligned} \quad (31)$$

where we employ the notation  $(\Delta^T)(t, t') = \Delta(t', t)$ . To find the equation determining the evolution of the diagonal component one uses  $\lim_{t' \rightarrow t} [B^A(t, t') - B^R(t, t')] = \lim_{t' \rightarrow t} [B^<(t, t') - B^>(t, t')] = -i[b^\dagger, b] = i$  and finds

$$\begin{aligned} \text{id}_t B^<(t, t) &= -i \sum_\sigma \kappa_\sigma G_\sigma^<(t, t) \\ &\quad + 2 \text{Re} \left\{ i \sum_\sigma \int_0^t d\bar{t} \left[ \Delta_\sigma^<(t - \bar{t}) B^<(t, \bar{t}) G_\sigma^A(\bar{t}, t) + \Delta_\sigma^>(t - \bar{t}) B^R(t, \bar{t}) G_\sigma^<(\bar{t}, t) \right] \right\} . \end{aligned} \quad (32)$$

It is easy to observe that the constraint is explicitly conserved:  $d_t Q(t) = \text{id}_t B^<(t, t) - \sum_\sigma \text{id}_t G_\sigma^<(t, t) = 0$ . So, if we take the appropriate initial condition  $B(0, 0) = -i[1 - \sum_\sigma n_{d\sigma}(0)]$ , we will always keep  $Q = 1$  during the dynamics, within the numerical accuracy of our computations.

*Numerical implementation* We solve the Equations above using the simple algorithm described in Ref. [13]. Conceptually, we have equations in the form  $\frac{d}{dt}f(t) = K[f](t)$ , where the right-hand side depends functionally on  $f$ . We first convert the equation to an integral one to obtain one integration step:  $\int_t^{t+\delta t} dt' \frac{d}{dt'} f(t') = f(t + \delta t) - f(t) = \int_t^{t+\delta t} dt' K[f](t')$ . The integrals on the right-hand side (i.e. the one for the time step and the one implied in  $K$ ) are computed by any quadrature rule. In our case, the simple trapezoidal rule with equally spaced points suffices. This choice leads to clearer equations, without sacrificing too much the accuracy (at the cost of needing a  $\delta t$  which cannot be too large). A key point is that the update rule for  $f(t + \delta t)$  is implicit, since  $f(t + \delta t)$  appears also in the discretization of the right-hand side. This makes the integration numerically stable. A nice feature of the implementation of Ref. [13] is that  $Q$  is exactly conserved also in the discretized dynamics, which implies that the deviations from 1 will be comparable by machine precision  $\sim 10^{-14 \div 15}$ .

The Dyson Eqs. (27) and (28) are integrated first, since they do not depend on the lesser functions. The equations for the fermionic and bosonic Green's functions are coupled, though, and need to be solved together. We only integrate the forward equations  $i\partial_t G_\sigma(t - t') = \dots$ , etc., and we take advantage of the time-translational invariance of the retarded Green's functions. The retarded functions thus obtained are then used to compute the lesser functions. The causal properties of the equations implies that we need to follow the path depicted in Fig. 2 on the discretized time grid  $(t, t')$ . Since lesser Green's functions are anti-Hermitian matrices in the time indices [15, 17], we compute only their values for  $t \leq t'$ .




FIG. 2. Sketch of the steps for the numerical integration for the determination of the lesser Green's functions. The first square on the diagonal is the initial condition  $G_{\sigma}^{<}(0, 0) = in_{\sigma}^0$ ,  $B^{<}(0, 0) = 1 - \sum_{\sigma} n_{\sigma}^0$ , and the subsequent integrations proceed along the columns, from  $t' = 0$  to  $t' = t$  (thick red arrows). The steps corresponding to the light gray squares make use of Eqs. (29) and (31), while the diagonal elements in darker gray employ Eqs. (30) and (32).

In most of our calculations we set the time step to  $\delta t = 0.5W^{-1}$ . We have verified that doubling or halving this value leads to differences of at most a few per mil in Green's functions and currents. The value of the observables after the initial transient, i.e. for times larger than a few  $\Gamma^{-1}$ , are less susceptible to changes of  $\delta t$ —a possible manifestation of the dynamically attractive nature of the local stationary state.

### Physical Green's function

In the subspace with no double occupancies  $d_{\sigma} = X_{0\sigma} = b^{\dagger}f_{\sigma}$ . Hence

$$G_{d\sigma}(t, t') = -i \langle \mathcal{T}_C d_{\sigma}(t) d_{\sigma}^{\dagger}(t') \rangle = -i \langle \mathcal{T}_C f_{\sigma}(t) b^{\dagger}(t) b(t') f_{\sigma}^{\dagger}(t') \rangle . \quad (33)$$

Within the non-crossing approximation [4, 6, 7, 9–13] one simply decouples the fermions from the bosons (i.e. one ignores vertices):

$$G_{d\sigma}(t, t') \approx iG_{\sigma}(t, t')B(t', t) . \quad (34)$$

This approximation can be obtained as the leading term in a large- $N$  expansion in the number of flavors of fermions (with the rescaling  $V_p \rightarrow V_P/N^{1/2}$ ). While the auxiliary particles' Green functions are approximated with a good accuracy by the NCA equations derived above, the neglect of vertices in the physical Green's function does introduce some spurious effects [18]. Nevertheless, it is known to provide the correct qualitative properties of the Kondo effect, with quantitative discrepancies of the order of 15% for observables such as the zero-bias conductance [10]. We obtain the lesser function as follows:

$$G_{d\sigma}^{<}(t, t') \approx iG_{\sigma}^{<}(t, t')B^{>}(t', t) \xrightarrow{\text{proj.}} iG_{\sigma}^{<}(t, t')[B^{>}(t', t) - B^{<}(t', t)] = G_{\sigma}^{<}(t, t')b(t', t) , \quad (35)$$

where the first approximate equality is the NCA, while the second equality is the projection onto the physical subspace  $Q = 1$ . We have introduced the functions

$$\begin{aligned} g_{\sigma}(t, t') &\equiv i[G_{\sigma}^{>}(t, t') - G_{\sigma}^{<}(t, t')] \\ b_{\sigma}(t, t') &\equiv i[B^{>}(t, t') - B^{<}(t, t')] \end{aligned} \quad (36)$$

so that

$$\begin{aligned} G_{\sigma}^R(t, t') &= -i\theta(t - t')g_{\sigma}(t - t') \\ G_{\sigma}^A(t, t') &= +i\theta(t' - t)g_{\sigma}(t, t') \end{aligned} \quad (37)$$



and analogously for the bosons. In particular,  $b(t, t) = 1$ , so  $\langle d_\sigma(t)^\dagger d_\sigma(t) \rangle \equiv -iG_{d\sigma}^<(t, t) = -iG_\sigma^<(t, t) = \langle f_\sigma(t)^\dagger f_\sigma(t) \rangle$ , where the equality  $G_{d\sigma}^<(t, t) = G_\sigma^<(t, t)$  is exact. Similarly,

$$G_{d\sigma}^>(t, t') \approx iG_\sigma^>(t, t')B^<(t', t) \xrightarrow{\text{proj.}} i[G_\sigma^>(t, t') - G_\sigma^<(t, t')]B^<(t', t) = g_\sigma(t, t')B^<(t', t) , \quad (38)$$

and finally

$$G_{d\sigma}^R(t, t') \equiv \theta(t - t')[G_{d\sigma}^>(t, t') - G_{d\sigma}^<(t, t')] = i[G_\sigma^R(t, t')B^<(t', t) + G_\sigma^<(t, t')B^A(t', t)] \quad (39)$$

At first sight, there appears to be a problem with the anticommutation relations  $\{d_\sigma, d_\sigma^\dagger\} = \mathbb{1}$ :  $G_{d\sigma}^>(t, t) = B^<(t, t) = -i + \sum_\sigma G_\sigma^<(t, t) \neq -i + G_\sigma^<(t, t)$ , therefore

$$G_{d\sigma}^R(t + 0, t) = B^<(t, t) - G_\sigma^<(t, t) = -i(1 - G_\sigma^<(t, t)) \neq -i\{d_\sigma, d_\sigma^\dagger\} = -i \quad (40)$$

This is not a shortcoming of the NCA, but rather of the Hilbert space truncation  $d_\sigma \rightarrow X_{0\sigma}$ . Indeed,  $-i\{X_{0\sigma}, X_{\sigma 0}\} = -i(X_{00} + X_{\sigma\sigma}) \neq \mathbb{1}$ . This modified anticommutation relation means that the spectral function will not integrate to 1, but rather to  $i(B^<(t, t) - G_\sigma^<(t, t)) = 1 - n_{-\sigma}(t)$ . The missing spectral weight is that of double occupancies, which would appear as a very broad and low peak at  $\omega = \varepsilon_d$ , with width  $\sim \gamma$  and height  $\sim \gamma^{-1}$ .

In the main text, we show the dot spectral function in the local stationary state. This function is defined as  $A_\sigma(\omega, t) = -\text{Im } G_{d\sigma}^R(\omega, t)/\pi$ , where  $G_{d\sigma}^R(\omega, t) \equiv \int_0^\infty d\tau e^{i\omega\tau} G_{d\sigma}^R(t + \tau, t)$ , choosing  $t \gtrsim 5\Gamma_T^{-1}$  such that the  $A_\sigma(\omega, t)$  has already saturated to its late-time value.

## Transport

As usual in impurity problems, [15] the Green's function of the leads  $g_{p\sigma\alpha, p'\sigma\alpha'}(t, t')$  is determined by a T-matrix equation

$$\hat{g} = \hat{g}_0 + \hat{g}_0 * \hat{P} * \hat{g}_0 , \quad (41)$$

where the hat indicates a matrix in  $p, \sigma, \alpha$  and times, and with the convolution symbol  $*$  representing a matrix multiplication in all of the above indices. The T-matrix  $\hat{P}$  on the contour can be obtained by deriving the Luttinger-Ward functional (Fig. 1) and reads

$$P_{p\sigma\alpha, p'\sigma'\alpha'}(t, t') = V_{p\alpha} V_{p'\alpha'} \delta_{\sigma\sigma'} \left[ iG_\sigma(t, t')B(t', t) + \frac{4}{\gamma} G_\sigma^<(t, t)\delta(t - t') \right] . \quad (42)$$

When applying Langreth's rules to determine the various components of  $\hat{P}$  we must apply the projection [12, 13] on the physical subspace  $Q = 1$ . In this case there is a subtlety: the rule of keeping at most one auxiliary bosonic or fermionic lesser function applies only if the projected quantity vanishes in the unphysical  $Q = 0$  space. This is not the case for  $\hat{g}$ , since  $\hat{g}_{Q=0} = \hat{g}_0 \neq 0$ . Therefore, the projection must be done on  $\hat{g} - \hat{g}_0$ , and we obtain that  $P = \mathcal{O}(G_\sigma^<, B^<)$  for all components. We can write the results in terms of a new function  $Q_\sigma(t, t')$  (not to be confused with the charge  $Q$ ) defined by

$$P_{p\sigma\alpha, p'\sigma'\alpha'}(t, t') = V_{p\alpha} V_{p'\alpha'} \delta_{\sigma\sigma'} Q_\sigma(t, t') . \quad (43)$$

Then, within the NCA we have  $Q_\sigma^{R, <}(t, t') = G_{d\sigma}^{R, <}(t, t') + Q_\sigma^{\text{diss}, R, <}(t)\delta(t - t')$ , with

$$\begin{aligned} Q_\sigma^{\text{diss}, R}(t) &= -\frac{2}{\gamma} G_\sigma^<(t, t) = -\frac{2i}{\gamma} n_\sigma(t) \\ Q_\sigma^{\text{diss}, <}(t) &= 0 \end{aligned} \quad (44)$$

The current leaving reservoir  $\alpha$  is defined as

$$I_\alpha \equiv -\frac{d}{dt} \sum_{p\sigma} \langle c_{p\sigma\alpha}^\dagger c_{p\sigma\alpha} \rangle_t = -\frac{d}{dt} \sum_{p\sigma} g_{p\sigma\alpha, p\sigma\alpha}^<(t, t) . \quad (45)$$

With the help of the appropriate T-matrix equation  $\hat{g}^< = \hat{g}_0^< + \hat{g}_0^< * \hat{P}^A * \hat{g}_0^A + \hat{g}_0^R * \hat{P}^< * \hat{g}_0^A + \hat{g}_0^R * \hat{P}^R * \hat{g}_0^<$  and the NCA expression for  $\hat{P}$  and  $\hat{Q}$  we find

$$\begin{aligned} I_\alpha(t) &= 2 \operatorname{Re} \int_0^t d\bar{t} \sum_\sigma \left[ Q_\sigma^R(t, \bar{t}) \Delta_{\sigma\alpha}^<(\bar{t} - t) + Q_\sigma^<(t, \bar{t}) \Delta_{\sigma\alpha}^A(\bar{t} - t) \right] = \\ &= 2 \operatorname{Re} \int_0^t d\bar{t} \sum_\sigma \left[ G_{d\sigma}^R(t, \bar{t}) \Delta_{\sigma\alpha}^<(\bar{t} - t) + G_{d\sigma}^<(t, \bar{t}) \Delta_{\sigma\alpha}^A(\bar{t} - t) \right] + \frac{4}{\gamma} \sum_\sigma n_\sigma(t) (-i) \Delta_{\sigma\alpha}^<(0) . \end{aligned} \quad (46)$$

The first term in the second line of the above Equation is the usual starting point of the Meir-Wingreen formula [15, 19], whereas the second term represents a new, dissipative contribution. We will show later that this latter contribution is related to the loss of particles from the system—a conclusion that can be anticipated by its proportionality to the dot occupancy. The loss term is proportional to the Green's function of the leads evaluated at coinciding times,  $-i\Delta_{\sigma\alpha}^<(0)$ , and is therefore non-universal, in the sense that it is sensitive to the full band shape of the leads,  $\xi(\omega)$ . For an infinite bandwidth  $W$ , this term would diverge [20].

We can then easily derive the conductance at zero bias, defined as  $g(0, t) \equiv \lim_{\Delta\mu \rightarrow 0} dI(t)/d\Delta\mu$ , where the transport current is  $I(t) \equiv (I_L(t) - I_R(t))/2$ , assuming that the left reservoir is at a higher chemical potential,  $\mu_{L,R} = \mu \pm \Delta\mu/2$ . Using

$$\frac{d}{d\Delta\mu} F_{L,R}(\omega) = \frac{d}{d\Delta\mu} F\left[\omega - \left(\mu \pm \frac{\Delta\mu}{2}\right)\right] = \mp \frac{1}{2} F'(\omega - \mu_\alpha) \quad (47)$$

in the expression for  $\Delta_{\sigma\alpha}^<$  (with the prime  $F'$  indicating the derivative), we obtain at zero temperature

$$g(0, t) = -\frac{\Gamma(\mu)}{2} \sum_\sigma \frac{1}{\pi} \operatorname{Im} \int_0^t d\bar{t} e^{i\mu\bar{t}} G_{d\sigma}^R(t, t - \bar{t}) + \frac{\Gamma(\mu)}{\pi\gamma} n_d(t) , \quad (48)$$

which is Eq. (2) quoted in the main text. As noted there, the first term directly probes the spectral function at the chemical potential, and at later times converges to the usual stationary formula [15, 19]  $g_\infty(0) = 2^{-1} \Gamma(\mu) \sum_\sigma A_\sigma(\mu)$ . Thus, the zero-bias conductance allows to detect directly the presence of the Kondo peak. Moreover, it is interesting to notice that the first term of Eq. (48) only probes the spectral function at the chemical potential of the leads, thus displaying universal behavior in the usual Hamiltonian case. On the other hand, the correction represented by the second term explicitly depends on the dot population  $n_d(t)$ , which is non-universal (i.e. explicitly dependent on the cutoff  $W$  and band shape  $\xi(\omega)$ ). On the other hand, it does not diverge for an infinite bandwidth, unlike the individual currents  $I_\alpha(t)$ . Indeed, the “loss” terms of the currents, i.e. the last term of (46), partially cancel out in the transported current even for a finite bias:

$$\frac{2}{\gamma} \sum_\sigma n_\sigma(t) (-i) [\Delta_{\sigma L}^<(0) - \Delta_{\sigma R}^<(0)] = \frac{2}{\gamma} n_d(t) \int \frac{d\omega}{2\pi} \Gamma(\omega) [F_L(\omega) - F_R(\omega)] = \frac{2}{\gamma} n_d(t) \int_{\mu_R}^{\mu_L} \frac{d\omega}{2\pi} \Gamma(\omega) , \quad (49)$$

where the last equality applies to zero temperature. In words, the losses contribute to the transport current only through the lead fermions present between the two chemical potentials. We need to remark that the loss correction to the zero-bias conductance is tiny with respect to the transport term (i.e. the first one of Eq. (48)): while the latter is of order  $10^{-1 \div 0}$ , the former is at most (for the maximal  $n_d(t) = 1$ )  $\Gamma/(\pi\gamma)$ , which in our calculation is always less than  $3 \cdot 10^{-3}$ . Thus, almost all of the effect of the residual dissipation affects the conductance through the impurity spectral function, and in particular the decrease of the height of the Kondo peak for decreasing dissipation rate.

It is interesting to compare the behavior of the transported current with the current of particles lost from the system,  $I_{\text{loss}}(t) = -d[N_R(t) + N_L(t) + n_d(t)]/dt = I_R(t) + I_L(t) - dn_d(t)/dt$ . Using the expressions (46), (30) and (39) we find

$$I_{\text{loss}}(t) = \frac{4}{\gamma} \sum_\sigma n_\sigma(t) (-i) \Delta_\sigma^<(0) = 2 \sum_\sigma \kappa_\sigma n_\sigma(t) . \quad (50)$$

As a check, we notice that in the limit  $\gamma \rightarrow +\infty$   $\kappa_\sigma$  vanishes and we recover the statement of conservation of the number of particles:  $I_{\text{loss}} = 0$ . In general, after a transient of a few  $\Gamma^{-1}$  the dot population saturates, and for a finite  $\gamma$  the loss current saturates to  $I_{\text{loss}}(t) \rightarrow I_{\text{loss}}^\infty = 2 \sum_\sigma \kappa_\sigma n_\sigma^\infty$ . As it could be expected on classical grounds, the stationary loss current is proportional to the dot population and to the density of leads fermions at the dot site, since  $\kappa_\sigma \propto \Delta_\sigma^<(0)$ . The factor of 2 accounts for the fact that each loss event entails the disappearance of two particles,




FIG. 3. Loss of scaling collapse in the nonlinear conductance in presence of a finite dissipation. The data are for  $\Gamma_T = 0.02W$ ,  $T_K = (2\Gamma_T W) \exp(\pi\varepsilon_d/\Gamma_T)$ . The full curves are for the AIM at  $\gamma \rightarrow +\infty$ , the dashed curves are for  $\gamma = 10W$  and the dotted curves are for  $\gamma = W$ . Decreasing  $\gamma$ , besides decreasing the conductance itself, spoils the data collapse as a function of  $\Delta\mu/T_K$ .

one from the dot and one from the leads. In the large- $\gamma$  regime that we are analyzing, the losses are suppressed by an explicit factor of  $\gamma$  (coming from  $\kappa_\sigma$ ), besides of the slow increase of  $n_d^\infty$  with  $\gamma$  (see later). As we have already remarked, the dot population does not show universal behavior, hence the loss current does not provide a good probe of the Kondo effect emerging at large dissipation.

### ADDITIONAL DATA

In this Section, we provide additional data on the relation between the loss rate  $\gamma$  and the presence of typical signatures of the Kondo effect.

In Fig. 3 we show the long-time nonlinear conductance  $g_\infty = \lim_{t \rightarrow \infty} I(t)/\Delta\mu$  as a function of the ratio  $\Delta\mu/T_K$ . A clear signature of the Kondo effect for  $\gamma \rightarrow +\infty$  (continuous lines) is that the curves from different dot energies  $\varepsilon_d$  collapse on a universal curve (except those in the mixed-valence regime  $\varepsilon_d \gtrsim -\Gamma_T$ ) [2, 4, 10, 11, 21]. This scaling collapse signals the presence of  $T_K$  as the only energy scale governing the low-energy transport properties, and can be used to detect the Kondo effect. In the Figure, dashed and dotted lines refer to the finite-dissipation cases of  $\gamma = 10W$  and  $\gamma = W$ , respectively. We observe that in this regime the scaling collapse is lost, since the new competing energy scale  $\kappa_\sigma$  emerges. Especially for the smaller  $\gamma = W$ , the conductance becomes scarcely dependent on the bias  $\Delta\mu$ . From the central panel of Fig. (1) in the main text, this mild dependence can be attributed to the complete loss of the Kondo peak in the spectral function, meaning that the dot behaves simply as a noninteracting dot, which has its maximal sensitivity to the bias only for  $\mu \approx \varepsilon_d$ .

In Fig. 4 we show the typical appearance of the spin decay in the mixed-valence regime  $\varepsilon_d = -\Gamma_T$  (left) and in the Kondo regime  $\varepsilon_d = -2\Gamma_T$  (right). While for  $|\varepsilon_d| \lesssim \Gamma_T$  the decay is slower than exponential, for deeper dot levels a purely exponential form is recovered. Regardless of the form of the decay, the effect of the two-body loss is to suppress the equilibration of the impurity spin, down to quite a small value. This suppression signals the crossover to the strongly dissipated regime in which the physics is dominated by the Kondo resonance (which corresponds precisely to a long-lived spin state [1]). The curves in the rightmost panel of Fig. (1) in the main text are obtained via a linear fit of  $\log \langle \sigma_d^z(t) \rangle$  versus time, excluding the initial data for  $t \leq 10\Gamma_T^{-1}$ . According to the literature [22–24], the exponential decay rate of  $\langle \sigma_d^z(t) \rangle$  in the Kondo regime should be proportional to the Kondo temperature—namely, it should scale exponentially with the ratio  $\varepsilon_d/\Gamma_T$  with a slope of  $\pi$ . While in the right panel of Fig. (1) in the main text we do observe an exponential scaling, we find that the slope is smaller, being close to  $0.56\pi$ . A similar behavior is observed with the variational method in the companion paper [25].




FIG. 4. Spin decay in real time for a dot in the mixed-valence regime (left) and in the Kondo regime (right). In both scenarios we can observe how the decay is suppressed by an increasing loss rate  $\gamma$ , thus signaling the regime of strong correlations. Both plots are for  $\Gamma_T = 10^{-2}W$ . The discrete steps in the  $\gamma = 100\Gamma_T$  curve of the right plot are caused by the reaching of machine precision  $\langle \sigma_d^z(t) \rangle \sim 10^{-13}$ .

### DISCUSSION OF THE MANY-SITES SETUP

In this Section, we provide a detailed discussion of the setup in which the dot is composed of more than one site, and on the conditions for the mapping of this dissipative model to a higher-spin Kondo model.

We consider the Lindblad master equation for a composite dot made of  $\ell_d$  sites, each one subject to a two-body decay rate of  $\gamma_j$ :

$$\frac{d}{dt}\rho(t) = -i[H, \rho(t)] + \sum_{j \in \text{dot}} \gamma_j \left( L_j \rho(t) L_j^\dagger - \frac{1}{2} \{ L_j^\dagger L_j, \rho(t) \} \right), \quad (51)$$

with jump operators  $L_j \equiv d_{j\downarrow} d_{j\uparrow}$  (the operator  $d_{j\sigma}$  annihilates a fermion with spin  $\sigma$  at the dissipative site  $j$ ) and a Hamiltonian having the usual form  $H = H_{\text{dot}} + H_{\text{leads}} + H_{\text{tun}}$ , with

$$\begin{aligned} H_{\text{dot}} &= \sum_{i,j \in \text{dot}, \sigma} h_{ij} d_{i\sigma}^\dagger d_{j\sigma} \\ H_{\text{leads}} &= \sum_{p\sigma\alpha} \varepsilon_{p\alpha} c_{p\sigma\alpha}^\dagger c_{p\sigma\alpha} \\ H_{\text{tun}} &= \sum_{pj\alpha\sigma} (V_{j,p\alpha} d_{j\sigma}^\dagger c_{p\sigma\alpha} + \text{H.c.}) \end{aligned}, \quad (52)$$

where now we do not need to specify the Hermitian matrix  $h_{ij}$ , the bath energies  $\varepsilon_{p\alpha}$  and the tunneling matrices  $V_{j,p\alpha}$ . The goal of the next paragraphs is to identify the dark states of the isolated dot, and this task does not rely on any specific form of the various parameters  $h_{ij}$ ,  $\varepsilon_{p\alpha}$ ,  $V_{j,p\alpha}$ . These dark states would then form the slow subspace (i.e. protected from dissipation in a suitable  $\gamma_j \rightarrow +\infty$  limit) in which the effective Hamiltonian dynamics induced by a nonzero tunneling  $V_{j,p\alpha} \neq 0$  would take place. We will show that for a suitable choice of the  $h_{ij}$  this effective Hamiltonian dynamics is that of higher-spin, possibly many-flavor (depending on the range of  $\alpha$ ) Kondo model.

We remind the reader that dark states are eigenstates  $|D\rangle$  of the Hamiltonian  $H_{\text{dot}}$ ,  $H_{\text{dot}}|D\rangle = \varepsilon_D|D\rangle$ , which are simultaneously annihilated by all jump operators  $L_j$ ,  $L_j|D\rangle = 0$  [26, 27]. Then, the states  $\rho_D = |D\rangle\langle D|$  are stationary states of the dynamics. In our case, following [27], the general construction of the dark states rests only on the strong rotational invariance of the Lindbladian dynamics of the isolated dot:

$$\frac{d}{dt}\rho(t) = -i[H_{\text{dot}}, \rho(t)] + \sum_{j \in \text{dot}} \gamma_j \left( L_j \rho(t) L_j^\dagger - \frac{1}{2} \{ L_j^\dagger L_j, \rho(t) \} \right), \quad (53)$$

in the sense that all components of the total dot spin  $S^a = \frac{1}{2} \sum_{j,\sigma\tau} (\sigma^a)_{\sigma\tau} d_{j\sigma}^\dagger d_{j\tau}$  (where  $\sigma^a$  are the Pauli matrices) commute both with the dot Hamiltonian and with all jump operators. Let us diagonalize the Hamiltonian  $H_{\text{dot}} =$



$\sum_{a\sigma} \varepsilon_a d_{a\sigma}^\dagger d_{a\sigma}$  in terms of the appropriate single-particle modes  $d_{a\sigma} = \sum_j \varphi_a(j) d_{j\sigma}$ , where  $\varphi_a(j)$  are the eigenfunctions of  $h_{ij}$ . Then, we can form an  $n$ -particle polarized state  $|\uparrow\uparrow\ldots\uparrow\rangle$  in which  $n$  fermions with spin  $\uparrow$  are put into  $n$  different single-body eigenstates of  $H_{\text{dot}}$  ( $n$  must be smaller or equal than the number of dot sites  $\ell_d$ ). This state corresponds to a spin eigenstate  $|S, M=S\rangle$  where  $S = n/2$ , and it is trivially a dark state since all fermions have the same spin projection and thus there cannot be any double occupancies. It will have energy  $\varepsilon_{\mathbf{a}} = \sum_k \varepsilon_{a_k}$ , where  $\mathbf{a} = (a_1, \dots, a_n)$  is the set of occupied single-body levels. The rest of the states of the spin- $S$  multiplet are generated by repeated application of the ladder operator for the total spin,  $S^- \equiv \sum_j d_{j\downarrow}^\dagger d_{j\uparrow} = \sum_a d_{a\downarrow}^\dagger d_{a\uparrow}$ . Since  $[L_j, S^-] = 0$ , the states generated in this way are all annihilated by  $L_j$  [28]. These multiplets of many-body eigenstates of a Hamiltonian are known as Dicke states [27, 29, 30]. We can see that for any non-extremal choice of the number of particles  $0 < n < \ell_d$  (namely, total spin  $S = n/2$ ) there will be  $\binom{\ell_d}{n}$  different multiplets with the same spin quantum numbers but different energies (barring degeneracies of  $\varepsilon_{\mathbf{a}}$  for different  $\mathbf{a}$ ), where  $\binom{\cdot}{\cdot}$  is the binomial coefficient.

In general, there is no guarantee that the set of dark states just described exhausts the possible stationary states. In the case of a dot consisting of a linear chain of nearest-neighbor hopping fermions (in either periodic or open boundary conditions), with dissipation acting on every site or on just one, we have verified via exact diagonalization (done with the QuTiP package [31, 32]) that this is indeed the case, at least for small chains. For one or two dissipative sites, we have backed up this conclusion with analytical calculations.

The eigenstates of the isolated dot dynamics (53) outside the dark subspace—the bright states—will have a finite dissipation rate, and their population will be depleted in time. For now, we assume that there is a parameter regime in which the decay rate of the all bright states can be made sufficiently large. Later, we will show that for more than one dot site, this requirement does not simply boil down to  $\gamma \rightarrow +\infty$ . In the presence of a large decay rate of the bright states, and in the spirit of adiabatic elimination [3] or the dissipative Schrieffer-Wolff transformation [33], to a first approximation the dynamics will be restricted to the dark subspace, and within this subspace it will be unitary. If we introduce back the coupling to the leads, the latter will mediate transitions between multiplets with neighboring values of  $S$ . Indeed, we can write

$$d_{a\sigma} = \sum_{S=0}^{\ell_d/2} \sum_{M=-S}^S \sum_{\mathbf{b} \in \mathcal{A}_S} \delta_{SM\mathbf{b}}^{a\sigma} |S-1, M-\sigma, \mathbf{b}-(a)\rangle \langle S, M, \mathbf{b}|, \quad (54)$$

for certain coefficients  $\delta_{SM\mathbf{b}}^{a\sigma}$ . The set  $\mathcal{A}_S = \{(a_1, \dots, a_{2S}) | a_k \neq a_l\}$  is the set of all unordered strings of  $2S$  distinct single-particle eigenstates of  $H_{\text{dot}}$ , and the notation  $\mathbf{b}-(a)$  means that the state  $a$  is removed from the string  $\mathbf{b}$  (if present). Substituting the above expression in  $H_{\text{tun}}$  we obtain the Hamiltonian in the dark subspace

$$\begin{aligned} H_{\text{eff}} = & \sum_{S=0}^{\ell_d/2} \sum_{M=-S}^S \sum_{\mathbf{a} \in \mathcal{A}_S} \varepsilon_{\mathbf{a}} |S, M, \mathbf{a}\rangle \langle S, M, \mathbf{a}| + \sum_{p\sigma\alpha} \varepsilon_{p\alpha} c_{p\sigma\alpha}^\dagger c_{p\sigma\alpha} \\ & + \sum_{p\sigma\alpha} \sum_{S=0}^{\ell_d/2} \sum_{M=-S}^S \sum_{\mathbf{a} \in \mathcal{A}_S} (V_{SM\mathbf{b},p\alpha}^{a\sigma} |S, M, \mathbf{b}\rangle \langle S-1, M-\sigma, \mathbf{b}-(a)| c_{p\sigma\alpha} + \text{H.c.}) \end{aligned} \quad (55)$$

with  $V_{SM\mathbf{b},p\alpha}^{a\sigma} \equiv \sum_j V_{j,p\alpha} \varphi_a^*(j) (\delta_{SM\mathbf{b}}^{a\sigma})^*$ . The above Hamiltonian belongs to the family of the ionic models [2], that have been extensively studied in the context of magnetic impurities in metals. The low-energy description of such models leads to higher-spin Kondo models in suitable regimes. In general, there will be a lowest-energy multiplet with spin  $S^*$  (usually, the highest-spin  $S^* = \ell_d/2$ ), separated from the next multiplet by an energy gap  $\Delta\varepsilon$ . If this gap is much larger than the level width induced by the leads,  $\Gamma_\alpha \sim \mathcal{N}_{\alpha F} V^2$  (where  $\mathcal{N}_{\alpha F}$  is the single-particle density of states at the chemical potential of lead  $\alpha$ , and  $V^2$  is the typical magnitude of the matrix elements  $V_{SM\mathbf{b},p\alpha}^{a\sigma}$ ), then  $H_{\text{eff}}$  can be mapped to a spin  $S^*$  Kondo model by the usual Schrieffer-Wolff transformation [1, 2]. On the other hand, there is no constraint on the number of possible leads, thus opening to the possibility of studying over-screened realizations of these exotic Kondo models, that are known to have non-Fermi liquid ground states [2, 34–37].

In any realistic realization of our setup the dissipation rate  $\gamma$  will be finite. Then, the coupling to the leads will introduce a residual dissipation, since by repeated tunneling events into the dot sites one ends up in the dissipated states. The quantitative description of the effective dissipation can be achieved by adiabatic elimination of the dissipated modes [3] or, equivalently, by a dissipative Schrieffer-Wolff transformation [33]. It is simple to guess that the maximal-spin multiplet  $S_m = \ell_d/2$  will suffer the largest effective dissipation, since any particle entering the dot will create a double occupancy. Thus, we can expect that the dissipation will couple  $S_m$  to  $S_m - 1$ , with a decay rate  $\mathcal{O}(V^2/\gamma)$ , where  $V$  is the tunneling rate to the leads. The higher-lying multiplets will need more tunneling events to reach the dissipative states, with a multiplet of spin  $S$  acquiring a dissipative rate only at order  $S_m - S + 1$  in  $V^2$ .

We now consider the concrete scenario of a dot composed of two dissipative sites, for which the dynamics (53) is amenable to an exact solution. We consider the Hamiltonian

$$H_{\text{dot}} = \varepsilon_d \sum_{j=0,\sigma}^1 d_{j\sigma}^\dagger d_{j\sigma} - J \sum_{\sigma} (d_{0\sigma}^\dagger d_{1\sigma} + \text{H.c.}) = \sum_{a,\sigma} \varepsilon_a d_{a\sigma}^\dagger d_{a\sigma} , \quad (56)$$

where the eigenstates are the symmetric and antisymmetric modes  $d_{a=\pm,\sigma} = (d_{0\sigma} \pm d_{1\sigma})/2^{1/2}$ , corresponding to the energies  $\varepsilon_{\pm} = \varepsilon_d \mp J$ . With these states, we obtain the dark subspace multiplet structure shown in Fig. (2) in the main text. In detail, we have the  $S = 0$  empty dot state  $|0\rangle$ , the two  $S = 1/2$  doublets  $d_{a\sigma}^\dagger |0\rangle$  with energies  $\varepsilon_a$  and the  $S = 1$  triplet  $\{d_{+\uparrow}^\dagger d_{-\uparrow}^\dagger |0\rangle, 2^{-1/2}(d_{+\uparrow}^\dagger d_{-\downarrow}^\dagger + d_{+\downarrow}^\dagger d_{-\uparrow}^\dagger) |0\rangle, d_{+\downarrow}^\dagger d_{-\downarrow}^\dagger |0\rangle\}$  with energy  $\varepsilon_{S=1} = \sum_a \varepsilon_a = 2\varepsilon_d$ . We are considering the case  $\varepsilon_d < 0$ ,  $|\varepsilon_d| > J$  which makes the spin triplet the lowest lying multiplet. In this case, following the construction detailed in the previous paragraph, the effective low-energy theory for a weakly coupled bath would yield a  $S = 1$  Kondo model. For  $-J < \varepsilon_d < 0$ , the doublet built out of the symmetric states would have the minimum energy  $\varepsilon_d - J$ , and the effective low-energy Hamiltonian would be the usual  $S = 1/2$  Kondo model.

We now build the full set of bright states of the  $\mathcal{L}_{\text{dot}} = -i[H_{\text{dot}}, \cdot] + \sum_{j \in \text{dot}} \gamma_j (L_j \cdot L_j^\dagger - \frac{1}{2}\{L_j^\dagger L_j, \cdot\})$  appearing in Eq. (53). Let us consider the non-Hermitian part of the Lindblad dynamics,  $K = H_{\text{dot}} - i\gamma/2 \sum_j L_j^\dagger L_j = H_{\text{dot}} - i/2 \sum_j \gamma_j n_{j\uparrow} n_{j\downarrow}$  (with  $n_{j\sigma} = d_{j\sigma}^\dagger d_{j\sigma}$  the number operator at site  $j$ ). For a uniform  $\gamma_j = \gamma$ , this Hamiltonian is a non-Hermitian version of the Hubbard model [1]. Since  $K$  conserves the number of fermions (and spin), we can consider each particle number  $N_f$  (and spin  $S$ ) sector separately. In the one-particle sector,  $K$  coincides with  $H_{\text{dot}}$ , since there are no double occupancies. In the two-particle sectors, it is easy to observe that the eigenstates of  $\mathcal{L}_{\text{dot}}$  can be inferred from the diagonalization of  $K$ . Indeed, let  $K|r_\alpha\rangle = \kappa_\alpha|r_\alpha\rangle$  be the right eigenstates (with  $|l_\alpha\rangle$  the left ones,  $\langle l_\alpha|K = \kappa_\alpha\langle l_\alpha|$ , with the normalization  $\langle l_\alpha|r_\beta\rangle = \delta_{\alpha\beta}$ ). If  $L_j|r_\alpha\rangle = 0$  for all  $j$ , then  $|r_\alpha\rangle$  is one of the dark states built above, and the eigenstates of  $\mathcal{L}_{\text{dot}}$  in the dark subspace are  $|r_\alpha\rangle\langle r_\beta|$ , with eigenvalue  $\lambda = -i(\kappa_\alpha - \kappa_\beta)$  (since dark states are eigenstates of the Hermitian Hamiltonian  $H_{\text{dot}}$ , right and left eigenstates coincide and the  $\kappa_\alpha$ s are real). This construction of the dark states is valid in all particle-number sectors. Let us consider the  $S = 0$ ,  $N_f = 2$  sector with two fermions of opposite spin. Since each jump operator removes precisely two opposite-spin fermions, we must have  $L_j|r_\alpha\rangle = \eta_{j\alpha}|0\rangle$ , and we can anticipate that

$$\sigma = p_0 |0\rangle\langle 0| + p_1 |r_\alpha\rangle\langle r_\beta| \quad (57)$$

will be a right eigenstate of the Lindblad superoperator. Since bright states must be traceless [as  $0 = \text{Tr}(\mathcal{L}_{\text{dot}}\sigma) = \lambda \text{Tr}\sigma$ ], we have  $p_0 = -p_1 \langle r_\beta|r_\alpha\rangle$ . Requiring  $\mathcal{L}_{\text{dot}}\sigma = \lambda\sigma$  we find

$$\lambda = -i(\kappa_\alpha - \kappa_\beta^*) , \quad (58)$$

provided that  $\sum_j \gamma_j \eta_{j\alpha} \eta_{j\beta}^* = -\lambda \langle r_\beta|r_\alpha\rangle$ . The latter is indeed satisfied: by taking the overlap of  $K|r_\alpha\rangle = \kappa_\alpha|r_\alpha\rangle$  with  $|r_\beta\rangle$  we find

$$\langle r_\beta|H_{\text{dot}}|r_\alpha\rangle = \kappa_\alpha \langle r_\beta|r_\alpha\rangle + \frac{i}{2} \sum_j \gamma_j \eta_{j\alpha}^* \eta_{j\beta} . \quad (59)$$

The Hermiticity of  $H_{\text{dot}}$ ,  $\langle r_\alpha|H_{\text{dot}}|r_\beta\rangle^* = \langle r_\beta|H_{\text{dot}}|r_\alpha\rangle$  yields the desired relation. From the above discussion, we infer that by diagonalizing  $K$  in the two-fermion sector we can find the decay rates  $-\text{Re}\lambda > 0$  by looking at the imaginary parts of the  $\kappa_\alpha$ s.

We now proceed to diagonalize  $K$  in the  $N_f = 2$ ,  $S = 0$  sector. We work in real space and we introduce the basis states [38]

$$\begin{cases} |s\rangle = \frac{1}{\sqrt{2}}(|\uparrow, \downarrow\rangle - |\downarrow, \uparrow\rangle) , \\ |d_0\rangle = |\uparrow\downarrow, 0\rangle , \\ |d_1\rangle = |0, \uparrow\downarrow\rangle , \end{cases} \quad (60)$$

where the notation  $|\uparrow, \downarrow\rangle = d_{0\uparrow}^\dagger d_{1\downarrow}^\dagger |0\rangle$  denotes a state with the  $\uparrow$  fermion sits on the  $j = 0$  site while the other sits at the  $j = 1$  site, and  $|\uparrow\downarrow, 0\rangle$  denotes that both fermions are at site 0 [39]. In the basis  $\{|s\rangle, |d_0\rangle, |d_1\rangle\}$  the non-Hermitian Hamiltonian  $K$  reads

$$\begin{pmatrix} 2\varepsilon_d & -\sqrt{2}J & -\sqrt{2}J \\ -\sqrt{2}J & 2\varepsilon_d - i\gamma_0/2 & 0 \\ -\sqrt{2}J & 0 & 2\varepsilon_d - i\gamma_1/2 \end{pmatrix} \quad (61)$$




FIG. 5. Real (left panel) and imaginary parts (right panel) of the eigenvalues of the non-Hermitian Hamiltonian  $K = H_{\text{dot}} - i\gamma/2 \sum_j n_{j\uparrow} n_{j\downarrow}$  in the  $N_f = 2$ ,  $S = 0$  sector, showing the presence of a state ( $|r_+\rangle$ ) which becomes effectively dark for infinite dissipation.

Let us consider the simpler case of a uniform dissipation  $\gamma_j = \gamma$ . Then, we can form the states  $|d_{\pm}\rangle \equiv (|d_0\rangle \pm |d_1\rangle)/2^{1/2}$  and we find that  $K|d_{-}\rangle = (2\varepsilon_d - i\gamma/2)|d_{-}\rangle$ . Hence, according to Eq. (58), eigenmatrices of  $\mathcal{L}_{\text{dot}}$  involving  $|d_{-}\rangle$  will have a decay rate of at least  $\gamma/2$ . In the remaining subspace spanned by  $\{|s\rangle, |d_{+}\rangle\}$  we have

$$K = \begin{pmatrix} 2\varepsilon_d & -2J \\ -2J & 2\varepsilon_d - i\gamma/2 \end{pmatrix}, \quad (62)$$

which is easily diagonalized to find the eigenvalues

$$\kappa_{\pm} = 2\varepsilon_d - \frac{i}{4}\gamma \pm \frac{1}{4}(64J^2 - \gamma^2)^{1/2} \quad (63)$$

and the eigenvectors

$$\begin{aligned} |r_{\pm}\rangle &= [4J^2 + (\kappa_{\pm} - 2\varepsilon_d)^2]^{-1/2} \begin{pmatrix} 2J \\ -\kappa_{\pm} + 2\varepsilon_d \end{pmatrix} \\ |l_{\pm}\rangle &= [4J^2 + (\kappa_{\pm} - 2\varepsilon_d)^2]^{-1/2} \begin{pmatrix} 2J \\ -\kappa_{\pm} + 2\varepsilon_d \end{pmatrix} \end{aligned} \quad (64)$$

Since  $K = K^T$  is symmetric,  $|l_{\pm}\rangle = (|r_{\pm}\rangle)^*$ . We sum up the spectral properties of the  $N_f = 2$ ,  $S = 0$  sector in Fig. 5. For the  $|r_{\pm}\rangle$  states we observe two distinct regimes, separated by  $\gamma^* = 8J$ . This value of  $\gamma$  marks an exceptional point [40] where the two eigenvectors become parallel (and  $\kappa_+ = \kappa_-$ ) and  $K$  is no longer diagonalizable. For  $\gamma < \gamma^*$ , the two eigenvalues  $\kappa_{\pm}$  have distinct real parts while the imaginary part of both is equal to  $-i\gamma/4$ . In this regime, a larger dissipation rate corresponds to a larger decay rate for the bright states. For larger  $\gamma > \gamma^*$  the two real parts coincide, while the imaginary parts start to diverge from each other: while  $-\text{Im}\kappa_- \sim i\gamma/2 + \mathcal{O}(J^2/\gamma)$  increases further,  $-\text{Im}\kappa_+$  decreases, ultimately as  $8J^2/\gamma$  for large  $\gamma \gg J$ . This phenomenon can be seen as an incarnation of the Zeno effect, and implies that for  $\gamma \rightarrow +\infty$  the  $|r_+\rangle$  state (which in this limit coincides with the singlet  $|s\rangle$ ) becomes effectively dark. A numerical diagonalization of the Lindbladian with QuTiP confirms that  $-2\text{Im}\kappa_+$  is indeed the lowest decay rate at large dissipation. Thus, in order to have an effective dynamics that involves the dark subspace only, we need to increase the intra-dot tunneling  $J$  at the same rate as the dissipation  $\gamma$ , i.e.  $J \propto \gamma$ . At the same time, to have an appreciable Kondo temperature for the effective  $S = 1$  (or  $S = 1/2$ ) Kondo model at low energy we must keep the gap  $\Delta\varepsilon = |\varepsilon_d + J|$  between the two lowest-lying multiplets finite: this requirement implies that  $\varepsilon_d$  must be scaled proportionally to  $\gamma$  as well. In other words, the adiabatic elimination of the bright states is possible only in the regime  $|\varepsilon_d| \sim J \sim \gamma \gg \Gamma$ , where  $\Gamma$  is the level width induced by the leads. See also the companion paper [25] for further discussions of these points.

To complete our description of the bright states, let us consider the completely filled state  $|F\rangle = |\uparrow\downarrow, \uparrow\downarrow\rangle$ , which is an eigenstate of  $K$ :  $K|F\rangle = (4\varepsilon_d - i\gamma)|F\rangle$ . The corresponding eigenmatrix of  $\mathcal{L}_{\text{dot}}$  can be constructed by analogy with Eq. (57), namely as a superposition of  $|F\rangle\langle F|$ ,  $|r_{\alpha}^{(2,0,0)}\rangle\langle r_{\beta}^{(2,0,0)}|$  and  $|0\rangle\langle 0|$ , where  $|r_{\alpha}^{(N_f, S, M)}\rangle$  indicates a right eigenstate of  $K$  in the sector with  $N_f$  particles, spin  $S$  and magnetization  $M$  [41]. We will not reproduce the full




FIG. 6. Real (left panel) and imaginary parts (right panel) of the eigenvalues of the non-Hermitian Hamiltonian  $K = H_{\text{dot}} - i\gamma/2 n_{0\uparrow} n_{0\downarrow}$  for the case of inhomogeneous dissipation in the  $N_f = 2$ ,  $S = 0$  sector, showing the presence of two states ( $|r_{1,2}\rangle$ ) which become effectively dark for infinite dissipation.

calculation here, but it is easy to understand that the real part of the eigenvalue of  $\mathcal{L}_{\text{dot}}$  will receive a contribution of  $-\text{Im}(\kappa_\alpha + \kappa_\beta)$  from each term  $|r_\alpha\rangle\langle r_\beta|$ . Hence, the resulting decay rate will be at least  $2\gamma$ —indeed, it is the state with the faster decay rate.

Finally, we consider the  $N_f = 3$ ,  $S = 1/2$  sector. Since all states in this sector have one unpaired spin, and by particle-hole correspondence with the  $N_f = 1$ ,  $S = 1/2$  states, we expect to find two  $S = 1/2$  doublets. Indeed, we can think of these states as featuring a “hole” on top of the full state  $|F\rangle$ : we have verified that the states  $|N_f = 3, a\sigma\rangle \equiv d_{-a\sigma} |F\rangle$  are the sought eigenstates, with eigenvalues  $\kappa = 4\varepsilon_d - \varepsilon_{-a} + i\gamma/2$ .

We remark that in the present case the diagonalization of  $K$  in all symmetry sectors allows to reconstruct all eigenmatrices of the Lindblad superoperator by suitably combining the projectors  $|r_\alpha^{(N_f, S, M)}\rangle\langle r_\beta^{(N'_f, S', M')}|$ ,  $|r_\alpha^{(N_f-2, S, M)}\rangle\langle r_\beta^{(N'_f-2, S', M')}|$  and  $|0\rangle\langle 0|$ . As observed above, each term will add a positive contribution  $-\text{Im}\kappa$  to the decay rate  $-\text{Re}\lambda$ . Since we have proven that the 3- and 4-particle sectors have all  $-\text{Im}\kappa \propto \gamma$ , any time an eigenmatrix features an eigenstate of  $K$  with more than 2 particles appears, the decay rate will be linear in  $\gamma$ . The upshot of this discussion is that the lowest decay rate, vanishing for  $\gamma \rightarrow +\infty$ , only comes from the  $N_f = 2$ ,  $S = 0$  sector considered before.

The conclusions drawn for the homogeneously dissipated case  $\gamma_j = \gamma$  are essentially unchanged also in the completely inhomogeneous one with all dissipation concentrated only on one site,  $\gamma_0 = \gamma$ ,  $\gamma_1 = 0$ . The dark subspace does not change, while the non-Hermitian Hamiltonian in the  $N_f = 2$ ,  $S = 0$  subspace becomes:

$$K = \begin{pmatrix} 2\varepsilon_d & -\sqrt{2}J & -\sqrt{2}J \\ -\sqrt{2}J & 2\varepsilon_d - i\gamma/2 & 0 \\ -\sqrt{2}J & 0 & 2\varepsilon_d \end{pmatrix}, \quad (65)$$

whose eigenvalues have to be found numerically. The results are shown in Fig. 6. As in the homogeneous case, there is one mode with  $\text{Re}\kappa_1 = 2\varepsilon_d$  and a monotonic decay rate  $-\text{Im}\kappa \sim \gamma/2$ , but both the other modes have the same non-monotonic imaginary part—which is generally smaller than in the homogeneous case. The decay rates of these states rises linearly  $-\text{Im}\kappa \sim \gamma/8$  for small dissipation, reaches a maximum for  $\gamma \approx 4J$  and then slowly decreases as  $-\text{Im}\kappa \sim 2J^2/\gamma$  for large dissipation. In this setup, there is no exceptional point. An exceptional point appears instead in the  $N_f = 3$  particle sector, in which

$$K = \begin{pmatrix} 3\varepsilon_d & J \\ J & 3\varepsilon_d - i\gamma/2 \end{pmatrix} \quad (66)$$

in the basis  $\{d_{0\sigma} |F\rangle, d_{1\sigma} |F\rangle\}$  has the same form of Eq. (62) [42]. We see that for a single-site dissipation, also the  $N_f = 3$  sector has a state with a non-monotonic decay rate, reaching a maximum for  $\gamma = 4J$ . The decay rate in this sector is always larger than the one in the  $N_f = 2$ ,  $S = 0$  sector, albeit they tend to coincide for large dissipation. The full state  $|F\rangle$  has a monotonic decay rate,  $K |F\rangle = (4\varepsilon_d - i\gamma/2) |F\rangle$ , which is obviously smaller than in the homogeneous

case. This brief analysis of the inhomogeneous setup allows to draw two conclusions: First, a single dissipative site is sufficient to provide all states outside the dark subspace with a finite dissipation rate, and so to implement the mapping to higher-spin Kondo models described above. The second conclusion is that this setup is more fragile than the homogeneous one, in the sense that more bright states become essentially dark for large dissipation  $\gamma \gg J$ , and that the overall decay rates are smaller than in the homogeneous case.

The ideas presented above can be extended to larger dot sizes, and we expect that the minimal decay rates of the bright states will have a non-monotonic behavior. We have explicitly verified this expectation for  $\ell_d = 3$ . Thus, in general, the fine-tuning  $J \sim |\varepsilon_d| \sim \gamma$  will be needed. This fine-tuning might be circumvented by choosing a different geometry for the dot sites, as for instance a potential well hosting more than one bound state.

- 
- [1] P. Coleman, *Introduction to Many-Body Physics* (Cambridge University Press, 2015).
  - [2] A. C. Hewson, *The Kondo Problem to Heavy Fermions*, Cambridge Studies in Magnetism (Cambridge University Press, 1993).
  - [3] J. J. García-Ripoll, S. Dürr, N. Syassen, D. M. Bauer, M. Lettner, G. Rempe, and J. I. Cirac, Dissipation-induced hard-core boson gas in an optical lattice, *New Journal of Physics* **11**, 013053 (2009).
  - [4] N. E. Bickers, Review of techniques in the large- $N$  expansion for dilute magnetic alloys, *Rev. Mod. Phys.* **59**, 845 (1987).
  - [5] E. M. Kessler, Generalized schrieffer-wolff formalism for dissipative systems, *Phys. Rev. A* **86**, 012126 (2012).
  - [6] S. E. Barnes, New method for the anderson model, *Journal of Physics F: Metal Physics* **6**, 1375 (1976).
  - [7] P. Coleman, New approach to the mixed-valence problem, *Phys. Rev. B* **29**, 3035 (1984).
  - [8] L. M. Sieberer, M. Buchhold, and S. Diehl, Keldysh field theory for driven open quantum systems, *Reports on Progress in Physics* **79**, 096001 (2016).
  - [9] M. H. Hettler, J. Kroha, and S. Hershfield, Nonequilibrium dynamics of the anderson impurity model, *Phys. Rev. B* **58**, 5649 (1998).
  - [10] N. S. Wingreen and Y. Meir, Anderson model out of equilibrium: Noncrossing-approximation approach to transport through a quantum dot, *Phys. Rev. B* **49**, 11040 (1994).
  - [11] N. Sivan and N. S. Wingreen, Single-impurity anderson model out of equilibrium, *Phys. Rev. B* **54**, 11622 (1996).
  - [12] D. C. Langreth and P. Nordlander, Derivation of a master equation for charge-transfer processes in atom-surface collisions, *Phys. Rev. B* **43**, 2541 (1991).
  - [13] H. Shao, D. C. Langreth, and P. Nordlander, Many-body theory for charge transfer in atom-surface collisions, *Phys. Rev. B* **49**, 13929 (1994).
  - [14] M. Fabrizio, *A Course in Quantum Many-Body Theory*, 1st ed., Graduate Texts in Physics (Springer Cham, 2022).
  - [15] H. Haug and A.-P. Jauho, *Quantum Kinetics in Transport and Optics of Semiconductors*, 2nd ed., Springer Series in Solid-State Sciences (Springer Berlin, Heidelberg, 2007).
  - [16] Since we are not interested in thermal transport, we assume that both baths have the same temperature.
  - [17] A. Kamenev, *Field Theory of Non-Equilibrium Systems* (Cambridge University Press, 2011).
  - [18] T. A. Costi, J. Kroha, and P. Wölfle, Spectral properties of the anderson impurity model: Comparison of numerical-renormalization-group and noncrossing-approximation results, *Phys. Rev. B* **53**, 1850 (1996).
  - [19] Y. Meir and N. S. Wingreen, Landauer formula for the current through an interacting electron region, *Phys. Rev. Lett.* **68**, 2512 (1992).
  - [20] This limit cannot be taken, however, since the derivation of the effective Hamiltonian assumes  $\gamma \gg W$ .
  - [21] N. S. Wingreen and Y. Meir, Anderson model out of equilibrium: Noncrossing-approximation approach to transport through a quantum dot, *Phys. Rev. B* **49**, 11040 (1994).
  - [22] D. Lobaskin and S. Kehrein, Crossover from nonequilibrium to equilibrium behavior in the time-dependent kondo model, *Phys. Rev. B* **71**, 193303 (2005).
  - [23] F. B. Anders and A. Schiller, Real-time dynamics in quantum-impurity systems: A time-dependent numerical renormalization-group approach, *Phys. Rev. Lett.* **95**, 196801 (2005).
  - [24] M. M. Wauters, C.-M. Chung, L. Maffi, and M. Burrello, Simulations of the dynamics of quantum impurity problems with matrix product states, *Phys. Rev. B* **109**, 115101 (2024).
  - [25] Y.-F. Qu, M. Stefanini, J. Marino, T. Esslinger, S. Gopalakrishnan, and E. Demler, tbd, *Phys. Rev. B* (2024).
  - [26] H.-P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, 2003).
  - [27] M. Foss-Feig, A. J. Daley, J. K. Thompson, and A. M. Rey, Steady-state many-body entanglement of hot reactive fermions, *Phys. Rev. Lett.* **109**, 230501 (2012).
  - [28] Of course, the same set of dark states would be found by starting with the  $\downarrow$  polarized state  $|\downarrow \uparrow \dots \downarrow\rangle$  in the same levels  $\mathbf{a}$ , by repeated application of the raising operator  $S^+ = (S^-)^\dagger$ .
  - [29] R. H. Dicke, Coherence in spontaneous radiation processes, *Phys. Rev.* **93**, 99 (1954).
  - [30] F. T. Arecchi, E. Courtens, R. Gilmore, and H. Thomas, Atomic coherent states in quantum optics, *Phys. Rev. A* **6**, 2211 (1972).
  - [31] J. Johansson, P. Nation, and F. Nori, QuTiP: An open-source Python framework for the dynamics of open quantum systems, *Computer Physics Communications* **183**, 1760 (2012).

- [32] J. Johansson, P. Nation, and F. Nori, QuTiP 2: A Python framework for the dynamics of open quantum systems, [Computer Physics Communications](#) **184**, 1234 (2013).
- [33] E. M. Kessler, Generalized schrieffer-wolff formalism for dissipative systems, [Phys. Rev. A](#) **86**, 012126 (2012).
- [34] A. O. Gogolin, A. A. Nersesyan, and A. M. Tsvelik, *Bosonization and Strongly Correlated Systems* (Cambridge University Press, Cambridge, 1998).
- [35] T. Giamarchi, *Quantum Physics in One Dimension*, 1st ed. (Clarendon Press, Oxford, 2003).
- [36] D. B. Karki, C. Mora, J. von Delft, and M. N. Kiselev, Two-color Fermi-liquid theory for transport through a multilevel Kondo impurity, [Phys. Rev. B](#) **97**, 195403 (2018).
- [37] D. B. Karki, A. I. Pavlov, and M. N. Kiselev, Multistage Kondo effect in a multiterminal geometry: A modular quantum interferometer, [Phys. Rev. B](#) **105**, L041410 (2022).
- [38] The “missing” basis state in the subspace with  $N_f = 2$  and total magnetization  $M = 0$  is the triplet state  $|t\rangle = \frac{1}{\sqrt{2}}(|\uparrow, \downarrow\rangle + |\downarrow, \uparrow\rangle)$ , which is a dark state belonging to the  $S = 1$  manifold.
- [39] We order the labels as  $0 \uparrow < 0 \downarrow < 1 \uparrow < 1 \downarrow < \dots$ .
- [40] Y. Ashida, Z. Gong, and M. Ueda, Non-Hermitian physics, [Advances in Physics](#) **69**, 249 (2020).
- [41] Since  $|F\rangle$  has  $S = M = 0$  and the jump operators conserve spin, the whole eigenmatrix must be composed only of states with the same spin quantum numbers.
- [42] The matrix reported is identical for the two possible magnetizations  $M = \pm 1$ .