EXCEPTIONAL FANO VARIETIES WITH SMALL MINIMAL LOG DISCREPANCY

LOUIS ESSER, JIHAO LIU, AND CHENGXI WANG

ABSTRACT. We construct exceptional Fano varieties with the smallest known minimal log discrepancies in all dimensions. These varieties are well-formed hypersurfaces in weighted projective space. Their minimal log discrepancies decay doubly exponentially with dimension, and achieve the optimal value in dimension 2.

CONTENTS

1. Introduction	1
2. Preliminaries	4
2.1. Weighted Projective Varieties	4
2.2. Singularities in Birational Geometry	5
3. Log Discrepancies of Hypersurface Singularities	6
4. The examples and their well-formedness	9
5. Computation of the minimal log discrepancy	14
6. Computation of the alpha invariant and exceptionality	20
References	27

1. INTRODUCTION

We work over the field of complex numbers \mathbb{C} .

Explicit birational geometry in high dimensions. Explicit birational geometry, often referred to as the "geography of birational geometry," studies the extreme values of algebraic invariants in birational geometry. Traditionally, the focus of explicit birational geometry has been on varieties of dimension at most 3 with at worst canonical singularities. Recent research, however, has expanded this focus to include higher-dimensional varieties, those with singularities worse than canonical, and new invariants such as Tian's α -invariant (also known as the global log canonical threshold in [28]), minimal log discrepancies, and the value *N* in the boundedness of *N*-complements.

A primary focus of recent studies is to construct examples with extreme invariants, either very small (when there is a positive lower bound) or very large (when there is an

²⁰²⁰ Mathematics Subject Classification. 14J40, 14J45 (primary); 14C20, 14E30, 14J17 (secondary).

upper bound). Although these examples are not yet known to be optimal in higher dimensions, they provide essential insights that guide the setting of optimization goals, which is important even in lower dimensions. For instance, the surface examples constructed in [2] have been crucial for optimizations in further studies [22, 23]. Thanks to the first author, the third author, B. Totaro, and others, numerous examples of varieties with extreme invariants have been established in arbitrary dimensions. These extreme values often show doubly exponential growth or decay with respect to the dimension of the ambient variety [9, 10, 11, 12, 28, 29, 31]. This paper continues this series of studies by focusing on the minimal log discrepancy of exceptional Fano varieties.

Explicit birational geometry for exceptional Fano varieties. Exceptional Fano varieties are Fano varieties whose \mathbb{R} -complements are klt; equivalently, these are Fano varieties with α -invariants strictly greater than 1 [7]. Birkar famously showed that exceptional Fano varieties form a bounded family [6], which is one crucial step in his proof of the boundedness of *N*-complements and the BAB conjecture. Additionally, since exceptional Fano varieties are K-stable [26, 25], studying the explicit geometry of exceptional Fano varieties is greatly beneficial for the construction of K-moduli spaces.

In the context of explicit birational geometry, the following four invariants are crucial for characterizing an exceptional Fano variety X of dimension n: the anti-canonical volume $vol(-K_X)$, Tian's alpha invariant $\alpha(X)$, complement index (i.e. the smallest positive integer N so that $N(K_X + B) \sim 0$ for some lc pair (X, B)), and the global minimal log discrepancy mld(X). Research on the explicit geometry of the first three invariants has been thorough:

- The well-known explicit upper bound for $vol(-K_X)$ is $(n+1)^n$ (cf. [24, Theorem 3]), and there are known examples with $vol(-K_X)$ doubly exponentially small; these examples are conjectured to attain the minimum [28, Theorem 3.1, Conjecture 3.2].
- $\alpha(X)$ has a trivial lower bound of 1, and there are known examples where $\alpha(X)$ increases doubly exponentially with dimension [28, Theorem 8.1, Question 8.2].
- The complement index has a trivial lower bound of 1, and there are known examples where the complement index increases doubly exponetially with dimension [28, Theorem 8.1, Question 8.2].

However, although the upper bound of mld(X) is trivially 1, its explicit lower bound remains unclear. Even for surfaces, the optimal lower bound was proven only very recently by the second author and V. V. Shokurov [22, Theorem 1.9]. This paper addresses this gap in all dimensions by establishing an explicit expectation for the lower bound of mld(X).

Theorem 1.1. Let s_n be the *n*th Sylvester number. Then there exists a sequence of exceptional Fano varieties $\{X_n\}_{n\geq 2}$, such that each X_n is of dimension n,

$$mld(X_n) = \frac{4(s_n - 1)}{s_n^3 - 9s_n}$$

if n is even, and

$$mld(X_n) = \frac{4(s_n - 3)}{s_n^3 - 19s_n + 14}$$

if n is odd. In particular, $mld(X_n)$ *is asymptotic to* $4/s_n^2$ *, as* $n \to \infty$ *, so that this value converges to zero doubly exponentially with n*.

The explicit construction of X_n can be found in Theorem 4.1 when n is even and in Theorem 4.2 when n is odd. In each case, X_n is a well-formed, but not quasismooth, hypersurface in a weighted projective space. We conjecture that the values obtained in Theorem 1.1 are exactly the smallest mld's among all exceptional Fano varieties of the same dimension:

Conjecture 1.2. For any even (resp. odd) integer $n \ge 2$, the smallest minimal log discrepancy of an exceptional Fano variety of dimension n is

$$\frac{4(s_n-1)}{s_n^3-9s_n}\left(\text{resp. } \frac{4(s_n-3)}{s_n^3-19s_n+14}\right).$$

Conjecture 1.2 holds in dimension 2 due to [22, Theorem 1.9(2)], where the optimal value is 3/35.

Idea of the proof of Theorem **1.1**. Similar to the construction of other extreme examples, the examples we construct in Theorem **1.1** are well-formed hypersurfaces in weighted projective spaces. The invariants central to some of those previous examples, such as anti-canonical volume and complement index, are easy to read off using the degree and weights. In contrast, there are two significant difficulties in proving Theorem **1.1**: the computation of the minimal log discrepancy and the proof that the varieties are exceptional.

- The computation of the minimal log discrepancy: The singularities of a quasismooth hypersurface in a weighted projective space may be determined in a straightforward way from the degree and weights. Since these are toric, the minimal log discrepancy is computable combinatorially (see Proposition 3.1). However, the computation of the mld at non-quasismooth points is not as straightforward, and we develop a method to resolve this issue. More precisely, we establish a formula for computing the minimal log discrepancies of hypersurfaces in toric varieties that satisfy certain properties, i.e., those defined by equations that are Newton non-degenerate (Theorem 3.3). This is essentially due to the fact that divisors computing the mlds in this case are obtained via toroidal resolutions. It is worth mentioning that we do not have similar formulas for arbitrary hypersurfaces. In our examples, we need to apply this result to the non-quasismooth point and also show that the resulting discrepancy value is indeed a global minimum. This involves lengthy but elementary arguments and is done in Theorem 5.1.
- Proving that the varieties are exceptional: The difficulty here is that there is no straightforward theoretical formula to control the lower bound of the α -invariant, so the explicit structure of the variety needs to be used. A natural idea here is to use similar arguments as the weighted cone construction in [28, Section 4]. At the end of the day, this idea works, and the proof is completed in Theorems 6.5 and 6.6. However, our arguments are more complicated than those in [28]. This is because the hypersurfaces in [28] have better inductive structure than ours, so we

need to have more accurate control over the multiplicities at every point that may attain the α -invariant.

Finally, we briefly mention here how we found the exceptional Fano varieties with small mlds. One key idea is that the exceptional del Pezzo surface with the smallest mld is already known [22, Table 8]. Although this surface was not created via a weighted hypersurface construction in [22], it has a very small volume [22, Table 8] and its singularities are explicitly characterized. Therefore, we can perform a thorough search for hypersurfaces *X* in weighted projective surfaces and determine that

$$X_{282} \coloneqq \{x_0^2 + x_1^3 + x_2^{19}x_3 + x_1x_2x_3^5 = 0\} \subset \mathbb{P}^3(141, 94, 13, 35),$$

is the desired surface. This is the starting point of the construction of examples. The higher-dimensional examples are constructed by carefully comparing examples with those in previous literature and choosing weights.

Sketch of the paper. In Section 2, we recall some preliminary definitions relating to subvarieties of weighted projective space and singularities in birational geometry. In Section 3, we establish a formula for the computation of minimal log discrepancies of hypersurface singularities in weighted projective spaces. In Section 4, we construct the examples and prove their basic properties. In Section 5, we compute the minimal log discrepancies of the examples and show that they are asymptotically proportional to $4/s_n^2$. In Section 6, we show that the examples we constructed are exceptional Fano, hence concluding the proof of Theorem 1.1.

Acknowledgement. The authors would like to thank Burt Totaro for useful discussions and comments.

2. Preliminaries

This section will introduce the necessary background on subvarieties in weighted projective space and the concepts from birational geometry that we use in the paper.

2.1. Weighted Projective Varieties. Let c_0, \ldots, c_n be positive integers. The weighted projective space $Y = \mathbb{P}(c_0, \ldots, c_n)$ is defined to be the quotient variety $(\mathbb{A}^{n+1} \setminus \{0\})/\mathbb{G}_m$ over \mathbb{C} , where the multiplicative group \mathbb{G}_m acts by $t \cdot (x_0, \ldots, x_n) = (t^{c_0}x_0, \ldots, t^{c_n}x_n)$ [15, section 6]. We can view the weighted projective space Y as the quotient stack $\mathcal{Y} = [(\mathbb{A}^{n+1} \setminus \{0\})/\mathbb{G}_m]$; it is a smooth Deligne-Mumford stack. We say that Y is well-formed if the stack \mathcal{Y} has trivial stabilizer in codimension 1, or equivalently if $gcd(c_0, \ldots, \widehat{c_j}, \ldots, c_n) = 1$ for each j. In the well-formed case, the canonical class of the variety Y is $K_Y = \mathcal{O}_Y(-\sum c_j)$.

When *Y* is well-formed, we denote by $\mathcal{O}_Y(1)$ the sheaf associated with the standard graded module shifted by 1 over the graded coordinate ring of the weighted projective space *Y* [27, Tag 01M3]. It is a reflexive sheaf associated to a Weil divisor. The divisor class $\mathcal{O}_Y(m)$ is Cartier if and only if *m* is a multiple of every weight c_j . We can also view $\mathcal{O}(1)$ as a line bundle on the stack \mathcal{Y} . The intersection number $\int_{\mathcal{Y}} c_1(\mathcal{O}(1))^n$ is $1/(c_0 \cdots c_n)$. The *degree* for an integral closed substack *T* of dimension *r* in \mathcal{Y} means $\int_T c_1(\mathcal{O}(1))^r$.

Note that if *X* is a hypersurface in \mathcal{Y} defined by a weighted-homogeneous polynomial of degree *d*, then its degree as a substack of \mathcal{Y} is $X \cdot c_1(\mathcal{O}(1))^{n-1} = d/(c_0 \cdots c_n)$.

We say that the subvariety *X* of *Y* is *well-formed* if *Y* is well-formed and $X \cap \text{Sing}(Y)$ has codimension at least 2 in *X*. The subvariety *X* is *quasismooth* if the affine cone over *X* is smooth, where the affine cone is the preimage of *X* in $\mathbb{A}^{n+1} \setminus \{0\}$.

2.2. Singularities in Birational Geometry. The minimal log discrepancy, or mld, of a variety (or pair) is a numerical measure of its singularities. We'll primarily be concerned with varieties in this paper, but will state the definition for a pair (X, D).

Definition 2.1. Let *X* be a normal complex variety and *D* an effective \mathbb{Q} -divisor on *X* with the property that $K_X + D$ is \mathbb{Q} -Cartier. Given a proper birational morphism $\mu : X' \to X$ from another normal variety *X* and $E \subset X'$ any irreducible divisor, the *log discrepancy* of *E* with respect to the morphism μ is defined as:

$$a_E(X, D) \coloneqq \operatorname{ord}_E(K_{X'} + E - \mu^*(K_X + D)).$$

This definition turns out to only depend on the valuation defined by *E* on the function field of *X*, and not on the specific choice of μ extracting this divisor. The *center* $c_X(E)$ of *E* is $\mu(E) \subset X$. For any point *x* of the scheme *X*, we define

$$\mathrm{mld}(X, D, x) \coloneqq \inf\{a_E(X, D) : c_X(E) = \bar{x}\},\$$

where the infimum runs over all divisors E on proper birational models of X with the indicated property. Finally, the (global) minimal log discrepancy of the variety X is

$$\operatorname{mld}(X, D) \coloneqq \inf_{x \in X} \operatorname{mld}(X, D, x).$$

where the infimum runs through all codimension ≥ 1 points x on X.

Various classes of singularities in birational geometry can be defined using the minimal log discrepancy.

Definition 2.2. The pair (X, D) as above is *log canonical* (*lc*) if $mld(X, D) \ge 0$. It is *Kawa-mata log terminal* (*klt*) if mld(X, D) > 0.

When a pair (X, D) is log canonical, we may compute the mld using only the finitely many irreducible divisors appearing in a single log resolution of singularities of (X, D); in particular, it is rational (see, e.g., [20]).

Definition 2.3. A klt Fano variety *X* is *exceptional* if for every effective \mathbb{Q} -divisor *D* that is \mathbb{Q} -linearly equivalent to $-K_X$, the pair (X, D) is klt.

Let *X* be a klt Fano variety. The *lc threshold* of an effective \mathbb{Q} -Cartier \mathbb{Q} -divisor *D* on *X* is defined to be

$$lct(X, D) := \sup\{t \in \mathbb{Q} : (X, tD) \text{ is } lc\}.$$

Let $|-K_X|_{\mathbb{Q}} := \{D \text{ is effective} | D \sim_{\mathbb{Q}} -K_X\}$ be the \mathbb{Q} -linear system of $-K_X$. The α -*invariant* (also called the global log canonical threshold) of X is defined to be

$$\alpha(X) := \inf\{ \operatorname{lct}(X, D) : D \in |-K_X|_{\mathbb{Q}} \}$$

which coincides with

$$\sup\{t \in \mathbb{Q} : (X, tD) \text{ is lc for all } D \in |-K_X|_{\mathbb{Q}}\}.$$

So the α -invariant in this paper is same as the global log canonical threshold in [28], but different from the global log canonical threshold in [22]. From the definitions, we see that α -invariant $\alpha(X) > 1$ implies X is exceptional. The converse holds by [7, Theorem 1.7].

3. LOG DISCREPANCIES OF HYPERSURFACE SINGULARITIES

In this section, we develop the tools needed to compute minimal log discrepancies later in the paper. The main result is a method of computing the minimal log discrepancy of the quotient of a hypersurface in affine space by a finite subgroup of the torus; this method requires that the defining equation for the hypersurface satisfy a Newton non-degeneracy condition.

First, we recall a simple formula for the minimal log discrepancy of a cyclic quotient singularity, which follows from the toric description of the singularity (see, e.g., [3, Section 2]):

Proposition 3.1. Let $X := \frac{1}{r}(b_1, \ldots, b_s)$ denote the quotient \mathbb{A}^s/μ_r , where μ_r acts on \mathbb{A}^s with the indicated weights. Suppose that the singularity is well-formed in the sense that $gcd(r, b_1, \ldots, \hat{b_i}, \ldots, b_s) = 1$ for each $1 \le i \le s$. Then the minimal log discrepancy of $\frac{1}{r}(b_1, \ldots, b_s)$ is given by

(1)
$$\operatorname{mld}(X) = \min\left\{1, \min_{1 \le j < r} \sum_{i=1}^{s} \left(\frac{jb_i}{r} - \left\lfloor \frac{jb_i}{r} \right\rfloor\right)\right\}.$$

Note that this is the global mld of this quotient indicated, not the mld of the point $0 \in \mathbb{A}^s/\mu_r$.

Proof. Since the quotient singularity is well-formed, no divisor in \mathbb{A}^s is fixed by a nontrivial subgroup of μ_r ; hence the pullback of the canonical divisor by the quotient morphism is again the canonical divisor. The quotient \mathbb{A}^s/μ_r is the affine toric variety corresponding to the cone σ generated by the standard basis vectors in \mathbb{R}^s , with toric lattice $N = \mathbb{Z}^s + \mathbb{Z} \cdot \frac{1}{r}(b_1, \ldots, b_s)$. The global minimal log discrepancy of \mathbb{A}^s/μ_r is therefore the minimum of the sum-of-coordinates functional over all points in $N \cap \sigma \setminus \{0\}$ [4, Section 2-A(f)]. Actually, it suffices to take the minimum over lattice points in the unit cube $[0, 1]^s$ (excluding the origin).

As *j* runs from 1 to r - 1, the point with coordinates

$$\frac{jb_i}{r} - \left\lfloor \frac{jb_i}{r} \right\rfloor$$

varies over every non-integral point in this unit cube. The minimum is taken with 1 to account for discrepancies of toric divisors on the original variety \mathbb{A}^s/μ_r , which correspond to the standard basis points.

Next, we develop a method for computing the minimal log discrepancies of certain hypersurfaces in toric varieties. The idea is to refine the fan of the ambient toric variety in such a way that the strict transform of the hypersurface is smooth. The method of resolving hypersurfaces in this way is well-studied and dates back at least to work of Varchenko [30] in the case of hypersurfaces of affine space. This procedure works in much greater generality, however, and has been extended to subvarieties defined by suitable ideals in any affine normal toric variety [5]. We will be concerned with hypersurfaces in the toric variety \mathbb{A}^{n+1}/G , where *G* is a finite cyclic subgroup of the torus acting freely in codimension 1.

Suppose that $Y \coloneqq \{f = 0\} \subset \mathbb{A}^{n+1}$ is a hypersurface passing through the origin and $G \coloneqq \mu_r$ is a finite cyclic group which acts by $\zeta \cdot (x_0, \ldots, x_n) = (\zeta^{a_0} x_0, \ldots, \zeta^{a_n} x_n)$, for any $\zeta \in \mu_r$. Assume further that the action of *G* is free in codimension 1 and preserves *Y*.

The toric variety \mathbb{A}^{n+1}/G corresponds to the fan Δ in the lattice $N = \mathbb{Z}^{n+1} + \mathbb{Z} \cdot \frac{1}{r}(a_0, \ldots, a_n)$ whose cones are the positive orthant σ in $N_{\mathbb{R}} \coloneqq N \otimes_{\mathbb{Z}} \mathbb{R}$ and all its faces. Let M be the dual lattice to N. Then we may write $f = \sum_{m \in M} c_m x^m$ (each $m \in M$ corresponds to a G-invariant monomial in x_0, \ldots, x_n). Denote by $\operatorname{supp}(f)$ the finite set of $m \in M$ for which $c_m \neq 0$.

Our methods will apply to hypersurfaces defined by suitably general polynomials *f*; the correct notion is Newton non-degeneracy (cf. [16, Definition 4.4.22], [5, Definition 1.2]):

Definition 3.2. The *Newton polytope* $\Gamma_+(f)$ of f in $M_{\mathbb{R}} := M \otimes_{\mathbb{Z}} \mathbb{R}$ is defined to be

$$\Gamma_+(f) \coloneqq \operatorname{conv} \left\{ \bigcup_{m \in \operatorname{supp}(f)} (m + \mathbb{R}^{n+1}_{\geq 0}) \right\}.$$

We say that f is *Newton non-degenerate at* 0 if for every compact face γ of $\Gamma_+(f)$, the polynomial $f_{\gamma} \coloneqq \sum_{m \in \gamma} c_m x^m$ satisfies $\operatorname{Sing}(V(f_{\gamma})) \cap T = \emptyset$, where T is the open torus defined by $x_0 \cdots x_n \neq 0$. (We note that the definitions of Newton polytope and Newton non-degeneracy do not depend on G, but only on the polynomial f.)

Given a point $\beta \in N$ and a polynomial $f \in \mathbb{C}[M \cap \sigma^{\vee}]$, we write $\beta(f) \coloneqq \min\{\beta(m) : m \in \operatorname{supp}(f)\}$, where $\beta(m)$ denotes the inner product. Now we can state the main result of this subsection:

Theorem 3.3. Let $(Y, 0) := \{f = 0\} \subset \mathbb{A}^{n+1}$ be a normal hypersurface singularity and $G := \mu_r$ a cyclic subgroup of the torus in \mathbb{A}^{n+1} acting freely in codimension 1. Assume that f is Newton non-degenerate at 0, set (X, x) := (Y, 0)/G, and suppose (X, x) is well-formed in the sense that $\operatorname{Sing}(\mathbb{A}^{n+1}/G) \cap X$ has codimension at least 2 in X. Then

$$\operatorname{mld}(X) = \inf_{\beta \in (N \cap |\Delta|) \setminus \{0\}} [\beta(x_0 \cdots x_n) - \beta(f)].$$

Note that this result computes the global mld of X, not just the mld at x. Moreover, the singularity need not be isolated, though we require that the origin 0 be contained in the hypersurface $\{f = 0\} \subset \mathbb{A}^{n+1}$. Before proving this result, we state a quick corollary, which gives an easy way to check that X is klt (cf. [17, Proposition 2.9]).

Corollary 3.4. Let (X, x) be a singularity satisfying the conditions of Theorem 3.3. If (1, ..., 1) is contained in the interior of the Newton polytope $\Gamma_+(f)$, then X is klt.

Proof. Let $\beta \in (N \cap |\Delta|) \setminus \{0\}$ be a lattice point. Notice that $\beta(x_0 \cdots x_n)$ is the same as the value of the functional $\beta : M_{\mathbb{R}} \to \mathbb{R}$ at the point $(1, \ldots, 1)$. By assumption, β has all nonnegative coordinates in $N_{\mathbb{R}}$, so the minimum value of β on the Newton polytope $\Gamma_+(f)$ must be attained by some monomial m of the equation f. Because $(1, \ldots, 1)$ is in the interior, we must have that $\beta(x_0 \cdots x_n) - \beta(m)$ is positive. Therefore, $\beta(x_0 \cdots x_n) - \beta(f)$ is positive. Since this holds for each β , it follows from Theorem 3.3 that X is klt. \Box

Proof of Theorem 3.3. Since f is Newton non-degenerate, it follows from [5, Theorem 1.3] that there exists a refinement Δ' of the fan Δ , corresponding to a toric birational morphism $\pi_{\Delta'}: V' \to V := \mathbb{A}^{n+1}/G$, such that V' is smooth and the irreducible components of the total transform $\pi_{\Delta'}^{-1}(X)$ are smooth and meet transversely. In particular, the strict transform \tilde{X} of X is smooth. More specifically, the fan Δ' can be any refinement of the dual fan Σ to $\Gamma_+(f)$ for which the corresponding toric variety is smooth. This dual fan Σ is called the *Groebner fan* in [5] (cf. [16, Theorem 4.4.23]). In particular, since any two refinements have a further common refinement, we can arrange that Δ' also refines any other subdivision of Δ .

Since X is a resolution of singularities of X, when X is lc, we can compute its mld using only exceptional divisors on \tilde{X} . In particular, in this case we only need to check the log discrepancies of exceptional divisors that arise from toric modifications of \mathbb{A}^{n+1}/G . Fix a primitive element β in $(N \cap |\Delta|) \setminus \{0\}$, the ray through which we may assume is in Δ' . We may refine Δ with a star-shaped subdivision $\Delta(\beta)$ centered at the ray through β ; this gives a birational morphism $\pi_{\beta} : V_{\beta} \to V$ extracting a unique exceptional divisor E_{β} . Choosing Δ' to also refine $\Delta(\beta)$, we have in our setup successive refinements $\Delta \subset \Delta(\beta) \subset \Delta'$ and corresponding birational morphisms $V' \to V_{\beta} \to V$.

Denote by X_{β} the strict transform of X in V_{β} . Then [16, Proposition 8.3.11] shows that

$$K_{V_{\beta}} = \pi_{\beta}^*(K_V) + (\beta(x_0 \cdots x_n) - 1)E_{\beta}$$

and

$$X_{\beta} = \pi_{\beta}^* X - \beta(f) E_{\beta}.$$

Combining these two equations gives that

$$K_{V_{\beta}} + X_{\beta} + E_{\beta} = \pi_{\beta}^*(K_V + X) + (\beta(x_0 \cdots x_n) - \beta(f))E_{\beta}.$$

Passing to the full refinement Δ' , a very similar equation still holds in a neighborhood of the generic point of the strict transform \tilde{E}_{β} of E_{β} :

$$K_{V'} + \tilde{X} + \tilde{E}_{\beta} = \pi^*_{\Delta'}(K_V + X) + (\beta(x_0 \cdots x_n) - \beta(f))\tilde{E}_{\beta}$$

Since X is well-formed and normal, the adjunction formulas $(K_V + X)|_X = K_X$ and $(K_{V'} + \tilde{X})|_{\tilde{X}} = K_{\tilde{X}}$ hold. Indeed, the canonical class of a normal variety is well-defined as a Weil divisor up to linear equivalence, so we can remove the singular locus from X, which has codimension at least 2 in X by the well-formedness assumptions, and observe that the equalities of divisors hold on the smooth locus. Thus, after restricting to \tilde{X} , we

obtain that the log discrepancy of each component of $\tilde{E}_{\beta} \cap \tilde{X}$ (which is an exceptional divisor of $\tilde{X} \to X$, as \tilde{E}_{β} intersects transversely with \tilde{X}) is $\beta(x_0 \cdots x_n) - \beta(f)$.

Suppose $\beta(x_0 \cdots x_n) - \beta(f)$ is negative for some β . Since this is the log discrepancy of some divisor over X, it follows that X is non-lc, so that the minimal log discrepancy is $-\infty$ (see, e.g., [21, Corollary 2.31]). This agrees with the infimum in Theorem 3.3, for we may replace β with positive multiples to make the difference $\beta(x_0 \cdots x_n) - \beta(f)$ arbitrarily negative. Otherwise, the infimum is a minimum and is nonnegative, so in particular all the log discrepancies of divisors on any resolution \tilde{X} constructed as above are nonnegative. This means X is lc. Since we can compute the mld in this case on just one resolution \tilde{X} , the minimum of $\beta(x_0 \cdots x_n) - \beta(f)$ must be the the minimal log discrepancy.

4. The examples and their well-formedness

The exceptional Fano varieties used to prove Theorem 1.1 are hypersurfaces in certain weighted projective spaces with weights defined using *Sylvester's sequence*. This sequence is defined by $s_0 \coloneqq 2$ and $s_{n+1} \coloneqq s_0 \cdots s_n + 1, n \ge 1$. The first several terms are 2,3,7,43,1807,.... All elements of the sequence are pairwise coprime and satisfy $s_n > 2^{2^{n-1}}$. Their reciprocals also have the following property for each n:

$$\frac{1}{s_0} + \dots + \frac{1}{s_n} = 1 - \frac{1}{s_{n+1} - 1}.$$

The hypersurfaces have slightly different equations in even vs odd dimensions. Theorem 4.1 and Theorem 4.2 summarize their properties in each case.

Theorem 4.1. For every even integer $n \ge 2$, let

$$a_{n} \coloneqq \frac{1}{4}(s_{n}^{2} + s_{n} - 4),$$

$$a_{n+1} \coloneqq \frac{1}{2}((s_{n} - 1)a_{n} - s_{n} - 1),$$

$$d \coloneqq (-1 + a_{n} + a_{n+1})(s_{n} - 1),$$

$$a_{i} \coloneqq \frac{d}{s_{i}}, i = 0, \dots, n - 1,$$

$$b \coloneqq \frac{1}{2}(s_{n}^{2} - s_{n} - 4),$$

$$c \coloneqq \frac{1}{2}(s_{n} + 3),$$

and let X be the hypersurface in $\mathbb{P} = \mathbb{P}(a_0, \ldots, a_{n+1})$ defined by the degree d equation

(2)
$$x_0^2 + x_1^3 + \dots + x_{n-1}^{s_{n-1}} + x_n^b x_{n+1} + x_1 \cdots x_{n-1} x_n x_{n+1}^c = 0.$$

Then X is a well-formed exceptional Fano variety with $-K_X = \mathcal{O}_X(1)$ and

$$\mathrm{mld}(X) = \frac{s_n - 1}{2a_{n+1}}.$$

Theorem 4.2. For every odd integer $n \ge 3$, Let

$$a_{n} \coloneqq \frac{1}{4}(s_{n}^{2} + 3s_{n} - 6),$$

$$a_{n+1} \coloneqq \frac{1}{4}((s_{n} - 3)a_{n} - s_{n} - 1),$$

$$d \coloneqq (-1 + a_{n} + a_{n+1})(s_{n} - 1),$$

$$a_{i} \coloneqq \frac{d}{s_{i}}, i = 0, \dots, n - 1,$$

$$b \coloneqq \frac{1}{4}(s_{n}^{2} - s_{n} - 2),$$

$$c \coloneqq \frac{1}{2}(s_{n} + 5).$$

and let X be the hypersurface in $\mathbb{P} = \mathbb{P}(a_0, \ldots, a_{n+1})$ defined by a degree d equation

(3)
$$x_0^2 + x_1^3 + \dots + x_{n-1}^{s_{n-1}} + x_n^b x_{n+1} + x_1 \cdots x_{n-1} x_n^2 x_{n+1}^c = 0.$$

Then X is a well-formed exceptional Fano variety with $-K_X = \mathcal{O}_X(1)$ and

$$\mathrm{mld}(X) = \frac{s_n - 3}{4a_{n+1}}.$$

We note that substituting in the definition of a_{n+1} in the expressions for the mld yields the form given in Theorem 1.1.

Remark 4.3. In the two theorems above, we choose equations with coefficients all equal to 1 to define X. However, since each equation has the same number of monomials as variables, we could in fact take the coefficients to be general nonzero complex numbers. This is because any hypersurface of this form is isomorphic to X by scaling the variables.

The hypersurfaces in Theorem 4.1 and Theorem 4.2 fail to be quasismooth at the coordinate point of the last variable x_{n+1} . We conjecture that each has the smallest mld among all exceptional Fano varieties of the same dimension n (Conjecture 1.2). As a comparison, [28, Theorem 3.1] defines non-quasismooth hypersurfaces with similar equations in weighted projective spaces with different weights; these are conjectured to have minimum anti-canonical volume among all exceptional Fano varieties of dimension n.

When n = 2, this example is the hypersurface $X_{282} \subset \mathbb{P}(141, 94, 13, 35)$ defined by the equation

$$x_0^2 + x_1^3 + x_2^{19}x_3 + x_1x_2x_3^5 = 0.$$

The mld of this surface is 3/35. This is known to be the smallest possible mld of an exceptional del Pezzo surface by [22, Theorem 1.9(2)]. In particular, this construction realizes the example in [22] as a non-quasismooth hypersurface in weighted projective space.

When n = 3, this example is the hypersurface $X_{227262} \subset \mathbb{P}^4(113631, 75754, 32466, 493, 4919)$ defined by the equation

$$x_0^2 + x_1^3 + x_2^7 + x_3^{451}x_4 + x_1x_2x_3^2x_4^{24} = 0.$$

11

The mld of this threefold is 10/4919. The best known explicit lower bound on the mld of an exceptional Fano threefold is $1/I_0$, where $I_0 \approx 10^{10^{10.8}}$ [8, Theorem 1.4]. We expect the example above to achieve the optimal bound. To the authors' knowledge, the Fano threefold with smallest mld previously known was Totaro's exceptional threefold of small anti-canonical volume [28, p. 8]. That example has mld 41/17629 > 10/4919.

When n = 4, this example is a hypersurface in a certain weighted projective space of dimension 5 defined by the equation

$$x_0^2 + x_1^3 + x_2^7 + x_3^{43} + x_4^{1631719}x_5 + x_1x_2x_3x_4x_5^{905} = 0.$$

The mld of this fourfold is 903/737536085.

Proof of Theorem 4.1 and Theorem 4.2. The proof of the properties of the examples defined in the two theorems will occupy the next few sections. By Lemma 4.4, the polynomials defining the hypersurfaces have degree d. By Lemma 4.5, the hypersurfaces are well-formed. Theorem 5.1 computes the mld. Theorem 6.5 and Theorem 6.6 show the hypersurfaces are exceptional.

We'll begin by confirming that the weights and equations above do in fact define a hypersurface of the indicated degree.

Lemma 4.4. For each $n \ge 2$, $a_0 + \cdots + a_{n+1} = d + 1$ and the polynomial in (2) or (3) is homogeneous of degree d.

Proof. First, the sum of the weights is d + 1 for all n:

$$a_0 + \dots + a_{n+1} = \frac{d}{s_0} + \dots + \frac{d}{s_{n-1}} + a_n + a_{n+1}$$
$$= \frac{s_n - 2}{s_n - 1} (-1 + a_n + a_{n+1})(s_n - 1) + (-1 + a_n + a_{n+1}) + 1$$
$$= (s_n - 2 + 1)(-1 + a_n + a_{n-1}) + 1 = d + 1.$$

It's clear that $x_0^2, x_1^3, \ldots, x_{n-1}^{s_{n-1}}$ are monomials of degree d in both the even and odd cases. We need to check that the last two monomials have degree d. We begin with the case of n even. For the monomial $x_n^b x_{n+1}$, observe that

$$d - a_{n+1} = (s_n - 1)a_n + (s_n - 2)a_{n+1} - s_n + 1$$

= $(s_n - 1)a_n + (s_n - 2)\frac{1}{2}((s_n - 1)a_n - s_n - 1) - s_n + 1$
= $(s_n - 1)a_n + \frac{1}{2}(s_n^2 - 3s_n + 2)a_n - \frac{1}{2}(s_n^2 - s_n - 2) - s_n + 1$
= $\frac{1}{2}(s_n^2 - s_n)a_n - \frac{1}{2}(s_n^2 + s_n - 4)$
= $\frac{1}{2}(s_n^2 - s_n)a_n - 2a_n = \frac{1}{2}(s_n^2 - s_n - 4)a_n = ba_n.$

Finally, for the monomial $x_1 \cdots x_{n-1} x_n x_{n+1}^c$, note that:

$$d - a_1 - \dots - a_n = \left(1 - \frac{1}{s_1} - \dots - \frac{1}{s_{n-1}}\right) d - a_n$$

= $\left(\frac{1}{2} + \frac{1}{s_n - 1}\right) d - a_n$
= $\frac{s_n + 1}{2}a_{n+1} + \frac{s_n - 1}{2}a_n - \frac{s_n + 1}{2}$
= $\frac{s_n + 1}{2}a_{n+1} + \frac{1}{2}((s_n - 1)a_n - s_n - 1))$
= $\frac{s_n + 1}{2}a_{n+1} + a_{n+1}$
= $\frac{1}{2}(s_n + 3)a_{n+1} = ca_{n+1}.$

This completes the proof that the equation in (2) is weighted homogeneous of degree d for n even. Next, we treat the case of odd n. For the monomial $x_n^b x_{n+1}$, we have

$$d - a_{n+1} = (s_n - 1)a_n + (s_n - 2)a_{n+1} - s_n + 1$$

= $(s_n - 1)a_n + (s_n - 2)\frac{1}{4}((s_n - 3)a_n - s_n - 1) - s_n + 1$
= $(s_n - 1)a_n + \frac{1}{4}(s_n^2 - 5s_n + 6)a_n - \frac{1}{4}(s_n^2 - s_n - 2) - s_n + 1$
= $\frac{1}{4}(s_n^2 - s_n + 2)a_n - \frac{1}{4}(s_n^2 + 3s_n - 6)$
= $\frac{1}{4}(s_n^2 - s_n + 2)a_n - a_n = \frac{1}{2}(s_n^2 - s_n - 2)a_n = ba_n.$

Finally, for the monomial $x_1 \cdots x_{n-1} x_n^2 x_{n+1}^c$, note that

$$d - a_1 - \dots - a_{n-1} - 2a_n = \left(1 - \frac{1}{s_1} - \dots - \frac{1}{s_{n-1}}\right) d - 2a_n$$

$$= \left(\frac{1}{2} + \frac{1}{s_n - 1}\right) d - 2a_n$$

$$= \frac{s_n + 1}{2}a_{n+1} + \frac{s_n - 3}{2}a_n - \frac{s_n + 1}{2}$$

$$= \frac{s_n + 1}{2}a_{n+1} + \frac{1}{2}((s_n - 3)a_n - s_n - 1))$$

$$= \frac{s_n + 1}{2}a_{n+1} + 2a_{n+1}$$

$$= \frac{1}{2}(s_n + 5)a_{n+1} = ca_{n+1}.$$

This completes the proof that the equation (3) is weighted homogeneous of degree d when n is odd.

Next, we'll prove that the hypersurface *X* is well-formed, normal, and Fano with $K_X = O_X(-1)$. The notations are as in Theorem 4.1 and Theorem 4.2.

Lemma 4.5. The weighted projective space \mathbb{P} and the hypersurface X are well-formed; in fact, we have $gcd(a_i, a_{n+1}) = 1$ for each i = 0, ..., n and $gcd(a_i, a_n) = 1$ for each i = 0, ..., n - 1. Moreover, X is quasismooth away from the coordinate point of the last variable x_{n+1} and $K_X = \mathcal{O}_X(-1)$.

Proof. We begin by verifying that the ambient weighted projective spaces \mathbb{P} are well-formed, meaning that $gcd(a_0, \ldots, \hat{a_j}, \ldots, a_{n+1}) = 1$ for all $j = 0, \ldots, n+1$. We'll actually prove the stronger gcd conditions in the lemma, which clearly imply well-formedness. Through the lemma, we use the notation $r := -1 + a_n + a_{n+1} = d/(s_n - 1)$.

Because of the factors of 1/2 and 1/4 in the definitions, we can first show that a_n and a_{n+1} are odd. For $n \ge 1$, we observe by induction that $s_n \equiv 7 \pmod{8}$ when n is even and that $s_n \equiv 3 \pmod{8}$ when n is odd. Then for n even, $s_n^2 + s_n - 4 \equiv 4 \pmod{8}$, and for n odd, $s_n^2 + 3s_n - 6 \equiv 4 \pmod{8}$. It follows that a_n is always odd. Since $ba_n + a_{n+1} = d$ and d is even, a_{n+1} is odd as well.

Next, we prove $gcd(r, a_n) = 1$. Suppose by way of contradiction that a prime p divides r and a_n . By the above, we may assume p is odd. The equation $r = -1 + a_n + a_{n+1}$ yields $a_{n+1} \equiv 1 \pmod{p}$. Now we split into two cases. For n even, $2a_{n+1} = (s_n - 1)a_n - s_n - 1$ yields $2 \equiv -s_n - 1 \pmod{p}$ so p divides $s_n + 3$. But $4a_n = s_n^2 + s_n - 4 = (s_n + 3)(s_n - 2) + 2 \equiv 2 \pmod{p}$, so $0 \equiv 2 \pmod{p}$, a contradiction. Similarly, for n odd, $4a_{n+1} \equiv (s_n - 3)a_n - s_n - 1$ yields $4 \equiv -s_n - 1 \pmod{p}$, so p divides $s_n + 5$. But $4a_n = s_n^2 + 3s_n - 6 = (s_n + 5)(s_n - 2) + 4 \equiv 4 \pmod{p}$, so $4 \equiv 0 \pmod{p}$, a contradiction.

To show $gcd(r, a_{n+1}) = 1$, suppose that p odd divides r and a_{n+1} . This gives $a_n \equiv 1 \pmod{p}$. For n even, $2a_{n+1} = (s_n - 1)a_n - s_n - 1 \equiv s_n - 1 - s_n - 1 = -2 \pmod{p}$, so $0 \equiv -2 \pmod{p}$, a contradiction. For n odd, we similarly get $4a_{n+1} = (s_n - 3)a_n - s_n - 1 \equiv s_n - 3 - s_n - 1 = -4 \pmod{p}$ so $0 \equiv -4 \pmod{p}$, a contradiction.

Next, we calculate $gcd(s_n - 1, a_n) = 1$. Assume an odd prime p divides both. When n is even, $4a_n = s_n^2 + s_n - 4 = (s_n - 1)(s_n + 2) - 2$ shows that $0 \equiv -2 \pmod{p}$. When n is odd, $4a_n = s_n^2 + 3s_n - 6 = (s_n - 1)(s_n + 4) - 2$ shows $0 \equiv -2 \pmod{p}$. This is a contradiction in either case.

Similarly, we can show $gcd(s_n - 1, a_{n+1}) = 1$. Assume an odd prime p divides both. When n is even, we use $2a_{n+1} = (s_n - 1)(-1 + a_n) - 2 \equiv -2 \pmod{p}$, implying $0 \equiv -2 \pmod{p}$. When n is odd, $4a_{n+1} = (s_n - 3)a_n - s_n - 1 = (s_n - 1)a_n - 2a_n - s_n - 1 \equiv -2a_n - 2 \pmod{p}$. Hence p divides $2a_n + 2$. But

$$2a_n + 2 = \frac{1}{2}(s_n^2 + 3s_n - 6) + 2 = \frac{1}{2}(s_n - 1)(s_n + 4) + 1 \equiv 1 \pmod{p}$$

This means $0 \equiv 1 \pmod{p}$. In either case, we have a contradiction.

Lastly, we'll show $gcd(a_n, a_{n+1}) = 1$. Assume *p* odd divides both. When *n* is even, notice that $2a_{n+1} \equiv -s_n - 1 \pmod{p}$ so *p* divides $s_n + 1$. But this would mean

$$a_n = \frac{1}{4}(s_n(s_n+1)-4) \equiv -1 \pmod{p}$$

a contradiction. When *n* is odd, we also get $p|s_n + 1$, but

$$a_n = \frac{1}{4}((s_n+1)(s_n+2)-8) \equiv -2 \pmod{p},$$

so $0 \equiv 2 \pmod{p}$, a contradiction.

This concludes the proof of all the gcd statements in the lemma. Indeed, $gcd(r, a_n) = gcd(s_n - 1, a_n) = 1$ implies $gcd(a_i, a_n) = 1$ for each i = 0, ..., n - 1, and similarly for a_{n+1} . In particular, \mathbb{P} is well-formed.

The hypersurface X is also well-formed because \mathbb{P} is well-formed, and the presence of the monomials $x_0^2, \ldots, x_{n-1}^{s_{n-1}}, x_n^b x_{n+1}$ means that only the coordinate points of x_n and x_{n+1} are contained in X among all the toric strata of X. Hence the hypersurface X does not contain any stratum of $Sing(\mathbb{P})$ of codimension 2 in \mathbb{P} . Each of these coordinate points point has codimension at least 2 in X since $dim(X) \ge 2$, so X is well-formed.

In light of Remark 4.3, we can take X to be general in the linear system defined by the monomials of the defining equation. The only base points for this linear system on \mathbb{A}^{n+2} are multiples of $(0, \ldots, 1, 0)$ and $(0, \ldots, 0, 1)$. Hence X is quasismooth away the coordinate points of the last two coordinates. Since $x_n^b x_{n+1}$ is a monomial in the equation for X, a general equation of the linear system will have a non-vanishing derivative at $(0, \ldots, 1, 0)$, so X is also quasismooth at the coordinate point of x_n . Hence only the coordinate point of x_{n+1} is a non-quasismooth point. Since X is a hypersurface and the non-quasismooth locus has codimension at least 2 in X, it follows that X is normal. Indeed, we can verify the normality of X in each affine chart $\{x_i \neq 0\} = \mathbb{A}^{n+1}/\mu_{a_i}$ of \mathbb{P} by Serre's criterion: hypersurfaces in \mathbb{A}^{n+1} are Cohen-Macaulay, the singular locus of the defining equation in this affine space has codimension at least 2, and the quotient by μ_{a_i} preserves normality.

We've shown that *X* is normal and well-formed. Therefore, the adjunction formula holds for *X* (cf. [28, p. 3]) and $K_X = \mathcal{O}_X(d - a_0 - \cdots - a_{n+1}) = \mathcal{O}_X(-1)$.

5. COMPUTATION OF THE MINIMAL LOG DISCREPANCY

In this section, we compute the exact value of mld(X) in each dimension, where X is the hypersurface from Theorems 4.1 and 4.2. We continue to use all notation from Section 4. The value of the mld comes from computing the difference $\beta(x_0 \cdots x_n) - \beta(f)$ from Theorem 3.3 for a particular hypersurface and lattice point β . Much of the length of the proof is then spent confirming that this is actually the smallest log discrepancy achieved by any divisor over X.

Theorem 5.1. *The minimal log discrepancy of the hypersurface X is*

$$\operatorname{mld}(X) = \begin{cases} \frac{s_n - 1}{2a_{n+1}} & n \text{ is even,} \\ \frac{s_n - 3}{4a_{n+1}} & n \text{ is odd.} \end{cases}$$

This mld is achieved at the unique non-quasismooth point of X.

Proof. For every n, we know from Lemma 4.5 that the hypersurface X is normal and that it is quasismooth away from the coordinate point of the last coordinate, x_{n+1} . Away from this point, X has only cyclic quotient singularities and we can use Proposition 3.1

to compute the mld. At the non-quasismooth point, we'll apply Theorem 3.3 to compute the mld; this will turn out to be the global minimum.

In the affine chart where $x_{n+1} \neq 0$, we may take $x_{n+1} = 1$, and X is locally a quotient of the hypersurface in affine space $\mathbb{A}_{x_0,\dots,x_n}^{n+1}$ defined by the equation

$$f \coloneqq x_0^2 + x_1^3 + \dots + x_{n-1}^{s_{n-1}} + x_n^b + x_1 \cdots x_{n-1} x_n = 0$$

if n is even, and

$$f \coloneqq x_0^2 + x_1^3 + \dots + x_{n-1}^{s_{n-1}} + x_n^b + x_1 \cdots x_{n-1} x_n^2 = 0$$

if *n* is odd. The quotient is by $\mu_{a_{n+1}}$, acting by $\zeta \cdot (x_0, \ldots, x_n) = (\zeta^{a_0} x_0, \ldots, \zeta^{a_n} x_n)$ for $\zeta \in \mu_{a_{n+1}}$. We already know \mathbb{P} and *X* are well-formed and normal, so this hypersurface of affine space satisfies the normality and well-formedness conditions of Theorem 3.3. Furthermore, by Remark 4.3, we can take the coefficients of the monomials in the example to be general, so that Newton non-degeneracy is also automatic (the base locus of any subset of the monomials in the equation is contained in the complement of the open torus orbit).

Therefore, Theorem 3.3 applies, and we can compute the mld of X in this affine chart by finding the minimum of $\beta(x_0 \cdots x_n) - \beta(f)$ as β varies over nonzero points of the lattice $N = \mathbb{Z}^{n+1} + \mathbb{Z} \cdot \frac{1}{a_{n+1}}(a_0, \ldots, a_n)$ which are contained in the positive orthant (minus the origin). It's not difficult to show that this difference is a positive integer on nonzero integral points, so we can reduce to looking at points $\beta_j := (\beta_j^0, \ldots, \beta_i^n)$ where

$$\beta_j^i = \left\{ \frac{ja_i}{a_{n+1}} \right\}.$$

for each i = 0, ..., n. (Here $\{x\}$ denotes the fractional part $x - \lfloor x \rfloor$.) and j is a positive integer satisfying $1 \le j \le a_{n+1} - 1$. These points of N are all in the interior of the cube $[0,1]^{n+1}$; this is because $gcd(a_i, a_{n+1}) = 1$ for each i = 0, ..., n, as shown in Lemma 4.5. Points in the interior correspond to exceptional divisors which have center equal to the non-quasismooth point, so the mld we will find in the proof occurs at that point. This is consistent with the fact that the non-quasismooth point is an isolated singularity of X. For the rest of the proof, we separately consider the cases of when n is even and when n is odd.

Case 1: *n* is even.

Since the equation of X is weighted homogeneous of degree d for the weights a_0, \ldots, a_{n+1} , each monomial m of the polynomial f above has degree d modulo a_{n+1} with the same weights. This implies that for each monomial m with nonzero coefficient in f, $\beta_j(m) \equiv jd/a_{n+1} \pmod{\mathbb{Z}}$. However,

$$d = (-1 + a_n + a_{n+1})(s_n - 1) \equiv (-1 + a_n)(s_n - 1) = 2a_{n+1} + 2 \equiv 2 \pmod{a_{n+1}}.$$

It follows that $\beta_j(m) \equiv 2j/a_{n+1} \pmod{\mathbb{Z}}$. We note that for $m = x_0^2$, $\beta_j(m)$ is actually equal to $2j/a_{n+1}$ for any $j = 1, \ldots, a_{n+1} - 1$. Indeed, the same reasoning as above gives that $a_0 = d/2 \equiv 1 \pmod{a_{n+1}}$ so $\beta_j^0 = j/a_{n+1}$. Then $\beta_j(x_0^2)$ is twice this value.

Similarly, $a_0 + \cdots + a_n \equiv d + 1 \equiv 3 \pmod{a_{n+1}}$, so $\beta_j(x_0 \cdots x_n)$ is congruent to $3j/a_{n+1}$ modulo \mathbb{Z} . The difference $\beta_j(x_0 \cdots x_n) - \beta_j(f)$ that we're interested in must therefore have

fractional part

$$\frac{3j}{a_{n+1}} - \frac{2j}{a_{n+1}} = \frac{j}{a_{n+1}}$$

Because $\beta(x_0 \cdots x_n) - \beta(x_1 \cdots x_n) = \beta^0 \ge 0$ for any β in the positive orthant, we furthermore know that the difference is positive. Hence $\beta_j(x_0 \cdots x_n) - \beta_j(f)$ is bounded below by j/a_{n+1} for each j. In particular, X is klt.

This lower bound is achieved when $j = j_0 \coloneqq (s_n - 1)/2$. Indeed, we already know $\beta_{j_0}(x_0^2) = 2j_0/a_{n+1} = s_{n-1}/a_{n+1}$, and since $2j_0 < a_{n+1}$, this is a fraction between 0 and 1. Thus, it actually equals the minimum $\beta_{j_0}(f)$. Next we verify $\beta_{j_0}(x_0 \cdots x_n) = 3j_0/a_{n+1} = (3s_n - 3)/(2a_{n+1})$. For $i = 0, \ldots, n - 1$:

$$j_0 a_i = \frac{s_n - 1}{s_i} \frac{d}{2} \equiv \frac{s_n - 1}{s_i} \pmod{a_{n+1}},$$

so

$$\beta_{j_0}^i = \frac{s_n - 1}{s_i a_{n+1}}.$$

For i = n, we have

$$j_0 a_n = \frac{1}{2}((s_n - 1)a_n) = a_{n+1} + \frac{1}{2}(s_n + 1) \equiv \frac{1}{2}(s_n + 1) \pmod{a_{n+1}}.$$

This proves

$$\beta_{j_0}^n = \frac{s_n + 1}{2a_{n+1}}.$$

Adding these together yields

$$\beta_{j_0}(x_0 \cdots x_n) = \frac{s_n - 1}{s_0 a_{n+1}} + \dots + \frac{s_n - 1}{s_{n-1} a_{n+1}} + \frac{s_n + 1}{2a_{n+1}}$$
$$= \frac{(s_n - 1)(s_n - 2)}{(s_n - 1)a_{n+1}} + \frac{s_n + 1}{2a_{n+1}}$$
$$= \frac{3s_n - 3}{2a_{n+1}},$$

as claimed. Subtracting,

$$\beta_{j_0}(x_0\cdots x_n) - \beta_{j_0}(f) = \frac{s_n - 1}{2a_{n+1}},$$

proving that there is a divisor over X with log discrepancy equal to $(s_n - 1)/(2a_{n+1})$.

It remains to see that this is actually the minimum value of $\beta_j(x_0 \cdots x_n) - \beta_j(f)$ for any $j = 1, \ldots, a_{n+1} - 1$. In light of the lower bound j/a_{n+1} for this difference, we need only verify that the difference is larger for $1 \le j < j_0$. For such a j,

$$\beta_j^0 + \dots + \beta_j^{n-1} \ge \left\{ \frac{1}{a_{n+1}} \left(\frac{dj}{s_0} + \dots + \frac{dj}{s_{n-1}} \right) \right\}$$
$$= \left\{ \frac{1}{a_{n+1}} \frac{(s_n - 2)dj}{s_n - 1} \right\}$$
$$= \left\{ \frac{1}{a_{n+1}} (s_n - 2)(-1 + a_n + a_{n+1})j \right\}$$

16

Modulo a_{n+1} we have

$$(s_n - 2)(-1 + a_n + a_{n+1}) \equiv (s_n - 2)(-1 + a_n) = (s_n - 1)(-1 + a_n) - (-1 + a_n)$$
$$= 2a_{n+1} + 2 - (-1 + a_n) \equiv 3 - a_n \pmod{a_{n+1}}.$$

Since $a_{n+1} > a_{n+1} + (3 - a_n)j > 3j$ for $1 \le j < j_0$ by definition of a_{n+1} ,

$$\beta_j(x_0\cdots x_n) = (\beta_j^0 + \cdots + \beta_j^{n-1}) + \beta_j^n \ge \frac{a_{n+1} + (3-a_n)j}{a_{n+1}} + \beta_j^n > \frac{3j}{a_{n+1}}.$$

Since $\beta_j(x_0 \cdots x_n)$ is equivalent to $3j/a_{n+1}$ modulo \mathbb{Z} , it follows that $\beta_j(x_0 \cdots x_n) \ge 1 + 3j/a_{n+1}$ for $1 \le j < j_0$. Therefore, the difference $\beta_j(x_0 \cdots x_n) - \beta_j(f)$ is greater than 1 for this range of j values. This completes the argument.

Case 2: *n* is odd.

The argument for odd *n* is very similar to the argument for even *n*. We note that

$$d = (-1 + a_n + a_{n+1})(s_n - 1) \equiv (-1 + a_n)(s_n - 3) + 2(-1 + a_n)$$
$$\equiv 4a_{n+1} + 4 + 2(-1 + a_n) \equiv 2 + 2a_n \pmod{a_{n+1}}.$$

For each monomial m in f, we have that $\beta_j(m)$ is equivalent to $(2 + 2a_n)/a_{n+1}$ modulo \mathbb{Z} . The sum of the weights is still d + 1, so it's again true that $\beta_j(x_0 \cdots x_n) - \beta_j(f) \equiv j/a_{n+1} \pmod{\mathbb{Z}}$. (mod \mathbb{Z}). For any j,

$$\beta_j(x_0\cdots x_n) - \beta_j(f) \ge \beta_j(x_0\cdots x_n) - \beta(x_0^2) = \beta_j^1 + \cdots + \beta_j^n - \beta_j^0.$$

But it's also true that

$$\beta_j(x_0\cdots x_n) - \beta_j(f) \ge \beta(x_0\cdots x_n) - \beta(x_1\cdots x_{n-1}x_n^2) = \beta_j^0 - \beta_j^n.$$

Since all the β_j^i are positive for $1 \le j \le a_{n+1} - 1$ (each point is in the interior of the unit cube), one of these two differences is positive. It follows that *X* is klt.

Next, we confirm that the claimed mld value is attained when $j = j_0 := (s_n - 3)/4$. First, for i = 0, ..., n - 1,

$$j_0 a_i = \frac{s_n - 3}{4} \frac{d}{s_i} = \frac{(s_n - 3)(s_n - 1)(-1 + a_n + a_{n+1})}{4s_i} \equiv \frac{(s_n - 3)(-1 + a_n)(s_n - 1)}{4s_i}$$
$$= \frac{(4a_{n+1} + 4)(s_n - 1)}{4s_i} \equiv \frac{s_n - 1}{s_i} \pmod{a_{n+1}}.$$

This shows

$$\beta_{j_0}^i = \frac{s_n - 1}{s_i a_{n+1}}.$$

Similarly, for i = n,

$$j_0 a_n = \frac{1}{4}(s_n - 3)a_n = a_{n+1} + \frac{1}{4}(s_n + 1) \equiv \frac{1}{4}(s_n + 1) \pmod{a_{n+1}}.$$

This proves

$$\beta_{j_0}^n = \frac{s_n + 1}{4a_{n+1}}.$$

Now we can compute

$$\beta_{j_0}(x_0 \cdots x_n) = \frac{s_n - 1}{s_0 a_{n+1}} + \dots + \frac{s_n - 1}{s_{n-1} a_{n+1}} + \frac{s_n + 1}{4a_{n+1}}$$
$$= \frac{(s_n - 1)(s_n - 2)}{(s_n - 1)a_{n+1}} + \frac{s_n + 1}{4a_{n+1}}$$
$$= \frac{5s_n - 7}{4a_{n+1}}.$$

Similarly,

$$\beta_{j_0}(x_1 \cdots x_{n-1} x_n^2) = \frac{s_n - 1}{s_1 a_{n+1}} + \dots + \frac{s_n - 1}{s_{n-1} a_{n+1}} + \frac{s_n + 1}{2a_{n+1}}$$
$$= \frac{(s_n - 3)(s_n - 1)}{2(s_n - 1)a_{n+1}} + \frac{s_n + 1}{2a_{n+1}}$$
$$= \frac{s_n - 1}{a_{n+1}}.$$

Subtracting these gives

$$\beta_{j_0}(x_0\cdots x_n) - \beta_{j_0}(x_1\cdots x_{n-1}x_n^2) = \frac{s_n - 3}{4a_{n+1}}$$

We already know that this is the correct value of the log discrepancy modulo \mathbb{Z} and this value is between 0 and 1, so in fact $\beta_{j_0}(x_0 \cdots x_n) - \beta_{j_0}(f) = (s_n - 3)/(4a_{n+1})$. It remains to check that this is in fact the lowest value attained for any j. As in the even case, we only need to verify that no lower value is attained in the range $1 \le j < j_0$. To see this, first note that,

$$\beta_j(x_0\cdots x_n) - \beta_j(f) \ge \beta_j(x_0\cdots x_n) - \beta_j(x_0^2) = \beta_j^n - \beta_j^0 + \beta_j^1 + \cdots + \beta_j^{n-1}.$$

Since $\beta_j^0 - \beta_j^n$ is also of the form $\beta_j(x_0 \cdots x_n) - \beta_j(m)$ for the monomial $m = x_1 \cdots x_{n-1} x_n^2$ of f, this difference is equivalent to j/a_{n+1} modulo \mathbb{Z} . Since both β_j^0 and β_j^n are between 0 and 1, it's simple to see that $\beta_j^n - \beta_j^0 \ge -j/a_{n+1}$ (and again is equivalent to it modulo \mathbb{Z}).

Therefore, the only way that the difference $\beta_j(x_0 \cdots x_n) - \beta_j(x_0^2)$ could achieve the minimum possible value j/a_{n+1} is if $\beta_j^1 + \cdots + \beta_j^{n-1} = 2j/a_{n+1}$. Suppose this holds, for some particular j. This would imply that

$$\beta_j(x_1^3) = 3\beta_j^1 \le 3\frac{2j}{a_{n+1}} = \frac{6j}{a_{n+1}}.$$

In comparison, we can calculate

$$\beta_j^0 = \frac{j(1+a_n)}{a_{n+1}}$$

for any j in this range we're considering, because $a_0 = d/2 \equiv 1 + a_n \pmod{a_{n+1}}$ and $j(1 + a_n) < a_{n+1}$ when $1 \le j < j_0$. But then

$$\beta_j(x_0\cdots x_n) - \beta_j(f) \ge \beta_j(x_0\cdots x_n) - \beta_j(x_1^3) \ge \frac{j(1+a_n)}{a_{n+1}} - \frac{6j}{a_{n+1}} = \frac{(a_n-5)j}{a_{n+1}}$$

This last expression is greater than j/a_{n+1} because $a_n > 6$. Since the discrepancy is equivalent to j/a_{n+1} modulo \mathbb{Z} , it must be greater than 1 for the given value of j.

So far, we've demonstrated that mld(X, x) equals the value from Theorem 5.1 when x is the unique non-quasismooth point of X. However, we also need to check that no smaller mld is attained at any quasismooth point of the variety X. The following lemma finishes the proof.

Lemma 5.2. Let $x \in X$ be the non-quasismooth point of the hypersurface X defined above. Then $mld(X \setminus x) \ge 2/a_n$.

Proof of Lemma 5.2: We can take hypersurface X to be a general member of a linear series and it is quasismooth away from x so the singularities on all toric strata apart from xare the quotient singularities described in, e.g., [11, Proposition 2.6]. In particular, the singularity type is constant along the intersection of X with any toric stratum of \mathbb{P} , and we need only check the mld of the singularities on 1-dimensional toric strata in \mathbb{P} (when neither coordinate point in its closure is in X) or at coordinate points of \mathbb{P} contained in X. This is in the sense of Proposition 3.1: every quotient singularity on a larger stratum of Xalready appears somewhere on the affine toric variety corresponding to the singularities on smaller strata. In particular, we look at quotient singularities at points y in X such that: 1) the only nonzero coordinates of y are x_{i_1} and x_{i_2} for some i_1, i_2 among $0, \ldots, n-1$, or 2) y is the coordinate point of x_n .

In case 1), the singularity type is always of type $\frac{1}{mr}(\ldots, a_n, a_{n+1})$, where $r = -1 + a_n + a_{n+1}$ and mr the gcd of two of the weights among a_0, \ldots, a_{n-1} . The multiple m here divides $s_n - 1$, but is at most $(s_n - 1)/6$. We omit the other weights in the singularity, because it will actually be enough to consider the contributions of the last two in the formula of Proposition 3.1. The sum of the weights in the singularity must equal $d+1 \equiv 1 \pmod{mr}$, so the sum in (1) is equivalent to $j/(mr) \mod \mathbb{Z}$ for each j.

When $j < mr/a_n$, $ja_n < mr$, so that

$$\frac{ja_n}{mr} - \left\lfloor \frac{ja_n}{mr} \right\rfloor = \frac{ja_n}{mr} > \frac{j}{mr}.$$

The left hand side is just one term in the sum (1), so the total sum is greater than 1 for this *j*. The first multiple of a_n which is greater than mr occurs when $j = m(s_n + 1)/2$, for *n* even, and when $j = m(s_n + 1)/4$, for *n* odd. Indeed, for the even case,

$$\frac{1}{2}m(s_n+1)a_n - mr = m\left(\frac{1}{2}(s_n+1)a_n + 1 - a_n - \frac{1}{2}((s_n-1)a_n - s_n - 1)\right) = \frac{1}{2}m(s_n+3),$$

which is between 0 and a_n , since $m \le (s_n - 1)/6$. Similarly, for the odd case,

$$\frac{1}{4}m(s_n+1)a_n - mr = m\left(\frac{1}{4}(s_n+1)a_n + 1 - a_n - \frac{1}{4}((s_n-3)a_n - s_n - 1)\right) = \frac{1}{4}m(s_n+5),$$

which has the same property. In both equations above, the right-hand side is the numerator of the difference $ja_n/(mr) - \lfloor ja_n/(mr) \rfloor$, for the indicated special value of j. Note that it is still greater than j in both cases, so the sum in (1) is greater than 1 for this j too. As j increases further, the contribution from a_n in the sum in (1) will continue to be larger than j/(mr) until, at least, ja_n reaches 2mr. This proves that the minimal log discrepancy of any quotient singularity of this type is at least

$$\frac{2mr/a_n}{mr} = \frac{2}{a_n},$$

as required.

The remaining singularity from case (2) is even simpler. At the coordinate point of x_n , the singularity type is $\frac{1}{a_n}(a_0, \ldots, a_{n-1})$. By Lemma 4.5, $gcd(a_i, a_n) = 1$ for each $i = 0, \ldots, n-1$, so the sum (1) always has n positive terms for each $j = 1, \ldots, a_n - 1$. This already proves that the mld is at least $n/a_n \ge 2/a_n$.

Theorem 5.1 is now proved by noting that $2/a_n$ is larger than the mld found at the non-quasismooth point for all *n*.

6. COMPUTATION OF THE ALPHA INVARIANT AND EXCEPTIONALITY

In this section, we demonstrate that the Fano varieties defined in Theorem 4.1 and Theorem 4.2 are exceptional.

Let X be a hypersurface in $\mathbb{A}^{n+1}_{\mathbb{C}}$ that contains the origin. For positive integers c_0, \ldots, c_n , we take $f : X' \to X$ to be the stack-theoretic weighted blow-up of $X \subset \mathbb{A}^{n+1}$ at the origin with the given weights $c = (c_0, \ldots, c_n)$. Section 3.4 in [1] gives explicit coordinate charts. We define the *weighted tangent cone* X^c to be the exceptional divisor in X'. In particular, we view X^c as a hypersurface in the stack $\mathbb{P}^n(c_0, \ldots, c_n) = [(\mathbb{A}^{n+1} - 0)/\mathbb{G}_m]$.

For positive integers c_0, \ldots, c_n , the *weighted multiplicity* $\operatorname{mult}_c S$ of a closed subscheme $S \subset \mathbb{A}^{n+1}$ at the origin is defined to be the degree of the weighted tangent cone S^c as a substack of the weighted projective space $\mathbb{P}^n(c_0, \ldots, c_n)$. Equivalently, let $F^l \mathcal{O}(S)$ be the linear span of the monomials x^I with $\sum_{j=0}^n c_j i_j \ge l$. This gives a decreasing filtration of the ring $\mathcal{O}(S)$ of regular functions. Then the weighted multiplicity is the limit

$$\lim_{l \to \infty} \frac{\dim(\mathcal{O}(S)/F^l\mathcal{O}(S))}{l^m/m!}$$

where m is the dimension of S. If all weights are 1, this limit equals the usual multiplicity of S at the origin [13, Example 4.3.1].

Lemma 6.1. For each even integer $n \ge 2$, the hypersurface S in \mathbb{A}^{n+1} defined by the equation

$$x_0^2 + x_1^3 + \dots + x_{n-1}^{s_{n-1}} + x_1 \cdots x_n = 0$$

is canonical, and in particular klt.

Proof. The number of monomials in this equation is the same as the number of variables, so by the same reasoning as in Remark 4.3, we may actually replace S with a hypersurface defined by a general linear combination of the monomials in the equation above (this new hypersurface is isomorphic to S). The base locus of the linear system generated by the monomials is the line spanned by (0, ..., 0, 1), so by Bertini's theorem, the surface S is smooth in codimension 1. It is also Cohen-Macaulay since it is a hypersurface in affine space, so it is normal by Serre's criterion.

20

Now we show that the convex hull of the points $(2, 0, ..., 0), (0, 3, 0, ..., 0), ..., (0, ..., 0, s_{n-1}, 0)$ and (0, 1, ..., 1) in \mathbb{R}^{n+1} contains a point with all coordinates less than 1. Indeed, the point

$$\frac{5}{12}(2,0,\ldots,0) + \frac{1}{6}(0,3,0,\ldots,0) + \frac{5}{12}(0,1,\ldots,1)$$

has all coordinates less than 1. So the Newton polytope of $x_0^2, x_1^3, \ldots, x_{n-1}^{s_{n-1}}$ and $x_1 \cdots x_n$ contains $(1, \ldots, 1)$ in the interior. Corollary 3.4 implies that *S* is klt. Since *S* is Gorenstein, it is canonical.

Lemma 6.2. For each odd integer $n \ge 3$, the hypersurface S in \mathbb{A}^{n+1} defined by the equation

$$x_0^2 + x_1^3 + \dots + x_{n-1}^{s_{n-1}} + x_1 \dots + x_n^2 = 0$$

is canonical, in particular klt.

Proof. The same argument from Lemma 6.1 shows that S is normal and that we may replace S by the hypersurface defined by the same equation but with general coefficients.

We show that the convex hull of the points $(2, 0, ..., 0), (0, 3, 0, ..., 0), ..., (0, ..., 0, s_{n-1}, 0)$ and (0, 1, ..., 1, 2) in \mathbb{R}^{n+1} contains a point with all coordinates less than 1. Indeed, the point

$$\frac{5}{12}(2,0,\ldots,0) + \frac{1}{6}(0,3,0,\ldots,0) + \frac{5}{12}(0,1,\ldots,1,2)$$

has all coordinates less than 1. So the Newton polytope of $x_0^2, x_1^3, \ldots, x_{n-1}^{s_{n-1}}$ and $x_1 \cdots x_n^2$ contains $(1, \ldots, 1)$ in the interior. Corollary 3.4 implies that *S* is klt, in particular canonical.

In weighted projective space $\mathbb{P} = \mathbb{P}(a_0, \ldots, a_n)$, the multiplicity of an irreducible closed substack (or an effective algebraic cycle) at a point is defined to be the multiplicity at a corresponding point of its inverse image in any orbifold chart $\mathbb{A}^n \to [\mathbb{A}^n/\mu_{a_i}] \cong \{x_i \neq 0\} \subset \mathbb{P}$. It is independent of the choice of *i* since different orbifold charts are étale-locally isomorphic. We can use the following criterion to check that a pair is canonical using multiplicity.

Theorem 6.3. [20, Claim 2.10.4] Let X be a regular scheme and Δ an effective \mathbb{Q} -divisor. Then (X, Δ) is canonical if $\operatorname{mult}_p \Delta \leq 1$ for every point p in X.

Johnson and Kollár gave the following bound for the multiplicity of an irreducible closed substack of weighted projective space.

Lemma 6.4. [18, Proposition 11] Let $a_0 \ge ... \ge a_n$ be positive integers and T be an irreducible closed substack of $\mathbb{P}^n(a_0,...,a_n)$ of dimension m < n. Then the multiplicity of T at every point is at most $(a_0 \cdots a_m) \deg(T)$.

Now we prove that the hypersurfaces from Theorem 4.1 and Theorem 4.2 are exceptional.

Theorem 6.5. For every even integer $n \ge 2$, the *n*-dimensional klt Fano variety X defined in Theorem 4.1 is exceptional. The Fano *n*-fold X has α -invariant

$$1 < \frac{a_n a_{n+1}}{d} \le \alpha(X) \le \frac{(s_n - 2)a_{n+1}}{s_n - 1},$$

22 LOUIS ESSER, JIHAO LIU, AND CHENGXI WANG where $a_n a_{n+1}/d \sim s_n/4$ and $(s_n - 2)a_{n+1}/(s_n - 1) \sim s_n^3/8$.

Proof. To show that X is exceptional, it is equivalent to show that the α -invariant of X is greater than 1. We now aim to show that there exists $\nu_n > 1$ such that the pair $(X, \nu_n D)$ is lc for every effective \mathbb{Q} -divisor $D \sim_{\mathbb{Q}} -K_X$. More precisely, we will show that if $\nu_n \leq a_n a_{n+1}/d$, then $(X, \nu_n D)$ is lc for every effective \mathbb{Q} -divisor $D \sim_{\mathbb{Q}} -K_X$.

Since $D \sim_{\mathbb{Q}} -K_X$ and $-K_X = \mathcal{O}(1)$, we have $\deg(D) = d/(a_0 \cdots a_{n+1})$. Note that $a_0 \geq \cdots \geq a_{n-1} \geq a_{n+1} \geq a_n$. Then by Lemma 6.4, the multiplicity of D at every point is at most $a_0 \cdots a_{n-1} \deg(D) = d/(a_n a_{n+1})$, where $a_0 \cdots a_{n-1}$ is the product of the n largest weights. If $\nu_n \leq a_n a_{n+1}/d$, then $\nu_n D$ has multiplicity at every point at most $\nu_n d/(a_n a_{n+1})$ which is at most 1. By Theorem 6.3, when

(4)
$$\nu_n \le \frac{a_n a_{n+1}}{d},$$

the pair $(X, \nu_n D)$ is lc at all smooth points of the stack X, hence at all points other than $[x_0 : \cdots : x_n : x_{n+1}] = [0 : \cdots : 0 : 1]$. Note that $a_n a_{n+1}/d \sim s_n/4$ since $a_n \sim s_n^2/4$, $a_{n+1} \sim s_n a_n/2 \sim s_n^3/8$ and $d \sim a_{n+1} s_n \sim s_n^4/8$. In particular, $a_n a_{n+1}/d > 1$ for all even $n \ge 2$.

In the affine chart $x_{n+1} = 1$, X is the hypersurface of \mathbb{A}^{n+1} given by $x_0^2 + x_1^3 + \cdots + x_{n-1}^{s_{n-1}} + x_n^b + x_1 \cdots x_n = 0$. We will show that $(X, \nu_n D)$ is lc near the origin $(0, \ldots, 0)$ for any $\nu_n \leq a_n a_{n+1}/d$. By [28, Lemma 4.4], we get

$$\operatorname{mult}_a D \le a_{n+1} \operatorname{deg}(D) = \frac{d}{a_0 \cdots a_n}$$

where mult_{*a*}*D* is the *a*-weighted multiplicity of *D* at the origin in \mathbb{A}^{n+1} with respect to the weights a_0, \ldots, a_n for x_0, \ldots, x_n , respectively.

Let $r \coloneqq -1 + a_n + a_{n+1}$. Let $\operatorname{mult}_u D$ be the *u*-weighted multiplicity of D at the origin with respect to the weights $u_i \coloneqq a_i$ for $i = 0, \ldots, n-1$, and $u_n \coloneqq (s_n + 1)r/2$. Since $u_n > a_n$, we have $\operatorname{mult}_u D \le \operatorname{mult}_a(D) \le d/(a_0 \cdots a_n)$. Let $b_i \coloneqq u_i/r$ for $i = 0, \ldots, n$. Then $\operatorname{mult}_b D = r^{n-1} \operatorname{mult}_u D \le r^{n-1} d/(a_0 \cdots a_n)$. Since $d = (s_n - 1)r$ and $a_i = b_i r$ for $i = 0, \ldots, n-1$, we get $\operatorname{mult}_b D \le (s_n - 1)/(b_0 \cdots b_{n-1}a_n)$.

Note that $b_i = (s_n - 1)/s_i$ for i = 0, ..., n - 1, and $b_n = (s_n + 1)/2$. Let X^c be the hypersurface in $\mathbb{P}(b_0, ..., b_n)$ of degree $s_n - 1$ defined by $x_0^2 + x_1^3 + \cdots + x_{n-1}^{s_{n-1}} + x_1 \cdots x_n = 0$, which is the weighted tangent cone to the hypersurface $X \subset \mathbb{A}^{n+1}$ at the origin with respect to the weights $b_0, ..., b_n$. The monomial x_n^b has disappeared since its degree in the weights $b_0, ..., b_n$ is larger than $s_n - 1$. Because the klt property is preserved by finite quotients, Lemma 6.1 means X^c is klt. Denote the weighted tangent cone of $D \subset \mathbb{A}^{n+1}$ at the origin by D^c . Then D^c is an effective \mathbb{Q} -divisor in X^c and is \mathbb{Q} -linearly equivalent to a rational multiple of $\mathcal{O}_{X^c}(1)$ since D is \mathbb{Q} -Cartier. By [28, Lemma 4.1], if $(X^c, \nu_n D^c)$ is lc and $K_{X^c} + \nu_n D^c \sim_{\mathbb{Q}} \mathcal{O}_{X^c}(l)$ with $l \leq 0$, then $(X, \nu_n D)$ is lc near the origin.

We have $-K_{X^c} = \mathcal{O}_{X^c}(-(s_n-1)+\sum_{j=0}^n b_j) = \mathcal{O}_{X^c}((s_n-1)/2)$ by the adjunction formula. Note that $l \leq 0$ is equivalent to $\nu_n \deg(D^c) \leq \deg(-K_{X^c}) = (-K_{X^c}) \cdot c_1(\mathcal{O}(1))^{n-2} = (s_n - C_{X^c}) \cdot c_1(\mathcal{O}(1))^{n-2} = (s_n - C$ $1)/2 \cdot (s_n - 1)/(b_0 \cdots b_n)$. That is to say, $l \leq 0$ iff

$$\nu_n \le \frac{(s_n - 1)^2}{2b_0 \cdots b_n \deg(D^c)}$$

The right side of this inequality is at least $(s_n - 1)a_n/(2b_n)$ since $\deg(D^c) = \operatorname{mult}_b D \leq (s_n - 1)/(b_0 \cdots b_{n-1}a_n)$. Thus we have $l \leq 0$ when

(5)
$$\nu_n \le \frac{(s_n - 1)a_n}{2b_n} = \frac{(s_n - 1)a_n}{s_n + 1},$$

where $(s_n - 1)a_n/(2b_n) \sim s_n^2/4$ since $a_n \sim s_n^2/4$ and $b_n \sim s_n/2$.

Now we want to find the inequality satisfied by ν_n so that $(X^c, \nu_n D^c)$ is lc. Note that $b_n > b_0 > \cdots > b_{n-1}$. By Lemma 6.4, since the dimension of D^c is n-2, the multiplicity of D^c at every point is at most $b_2 \text{deg}(D^c)$ when n = 2, or $b_n b_0 \cdots b_{n-3} \text{deg}(D^c)$ when $n \ge 4$, where $b_n b_0 \cdots b_{n-3}$ is the product of the n-1 largest weights. Since $\text{deg}(D^c) \le (s_n - 1)/(b_0 \cdots b_{n-1}a_n)$, the multiplicity of D^c at every point is at most $b_2(s_2 - 1)/(b_0 b_1 a_2) = 4/13$ when n = 2, or $b_n(s_n - 1)/(b_{n-2}b_{n-1}a_n)$ when $n \ge 4$. Therefore, for each even integer $n \ge 2$, the multiplicity of $\nu_n D^c$ at every point is at most 1 if

(6)
$$\nu_n \le \frac{b_{n-2}b_{n-1}a_n}{b_n(s_n-1)} = \frac{2a_n(s_n-1)}{s_{n-2}s_{n-1}(s_n+1)}.$$

We have $2a_n(s_n-1)/(s_{n-2}s_{n-1}(s_n+1)) \sim s_{n-2}^5/2$ since $s_n-1 \sim s_n \sim s_{n-1}^2 \sim s_{n-2}^4$ and $a_n \sim s_n^2/4 \sim s_{n-2}^8/4$. Since the stack X^c is a smooth outside the point $[x_0 : \cdots : x_{n-1} : x_n] = [0 : \cdots : 0 : 1]$, by Theorem 6.3, $(X^c, \nu_n D^c)$ is lc outside this point if ν satisfies inequality (6).

Now we want to find the inequality satisfied by ν_n such that $(X^c, \nu_n D^c)$ is lc near the point $[x_0 : \cdots : x_{n-1} : x_n] = [0 : \cdots : 0 : 1]$ in $\mathbb{P}(b_0, \ldots, b_n)$. In the affine coordinate chart $x_n = 1$, the hypersurface X^c is defined by $x_0^2 + x_1^3 + \cdots + x_{n-1}^{s_{n-1}} + x_1 \cdots x_{n-1} = 0$. Note that $b_i = c_i$ for $i = 0, \ldots, n-1$, for the values of c_i appearing in [28, Lemma 5.1]. Hence the pair $(X^c, \nu_n D^c)$ is lc near the point $[x_0 : \cdots : x_{n-1} : x_n] = [0 : \cdots : 0 : 1]$ if

$$\nu_n \leq \begin{cases} \frac{1}{\mathrm{mult}_b D^c} \text{ for } n = 2, \\ \\ \frac{2}{s_{n-1}^{n-2} (s_{n-1}+1)^2 (s_{n-1}-1)^{n-4} \mathrm{mult}_b D^c} \text{ for } n \geq 4. \end{cases}$$

We know D^c corresponds to a cycle with codimension 2 in this affine coordinate chart \mathbb{A}^n and by [28, Lemma 4.4]

$$\operatorname{mult}_b D^c \le b_n \operatorname{deg}(D^c) = \frac{b_n(s_n - 1)}{b_0 \cdots b_{n-1}a_n} = \frac{s_n + 1}{2(s_n - 1)^{n-2}a_n}.$$

Thus $(X^c, \nu_n D^c)$ is lc near the point $[x_0 : \cdots : x_{n-1} : x_n] = [0 : \cdots : 0 : 1]$ if

(7)
$$\nu_n \leq \begin{cases} \frac{2a_2}{s_2+1} = \frac{13}{4} \text{ for } n = 2, \\ \frac{4a_n(s_n-1)^{n-2}}{s_{n-1}^{n-2}(s_{n-1}+1)^2(s_{n-1}-1)^{n-4}(s_n+1)} \sim s_{n-1}^2 \text{ for } n \geq 4 \end{cases}$$

since $4a_n(s_n-1)^{n-2} \sim 4 \cdot s_{n-1}^4(s_n-1)^{n-2}/4 = s_{n-1}^{2n}$ and $s_{n-1}^{n-2}(s_{n-1}+1)^2(s_{n-1}-1)^{n-4}(s_n+1) \sim s_{n-1}^{2n-4}s_{n-1}^2 = s_{n-1}^{2n-2}$. Among all the right sides of inequalities (4), (5), (6) and (7), the one in inequality (4) is the smallest one and greater than 1. We know α -invariant of X is the supremum of the real numbers ν_n such that $(X, \nu_n D)$ is lc for every effective \mathbb{Q} -divisor $D \sim_{\mathbb{Q}} -K_X$. Hence the α -invariant $\alpha(X) \geq a_n a_{n+1}/d > 1$.

Now we show $\alpha(X) \leq (s_n - 2)a_{n+1}/(s_n - 1)$. Indeed, there exists a Q-divisor $D \in |-K_X|_Q$ such that $\operatorname{lct}(X, D) = (s_n - 2)a_{n+1}/(s_n - 1)$. Let $D \coloneqq (1/a_{n+1})E$, where E is the hyperplane section in X defined by the equations $\{x_{n+1} = 0, x_0^2 + \cdots + x_{n-1}^{s_{n-1}} = 0\}$ in $\mathbb{P}(a_0, \ldots, a_n)$. The stack E has one singular point $p_n \coloneqq [x_0 : \cdots : x_{n-1} : x_n : x_{n+1}] = [0 : \cdots : 0 : 1 : 0]$, which is a smooth point of stack X. In the affine chart $x_n = 1$, the stack E is étale-locally isomorphic to the Fermat-type hypersurface singularity $x_0^2 + \cdots + x_{n-1}^{s_{n-1}} = 0$ in \mathbb{A}^n . By [19, Example 8.15], the log canonical threshold of the pair (X, E) at the point p_n is $\min\{1/2 + \cdots + 1/s_{n-1}, 1\} = (s_n - 2)/(s_n - 1)$. Hence the log canonical threshold of (X, D) at p_n is $(s_n - 2)a_{n+1}/(s_n - 1)$. Since p_n is the only singular point in E, we get $\operatorname{lct}(X, D) = (s_n - 2)a_{n+1}/(s_n - 1)$. So $\alpha(X) \leq \operatorname{lct}(X, D) = (s_n - 2)a_{n+1}/(s_n - 1)$. We have $(s_n - 2)a_{n+1}/(s_n - 1) \sim s_n^3/8$ since $a_{n+1} \sim s_n a_n/2 \sim s_n^3/8$.

Theorem 6.6. For every odd integer $n \ge 3$, the *n*-dimensional klt Fano variety X defined in Theorem 4.2 is exceptional. The Fano *n*-fold X has α -invariant

$$1 < \frac{a_n a_{n+1}}{d} \le \alpha(X) \le \frac{(s_n - 2)a_{n+1}}{s_n - 1}$$

where $a_n a_{n+1}/d \sim s_n/4$ and $(s_n - 2)a_{n+1}/(s_n - 1) \sim s_n^3/16$.

Proof. To show X is exceptional, it is equivalent to show that the α -invariant of X is greater than 1. We now aim to show that there exists $\nu_n > 1$ such that the pair $(X, \nu_n D)$ is lc for every effective \mathbb{Q} -divisor $D \sim_{\mathbb{Q}} -K_X$. In particular, we can show that if $\nu_n \leq a_n a_{n+1}/d$, then $(X, \nu_n D)$ is lc for every effective \mathbb{Q} -divisor $D \sim_{\mathbb{Q}} -K_X$.

Since $D \sim_{\mathbb{Q}} -K_X$ and $-K_X = \mathcal{O}(1)$, we have $\deg(D) = d/(a_0 \cdots a_{n+1})$. Note that $a_0 \geq \cdots \geq a_{n-1} \geq a_{n+1} \geq a_n$. Then by Lemma 6.4, the multiplicity of D at every point is at most $a_0 \cdots a_{n-1} \deg(D) = d/(a_n a_{n+1})$, where $a_0 \cdots a_{n-1}$ is the product of the n largest weights. If $\nu_n \leq a_n a_{n+1}/d$, then $\nu_n D$ has multiplicity at every point at most $\nu_n d/(a_n a_{n+1})$ which is at most 1. By Theorem 6.3, when

(8)
$$\nu_n \le \frac{a_n a_{n+1}}{d},$$

the pair $(X, \nu_n D)$ is lc at all smooth points of the stack X, hence at all points other than $[x_0 : \cdots : x_n : x_{n+1}] = [0 : \cdots : 0 : 1]$. Note that $(a_n a_{n+1})/d \sim s_n/4$ since $a_n \sim s_n^2/4$, $a_{n+1} \sim (s_n a_n)/4 \sim s_n^3/16$ and $d \sim a_{n+1}s_n \sim s_n^4/16$. In particular, $a_n a_{n+1}/d > 1$ for all odd integers $n \ge 3$.

In the affine chart $x_{n+1} = 1$, the hypersurface $X \subset \mathbb{A}^{n+1}$ is given by $x_0^2 + x_1^3 + \cdots + x_{n-1}^{s_{n-1}} + x_n^b + x_1 \cdots x_n^2 = 0$. We will show that $(X, \nu_n D)$ is lc near the origin $(0, \ldots, 0)$ for any $\nu_n \leq a_n a_{n+1}/d$. By [28, Lemma 4.4], we get $\operatorname{mult}_a D \leq a_{n+1} \operatorname{deg}(D) = d/(a_0 \cdots a_n)$, where $\operatorname{mult}_a D$ is the *a*-weighted multiplicity of D at the origin in \mathbb{A}^{n+1} with respect to the weights a_0, \ldots, a_n for x_0, \ldots, x_n respectively.

Let $r \coloneqq -1 + a_n + a_{n+1}$. Let $\operatorname{mult}_u D$ be the *u*-weighted multiplicity of D at the origin with respect to the weights $u_i \coloneqq a_i$ for $i = 0, \ldots, n-1$, and $u_n \coloneqq (s_n + 1)r/4$. Since $u_n > a_n$, we have $\operatorname{mult}_u D \le \operatorname{mult}_a(D) \le d/(a_0 \cdots a_n)$. Let $b_i \coloneqq u_i/r$ for $i = 0, \ldots, n$. Then $\operatorname{mult}_b D = r^{n-1} \operatorname{mult}_u D \le r^{n-1} d/(a_0 \cdots a_n)$. Since $d = (s_n - 1)r$ and $a_i = b_i r$ for $i = 0, \ldots, n-1$, we get $\operatorname{mult}_b D \le (s_n - 1)/(b_0 \cdots b_{n-1}a_n)$.

Note that $b_i = (s_n - 1)/s_i$ for i = 0, ..., n - 1, and $b_n = (s_n + 1)/4$. Let X^c be the hypersurface in $\mathbb{P}(b_0, ..., b_n)$ of degree $s_n - 1$ defined by $x_0^2 + x_1^3 + \cdots + x_{n-1}^{s_{n-1}} + x_1 \cdots x_{n-1} x_n^2 = 0$, which is the weighted tangent cone to the hypersurface $X \subset \mathbb{A}^{n+1}$ at the origin with respect to the weights $b_0, ..., b_n$. The monomial x_n^b has disappeared since its degree in the weights $b_0, ..., b_n$ is larger than $s_n - 1$. Because the klt property is preserved by finite quotients, Lemma 6.2 means X^c is klt. Denote the weighted tangent cone of $D \in \mathbb{A}^{n+1}$ at the origin by D^c . Then D^c is an effective \mathbb{Q} -divisor in X^c and is \mathbb{Q} -linearly equivalent to a rational multiple of $\mathcal{O}_{X^c}(1)$ since D is \mathbb{Q} -Cartier. By [28, Lemma 4.1], if $(X^c, \nu_n D^c)$ is lc and $K_{X^c} + \nu_n D^c \sim_{\mathbb{Q}} \mathcal{O}_{X^c}(l)$ with $l \leq 0$, then $(X, \nu_n D)$ is lc near the origin.

We have $-K_{X^c} = \mathcal{O}_{X^c}(-(s_n-1) + \sum_{j=0}^n b_j) = \mathcal{O}_{X^c}((s_n-3)/4)$ by adjunction formula. Note that $l \leq 0$ is equivalent to $\nu_n \deg(D^c) \leq \deg(-K_{X^c}) = (-K_{X^c}) \cdot c_1(\mathcal{O}(1))^{n-2} = (s_n - 3)/4 \cdot (s_n - 1)/(b_0 \cdots b_n)$. That is to say, $l \leq 0$ iff

$$\nu_n \le \frac{(s_n - 3)(s_n - 1)}{4b_0 \cdots b_n \deg(D^c)}.$$

The left side of this inequality is at least $(s_n - 3)a_n/(4b_n)$ since $\deg(D^c) = \operatorname{mult}_b D \le (s_n - 1)/(b_0 \cdots b_{n-1}a_n)$. Thus we have $l \le 0$ when

(9)
$$\nu_n \le \frac{(s_n - 3)a_n}{4b_n} = \frac{(s_n - 3)a_n}{s_n + 1}$$

where $(s_n - 3)a_n/(s_n + 1) \sim s_n^2/4$ since $a_n \sim s_n^2/4$.

Now we want to find the inequality satisfied by ν_n so that $(X^c, \nu_n D^c)$ is lc. Note that $b_0 > b_1 > b_n > b_2 > \cdots > b_{n-1}$. By Lemma 6.4, since dimension of D^c is n-2, the multiplicity of D^c at every point is at most $b_0b_1\text{deg}(D^c)$ when n = 3, or $b_nb_0b_1\cdots b_{n-3}\text{deg}(D^c)$ when $n \ge 5$, where $b_nb_0\cdots b_{n-3}$ is the product of the n-1 largest weights.. Since $\text{deg}(D^c) \le (s_n - 1)/(b_0\cdots b_{n-1}a_n)$, the multiplicity of D^c at every point is at most $(s_3 - 1)/(b_2a_3) = 7/493$ when n = 3, or $(b_n(s_n - 1))/(b_{n-2}b_{n-1}a_n)$ when $n \ge 5$. Therefore, the multiplicity of $\nu_n D^c$ at every point is at most 1 if

(10)
$$\nu_n \leq \begin{cases} \frac{493}{7} \text{ for } n = 3, \\ \frac{b_{n-2}b_{n-1}a_n}{b_n(s_n - 1)} = \frac{4a_n(s_n - 1)}{s_{n-2}s_{n-1}(s_n + 1)} \text{ for } n \geq 5 \end{cases}$$

We have $4a_n(s_n - 1)/(s_{n-2}s_{n-1}(s_n + 1)) \sim s_n^2/(s_{n-2}s_{n-1}) \sim s_{n-2}^5$ since $s_n - 1 \sim s_n \sim s_{n-1}^2 \sim s_{n-2}^4$ and $a_n \sim s_n^2/4 \sim s_{n-2}^8/4$. Since the stack X^c is a smooth outside the point $[x_0 : \cdots : x_{n-1} : x_n] = [0 : \cdots : 0 : 1]$, then by Theorem 6.3, $((X^c, \nu_n D^c))$ is lc outside this point if ν satisfies inequality (6).

Now we want to find the inequality satisfied by ν_n such that $(X^c, \nu_n D^c)$ is lc near the point $[x_0 : \cdots : x_n] = [0 : \cdots : 0 : 1]$ in $\mathbb{P}(b_0, \ldots, b_n)$. In the affine coordinate chart $x_n = 1$,

the hypersurface X^c is defined by $x_0^2 + x_1^3 + \cdots + x_{n-1}^{s_{n-1}} + x_1 \cdots x_{n-1} = 0$. Note that $b_i = c_i$ for $i = 0, \ldots, n-1$, for the values of c_i appearing in [28, Lemma 5.1]. Hence the pair $(X^c, \nu_n D^c)$ is lc near the point $[x_0 : \cdots : x_{n-1} : x_n] = [0 : \cdots : 0 : 1]$ if

$$\nu_n \le \frac{2}{s_{n-1}^{n-2}(s_{n-1}+1)^2(s_{n-1}-1)^{n-4} \operatorname{mult}_b D^c} \text{ for } n \ge 3.$$

We know D^c corresponds to a cycle with codimension 2 in this affine coordinate chart \mathbb{A}^n and by [28, Lemma 4.4]

$$\operatorname{mult}_b D^c \le b_n \operatorname{deg}(D^c) = \frac{b_n(s_n-1)}{b_0 \cdots b_{n-1}a_n} = \frac{s_n+1}{4(s_n-1)^{n-2}a_n}$$

Thus $(X^c, \nu_n D^c)$ is lc near the point $[x_0 : \cdots : x_n] = [0 : \cdots : 0 : 1]$ if

(11)
$$\nu_n \le \frac{8a_n(s_n-1)^{n-2}}{s_{n-1}^{n-2}(s_{n-1}+1)^2(s_{n-1}-1)^{n-4}(s_n+1)} \sim 2s_{n-1}^2 \text{ for } n \ge 3$$

since $8a_n(s_n-1)^{n-2} \sim 8s_{n-1}^4(s_n-1)^{n-2}/4 = 2s_{n-1}^{2n}$ and $s_{n-1}^{n-2}(s_{n-1}+1)^2(s_{n-1}-1)^{n-4}(s_n+1) \sim s_{n-1}^{2n-4}s_{n-1}^2 = s_{n-1}^{2n-2}$. Among all the right sides of inequalities (8), (9), (10) and (11), the one in inequality (8) is the smallest one and greater than 1. We know α -invariant of X is the supremum of the real numbers ν_n such that $(X, \nu_n D)$ is lc for every effective \mathbb{Q} -divisor $D \sim_{\mathbb{Q}} -K_X$. Hence the α -invariant $\alpha(X) \geq (a_n a_{n+1})/d > 1$.

Now we show $\alpha(X) \leq (s_n - 2)a_{n+1}/(s_n - 1)$. Indeed, there exists a \mathbb{Q} -divisor $D \in |-K_X|_{\mathbb{Q}}$ such that $\operatorname{lct}(X, D) = (s_n - 2)a_{n+1}/(s_n - 1)$. Let $D \coloneqq (1/a_{n+1})E$, where E is the hyperplane section in X given by $\{x_{n+1} = 0, x_0^2 + \cdots + x_{n-1}^{s_{n-1}} = 0\}$. The stack E has one singular point $p_n \coloneqq [x_0 : \cdots : x_{n-1} : x_n : x_{n+1}] = [0 : \cdots : 0 : 1 : 0]$ which is a smooth point of stack X. In the affine chart $x_n = 1$, the stack E is étale-locally isomorphic to the Fermat-type hypersurface singularity $x_0^2 + \cdots + x_{n-1}^{s_{n-1}} = 0$ in \mathbb{A}^n . By [19, Example 8.15], the log canonical threshold of the pair (X, E) at the point p_n is $\min\{1/2 + \cdots + 1/s_{n-1}, 1\} = (s_n - 2)/(s_n - 1)$. Hence the log canonical threshold of (X, D) at p_n is $(s_n - 2)a_{n+1}/(s_n - 1)$. Since p_n is the only singular point in E, we get $\operatorname{lct}(X, D) = (s_n - 2)a_{n+1}/(s_n - 1)$. So $\alpha(X) \leq \operatorname{lct}(X, D) = (s_n - 2)a_{n+1}/(s_n - 1)$. We have $(s_n - 2)a_{n+1}/(s_n - 1) \sim s_n^3/16$ since $a_{n+1} \sim s_n a_n/4 \sim s_n^3/16$.

Remark 6.7. When n = 2, our example is the hypersurface $X_{282} \subset \mathbb{P}(141, 94, 13, 35)$ defined by the equation

$$x_0^2 + x_1^3 + x_2^{19}x_3 + x_1x_2x_3^5 = 0.$$

Let *C* be the curve in X_{282} defined by $\{x_2 = 0, x_0^2 + x_1^3 = 0\}$. By taking a resolution, we can compute lct(X, C) = 3/4. Hence for $D = (1/13)C \in |-K_X|_{\mathbb{Q}}$, we have lct(X, D) = 39/4, which gives a sharper upper bound for $\alpha(X)$ than in Theorem 6.5. This upper bound is smaller than the α -invariant of 21/2 achieved by another exceptional del Pezzo surface constructed by Totaro [28, Section 8], which he conjectures to be optimal. That example is another non-quasismooth hypersurface $X_{154} \subset \mathbb{P}(77, 45, 19, 14)$.

Unlike in the small anti-canonical volume examples from [28], the divisor computing the α -invariant in our examples likely passes through the non-quasismooth point, and the exact value of the invariant seems to be very difficult to compute in higher dimensions.

Since our examples do not appear to have optimal α -invariant (this is true for n = 2, for instance, from the last paragraph), we only demonstrate bounds on $\alpha(X)$ in this section.

REFERENCES

- D. Abramovich, M. Temkin, and J. Włodarczyk. Functorial embedded resolution via weighted blowings up. Algebra Number Theory, to appear. arXiv:1906.07106 20
- [2] V. Alexeev and W. Liu. Open surfaces of small volume. Algebraic Geom. 6 (2019), no. 3, 312–327. 2
- [3] F. Ambro. The set of toric minimal log discrepancies. Cent. Eur. J. Math. 4 (2006), no. 3, 358–370. 6
- [4] F. Ambro. The minimal log discrepancy. In: Proceedings of the workshop "Multiplier ideals and arc spaces" (RIMS 2006), K. Watanabe (Ed.). RIMS Koukyuuroku 1550 (2006), 121–130. 6
- [5] F. Aroca, M. Gomez-Morales, and H. Mourtada. Groebner fan and embedded resolutions of ideals on toric varieties. *Beiträge zur Algebra und Geometrie/Contributions to Algebra and Geometry* 65 (2024), 217–228. 7, 8
- [6] C. Birkar. Anti-pluricanonical linear systems on Fano varieties. Ann. of Math. 190 (2019), 345-463.2
- [7] C. Birkar. Singularities of linear systems and boundedness of Fano varieties. Ann. of Math. 193 (2021), no. 2, 347–405. 2, 6
- [8] C. Birkar and J. Liu. On explicit bounds of Fano threefolds. arXiv: 2311.06732. 11
- [9] L Esser. Minimal log discrepancies of hypersurface mirrors. *Forum. Math. Sigma.* **12** (2024), e23, 1–17. 2
- [10] L. Esser and B. Totaro. Log canonical pairs with conjecturally minimal volume. arXiv: 2308.08034.2
- [11] L. Esser, B. Totaro, and C. Wang. Varieties of general type with doubly exponential asymptotics. *Trans. Amer. Math. Soc. Ser. B* **10** (2023), 288–309. **2**, 19
- [12] L. Esser, B. Totaro, and C. Wang. Calabi-Yau varieties of large index. arXiv: 2209.04597.2
- [13] W. Fulton. Intersection theory. 2nd ed. Springer, Berlin (1998). 20
- [14] C. Hacon, J. M^cKernan, and C. Xu. ACC for log canonical thresholds. Ann. of Math. 180 (2014), no. 2, 523–571.
- [15] A. R. Iano-Fletcher. Working with weighted complete intersections. *Explicit birational geometry of 3-folds*, 101–173. London Math. Soc. Lecture Notes Ser. 281, Cambridge Univ. Press, Cambridge (2000).
 4
- [16] S. Ishii. Introduction to singularities. Springer, Tokyo, 2014. 7, 8
- [17] S. Ishii and Y. Prokhorov. Hypersurface exceptional singularities. *Internat. J. Math.* 12 (2001), 661–687.
 7
- [18] J. Johnson and J. Kollár. Kähler-Einstein metrics on log del Pezzo surfaces in weighted projective 3-spaces. Ann. Inst. Fourier (Grenoble) 51 (2001), 69–79. 21
- [19] J. Kollár. Singularities of pairs. *Algebraic geometry* (Santa Cruz, 1995), 221–287. Proc. Symp. Pure Math. 62, Amer. Math. Soc. (1997). 24, 26
- [20] J. Kollár. *Singularities of the minimal model program.* With the collaboration of Sándor Kovács. Cambridge, 2013. 5, 21
- [21] J. Kollár and S. Mori. *Birational geometry of algebraic varieties*. Cambridge Tracts in Mathematics, vol. 134, Cambridge University Press, Cambridge, 1998, With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original. 9
- [22] J. Liu and V. V. Shokurov. Optimal bounds on surfaces. arXiv: 2305.19248 2, 3, 4, 6, 10
- [23] J. Liu and W. Liu. The minimal volume of surfaces of log general type with non-empty non-klt locus. arXiv:2308.14268.2
- [24] Y. Liu. The volume of singular Kähler-Einstein Fano varieties. Compos. Math. 154 (2018), no. 6, 1131– 1158. 2
- [25] Y. Odaka and Y. Sano. Alpha invariant and K-stability of Q-Fano varieties. Adv. Math. 229 (2012), no. 5, 2018–2834. 2
- [26] G. Tian. On Kähler–Einstein metrics on certain Kähler manifolds with $c_1(M) > 0$. *Invent. Math.* 89 (1987), 225–246. 2

LOUIS ESSER, JIHAO LIU, AND CHENGXI WANG

- [27] The Stacks project authors. *The Stacks project*. https://stacks.math.columbia.edu, 2024. 4
- [28] B. Totaro. Klt varieties with conjecturally minimal volume. *Int. Math. Res. Not. IMRN* 2024, no. 1, 462–491. 1, 2, 3, 6, 10, 11, 14, 22, 23, 24, 25, 26
- [29] B. Totaro and C. Wang. Klt varieties of general type with small volume. *Annali della Scuola Normale Superiore di Pisa* **24** (2023), 1557–1573. **2**
- [30] A. Varchenko. Zeta-function of monodromy and Newton's diagram. *Invent. Math.* **37** (1976), no. 3, 253–262. 7
- [31] C. Wang. Fano varieties with conjecturally largest Fano index. arXiv: 2308.06563.2

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, FINE HALL, WASHINGTON ROAD, PRINCE-TON, NJ 08544-1000, USA

Email address: esserl@math.princeton.edu

DEPARTMENT OF MATHEMATICS, NORTHWESTERN UNIVERSITY, 2033 SHERIDAN RD, EVANSTON, IL 60208, USA

Email address: jliu@northwestern.edu

UCLA MATHEMATICS DEPARTMENT, BOX 951555, LOS ANGELES, CA 90095-1555, USA

Email address: chwang@math.ucla.edu