Field Theory of Active Brownian Particles with Dry Friction

Ziluo Zhang,^{1,2,*} Shurui Yuan,^{3,1} and Shigeyuki Komura^{1,4,5,†}

¹Wenzhou Institute, University of Chinese Academy of Sciences, Wenzhou, Zhejiang 325001, China

²Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190, China

³Harbin Institute of Technology, Harbin, 150001, China

⁴Oujiang Laboratory, Wenzhou, Zhejiang 325000, China

⁵Department of Chemistry, Graduate School of Science,

Tokyo Metropolitan University, Tokyo 192-0397, Japan

We present a field theoretic approach to capture the motion of a particle with dry friction for one- and two-dimensional diffusive particles, and further expand the framework for two-dimensional active Brownian particles. Starting with the Fokker-Planck equation and introducing the Hermite polynomials as the corresponding eigen-functions, we obtain the actions and propagators. Using a perturbation expansion, we calculate the effective diffusion coefficient in the presence of both wet and dry frictions in a perturbative way via the Green-Kubo relation. We further compare the analytical result with the numerical simulation. Our result can be used to estimate the values of dry friction coefficient in experiments.

I. INTRODUCTION

Active matter [1–4] refers to particles or creatures that can absorb energy from the environment and change their mechanical state by using the energy. The collective behaviors of animals such as swarming [5, 6], flocks [7– 11], and bacteria [12] can be described effectively by appropriate models. Furthermore, artificial active colloidal particles [13, 14] are created experimentally to explore active systems. There are plenty of analytical works from the Stokes–Einstein–Sutherland relation of Brownian motion [15] to active systems [16, 17]. However, fitting analytical results with experimental ones is always a challenge. For example, the restricted confinement can significantly affect the diffusion coefficient [18]. Furthermore, the motion of particles can be influenced by dry friction (Coulomb friction) with the surface if the mass of the particle is effectively large. Unlike the wet friction that is always proportional to the velocity, dry friction depends only on the velocity direction.

In the most of the previous studies, dry and wet frictions are not considered together. The particles experiencing dry friction are mostly unaffected by the thermal noise, since the particles are large enough. Moreover, the gravity acting on the particles is usually smaller than thermal noise, and the particles hardly reach the lower surface. Hence, dry friction of such particles can also be neglected. For mesoscopic particles, however, both wet and dry frictions can not be neglected because the gravity drags the particles and their random motions take place at the bottom surface [19, 20]. A one-dimensional (1D) passive particle with dry friction has been discussed [21–23], and further two-dimensional (2D) problem has been also addressed [24].

In this work, we are motivated to explore the motion of an active particle with dry friction. In particular, we obtain the effective diffusion coefficient of a 2D active Brownian particle (ABP) with both dry and wet frictions by using the Doi-Peliti field theory [25–27]. A 2D ABP moves with a finite velocity, and its director undergoes rotational diffusion [28–30]. We first introduce the Langevin equations to describe the motion of an ABP, and later convert it to the corresponding Fokker-Planck equation. Since the effective diffusion coefficient can be calculated via the velocity-velocity correlation function by using the Green-Kubo relation [15], we only consider the particle velocity in this work.

In Sec. II, we introduce the mathematical basis of this work, such as the equation of motion and its eigen-system to simplify the calculation. In Sec. III, we use the field-theoretic framework to capture the motion of a 1D diffusing particle with dry friction in a perturbative way. We expand our framework to 2D space in Sec. IV by approximating dry friction to be isotropic. By treating the self-propulsion of an ABP as a perturbative part, we obtain the corresponding effective diffusion coefficient in Sec. V. We also compare the 2D result with the simulation result. A summary of our work and some discussion are given in Sec. VI.

^{*} zhangzl@wiucas.ac.cn

[†] komura@wiucas.ac.cn

II. MODEL

In this section, we derive the field theory of a particle with dry friction. Our main aim is to calculate the effective diffusion coefficient D_{eff} as a function of dry friction coefficient μ . The diffusion coefficient D_{eff} can be obtained by the Green-Kubo relation [15],

$$D_{\text{eff}} = \frac{1}{d} \int_0^\infty dt \, \left\langle \mathbf{v}(t) \cdot \mathbf{v}(0) \right\rangle \,, \tag{1}$$

where the $\langle \bullet \rangle$ is the ensemble average and d is the space dimension.

We describe the motion of an ABP with dry friction by the Langevin equations

$$m\dot{\mathbf{v}}(t) = -\Gamma\mathbf{v}(t) - F\frac{\mathbf{v}(t)}{||\mathbf{v}(t)||} + \Gamma\mathbf{w}_{\theta}(t) + \mathbf{\Xi}(t) \qquad \text{with} \quad \left\langle \mathbf{\Xi}(t)\mathbf{\Xi}^{\mathrm{T}}(t') \right\rangle = 2D\Gamma^{2}\mathbb{1}_{2}\delta(t-t') \quad \text{and} \quad \left\langle \mathbf{\Xi} \right\rangle = \mathbf{0} , \quad (2a)$$

$$\dot{\theta}(t) = \zeta(t) \qquad \text{with} \quad \langle \zeta(t)\zeta(t') \rangle = 2D_{\rm r}\delta(t-t') \quad \text{and} \quad \langle \zeta(t) \rangle = 0 , \qquad (2b)$$

$$\dot{\mathbf{x}} = \mathbf{v} , \qquad (2c)$$

where $||\mathbf{v}||$ is the norm of the vector \mathbf{v} , $\mathbf{w}_{\theta}(t) = w[\cos\theta, \sin\theta]^{\mathrm{T}}$ is the self-propelled velocity of the particle which stays on a ring of a radius of w, and m is the mass of the particle. Furthermore, $\Gamma \mathbf{v}(t)$ is the viscous force proportional to the velocity and $F\mathbf{v}(t)/||\mathbf{v}(t)||$ is dry friction force whose magnitude is a constant. And $\mathbb{1}_2$ is a 2D unit matrix. Each component of Ξ and also ζ are Gaussian white noise with zero mean and variance $D\Gamma^2$ and D_r , respectively, where D and D_r are the diffusion and rotational diffusion coefficients respectively. And each component of Ξ has zero correlation with ζ . Finally, \mathbf{x} is the position of the particle.

Introducing the rescaled friction coefficients by the mass $\gamma = \Gamma/m$ and $\mu = F/m$, we modify the Langevin equation as

$$\dot{\mathbf{v}}(t) = -\gamma \mathbf{v}(t) - \mu \frac{\mathbf{v}(t)}{||\mathbf{v}(t)||} + \gamma \mathbf{w}_{\theta}(t) + \boldsymbol{\xi}(t) \qquad \text{with} \quad \left\langle \boldsymbol{\xi}(t) \boldsymbol{\xi}^{\mathrm{T}}(t') \right\rangle = 2D\gamma^{2} \mathbb{1}_{2}\delta(t - t') \quad \text{and} \quad \left\langle \boldsymbol{\xi}(t) \right\rangle = \mathbf{0} , \quad (3a)$$

$$\dot{\theta}(t) = \zeta(t)$$
 with $\langle \zeta(t)\zeta(t')\rangle = 2D_{\rm r}\delta(t-t')$ and $\langle \zeta(t)\rangle = 0$, (3b)

whose corresponding Fokker-Planck equation is [31, 32]

$$\partial_t P(\mathbf{v},\theta,t) = \mathcal{L}P(\mathbf{v},\theta,t) \quad \text{with} \quad \mathcal{L} = D\gamma^2 \nabla_{\mathbf{v}}^2 + \gamma \nabla_{\mathbf{v}} \cdot \mathbf{v} + \mu \nabla_{\mathbf{v}} \cdot \frac{\mathbf{v}}{||\mathbf{v}||} - \mathbf{w}_\theta \cdot \nabla_{\mathbf{v}} + D_r \partial_\theta^2 . \tag{4}$$

Since the diffusion coefficient can be extracted from the velocity-velocity correlation function, we drop the positional variable \mathbf{x} in the Fokker-Planck equation, and we only consider the particle in the velocity space. The effective diffusion coefficient of an isolated 2D ABP in free space is well known [33, 34], in our case, it presents the zero dry friction limit $\mu = 0$. We write it below for the later comparison

$$D_0 = D + \frac{w^2}{2D_{\rm r}} \ . \tag{5}$$

The following functions are used later to simplify the calculation,

$$u_n(v) = e^{-\frac{v^2}{2\Omega^2}} He_n\left(\frac{v}{\Omega}\right) , \qquad (6a)$$

$$\tilde{u}_n(v) = \frac{1}{\sqrt{2\pi}n!} He_n\left(\frac{v}{\Omega}\right) , \qquad (6b)$$

where $He_n(x)$ is the *n*-th order of the probabilist's Hermite polynomials and $\Omega^2 = D\gamma$ [35, 36]. Our definition of $He_n(x)$ corresponds to $2^{-\frac{n}{2}}$ HermiteH $[n, x/\sqrt{2}]$ [37].

The orthogonality relation between $u_n(v)$ and $\tilde{u}(v)$ is

$$\int \mathrm{d}v \, u_n(v) \tilde{u}_m(v) = \Omega \delta_{n,m} \,\,, \tag{7}$$

where $\delta_{n,m}$ is the Kronecker delta, and $u_n(v)$ are the eigenfunctions of the operator

$$\Omega \partial_v^2 u_n(v) + \partial_v \left[v u_n(v) \right] = -n u_n(v) .$$
(8)

The full propagator is the probability density of finding a particle at velocity \mathbf{v} with the self-propulsion direction θ at time t, when the initial state is given at velocity \mathbf{v}_0 with the propulsion director θ_0 at time t_0 ,

$$\mathcal{G}(\mathbf{v},\theta,t|\mathbf{v}_0,\theta_0,t_0) = \left\langle \phi(\mathbf{v},\theta,t)\tilde{\phi}(\mathbf{v}_0,\theta_0,t_0) \right\rangle , \qquad (9)$$

where ϕ and $\tilde{\phi}$ are the annihilation and Doi-shifted [27] creation fields, respectively.

The velocity-velocity correlation function can be calculated by

$$\langle \mathbf{v}(t)\mathbf{v}(0)\rangle = \int \mathrm{d}^2 v(t) \mathrm{d}^2 v(0) \int_0^{2\pi} \mathrm{d}\theta \mathrm{d}\theta' \,\mathbf{v}(t) \mathcal{G}\big(\mathbf{v}(t), \theta, t | \mathbf{v}(0), \theta', 0\big) \mathbf{v}(0) \mathcal{G}\big(\mathbf{v}(0), \theta', 0\big) \,. \tag{10}$$

Since we choose the stationary state as the initial state, the most right propagator $\mathcal{G}(\mathbf{v}(0), \theta', 0)$ does not dependent on the initial condition. By using the orthogonality of the Hermite polynomials [38], the above integral can be simplified further, and the result will be presented in the following sections.

III. DIFFUSIVE PARTICLE IN 1D SPACE

We start with the simplest case, i.e., a diffusive particle with dry friction in 1D space. The corresponding Fokker-Planck equation is

$$\partial_t P(v,t) = D\gamma^2 \partial_v^2 P(v,t) + \gamma \partial_v v P(v,t) + \mu \partial_v \frac{v}{|v|} P(v,t) , \qquad (11)$$

where in 1D space, the unit vector $\mathbf{v}/||\mathbf{v}||$ becomes the sign function v/|v| where |v| is the absolute value of v. We further introduce the notations

$$\mathcal{L}_0 = D\gamma^2 \partial_v^2 + \gamma \partial_v v , \qquad (12a)$$

$$\mathcal{L}_{\mu} = \mu \partial_{v} \frac{v}{|v|} . \tag{12b}$$

1. Action

The corresponding bilinear action and perturbative action are

$$\mathcal{A} = \mathcal{A}_0 + \mathcal{A}_{\text{pert}} , \qquad (13a)$$

$$\mathcal{A}_{0}(\phi,\tilde{\phi}) = \int \mathrm{d}t \, \int \mathrm{d}v \,\tilde{\phi}(v,t) \big(-\partial_{t} + \mathcal{L}_{0} - r \big) \phi(v,t) \,, \tag{13b}$$

$$\mathcal{A}_{\text{pert}}(\phi, \tilde{\phi}) = \int dt \int dv \, \tilde{\phi}(v, t) \mathcal{L}_{\mu} \phi(v, t) \,. \tag{13c}$$

where r is the death rate to maintain the causality, and will be taken to zero after the inverse Fourier transform. We have introduced the annihilation field ϕ and the Doi-shifted creation field $\tilde{\phi}$ as [25, 27]

$$\phi(v,t) = \frac{1}{\Omega} \int d\omega e^{-i\omega t} \sum_{n=0}^{\infty} u_n(v)\phi_n(\omega) , \qquad (14a)$$

$$\tilde{\phi}(v,t) = \frac{1}{\Omega} \int d\omega e^{-i\omega t} \sum_{n=0}^{\infty} \tilde{u}_n(v) \tilde{\phi}_n(\omega) .$$
(14b)

Here $\Omega^2 = D\gamma$, $d\omega = d\omega/(2\pi)$ and further $\delta(2\pi) = 2\pi\delta(\omega)$, while u and \tilde{u} are consistent with Eq. (6). Any expectation value can be calculated perturbatively by a path integral [39]

$$\langle \bullet \rangle = \int \mathcal{D}\phi \, \mathcal{D}\tilde{\phi} \, \bullet \, \mathrm{e}^{\mathcal{A}[\tilde{\phi},\phi]} = \left\langle \bullet \, \mathrm{e}^{\mathcal{A}_{\mathrm{pert}}[\tilde{\phi},\phi]} \right\rangle_{0} \quad \text{with} \quad \left\langle \bullet \right\rangle_{0} = \int \mathcal{D}\phi \, \mathcal{D}\tilde{\phi} \, \bullet \, \mathrm{e}^{\mathcal{A}_{0}[\tilde{\phi},\phi]} \,, \tag{15}$$

where we split the action via Eq. (13a). Expanding the exponential with respect to the perturbative part A_{pert} , we obtain

$$\langle \bullet \rangle = \int \mathcal{D}\phi \, \mathcal{D}\tilde{\phi} \, \mathrm{e}^{\mathcal{A}_0[\tilde{\phi},\phi]} \bullet \sum_{n=0}^{\infty} \frac{\mathcal{A}_{\mathrm{pert}}^n}{n!} = \left\langle \bullet \sum_{n=0}^{\infty} \frac{\mathcal{A}_{\mathrm{pert}}^n}{n!} \right\rangle_0 \,. \tag{16}$$

By plugging the fields Eq. (14) into the action Eq. (13b), we obtain

$$\mathcal{A}_{0} = -\frac{1}{\Omega^{2}} \int d\omega \int d\omega' \sum_{n,m=0}^{\infty} \tilde{\phi}_{m}(\omega') \Big(-i\omega + \gamma n + r \Big) \phi_{n}(\omega) \delta(\omega + \omega') \Omega \delta_{n,m} , \qquad (17)$$

where $\Omega \delta_{n,m}$ comes from the orthogonality relation Eq. (7) between u_n and u_m . The bare propagator can be read off as

$$\left\langle \phi_n(\omega)\tilde{\phi}_m(\omega')\right\rangle_0 = \frac{\Omega\delta_{n,m}\delta(\omega+\omega')}{-i\omega+\gamma n+r} = \Omega\delta_{n,m}\delta(\omega+\omega')G(n,\omega) \triangleq \frac{n,\omega}{m,\omega'} \quad . \tag{18}$$

Similarly, we substitute the fields in Eq. (14) into the perturbative action in Eq. (13c),

$$\mathcal{A}_{\text{pert}} = \sum_{n,m} \frac{\mu}{\Omega^2} \int \mathrm{d}\omega \mathrm{d}\omega' \tilde{\phi}_m(\omega') \phi_n(\omega) \int \mathrm{d}v \, \tilde{u}_m(v) \partial_v \big[\frac{v}{|v|} u_n(v) \big] \delta(\omega + \omega') \tag{19a}$$

$$=\sum_{n,m}\frac{\mu}{\Omega^2}\int \mathrm{d}\omega\tilde{\phi}_m(-\omega)\phi_n(\omega)\int \mathrm{d}v\left(2\delta(0)\tilde{u}_m(v)u_n(v)+\frac{v}{|v|}\tilde{u}_m(v)\partial_v u_n(v)\right)\,,\tag{19b}$$

where $\delta(0)$ comes from the derivative of the sign function

~

$$\partial_v \left(\frac{v}{|v|}\right) = 2\delta(0) \ . \tag{20}$$

We diagrammatically write the perturbative part of the action as

$$\frac{\mu}{\Omega^2} \int dv \left(2\delta(0)\tilde{u}_m(v)u_n(v) + \frac{v}{|v|}\tilde{u}_m(v)\partial_v u_n(v) \right) = \Lambda^{m,n} \triangleq \frac{m-n}{|v|} \quad .$$
⁽²¹⁾

By using the following properties of the Hermite polynomials,

$$He_{n}(v) He_{m}(v) = \sum_{k=0}^{\min(n,m)} \frac{n!m!}{(n-k)!(m-k)!k!} He_{n+m-2k}(v) , \qquad (22a)$$

$$\partial_v \left(He_n(v) e^{-\frac{v^2}{2}} \right) = -He_{n+1}(v) e^{-\frac{v^2}{2}},$$
(22b)

the analytic form of the above vertex becomes

$$\frac{m}{\Omega^2} = \frac{\mu}{\Omega^2} \frac{2}{\sqrt{2\pi}m!} \left(He_n(0) He_m(0) - \sum_{k=0}^{\min(n+1,m)} \frac{(n+1)!m!}{(n+1-k)!(m-k)!k!} He_{n+m-2k}(0) \right),$$
(23)

where $He_n(0)$ is the "Hermite zero", which is probabilist's Hermite polynomials evaluated at zero with respect to the *n*-th order. It is trivial to see that

$$\Lambda^{0,n} = 0 \quad \text{for} \quad n \in \mathbb{Z}_0^+ \ , \tag{24}$$

which indicates that a propagator does not have an outgoing index 0 unless it is a bare one. Moreover, since the "Hermite zero" of arbitrary odd order is zero, $He_{2n+1}(0) = 0$, we find

$$\Lambda^{2m+1,2n} = \Lambda^{2m,2n+1} = 0 \quad \text{for} \quad n,m \in \mathbb{Z}_0^+ \ , \tag{25}$$

which shows the outgoing and incoming indices should have the same parity. Otherwise, the corresponding combinations become zero.

In the following, we only consider the perturbative vertices with the limited indices n, m = 0, 1, 2, which are

$$\Lambda^{2,0} = \Lambda^{1,1} = \Lambda^{2,2} = -\sqrt{\frac{2}{\pi} \frac{\mu}{\Omega^2}} , \qquad (26a)$$

$$\Lambda^{0,0} = \Lambda^{0,1} = \Lambda^{1,0} = \Lambda^{0,2} = \Lambda^{1,2} = \Lambda^{2,1} = 0.$$
(26b)

2. Propagator

By using the bare propagator and perturbative vertices, the full propagator can be written in a perturbative way as

$$\left\langle \phi_{n}(\omega)\tilde{\phi}_{m}(\omega')\right\rangle \triangleq \frac{n,\omega-m,\omega'}{m,\omega'} + \frac{n,\omega-m,\omega'}{m,\omega'} + \frac{n,\omega}{m,\omega'} + \frac{n,\omega'}{m,\omega'} + \dots \quad (27a)$$
$$= \delta(\omega+\omega')\left\{\Omega\delta_{n,m}G(n,\omega) + \Omega^{2}G(n,\omega)\Lambda^{n,m}G(m,\omega) + \dots\right\}.$$
(27b)

The full propagators is therefore

$$\left\langle \phi_n(\omega)\tilde{\phi}_m(\omega')\right\rangle = \delta\left(\omega + \omega'\right)\sum_{j=0}^{\infty} E_j(n,m,\omega) ,$$
 (28)

and the corresponding E_j are given by

$$E_0(n,m,\omega) = \Omega \delta_{n,m} G(n,\omega) \tag{29a}$$

$$E_1(n,m,\omega) = \Omega^2 G(n,\omega) \Lambda^{n,m} G(m,\omega)$$
(29b)

$$E_2(n,m,\omega) = \Omega^3 G(n,\omega) \sum_{q=0}^{\infty} \Lambda^{n,q} G(q,\omega) \Lambda^{q,m} G(m,\omega) = \Omega G(n,\omega) \sum_{q=0}^{\infty} \Lambda^{n,q} E_1(q,m,\omega) .$$
(29c)

We further introduce the recurrence relation between E_j and E_{j+1}

$$E_{j+1}(n,m,\omega) = \Omega G(n,\omega) \sum_{q=0}^{\infty} \Lambda^{n,q} E_j(q,m,\omega)$$
(30)

The probability density of velocity v at time t with a given initial state v' at time t' is

$$\mathcal{G}(v,t|v',t') = \frac{1}{\Omega^2} \sum_{n,m=0}^{\infty} u_n(v) \tilde{u}_m(v') \lim_{r \downarrow 0} \int d\omega d\omega' e^{-i\omega t} e^{-i\omega' t'} \left\langle \phi_n(\omega) \tilde{\phi}_m(\omega') \right\rangle$$
(31a)

$$= \frac{1}{\Omega^2} \sum_{n,m=0}^{\infty} u_n(v) \tilde{u}_m(v') \lim_{r \downarrow 0} \int \mathrm{d} \omega \mathrm{e}^{-i\omega(t-t')} \sum_j E_j(n,m,\omega) \;. \tag{31b}$$

3. Stationary-state correlation function

By taking the limit $t_0 \to -\infty$, we obtain the stationary density

$$\mathcal{G}(v) = \lim_{t_0 \to -\infty} \left\langle \phi(v, t) \tilde{\phi}(v_0, t_0) \right\rangle .$$
(32)

Previous work shows that taking the stationary limit $t_0 \to -\infty$ and the zero death rate limit $r \downarrow 0$ replaces the incoming index by $\delta_{m,0}$ [34], which physically indicates that the steady-state is independent of the initial condition. Therefore, Eq. (30) is

where we have dropped the arguments m, ω , since the observable no longer depends on time and initial state. The form of the stationary density is

$$\mathcal{G}(v) = \frac{1}{\Omega^2} \sum_{n=0}^{\infty} u_n(v) \tilde{u}_0 \sum_{j=0}^{\infty} E_j(n) .$$
(34)

$$E_0(n) = \Omega \delta_{n,0} , \qquad (35a)$$

$$E_1(n) = \frac{\Lambda^{n,0} \Omega^2}{n\gamma} , \qquad (35b)$$

$$E_{j+1}(n) = \frac{\Lambda^{n,q}\Omega}{n\gamma} \sum_{q=1} E_j(q) .$$
(35c)

Since $\Lambda^{0,n}$ is always zero in Eq. (24) which vanishes all the propagators with outgoing index n = 0 before taking stationary limit, we define $\lim_{n\to 0} \Lambda^{n,m}/n = 0$ for arbitrary m and perform the summation from q = 1 to avoid the zero denominators in Eq. (35).

Our aim is to use the Green-Kubo relation Eq. (1) to calculate the effective diffusion coefficient. Now, we perform the integral of the velocity-velocity correlation function in the present field theory framework

$$\langle v(t)v(t')\rangle = \int dv \int dv' \, v \mathcal{G}(v,t|v',t')v' \mathcal{G}(v') , \qquad (36)$$

where we simplify the notation v = v(t) and v' = v(t'). By using Eqs. (10), (31), (34) and the orthogonality of the Hermite polynomials [38], the correlation function becomes

$$\langle v(t)v(t')\rangle = \int d\omega e^{-i\omega(t-t')} \left\langle \phi_1(\omega)\tilde{\phi}_1(-\omega) \right\rangle \left(2\left\langle \phi_2\tilde{\phi}_0 \right\rangle + \left\langle \phi_0\tilde{\phi} \right\rangle \right) \,. \tag{37}$$

We first calculate the time-independent observable in the brackets as

$$\left\langle \phi_2 \tilde{\phi}_0 \right\rangle = \sum_{j=0} E_j(2) \approx -\Omega \frac{\beta}{2+\beta} ,$$
 (38a)

$$\left\langle \phi_0 \tilde{\phi}_0 \right\rangle = \Omega$$
, (38b)

where we only consider the finite number of perturbative vertices up to the second order, and the details are shown in Appendix A_1 . We introduce a dimensionless parameter

$$\beta = \sqrt{\frac{2}{\pi}} \frac{\mu}{\Omega \gamma} \ . \tag{39}$$

to simplify the notation. Then we have

$$2\left\langle\phi_2(0)\tilde{\phi}_0(0)\right\rangle + \left\langle\phi_0(0)\tilde{\phi}(0)\right\rangle \approx \Omega\left(1 - \frac{\beta}{1 + \frac{\beta}{2}}\right).$$

$$\tag{40}$$

To calculate the time-dependent observable, we only consider the perturbative part with both incoming and outgoing indices fixed to unity,

$$\int d\omega e^{-i\omega(t-t')} \left\langle \phi_1(\omega)\tilde{\phi}_1(-\omega) \right\rangle$$

$$\approx \int d\omega e^{-i\omega(t-t')} \frac{1}{\omega} \frac{1}{\omega} \frac{1}{\omega} + \frac{1}{\omega} \frac{1}{\omega} \frac{1}{\omega} + \frac{1}{\omega} \frac{1}{\omega} \frac{1}{\omega} + \frac{1}{\omega} \frac{1}{\omega} \frac{1}{\omega} \frac{1}{\omega} + \dots \quad (41a)$$

$$\int d\omega e^{-i\omega(t-t')} \Omega G(1,\omega) \sum_{\ell=0}^{\infty} \left(\Omega G(1,\omega) \Lambda^{1,1} \right)^{\ell}$$
(41b)

$$= \int d\omega e^{-i\omega(t-t')} \frac{\Omega G(1,\omega)}{1 - \Omega G(1,\omega)\Lambda^{1,1}}$$
(41c)

$$= \int d\omega \, e^{-i\omega(t-t')} \frac{\Omega}{-i\omega + \gamma + \sqrt{\frac{2}{\pi}} \frac{\mu}{\Omega}} = \Omega e^{-(\gamma + \gamma\beta)(t-t')}$$
(41d)

Then, the time-correlation function is obtained by substituting Eqs. (40) and (41d) into Eq. (37),

$$\langle v(t)v(t')\rangle \approx \Omega^2 \mathrm{e}^{-\gamma(1+\beta)(t-t')} \left(1 - \frac{\beta}{1+\frac{\beta}{2}}\right).$$
 (42)

By using the Green-Kubo relation in Eq. (1) and recalling $\Omega^2 = D\gamma$, the effective diffusion coefficient is

$$D_{\text{eff}} = \int_{t'}^{\infty} \mathrm{d}t \, \left\langle v(t)v(t')\right\rangle \approx \frac{D}{1+\beta} \left(1 - \frac{\beta}{1+\frac{\beta}{2}}\right) \,. \tag{43}$$

If there is no dry friction, $\beta \to 0$, we recover the normal translational diffusion coefficient $D_{\text{eff}}|_{\beta\to 0} = D$, as it should. By expanding the fractions, we obtain the first order correction with respect to the parameter β as

$$D_{\text{eff}} = D(1 - 2\beta) + \mathcal{O}(\beta^2) . \tag{44}$$

IV. DIFFUSIVE PARTICLE IN 2D SPACE

In 2D space, the corresponding Fokker-Planck equation of a diffusive particle with dry friction is

$$\partial_t P(\mathbf{v}, t) = D\gamma^2 \nabla_{\mathbf{v}}^2 P(\mathbf{v}, t) + \gamma \nabla_{\mathbf{v}} \cdot \left[\mathbf{v} P(\mathbf{v}, t) \right] + \mu \nabla_{\mathbf{v}} \cdot \left[\frac{\mathbf{v}}{||\mathbf{v}||} P(\mathbf{v}, t) \right] \,. \tag{45}$$

Similar to the 1D case, we split the operator into two parts,

$$\mathcal{L}_0 = D\gamma^2 \nabla_{\mathbf{v}}^2 + \gamma \nabla_{\mathbf{v}} \cdot \mathbf{v} , \qquad (46a)$$

$$\mathcal{L}_{\mu} = \mu \nabla_{\mathbf{v}} \cdot \frac{\mathbf{v}}{||\mathbf{v}||} . \tag{46b}$$

Applying the isotropic property of the first operator \mathcal{L}_0 , one can simplify the calculation into 1D space with an extra prefactor 2. However, because of the anisotropy of dry friction, there is no such a way to simplify the operator \mathcal{L}_{μ} . Hence, we use an approximation of the "sign" function of the velocity **v**

$$\frac{\mathbf{v}}{||\mathbf{v}||} = \begin{bmatrix} \frac{v_1}{\sqrt{v_1^2 + v_2^2}} \\ \frac{v_2}{\sqrt{v_2^2 + v_2^2}} \end{bmatrix} = \begin{bmatrix} \frac{v_1}{|v_1|\sqrt{1 + \frac{v_2^2}{v_1^2}}} \\ \frac{v_2}{|v_2|\sqrt{1 + \frac{v_1^2}{v_2^2}}} \end{bmatrix} \approx e^{-\frac{1}{4}} \begin{bmatrix} \frac{v_1}{|v_1|} \\ \frac{v_2}{|v_2|} \end{bmatrix} , \qquad (47)$$

where we use the following Hermite expansion of the square root in the denominator

$$\frac{1}{\sqrt{1+z^2}} = e^{-\frac{1}{4}} \sum_{k=0}^{\infty} \frac{(-1)^k}{2^k (2k)!} He_{2k} \left(\sqrt{2}\theta\right) , \qquad (48)$$

with $\theta = \arctan(z)$. By considering only the zeroth order Hermite polynomial and dropping all the higher order terms, we find the approximation in Eq. (47). The operator \mathcal{L}_{μ} is rewritten as

$$\mathcal{L}_{\mu} \approx \mathcal{L}_{\mu}' = \mu \mathrm{e}^{-\frac{1}{4}} \left(\partial_{v_1} \frac{v_1}{|v_1|} + \partial_{v_2} \frac{v_2}{|v_2|} \right) \,, \tag{49}$$

which is now isotropic. Then, we can simplify this 2D problem to a 1D problem whose corresponding Fokker-Planck equation is

$$\partial_t P^{(2)}(v,t) = D\gamma^2 \partial_v^2 P^{(2)}(v,t) + \gamma \partial_v v P^{(2)}(v,t) + e^{-\frac{1}{4}} \mu \partial_v \frac{v}{|v|} P^{(2)}(v,t) , \qquad (50)$$

where we have introduced an upper index $^{(2)}$ to indicate that the above Fokker-Planck equation is obtained from the 2D equation. The corresponding velocity-velocity correlation function is

$$\langle \mathbf{v}(t) \cdot \mathbf{v}(t') \rangle = 2 \langle v(t)v(t') \rangle \quad (51)$$

where the prefactor 2 in the RHS is the dimension factor.

The only difference between the Fokker-Planck equation in Eq. (50) and 1D case in Eq. (11) is an extra prefactor $\exp(-1/4)$ in the friction term. Therefore, by following the same steps presented in 1D case in Sec. III, we immediately obtain the velocity-velocity correlation and the effective diffusion coefficient by introducing the modified dimensionless parameter as

$$\tilde{\beta} = e^{-\frac{1}{4}} \sqrt{\frac{2}{\pi}} \frac{\mu}{\Omega \gamma} , \qquad (52)$$

and they are

$$\langle \mathbf{v}(t) \cdot \mathbf{v}(t') \rangle \approx 2D\gamma \mathrm{e}^{-\gamma(1+\tilde{\beta})(t-t')} \left(1 - \frac{\tilde{\beta}}{1+\frac{\tilde{\beta}}{2}}\right),$$
(53a)

$$D_{\text{eff}}^{(2)} \approx \frac{D}{1+\tilde{\beta}} \left(1 - \frac{\tilde{\beta}}{1+\frac{\tilde{\beta}}{2}} \right) .$$
 (53b)

Here we use an upper index ⁽²⁾ to distinguish the effective diffusion coefficient in 2D from the 1D result in Eq. (43), where the parameter $\tilde{\beta}$ in the 2D problem is changed from the parameter β in the 1D case.

We compare our result in Eq. (53b) with the numerical simulation. In this work, only finite types of the perturbation vertices are considered, the friction operator is approximated by Eq. (49), and the result is a perturbation calculation using the first three orders of the Hermite polynomials. We compare the effective diffusion coefficient with the numerical simulation and our field theory approach in Fig. 1(a), showing a good agreement between them even $\tilde{\beta}$ is large. Additionally, the fourth order correction of the time-independent observable $\langle \phi_2 \tilde{\phi}_0 \rangle$ is also calculated in Appendix A 1.

V. ACTIVE BROWNIAN PARTICLE

ABPs are particles that move with constant speed $|\mathbf{w}| = w$ but whose director θ diffuses with the rotational diffusion coefficient $D_{\rm r}$. The corresponding Fokker-Planck equation of a 2D ABP with dry friction is

$$\partial_t P(\mathbf{v},\theta,t) = \mathcal{L}P(\mathbf{v},\theta,t) \quad \text{with} \quad \mathcal{L} = D\gamma^2 \nabla_{\mathbf{v}} + \gamma \nabla_{\mathbf{v}} \cdot \mathbf{v} + \mu \nabla \cdot \frac{\mathbf{v}}{||\mathbf{v}||} - \gamma \mathbf{w}_\theta \cdot \nabla + D_r \partial_\theta^2 , \tag{54}$$

where $\mathbf{w}_{\theta} = w[\cos\theta, \sin\theta]^{\mathrm{T}}$.

Similar to the approach presented in Sec. IV, we first use the approximation of the friction term in Eqs. (47) and (49), to approximate the anisotropic operator by an isotropic one. Second, we reduce the problem to 1D and choose the component of \mathbf{v} on the x-axis of the Cartesian plane. The Fokker-Planck equation is then

$$\partial_t P^{(2)}(v,\theta,t) = \left(\mathcal{L}_0 + \mathcal{L}_\mu + \mathcal{L}_w\right) P^{(2)}(v,\theta,t) \tag{55}$$

with

$$\mathcal{L}_0 = D\gamma^2 \partial_v^2 + \gamma \partial_v v + D_r \partial_\theta^2 , \qquad (56a)$$

$$\mathcal{L}_{\mu} \approx \mathcal{L}_{\mu}' = \mu \mathrm{e}^{-\frac{1}{4}} \partial_{v} \frac{v}{|v|} , \qquad (56b)$$

$$\mathcal{L}_w = -\gamma w \cos \theta \partial_v \ . \tag{56c}$$

A. Action

Accordingly, the bilinear action and perturbative actions are

$$\mathcal{A}_{0}(\phi,\tilde{\phi}) = \int \mathrm{d}t \, \int \mathrm{d}v \, \int_{0}^{2\pi} \mathrm{d}\theta \, \tilde{\phi}(v,\theta,t) \big(-\partial_{t} + \mathcal{L}_{0} - r \big) \phi(v,\theta,t) \,, \tag{57a}$$

$$\mathcal{A}_{\mu}(\phi,\tilde{\phi}) = \int \mathrm{d}t \, \int \mathrm{d}v \, \int_{0}^{2\pi} \mathrm{d}\theta \, \tilde{\phi}(v,\theta,t) \mathcal{L}_{\mu}'\phi(v,\theta,t) \,, \tag{57b}$$

$$\mathcal{A}_w(\phi, \tilde{\phi}) = \int \mathrm{d}t \, \int \mathrm{d}v \, \int_0^{2\pi} \mathrm{d}\theta \, \tilde{\phi}(v, \theta, t) \mathcal{L}_w \phi(v, \theta, t) \;. \tag{57c}$$

Since there is a new variable θ , the fields in this case are introduced as

$$\phi(v,\theta,t) = \frac{1}{\Omega} \sum_{n=0}^{\infty} \sum_{\alpha=-\infty}^{\infty} \int d\omega e^{-i\omega t} u_n(v) e^{-i\alpha \theta} \phi_{n,\alpha}(\omega) , \qquad (58a)$$

$$\tilde{\phi}(v,\theta,t) = \frac{1}{2\pi\Omega} \sum_{n=0}^{\infty} \sum_{\alpha=-\infty}^{\infty} \int d\omega e^{-i\omega t} \tilde{u}_n(v) e^{i\alpha\theta} \tilde{\phi}_{n,\alpha}(\omega) .$$
(58b)

Here, we use Roman letters for the velocity indices and Greek letters for the director indices. Similar to the 1D case, the corresponding bilinear action and the μ -dependent perturbative action are

$$\mathcal{A}_{0} = -\frac{1}{2\pi\Omega^{2}} \sum_{n,m=0}^{\infty} \sum_{\alpha,\alpha'=-\infty}^{\infty} \int d\omega d\omega' \tilde{\phi}_{m,\alpha'}(\omega') \left(-i\omega + \gamma n + D_{r}\alpha^{2} + r\right) \phi_{n,\alpha}(\omega) \delta(\omega + \omega') \Omega \delta_{n,m} 2\pi \delta_{\alpha,\alpha'} , \qquad (59a)$$
$$\mathcal{A}_{\mu} = \frac{\mu \exp\left(-1/4\right)}{2\pi\Omega^{2}} \sum_{n,m=0}^{\infty} \sum_{\alpha,\alpha'=-\infty}^{\infty} \int d\omega \tilde{\phi}_{m,\alpha'}(-\omega) \phi_{n,\alpha}(\omega) \int dv \left(2\delta(0)\tilde{u}_{m}(v)u_{n}(v) + \frac{v}{|v|}\tilde{u}_{m}(v)\partial_{v}u_{n}(v)\right) 2\pi \delta_{\alpha,\alpha'} ,$$

and the self-motility dependent action \mathcal{A}_w is

$$\mathcal{A}_{w} = -\frac{\gamma w}{2\pi\Omega^{2}} \sum_{n,m=0}^{\infty} \sum_{\alpha,\alpha'=-\infty}^{\infty} \int \mathrm{d}\omega \tilde{\phi}_{m,\alpha'}(-\omega) \phi_{n,\alpha}(\omega) \int \mathrm{d}v \,\tilde{u}_{m}(v) \partial_{v} u_{n}(v) \int_{0}^{2\pi} \mathrm{d}\theta \,\mathrm{e}^{-i(\alpha-\alpha')\theta} \cos\theta \tag{60a}$$

$$= \frac{\gamma w}{2\pi\Omega^2} \sum_{n,m=0}^{\infty} \sum_{\alpha,\alpha'=-\infty}^{\infty} \int d\omega \tilde{\phi}_{m,\alpha'}(-\omega) \phi_{n,\alpha}(\omega) \delta_{n,m-1} 2\pi \frac{\delta_{\alpha+1,\alpha'} + \delta_{\alpha-1,\alpha'}}{2} .$$
(60b)

There are two different types of perturbative vertices. We diagrammatically write the friction-dependent and self-propulsion dependent vertices in the following way

$$\frac{\mu \exp\left(-\frac{1}{4}\right)}{\Omega^2} \int dv \left(2\delta(0)\tilde{u}_m(v)u_n(v) + \frac{v}{|v|}\tilde{u}_m(v)\partial_v u_n(v)\right) \delta_{\alpha,\alpha'} \triangleq \Lambda^{m,n}_{\alpha',\alpha} \triangleq \frac{m}{\alpha' \cdot \alpha} n$$
(61a)

$$\gamma w \frac{\delta_{\alpha+1,\alpha'} + \delta_{\alpha-1,\alpha'}}{2\Omega^2} \delta_{n,m-1} \triangleq \Upsilon^{m,n}_{\alpha',\alpha} \triangleq \overset{m}{\underset{\alpha'}{\longleftarrow}} \overset{n}{\alpha} , \qquad (61b)$$

and the bare propagator is

$$\frac{\Omega\delta_{n,m}\delta_{\alpha,\alpha'}\delta(\omega+\omega')}{-\iota\omega+\gamma n+D_{r}\alpha^{2}+r} = \Omega\delta_{n,m}\delta_{\alpha,\alpha'}\delta(\omega+\omega')G(n,\alpha,\omega) \triangleq \frac{n,\omega}{\alpha} \frac{m,\omega'}{\alpha'} .$$
(62)

B. Propagator

The full propagator is the summation of all the possible combination of the bare propagator in Eq. (62) and the perturbative vertices in Eq. (61)

$$\left\langle \phi_{n,\alpha}(\omega)\tilde{\phi}_{m,\alpha'}(\omega') \right\rangle \triangleq \begin{array}{c} n, \omega & m, \omega' \\ \alpha & \alpha' \end{array} + \begin{array}{c} n, \omega & m, \omega' \\ \alpha' & \alpha' \end{array} + \begin{array}{c} n, \omega & m, \omega' \\ \alpha' & \alpha' \end{array} + \begin{array}{c} n, \omega & m, \omega' \\ \alpha' & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{q} & m, \omega' \\ \alpha' & \sum_{\nu} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{q} & m, \omega' \\ \alpha' & \sum_{\nu} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{q} & m, \omega' \\ \alpha' & \sum_{\nu} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{q} & m, \omega' \\ \alpha' & \sum_{\nu} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{q} & m, \omega' \\ \alpha' & \sum_{\nu} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{q} & m, \omega' \\ \alpha' & \sum_{\nu} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{q} & m, \omega' \\ \alpha' & \sum_{\nu} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{q} & m, \omega' \\ \alpha' & \sum_{\nu} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{q} & m, \omega' \\ \alpha' & \sum_{\nu} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{q} & m, \omega' \\ \alpha' & \sum_{\nu} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{q} & m, \omega' \\ \alpha' & \sum_{\nu} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{q} & m, \omega' \\ \alpha' & \sum_{\nu} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{q} & m, \omega' \\ \alpha' & \sum_{\nu} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{q} & m, \omega' \\ \alpha' & \sum_{\nu} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{q} & m, \omega' \\ \alpha' & \sum_{\nu} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{q} & m, \omega' \\ \alpha' & \sum_{\nu} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{q} & m, \omega' \\ \alpha' & \sum_{\nu} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{q} & m, \omega' \\ \alpha' & \sum_{\nu} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & m, \omega' \\ \alpha' & \sum_{\nu} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & m, \omega' \\ \alpha' & \sum_{\nu} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & m, \omega' \\ \alpha' & \sum_{\nu} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & m, \omega' \\ \alpha' & \sum_{\nu} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & m, \omega' \\ \alpha' & \sum_{\nu} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & m, \omega' \\ \alpha' & \sum_{\nu} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & \alpha' \end{array} + \begin{array}{c} n, \omega & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & \alpha' \end{array} + \begin{array}{c} n, \omega & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & \alpha' \end{array} + \begin{array}{c} n, \omega & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & \alpha' \end{array} + \begin{array}{c} n, \omega & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum_{\alpha'} & \alpha' \end{array} + \begin{array}{c} n, \omega & \alpha' \end{array} + \begin{array}{c} n, \omega & \sum$$

(59b)

Similar to Eq. (28), we write the full propagator as,

$$\left\langle \phi_{n,\alpha}(\omega)\tilde{\phi}_{m,\alpha'}(\omega')\right\rangle = \delta(\omega+\omega')\sum_{j=0}^{\infty} E_j(n,m,\alpha,\alpha',\omega) ,$$
 (64)

where the terms shown in Eq. (63b) are exactly the first three orders of $E_j(n, m, \alpha, \alpha', \omega)$. We list the recurrence relation as

$$E_{j+1}(n,m,\alpha,\alpha',\omega) = \Omega G(n,\alpha,\omega) \sum_{q=0}^{\infty} \sum_{\nu=-\infty}^{\infty} \left(\Upsilon_{n,q}^{\alpha,\nu} + \Lambda_{n,q}^{\alpha,\nu}\right) E_j(q,m,\nu,\alpha',\omega) .$$
(65)

We therefore obtain the probability density of the velocity at v with director θ at time t with the corresponding initial state (v', θ', t') as

$$\mathcal{G}(v,\theta,t|v',\theta',t') = \frac{1}{2\pi\Omega^2} \sum_{n,m=0}^{\infty} \sum_{\alpha,\alpha'=-\infty}^{\infty} u_n(v)\tilde{u}_m(v') \mathrm{e}^{-i\alpha\theta} \mathrm{e}^{i\alpha'\theta'} \lim_{r\downarrow 0} \int \mathrm{d}\omega \mathrm{d}\omega' \delta(\omega+\omega') \left\langle \phi_{n,\alpha}(\omega)\tilde{\phi}_{m,\alpha'}(\omega') \right\rangle \quad (66a)$$

$$= \frac{1}{2\pi\Omega^2} \sum_{n,m=0}^{\infty} \sum_{\alpha,\alpha'=-\infty}^{\infty} u_n(v) \tilde{u}_m(v') \mathrm{e}^{-i\alpha\theta} \mathrm{e}^{i\alpha'\theta'} \lim_{r\downarrow 0} \int \mathrm{d}\omega \mathrm{e}^{-i\omega(t-t')} \sum_{j=0} E_j(n,m,\alpha,\alpha',\omega) \;. \tag{66b}$$

C. Velocity-velocity correlation function

By substituting Eq. (66b) into Eq. (10), and using the orthogonality of the Hermite polynomials and the exponential terms, we have

$$\left\langle v(t)v(t')\right\rangle = \left\langle \phi_{1,0}(t)\tilde{\phi}_{1,0}(t')\right\rangle \left(2\left\langle \phi_{2,0}\tilde{\phi}_{0,0}\right\rangle + \left\langle \phi_{0,0}\tilde{\phi}_{0,0}\right\rangle\right) + \sum_{\nu=-\infty}^{\infty} \left\langle \phi_{1,0}(t)\tilde{\phi}_{0,\nu}(t')\right\rangle \left\langle \phi_{1,\nu}\tilde{\phi}_{0,0}\right\rangle , \qquad (67)$$

where the second term comes from the integral over θ' . Since we only consider the first three orders of the speed indices n, m = 0, 1, 2 of the perturbation vertices in Eq. (61), only $\nu = \pm 1$ in the summation are concerned in the second term. We calculate the time-independent observable first by the inverse Fourier transform and take the limits $t_0 \rightarrow -\infty$ and $r \downarrow 0$

$$\left\langle \phi_{n,\alpha} \tilde{\phi}_{m,\alpha'} \right\rangle = \lim_{r \downarrow 0} \lim_{t_0 \to -\infty} \int \mathrm{d}\omega \mathrm{e}^{-i\omega(t-t_0)} \left\langle \phi_{n,\alpha}(\omega) \tilde{\phi}_{m,\alpha'}(-\omega) \right\rangle \ . \tag{68}$$

We obtain

$$\left\langle \phi_{0,0}\tilde{\phi}_{0,0}\right\rangle = \Omega$$
, (69a)

$$\left\langle \phi_{2,0}\tilde{\phi}_{0,0}\right\rangle \approx -\Omega \frac{\tilde{\beta}}{2+\tilde{\beta}} + \frac{w^2\gamma}{4\Omega(\gamma+D_{\rm r}+\tilde{\beta}\gamma)(1+\frac{\tilde{\beta}}{2})} ,$$
 (69b)

$$\left\langle \phi_{1,1}\tilde{\phi}_{0,0}\right\rangle = \left\langle \phi_{1,-1}\tilde{\phi}_{0,0}\right\rangle \approx \frac{w\gamma}{2(\gamma + D_{\rm r} + \tilde{\beta}\gamma)} ,$$
 (69c)

where the details are presented in Appendix A 2.

For the time-dependent correlation function, we only consider the lowest perturbation vertices, similar to Sec. III. The first one is

$$\left\langle \phi_{1,0}(t)\tilde{\phi}_{1,0}(t')\right\rangle = \int \mathrm{d}\omega \mathrm{e}^{-i\omega(t-t')} \left\langle \phi_{1,0}(\omega)\tilde{\phi}_{1,0}(-\omega)\right\rangle \tag{70a}$$

$$\approx \int d\omega e^{-i\omega(t-t')} \frac{1,\omega}{0} \frac{1,-\omega}{0} + \frac{1,\omega}{0} \frac{1,-\omega}{0} + \frac{1,\omega}{0} \frac{1,-\omega}{0} + \dots$$
(70b)

$$= \int \mathrm{d}\omega \mathrm{e}^{-i\omega(t-t')}\Omega G(1,0,\omega) \sum_{\ell=0}^{\infty} \left(\Omega G(1,0,\omega) \Lambda_{0,0}^{1,1}\right)^{\ell}$$
(70c)

$$= \int d\omega \, e^{-i\omega(t-t')} \frac{\Omega}{-i\omega + \gamma + \exp\left(-\frac{1}{4}\right)\sqrt{\frac{2}{\pi}\frac{\mu}{\Omega}}} = \Omega e^{-(\gamma + \gamma\tilde{\beta})(t-t')} , \qquad (70d)$$

where $\tilde{\beta}$ is the corresponding dimensionless parameter introduced in Eq. (52). The second term is

$$\left\langle \phi_{1,0}(t)\tilde{\phi}_{0,1}(t')\right\rangle = \left\langle \phi_{1,0}(t)\tilde{\phi}_{0,-1}(t')\right\rangle = \int d\omega e^{-i\omega(t-t')} \left\langle \phi_{1,0}(\omega)\tilde{\phi}_{0,1}(-\omega)\right\rangle$$
(71a)

$$+ \frac{1,\omega}{0} \qquad 1 \qquad 1 \qquad 0,-\omega$$
 (71b)

$$= \int \mathrm{d}\omega \mathrm{e}^{-i\omega(t-t')} \Omega^2 G(1,0,\omega) \Upsilon_{0,1}^{1,0} G(0,1,\omega) \sum_{\ell=0}^{\infty} \left(\Omega G(1,0,\omega) \Lambda_{0,0}^{1,1} \right)^{\ell}$$
(71c)

$$= \frac{w\gamma}{2(\gamma + \tilde{\beta}\gamma - D_{\rm r})} \left(e^{-D_{\rm r}(t-t')} - e^{-\gamma(1+\tilde{\beta})(t-t')} \right).$$
(71d)

Then, the velocity-velocity correlation function is

$$\langle v(t)v(t')\rangle \approx \Omega^2 \mathrm{e}^{-\gamma(1+\tilde{\beta})(t-t')} \left(1 - \frac{\tilde{\beta}}{1+\frac{\tilde{\beta}}{2}}\right) + \frac{\gamma w^2 \mathrm{e}^{-\gamma(1+\tilde{\beta})(t-t')}}{2(\gamma+D_\mathrm{r}+\gamma\tilde{\beta})(1+\frac{\tilde{\beta}}{2})} + \frac{\mathrm{e}^{-D_\mathrm{r}(t-t')} - \mathrm{e}^{-\gamma(1+\tilde{\beta})(t-t')}}{\gamma+\tilde{\beta}\gamma-D_\mathrm{r}} \times \frac{w^2\gamma^2}{2(\gamma+D_\mathrm{r}+\tilde{\beta}\gamma)} \,. \tag{72}$$

By applying the Green-Kubo relation in Eq. (1), we obtain the effective diffusion coefficient for an ABP with dry friction as

$$D_{\text{eff}}^{(2)} \approx \frac{D}{1+\tilde{\beta}} \left(1 - \frac{\tilde{\beta}}{1+\frac{\tilde{\beta}}{2}}\right) + \frac{w^2}{2(\gamma + D_{\text{r}} + \gamma\tilde{\beta})(1+\tilde{\beta})(1+\frac{\tilde{\beta}}{2})} + \frac{w^2\gamma}{2D_{\text{r}}(\gamma + D_{\text{r}} + \gamma\tilde{\beta})(1+\tilde{\beta})}$$
(73)

When there is no friction, $\tilde{\beta} = 0$, we recover the effective diffusion for an isolated ABP in Eq. (5),

$$D_{\rm eff}^{(2)}\big|_{\tilde{\beta}\to 0} = D + \frac{w^2}{2D_{\rm r}} = D_0 \ . \tag{74}$$

Similar to Eq. (44), by expanding the fractions, we obtain the first order correction of the effective diffusion coefficient for an ABP with dry friction as

$$D_{\rm eff}^{(2)} = D(1 - 2\tilde{\beta}) + \frac{w^2}{2D_{\rm r}} \left(1 - \frac{3D_{\rm r} + 4\gamma}{2(D_{\rm r} + \gamma)}\tilde{\beta} \right) + \mathcal{O}(\tilde{\beta}^2) .$$
(75)

We further compare the analytical result in Eq. (73) with the numerical simulation in Fig. 1(b). Even with the existence of the self-propulsion, the agreement between the field theoretic approach and the simulation is still good.



FIG. 1. Comparison between the analytical result and numerical simulation. The horizontal axis is the dimensionless parameter $\tilde{\beta}$ in Eq. (52), and the vertical axis is the ratio between the effective diffusion coefficient $D_{\text{eff}}^{(2)}$ in Eqs. (53b), (73) and the diffusion coefficient D_0 in the absence of dry friction in Eq. (74). The numerical result of $D_{\text{eff}}^{(2)}$ is obtained from the Langevin equation in Eq. (3) via the Euler-Maruyama method, where the deatils are explained in Appendix B. We have used the following parameters: number of the particles N = 20000, total time T = 400, and time step $\Delta t = 0.05$. The initial state of the simulation is $\mathbf{x}(-50) = \mathbf{0}$ and $\theta(-50) = 0$, where we have spent $T_0 = 50$ to wait for the system to reach the stationary state and then started to average the MSD.

VI. SUMMARY AND DISCUSSION

In this paper, we first discussed the 1D diffusing particle with dry friction. From the velocity-corresponding Fokker-Planck equation in Eq. (11), we determined the bilinear and perturbative actions for a diffusive particle with dry friction in Eqs. (17) and (19), respectively. The full propagator was derived in Eq. (27). Considering the first three orders of the Hermite polynomials, we only count three types of perturbative vertices with non-zero contributions Eq. (26). Therefore, the summation in the propagator is reduced to a single term, and the geometric sum is used to obtain the velocity-velocity correlation function in Eq. (42). Equation (43) was further obtained by using the Green-Kubo relation Eq. (1).

Then, we have extended this framework to the 2D space. For wet friction, we can always reduce the higher dimensional problem to a 1D problem because the system is isotropic. For dry friction, however, such a treatment is no longer possible because it is anisotropic. We then introduced a Hermite expansion of dry friction, and transformed dry friction into an isotropic operator in Eq. (47) by considering the leading order. With a prefactor difference, we obtained the 2D effective diffusion coefficient Eq. (53b) from the 1D result in Eq. (43).

Studying the ABP problem is more complicated since the self-propulsion of the particle provides another type of perturbation vertex. By using the same treatment as in the 1D case, we only considered the limited Hermite order $(n \leq 2)$. Then, we applied the geometric sums to calculate the velocity-velocity correlation function in Eq. (72) and further obtained the effective diffusion coefficient in Eq. (73). Since we have neglected higher orders of the Hermite expansion for both fields and dry friction operator, the diffusion coefficient is a perturbative result. However, the analytical result in Eq. (73) recovers the numerical simulation very well.

Dry friction force F in the experiment can be estimated by using Eq. (73) or the first order expansion in Eq. (75). Substituting back $\Gamma = m\gamma$ and $F = m\mu$, and using the Stokes–Einstein–Sutherland relations $\Gamma = 6\pi\eta a$ and $D = k_{\rm B}T/\Gamma$, where η is the fluid viscosity, a is the radius of the particle, $k_{\rm B}$ is the Boltzmann constant and T is the temperature, we rewrite the dimensionless parameter $\hat{\beta}$ as

$$\tilde{\beta} = e^{-\frac{1}{4}} F \sqrt{\frac{2m}{\pi D \Gamma^3}} = e^{-\frac{1}{4}} \frac{F}{6\pi \eta a} \sqrt{\frac{2m}{\pi k_{\rm B} T}} .$$
(76)

Considering a colloidal particle with a mass $m \approx 10^{-15}$ kg and a radius $a \approx 10^{-6}$ m in water with the room temperature, and assuming $F \approx 0.1 \times mg$, where g is the gravitational constant, we estimate $\tilde{\beta} \approx 5 \times 10^{-4}$. If the radius is increased to $a \approx 5 \times 10^{-6}$ m, the corresponding parameter becomes $\tilde{\beta} \approx 0.14$, which leads to a substantial decrease of the effective diffusion coefficient.

The present work provides a basis for the characterization of the motion of active matter in velocity space via field theory, especially for mesoscopic particles whose diffusion and mass can not be neglected. Because of the wet friction γ , the calculation is simplified by using Hermite expansion of the fields [36] rather than the Fourier transform [34]. This work also provides a basis for ABPs with harmonic interactions. With the aid of the previous work on field theoretic approach of interacting diffusive particles [39, 40], we are currently working on ABPs with non-reciprocal harmonic interactions.

ACKNOWLEDGEMENTS

S.K. acknowledges the support by National Natural Science Foundation of China (Nos. 12274098 and 12250710127) and the startup grant of Wenzhou Institute, University of Chinese Academy of Sciences (No. WIUCASQD2021041). This work was supported by the JSPS Core-to-Core Program "Advanced core-to-core network for the physics of self-organizing active matter" (JPJSCCA20230002).

- M.E. Cates. Diffusive transport without detailed balance in motile bacteria: does microbiology need statistical physics? *Rep. Prog. Phys*, 75(4):042601, 2012.
- [2] C. Bechinger, R. Di Leonardo, H. Löwen, C. Reichhardt, G. Volpe, and G. Volpe. Active particles in complex and crowded environments. *Rev. Mod. Phys.*, 88:045006, 2016.
- [3] M. C. Marchetti, J. F. Joanny, S. Ramaswamy, T. B. Liverpool, J. Prost, Madan Rao, and R. Aditi Simha. Hydrodynamics of soft active matter. *Rev. Mod. Phys.*, 85:1143–1189, 2013.
- [4] S. Ramaswamy. The mechanics and statistics of active matter. Annu. Rev. Condens. Matter Phys, 1(1):323–345, 2010.
- [5] R. Mach and F. Schweitzer. Modeling vortex swarming in daphnia. Bull. Math. Biol., 69(2):539–562, 2007.
- [6] J. Vollmer, A. Vegh, C. Lange, and B. Eckhardt. Vortex formation by active agents as a model for daphnia swarming. *Phys. Rev. E*, 73:061924, 2006.
- [7] A. Cavagna, I. Giardina, and T. S. Grigera. The physics of flocking: Correlation as a compass from experiments to theory. *Phys. Rep.*, 728:1–62, 2018.
- [8] C. Reynolds. Flocks, herds, and schools: A distributed behavioral model. ACM SIGGRAPH Computer Graphics, 21:25–34, 1987.
- J. Toner and Y. Tu. Long-range order in a two-dimensional dynamical XY model: How birds fly together. Phys. Rev. Lett., 75:4326-4329, 1995.
- [10] J. Toner and Y. Tu. Flocks, herds, and schools: A quantitative theory of flocking. Phys. Rev. E, 58:4828–4858, 1998.
- [11] T. Vicsek, A. Czirók, E. Ben-Jacob, I. Cohen, and O. Shochet. Novel type of phase transition in a system of self-driven particles. *Phys. Rev. Lett.*, 75:1226–1229, 1995.
- [12] H. C. Berg. E. coli in Motion. Springer New York, 2004.
- [13] Yaouen Fily and M. Cristina Marchetti. Athermal phase separation of self-propelled particles with no alignment. Phys. Rev. Lett., 108:235702, Jun 2012.
- [14] Y. Huang, Z. Zhang, L. Yang, and Q. He. Typhoon eye-shaped global convective flow field-induced colloidal motor swarm. Colloids and Surfaces A: Physicochemical and Engineering Aspects, 689:133692, 2024.
- [15] M. Doi. Soft Matter Physics. OUP Oxford, 2013.
- [16] J. Tailleur and M. E. Cates. Statistical mechanics of interacting run-and-tumble bacteria. Phys. Rev. Lett., 100:218103, 2008.
- [17] C. Roberts and G. Pruessner. Exact solution of a boundary tumbling particle system in one dimension. Phys. Rev. Res., 4:033234, 2022.
- [18] Yael Avni, Shigeyuki Komura, and David Andelman. Brownian motion of a charged colloid in restricted confinement. Phys. Rev. E, 103:042607, Apr 2021.
- [19] Xie Hui, Sun Mengmeng, Fan Xinjian, Lin Zhihua, Chen Weinan, Wang Lei, Dong Lixin, and He Qiang. Reconfigurable magnetic microrobot swarm: Multimode transformation, locomotion, and manipulation. *Science Robotics*, 4(28):eaav8006, 2019.

- [20] Zhihua Lin, Tieyan Si, Zhiguang Wu, Changyong Gao, Xiankun Lin, and Qiang He. Light-activated active colloid ribbons. Angewandte Chemie International Edition, 56(43):13517–13520, 2017.
- [21] P.G. de Gennes. Brownian motion with dry friction. J Stat Phys, 119:953–962, 2005.
- [22] Hisao Hayakawa. Langevin equation with coulomb friction. Physica D: Nonlinear Phenomena, 205(1):48–56, 2005.
- [23] Hugo Touchette, Erik Van der Straeten, and Wolfram Just. Brownian motion with dry friction: Fokker-planck approach. Journal of Physics A: Mathematical and Theoretical, 43(44):445002, 2010.
- [24] P. Das, S. Puri, and M. Schwartz. Single particle brownian motion with solid friction. Eur. Phys. J. E, 2017.
- [25] M. Doi. Second quantization representation for classical many-particle system. J. Phys. A: Math. Gen, 9(9):1465–1477, 1976.
- [26] L. Peliti. Path integral approach to birth-death processes on a lattice. J. Phys.-Paris, 46:1469–1483, 1985.
- [27] J. Cardy. Reaction-diffusion processes. In Non-equilibrium Statistical Mechanics and Turbulence, pages 108–161. Cambridge University Press, Cambridge, UK, 2008.
- [28] M. E. Cates and J. Tailleur. When are active Brownian particles and run-and-tumble particles equivalent? Consequences for motility-induced phase separation. *Europhys. Lett.*, 101(2):20010, 2013.
- [29] C. Kurzthaler, S. Leitmann, and T. Franosch. Intermediate scattering function of an anisotropic active brownian particle. Sci Rep, 6:36702, 2016.
- [30] J. Stenhammar, D. Marenduzzo, R. J. Allen, and M. E. Cates. Phase behaviour of active Brownian particles: the role of dimensionality. Soft Matter, 10(10):1489–1499, 2014.
- [31] H. Risken. The Fokker-Planck Equation: Methods of Solutions and Applications, 2nd ed. Springer Verlag, Berlin, Germany, 1989.
- [32] G. A. Pavliotis. Stochastic Processes and Applications Diffusion Processes, the Fokker-Planck and Langevin Equations. Springer, Berlin, Germany, 2014.
- [33] M. Cates and J. Tailleur. Motility-induced phase separation. Annu. Rev. Condens. Matter Phys., 6(1):219–244, 2015.
- [34] Z. Zhang, L. Fehértói-Nagy, M. Polackova, and G. Pruessner. Field theory of active brownian particles in potentials. New Journal of Physics, 26(1):013040, 2024.
- [35] M. Abramowitz and I. A. Stegun, editors. Handbook of Mathematical Functions. Dover Publications, Inc., New York, NY, USA, 1970.
- [36] R. Garcia-Millan and G. Pruessner. Run-and-tumble motion in a harmonic potential: field theory and entropy production. J. Stat. Mech.: Theory Exp., 2021(6):063203, 2021.
- [37] Wolfram Research Inc. Mathematica. Wolfram Research, Inc., Champaign, IL, USA, 2014. Version 10.0.2.0.
- [38] R. Garcia-Millan. Interactions, correlations and collective behaviour in nonequilibrium systems. PhD thesis, Imperial College London, 2020.
- [39] G. Pruessner and R. Garcia-Millan. Field theories of active particle systems and their entropy production. preprint arXiv:2211.11906, 2022.
- [40] Z. Zhang and R. Garcia-Millan. Entropy production of nonreciprocal interactions. Phys. Rev. Res., 5:L022033, 2023.

Appendix A: Observables

In the following, we show the calculation of the observable $\langle \bullet \rangle$ in detail.

1. Purely diffusive case

We first show Eq. (38). By using Eq. (24), we immediately obtain Eq. (38b) since the bare propagator is the only contribution. For the observable in the LHS of Eq. (38a), we only consider the limited perturbation vertices listed in Eq. (26). Diagrammatically, it is

$$\left\langle \phi_2 \tilde{\phi}_0 \right\rangle \approx \frac{2}{2} \qquad 0 + \frac{2}{2} \qquad 2 \qquad 0 + \frac{2}{2} \qquad 2 \qquad 0 + \dots$$
 (A1a)
$$= \frac{2}{2} \qquad 0 \qquad \sum_{k=1}^{\infty} \left(\frac{2}{2} \qquad 2\right)^k$$
 (A1b)

$$= \frac{2}{2} \qquad 0 \qquad 1 \qquad (A1c)$$

$$1 - \frac{2}{1}$$

$$= \Omega \frac{\Lambda^{-1} \Omega}{2\gamma} \frac{1}{1 - \frac{\Lambda^{2,2} \Omega}{2\gamma}} .$$
(A1d)

We further include higher order terms up to the fourth order, and the above observable becomes

$$\left\langle \phi_{2}\tilde{\phi}_{0} \right\rangle \approx \frac{2}{2} \underbrace{0}_{k=0}^{\infty} \left(2 \underbrace{2}_{k=0}^{2} \right)^{k} \left(1 + \sum_{\ell=1}^{\infty} \left(2 \underbrace{4}_{\ell=1}^{2} \right)^{\ell} \sum_{j=0}^{\infty} \left(4 \underbrace{4}_{\ell=1}^{4} \right)^{j} \right)$$

$$+ \frac{2}{4} \underbrace{4}_{k=0}^{0} \left(2 \underbrace{2}_{k=0}^{2} \right)^{k} \sum_{\ell=0}^{\infty} \left(2 \underbrace{4}_{\ell=0}^{2} \right)^{\ell} \sum_{j=0}^{\infty} \left(4 \underbrace{4}_{\ell=0}^{4} \right)^{j}$$

$$= \Omega \frac{\Lambda^{2,0}\Omega}{2\gamma} \frac{1}{1 - \frac{\Lambda^{2,2}\Omega}{2\gamma}} \left(1 + \frac{\frac{\Lambda^{2,4}\Lambda^{4,2}\Omega^{2}}{8\gamma^{2}}}{1 - \frac{\Lambda^{2,4}\Lambda^{4,2}\Omega^{2}}{8\gamma^{2}}} \frac{1}{1 - \frac{\Lambda^{4,4}\Omega}{4\gamma}} \right) + \Omega \frac{\Lambda^{2,4}\Lambda^{4,0}\Omega^{2}}{8\gamma^{2}} \frac{1}{1 - \frac{\Lambda^{2,2}\Omega}{2\gamma}} \frac{1}{1 - \frac{\Lambda^{2,4}\Lambda^{4,2}\Omega^{2}}{8\gamma^{2}}} \frac{1}{1 - \frac{\Lambda^{4,4}\Omega}{4\gamma}} ,$$

$$(A2b)$$

where the first line of Eq. (A2a) indicates that the propagator starts with a vertex $\begin{pmatrix} 2 & 0 \\ 0 & 0 \end{pmatrix}$ from most left, and the first summation shows that there can be arbitrary number of the propagator $\begin{pmatrix} 2 & -2 & 2 \\ 0 & 0 \end{pmatrix}$ connecting to the left of the propagator. The terms in the bracket indicates that, if there is at least one $\begin{pmatrix} 2 & 4 & 2 \\ 0 & 0 \end{pmatrix}$ connected, the propagator $\begin{pmatrix} 4 & -4 \\ 0 & 0 \end{pmatrix}$ can also appear as many as possible. Similarly, in the second line of Eq. (A2a), the propagator $\begin{pmatrix} 2 & 4 & 0 \\ 0 & 0 \end{pmatrix}$ is the leading order, and there can be arbitrary number of the three types of the three types of the propagator $\begin{pmatrix} 2 & 4 & 0 \\ 0 & 0 \end{pmatrix}$ is the leading order, and there can be arbitrary number of the three types of the three types of the propagator $\begin{pmatrix} 2 & 4 & 0 \\ 0 & 0 \end{pmatrix}$ is the leading order, and there can be arbitrary number of the three types of the three types of the propagator $\begin{pmatrix} 2 & 4 & 0 \\ 0 & 0 \end{pmatrix}$ is the leading order, and there can be arbitrary number of the three types of the three types of the propagator $\begin{pmatrix} 2 & 4 & 0 \\ 0 & 0 \end{pmatrix}$ is the leading order, and there can be arbitrary number of the three types of the three types of the propagator $\begin{pmatrix} 2 & 4 & 0 \\ 0 & 0 \end{pmatrix}$ is the leading order.

propagators appearing in the summations. We perform the geometric sum and obtain Eq. (A2b).

Since the term $\Lambda\Omega/\gamma$ is proportional to the dimensionless parameters β and $\tilde{\beta}$ introduced in Eq. (39) and Eq. (52) for 1D and 2D space, respectively, we obtain the effective diffusion coefficient with the fourth order correction as

$$D_{\text{eff}} \approx \frac{D}{1+\beta} \left[1 - \frac{\beta}{1+\frac{\beta}{2}} \left(1 - \frac{\frac{\beta^2}{16}}{1+\frac{\beta^2}{16}} \frac{1}{1+\frac{3\beta}{8}} \right) - \frac{\beta^2}{24} \frac{1}{1+\frac{\beta}{2}} \frac{1}{1+\frac{\beta^2}{16}} \frac{1}{1+\frac{3\beta}{8}} \right], \text{ for 1D}$$
(A3a)

$$D_{\text{eff}}^{(2)} \approx \frac{D}{1+\tilde{\beta}} \left[1 - \frac{\tilde{\beta}}{1+\frac{\tilde{\beta}}{2}} \left(1 - \frac{\frac{\tilde{\beta}^2}{16}}{1+\frac{\tilde{\beta}^2}{16}} \frac{1}{1+\frac{3\tilde{\beta}}{8}} \right) - \frac{\tilde{\beta}^2}{24} \frac{1}{1+\frac{\tilde{\beta}}{2}} \frac{1}{1+\frac{\tilde{\beta}^2}{16}} \frac{1}{1+\frac{3\tilde{\beta}}{8}} \right], \text{ for } 2\text{D}.$$
(A3b)

2. ABP case

In the following, we obtain Eq. (69). Similar to the pure diffusive case, $\langle \phi_{0,0} \tilde{\phi}_{0,0} \rangle$ in Eq. (69a) is a bare propagator without any perturbative vertices. We write the LHS of Eq. (69b) diagrammatically

where the first line is exactly Eq. (A1), which is a combination of the bare propagator and the friction dependent perturbation only. The second and third lines includes the self-propulsion perturbation part, where the first term in Eq. (A4b) is the leading order. The rest terms show that arbitrary number of $\frac{1}{\pm 1}$ and $\frac{2}{0}$ $\frac{2}{0}$

can be inserted into the diagram in a suitable position. Therefore, we write Eq. (A4)

$$\left\langle \phi_{2,0}\tilde{\phi}_{0,0} \right\rangle \approx \frac{2}{1} \left(2 - \frac{2}{1} \right)^{k} + 2 \times \frac{2}{0} \left(2 - \frac{1}{1} \right)^{k} + 2 \times \frac{2}{0} \left(2 - \frac{1}{1} \right)^{k} = 0 \left(2 - \frac{2}{1} \right)^{k} \sum_{k=0}^{\infty} \left(2 - \frac{2}{0} \right)^{k} \sum_{\ell=0}^{\infty} \left(2 - \frac{1}{1} \right)^{\ell} + \dots$$

$$= \Omega \frac{\Lambda^{2,0}\Omega}{2\gamma} \frac{1}{1 - \frac{\Lambda^{2,2}\Omega}{2\gamma}} + 2\Omega \frac{\Upsilon^{2,1}\Omega_{1,0}^{2,1}\Omega_{1,0}^{2}\Omega}{2\gamma(\gamma + D_{r})} \frac{1}{1 - \frac{\Lambda^{2,2}\Omega}{2\gamma}} \frac{1}{1 - \frac{\Lambda^{1,1}\Omega}{\gamma + D_{r}}},$$
(A5a)
(A5b)

where the prefactor 2 is a symmetry factor that comes from ± 1 .

Similary, for the observable in Eq. (69c), we write it by the diagrams as

$$\left\langle \phi_{1,1}\tilde{\phi}_{0,0} \right\rangle \approx \frac{1}{1} \qquad \begin{array}{c} 0 \\ 1 \\ 0 \\ 0 \\ \end{array} + \frac{1}{1} \\ \begin{array}{c} 1 \\ 0 \\ 0 \\ \end{array} + \frac{1}{1} \\ \begin{array}{c} 1 \\ 0 \\ 0 \\ \end{array} + \frac{1}{1} \\ \begin{array}{c} 1 \\ 0 \\ 0 \\ \end{array} + \frac{1}{1} \\ \begin{array}{c} 1 \\ 1 \\ 0 \\ \end{array} + \frac{1}{1} \\ \begin{array}{c} 1 \\ 0 \\ 0 \\ \end{array} + \frac{1}{1} \\ \end{array} + \frac{1}{1} \\ \begin{array}{c} 1 \\ 0 \\ 0 \\ \end{array} + \frac{1}{1} \\ \begin{array}{c} 1 \\ 0 \\ 0 \\ \end{array} + \frac{1}{1} \\ \end{array} + \frac{1}{1} \\ \begin{array}{c} 1 \\ 0 \\ 0 \\ \end{array} + \frac{1}{1} \\ \end{array} + \frac{1}{1} \\ \end{array} + \frac{1}{1} \\ \begin{array}{c} 1 \\ 0 \\ 0 \\ \end{array} + \frac{1}{1} \\ = \frac{1}{1} \\ \end{array} + \frac{1}{1} \\ \end{array} + \frac{1}{1} \\ \end{array} + \frac{1}{1} \\ = \frac{1}{1} \\ = \frac{1}{1} \\ \end{array} + \frac{1}{1} \\ = \frac{1}{1}$$

$$=\Omega \frac{\Upsilon_{1,0}^{0,\Omega}\Omega}{\gamma + D_{\rm r}} \frac{1}{1 - \frac{\Lambda_{1,1}^{1,1}\Omega}{\gamma + D_{\rm r}}},\tag{A6c}$$

By plugging Eq. (61) and the dimensionless parameter Eq. (52) into Eqs. (A5b) and (A6c), we obtain Eqs. (69b) and (69c), respectively.

Appendix B: Numerical simulation

In this paper, we use the Euler-Maruyama method to simulate the system via the corresponding Langevin equation Eq. (3),

$$\mathbf{v}_{i+1} - \mathbf{v}_i = -\gamma \mathbf{v}_i \Delta t - \mu \frac{\mathbf{v}_i}{||\mathbf{v}_i||} \Delta t + \gamma \mathbf{w}_{\theta_i} \Delta t + \sqrt{2D\gamma^2 \Delta t} \boldsymbol{\xi}_i , \qquad (B1a)$$

$$\theta_{i+1} - \theta_i = \sqrt{2D_r \Delta t} \zeta_i , \qquad (B1b)$$

$$\mathbf{x}_{i+1} - \mathbf{x}_i = \mathbf{v}_i \Delta t , \qquad (B1c)$$

with a suitable initial condition. In the above, ξ_i is a 2D vector and the components are Gaussian random variables with zero expected value and unit variance. Similarly, ζ_i are also Gaussian random variables with zero expected value and unit variance. The effective diffusion coefficient is obtained from the mean-squared displacement

$$D_{\text{eff}}^{(2)} = \frac{1}{4NM\Delta t} \sum_{i=1}^{N} (\mathbf{x}_{M}^{(i)} - \mathbf{x}_{0}^{(i)})^{2} , \qquad (B2)$$

where M is the total steps and $M\Delta t = T$ is the total time, and N is the number of the particles to obtain the ensemble average.