

THE GÖDEL UNIVERSE AS A LIE GROUP WITH LEFT-INVARIANT LORENTZ METRIC AND THE IWASAWA DECOMPOSITION

V. N. BERESTOVSKII

ABSTRACT. We discuss models of the Gödel Universe as Lie groups with left-invariant Lorentz metric for two simply connected four-dimensional Lie groups, the Iwasawa decomposition for semisimple Lie groups, and left-invariant Lorentz metric on $SL(2, \mathbb{R})$, following K.-H. Neeb. Also we show that the isometry between two non-isomorphic sub-Riemannian Lie group, constructed by A. Agrachev and D. Barilari, is induced by some Iwasawa decomposition of $SL(2, \mathbb{R})$.

Keywords and phrases: Gödel Universe, Iwasawa decomposition, left-invariant Lorentz metric, left-invariant sub-Riemannian metric, Lie algebra, Lie group.

1. INTRODUCTION

Kurt Gödel in paper [1] of 1949 introduced the Lorentz metric (1) of the signature $(+, -, -, -)$ on the space \mathbb{R}^4 . The Gödel Universe (space-time) S is a solution of the General Relativity Theory (the Einstein gravitation equations).

In paper [2], the author found timelike and isotropic geodesics of the Gödel Universe, considered as the Lie group $G = (\mathbb{R}, +) \times A^+(\mathbb{R}) \times (\mathbb{R}, +)$ with left invariant Lorentz metric. Here $A^+(\mathbb{R})$ is connected Lie group of affine transformations on \mathbb{R} .

Professor Karl–Hermann Neeb wrote to the author that it is possible to realize the Gödel Universe otherwise and sent an electronic version of his joint with Joachim Hilgert book [3], where in section 2.7 "Gödel's cosmological model and universal covering of $SL(2, \mathbb{R})$ " is suggested a left-invariant Lorentz metric on this Lie group.

In this connection, it is useful to mention the paper [4] by A. Agrachev and D. Barilari, where the authors obtained a full classification of left-invariant sub-Riemannian metrics on three-dimensional Lie groups and "explicitly find a sub-Riemannian isometry between nonisomorphic Lie groups $SL(2, \mathbb{R})$ and $SO(2) \times A^+(\mathbb{R})$ " [4].

The existence of such isometry was indicated earlier in [5] by Falbel and Gorodski.

In a message to the author, Professor Neeb explains this by a diffeomorphism of Lie groups $SL(2, \mathbb{R})$ and $SO(2) \times A^+(\mathbb{R})$ by means of the Iwasawa decomposition for $SL(2, \mathbb{R})$. Let us cite now only Theorems 6.5.1 and 9.1.3 from [6].

In p. 10.6.4 (i) from [6] are indicated the following isomorphisms of Lie algebras:

$$\mathfrak{sl}(2, \mathbb{R}) \cong \mathfrak{su}(1, 1) \cong \mathfrak{so}(2, 1) \cong \mathfrak{sp}(1, \mathbb{R}).$$

Consequently, simply connected Lie groups with these Lie algebras are isomorphic.

The work was carried out within the framework of the State Contract to the IM SB RAS, project FWNF-2022-0006.

In Theorem 3 we prove some properties of special left-invariant Lorentz metrics on three-dimensional Lie groups. Also we show in Proposition 3 that the isometry between two non-isomorphic sub-Riemannian Lie group, constructed by A. Agrachev and D. Barilari, is induced by some Iwasawa decomposition of $SL(2, \mathbb{R})$.

The author expresses his gratitudes to Professor Neeb for fruitful discussions.

2. THE GÖDEL UNIVERSE AS A LIE GROUP WITH LEFT-INVARIANT LORENTZ METRIC

Gödel introduced in [1] his space-time S as \mathbb{R}^4 with the linear element

$$ds^2 = a^2 \left(dx_0^2 + 2e^{x_1} dx_0 dx_2 + \frac{e^{2x_1}}{2} dx_2^2 - dx_1^2 - dx_3^2 \right), \quad a > 0. \quad (1)$$

Gödel noticed in [1] that it is possible to rewrite this quadratic form in view of

$$ds^2 = a^2 \left[(dx_0 + e^{x_1} dx_2)^2 - dx_1^2 - \frac{e^{2x_1}}{2} dx_2^2 - dx_3^2 \right], \quad (2)$$

which shows obvious that its signature is equal everywhere to $(+, -, -, -)$.

We shall assume that $a = 1$.

Gödel's noticed in [1] that on (S, ds^2) acts simply transitively a four-dimensional isometry Lie group. It is easy to see that such action could be written as

$$x_0 = x'_0 + a, \quad x_1 = x'_1 + b, \quad x_2 = x'_2 e^{-b} + c, \quad x_3 = x'_3 + d \quad (3)$$

with arbitrary $a, b, c, d \in \mathbb{R}$. This implies that corresponding Lie group G is the simplest simply connected noncommutative four-dimensional Lie group of the view

$$G \cong [(\mathbb{R}, +) \times G_2] \times (\mathbb{R}, +), \quad (4)$$

where G_2 is unique up to isomorphism, necessary isomorphic to \mathbb{R}^2 , two-dimensional noncommutative Lie group. The Lie group G_2 is isomorphic to the Lie group $A^+(\mathbb{R})$ of preserving orientation affine transformations of the real direct line $(\mathbb{R}, +)$.

In case under consideration, identifying the quad (x'_0, x'_1, x'_2, x'_3) with the vector $(x'_2, x'_1, x'_0, x'_3, 1)^T$, where T is the sign of transposition, the action of the group G on \mathbb{R}^4 by formula (3) has the view $(x_2, x_1, x_0, x_3, 1)^T = A(x'_2, x'_1, x'_0, x'_3, 1)^T$, where

$$A = \begin{pmatrix} e^{-b} & 0 & 0 & 0 & c \\ 0 & 1 & 0 & 0 & b \\ 0 & 0 & 1 & 0 & a \\ 0 & 0 & 0 & 1 & d \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}. \quad (5)$$

Under this the equality

$$\begin{pmatrix} e^{-x_1} & 0 & 0 & 0 & x_2 \\ 0 & 1 & 0 & 0 & x_1 \\ 0 & 0 & 1 & 0 & x_0 \\ 0 & 0 & 0 & 1 & x_3 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix} (0, 0, 0, 0, 1)^T = (x_2, x_1, x_0, x_3, 1)^T \quad (6)$$

sets the bijection of the group G onto \mathbb{R}^4 and the unit of G corresponds to the zero-vector $(0, 0, 0, 0) \in \mathbb{R}^4$. On base of this, (4) and (1), we can identify (S, ds^2) with the Lie group G equipped with left-invariant Lorentz metric. Let

$$e_0 = \frac{\partial}{\partial x_0}(0), e_1 = \frac{\partial}{\partial x_1}(0), e_2 = \frac{\partial}{\partial x_2}(0), e_3 = \frac{\partial}{\partial x_3}(0)$$

be the basis of the Lie algebra \mathfrak{g} of the Lie group G at the unit of G , corresponding to coordinates (x_0, x_1, x_2, x_3) . Then, according to what has been said and (1), the components of the linear element ds^2 with respect to this basis are equal to

$$g_{00} = 1, g_{02} = g_{20} = 1, g_{22} = \frac{1}{2}, g_{11} = -1, g_{33} = -1, g_{ij} = g_{ji} = 0, i \neq j, j \neq 2. \quad (7)$$

According to (6), the Lie subgroup G_3 can be identified with the matrix Lie group

$$\left(\begin{array}{cccc} e^{-x_1} & 0 & 0 & x_2 \\ 0 & 1 & 0 & x_1 \\ 0 & 0 & 1 & x_0 \\ 0 & 0 & 0 & 1 \end{array} \right), \quad (x_0, x_1, x_2) \in \mathbb{R}^3. \quad (8)$$

It is obvious that $(S, ds^2) = (S_0, ds_0^2) \times (S_1, ds_1^2)$, where $S_0 = \mathbb{R}^3$, $S_1 = \mathbb{R}$,

$$ds_0^2 = dx_0^2 + 2e^{x_1} dx_0 dx_2 + \frac{e^{2x_1}}{2} dx_2^2 - dx_1^2, \quad ds_1^2 = -dx_3^2. \quad (9)$$

Also it is clear that we can consider (S_0, ds_0^2) as the matrix Lie group (8) with left-invariant Lorentz metric, which according to (7) has components

$$g_{00} = 1, g_{02} = g_{20} = 1, g_{22} = \frac{1}{2}, g_{11} = -1, g_{ij} = g_{ji} = 0, i \neq j, j \neq 2; \quad (10)$$

with respect to the basis e_0, e_1, e_2 of the Lie algebra \mathfrak{g}_3 of the matrix Lie group (8).

In consequence of (6), for the Lie algebra \mathfrak{g}_3 of the Lie group G_3 ,

$$e_0 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad e_1 = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}. \quad (11)$$

Then in the Lie algebra \mathfrak{g}_3 ,

$$[e_1, e_2] = e_1 e_2 - e_2 e_1 = -e_2, \quad [e_0, e_1] = [e_0, e_2] = 0. \quad (12)$$

3. THE IWASAWA DECOMPOSITIONS OF LIE ALGEBRAS AND LIE GROUPS

Let \mathfrak{g} be a semisimple real Lie algebra, σ be some Cartan involution of \mathfrak{g} , and $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ be the corresponding Cartan decomposition (\mathfrak{k} is the Lie subalgebra of \mathfrak{g} , consisting of fixed points relative to σ). Let us denote by \mathfrak{a} a maximal commutative subspace in \mathfrak{p} . Then there is the following *Iwasawa decomposition of Lie algebra \mathfrak{g}* .

Theorem 1. (4.7.2) in [7]. *Let \mathfrak{g} be a semisimple real Lie algebra.*

Then there exists a direct sum of vector subspaces in \mathfrak{g}

$$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{a} \oplus \mathfrak{n}, \quad (13)$$

where \mathfrak{n} is a nilpotent subalgebra in \mathfrak{g} such that the endomorphism $\text{ad } X$ is nilpotent for every $X \in \mathfrak{n}$, and $\mathfrak{a} \oplus \mathfrak{n}$ is a solvable subalgebra in \mathfrak{g} .

As an example, the authors of [7] give the decomposition (13) for $\mathfrak{g} = \mathfrak{sl}(n, \mathbb{R})$. In this case \mathfrak{k} is the Lie subalgebra of skew-symmetric matrices, \mathfrak{a} is the Lie subalgebra of diagonal matrices (with zero trace), and \mathfrak{n} is the Lie subalgebra of strictly upper triangular matrices. In particular, for $\mathfrak{g} = \mathfrak{sl}(2, \mathbb{R})$ we have

$$\mathfrak{k} = \left\{ \begin{pmatrix} 0 & t \\ -t & 0 \end{pmatrix} \right\}, \quad \mathfrak{a} = \left\{ \begin{pmatrix} t & 0 \\ 0 & -t \end{pmatrix} \right\}, \quad \mathfrak{n} = \left\{ \begin{pmatrix} 0 & t \\ 0 & 0 \end{pmatrix} \right\}, \quad t \in \mathbb{R}, \quad (14)$$

with natural basis

$$f_0 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad f_1 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad f_2 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \quad (15)$$

and Lie brackets for this basis

$$[f_0, f_1] = 2f_0 - 3f_2, \quad [f_0, f_2] = f_1, \quad [f_1, f_2] = 2f_2. \quad (16)$$

Let $K = \exp(\mathfrak{k})$, $A = \exp(\mathfrak{a})$, $N = \exp(\mathfrak{n})$ be Lie subgroups of the semisimple Lie group G , corresponding to the decomposition (13).

Theorem 2. (Theorem 9.1.3 in [6]) *Let G be a connected semisimple real Lie group. Then $G = KAN$ and the mapping*

$$(k, a, n) \rightarrow kan \quad (17)$$

is the diffeomorphism of manifold $K \times A \times N$ onto the Lie group G .

Corollary 1. *The Lie group G is diffeomorphic to Lie groups $K \times AN$ and $K \times A \times N$.*

Theorem 1 and Theorem 6.1.1 from [6] imply the following

Proposition 1. *The sets K , A , N , and AN are connected closed Lie subgroups of the Lie group G , where $\text{Ad}_G(K)$ is compact, A is commutative, N is nilpotent, and AN is solvable. The subgroup K contains the center Z of the Lie group G . In addition, K is compact if and only if the center Z of G is finite; in this case K is a maximal compact subgroup of the Lie group G .*

The statements above imply the following Corollary.

Corollary 2. *If $G = \text{SL}(n, \mathbb{R})$, then $K = \text{SO}(n)$, A is the group of all real diagonal $(n \times n)$ -matrices with unit determinant, N could be considered as the group of all real upper triangular $(n \times n)$ -matrices with units on the main diagonal, and $\text{Sol}(n) := AN$ as the group of all real upper triangular $(n \times n)$ -matrices with unit determinant.*

Corollary 3. *The Lie group $\text{SL}(n, \mathbb{R})$ is diffeomorphic to Lie groups $\text{SO}(n) \times AN$ and $\text{SO}(n) \times A \times N$. As a consequence, $\text{SL}(2, \mathbb{R})$ is diffeomorphic to the commutative Lie group $\text{SO}(2) \times A \times N$.*

4. APPLICATIONS OF IWASAWA DECOMPOSITIONS TO $\mathfrak{sl}(2, \mathbb{R})$ AND $\mathrm{SL}(2, \mathbb{R})$

This is a preparatory section.

Proposition 2. *There exist an isomorphism of the Lie group G_3 onto the Lie group $(\mathbb{R}, +) \times \mathrm{Sol}(2)$ and corresponding realization of (S_0, ds_0^2) as Lie group $(\mathbb{R}, +) \times \mathrm{Sol}(2)$ with left-invariant Lorentz metric.*

Proof. Comparing (12) and (16), we see that the linear map $\varphi : \mathfrak{a} \oplus \mathfrak{n} \rightarrow \mathfrak{g}$ such that

$$\varphi\left(\frac{-f_1}{2}\right) = e_1, \quad \varphi(f_2) = e_2 \quad (18)$$

is an isomorphism of Lie algebras. Then (18) defines isomorphism of Lie groups $\psi : \mathrm{Sol}(2) \rightarrow G_2$:

$$\psi\left(\left(\begin{array}{cc} e^{-s/2} & 0 \\ 0 & e^{s/2} \end{array}\right)\left(\begin{array}{cc} 1 & r \\ 0 & 1 \end{array}\right)\right) = \left(\begin{array}{cccc} e^{-s} & 0 & 0 & e^{-s}r \\ 0 & 1 & 0 & s \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{array}\right), \quad (19)$$

$$\psi\left(\left(\begin{array}{cc} 1 & r \\ 0 & 1 \end{array}\right)\left(\begin{array}{cc} e^{-s/2} & 0 \\ 0 & e^{s/2} \end{array}\right)\right) = \left(\begin{array}{cccc} e^{-s} & 0 & 0 & r \\ 0 & 1 & 0 & s \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{array}\right). \quad (20)$$

We can consider $(\mathbb{R}, +)$ as a Lie algebra and as a Lie group. Then the mappings

$$t \in (\mathbb{R}, +) \rightarrow \left(\begin{array}{cc} 0 & t \\ -t & 0 \end{array}\right), \quad t \in (\mathbb{R}, +) \rightarrow \left(\begin{array}{cc} \cos t & \sin t \\ -\sin t & \cos t \end{array}\right) \quad (21)$$

are correspondingly the isomorphism of Lie algebras and respective universal covering epimorphism of Lie groups. Then there exists unique isomorphism ψ of the Lie group $(\mathbb{R}, +) \times \mathrm{Sol}(2)$ onto the Lie group G_3 , with properties (19), (20), and $\psi(t) = t$ for $t \in (\mathbb{R}, +)$. It follows from previous considerations that we shall realize (S_0, ds_0^2) as the Lie group $(\mathbb{R}, +) \times \mathrm{Sol}(2)$ with left-invariant Lorentz metric if components of this metric in the basis $\{f_0, -f_1/2, f_2\}$ of its Lie algebra will be as in (10).

The corresponding orthonormal basis is

$$X = f_0, \quad Y = -f_1/2, \quad Z = \sqrt{2}(f_0 - f_2). \quad (22)$$

□

Now we shall consider the Iwasawa decompositions for Lie groups $\mathrm{SO}(2) \times \mathrm{Sol}(2)$ and $\mathrm{SL}(2, \mathbb{R})$ and corresponding mutual expressions of partial derivatives of these Lie groups by parameters r, s, t and natural left-invariant vector fields on the above Lie groups which represent some interest.

Let $kan = kan(t, s, r)$ denotes the following matrix from $\mathrm{SL}(2, \mathbb{R})$:

$$\left(\begin{array}{cc} \cos t & \sin t \\ -\sin t & \cos t \end{array}\right)\left(\begin{array}{cc} e^{-s/2} & 0 \\ 0 & e^{s/2} \end{array}\right)\left(\begin{array}{cc} 1 & r \\ 0 & 1 \end{array}\right) =$$

$$\begin{pmatrix} \cos te^{-s/2} & r \cos te^{-s/2} + \sin te^{s/2} \\ -\sin te^{-s/2} & -r \sin te^{-s/2} + \cos te^{s/2} \end{pmatrix}.$$

Then we have the following partial derivatives

$$(kan)'_t = \begin{pmatrix} -\sin te^{-s/2} & -r \sin te^{-s/2} + \cos te^{s/2} \\ -\cos te^{-s/2} & -r \cos te^{-s/2} - \sin te^{s/2} \end{pmatrix}, \quad (23)$$

$$(kan)'_s = \frac{-1}{2} \begin{pmatrix} \cos te^{-s/2} & r \cos te^{-s/2} - \sin te^{s/2} \\ -\sin te^{-s/2} & -r \sin te^{-s/2} - \cos te^{s/2} \end{pmatrix}, \quad (24)$$

$$(kan)'_r = \begin{pmatrix} 0 & \cos te^{-s/2} \\ 0 & -\sin te^{-s/2} \end{pmatrix}. \quad (25)$$

The left-invariant vector fields on $\mathrm{SL}(2, \mathbb{R})$, corresponding to f_0 , $-f_1/2$, and f_2 , are

$$v_0 = (kan) \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = \begin{pmatrix} -(r \cos te^{-s/2} + \sin te^{s/2}) & \cos te^{-s/2} \\ -(-r \sin te^{-s/2} + \cos te^{s/2}) & -\sin te^{-s/2} \end{pmatrix}, \quad (26)$$

$$v_1 = \frac{1}{2}(kan) \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} = \frac{1}{2} \begin{pmatrix} -\cos te^{-s/2} & r \cos te^{-s/2} + \sin te^{s/2} \\ \sin te^{-s/2} & -r \sin te^{-s/2} + \cos te^{s/2} \end{pmatrix}, \quad (27)$$

$$v_2 = (kan) \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & \cos te^{-s/2} \\ 0 & -\sin te^{-s/2} \end{pmatrix} = (kan)'_r. \quad (28)$$

Comparing (23), (24), (25) and (26), (27), (28), we see that

$$(kan)'_s = v_1 - rv_2, \quad (kan)'_t = e^{-s}(v_0 - 2rv_1 + (e^{2s} + r^2 - 1)v_2). \quad (29)$$

The equalities imply that

$$v_2 = (kan)'_r, v_1 = (kan)'_s + r(kan)'_r, v_0 = e^s(kan)'_t + 2r(kan)'_s + (r^2 + 1 - e^{2s})(kan)'_r. \quad (30)$$

Let us consider now the Lie group $K \times AN = \mathrm{SO}(2) \times \mathrm{Sol}(2)$ with the Lie algebra $\mathfrak{k} \oplus (\mathfrak{a} + \mathfrak{n})$ and the same basis $(f_0, -f_1/2, f_2)$. Then

$$k(t) = \begin{pmatrix} \cos t & \sin t \\ -\sin t & \cos t \end{pmatrix}, \quad an(s, r) = \begin{pmatrix} e^{-s/2} & 0 \\ 0 & e^{s/2} \end{pmatrix} \begin{pmatrix} 1 & r \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} e^{-s/2} & re^{-s/2} \\ 0 & e^{s/2} \end{pmatrix},$$

$$k'(t) = \begin{pmatrix} -\sin t & \cos t \\ -\cos t & -\sin t \end{pmatrix}, (an)'_s = \frac{-1}{2} \begin{pmatrix} e^{-s/2} & re^{-s/2} \\ 0 & -e^{s/2} \end{pmatrix}, (an)'_r = \begin{pmatrix} 0 & e^{-s/2} \\ 0 & 0 \end{pmatrix},$$

$$v_0 = \begin{pmatrix} \cos t & \sin t \\ -\sin t & \cos t \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = \begin{pmatrix} -\sin t & \cos t \\ -\cos t & -\sin t \end{pmatrix} = k'(t),$$

$$v_1 = \frac{1}{2} \begin{pmatrix} e^{-s/2} & re^{-s/2} \\ 0 & e^{s/2} \end{pmatrix} \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} = \frac{1}{2} \begin{pmatrix} -e^{-s/2} & re^{-s/2} \\ 0 & e^{s/2} \end{pmatrix},$$

$$v_2 = \begin{pmatrix} e^{-s/2} & re^{-s/2} \\ 0 & e^{s/2} \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & e^{-s/2} \\ 0 & 0 \end{pmatrix} = (an)'_r, \quad (an)'_s = v_1 - rv_2,$$

$$k'(t) = v_0, \quad (an)'_s = v_1 - rv_2, \quad (an)'_r = v_2, \quad (31)$$

$$v_0 = k'(t), \quad v_1 = (an)'_s + r(an)'_r, \quad v_2 = (an)'_r. \quad (32)$$

5. LEFT-INVARIANT LORENTZ METRICS ON $\text{SO}(2) \times \text{Sol}(2)$ AND $\text{SL}(2, \mathbb{R})$

Theorem 3. *the Lorentz metric on $\widetilde{\text{SL}(2, \mathbb{R})}$ from [3] is not isometric to the subspace (S_0, ds_0^2) of the Gödel Universe and the Iwasawa diffeomorphism of $\text{SL}(2, \mathbb{R})$ onto $\text{SO}(2) \times \text{Sol}(2)$ is not isometry for Lorentz metric on $\text{SL}(2, \mathbb{R})$ from [3].*

Proof. For any (pseudo)-Riemannian manifold M with (pseudo)-metric tensor (\cdot, \cdot) , the Levi-Civita connection ∇ , and smooth vector fields X, Y, Z is valid the following equation (3.5.(7)) from [8]:

$$(\nabla_X Y, Z) = \frac{1}{2}[X(Y, Z) + Y(Z, X) - Z(X, Y) + (Z, [X, Y]) + (Y, [Z, X]) - (X, [Y, Z])]. \quad (33)$$

As a consequence, if $(M, (\cdot, \cdot))$ is a Lie group G with left-invariant (pseudo)-metric (\cdot, \cdot) and X, Y, Z are left-invariant, then

$$(\nabla_X Y, Z) = \frac{1}{2}[(Z, [X, Y]) + (Y, [Z, X]) - (X, [Y, Z])]. \quad (34)$$

Let assume now that $G = \text{SO}(2) \times \text{Sol}(2)$ with left-invariant Lorentz metric and (X, Y, Z) be an orthonormal basis of left-invariant vector fields such that X be tangent to $\text{SO}(2)$ and $(X, X) = 1$. Then $Y = \alpha X + Y_1$, $Z = \beta X + Z_1$, where $Y_1, Z_1 \in \mathfrak{sol}(2)$ and

$$(\nabla_X Y, Z) = \frac{-1}{2}(X, [Y, Z]) = \frac{-1}{2}(X, W_1) := -\gamma_1, \quad W_1 \in \mathfrak{n}, \quad (35)$$

$$(\nabla_X Y, X) = 0, \quad (\nabla_X Y, Y) = 0, \quad \nabla_X Y = \gamma_1 Z.$$

We want to calculate

$$R(X, Y)Y = \nabla_X \nabla_Y Y - \nabla_Y \nabla_X Y - \nabla_{[X, Y]} Y = \nabla_X \nabla_Y Y - \nabla_Y \nabla_X Y.$$

It follows from (34) that

$$(\nabla_Y Y, X) = (\nabla_Y Y, Y) = 0, \quad (\nabla_Y Y, Z) = -(Y, W_1) := -\gamma_2, \quad \nabla_Y Y = \gamma_2 Z,$$

$$(\nabla_X Z, X) = (\nabla_X Z, Z) = 0, \quad (\nabla_X Z, Y) = \frac{-1}{2}(X, [Z, Y]) = \gamma_1, \quad \nabla_X Z = -\gamma_1 Y,$$

$$(\nabla_Y Z, X) = \gamma_1, \quad (\nabla_Y Z, Y) = \gamma_2, \quad (\nabla_Y Z, Z) = 0, \quad \nabla_Y Z = \gamma_1 X - \gamma_2 Y,$$

$$R(X, Y)Y = \nabla_X(\gamma_2 Z) - \nabla_Y(\gamma_1 Z) = -\gamma_1 \gamma_2 Y - \gamma_1^2 X + \gamma_1 \gamma_2 Y = -\gamma_1^2 X.$$

Let us calculate yet an analogue of the sectional curvature $(R(X, Y)Y, X) = -\gamma_1^2$.

In [3], the authors consider the Lie group $\text{SL}(2, \mathbb{R})$ with left-invariant Lorentz metric and orthonormal basis of opposite to above signature of the form

$$X = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, Y = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, Z = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \in \mathfrak{sl}(2, \mathbb{R}). \quad (36)$$

Then $[Y, Z] = 2\sqrt{2}X$ in $\mathfrak{sl}(2, \mathbb{R})$, but $[Y, Z] = 0$ in the Lie algebra $\mathfrak{so}(2) \oplus \mathfrak{sol}(2)$, and according to the previous calculations, $\gamma_1 = 0$ in $\text{SO}(2) \times \text{Sol}(2)$ and

$$R(X, Y)Y = 0. \quad (37)$$

Now let us calculate $R(X, Y)Y$ and $(R(X, Y)Y, X)$ for left-invariant Lorentz metric on the Lie group $\text{SO}(2) \times \text{Sol}(2)$, locally isometric to the three-dimensional subspace (S_0, ds_0^2) of the Gödel Universe. According to (43) and (10), we have

$$W_1 = [Y, Z] = \sqrt{2}[-f_1/2, (f_0 - f_2)] = -\sqrt{2}f_2, \quad \gamma_1 = -\frac{1}{2}(X, W_1) = \sqrt{2}.$$

So, according to previous calculations,

$$R(X, Y)Y = -2X, \quad (R(X, Y)Y, X) = -2. \quad (38)$$

Let us compute now $R(X, Y)Y$ for left-invariant Lorentz metric on $\text{SL}(2, \mathbb{R})$, defined in [3]. It follows from (36) that

$$[X, Y] = \sqrt{2}Z, \quad [X, Z] = \sqrt{2}Y, \quad [Y, Z] = 2\sqrt{2}X. \quad (39)$$

Let us apply (34) and (39) in the further computations.

$$(\nabla_X Y, Z) = -\sqrt{2}, \quad (\nabla_X Y, X) = (\nabla_X Y, Y) = 0, \quad \nabla_X Y = \sqrt{2}Z,$$

$$(\nabla_Y Y, X) = (\nabla_Y Y, Y) = (\nabla_Y Y, Z) = 0, \quad \nabla_Y Y = 0,$$

$$(\nabla_Z Y, X) = -2\sqrt{2}, \quad (\nabla_Z Y, Y) = (\nabla_Z Y, Z) = 0, \quad \nabla_Z Y = -2\sqrt{2}X,$$

$$\nabla_Y Z = \nabla_Z Y + [Y, Z] = 0,$$

$$R(X, Y)Y = \nabla_X \nabla_Y Y - \nabla_Y \nabla_X Y - \nabla_{[X, Y]} Y = -\sqrt{2}(\nabla_Y Z - \nabla_Z Y) = -4X. \quad (40)$$

All equalities (37), (38), and (40) are pairwise different.

This implies all statements of Theorem 3. \square

6. ISOMETRY OF NON-ISOMORPHIC SUB-RIEMANNIAN LIE GROUPS

Let us change notation $x_1 \leftrightarrow x_2$, $x_0 \rightarrow x_3$. Then the mapping

$$\begin{pmatrix} e^{-x_2} & 0 & 0 & x_1 \\ 0 & 1 & 0 & x_2 \\ 0 & 0 & 1 & x_3 \\ 0 & 0 & 0 & 1 \end{pmatrix} \rightarrow \begin{pmatrix} e^{-x_2} & 0 & x_1 \\ 0 & 1 & x_3 \\ 0 & 0 & 1 \end{pmatrix} \in A^+(\mathbb{R}) \times (\mathbb{R}, +) \quad (41)$$

is an isomorphism of matrix Lie groups with the basis of the Lie algebra

$$e_1 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad e_3 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix},$$

such that only $[e_1, e_2] = -[e_2, e_1] = e_1$ are unique nonzero Lie brackets.

Analogously to (19) and (20), the mapping

$$F \left(\begin{pmatrix} e^{-x_2} & 0 & x_1 \\ 0 & 1 & x_3 \\ 0 & 0 & 1 \end{pmatrix} \right) = \left(\begin{pmatrix} e^{-x_2/2} & x_1 \\ 0 & e^{x_2/2} \end{pmatrix}, \begin{pmatrix} \cos x_3 & \sin x_3 \\ -\sin x_3 & \cos x_3 \end{pmatrix} \right) \quad (42)$$

is the universal covering epimorphism of $A^+(\mathbb{R}) \times (\mathbb{R}, +)$ onto $\text{Sol}(2) \times \text{SO}(2)$.

The standard left-invariant sub-Riemannian structure on $A^+(\mathbb{R}) \times (\mathbb{R}, +)$ is defined in [4] by the orthonormal frame $\Delta = \text{span}\{e_2, e_1 + e_3\}$. Then there is unique sub-Riemannian structure on $\text{Sol}(2) \times \text{SO}(2)$ such that F is a local isometry; it is defined by the orthonormal frame $\overline{\Delta} = \text{span}\{\overline{e}_2, \overline{e}_1 + \overline{e}_3\}$, where

$$\overline{e}_1 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad \overline{e}_2 = \begin{pmatrix} -1/2 & 0 \\ 0 & 1/2 \end{pmatrix}, \quad \overline{e}_3 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \quad (43)$$

Now we follow [4]. Let $a = e^{-x_2}$ and $b = x_1$ in the second matrix of (41).

The subgroup $A^+(\mathbb{R})$ is diffeomorphic to the half-plane $\{(a, b) \in \mathbb{R}^2, a > 0\}$, which is described in the standard polar coordinates as $\{(\rho, \theta) | \rho > 0, -\pi/2 < \theta < \pi/2\}$.

Theorem 4. [4]. *The diffeomorphism $\Psi : A^+(\mathbb{R}) \times S^1 \rightarrow \text{SL}(2, \mathbb{R})$ defined by*

$$\Psi(\rho, \theta, \varphi) = \frac{1}{\sqrt{\rho \cos \theta}} \begin{pmatrix} \cos \varphi & \sin \varphi \\ \rho \sin(\theta - \varphi) & \rho \cos(\theta - \varphi) \end{pmatrix}, \quad (44)$$

where $(\rho, \theta) \in A^+(\mathbb{R})$ and $\varphi \in S^1$, is a global sub-Riemannian isometry.

Remark 1. *Using the above locally isometric covering F , we can and will understand Ψ as the global isometry between $\text{Sol}(2) \times \text{SO}(2)$ and $\text{SL}(2, \mathbb{R})$ supplied with sub-Riemannian metrics defined by the same frame $\overline{\Delta}$.*

Corollary 4. *$A^+(\mathbb{R}) \times (\mathbb{R}, +)$ with sub-Riemannian metric, defined by the frame Δ , is isometric to the universal covering $\widetilde{\text{SL}(2, \mathbb{R})}$ of $\text{SL}(2, \mathbb{R})$ with sub-Riemannian metric such that the natural universal covering epimorphism of $\widetilde{\text{SL}(2, \mathbb{R})}$ onto $\text{SL}(2, \mathbb{R})$ with sub-Riemannian metric, defined by the frame $\overline{\Delta}$, is a local isometry.*

Proposition 3. *The global isometry Ψ in the sense of Remark 1 is the Iwasawa diffeomorphism of $\text{Sol}(2) \times \text{SO}(2)$ onto $\text{SL}(2, \mathbb{R})$ of the view $(n\overline{a}, k) \in NA \times \text{SO}(2) \rightarrow n\overline{a}k \in NAK = \text{SL}(2, \mathbb{R})$, where*

$$n = \begin{pmatrix} 1 & 0 \\ b & 1 \end{pmatrix}, \quad \overline{a} = \begin{pmatrix} a^{-1/2} & 0 \\ 0 & a^{1/2} \end{pmatrix}, \quad k = \begin{pmatrix} \cos \varphi & \sin \varphi \\ -\sin \varphi & \cos \varphi \end{pmatrix}, \quad a = \rho \cos \theta, \quad b = \rho \sin \theta.$$

Proof. One needs simply to check that $n\overline{a}k$ is equal to the matrix in (44). \square

Remark 2. *Notice that $n = \exp(t\tilde{e}_1)$, $\overline{a} = \exp(s\overline{e}_2)$, where $\tilde{e}_1 = (\overline{e}_1)^T$, T is the sign of transposition, $b = t$, and $a^{1/2} = e^{s/2}$. Also $[\tilde{e}_1, \overline{e}_2] = -\tilde{e}_1$.*

REFERENCES

- [1] Gödel K. An example of a new type of cosmological solutions of Einstein's field equations of gravitation. *Rev. Mod. Phys.*, 21:3 (1949), 447–450.
- [2] Berestovskii V.N. The Gödel Universe as the Lie group with left-invariant Lorentz metric. Preprint arXiv: 2404.06866v1 [math.DG] 10 Apr. 2024. 16 p.
- [3] Hilgert Jo., Neeb K.-H. Lie semigroups and their Applications. *Lect. Notes Math.*, 1552. Springer-Verlag: Berlin, Heidelberg, 1993.
- [4] Agrachev A., Barilari D. Sub-Riemannian structures on 3d Lie groups. *Journal of Dynamical and Control Systems*, 18:1 (2012), 21–44. DOI:10.1007/s10883-012-9133-8
- [5] Falbel E., Gorodski C. Sub-Riemannian homogeneous spaces in dimensions 3 and 4. *Geom.Dedicata* 62:3(1996), 227–252.

- [6] *Helgason S.* Differential Geometry, Lie groups and Symmetric Spaces. Graduate Studies in Mathematics, Vol. 34. Providence, R.I.: AMS, 2001.
- [7] *Goto M., Grosshans F.D.* Semisimple Lie algebras. Lecture Notes in Pure and Applied Mathematics, Vol. 38. Marcel Dekker, Inc. New York and Basel, 1978.
- [8] *Gromoll D., Klingenberg W., Meyer W.* Riemannsche Geometrie im Grossen. Lect. Notes Math., 55. Springer-Verlag, Heidelberg, 1968.

SOBOLEV INSTITUTE OF MATHEMATICS OF THE SB RAS,
4 ACAD. KOPTYUG AVE., NOVOSIBIRSK 630090, RUSSIA
Email address: `vberestov@inbox.ru`