On the Diameter of Undirected Cayley Graphs of Finite Abelian Groups

Béla Bajnok* W. Kyle Beatty†

April 1, 2024

Abstract

Let s be a positive integer. Our goal is to find all finite abelian groups G that contain a 2-subset A for which the undirected Cayley graph $\Gamma(G,A)$ has diameter at most s. We provide a complete answer when G is cyclic, and a conjecture and some partial answers when G is noncyclic.

2020 AMS Mathematics Subject Classification: Primary: 11B13; Secondary: 05B10, 05C35, 11P70, 11B75, 20K01. Key words: Abelian group, Cayley graph, diameter, sumset, signed sumset.

1 Introduction

1 Introduction

3 Introduction

3 Introduction

3 Given a finite abelian group G and a subset $A \subseteq G$, the undirected Cayley graph $\Gamma(G,A)$ has its vertex set formed by the elements of G, and has an edge between two vertices whenever their difference is an element of A, that is, $g_1 \in G$ and $g_2 \in G$ are connected when $g_2 = g_1 + a$ or $g_1 = g_2 + a$ for some $a \in A$. If A is a generating set of G, then $\Gamma(G,A)$ is connected; in particular, if A is an s-spanning set of G, meaning that every element of G can be written as the signed sum of at most s (not-necessarily distinct) elements of A, then $\Gamma(G,A)$ has diameter at most s.

Suppose now that A is an m-subset of G. It is easy to compute that there are at most $\sum_{i \ge 0} \binom{n}{i} \binom{n-1}{i-1} 2^i$ vertices of $\Gamma(G,A)$ that are of distance h away from any given vertex, and thus if A is an s-spanning set in G, then the order of G can be at most

be at most

$$\sum_{h=0}^{s} \sum_{i>0} {m \choose i} {h-1 \choose i-1} 2^{i} = \sum_{i>0} {m \choose i} {s \choose i} 2^{i}.$$

Therefore, we have the following bound.

Proposition 1.1 For positive integers m and s, define

$$a(m,s) = \sum_{i \ge 0} {m \choose i} {s \choose i} 2^i. \tag{1}$$

If an abelian group G has an m-subset A for which the diameter of $\Gamma(G,A)$ is s, then the order of G is at most a(m,s).

^{*}Department of Mathematics, Gettysburg College, U.S.A. Email: bbajnok@gettysburg.edu.

[†]Department of Applied Mathematics and Statistics, Johns Hopkins University, U.S.A. Email: wbeatty1@jhu.edu.

We note, in passing, that the values of a(m,s) are known as *Delannoy numbers* (see [3]), and that they satisfy the recursion

$$a(m,s) = a(m,s-1) + a(m-1,s) + a(m-1,s-1).$$

It is a natural question to ask for the largest possible order of a group G with an m-subset A for which the diameter of $\Gamma(G, A)$ is at most s. Before we discuss our results on this question, let us review the analogous question in the directed case.

Given a finite abelian group G and a subset $A \subseteq G$, the directed Cayley graph $\vec{\Gamma}(G, A)$ has its vertex set formed by the elements of G, and has a directed edge from $g_1 \in G$ to $g_2 \in G$ when $g_2 = g_1 + a$ for some $a \in A$. We say that A is an s-basis of G if every element of G can be written as the sum of at most s (not-necessarily distinct) elements of G; in this case, the diameter of $\vec{\Gamma}(G, A)$ is at most s.

For an *m*-subset *A* of *G*, we see that there are at most $\binom{m+h-1}{h}$ vertices of $\vec{\Gamma}(G,A)$ that are of distance *h* away from any given vertex, so if *A* is an *s*-basis of *G*, then we must have

$$|G| \le \sum_{h=0}^{s} {m+h-1 \choose h} = {m+s \choose s}.$$

When equality holds, we say that A is a *perfect s-basis* of G: in this case, every element of G arises as the sum of at most s (not-necessarily distinct) elements of A in a unique way. Trivially, the set of nonzero elements is a perfect 1-basis in any group G, and the 1-element set consisting of a generator of G is a perfect s-basis in the cyclic group of order s+1. It turns out that there are no others:

Theorem 1.2 (Bajnok, Berson, and Just; cf. [2]) If a subset A of a finite abelian group G is a perfect s-basis, then s = 1 and $A = G \setminus \{0\}$, or $G \cong \mathbb{Z}_{s+1}$ and |A| = 1.

Given the scarcity of perfect bases, it is natural to ask for the largest groups with a given diameter and size of generating set. The paper [9] asks the following questions:

- What is the largest possible order of an abelian group G with an m-subset A for which the diameter of $\vec{\Gamma}(G,A)$ is at most s?
- What is the largest possible order of a cyclic group G with an m-subset A for which the diameter of $\vec{\Gamma}(G,A)$ is at most

After a series of papers [10, 5, 8, 9, 4], these two questions have been completely solved for the case of m = 2. We have the following two results.

Theorem 1.3 (Morillo, Fiol, and Fàbrega; cf. [10]) Let s be a positive integer. The largest possible order of a cyclic group G with a 2-subset for which the diameter of $\vec{\Gamma}(G, A)$ is at most s is given by $\lfloor (s^2 + 4s + 3)/3 \rfloor$.

Theorem 1.4 (Fiol; cf. [4]) Let s be any positive integer. The largest possible order of an abelian group G with a 2-subset for which the diameter of $\vec{\Gamma}(G, A)$ is at most s is given by $\lfloor (s^2 + 4s + 4)/3 \rfloor$.

Observe that the quantities in Theorems 1.3 and 1.4 agree, unless $s \equiv 1 \mod 3$; for that case one can verify that the group $G = \mathbb{Z}_k \times \mathbb{Z}_{3k}$ has the s-basis $\{(0,1), (1,3k-1)\}$, where k = (s+2)/3. However, the question specifically for noncyclic groups is still open when $s \not\equiv 1 \mod 3$.

Let us now return to the undirected case. When equality occurs in Proposition 1.1, then A is called a *perfect s-spanning* set in G. Perfect spanning sets can be easily characterized for m = 1 and s = 1. Note that a(1, s) = 2s + 1 and, clearly, for

any $a \in \mathbb{Z}_{2s+1}$ with gcd(a, 2s+1) = 1, the set $\{a\}$ is a perfect s-spanning set in \mathbb{Z}_{2s+1} . Also, a(m, 1) = 2m + 1, and for any group G of order 2m + 1, every asymmetric subset of size m (that is, a set containing exactly one of any nonzero element or its inverse) is a perfect 1-spanning set in G.

From (1) we get $a(2, s) = 2s^2 + 2s + 1$, and below we establish the following result.

Theorem 1.5 Let s be a positive integer. The largest possible order of a cyclic group G with a 2-subset for which the diameter of $\Gamma(G,A)$ is at most s is given by $2s^2 + 2s + 1$.

For noncyclic groups the situation seems a bit more nuanced. We have the following conjecture.

Conjecture 1.6 Let $s \ge 3$ be an integer.

- If 2s + 1 is prime, then the largest noncyclic group with an s-spanning set of size 2 has order $2s^2$.
- If 2s+1 is composite and p is its smallest prime divisor, then the largest noncyclic group with an s-spanning set of size 2 has order $2s^2 + 2s (p^2 1)/2$.

In particular, for any noncyclic group G with an s-spanning set of size 2 we have $|G| \le 2s^2 + 2s - 4$.

(We add that the largest noncyclic group with a 2-spanning set of size 2 is $\mathbb{Z}_3 \times \mathbb{Z}_3$.)

Below we prove that the quantities in Conjecture 1.6 are lower bounds.

Theorem 1.7 Let s be a positive integer.

- The largest noncyclic group with an s-spanning set of size 2 has order at least $2s^2$.
- If 2s + 1 is composite and p is any prime divisor of 2s + 1, then the largest noncyclic group with an s-spanning set of size 2 has order at least $2s^2 + 2s (p^2 1)/2$.

Establishing upper bounds appears to be more difficult, and we only have the partial results that we summarize here. We first note that a noncyclic group that has a spanning set of size 2 must have rank 2 and thus is isomorphic to $\mathbb{Z}_c \times \mathbb{Z}_{kc}$ for some integers $c \geq 2$ and $k \geq 1$.

We say that the group $\mathbb{Z}_c \times \mathbb{Z}_{kc}$ is s-regular for some positive integer s, if it possesses an s-spanning set of the form $\{(1,u),(1,v)\}$ for some $u,v \in \mathbb{Z}_{kc}$. While we have no precise count on the number of s-regular groups, it appears that most groups of rank 2 that have an s-spanning set of size 2 are s-regular. For example, for s = 10, of the 105 groups of rank 2 that have an s-spanning set of size 2, 103 are s-regular. (The two exceptions are $\mathbb{Z}_8 \times \mathbb{Z}_{16}$ and $\mathbb{Z}_{10} \times \mathbb{Z}_{20}$.) In Theorem 3.1 below we prove for every odd prime c and positive integer k that if $\mathbb{Z}_c \times \mathbb{Z}_{kc}$ has an s-spanning set at all, then it is s-regular.

Our results for upper bounds can be summarized as follows.

Theorem 1.8 Let s be a positive integer, and let $G = \mathbb{Z}_c \times \mathbb{Z}_{kc}$ be an s-regular group for some integers $c \geq 2$ and $k \geq 1$.

- If c=3 and s+1 is divisible by 3, then $|G| \leq 2s^2 + 1$.
- If c is odd and s is divisible by c, then $|G| \le 2s^2 + s$, and if 2s + 1 is divisible by c, then $|G| \le 2s^2 + 2s (c^2 1)/2$.
- In all other cases, $|G| \leq 2s^2$.

As a consequence of our results, we can prove that for $s \geq 3$, every s-regular abelian group of rank 2 has order at most $2s^2 + 2s - 4$, with equality if, and only if, it is isomorphic to $\mathbb{Z}_3 \times \mathbb{Z}_{(2s^2 + 2s - 4)/3}$ for some $s \equiv 1 \mod 3$ (See Theorem 3.3 below).

For c=2 we have exact results, even without the assumption that the group is s-regular:

Theorem 1.9 Let k and s be positive integers. The largest possible order of a group of the form $\mathbb{Z}_2 \times \mathbb{Z}_{2k}$ that has an s-spanning set of size 2 is $2s^2$ when s is even and $2s^2 - 2$ when s is odd.

In Section 4 below we provide some further results on groups of the form $\mathbb{Z}_2 \times \mathbb{Z}_{2k}$.

For $m \ge 3$, the largest order of abelian groups with an m-subset for which the diameter of their Cayley graphs is at most s is generally unknown, though some small cases for cyclic groups were displayed by Haanpää in [7], by Hsu and Jia in [8], and by Graham and Sloane in [6].

Of particular interest are perfect spanning sets. An m-subset A of G is called a perfect s-spanning set, if every element of G arises as a signed sum of at most s terms in a unique way; in this case we have |G| = a(m, s). Above we mentioned perfect spanning sets for m = 1, for s = 1, and for m = 2; we believe that there are no others.

Conjecture 1.10 If an m-subset A of a finite abelian group G is a perfect s-spanning set, then one of the following holds:

- $m = 1, G \cong \mathbb{Z}_{2s+1}$, and $A = \{a\}$ for some $a \in \mathbb{Z}_{2s+1}$ with gcd(a, 2s+1) = 1;
- s = 1, and G is the disjoint union of $\{0\}$, A, and -A; or
- m = 2, $G \cong \mathbb{Z}_{2s^2+2s+1}$, and $A = d \cdot \{s, s+1\}$ for some $d \in \mathbb{Z}_{2s+1}$ with $gcd(d, 2s^2+2s+1) = 1$.

In Sections 2 and 3 below, we present our results on lower bounds and upper bounds, respectively. In Section 4 we treat the case of the groups $\mathbb{Z}_2 \times \mathbb{Z}_{2k}$ in more detail.

2 Constructions and lower bounds

In this section we provide some explicit constructions for s-spanning sets of size 2. We denote the s-span of a 2-subset $A = \{a_1, a_2\}$ of G by

$$\langle A \rangle_s = \{ \lambda_1 a_1 + \lambda_2 a_2 \mid \lambda_1, \lambda_2 \in \mathbb{Z}, |\lambda_1| + |\lambda_2| < s \}.$$

It will be helpful to have the following facts available.

Lemma 2.1 For a given positive integer s, let

$$I(s) = \{(\lambda_1, \lambda_2) \mid \lambda_1, \lambda_2 \in \mathbb{Z}, |\lambda_1| + |\lambda_2| \le s\}.$$

- 1. For a given integer h with $|h| \le s$, the number of ordered pairs $(\lambda_1, \lambda_2) \in I(s)$ with $\lambda_1 + \lambda_2 = h$ equals s + 1 when s and h have the same parity, and s otherwise.
- 2. There are $(s+1)^2$ ordered pairs $(\lambda_1, \lambda_2) \in I(s)$ where $\lambda_1 + \lambda_2$ and s have the same parity, and s^2 where $\lambda_1 + \lambda_2$ and s have opposite parity.
- 3. There are $2s^2 + 2s + 1$ elements in I(s).

Proof. By symmetry, we may assume that $h \ge 0$. Observe that the 'leftmost' integer point (λ_1, λ_2) on the line x + y = h within $|\lambda_1| + |\lambda_2| \le s$ is

$$(-\lfloor (s-h)/2\rfloor, h+\lfloor (s-h)/2\rfloor),$$

and the 'rightmost' such point is

$$(|(s+h)/2|, h-|(s+h)/2|),$$

a set of s+1 points when s and h have the same parity, and s points otherwise. This establishes the first claim, from which our other two claims follow easily.

Proposition 2.2 For all positive integers s and n with

$$2 \le n \le 2s^2 + 2s + 1,$$

the set $\{s, s+1\}$ is an s-spanning set in \mathbb{Z}_n .

Proof. The fact that n cannot be larger than $2s^2 + 2s + 1$ follows from Proposition 1.1. We verify that for every n in the range given above, we have $\langle \{s, s+1\} \rangle_s = \mathbb{Z}_n$.

Considering the elements of

$$S = \{\lambda_1 s + \lambda_2 (s+1) \mid \lambda_1, \lambda_2 \in \mathbb{Z}, |\lambda_1| + |\lambda_2| \le s\}$$

in \mathbb{Z} (rather than \mathbb{Z}_n), we see that they lie in the interval $[-(s^2+s),(s^2+s)]$. Since the index set

$$I(s) = \{(\lambda_1, \lambda_2) \mid \lambda_1, \lambda_2 \in \mathbb{Z}, |\lambda_1| + |\lambda_2| \le s\}$$

contains exactly $2s^2 + 2s + 1$ elements (see Lemma 2.1), it suffices to prove that no integer in $[-(s^2 + s), (s^2 + s)]$ can be written as an element of S in two different ways.

For that, suppose that

$$\lambda_1 s + \lambda_2 (s+1) = \lambda_1' s + \lambda_2' (s+1)$$

for some $(\lambda_1, \lambda_2) \in I(s)$ and $(\lambda'_1, \lambda'_2) \in I(s)$; without loss of generality, we can assume that $\lambda_2 \geq \lambda'_2$. Our equation implies that $\lambda_2 - \lambda'_2$ is divisible by s and is, therefore, equal to 0, s, or 2s.

If $\lambda_2 - \lambda_2' = 2s$, then $\lambda_2 = s$ and $\lambda_2' = -s$; since $|\lambda_1| + |\lambda_2| \le s$ and $|\lambda_1'| + |\lambda_2'| \le s$, this can only happen if $\lambda_1 = \lambda_1' = 0$, but this case leads to a contradiction with our equation.

Assume, next, that $\lambda_2 - \lambda_2' = s$. Our equation then yields $\lambda_1 - \lambda_1' = -s - 1$. In this case, we have $\lambda_1 \leq 0$, $\lambda_1' \geq 0$, $\lambda_2 \geq 0$, and $\lambda_2' \leq 0$, and we see that

$$|\lambda_1| + |\lambda_2| = -\lambda_1 + \lambda_2 = (s+1-\lambda_1') + (s+\lambda_2') = 2s+1 - (|\lambda_1'| + |\lambda_2'|) > s+1,$$

but that is a contradiction.

This leaves us with the case that $\lambda_2 = \lambda_2'$, which implies $\lambda_1 = \lambda_1'$, as claimed, and therefore the set $\{s, s+1\}$ is s-spanning in \mathbb{Z}_n .

Proposition 2.3 For every positive integer s, the set $\{(0,1),(1,1)\}$ is an s-spanning set in $\mathbb{Z}_s \times \mathbb{Z}_{2s}$.

Proof. We need to prove that for any $(a,b) \in \mathbb{Z}_s \times \mathbb{Z}_{2s}$, we have $(a,b) \in \langle \{(0,1),(1,1)\} \rangle_s$. Note that by symmetry we may assume that $0 \le b \le s$. We consider three subcases.

When $b \ge a$, we let $\lambda_1 = b - a$ and $\lambda_2 = a$. We then have

$$\lambda_1(0,1) + \lambda_2(1,1) = (a,b)$$

and

$$|\lambda_1| + |\lambda_2| = \lambda_1 + \lambda_2 = b \le s,$$

so $(a,b) \in \langle \{(0,1),(1,1)\} \rangle_s$.

Next, when $2a - s \le b < a$, we again let $\lambda_1 = b - a$ and $\lambda_2 = a$. In this case, we have

$$|\lambda_1| + |\lambda_2| = -\lambda_1 + \lambda_2 = 2a - b \le s$$
,

so again $(a, b) \in \langle \{(0, 1), (1, 1)\} \rangle_s$.

Finally, when b < 2a - s, we let $\lambda_1 = b - a + s$ and $\lambda_2 = a - s$. Here we have

$$\lambda_1(0,1) + \lambda_2(1,1) = (a-s,b),$$

where $a - s \equiv a \mod s$. Furthermore,

$$|\lambda_1| + |\lambda_2| = \lambda_1 - \lambda_2 = (b - a + s) + (s - a) < s,$$

so again $(a,b) \in \langle \{(0,1),(1,1)\} \rangle_s$. This completes our proof.

Theorem 2.4 Let c and s be positive integers so that the remainder of s mod c is at least (c-1)/2. Let $u = \lfloor s/c \rfloor$, $v = \lceil s/c \rceil$, and k = 2uv. Then $A = \{(1, u), (1, v)\}$ is an s-spanning set in $G = \mathbb{Z}_c \times \mathbb{Z}_{ck}$.

Proof. Let $H = \langle (1, u) \rangle$ be the subgroup of G generated by the element (1, u). Since H has order 2cv, it has cu cosets in G. We claim that these cosets are $\lambda_2 \cdot (1, v) + H$, where $\lambda_2 = 0, 1, \ldots, cu - 1$. In order to prove this, we need to show that these cu cosets are pairwise distinct.

Suppose, then, that $\mu_1(1,v)$ and $\mu_2(1,v)$ are in the same coset of H for some $0 \le \mu_1, \mu_2 \le cu - 1$. This implies that $(\mu_2 - \mu_1) \cdot (1,v)$ is an element of H, so there exists an integer μ for which

$$(\mu_2 - \mu_1) \cdot (1, v) = \mu \cdot (1, u).$$

From the first coordinates, this means that $\mu_2 - \mu_1 - \mu$ is divisible by c, and from the second coordinates we see that $(\mu_2 - \mu_1)v - \mu u$, which, since v = u + 1, equals $(\mu_2 - \mu_1 - \mu)u + (\mu_2 - \mu_1)$, is divisible by ck and thus by cu. Combining these, we get that $\mu_2 - \mu_1$ must also be divisible by cu, but that is only possible if $\mu_1 = \mu_2$.

Let $g \in G$ be arbitrary. According to our claim, we have integers $\lambda_1 \in [-cv+1, cv]$ and $\lambda_2 \in [0, cu-1]$ for which

$$g = \lambda_1(1, u) + \lambda_2(1, v).$$

By assumption, we have integers q (quotient) and r (remainder) with $(c-1)/2 \le r \le c-1$ for which s = qc + r. We then have

$$|\lambda_1| + |\lambda_2| \le cv + (cu - 1) = c(q + 1) + (cq - 1) = 2cq + c - 1 = 2s + (c - 2r - 1) \le 2s. \tag{2}$$

If $|\lambda_1| + |\lambda_2| \le s$, then $g \in \langle A \rangle_s$, and we are done.

Assume then that $|\lambda_1| + |\lambda_2| \ge s + 1$. In this case, let $\lambda_1' = \lambda_1 - \epsilon cv$ and $\lambda_2' = \lambda_2 - cu$, where ϵ equals 1 when $\lambda_1 \ge 0$, and -1 when $\lambda_1 < 0$. We can compute that

$$\lambda_1'(1,u) + \lambda_2'(1,v) = (\lambda_1 - \epsilon cv)(1,u) + (\lambda_2 - cu)(1,v) = \lambda_1(1,u) + \lambda_2(1,v) - (\epsilon cv + cu, (\epsilon + 1)cuv),$$

and since $\epsilon + 1$ equals 0 or 2, in G we have

$$\lambda_1'(1, u) + \lambda_2'(1, v) = \lambda_1(1, u) + \lambda_2(1, v) = g.$$

Finally, we show that $|\lambda'_1| + |\lambda'_2| \leq s$. Indeed, we have

$$|\lambda_1'| + |\lambda_2'| = |\lambda_1 - \epsilon cv| + |\lambda_2 - cu| = cv - |\lambda_1| + cu - |\lambda_2|.$$

From (2) above we see that $cv + cu \le 2s + 1$, and by our current assumption we have $|\lambda_1| + |\lambda_2| \ge s + 1$. Therefore, $|\lambda_1'| + |\lambda_2'| \le s$, as claimed, and our proof is now complete.

Corollary 2.5 Let s be a positive integer for which 2s + 1 is composite, and let p be any prime divisor of 2s + 1. Then the largest noncyclic group with an s-spanning set of size 2 has order at least $2s^2 + 2s - (p^2 - 1)/2$.

Proof. Let us write 2s+1=pt for some integer $t\geq 2$. Note that p and t must be odd, and we have

$$s = \frac{t-1}{2}p + \frac{p-1}{2},$$

 $\lfloor s/p \rfloor = (t-1)/2$, and $\lceil s/p \rceil = (t+1)/2$. Let $G = \mathbb{Z}_p \times \mathbb{Z}_{pk}$ where $k = (t^2-1)/2$. Then G has order $2s^2 + 2s - (p^2-1)/2$ and, according to Theorem 2.4, it has an s-spanning set of size 2.

3 s-regular groups and upper bounds

Recall that we say that the group $\mathbb{Z}_c \times \mathbb{Z}_{kc}$ is *s-regular* for some positive integer *s*, if it possesses an *s*-spanning set of the form $\{(1,u),(1,v)\}$ for some $u,v \in \mathbb{Z}_{kc}$. In this section we provide some upper bounds for the largest possible order of an *s*-regular group.

Theorem 3.1 Let $G = \mathbb{Z}_c \times \mathbb{Z}_{ck}$ for some integers $c \geq 2$ and $k \geq 1$ that has an s-spanning set for some positive integer s. If c is an odd prime, then G is s-regular.

Proof. First, we prove that G has an s-spanning set of the form $A = \{(a, u), (b, v)\}$ for some $a, b \in \mathbb{Z}_c$ and $u, v \in \mathbb{Z}_{ck}$ with $u \not\equiv v \mod c$. Note that if $\{(a, u), (b, v)\}$ is an s-spanning set, then so is $\{(a, u), (-b, -v)\}$, so if our claim were false, then we would have $u \equiv v \equiv -v \mod c$ for each s-spanning set $A = \{(a, u), (b, v)\}$ of G. But since c is an odd prime, that would imply that $u \equiv v \equiv 0 \mod c$, and that is a contradiction with A being a spanning set.

Suppose then that $A = \{(a, u), (b, v)\}$ is an s-spanning in G with $u \not\equiv v \mod c$. We will prove that, in this case, $B = \{(1, u), (1, v)\}$ is also an s-spanning in G.

Let

$$I(s) = \{(\lambda_1, \lambda_2) \mid \lambda_1, \lambda_2 \in \mathbb{Z}, |\lambda_1| + |\lambda_2| \le s\},\$$

and consider the function $f: I(s) \to G$, given by

$$(\lambda_1, \lambda_2) \mapsto \lambda_1(a, u) + \lambda_2(b, v).$$

Since A is an s-spanning set, f is surjective; we can thus find a subset I_0 of I(s) on which the restriction $f_0: I_0 \to G$ of f is bijective. We will prove that $g_0: I_0 \to G$, given by

$$(\lambda_1, \lambda_2) \mapsto \lambda_1(1, u) + \lambda_2(1, v)$$

is also bijective. It suffices to show that g_0 is injective. In order to do so, let us assume that the equation

$$\lambda_1(1, u) + \lambda_2(1, v) = \mu_1(1, u) + \mu_2(1, v)$$

holds in G for some (λ_1, λ_2) and (μ_1, μ_2) of I_0 . We then have

$$\lambda_1 + \lambda_2 \equiv \mu_1 + \mu_2 \bmod c, \tag{3}$$

and

$$\lambda_1 u + \lambda_2 v \equiv \mu_1 u + \mu_2 v \bmod ck. \tag{4}$$

From (4) we find that

$$(\lambda_1 + \lambda_2)u + \lambda_2(v - u) \equiv (\mu_1 + \mu_2)u + \mu_2(v - u) \mod ck;$$

considering (3), this gives us that

$$\lambda_2(v-u) \equiv \mu_2(v-u) \bmod c.$$

Therefore, since c is prime and $u \not\equiv v \mod c$, we get that $\lambda_2 \equiv \mu_2 \mod c$. From (3) then we get $\lambda_1 \equiv \mu_1 \mod c$, and thus

$$\lambda_1 a + \lambda_2 b \equiv \mu_1 a + \mu_2 b \bmod c. \tag{5}$$

But f_0 is an injection, so (5) and (4) can hold together only when $(\lambda_1, \lambda_2) = (\mu_1, \mu_2)$. This completes our proof.

Theorem 3.2 Let $G = \mathbb{Z}_c \times \mathbb{Z}_{ck}$ for some integers $c \geq 2$ and $k \geq 1$, and suppose that G is s-regular for some positive integer s.

- 1. If c is even, then $|G| \leq 2s^2$, and equality may occur only when $s \equiv 0 \mod c$.
- 2. If c is odd and
 - $s \equiv 0 \mod c$, then $|G| < 2s^2 + s$:
 - $s \equiv (c-1)/2 \mod c$, then $|G| < 2s^2 + 2s (c^2 1)/2$:
 - c = 3 and $s \equiv 2 \mod 3$, then $|G| \le 2s^2 + 1$.

In all other cases, $|G| < 2s^2$.

Proof. Let $A = \{(1, u), (1, v)\}$ be an s-spanning set in G, in which case

$$\{(\lambda_1 + \lambda_2, \lambda_1 u + \lambda_2 v) \mid (\lambda_1, \lambda_2) \in \mathbb{Z}^2, |\lambda_1| + |\lambda_2| < s\} = G.$$

Recall from Lemma 2.1 that, for a given integer h with $|h| \leq s$, the number $\Lambda(h)$ of ordered pairs $(\lambda_1, \lambda_2) \in \mathbb{Z}^2$ with $|\lambda_1| + |\lambda_2| \leq s$ and $\lambda_1 + \lambda_2 = h$ equals s+1 when s and h have the same parity, and s otherwise. For each $i=0,1,\ldots,c-1$, we let N(i) denote the sum of all $\Lambda(h)$ with $|h| \leq s$ and $h \equiv i \mod c$. Since A is an s-spanning set in G, we must have

$$\{(\lambda_1 + \lambda_2, \lambda_1 u + \lambda_2 v) \mid (\lambda_1, \lambda_2) \in \mathbb{Z}^2, |\lambda_1| + |\lambda_2| \le s\} = \mathbb{Z}_c \times \mathbb{Z}_{ck},$$

and thus for each $i = 0, 1, \ldots, c - 1$,

$$\{(\lambda_1 + \lambda_2, \lambda_1 u + \lambda_2 v) \mid (\lambda_1, \lambda_2) \in \mathbb{Z}^2, |\lambda_1| + |\lambda_2| \le s, \lambda_1 + \lambda_2 \equiv i \bmod c\} = \{i\} \times \mathbb{Z}_{ck}.$$

In particular,

$$|G| \le cN(i) \tag{6}$$

must hold for each $i = 0, 1, \dots, c - 1$. We now inspect (6) in several cases.

When c is even, we consider i = r + 1, where r is the nonnegative remainder of s mod c (with r + 1 understood to be 0 when r = c - 1). To compute N(r + 1), we need to identify the values of h for which $|h| \le s$ and $h \equiv r + 1 \mod c$, and these values are h = s + 1 - jc for $j = 1, 2, \ldots, j_{\text{max}}$, where

$$j_{\text{max}} = |(2s+1)/c| = (2s-2r)/c + |(2r+1)/c|.$$

Note that $\lfloor (2r+1)/c \rfloor$ equals 0 when $0 \le r \le c/2 - 1$, and 1 when $c/2 \le r \le c - 1$. Since s and h = s + 1 - jc have different parity, and thus $\Lambda(h) = s$ for each j, we have

$$cN(r+1) = (2s - 2r + c|(2r+1)/c|)s$$
,

and we see that this quantity is at most $2s^2$, with equality when r = 0 or r = c/2. But |G| must be divisible by c^2 , so $|G| = 2s^2$ may only occur when r = 0.

When c is odd and $r \le c - 2$, we consider i = r + 2 (where r + 2 is understood to be 0 when r = c - 2). In this case the values of h with $|h| \le s$ and $h \equiv r + 2 \mod c$ are h = s + 2 - tc for $t = 1, 2, ..., t_{\text{max}}$, where

$$t_{\text{max}} = |(2s+2)/c| = (2s-2r)/c + |(2r+2)/c|.$$

Note that $\lfloor (2r+2)/c \rfloor$ equals 0 when $0 \le r \le (c-3)/2$, and 1 when $(c-1)/2 \le r \le c-2$. Since s and h=s+2-tc have the same parity when t is even and different parity when t is odd, we have

$$cN(r+2) = (s-r+c\lfloor (2r+2)/c\rfloor) s + (s-r)(s+1).$$

We consider four subcases. For r = 0, this quantity equals $2s^2 + s$, and when $1 \le r \le (c-3)/2$, it equals (s-r)s + (s-r)(s+1), which is less than $2s^2$. When $(c+1)/2 \le r \le c-2$, we have

$$cN(r+2) = (s-r+c)s + (s-r)(s+1) = (s-r)(2s+1) + cs < (s-(c+1)/2)(2s+1) + cs$$

which again is less than $2s^2$. Our last subcase is when r = (c-1)/2, when we have

$$cN(r+2) = (s - (c-1)/2)(2s+1) + cs = 2s^2 + 2s - (c-1)/2.$$

Observe that when $s \equiv (c-1)/2 \mod c$, then $2s^2 + 2s - (c^2 - 1)/2 \equiv 0 \mod c^2$, and since |G| must be divisible by c^2 , this is the largest it can then be.

Our final case is when c is odd and r = c - 1; we consider i = 0. In this case the values of h with $|h| \le s$ and $h \equiv 0 \mod c$ are h = s + 1 - zc for $z = 1, 2, \ldots, z_{\max}$, where

$$z_{\text{max}} = |(2s+1)/c| = (2s+2-c)/c.$$

Note that z_{max} is odd. Since s and h = s + 1 - zc have the same parity when z is odd and different parity when z is even, we have

$$N(0) = (z_{\text{max}} - 1)/2 \cdot s + (z_{\text{max}} + 1)/2 \cdot (s + 1),$$

and thus

$$cN(0) = (s+1-c)s + (s+1)(s+1) = 2s^2 - (c-3)s + 1,$$

which is less than $2s^2$, unless c=3. This completes our proof.

Theorem 3.3 For $s \ge 3$, every s-regular abelian group of rank 2 has order at most $2s^2 + 2s - 4$, with equality if, and only if, it is isomorphic to $\mathbb{Z}_3 \times \mathbb{Z}_{(2s^2+2s-4)/3}$ for some $s \equiv 1 \mod 3$.

Proof. Suppose that $G = \mathbb{Z}_c \times \mathbb{Z}_{ck}$ is an s-regular group of order at least $2s^2 + 2s - 4$ for some integers $s \geq 3$, $c \geq 2$, and $k \geq 1$. Then, by Theorem 3.2, $c \geq 2$ is odd; furthermore,

- $s \equiv 0 \mod c$, and $2s^2 + 2s 4 < c^2k < 2s^2 + s$, or
- $s \equiv (c-1)/2 \mod c$, and $2s^2 + 2s 4 \le c^2 k \le 2s^2 + 2s (c^2 1)/2$.

Since the first possibility leads to no solutions, our claim is established.

4 The groups $\mathbb{Z}_2 \times \mathbb{Z}_{2k}$

In this section we provide some results for s-spanning sets in groups of the form $\mathbb{Z}_2 \times \mathbb{Z}_{2k}$, including the exact value of the largest possible order of such groups. Our general conjecture is as follows.

Conjecture 4.1 Let k and s be positive integers. The group $\mathbb{Z}_2 \times \mathbb{Z}_{2k}$ has an s-spanning set of size 2 if, and only if, one of the following holds:

- s is odd and $k \le (s^2 1)/2$;
- s is even and $k \leq (s^2 s)/2$;
- $s \equiv 0 \mod 4$, k is even, and $(s^2 s)/2 < k \le s^2/2$; or
- $s \equiv 2 \mod 4$, $k \equiv 2 \mod 4$, and $(s^2 s)/2 < k \le s^2/2$.

Theorem 4.2 Let k and s be positive integers. If the group $G = \mathbb{Z}_2 \times \mathbb{Z}_{2k}$ has an s-spanning set of size 2, then it has order at most $2s^2$.

Proof. If G is s-regular, that is, it has an s-spanning set of the form $\{(1,u),(1,v)\}$ for some $u,v\in\mathbb{Z}_{2k}$, then our claim follows from Theorem 3.2. Since no set of the form $\{(0,u),(0,v)\}$ can be a spanning set, we can suppose that G has an s-spanning set of the form $A = \{(0,u),(1,v)\}$ for some $u,v\in\mathbb{Z}_{2k}$. Therefore, each element in $\langle A\rangle_s$ has the form $(\lambda_2,\lambda_1u+\lambda_2v)$ for some integers λ_1 and λ_2 with $|\lambda_1|+|\lambda_2|\leq s$.

Clearly, if u and v are both even, then A can only generate elements of G with an even second component, and thus it cannot be a spanning set.

Now if u and v are both odd, then $\lambda_1 u + \lambda_2 v$ has the same parity as $\lambda_1 + \lambda_2$ does. According to Lemma 2.1, the parity of $\lambda_1 + \lambda_2$ differs from the parity of s exactly s^2 times (and agrees with it $(s+1)^2$ times) within $|\lambda_1| + |\lambda_2| \le s$. This means that A cannot generate more than s^2 elements in $\mathbb{Z}_2 \times \mathbb{Z}_{2k}$ whose second coordinate has different parity than s does, so if it is an s-spanning set, then we must have $|G| \le 2s^2$.

Next, if u is even and v is odd, then for each element of G that A generates, its two components will have the same parity (namely, the parity of λ_2). Therefore, A cannot be a spanning set.

Finally, we prove that if $A = \{(0, u), (1, v)\}$ is an s-spanning set of G with u odd and v even, then $A' = \{(1, u), (1, v)\}$ is an s-spanning set as well, meaning that G is s-regular and thus has order at most $2s^2$ by Theorem 3.2.

Let $a \in \mathbb{Z}_2$ and $b \in \mathbb{Z}_{2k}$. Since $\langle A \rangle_s = G$, there are integers λ_1 and λ_2 with $|\lambda_1| + |\lambda_2| \leq s$ for which

$$\lambda_1(0, u) + \lambda_2(1, v) = (\lambda_2, \lambda_1 u + \lambda_2 v) = (a', b),$$

where a' = a if b is even and a' = 1 + a when b is odd. Observe that when b is even, then (since u is odd and v is even), λ_1 must be even, and thus

$$\lambda_1(1, u) + \lambda_2(1, v) = (\lambda_1 + \lambda_2, \lambda_1 u + \lambda_2 v) = (\lambda_2, \lambda_1 u + \lambda_2 v) = (a', b) = (a, b).$$

Similarly, when b is odd, λ_1 must be odd, and

$$\lambda_1(1, u) + \lambda_2(1, v) = (1 + a', b) = (a, b).$$

Therefore, $\langle A' \rangle_s = G$, as claimed, which completes our proof.

Theorem 4.3 Let k and s be positive integers with s odd. Then the group $G = \mathbb{Z}_2 \times \mathbb{Z}_{2k}$ has an s-spanning set of size 2 if, and only if, $k \leq (s^2 - 1)/2$.

Proof. The fact that $k \leq (s^2 - 1)/2$ is necessary follows from Theorem 4.2. We may also recall that Theorem 2.4, applied with c = 2 and odd s, yields that with u = (s - 1)/2 and v = (s + 1)/2, the set $\{(1, u), (1, v)\}$ is an s-spanning set in $\mathbb{Z}_2 \times \mathbb{Z}_{s^2 - 1}$. This takes care of the case $k = (s^2 - 1)/2$.

Our goal below is to show that the same values of u = (s-1)/2 and v = (s+1)/2 as integers provide an s-span for the set $\mathbb{Z}_2 \times \{0, 1, 2, \dots, k\}$ when $k \leq (s^2 - 3)/2$. (Here we consider the operation in the first component in \mathbb{Z}_2 and in the second component in \mathbb{Z} .) By symmetry, we can then conclude that $\mathbb{Z}_2 \times \mathbb{Z}_{2k}$ has an s-spanning set of size 2 for $k \leq (s^2 - 3)/2$ as well.

We thus need to establish that for every $a \in \mathbb{Z}_2$ and $b \in \{0, 1, 2, ..., k\}$, there are integers μ_1 and μ_2 with $|\mu_1| + |\mu_2| \le s$ for which

$$\mu_1(1, (s-1)/2) + \mu_2(1, (s+1)/2) = (a, b)$$
 (7)

holds in $\mathbb{Z}_2 \times \{0, 1, 2, \dots, k\}$.

Let us recall from the proof of Theorem 2.4 (applied again for odd s and c=2) that there are integers $\lambda_1 \in [-s, s+1]$ and $\lambda_2 \in [0, s-2]$ for which one of the following two sets of conditions holds:

- (i) $|\lambda_1| + |\lambda_2| \le s$, and equation (7) is satisfied in $\mathbb{Z}_2 \times \mathbb{Z}_{s^2-1}$ with $(\mu_1, \mu_2) = (\lambda_1, \lambda_2)$; or
- (ii) $|\lambda_1| + |\lambda_2| \ge s + 1$, and equation (7) is satisfied in $\mathbb{Z}_2 \times \mathbb{Z}_{s^2 1}$ with $(\mu_1, \mu_2) = (\lambda_1 \epsilon(s + 1), \lambda_2 s + 1)$, where ϵ equals 1 when $\lambda_1 \ge 0$, and -1 when $\lambda_1 < 0$.

To complete our proof, we will show that, considering

$$B = \mu_1(s-1)/2 + \mu_2(s+1)/2$$

as an integer (rather than as an element of \mathbb{Z}_{s^2-1}), we have B=b. Since $0 \le b \le k < (s^2-1)/2$, it suffices to prove that

$$-(s^2 - 1)/2 \le B < s^2 - 1. \tag{8}$$

In case (i), we have $\mu_1 = \lambda_1 \in [-s, s+1]$ and $\mu_2 = \lambda_2 \in [0, s-2]$, from which (8) readily follows. For case (ii), we find that

$$|\mu_1| + |\mu_2| = (s+1-|\lambda_1|) + (s-1-|\lambda_2|) \le 2s - (s+1) = s-1$$

and thus

$$|B| = |\mu_1(s-1)/2 + \mu_2(s+1)/2| \le |\mu_1|(s-1)/2 + |\mu_2|(s+1)/2 \le (|\mu_1| + |\mu_2|)(s+1)/2 \le (s^2 - 1)/2,$$

so (8) holds again. This completes our proof.

Theorem 4.4 Let k and s be positive integers with s even. Then the group $G = \mathbb{Z}_2 \times \mathbb{Z}_{2k}$ has an s-spanning set of size 2 if $k \leq (s^2 - s)/2$.

Proof. Like in the proof of Theorem 4.3, we will prove that there are integers u and v for which the s-span of $A = \{(1, u), (1, v)\}$ contains the set $\mathbb{Z}_2 \times \{0, 1, 2, \dots, k\}$. (Here too we consider the operation in the first component in \mathbb{Z}_2 and in the second component in \mathbb{Z} .)

Since s-1 is odd, from the proof of Theorem 4.3 we see that with u=(s-2)/2 and v=s/2, $\langle A \rangle_{s-1}$ contains the set $\mathbb{Z}_2 \times \{0,1,2,\ldots,k\}$ when $k \leq ((s-1)^2-1)/2$. Therefore, $(1,v)+\langle A \rangle_{s-1}$ contains $\mathbb{Z}_2 \times \{s/2,s/2+1,\ldots,s/2+k\}$ for each $k \leq (s^2-2s)/2$, and since $(1,v)+\langle A \rangle_{s-1} \subseteq \langle A \rangle_s$, our claim follows for all $k \leq (s^2-s)/2$.

References

- [1] B. Bajnok, Additive Combinatorics: A Menu of Research Problems. CRC Press, Boca Raton, 2018, xix+390 pp.
- [2] B. Bajnok, C. Berson, H. A. Just, On perfect bases in finite Abelian groups. Involve 15 (2022), no. 3, 525–536.
- [3] H. Delannoy, Emploi de l'échiquier pour la résolution de certains problèmes de probabilités. Assoc. Franc. Bordeaux 24 (1895), 70–90.
- [4] M. A. Fiol, Comments on "Extremal Cayley digraphs of finite abelian groups" [Intercon. Networks 12 (2011), no. 1–2, 125–135]. J. Inter. Net. 14 (2013), no. 4, 1350016.
- [5] M. A. Fiol, J. L. A. Yebra, I. Alegre, and M. Valero, A discrete optimization problem in local networks and data alignment. *IEEE Trans. Comput.* **36** (1987), no. 6, 702–713.
- [6] R. L. Graham and N. J. A. Sloane, On additive bases and harmonious graphs. SIAM J. Algebraic Discrete Methods 1 (1980), no. 4, 382–404.
- [7] H. Haanpää, Minimum sum and difference covers of abelian groups. J. Integer Seq. 7 (2004), no. 2, Article 04.2.6, 10 pp.
- [8] D. F. Hsu and X. Jia, Extremal problems in the construction of distributed loop networks. SIAM J. Discrete Math. 7 (1994), no. 1, 57–71.
- [9] A.G. Mask, J. Schneider, X. Jia, Extremal Cayley digraphs of finite Abelian groups, J. Intercon. Networks 12 (2011), no. 1–2, 125–135.
- [10] P. Morillo, M. A. Fiol, and J. Fàbrega, The diameter of directed graphs associated to plane tessellations. Tenth British combinatorial conference (Glasgow, 1985). Ars Combin. 20-A (1985), A, 17–27.