Floquet Theory in an Irradiated Nodal Surface Semimetal

Bhaskar Pandit¹, Satyaki Kar^{2*}

¹Netaji Mahavidyalaya, Arambagh, West Bengal - 712601, India

 2AKPC Mahavidyalaya, Bengai, West Bengal -712611, India

A nodal surface semimetal (NSSM) features symmetry enforced band crossings along a surface within the three-dimensional (3D) Brillouin zone. The topological robustness of the same does not always come with nonzero Berry fluxes around nodal surfaces. Irrespective of that, however, light irradiation on such system can result in interesting dynamic behavior. We find that depending on the state of polarization, one can obtain additional Weyl points/ nodal surfaces in the Floquet Hamiltonian that the time periodic system gives rise to. For simplicity we only consider two band spinless models without any spin orbit coupling, the main emphasis being to understand the low energy behavior close to the band crossings and its evolution in a Floquet system in the high frequency limit. There we find the nodal surfaces to perish or get duplicated or triplicated for different polarization scenario or different NSSM Hamiltonians. One findings open up important directions on what out of equilibrium NSSM systems has to offer in many active fields including quantum computations.

I. INTRODUCTION

Band crossing along a two dimensional (2D) surface with linear dispersion away from it is what it takes to form a nodal surface[1] or Weyl surface[2] or Weyl nodal surface[3] which generally indicates topological stability in presence of small symmetry preserving perturbations of the non-interacting Bloch Hamiltonians[3, 4]. A 2D nodal surface can be a closed surface forming a Dirac nodal sphere or pseudo Dirac nodal sphere[5]. But it can also be distinct from conventional Fermi surfaces of ordinary metals as a nodal surface often represents a torus (and hence non-deformable continually to a sphere)[1] with opposite edges connected via periodic boundaries of the Brillouin zone (BZ).

A nodal surface semimetal (NSSM) can be obtained as compulsion from the global internal symmetry providing topological protection because the crossed bands with different eigenvalues for symmetry operator do not hybridize to open a gap[3]. One can also consider a combination of unitary and antiunitary symmetries, namely a space-time inversion symmetry \mathcal{PT} . This in addition to a sublattice symmetry \mathcal{S} can carry a \mathbb{Z}_2 topological index[1] (identifying whether the band gap is inverted or not[2]). For independent \mathcal{PT} and \mathcal{S} operators with $[\mathcal{PT}, \mathcal{S}] = 0$, if one takes $\mathcal{PT} = \mathcal{K}$, the complex conjugation, H(k) is bound to be real whereas due to sublattice symmetry, the Hamiltonian can be transformed to a block anti-diagonal form, which can be unitary transformed to a real matrix[1]. Generically, band-crossing points can spread out a compact surface protected by the symmetries and one can find gapped points $k_{1,2}$ above and below that surface where opposite signs of the Pfaffian of the anti-diagonal Hamiltonian matrix, indicating topological charges, are realized [1, 3]. But the requirement of this kind is quite stringent for it also requires regions within BZ having inverted band ordering with different \mathbb{Z}_2 class[1]. Easier is to find the essential nodal surfaces in a system obeying a nonsymmorphic symmetry where linear band crossings occur at sub-BZ boundaries[1].

A non-symmorphic symmetry [6] is a combination of a point group symmetry and half-lattice translation (but not as a product of a lattice translation and a point group operation [7]) that restrict the form of the band structure both locally and globally leading to unavoidable band crossings in the bulk[9]. A material with such symmetry generally possess a two fold screw rotation and glide reflection symmetry given symbolically as $S_z = \{C_{2z} | \mathbf{T}_z = c/2\}$. A combination of $C = \mathcal{PT}$ and $D = \mathcal{P}S_z$ symmetry can give rise to nodal surfaces. The compound symmetry D turns $k_z \rightarrow -k_z$ and is respected only where $k_z = -k_z$, *i.e.*, at $k_z = 0$ and $k_z = \pi$ where D anticommutes with $\mathcal{PT}[7]$. This protects the twofold degeneracy there. If we instead consider the compound symmetry $D' = \mathcal{T}S_z$ then also, for a spinless case, one gets Kramer's two-fold degeneracy at each point of the $k_z = \pi$ plane and thus the energy bands stick together at the boundary of the BZ[1, 7]. These band crossings are topologically robust both locally and globally[9]. This combination of two-fold screw rotation symmetry and time reversal symmetry features essential band crossings solely determined by symmetry. The system can be considered as a collection of 1D chains parameterized by transverse quantum numbers (k_x, k_y) [1]. Each 1D subsystems has a twisted band structure (like in a Mobius strip) with single band crossing at the boundaries. Here nodal surfaces can be protected even in presence of SOC provided space-time inversion symmetry is violated[1].

A Nielsen-Ninomiya no-go theorem[8] predicts overall zero topological charge within a BZ. But a nodal surface (NS) at the BZ sub-boundaries does not get any symmetrically positioned partner with opposite topological charge to make the overall charge zero and so even being topologically protected (due to having twisted band crossings like in a Mobius strip) by a twofold screw axis and time-reversal symmetry[1, 9], these NSs possess zero \mathbb{Z} charge of Berry fluxes[10]. A non-symmorphic symmetry features a global topology with \mathbb{Z}_2 classification where the band crossing can be avoided only using doubled Hamiltonians in four-band theories [1, 9]. But in a two-band theory, one can also have NS not at such high symmetry points and get characterized by a \mathbb{Z}_2 topological index in presence of \mathcal{PT} and \mathcal{S} symmetry [1]. Besides, it is possible to attach 1D or 2D topological charges leading to nontrivial surface states in these systems. M. Xiao *et al.*[10] showed that even in presence of additional Weyl points within the 3D BZ, one can have topologically charged nodal surfaces at BZ boundaries (*i.e.*, at $k_z = \pi$) yet obeying the no-go theorem.

In this report we study both examples of neutral and topologically charged NSSM under light irradiation and investigate their dynamic behavior using a Floquet-Magnus analysis[14].

II. FORMULATION

Let us first consider an one dimensional chain. A twofold unitary nonsymmorphic symmetry operator[9] for a 2 band model there can be given as

$$G(k_z) = \left(\begin{array}{cc} 0 & e^{-ik_z} \\ 1 & 0 \end{array}\right)$$

with $G^2(k_z) = e^{-ik_z}\sigma_0$ and thereby having $G(k_z)$ eigenvalues of $\pm e^{-ik_z/2}$. The relation $GHG^{-1} = H$ requires the Hamiltonian to be of form

$$H(k_z) = \left(\begin{array}{cc} 0 & Q(k_z) \\ Q^{\star}(k_z) & 0 \end{array}\right)$$

with constraint $Q(k_z)e^{ik_z} = Q^*(k_z)$. But that again requires $Q(k_z)$ and hence energy mode to become zero for some momentum or k_z . Additionally if we also consider an inversion symmetry via operator $\hat{P} = \sigma_2 \hat{i}$ ($\hat{P} = \sigma_1 \hat{i}$), where \hat{i} inverses the momentum, we find the symmetry enforced band crossings to occur at $k_z = 0$ (at BZ boundaries *i.e.*, $k_z = \pm \pi$)[9], where the protected crossed energy bands have different \hat{P} eigenvalues. In cases with band crossings at $k_z = \pm \pi$, one gets a twisted band structure of Mobius strip featuring crossed bands at the BZ sub-boundaries in this nonsymmorphic set-up. And hence this is an essential band crossing[1].

Next for a magnetic nonsymmorphic symmetry (MNS), which is a combination of $G(k_z)$ and time reversal symmetry T, we find that for $T = \mathcal{K}\hat{i}$, two-fold band degeneracy among the Kramer's pairs occurs at every point with $k_z = \pi$. One can see this from the commutation relation [GT, H] = 0 leading to $e^{ik_z}Q(k_z) = Q(-k_z)$. This makes $Q(\pi) = 0$ and thus $k_z = \pi$ a symmetry enforced band crossing point. One can also add a term $f(k_z)\sigma_z$ to the Hamiltonian with $f^*(-k_z) = -f(k_z)$ satisfying the GT symmetry. This demands $Re[f(\pi)] = 0$. Interestingly, the presence of nodal surfaces does not lead to any special gapless boundary mode for they have zero spatial codimensionality[1]. Typically a nodal surface is a torus in a 3D BZ.

We find that a light irradiation on such a system can cause Floquet Hamiltonians that contain Weyl points or nodal lines.

Without loss of generality, we can consider a low energy continuum model about a flattened NSSM (nodal surface semimetal) to be defined about a point $\mathbf{k_0} = (k_{x0}, k_{y0}, \pi)$ in the nodal surface $k_z = \pi$ with $k \to k_0 + q$. Only considering the 1D subsystem along \hat{z} , one can write $H_{np}(q_z) = H(\pi + q_z)$ for which MNS requires $-e^{iq_z}Q(\pi + q_z) = Q(\pi - q_z)$ and $f^*(\pi + q_z) = -f(\pi - q_z)$. Here a Taylor's expansion upto 2nd order gives $Q''(\pi) = -iQ'(\pi)$ (and also $f(\pi + q_z) \sim \sin q_z$) for which a possible solution is $Q(\pi + q_z) \sim 1 - e^{-iq_z}$. This gives a low energy Hamiltonian $H_{np} = v(1 - \cos q_z) \sigma_x + v \sin q_z \sigma_y + v_z \sin q_z \sigma_z$ which for small q_z becomes $H_{np} = v \frac{q_z^2}{2} \sigma_x + v q_z \sigma_y + v_z q_z \sigma_z$.

Now for a 3D system with nodal surfaces protected by nonsymmorphic symmetry, the 3D Hamiltonian can be considered as a collection of 1D k_z subsystems parameterized by (k_x, k_y) so that one can write $H(k_x, k_y, k_z) =$ $H_{k_x,k_y}^{1D}(k_z)$. A 3D Dirac Hamiltonian is expressed in terms of 4×4 matrices, though for our spinless system, a 2×2 Hamiltonian matrix suffices to describe the physics for merely an outer product of $\mathbb{I}_{2\times 2}$ comes into play due to the spin subspace (or spin degeneracy). Thus we can, in general, write the Hamiltonian as[10]

$$H_{NS}^{(1)} = A(k_x, k_y) [\frac{q_z^2}{2}\sigma_x + q_z\sigma_y] + B(k_x, k_y)q_z\sigma_z.$$
 (1)

For simplicity, one can consider constant prefactors A = B = 1 independent of k_x and k_y yielding a Hamiltonian in terms of variable q_z alone.

This NSSM not necessarily feature nonzero topological charge. However, one can choose smart tight binding models which feature both NS and Weyl points[10] and in order to compensate the Weyl point charges, the NS requires to have nonzero topological charges (as per the no-go theorem). Xiao *et. al.* designed a phononic crystal which features a topologically charged NSSM[10]. Likewise one can write a continuum Hamiltonian about a flattened NSSM, more particularly about a point $k_0 =$ (k_{x0}, k_{y0}, π) of the nodal surface $k_z = \pi$, with $k \to k_0 + q$ to be

$$H_{NS}^{(2)} = Aq_z(q_x\sigma_x + q_y\sigma_y) + Bq_z\sigma_z \tag{2}$$

The nodal surface is given by the $q_x - q_y$ plane for $q_z = 0$. The prefactors A and B are compatible with dimensionality and can be taken as unity for simplicity. To accommodate for nonzero charges, the 3D Brillouin zone (BZ) need to also incorporate Weyl points with compensating topological charges as per the Nielsen-Ninomiya no-go theorem[8]. In Ref.[10] there are a pair of Weyl points at $(q_x = \pm 4\pi/3, q_y = 0, q_z = \pi/2)$ apart from a NS at $q_z = \pi$.

III. TOPOLOGY

Let us now discuss the topological characteristics of the nodal surfaces concerned. For obtaining the corresponding topological charge it is customary to consider a manifold S^c enclosing the nodal surface where all bands are gapped[3]. The topological invariant is defined on this manifold of lower dimensionality (called codimensionality) $d_c = d - d_{NS} - 1[11]$. As the dimension of NS, $d_{NS} = 2$, the enclosing manifold in a 3D system corresponds to the codimension, $d_c = 0$. There are different topological invariants one can look into. One can calculate Chern number in surfaces above and below the NS and take a difference to find the topological charge and consequently the topological stability of the NS[3]. Again one can consider two points on two sides of the nodal surface and a difference in the estimate of a symmetry preserving perturbation in an eigenstate calculated at two points on two sides of the NS can constitute a topological invariant of the nodal surface[3]. We know that the total topological charge should add up to zero within the BZ, as per the no-go theorem[8]. So unless there are more than one NSs within the BZ, one can't expect nonzero topological charge for the same.

The Chern number or the Berry phase calculation involves obtaining the line integral of Berry connection or surface integral of Berry curvatures over the whole BZ. But for a NS, we need to consider the 2D cuts above and below the NS and calculate these line and surface integrals there. The topological charge or Chern numbers of Bloch wavefunctions on a surface enclosing the NS can be obtained using Wilson loop method[12, 13]. The Berry connection is given as $A = \langle \psi_k | -i\nabla_k | \psi_k \rangle$, $|\psi_k \rangle$ being the Bloch wavevectors. For a chiral symmetric Hamiltonian (1) with A = 1, B = 0, one gets $|\psi_k \rangle_{\pm} = (\frac{\pm\sqrt{4+q_z^2}}{q_z+2i}, 1)^T$ giving $A(q_z) = A(-q_z)$ leading to zero topological charge for the NS. This is true even considering the full lattice model as well.

For the Hamiltonian 2 with A = B = 1, the energy and wavevectors come out to be $E_{\pm} = \pm sgn(q_z)q_z\sqrt{1+q_{\perp}^2}$ and $\psi_{\pm} = [e^{-i\theta}\frac{1\pm sgn(q_z)\sqrt{1+q_{\perp}^2}}{q_{\perp}}, 1]^T$ respectively with $\theta = tan^{-1}(q_y/q_x)$ and $q_{\perp}^2 = q_x^2 + q_y^2$. For each band we get $v_z = -v_{-z}$ for $q_z \neq 0$. Notice that ψ_{\pm} and hence A do not depend on z. A doesn't depend of θ either. We get $A_z = 0$ and $A_{\theta} = \frac{(1\pm sgn(q_z)\sqrt{1+q_{\perp}^2})^2}{q_{\perp}^3}$. The Berry curvature $\Omega = \nabla \times A$ is independent of the magnitude of q_z but contains only the z component:

 $\Omega = \frac{\mp 2sgn(q_z)}{\sqrt{1+q_{\perp}^2(1\mp sgn(q_z)\sqrt{1+q_{\perp}^2})^2}}\hat{z}.$ Interestingly, Ω blows up at $q_{\perp} = 0$ in the E_+ (E_-) band for $q_z > 0$ ($q_z < 0$) irrespective of the magnitude of q_z (unlike reported in Ref.[10]). Considering all possible \mathbf{k}_0 points, one finds singular Ω in the whole nodal surface as well as above (below) it. Thus it represents a nontrivial topology.

As mentioned before, the calculation of Berry phase involves considering a manifold/surface enclosing the gapless regime of nodal surface. As these nodal surfaces are not simple closed surfaces with genus 0 (like a sphere), we need to consider two surfaces above and below to enclose the nodal surface and then calculate the topological invariant in both of them. Their difference gives the Chern number or Berry phase for the nodal surface. Due to the symmetry of the problem we consider a cylindrical unit cell and its 2D projection transverse to the k_z direction is a circle in the $q_x - q_y$ plane with radius $2\sqrt{\pi}$. So the Berry flux, which is the line integral over this 2D projection of BZ is given by

$$\gamma_B = \left(\int q_\perp A_\theta d\theta\right)_{q_\perp = 2\sqrt{\pi}} = \frac{(1 \pm sgn(q_z)\sqrt{1 + 4\pi})^2}{2}$$
(3)

Notice that it yields $\gamma_B = 2\pi$ for $q_{\perp} \to \infty$, as also mentioned in Ref.[10]. Here $\gamma_B|_{q_z\to 0^+} - \gamma_B|_{q_z\to 0^-} = \pm 2\sqrt{1+4\pi}$ denote the Chern number for E_{\pm} bands signifying topological charge in the nodal surface. As long as Ω is nonsingular, one can also calculate the surface integral across the full 2D BZ that gives

$$\gamma_{B-}|_{q_z \to 0^+} = \gamma_{B+}|_{q_z \to 0^-} = \int_{BZ} \Omega dS = -4\pi \int_0^{2\sqrt{\pi}} \frac{q_\perp dq_\perp}{\sqrt{1+q_\perp^2}(1+\sqrt{1+q_\perp^2})^2} = \frac{1}{2}(1-\sqrt{1+4\pi})^2$$

in compatible with Eq.3. But $\gamma_{B-}|_{q_z \to 0^-}$ and $\gamma_{B+}|_{q_z \to 0^+}$ are not calculable in a similar manner as Ω blows up at $q_{\perp} = 0$ in those limit. Notice that similar calculations about a straight forward lattice model version of Eq.(1), namely,

$$H_L = \sin q_z (\sin q_x \sigma_x + \sin q_y \sigma_y) + q_z \sigma_z \tag{4}$$

would give a zero Berry phase because the information about the Weyl points are missing from it. Including that (like in the tight binding model used in Ref. [10]) one is supposed to get nonzero Chern number although the calculations can be very cumbersome.

IV. IRRADIATION - FLOQUET THEORY

The dynamics of a time periodic Hamiltonian is well expressed via quasi-energies and periodic Floquet modes and the stroboscopic time evolution can be described by a stationary Hamiltonian called a Floquet Hamiltonian[14]. Be it a graphene sheet, a topological insulator or a topological semimetals with time periodic couplings, such Floquet Hamiltonian often displays nontrivialities different from its time independent counterpart[15–18]. Such analysis can be performed on an irradiated NS system where the time periodic E-M waves couple with the system parameters.

We consider both the cases of Hamiltonian (1) (with A = B = 1) and (2) to investigate the Floquet theory. A electric field along \hat{x} or \hat{y} direction does not add any time

dependence in the Hamiltonian (1). But irradiation via linearly polarized wave of frequency ω and electric field $E = -E_0 cos(\omega t)\hat{z}$ brings in an vector potential, which in the Coulomb gauge, can be written as $\mathcal{A} = \frac{E_0}{\omega} sin(\omega t)\hat{z}$. A Peierl's substitution: $\hbar k \to \hbar (k_0 + q) + e\mathcal{A}$ for Hamiltonian (1) leads to the time dependent Hamiltonian H(t)given by

$$H(t) = H_{NS}^{(1)} + \frac{eE_0}{\hbar\omega} sin(\omega t)[q_z \sigma_x + \sigma_y + \sigma_z] \qquad (5)$$
$$+ \frac{e^2 E_0^2}{2\hbar^2 \omega^2} sin^2(\omega t)\sigma_x$$

One can see that such a time periodic system leads to an effective Floquet Hamiltonian indicating band crossings other than the $q_z = 0$ plane. Consider the Fourier modes:

$$H^{(n)} = \frac{1}{T} \int_0^T dt e^{-in\omega t} H(t) \tag{6}$$

leading to $H^{(0)} = H_{NS}^{(1)} + \frac{e^2 E_0^2}{4\hbar^2 \omega^2} \sigma_x$, $H^{(1)}(H^{(-1)}) = -(+)i\frac{eE_0}{2\hbar\omega}[q_z\sigma_x + \sigma_y + \sigma_z]$ and $H^{(n)}, H^{-(n)} = 0$ for |n| > 1. Thus one gets $[H^{(0)}, H^{(1)}] = \frac{eE_0}{2\hbar\omega}(q_z^2 - \frac{e^2 E_0^2}{2\hbar^2 \omega^2})[\sigma_y - \sigma_z],$ $[H^{(-1)}, [H^{(0)}, H^{(1)}]] = \frac{e^2 E_0^2}{2\hbar^2 \omega^2}(q_z^2 - \frac{e^2 E_0^2}{2\hbar^2 \omega^2})[2\sigma_x - q_z(\sigma_y + \sigma_z)].$

The effective stationary Hamiltonian at high frequency limit is given by [14]

$$H_F = H^{(0)} + \sum_{n>0} \frac{[H^{(n)}, H^{-(n)}]}{n\omega} + \sum_{n\neq 0} \frac{[H^{(-n)}, [H^{(0)}, H^{(n)}]]}{2(n\hbar\omega)^2} + \sum_{m\neq n} \frac{[H^{(-n)}, [H^{(n-m)}, H^{(m)}]]}{3nm(\hbar\omega)^2} + O(\frac{1}{\omega^2}) \simeq H_{NS}^{(1)} + \frac{e^2 E_0^2}{4\hbar^2 \omega^2} \sigma_x + (\frac{eE_0}{2\hbar\omega})^2 (2q_z^2 - \frac{e^2 E_0^2}{\hbar^2 \omega^2}) [2\sigma_x - q_z(\sigma_y + \sigma_z)].$$
(7)

Thus the Floquet system does not see band crossings anymore for the zero of the spectrum would now require to satisfy simultaneously $\frac{q_z}{2} + \frac{e^2 E_0^2}{4\hbar^2 \omega^2} + (\frac{eE_0}{\hbar \omega})^2 (q_z^2 - \frac{e^2 E_0^2}{2\hbar^2 \omega^2}) = 0$ and $q_z [1 - (\frac{eE_0}{2\hbar \omega})^2 (2q_z^2 - \frac{e^2 E_0^2}{\hbar^2 \omega^2})] = 0$ which is not possible for a real q_z value. The Floquet system thus become gapped.

As we did not consider any k_x or k_y dependence in the Hamiltonian, the variation in this Floquet system is not rich enough. But when we consider Hamiltonian (2), we witness interesting dynamic behavior. For a linearly polarized light with $E = -E_0 cos(\omega t)\hat{x}$ or $\mathcal{A} = \frac{E_0}{\omega} sin(\omega t)\hat{x}$ the time dependent Hamiltonian becomes

$$H(t) = H_{NP}^{(2)} + \frac{eE_0}{\hbar\omega} \sin(\omega t) Aq_z \sigma_x.$$
 (8)

Among the time-Fourier modes we get $H^{(0)} = H_{NP}^{(2)}$, $H^{(1)}(H^{(-1)}) = -(+)i\frac{eE_0}{2\hbar\omega}Aq_z\sigma_x$ and $H^{(n)}, H^{-(n)} = 0$ for all integer |n| > 1. Thus one gets $[H^{(0)}, H^{(-1)}] = Aq_z^2 \frac{eE_0}{\hbar\omega}[Aq_y\sigma_z - B\sigma_y]$ and consequently

$$\begin{split} [H^{(1)}, [H^{(0)}, H^{(-1)}]] &= -A^2 q_z^3 (\frac{eE_0}{\hbar\omega})^2 [B\sigma_z + Aq_y\sigma_y]. \end{split}$$
 The Floquet Hamiltonian thus takes the form[14]

$$H_F \simeq H_{NP}^{(2)} - A^2 q_z^3 \frac{(eE_0)^2}{2(\hbar\omega)^4} [B\sigma_z + Aq_y\sigma_y] \qquad (9)$$

which indicates a pair of additional Weyl points at $(k_{x0}, k_{y0}, \pm \pi \mp \sqrt{\frac{2}{B}} \frac{(\hbar \omega)^2}{A e E_0})$, in case it falls within the BZ boundary despite the high frequency limit. Notice that this WP turns into a NS if the Hamiltonian 2 remains the same for arbitrary choice of k_{x0} and k_{y0} .

If we change the angle of polarization resulting in $\mathcal{A} = \frac{E_0}{\omega} \sin(\omega t) \hat{z}$, we get

$$H(t) = H_{NS}^{(2)} + \frac{eE_0}{\hbar\omega} sin(\omega t) [A(q_x \sigma_x + q_y \sigma_y) + B\sigma_z]$$
(10)

In this case the energy zeros corresponding to H(t) appear at $(qx, qy, -\frac{eE_0}{\hbar\omega}sin(\omega t))$, showing the nodal plane to sinusoidally fluctuate in time with q_z varying between $(-\frac{eE_0}{\hbar\omega}:\frac{eE_0}{\hbar\omega})$. But the high frequency expansion does not lead to any new terms and the Floquet Hamiltonian remains at the bare zeroth level as: $H_F = H_{NS}^{(2)}$.

Next we consider a circularly polarized light for which we take $\mathcal{A} = \frac{E_0}{\omega} [sin(\omega t)\hat{x} + cos(\omega t)\hat{y}]$ and one gets

$$H(t) = H_{NS}^{(2)} + \frac{eE_0}{\hbar\omega} q_z A[\sin(\omega t)\sigma_x + \cos(\omega t)\sigma_y].$$
(11)

Again we obtain $H^{(0)} = H_{NS}^{(2)}$. But now $H^{(1)}(H^{(-1)}) = \frac{eE_0}{2\hbar\omega}q_z A[-(+)i\sigma_x + \sigma_y]$. Thus $[H^{(1)}, H^{(-1)}] = (\frac{eE_0}{\hbar\omega})^2 q_z^2 A^2 \sigma_z$. And we obtain nonzero correction even in the 1st order of the expansion. More precisely, the Floquet Hamiltonian is given by,

$$H_F = H^{(0)} + \sum_{n>0} \frac{[H^{(n)}, H^{-(n)}]}{n\omega} + O(\frac{1}{\omega^2})$$
$$\simeq H^{(2)}_{NS} + \frac{(eE_0q_zA)^2}{\hbar^2\omega^3}\sigma_z$$
(12)

Thus this time, the Floquet Hamiltonian indicates one single additional nodal plane at $q_z = -\frac{B\hbar^2 \omega^3}{(eE_0 A)^2}$ in the high frequency limit. Again notice that though these additional planes appear as Weyl points in the Hamiltonian, they in fact are indicative of nodal planes if the Hamiltonian remains the same irrespective of the choice of the point (k_{x0}, k_{y0}) in the nodal plane. More generally, for an elliptically polarized light with $\mathcal{A} = \frac{E_0}{\omega} [sin(\omega t + \phi)\hat{x} + cos(\omega t)\hat{y}]$ we get

$$H(t) = H_{NS} + \frac{eE_0}{\hbar\omega} q_z A[\sin(\omega t + \phi)\sigma_x + \cos(\omega t)\sigma_y]$$
(13)

Again $H^{(0)} = H^{(2)}_{NS}$. But now $H^{(1)}(H^{(-1)}) = \frac{eE_0}{2\hbar\omega}q_z A[-(+)ie^{+(-)i\phi}\sigma_x + \sigma_y]$. Thus $[H^{(1)}, H^{(-1)}] =$

 $\left(\frac{eE_0}{\hbar\omega}\right)^2 q_z^2 A^2 \cos(\phi) \sigma_z$. So other than the nodal plane at $q_z = 0$ we get a 2nd nodal plane at $q_z = -\frac{B\hbar^2\omega^3}{(eE_0A)^2\cos\phi}$.

V. SUMMARY

In this brief report, we have studied simple two band models of NSSM to understand its low energy behavior near the band crossings. While some nodal surfaces appear due to additional internal symmetries and carry \mathbb{Z}_2 topological charges with them, some others are outcome of composite symmetries like nonsymmorphic symmetry that make the nodal surface essentially pinned at the BZ sub-boundaries. Charged NS, in compatible with the no-go theorem can be designed with the presence of additional Weyl points within the 3D BZ. Under irradiation, such systems, both with or without a topological charge, show interesting time periodic behavior which for stroboscopic time variation can be well analyzed using Floquet-Magnus theory. We find the nodal surface to vanish in the Floquet system corresponding to a NSSM Hamiltonian constituting zero topological charges for the NS. But in presence of a charged NS, one can get more interesting dynamical phenomena namely doubling or tripling of nodal surfaces in the Floquet Hamiltonian under different choice of polarization directions.

5

but are of more practical interest when situated close to Fermi levels. Our calculations have scope to delve into further deep analysis on dynamic behavior of synthesized materials exhibiting nodal planes close to Fermi surfaces. For example, de Haas-van Alphen spectroscopy in combination with density function theory calculations can show nonsymmorphic symmetry in chiral, ferromagnetic compound MnSi and consequent nodal planes enforces topological protectorates with large Ω at the intersection of NS and the Fermi surface[19]. An irradiation on such system and a consequent Floquet analysis can be performed to explore what dynamic features it has to offer. It will also be interesting to probe the effect of time periodic perturbation on the semimetal ZrSiS where a coexistence of nodal planes and nodal lines are observed^[20] at low energies. Floquet nodal rings and Floquet nodal spheres have also been studied using four-band theories where topological invariants are captured using quantum metric measurements[21]. In future we plan to repeat similar calculations for the non-symmorphic symmetry protected nodal surfaces pinned at Floquet BZ subboundaries. Such time periodic drives have the potential to design tunable topological qubits[22].

ACKNOWLEDGEMENT

Nodal surfaces are not necessarily equi-energy surfaces

SK acknowledges financial help from DST-SERB, Government of India via grant no. CRG/2022/002781.

- W. Wu et al., Phys. Rev. B97, 115125 (2018).
- [2] C. Zhong *et al.*, Nanoscale **8**, 7232 (2016).
- [3] O. Turker *et al.*, Phys. Rev. B**97**, 075120 (2018).
- [4] P. A. volkov, and S. Moroz, Phys. Rev. B98, 241107 (2018).
- [5] J. Wang et al., Phys. Rev. B98, 201112(R) (2018).
- [6] A.Furusaki, Science Bulletin **62**, 788-794 (2017).
- [7] Q.-F. Liang et al., Phys. Rev. B 93, 085427 (2016).
- [8] H. Nielsen, and M. Ninomiya, Nuc. Phys. B 193, 173 (1981).
- [9] Y. X. Zhao et al., Phys. Rev. B94, 195109 (2016).
- [10] M. Xiao et al., arXiv:1709.02363 (2017); M. Xiao et al., Sci. Adv.6, eaav2360 (2020).

- [11] Y. X. Zhao et al., Phys. Rev. Lett.116 156402 (2016).
- [12] Y. Yang et al., Nat. com. 10, 5185 (2019).
- [13] G. Chang et al., Nat. Mat. 17, 978 (2018).
- [14] A. Eckardt et al., New Jour. Phys. 17, 093039 (2015).
- [15] J. Cayssol *et al.*, Phys. Stat. Solidi RRL7, No.1-2, 101 (2013).
- [16] Z. Yan et al., Phys. Rev. Lett.117, 087402 (2016).
- [17] D. Sinha et al., Cur. App. Phys. 18, 1087 (2018).
- [18] S. Kar et al., Phys. Rev. B98, 245119 (2018).
- [19] A. Wilde *et al.*, Nature **594**, 374 (2021).
- [20] B. -B. Fu et al., Sci. Adv. 5, eaau6459 (2019).
- [21] G. Salerno et al., Phys. Rev. Res.2, 013224 (2020).
- [22] M. Yu et al., Natl. Sc. Rev., 7, 254-260 (2020).