ON THE TENSOR PROPERTY OF BERNSTEIN-SATO POLYNOMIAL

QUAN SHI AND HUAIQING ZUO

ABSTRACT. We prove the multiplicative Thom-Sebastiani rule for Bernstein-Sato polynomials, answering the longstanding questions of Budur and Popa. We generalize the result to the tensor of two effective divisors on the product of two arbitrary non-singular complex varieties. This also leads to a multiplicative property related to Igusa's strong monodromy conjecture. Moreover, we propose an extension of our result to Bernstein-Sato polynomials for ideals and prove it for monomial ideals.

2020 Mathematics Subject Classification. 14F10, 13N10, 13A35, 16S32. Keywords: Bernstein-Sato polynomial, roots, singularity.

Content

Convention	1
1. Introduction	1
1.1. Announcement	4
2. Preliminary	4
2.1. Algebraic \mathcal{D} -module over \mathbb{C} and $\mathbb{C}(s)$	4
2.2. Bernstein-Sato polynomial, Existence, and Some Results	7
2.3. Igusa Zeta Function and Strong Monodromy Conjecture	9
2.4. Bernstein-Sato polynomial for Ideals	10
2.5. Combinatoric Description in Monomial Ideal Case	10
3. Tensor Property of Bernstein-Sato Polynomials	11
3.1. Probable Motivations	11
3.2. Proof of $b_f(s)b_g(s) = b_{f \cdot g}(s)$	12
3.3. Generalization to Ideals	15
References	17

Convention. In this paper, we adopt multi-index. We use \boldsymbol{x} to denote an array of variables or numbers depending on the context. For $\boldsymbol{x} = (x_1, ..., x_n)$ a sequence of variables and $\boldsymbol{\alpha} = (\alpha_1, ..., \alpha_n) \in \mathbb{Z}^n$, we use $\boldsymbol{x}^{\boldsymbol{\alpha}}$ to mean $\prod_i x_i^{\alpha_i}$. For the convention of sheaf, we use $f \in \mathcal{F}$ to mean a section in some open set. If \mathcal{F} is a quasi-coherent sheaf over an affine scheme X, we may sometimes mix \mathcal{F} with its module of global sections deliberately.

1. INTRODUCTION

Suppose X is a non-singular complex variety and f is a non-zero regular function. Let \mathcal{D}_X be the sheaf of differential operators of X, which is locally the non-commutative polynomial

Zuo is supported by NSFC Grant 12271280.

ring in partial derivations of X. The Bernstein-Sato polynomial $b_f(s)$ is defined as the non-zero polynomial in s of the minimal degree such that

$$b_f(s) \cdot f^s \in \mathcal{D}_X[s] \cdot f^{s+1},\tag{1.1}$$

where s is a formal indeterminant and the $\mathcal{D}_X[s]$ action on the right-hand side is in the expected way. We have demonstrated the complete definition and related machinery in **Subsection 2.2**. The existence of b_f is a non-trivial issue and is due to the holonomicity of \mathcal{D}_{X_s} -module $\mathcal{O}_{X_s}[\frac{1}{f}] \cdot f^s$, where $X_s = X \times_{\mathbb{C}} \text{Spec } \mathbb{C}(s)$. The case for $X = \mathbb{A}^n_{\mathbb{C}}$ is proven in [Ber71] and for the general case, we refer to [Pop18].

The definition of b_f can be extended to a log pair (X, D), where D is an effective divisor. For an affine covering $\{U_i\}_i$ and sections $f_i \in \Gamma(U_i, \mathcal{O}_X)$ such that $D|_{U_i} = \operatorname{div} f_i$, we have $b_D(s) =$ $\operatorname{lcm}_i b_{f_i}(s)$. b_D is an important invariant of the log pair (X, D). It has a subtle connection with multiplier ideals (for definition, see [Laz04]), and we refer to [Kol97] and [ELSV04]. Also, it plays a role in the classification of singularities (see [KS11], [Sai93], and [Sai09]). There are numerous other applications, we refer to [Pop18].

For $X = \mathbb{A}^n_{\mathbb{C}}$, (1.1) can be rewritten as

$$b_f(s) \cdot f^s = P(s, \boldsymbol{x}, \boldsymbol{\partial}_{\boldsymbol{x}}) \cdot f^{s+1}$$

where P is an element in the Weyl algebra (tensoring $\mathbb{C}[s]$) $A_n(\mathbb{C})[s]$ (see [Sch19]).

For all invariants about functions, it is always natural to ask how will they change if the functions are put into some algebraic operations. For $f \in \mathbb{C}[\mathbf{x}]$ and $g \in \mathbb{C}[\mathbf{y}] = \mathbb{C}[y_1..., y_m]$, Nero Budur proposed the following problem.

Problem 1.1 ([Bud12], 2.11). Is there a multiplicative Thom-Sebastiani rule for b-functions? That is, is there any relation between b_f , b_g , and b_{fg} ?

Years later, Mihnea Popa further asked the following.

Problem 1.2 ([Pop18], page 33). Do the Bernstein-Sato polynomials of f, g, and $f \cdot g$ satisfy

$$b_f(s)b_q(s) = b_{f \cdot q}(s)?$$

Honestly, they are the problems that everyone knows Bernstein-Sato polynomial may consider and even thought clear. However, although one side of the division i.e. $b_{f \cdot g}(s) \mid b_f(s)b_g(s)$ is obvious, the other side is highly non-trivial.

In literature, only few clues related to this problem can be found. In [Wal15], Proposition 2.6.1. tells us $b_f b_g = b_{f \cdot g}$ holds for f and g whose Bernstein-Sato polynomials can be attained by differential operators consisting only of s and ∂_x (or ∂_y). In [Pop18], the formula of Bernstein-Sato polynomial for monomial helps prove the identity, after some works, when either f or g is a monomial. Apart from them, useful results can hardly be found in literature, much less complete proof.

The problems are about the tensor property of Bernstein-Sato polynomials since $f \cdot g \in \mathbb{C}[x, y]$ can be identified as $f \otimes g \in \mathbb{C}[x] \otimes_{\mathbb{C}} \mathbb{C}[y]$. One will also see that tensors appear everywhere in the later sections. In this paper, we prove a strong version of the problems.

Theorem A (Theorem 3.2). Suppose A and B are finitely generated integral regular k-algebras. $C = A \otimes_{\mathbb{C}} B$ is the tensor product, with $f \in A$ and $g \in B$ non-zero. Let $b_f(s), b_g(s), b_{f \cdot g}(s)$ be the corresponding Bernstein-Sato polynomials, then the following equality holds.

$$b_{f \cdot g}(s) = b_f(s)b_g(s).$$

Generalizing to divisors, we have the following global version.

Theorem B (Theorem 3.6). Suppose X_1 and X_2 non-singular complex varieties and D_1, D_2 are effective Cartier divisors on X_1, X_2 , then we have

$$b_{D_1 \otimes D_2}(s) = b_{D_1}(s) \cdot b_{D_2}(s).$$

The theorem turns out to give a product result of the Igusa's strong monodromy conjecture.

Conjecture 2.11 (Strong Monodromy Conjecture, Igusa). Given $f \in \mathbb{Z}[x]$, for every prime p large enough, if s_0 is a pole of $Z_p(f;s)$, then $\Re s_0$ is a root of $b_f(s)$. Moreover, if the order of s_0 as a pole is m, then s_0 is a root of $b_f(s)$ of multiplicity $\geq m$.

Corollary 3.4. Suppose $f \in \mathbb{Z}[x]$ and $g \in \mathbb{Z}[y]$ are non-zero polynomials. Suppose strong monodromy conjecture (Conjecture 2.11) holds for f and g, then it also holds for $f \cdot g \in \mathbb{Z}[x, y]$.

The Bernstein-Sato polynomial can be generalized to ideals (see [BMS06b] and [Mus22]). Let $\mathfrak{a} \subset \mathbb{C}[\mathbf{x}]$ with non-zero generators $f_1, ..., f_r$, then $b_{\mathfrak{a}}(|\mathbf{s}|)$ is the the minimal polynomial in $|\mathbf{s}|$ such that

$$b_{\mathfrak{a}}(|\boldsymbol{s}|)f_{1}^{s_{1}}f_{2}^{s_{2}}...f_{r}^{s_{r}} \in \sum_{\boldsymbol{u}\in\mathbb{Z}^{r},|\boldsymbol{u}|=1} D(A)[\boldsymbol{s}]\cdot(\prod_{u_{i}<0}\binom{s_{i}}{-u_{i}})f_{1}^{s_{1}+u_{1}}f_{2}^{s_{2}+u_{2}}...f_{r}^{s_{r}+u_{r}}.$$

We write more information in **Subsection 2.3**. The general computation is very difficult. But in monomial ideal case, we have a combinatoric description (see [BMS06c]).

One may be interested in ideal analogues of **Problems 1.1** and **Problems 1.2** i.e. for $\mathfrak{a} \subseteq \mathbb{C}[\mathbf{x}]$ and $\mathfrak{b} \subseteq \mathbb{C}[\mathbf{y}]$, whether one can find relations between $b_{\mathfrak{a}}, b_{\mathfrak{b}}$, and $b_{\mathfrak{a}\cdot\mathfrak{b}}$, or even has $b_{\mathfrak{a}}b_{\mathfrak{b}} = b_{\mathfrak{a}\cdot\mathfrak{b}}$. Unfortunately, the general answer to the second is no, even if we do not care about the multiplicity of roots (see **Example 3.8**). But if one of the ideals is principal, we have the following.

Proposition 3.9. Let $\mathfrak{a} \subseteq \mathbb{C}[\mathbf{x}]$ be an ideal and $g \in \mathbb{C}[\mathbf{y}]$, then $b_{\mathfrak{a}}b_g = b_{\mathfrak{a} \cdot (g)}$.

For monomial ideals, we find $W_{\mathfrak{a}} \cup W_{\mathfrak{b}}$ is not that far from $W_{\mathfrak{a}\cdot\mathfrak{b}}$, where $W_{\mathfrak{a}}$ means the roots of $b_{\mathfrak{a}}$, not counting multiplicity.

Theorem C (Theorem 3.10). Let $\mathfrak{a} \subseteq \mathbb{C}[\boldsymbol{x}]$ and $\mathfrak{b} \subseteq \mathbb{C}[\boldsymbol{y}]$ be non-zero monomial ideals, then $W_{\mathfrak{a}} \cup W_{\mathfrak{b}} \subseteq W_{\mathfrak{a}\mathfrak{b}}.$

Theorem D (Theorem 2.16). Let $\mathfrak{a} \subseteq \mathbb{C}[\boldsymbol{x}]$ and $\mathfrak{b} \subseteq \mathbb{C}[\boldsymbol{y}]$ be non-zero monomial ideals, then $W_{\mathfrak{a}} \cup W_{\mathfrak{b}} = W_{\mathfrak{a}\mathfrak{b}} \mod \mathbb{Z}.$

Consequently, we conjecture the following multiplicative Thom-Sebastiani rule for all ideals. Conjecture 3.15. Let $\mathfrak{a} \subseteq \mathbb{C}[x]$ and $\mathfrak{b} \subseteq \mathbb{C}[y]$ be non-zero ideals, then

$$W_{\mathfrak{a}} \cup W_{\mathfrak{b}} \subseteq W_{\mathfrak{ab}}$$

Moreover, we have

$$W_{\mathfrak{a}} \cup W_{\mathfrak{b}} = W_{\mathfrak{a}\mathfrak{b}} \mod \mathbb{Z}.$$

The organization of this paper comes as follows.

In Section 2, we provide preliminary knowledge for our study. In Subsection 2.1, we review the general picture of algebraic D-modules and some variations needed to the proof of the existence of Bernstein-Sato polynomial. In the subsequent subsection of Section 2, the content comes as the subsection title.

In Section 3, we discuss the proof, generalization, and some consequences of Problem 1.1 and Problem 1.2. In Subsection 3.1, we give two sound motivations. In Subsection

QUAN SHI AND HUAIQING ZUO

3.2, we prove a generalized version of the problems and give one of consequences about strong monodromy conjecture, we discuss analogous problems of **Problem 1.1** and **Problem 1.2** about ideals and prove two results for monomial ideals.

1.1. Announcement. This is a paper finished at the beginning of January 2024. It was first submitted to a journal on January 31. In June, we were told there was a preprint that appeared on arxiv on February 7, a week after our submission (see [Lee24],https://arxiv.org/pdf/2402.04512). It also solves the problems in the abstract and has some overlaps with our paper. Two solutions are completed independently. Besides, the methods of the two papers are different. Our solution comes from a definition of b_f from commutative algebra and seems more direct towards the original problems.

2. Preliminary

2.1. Algebraic \mathcal{D} -module over \mathbb{C} and $\mathbb{C}(s)$. In this subsection, we will pay a brief review of the theory of algebraic \mathcal{D} -module related to our theme. Let X be a n-dimensional nonsingular complex algebraic variety i.e. a separated scheme of finite type over \mathbb{C} . To make the definition explicit, the sheaf of differential operators on X, \mathcal{D}_X , is defined to be the subalgebra of $\mathcal{E}nd_{\mathbb{C}}(\mathcal{O}_X)$ generated by \mathcal{O}_X and $\mathcal{D}er_{\mathbb{C}}(\mathcal{O}_X)$.

On an affine open subset U small enough, one can choose local coordinates $x_1, ..., x_n$ i.e.

$$\Omega_{U/\mathbb{C}} = \bigoplus_{i=1}^{n} \mathcal{O}_U \cdot dx_i$$

where $d: \mathcal{O}_X \to \Omega_{X/\mathbb{C}}$ is the universal derivation (Kähler differential). Then on U, the sheaf of differential operators is defined to be the quasi-coherent \mathcal{O}_U -non-commutative algebra

$$\mathfrak{D}_X|_U = \bigoplus_{\boldsymbol{lpha} \in \mathbb{N}^n} \mathcal{O}_U \cdot \boldsymbol{\partial}^{\boldsymbol{lpha}},$$

where $\partial_i : \mathcal{O}_U \to \mathcal{O}_U$ is the dual derivation of dx_i . To distinguish the action of ∂_i on \mathcal{O}_U and its multiplication in \mathcal{D}_X , we use $\frac{\partial f}{\partial x_i}$ to mean $\partial_i(f)$. Here $f \in \mathcal{O}_U$ is a local section. The multiplication of \mathcal{D}_X is defined as

$$[\partial_i, \partial_j] = \delta_{ij}, \ [\partial_i, f] = \frac{\partial f}{\partial x_i}.$$

There is a natural filtration on \mathcal{D}_X , the order filtration $F_{\bullet}\mathcal{D}_X$. The increasing chain $\mathcal{O}_X = F_0\mathcal{D}_X \subseteq F_1\mathcal{D}_X \subseteq \dots$ is a sequence of quasi-coherent \mathcal{O}_X -modules (see [Pop18] or [HTT08]). Locally, we have

$$F_j \mathcal{D}_X|_U = \bigoplus_{\alpha \in \mathbb{N}^n, |\alpha| \le j} \mathcal{O}_U \cdot \partial^{\alpha}.$$

We can define \mathcal{D}_X more intrinsically. Let R be a ring and A an R-algebra, we can define the (non-commutative) subalgebra $F_i D_R(A) \subseteq \operatorname{End}_R(A)$ inductively. Set

$$F_0 D_R(A) := A,$$

$$F_j D_R(A) := \{ P \in \operatorname{End}_R(A) \mid [P, f] \in F_{j-1} D_R(A), \text{ for all } f \in A \}$$

In our case, $R = \mathbb{C}$ and A is a finitely generated regular \mathbb{C} -algebra. In our assumption, for each multiplicative closed subset $S \subseteq A$, there is a canonical isomorphism

$$\pi_j: S^{-1}F_j D_{\mathbb{C}}(A) \to F_j D_{\mathbb{C}}(S^{-1}A).$$

Taking S to be all localizations to a principal open set of A, one finds $F_j D_{\mathbb{C}}(A)$, with A running through all affine open sets of X, defines a quasi-coherent sheaf on X. The sheaf is exactly the

 $F_i \mathcal{D}_X$ defined before (see [Sch19]). In particular, we have

$$F_j D_{\mathbb{C}}(A) = \operatorname{span}_{\mathbb{C}} \{ \delta_1 \delta_2 \dots \delta_j \mid \delta_1, \dots, \delta_j \in F_1 D_{\mathbb{C}}(A) = A \oplus \operatorname{Der}_{\mathbb{C}}(A) \}.$$

In the rest of the paper, we use $F_i D(A)$ instead of $F_i D_{\mathbb{C}}(A)$ if no confusion arises.

A (left) algebraic \mathcal{D}_X -module is a quasi-coherent \mathcal{O}_X -module \mathcal{M} equipped with a (left) \mathcal{D}_X multiplication compatible with the \mathcal{O}_X -multiplication. The quasi-coherence condition is crucial, without which we cannot deal with \mathcal{D}_X -modules locally.

The graded ring of $F_{\bullet}\mathcal{D}_X$ is $\operatorname{Sym}_{\mathcal{O}_X}\mathcal{T}_X$, where $\mathcal{T}_X = \mathcal{H}om(\Omega_{X/\mathbb{C}}, \mathcal{O}_X)$ is the tangent sheaf. Hence, $\operatorname{Spec}_X(\operatorname{gr}^F\mathcal{D}_X) = T^*X$, the cotangent bundle of X. This fact leads to the filtration study of \mathcal{D} -modules.

A compatible filtration on \mathcal{M} is an \mathcal{O}_X -module filtration $G_{\bullet}\mathcal{M}$ such that $F_i\mathcal{D}_X \cdot G_j\mathcal{M} \subseteq G_{i+j}\mathcal{M}$. The graded module $\operatorname{gr}^G\mathcal{M}$ is a $\operatorname{gr}^F\mathcal{D}_X$ -module and hence can be identified with an \mathcal{O}_{T^*X} -module. We also call this module $\operatorname{gr}^G\mathcal{M}$. We call the filtration G_{\bullet} good if $\operatorname{gr}^G\mathcal{M}$ is a coherent \mathcal{O}_{T^*X} -module. \mathcal{M} is called a *coherent* \mathcal{D}_X -module if it admits a good filtration. From [Sch19], we know \mathcal{M} is coherent if and only if it is locally a finitely generated \mathcal{D}_X -module. and all good filtrations are equivalent i.e. bounded by a shift of each other.

For a coherent \mathcal{D}_X -module \mathcal{M} , choose a good filtration $G_{\bullet}\mathcal{M}$. The *characteristic variety* of \mathcal{M} is defined to be

$$\mathrm{Ch}(\mathcal{M}) = V(\mathrm{Ann}_{\mathcal{O}_{T^*X}}\mathrm{gr}^G\mathcal{M}),$$

equipped with reduced scheme structure. $Ch(\mathcal{M})$ is independent of the choice of good filtration, which follows from the fact that all good filtrations are equivalent. Through arguments of symplectic algebraic geometry or reducing to affine spaces, one can prove the dimension of $Ch(\mathcal{M}) \geq n$ if $\mathcal{M} \neq 0$ (see [HTT08], [Pop18] or [Sch19]). As a result, \mathcal{M} is called holonomic holonomic if dim $Ch(\mathcal{M}) = n$ or $\mathcal{M} = 0$. Submodules and quotient modules of holonomic modules are again holonomic since the the graded modules with induced good filtration have larger annihilators.

There are also push-forward and pulling-back operations for \mathcal{D} -modules. The general construction is hard to describe in a few pages since it involves many statements in derived category and the transformation between left and right \mathcal{D}_X -modules. Roughly speaking, let $f : X \to Y$ be a morphism between non-singular varieties, the push-forward is the functor $f_+ := Rf_*(\mathcal{D}_{Y \leftarrow X} \otimes_{\mathcal{D}_X} -) : D^b(\mathcal{D}_X) \to D^b(\mathcal{D}_Y)$ between derived categories (see [HTT08] and [Pop18]).

Here, we only describe the push-forward functor of closed immersion and affine open immersion, in which the performance of the functor is simple.

1. Closed immersion.

Let $i : X \to Y$ be a closed immersion of non-singular varieties and \mathcal{M} be a \mathcal{D}_X -module. Restricting our attention to some small affine open set, we may assume $Y = \operatorname{Spec} A$ with local coordinate $x_1, ..., x_n$ with $X = \operatorname{Spec} A/(x_{r+1}, ..., r_n)$. Let $\tilde{M} = M^{\sim}$ and $i_+ \mathcal{M} = (\overline{M})^{\sim}$ is a \mathcal{D}_Y -module such that

$$\overline{M} = A[\partial_{r+1}, ..., \partial_n] \otimes_A M = \mathbb{C}[\partial_{r+1}, ..., \partial_n] \otimes_{\mathbb{C}} M$$

The \mathcal{D}_Y -module action on $i_+\mathcal{M}$ (equivalent, D(A)-module action on \overline{M}) is described as follows.

Let $P \otimes m \in \mathbb{C}[\partial_{r+1}, ..., \partial_n] \otimes_{\mathbb{C}} M$, we only need to describe the multiplication of (1) $\partial_i, 1 \leq i \leq r$, (2) $\partial_j, r+1 \leq j \leq n$, and (3) $g \in A$ on $P \otimes m$. For (1), one simply defines $\partial_i \cdot (P \otimes m) = P \otimes (\partial_i m)$. For (2), one defines $\partial_j \cdot (P \otimes m) = (\partial_j P) \otimes m$. (3) is a bit complicated. Suppose $A[\partial_{r+1}, ..., \partial_n] \ni f \cdot P = \sum_{\alpha' \in \mathbb{N}^{n-r}} \partial'^{\alpha'} f_{\alpha'}$, where $\alpha' = (\alpha_{r+1}, ..., \alpha_n)$ and $f_{\alpha'} \in A$. Then $f \cdot (P \otimes m)$ is defined to be $\sum_{\alpha' \in \mathbb{N}^{n-r}} \partial'^{\alpha'} \otimes f_{\alpha'} m$.

It is easy to see i_+ is an exact functor and $\mathcal{M} = 0$ iff $i_+\mathcal{M} = 0$. Moreover, for a good filtration G_{\bullet} on \mathcal{M} , we have an induced good filtration on $i_+\mathcal{M}$ locally. That is,

$$G_l \overline{M} := \sum_{i+j=l} (\sum_{|\boldsymbol{\alpha}'| \leq i} \boldsymbol{\partial}'^{\boldsymbol{\alpha}'}) \otimes G_j M.$$

Hence, by definition, one has dim $\operatorname{Ch}(i_+\mathcal{M}) - \dim Y = \dim \operatorname{Ch}(\mathcal{M}) - \dim X$. As a direct corollary, we see that \mathcal{M} is holonomic iff $i_+\mathcal{M}$ is holonomic.

2. Affine open immersion

In this case, let $f : U \hookrightarrow X$ be an affine open immersion and \mathcal{M} be a \mathcal{D}_U -module, then $f_+\mathcal{M} = f_*\mathcal{M}$, equipped with the natural \mathcal{D}_X -module structure given by the restriction maps of sheaves.

The general theorem tells us that holonomicity is preserved by the push-forward of an arbitrary morphism. Applied to the two special cases we introduced, it tells us that i_+ and f_* send holonomic modules to holonomic modules. (Although we have already known the result for i_+ .)

Theorem 2.1 ([HTT08] or [Sch19]). Let $X \to Y$ be a morphism of non-singular varieties, then $f_+: D^b(\mathcal{D}_X) \to D^b(\mathcal{D}_Y)$ induces $f_+: D^b_h(\mathcal{D}_X) \to D^b(\mathcal{D}_X)$. Here $D^b_h(\mathcal{D})$ is the subcategory in which the cohomology modules of every complex are holonomic.


For further use, we need to give a brief sketch of \mathcal{D}_{X_s} -modules. Let $\mathbb{C}(s)$ be the function field of \mathbb{C} and $S = \operatorname{Spec} \mathbb{C}(s)$. Then $X \mapsto X_s := X \times_{\mathbb{C}} S$ is a functor from smooth \mathbb{C} -varieties to smooth $\mathbb{C}(s)$ -varieties of the same dimension, since flatness of morphisms, local freeness of Kählere differential, and local transcendental degree are all preserved. For $X = \operatorname{Spec} A$, one can also define $F_j \mathcal{D}_{X_s} = (F_j D_{\mathbb{C}(s)}(A(s)))^{\sim}$. One can compute $F_j D_{\mathbb{C}(s)}(A(s))$ through $F_j D_{\mathbb{C}}A[s]$ and using the compatibility of $F_j D_{\mathbb{C}}(A[s])$ with the localization $S = \mathbb{C}[s] \setminus \{0\}$. Consequently, $F_j D_{\mathbb{C}(s)}(A(s)) = F_j D_{\mathbb{C}}(A) \otimes_{\mathbb{C}} \mathbb{C}(s)$. For a general variety X, just glue the local module and we have \mathcal{D}_{X_s} one the induced variety X_s .

Again, a \mathcal{D}_{X_s} -module is a quasi-coherent \mathcal{O}_{X_s} -module \mathcal{M} equipped with a \mathcal{D}_{X_s} -multiplication. One can easily define notions of coherence, good filtration, and characteristic varieties. Furthermore, we have $\operatorname{Spec}_{X_s}(\operatorname{gr}^F \mathcal{D}_{X_s}) = T^*X_s = (T^*X)_s$, we also define \mathcal{M} to be holonomic if $\dim \operatorname{Ch}(\mathcal{M}) = \dim X = \dim X_s$ or \mathcal{M} .

Suppose $f: X \to Y$ is a morphism of non-singular \mathbb{C} -varieties, then we can also define the functor $(f_s)_+: X_s \to Y_s$. If f is a closed immersion or an affine open immersion, the construction is the same as before, just substitute \mathbb{C} -tensor for $\mathbb{C}(s)$ -tensor. For a closed immersion, we still have dim $\operatorname{Ch}(i_+\mathcal{M}) - \dim Y = \dim \operatorname{Ch}(\mathcal{M}) - \dim X$ if $\mathcal{M} \neq 0$. In particular, holonomicity is preserved.

Locally, there is a closed immersion $X \hookrightarrow \mathbb{A}^m_{\mathbb{C}(s)}$, then we have dim $\operatorname{Ch}(\mathcal{M}) - \dim X = \dim \operatorname{Ch}(i_+\mathcal{M}) - m \geq 0$, by the Bernstein inequality on $\mathbb{A}^m_{\mathbb{C}(s)}$ (see [Sch19]). Hence we still have dim $\operatorname{Ch}(\mathcal{M}) \geq \dim X$. Besides, since $(i_s)_+$ is exact and faithful $(\mathcal{M} = 0 \iff i_+\mathcal{M} = 0)$, we have holonomic \mathcal{D}_{X_s} -module satisfies a.c.c. and d.c.c. condistion, also applying the result on $\mathbb{A}^m_{\mathbb{C}(s)}$ (see [Sch19]).

For $f: U \to X$ an affine open immersion, $(f_s)_+$ also preserves holonomicity. Here we only sketch the proof for a principal localization $U = \operatorname{Spec} A_g \to X = \operatorname{Spec} A$. Suppose $A = k[z_1, ..., z_m]/I = k[\mathbf{z}]/I$, then $A_g = k[\mathbf{z}, w]/(wg-1, I)$, hence we have the following commutative diagram.



Vertical arrows are both closed immersion and the arrow in the second row is the coordinate projection. Since closed immersion preserves holonomicity, the problem is reduced to $\mathbb{A}^{m+1} \to \mathbb{A}^n$. This can be proven using the Fourier transformation for \mathcal{D} -modules on $\mathbb{A}^m_{\mathbb{C}(s)}$ (see [Sch19]).

In the next subsection, we will apply these tools to sketch the existence of Bernstein-Sato polynomial.

2.2. Bernstein-Sato polynomial, Existence, and Some Results. The Bernstein-Sato polynomial $b_f(s)$ is an important product of the \mathcal{D} -module theory. Let us first give a brief definition of it. Let $X = \operatorname{Spec} A$ be a non-singular affine complex variety (WLOG integral) and $f \in A$ be a non-zero section.

We define a $D_{\mathbb{C}(s)}(A(s)) = D(A)(s)$ -module $M = A_f \cdot f^s$ as follows.

(1) $M = A_f(s)$ as an A-module;

(2) the multiplication of A(s) on M is the natural one i.e.

$$a(b \cdot f^s) := (ab) \cdot f^s, a \in A(s), b \in A_f(s);$$

(3) the $\operatorname{Der}_{\mathbb{C}(s)}(A(s))$ multiplication is operated by considering s a formal power i.e.

$$\delta(c \cdot f^s) = [\delta(c) + (\delta(f) \cdot c)] \cdot f^s, c \in A_f(s), \delta \in \operatorname{Der}_{\mathbb{C}(s)}(A(s));$$

There are some D(A)(s)-submodules of M e.g. $D(A)(s) \cdot f^{k+s} := D(A)(s) \cdot (f^k \cdot f^s)$ and D(A)[s]-submodules e.g. $D(A)[s] \cdot f^{s+k} := D(A)[s] \cdot (f^k \cdot f^s)$. It is clear that $D(A)(s) \cdot f^{s+k} \supseteq D(A)(s) \cdot f^{s+k+1}$ and $D(A)[s] \cdot f^{s+k} \supseteq D(A)[s] \cdot f^{s+k+1}$. The Bernstein-Sato polynomial of f is defined to be the polynomial $b_f(s) \in \mathbb{C}[s] \setminus \{0\}$ such that there exists $P \in D(A)[s]$ satisfying

$$P \cdot f^{s+1} = b_f(s) \cdot f^s. \tag{B-S}$$

For a general non-singular variety X (WLOG integral) and $0 \neq f \in \Gamma(X, \mathcal{O}_X)$, we can also define these two kinds of modules from this local construction. Note that $i_*(\mathcal{O}_{X_s}|_{(X_s)_f}) \cdot f^s = \mathcal{O}_{X_s}[\frac{1}{f}] \cdot f^s$ has the natural \mathcal{D}_{X_s} -module structure, on each affine open subset of which is the same as above. For the latter, we simply glue the locally defined modules $D(A)[s] \cdot f^{s+k}$ to be a $\mathcal{D}_X[s]$ -module. We can also define $b_f(s)$ to be the minimal polynomial such that $b_f(s) \cdot f^s \in \mathcal{D}_X[s] \cdot f^{s+1}$. In affine cases, the two definitions coincide since every module is quasi-coherent, and hence inclusion is determined by global sections. Furthermore, $b_f(s)$ defined on X is the least common multiple of the ones on all affine open sets (if the existence is guaranteed).

Historically, Joseph Bernstein and Mikio Sato discovered $b_f(s)$ independently in the 1970s. Bernstein found it when studying meromorphic extension of distributions and Sato found it when studying prehomogeneous vector spaces (see [AMHJNBTW22]).

In the 50 years since, $b_f(s)$ have been applied to many fields of mathematics, especially in singularity theory. For example, taking $X = \mathbb{A}^n_{\mathbb{C}}$, then the negative of the maximal root $b_f(s)$, α_f , is equal to the log canonical threshold of f (see [Kol97]).

Also, we can define the analytic Bernstein-Sato polynomial, simply by substituting ring A to the ring of holomorphic germ $\mathbb{C}\{x\}$ (for definition, see [GLS07]). Morihiko Saito proved α_f coincides with the minimal spectrum number (see [Sai06] and [Var81]). More surprisingly, roots of $b_f(s)$ happen to be related to two other important invariants, poles of topological zeta function $Z_{top,f}(s)$ and eigenvalues of monodromy action T_f (for definition, see [DL92] and [vS20] respectively). This relation is the well-known *Monodromy Conjecture*.

Conjecture 2.2 (Monodromy Conjecture, [DL92]). Assume $s_0 \in \mathbb{C}$ is a pole of $Z_{f,top}(s)$, then $e^{2\pi i s_0}$ is an eigenvalue of T_f and s_0 is a zero of $b_f(s)$.

These are the general stories of Bernstein-Sato polynomials. Next, we will show some important properties of the Bernstein-Sato polynomials. The existence of $b_f(s)$ is itself non-trivial. For $X = \mathbb{A}^n_{\mathbb{C}}$, the existence is proven by using the Bernstein filtration on the Weyl algebra $A_n(\mathbb{C}(s))$ and some holonomicity arguments (see [Ber71]). For a general non-singular variety X, there is no such filtration. But fortunately, we can use an exquisite construction to circumvent this obstacle.

Theorem 2.3 (see also [Pop18]). Let X be a non-singular complex variety (WLOG integral) and $0 \neq f \in \Gamma(X, \mathcal{O}_X)$. Then the Bernstein-Sato polynomial $b_f(s)$ exists.

Proof. The problem is local, we may assume $X = \operatorname{Spec} A$ is affine. Let U = D(f) be an open affine subset of X and $j : U \hookrightarrow X$ be the open immersion. One can find $\mathcal{O}_U \cdot f^s$ is a holonomic \mathcal{D}_{U_s} -module since its characteristic variety is the zero section of T^*X_s . Hence, we have $(j_s)_+(\mathcal{O}_U \cdot f^s) = \mathcal{O}_X[\frac{1}{f}] \cdot f^s$ is holonomic. Considering the following descending chain of its submodules.

$$\mathcal{D}_{X_s} \cdot f^s \supseteq \mathcal{D}_{X_s} \cdot f^{s+1} \supseteq \mathcal{D}_{X_s} \cdot f^{s+2} \supseteq \dots$$

Since holonomic modules satisfy d.c.c. considtion, there is some $k \ge 0$ such that $\mathcal{D}_{X_s} \cdot f^{s+k} = \mathcal{D}_{X_s} \cdot f^{s+k+1}$. Apply the automorphism $s \mapsto s+k$ of $\mathcal{O}_X[\frac{1}{f}] \cdot f^s$, we have $\mathcal{D}_{X_s} \cdot f^s \supseteq \mathcal{D}_{X_s} \cdot f^{s+1}$. Hence, there is some $P \in \Gamma(X, \mathcal{D}_{X_s}) = D(A)(s)$ such that $f^s = P \cdot f^{s+1}$. Killing the denominators of P (polynomials in $\mathbb{C}[s]$), we are done.

More than existence, Kashiwara proved the following.

Theorem 2.4 ([Kas77]). Roots of $b_f(s)$ are all negative rational numbers.

We can give an explicit description of $b_f(s)$ in commutative algebra, as below. The action of $F_j D(A)[s]$ on $A_f[s]f^s$ gives an ideal I_j as follows.

$$F_j D(A)[s] \cdot f^{s+1} = I_j \cdot f^{s+1-j}$$

 $I_j \subset A[s]$ is an ideal and it is related to I_{j-1} by the identity below.

$$I_{j} = I_{j-1} \cdot f + A[s] \cdot \{\delta(a)f + (s+2-j)a\delta(f) \mid a \in I_{j-1}, \delta \in \text{Der}_{k}(A)\}$$

This is simply because $F_j D(A) = F_1 D_k(A) \cdot F_{j-1} D_k(A) = (A \oplus \text{Der}_{\mathbb{C}}(A)) \cdot F_{j-1} D(A).$

Consequently, $0 \neq b_f(s)$ is the polynomial with the minimal degree such that $b_f(s) \cdot f^{j-1} \cdot I_j$ for some j. We point out this description is useful in proving our main theorem (**Theorem 3.2**).

The following two propositions show $b_f(s)$ is invariant under the multiplication of units and can be locally glued. As a result, for an effective divisor D on non-singular complex variety X, one can define the Bernstein-Sato polynomial of D as

$$b_D(s) = \lim b_{f_i}(s),$$

where $\{U_i\}_i$ is an affine open covering of $X, f_i \in \Gamma(U_i, \mathcal{O}_X)$, and $D|_{U_i} = \operatorname{div}(f_i)$.

We will give the proof of the first proposition and only give reference to the second.

Proposition 2.5. Suppose X = Spec A is an integral non-singular complex affine variety. Let $0 \neq f \in A$ and $g \in A^*$, then $b_f(s) = b_{f \cdot g}(s)$.

Proof. For $f \cdot g$, we define the ideals J_j as the I_j above i.e.

$$F_j D(A)[s] \cdot (f \cdot g)^{s+1} = J_j \cdot (f \cdot g)^{s+1-j}.$$

Since g is invertible, it suffices to prove $I_j = J_j$ for all j. We prove it by induction. For j = 0, both sides are A[s]. Suppose the identity holds for j - 1. We can present I_j by I_{j-1} i.e.

$$I_j = I_{j-1} \cdot f + A[s] \cdot \{\delta(a)f + (s+2-j)a\delta(f) \mid a \in I_{j-1}, \delta \in \operatorname{Der}_{\mathbb{C}}(A)\}$$

Similar presentation holds for J_j i.e.

$$J_j = J_{j-1} \cdot gf + A[s] \cdot \{\delta(a)gf + (s+2-j)a[g\delta(f) + \delta(g)f] \mid a \in J_{j-1}, \delta \in \operatorname{Der}_{\mathbb{C}}(A)\}.$$

Since g is invertible, $(s+2-j)a\delta(g)f \in J_{j-1} \cdot gf$. Then we can rewrite J_j as

$$J_j = J_{j-1} \cdot gf + A[s] \cdot \{\delta(a)gf + (s+2-j)ag\delta(f) \mid a \in J_{j-1}, \delta \in \operatorname{Der}_{\mathbb{C}}(A)\}.$$

By the induction hypothesis, $I_{j-1} = J_{j-1}$, we then have $I_j = J_j$.

Remark 2.6. The argument can also be applied to analytic $b_f(s)$ for $f \in \mathbb{C}\{x\}$. Since an automorphism of $\mathbb{C}\{x\}$ gives an isomorphism of its ring of differential operators, the result tells us in a direct way that $b_f(s)$ is a contact invariant of hypersurface singularity.

Proposition 2.7 (see [Pop18]). Suppose X = Spec A is an integral non-singular affine complex variety and $f \in A$, then the global and local Bernstein-Sato polynomials are related by the formula

$$b_f(s) = \lim_{i \to 0} b_{f|_{U_i}}(s).$$

where $\{U_i\}_i$ is an arbitrary affine open covering of X and $b_{f|_{U_i}}(s)$ the Bernstein-Sato polynomial of $f|_{U_i}$ on U_i .

The computation of $b_f(s)$ is a rather difficult problem and there has not been a general formula so far. In the rest of this subsection, we present some computational results of $b_f(s)$.

Theorem 2.8 (see [Pop18]). Let $f = \det(x_{ij}) \in \mathbb{C}[x_{ij}]_{1 \leq i,j \leq n}$ be the determinant of variables, then

$$b_f(s) = (s+1)(s+2)...(s+n).$$

Theorem 2.9 ([Sai16]). The Bernstein-Sato polynomial of a generic central hyperplane arrangement $f \in \mathbb{C}[\mathbf{x}]$ of degree $l \ge n$ is

$$b_f(s) = (s+1)^{n-1} \prod_{j=0}^{2l-n-2} (s+\frac{j+n}{l}).$$

Theorem 2.10 (see [Gra10] or [Pop18]). Suppose $f \in \mathbb{C}[\mathbf{x}]$ is a weighted homogeneous polynomial defining an isolated singularity at the origin, with weight type $\mathbf{w} = (w_1, ..., w_n)$. Let $M_f := \mathbb{C}\{\mathbf{x}\}/(\frac{\partial f}{\partial x_1}, ..., \frac{\partial f}{\partial x_1})$ be the Milnor algebra of f and $\{\mathbf{x}^{\alpha_i}\}_{1 \leq i \leq \mu}$ be a monomial basis of M_f . Here $\mu = \dim_{\mathbb{C}} M_f$ is the Milnor number of f. Let Δ be the set (not counting multiplicity) consisting of all rational numbers $\mathbf{w} \cdot \mathbf{\alpha}_i$.

The Bernstein-Sato polynomial of f is

$$b_f(s) = (s+1) \prod_{\rho \in \Delta} (s+|\boldsymbol{w}|+\rho).$$

2.3. Igusa Zeta Function and Strong Monodromy Conjecture. In this section, we review the *p*-adic Igusa zeta function and the strong monodromy conjecture. For a reference, we recommend [Nic10].

Let p be a rational prime number, then we have the p-adic number field \mathbb{Q}_p is a locally compact group and admits a unique Haar measure μ_p up to scalar (see [Wei67]). After normalizing, we have $\mu_p(p^r \mathbb{Z}_p) = p^{-r}$ for all $r \in \mathbb{Z}$.

Suppose $f \in \mathbb{Z}[x]$ is a non-zero polynomial, the *p*-adic Igusa zeta function is defined as

$$Z_p(f;s) = \int_{\mathbb{Z}_p^n} |f|_p^s \, d\mu_p$$

It was shown by Igusa that $Z_p(f;s)$ is a rational function of p^{-s} (see [Igu75]). There is a well-known conjecture about the poles of $Z_p(f;s)$ (see [Nic10] or [Mus22]).

Conjecture 2.11 (strong monodromy conjecture, Igusa). Given $f \in \mathbb{Z}[x]$, for every prime p large enough, if s_0 is a pole of $Z_p(f;s)$, then s_0 is a root of $b_f(s)$. Moreover, if the order of s_0 as a pole is m, then $\Re s_0$ is a root of $b_f(s)$ of multiplicity $\geq m$.

2.4. Bernstein-Sato polynomial for Ideals. The Bernstein-Sato polynomial can be generalized to an arbitrary variety (see [BMS06b] or [Mus22]). Definitions in the two papers are stated differently but equivalent in fact. We first review the definition.

Suppose X = Spec A is an affine non-singular complex variety. Let $\mathfrak{a} = (f_1, ..., f_r)$ $(f_1, ..., f_r \neq 0)$ be an ideal and $s = (s_1, ..., s_r)$ be a set of indeterminant. We consider the following equation:

$$b_{\mathfrak{a}}(|\boldsymbol{s}|)f_{1}^{s_{1}}f_{2}^{s_{2}}...f_{r}^{s_{r}} \in \sum_{\boldsymbol{u}\in\mathbb{Z}^{r},|\boldsymbol{u}|=1} D(A)[\boldsymbol{s}]\cdot(\prod_{u_{i}<0}\binom{s_{i}}{-u_{i}})f_{1}^{s_{1}+u_{1}}f_{2}^{s_{2}+u_{2}}...f_{r}^{s_{r}+u_{r}}.$$

The action of D(A) is the natural in the same sense as (B-S) and $b_{\alpha} \in \mathbb{C}[t]$ is a polynomial. The existence and independence of choice of generators of α was proven in [BMS06b] or one may turn to the following result in [Mus22].

Theorem 2.12 (see [Mus22]). Let X be a smooth complex algebraic variety. If $f_1, ..., f_r \in \Gamma(X, \mathcal{O}_X)$ are nonzero, generating the ideal sheaf \mathfrak{a} , and if $g = \sum_{i=1}^r f_i y_i \in \Gamma(X \times \mathbb{A}^r, \mathcal{O}_{X \times \mathbb{A}^r})$, then $b_{\mathfrak{a}}(s) = b_g(s)/(s+1)$.

The theorem also provides a method to compute $b_{\mathfrak{a}}$, but the general computation is still difficult so far. Even if we do not consider the multiplicity, the determination of roots is also a hard issue. But in monomial ideal cases, we can find the roots in a combinatoric way (see the next subsection). $b_{\mathfrak{a}}$ satisfies the following union property.

Theorem 2.13. Suppose X and Y are non-singular complex algebraic varieties. Let $f_1, ..., f_r \in \Gamma(X, \mathcal{O}_X) \setminus \{0\}$ and $g_1, ..., g_t \in \Gamma(Y, \mathcal{O}_Y)$ define ideal sheaves \mathfrak{a} and \mathfrak{b} . We have an ideal sheaf \mathfrak{c} of $X \times Y$ defined by $f_1, ..., f_r, g_1, ..., g_t \in \gamma(X \times Y, \mathcal{O}_{X \times Y})$, then their Bernsetin-Sato polynomial satisfies the following:

$$b_{\mathfrak{a}}(s) = \prod_{\alpha} (s+\alpha)^{n_{\alpha}}, b_{\mathfrak{b}}(s) = \prod_{\beta} (s+\beta)^{m_{\beta}}, \text{ and } b_{\mathfrak{c}}(s) = \prod_{\gamma} (s+\gamma)^{n_{\gamma}},$$

where $q_{\gamma} = \max\{n_{\alpha} + m_{\beta} - 1 \mid n_{\alpha} > 0, m_{\beta} > 0, \alpha + \beta = \gamma\}.$

2.5. Combinatoric Description in Monomial Ideal Case. In this subsection, we review the combinatoric description of roots of $b_{\mathfrak{a}}$ when $\mathfrak{a} \subseteq \mathbb{C}[\mathbf{x}]$ is a monomial ideal. For reference, we recommend [BMS06a] and [BMS06c].

For a monomial ideal $\mathfrak{a} \triangleleft \mathbb{C}[\boldsymbol{x}]$, generated by $\boldsymbol{x}^{\boldsymbol{v}_i}, \boldsymbol{v}_i = (v_i^1, v_i^2, ..., v_i^n) \in \mathbb{N}^n, i = 1, 2...r$. Let

$$\Gamma_{\mathfrak{a}} := \{ \boldsymbol{u} \in \mathbb{N}^n \mid \boldsymbol{x}^{\boldsymbol{u}} \in \mathfrak{a} \}$$

be the semigroup associated with \mathfrak{a} . The Newton polyhedron of \mathfrak{a} , $P_{\mathfrak{a}}$, is defined to be the convex hull of $\Gamma_{\mathfrak{a}}$. A face Q of $P_{\mathfrak{a}}$ refers to (a) $P_{\mathfrak{a}}$ itself, or (b) the intersection of an affine hyperplane H and $P_{\mathfrak{a}}$ with $P_{\mathfrak{a}}$, where $P_{\mathfrak{a}}$ lie in one of the half space given H. We call a face of (b) type a proper face.

Let $\boldsymbol{e} = (1, 1, ..., 1) \in \mathbb{N}^n$. If Q is a face, let

$$M_Q := \boldsymbol{e} + \mathbb{N}\{\boldsymbol{u} - \boldsymbol{v} \mid \boldsymbol{u} \in \Gamma_{\mathfrak{a}}, \boldsymbol{v} \in \Gamma_{\mathfrak{a}} \cap Q\}$$

be the translation of the semigroup generated by $\boldsymbol{u} - \boldsymbol{v}, \boldsymbol{u} \in \Gamma_{\mathfrak{a}}, \boldsymbol{v} \in Q \cap \Gamma_{\mathfrak{a}}$ by \boldsymbol{e} . Take $M'_Q = \boldsymbol{v}_0 + M_Q$, where $\boldsymbol{v}_0 \in Q \cap \Gamma_{\mathfrak{a}}$. M'_Q does not depend on the choice of \boldsymbol{v}_0 . If Q is not contained in any coordinate hyperplane, there is some linear function L_Q over \mathbb{R}^n s.t. $L|_Q \equiv 1$. Let V_Q denote the linear span of Q, then L_Q is uniquely determined on V_Q . Hence the set

$$R_Q := \{-L_Q(u) \mid u \in (M_Q \setminus M'_Q) \cap V_Q\}$$

is well-defined.

Particularly, if Q is a facet i.e. codim Q = 1, then $V_Q = \mathbb{R}^n$ and hence L_Q is unique. Let m_Q be the minimal positive integer such that the coefficients of $m_Q L_Q$ are all integers.

The following are **Theorem 1.1** and **Theorem 1.2** of [BMS06c].

Theorem 2.14. The roots of the Bernstein-Sato polynomial of \mathfrak{a} is the union of all R_Q , where Q runs through all faces not contained in any coordinate hyperplanes.

Theorem 2.15. The set of classes in \mathbb{Q}/\mathbb{Z} of the roots of the Bernstein-Sato polynomial of \mathfrak{a} is equal to the union of subgroups generated by $\frac{1}{m_Q}$, with Q running over the facets of $P_{\mathfrak{a}}$ that are not contained in any coordinate hyperplanes.

We give an example of **Theorem 2.14**.

Example 2.16. Let $\mathfrak{a} = (x^{a_1}y^{b_1}, x^{a_2}y^{b_2}) \subseteq \mathbb{C}[x, y]$ be a monomial ideal, where $a_1 < a_2$ and $b_1 > b_2$. There are five proper faces of $P_{\mathfrak{a}}$. They are L, the diagonal, X the horizontal, Y the vertical, and points P_1, P_2 .

The linear function L_L defined by L is

$$L_L(x,y) = -\frac{b_2 - b_1}{b_1 a_2 - a_1 b_2} x + \frac{a_2 - a_1}{b_1 a_2 - a_1 b_2} y$$

We have $M_L = (1, 1) + \mathbb{N}^2 + \mathbb{Z}(a_1 - a_2, b_1 - b_2)$. After a elementary calculation, one finds $M_L \setminus M'_L$ consisting of those points $(s + 1, t + 1) + \mathbb{Z}(a_1 - a_2, b_1 - b_2), s, t \ge 0$ such that

$$\mathbb{Z} \cap \left[\frac{t-b_1}{b_2-b_1}, \frac{s-a_1}{a_2-a_1}\right] = \emptyset.$$

Consequently, R_L is provided.

For X and Y, one may find they contribute to $W_{\mathfrak{a}}$ the following roots.

$$\frac{1}{a_1}, ..., \frac{a_1 - 1}{a_1}, ..., 1, \frac{1}{b_2}, ..., \frac{b_2 - 1}{b_2}, 1$$

For P_1 and P_2 , L_{P_1} can be chosen as $\frac{x}{a_1}$. It is easy to see all $M_{P_1} \cap V_Q \cap \{x \ge a_1\} \subseteq M'_{P_1}$. Then no new roots emerge. A similar result holds for P_2 .

In particular if $(a_1, b_1, a_2, b_2) = (0, b, a, 0)$, we have

$$L_L = \frac{x}{a} + \frac{y}{b},$$

$$M_L \setminus M'_L = \{(s+1,t+1) \mid 0 \le s < a, 0 \le t < b\} + \mathbb{Z}(-a,b),$$

$$R_L = \{-\frac{ai+bj}{ab} \mid 1 \le i \le b, 1 \le j \le a\}.$$

Hence, for $\mathfrak{a} = (x^a, y^b)$, we have

$$W_{\mathfrak{a}} = \{ -\frac{ai+bj}{ab} \mid 1 \le i \le b, 1 \le j \le a \}.$$

3. Tensor Property of Bernstein-Sato Polynomials

3.1. Probable Motivations. Let $f \in \mathbb{C}[\mathbf{x}]$ and $g \in \mathbb{C}[\mathbf{y}]$ be two non-zero polynomials. It is obvious that $b_{f \cdot g}(s)$ is a divisor of $b_f(s) \cdot b_g(s)$ since if $P_f \in \mathbb{C}[\mathbf{x}, \partial_{\mathbf{x}}, s]$ and $P_g \in \mathbb{C}[\mathbf{y}, \partial_{\mathbf{y}}, s]$ satisfies

$$P_f \cdot f^{s+1} = b_f(s) \cdot f^s, \ P_g \cdot g^{s+1} = b_g(s) \cdot g^s,$$

then we have

$$(P_f \cdot P_g) \cdot (f \cdot g)^{s+1} = b_f(s)b_g(s) \cdot (f \cdot g)^s$$

So one may be interested in the converse divisibility. In our preliminary work, we proved at least (s + 1) timing lcm $(b_f(s), b_a(s))$ is a divisor i.e.

$$(s+1)\operatorname{lcm}(b_f(s), b_g(s)) \mid b_{f \cdot q}(s).$$

Hence, if we only consider the cases when f and g are weighted homogeneous with isolated singularity, and all denominators of weights of f, g are coprime, we have $b_f(s)b_g(s) = b_{f\cdot g}(s)$ by **Theorem 2.10**.

Another motivation seems more profound. We further assume coefficients of f and g are rational integers. Then we have a trivial relation among zeta functions:

$$Z_p(f \cdot g; s) = Z_p(f; s) \cdot Z_p(g; s).$$

If the strong monodromy conjecture, especially the assertion about the multiplicity, holds, it is natural to ask if $b_{f \cdot q}(s) = b_f(s) \cdot b_g(s)$.

3.2. **Proof of** $b_f(s)b_g(s) = b_{f \cdot g}(s)$. In this subsection, we give a positive answer to a generalized version of " $b_f(s)b_g(s) = b_{f \cdot g}(s)$ ". More precisely, we generalized polynomials f, g to effective divisors on non-singular varieties. We first prove a simple lemma below.

Lemma 3.1. Let R be a PID and M, N be free R-modules. Suppose $M_1, M_2 \subseteq M$ and $N_1, N_2 \subseteq N$ are submodules (hence free). For free submodules $M' \subseteq M$ and $N' \subseteq N$, we have canonical injection $M' \otimes N' \to M \otimes N$, and hence identify $M' \otimes N'$ with its image in $M \otimes N$. In this sense, we have the following identity:

$$(M_1 \cap M_2) \otimes (N_1 \cap N_2) = \bigcap_{i,j \in \{1,2\}} (M_i \otimes N_j).$$

Proof. Let us first prove the following:

$$(M_1 \otimes N_1) \cap (M_2 \otimes N_1) = (M_1 \cap M_2) \otimes N_1.$$
(3.1.1)

Since N_1 is free, we set $N_1 = A^{\oplus Q}$. Take a basis $\{b_i\}_{i \in Q}$ of N_1 , then elements of $M \otimes N_1$ can be uniquely written in a finite sum $\sum_{i \in Q} m_i \otimes b_i, m_i \in M$. Then one can easily deduce (3.1.1).

In parallel, we have

$$(M_1 \otimes N_2) \cap (M_2 \otimes N_2) = (M_1 \cap M_2) \otimes N_2.$$

Applying the same argument to $(M_1 \cap M_2)$ (also free), N_1 , and N_2 , we are done.

Theorem 3.2. Suppose A and B are finitely generated integral regular k-algebras. $C = A \otimes_{\mathbb{C}} B$ is the tensor product, with $f \in A$ and $g \in B$ non-zero. Let $b_f(s), b_g(s), b_{f \cdot g}(s)$ be the corresponding Bernstein-Sato polynomials, then the following equality holds.

$$b_{f \cdot g}(s) = b_f(s)b_g(s).$$

Here we use \cdot instead of \otimes in the multiplication of C.

Remark 3.3. One side of the division i.e. $b_{f \cdot g}(s) | b_f(s)b_g(s)$ is trivial since we simply need to multiplicate the operators to attain $b_f(s)$ and b_g . Hence, we only need to consider the other side of the division. Moreover, the case with f or g invertible is trivial, so we may assume f and g are both non-invertible.

Proof. As in Subsection 2.2, we define $I_j \subseteq A[s]$ to be an ideal such that $F_j D(A) f^{s+1} = I_j \cdot f^{s+1-j}$.

Similarly, we define $J_j \subseteq B[s]$ and $L_j \subseteq C[s]$ for $F_jD(B) \cdot g^s$ and $F_jD(C) \cdot (f \cdot g)^s$ respectively. Furthermore, we define $\mathbb{C}[s]$ -modules

$$M_i := I_i \cdot f^{-i+1} \subseteq A_f[s],$$

$$N_j := J_j \cdot g^{-j+1} \subseteq B_g[s],$$

$$\Lambda_l := L_l \cdot (f \cdot g)^{-l+1} \subseteq C_{f \cdot g}[s].$$

Since $I_j \supseteq I_{j-1} \cdot f$, we see $M_j \supseteq M_{j-1}$. Similarly, we have $N_j \supseteq N_{j-1}$ and $L_l \supseteq N_{l-1}$. By definition, we obtain a way to characterize the Bernstein-Sato polynomials as

$$b_f(s) \cdot \mathbb{C}[s] = \bigcup_{i=1}^{\infty} (\mathbb{C}[s] \cap M_i),$$
$$b_g(s) \cdot \mathbb{C}[s] = \bigcup_{j=1}^{\infty} (\mathbb{C}[s] \cap N_j),$$
$$b_{f \cdot g}(s) \cdot \mathbb{C}[s] = \bigcup_{l=1}^{\infty} (\mathbb{C}[s] \cap \Lambda_l).$$

It is worth mentioning that the intersections happen in $A_f[s]$, $B_g[s]$, and $C_{f \cdot g}[s]$, where $\mathbb{C}[s]$ can be naturally embedded. The identities tell us that for all $i, j, l \gg 0$, we have

$$M_i \cap \mathbb{C}[s] = b_f(s) \cdot \mathbb{C}[s], \ N_i \cap \mathbb{C}[s] = b_g(s) \cdot \mathbb{C}[s], \ \Lambda_l \cap \mathbb{C}[s] = b_{f \cdot g}(s) \cdot \mathbb{C}[s].$$

Since $\operatorname{Der}_{\mathbb{C}} C = (\operatorname{Der}_{\mathbb{C}}(A) \otimes_{\mathbb{C}} B) \oplus (A \otimes_{\mathbb{C}} \operatorname{Der}_{\mathbb{C}}(B))$ i.e. derivations of C are induced by those of A and B. Since derivations of A and B commutes, we have the following identity:

$$F_l D(C) = \sum_{i+j=l} F_i D(A) \cdot F_j D(B).$$

Here we identify $F_iD(A)$ and $F_jD(B)$ with their images under canonical embeddings $F_iD_A \hookrightarrow F_iD_C$ and $F_jD_B \hookrightarrow F_jD_C$. As a result, we have a subsequent identity for I_i, J_j and L_l as below:

$$L_l = \sum_{i+j=l} f^{l-i} \cdot I_i \cdot g^{l-j} \cdot J_j$$

Dividing $(f \cdot g)^{l-1}$ on both sides, we have

$$\Lambda_l = \sum_{i+j=l} M_i \cdot N_j \subseteq M_l \cdot N_l \subseteq C_{f \cdot g}[s].$$

Take $l \in \mathbb{N}$ sufficiently large, such that


$$\Lambda_l \cap \mathbb{C}[s] = b_{f \cdot g}(s) \cdot \mathbb{C}[s], \ M_l \cap \mathbb{C}[s] = b_f(s) \cdot \mathbb{C}[s], \ N_l \cap \mathbb{C}[s] = b_g(s) \cdot \mathbb{C}[s].$$

It suffices to prove $(M_l \cdot N_l) \cap \mathbb{C}[s] = b_f(s)b_g(s) \cdot \mathbb{C}[s]$. As a result, we have $\Lambda_l \cap [s] \subseteq b_f(s)b_g(s) \cdot \mathbb{C}[s]$ i.e. $b_f(s)b_g(s)$ divides $b_{f \cdot g}(s)$. Now let us focus on the category of $\mathbb{C}[s]$ -modules. The following two diagrams are Cartesian products.

$$b_f(s) \cdot \mathbb{C}[s] \longrightarrow \mathbb{C}[s]$$

$$\downarrow \qquad \qquad \downarrow$$

$$M_l \longrightarrow A_f[s]$$



All of the eight arrows are $\mathbb{C}[s]$ -module injections. Tensoring ($\mathbb{C}[s]$ -module tensor) the two squares, we have a commutative diagram.

A simple observation shows $M_l, N_l, A_f[s]$ and $B_g[s]$ are all $\mathbb{C}[s]$ -free modules. In fact, $A_f[s] = A_f \otimes_{\mathbb{C}} \mathbb{C}[s]$ is a linear space tensored with $\mathbb{C}[s]$, and hence free. Moreover, since $\mathbb{C}[s]$ is a PID and M_l is a $\mathbb{C}[s]$ -submodule of $A_f[s]$, M_l is also free. Similarly, N_l and $B_g[s]$ are both free. Consequently, one deduce the four arrows in the diagram above are all injections.

Under the canonical isomorphism $A_f[s] \otimes_{\mathbb{C}[s]} B_g[s] \simeq C_{f \cdot g}[s]$, the diagram above is identified with the following.

Therefore, to prove **Theorem 3.2**, we only need to show (3.2.1) is a Cartesian product.

Applying **Lemma 3.1**, we have:

 $(b_f(s) \cdot \mathbb{C}[s]) \otimes_{\mathbb{C}[s]} (b_g(s) \cdot \mathbb{C}[s]) = (M_l \otimes_{\mathbb{C}[s]} N_l) \cap (\mathbb{C}[s] \otimes_{\mathbb{C}[s]} \mathbb{C}[s]) \cap (M_l \otimes_{\mathbb{C}[s]} \mathbb{C}[s]) \cap (\mathbb{C}[s] \otimes_{\mathbb{C}[s]} N_l)$ Since all arrows in (3.2.1) are injective, it suffices to show

$$(M_l \otimes_{\mathbb{C}[s]} N_l) \cap (\mathbb{C}[s] \otimes_{\mathbb{C}[s]} \mathbb{C}[s]) \subseteq M_l \otimes_{\mathbb{C}[s]} \mathbb{C}[s].$$

$$(3.2.2)$$

And similarly, the corresponding containing holds for N_l . As a result (3.2.1) is a Cartesian product.

To prove (3.2.2), we use the canonical isomorphism $A_f[s] \otimes_{\mathbb{C}[s]} B_g[s] \simeq C_{f \cdot g}[s]$ to pass everything to $C_{f \cdot g}[s]$. It is equivalent to show

$$(M_l \cdot N_l) \cap \mathbb{C}[s] \subseteq M_l.$$

Let $\sum_{i} a_i \cdot f^{-l+1} \cdot b_i \cdot g^{-l+1} = c$ be an element in the left-hand side, where $a_i \in I_l, b_i \in J_l$, and $c(s) \in \mathbb{C}[s]$. That is,

$$\sum_{i} a_i b_i = c(s) (f \cdot g)^{l-1}.$$
(3.2.3)

The equality happens in C[s].

We will localize B to show $c(s) \in M_l$. Let $\mathfrak{m}_y \subset B$ be a maximal ideal such that $g \in \mathfrak{m}_y$ and $y = (y_1, ..., y_m)$ be a set of local coordinates of B at \mathfrak{m}_y . Let $\tilde{B} = B_{\mathfrak{m}_y}$ be the localization and we still write \mathfrak{m}_y the maximal ideal of \tilde{B} . Tensoring A and $\mathbb{C}[s]$ to the natural embedding $B \hookrightarrow \tilde{B}$, we have $C[s] \hookrightarrow (A \otimes_{\mathbb{C}} \tilde{B}) \otimes_{\mathbb{C}} \mathbb{C}[s] = (A \otimes_{\mathbb{C}} \tilde{B})[s]$. Suppose the order of g in \tilde{B} is t > 0i.e. $g \in \mathfrak{m}_y^t$ but $g \notin \mathfrak{m}_y^{t-1}$. Since B is a regular local ring, we have

$$\operatorname{gr}_{\mathfrak{m}_{\boldsymbol{y}}} \tilde{B} = \bigoplus_{j \ge 0} \mathfrak{m}_{\boldsymbol{y}}^{j} / \mathfrak{m}_{\boldsymbol{y}}^{j+1}.$$

is isomorphic to the polynomial ring of *m*-variables, with indeterminants being images of \boldsymbol{y} in $\operatorname{gr}_{\mathfrak{m}_{\boldsymbol{y}}}\tilde{B}$. Suppose \boldsymbol{y}^{α} appears in g i.e. in the image of g in $\tilde{B}/\mathfrak{m}_{\boldsymbol{y}}^{t+1} = \bigoplus_{0 \leq j \leq t} \mathfrak{m}_{\boldsymbol{y}}^{j}/\mathfrak{m}_{\boldsymbol{y}}^{j+1}$. We may assume the coefficient of \boldsymbol{y}^{α} is 1.

Let us treat (3.2.3) in $(A \otimes_k \tilde{B})[s]$. Moduling $\mathfrak{m}_{\boldsymbol{y}}^{(l-1)t+1}$, we know the identity also holds in $A[s] \otimes_{\mathbb{C}[s]} (\tilde{B}/\mathfrak{m}_{\boldsymbol{y}}^{(l-1)t+1})[s]$. Since $(\tilde{B}/\mathfrak{m}_{\boldsymbol{y}}^{(l-1)t+1})[s]$ is a $\mathbb{C}[s]$ -free module, we can still compare the coefficient of $\boldsymbol{y}^{(l-1)\alpha}$ on both sides (coefficients in A[s]). On one hand, the coefficient of $\boldsymbol{y}^{(l-1)\alpha}$ in the right-hand side is $c(s)f^{l-1}$. On the other hand, suppose the coefficient of $\boldsymbol{y}^{(l-1)\alpha}$ in b_i is $\tilde{b}_i(s) \in \mathbb{C}[s]$. Hence, we have $\sum_i \tilde{b}_i(s)a_i = c(s)f^{l-1}$. Consequently, $c(s) = f^{-l+1} \cdot (\sum_i \tilde{b}_i(s)a_i) \in M_l$. We are done.

This theorem immediately implies the following.

Corollary 3.4. Suppose $f \in \mathbb{Z}[\mathbf{x}]$ and $g \in \mathbb{Z}[\mathbf{y}]$ are non-zero polynomials. Suppose strong monodromy conjecture (Conjecture 2.11) holds for f and g, then it also holds for $f \cdot g \in \mathbb{Z}[\mathbf{x}, \mathbf{y}]$.

As another corollary, we can generalize the theorem for affine varieties to a global theorem. We first define the tensor of divisors, as a generalization of $f \cdot g$.

Definition 3.5. Suppose X_1, X_2 are non-singular complex varieties and D_1, D_2 are effective Cartier divisors on X_1, X_2 . We define $D_1 \otimes D_2$ to be an effective Cartier divisor on $X_1 \times X_2$, as follows. Take affine coverings $\{U_i\}, \{V_j\}$ of X_1 and X_2 respectively, such that D_1 is represented by f_i on U_i and D_2 is represented by g_j on V_j . Define $D_1 \otimes D_2$ to be the Cartier divisor represented by $f_i \otimes g_j$ on each $U_i \times V_j$.

Theorem 3.6. Suppose X_1 and X_2 non-singular complex varieties and D_1, D_2 are effective Cartier divisors on X_1, X_2 , then we have

$$b_{D_1 \otimes D_2}(s) = b_{D_1}(s) \cdot b_{D_2}(s).$$

Proof. As in **Definition 3.5**, take open affine coverings $\{U_i\}, \{V_i\}$. By **Theorem 3.2**, we have

$$b_{D_1 \otimes D_2|_{U_i \times V_j}}(s) = b_{D_1|_{U_i}}(s) \cdot b_{D_2|_{V_j}}(s).$$

Since $X_1 \times V_j = \bigcup_i U_i \times V_j$, we have

$$b_{D_1 \otimes D_2|_{X \times V_j}}(s) = \lim_i b_{D_1|_{U_i}}(s) \cdot b_{D_2|_{V_j}}(s) = [\lim_i b_{D_1|_{U_i}}(s)] \cdot b_{D_2|_{V_j}}(s) = b_{D_1}(s) \cdot b_{D_2|_{V_j}}(s).$$

Furthermore, by $X_1 \times X_2 = \bigcup_i X_1 \times V_j$, we have

$$b_{D_1 \otimes D_2}(s) = \lim_{j} b_{D_1}(s) \cdot b_{D_2|_{V_j}}(s) = b_{D_1}(s) \cdot [\lim_{j} b_{D_2|_{V_j}}(s)] = b_{D_1}(s) \cdot b_{D_2}(s).$$

So we are done.

3.3. Generalization to Ideals. Having done the problem about the product of two regular functions, it is natural to ask the following.

Question 3.7. For $\mathfrak{a} \subseteq \mathbb{C}[x]$ and $\mathfrak{b} \subseteq \mathbb{C}[y]$, is $b_{\mathfrak{a} \cdot \mathfrak{b}}(s) = b_{\mathfrak{a}}(s) \cdot b_{\mathfrak{b}}(s)$?

The general answer is no. Here is a counter-example.

Example 3.8. Consider $\mathfrak{a} = (x^2, y^7) \subseteq \mathbb{C}[x, y]$ and $\mathfrak{b} = (z^{14}, w) \subset \mathbb{C}[z, w]$. By **Example 2.16**, we have

$$W_{\mathfrak{a}} = \{\frac{7i+2j}{14} \mid 1 \le i \le 2, 1 \le j \le 7\},\$$
$$W_{\mathfrak{b}} = \{1 + \frac{j}{14} \mid 1 \le j \le 14\}.$$

By **Theorem 2.14**, considering the product of the two diagonals, we have $\frac{15}{7}$ is contained in W_{ab} .

But, if \mathfrak{a} or \mathfrak{b} is principal, we have the following.

Proposition 3.9. Let $\mathfrak{a} \subseteq \mathbb{C}[\mathbf{x}]$ be an ideal and $g \in \mathbb{C}[\mathbf{y}]$, then $b_{\mathfrak{a}}b_g = b_{\mathfrak{a} \cdot (g)}$.

Proof. By Theorem 2.12, $b_{\mathfrak{a}} = b_h/(s+1)$, where $h = \sum f_i z_i$, z are indeterminants. Again by Theorem 2.12, $b_{\mathfrak{a}}(g) = b_{h'}/(s+1)$, where $h' = \sum f_i g z_i$. Applying Theorem 3.2, we get $b_{h'} = b_g b_h$, hence $b_{\mathfrak{a}}(g) = b_{\mathfrak{a}} b_g$.

If we neglect the multiplicity of roots, there are still some interesting things. Let W_n be the roots of b_n for an ideal \mathfrak{n} contained in some polynomial ring. We find the following.

Theorem 3.10. Let $\mathfrak{a} \subseteq \mathbb{C}[x]$ and $\mathfrak{b} \subseteq \mathbb{C}[y]$ be non-zero monomial ideals, then

 $W_{\mathfrak{a}} \cup W_{\mathfrak{b}} \subseteq W_{\mathfrak{ab}}.$

Despite the sets of roots are not equal totally, it is surprising that they are equal when modulo \mathbb{Z} . More precisely, we have the following.

Theorem 3.11. Let $\mathfrak{a} \subseteq \mathbb{C}[x]$ and $\mathfrak{b} \subseteq \mathbb{C}[y]$ be non-zero monomial ideals, then

 $W_{\mathfrak{a}} \cup W_{\mathfrak{b}} = W_{\mathfrak{a}\mathfrak{b}} \mod \mathbb{Z}.$

We need some preparations before proving the theorem.

For the convenience of further discussion, we shall first develop some properties of Newton polyhedron of the product ideal of two monomial ideals. The ideals considered are all non-trivial.

Let $\mathfrak{a} \triangleleft \mathbb{C}[x_1, x_2, ..., x_n], \mathfrak{b} \triangleleft \mathbb{C}[y_1, y_2, ..., y_m]$ be monomial ideals. Suppose

$$\mathbf{a} = (\mathbf{x}^{\mathbf{v}_i}), \mathbf{v}_i = (v_i^1, ..., v_i^n) \in \mathbb{N}^n, i = 1, 2, ..., r, \text{ and}$$

 $\mathbf{b} = (\mathbf{y}^{\mathbf{w}_i}), \mathbf{w}_i = (w_i^1, ..., w_i^m), i = 1, 2, ..., s.$

We have the following lemma.

Lemma 3.12. (a) $\Gamma_{\mathfrak{ab}} = \Gamma_{\mathfrak{a}} \times \Gamma_{\mathfrak{b}}$.

(b) $P_{\mathfrak{ab}} = P_{\mathfrak{a}} \times P_{\mathfrak{b}}.$

(c) Faces of $P_{\mathfrak{ab}}$ are exactly those of $Q_1 \times Q_2$, where $Q_1(resp. Q_2)$ is a face of $P_{\mathfrak{a}}$ (resp. $P_{\mathfrak{b}}$). (d) For face Q_1 of $P_{\mathfrak{a}}$ and Q_2 of $P_{\mathfrak{b}}$, we have

$$M_{Q_1 \times Q_2} \setminus M'_{Q_1 \times Q_2} = [(M_{Q_1} \setminus M'_{Q_1}) \times M_{Q_2}] \cup [M_{Q_1} \times (M_{Q_2} \setminus M'_{Q_2})].$$

Proof. We only prove (c) since (a)(b) are trivial and (d) is merely a calculation for sets.

It suffices to consider proper faces. Let H be a hyperplane such that one of the corresponding half spaces contains P_{ab} . Suppose H is defined by the equation:

$$\lambda_1 x_1 + \dots \lambda_n x_n + \mu_1 y_1 + \dots + \mu_m y_m - C = 0.$$

It is easy to find that all $\lambda_i, \mu_j, C \in \mathbb{Q}$ have the same sign since there exists positive integer R sufficiently large s.t. $(R, R, ..., R) + \mathbb{Z}_{\geq 0} \subset P_{\mathfrak{ab}}$ and difference of signs among $\{\lambda_i, \mu_j\}$ implies that $P_{\mathfrak{ab}}$ does not lies in one side of H, a contradiction. We may assume $\lambda_i, \mu_j > 0$ for all i, j, then the nonemptiness of $H \cap P_{\mathfrak{ab}}$ implies C > 0. Since the minimum of $X = \lambda_1 x_1 + ... \lambda_n x_n$ and $Y = \mu_1 y_1 + ... + \mu_m y_m$ can be attained in $P_{\mathfrak{a}}$ and $P_{\mathfrak{b}}$ respectively, denoted by C_1 and C_2 , then we have $C = C_1 + C_2$ and $H \cap P_{\mathfrak{ab}} = H_1 \times H_2$, where H_1 (resp. H_2) is the face determined by $X - C_1$ (resp. $Y - C_2$).

The following is the proof of **Theorem 3.10**.

Proof. For any proper face Q of $P_{\mathfrak{a}}$ which is not contained in any coordinate hyperplane, $Q \times P_{\mathfrak{b}}$ is a proper face of $P_{\mathfrak{a}\mathfrak{b}}$. It is not contained in any coordinate hyperplane as well. Let L_Q be a linear function on \mathbb{R}^n s.t. $L_Q|_Q \equiv 1$, then L_Q is naturally extended to a linear function whose restriction on $Q \times P_{\mathfrak{b}}$ is 1. One can see $M_{Q \times P_{\mathfrak{b}}} = (M_Q \setminus M'_Q) \times \mathbb{Z}^m$ by **Lemma 3.12** and the simple fact that $M_{P_{\mathfrak{a}}} = \mathbb{Z}^m$. Since the linear span of $Q \times P_{\mathfrak{b}}$ is $V_Q \times \mathbb{R}^m$, we have $R_{Q \times P_{\mathfrak{b}}} = R_Q$. Therefore, by **Theorem 2.14** roots of $b_{\mathfrak{a}}$ are contained in those of $b_{\mathfrak{a}}$. Symmetrically, roots of $b_{\mathfrak{b}}$ are also contained in $W_{\mathfrak{a}\mathfrak{b}}$.

Remark 3.13. Through the preceding proof, we can see b_{ab} can have roots other than those of b_a and b_b since it has faces other than those of the form $Q \times P_b$ and $P_a \times Q$. This is why we construct **Example 3.8**.

The following is the proof of **Theorem 3.11**.

Proof. By Theorem 3.11, it suffices to show all facets of $P_{\mathfrak{a}} \times P_{\mathfrak{b}}$ are exactly those of the form $Q_1 \times P_{\mathfrak{b}}$ or $P_{\mathfrak{a}} \times Q_2$, where Q_1, Q_2 are facets since we have shown in the proof of Theorem 3.10 that $R_{Q_1 \times P_{\mathfrak{a}}} = R_{Q_1}$ and $R_{P_{\mathfrak{b}} \times Q_2} = R_{Q_2}$.

Suppose $Q_1 \subseteq P_{\mathfrak{a}}$ and $Q_2 \subseteq P_{\mathfrak{b}}$ are both proper faces. By the following lemma, we have

 $\dim V_{Q_1 \times Q_2} \le \dim V_{Q_1} + \dim V_{Q_2} - 1 < n + m.$

Therefore, the dimension of $V_{Q_1 \times Q_2}$ is n + m - 1. If $Q_1 \times Q_2$ is a facet, it must pass through the origin. However, this tells us that $Q_1 \times Q_2$ is contained in some coordinate hyperplane. Otherwise, one can see the dimension of the linear span of $\mathbb{R}^{m+n}_{\geq 0} \cap Q_1 \times Q_2 < m + n - 1$. So we are done.

Lemma 3.14. For affine hyperplanes $Q_1 \in \mathbb{R}^n$ and $Q_2 \in \mathbb{R}^m$, then the dimension of the linear space generated by $Q_1 \times Q_2$ is not greater than n + m - 1.

Proof. We may assume both Q_1 and Q_2 do not pass through the origin, since otherwise we can replace the total space with Q_i . Let $Q_i = Q'_i + w_i$, where Q'_i passes through the origin and $w_i \in Q_i$. Let $u_1, u_2, ..., u_{n-1}$ be a basis of Q'_1 and $v_1, v_2, ..., v_{m-1}$ be a basis of Q'_2 . Then every point in Q_1 can be written uniquely in the form $\sum a_i u_i$ with $\sum a_i = 1$. A similar result holds for Q_2 . One can easily deduce that $Q_1 \times Q_2$ is contained in the linear space generated by $(u_i, v_j)_{1 \le i \le n-1, 1 \le j \le m-1}$. Furthermore, the following is a basis of this space:

$$(u_1, v_1) (u_2, v_1) \dots (u_{n-1}, v_1) (u_{n-1}, v_2) \dots (u_{n-1}, v_{m-1}).$$

Hence we have the result.

Taking both **Proposition 3.9**, **Theorem 3.10**, and **Theorem 3.11** into consideration, we hence make the following conjecture.

Conjecture 3.15. Let $\mathfrak{a} \subseteq \mathbb{C}[x]$ and $\mathfrak{b} \subseteq \mathbb{C}[y]$ be non-zero ideals, then

$$W_{\mathfrak{a}} \cup W_{\mathfrak{b}} \subseteq W_{\mathfrak{a}\mathfrak{b}}$$

Moreover, we have

$$W_{\mathfrak{a}} \cup W_{\mathfrak{b}} = W_{\mathfrak{a}\mathfrak{b}} \mod \mathbb{Z}.$$

References

- [AMHJNBTW22] J. Àlvarez Montaner, D. Hernández, J. Jeffries, L. Núñez Betancourt, P. Teixeira, and E. E. Witt. Bernstein-sato functional equations, v-filtrations, and multiplier ideals of direct summands. Communications in Contemporary Mathematics, 24(10):2150083, 2022.
- [Ber71] I. N. Bernštein. Modules over a ring of differential operators. An investigation of the fundamental solutions of equations with constant coefficients. *Funkcional. Anal. i Priložen.*, 5(2):1–16, 1971.

- [BMS06a] N. Budur, M. Mustață, and M. Saito. Roots of Bernstein-Sato polynomials for monomial ideals: a positive characteristic approach. *Math. Res. Lett.*, 13(1):125–142, 2006.
- [BMS06b] N. Budur, M. Mustaţă, and M. Saito. Bernstein-Sato polynomials of arbitrary varieties. Compos. Math., 142(3):779–797, 2006.
- [BMS06c] N. Budur, M. Mustaţă, and M. Saito. Combinatorial description of the roots of the Bernstein-Sato polynomials for monomial ideals. Comm. Algebra, 34(11):4103–4117, 2006.
- [Bud12] Nero Budur. Singularity invariants related to Milnor fibers: survey. In Zeta functions in algebra and geometry, volume 566 of Contemp. Math., pages 161–187. Amer. Math. Soc., Providence, RI, 2012.
- [DL92] J. Denef and F. Loeser. Caractéristiques d'Euler-Poincaré, fonctions zêta locales et modifications analytiques. J. Amer. Math. Soc., 5(4):705–720, 1992.
- [ELSV04] L. Ein, R. Lazarsfeld, K. E. Smith, and D. Varolin. Jumping coefficients of multiplier ideals. Duke Math. J., 123(3):469–506, 2004.
- [GLS07] G.-M. Greuel, C. Lossen, and E. Shustin. Introduction to singularities and deformations. Springer Monographs in Mathematics. Springer, Berlin, 2007.
- [Gra10] M. Granger. Bernstein-Sato polynomials and functional equations. In Algebraic approach to differential equations, pages 225–291. World Sci. Publ., Hackensack, NJ, 2010.
- [HTT08] R. Hotta, K. Takeuchi, and T. Tanisaki. D-modules, perverse sheaves, and representation theory, volume 236 of Progress in Mathematics. Birkhäuser Boston, Inc., Boston, MA, japanese edition, 2008.
- [Igu75] J. Igusa. Complex powers and asymptotic expansions. II. Asymptotic expansions. J. Reine Angew. Math., 278/279:307–321, 1975.
- [Kas77] M. Kashiwara. B-functions and holonomic systems. Rationality of roots of B-functions. Invent. Math., 38(1):33–53, 1976/77.
- [Kol97] J. Kollár. Singularities of pairs. In Algebraic geometry—Santa Cruz 1995, volume 62, Part 1 of Proc. Sympos. Pure Math., pages 221–287. Amer. Math. Soc., Providence, RI, 1997.
- [KS11] Sándor J. Kovács and Karl E. Schwede. Hodge theory meets the minimal model program: a survey of log canonical and Du Bois singularities. In *Topology of stratified spaces*, volume 58 of *Math. Sci. Res. Inst. Publ.*, pages 51–94. Cambridge Univ. Press, Cambridge, 2011.
- [Laz04] R. Lazarsfeld. Positivity in algebraic geometry. I, volume 48 of Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]. Springer-Verlag, Berlin, 2004. Classical setting: line bundles and linear series.
- [Lee24] Jonghyun Lee. Multiplicative thom-sebastiani for bernstein-sato polynomials, 2024.
- [Mus22] M. Mustaţă. Bernstein-Sato polynomials for general ideals vs. principal ideals. Proc. Amer. Math. Soc., 150(9):3655–3662, 2022.
- [Nic10] J. Nicaise. An introduction to p-adic and motivic zeta functions and the monodromy conjecture. In Algebraic and analytic aspects of zeta functions and L-functions, volume 21 of MSJ Mem., pages 141–166. Math. Soc. Japan, Tokyo, 2010.
- [Pop18] M. Popa. D-modules in birational geometry. https://people.math.harvard.edu/~mpopa/notes/DMBG-posted.pdf, 2018.
- [Sai93] M. Saito. On b-function, spectrum and rational singularity. Math. Ann., 295(1):51–74, 1993.
- [Sai06] M. Saito. Introduction to a theory of b-functions. https://arxiv.org/abs/math/0610783v1, 2006.
- [Sai09] M. Saito. On the Hodge filtration of Hodge modules. Mosc. Math. J., 9(1):161–191, back matter, 2009.
- [Sai16] M. Saito. Bernstein-Sato polynomials of hyperplane arrangements. Selecta Math. (N.S.), 22(4):2017–2057, 2016.
- [Sch19] C. Schnell. Lecture notes on algebraic d-modules (stonybrook 2019). https://www.math.stonybrook.edu/~cschnell/pdf/notes/d-modules.pdf, 2019.
- [Var81] A. N. Varčenko. Asymptotic Hodge structure on vanishing cohomology. Izv. Akad. Nauk SSSR Ser. Mat., 45(3):540–591, 688, 1981.
- [vS20] D. van Straten. The spectrum of hypersurface singularities. https://arxiv.org/abs/2003.00519, 2020.
- [Wal15] R. Walters. An application of D-modules: The Bernstein-Sato polynomial. ProQuest LLC, Ann Arbor, MI, 2015. Thesis (Ph.D.)–The University of Chicago.
- [Wei67] A. Weil. Basic number theory, volume Band 144 of Die Grundlehren der mathematischen Wissenschaften. Springer-Verlag New York, Inc., New York, 1967.

ZHILI COLLEGE, TSINGHUA UNIVERSITY, BEIJING, 100084, P. R. CHINA. *Email address:* shiq20@mails.tsinghua.edu.cn

DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, BEIJING, 100084, P. R. CHINA. *Email address:* hqzuo@mail.tsinghua.edu.cn