

# SYSTOLIC INEQUALITIES AND THE HOROWITZ-MYERS CONJECTURE

SIMON BRENDLE AND PEI-KEN HUNG

ABSTRACT. Let  $3 \leq n \leq 7$ , and let  $g$  be a Riemannian metric on  $B^2 \times T^{n-2}$  with scalar curvature at least  $-n(n-1)$ . We establish an inequality relating the systole of the boundary to the infimum of the mean curvature on the boundary. As a consequence, we confirm a conjecture of Horowitz and Myers.

## 1. INTRODUCTION

Our goal in this paper is to prove the following geometric inequality for two-dimensional surfaces.

**Theorem 1.1.** *Let us fix an integer  $n \geq 3$ . Let  $\Sigma$  be a compact, connected, orientable surface with boundary  $\partial\Sigma \neq \emptyset$ , and let  $g$  be a Riemannian metric on  $\Sigma$ . We denote by  $K$  the Gaussian curvature of  $\Sigma$ , by  $\kappa$  the geodesic curvature of the boundary  $\partial\Sigma$ , and by  $\eta$  the outward-pointing unit normal vector field to  $\partial\Sigma$ . Let  $\psi$  be a smooth function on  $\Sigma$  such that*

$$-2\Delta\psi - \frac{n-1}{n-2}|\nabla\psi|^2 + n(n-1) + 2K \geq 0$$

at each point in  $\Sigma$ . If  $\Sigma$  is diffeomorphic to  $B^2$ , then

$$2|\partial\Sigma|^n \inf_{\partial\Sigma} (\langle \nabla\psi, \eta \rangle + \kappa - (n-1)) \leq \left(\frac{4\pi}{n}\right)^n.$$

If  $\Sigma$  is not diffeomorphic to  $B^2$ , then

$$\inf_{\partial\Sigma} (\langle \nabla\psi, \eta \rangle + \kappa - (n-1)) \leq 0.$$

The proof of Theorem 1.1 consists of two parts. If  $\Sigma$  is diffeomorphic to  $B^2$ , the proof relies on a monotonicity formula (see Corollary 2.10 below). This argument shares some common features with the proof of the fill radius estimate in the groundbreaking work of Gromov and Lawson (see [10], Section 10). If  $\Sigma$  is not diffeomorphic to  $B^2$ , we minimize a weighted length functional and apply the stability inequality. The latter argument is inspired by work of Schoen and Yau [16].

---

The authors are grateful to Professors Nick Edelen, Gerhard Huisken, Christos Mantoulidis, and Leon Simon for discussions. The first author was supported by the National Science Foundation under grant DMS-2403981 and by the Simons Foundation. He acknowledges the hospitality of Tübingen University and CRM Montréal, where part of this work was carried out.

In higher dimensions, we prove the following estimate.

**Theorem 1.2.** *Let  $3 \leq n \leq 7$ . Let us consider the manifold  $M = B^2 \times T^{n-2}$ . Let  $\Xi$  denote the pull-back of the volume form on  $S^1$  under the projection*

$$\partial M = S^1 \times T^{n-2} \rightarrow S^1, (\xi, \theta_1, \dots, \theta_{n-2}) \mapsto \xi.$$

*Note that  $\Xi$  is a closed one-form on  $\partial M$ . Let  $g$  be a Riemannian metric on  $M$  with scalar curvature at least  $-n(n-1)$ . Then*

$$2\sigma^n \inf_{\partial M} (H_{\partial M} - (n-1)) \leq \left(\frac{4\pi}{n}\right)^n,$$

*where  $\sigma$  denotes the length of the shortest closed curve  $\alpha$  in  $(\partial M, g)$  satisfying  $\int_{\alpha} \Xi \neq 0$ .*

To deduce Theorem 1.2 from Theorem 1.1, we construct a minimal slicing with free boundary. The minimal slicing technique was pioneered in the fundamental work of Schoen and Yau [15]. This argument is closely related to the torical symmetrization procedure described in Section 12 of Gromov and Lawson's paper [10] (see also [7] and [9]).

As a consequence of Theorem 1.2, we obtain the following result, which confirms a conjecture of Horowitz and Myers [13].

**Theorem 1.3.** *Let  $3 \leq n \leq 7$ . Let us fix a flat metric  $\gamma$  on the torus  $S^1 \times T^{n-2}$ . Let  $\Xi$  denote the pull-back of the volume form on  $S^1$  under the projection*

$$S^1 \times T^{n-2} \rightarrow S^1, (\xi, \theta_1, \dots, \theta_{n-2}) \mapsto \xi.$$

*Note that  $\Xi$  is a closed one-form on  $S^1 \times T^{n-2}$ . Let  $Q$  be a symmetric  $(0, 2)$ -tensor on  $S^1 \times T^{n-2}$ . We fix a real number  $r_0 > 0$  and consider the hyperbolic metric*

$$\bar{g} = r^{-2} dr \otimes dr + r^2 \gamma$$

*on  $(r_0, \infty) \times S^1 \times T^{n-2}$ . Let  $M = \mathbb{R}^2 \times T^{n-2}$ . We identify  $(r_0, \infty) \times S^1 \times T^{n-2}$  with a subset of  $M$  via the embedding*

$$(r_0, \infty) \times S^1 \times T^{n-2} \rightarrow M, (r, \xi, \theta_1, \dots, \theta_{n-2}) \mapsto (r\xi, \theta_1, \dots, \theta_{n-2}).$$

*Let  $g$  be a Riemannian metric on  $M$  with scalar curvature at least  $-n(n-1)$ . Moreover, suppose that  $g$  satisfies*

$$|g - \bar{g} - r^{2-n} Q|_{\bar{g}} \leq o(r^{-n})$$

*and*

$$|\bar{D}(g - \bar{g} - r^{2-n} Q)|_{\bar{g}} \leq o(r^{-n})$$

*as  $r \rightarrow \infty$ . Then*

$$\int_{S^1 \times T^{n-2}} \left( n \operatorname{tr}_{\gamma}(Q) + \left(\frac{4\pi}{n\sigma}\right)^n \right) d\operatorname{vol}_{\gamma} \geq 0,$$

*where  $\sigma$  denotes the length of the shortest closed curve  $\alpha$  in  $(S^1 \times T^{n-2}, \gamma)$  satisfying  $\int_{\alpha} \Xi \neq 0$ .*

The inequality in Theorem 1.3 is sharp for the Horowitz-Myers metrics. For a discussion of the background of the Horowitz-Myers conjecture, we refer to [1], [3], [4], [8], [13], [14], and [20].

**Remark 1.4.** The proof of Theorem 1.3 does not actually use the assumption that  $\gamma$  is flat. If  $\gamma$  is a non-flat metric on  $S^1 \times T^{n-2}$ , then, by the solution of Geroch's conjecture, there exists a point on  $S^1 \times T^{n-2}$  where the scalar curvature of  $\gamma$  is strictly negative. The scalar curvature of the metric  $\bar{g}$  is related to the scalar curvature of  $\gamma$  by the formula  $R_{\bar{g}} = -n(n-1) + r^{-2} R_{\gamma}$ . Hence, if  $\gamma$  is a non-flat metric on  $S^1 \times T^{n-2}$ , then there are points where the scalar curvature of  $\bar{g}$  is strictly less than  $-n(n-1)$ .

## 2. PROOF OF THEOREM 1.1 – THE CASE WHEN $\Sigma$ IS DIFFEOMORPHIC TO $B^2$

In this section, we prove Theorem 1.1 in the special case when  $\Sigma$  is diffeomorphic to  $B^2$ . We define a function  $u : \Sigma \rightarrow [0, \infty)$  by  $u(x) = d(x, \partial\Sigma)$  for  $x \in \Sigma$ . Moreover, we define a function  $\rho : \partial\Sigma \rightarrow (0, \infty)$  by

$$\rho(x) = \sup\{t > 0 : \exp_x(-t\eta(x)) \text{ is defined and } u(\exp_x(-t\eta(x))) = t\}$$

for each point  $x \in \partial\Sigma$ . Let

$$D = \{(x, t) \in \partial\Sigma \times [0, \infty) : t \in [0, \rho(x)]\}$$

and

$$D^0 = \{(x, t) \in \partial\Sigma \times [0, \infty) : t \in [0, \rho(x))\}.$$

Clearly,  $D$  is a compact subset of  $\partial\Sigma \times [0, \infty)$ . Moreover, it is well known that  $D^0$  is a relatively open subset of  $\partial\Sigma \times [0, \infty)$ . Putting these facts together, it follows that  $\rho$  is a continuous function (see also [18], Proposition 4.2.1).

We define a map  $\Phi : D \rightarrow \Sigma$  by

$$\Phi(x, t) = \exp_x(-t\eta(x)).$$

Note that  $\Phi$  is surjective. Moreover, the restriction  $\Phi|_{D^0}$  is injective. Indeed, if  $(x, t) \in D^0$  and  $(\tilde{x}, \tilde{t}) \in D$  satisfy  $\Phi(x, t) = \Phi(\tilde{x}, \tilde{t})$ , then  $(x, t) = (\tilde{x}, \tilde{t})$ . Finally, with a suitable choice of orientation, we have  $\det(D\Phi)_{(x,t)} > 0$  for each point  $(x, t) \in D^0$ .

Let  $l = \sup_{y \in \Sigma} u(y)$ . For each  $s \in (0, l)$ , we denote by  $A(s)$  the area of the tubular neighborhood  $\{y \in \Sigma : u(y) \leq s\}$ . We may write

$$A(s) = \int_{\partial\Sigma} \left( \int_0^{\min\{\rho(x), s\}} \det D\Phi(x, t) dt \right) d\text{vol}(x)$$

for each  $s \in (0, l)$ . Moreover, we define

$$L(s) = \int_{\partial\Sigma} 1_{\{\rho(x) \geq s\}} \det D\Phi(x, s) d\text{vol}(x)$$

for each  $s \in (0, l)$ . Note that the function  $s \mapsto L(s)$  is not necessarily continuous.

**Lemma 2.1.** *We can find a large constant  $C$  such that the function  $s \mapsto A(s) - Cs$  is monotone decreasing for  $s \in (0, l)$ . Moreover,  $|\Sigma| - A(s) \leq C(l - s)$  for each  $s \in (0, l)$ .*

**Proof.** Let us fix a large constant  $C$  such that  $\det(D\Phi)_{(x,t)} \leq C$  for each point  $x \in \partial\Sigma$  and each  $t \in [0, \rho(x)]$ . This implies that the function  $s \mapsto A(s) - C|\partial\Sigma|s$  is monotone decreasing for  $s \in (0, l)$ . Moreover, using the identity

$$|\Sigma| = \int_{\partial\Sigma} \left( \int_0^{\rho(x)} \det D\Phi(x, t) dt \right) d\text{vol}(x),$$

we obtain

$$|\Sigma| - A(s) = \int_{\partial\Sigma} \left( \int_{\min\{\rho(x), s\}}^{\rho(x)} \det D\Phi(x, t) dt \right) d\text{vol}(x)$$

for each  $s \in (0, l)$ . Since  $\rho(x) \leq l$  for all  $x \in \partial\Sigma$ , it follows that  $|\Sigma| - A(s) \leq C|\partial\Sigma|(l - s)$  for each  $s \in (0, l)$ . This completes the proof of Lemma 2.1.

**Lemma 2.2.** *The function  $s \mapsto L(s)$  is uniformly bounded from above for  $s \in (0, l)$ . Moreover, we can find a large constant  $C$  such that the function  $s \mapsto L(s) - Cs$  is monotone decreasing for  $s \in (0, l)$ .*

**Proof.** Standard results in comparison geometry imply that we can find a large constant  $C$  with the following property. For each point  $x \in \partial\Sigma$ , the function  $t \mapsto e^{-Ct} \det(D\Phi)_{(x,t)}$  is monotone decreasing for  $t \in [0, \rho(x)]$ . Consequently, the function  $s \mapsto e^{-Cs} L(s)$  is monotone decreasing for  $s \in (0, l)$ . From this, the assertion follows. This completes the proof of Lemma 2.2.

We define a function  $F : (1, \infty) \rightarrow (0, \infty)$  by

$$F(z) = \int_1^z \zeta^{-1} (1 - \zeta^{-n})^{-\frac{1}{2}} d\zeta$$

for each  $z \in (1, \infty)$ . Note that  $F$  is smooth and strictly monotone increasing. We denote by  $G : (0, \infty) \rightarrow (1, \infty)$  the inverse of  $F$ . Clearly,  $G$  is smooth and strictly monotone increasing. A straightforward calculation gives

$$\frac{d}{ds} G(s) = G(s) (1 - G(s)^{-n})^{\frac{1}{2}}$$

for each  $s \in (0, \infty)$ . This implies

$$(1) \quad \frac{d}{ds} (G(s)^{n-1}) = (n-1) G(s)^{n-1} (1 - G(s)^{-n})^{\frac{1}{2}}$$

and

$$(2) \quad \frac{d}{ds} ((1 - G(s)^{-n})^{\frac{1}{2}}) = \frac{n}{2} G(s)^{-n}$$

for each  $s \in (0, \infty)$ .

For each  $s \in (0, l)$ , we define

$$\Omega(s) = \{y \in \Sigma : u(y) > s\}.$$

For each  $s \in (0, l)$ ,  $\Omega(s)$  is a non-empty open subset of  $\Sigma$  with area  $|\Sigma| - A(s)$ . Finally, we define

$$I(s) = 2\pi - (n-1)(1 - G(l-s)^{-n})^{\frac{1}{2}} L(s) + \int_{\Omega(s)} (\Delta\psi - K)$$

and

$$J(s) = G(l-s)^{n-1} I(s)$$

for each  $s \in (0, l)$ .

**Lemma 2.3.** *The function  $s \mapsto |I(s)|$  is uniformly bounded from above for  $s \in (0, l)$ . Moreover, we can find a large constant  $C$  such that the function  $s \mapsto I(s) + Cs$  is monotone increasing for  $s \in (0, l)$ .*

**Proof.** Note that

$$|I(s)| \leq 2\pi + (n-1)L(s) + \int_{\Sigma} |\Delta\psi - K|$$

for all  $s \in (0, l)$ . Since the function  $s \mapsto L(s)$  is uniformly bounded from above for  $s \in (0, l)$ , it follows that the function  $s \mapsto |I(s)|$  is uniformly bounded from above for  $s \in (0, l)$ .

In view of Lemma 2.1 and Lemma 2.2, we can find a large constant  $C$  such that the functions  $s \mapsto A(s) - Cs$  and  $s \mapsto L(s) - Cs$  are monotone decreasing for  $s \in (0, l)$ . Since the function  $G$  is monotone increasing, we obtain

$$(1 - G(l-s_0)^{-n})^{\frac{1}{2}} - (1 - G(l-s_1)^{-n})^{\frac{1}{2}} \geq 0$$

for  $0 < s_0 < s_1 < l$ . This implies

$$\begin{aligned} I(s_1) - I(s_0) &= -(n-1)(1 - G(l-s_0)^{-n})^{\frac{1}{2}} (L(s_1) - L(s_0)) \\ &\quad + (n-1) \left( (1 - G(l-s_0)^{-n})^{\frac{1}{2}} - (1 - G(l-s_1)^{-n})^{\frac{1}{2}} \right) L(s_1) \\ &\quad - \int_{\Omega(s_0) \setminus \Omega(s_1)} (\Delta\psi - K) \\ &\geq -(n-1)(1 - G(l-s_0)^{-n})^{\frac{1}{2}} (L(s_1) - L(s_0)) \\ &\quad - \sup_{\Sigma} |\Delta\psi - K| (A(s_1) - A(s_0)) \\ &\geq -C(n-1)(s_1 - s_0) \\ &\quad - C \sup_{\Sigma} |\Delta\psi - K| (s_1 - s_0) \end{aligned}$$

for  $0 < s_0 < s_1 < l$ . This completes the proof of Lemma 2.3.

**Lemma 2.4.** *The function  $s \mapsto |J(s)|$  is uniformly bounded from above for  $s \in (0, l)$ . Moreover, we can find a large constant  $C$  such that the function  $s \mapsto J(s) + Cs$  is monotone increasing for  $s \in (0, l)$ .*

**Proof.** The first statement follows immediately from Lemma 2.3. To prove the second statement, let us fix a large constant  $C$  such that  $|I(s)| \leq C$  for all  $s \in (0, l)$  and the function  $s \mapsto I(s) + Cs$  is monotone increasing for  $s \in (0, l)$ . Using (1), we obtain

$$0 \leq G(l - s_0)^{n-1} - G(l - s_1)^{n-1} \leq (n-1)G(l)^{n-1}(s_1 - s_0)$$

for  $0 < s_0 < s_1 < l$ . This implies

$$\begin{aligned} & J(s_1) - J(s_0) \\ &= G(l - s_0)^{n-1}(I(s_1) - I(s_0)) - (G(l - s_0)^{n-1} - G(l - s_1)^{n-1})I(s_1) \\ &\geq -CG(l - s_0)^{n-1}(s_1 - s_0) - C(G(l - s_0)^{n-1} - G(l - s_1)^{n-1}) \\ &\geq -CnG(l)^{n-1}(s_1 - s_0) \end{aligned}$$

for  $0 < s_0 < s_1 < l$ . This completes the proof of Lemma 2.4.

The following result was proved by Fiala [6] in the real-analytic case and by Hartman [12] in the smooth case.

**Theorem 2.5** (F. Fiala [6]; P. Hartman [12]). *We can find a set  $\mathcal{E} \subset (0, l)$  of measure 0 with the following properties:*

- (i) *Suppose that  $s \in (0, l) \setminus \mathcal{E}$ . Moreover, suppose that  $x$  is a point in  $\partial\Sigma$  with  $\rho(x) \geq s$ . Then  $\det(D\Phi)_{(x,s)} \neq 0$ .*
- (ii) *Suppose that  $s \in (0, l) \setminus \mathcal{E}$ . Moreover, suppose that  $y$  is a point in  $\Sigma$  with  $u(y) = s$ . Then the set  $\{x \in \partial\Sigma : \rho(x) \geq s \text{ and } \Phi(x, s) = y\}$  consists of at most two elements.*
- (iii) *Suppose that  $s \in (0, l) \setminus \mathcal{E}$ . Then the set  $\{x \in \partial\Sigma : \rho(x) = s\}$  is finite.*
- (iv) *Suppose that  $s \in (0, l) \setminus \mathcal{E}$ . Then the domain  $\Omega(s)$  has piecewise smooth boundary. The length of the boundary  $\partial\Omega(s)$  is given by  $L(s)$ . Moreover,*

$$\limsup_{\delta \searrow 0} \frac{L(s + \delta) - L(s)}{\delta} \leq -\Lambda(s),$$

where  $\Lambda(s)$  denotes the total geodesic curvature of the boundary  $\partial\Omega(s)$  (including angle contributions).

**Proof.** These statements are proved in [18], Chapter 4. To be more specific, let  $\mathcal{E}$  denote the set of exceptional values defined in Definition 4.3.1 in [18]. By Lemma 4.3.6 in [18],  $\mathcal{E}$  is a set of measure zero. Properties (i) and (ii) follow directly from the definition of the set  $\mathcal{E}$ . Property (iii) follows from Lemma 4.4.1 in [18]. Finally, property (iv) follows from Theorem 4.4.1 in [18], keeping in mind that  $\tan \frac{\theta}{2} \geq \frac{\theta}{2}$  for  $\theta \in [0, \pi)$ .

**Lemma 2.6.** *For each  $s \in (0, l) \setminus \mathcal{E}$ , the domain  $\Omega(s)$  has Euler characteristic at least 1.*

**Proof.** Let us fix a real number  $s \in (0, l) \setminus \mathcal{E}$ . Since  $\partial\Sigma$  is connected, the complement  $\Sigma \setminus \Omega(s)$  is connected. Moreover,  $\Sigma \setminus \Omega(s)$  contains a collar neighborhood of  $\partial\Sigma$ . Since  $\Sigma$  is diffeomorphic to  $B^2$ , it follows that each connected component of  $\Omega(s)$  is simply connected. Thus, the Euler characteristic of  $\Omega(s)$  equals the number of connected components of  $\Omega(s)$ . This completes the proof of Lemma 2.6.

**Lemma 2.7.** *For each  $s \in (0, l) \setminus \mathcal{E}$ , we have*

$$2\pi - \int_{\Omega(s)} K \leq \Lambda(s)$$

and

$$\int_{\Omega(s)} \Delta\psi \leq \int_{\partial\Omega(s)} |\nabla\psi|.$$

As above,  $\Lambda(s)$  denotes the total geodesic curvature of the boundary  $\partial\Omega(s)$  (including angle contributions).

**Proof.** Let us fix a real number  $s \in (0, l) \setminus \mathcal{E}$ . By Lemma 2.6, the domain  $\Omega(s)$  has Euler characteristic at least 1. Hence, the first statement follows from the Gauss-Bonnet theorem. The second statement follows from the divergence theorem. This completes the proof of Lemma 2.7.

**Proposition 2.8.** *For each  $s \in (0, l) \setminus \mathcal{E}$ , we have*

$$\liminf_{\delta \searrow 0} \frac{I(s+\delta) - I(s)}{\delta} - (n-1)(1 - G(l-s)^{-n})^{\frac{1}{2}} I(s) \geq 0.$$

**Proof.** Let us fix a real number  $s \in (0, l) \setminus \mathcal{E}$ . By Theorem 2.5 (iv), we know that

$$\limsup_{\delta \searrow 0} \frac{L(s+\delta) - L(s)}{\delta} \leq -\Lambda(s).$$

Using (2), we obtain

$$\begin{aligned} \liminf_{\delta \searrow 0} \frac{I(s+\delta) - I(s)}{\delta} &\geq \frac{n(n-1)}{2} G(l-s)^{-n} L(s) \\ &\quad + (n-1)(1 - G(l-s)^{-n})^{\frac{1}{2}} \Lambda(s) \\ &\quad + \int_{\partial\Omega(s)} (-\Delta\psi + K). \end{aligned}$$

By assumption,

$$-\Delta\psi - \frac{n-1}{2(n-2)} |\nabla\psi|^2 + \frac{n(n-1)}{2} + K \geq 0.$$

This implies

$$\begin{aligned} \liminf_{\delta \searrow 0} \frac{I(s+\delta) - I(s)}{\delta} &\geq -\frac{n(n-1)}{2} (1 - G(l-s)^{-n}) L(s) \\ &\quad + (n-1) (1 - G(l-s)^{-n})^{\frac{1}{2}} \Lambda(s) \\ &\quad + \frac{n-1}{2(n-2)} \int_{\partial\Omega(s)} |\nabla\psi|^2. \end{aligned}$$

On the other hand, Lemma 2.7 gives

$$I(s) \leq -(n-1) (1 - G(l-s)^{-n})^{\frac{1}{2}} L(s) + \Lambda(s) + \int_{\partial\Omega(s)} |\nabla\psi|.$$

Putting these facts together, we conclude that

$$\begin{aligned} &\liminf_{\delta \searrow 0} \frac{I(s+\delta) - I(s)}{\delta} - (n-1) (1 - G(l-s)^{-n})^{\frac{1}{2}} I(s) \\ &\geq \frac{(n-2)(n-1)}{2} (1 - G(l-s)^{-n}) L(s) \\ &\quad - (n-1) (1 - G(l-s)^{-n})^{\frac{1}{2}} \int_{\partial\Omega(s)} |\nabla\psi| + \frac{n-1}{2(n-2)} \int_{\partial\Omega(s)} |\nabla\psi|^2 \\ &= \frac{n-1}{2(n-2)} \int_{\partial\Omega(s)} \left( (n-2) (1 - G(l-s)^{-n})^{\frac{1}{2}} - |\nabla\psi| \right)^2. \end{aligned}$$

The expression on the right hand side is nonnegative. This completes the proof of Proposition 2.8.

**Proposition 2.9.** *For each  $s \in (0, l) \setminus \mathcal{E}$ , we have*

$$\liminf_{\delta \searrow 0} \frac{J(s+\delta) - J(s)}{\delta} \geq 0.$$

**Proof.** Let us fix a real number  $s \in (0, l) \setminus \mathcal{E}$ . Using (1), we obtain

$$\begin{aligned} \liminf_{\delta \searrow 0} \frac{J(s+\delta) - J(s)}{\delta} &= G(l-s)^{n-1} \liminf_{\delta \searrow 0} \frac{I(s+\delta) - I(s)}{\delta} \\ &\quad - (n-1) G(l-s)^{n-1} (1 - G(l-s)^{-n})^{\frac{1}{2}} I(s), \end{aligned}$$

and the expression on the right hand side is nonnegative by Proposition 2.8. This completes the proof of Proposition 2.9.

**Corollary 2.10.** *The function  $s \mapsto J(s)$  is monotone increasing for  $s \in (0, l)$ .*

**Proof.** It follows from Lemma 2.4 and the monotone differentiation theorem that the function  $s \mapsto J(s)$  is differentiable almost everywhere and

$$J(s_1) - J(s_0) \geq \int_{s_0}^{s_1} J'(s) ds$$

for all  $0 < s_0 < s_1 < l$ . Moreover, since  $\mathcal{E}$  is a set of measure zero, Proposition 2.9 implies that  $J'(s) \geq 0$  almost everywhere. Putting these facts together, we conclude that  $J(s_1) - J(s_0) \geq 0$  for  $0 < s_0 < s_1 < l$ . This completes the proof of Corollary 2.10.

**Corollary 2.11.** *We have*

$$G(l)^{n-1} \left( - (n-1) (1 - G(l)^{-n})^{\frac{1}{2}} |\partial\Sigma| + \int_{\partial\Sigma} (\langle \nabla\psi, \eta \rangle + \kappa) \right) \leq 2\pi.$$

**Proof.** Using the Gauss-Bonnet theorem and the divergence theorem, we obtain

$$\begin{aligned} \liminf_{s \searrow 0} I(s) &= 2\pi - (n-1) (1 - G(l)^{-n})^{\frac{1}{2}} |\partial\Sigma| + \int_{\Sigma} \Delta\psi - \int_{\Sigma} K \\ &= -(n-1) (1 - G(l)^{-n})^{\frac{1}{2}} |\partial\Sigma| + \int_{\partial\Sigma} \langle \nabla\psi, \eta \rangle + \int_{\partial\Sigma} \kappa. \end{aligned}$$

This gives

$$\liminf_{s \searrow 0} J(s) = G(l)^{n-1} \left( - (n-1) (1 - G(l)^{-n})^{\frac{1}{2}} |\partial\Sigma| + \int_{\partial\Sigma} (\langle \nabla\psi, \eta \rangle + \kappa) \right).$$

On the other hand,

$$I(s) \leq 2\pi + \int_{\Omega(s)} |\Delta\psi - K| \leq 2\pi + \sup_{\Sigma} |\Delta\psi - K| (|\Sigma| - A(s))$$

for each  $s \in (0, l)$ . Moreover, Lemma 2.1 implies that  $|\Sigma| - A(s) \leq C(l - s)$  for each  $s \in (0, l)$ . Thus,

$$\limsup_{s \nearrow l} I(s) \leq 2\pi.$$

Since  $\lim_{s \nearrow l} G(l - s) = 1$ , we conclude that

$$\limsup_{s \nearrow l} J(s) \leq 2\pi.$$

Hence, the assertion follows from Corollary 2.10. This completes the proof of Corollary 2.11.

**Corollary 2.12.** *We have*

$$2|\partial\Sigma|^{n-1} \int_{\partial\Sigma} (\langle \nabla\psi, \eta \rangle + \kappa - (n-1)) \leq \left( \frac{4\pi}{n} \right)^n.$$

**Proof.** We may assume that

$$\int_{\partial\Sigma} (\langle \nabla\psi, \eta \rangle + \kappa - (n-1)) > 0,$$

for otherwise the assertion is trivial. Using Corollary 2.11, we obtain

$$G(l)^{n-1} \left( - 2(n-1) (1 - G(l)^{-n})^{\frac{1}{2}} |\partial\Sigma| + \int_{\partial\Sigma} (2\langle \nabla\psi, \eta \rangle + 2\kappa) \right) \leq 4\pi.$$

Using the inequality  $2(1-a)^{\frac{1}{2}} \leq 2-a$  for  $0 \leq a \leq 1$ , we deduce that

$$G(l)^{n-1} \left( (n-1)G(l)^{-n} |\partial\Sigma| + \int_{\partial\Sigma} (2\langle \nabla\psi, \eta \rangle + 2\kappa - 2(n-1)) \right) \leq 4\pi.$$

Using the inequality  $(n-1)a^n + b^n \geq na^{n-1}b$  for  $a, b \geq 0$ , we obtain

$$\begin{aligned} & (n-1)G(l)^{-n} |\partial\Sigma| + \int_{\partial\Sigma} (2\langle \nabla\psi, \eta \rangle + 2\kappa - 2(n-1)) \\ & \geq nG(l)^{-(n-1)} |\partial\Sigma|^{\frac{n-1}{n}} \left( \int_{\partial\Sigma} (2\langle \nabla\psi, \eta \rangle + 2\kappa - 2(n-1)) \right)^{\frac{1}{n}}. \end{aligned}$$

Putting these facts together, we conclude that

$$n |\partial\Sigma|^{\frac{n-1}{n}} \left( \int_{\partial\Sigma} (2\langle \nabla\psi, \eta \rangle + 2\kappa - 2(n-1)) \right)^{\frac{1}{n}} \leq 4\pi.$$

This completes the proof of Corollary 2.12.

Corollary 2.12 implies

$$2 |\partial\Sigma|^n \inf_{\partial\Sigma} (\langle \nabla\psi, \eta \rangle + \kappa - (n-1)) \leq \left( \frac{4\pi}{n} \right)^n,$$

as desired.

### 3. PROOF OF THEOREM 1.1 – THE CASE WHEN $\Sigma$ IS NOT DIFFEOMORPHIC TO $B^2$

In this section, we prove Theorem 1.1 in the special case when  $\Sigma$  is not diffeomorphic to  $B^2$ .

**Lemma 3.1.** *The relative homotopy group  $\pi_1(\Sigma, \partial\Sigma)$  is non-trivial.*

**Proof.** We argue by contradiction. Suppose that  $\pi_1(\Sigma, \partial\Sigma)$  is trivial. This implies that  $\partial\Sigma$  is connected, hence diffeomorphic to  $S^1$ . We glue  $\Sigma$  with the disk  $B^2$  along their common boundary. In this way, we produce a compact, connected, orientable surface  $\hat{\Sigma}$  without boundary. Since  $\pi_1(\Sigma, \partial\Sigma)$  is trivial, it follows that  $\hat{\Sigma}$  is simply connected. Thus,  $\hat{\Sigma}$  is diffeomorphic to  $S^2$ . Consequently,  $\Sigma$  is diffeomorphic to  $B^2$ , contrary to our assumption. This completes the proof of Lemma 3.1.

In view of Lemma 3.1, we can find a stable free boundary geodesic  $\Gamma$  in  $(\Sigma, e^{2\psi}g)$ . In particular,  $\partial\Gamma \subset \partial\Sigma$  and  $\Gamma$  meets  $\partial\Sigma$  orthogonally along  $\partial\Gamma$ . Let  $\gamma : [0, l] \rightarrow \Gamma$  be a unit speed parametrization of  $\Gamma$ . Let  $\nu(s) \in T_{\gamma(s)}\Sigma$  denote the unit normal at the point  $\gamma(s)$  and let  $H(s)$  denote the geodesic curvature at the point  $\gamma(s)$ . Then  $H(s) + \langle \nabla\psi|_{\gamma(s)}, \nu(s) \rangle = 0$  for each

$s \in [0, l]$ . Applying Theorem A.1 with  $\rho = e^\psi$  gives

$$\begin{aligned} & \int_0^l e^{\psi(\gamma(s))} \zeta'(s)^2 ds - \int_0^l e^{\psi(\gamma(s))} K(\gamma(s)) \zeta(s)^2 - \int_0^l e^{\psi(\gamma(s))} H(s)^2 \zeta(s)^2 \\ & + \int_0^l e^{\psi(\gamma(s))} (D_{\Sigma}^2 \psi)_{\gamma(s)}(\nu(s), \nu(s)) \zeta(s)^2 \\ & - e^{\psi(\gamma(0))} \kappa(\gamma(0)) \zeta(0)^2 - e^{\psi(\gamma(l))} \kappa(\gamma(l)) \zeta(l)^2 \geq 0 \end{aligned}$$

for every test function  $\zeta \in C^\infty([0, l])$ . We next consider the first eigenfunction of the stability operator. This gives a positive function  $v \in C^\infty([0, l])$  such that

$$(3) \quad -v''(s) - K(\gamma(s))v(s) - H(s)^2v(s) + (D_{\Sigma}^2 \psi)_{\gamma(s)}(\nu(s), \nu(s))v(s) - \langle \nabla \psi|_{\gamma(s)}, \gamma'(s) \rangle v'(s) = \lambda v(s)$$

for each  $s \in [0, l]$  with Neumann boundary conditions

$$-v'(0) = \kappa(\gamma(0))v(0)$$

and

$$v'(l) = \kappa(\gamma(l))v(l).$$

Here,  $\lambda$  is a nonnegative constant. We next define

$$w(s) = \psi(\gamma(s)) + \log v(s)$$

for each  $s \in [0, l]$ .

**Lemma 3.2.** *The function  $w$  satisfies*

$$-w''(s) - \frac{n}{2(n-1)} w'(s)^2 + \frac{n(n-1)}{2} \geq 0$$

for each  $s \in [0, l]$ .

**Proof.** Note that

$$(4) \quad -\frac{d^2}{ds^2} \psi(\gamma(s)) - H(s) \langle \nabla \psi|_{\gamma(s)}, \nu(s) \rangle + (D_{\Sigma}^2 \psi)_{\gamma(s)}(\gamma'(s), \gamma'(s)) = 0$$

for each  $s \in [0, l]$ . Using (3) and (4) together with the identity  $H(s) + \langle \nabla \psi|_{\gamma(s)}, \nu(s) \rangle = 0$ , we obtain

$$\begin{aligned} & -w''(s) - \frac{n}{2(n-1)} w'(s)^2 \\ & = -\Delta \psi(\gamma(s)) + K(\gamma(s)) - \frac{n-1}{2(n-2)} \langle \nabla \psi|_{\gamma(s)}, \gamma'(s) \rangle^2 \\ & + \frac{1}{2(n-2)(n-1)} \left( \langle \nabla \psi|_{\gamma(s)}, \gamma'(s) \rangle - (n-2)v(s)^{-1}v'(s) \right)^2 + \lambda \end{aligned}$$

for each  $s \in [0, l]$ . By assumption,

$$-\Delta \psi - \frac{n-1}{2(n-2)} |\nabla \psi|^2 + \frac{n(n-1)}{2} + K \geq 0$$

at each point in  $\Sigma$ . Moreover,  $\lambda$  is nonnegative. Putting these facts together, the assertion follows. This completes the proof of Lemma 3.2.

**Lemma 3.3.** *We have  $\min\{-w'(0), w'(l)\} \leq n - 1$ .*

**Proof.** If  $-w'(0) \leq n - 1$ , the assertion is clearly true. Suppose next that  $-w'(0) > n - 1$ . Using Lemma 3.2 and standard ODE arguments, we conclude that  $-w'(s) > n - 1$  for each  $s \in [0, l]$ . In particular,  $w'(l) < -(n - 1) < n - 1$ . This completes the proof of Lemma 3.3.

Finally, we observe that

$$-w'(0) = -\langle \nabla\psi|_{\gamma(0)}, \gamma'(0) \rangle + \kappa(\gamma(0))$$

and

$$w'(l) = \langle \nabla\psi|_{\gamma(l)}, \gamma'(l) \rangle + \kappa(\gamma(l)).$$

Using Lemma 3.3, we conclude that

$$\min \left\{ -\langle \nabla\psi|_{\gamma(0)}, \gamma'(0) \rangle + \kappa(\gamma(0)), \langle \nabla\psi|_{\gamma(l)}, \gamma'(l) \rangle + \kappa(\gamma(l)) \right\} \leq n - 1.$$

Thus,

$$\inf_{\partial\Sigma} (\langle \nabla\psi, \eta \rangle + \kappa) \leq n - 1,$$

as claimed.

#### 4. PROOF OF THEOREM 1.2

In this section, we discuss how Theorem 1.2 can be deduced from Theorem 1.1. Let  $\eta$  denote the outward-pointing unit normal vector field to  $\partial M$ . We denote by  $h_{\partial M}$  the second fundamental form of  $\partial M$  and by  $H_{\partial M}$  the mean curvature of  $\partial M$ . Throughout this section, we assume that

$$\inf_{\partial M} (H_{\partial M} - (n - 1)) > 0,$$

for otherwise the assertion is trivial.

For each  $k \in \{1, \dots, n - 2\}$ , we denote by  $\Theta_k$  the pull-back of the volume form on  $S^1$  under the projection

$$\partial M = S^1 \times T^{n-2} \rightarrow S^1, (\xi, \theta_1, \dots, \theta_{n-2}) \mapsto \theta_k.$$

For each  $k \in \{1, \dots, n - 2\}$ ,  $\Theta_k$  is a closed one-form on  $\partial M$ .

**Proposition 4.1.** *We can find a collection of compact, connected, orientable submanifolds  $\Sigma_k$ ,  $k \in \{0, \dots, n - 2\}$ , a collection of positive functions  $v_k \in C^\infty(\Sigma_k)$ ,  $k \in \{1, \dots, n - 2\}$ , and a collection of positive functions  $\rho_k \in C^\infty(\Sigma_k)$ ,  $k \in \{0, \dots, n - 2\}$  with the following properties.*

- (i)  $\Sigma_0 = M$  and  $\rho_0 = 1$ .
- (ii) For each  $k \in \{0, \dots, n - 2\}$ , we have  $\dim \Sigma_k = n - k$ .
- (iii) For each  $k \in \{1, \dots, n - 2\}$ ,  $\Sigma_k$  is a compact, connected, embedded, orientable hypersurface in  $\Sigma_{k-1}$  satisfying  $\partial\Sigma_k \subset \partial\Sigma_{k-1}$ . Moreover,  $\Sigma_k$  meets  $\partial\Sigma_{k-1}$  orthogonally along  $\partial\Sigma_k$ .

- (iv) For each  $k \in \{1, \dots, n-2\}$ , the outward-pointing unit normal vector field to  $\partial\Sigma_k$  in  $\Sigma_k$  equals  $\eta$ . Moreover, the second fundamental form of  $\partial\Sigma_k$  in  $\Sigma_k$  equals the restriction of  $h_{\partial M}$  to  $T(\partial\Sigma_k)$ .
- (v) For each  $k \in \{1, \dots, n-2\}$ ,  $\Sigma_k$  is a stable free boundary minimal hypersurface in  $(\Sigma_{k-1}, \rho_{k-1}^{\frac{2}{n-k}} g_{\Sigma_{k-1}})$ .
- (vi) For each  $k \in \{0, \dots, n-2\}$ , we have

$$\int_{\partial\Sigma_k} \Xi \wedge \Theta_{k+1} \wedge \dots \wedge \Theta_{n-2} \neq 0.$$

- (vii) For each  $k \in \{1, \dots, n-2\}$ , the function  $v_k \in C^\infty(\Sigma_k)$  satisfies

$$\begin{aligned} & -\Delta_{\Sigma_k} v_k - \text{Ric}_{\Sigma_{k-1}}(\nu_{\Sigma_k}, \nu_{\Sigma_k}) v_k - |h_{\Sigma_k}|^2 v_k \\ & + (D_{\Sigma_{k-1}}^2 \log \rho_{k-1})(\nu_{\Sigma_k}, \nu_{\Sigma_k}) v_k - \langle \nabla^{\Sigma_k} \log \rho_{k-1}, \nabla^{\Sigma_k} v_k \rangle = \lambda_k v_k \end{aligned}$$

on  $\Sigma_k$  with Neumann boundary condition

$$\langle \nabla^{\Sigma_k} v_k, \eta \rangle - h_{\partial M}(\nu_{\Sigma_k}, \nu_{\Sigma_k}) v_k = 0$$

on  $\partial\Sigma_k$ . Here,  $\lambda_k$  is a nonnegative constant.

- (viii) For each  $k \in \{1, \dots, n-2\}$ , the function  $\rho_k \in C^\infty(\Sigma_k)$  is given by  $\rho_k = \rho_{k-1}|_{\Sigma_k} \cdot v_k$ .

- (ix) For each  $k \in \{1, \dots, n-2\}$ , the normal derivative of  $\rho_k$  satisfies

$$\langle \nabla^{\Sigma_k} \rho_k, \eta \rangle - \sum_{j=1}^k h_{\partial M}(\nu_{\Sigma_j}, \nu_{\Sigma_j}) \rho_k = 0$$

at each point on  $\partial\Sigma_k$ .

- (x) For each  $k \in \{1, \dots, n-2\}$ , the manifold  $(\Sigma_k, \rho_k^{\frac{2}{n-k-1}} g_{\Sigma_k})$  has boundary mean curvature equal to  $\rho_k^{-\frac{1}{n-k-1}} H_{\partial M}$ . In particular, the manifold  $(\Sigma_k, \rho_k^{\frac{2}{n-k-1}} g_{\Sigma_k})$  has strictly mean convex boundary.

**Proof.** We argue by induction on  $k$ . For  $k = 0$ , we define  $\Sigma_0 = M$  and  $\rho_0 = 1$ . We now turn to the inductive step. Suppose that  $k \in \{1, \dots, n-2\}$ , and that we have constructed submanifolds  $\Sigma_0, \dots, \Sigma_{k-1}$ , positive functions  $v_1 \in C^\infty(\Sigma_1), \dots, v_{k-1} \in C^\infty(\Sigma_{k-1})$ , and positive functions  $\rho_0 \in C^\infty(\Sigma_0), \dots, \rho_{k-1} \in C^\infty(\Sigma_{k-1})$  satisfying the conditions (i)–(x) above. The inductive hypothesis implies

$$\int_{\partial\Sigma_{k-1}} \Xi \wedge \Theta_k \wedge \dots \wedge \Theta_{n-2} \neq 0.$$

By taking the intersection of  $\Sigma_{k-1}$  with a generic level set of the function  $\theta_k : M \rightarrow S^1$ , we obtain a compact, embedded, orientable hypersurface  $\tilde{\Sigma}_k$  in  $\Sigma_{k-1}$  with the property that

$$\int_{\partial\tilde{\Sigma}_k} \Xi \wedge \Theta_{k+1} \wedge \dots \wedge \Theta_{n-2} \neq 0.$$

Note that  $\tilde{\Sigma}_k$  may be disconnected, but this does not affect the subsequent arguments.

The inductive hypothesis implies that the manifold  $(\Sigma_{k-1}, \rho_{k-1}^{\frac{2}{n-k}} g_{\Sigma_{k-1}})$  has strictly mean convex boundary. By Theorem B.1, we can find a compact, connected, embedded, orientable hypersurface  $\Sigma_k$  in  $\Sigma_{k-1}$  with the following properties:

- The boundary  $\partial\Sigma_k$  is contained in  $\partial\Sigma_{k-1}$ . Moreover,  $\Sigma_k$  meets  $\partial\Sigma_{k-1}$  orthogonally along  $\partial\Sigma_k$ .
- The submanifold  $\Sigma_k$  is a stable free boundary minimal hypersurface in  $(\Sigma_{k-1}, \rho_{k-1}^{\frac{2}{n-k}} g_{\Sigma_{k-1}})$ .
- We have  $\int_{\partial\Sigma_k} \Xi \wedge \Theta_{k+1} \wedge \dots \wedge \Theta_{n-2} \neq 0$ .

In view of the inductive hypothesis, the outward-pointing unit normal vector field to  $\partial\Sigma_{k-1}$  in  $\Sigma_{k-1}$  equals  $\eta$ . Consequently, the outward-pointing unit normal vector field to  $\partial\Sigma_k$  in  $\Sigma_k$  equals  $\eta$ . From this, we deduce that the second fundamental form of  $\partial\Sigma_k$  in  $\Sigma_k$  equals the restriction of  $h_{\partial M}$  to  $T(\partial\Sigma_k)$ . In particular, the mean curvature of  $\partial\Sigma_k$  in  $\Sigma_k$  is given by

$$(5) \quad \text{tr}_{\partial\Sigma_k}(h_{\partial M}) = H_{\partial M} - \sum_{j=1}^k h_{\partial M}(\nu_{\Sigma_j}, \nu_{\Sigma_j}).$$

To summarize, we have shown that properties (ii)–(vi) hold for  $\Sigma_k$ .

The stability inequality implies that

$$\begin{aligned} & \int_{\Sigma_k} \rho_{k-1} |\nabla^{\Sigma_k} \zeta|^2 - \int_{\Sigma_k} \rho_{k-1} \text{Ric}_{\Sigma_{k-1}}(\nu_{\Sigma_k}, \nu_{\Sigma_k}) \zeta^2 - \int_{\Sigma_k} \rho_{k-1} |h_{\Sigma_k}|^2 \zeta^2 \\ & + \int_{\Sigma_k} \rho_{k-1} (D_{\Sigma_{k-1}}^2 \log \rho_{k-1})(\nu_{\Sigma_k}, \nu_{\Sigma_k}) \zeta^2 \\ & - \int_{\partial\Sigma_k} \rho_{k-1} h_{\partial\Sigma_{k-1}}(\nu_{\Sigma_k}, \nu_{\Sigma_k}) \zeta^2 \geq 0 \end{aligned}$$

for every test function  $\zeta \in C^\infty(\Sigma_k)$  (see Theorem A.1 below). Here,  $h_{\partial\Sigma_{k-1}}$  denotes the second fundamental form of  $\partial\Sigma_{k-1}$  in  $\Sigma_{k-1}$ . The inductive hypothesis implies that  $h_{\partial\Sigma_{k-1}}(\nu_{\Sigma_k}, \nu_{\Sigma_k}) = h_{\partial M}(\nu_{\Sigma_k}, \nu_{\Sigma_k})$ . Hence, if we define  $v_k \in C^\infty(\Sigma_k)$  to be a first eigenfunction of the stability operator, then property (vii) holds for  $v_k$ .

We next define the function  $\rho_k \in C^\infty(\Sigma_k)$  by  $\rho_k = \rho_{k-1}|_{\Sigma_k} \cdot v_k$ . Then property (viii) holds for  $\rho_k$ . Moreover, the inductive hypothesis implies

$$\langle \nabla^{\Sigma_{k-1}} \rho_{k-1}, \eta \rangle - \sum_{j=1}^{k-1} h_{\partial M}(\nu_{\Sigma_j}, \nu_{\Sigma_j}) \rho_{k-1} = 0$$

at each point on  $\partial\Sigma_{k-1}$ . Since

$$\langle \nabla^{\Sigma_k} v_k, \eta \rangle - h_{\partial M}(\nu_{\Sigma_k}, \nu_{\Sigma_k}) v_k = 0$$

at each point on  $\partial\Sigma_k$ , we conclude that

$$(6) \quad \langle \nabla^{\Sigma_k} \rho_k, \eta \rangle - \sum_{j=1}^k h_{\partial M}(\nu_{\Sigma_j}, \nu_{\Sigma_j}) \rho_k = 0$$

at each point on  $\partial\Sigma_k$ . Therefore, property (ix) holds for  $\rho_k$ .

Finally, combining (5) and (6), we conclude that the manifold  $(\Sigma_k, \rho_k^{\frac{2}{n-k-1}} g_{\Sigma_k})$  has boundary mean curvature

$$\rho_k^{-\frac{1}{n-k-1}} \left( H_{\partial M} - \sum_{j=1}^k h_{\partial M}(\nu_{\Sigma_j}, \nu_{\Sigma_j}) + \langle \nabla^{\Sigma_k} \log \rho_k, \eta \rangle \right) = \rho_k^{-\frac{1}{n-k-1}} H_{\partial M}.$$

Thus, property (x) holds for  $\Sigma_k$ . This completes the proof of Proposition 4.1.

The following identity plays a central role in the dimension reduction argument of Schoen and Yau [15],[17] (see also [10], Section 12).

**Proposition 4.2** (cf. R. Schoen, S.T. Yau [15],[17]). *For each  $k \in \{0, \dots, n-2\}$ , the scalar curvature of  $\Sigma_k$  satisfies*

$$\begin{aligned} & R_{\Sigma_k} - R_M - 2 \Delta_{\Sigma_k} \log \rho_k - |\nabla^{\Sigma_k} \log \rho_k|^2 \\ & - \sum_{j=1}^k |\nabla^{\Sigma_j} \log v_j|^2 - \sum_{j=1}^k |h_{\Sigma_j}|^2 = \sum_{j=1}^k \lambda_j \end{aligned}$$

at each point on  $\Sigma_k$ .

**Proof.** We include the details for the convenience of the reader. For  $k=0$ , the assertion is trivial. Suppose next that  $k \in \{1, \dots, n-2\}$  and the assertion is true for  $k-1$ . In other words,

$$(7) \quad \begin{aligned} & R_{\Sigma_{k-1}} - R_M - 2 \Delta_{\Sigma_{k-1}} \log \rho_{k-1} - |\nabla^{\Sigma_{k-1}} \log \rho_{k-1}|^2 \\ & - \sum_{j=1}^{k-1} |\nabla^{\Sigma_j} \log v_j|^2 - \sum_{j=1}^{k-1} |h_{\Sigma_j}|^2 = \sum_{j=1}^{k-1} \lambda_j \end{aligned}$$

at each point on  $\Sigma_{k-1}$ . Using the Gauss equations, we obtain

$$(8) \quad R_{\Sigma_k} - R_{\Sigma_{k-1}} + 2 \operatorname{Ric}_{\Sigma_{k-1}}(\nu_{\Sigma_k}, \nu_{\Sigma_k}) - H_{\Sigma_k}^2 + |h_{\Sigma_k}|^2 = 0$$

at each point on  $\Sigma_k$ . In view of property (vii) in Proposition 4.1, we know that

$$(9) \quad \begin{aligned} & -2 \Delta_{\Sigma_k} \log v_k - 2 |\nabla^{\Sigma_k} \log v_k|^2 - 2 \operatorname{Ric}_{\Sigma_{k-1}}(\nu_{\Sigma_k}, \nu_{\Sigma_k}) - 2 |h_{\Sigma_k}|^2 \\ & + 2 (D_{\Sigma_{k-1}}^2 \log \rho_{k-1})(\nu_{\Sigma_k}, \nu_{\Sigma_k}) - 2 \langle \nabla^{\Sigma_k} \log \rho_{k-1}, \nabla^{\Sigma_k} \log v_k \rangle = \lambda_k \end{aligned}$$

at each point on  $\Sigma_k$ . Moreover,

$$(10) \quad \begin{aligned} & 2 \Delta_{\Sigma_{k-1}} \log \rho_{k-1} - 2 (D_{\Sigma_{k-1}}^2 \log \rho_{k-1})(\nu_{\Sigma_k}, \nu_{\Sigma_k}) \\ & - 2 H_{\Sigma_k} \langle \nabla^{\Sigma_{k-1}} \log \rho_{k-1}, \nu_{\Sigma_k} \rangle - 2 \Delta_{\Sigma_k} \log \rho_{k-1} = 0 \end{aligned}$$

at each point on  $\Sigma_k$ . Finally,

$$(11) \quad \begin{aligned} & H_{\Sigma_k}^2 + 2 H_{\Sigma_k} \langle \nabla^{\Sigma_{k-1}} \log \rho_{k-1}, \nu_{\Sigma_k} \rangle \\ & + |\nabla^{\Sigma_{k-1}} \log \rho_{k-1}|^2 - |\nabla^{\Sigma_k} \log \rho_{k-1}|^2 \\ & = (H_{\Sigma_k} + \langle \nabla^{\Sigma_{k-1}} \log \rho_{k-1}, \nu_{\Sigma_k} \rangle)^2 = 0 \end{aligned}$$

at each point on  $\Sigma_k$ . In the next step, we add (7) – (11). This gives

$$\begin{aligned} & R_{\Sigma_k} - R_M - 2 \Delta_{\Sigma_k} \log \rho_{k-1} - 2 \Delta_{\Sigma_k} \log v_k \\ & - |\nabla^{\Sigma_k} \log \rho_{k-1}|^2 - 2 \langle \nabla^{\Sigma_k} \log \rho_{k-1}, \nabla^{\Sigma_k} \log v_k \rangle - |\nabla^{\Sigma_k} \log v_k|^2 \\ & - \sum_{j=1}^k |\nabla^{\Sigma_j} \log v_j|^2 - \sum_{j=1}^k |h_{\Sigma_j}|^2 = \sum_{j=1}^k \lambda_j \end{aligned}$$

at each point on  $\Sigma_k$ . Since  $\log \rho_k = \log \rho_{k-1} + \log v_k$  at each point on  $\Sigma_k$ , we conclude that

$$\begin{aligned} & R_{\Sigma_k} - R_M - 2 \Delta_{\Sigma_k} \log \rho_k - |\nabla^{\Sigma_k} \log \rho_k|^2 \\ & - \sum_{j=1}^k |\nabla^{\Sigma_j} \log v_j|^2 - \sum_{j=1}^k |h_{\Sigma_j}|^2 = \sum_{j=1}^k \lambda_j \end{aligned}$$

at each point on  $\Sigma_k$ . This completes the proof of Proposition 4.2.

**Corollary 4.3.** *For each  $k \in \{0, \dots, n-2\}$ , the scalar curvature of  $\Sigma_k$  satisfies*

$$R_{\Sigma_k} - R_M - 2 \Delta_{\Sigma_k} \log \rho_k - \frac{k+1}{k} |\nabla^{\Sigma_k} \log \rho_k|^2 \geq 0$$

at each point on  $\Sigma_k$ .

**Proof.** It follows from properties (i) and (viii) in Proposition 4.1 that  $\rho_k = \prod_{j=1}^k v_j$  at each point on  $\Sigma_k$ . This implies

$$\begin{aligned} \sum_{j=1}^k |\nabla^{\Sigma_j} \log v_j|^2 & \geq \sum_{j=1}^k |\nabla^{\Sigma_k} \log v_j|^2 \\ & \geq \frac{1}{k} \left| \sum_{j=1}^k \nabla^{\Sigma_k} \log v_j \right|^2 \\ & = \frac{1}{k} |\nabla^{\Sigma_k} \log \rho_k|^2 \end{aligned}$$

at each point on  $\Sigma_k$ . Moreover,  $\sum_{j=1}^k \lambda_j \geq 0$ . Hence, the assertion follows from Proposition 4.2. This completes the proof of Corollary 4.3.

After these preparations, we now complete the proof of Theorem 1.2. To that end, we consider the surface  $\Sigma := \Sigma_{n-2}$ . Moreover, we define a function  $\psi \in C^\infty(\Sigma)$  by  $\psi := \log \rho_{n-2}$ . It follows from property (vi) in Proposition

4.1 that  $\int_{\partial\Sigma} \Xi \neq 0$ . In the next step, we apply Corollary 4.3 with  $k = n - 2$ . Using the inequality  $R_M \geq -n(n - 1)$ , we conclude that

$$2K + n(n - 1) - 2\Delta\psi - \frac{n-1}{n-2} |\nabla\psi|^2 \geq 0$$

at each point on  $\Sigma$ . In view of property (iv) in Proposition 4.1, the geodesic curvature of  $\partial\Sigma$  in  $\Sigma$  is given by

$$(12) \quad \kappa = h_{\partial M}(\tau, \tau) = H_{\partial M} - \sum_{j=1}^{n-2} h_{\partial M}(\nu_{\Sigma_j}, \nu_{\Sigma_j}),$$

where  $\tau$  denotes a unit vector tangential to  $\partial\Sigma$ . By property (ix) in Proposition 4.1, the normal derivative of  $\psi$  is given by

$$(13) \quad \langle \nabla\psi, \eta \rangle = \sum_{j=1}^{n-2} h_{\partial M}(\nu_{\Sigma_j}, \nu_{\Sigma_j})$$

at each point on  $\partial\Sigma$ . Combining (12) and (13), we obtain

$$\kappa + \langle \nabla\psi, \eta \rangle = H_{\partial M}$$

at each point on  $\partial\Sigma$ . Applying Theorem 1.1, we conclude that

$$2|\partial\Sigma|^n \inf_{\partial\Sigma} (H_{\partial M} - (n-1)) \leq \left(\frac{4\pi}{n}\right)^n.$$

On the other hand, since  $\int_{\partial\Sigma} \Xi \neq 0$ , it follows that  $|\partial\Sigma| \geq \sigma$  by definition of  $\sigma$ . This completes the proof of Theorem 1.2.

## 5. PROOF OF THEOREM 1.3

In this section, we explain how Theorem 1.3 follows from Theorem 1.2. Let us fix a smooth function  $u : S^1 \times T^{n-2} \rightarrow \mathbb{R}$  and a constant  $\mu$  such that

$$(14) \quad \Delta_\gamma u + \frac{n}{2} \text{tr}_\gamma(Q) + \mu = 0$$

at each point on  $S^1 \times T^{n-2}$ . The function  $u$  is unique up to additive constants. We can make  $u$  unique by requiring that  $\int_{S^1 \times T^{n-2}} u \, d\text{vol}_\gamma = 0$ .

In the following, we assume that  $\hat{r}$  is chosen sufficiently large. We define  $\hat{M} = M \setminus \{r > \hat{r} + \hat{r}^{3-n} u\}$ . Note that  $\hat{M}$  is a compact subset of  $M$ , and  $\hat{M}$  is diffeomorphic to  $B^2 \times S^1$ .

**Lemma 5.1.** *The mean curvature of the boundary  $\partial\hat{M} = \{r = \hat{r} + \hat{r}^{3-n} u\}$  with respect to the metric  $g$  is given by  $(n-1) + \hat{r}^{-n} \mu + o(\hat{r}^{-n})$ .*

**Proof.** For abbreviation, we define

$$\hat{g} = \bar{g} + r^{2-n} Q = r^{-2} dr \otimes dr + r^2 \gamma + r^{2-n} Q.$$

Moreover, we consider the vector field  $V = r^2 \frac{\partial}{\partial r}$ . Then

$$\mathcal{L}_V(\bar{g}) - 2r \bar{g} = 0$$

and

$$\mathcal{L}_V(\hat{g}) - 2r \hat{g} + n r^{3-n} Q = 0.$$

Note that  $|V|_{\bar{g}} = r$  and  $|\bar{D}V|_{\bar{g}} = \frac{1}{2} |\mathcal{L}_V(\bar{g})|_{\bar{g}} \leq O(r)$ . Since  $|g - \hat{g}|_{\bar{g}} \leq o(r^{-n})$  and  $|\bar{D}(g - \hat{g})|_{\bar{g}} \leq o(r^{-n})$ , we obtain

$$|\mathcal{L}_V(g - \hat{g})|_{\bar{g}} \leq C |V|_{\bar{g}} |\bar{D}(g - \hat{g})|_{\bar{g}} + C |\bar{D}V|_{\bar{g}} |g - \hat{g}|_{\bar{g}} \leq o(r^{1-n}).$$

This implies

$$(15) \quad |\mathcal{L}_V(g) - 2r g + n r^{3-n} Q|_{\bar{g}} \leq o(r^{1-n}).$$

Let  $W$  denote the gradient of the function  $r$  with respect to the metric  $g$ , so that  $W^i = g^{ij} \partial_j r$ . Using the identity  $\partial_j r = \bar{g}_{jk} V^k = \hat{g}_{jk} V^k$ , we obtain

$$V^i - W^i = g^{ij} (g_{jk} - \hat{g}_{jk}) V^k.$$

Since  $|g - \hat{g}|_{\bar{g}} \leq o(r^{-n})$  and  $|\bar{D}(g - \hat{g})|_{\bar{g}} \leq o(r^{-n})$ , it follows that

$$|V - W|_{\bar{g}} \leq C |g - \hat{g}|_{\bar{g}} |V|_{\bar{g}} \leq o(r^{1-n})$$

and

$$\begin{aligned} & |\bar{D}(V - W)|_{\bar{g}} \\ & \leq C |\bar{D}(g - \hat{g})|_{\bar{g}} |V|_{\bar{g}} + C |\bar{D}g|_{\bar{g}} |g - \hat{g}|_{\bar{g}} |V|_{\bar{g}} + C |g - \hat{g}|_{\bar{g}} |\bar{D}V|_{\bar{g}} \leq o(r^{1-n}). \end{aligned}$$

This implies

$$|\mathcal{L}_{V-W}(g)|_{\bar{g}} \leq C |V - W|_{\bar{g}} |\bar{D}g|_{\bar{g}} + C |\bar{D}(V - W)|_{\bar{g}} |g|_{\bar{g}} \leq o(r^{1-n}).$$

Using (15), we conclude that

$$(16) \quad |\mathcal{L}_W(g) - 2r g + n r^{3-n} Q|_{\bar{g}} \leq o(r^{1-n}).$$

We next consider the hypersurface  $\partial \hat{M} = \{r = \hat{r} + \hat{r}^{3-n} u\}$ . We denote by  $\eta$  denote the outward-pointing unit normal vector field to  $\partial \hat{M}$  with respect to  $g$  and by  $H_{\partial \hat{M}}$  the mean curvature of  $\partial \hat{M}$  with respect to  $g$ . Then

$$(17) \quad H_{\partial \hat{M}} \langle W, \eta \rangle = \text{tr}_{\partial \hat{M}}(DW) - \text{div}_{\partial \hat{M}}(W^{\text{tan}}).$$

Using (16), we obtain

$$\begin{aligned} & \text{tr}_{\partial \hat{M}}(DW) \\ (18) \quad & = \frac{1}{2} \text{tr}_{\partial \hat{M}}(\mathcal{L}_W(g)) \\ & = (n-1) (\hat{r} + \hat{r}^{3-n} u) - \frac{n}{2} (\hat{r} + \hat{r}^{3-n} u)^{3-n} \text{tr}_{\partial \hat{M}}(Q) + o(\hat{r}^{1-n}). \end{aligned}$$

We next consider the map

$$\begin{aligned} \varphi : S^1 \times T^{n-2} & \rightarrow \partial \hat{M}, \\ (\xi, \theta_1, \dots, \theta_{n-2}) & \mapsto (\hat{r} + \hat{r}^{3-n} u(\xi, \theta_1, \dots, \theta_{n-2}), \xi, \theta_1, \dots, \theta_{n-2}). \end{aligned}$$

The pull-back of the metric  $\hat{g}$  under the map  $\varphi$  is given by

$$\varphi^*(\hat{g}) = (\hat{r} + \hat{r}^{3-n} u)^2 \gamma + (\hat{r} + \hat{r}^{3-n} u)^{2-n} Q + (\hat{r} + \hat{r}^{3-n} u)^{-2} \hat{r}^{6-2n} du \otimes du.$$

Since  $|g - \hat{g}|_{\hat{g}} \leq o(r^{-n})$ , it follows that

$$\varphi^*(g) = (\hat{r} + \hat{r}^{3-n} u)^2 \gamma + \chi,$$

where  $\chi$  is a symmetric  $(0, 2)$ -tensor on  $S^1 \times T^{n-2}$  satisfying  $|\chi|_{\gamma} \leq O(\hat{r}^{2-n})$ . This implies

$$(19) \quad \text{tr}_{\partial \hat{M}}(Q) = (\hat{r} + \hat{r}^{3-n} u)^{-2} \text{tr}_{\gamma}(Q) + O(\hat{r}^{-n-2}).$$

Substituting (19) into (18), we obtain

$$(20) \quad \text{tr}_{\partial \hat{M}}(DW) = (n-1) (\hat{r} + \hat{r}^{3-n} u) - \frac{n}{2} (\hat{r} + \hat{r}^{3-n} u)^{1-n} \text{tr}_{\gamma}(Q) + o(\hat{r}^{1-n}).$$

Moreover, since  $r = \hat{r} + \hat{r}^{3-n} u$  along  $\partial \hat{M}$ , we know that

$$(21) \quad \begin{aligned} \text{div}_{\partial \hat{M}}(W^{\text{tan}}) &= \Delta_{\partial \hat{M}} r \\ &= \hat{r}^{3-n} \Delta_{\partial \hat{M}} u \\ &= (\hat{r} + \hat{r}^{3-n} u)^{-2} \hat{r}^{3-n} \Delta_{\gamma} u + o(\hat{r}^{1-n}). \end{aligned}$$

Substituting (20) and (21) into (17), we obtain

$$\begin{aligned} H_{\partial \hat{M}} \langle W, \eta \rangle &= (n-1) (\hat{r} + \hat{r}^{3-n} u) - \frac{n}{2} (\hat{r} + \hat{r}^{3-n} u)^{1-n} \text{tr}_{\gamma}(Q) \\ &\quad - (\hat{r} + \hat{r}^{3-n} u)^{-2} \hat{r}^{3-n} \Delta_{\gamma} u + o(\hat{r}^{1-n}). \end{aligned}$$

In view of (14), we conclude that

$$(22) \quad H_{\partial \hat{M}} \langle W, \eta \rangle = (n-1) (\hat{r} + \hat{r}^{3-n} u) + (\hat{r} + \hat{r}^{3-n} u)^{1-n} \mu + o(\hat{r}^{1-n}).$$

It remains to compute the inner product  $\langle W, \eta \rangle$ . Note that

$$(23) \quad |W|^2 = |W|_{\hat{g}}^2 + o(\hat{r}^{2-n}) = (\hat{r} + \hat{r}^{3-n} u)^2 + o(\hat{r}^{2-n}).$$

Moreover, since  $r = \hat{r} + \hat{r}^{3-n} u$  along  $\partial \hat{M}$ , we obtain

$$(24) \quad |W^{\text{tan}}|^2 = |\nabla^{\partial \hat{M}} r|^2 = \hat{r}^{6-2n} |\nabla^{\partial \hat{M}} u|^2 = O(\hat{r}^{4-2n}).$$

Subtracting (24) from (23) gives

$$\langle W, \eta \rangle^2 = |W|^2 - |W^{\text{tan}}|^2 = (\hat{r} + \hat{r}^{3-n} u)^2 + o(\hat{r}^{2-n}),$$

hence

$$(25) \quad \langle W, \eta \rangle = (\hat{r} + \hat{r}^{3-n} u) + o(\hat{r}^{1-n}).$$

Combining (22) and (25), we conclude that

$$H_{\partial \hat{M}} = (n-1) + (\hat{r} + \hat{r}^{3-n} u)^{-n} \mu + o(\hat{r}^{-n}).$$

This completes the proof of Lemma 5.1.

After these preparations, we now describe the proof of Theorem 1.3. We argue by contradiction. Suppose that

$$\int_{S^1 \times T^{n-2}} \left( n \text{tr}_{\gamma}(Q) + \left( \frac{4\pi}{n\sigma} \right)^n \right) d\text{vol}_{\gamma} < 0.$$

Let us fix a real number  $\varepsilon \in (0, 1)$  such that

$$(26) \quad \int_{S^1 \times T^{n-2}} \left( n \operatorname{tr}_\gamma(Q) + (1 - \varepsilon)^{-n-1} \left( \frac{4\pi}{n\sigma} \right)^n \right) d\operatorname{vol}_\gamma \leq 0.$$

On the other hand, using (14) and the divergence theorem, we obtain

$$(27) \quad \int_{S^1 \times T^{n-2}} (n \operatorname{tr}_\gamma(Q) + 2\mu) d\operatorname{vol}_\gamma = 0.$$

Combining (26) and (27), we conclude that

$$(28) \quad 2(1 - \varepsilon)^{n+1} \sigma^n \mu \geq \left( \frac{4\pi}{n} \right)^n.$$

In particular,  $\mu$  is a positive real number.

If  $\hat{r}$  is sufficiently large (depending on  $\varepsilon$ ), then the mean curvature of  $\partial\hat{M}$  is bounded from below by  $(n - 1) + (1 - \varepsilon) \hat{r}^{-n} \mu$ . Using Theorem 1.2, we obtain

$$(29) \quad 2(1 - \varepsilon) \hat{\sigma}^n \hat{r}^{-n} \mu \leq \left( \frac{4\pi}{n} \right)^n,$$

where  $\hat{\sigma}$  denotes the length of the shortest closed curve  $\alpha$  in  $(\partial\hat{M}, g)$  satisfying  $\int_\alpha \Xi \neq 0$ . Combining (28) and (29), we conclude that

$$\hat{\sigma} \hat{r}^{-1} \leq (1 - \varepsilon) \sigma.$$

Since  $\varepsilon$  is independent of  $\hat{r}$ , this leads to a contradiction if  $\hat{r}$  is sufficiently large. This completes the proof of Theorem 1.3.

#### APPENDIX A. THE SECOND VARIATION FORMULA FOR WEIGHTED AREA

In this section, we derive the stability inequality for free boundary minimal hypersurfaces with respect to a conformally modified metric.

**Theorem A.1.** *Let  $M$  be a compact, orientable manifold of dimension  $n$  with boundary  $\partial M$ . Let  $g$  be a Riemannian metric on  $M$ , and let  $\rho$  be a smooth positive function on  $M$ . Suppose that  $\Sigma$  is an orientable hypersurface in  $M$  such that  $\partial\Sigma \subset \partial M$  and  $\Sigma$  meets  $\partial M$  orthogonally along  $\partial\Sigma$ . If  $\Sigma$  is a stable free boundary minimal hypersurface in  $(M, \rho^{\frac{2}{n-1}} g)$ , then*

$$\begin{aligned} & \int_\Sigma \rho |\nabla^\Sigma \zeta|^2 - \int_\Sigma \rho \operatorname{Ric}_M(\nu_\Sigma, \nu_\Sigma) \zeta^2 - \int_\Sigma \rho |h_\Sigma|^2 \zeta^2 \\ & + \int_\Sigma (D_M^2 \rho)(\nu_\Sigma, \nu_\Sigma) \zeta^2 - \int_\Sigma \rho^{-1} \langle \nabla^M \rho, \nu_\Sigma \rangle^2 \zeta^2 \\ & - \int_{\partial\Sigma} \rho h_{\partial M}(\nu_\Sigma, \nu_\Sigma) \zeta^2 \geq 0 \end{aligned}$$

for every test function  $\zeta \in C^\infty(\Sigma)$ .

**Proof.** We consider the conformal metric  $\tilde{g} = \rho^{\frac{2}{n-1}} g$ . Let  $\eta$  denote the outward-pointing unit normal vector field to  $\partial M$  with respect to  $g$ . The unit normal vector field to  $\partial M$  with respect to  $\tilde{g}$  is given by  $\tilde{\eta} = \rho^{-\frac{1}{n-1}} \eta$ , and the second fundamental form of  $\partial M$  with respect to  $\tilde{g}$  is given by

$$\tilde{h}_{\partial M} = \rho^{\frac{1}{n-1}} \left( h_{\partial M} + \frac{1}{n-1} \rho^{-1} d\rho(\eta) g \right).$$

The unit normal vector field to  $\Sigma$  with respect to  $\tilde{g}$  is given by  $\tilde{\nu}_\Sigma = \rho^{-\frac{1}{n-1}} \nu_\Sigma$ , and the second fundamental form of  $\Sigma$  with respect to  $\tilde{g}$  is given by

$$\tilde{h}_\Sigma = \rho^{\frac{1}{n-1}} \left( h_\Sigma + \frac{1}{n-1} \rho^{-1} d\rho(\nu_\Sigma) g \right).$$

Since  $\Sigma$  is a minimal hypersurface in  $(M, \tilde{g})$ , it follows that  $H_\Sigma + \rho^{-1} d\rho(\nu_\Sigma) = 0$ . This implies

$$|\tilde{h}_\Sigma|_{\tilde{g}}^2 = \rho^{-\frac{2}{n-1}} \left( |h_\Sigma|^2 - \frac{1}{n-1} \rho^{-2} d\rho(\nu_\Sigma)^2 \right).$$

The Ricci tensor of  $\tilde{g}$  is related to the Ricci tensor of  $g$  by the formula

$$\begin{aligned} \text{Ric}_{\tilde{g}} &= \text{Ric} - \left( 1 - \frac{1}{n-1} \right) \rho^{-1} D^2 \rho - \frac{1}{n-1} \rho^{-1} \Delta \rho g \\ &\quad + \left( 1 - \frac{1}{(n-1)^2} \right) \rho^{-2} d\rho \otimes d\rho + \frac{1}{(n-1)^2} \rho^{-2} |d\rho|^2 g \end{aligned}$$

(see [2], Theorem 1.159). Since  $\Sigma$  is a stable free boundary minimal hypersurface in  $(M, \tilde{g})$ , we know that

$$\begin{aligned} &\int_\Sigma |d\zeta|_{\tilde{g}}^2 d\text{vol}_{\tilde{g}} - \int_\Sigma \text{Ric}_{\tilde{g}}(\tilde{\nu}_\Sigma, \tilde{\nu}_\Sigma) \zeta^2 d\text{vol}_{\tilde{g}} - \int_\Sigma |\tilde{h}_\Sigma|_{\tilde{g}}^2 \zeta^2 d\text{vol}_{\tilde{g}} \\ &\quad - \int_{\partial\Sigma} \tilde{h}_{\partial M}(\tilde{\nu}_\Sigma, \tilde{\nu}_\Sigma) \zeta^2 d\text{vol}_{\tilde{g}} \geq 0 \end{aligned}$$

for every test function  $\zeta \in C^\infty(\Sigma)$ . In the next step, we replace  $\zeta$  by  $\rho^{\frac{1}{n-1}} \zeta$ . Moreover, we convert all the geometric quantities back to the metric  $g$ . This gives

$$\begin{aligned} &\int_\Sigma \rho^{1-\frac{2}{n-1}} |\nabla^\Sigma(\rho^{\frac{1}{n-1}} \zeta)|^2 - \int_\Sigma \rho \text{Ric}_M(\nu_\Sigma, \nu_\Sigma) \zeta^2 - \int_\Sigma \rho |h_\Sigma|^2 \zeta^2 \\ &\quad + \left( 1 - \frac{1}{n-1} \right) \int_\Sigma (D_M^2 \rho)(\nu_\Sigma, \nu_\Sigma) \zeta^2 + \frac{1}{n-1} \int_\Sigma \Delta_M \rho \zeta^2 \\ &\quad - \left( 1 - \frac{1}{n-1} - \frac{1}{(n-1)^2} \right) \int_\Sigma \rho^{-1} \langle \nabla^M \rho, \nu_\Sigma \rangle^2 \zeta^2 \\ &\quad - \frac{1}{(n-1)^2} \int_\Sigma \rho^{-1} |\nabla^M \rho|^2 \zeta^2 \\ &\quad - \int_{\partial\Sigma} \rho h_{\partial M}(\nu_\Sigma, \nu_\Sigma) \zeta^2 - \frac{1}{n-1} \int_{\partial\Sigma} \langle \nabla^\Sigma \rho, \eta \rangle \zeta^2 \geq 0 \end{aligned}$$

for every test function  $\zeta \in C^\infty(\Sigma)$ . Note that

$$\begin{aligned} & \int_{\Sigma} \rho^{1-\frac{2}{n-1}} |\nabla^\Sigma(\rho^{\frac{1}{n-1}} \zeta)|^2 \\ &= \int_{\Sigma} \rho |\nabla^\Sigma \zeta|^2 + \frac{2}{n-1} \int_{\Sigma} \langle \nabla^\Sigma \rho, \nabla^\Sigma \zeta \rangle \zeta + \frac{1}{(n-1)^2} \int_{\Sigma} \rho^{-1} |\nabla^\Sigma \rho|^2 \zeta^2 \end{aligned}$$

for every test function  $\zeta \in C^\infty(\Sigma)$ . Using the divergence theorem, we obtain

$$\int_{\partial\Sigma} \langle \nabla^\Sigma \rho, \eta \rangle \zeta^2 = \int_{\Sigma} \Delta_\Sigma \rho \zeta^2 + 2 \int_{\Sigma} \langle \nabla^\Sigma \rho, \nabla^\Sigma \zeta \rangle \zeta$$

for every test function  $\zeta \in C^\infty(\Sigma)$ . Putting these facts together, we conclude that

$$\begin{aligned} & \int_{\Sigma} \rho |\nabla^\Sigma \zeta|^2 - \int_{\Sigma} \rho \operatorname{Ric}_M(\nu_\Sigma, \nu_\Sigma) \zeta^2 - \int_{\Sigma} \rho |h_\Sigma|^2 \zeta^2 \\ &+ \left(1 - \frac{1}{n-1}\right) \int_{\Sigma} (D_M^2 \rho)(\nu_\Sigma, \nu_\Sigma) \zeta^2 + \frac{1}{n-1} \int_{\Sigma} (\Delta_M \rho - \Delta_\Sigma \rho) \zeta^2 \\ &- \left(1 - \frac{1}{n-1} - \frac{1}{(n-1)^2}\right) \int_{\Sigma} \rho^{-1} \langle \nabla^M \rho, \nu_\Sigma \rangle^2 \zeta^2 \\ &- \frac{1}{(n-1)^2} \int_{\Sigma} \rho^{-1} (|\nabla^M \rho|^2 - |\nabla^\Sigma \rho|^2) \zeta^2 \\ &- \int_{\partial\Sigma} \rho h_{\partial M}(\nu_\Sigma, \nu_\Sigma) \zeta^2 \geq 0 \end{aligned}$$

for every test function  $\zeta \in C^\infty(\Sigma)$ . The assertion now follows from the identities

$$\begin{aligned} \Delta_M \rho - \Delta_\Sigma \rho &= (D_M^2 \rho)(\nu_\Sigma, \nu_\Sigma) + H_\Sigma \langle \nabla^M \rho, \nu_\Sigma \rangle \\ &= (D_M^2 \rho)(\nu_\Sigma, \nu_\Sigma) - \rho^{-1} \langle \nabla^M \rho, \nu_\Sigma \rangle^2 \end{aligned}$$

and

$$|\nabla^M \rho|^2 - |\nabla^\Sigma \rho|^2 = \langle \nabla^M \rho, \nu_\Sigma \rangle^2.$$

This completes the proof of Theorem A.1.

## APPENDIX B. EXISTENCE AND REGULARITY OF FREE BOUNDARY MINIMAL HYPERSURFACES

In this section, we recall some well known results concerning the existence and regularity of free boundary minimal hypersurfaces.

**Theorem B.1** (cf. H. Federer [5]; M. Grüter [11]). *Let us fix an integer  $3 \leq n \leq 7$ . Let  $M$  be a compact, orientable manifold of dimension  $n$  with boundary  $\partial M$ . Let  $g$  be a Riemannian metric on  $M$ . We assume that the mean curvature of  $\partial M$  with respect to  $g$  is strictly positive. Let  $\Omega$  be a closed  $(n-2)$ -form on  $\partial M$ . Let  $\tilde{\Sigma}$  be a compact, embedded, orientable hypersurface in  $M$  such that  $\partial \tilde{\Sigma} \subset \partial M$  and  $\int_{\partial \tilde{\Sigma}} \Omega \neq 0$ . Then we can find a compact, connected, embedded, orientable hypersurface  $\Sigma$  with the following properties:*

- The boundary  $\partial\Sigma$  is contained in  $\partial M$ . Moreover,  $\Sigma$  meets  $\partial M$  orthogonally along  $\partial\Sigma$ .
- $\Sigma$  is a stable free boundary minimal hypersurface.
- $\int_{\partial\Sigma} \Omega \neq 0$ .

In the remainder of this section, we explain how Theorem B.1 follows from the results of Federer [5] and Grüter [11]. Let  $\hat{M}$  be a compact manifold which contains the given manifold  $M$  in its interior. We extend the given Riemannian metric  $g$  on  $M$  to a Riemannian metric on  $\hat{M}$ . If  $\varepsilon_0 > 0$  is sufficiently small, then the set  $\{x \in \hat{M} : d_{\hat{M}}(x, \partial M) \leq \varepsilon_0\}$  can be identified with  $\partial M \times [-\varepsilon_0, \varepsilon_0]$  via the normal exponential map. For  $\varepsilon \in (0, \varepsilon_0]$ , we define  $\hat{M}_\varepsilon = \{x \in \hat{M} : d_{\hat{M}}(x, M) \leq \varepsilon\}$ . If we choose  $\varepsilon_0 > 0$  sufficiently small, then the boundary  $\partial\hat{M}_\varepsilon$  is strictly mean convex for each  $\varepsilon \in (0, \varepsilon_0]$ . The nearest point projection gives a Lipschitz continuous map  $f : \hat{M}_{\varepsilon_0} \rightarrow M$  with  $f(x) = x$  for each point  $x \in M$ . For each  $\varepsilon \in (0, \varepsilon_0]$ , the restriction  $f|_{\partial\hat{M}_\varepsilon}$  is a volume-decreasing map from  $\partial\hat{M}_\varepsilon$  to  $\partial M$ .

**Lemma B.2.** *Let  $x \in \hat{M}_{\varepsilon_0} \setminus M$ . Then*

$$|df_x(v_1) \wedge \dots \wedge df_x(v_{n-1})| \leq |v_1 \wedge \dots \wedge v_{n-1}|$$

for all tangent vectors  $v_1, \dots, v_{n-1} \in T_x \hat{M}$ .

**Proof.** Let us fix a real number  $\varepsilon \in (0, \varepsilon_0]$ , a point  $x \in \partial\hat{M}_\varepsilon$ , and tangent vectors  $v_1, \dots, v_{n-1} \in T_x \hat{M}$ . For each  $1 \leq i \leq n-1$ , we denote by  $w_i$  the orthogonal projection of  $v_i$  to the tangent space  $T_x(\partial\hat{M}_\varepsilon)$ . Since the restriction  $f|_{\partial\hat{M}_\varepsilon}$  is a volume-decreasing map from  $\partial\hat{M}_\varepsilon$  to  $\partial M$ , we know that

$$|df_x(w_1) \wedge \dots \wedge df_x(w_{n-1})| \leq |w_1 \wedge \dots \wedge w_{n-1}|.$$

Since the matrix  $\{\langle v_i, v_j \rangle - \langle w_i, w_j \rangle\}_{1 \leq i, j \leq n-1}$  is weakly positive definite, we obtain

$$\det\{\langle w_i, w_j \rangle\}_{1 \leq i, j \leq n-1} \leq \det\{\langle v_i, v_j \rangle\}_{1 \leq i, j \leq n-1},$$

hence

$$|w_1 \wedge \dots \wedge w_{n-1}| \leq |v_1 \wedge \dots \wedge v_{n-1}|.$$

Finally, it follows from the definition of  $w_i$  that  $df_x(w_i) = df_x(v_i)$  for  $1 \leq i \leq n-1$ . This implies

$$|df_x(w_1) \wedge \dots \wedge df_x(w_{n-1})| = |df_x(v_1) \wedge \dots \wedge df_x(v_{n-1})|.$$

This completes the proof of Lemma B.2.

By the Nash embedding theorem, the manifold  $(\hat{M}, g)$  can be isometrically embedded into  $\mathbb{R}^N$  for some large integer  $N$ . We recall the definition of an integer multiplicity rectifiable current from Leon Simon's notes [19].

**Definition B.3** (cf. L. Simon [19], Definition 3.1 in Chapter 6). Let  $m$  be a positive integer, and let  $T$  be an  $m$ -dimensional current in  $\mathbb{R}^N$ . We

say that  $T$  is an integer multiplicity rectifiable current if there exist a  $\mathcal{H}^m$ -measurable countably  $m$ -rectifiable set  $A$ , a locally  $\mathcal{H}^m$ -integrable positive function  $\theta : A \rightarrow \mathbb{Z}$ , and an  $\mathcal{H}^m$ -measurable function  $\xi : A \rightarrow \Lambda^m(\mathbb{R}^N)$  with the following properties:

- For  $\mathcal{H}^m$ -a.e. point  $x \in A$ ,  $\xi(x)$  can be expressed in the form  $v_1 \wedge \dots \wedge v_m$ , where  $v_1, \dots, v_m$  form an orthonormal basis for the approximate tangent space  $T_x A$ .
- We have

$$T(\omega) = \int_A \langle \omega(x), \xi(x) \rangle \theta(x) d\mathcal{H}^m(x)$$

for every smooth  $m$ -form  $\omega$  with compact support.

Let  $\delta$  be a positive real number, and let  $U \subset \mathbb{R}^N$  denote the set of all points in  $\mathbb{R}^N$  that have distance less than  $\delta$  from  $M$ . If  $\delta > 0$  is chosen sufficiently small, then the nearest point projection gives a smooth map  $\pi : U \rightarrow \hat{M}_{\varepsilon_0}$ . We define a Lipschitz map  $F : U \rightarrow M$  by  $F = f \circ \pi$ .

**Lemma B.4.** *Suppose that  $T$  is an integer multiplicity rectifiable current with  $\text{supp}(T) \subset \hat{M}_{\varepsilon_0} \cap U$ . Then  $\mathbf{M}(F_{\#}(T)) \leq \mathbf{M}(T)$ .*

**Proof.** We use the notation of Definition B.3. Then

$$\mathbf{M}(T) = \int_A \theta(x) d\mathcal{H}^{n-1}(x).$$

Let  $J_A F$  denote the Jacobian determinant defined in §2 of Chapter in [19]. It follows from Lemma 3.9 in Chapter 6 of [19] that

$$\mathbf{M}(F_{\#}(T)) \leq \int_{F(A)} N(y) d\mathcal{H}^{n-1}(y),$$

where  $N(y)$  is defined by

$$N(y) = \sum_{x \in A, F(x)=y, J_A F(x) > 0} \theta(x)$$

for  $y \in F(A)$ . This implies

$$\mathbf{M}(F_{\#}(T)) \leq \int_A J_A F(x) \theta(x) d\mathcal{H}^{n-1}(x).$$

The assertion follows now from Lemma B.2. This completes the proof of Lemma B.4.

In the following, we recall some basic definitions from Federer's work [5]. Given a positive integer  $m$  and a compact set  $K \subset \mathbb{R}^N$ , let  $\mathcal{R}_{m,K}(\mathbb{R}^N)$  be defined as in [5], Section 4.1.24. It follows directly from the definition of  $\mathcal{R}_{m,K}(\mathbb{R}^N)$  that  $\text{supp}(T) \subset K$  for all  $T \in \mathcal{R}_{m,K}(\mathbb{R}^N)$ . For each positive integer  $m$ , we define

$$\mathcal{R}_m(\mathbb{R}^N) = \bigcup_K \mathcal{R}_{m,K}(\mathbb{R}^N),$$

where the union is taken over all compact sets  $K \subset \mathbb{R}^N$ . In particular, if  $T \in \mathcal{R}_m(\mathbb{R}^N)$ , then  $T$  has compact support.

**Proposition B.5** (cf. H. Federer [5]). *Let  $T$  be an  $m$ -dimensional current in  $\mathbb{R}^N$  with compact support. Then  $T \in \mathcal{R}_m(\mathbb{R}^N)$  if and only if  $T$  is an integer multiplicity rectifiable current in the sense of Definition B.3.*

**Proof.** This follows from Theorem 4.1.28 in [5]. Specifically, we use the equivalence of statements (1) and (4). This completes the proof of Proposition B.5.

For each positive integer  $m$ , the space of  $m$ -dimensional flat currents on  $\mathbb{R}^N$  is defined by

$$\mathcal{F}_m(\mathbb{R}^N) = \{P + \partial Q : P \in \mathcal{R}_m(\mathbb{R}^N), Q \in \mathcal{R}_{m+1}(\mathbb{R}^N)\}$$

(see [5], Section 4.1.24). Note that  $\mathcal{R}_m(\mathbb{R}^N) \subset \mathcal{F}_m(\mathbb{R}^N)$ . For each positive integer  $m$ , the space of  $m$ -dimensional integral flat cycles is defined by

$$\mathcal{Z}_m(M, \partial M) = \{T \in \mathcal{F}_m(\mathbb{R}^N) : \text{supp}(T) \subset M, \text{supp}(\partial T) \subset \partial M\}$$

(see [5], Section 4.4.1). For each positive integer  $m$ , the space of  $m$ -dimensional integral flat boundaries is defined by

$$\mathcal{B}_m(M, \partial M)$$

$$= \{P + \partial Q : P \in \mathcal{F}_m(\mathbb{R}^N), Q \in \mathcal{F}_{m+1}(\mathbb{R}^N), \text{supp}(P) \subset \partial M, \text{supp}(Q) \subset M\}$$

(see [5], Section 4.4.1). Note that  $\mathcal{B}_m(M, \partial M) \subset \mathcal{Z}_m(M, \partial M)$ .

We now apply Federer's results to our situation. By assumption,  $\Omega$  is a closed  $(n-2)$ -form on  $\partial M$ . We may extend  $\Omega$  to an  $(n-2)$ -form defined on  $\mathbb{R}^N$  such that  $\Omega$  has compact support and  $d\Omega$  vanishes in an open neighborhood of  $\partial M$ . The given hypersurface  $\tilde{\Sigma}$  defines a current  $\tilde{S} \in \mathcal{Z}_{n-1}(M, \partial M)$  such that  $\partial\tilde{S}(\Omega) \neq 0$ . Since  $d\Omega$  vanishes in an open neighborhood of  $\partial M$ , we know that  $\partial P(\Omega) = P(d\Omega) = 0$  for all currents  $P \in \mathcal{F}_{n-1}(\mathbb{R}^N)$  with  $\text{supp}(P) \subset \partial M$ . This implies  $\partial T(\Omega) = 0$  for all currents  $T \in \mathcal{B}_{n-1}(M, \partial M)$ .

It follows from results in Section 5.1.6 in [5] that we can find a current  $S \in \mathcal{R}_{n-1}(\mathbb{R}^N)$  with the following properties:

- We have  $S - \tilde{S} \in \mathcal{B}_{n-1}(M, \partial M)$ . In particular,  $S \in \mathcal{Z}_{n-1}(M, \partial M)$ .
- The current  $S$  is homologically area-minimizing in the sense that  $\mathbf{M}(S) \leq \mathbf{M}(S + X)$  for all currents  $X \in \mathcal{B}_{n-1}(M, \partial M) \cap \mathcal{R}_{n-1}(\mathbb{R}^N)$ .

By Proposition B.5,  $S$  is an  $(n-1)$ -dimensional integer multiplicity rectifiable current. Since  $S - \tilde{S} \in \mathcal{B}_{n-1}(M, \partial M)$ , it follows that  $\partial S(\Omega) = \partial\tilde{S}(\Omega)$ . In particular,  $\partial S(\Omega) \neq 0$ .

**Lemma B.6.** *Let  $p$  be an arbitrary point in  $M$ . Then we can find an open set  $O \subset \mathbb{R}^N$  containing  $p$  such that  $\mathbf{M}(S) \leq \mathbf{M}(S + X)$  for all  $(n-1)$ -dimensional integer multiplicity rectifiable currents  $X$  satisfying  $\text{supp}(X) \subset M \cap O$  and  $\text{supp}(\partial X) \subset \partial M \cap O$ .*

**Proof.** Since  $S$  is homologically area-minimizing, results in Section 5.1.6 in [5] imply that  $S$  is locally area-minimizing with respect to  $(M, \partial M)$ . Hence, we can find an open set  $O \subset \mathbb{R}^N$  containing  $p$  such that  $\mathbf{M}(S) \leq \mathbf{M}(S + X)$  for all currents  $X \in \mathcal{R}_{n-1}(\mathbb{R}^N)$  satisfying  $\text{supp}(X) \subset M \cap O$  and  $\text{supp}(\partial X) \subset \partial M \cap O$ . On the other hand, by Proposition B.5, every  $(n-1)$ -dimensional integer multiplicity rectifiable current with compact support belongs to  $\mathcal{R}_{n-1}(\mathbb{R}^N)$ . This completes the proof of Lemma B.6.

**Lemma B.7.** *Let  $p$  be an arbitrary point in  $M$ . Then we can find an open set  $\hat{O} \subset \mathbb{R}^N$  containing  $p$  such that  $\mathbf{M}(S) \leq \mathbf{M}(S + X)$  for all  $(n-1)$ -dimensional integer multiplicity rectifiable currents  $X$  satisfying  $\text{supp}(X) \subset \hat{M}_{\varepsilon_0} \cap \hat{O}$  and  $\text{supp}(\partial X) \subset \partial M \cap \hat{O}$ .*

**Proof.** By Lemma B.6, we can find an open set  $O \subset \mathbb{R}^N$  containing  $p$  such that  $\mathbf{M}(S) \leq \mathbf{M}(S + X)$  for all  $(n-1)$ -dimensional integer multiplicity rectifiable currents  $X$  satisfying  $\text{supp}(X) \subset M \cap O$  and  $\text{supp}(\partial X) \subset \partial M \cap O$ . We define  $\hat{O} = U \cap F^{-1}(O)$ . Clearly,  $\hat{O} \subset \mathbb{R}^N$  is an open set containing  $p$ .

We claim that  $\hat{O}$  has the desired property. To see this, suppose that  $X$  is an  $(n-1)$ -dimensional integer multiplicity rectifiable current satisfying  $\text{supp}(X) \subset \hat{M}_{\varepsilon_0} \cap \hat{O}$  and  $\text{supp}(\partial X) \subset \partial M \cap \hat{O}$ . Since  $\text{supp}(X) \subset \hat{O} \subset U$ , the push-forward  $F_{\#}(X)$  is well-defined. The results in Section 4.1.14 in [5] imply that

$$\text{supp}(F_{\#}(X)) \subset F(\text{supp}(X)) \subset F(\hat{O}) \subset M \cap O$$

and

$$\text{supp}(\partial F_{\#}(X)) = \text{supp}(F_{\#}(\partial X)) \subset F(\text{supp}(\partial X)) \subset F(\partial M \cap \hat{O}) \subset \partial M \cap O.$$

This implies

$$\mathbf{M}(S) \leq \mathbf{M}(S + F_{\#}(X)).$$

On the other hand, since  $\text{supp}(S + X) \subset \hat{M}_{\varepsilon_0} \cap U$ , Lemma B.4 gives

$$\mathbf{M}(F_{\#}(S + X)) \leq \mathbf{M}(S + X).$$

Finally, since  $\text{supp}(S) \subset M$  and the restriction of  $F$  to  $M$  is the identity, it follows that  $F_{\#}(S) = S$  (see [5], Section 4.1.15). Putting these facts together, we conclude that

$$\mathbf{M}(S) \leq \mathbf{M}(S + X).$$

This completes the proof of Lemma B.7.

Lemma B.7 allows us to apply Grüter's regularity theorem [11] (which was stated in Euclidean space, but extends to the manifold setting). Since  $n \leq 7$ , it follows that the singular set of  $S$  is empty. Since  $\partial S(\Omega) \neq 0$ , we can find a connected component of  $\text{supp}(S)$ , denoted by  $\Sigma$ , such that  $\int_{\partial \Sigma} \Omega \neq 0$ . This completes the proof of Theorem B.1.

## REFERENCES

- [1] H. Barzegar, P.T. Chruściel, M. Hörzinger, M. Maliborski, and L. Nguyen, *Remarks on the energy of asymptotically Horowitz-Myers metrics*, Phys. Rev. D 101, 024007 (2020)
- [2] A. Besse, *Einstein manifolds*, Springer-Verlag, Berlin, 1987
- [3] P.T. Chruściel, E. Delay, and R. Wutte, *Hyperbolic energy and Maskit gluing*, Adv. Theor. Math. Phys. (to appear)
- [4] P.T. Chruściel, G.J. Galloway, L. Nguyen, and T.-T. Paetz, *On the mass aspect function and positive energy theorems for asymptotically hyperbolic manifolds*, Class. Quantum Grav. 35, 115015 (2018)
- [5] H. Federer, *Geometric Measure Theory*, Grundlehren Math. Wiss. vol. 153, Springer-Verlag, New York (1969)
- [6] F. Fiala, *Le problème des isopérimètres sur les surfaces ouvertes à courbure positive*, Comm. Math. Helv. 13, 293–346 (1941)
- [7] D. Fischer-Colbrie and R. Schoen, *The structure of complete stable minimal surfaces in 3-manifolds of nonnegative scalar curvature*, Comm. Pure Appl. Math. 33, 199–211 (1980)
- [8] G.J. Galloway and E. Woolgar, *On static Poincaré-Einstein metrics*, J. High Energy Phys. 2015, 51 (2015)
- [9] M. Gromov, *Metric inequalities with scalar curvature*, Geom. Funct. Anal. 28, 645–726 (2018)
- [10] M. Gromov and H.B. Lawson, Jr., *Positive scalar curvature and the Dirac operator on complete Riemannian manifolds*, Publ. Math. IHÉS 58, 83–196 (1984)
- [11] M. Grüter, *Optimal regularity for codimension one minimal surfaces with a free boundary*, Manuscripta Math. 58, 295–343 (1987)
- [12] P. Hartman, *Geodesic parallel coordinates in the large*, Amer. J. Math. 86, 705–727 (1964)
- [13] G.T. Horowitz and R.C. Myers, *AdS-CFT correspondence and a new positive energy conjecture for general relativity*, Phys. Rev. D 59, 026005 (1998)
- [14] Z. Liang and X. Zhang, *The positive energy conjecture for a class of AHM metrics on  $\mathbb{R}^2 \times \mathbb{T}^{n-2}$* , Sci. China Math. 66, 1041–1052 (2023)
- [15] R. Schoen and S.T. Yau, *On the structure of manifolds with positive scalar curvature*, Manuscripta Math. 28, 159–183 (1979)
- [16] R. Schoen and S.T. Yau, *The existence of a black hole due to condensation of matter*, Comm. Math. Phys. 90, 575–579 (1983)
- [17] R. Schoen and S.T. Yau, *Positive scalar curvature and minimal hypersurface singularities*, arXiv:1704.05490
- [18] K. Shiohama, T. Shioya, and M. Tanaka, *The geometry of total curvature on complete open surfaces*, Cambridge Tracts in Mathematics vol. 159, Cambridge University Press, Cambridge, 2003
- [19] L. Simon, *Introduction to Geometric Measure Theory*, available at <https://math.stanford.edu/~lms/ntu-gmt-text.pdf>
- [20] E. Woolgar, *The rigid Horowitz-Myers conjecture*, J. High Energy Phys. 2017, 104 (2017)

COLUMBIA UNIVERSITY, 2990 BROADWAY, NEW YORK NY 10027, USA

UNIVERSITY OF ILLINOIS, 1409 W. GREEN STREET, URBANA IL 61801, USA