

A study of the spin 1 Unruh-De Witt detectors

F. M. Guedes,^{1,*} M. S. Guimaraes,^{1,†} I. Roditi,^{2,‡} and S. P. Sorella^{1,§}

¹*UERJ – Universidade do Estado do Rio de Janeiro,
Instituto de Física – Departamento de Física Teórica – Rua São
Francisco Xavier 524, 20550-013, Maracanã, Rio de Janeiro, Brazil*

²*CBPF – Centro Brasileiro de Pesquisas Físicas,
Rua Dr. Xavier Sigaud 150, 22290-180, Rio de Janeiro, Brazil*

A study of the spin 1 Unruh-De Witt detectors interacting with a relativistic scalar quantum field is presented. After tracing out the field modes, the resulting density matrix for a bipartite qutrit system is employed to investigate the violation of the Bell-CHSH inequality. Unlike the case of spin 1/2, for which the effects of the quantum field result in a decreasing of the size of violation, in the case of spin 1 both decreasing and increasing of the violation may occur. This effect is ascribed to the fact that Tsirelson's bound is not saturated in the case of qutrits.

I. INTRODUCTION

The Unruh-De Witt (UDW) detectors are useful models, broadly employed in the study of relativistic quantum information, see [1–3] and refs. therein.

In this work, we shall employ spin 1 Unruh-De Witt detectors to investigate the effects of a quantum relativistic scalar field on the Bell-CHSH inequality [4, 5], following the setup already outlined for spin 1/2 detectors [6]. More precisely, we consider the interaction of a pair of qutrits with a real Klein-Gordon field in Minkowski spacetime, by taking as initial field configuration the vacuum state $|0\rangle$. On the other hand, as starting state for the qutrit, we choose the entangled singlet state

$$|\psi\rangle_{AB} = \frac{1}{\sqrt{3}} (|1\rangle_A \otimes |-1\rangle_B - |0\rangle_A \otimes |0\rangle_B + |-1\rangle_A \otimes |1\rangle_B), \quad (1)$$

where, as usual, A, B stand for Alice and Bob, which are meant to be located in the right and left Rindler wedges respectively¹, according to the relativistic causality requirement.

Due to the use of the state $|0\rangle$ for the quantum field, this framework is referred as the extraction of the entanglement from the vacuum. Nevertheless, as we shall see in the sequel, there are remarkable differences between the spin 1/2 and the spin 1 cases. As far as the Bell-CHSH inequality is concerned, for spin 1/2, the effects induced by the quantum field result in a decreasing of the size of the violation, due to the fact that the Tsirelson bound [8], *i.e.* $2\sqrt{2} \approx 2.83$ is already attained in the absence of the field. As the Tsirelson bound is the maximum allowed value for the violation, one can easily figure out that the presence of a quantum field can only induce a decreasing of the size of the violation, see [6] for more details. Instead, in the case of spin 1, the situation looks rather different. Here, it is known that Tsirelson's bound is never attained [9, 10]. The maximum value for the Bell-CHSH inequality is ≈ 2.55 . As such, depending on the choice of the parameters, the effects of the quantum field may give rise either to a decreasing or to an increasing of the violation, while remaining compatible with Tsirelson's bound $2\sqrt{2}$.

Let us begin by specifying the initial density matrix

$$\rho_{AB\varphi}(0) = \rho_{AB}(0) \otimes |0\rangle\langle 0|, \quad (2)$$

where

$$\rho_{AB}(0) = |\psi\rangle_{AB} {}_{AB}\langle\psi|. \quad (3)$$

* fmkguedes@gmail.com

† msguimaraes@uerj.br

‡ roditi@cbpf.br

§ silvio.sorella@gmail.com

¹ The Rindler geometry is that observed by an accelerated observer in Minkowski space.

The time evolution of the initial density matrix, Eq.(2), is governed by the unitary operator

$$\mathcal{U} = e^{-i[J_A^z \otimes \varphi(f_A) + J_B^z \otimes \varphi(f_B)]}, \quad (4)$$

where the operator J^z corresponds to the component of spin along the z -axis, and $\varphi(f_j)$, $j = A, B$, is the smeared field [7]:

$$\varphi(f_j) = \int d^4x \varphi(x) f_j(x), \quad j = A, B, \quad (5)$$

where $f_j(x)$ are smooth test functions with compact support. The role of $f_j(x)$ is that of restricting the field φ to suitable regions of the Minkowski spacetime, so as to ensure that Alice and Bob are space-like separated. For the quantum field φ , one writes

$$\begin{aligned} \varphi(x) &= \int \frac{d^3p}{(2\pi)^3} \frac{1}{2\omega_p} (e^{-ipx} a_p + e^{ipx} a_p^\dagger), \quad \omega_p = \sqrt{p^2 + m^2} \\ [a_p, a_q^\dagger] &= (2\pi)^3 (2\omega_p) \delta^3(p - q), \quad [a_p, a_q] = 0. \end{aligned} \quad (6)$$

At large time, the density matrix is written as

$$\rho_{AB\varphi} = \rho_{AB\varphi}(t \rightarrow \infty) = \mathcal{U} \rho_{AB\varphi}(0) \mathcal{U}^\dagger. \quad (7)$$

The subsequent stage involves deriving the density matrix $\hat{\rho}_{AB}$ for the qutrit system through the process of tracing out the field modes:

$$\hat{\rho}_{AB} = \text{Tr}_\varphi(\rho_{AB\varphi}). \quad (8)$$

Finally, once the density matrix $\hat{\rho}_{AB}$ is known, one is capable of evaluating the Bell-CHSH correlator

$$\langle \mathcal{C} \rangle = \text{Tr}(\hat{\rho}_{AB} \mathcal{C}), \quad (9)$$

where

$$\mathcal{C} = (A + A')B + (A - A')B', \quad (10)$$

with (A, A') , (B, B') being the Bell operators, namely

$$\begin{aligned} A &= A^\dagger, \quad A' = A'^\dagger, \quad B = B^\dagger, \quad B' = B'^\dagger \\ A^2 &= A'^2 = B^2 = B'^2 = 1 \\ [A, B] &= [A, B'] = [A', B] = [A', B'] = 0. \end{aligned} \quad (11)$$

The Bell-CHSH inequality is said to be violated whenever

$$2 < |\langle \mathcal{C} \rangle| \leq 2\sqrt{2}. \quad (12)$$

This work is organized as follows. In Sec.(II), we evaluate the qutrit density matrix by considering the dephasing coupling regime. In Sec.(III) we provide an overview of the fundamental characteristics of the Weyl operators W_{f_j}

$$W_{f_j} = e^{i\varphi(f_j)}, \quad j = A, B, \quad (13)$$

and their von Neumann algebra, introducing key concepts that will be employed throughout this study. In Sec.(IV), we discuss how the effects of the quantum field φ on the violation of the Bell-CHSH inequality, which can be obtained in closed form by using the powerful modular theory of Tomita-Takesaki [11–15]. Notably, it turns out that the violation of the Bell-CHSH inequality exhibits both an increasing and a decreasing behavior as compared to the case in which the field φ is absent, Section (V) collects our conclusion.

II. EVALUATION OF THE QUTRIT DENSITY MATRIX IN THE CASE OF THE DEPHASING COUPLING DETECTORS

We shall consider the density matrix $\hat{\rho}_{AB}$ in the so-called dephasing coupling regime [2, 3], for which the evolution operator is given by $\mathcal{U} = \mathcal{U}_A \otimes \mathcal{U}_B$, where the unitary operator for the detector $j = A, B$ is

$$\mathcal{U}_j = e^{-iJ_j^z \otimes \varphi(f_j)}, \quad (14)$$

with the commutation relation

$$[\mathcal{U}_A, \mathcal{U}_B] = 0. \quad (15)$$

Using the algebra of the spin 1 matrices, the expression (14) can be written as

$$\mathcal{U}_j = \mathbb{1}_j - iJ_j^z s_j + (J_j^z)^2 (c_j - 1), \quad (16)$$

where $c_j \equiv \cos \varphi(f_j)$ and $s_j \equiv \sin \varphi(f_j)$. With the initial density matrix $\rho_{AB\varphi}(0)$ given in Eq.(2), its evolution is described as follows:

$$\begin{aligned} \rho_{AB\varphi} &= (\mathcal{U}_A \otimes \mathcal{U}_B) \rho_{AB\varphi}(0) (\mathcal{U}_A^\dagger \otimes \mathcal{U}_B^\dagger) \\ &= [\mathbb{1}_A - iJ_A^z s_A (J_A^z)^2 (c_A - 1)] \otimes [\mathbb{1}_B - iJ_B^z s_B (J_B^z)^2 (c_B - 1)] \rho_{AB}(0) |0\rangle\langle 0| \\ &\quad \times [\mathbb{1}_A + iJ_A^z s_A (J_A^z)^2 (c_A - 1)] \otimes [\mathbb{1}_B + iJ_B^z s_B (J_B^z)^2 (c_B - 1)]. \end{aligned} \quad (17)$$

Tracing over φ , we get a rather lengthy expression for $\hat{\rho}_{AB}$, namely

$$\begin{aligned} \hat{\rho}_{AB} &= \rho_{AB}(0) + \rho_{AB}(0) (J_B^z)^2 \langle c_B - 1 \rangle - \rho_{AB}(0) J_A^z \otimes J_B^z \langle s_A s_B \rangle + \rho_{AB}(0) (J_A^z)^2 \langle c_A - 1 \rangle \\ &\quad + \rho_{AB}(0) (J_A^z)^2 \otimes (J_B^z)^2 \langle (c_A - 1)(c_B - 1) \rangle + (J_B^z)^2 \rho_{AB}(0) J_B^z \langle s_B^2 \rangle + (J_B^z)^2 \rho_{AB}(0) J_A^z \langle s_B s_A \rangle \\ &\quad + J_B^z \rho_{AB}(0) J_A^z \otimes (J_B^z)^2 \langle s_B s_A (c_B - 1) \rangle + J_B^z \rho_{AB}(0) (J_A^z)^2 \otimes J_B^z \langle s_B^2 (c_A - 1) \rangle \\ &\quad + (J_B^z)^2 \rho_{AB}(0) \langle c_B - 1 \rangle + (J_B^z)^2 \rho_{AB}(0) (J_B^z)^2 \langle (c_B - 1)^2 \rangle - (J_B^z)^2 \rho_{AB}(0) J_z^B \otimes J_z^A \langle (c_B - 1) s_A s_B \rangle \\ &\quad + (J_B^z)^2 \rho_{AB}(0) J_z^A \langle (c_B - 1)(c_A - 1) \rangle + (J_B^z)^2 \rho_{AB}(0) (J_B^z)^2 \otimes (J_A^z)^2 \langle (c_B - 1)^2 (c_A - 1) \rangle \\ &\quad + J_A^z \rho_{AB}(0) J_B^z \langle s_A s_B \rangle + J_A^z \rho_{AB}(0) J_A^z \langle s_A^2 \rangle + J_A^z \rho_{AB}(0) J_A^z \otimes (J_B^z)^2 \langle s_A^2 (c_B - 1) \rangle \\ &\quad + J_A^z \rho_{AB}(0) (J_A^z)^2 \otimes J_B^z \langle s_A s_B (c_A - 1) \rangle - J_A^z \otimes J_B^z \rho_{AB}(0) \langle s_A s_B \rangle - J_A^z \otimes J_B^z \rho_{AB}(0) (J_B^z)^2 \langle s_A s_B (c_B - 1) \rangle \\ &\quad + J_A^z \otimes J_B^z \rho_{AB}(0) J_A^z \otimes J_B^z \langle s_A^2 s_B^2 \rangle - J_A^z \otimes J_B^z \rho_{AB}(0) (J_A^z)^2 \langle s_A s_B (c_A - 1) \rangle \\ &\quad - J_A^z \otimes J_B^z \rho_{AB}(0) (J_A^z)^2 \otimes (J_B^z)^2 \langle s_A s_B (c_A - 1)(c_B - 1) \rangle + J_A^z \otimes (J_B^z)^2 \rho_{AB}(0) J_B^z \langle s_A s_B (c_B - 1) \rangle \\ &\quad + J_A^z \otimes (J_B^z)^2 \rho_{AB}(0) J_A^z \langle s_A^2 (c_B - 1) \rangle + J_A^z \otimes (J_B^z)^2 \rho_{AB}(0) J_A^z \otimes (J_B^z)^2 \langle s_A^2 (c_B - 1)^2 \rangle \\ &\quad + J_A^z \otimes (J_B^z)^2 \rho_{AB}(0) (J_A^z)^2 \otimes J_B^z \langle s_A s_B (c_B - 1)(c_A - 1) \rangle + (J_A^z)^2 \rho_{AB}(0) \langle (c_A - 1) \rangle \\ &\quad + (J_A^z)^2 \rho_{AB}(0) (J_B^z)^2 \langle (c_A - 1)(c_B - 1) \rangle - (J_A^z)^2 \rho_{AB}(0) J_A^z \otimes J_B^z \langle (c_A - 1) s_A s_B \rangle \\ &\quad + (J_A^z)^2 \rho_{AB}(0) (J_A^z)^2 \langle (c_A - 1)^2 \rangle + (J_A^z)^2 \rho_{AB}(0) (J_A^z)^2 \otimes (J_B^z)^2 \langle (c_A - 1)^2 (c_B - 1) \rangle \\ &\quad + (J_A^z)^2 \otimes J_B^z \rho_{AB}(0) J_B^z \langle (c_A - 1) s_B^2 \rangle + (J_A^z)^2 \otimes J_B^z \rho_{AB}(0) J_A^z \langle (c_A - 1) s_B s_A \rangle \\ &\quad + (J_A^z)^2 \otimes J_B^z \rho_{AB}(0) J_A^z \otimes (J_B^z)^2 \langle (c_A - 1)(c_B - 1) s_B s_A \rangle + (J_A^z)^2 \otimes J_B^z \rho_{AB}(0) (J_A^z)^2 \otimes J_B^z \langle (c_A - 1)^2 s_B^2 \rangle \\ &\quad + (J_A^z)^2 \otimes (J_B^z)^2 \rho_{AB}(0) \langle (c_A - 1)(c_B - 1) \rangle + (J_A^z)^2 \otimes J_B^z \rho_{AB}(0) (J_B^z)^2 \langle (c_A - 1)(c_B - 1)^2 \rangle \\ &\quad - (J_A^z)^2 \otimes J_B^z \rho_{AB}(0) J_A^z \otimes J_B^z \langle (c_A - 1)(c_B - 1) s_A s_B \rangle + (J_A^z)^2 \otimes J_B^z \rho_{AB}(0) (J_A^z)^2 \langle (c_A - 1)^2 (c_B - 1) \rangle \\ &\quad + (J_A^z)^2 \otimes J_B^z \rho_{AB}(0) (J_A^z)^2 \otimes (J_B^z)^2 \langle (c_A - 1)^2 (c_B - 1)^2 \rangle. \end{aligned} \quad (18)$$

where $\langle s_A s_B (c_A - 1)(c_B - 1) \rangle$, etc., denotes the expectation value of the Weyl operators, namely

$$\langle s_A s_B (c_A - 1)(c_B - 1) \rangle = \langle 0 | s_A s_B (c_A - 1)(c_B - 1) | 0 \rangle. \quad (19)$$

In the next section we shall see how can these correlation functions be addressed in closed form, using the Tomita-Takesaki theory.

III. TOMITA-TAKESAKI MODULAR THEORY AND THE VON NEUMANN ALGEBRA OF THE WEYL OPERATORS

To calculate the correlation functions of the Weyl operators, Eq. (19), it is worth providing a compact review of the properties of the von Neumann algebra related to such operators. For a more detailed review, one can check ref.[15].

Let us begin by recalling the commutator between the scalar fields, for arbitrary spacetime separation

$$[\varphi(x), \varphi(y)] = i\Delta_{PJ}(x-y), \quad (20)$$

where the Lorentz-invariant causal Pauli-Jordan distribution $\Delta_{PJ}(x-y)$ is defined by

$$i\Delta_{PJ}(x-y) = \int \frac{d^4k}{(2\pi)^3} \varepsilon(k^0) \delta(k^2 - m^2) e^{-ik(x-y)}, \quad (21)$$

with $\varepsilon(x) \equiv \theta(x) - \theta(-x)$. The Pauli-Jordan distribution $\Delta_{PJ}(x-y)$ vanishes outside of the light cone, guaranteeing that measurements at points separated by space-like intervals do not interfere, that is

$$\Delta_{PJ}(x-y) = 0, \quad \text{for } (x-y)^2 < 0. \quad (22)$$

Now, let \mathcal{O} be an open region of the Minkowski spacetime and let $\mathcal{M}(\mathcal{O})$ be the space of smooth test functions with support contained in \mathcal{O} , namely

$$\mathcal{M}(\mathcal{O}) = \{f | \text{supp}(f) \subseteq \mathcal{O}\}. \quad (23)$$

Following [13, 14] one introduces the symplectic complement of $\mathcal{M}(\mathcal{O})$ as

$$\mathcal{M}'(\mathcal{O}) = \{g | \Delta_{PJ}(g, f) = 0, \forall f \in \mathcal{M}(\mathcal{O})\}. \quad (24)$$

This symplectic complement $\mathcal{M}'(\mathcal{O})$ comprises all test functions for which the smeared Pauli-Jordan expression $\Delta_{PJ}(f, g)$ vanishes for any f belonging to $\mathcal{M}(\mathcal{O})$,

$$[\varphi(f), \varphi(g)] = i\Delta_{PJ}(f, g), \quad (25)$$

allowing us to rephrase causality, Eq.(22), as [13, 14]

$$[\varphi(f), \varphi(g)] = 0, \quad (26)$$

whenever $f \in \mathcal{M}(\mathcal{O})$ and $g \in \mathcal{M}'(\mathcal{O})$.

As already mentioned in Sec.(I), the so-called Weyl operators [13–15] play an important role in the study of the Bell-CHSH inequality. This class of unitary operators is obtained by exponentiating the smeared field

$$W_h = e^{i\varphi(h)}. \quad (27)$$

By applying the Baker–Campbell–Hausdorff formula together with the commutation relation (21), one finds that the Weyl operators lead to the following algebraic structure:

$$\begin{aligned} W_f W_g &= e^{-\frac{i}{2}\Delta_{PJ}(f,g)} W_{(f+g)}, \\ W_f^\dagger W_f &= W_f W_f^\dagger = 1, \\ W_f^\dagger &= W_{(-f)}. \end{aligned} \quad (28)$$

Moreover, if f and g are space-like, the Weyl operators W_f and W_g commute. By expanding the field φ in terms of creation and annihilation operators, one can evaluate the expectation value of the Weyl operator, finding

$$\langle 0 | W_h | 0 \rangle = e^{-\frac{1}{2}\|h\|^2}, \quad (29)$$

where $\|h\|^2 = \langle h | h \rangle$ and

$$\langle f | g \rangle = \int \frac{d^3k}{(2\pi)^3} \frac{1}{2\omega_k} f(\omega_k, \vec{k})^* g(\omega_k, \vec{k}), \quad (30)$$

is the Lorentz invariant inner product between the test functions (f, g) [13–15], with the usual relation $\omega_k^2 = \vec{k}^2 + m^2$ and

$$f(\omega_k, \vec{k}) = \int d^4x e^{ikx} f(x), \quad k_0 = \omega_k$$

A von Neumann algebra $\mathcal{A}(\mathcal{M})$ arises by taking all possible products and linear combinations of the Weyl operators defined on $\mathcal{M}(\mathcal{O})$. In particular, the Reeh-Schlieder theorem [7, 11, 13, 14], states that the vacuum state $|0\rangle$ is both cyclic and separating for the von Neumann algebra \mathcal{A} . Consequently, we can apply the Tomita-Takesaki modular theory [11–15] and introduce the anti-linear unbounded operator S , whose action on the von Neumann algebra $\mathcal{A}(\mathcal{M})$ is defined as

$$S a|0\rangle = a^\dagger|0\rangle, \quad \forall a \in \mathcal{A}(\mathcal{M}), \quad (31)$$

from which it follows that $S^2 = 1$ and $S|0\rangle = |0\rangle$. The operator S has a unique polar decomposition [12]:

$$S = J\Delta^{1/2}, \quad (32)$$

where J is anti-unitary and Δ is positive and self-adjoint. These operators are characterized by the following set of properties [11–15]:

$$\begin{aligned} \Delta &= S^\dagger S, & J\Delta^{1/2}J &= \Delta^{-1/2}, \\ J^2 &= 1, & S^\dagger &= J\Delta^{-1/2}, \\ J^\dagger &= J, & \Delta^{-1} &= SS^\dagger. \end{aligned} \quad (33)$$

From the Tomita-Takesaki theorem [11–15], it follows that $J\mathcal{A}(\mathcal{M})J = \mathcal{A}'(\mathcal{M})$, meaning that, upon conjugation by the operator J , the algebra $\mathcal{A}(\mathcal{M})$ is mapped onto its commutant $\mathcal{A}'(\mathcal{M})$, namely:

$$\mathcal{A}'(\mathcal{M}) = \{ a' \mid [a, a'] = 0, \forall a \in \mathcal{A}(\mathcal{M}) \}. \quad (34)$$

Furthermore, the theorem states that there is a one-parameter family of operators Δ^{it} , $t \in \mathbb{R}$, that leave the algebra $\mathcal{A}(\mathcal{M})$ invariant, such that the the following equation holds

$$\Delta^{it}\mathcal{A}(\mathcal{M})\Delta^{-it} = \mathcal{A}(\mathcal{M}).$$

The Tomita-Takesaki modular theory is particularly well-suited for analyzing the Bell-CHSH inequality within the framework of relativistic Quantum Field Theory [13, 14]. As demonstrated in [15], it provides a purely algebraic method for constructing Bob's operators from Alice's ones by using the modular conjugation J . Given Alice's operator A_f , one can assign to Bob the operator $B_f = JA_fJ$, ensuring their mutual commutativity due to the Tomita-Takesaki theorem, as $B_f = JA_fJ$ belongs to the commutant $\mathcal{A}'(\mathcal{M})$ [15].

An important outcome of the Tomita-Takesaki modular theory, established by [16, 17], allows the extension of the action of the modular operators (J, Δ) to the space of the test functions. In fact, when equipped with the Lorentz-invariant inner product $\langle f|g\rangle$, Eq.(30), the set of test functions forms a complex Hilbert space \mathcal{F} that possesses a variety of properties. To be more precise, it is found that the subspaces \mathcal{M} and $i\mathcal{M}$ are standard subspaces for \mathcal{F} [16]. This implies that:

- i. $\mathcal{M} \cap i\mathcal{M} = \{0\}$;
- ii. $\mathcal{M} + i\mathcal{M}$ is dense in \mathcal{F} .

As shown in to [16], for such subspaces, it's viable to establish a modular theory similar to that of the Tomita-Takesaki theory. This involves introducing an operator s acting on $\mathcal{M} + i\mathcal{M}$ such that

$$s(f + ih) = f - ih, \quad (35)$$

for $f, h \in \mathcal{M}$. With this definition, it's worth noting that $s^2 = 1$. Employing the polar decomposition, one obtains:

$$s = j\delta^{1/2}, \quad (36)$$

where j is an anti-unitary operator, while δ is positive and self-adjoint. Similarly to the operators (J, Δ) , the operators (j, δ) fulfill the following properties [16]:

$$j\delta^{1/2}j = \delta^{-1/2}, \quad \delta^\dagger = \delta$$

$$\begin{aligned} s^\dagger &= j\delta^{-1/2}, \quad j^\dagger = j \\ \delta &= s^\dagger s, \quad j^2 = 1. \end{aligned} \quad (37)$$

Further, one can show [14, 16] that a test function f belongs to \mathcal{M} if and only if

$$sf = f. \quad (38)$$

Indeed, let us suppose that $f \in \mathcal{M}$. From Eq.(35), one can express

$$sf = h_1 + ih_2, \quad (39)$$

for some (h_1, h_2) . Since $s^2 = 1$ it follows that

$$f = s(h_1 + ih_2) = h_1 - ih_2, \quad (40)$$

so that $h_1 = f$ and $h_2 = 0$. Similarly, one has that $f' \in \mathcal{M}'$ if and only if $s^\dagger f' = f'$.

Thus, the lifting of the action of the operators (J, Δ) to the space of test functions is accomplished by [17]

$$J e^{i\varphi(f)} J = e^{-i\varphi(jf)}, \quad \Delta e^{i\varphi(f)} \Delta^{-1} = e^{i\varphi(\delta f)}. \quad (41)$$

Also, it is important to note that if $f \in \mathcal{M} \implies jf \in \mathcal{M}'$. This property follows from

$$s^\dagger(jf) = j\delta^{-1/2}jf = \delta f = j(j\delta f) = j(sf) = jf. \quad (42)$$

It is worth reminding here that, for wedge regions in Minkowski spacetime, the spectrum of δ coincides with the positive real line, *i.e.*, $\log(\delta) = \mathbb{R}$ [18]. δ is an unbounded operator with continuous spectrum.

We have now all the necessary ingredients to evaluate the correlation functions of the Weyl operators. By examining expression (18), one recognizes that the fundamental quantity to be computed is of the form

$$\langle e^{i\varphi(f_A)} e^{\pm i\varphi(f_B)} \rangle = \langle e^{i(\phi(f_A) \pm \phi(f_B))} \rangle = e^{-\frac{1}{2}\|f_A \pm f_B\|^2}, \quad (43)$$

so that we need to evaluate the following norms ($\|f_A\|^2, \|f_B\|^2$) and the inner product $\langle f_A | f_B \rangle$. We begin by focusing on Alice's test function f_A . We require that $f_A \in \mathcal{M}(\mathcal{O})$ where \mathcal{O} is located in the right Rindler wedge. Following [13–15], the test function f_A can be further specified by considering the spectrum of the operator δ . By selecting the spectral subspace defined by $[\lambda^2 - \varepsilon, \lambda^2 + \varepsilon] \subset (0, 1)$ and introducing the normalized vector ϕ belonging to this subspace, one writes

$$f_A = \eta(1 + s)\phi, \quad (44)$$

where η is an arbitrary parameter. As required by the setup outlined above, Eq.(44) ensures that

$$sf_A = f_A. \quad (45)$$

We observe that $j\phi$ is orthogonal to ϕ , *i.e.*, $\langle \phi | j\phi \rangle = 0$. In fact, from

$$\delta^{-1}(j\phi) = j(j\delta^{-1}j)\phi = j(\delta\phi), \quad (46)$$

it follows that the modular conjugation j exchanges the spectral subspace $[\lambda^2 - \varepsilon, \lambda^2 + \varepsilon]$ with $[1/\lambda^2 - \varepsilon, 1/\lambda^2 + \varepsilon]$. Regarding Bob's test function f_B , we use the modular conjugation operator j and define

$$f_B = jf_A, \quad (47)$$

ensuring that

$$s^\dagger f_B = f_B \quad (48)$$

This implies that, as required by the relativistic causality, f_B belongs to the symplectic complement $\mathcal{M}'(\mathcal{O})$, located in the left Rindler wedge, namely: $f_B \in \mathcal{M}'(\mathcal{O})$. Finally, considering that ϕ belongs to the spectral subspace $[\lambda^2 - \varepsilon, \lambda^2 + \varepsilon]$, it follows that [15],

$$\begin{aligned} \|f_A\|^2 &= \|jf_A\|^2 = \eta^2(1 + \lambda^2), \\ \langle f_A | jf_A \rangle &= 2\eta^2\lambda, \end{aligned} \quad (49)$$

which provide us the needed inner products.

IV. THE BELL-CHSH INEQUALITY

We face now the Bell-CHSH inequality, Eq.(9). We begin by defining the Bell operators [13, 14, 19]:

$$\begin{aligned}
A|-1\rangle &= e^{i\alpha}|1\rangle, & A|0\rangle &= |0\rangle, & A|1\rangle &= e^{-i\alpha}|-1\rangle \\
A'|-1\rangle &= e^{i\alpha'}|1\rangle, & A'|0\rangle &= |0\rangle, & A'|1\rangle &= e^{-i\alpha'}|-1\rangle \\
B|-1\rangle &= e^{-i\beta}|1\rangle, & B|0\rangle &= |0\rangle, & B|1\rangle &= e^{i\beta}|-1\rangle \\
B'|-1\rangle &= e^{-i\beta'}|1\rangle, & B'|0\rangle &= |0\rangle, & B'|1\rangle &= e^{i\beta'}|-1\rangle,
\end{aligned} \tag{50}$$

which fulfill the whole set of conditions (11). The free parameters $(\alpha, \alpha', \beta, \beta')$, which will be chosen at the best convenience, correspond to the four Bell's angles.

Reminding that the initial state for AB is

$$|\psi\rangle_{AB} = \frac{1}{\sqrt{3}} (|-1\rangle_A \otimes |-1\rangle_B - |0\rangle_A \otimes |0\rangle_B + |1\rangle_A \otimes |1\rangle_B),$$

and using Eq (9), one gets the Bell-CHSH correlator

$$\begin{aligned}
\langle \mathcal{C} \rangle &= \frac{1}{3} [1 + 2 \cos(\alpha + \beta)] + \frac{2}{3} \cos(\alpha + \beta) [2\langle (c_A - 1) \rangle + 2\langle (c_B - 1) \rangle + 4\langle s_A s_B \rangle \\
&\quad - \langle s_A^2 \rangle - \langle s_B^2 \rangle + 4\langle s_A s_B (c_B - 1) \rangle + 4\langle s_A s_B (c_A - 1) \rangle - 2\langle s_B^2 (c_A - 1) \rangle \\
&\quad - 2\langle s_A^2 (c_B - 1) \rangle + 4\langle (c_A - 1)(c_B - 1) \rangle + \langle (c_A - 1)^2 \rangle + \langle (c_B - 1)^2 \rangle \\
&\quad + 2\langle (c_A - 1)(c_B - 1)^2 \rangle + 2\langle (c_A - 1)^2 (c_B - 1) \rangle + \langle s_A^2 s_B^2 \rangle \\
&\quad + 4\langle s_A s_B (c_A - 1)(c_B - 1) \rangle - \langle s_A^2 (c_B - 1)^2 \rangle - \langle s_B^2 (c_A - 1)^2 \rangle \\
&\quad + \langle (c_A - 1)^2 (c_B - 1)^2 \rangle] + (\alpha \rightarrow \alpha') + (\beta \rightarrow \beta') - (\alpha \rightarrow \alpha', \beta \rightarrow \beta').
\end{aligned} \tag{51}$$

The expression above is written in terms of the inner products between test functions, which can be evaluated by employing the expressions (49). The final expression reads

$$\begin{aligned}
\langle \mathcal{C} \rangle &= \frac{2}{3} \{1 + [\cos(\alpha + \beta) + \cos(\alpha' + \beta) + \cos(\alpha + \beta') - \cos(\alpha' + \beta')]\} \\
&\quad - \frac{4}{3} f(\eta, \lambda) [\cos(\alpha + \beta) + \cos(\alpha' + \beta) + \cos(\alpha + \beta') - \cos(\alpha' + \beta')],
\end{aligned} \tag{52}$$

where the function $f(\eta, \lambda)$ is

$$f(\eta, \lambda) = e^{-2\eta^2(1+\lambda^2)} - e^{-4\eta^2(1-\lambda)^2}, \tag{53}$$

with $\eta \neq 0$ and $0 < \lambda < 1$. From (52), one learns several things:

- When the quantum field φ is removed, *i.e.* $\eta^2 \rightarrow 0$, and therefore, $f(\eta, \lambda) = 0$, we recover the Bell-CHSH inequality of Quantum Mechanics for qutrits, whose maximum value is [9, 10]

$$\langle \mathcal{C} \rangle_{f=0} = \frac{2}{3} (1 + 2\sqrt{2}) \approx 2.55 > 2. \tag{54}$$

One notices that this value is lower than Tsirelson's bound, as we are dealing with a spin 1 system.

- The contributions arising from the scalar field φ are emcoded in the exponential terms $e^{-4\eta^2(1-\lambda)^2}$ and $e^{-2\eta^2(1+\lambda^2)}$. It is worth reminding here that the parameter η^2 is related to the norm of the test function f_A , Eqs.(49), that is, this parameter reflects the freedom one has in defining the test function f_A through the operator s . As pointed out in [15, 20], η is a free parameter appearing in the Quantum Field Theory formulation of the Bell-CHSH inequality in terms of Weyl operators, playing a similar role of the free Bell's angles and it can be chosen in the most suitable way.
- It remains to investigate what happens when $f(\eta, \lambda) \neq 0$. From $e^{-2\eta^2(1+\lambda^2)} > e^{-4\eta^2(1-\lambda)^2}$, we get the roots $\lambda_{\pm} = 2 \pm \sqrt{3}$. Since $\lambda \in [0, 1]$, one can distinguish two possibilities. The first one is when $0 < \lambda < 2 - \sqrt{3}$. In this case, the quantum field produces a damping, resulting in a decreasing of the violation of the Bell-CHSH inequality, as compared to the pure Quantum Mechanical case. The second possibility takes place when $2 - \sqrt{3} < \lambda < 1$, resulting in an improvement of the size of the violation.

- The possible values of η and λ are restricted to be such that $|\langle \mathcal{C} \rangle|$ does not exceed Tsirelson's bound. Thus, observing that the maximum value of the angular part of Eq.(52) is $2\sqrt{2}$, attained for the following values of the Bell's angles:

$$\alpha = 0, \quad \alpha' = \frac{\pi}{2}, \quad \beta = -\frac{\pi}{4}, \quad \beta' = \frac{\pi}{4}, \quad (55)$$

it follows that

$$\langle \mathcal{C} \rangle = \frac{2}{3} \{1 + 2\sqrt{2}[1 - 2f(\eta, \lambda)]\}. \quad (56)$$

Accordingly, we shall require that $\langle \mathcal{C} \rangle \leq 2\sqrt{2}$. By doing so, we find that

$$f(\eta, \lambda) \geq \frac{1 - \sqrt{2}}{4\sqrt{2}}. \quad (57)$$

That is, the possible values for η and λ are those for which (57) holds, as shown in Fig. (1).




FIG. 1. Region Plot showing the function $f(\eta, \lambda)$. The possible values for η and λ for which Eq.(57) is fulfilled are those in the blue region.

- The whole effects produced by the quantum field can be captured in Fig. (2). The orange surface represents the maximum value of $\langle \mathcal{C} \rangle$ without the presence of φ , *i.e.* $\langle \mathcal{C} \rangle = 2.55$. One notices the existence of a small region in blue, above the orange surface. This region corresponds to values of (η, λ) for which the size of the violation is improved, almost till ≈ 2.7 . This phenomenon occurs when $\lambda > 0.26$.

V. CONCLUSIONS

In this work, we have analyzed the interaction between spin 1 Unruh-De Witt detectors, *i.e.* a pair of qutrits, and a relativistic quantum scalar field φ . The effects of the quantum field on the Bell-CHSH inequality have been scrutinized in detail by making use of the dephasing channel for the evolution operator. By employing the Tomita-Takesaki modular theory and the properties of the Weyl operators, these effects have been evaluated in closed form, as expressed by Eq. (52).

The main finding of the present study is that the presence of a scalar quantum field may induce both a damping as well as an improvement effect, resulting, respectively, in a decreasing and an increasing of the size of the violation of the Bell-CHSH inequality as compared to the case in which the field is absent.

As such, the case of spin 1 looks much different from that of spin 1/2, for which only a decreasing of the violation has been detected [6]. As already underlined, the existence of an improvement of the size of the violation of




FIG. 2. Behavior of the Bell-CHSH correlator $\langle C \rangle$ as a function of the parameters η and λ . The orange surface represents the maximum value of $\langle C \rangle$ without the presence of φ , *i.e.* $\langle C \rangle = 2.55$. The blue surface above the orange one corresponds to the region in which the effects of the quantum field result in an increasing of the size of the violation of the Bell-CHSH inequality.

the Bell-CHSH inequality can be ascribed to the fact that, for spin 1, the Tsirelson bound $2\sqrt{2}$ is never saturated. Instead, the maximum value obtained in Quantum Mechanics is $[\frac{2}{3}(1 + 2\sqrt{2}), 2\sqrt{2}]$. As such, in the presence of a quantum field φ , there exists a permissible interval, $[\frac{2}{3}(1 + 2\sqrt{2}), 2\sqrt{2}]$, where an increasing of the size of the violation occurs.

ACKNOWLEDGMENTS

The authors would like to thank the Brazilian agencies CNPq and FAPERJ, for financial support. S.P. Sorella, I. Roditi, and M.S. Guimaraes are CNPq researchers under contracts 301030/2019-7, 311876/2021-8, and 309793/2023-8, respectively. F.M. Guedes acknowledges FAPERJ for financial support under the contract SEI-260003/007871/2024.

-
- [1] B. Reznik, *Entanglement from the vacuum*, Found. Phys. **33**, 167 (2003).
 - [2] E. Tjoa, *Quantum teleportation with relativistic communication from first principles*, Phys. Rev. A **106**, 032432 (2022).
 - [3] E. Tjoa, *Nonperturbative simple-generated interactions with a quantum field for arbitrary Gaussian states*, Phys. Rev. D **108**, 045003 (2023).
 - [4] J. S. Bell, *On the Einstein Podolsky Rosen paradox*, Physics Physique Fizika **1**, 195 (1964).
 - [5] J. F. Clauser, M. A. Horne, A. Shimony and R. A. Holt, *Proposed Experiment to Test Local Hidden-Variable Theories*, Phys. Rev. Lett. **23**, 880 (1969).
 - [6] F. M. Guedes, M. S. Guimaraes, I. Roditi and S. P. Sorella, *Unruh-De Witt detectors, Bell-CHSH inequality and Tomita-Takesaki theory*, arXiv:2401.03313v2, to appear in JHEP
 - [7] R. Haag, *Local quantum physics: Fields, particles, algebras*, Springer-Verlag, 1992.
 - [8] B. S. Tsirelson, J. Math. Sci. **36**, 557 (1987).
 - [9] N. Gisin and A. Peres, *Maximal violation of Bell's inequality for arbitrarily large spin*, Phys. Lett. A **162**, 15 (1992).
 - [10] G. Peruzzo and S. P. Sorella, *Entanglement and maximal violation of the CHSH inequality in a system of two spins j : A novel construction and further observations*, Phys. Lett. A **474**, 128847 (2023).
 - [11] E. Witten, *APS Medal for Exceptional Achievement in Research: Invited article on entanglement properties of quantum field theory*, Rev. Mod. Phys. **90**, 045003 (2018).
 - [12] O. Bratteli and D. W. Robinson, *Operator Algebras and Quantum Statistical Mechanics 1*, Springer, 1997.
 - [13] S. J. Summers and R. Werner, *Bell's Inequalities and Quantum Field Theory. 1. General Setting*, J. Math. Phys. **28**, 2440 (1987).
 - [14] S. J. Summers and R. Werner, *Bell's inequalities and quantum field theory. II. Bell's inequalities are maximally violated in the vacuum*, J. Math. Phys. **28**, 2448 (1987).

- [15] P. De Fabritiis, F. M. Guedes, M. S. Guimaraes, G. Peruzzo, I. Roditi, and S. P. Sorella, *Weyl operators, Tomita-Takesaki theory, and Bell-Clauser-Horne-Shimony-Holt inequality violations*, Phys. Rev. D **108**, 085026 (2023).
- [16] M. A. Rieffel and A. Van Daele, *A bounded operator approach to Tomita-Takesaki theory*, Pacific J. Math. **69**, 187 (1977).
- [17] J-P. Eckmann and K. Osterwalder, *An application of Tomita's theory of modular Hilbert algebras: Duality for free Bose fields*, J. Funct. Anal. **13**, 1 (1973).
- [18] J. J. Bisognano and E. H. Wichmann, *On the Duality Condition for a Hermitian Scalar Field*, J. Math. Phys. **16**, 985 (1975).
- [19] S. P. Sorella, *On the Representations of Bell's Operators in Quantum Mechanics*, Found. Phys. **53**, 59 (2023).
- [20] M. S. Guimaraes, I. Roditi and S. P. Sorella, *Bell-CHSH inequality and unitary operators*, arXiv:2403.15276.