# DISTRIBUTION OF THE NUMBER OF PRIME FACTORS WITH A GIVEN MULTIPLICITY

### ERTAN ELMA AND GREG MARTIN

ABSTRACT. Given an integer  $k \geq 2$ , let  $\omega_k(n)$  denote the number of primes that divide n with multiplicity exactly k. We compute the density  $e_{k,m}$  of those integers n for which  $\omega_k(n) = m$  for every integer  $m \geq 0$ . We also show that the generating function  $\sum_{m=0}^{\infty} e_{k,m} z^m$  is an entire function that can be written in the form  $\prod_p (1 + (p-1)(z-1)/p^{k+1})$ ; from this representation we show how to both numerically calculate the  $e_{k,m}$  to high precision and provide an asymptotic upper bound for the  $e_{k,m}$ . We further show how to generalize these results to all additive functions of the form  $\sum_{j=2}^{\infty} a_j \omega_j(n)$ ; when  $a_j = j-1$  this recovers a classical result of Rényi on the distribution of  $\Omega(n) - \omega(n)$ .

#### 1. Introduction

Let  $\omega(n)$  be the number of distinct prime factors of a positive integer n, and let  $\Omega(n)$  be the number of prime factors of n counted with multiplicity. Average behaviours of such arithmetic functions are understood via their summatory functions. It is known [5] (see also [6, Theorems 427–430]) that

(1) 
$$\sum_{n \leq x} \omega(n) = x \log \log x + bx + O\left(\frac{x}{\log x}\right)$$
$$\sum_{n \leq x} \Omega(n) = x \log \log x + \left(b + \sum_{p} \frac{1}{p(p-1)}\right)x + O\left(\frac{x}{\log x}\right);$$

here the constant b is defined by

(2) 
$$b = \gamma_0 + \sum_p \sum_{j=2}^{\infty} \frac{1}{jp^j}$$

where  $\gamma_0$  denotes the Euler–Mascheroni constant. (In this paper,  $\sum_p$  and  $\prod_p$  always denote sums and products running over all prime numbers.) The celebrated Erdős–Kac theorem tells us that both  $\omega(n)$  and  $\Omega(n)$  can be normalized to have Gaussian limiting distribution functions.

By the asymptotic formulas (1), the difference  $\Omega(n) - \omega(n)$  has an average value, namely the constant

$$\lim_{x \to \infty} \frac{1}{x} \sum_{n \leqslant x} (\Omega(n) - \omega(n)) = \sum_{p} \frac{1}{p(p-1)},$$

<sup>2020</sup> Mathematics Subject Classification. 11N37, 11N60.

Key words and phrases. Prime factors, additive functions, limiting distributions.

which provides motivation to study the frequency of each possible value of  $\Omega(n) - \omega(n)$ . For any integer  $m \ge 0$ , define

$$\mathcal{N}_m(x) = \{ n \leqslant x \colon \Omega(n) - \omega(n) = m \}.$$

Rényi [8] (see also [7, Section 2.4]) proved that the (natural) densities

(3) 
$$d_m = \lim_{x \to \infty} \frac{\# \mathcal{N}_m(x)}{x} = \frac{6}{\pi^2} \sum_{\substack{f \in \mathcal{F} \\ \Omega(f) - \omega(f) = m}} \frac{1}{f} \prod_{p|f} \left(1 + \frac{1}{p}\right)^{-1}$$

exist for every  $m \ge 0$ , where  $\mathcal{F}$  is the set of powerful numbers (the set of positive integers all of whose prime factors have multiplicity  $\ge 2$ ). Furthermore, he showed that these densities have the generating function

(4) 
$$\sum_{m=0}^{\infty} d_m z^m = \prod_p \left( 1 - \frac{1}{p} \right) \left( 1 + \frac{1}{p-z} \right) \qquad (|z| < 2).$$

(Note the special case  $d_0 = \prod_p (1 - \frac{1}{p})(1 + \frac{1}{p}) = \frac{1}{\zeta(2)} = \frac{6}{\pi^2}$  for the density of squarefree numbers, which can also be confirmed by realizing that the sum in equation (3) contains only the single term f = 1 when m = 0.) In particular, the smaller function  $\Omega(n) - \omega(n)$  already has a (discrete) limiting distribution function, without needing normalization in the way that the larger functions  $\omega(n)$  and  $\Omega(n)$  individually do.

As a refinement of the function  $\omega(n)$ , Liu and the first author introduced the functions

$$\omega_k(n) = \sum_{p^k || n} 1$$

for each integer  $k \ge 1$ , so that  $\omega_k(n)$  counts the number of prime factors of n with multiplicity k and thus  $\omega(n) = \sum_{k=1}^{\infty} \omega_k(n)$ . They showed [2] that

$$\sum_{n \le x} \omega_1(n) = x \log \log x + \left(b - \sum_{n} \frac{1}{p^2}\right) x + O\left(\frac{x}{\log x}\right)$$

where b is the constant from equation (2), while

(5) 
$$\sum_{n \le x} \omega_k(n) = x \sum_{n} \frac{p-1}{p^{k+1}} + O\left(x^{(k+1)/(3k-1)} \log^2 x\right) \qquad (k \ge 2).$$

They also showed that the larger function  $\omega_1(n)$  has a Gaussian limiting distribution function after being normalized in the same way as the classical  $\omega(n)$  and  $\Omega(n)$ . However, since equation (5) shows that  $\omega_k(n)$  has an average value for each  $k \ge 2$ , we might expect these smaller functions to have limiting distributions without needing to be normalized.

In this paper, we obtain the limiting distribution for the functions  $\omega_k(n)$  for  $k \geq 2$ , analogous to the results of Rényi described above. For integers  $m \geq 0$ , define

(6) 
$$\mathcal{N}_{k,m}(x) = \{ n \leqslant x \colon \omega_k(n) = m \}$$

to be the set of positive integers  $n \leq x$  with exactly m prime factors of multiplicity k. Our main result establishes the existence of the densities

$$e_{k,m} = \lim_{x \to \infty} \frac{\# \mathcal{N}_{k,m}(x)}{x}$$

and provides a closed-form expression for them.

**Theorem 1.** Uniformly for all integers  $k \ge 2$  and  $m \ge 0$ ,

$$\#\mathcal{N}_{k,m}(x) = e_{k,m}x + O(x^{1/2}\log x)$$

with

$$e_{k,m} = \frac{6}{\pi^2} \sum_{\substack{f \in \mathcal{F} \\ \omega_k(f) = m}} \frac{1}{f} \prod_{p|f} \left(1 + \frac{1}{p}\right)^{-1}.$$

**Remark 2.** Note that the  $e_{k,m}$  are all nonnegative, and we can check that they do sum to 1:

$$\sum_{m=0}^{\infty} e_{k,m} = \sum_{m=0}^{\infty} \frac{6}{\pi^2} \sum_{\substack{f \in \mathcal{F} \\ \omega_k(f) = m}} \frac{1}{f} \prod_{p|f} \left( 1 + \frac{1}{p} \right)^{-1} = \frac{6}{\pi^2} \sum_{f \in \mathcal{F}} \frac{1}{f} \prod_{p|f} \left( 1 + \frac{1}{p} \right)^{-1}.$$

Since the summand is a multiplicative function of f, as is the indicator function of  $\mathcal{F}$ , the right-hand side equals its Euler product

$$\frac{6}{\pi^2} \prod_{p} \left( 1 + 0 + \left( 1 + \frac{1}{p} \right)^{-1} \left( \frac{1}{p^2} + \frac{1}{p^3} + \dots \right) \right) = \frac{6}{\pi^2} \prod_{p} \left( 1 - \frac{1}{p^2} \right)^{-1} = 1.$$

The same remark applies to the densities in equation (10) below.

Moreover, we obtain an identity analogous to equation (4) for the generating function of the densities  $e_{k,m}$  for fixed  $k \ge 2$ , from which we can derive an upper bound for the densities  $e_{k,m}$  when  $k \ge 2$  is fixed and  $m \to \infty$ .

**Theorem 3.** Let  $k \ge 2$  be an integer. For all  $z \in \mathbb{C}$  with  $|z| \le 1$ ,

(7) 
$$\sum_{m=0}^{\infty} e_{k,m} z^m = \prod_{p} \left( 1 + \frac{(p-1)(z-1)}{p^{k+1}} \right).$$

Corollary 4. For each fixed  $k \ge 2$ , we have  $e_{k,m} \le m^{-(k-o(1))m}$  as  $m \to \infty$ .

Remark 5. The proof of the upper bound in Corollary 4 (see Section 3) shows that for each  $k \ge 2$ , the bound is attained for infinitely many m; it would be interesting to try to show that  $e_{k,m} = m^{-(k-o(1))m}$  for all k and m. Moreover, the corollary and its proof show that both sides of equation (7) converge to entire functions, and thus Theorem 3 actually holds for all  $z \in \mathbb{C}$  by uniqueness of analytic continuation. The same remarks apply to the generating functions in Corollary 13 and the upper bounds in Corollary 14 below.

Some numerical values of  $e_{k,m}$  are given in Table 1. The numbers in the first column corresponding to m=0 are increasing as k increases, whereas the numbers in other columns are decreasing. This behaviour stems from the fact that the case m=0 indicates the nonexistence of prime factors with multiplicity k, which becomes more probable as k increases. (Note also that each number in the first column exceeds  $\frac{6}{\pi^2} \approx 0.608$ , since every squarefree number n certainly has  $\omega_k(n) = 0$  for all  $k \ge 2$ .) On the other hand, for  $m \ge 1$ , the criterion  $\omega_k(n) = m$  indicates the existence of prime factors with multiplicity k, which becomes less probable as k increases. Details of the calculations of these values are given in Section 4,

Table 1. Some values of  $e_{k,m}$ 

	m = 0	m = 1	m=2	m=3
	0.748535831			
	0.904708927			
	0.959088654			
k = 5	0.981363751	0.018587581	0.000048654	0.000000014

although we do note here that the calculations use the generating function in Theorem 3 rather than the formula for  $e_{k,m}$  in Theorem 1.

A consequence of Theorem 1 and Remark 2 is that  $\omega_k(n)$  has a limiting distribution, which is the same as the distribution of the nonnegative integer-valued random variable  $X_k$  that takes the value m with probability  $e_{k,m}$ . While it is straightforward to calculate the expectation and variance of this limiting distribution via the expressions

$$\lim_{x \to \infty} \frac{1}{x} \sum_{n \le x} \omega_k(n) \quad \text{and} \quad \lim_{x \to \infty} \frac{1}{x} \sum_{n \le x} \omega_k(n)^2 - \left(\lim_{x \to \infty} \frac{1}{x} \sum_{n \le x} \omega_k(n)\right)^2,$$

we can observe that the generating function from Theorem 3 provides a quick way to obtain the answers with no further input from number theory.

Corollary 6. The limiting distribution of  $\omega_k(n)$  has expectation  $\sum_{p} \frac{p-1}{p^{k+1}}$  and variance

$$\sum_{p} \frac{p-1}{p^{k+1}} \left( 1 - \frac{p-1}{p^{k+1}} \right).$$

**Remark 7.** Not surprisingly, these quantities are the expectation and variance of the sum of infinitely many Bernoulli random variables  $B_p$ , indexed by primes p, where  $B_p$  takes the value 1 with probability  $(p-1)/p^{k+1}$  (the density of those integers exactly divisible by  $p^k$ ).

These quantities are easy to calculate to reasonably high precision (see Section 4 for details); we record some numerical values in Table 2. The reader can confirm that the listed expectations are in good agreement with the quantities  $0e_{k,0}+1e_{k,1}+2e_{k,2}+3e_{k,3}$  as calculated from Table 1.

Table 2. Statistics of the limiting distribution of  $\omega_k(n)$ 

	expectation of $\omega_k(n)$	variance of $\omega_k(n)$
k=2	0.277484775	0.254931583
k = 3	0.097769500	0.093205673
k=4	0.041238122	0.040192048
k = 5	0.018684931	0.018433195

1.1. **Generalizations.** It turns out that our proof of Theorem 1 goes through for a far larger class of additive functions than just the  $\omega_k(n)$ . Given any sequence  $A = (a_1, a_2, a_3, \dots)$  of

complex numbers, define the additive function

(8) 
$$\omega_A(n) = \sum_{j=1}^{\infty} a_j \omega_j(n),$$

which is of course a finite sum for each integer n.

Remark 8. This definition generalizes all the examples we have seen so far:

- if  $a_i = 1$  always then  $\omega_A(n) = \omega(n)$ ;
- if  $a_i = j$  always then  $\omega_A(n) = \Omega(n)$ ;
- if  $a_j = j 1$  always then  $\omega_A(n) = \Omega(n) \omega(n)$ ;
- for a fixed positive integer k, if  $a_k = 1$  while  $a_j = 0$  for  $j \neq k$ , then  $\omega_A(n) = \omega_k(n)$ .

When  $a_1 \neq 0$ , classical techniques show that the large function  $\frac{1}{a_1}\omega_A(n)$  has the same Gaussian limiting distribution as  $\omega(n)$  and  $\Omega(n)$  when properly normalized (at least if the  $a_j$  do not grow too quickly). Therefore we restrict our attention to the smaller functions  $\omega_A(n)$  where  $a_1 = 0$ , which we expect to have limiting distributions without needing normalization. For  $m \in \mathbb{C}$ , define

$$\mathcal{N}_{A,m}(x) = \{ n \leqslant x \colon \omega_A(n) = m \}.$$

Our next result, which generalizes both equation (3) and Theorem 1, establishes the existence of the densities

(9) 
$$e_{A,m} = \lim_{x \to \infty} \frac{\# \mathcal{N}_{A,m}(x)}{x}$$

and provides a closed-form expression for them.

**Theorem 9.** Uniformly for all sequences  $A = (0, a_2, a_3, ...)$  of complex numbers with  $a_1 = 0$  and for all  $m \in \mathbb{C}$ ,

$$\#\mathcal{N}_{Am}(x) = e_{Am}x + O(x^{1/2}\log x)$$

with

(10) 
$$e_{A,m} = \frac{6}{\pi^2} \sum_{\substack{f \in \mathcal{F} \\ \omega_A(f) = m}} \frac{1}{f} \prod_{p|f} \left( 1 + \frac{1}{p} \right)^{-1}.$$

If we now restrict to the case where the  $a_j$  (and thus all values of  $\omega_A(n)$ ) are nonnegative integers, it once again makes sense to consider generating functions. Our next result generalizes both equation (4) and Theorem 3 in light of Remark 8.

**Theorem 10.** Let  $A = (0, a_2, a_3, ...)$  be a sequence of nonnegative integers. For all  $z \in \mathbb{C}$  with  $|z| \leq 1$ ,

(11) 
$$\sum_{m=0}^{\infty} e_{A,m} z^m = \prod_{p} \left( 1 - \frac{1}{p^2} + \sum_{j=2}^{\infty} z^{a_j} \left( \frac{1}{p^j} - \frac{1}{p^{j+1}} \right) \right).$$

Again this theorem shows that  $\omega_A(n)$  has a limiting distribution when the  $a_j$  are nonnegative integers, and we can therefore generalize Corollary 6; we record only the expectation for simplicity.

Corollary 11. Let  $A = (0, a_2, a_3, ...)$  be a sequence of nonnegative integers. The limiting distribution of  $\omega_A(n)$  has expectation  $\sum_{p} \sum_{j=2}^{\infty} a_j \left( \frac{1}{p^j} - \frac{1}{p^{j+1}} \right)$ .

**Remark 12.** It is certainly possible for this expectation to be infinite, as the example A = (0, 2, 4, 8, 16, ...) shows. In such cases  $\frac{1}{x} \sum_{n \leq x} \omega_A(n)$  grows too quickly for the mean value of  $\omega_A(n)$  to exist. Note, however, that Theorems 9 and 10 hold no matter how quickly the sequence A might grow.

We examine three specific examples of such sequences for the purposes of illustration: set S = (0, 1, 1, ...) and E = (0, 1, 0, 1, ...) and O = (0, 0, 1, 0, 1, 0, 1, ...). Then the corresponding omega functions are

$$\omega_S(n) = \sum_{j \geqslant 2} \omega_j(n)$$
 and  $\omega_E(n) = \sum_{\substack{j \geqslant 2 \ j \text{ even}}} \omega_j(n)$  and  $\omega_O(n) = \sum_{\substack{j \geqslant 3 \ j \text{ odd}}} \omega_j(n)$ 

which count, respectively, the number of primes dividing the powerful part of n (that is, the number of primes dividing n at least twice), the number of primes dividing n with even multiplicity, and the number of primes dividing n with odd multiplicity exceeding 1. For integers  $m \ge 0$ , let  $e_{S,m}$  and  $e_{E,m}$  and  $e_{O,m}$  be the corresponding densities defined in equation (9). An easy calculation of the right-hand side of equation (11) in these cases (for which each factor becomes a geometric series) yields the following generating functions:

Corollary 13. For all  $z \in \mathbb{C}$  with  $|z| \leq 1$ ,

$$\sum_{m=0}^{\infty} e_{S,m} z^m = \prod_{p} \left( 1 + \frac{z-1}{p^2} \right)$$

$$\sum_{m=0}^{\infty} e_{E,m} z^m = \prod_{p} \left( 1 + \frac{z-1}{p(p+1)} \right)$$

$$\sum_{m=0}^{\infty} e_{O,m} z^m = \prod_{p} \left( 1 + \frac{z-1}{p^2(p+1)} \right).$$

**Corollary 14.** For each fixed  $k \ge 2$ , we have  $e_{S,m} \le m^{-(2-o(1))m}$  and  $e_{E,m} \le m^{-(2-o(1))m}$  and  $e_{O,m} \le m^{-(3-o(1))m}$  as  $m \to \infty$ .

Remark 15. One interesting class of functions for which our methods accomplish less than desired are functions of the form  $\omega_A(n)$  where A contains integers but not necessarily only nonnegative integers. For example, if A = (0, 1, -1, 0, 0, ...) then  $\omega_A(n) = \omega_2(n) - \omega_3(n)$ , while if A = (0, 1, -1, 1, -1, ...) then  $\omega_A(n) = \omega_E(n) - \omega_O(n)$ . The target m = 0 is natural to investigate, as  $\omega_A(n) = 0$  in these two examples translates into  $\omega_2(n) = \omega_3(n)$  and  $\omega_E(n) = \omega_O(n)$ , respectively. While Theorem 9 gives a formula for the density of those integers n satisfying each of these equalities, our numerical techniques in Section 4 (which ultimately rely on being able to find the values of the derivatives of the appropriate generating function at z = 0) are not able to approach the question of good numerical approximations to these densities.

In Section 2 we establish Theorems 9 and 10, the formula and generating function for  $e_{A,m}$ , from which Theorems 1 and 3 follow as special cases. In Section 3 we deduce Corollaries 4 and 14 (the decay rates of  $e_{k,m}$  and certain variants) from Theorem 3 and Corollary 13. Finally, in Section 4 we describe the computations leading to the numerical values in Tables 1 and 2, as well as establishing Corollaries 6 and 11 concerning the expectation and variance of the additive functions under examination.

## 2. Closed form and generating function for the densities

We first prove Theorem 9, which will also establish the special case that is Theorem 1, by following the exposition of Rényi's result (3) in [7, Section 2.4]. Recall the notation of equation (8), and recall that  $\mathcal{F}$  denotes the set of powerful numbers.

**Lemma 16.** Uniformly for all sequences  $A = (a_1, a_2, ...)$  of complex numbers and all  $m \in \mathbb{C}$ ,

$$\sum_{\substack{f \leqslant x \\ f \in \mathcal{F} \\ \omega_A(f) = m}} \frac{1}{f^{1/2}} \prod_{p|f} (1 - p^{-1/2})^{-1} \ll \log x.$$

*Proof.* By dropping the condition  $\omega_A(f) = m$  and noting that  $f \leq x$  implies that all prime factors of f are at most x, we have by positivity

$$\sum_{\substack{f \leqslant x \\ f \in \mathcal{F} \\ \omega_A(f) = m}} \frac{1}{f^{1/2}} \prod_{p|f} (1 - p^{-1/2})^{-1} \leqslant \sum_{\substack{f \leqslant x \\ f \in \mathcal{F}}} \frac{1}{f^{1/2}} \prod_{p|f} (1 - p^{-1/2})^{-1} \leqslant \sum_{\substack{f \in \mathcal{F} \\ p|f \implies p \leqslant x}} \frac{1}{f^{1/2}} \prod_{p|f} (1 - p^{-1/2})^{-1}.$$

The right-hand side has an Euler product whose factors involve geometric series with common ratio  $p^{-1/2}$ :

$$\sum_{\substack{f \in \mathcal{F} \\ p \mid f \implies p \leqslant x}} \frac{1}{f^{1/2}} \prod_{p \mid f} (1 - p^{-1/2})^{-1} = \prod_{p \leqslant x} \left( 1 + \frac{(1 - p^{-1/2})^{-1}}{(p^2)^{1/2}} + \frac{(1 - p^{-1/2})^{-1}}{(p^3)^{1/2}} + \cdots \right)$$

$$= \prod_{p \leqslant x} \left( 1 + \frac{1}{(p^{1/2} - 1)^2} \right)$$

$$= \prod_{p \leqslant x} \left( 1 - \frac{1}{p} \right)^{-1} \prod_{p \leqslant x} \left( 1 + \frac{2}{p(p^{1/2} - 1)} \right);$$

this establishes the lemma, since the first product is asymptotic to a multiple of  $\log x$  as shown by Mertens, while the second is a convergent product of the form  $\prod_{p} (1 + O(p^{-3/2}))$ .

**Lemma 17.** Uniformly for all sequences  $A = (a_1, a_2, ...)$  of complex numbers and all  $m \in \mathbb{C}$ ,

$$\sum_{\substack{f>x\\f\in\mathcal{F}\\\omega_A(f)=m}}\frac{1}{f}\prod_{p\mid f}\left(1+\frac{1}{p}\right)^{-1}\ll x^{-1/2}.$$

*Proof.* Golomb [4] proved that the number of powerful numbers up to y is asymptotic to a constant times  $y^{1/2}$ . Thus for each integer  $r \ge 0$ ,

$$\sum_{\substack{2^r x < f \leqslant 2^{r+1} x \\ f \in \mathcal{F} \\ \omega_A(f) = m}} \frac{1}{f} \prod_{p|f} \left( 1 + \frac{1}{p} \right)^{-1} < \frac{1}{2^r x} \sum_{\substack{2^r x < f \leqslant 2^{r+1} x \\ f \in \mathcal{F}}} 1 \ll \frac{1}{2^r x} (2^{r+1} x)^{1/2} \ll 2^{-r/2} x^{-1/2},$$

and consequently

$$\sum_{\substack{f>x\\f\in\mathcal{F}\\\omega_A(f)=m}}\frac{1}{f}\prod_{p\mid f}\left(1+\frac{1}{p}\right)^{-1}=\sum_{r=0}^{\infty}\sum_{\substack{2^rx< f\leqslant 2^{r+1}x\\f\in\mathcal{F}\\\omega_A(f)=m}}\frac{1}{f}\prod_{p\mid f}\left(1+\frac{1}{p}\right)^{-1}\ll\sum_{r=0}^{\infty}2^{-r/2}x^{-1/2}\ll x^{-1/2}.\quad \Box$$

Proof of Theorem 9. Fix a sequence  $A = (0, a_2, a_3, ...)$  of complex numbers and a target  $m \in \mathbb{C}$ . Every positive integer n can be written uniquely as n = qf where q is squarefree, f is powerful, and (q, f) = 1 (indeed, q is the product of the primes dividing n exactly once). In this notation, the condition  $\omega_A(n) = m$  is equivalent to  $\omega_A(f) = m$  (since  $a_1 = 0$ ), and thus

(12) 
$$\# \mathcal{N}_{A,m}(x) = \sum_{\substack{f \leqslant x \\ f \in \mathcal{F} \\ \omega_A(f) = m}} \sum_{\substack{q \leqslant x/f \\ (q,f) = 1}} \mu^2(q).$$

To estimate the inner sum above, we use [7, Lemma 2.17] which says that for any  $y \ge 1$  and any positive integer f,

$$\sum_{\substack{n \le y \\ (n,f)=1}} \mu^2(n) = \frac{6}{\pi^2} y \prod_{p|f} \left(1 + \frac{1}{p}\right)^{-1} + O\left(y^{1/2} \prod_{p|f} (1 - p^{-1/2})^{-1}\right).$$

Inserting this asymptotic formula into equation (12) yields

$$\#\mathcal{N}_{A,m}(x) = \frac{6}{\pi^2} x \sum_{\substack{f \leqslant x \\ f \in \mathcal{F} \\ \omega_A(f) = m}} \frac{1}{f} \prod_{p|f} \left( 1 + \frac{1}{p} \right)^{-1} + O\left( x^{1/2} \sum_{\substack{f \leqslant x \\ f \in \mathcal{F} \\ \omega_A(f) = m}} \frac{1}{f^{1/2}} \prod_{p|f} (1 - p^{-1/2})^{-1} \right) \\
= \frac{6}{\pi^2} x \left( \sum_{\substack{f \in \mathcal{F} \\ \omega_A(f) = m}} \frac{1}{f} \prod_{p|f} \left( 1 + \frac{1}{p} \right)^{-1} - \sum_{\substack{f > x \\ f \in \mathcal{F} \\ \omega_A(f) = m}} \frac{1}{f} \prod_{p|f} \left( 1 + \frac{1}{p} \right)^{-1} \right) + O(x^{1/2} \log x) \\
= \frac{6}{\pi^2} x \left( \sum_{\substack{f \in \mathcal{F} \\ \omega_A(f) = m}} \frac{1}{f} \prod_{p|f} \left( 1 + \frac{1}{p} \right)^{-1} + O(x^{-1/2}) \right) + O(x^{1/2} \log x)$$

by Lemmas 16 and 17, which completes the proof of the theorem.

With Theorem 9 now established, it is a simple matter to prove Theorem 10, which will also establish the special case that is Theorem 3.

Proof of Theorem 10. Fix a sequence  $A = (0, a_2, a_3, ...)$  of nonnegative integers. Note that  $\sum_{m=0}^{\infty} e_{A,m} = 1$  (by the argument in Remark 2), and therefore  $\sum_{m=0}^{\infty} e_{A,m} z^m$  converges absolutely for any complex number z with  $|z| \leq 1$ . By Theorem 9,

$$\sum_{m=0}^{\infty} e_{A,m} z^m = \frac{6}{\pi^2} \sum_{m=0}^{\infty} z^m \sum_{\substack{f \in \mathcal{F} \\ \omega_A(f) = m}} \frac{1}{f} \prod_{p|f} \left( 1 + \frac{1}{p} \right)^{-1} = \frac{6}{\pi^2} \sum_{f \in \mathcal{F}} \frac{z^{\omega_A(f)}}{f} \prod_{p|f} \left( 1 + \frac{1}{p} \right)^{-1}.$$

Since  $z^{\omega_A(f)}/f = \prod_{p^j ||f} z^{\omega_A(p^j)}/p^j = \prod_{p^j ||f} z^{a_j}/p^j$ , the right-hand side equals its Euler product

$$\frac{6}{\pi^2} \sum_{f \in \mathcal{F}} \frac{z^{\omega_A(f)}}{f} \prod_{p|f} \left( 1 + \frac{1}{p} \right)^{-1} = \frac{6}{\pi^2} \prod_p \left( 1 + \left( 1 + \frac{1}{p} \right)^{-1} \sum_{j=2}^{\infty} \frac{z^{a_j}}{p^j} \right) \\
= \prod_p \left( 1 - \frac{1}{p^2} \right) \left( 1 + \left( 1 + \frac{1}{p} \right)^{-1} \sum_{j=2}^{\infty} \frac{z^{a_j}}{p^j} \right) \\
= \prod_p \left( 1 - \frac{1}{p^2} + \left( 1 - \frac{1}{p} \right) \sum_{j=2}^{\infty} \frac{z^{a_j}}{p^j} \right),$$

which is equal to the right-hand side of equation (11), thus establishing the theorem.

## 3. Decay rates of the densities

In this section, we deduce Corollary 4 from Theorem 3 and Corollary 14 from Theorem 10. The key step is to give a proposition establishing the rate of growth of infinite products such as those appearing in Theorems 3 and 10, which we do after the following simple lemma for the prime-counting function  $\pi(y)$  and its logarithmically weighted version  $\theta(y)$ .

**Lemma 18.**  $\pi(y) \log y - \theta(y) \sim y/\log y \text{ as } y \to \infty.$ 

*Proof.* By the prime number theorem,

$$\begin{split} \pi(y)\log y - \theta(y) &= \left(\mathrm{li}(y) + O(ye^{-c\sqrt{\log y}})\right)\log y - \left(y + O(ye^{-c\sqrt{\log y}})\right) \\ &= \left(\left(\frac{y}{\log y} + \frac{y}{\log^2 y} + O\left(\frac{y}{\log^3 y}\right)\right) + O\left(\frac{y}{\log^3 y}\right)\right)\log y - \left(y + O\left(\frac{y}{\log^2 y}\right)\right) \\ &= \frac{y}{\log y} + O\left(\frac{y}{\log^2 y}\right). \end{split}$$

**Proposition 19.** Fix a real number  $\kappa > 1$ , and let R(p) be a positive function defined on primes p such that  $R(p) \sim p^{-\kappa}$  as  $p \to \infty$ . Define the function  $P(x) = \prod_p (1 + R(p)x)$ . Then  $\log P(x) \approx x^{1/\kappa} / \log x$  as  $x \to \infty$ .

*Proof.* All implicit constants in this proof may depend on R(p) and  $\kappa$ . Choose  $p_0$  so that  $\frac{1}{2}p^{-\kappa} < R(p) < 2p^{-\kappa}$  for all  $p > p_0$ . We write

$$\log P(x) = \sum_{p \le p_0} \log (1 + R(p)x) + \sum_{p_0 x^{1/\kappa}} \log (1 + R(p)x)$$

$$= O(\log x) + \sum_{p_0 x^{1/\kappa}} \log (1 + R(p)x),$$
(13)

since the number of terms in the first sum, and the largest value of R(p) appearing in that sum, are both bounded in terms of the function R.

In the first sum in equation (13),

$$\frac{1}{2}p^{-\kappa}x < R(p)x < 1 + R(p)x < (x^{1/\kappa}/p)^{\kappa} + 2p^{-\kappa}x = 3p^{-\kappa}x.$$

Therefore

$$\sum_{p_0$$

The right-hand inequality is the same as

$$\sum_{p_0 
$$= \kappa \left( \pi(x^{1/\kappa}) \log(x^{1/\kappa}) - \theta(x^{1/\kappa}) \right) + \pi(x^{1/\kappa}) \log 3 + O(\log x)$$

$$\sim (\kappa + \log 3) \frac{x^{1/\kappa}}{\log(x^{1/\kappa})} = \kappa (\kappa + \log 3) \frac{x^{1/\kappa}}{\log x}$$$$

by Lemma 18. By the same calculation with  $\log \frac{1}{2}$  in place of  $\log 3$ ,

$$\sum_{p_0$$

(note that  $\kappa - \log 2 > 1 - \log 2$  is bounded away from 0). We conclude that

(14) 
$$\sum_{p_0$$

In the second sum in equation (13),

$$0 \leqslant \log(1 + R(p)x) \leqslant R(p)x < 2p^{-\kappa}x,$$

and thus by partial summation.

$$0 \leqslant \sum_{p>x^{1/\kappa}} \log(1+R(p)x) \leqslant \sum_{p>x^{1/\kappa}} 2p^{-\kappa}x$$

$$= 2x \int_{x^{1/\kappa}}^{\infty} t^{-\kappa} d\pi(t)$$

$$= 2x \left(\pi(t)t^{-\kappa}\right)_{x^{1/\kappa}}^{\infty} + \int_{x^{1/\kappa}}^{\infty} \kappa t^{-\kappa-1}\pi(t) dt.$$

The boundary term is well defined (since  $\kappa > 1$ ) and negative, and thus by the prime number theorem,

$$0 \leqslant \sum_{p > x^{1/\kappa}} \log \left( 1 + R(p)x \right) \ll x \left( 0 + \int_{x^{1/\kappa}}^{\infty} t^{-\kappa - 1} \frac{t}{\log t} dt \right)$$
$$\ll \frac{x}{\log x} \int_{x^{1/\kappa}}^{\infty} t^{-\kappa} dt = \frac{x}{\log x} \frac{(x^{1/\kappa})^{1-\kappa}}{\kappa - 1} \ll \frac{x^{1/\kappa}}{\log x}.$$

The proposition now follows by combining these inequalities with equations (13) and (14).

All that is left is to connect the rates of growth of the generating functions in Theorems 3 and 10 to the decay rate of their Maclaurin coefficients. We use the following classical information about entire functions [1, Definition 2.1.1 and Theorem 2.2.2]:

**Definition 20.** An entire function f(z) is said to be of order  $\rho$  if

$$\limsup_{r \to \infty} \frac{\log \log M_f(r)}{\log r} = \rho$$

where  $M_f(r) = \max_{|z|=r} |f(z)|$ . It is of finite order if it is of order  $\rho$  for some  $\rho \in \mathbb{R}$ .

**Lemma 21.** Let  $f(z) = \sum_{m=0}^{\infty} b_m z^m$  be an entire function. The function f(z) is of finite order if and only if

$$\mu = \limsup_{\substack{m \to \infty \\ b_m \neq 0}} \frac{m \log m}{\log(1/|b_m|)}$$

is finite, and in this case f(z) is of order  $\mu$ .

Proof of Corollary 4. Set

$$P(x) = \prod_{p} \left( 1 + \frac{p-1}{p^{k+1}} x \right)$$
 and  $Q(z) = \sum_{m=0}^{\infty} e_{k,m} z^m = \prod_{p} \left( 1 + \frac{p-1}{p^{k+1}} (z-1) \right)$ .

When |z| = r, note that

$$|Q(z)| \le \prod_{p} \left(1 + \frac{p-1}{p^{k+1}}(|z|+1)\right) = P(r+1);$$

thus by Proposition 19 with  $\kappa = k$  and  $R(p) = (p-1)/p^{k+1}$ ,

$$\log |Q(z)| \ll \frac{(r+1)^{1/k}}{\log(r+1)} \ll \frac{r^{1/k}}{\log r}.$$

On the other hand, when z = r > 3 is real, then

$$\log |Q(r)| = \log P(r-1) \gg \frac{(r-1)^{1/k}}{\log(r-1)} \gg \frac{r^{1/k}}{\log r}$$

again by Proposition 19. Together these last estimates show that  $\log M_Q(r) \approx r^{1/k}/\log r$ , which implies that

$$\limsup_{r \to \infty} \frac{\log \log M_Q(r)}{\log r} = \limsup_{r \to \infty} \frac{\log(r^{1/k}) - \log \log r + O(1)}{\log r} = \frac{1}{k}.$$

In particular, Q(z) has order  $\frac{1}{k}$  by Definition 20; consequently, by Lemma 21,

$$\limsup_{\substack{m \to \infty \\ e_{k,m} \neq 0}} \frac{m \log m}{\log(1/|e_{k,m}|)} = \frac{1}{k}.$$

We know that  $e_{k,m} > 0$  by Theorem 1, and so

$$\frac{m\log m}{\log(1/e_{k,m})} \leqslant \frac{1}{k} + o(1)$$

with asymptotic equality for infinitely many m; we conclude that

$$e_{k,m} \leqslant m^{-(k-o(1))m}$$

which completes the proof of the corollary.

*Proof of Corollary 13.* The proof is the same as the proof of Corollary 4, except that Q(z) is changed to each of the three products

$$\sum_{m=0}^{\infty} e_{S,m} z^m = \prod_{p} \left( 1 + \frac{z-1}{p^2} \right)$$

$$\sum_{m=0}^{\infty} e_{E,m} z^m = \prod_{p} \left( 1 + \frac{z-1}{p(p+1)} \right)$$

$$\sum_{m=0}^{\infty} e_{O,m} z^m = \prod_{p} \left( 1 + \frac{z-1}{p^2(p+1)} \right)$$

in turn, with corresponding modifications to P(x) and R(p); instead of with  $\kappa = k$ , the appeal to Proposition 19 is made with  $\kappa = 2$  in the first two cases and  $\kappa = 3$  in the last case, and the rest of the proof goes through in exactly the same way.

## 4. Numerical calculations of densities, expectations, and variances

We now describe how we used the generating functions in Theorem 3 to facilitate the calculation of the densities  $e_{k,m}$  in Table 1 to the indicated high level of precision. Our approach is based on observations of Marcus Lai (private communication).

**Proposition 22.** Let P(z) be any function with Maclaurin series

$$P(z) = \sum_{n=0}^{\infty} C(n)z^n,$$

so that  $C(n) = \frac{1}{n!}P^{(n)}(0)$  for every  $n \geq 0$ . Define

$$S(0,z) = \log P(z)$$
 and  $S(n,z) = \frac{d^n}{dz^n} S(0,z)$ 

for all  $n \ge 1$ ; and define S(n) = S(n,0) and  $\tilde{S}(n) = \frac{1}{(n-1)!}S(n)$ . Then for any  $n \ge 1$ ,

$$P^{(n)}(z) = \sum_{k=0}^{n-1} \binom{n-1}{k} P^{(k)}(z) S(n-k, z).$$

In particular, for  $n \geq 1$ ,

(15) 
$$C(n) = \frac{1}{n} \sum_{k=0}^{n-1} C(k)\tilde{S}(n-k),$$

so that for example

$$\begin{split} &C(1) = C(0)\tilde{S}(1) = P(0)\tilde{S}(1), \\ &C(2) = \frac{1}{2} \left( C(0)\tilde{S}(2) + C(1)\tilde{S}(1) \right) = \frac{P(0)}{2} \left( \tilde{S}(1)^2 + \tilde{S}(2) \right), \\ &C(3) = \frac{1}{3} \left( C(0)\tilde{S}(3) + C(1)\tilde{S}(2) + C(2)\tilde{S}(1) \right) = \frac{P(0)}{6} \left( \tilde{S}(1)^3 + 3\tilde{S}(1)\tilde{S}(2) + 2\tilde{S}(3) \right). \end{split}$$

*Proof.* We first verify that

$$P'(z) = P(z)\frac{P'(z)}{P(z)} = P(z)\frac{d}{dz}\log P(z) = P(z)\frac{d}{dz}S(0,z) = P(z)S(1,z),$$

which is the case n=1 of the first identity. The general case of the first identity now follows from using the product rule n-1 times in a row on this initial identity P'(z) = P(z)S(1,z). The second identity follows by plugging in z=0 into the first identity and recalling that  $C(n) = \frac{1}{n!}P^{(n)}(0)$ .

We apply this recursive formula (with subscripts inserted throughout the notation for clarity) with  $C(m) = e_{k,m}$ , so that

(16) 
$$P_k(z) = \sum_{m=0}^{\infty} e_{k,m} z^m = \prod_p \left( 1 + \frac{(p-1)(z-1)}{p^{k+1}} \right)$$

by Theorem 3. We compute

$$S_{k}(0,z) = \sum_{p} \log \left( 1 + \frac{(p-1)(z-1)}{p^{k+1}} \right)$$

$$(17) \quad S_{k}(1,z) = \frac{d}{dz} \sum_{p} \log \left( 1 + \frac{(p-1)(z-1)}{p^{k+1}} \right) = \sum_{p} \left( z + \frac{p^{k+1}}{p-1} - 1 \right)^{-1}$$

$$S_{k}(n,z) = \frac{d^{n-1}}{dz^{n-1}} \sum_{p} \left( z + \frac{p^{k+1}}{p-1} - 1 \right)^{-1} = \sum_{p} (-1)^{n-1} (n-1)! \left( z + \frac{p^{k+1}}{p-1} - 1 \right)^{-n},$$

so that

(18) 
$$\tilde{S}_k(n) = (-1)^{n-1} \sum_{p} \left( \frac{p^{k+1}}{p-1} - 1 \right)^{-n}.$$

Therefore equation (15) becomes

$$e_{k,m} = \frac{1}{m} \sum_{j=0}^{m-1} e_{k,j} \tilde{S}_k(m-j),$$

and in particular we have

$$e_{k,0} = \prod_{p} \left( 1 - \frac{p-1}{p^{k+1}} \right),$$

$$e_{k,1} = e_{k,0} \tilde{S}_k(1),$$

$$e_{k,2} = \frac{e_{k,0}}{2} \left( \tilde{S}_k(1)^2 + \tilde{S}_k(2) \right),$$

$$e_{k,3} = \frac{e_{k,0}}{6} \left( \tilde{S}_k(1)^3 + 3\tilde{S}_k(1)\tilde{S}_k(2) + 2\tilde{S}_k(3) \right).$$

Remark 23. We coded these formulas, and the one in equation (18), into SageMath and calculated approximations to them where we truncated the infinite product and sums to run over primes  $p \leq 10^7$ , resulting in the densities appearing in Table 1 (the cases k = 2, 3, 4, 5 and m = 0, 1, 2, 3). While we do not include a formal analysis of the error arising from these truncations, we have listed the densities to nine decimal places to display our confidence in that level of precision.

Finally, we extract the expectations and variances of various limiting distributions from their generating functions by relating those quantities to derivatives of their generating functions.

Proof of Corollary 6. Let  $X_k$  be the discrete random variable whose distribution is the same as the limiting distribution of  $\omega_k(n)$ ; then the generating function of this distribution is the function  $P_k(z)$  in equation (16). Proposition 22 and equations (16)–(17) tell us that

$$P'_{k}(1) = P_{k}(1)S_{k}(1,1) = 1 \cdot \sum_{p} \frac{p-1}{p^{k+1}} = \sum_{p} \frac{p-1}{p^{k+1}}$$

$$P''_{k}(1) = P'_{k}(1)S_{k}(1,1) + P_{k}(1)S_{k}(2,1)$$

$$= \sum_{p} \frac{p-1}{p^{k+1}} \cdot \sum_{p} \frac{p-1}{p^{k+1}} + 1\left(-\sum_{p} \left(\frac{p-1}{p^{k+1}}\right)^{2}\right) = \left(\sum_{p} \frac{p-1}{p^{k+1}}\right)^{2} - \sum_{p} \left(\frac{p-1}{p^{k+1}}\right)^{2}$$

But now by standard results from probability [3, Chapter XI, Theorems 2–3],

$$\mathbb{E}[X_k] = P_k'(1) = \sum_{p} \frac{p-1}{p^{k+1}}$$

$$\sigma^2[X_k] = P_k''(1) + P_k'(1) - P_k'(1)^2 = \sum_{p} \frac{p-1}{p^{k+1}} - \sum_{p} \left(\frac{p-1}{p^{k+1}}\right)^2$$

which is equivalent to the statement of the corollary.

**Remark 24.** As before, we used SageMath to calculate truncations of these infinite sums, running over primes  $p \leq 10^7$ , to generate the approximate expectations and variances listed in Table 2 for k = 2, 3, 4, 5.

Proof of Corollary 11. Let  $X_A$  be the random variable whose distribution is the same as the limiting distribution of  $\omega_A(n)$ . Using the same approach starting from the generating

function (10), we see that

$$\mathbb{E}[X_A] = P_A'(1) = \frac{d}{dz} \prod_p \left( 1 - \frac{1}{p^2} + \sum_{j=2}^{\infty} z^{a_j} \left( \frac{1}{p^j} - \frac{1}{p^{j+1}} \right) \right) \Big|_{z=1}$$

$$= \left\{ \prod_p \left( 1 - \frac{1}{p^2} + \sum_{j=2}^{\infty} z^{a_j} \left( \frac{1}{p^j} - \frac{1}{p^{j+1}} \right) \right) \sum_p \frac{d}{dz} \log \left( 1 - \frac{1}{p^2} + \sum_{j=2}^{\infty} z^{a_j} \left( \frac{1}{p^j} - \frac{1}{p^{j+1}} \right) \right) \right\} \Big|_{z=1}$$

$$= 1 \sum_p \left( 0 + \sum_{j=2}^{\infty} a_j z^{a_j - 1} \left( \frac{1}{p^j} - \frac{1}{p^{j+1}} \right) \right) \Big/ \left( 1 - \frac{1}{p^2} + \sum_{j=2}^{\infty} z^{a_j} \left( \frac{1}{p^j} - \frac{1}{p^{j+1}} \right) \right) \Big|_{z=1}$$

$$= \sum_p \sum_{j=2}^{\infty} a_j \left( \frac{1}{p^j} - \frac{1}{p^{j+1}} \right) \Big/ 1$$

as claimed.

#### ACKNOWLEDGMENTS

We thank Marcus Lai and Paul Péringuey for helpful discussions related to the calculations in Section 4, and we are grateful to the Pacific Institute for the Mathematical Sciences (PIMS) whose welcoming environment contributed to this project. The first author was supported by a University of Lethbridge postdoctoral fellowship. The second author was supported in part by a Natural Sciences and Engineering Council of Canada Discovery Grant.

#### References

- [1] R. P. Boas, Jr., Entire functions, Academic Press Inc. Publishers, New York, 1954.
- [2] E. Elma and Y-R. Liu, Number of prime factors with a given multiplicity, Canad. Math. Bull. 65 (2022), no. 1, 253–269.
- [3] W. Feller, An introduction to probability theory and its applications. Vol. I, third edition, John Wiley & Sons, Inc., New York–London–Sydney, 1968.
- [4] S. W. Golomb, *Powerful numbers*, Amer. Math. Monthly **77** (1970), 848–855.
- [5] G. H. Hardy and S. Ramanujan, *The normal number of prime factors of a number n*, Quart. J. Math. 48 (1917), 76–92. In *Collected papers of Srinivasa Ramanujan*, AMS Chelsea Publishing, Providence, RI, 2000, pp. 562–575.
- [6] G. H. Hardy and E. M. Wright, An introduction to the theory of numbers, sixth edition, revised by D. R. Heath-Brown and J. H. Silverman, Oxford University Press, Oxford, 2008.
- [7] H. L. Montgomery and R. C. Vaughan, *Multiplicative number theory I, Classical theory*, Cambridge Studies in Advanced Mathematics **97**, Cambridge University Press, Cambridge, 2007.
- [8] A. Rényi, On the density of certain sequences of integers, Acad. Serbe Sci. Publ. Inst. Math. 8 (1955), 157–162.

Department of Mathematics & Computer Science, University of Lethbridge, Faculty of Arts and Science, 4401 University Drive, Lethbridge, Alberta T1K 3M4, Canada

Email address: ertan.elma@uleth.ca

Department of Mathematics, University of British Columbia, Vancouver, British Columbia V6T 1Z2, Canada

Email address: gerg@math.ubc.ca