

Nonsmooth folds are tipping points.

D.J.W. Simpson

School of Mathematical and Computational Sciences
Massey University
Palmerston North, 4410
New Zealand

June 10, 2024

Abstract

A nonsmooth fold is where an equilibrium or limit cycle of a nonsmooth dynamical system hits a switching manifold and collides and annihilates with another solution of the same type. We show that beyond the bifurcation the leading-order truncation to the system in general has no bounded invariant set. This is proved for boundary equilibrium bifurcations of Filippov systems, hybrid systems, and continuous piecewise-smooth ODEs, and grazing-type events for which the truncated form is a continuous piecewise-linear map. The omitted higher-order terms are expected to be incapable of altering the local dynamics qualitatively, implying the system has no local invariant set on one side of a nonsmooth fold, and we demonstrate this with an example. Thus if the equilibrium or limit cycle is attracting the bifurcation causes the local attractor of the system to tip to a new state. The results also help explain global aspects of the bifurcation structures of the truncated systems.

1 Introduction

Bifurcations are critical parameter values at which the dynamical behaviour of a system changes in a fundamental way. For example oscillations in chemical kinetics are often onset by Hopf bifurcations [20], turbulent fluid flow is usually generated through period-doubling cascades [15], and Type I excitability in neurons is caused by SNIC bifurcations [21]. These examples require local smoothness in the equations of motion. If the equations are nonsmooth, additional bifurcations are possible [8].

Nonsmooth models commonly belong to one of three classes: piecewise-smooth continuous ODEs, Filippov systems, and hybrid systems. The phase space of these systems contain switching manifolds where the ODEs are non-differentiable, discontinuous, or a map is applied, Fig. 1. In each case a *boundary equilibrium bifurcation* (BEB) occurs when an equilibrium of a smooth component of the ODEs hits a switching manifold as parameters

are varied. Functionally the equilibrium is well-defined on both sides of the bifurcation, but only on one side of the bifurcation is it an equilibrium of the system where it is said to be *admissible*; on the other side of the bifurcation it is *virtual*.

Generic BEBs involve two equilibria, where for Filippov and hybrid systems the second equilibrium is a *pseudo-equilibrium* (an equilibrium of the sliding or sticking motion) [10]. Both equilibria are admissible on exactly one side of the bifurcation, and this immediately presents us with two cases: Either the equilibria are admissible on different sides of the bifurcation, as in Fig. 2, or the equilibria are admissible on the same side of the bifurcation, as in Fig. 3. The first case is termed *persistence*. The second case is termed a *nonsmooth fold* because if we consider only admissible solutions then at the bifurcation two equilibria collide and annihilate, analogous to a saddle-node bifurcation or fold.

BEBs readily generate other invariant sets growing out of the bifurcation point, such as limit cycles [17, 28] and chaotic sets [10, 25, 26], and the various transitions that BEBs bring about have been heavily studied. It is well known that the local dynamics is usually captured by the leading-order terms of a Taylor expansion of each smooth component of the system centred at the bifurcation point. This leads to a truncated form that is more amenable to an exact analysis. Indeed when analysing families of BEBs in mathematical models it is usually convenient to work with the truncated form in order to accurately identify codimension-two points.

In this paper we describe a simple property of BEBs that appears to have been overlooked. We show that on the side of a nonsmooth fold where both equilibria are virtual the system has no local invariant sets, at least for the truncated form. This result applies to piecewise-smooth continuous ODEs, Filippov systems, and hybrid systems, and with no restriction on the dimension of the system. Also, if a limit cycle has a degenerate interaction with a switching manifold, the same result applies when the associated Poincaré map is continuous and asymptotically piecewise-linear. This occurs for grazing-sliding bifurcations [9], corner




Figure 1: Schematic phase portraits of two-dimensional nonsmooth systems with one switching manifold (vertical line). In (a) the ODEs are continuous but non-differentiable on the switching manifold. In (b) the ODEs are discontinuous on the switching manifold and Filippov’s convention is used to specify sliding motion on the switching manifold. In (c) the system involves ODEs and a map (reset law) that instantaneously transports the system state from the top half of the switching manifold to the bottom half of the switching manifold. Such hybrid systems are commonly used to model mechanical systems with hard impacts [3, 4, 13].

collisions [7], and event collisions for piecewise-smooth ODEs with time-delay in the switching condition [24]. To prove the result, different calculations are required for each setting, but in each case there is a certain direction in which solutions evolve monotonically and hence eventually diverge. This is a consequence of piecewise-linearity, the absence of admissible equilibria, and a continuity constraint on the vector field or map (for Filippov systems the vector field can be viewed as continuous through tangency points).

We expect that in all settings if the truncated form has no local invariant set then neither




Figure 2: A bifurcation diagram and sample phase portraits of a piecewise-smooth continuous ODE system that experiences a persistence-type BEB as a parameter μ is varied. The blue curves indicate the location of the admissible equilibrium in relation to the switching manifold.




Figure 3: A bifurcation diagram and sample phase portraits of a piecewise-smooth continuous ODE system that experiences a nonsmooth fold BEB as a parameter μ is varied. To the left of the bifurcation the system has two admissible equilibria; to the right of the bifurcation it has no admissible equilibria. Throughout this paper stable solutions are coloured blue and unstable solutions are coloured red.

does the full system because the terms that have been omitted are higher order. In §4 we verify this for an example and outline the technical difficulties that need to be overcome in order to prove this in general. If true then any nonsmooth fold for which one of the two solutions is attracting has the features of a tipping point: By allowing the parameter that controls the bifurcation to vary slowly in time, the system state changes rapidly after it passes the nonsmooth fold. Fig. 4 illustrates this for the system

$$\begin{aligned}\dot{T} &= 1 - T - k(T, S)T, \\ \dot{S} &= \beta(\mu - S) - k(T, S)S,\end{aligned}\tag{1.1}$$

where $k(T, S) = \alpha\beta|T - S|$. This is a version of Stommel’s two-box model [29] for ocean circulation (T and S represent temperature and salinity differences). If μ is a fixed parameter, the system has a nonsmooth fold at $\mu = 1$. If μ allowed to decrease slowly, the system state heads rapidly to the upper equilibrium branch after passing $\mu = 1$. This has been studied further by Roberts and coworkers [22, 23] and Budd *et al.* [5].

It should be stressed that in general the absence of an attractor only occurs locally. For example in Fig. 5 the nonsmooth fold creates a large amplitude limit cycle. This is analogous to a SNIC bifurcation for smooth systems. Also the occurrence of two virtual stable equilibria can create stable oscillations [19, 32, 31]. For example Fig. 6 uses again (1.1), but now with $k(T, S) = 1$ if $-\alpha T + S > \varepsilon$, and $k(T, S) = 0$ otherwise (this is Welander’s ocean circulation model [33] in the discontinuous limit [18]). Orbits move towards one equilibrium until crossing the switching manifold, then move towards the other equilibrium until crossing the switching manifold, and so on, leading to a stable limit cycle.

The results of this paper were inspired by a 2002 paper of Carmona *et al.* [6]. Their work provided canonical forms for piecewise-linear ODEs arising in control circuitry. As a side note (their Proposition 19) they proved that if their two-piece canonical form has no admissible equilibria then it has no periodic solutions. We generalise their result beyond the




Figure 4: A bifurcation diagram of Stommel’s ocean circulation model with $\alpha = 5$ and $\beta = 0.2$. The blue and red curves are branches of stable and unstable equilibria. The black solution uses $\dot{\mu} = -0.01$ and experiences a tipping point by passing the nonsmooth fold at $\mu = 1$.

canonical form, from periodic solutions to arbitrary bounded invariant sets, place the result in the context of BEBs, and extend the result to Filippov systems, hybrid systems, and maps.

The remainder of the paper is organised as follows. In §2 we describe adjugate matrices as these are central to our main proofs. In §3 we prove the result for piecewise-linear maps, then in §4 explore the influence of higher order terms.

In §5 we treat piecewise-smooth continuous ODEs for which the result follows similarly to the map case. In §6 we treat piecewise-smooth discontinuous ODEs using Filippov’s convention, as is standard [8, 12, 14], to define evolution on switching manifolds. Then in §7 we treat hybrid systems. Specifically we consider hybrid models of impacting systems where the reset law models impact events. This is because such models have constraints that




Figure 5: A bifurcation diagram and sample phase portraits of a piecewise-smooth continuous ODE system that experiences a nonsmooth fold BEB as a parameter μ is varied. The bifurcation creates a large amplitude limit cycle due to global features of the dynamics.




Figure 6: A phase portrait of Welander’s ocean circulation model with $\alpha = 1.3$, $\beta = 0.2$, $\varepsilon = -0.4$, and $\mu = 1$. There are two virtual equilibria and a stable limit cycle (blue).

mean BEBs occur in a local fashion and correspond to either persistence or a nonsmooth fold [10]. Finally §8 presents a summary and discusses consequences to global bifurcations of parameterised families of truncated forms.

2 Adjugate matrices

Here we define adjugate matrices and briefly describe some of their properties. Further discussion can be found in standard textbooks [2, 16]. Throughout the paper I denotes the $n \times n$ identity matrix and e_1 denotes the first standard basis vector of \mathbb{R}^n .

Definition 1. The *adjugate* of a matrix $A \in \mathbb{R}^{n \times n}$ is the $n \times n$ matrix defined by $\text{adj}(A)_{ij} = (-1)^{i+j} m_{ji}$, where m_{ji} is the determinant of the $(n-1) \times (n-1)$ matrix obtained by removing the j^{th} row and i^{th} column from A .

The key property of the adjugate matrix is that for any $A \in \mathbb{R}^{n \times n}$,

$$A \text{adj}(A) = \text{adj}(A)A = \det(A)I. \quad (2.1)$$

So if A is invertible,

$$A^{-1} = \frac{\text{adj}(A)}{\det(A)}. \quad (2.2)$$

The first row of $\text{adj}(A)$ contains the values $(-1)^{i+j} m_{j1}$ which are independent of the entries in the first column of A . Thus if two $n \times n$ matrices differ only in their first columns, then the first rows of their adjugates will be identical. Algebraically this means

$$e_1^{\text{T}} \text{adj}(A) = e_1^{\text{T}} \text{adj}(A + ce_1^{\text{T}}) \quad (2.3)$$

for any $A \in \mathbb{R}^{n \times n}$ and $c \in \mathbb{R}^n$.

3 Maps

Let f be a piecewise- C^1 continuous map with vector variable $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ and scalar parameter $\mu \in \mathbb{R}$. Suppose f has one switching manifold and coordinates are chosen so that this manifold is $x_1 = 0$. Then f has the form

$$f(x; \mu) = \begin{cases} f^L(x; \mu), & x_1 \leq 0, \\ f^R(x; \mu), & x_1 \geq 0, \end{cases} \quad (3.1)$$

where f^L and f^R are C^1 .

As μ is varied a border-collision bifurcation occurs when a fixed point of f^L or f^R hits the switching manifold. If f is a Poincaré map of an piecewise-smooth ODE system, its fixed points correspond to periodic orbits and the border-collision bifurcation corresponds to a periodic orbit that interacts degenerately with a switching manifold of the ODEs [8]. Examples of this include grazing-sliding bifurcations [9], corner collisions [7], and event collisions [24].

Suppose f has a border-collision bifurcation at $x = 0$ when $\mu = 0$. Then $f^L(0; 0) = f^R(0; 0) = 0$, so we can write

$$f(x; \mu) = \begin{cases} A_L x + b\mu + E^L(x; \mu), & x_1 \leq 0, \\ A_R x + b\mu + E^R(x; \mu), & x_1 \geq 0, \end{cases} \quad (3.2)$$

where $A_L, A_R \in \mathbb{R}^{n \times n}$ differ only in their first columns (by continuity), $b \in \mathbb{R}^n$, and $E^L(x; \mu)$ and $E^R(x; \mu)$ contain only higher order terms. By dropping E^L and E^R we obtain the truncated form

$$g(x; \mu) = \begin{cases} A_L x + b\mu, & x_1 \leq 0, \\ A_R x + b\mu, & x_1 \geq 0, \end{cases} \quad (3.3)$$

which approximates f in a neighbourhood of $(x; \mu) = (0; 0)$. In the remainder of this section we study the truncated form g ; the general map f is revisited in the next section.

If $\det(I - A_L) \neq 0$ and $\det(I - A_R) \neq 0$, as is usually the case, the pieces of g have the unique fixed points

$$\begin{aligned} x^L(\mu) &= (I - A_L)^{-1} b\mu, \\ x^R(\mu) &= (I - A_R)^{-1} b\mu. \end{aligned} \quad (3.4)$$

The point $x^L(\mu)$ is admissible if $x_1^L(\mu) < 0$ and virtual if $x_1^L(\mu) > 0$. Similarly $x^R(\mu)$ is admissible if $x_1^R(\mu) > 0$ and virtual if $x_1^R(\mu) < 0$. We can now state the main result for maps.

Theorem 1. *If $\det(I - A_L) \neq 0$, $\det(I - A_R) \neq 0$, and both fixed points of (3.3) are virtual, then every forward orbit of (3.3) diverges.*

Below we prove Theorem 1 by showing there exists $s > 0$ and $p \in \mathbb{R}^n$ such that

$$p^\top g(x; \mu) \geq p^\top x + s, \quad (3.5)$$

for all $x \in \mathbb{R}^n$. This shows that the value of $p^\top x$ increases without bound along forward orbits of g , hence all forward orbits diverge.

As an example for which this can be seen graphically, consider (3.3) with

$$\begin{aligned} A_L &= \begin{bmatrix} \delta_L + 1 - \alpha & 1 \\ & -\delta_L & 0 \end{bmatrix}, \\ A_R &= \begin{bmatrix} \delta_R + 1 + \alpha & 1 \\ & -\delta_R & 0 \end{bmatrix}, \\ b &= \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \end{aligned} \quad (3.6)$$

where $\delta_L, \delta_R \in \mathbb{R}$ and $\alpha > 0$ are additional parameters. The fixed points are

$$\begin{aligned} x^L(\mu) &= \left(\frac{\mu}{\alpha}, \frac{-\delta_L \mu}{\alpha} \right), \\ x^R(\mu) &= \left(\frac{-\mu}{\alpha}, \frac{\delta_R \mu}{\alpha} \right), \end{aligned}$$

so are both virtual when $\mu > 0$. The inequality (3.5) holds for all $x \in \mathbb{R}^n$ using $s = \mu$ and $p^\top = [1 \ 1]$ (obtained by applying the formulas below to this example). Fig. 7 uses $\mu = 1$ and shades regions between integer values of $p^\top x$. The inequality (3.5) implies that no two points of an orbit of g belong to the same region, and this is evident for the sample orbit shown.

Proof of Theorem 1. By (2.2) applied to (3.4),

$$x_1^L(\mu) = \frac{p^\top b \mu}{\det(I - A_L)}, \quad (3.7)$$

where we let

$$p^\top = e_1^\top \text{adj}(I - A_L)$$

denote the first row of $\text{adj}(I - A_L)$. By (2.3) this is also the first row of $\text{adj}(I - A_R)$, so

$$x_1^R(\mu) = \frac{p^\top b \mu}{\det(I - A_R)}. \quad (3.8)$$

Let

$$s = p^\top b \mu.$$

Notice $s \neq 0$, for otherwise the fixed points are not virtual. Without loss of generality suppose $s > 0$. Then $\det(I - A_L) > 0$ and $\det(I - A_R) < 0$ because the fixed points are virtual.




Figure 7: A forward orbit of the piecewise-linear map (3.3) with (3.6) and $\delta_L = 1.2$, $\delta_R = -2.4$, $\alpha = 0.1$, and $\mu = 1$. The shaded regions are bounded by integer values of $p^\top x = x_1 + x_2$.

For any $x \in \mathbb{R}^n$ with $x_1 \leq 0$,

$$\begin{aligned} p^\top g(x; \mu) &= p^\top (A_L x + b\mu) \\ &= p^\top (-x + A_L x + x + b\mu) \\ &= -p^\top (I - A_L)x + p^\top x + s. \end{aligned}$$

By (2.1) we have $p^\top (I - A_L)x = \det(I - A_L)x_1$. Importantly $\det(I - A_L) > 0$ and $x_1 \leq 0$, hence

$$p^\top g(x; \mu) \geq p^\top x + s. \quad (3.9)$$

Similarly for any $x \in \mathbb{R}^n$ with $x_1 \geq 0$,

$$p^\top g(x; \mu) = -p^\top (I - A_R)x + p^\top x + s,$$

where $p^\top (I - A_R)x = \det(I - A_R)x_1 \leq 0$ because $\det(I - A_R) < 0$, so again we have (3.9). Thus for any forward orbit of g , the value of $p^\top x$ increases by at least $s > 0$ every iteration, so the orbit diverges. \square

4 The effect of higher order terms

The quantities E^L and E^R in (3.2) that were omitted to produce the truncated form (3.3) are higher order. To be precise, $\frac{\|E^L(x; \mu)\|}{\|x\| + |\mu|} \rightarrow 0$ and $\frac{\|E^R(x; \mu)\|}{\|x\| + |\mu|} \rightarrow 0$ as $(x; \mu) \rightarrow (0; 0)$, where $\|\cdot\|$ denotes the Euclidean norm on \mathbb{R}^n . For this reason if (3.3) satisfies the conditions of Theorem 1, we expect (3.2) has no local invariant set on one side of the nonsmooth fold. The following conjecture formalises this claim. Without loss of generality we suppose $p^\top b > 0$ and consider $\mu > 0$ in which case $\det(I - A_L) > 0$ and $\det(I - A_R) < 0$ are needed for both fixed points to be virtual by (3.7) and (3.8). We write $B_\eta = \{x \in \mathbb{R}^n \mid \|x\| \leq \eta\}$ for the closed ball of radius $\eta > 0$ centred at the origin.

Conjecture 2. *Let f be a map of the form (3.2) with $\det(I - A_L) > 0$, $\det(I - A_R) < 0$, and $p^\top b > 0$. Then there exists $\mu_0 > 0$ and $\eta > 0$ such that for all $\mu \in (0, \mu_0)$ and all $x \in B_\eta$ there exists $m \geq 1$ such that $\|f^m(x; \mu)\| > \eta$.*

That is, f has no invariant set in B_η for all $\mu \in (0, \mu_0)$. To illustrate Conjecture 2 we add the quadratic terms $E^L(x; \mu) = E^R(x; \mu) = \begin{bmatrix} 0 \\ -x_2^2 \end{bmatrix}$ to our previous example of (3.3) with (3.6) to obtain the map

$$f(x; \mu) = \begin{cases} \begin{bmatrix} (\delta_L + 1 - \alpha)x_1 + x_2 + \mu \\ -\delta_L x_1 - x_2^2 \end{bmatrix}, & x_1 \leq 0, \\ \begin{bmatrix} (\delta_R + 1 + \alpha)x_1 + x_2 + \mu \\ -\delta_R x_1 - x_2^2 \end{bmatrix}, & x_1 \geq 0. \end{cases} \quad (4.1)$$

Conjecture 2 is true for this map with the parameter values of Fig. 7, but to prove this it is insufficient to use the linear function $p^\top x = x_1 + x_2$. This is because

$$p^\top f(x; \mu) - p^\top x = \begin{cases} \mu - \alpha x_1 - x_2^2, & x_1 \leq 0, \\ \mu + \alpha x_1 - x_2^2, & x_1 \geq 0, \end{cases}$$

so for any $\eta > 0$ and $\mu \in (0, \eta^2)$, at the point $x = (0, \eta) \in B_\eta$ the difference $p^\top f(x; \mu) - p^\top x = \mu - \eta^2$ is negative. To prove Conjecture 2 for (4.1) we can instead use $\Phi(x) = x_1 + x_2 - 2x_2^2$, as shown in Appendix A. It remains for future work to show that such a function Φ can always be constructed, prove Conjecture 2 in different way, or find a counter-example.

5 Piecewise-smooth continuous ODEs

We now consider BEBs of piecewise-continuous ODEs. Our main result is a continuous-time analogue of Theorem 1.

Consider n -dimensional ODE systems of the form

$$\dot{x} = \begin{cases} f^L(x; \mu), & x_1 \leq 0, \\ f^R(x; \mu), & x_1 \geq 0, \end{cases} \quad (5.1)$$

where f^L and f^R are C^1 and $f^L(x; \mu) = f^R(x; \mu)$ at all points on the switching manifold $x_1 = 0$. Suppose (5.1) has a BEB at $x = 0$ when $\mu = 0$. Then by replacing f^L and f^R with their linearisations we obtain

$$\dot{x} = \begin{cases} A_L x + b\mu, & x_1 \leq 0, \\ A_R x + b\mu, & x_1 \geq 0, \end{cases} \quad (5.2)$$

where $A_L, A_R \in \mathbb{R}^{n \times n}$ differ in only their first columns (by continuity) and $b \in \mathbb{R}^n$. Solutions to (5.2) are well-defined for all $x \in \mathbb{R}^n$ and $t \in \mathbb{R}$ because (5.2) is Lipschitz and each piece is linear.

If $\det(A_L) \neq 0$ and $\det(A_R) \neq 0$ the pieces of (5.2) have the unique equilibria

$$\begin{aligned} x^L(\mu) &= -A_L^{-1}b\mu, \\ x^R(\mu) &= -A_R^{-1}b\mu. \end{aligned} \quad (5.3)$$

The equilibrium $x^L(\mu)$ is admissible if $x_1^L(\mu) < 0$ and virtual if $x_1^L(\mu) > 0$, while $x^R(\mu)$ is admissible if $x_1^R(\mu) > 0$ and virtual if $x_1^R(\mu) < 0$.

Theorem 3. *If $\det(A_L) \neq 0$, $\det(A_R) \neq 0$, and both equilibria of (5.2) are virtual, then every solution to (5.2) diverges as $t \rightarrow \pm\infty$.*

Proof. By (2.2) applied to (5.3),

$$x_1^L(\mu) = -\frac{q^\top b\mu}{\det(A_L)}, \quad (5.4)$$

where we let

$$q^\top = e_1^\top \text{adj}(A_L).$$

By (2.3) this is also the first row of $\text{adj}(A_R)$, so

$$x_1^R(\mu) = -\frac{q^\top b \mu}{\det(A_R)}. \quad (5.5)$$

The value

$$s = q^\top b \mu$$

cannot be zero because the equilibria are virtual. Without loss of generality suppose $s > 0$; then $\det(A_L) < 0$ and $\det(A_R) > 0$.

For any $x \in \mathbb{R}^n$ with $x_1 \leq 0$,

$$q^\top f^L(x; \mu) = q^\top (A_L x + b \mu) = q^\top A_L x + s.$$

Notice $q^\top A_L x = \det(A_L) x_1 \geq 0$, thus

$$q^\top \dot{x} \geq s. \quad (5.6)$$

Similarly for any $x \in \mathbb{R}^n$ with $x_1 \geq 0$,

$$q^\top f^R(x; \mu) = q^\top (A_R x + b \mu) = q^\top A_R x + s,$$

where $q^\top A_R x = \det(A_R) x_1 \geq 0$, so again we have (5.6). Thus any solution $\phi(t)$ to (5.2) has $\frac{d}{dt} q^\top \phi(t) \geq s > 0$ for all $t \in \mathbb{R}$, and so $\|\phi(t)\| \rightarrow \infty$ as $t \rightarrow \pm\infty$. \square

6 Filippov systems

Here we treat systems

$$\dot{x} = \begin{cases} f^L(x; \mu), & x_1 < 0, \\ f^R(x; \mu), & x_1 > 0, \end{cases} \quad (6.1)$$

where again f^L and f^R are C^1 , but now (6.1) is not necessarily continuous on the switching manifold $\Sigma = \{x \in \mathbb{R}^n \mid x_1 = 0\}$. To describe the dynamics of (6.1) it is useful to partition Σ into regions throughout which f^L and f^R each either direct solutions into Σ or away from Σ :

Definition 2. Consider a system (6.1) with fixed $\mu \in \mathbb{R}$. A subset $S \subset \Sigma$ is a

- i) a *crossing region* if $f_1^L(x; \mu) f_1^R(x; \mu) > 0$ for all $x \in S$, and
- ii) a *sliding region* if $f_1^L(x; \mu) f_1^R(x; \mu) < 0$ for all $x \in S$.

A sliding region is *attracting* if $f_1^L(x; \mu) > 0$ and $f_1^R(x; \mu) < 0$, and *repelling* if $f_1^L(x; \mu) < 0$ and $f_1^R(x; \mu) > 0$.

For example in Fig. 8 the top half of Σ is an attracting sliding region, while the bottom half of Σ is a crossing region.

To define solutions on sliding regions we use Filippov's convention [11, 12]. That is, solutions on sliding regions evolve according to $\dot{x} = f^S(x; \mu)$, where f^S is the *sliding vector field*

$$f^S(x; \mu) = \frac{f_1^L(x; \mu)f^R(x; \mu) - f_1^R(x; \mu)f^L(x; \mu)}{f_1^L(x; \mu) - f_1^R(x; \mu)}, \quad (6.2)$$

defined as the unique convex combination of f^L and f^R that is tangent to Σ . With this convention (6.1) is termed a *Filippov system*. Solutions are termed *Filippov solutions* and are concatenations of smooth segments of motion in $x_1 < 0$ under f^L , in $x_1 > 0$ under f^R , and on sliding regions under f^S .

Equilibria of f^L and f^R are admissible or virtual as in the previous section. They are sometimes called *regular equilibria* to distinguish them from equilibria of f^S :

Definition 3. A point $x \in \Sigma$ is a *pseudo-equilibrium* of (6.1) if $f^S(x; \mu) = 0$. It is *admissible* if it belongs to a sliding region, and *virtual* if it belongs to a crossing region.

Now suppose (6.1) has a BEB caused by an equilibrium of f^L hitting Σ at $x = 0$ when $\mu = 0$. Then $f^L(0; 0) = 0$, but unlike in the previous section typically $f^R(0; 0) \neq 0$. By replacing $f^L(x; \mu)$ with its linearisation about $(x; \mu) = (0; 0)$ and $f^R(x; \mu)$ with its value at $(x; \mu) = (0; 0)$, we obtain the truncated form

$$\dot{x} = \begin{cases} Ax + b\mu, & x_1 < 0, \\ c, & x_1 > 0, \end{cases} \quad (6.3)$$

where $A \in \mathbb{R}^{n \times n}$ and $b, c \in \mathbb{R}^n$. If $\det(A) \neq 0$ then f^L has the unique equilibrium

$$x^L(\mu) = -A^{-1}b\mu, \quad (6.4)$$




Figure 8: A schematic phase portrait of a two-dimensional Filippov system of the form (6.1).

which is admissible if $x_1^L < 0$ and virtual if $x_1^L > 0$. If $c_1 \neq 0$ and $q^\top c \neq 0$, where again

$$q^\top = e_1^\top \text{adj}(A),$$

then (6.3) has a unique pseudo-equilibrium $x^S(\mu)$ with $x^S(0) = 0$ [10, 26].

Theorem 4. *If $\det(A) \neq 0$, $c_1 \neq 0$, $q^\top c \neq 0$, and $x^L(\mu)$ and $x^S(\mu)$ are virtual, then every Filippov solution to (6.3) diverges as $t \rightarrow \infty$.*

Let us first give some intuition behind Theorem 4. As in the previous proof, below we obtain $q^\top f^L(x; \mu) > 0$ for points with $x_1 \leq 0$; but instead of f^R we work with f^S because the second equilibrium is a zero of f^S not f^R . Filippov solutions switch from following f^S to f^L by passing through points $x \in \Sigma$ for which $f_1^L(x; \mu) = 0$, and notice $f^S(x; \mu) = f^L(x; \mu)$ at such points by (6.2). This is a type of continuity constraint that together with the assumption that $x^S(\mu)$ is virtual, leads to the inequality $q^\top f^S(x; \mu) > 0$ throughout sliding regions and the conclusion that all Filippov solutions diverge.

Proof. By (2.2) applied to (6.4),

$$x_1^L(\mu) = -\frac{q^\top b\mu}{\det(A)}. \quad (6.5)$$

Thus the admissibility of $x^L(\mu)$ is determined by the signs of $\det(A)$ and

$$s = q^\top b\mu.$$

Notice $s \neq 0$ because $x^L(\mu)$ is assumed to be virtual.

We now derive an analogous characterisation for the admissibility of $x^S(\mu)$. Since $x^S(\mu)$ is zero of f^S ,

$$f_1^L(x^S(\mu); \mu) f^R(x^S(\mu); \mu) - f_1^R(x^S(\mu); \mu) f^L(x^S(\mu); \mu) = 0.$$

By multiplying both sides of this equation by q^\top , then solving for f_1^L , we obtain

$$f_1^L(x^S(\mu); \mu) = \frac{f_1^R(x^S(\mu); \mu) q^\top f^L(x^S(\mu); \mu)}{q^\top f^R(x^S(\mu); \mu)}. \quad (6.6)$$

Notice $f_R(x^S(\mu); \mu) = c$ and

$$\begin{aligned} q^\top f^L(x^S(\mu); \mu) &= q^\top (Ax^S(\mu) + b\mu) \\ &= \det(A)x_1^S(\mu) + s \\ &= s, \end{aligned}$$

where $x_1^S(\mu) = 0$ because $x^S(\mu) \in \Sigma$. Inserting these into (6.6) produces

$$f_1^L(x^S(\mu); \mu) = \frac{sc_1}{q^\top c}.$$

Thus the admissibility of $x^S(\mu)$ is determined by the sign of

$$f_1^L(x^S(\mu); \mu) f_1^R(x^S(\mu); \mu) = \frac{sc_1^2}{q^\top c}. \quad (6.7)$$

Without loss of generality suppose $s > 0$. Then $\det(A) < 0$, because $x^L(\mu)$ is virtual, and $q^\top c > 0$, because $x^S(\mu)$ is virtual. For any $x \in \mathbb{R}^n$ with $x_1 \leq 0$,

$$q^\top f^L(x; \mu) \geq s,$$

as in the previous section. For any $x \in \Sigma$ for which the denominator of (6.2) is nonzero,

$$\begin{aligned} q^\top f^S(x; \mu) &= \frac{f_1^L(x; \mu) q^\top c - c_1 q^\top f^L(x; \mu)}{f_1^L(x; \mu) - c_1} \\ &= \frac{s + r(x; \mu) q^\top c}{1 + r(x; \mu)}, \end{aligned}$$

where $q^\top f^L(x; \mu) = s$ because $x_1 = 0$, and we have let $r(x; \mu) = -\frac{f_1^L(x; \mu)}{c_1}$. If $f_1^L(x; \mu) c_1 \leq 0$, i.e. x does not belong to a crossing region, then $r(x; \mu) \geq 0$. Notice $q^\top f^S(x; \mu)$ ranges monotonically from $s > 0$ at $r = 0$ to $q^\top c > 0$ as $r \rightarrow \infty$. Thus at all $x \in \Sigma$ not on a crossing region the denominator of (6.2) is nonzero and

$$q^\top f^S(x; \mu) \geq \min(s, q^\top c) > 0.$$

Finally consider the behaviour of an arbitrary Filippov solution $\phi(t)$ to (6.3) as $t \rightarrow \infty$. If $c_1 < 0$ then $\phi(t)$ eventually evolves exclusively under f^L and f^S and diverges, while if $c_1 > 0$ then $\phi(t)$ either evolves exclusively under f^L and f^S and diverges, or eventually evolves exclusively under f^R and diverges. \square

7 Impacting hybrid systems

Finally we consider hybrid systems of the form

$$\begin{aligned} \dot{x} &= f(x; \mu), & \text{for } x_1 < 0, \\ x &\mapsto g(x; \mu), & \text{whenever } x_1 = 0, \end{aligned} \quad (7.1)$$

where f and g are C^1 . The following assumptions are motivated by the view that $x_1(t) < 0$ represents the displacement of a rigid object relative to a wall located at $x_1 = 0$. The map g is a reset law that models impacts as instantaneous events with velocity reversal and possibly energy loss.

Let $\Sigma = \{x \in \mathbb{R}^n \mid x_1 = 0\}$ denote the switching manifold, and

$$v(x; \mu) = f_1(x; \mu) \quad (7.2)$$

denote the velocity of the object relative to the wall. We partition Σ into the *incoming set* $\Sigma_{\text{in}} = \{x \in \Sigma \mid v(x; \mu) > 0\}$, the *outgoing set* $\Sigma_{\text{out}} = \{x \in \Sigma \mid v(x; \mu) < 0\}$, and the *grazing set*

$\Gamma = \{x \in \Sigma \mid v(x; \mu) = 0\}$, Fig. 9. We assume g maps Σ_{in} to Σ_{out} and is the identity map on Γ , thus

$$g(x; \mu) = x + v(x; \mu)h(x; \mu), \quad (7.3)$$

for some C^1 function h . The following definition uses

$$a(x; \mu) = \nabla v(x; \mu)^\top f(x; \mu), \quad (7.4)$$

which represents the acceleration of the object relative to the switching manifold.

Definition 4. Consider a system (7.1) with fixed $\mu \in \mathbb{R}$. A subset $S \subset \Gamma$ is a

- i) a *sticking region* if $a(x; \mu) > 0$ for all $x \in S$, and
- ii) a *detaching region* if $a(x; \mu) < 0$ for all $x \in S$.

Sticking regions and detaching regions are analogous to sliding regions and crossing regions of Filippov systems. On sticking regions it is natural to define the *sticking vector field*

$$f^{St}(x; \mu) = \left(I - \frac{h(x; \mu) \nabla v(x; \mu)^\top}{\nabla v(x; \mu)^\top h(x; \mu)} \right) f(x; \mu), \quad (7.5)$$

given by projecting f onto Γ by translating in the direction h (see Section 2.2.4 of di Bernardo *et al.* [8]). Solutions to (7.1) are concatenations of smooth segments of motion in $x_1 < 0$ under f , instantaneous jumps defined by g , and smooth segments of motion on sticking regions under f^{St} . Regular equilibria of (7.1) are zeros of f ; pseudo-equilibria of (7.1) are zeros of f^{St} :

Definition 5. A point $x \in \Gamma$ is a *pseudo-equilibrium* of (7.1) if $f^{St}(x; \mu) = 0$. It is *admissible* if it belongs to a sticking region, and *virtual* if it belongs to a detaching region.




Figure 9: A schematic phase portrait of a two-dimensional hybrid system of the form (7.1).

Now suppose (7.1) has a BEB caused by a regular equilibrium hitting Σ at $x = 0$ when $\mu = 0$. By replacing $f(x; \mu)$ with its linearisation about $(x; \mu) = (0; 0)$, and $h(x; \mu)$ with its value at $(x; \mu) = (0; 0)$, we obtain the truncated form

$$\begin{aligned} \dot{x} &= Ax + b\mu, \quad \text{for } x_1 < 0, \\ x &\mapsto x + v(x; \mu)c, \quad \text{whenever } x_1 = 0. \end{aligned} \tag{7.6}$$

where $A \in \mathbb{R}^{n \times n}$ and $b, c \in \mathbb{R}^n$ with $c_1 = 0$; also

$$v(x; \mu) = e_1^\top (Ax + b\mu). \tag{7.7}$$

The regular equilibrium $x^L(\mu)$ is given by (6.4), assuming $\det(A) \neq 0$. Note that some authors prefer an equivalent truncated form for which the regular equilibrium is fixed at origin [30].

Let $x' = x + v(x; \mu)c$ denote image of the reset law. Then

$$v(x'; \mu) = (1 + e_1^\top Ac)v(x; \mu),$$

and hence the condition

$$e_1^\top Ac < -1 \tag{7.8}$$

ensures the reset law maps the incoming set to the outgoing set. If $q^\top c \neq 0$ then (7.6) has a unique pseudo-equilibrium $x^{St}(\mu)$ with $x^{St}(0) = 0$ [10].

Theorem 5. *If $\det(A) \neq 0$, $e_1^\top Ac < -1$, $q^\top c \neq 0$, and $x^L(\mu)$ and $x^{St}(\mu)$ are virtual, then every solution to (7.6) diverges as $t \rightarrow \infty$.*

Notice $f^{St}(x; \mu) = f(x; \mu) - \frac{a(x; \mu)h(x; \mu)}{\nabla v(x; \mu)^\top h(x; \mu)}$, so at any point $x \in \Gamma$ for which $a(x; \mu) = 0$ we have $f^{St}(x; \mu) = f(x; \mu)$. This continuity underpins the following proof. The proof is similar to that of the previous section except we also need to consider the action of the reset law relative to the direction q (see (7.14)).

Proof. We first characterise the admissibility of $x^{St}(\mu)$. Observe $h(x; \mu) = c$ and $\nabla v(x; \mu)^\top = e_1^\top A$, so the sticking vector field (7.5) can be written as

$$f^{St}(x; \mu) = \frac{1}{e_1^\top Ac} (e_1^\top Ac f(x; \mu) - a(x; \mu)c). \tag{7.9}$$

Since $x^{St}(\mu)$ is a zero of f^{St} ,

$$0 = e_1^\top Ac f(x^{St}(\mu); \mu) - a(x^{St}(\mu); \mu)c. \tag{7.10}$$

By multiplying both sides of this equation by $q^\top = e_1^\top \text{adj}(A)$, then solving for a , we obtain

$$a(x^{St}(\mu); \mu) = \frac{e_1^\top Ac q^\top f(x^{St}(\mu); \mu)}{q^\top c}.$$

But $x_1^{St}(\mu) = 0$, so

$$\begin{aligned} q^\top f(x^{St}(\mu); \mu) &= q^\top (Ax^{St}(\mu) + b\mu) \\ &= \det(A)x_1^{St}(\mu) + q^\top b\mu \\ &= s, \end{aligned}$$

where we again let $s = q^\top b\mu$, hence

$$a(x^{St}(\mu); \mu) = \frac{e_1^\top Acs}{q^\top c}. \quad (7.11)$$

The first component of $x^L(\mu)$ is given by (6.5), so we require $s \neq 0$ because $x^L(\mu)$ is virtual. Without loss of generality suppose $s > 0$; then $\det(A) < 0$. Also $x^{St}(\mu)$ is virtual, meaning $a(x^{St}(\mu); \mu) < 0$, thus by (7.11) we have $q^\top c > 0$ (notice $e_1^\top Ac < 0$ by the theorem statement).

For any $x \in \mathbb{R}^n$ with $x_1 \leq 0$,

$$q^\top f(x; \mu) \geq s, \quad (7.12)$$

as in the proof of Theorem 3. By (7.9), for any $x \in \Gamma$

$$q^\top f^{St}(x; \mu) = q^\top f(x; \mu) - \frac{a(x; \mu)q^\top c}{e_1^\top Ac}.$$

So since $q^\top c > 0$ and $e_1^\top Ac < 0$, for any $x \in \Gamma$ with $a(x; \mu) \geq 0$, i.e. not in a detaching region, we have

$$q^\top f^{St}(x; \mu) \geq s, \quad (7.13)$$

using also (7.12). Finally, for any $x \in \Sigma_{\text{in}}$, let $x' = x + v(x; \mu)c$ and observe

$$q^\top (x' - x) = v(x; \mu)q^\top c > 0, \quad (7.14)$$

because $v(x; \mu) > 0$.

By (7.12), (7.13), and (7.14) any solution $\phi(t)$ to (7.6) satisfies $\frac{d}{dt}q^\top \phi(t) \geq s > 0$ while in $x_1 < 0$ and on sticking regions, plus whenever the reset law is applied the value of $q^\top \phi(t)$ increases, hence $q^\top \phi(t) \rightarrow \infty$ as $t \rightarrow \infty$, so the solution diverges. \square

8 Discussion

This paper has considered previously established truncated forms for BEBs and border-collision bifurcations and shown that if two equilibria or fixed points are virtual then all orbits diverge. It remains to determine if the same result holds for general systems, formulated as Conjecture 2 in the case of the maps. Implications to tipping points were discussed in §1. We finish by discussing implications to bifurcation structures of truncated forms.

The ‘robust chaos’ paper of Banerjee *et al.* [1] considered a family of two-dimensional piecewise-linear maps of the form (3.3) having $p^\top b\mu = 1$ fixed. Their main result was an identification of a parameter region where the fixed points $x^L(\mu)$ and $x^R(\mu)$ are admissible

saddles and the map has a chaotic attractor. One boundary of this region is $\det(I - A_L) = 0$ where $x^L(\mu)$ changes from admissible to virtual by (3.7). However, the attractor is unrelated to $x^L(\mu)$ so in fact it persists beyond this boundary [27].

Similarly crossing $\det(I - A_R) = 0$ rarely affects invariant sets other than $x^R(\mu)$. But once $\det(I - A_L) = 0$ and $\det(I - A_R) = 0$ have both been crossed Theorem 1 shows that the map can no longer have an attractor, or in fact any bounded invariant set. In this way the boundaries $\det(I - A_L) = 0$ and $\det(I - A_R) = 0$ *together* have a *global* effect on the dynamics. To illustrate this, Fig. 10 shows how the attractor of the three-dimensional border-collision normal form varies with two parameters (white represents the absence of an attractor). This form consists of (3.3) with

$$\begin{aligned} A_L &= \begin{bmatrix} \tau_L & 1 & 0 \\ -\sigma_L & 0 & 1 \\ \delta_L & 0 & 0 \end{bmatrix}, \\ A_R &= \begin{bmatrix} \tau_R & 1 & 0 \\ -\sigma_R & 0 & 1 \\ \delta_R & 0 & 0 \end{bmatrix}, \\ b &= \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}. \end{aligned} \tag{8.1}$$

Notice an attractor exists for parameter values arbitrarily close to the corner of the top-left quadrant where no attractor exists by Theorem 1. Assuming Fig. 10 is a typical two-dimensional slice of six-dimensional parameter space, as appears to be the case from numerical explorations, we can make the following conclusion. Individually the boundaries $\det(I - A_L) = 0$ and $\det(I - A_R) = 0$ do not affect the attractor, but together they form a codimension-two surface where the attractor is destroyed.

Acknowledgements

This work was supported by Marsden Fund contract MAU2209 managed by Royal Society Te Apārangi.

A Details of the higher order terms example

Here we prove Conjecture 2 for the example map (4.1). Specifically we show the following.

Proposition 6. *Consider the map (4.1) with $\delta_L > 0$, $\delta_R < 0$, and $\alpha > 0$. Then there exists $\eta > 0$ such that for any $\mu > 0$ and $x \in B_\eta$ there exists positive $m \leq \frac{6\eta}{\mu}$ such that $\|f^m(x; \mu)\| > \eta$.*

Proof. Let $\eta = \min\left(\frac{\alpha}{2\delta_L^2}, \frac{\alpha}{2\delta_R^2}, \frac{1}{\sqrt{2}}\right)$ and $\Phi(x) = x_1 + x_2 - 2x_2^2$. Notice $-4\eta \leq \Phi(x) \leq 2\eta$ for any $x \in B_\eta$, so over B_η the value of $\Phi(x)$ ranges by at most 6η . Using (4.1) direct calculations




Figure 10: A two-parameter bifurcation diagram of (3.3) with (8.1) and $\sigma_L = 0$, $\delta_L = 0.5$, $\sigma_R = 1$, and $\delta_R = 1.5$. This diagram was produced by computing 10^6 iterates of the forward orbit of $x = 0$ over a 500×500 equi-spaced grid of (τ_L, τ_R) values and using the last few iterates to predict the existence of an attractor and its period. The numerics identified a stable period- T solution with $T \leq 50$ in the coloured regions, a higher period or aperiodic solution in the black regions, and no attractor in the white regions.

give

$$\Phi(f(x; \mu)) - \Phi(x) = \begin{cases} \mu - (\alpha + 2\delta_L^2 x_1)x_1 - 4\delta_L x_1 x_2^2 + (1 - 2x_2^2)x_2^2, & x_1 \leq 0, \\ \mu + (\alpha - 2\delta_R^2 x_1)x_1 - 4\delta_R x_1 x_2^2 + (1 - 2x_2^2)x_2^2, & x_1 \geq 0. \end{cases} \quad (\text{A.1})$$

Notice $\Phi(f(x; \mu)) - \Phi(x) \geq \mu$ for any $\mu > 0$ and $x \in B_\eta$, using $\delta_L > 0$, $\delta_R < 0$, and $\alpha > 0$. So in B_η the value of Φ increases by at least μ under every iteration of f , thus forward orbits of f escape B_η within at most $\frac{6\eta}{\mu}$ iterations. \square

References

- [1] S. Banerjee, J.A. Yorke, and C. Grebogi. Robust chaos. *Phys. Rev. Lett.*, 80(14):3049–3052, 1998.
- [2] S.K. Berberian. *Linear Algebra*. Oxford University Press, New York, 1992.
- [3] B. Blazejczyk-Okolewska, K. Czolczynski, T. Kapitaniak, and J. Wojewoda. *Chaotic Mechanics in Systems with Impacts and Friction*. World Scientific, Singapore, 1999.
- [4] B. Brogliato. *Nonsmooth Mechanics: Models, Dynamics and Control*. Springer-Verlag, New York, 1999.

- [5] C. Budd, C. Griffith, and R. Kuske. Dynamic tipping in the non-smooth Stommel-box model, with fast oscillatory forcing. *Phys. D*, 432:132948, 2022.
- [6] V. Carmona, E. Freire, E. Ponce, and F. Torres. On simplifying and classifying piecewise-linear systems. *IEEE Trans. Circuits Systems I Fund. Theory Appl.*, 49(5):609–620, 2002.
- [7] M. di Bernardo, C.J. Budd, and A.R. Champneys. Corner collision implies border-collision bifurcation. *Phys. D*, 154:171–194, 2001.
- [8] M. di Bernardo, C.J. Budd, A.R. Champneys, and P. Kowalczyk. *Piecewise-smooth Dynamical Systems. Theory and Applications*. Springer-Verlag, New York, 2008.
- [9] M. di Bernardo, P. Kowalczyk, and A. Nordmark. Bifurcations of dynamical systems with sliding: Derivation of normal-form mappings. *Phys. D*, 170:175–205, 2002.
- [10] M. di Bernardo, A. Nordmark, and G. Olivar. Discontinuity-induced bifurcations of equilibria in piecewise-smooth and impacting dynamical systems. *Phys. D*, 237:119–136, 2008.
- [11] A.F. Filippov. Differential equations with discontinuous right-hand side. *Mat. Sb.*, 51(93):99–128, 1960. English transl. *Amer. Math. Soc. Transl.* 42(2):199–231, 1964.
- [12] A.F. Filippov. *Differential Equations with Discontinuous Righthand Sides*. Kluwer Academic Publishers., Norwell, 1988.
- [13] R.A. Ibrahim. *Vibro-Impact Dynamics.*, volume 43 of *Lecture Notes in Applied and Computational Mechanics*. Springer, New York, 2009.
- [14] M.R. Jeffrey. *Hidden Dynamics. The Mathematics of Switches, Decisions and Other Discontinuous Behaviour*. Springer, New York, 2018.
- [15] G.E. Karniadakis and G.S. Triantafyllou. Three-dimensional dynamics and transition to turbulence in the wake of bluff objects. *J. Fluid. Mech.*, 238:1–30, 1992.
- [16] B. Kolman. *Elementary Linear Algebra*. Prentice Hall, Upper Saddle River, NJ, 1996.
- [17] Yu.A. Kuznetsov, S. Rinaldi, and A. Gragnani. One-parameter bifurcations in planar Filippov systems. *Int. J. Bifurcation Chaos*, 13(8):2157–2188, 2003.
- [18] J. Leifeld. Non-smooth homoclinic bifurcation in a conceptual climate model. *Euro. J. Appl. Math.*, 29:891–904, 2018.
- [19] K.S. Morupisi and C. Budd. An analysis of the periodically forced PP04 climate model, using the theory of non-smooth dynamical systems. *IMA J. Appl. Math.*, 86(1):76–120, 2021.
- [20] P.H. Richter, P. Rehmus, and J. Ross. Control and dissipation in oscillatory chemical engines. *Prog. Theor. Phys.*, 66:385–405, 1981.

- [21] J. Rinzel and B. Ermentrout. Analysis of neural excitability and oscillations. In C. Koch and I. Segev, editors, *Methods in Neuronal Modeling.*, pages 251–291. MIT Press, Cambridge, MA, 1989.
- [22] A. Roberts and P. Glendinning. Canard-like phenomena in piecewise-smooth Van der Pol systems. *Chaos*, 24:023138, 2014.
- [23] A. Roberts and R. Saha. Relaxation oscillations in an idealized ocean circulation model. *Clim. Dyn.*, 48(7-8):2123–2134, 2017.
- [24] J. Sieber. Dynamics of delayed relay systems. *Nonlinearity*, 19(11):2489–2527, 2006.
- [25] D.J.W. Simpson. The instantaneous local transition of a stable equilibrium to a chaotic attractor in piecewise-smooth systems of differential equations. *Phys. Lett. A*, 380(38):3067–3072, 2016.
- [26] D.J.W. Simpson. A general framework for boundary equilibrium bifurcations of Filippov systems. *Chaos*, 28(10):103114, 2018.
- [27] D.J.W. Simpson. Detecting invariant expanding cones for generating word sets to identify chaos in piecewise-linear maps. *J. Difference Eq. Appl.*, 29:1094–1126, 2023.
- [28] D.J.W. Simpson and J.D. Meiss. Aspects of bifurcation theory for piecewise-smooth, continuous systems. *Phys. D*, 241(22):1861–1868, 2012.
- [29] H. Stommel. Thermohaline convection with two stable regimes of flow. *Tellus*, 13(2):224–230, 1961.
- [30] H. Tang and A. Champneys. Bifurcation of limit cycles from boundary equilibria in impacting hybrid systems. *SIAM J. Appl. Dyn. Sys.*, 22(4):3320–3357, 2023.
- [31] J. Walsh and E. Widiasih. A discontinuous ODE model of the glacial cycles with diffusive heat transport. *Mathematics*, 8:316, 2020.
- [32] J. Walsh, E. Widiasih, J. Hahn, and R. McGehee. Periodic orbits for a discontinuous vector field arising from a conceptual model of glacial cycles. *Nonlinearity*, 29(6):1843–1864, 2016.
- [33] P. Welander. A simple heat-salt oscillator. *Dyn. Atmos.*, 6(4):233–242, 1982.