

Convergence Analysis of Adaptive Gradient Methods under Refined Smoothness and Noise Assumptions

Devyani Maladkar*

Ruichen Jiang[†]

Aryan Mokhtari[†]

Abstract

Adaptive gradient methods are arguably the most successful optimization algorithms for neural network training. While it is well-known that adaptive gradient methods can achieve better dimensional dependence than stochastic gradient descent (SGD) under favorable geometry for stochastic convex optimization, the theoretical justification for their success in stochastic non-convex optimization remains elusive. In this paper, we aim to close this gap by analyzing the convergence rates of AdaGrad measured by the ℓ_1 -norm of the gradient. Specifically, when the objective has L -Lipschitz gradient and the stochastic gradient variance is bounded by σ^2 , we prove a worst-case convergence rate of $\tilde{O}(\frac{\sqrt{d}L}{\sqrt{T}} + \frac{\sqrt{d}\sigma}{T^{1/4}})$, where d is the dimension of the problem. We also present a lower bound of $\Omega(\frac{\sqrt{d}}{\sqrt{T}})$ for minimizing the gradient ℓ_1 -norm in the deterministic setting, showing the tightness of our upper bound in the noiseless case. Moreover, under more fine-grained assumptions on the smoothness structure of the objective and the gradient noise and under favorable gradient ℓ_1/ℓ_2 geometry, we show that AdaGrad can potentially shave a factor of \sqrt{d} compared to SGD. To the best of our knowledge, this is the first result for adaptive gradient methods that demonstrates a provable gain over SGD in the non-convex setting.

*Department of Computer Science, The University of Texas at Austin, Austin, TX, USA
{devyani.maladkar@utexas.edu}

[†]Department of Electrical and Computer Engineering, The University of Texas at Austin, Austin, TX, USA
{rjiang@utexas.edu, mokhtari@austin.utexas.edu}

1 Introduction

Adaptive methods, such as variants of AdaGrad [MS10; DHS11], are the workhorses of training massive neural networks and large language models. Their extensive applicability has spurred significant effort in identifying their convergence properties and complexity bounds. The major advantage of these algorithms, compared to classic stochastic gradient descent (SGD) methods [RM51], is that they do not require knowledge of any problem parameters, such as the gradient’s Lipschitz constant or the variance of stochastic gradients.

In this paper, our main focus is on the analysis of AdaGrad-type methods, which were proposed simultaneously by [MS10] and [DHS11], in the non-convex setting. Perhaps the most commonly analyzed AdaGrad-type method in the literature is the AdaGrad-Norm, which was first considered in [MS10]. Specifically, when we aim to minimize an objective function $F : \mathbb{R}^d \rightarrow \mathbb{R}$, AdaGrad-Norm updates the iterates \mathbf{w}_t according to the following update rule:

$$\mathbf{w}_{t+1} = \mathbf{w}_t - \frac{\eta}{b_t + \delta} \mathbf{g}_t, \quad (\text{AdaGrad-Norm})$$

where \mathbf{g}_t is the stochastic gradient of F at \mathbf{w}_t , the scalar η is a free constant, $\delta > 0$ is a small constant to ensure numerical stability, and $b_t^2 = \sum_{s=0}^t \|\mathbf{g}_s\|^2$. While this method has been extensively used in practice, most prior works demonstrated convergence similar to the guarantees obtained by SGD under stronger assumptions.

Specifically, it is well-known that in the non-convex setting, after T iterations, SGD can find a point such that the squared gradient ℓ_2 -norm is bounded above by $\mathcal{O}(1/\sqrt{T})$ (in expectation), when the objective function is smooth and the stochastic gradients are unbiased with bounded variance [GL13; BCN18]. This rate is, in fact, known to be optimal for any algorithm with only first-order oracle access under these assumptions [ACDFSW23]. Several works have managed to prove a similar convergence rate for AdaGrad-Norm, but under stronger assumptions, such as bounded gradients [WWB20; KLC21; GG22], the step-size being (conditionally) independent of the stochastic gradient [LO19; LO20], or sub-Gaussian noise [LO20; KLC21].

Recently, [FTCMSW22] addressed this issue and showed that under standard assumptions—Lipschitz gradients and bounded variance—AdaGrad-Norm achieves the complexity of SGD for finding an approximate first-order stationary point (up to a logarithmic factor). They further explored the setting where the stochastic gradient has affine variance. The extension to the generalized smoothness setting, where the Lipschitz constant for the gradient can grow with the norm of the gradient, was shown in [FRCS23].

However, all these results focus solely on the AdaGrad-Norm algorithm and do not offer any guarantees for the more commonly used variant known as AdaGrad. In AdaGrad, instead of selecting a universal step size for all coordinates by accumulating all gradient norms over time, a coordinate-specific step size is used. This is obtained by accumulating the squared norms of a specific coordinate of the gradient for the update of each coordinate. The updates are given by

$$w_{t+1,i} = w_{t,i} - \eta \frac{g_{t,i}}{b_{t,i} + \delta}, \quad \text{where } b_{t,i}^2 = b_{t-1,i}^2 + g_{t,i}^2 \quad \forall i \in \{1, 2, \dots, d\}, \quad (\text{AdaGrad})$$

where constant δ is introduced to ensure numerical stability. It should be noted that in some literature, the update in (AdaGrad) is referred to as diagonal AdaGrad or coordinate-wise AdaGrad, while the variant with full matrix inversion is called AdaGrad. Here, we refer to the diagonal version as AdaGrad, as it is the commonly used variant in practice.

Since in (AdaGrad) we tune a specific step size for each coordinate, unlike AdaGrad-Norm, it is expected to provide better convergence guarantees. However, the current convergence rate guarantees for these methods in the “non-convex setting” do not demonstrate this advantage. In fact, most results present a dimension-dependent bound for achieving a point with a small gradient ℓ_2 -norm, showing worse complexity compared to AdaGrad-Norm or SGD, even under stronger assumptions. Specifically, [CLSH19] showed that if the gradient and stochastic gradient ℓ_2 -norms are both uniformly bounded above, then the squared ℓ_2 -norm of the gradient converges at the rate of $\tilde{O}(d/\sqrt{T})$. The work in [DBBU22], under the assumption that the ℓ_∞ -norm of the stochastic gradients is uniformly almost surely bounded, showed that the squared ℓ_2 -norm of the gradient converges at a rate of $\tilde{O}(d/\sqrt{T})$. The work in [ZCCTYG18] studied the rate for the case that the ℓ_∞ -norm of the stochastic gradient is uniformly bounded above and showed that the squared ℓ_2 norm of the gradient is bounded above by $\tilde{O}(d/\sqrt{T})$. In this paper, we argue that since these analyses are not specific to coordinate-wise variations of adaptive methods, they fail to prove any improvement for AdaGrad compared to SGD or AdaGrad-Norm, and they are consequently doomed to a worse complexity bound.

Contributions. In this paper, we address this issue and provide a novel analysis for AdaGrad that takes into account its coordinate-specific structure. Specifically, we argue that the ℓ_1 -norm of the gradient is a more appropriate metric for capturing the convergence behavior of AdaGrad, leading to a tighter convergence guarantee than using the ℓ_2 -norm. We show that when the objective function has an L -Lipschitz gradient and the variance of the stochastic gradient is bounded above by σ^2 , in the worst case, after T iterations, AdaGrad ensures that ℓ_1 -norm of gradient is bounded above by $\tilde{O}\left(\frac{\sqrt{d}L}{\sqrt{T}} + \frac{\sqrt{d}\sigma}{T^{1/4}}\right)$, where d is the problem’s dimension. To the best of our knowledge, this is the best-known result for AdaGrad in the non-convex setting in terms of the ℓ_1 -norm. Additionally, we discuss the conditions under which the complexity of AdaGrad could be better than that of AdaGrad-Norm and SGD. To the best of our knowledge, this is the first result to show a provable gain for AdaGrad over AdaGrad-Norm and SGD in the non-convex setting. Finally, we establish a lower bound of $\Omega\left(\frac{\sqrt{d}}{\sqrt{T}}\right)$ for minimizing the ℓ_1 -norm of the gradient in the deterministic setting, demonstrating that a dimensional dependence bound is unavoidable. This result also illustrates the tightness of our analysis in the noiseless setting.

1.1 Related work

Recently, few papers have examined the convergence of adaptive gradient methods with respect to the gradient’s ℓ_1 -norm, which are more closely related to our work. In [LNNEN23], the authors provided a high probability convergence guarantee for (AdaGrad) under the stronger distributional assumption that the gradient noise is coordinate-wise subgaussian. They established a convergence rate of $\tilde{O}\left(\frac{dL}{\sqrt{T}} + \frac{d\sigma}{T^{1/4}}\right)$ in terms of the gradient ℓ_1 -norm, which is worse than ours by a factor of \sqrt{d} . Moreover, in a concurrent work [LL24], the authors analyzed the RMSprop method, another popular variant of adaptive gradient methods, under the standard smoothness assumption and coordinate-wise bounded noise variance assumption. They established a convergence rate of $\tilde{O}\left(\frac{\sqrt{d}}{\sqrt{T}} + \frac{\sqrt{d}}{T^{1/4}}\right)$, which matches our bound in the worst case setting. However, their convergence result only showed the possibility of matching the convergence rate of SGD instead of surpassing it, and thus it did not fully explain the advantage of adaptive gradient methods. In our work, we present a more fine-grained analysis of Adagrad under the refined smoothness assumption, which enables us to demonstrate a provable gain for AdaGrad over SGD. To the best of our knowledge, our work

presents the first theoretical result in non-convex optimization showing the advantage of coordinate-wise adaptive methods over SGD.

Notation. In this paper, we use boldface letters for vectors and normal font letters for scalars. The Euclidean or ℓ_2 norm of a vector \mathbf{w} is denoted by $\|\mathbf{w}\|_2$ and its ℓ_1 norm is indicated by $\|\mathbf{w}\|_1$. For a vector $\mathbf{w} \in \mathbb{R}^d$, we denote its i -th coordinate by w_i . For any integer $n \geq 1$, We use $[n]$ to denote the set $\{1, 2, \dots, n\}$. Further, we use \mathcal{F}_t as the sigma algebra generated after time index t . In our case, \mathcal{F}_t contains all iterates $\mathbf{w}_0, \dots, \mathbf{w}_{t+1}$ and all stochastic gradients $\mathbf{g}_0, \dots, \mathbf{g}_t$. In this paper, the notation $\hat{\mathcal{O}}$ hides logarithmic dependence.

2 Preliminaries

In this section, we present the assumptions and definitions required to outline our results. As mentioned earlier, our objective is to identify an approximate stationary point of a non-convex function $F : \mathbb{R}^d \rightarrow \mathbb{R}$ over the unbounded domain \mathbb{R}^d . To characterize the complexity of AdaGrad in achieving this goal, we require some mild conditions on the objective function F and its stochastic gradient, which are customary for the analysis of stochastic gradient-based methods. We introduce these assumptions and some necessary definitions here. Specifically, we adopt fine-grained assumptions similar to those in [BWAA18], which allow for a clearer analysis of how coordinate-wise AdaGrad exploits structural features.

Assumption 2.1. *The stochastic gradient \mathbf{g}_t is an unbiased estimate of the exact gradient $\nabla F(\mathbf{w}_t)$, i.e., $\mathbb{E}[\mathbf{g}_t | \mathcal{F}_{t-1}] = \nabla F(\mathbf{w}_t)$.*

Assumption 2.2. *The stochastic gradient \mathbf{g}_t with elements $[g_{t,1}, \dots, g_{t,d}]$ has a coordinate-wise bounded variance. In other words, for all $i \in [d]$, we have*

$$\mathbb{E}[|g_{t,i} - \nabla_i F(\mathbf{w}_t)|^2 | \mathcal{F}_{t-1}] \leq \sigma_i^2,$$

where σ_i is a fixed non-negative constant corresponding to the variance of the i -th coordinate of stochastic gradient \mathbf{g}_t , and $\nabla_i F(\mathbf{w}_t)$ represents the i -th coordinate of the gradient $\nabla F(\mathbf{w}_t)$. Moreover, we define the vector $\boldsymbol{\sigma}$ as $\boldsymbol{\sigma} = [\sigma_1, \sigma_2, \dots, \sigma_d] \in \mathbb{R}^d$.

The above condition on the variance of the stochastic gradient is a more fine-grained assumption compared to the standard assumptions used in the literature to analyze stochastic gradient methods. The common assumption is given by $\mathbb{E}[\|\mathbf{g}_t - \nabla F(\mathbf{w}_t)\|^2 | \mathcal{F}_{t-1}] \leq \sigma^2$. Indeed, our considered assumption is more refined as it implies the standard assumption when we consider $\sigma^2 = \sum_{i=1}^d \sigma_i^2$. As discussed earlier, since we aim to study an algorithm with a coordinate-specific update, the above assumption better captures the convergence behavior of the method of interest.

Assumption 2.3. *The function $F(\cdot)$ is coordinate wise smooth, i.e., for any vectors $\mathbf{x}, \mathbf{y} \in \mathbb{R}^d$*

$$|F(\mathbf{x}) - F(\mathbf{y}) - \langle \nabla F(\mathbf{x}), \mathbf{x} - \mathbf{y} \rangle| \leq \sum_{i=1}^d \frac{L_i}{2} |x_i - y_i|^2,$$

where the constant $L_i \geq 0$ is the Lipschitz constant of the gradient associated with the i -th coordinate. Moreover, we define the vector \mathbf{L} as $\mathbf{L} = [L_1, L_2, \dots, L_d] \in \mathbb{R}^d$.

This is similar to the fine-grained assumptions made in the literature for coordinate-wise analysis of algorithms [RT11; BWAA18]. We recover the standard smoothness assumption by considering the Lipschitz constant as $L := \max_i L_i = \|\mathbf{L}\|_\infty$.

3 Convergence analysis

In this section, we present our convergence guarantees for (AdaGrad). Specifically, we upper bound the number of iterations required for both algorithms to find a near-stationary point in terms of ℓ_1 -norm, instead of the more conventional ℓ_2 -norm. As discussed earlier, this measure of optimality is more suitable given the coordinate-specific structure of (AdaGrad) and it better highlights the advantages compared to the AdaGrad-Norm method, as we will demonstrate.

Our starting point is applying Assumption 2.3 to two consecutive iterates \mathbf{w}_t and \mathbf{w}_{t+1} to obtain

$$\begin{aligned} F(\mathbf{w}_{t+1}) &\leq F(\mathbf{w}_t) + \langle \nabla F(\mathbf{w}_t), \mathbf{w}_{t+1} - \mathbf{w}_t \rangle + \sum_{i=1}^d \frac{L_i}{2} |w_{t+1,i} - w_{t,i}|^2 \\ &= F(\mathbf{w}_t) - \sum_{i=1}^d \eta_{t,i} \nabla_i F(\mathbf{w}_t) g_{t,i} + \sum_{i=1}^d \frac{L_i}{2} \eta_{t,i}^2 g_{t,i}^2, \end{aligned} \quad (1)$$

where we used the fact that $w_{t+1,i} = w_{t,i} - \eta_{t,i} g_{t,i}$ in the last equality. If the step size $\eta_{t,i}$ were conditionally independent of the stochastic gradient $g_{t,i}$, then by taking the conditional expectation with respect to \mathcal{F}_{t-1} , the second term in the right-hand side of (1) would result in $-\eta_{t,i} \nabla_i F(\mathbf{w}_t) \mathbb{E}[g_{t,i} | \mathcal{F}_{t-1}] = -\eta_{t,i} \nabla_i F(\mathbf{w}_t)^2$ by Assumption 2.1. However, one of the main challenges of analyzing adaptive gradient methods is the correlation between $\eta_{t,i}$ and $g_{t,i}$. Specifically, by the update rule in (AdaGrad) we can also write the step sizes as

$$\eta_{t,i} := \frac{\eta}{\sqrt{b_{t-1,i}^2 + g_{t,i}^2 + \delta}} \quad (2)$$

which depends on the stochastic gradient at the current iterate \mathbf{w}_t . Consequently, this dependence implies that $\mathbb{E}[\eta_{t,i} g_{t,i} | \mathcal{F}_{t-1}] \neq \eta_{t,i} \mathbb{E}[g_{t,i} | \mathcal{F}_{t-1}]$ in general, which complicates the analysis. Therefore, it is crucial to introduce a “proxy” of the step size that is decorrelated from \mathbf{g}_t , ensuring that with respect to the filtration \mathcal{F}_{t-1} , it remains deterministic. We refer to this as the “decorrelated step size” [WWB20; FTCMSW22], which we will formally define next.

Definition 3.1 (Decorrelated step sizes). *For any $i \in [d]$ and $t \geq 1$, the decorrelated step size is defined as*

$$\hat{\eta}_{t,i} = \frac{\eta}{\sqrt{b_{t-1,i}^2 + \sigma_i^2 + \nabla_i F(\mathbf{w}_t)^2 + \delta}},$$

Based on Definition 3.1, in the decorrelated step size $\hat{\eta}_{t,i}$ we replace the stochastic gradient $g_{t,i}^2$ by $\nabla_i F(\mathbf{w}_t)^2 + \sigma_i^2$. Consequently, $\hat{\eta}_{t,i}$ belongs to the filtration \mathcal{F}_{t-1} and thus $\mathbb{E}[\hat{\eta}_{t,i} \nabla_i F(\mathbf{w}_t) g_{t,i} | \mathcal{F}_{t-1}] = \hat{\eta}_{t,i} \nabla_i F(\mathbf{w}_t)^2$, leading to the desired squared gradient that we aim to bound. Equipped with the decorrelated step size, in the following lemma we prove an upper bound on a (weighted) gradient square norm at the current iterate \mathbf{w}_t . Its proof can be found in Appendix A.1.

Lemma 3.1. *Suppose Assumptions 2.1 and 2.3 hold. Consider the update rules given in (AdaGrad) and recall the decorrelated step sizes defined in Definition 3.1. Then we have*

$$\begin{aligned} \sum_{i=1}^d \hat{\eta}_{t,i} \nabla_i F(\mathbf{w}_t)^2 &\leq \mathbb{E}[F(\mathbf{w}_{t+1}) | \mathcal{F}_{t-1}] - F(\mathbf{w}_t) \\ &\quad + \sum_{i=1}^d \mathbb{E}[(\hat{\eta}_{t,i} - \eta_{t,i}) \nabla_i F(\mathbf{w}_t) g_{t,i} | \mathcal{F}_{t-1}] + \sum_{i=1}^d \frac{L_i}{2} \mathbb{E}[\eta_{t,i}^2 g_{t,i}^2 | \mathcal{F}_{t-1}]. \end{aligned} \quad (3)$$

In Lemma 3.1, the left-hand side is a weighted version of the squared gradient norm at \mathbf{w}_t , where the weights for each coordinate are given by the decorrelated step sizes $\hat{\eta}_{t,i}$. Note that this is the key difference compared to the analysis of (AdaGrad-Norm) in [FTCMSW22]. Indeed, for (AdaGrad-Norm), the left-hand side will become $\hat{\eta}_t \|\nabla F(\mathbf{w}_t)\|^2$, and thus the squared ℓ_2 -norm of the gradient naturally arises from the analysis. On the other hand, as we shall see later, in our case ℓ_2 -norm is not the best choice of the norm and instead we will relate the left-hand side in (3) to the ℓ_1 -norm of the gradient.

In light of Lemma 3.1, we need to manage the *bias term* $\sum_{i=1}^d \mathbb{E}[(\hat{\eta}_{t,i} - \eta_{t,i}) \nabla_i F(\mathbf{w}_t) g_{t,i} \mid \mathcal{F}_{t-1}]$, which is due to the difference between the step size $\eta_{t,i}$ and its decorrelated version $\hat{\eta}_{t,i}$, and a *quadratic term* $\sum_{i=1}^d \mathbb{E}[\eta_{t,i}^2 g_{t,i}^2]$, which comes from Assumption 2.3. The following lemma addresses these two terms and the proofs for these two results are presented in Appendix A.2. Note that similar terms also appear in the analysis of (AdaGrad-Norm) [FTCMSW22], and ours can be viewed as a refined coordinate-wise counterpart of the previous analysis.

Lemma 3.2. *Consider the update rule in (AdaGrad). For any $t \in [T]$ and $i \in [d]$, we have*

$$\mathbb{E}[(\hat{\eta}_{t,i} - \eta_{t,i}) \nabla_i F(\mathbf{w}_t) g_{t,i} \mid \mathcal{F}_{t-1}] \leq \frac{\hat{\eta}_{t,i}}{2} \nabla_i F(\mathbf{w}_t)^2 + \frac{2\sigma_i}{\eta} \mathbb{E}[\eta_{t,i}^2 g_{t,i}^2 \mid \mathcal{F}_{t-1}]. \quad (4)$$

Moreover, we have

$$\mathbb{E} \left[\sum_{t=1}^T \eta_{t,i}^2 g_{t,i}^2 \right] \leq \eta^2 \log h_i(T), \quad (5)$$

where $h_i(T) = 1 + \frac{T\sigma_i^2}{\delta^2} + \frac{T(\|\nabla F(\mathbf{w}_1)\| + \eta\sqrt{d}L_i T)^2}{\delta^2} = \mathcal{O}(dT^3)$.

The first result in Lemma 3.2 shows that for each coordinate $i \in [d]$, we can upper bound the bias term in terms of the squared gradient $\frac{\hat{\eta}_{t,i}}{2} \nabla_i F(\mathbf{w}_t)^2$ and the quadratic term $\mathbb{E}[\eta_{t,i}^2 g_{t,i}^2]$. The second result in the above lemma shows that the accumulation of the quadratic terms $\eta_{t,i}^2 g_{t,i}^2$ over T iterations can be bounded in expectation by $\mathcal{O}(\eta^2 \log(dT))$. By combining Lemma 3.2 with Lemma 3.1, we obtain the following key corollary.

Corollary 3.3. *Define $h(T) = 1 + \frac{T\|\sigma\|_\infty^2}{\delta^2} + \frac{T(\|\nabla F(\mathbf{w}_1)\| + \eta\sqrt{d}\|\mathbf{L}\|_\infty T)^2}{\delta^2}$. For (AdaGrad), we have*

$$\mathbb{E} \left[\sum_{t=1}^T \sum_{i=1}^d \frac{\hat{\eta}_{t,i}}{2} \nabla_i F(\mathbf{w}_t)^2 \right] \leq F(\mathbf{w}_1) - F^* + \left(2\eta\|\sigma\|_1 + \frac{\eta^2\|\mathbf{L}\|_1}{2} \right) \log h(T). \quad (6)$$

To simplify the notation, let us denote the right-hand side of (6) by Q . This implies that, if we ignore the logarithmic term, we have $Q = \tilde{\mathcal{O}}(F(\mathbf{w}_1) - F^* + \eta\|\sigma\|_1 + \eta^2\|\mathbf{L}\|_1)$. Corollary 3.3 shows that the sum of weighted squared gradient norms is bounded by a constant depending on problem parameters, up to log factors. Hence, the remaining task is to establish lower bounds on the step sizes $\hat{\eta}_{t,i}$. For instance, if we were able to show that all the step sizes $\hat{\eta}_{t,i}$ are uniformly lower bounded by $\tilde{\Omega}(\frac{1}{\sqrt{T}})$, then Corollary 3.3 would immediately imply a rate of $\tilde{\mathcal{O}}(\frac{1}{T^{1/4}})$ in terms of the gradient ℓ_2 -norm $\|\nabla F(\mathbf{w}_t)\|_2$. However, there are several challenges: (i) The step sizes $\hat{\eta}_{t,i}$ are determined by the observed stochastic gradient rather than specified by the user. (ii) To further complicate the issue, due to correlation between the step size $\hat{\eta}_{t,i}$ and the iterate \mathbf{w}_t , this implies that $\mathbb{E}[\hat{\eta}_{t,i} \nabla_i F(\mathbf{w}_t)^2] \neq \mathbb{E}[\hat{\eta}_{t,i}] \mathbb{E}[\nabla_i F(\mathbf{w}_t)^2]$ and hence a lower bound on $\mathbb{E}[\hat{\eta}_{t,i}]$ would not suffice. (iii) Finally, since the step sizes for each coordinate are updated independently, it is unclear how to construct a uniform lower bound across all the coordinates.

To address the last challenge, we study each coordinate and construct a uniform lower bound on $\hat{\eta}_{t,i}$ for $t \in [T]$. Specifically, for each coordinate $i \in [d]$, we define a new auxiliary step size $\tilde{\eta}_{T,i}$ as

$$\tilde{\eta}_{T,i} = \frac{\eta}{\sqrt{\sum_{i=1}^{T-1} g_{t,i}^2 + \sum_{t=1}^T \nabla_i F(\mathbf{w}_t)^2 + \sigma_i^2 + \delta}}. \quad (7)$$

From Definition 3.1 and $b_{t-1,i} = \sum_{s=1}^{t-1} g_{s,i}^2$ in (AdaGrad), it can be shown that $\hat{\eta}_{t,i} \geq \tilde{\eta}_{T,i}$ for all $t \in [T]$. To address the second issue, we separate the step sizes from the gradients as follows:

$$\mathbb{E} \left[\sum_{t=1}^T \frac{\hat{\eta}_{t,i}}{2} \nabla_i F(\mathbf{w}_t)^2 \right] \geq \mathbb{E} \left[\frac{\tilde{\eta}_{T,i}}{2} \sum_{t=1}^T \nabla_i F(\mathbf{w}_t)^2 \right] \geq \frac{\mathbb{E} \left[\sqrt{\sum_{t=1}^T \nabla_i F(\mathbf{w}_t)^2} \right]^2}{\mathbb{E} \left[\frac{2}{\tilde{\eta}_{T,i}} \right]}, \quad (8)$$

where we used the elementary inequality that $\mathbb{E} \left[\frac{X^2}{Y} \right] \geq \frac{\mathbb{E}[X]^2}{\mathbb{E}[Y]}$ for any two random variables X and Y . Hence, in the following lemma, we will establish an upper bound on $\mathbb{E} \left[\frac{1}{\tilde{\eta}_{T,i}} \right]$, instead of directly lower bounding $\mathbb{E}[\tilde{\eta}_{T,i}]$. The proof is in Appendix A.4.

Lemma 3.4. *Consider the step size $\tilde{\eta}_{T,i}$ defined in (7). For any $i \in [d]$, we have*

$$\mathbb{E} \left[\frac{1}{\tilde{\eta}_{T,i}} \right] \leq \frac{\sigma_i \sqrt{2T} + \delta}{\eta} + \frac{\sqrt{3}}{\eta} \mathbb{E} \left[\sqrt{\sum_{t=1}^T \nabla_i F(\mathbf{w}_t)^2} \right].$$

Lemma 3.4 establishes an upper bound on $\mathbb{E} \left[\frac{1}{\tilde{\eta}_{T,i}} \right]$ in terms of the sum $\mathbb{E} \left[\sqrt{\sum_{t=1}^T \nabla_i F(\mathbf{w}_t)^2} \right]$, which also appears on the right hand side of (8). By combining Corollary 3.3, (8) and Lemma 3.4, we arrive at the following lemma. The proof is in Appendix A.5.

Lemma 3.5. *Consider the update in (AdaGrad) and recall that Q denotes the right-hand side in (6). It holds that*

$$\mathbb{E} \left[\sum_{i=1}^d \sqrt{\sum_{t=1}^T \nabla_i F(\mathbf{w}_t)^2} \right] \leq \frac{2\sqrt{3}}{\eta} Q + \sqrt{\frac{2d\delta Q}{\eta}} + 2\sqrt{\frac{\|\sigma\|_1 Q}{\eta}} T^{\frac{1}{4}}. \quad (9)$$

Having established Lemma 3.5, the last step is to relate the left-hand side of (9) to the ℓ_1 -norm of the gradients. Specifically, we can write:

$$\frac{1}{T} \sum_{t=1}^T \|\nabla F(\mathbf{w}_t)\|_1 = \frac{1}{T} \sum_{t=1}^T \sum_{i=1}^d |\nabla_i F(\mathbf{w}_t)| = \frac{1}{T} \sum_{i=1}^d \sum_{t=1}^T |\nabla_i F(\mathbf{w}_t)| \leq \frac{1}{\sqrt{T}} \sum_{i=1}^d \sqrt{\sum_{t=1}^T |\nabla_i F(\mathbf{w}_t)|^2},$$

where we switched the order of the two summations in the second equality and used the Cauchy-Schwarz inequality in the last inequality. This leads to our main theorem below.

Remark 3.1. *We choose to use the ℓ_1 -norm of the gradient as the measure of convergence since it is the tightest bound we can derive from the inequality in (9). Indeed, note that the ℓ_1 -norm is always an upper bound on the ℓ_2 -norm, and thus the above bound also immediately implies an upper bound on $\frac{1}{T} \sum_{t=1}^T \|\nabla F(\mathbf{w}_t)\|_2$. However, this relaxation will ruin the advantage of (AdaGrad) when compared to SGD or (AdaGrad-Norm).*

Theorem 3.6 (Main result). *Let $\{\mathbf{w}_t\}_{t=1}^T$ be the iterates generated by AdaGrad in (AdaGrad) and suppose that Assumptions 2.1, 2.2, and 2.3 hold. Recall that Q denotes the right-hand side in (6) and further assume that $\delta < \frac{1}{d}$. Then, we have*

$$\mathbb{E} \left[\frac{1}{T} \sum_{t=1}^T \|\nabla F(\mathbf{w}_t)\|_1 \right] \leq \frac{2\sqrt{3}Q}{\eta\sqrt{T}} + \sqrt{\frac{2Q}{\eta T}} + 2\sqrt{\frac{\|\boldsymbol{\sigma}\|_1 Q}{\eta}} \frac{1}{T^{\frac{1}{4}}}.$$

Recall that $Q = \tilde{\mathcal{O}}(F(\mathbf{w}_1) - F^* + \eta\|\boldsymbol{\sigma}\|_1 + \eta^2\|\mathbf{L}\|_1)$ if we ignore logarithmic terms. Thus, Theorem 3.6 implies that the averaged gradient ℓ_1 -norm can be bounded by

$$\tilde{\mathcal{O}} \left(\frac{F(\mathbf{w}_1) - F^*}{\eta\sqrt{T}} + \frac{\eta\|\mathbf{L}\|_1}{\sqrt{T}} + \frac{\sqrt{\|\boldsymbol{\sigma}\|_1(F(\mathbf{w}_1) - F^*)}}{\sqrt{\eta}T^{\frac{1}{4}}} + \frac{\|\boldsymbol{\sigma}\|_1}{T^{\frac{1}{4}}} + \frac{\sqrt{\eta\|\boldsymbol{\sigma}\|_1\|\mathbf{L}\|_1}}{T^{\frac{1}{4}}} \right) \quad (10)$$

An important feature of the upper bound in (10) is that, except inside the logarithmic term, it does not depend on the dimension d explicitly. Instead, the dependence is implicit via the noise variance vector $\boldsymbol{\sigma}$ and the Lipschitz constant vector \mathbf{L} introduced in Assumptions 2.2 and 2.3.

3.1 Discussion about the main result

As mentioned above the upper bound in (10) is independent of problem dimension, as such dependence is absorbed in $\|\boldsymbol{\sigma}\|_1$ and $\|\mathbf{L}\|_1$. To facilitate our discussions and the comparison with the existing results in Section 4, we will rewrite our bound in terms of the gradient's Lipschitz constants and the gradient noise variance, which are more standard in the existing works. Specifically, note that Assumption 2.3 implies that the function F is $\|\mathbf{L}\|_\infty$ -Lipschitz, i.e., $\|\nabla F(\mathbf{x}) - \nabla F(\mathbf{y})\|_2 \leq \|\mathbf{L}\|_\infty\|\mathbf{x} - \mathbf{y}\|_2$. Moreover, Assumption 2.2 implies that $\mathbb{E}[\|\mathbf{g}_t - \nabla F(\mathbf{w}_t)\|_2^2] \leq \sum_{i=1}^d \sigma_i^2 = \|\boldsymbol{\sigma}\|_2^2$.

Therefore, when we translate our bounds to the standard assumptions that are not tailored for coordinate-wise analysis, the ratios of $\|\mathbf{L}\|_1/\|\mathbf{L}\|_\infty$ as well as $\|\boldsymbol{\sigma}\|_1/\|\boldsymbol{\sigma}\|_2$ will appear in the upper bound. Given the behavior of these ratios, the dependence of our final bound on d could change as we describe in the following cases.

- **Worst case:** In this case, we have $\frac{\|\mathbf{L}\|_1}{\|\mathbf{L}\|_\infty} = \Theta(d)$ and $\frac{\|\boldsymbol{\sigma}\|_1}{\|\boldsymbol{\sigma}\|_2} = \Theta(\sqrt{d})$. To achieve the best dependence on d , we set $\eta = \frac{\eta'}{\sqrt{d}}$, where η' is a constant. The bound in (10) reduces to

$$\tilde{\mathcal{O}} \left(\frac{\sqrt{d}(F(\mathbf{w}_1) - F^*)}{\eta'\sqrt{T}} + \frac{\eta'\sqrt{d}L}{\sqrt{T}} + \frac{\sqrt{d\sigma(F(\mathbf{w}_1) - F^*)}}{\sqrt{\eta'}T^{\frac{1}{4}}} + \frac{\sqrt{d}\sigma}{T^{\frac{1}{4}}} + \frac{\sqrt{\eta'd\sigma L}}{T^{\frac{1}{4}}} \right). \quad (11)$$

Focusing on the dependence on the dimension d , we obtain the rate of $\tilde{\mathcal{O}}(\frac{\sqrt{d}}{\sqrt{T}} + \frac{\sqrt{d}}{T^{\frac{1}{4}}})$. Moreover, this bound is adaptive to the noise level σ : when the noise level is relatively small, i.e., $\sigma \ll \frac{1}{\sqrt{T}}$, then (AdaGrad) will automatically achieve a faster rate of $\tilde{\mathcal{O}}(\frac{\sqrt{d}}{\sqrt{T}})$. As we shall prove in Section 5, this rate is optimal and it matches our lower bound.

- **Well structured case:** In this case, we have $\frac{\|\mathbf{L}\|_1}{\|\mathbf{L}\|_\infty} = \mathcal{O}(1)$ and $\frac{\|\boldsymbol{\sigma}\|_1}{\|\boldsymbol{\sigma}\|_2} = \mathcal{O}(1)$. This means that the curvature and the noises of the gradient are heterogeneous and dominated by a few coordinates. Under such circumstances, we can set η as a constant and our convergence rate in (10) becomes a dimensional-independent rate of $\mathcal{O}(\frac{1}{\sqrt{T}} + \frac{1}{T^{\frac{1}{4}}})$.

Comparison with existing results on (AdaGrad). Most of the existing works use the ℓ_2 -norm as a measure of convergence [SCZJSL23; DBBU22; WZMC23; HL24]. The state-of-the-art result is [HL24]: with a fine-tuned step size, the authors show that, with high probability, (AdaGrad) satisfies $\frac{1}{T} \sum_{t=1}^T \|\nabla F(\mathbf{w}_t)\|_2^2 = \tilde{\mathcal{O}}\left(\frac{d}{T^{1/2}} + \frac{d^2}{T}\right)$. If we use this result to show a bound for the ℓ_1 norm, since $\|\nabla F(\mathbf{w}_t)\|_1 = \Theta(\sqrt{d}\|\nabla F(\mathbf{w}_t)\|_2)$ in the worst case, the upper bound becomes worse by a factor of \sqrt{d} and leads to

$$\min_{t=1,\dots,T} \|\nabla F(\mathbf{w}_t)\|_1 = \mathcal{O}\left(\frac{d}{T^{1/4}} + \frac{d^{1.5}}{\sqrt{T}}\right), \quad (12)$$

which is worse than the bound that we presented.

Also, in [LNNEN23], the authors considered the case that the function is L -smooth and the noise of gradient is coordinate-wise subgaussian, i.e., $\mathbb{E}[\exp(\lambda^2(g_{t,i} - \nabla_i F(\mathbf{w}_t))^2)] \leq \exp(\lambda^2 \sigma_i^2)$ for all λ such that $|\lambda| < \frac{1}{\sigma_i}$. Note that the subgaussian noise assumption is stronger than the bounded variance assumption in Assumption 2.2. Under these assumptions, they characterized the convergence rate of (AdaGrad) in terms of ℓ_1 -norm of the gradient as

$$\frac{1}{T} \sum_{t=1}^T \|\nabla F(\mathbf{w}_t)\|_1^2 \leq \tilde{\mathcal{O}}\left((\Delta_1 + \|\boldsymbol{\sigma}\|_3^3 + d\|\boldsymbol{\sigma}\|_1 + dL) \left(\frac{\|\boldsymbol{\sigma}\|_1}{\sqrt{T}} + \frac{\Delta_1 + Ld + \|\boldsymbol{\sigma}\|_2^2 + \|\boldsymbol{\sigma}\|_1}{T}\right)\right),$$

where $\Delta_1 = F(\mathbf{w}_1) - F^*$ is the initial function gap. Hence, in terms of the averaged *unsquared* gradient ℓ_1 -norm, their result is no better than $\tilde{\mathcal{O}}\left(\frac{\Delta_1}{\sqrt{T}} + \frac{dL}{\sqrt{T}} + \frac{\sqrt{\Delta_1\|\boldsymbol{\sigma}\|_1}}{T^{1/4}} + \frac{\sqrt{d}\|\boldsymbol{\sigma}\|_1}{T^{1/4}} + \frac{\sqrt{dL\|\boldsymbol{\sigma}\|_1}}{T^{1/4}}\right)$. Compared to our bounds in (10), we observe that their term $\frac{\sqrt{d}\|\boldsymbol{\sigma}\|_1}{T^{1/4}}$ is worse than ours by a factor of \sqrt{d} . Moreover, in the worst case where $\frac{\|\mathbf{L}\|_1}{\|\mathbf{L}\|_\infty} = \Theta(d)$ and $\frac{\|\boldsymbol{\sigma}\|_1}{\|\boldsymbol{\sigma}\|_2} = \Theta(\sqrt{d})$, their overall bound is worse than ours in (11) by a factor of \sqrt{d} .

4 Comparison with AdaGrad-Norm and SGD

In this section, we compare the rate obtained in Theorem 3.6 for (AdaGrad) with the existing rates of SGD and (AdaGrad-Norm). Note that the existing convergence results for these methods typically work under the assumption that the function F is L -smooth and the gradient noise variance is bounded by σ^2 . Under Assumptions 2.2 and 2.3, the function F is $\|\mathbf{L}\|_\infty$ -smooth and the gradient noise variance is bounded by $\mathbb{E}[\|\mathbf{g}_t - \nabla F(\mathbf{w}_t)\|_2^2] \leq \|\boldsymbol{\sigma}\|_2^2$. Therefore, we will set $L = \|\mathbf{L}\|_\infty$ and $\sigma = \|\boldsymbol{\sigma}\|_2$ to report their results.

With a constant step size η , SGD achieves the following rate [GL13; BCN18]:

$$\frac{1}{T} \sum_{t=1}^T \mathbb{E}[\|\nabla F(\mathbf{w}_t)\|_2^2] \leq \eta\|\mathbf{L}\|_\infty\|\boldsymbol{\sigma}\|_2^2 + \frac{2(F(\mathbf{w}_1) - F^*)}{\eta T} \quad (13)$$

for any $\eta \leq \frac{1}{\|\mathbf{L}\|_\infty}$. Specifically, if one chooses $\eta = \min\left\{\frac{1}{\|\mathbf{L}\|_\infty}, \frac{\sqrt{2((F(\mathbf{w}_1) - F^*))}}{\|\boldsymbol{\sigma}\|_2\sqrt{\|\mathbf{L}\|_\infty T}}\right\}$ to optimize the bound, the right-hand side of the above expression becomes

$$\min_{t=1,\dots,T} \mathbb{E}[\|\nabla F(\mathbf{w}_t)\|_2^2] = \mathcal{O}\left(\frac{\|\mathbf{L}\|_\infty(F(\mathbf{w}_0) - F^*)}{T} + \frac{\|\boldsymbol{\sigma}\|_2\sqrt{\|\mathbf{L}\|_\infty(F(\mathbf{w}_0) - F^*)}}{\sqrt{T}}\right) \quad (14)$$

In addition, note that (**AdaGrad-Norm**) achieves a similar rate as in (14) up to logarithmic factors [FTCMSW22]. Hence, in the following, we focus on the comparisons with the rate in (14). Note that the above rate for SGD and the one we presented in (10) for AdaGrad are not directly comparable, since one is established for the ℓ_2 -norm of the gradient and the other for the ℓ_1 -norm. Hence, we proceed to obtain a result for ℓ_1 norm using the bound in (14).

Inspired by the analysis in [BWAA18], we introduce the density function to do this conversion. Specifically, we define the two density functions $\phi, \tilde{\phi} : \mathbb{R}^d \rightarrow [0, 1]$ as follows:

$$\phi(\mathbf{v}) := \frac{\|\mathbf{v}\|_1^2}{d\|\mathbf{v}\|_2^2} \in \left[\frac{1}{d}, 1\right] \quad \text{and} \quad \tilde{\phi}(\mathbf{v}) := \frac{\|\mathbf{v}\|_1}{d\|\mathbf{v}\|_\infty} \in \left[\frac{1}{d}, 1\right]. \quad (15)$$

Specifically, a larger value of $\phi(\mathbf{v})$ or $\tilde{\phi}(\mathbf{v})$ indicates that the vector \mathbf{v} is denser. Moreover, we let $\phi(\nabla F)$ as a uniform upper bound such that $\phi(\nabla F(\mathbf{w}_t)) \leq \phi(\nabla F)$ for all $t \in [T]$. Under these notations, we can write $\|\boldsymbol{\sigma}\|_2 = \frac{\|\boldsymbol{\sigma}\|_1}{\sqrt{d\phi(\boldsymbol{\sigma})}}$ and $\|\mathbf{L}\|_\infty = \frac{\|\mathbf{L}\|_1}{d\tilde{\phi}(\mathbf{L})}$. Moreover, we can write $\|\nabla F(\mathbf{w}_t)\|_1 = \sqrt{d\phi(\nabla F(\mathbf{w}_t))}\|\nabla F(\mathbf{w}_t)\|_2 \leq \sqrt{d\phi(\nabla F)}\|\nabla F(\mathbf{w}_t)\|_2$. Using these expressions, the bound in (14) leads to

$$\min_{t=1, \dots, T} \mathbb{E} [\|\nabla F(\mathbf{w}_t)\|_1] = \mathcal{O} \left(R_1 \sqrt{\frac{\|\mathbf{L}\|_1(F(\mathbf{w}_1) - F^*)}{T}} + R_2 \frac{\|\boldsymbol{\sigma}\|_1^{\frac{1}{2}} \|\mathbf{L}\|_1^{\frac{1}{4}} (F(\mathbf{w}_1) - F^*)^{\frac{1}{4}}}{T^{\frac{1}{4}}} \right) \quad (16)$$

where

$$R_1 = \sqrt{\frac{\phi(\nabla F)}{\tilde{\phi}(\mathbf{L})}} \quad \text{and} \quad R_2 = \frac{\sqrt{\phi(\nabla F)}}{(\phi(\boldsymbol{\sigma})\tilde{\phi}(\mathbf{L}))^{\frac{1}{4}}}. \quad (17)$$

For the sake of argument, assume that we fine tune the step size parameter η in (**AdaGrad**) to achieve the best convergence bound. Then (10) will become

$$\tilde{\mathcal{O}} \left(\sqrt{\frac{\|\mathbf{L}\|_1(F(\mathbf{w}_1) - F^*)}{T}} + \frac{\|\boldsymbol{\sigma}\|_1^{\frac{1}{2}} \|\mathbf{L}\|_1^{\frac{1}{4}} (F(\mathbf{w}_1) - F^*)^{\frac{1}{4}}}{T^{\frac{1}{4}}} + \frac{\|\boldsymbol{\sigma}\|_1}{T^{\frac{1}{4}}} \right). \quad (18)$$

Except the last additional term $\frac{\|\boldsymbol{\sigma}\|_1}{T^{\frac{1}{4}}}$ in (18), we observe that the two bounds in (16) and (18) are similar. If we assume that the noise is relatively small, i.e., $\|\boldsymbol{\sigma}\|_1 \ll \sqrt{\|\mathbf{L}\|_1(F(\mathbf{w}_1) - F^*)}$, then the first two terms dominate. Then we can consider two extreme cases:

- At one extreme, if the gradients are dense and both the smoothness vector \mathbf{L} and the noise vector $\boldsymbol{\sigma}$ are sparse, then $R_1 = \Theta(\sqrt{d})$ and $R_2 = \Theta(\sqrt{d})$. Under this setting, the convergence rate of (**AdaGrad**) is better than that of SGD by a factor of \sqrt{d} .
- Conversely, if the gradients are sparse and both the smoothness vector \mathbf{L} and the noise vector $\boldsymbol{\sigma}$ are dense, then $R_1 = \Theta(\frac{1}{\sqrt{d}})$ and $R_2 = \Theta(\frac{1}{\sqrt{d}})$. Under this setting, the convergence rate of (**AdaGrad**) is worse than that of SGD by a factor of \sqrt{d} .

To the best of our knowledge, our results provide the first problem setting where (**AdaGrad**) provably achieves a better dimensional dependence than SGD in the non-convex setting. Interestingly, our discussions here mirror the comparison between AdaGrad and Online Gradient Descent in

[MS10; DHS11] regarding online convex optimization problems. Similarly, depending on the geometry of the feasible set and the density of the gradient vectors, it is shown that the rate of AdaGrad can be better or worse by a factor of \sqrt{d} . In this sense, our result complements this classical result and demonstrates that a similar phenomenon also occurs in the non-convex setting.

5 Lower bound result

In this section, we give a lower bound result for the query complexity of finding a stationary point for a non-convex but smooth function $p : \mathbb{R}^d \rightarrow \mathbb{R}$. We consider deterministic algorithms with access to the first-order oracle, i.e., the gradient $\nabla p(\mathbf{x})$, to obtain an ϵ -stationary point measured using ℓ_1 -norm given by $\|\nabla p(\mathbf{x})\|_1 \leq \epsilon$.

Theorem 5.1. *There exists a function $p : \mathbb{R}^d \rightarrow \mathbb{R}$ such that: (i) The gradient of p is 1-Lipschitz; (ii) $p(0) - \inf p \leq 1$; (iii) For any first-order methods, it takes at least $\Omega(d/\epsilon^2)$ queries to find a point $\hat{\mathbf{x}}$ with $\|\nabla p(\hat{\mathbf{x}})\|_1 \leq \epsilon$.*

Proof sketch: For constructing the function $p : \mathbb{R}^d \rightarrow \mathbb{R}$ in our lower bound proof, we use the result presented in Theorem 4 of [CBS23]. In this work, a function $f : \mathbb{R} \rightarrow \mathbb{R}$ of dimension $d = 1$ was proposed that is 1-smooth and satisfies $f(0) - \inf f \leq 1$. It was shown that any deterministic algorithm with access to a first-order oracle (i.e., one that provides $f'(x)$) requires at least $\frac{1}{32\epsilon^2}$ oracle calls to find a point where $|f'(x)| \leq \epsilon$ for any $\epsilon \in (0, 1)$. The details of the function f are given in Appendix B.

Given the above scalar function, we construct the function $p : \mathbb{R}^d \rightarrow \mathbb{R}$ corresponding to our lower bound in \mathbb{R}^d as

$$p(\mathbf{x}) := \frac{1}{d} \sum_{i=1}^d p_i(x_i) \quad \text{with} \quad p_i(x_i) = f(\sqrt{d}x_i), \quad (19)$$

where x_i is the i -th coordinate of \mathbf{x} . This function is coordinate-wise decomposable and the function associated with each coordinate is the one introduced in [CBS23]. It can be verified that each coordinate of the function p is 1-smooth, and it satisfies the condition $p(0) - \inf p \leq 1$.

Now consider any deterministic first-order method with access to the gradient $\nabla p(\mathbf{x})$, and suppose that it outputs a point $\hat{\mathbf{x}}$ after making fewer than $\frac{d}{32\epsilon^2}$ first-order oracle queries. Note that the constant 32 here is the same constant as in the lower bound of [CBS23, Theorem 4]. We claim that the condition $\|\nabla p(\hat{\mathbf{x}})\|_1 > \epsilon$ must hold. To prove this, let us focus on the first coordinate x_1 and the associated function $p_1(x_1)$. Since each query to the oracle $\nabla p(\mathbf{x})$ reveals exactly one value of p'_1 , the considered first-order method also makes fewer than $\frac{d}{32\tilde{\epsilon}^2} = \frac{1}{32\tilde{\epsilon}^2}$ queries to p'_1 , where $\tilde{\epsilon} = \frac{\epsilon}{\sqrt{d}}$. By definition, we have $p'_1(x_1) = \sqrt{d}f'(\sqrt{d}x_1)$. Moreover, using the property of f constructed in [CBS23], we obtain that $|f'(\sqrt{d}\hat{x}_1)| > \tilde{\epsilon} = \frac{\epsilon}{\sqrt{d}}$, which further implies $|p'_1(\hat{x}_1)| = \sqrt{d}|f'(\sqrt{d}\hat{x}_1)| > \epsilon$. With the same arguments, we can also show that $|p'_i(\hat{x}_i)| > \epsilon$ for all $i \in [d]$. Thus, we conclude that

$$\|\nabla p(\hat{\mathbf{x}})\|_1 = \frac{1}{d} \sum_{i=1}^d |p'_i(\hat{x}_i)| > \epsilon.$$

Therefore, any deterministic methods require at least $\frac{d}{32\epsilon^2}$ queries to the first-order oracle to find an ϵ -stationary point of the function p , in terms of ℓ_1 -norm. \square

Note that our upper bound in (10) can be simplified in the noiseless case to $\tilde{\mathcal{O}}\left(\frac{F(\mathbf{w}_1) - F^*}{\eta\sqrt{T}} + \frac{\eta\|\mathbf{L}\|_1}{\sqrt{T}}\right)$. In the worst case, it becomes $\tilde{\mathcal{O}}\left(\frac{F(\mathbf{w}_1) - F^*}{\eta\sqrt{T}} + \frac{\eta dL}{\sqrt{T}}\right)$. Setting $\eta = 1/\sqrt{d}$, it simplifies to $\tilde{\mathcal{O}}\left(\frac{\sqrt{d}}{\sqrt{T}}\right)$, assuming $(F(\mathbf{w}_1) - F^*) = \mathcal{O}(1)$ and $L = \mathcal{O}(1)$. This result is equivalent to a complexity of $\tilde{\mathcal{O}}\left(\frac{d}{\epsilon^2}\right)$ and matches the lower bound established in Theorem 5.1 up to logarithmic terms. Hence, in the noiseless case, our upper bound is tight up to a logarithmic factor.

6 Conclusion and limitations

In this work, we presented the convergence analysis of adaptive gradient methods, specifically AdaGrad, measured by the ℓ_1 -norm of the gradient. Under the general assumptions of an L -Lipschitz gradient and bounded stochastic gradient variance, we proved a worst-case convergence rate of $\mathcal{O}\left(\frac{\sqrt{d}L}{\sqrt{T}} + \frac{\sqrt{d}\sigma}{T^{1/4}}\right)$, where d is the dimension. Under more refined assumptions on the structure of the gradient and noise, we analyzed different ℓ_1/ℓ_2 geometries where AdaGrad outperforms SGD and AdaGrad-Norm in terms of dimensional dependence. Moreover, we provided a lower bound result of $\Omega(\sqrt{d/T})$ for deterministic algorithms with access to first-order information when minimizing the ℓ_1 norm of the gradient for a smooth non-convex function.

Limitations: We focused on the setting where the Lipschitz constant of the gradient is fixed. Extending our theory to scenarios where the Lipschitz constant grows with the gradient norm is a potential direction for future work. Additionally, we considered the classic setting with bounded stochastic gradient variance. A more general analysis could extend our theory to the affine variance assumption explored in the literature.

References

- [ACDFSW23] Y. Arjevani, Y. Carmon, J. C. Duchi, D. J. Foster, N. Srebro, and B. Woodworth. “Lower bounds for non-convex stochastic optimization”. *Mathematical Programming* 199.1 (2023), pp. 165–214 (page 2).
- [BWAA18] J. Bernstein, Y.-X. Wang, K. Azizzadenesheli, and A. Anandkumar. *signSGD: Compressed Optimisation for Non-Convex Problems*. 2018. arXiv: 1802.04434 [cs.LG] (pages 4, 10).
- [BCN18] L. Bottou, F. E. Curtis, and J. Nocedal. “Optimization methods for large-scale machine learning”. *SIAM review* 60.2 (2018), pp. 223–311 (pages 2, 9).
- [CLSH19] X. Chen, S. Liu, R. Sun, and M. Hong. “On the convergence of a class of Adam-type algorithms for non-convex optimization”. In: *7th International Conference on Learning Representations, ICLR 2019*. 2019 (page 3).
- [CBS23] S. Chewi, S. Bubeck, and A. Salim. *On the complexity of finding stationary points of smooth functions in one dimension*. 2023. arXiv: 2209.07513 [math.OC] (pages 11, 21).
- [DBBU22] A. Défossez, L. Bottou, F. Bach, and N. Usunier. “A Simple Convergence Proof of Adam and Adagrad”. *Transactions on Machine Learning Research* (2022) (pages 3, 9).
- [DHS11] J. Duchi, E. Hazan, and Y. Singer. “Adaptive Subgradient Methods for Online Learning and Stochastic Optimization”. *Journal of Machine Learning Research* 12.61 (2011), pp. 2121–2159. URL: <http://jmlr.org/papers/v12/duchi11a.html> (pages 2, 11).
- [FRCS23] M. Faw, L. Rout, C. Caramanis, and S. Shakkottai. “Beyond uniform smoothness: A stopped analysis of adaptive sgd”. In: *The Thirty Sixth Annual Conference on Learning Theory*. PMLR. 2023, pp. 89–160 (page 2).
- [FTCMSW22] M. Faw, I. Tziotis, C. Caramanis, A. Mokhtari, S. Shakkottai, and R. Ward. *The Power of Adaptivity in SGD: Self-Tuning Step Sizes with Unbounded Gradients and Affine Variance*. 2022. arXiv: 2202.05791 [stat.ML] (pages 2, 5, 6, 10, 17).
- [GG22] S. Gadat and I. Gavra. “Asymptotic study of stochastic adaptive algorithms in non-convex landscape”. *Journal of Machine Learning Research* 23.228 (2022), pp. 1–54 (page 2).
- [GL13] S. Ghadimi and G. Lan. “Stochastic first-and zeroth-order methods for nonconvex stochastic programming”. *SIAM journal on optimization* 23.4 (2013), pp. 2341–2368 (pages 2, 9).
- [HL24] Y. Hong and J. Lin. “Revisiting Convergence of AdaGrad with Relaxed Assumptions”. *arXiv preprint arXiv:2402.13794* (2024) (page 9).
- [KLC21] A. Kavis, K. Y. Levy, and V. Cevher. “High Probability Bounds for a Class of Nonconvex Algorithms with AdaGrad Stepsize”. In: *International Conference on Learning Representations*. 2021 (page 2).
- [LL24] H. Li and Z. Lin. “On the $O(\frac{\sqrt{d}}{T^{1/4}})$ Convergence Rate of RMSProp and Its Momentum Extension Measured by ℓ_1 Norm: Better Dependence on the Dimension”. *arXiv preprint arXiv:2402.00389* (2024) (page 3).

- [LO20] X. Li and F. Orabona. “A High Probability Analysis of Adaptive SGD with Momentum”. In: *Workshop on Beyond First Order Methods in ML Systems at ICML’20*. 2020 (page 2).
- [LO19] X. Li and F. Orabona. “On the convergence of stochastic gradient descent with adaptive stepsizes”. In: *The 22nd international conference on artificial intelligence and statistics*. PMLR. 2019, pp. 983–992 (page 2).
- [LNNEN23] Z. Liu, T. D. Nguyen, T. H. Nguyen, A. Ene, and H. L. Nguyen. *High Probability Convergence of Stochastic Gradient Methods*. 2023. arXiv: [2302.14843](#) [math.OC] (pages 3, 9).
- [MS10] H. B. McMahan and M. Streeter. “Adaptive Bound Optimization for Online Convex Optimization”. *COLT 2010* (2010), p. 244 (pages 2, 11).
- [RT11] P. Richtárik and M. Takáč. *Iteration Complexity of Randomized Block-Coordinate Descent Methods for Minimizing a Composite Function*. 2011. arXiv: [1107.2848](#) [math.OC] (page 4).
- [RM51] H. Robbins and S. Monro. “A stochastic approximation method”. *The annals of mathematical statistics* (1951), pp. 400–407 (page 2).
- [SCZJSL23] L. Shen, C. Chen, F. Zou, Z. Jie, J. Sun, and W. Liu. “A unified analysis of AdaGrad with weighted aggregation and momentum acceleration”. *IEEE Transactions on Neural Networks and Learning Systems* (2023) (page 9).
- [WZMC23] B. Wang, H. Zhang, Z. Ma, and W. Chen. “Convergence of adagrad for non-convex objectives: Simple proofs and relaxed assumptions”. In: *The Thirty Sixth Annual Conference on Learning Theory*. PMLR. 2023, pp. 161–190 (page 9).
- [WWB20] R. Ward, X. Wu, and L. Bottou. “Adagrad stepsizes: Sharp convergence over non-convex landscapes”. *Journal of Machine Learning Research* 21.219 (2020), pp. 1–30 (pages 2, 5).
- [ZCCTYG18] D. Zhou, J. Chen, Y. Cao, Y. Tang, Z. Yang, and Q. Gu. “On the convergence of adaptive gradient methods for nonconvex optimization”. *arXiv preprint arXiv:1808.05671* (2018) (page 3).

Appendix

A Results from Section 3

A.1 Proof of Lemma 3.1

We begin with the L-smoothness Assumption 2.3 and the definition of the Algorithm,

$$\begin{aligned} F(\mathbf{w}_{t+1}) - F(\mathbf{w}_t) &\leq \langle \nabla F(\mathbf{w}_t), \mathbf{w}_{t+1} - \mathbf{w}_t \rangle + \sum_{i=1}^d \frac{L_i}{2} |w_{t+1,i} - w_{t,i}|^2 \\ &= \sum_{i=1}^d \nabla_i F(\mathbf{w}_t) (w_{t+1,i} - w_{t,i}) + \sum_{i=1}^d \frac{L_i}{2} |w_{t+1,i} - w_{t,i}|^2 \end{aligned}$$

Since $w_{t+1,i} = w_{t,i} - \eta_{t,i} g_{t,i}$, we further have

$$F(\mathbf{w}_{t+1}) - F(\mathbf{w}_t) = \sum_{i=1}^d -\eta_{t,i} \nabla_i F(\mathbf{w}_t) g_{t,i} + \frac{L_i}{2} \eta_{t,i}^2 g_{t,i}^2.$$

Taking the expectation with respect to \mathcal{F}_{t-1} , we obtain:

$$\mathbb{E}[F(\mathbf{w}_{t+1}) | \mathcal{F}_{t-1}] - F(\mathbf{w}_t) = \sum_{i=1}^d -\mathbb{E}[\eta_{t,i} \nabla_i F(\mathbf{w}_t) g_{t,i} | \mathcal{F}_{t-1}] + \frac{L_i}{2} \mathbb{E}[\eta_{t,i}^2 g_{t,i}^2 | \mathcal{F}_{t-1}].$$

Since $\mathbb{E}[-\hat{\eta}_{t,i} \nabla_i F(\mathbf{w}_t) (\nabla_i F(\mathbf{w}_t) - g_{t,i}) | \mathcal{F}_{t-1}] = 0$, we further have

$$\begin{aligned} &\mathbb{E}[F(\mathbf{w}_{t+1}) | \mathcal{F}_{t-1}] - F(\mathbf{w}_t) \\ &\leq \sum_{i=1}^d \left[-\hat{\eta}_{t,i} \nabla_i F(\mathbf{w}_t)^2 - \mathbb{E}[(\eta_{t,i} - \hat{\eta}_{t,i}) \nabla_i F(\mathbf{w}_t) g_{t,i} | \mathcal{F}_{t-1}] + \frac{L_i}{2} \mathbb{E}[\eta_{t,i}^2 g_{t,i}^2 | \mathcal{F}_{t-1}] \right] \end{aligned}$$

Hence we have the stated result.

A.2 Proof of Lemma 3.2

We consider (AdaGrad). Recall from the definition that

$$\eta_{t,i} = \frac{\eta}{\sqrt{b_{t-1,i}^2 + g_{t,i}^2 + \delta}} \quad \text{and} \quad \hat{\eta}_{t,i} = \frac{\eta}{\sqrt{b_{t-1,i}^2 + \nabla_i F(\mathbf{w}_t)^2 + \sigma_i^2 + \delta}}.$$

Let $a = b_{t-1,i}^2 + g_{t,i}^2$ and $b = b_{t-1,i}^2 + \nabla_i F(\mathbf{w}_t)^2 + \sigma_i^2$. Then

$$\begin{aligned} |\eta_{t,i} - \hat{\eta}_{t,i}| &= \eta \left| \frac{1}{\sqrt{a} + \delta} - \frac{1}{\sqrt{b} + \delta} \right| = \eta \left| \frac{b - a}{(\sqrt{a} + \delta)(\sqrt{b} + \delta)(\sqrt{a} + \sqrt{b} + 2\delta)} \right| \\ &= \eta \left| \frac{\nabla_i F(\mathbf{w}_t)^2 + \sigma_i^2 - g_{t,i}^2}{(\sqrt{a} + \delta)(\sqrt{b} + \delta)(\sqrt{a} + \sqrt{b} + 2\delta)} \right| \\ &\leq \frac{\eta |\nabla_i F(\mathbf{w}_t)^2 - g_{t,i}^2|}{(\sqrt{a} + \delta)(\sqrt{b} + \delta)(\sqrt{a} + \sqrt{b})} + \frac{\eta \sigma_i^2}{(\sqrt{a} + \delta)(\sqrt{b} + \delta)(\sqrt{a} + \sqrt{b})}. \end{aligned}$$

Since $\sqrt{a} \geq |g_{t,i}|$, $\sqrt{b} \geq \max\{|\nabla_i F(\mathbf{w}_t)|, \sigma_i\}$, and $|\nabla_i F(\mathbf{w}_t)^2 - g_{t,i}^2| \leq |\nabla_i F(\mathbf{w}_t) - g_{t,i}|(|\nabla_i F(\mathbf{w}_t)| + |g_{t,i}|)$, we further have

$$|\eta_{t,i} - \hat{\eta}_{t,i}| \leq \frac{\eta|\nabla_i F(\mathbf{w}_t) - g_{t,i}| + \eta\sigma_i}{(\sqrt{a} + \delta)(\sqrt{b} + \delta)} = \frac{1}{\eta} (|\nabla_i F(\mathbf{w}_t) - g_{t,i}| + \sigma_i) \eta_{t,i} \tilde{\eta}_{t,i}.$$

Hence we have,

$$\begin{aligned} |\eta_{t,i} - \hat{\eta}_{t,i}| |\nabla_i F(\mathbf{w}_t)| |g_{t,i}| &\leq \frac{1}{\eta} \eta_{t,i} \tilde{\eta}_{t,i} (|\nabla_i F(\mathbf{w}_t) - g_{t,i}| + \sigma_i) |\nabla_i F(\mathbf{w}_t)| |g_{t,i}| \\ &= \frac{1}{\eta} \eta_{t,i} \tilde{\eta}_{t,i} |\nabla_i F(\mathbf{w}_t) - g_{t,i}| |\nabla_i F(\mathbf{w}_t)| |g_{t,i}| + \frac{1}{\eta} \sigma_i \eta_{t,i} \tilde{\eta}_{t,i} |\nabla_i F(\mathbf{w}_t)| |g_{t,i}| \end{aligned}$$

Using the Cauchy-Schwartz inequality, we further have

$$\begin{aligned} &\mathbb{E} [\eta_{t,i} \tilde{\eta}_{t,i} |\nabla_i F(\mathbf{w}_t) - g_{t,i}| |\nabla_i F(\mathbf{w}_t)| |g_{t,i}| \mid \mathcal{F}_{t-1}] \\ &\leq \tilde{\eta}_{t,i} |\nabla_i F(\mathbf{w}_t)| \sqrt{\mathbb{E} [|\nabla_i F(\mathbf{w}_t) - g_{t,i}|^2 \mid \mathcal{F}_{t-1}] \mathbb{E} [\eta_{t,i}^2 g_{t,i}^2 \mid \mathcal{F}_{t-1}]} \\ &\leq \sigma_i \tilde{\eta}_{t,i} |\nabla_i F(\mathbf{w}_t)| \sqrt{\mathbb{E} [\eta_{t,i}^2 g_{t,i}^2 \mid \mathcal{F}_{t-1}]} \end{aligned}$$

where the last step follows from the bounded-variance in Assumption 2.2. We proceed to bound the second term in a similar manner,

$$\mathbb{E} [\sigma_i \eta_{t,i} \tilde{\eta}_{t,i} |\nabla_i F(\mathbf{w}_t)| |g_{t,i}| \mid \mathcal{F}_{t-1}] \leq \sigma_i \tilde{\eta}_{t,i} |\nabla_i F(\mathbf{w}_t)| \sqrt{\mathbb{E} [\eta_{t,i}^2 g_{t,i}^2 \mid \mathcal{F}_{t-1}]}$$

Combining the results, the term $\mathbb{E} [(\eta_{t,i} - \hat{\eta}_{t,i}) \langle \nabla_i F(\mathbf{w}_t), g_{t,i} \rangle \mid \mathcal{F}_{t-1}]$ is bounded as follows,

$$\begin{aligned} \mathbb{E} [|\eta_{t,i} - \hat{\eta}_{t,i}| |\nabla_i F(\mathbf{w}_t)| |g_{t,i}| \mid \mathcal{F}_{t-1}] &\leq \frac{2\sigma_i \tilde{\eta}_{t,i} |\nabla_i F(\mathbf{w}_t)|}{\eta} \sqrt{\mathbb{E} [\eta_{t,i}^2 g_{t,i}^2 \mid \mathcal{F}_{t-1}]} \\ &\leq \frac{1}{2} \tilde{\eta}_{t,i} \|\nabla_i F(\mathbf{w}_t)\|^2 + \frac{2\tilde{\eta}_{t,i} \sigma_i^2}{\eta^2} \mathbb{E} [\eta_{t,i}^2 g_{t,i}^2 \mid \mathcal{F}_{t-1}] \end{aligned} \quad (20)$$

where we used Young's inequality in (20) in the last inequality. Finally, note that we have $\tilde{\eta}_{t,i} \leq \frac{\eta}{\sigma_i}$.

It remains to prove the bound in 5 for Lemma 3.2. We first prove some intermediate results.

Lemma A.1. *Suppose that Assumption 2.3 holds. Then for any two iteration indices $t_2 \geq t_1$,*

$$\left| \|\nabla F(\mathbf{w}_{t_2})\| - \|\nabla F(\mathbf{w}_{t_1})\| \right| \leq \eta \sqrt{d} L (t_2 - t_1),$$

In particular, this implies that

$$\|\nabla F(\mathbf{w}_t)\| \leq \|\nabla F(\mathbf{w}_1)\| + \eta \sqrt{d} L t,$$

and it further follows that for all i ,

$$\nabla_i F(\mathbf{w}_t)^2 \leq \|\nabla F(\mathbf{w}_t)\|^2 \leq \left(\|\nabla F(\mathbf{w}_1)\| + \eta \sqrt{d} L t \right)^2$$

Proof. Using the triangle inequality we have,

$$\left| \|\nabla F(\mathbf{w}_{t_2})\| - \|\nabla F(\mathbf{w}_{t_1})\| \right| \leq \|\nabla F(\mathbf{w}_{t_2}) - \nabla F(\mathbf{w}_{t_1})\| = \left\| \sum_{s=t_1}^{t_2-1} \nabla F(\mathbf{w}_{s+1}) - \nabla F(\mathbf{w}_s) \right\|.$$

Given the coordinate-wise smoothness assumption, the general smoothness assumption also holds by considering the Lipschitz constant as $L := \max_i L_i = \|\mathbf{L}\|_\infty$. This can be used to bound the term above,

$$\begin{aligned} \left\| \sum_{s=t_1}^{t_2-1} \nabla F(\mathbf{w}_{s+1}) - \nabla F(\mathbf{w}_s) \right\| &\leq \sum_{s=t_1}^{t_2-1} \|\nabla F(\mathbf{w}_{s+1}) - \nabla F(\mathbf{w}_s)\| \\ &\leq \sum_{s=t_1}^{t_2-1} L \|\mathbf{w}_{s+1} - \mathbf{w}_s\| = \sum_{s=t_1}^{t_2-1} L \sqrt{\sum_{i=1}^d |w_{s+1,i} - w_{s,i}|^2} \end{aligned}$$

Using the update rule $w_{t+1,i} = w_{t,i} - \eta_{t,i} g_{t,i}$,

$$\left| \|\nabla F(\mathbf{w}_{t_2})\| - \|\nabla F(\mathbf{w}_{t_1})\| \right| \leq \sum_{s=t_1}^{t_2-1} L \sqrt{\sum_{i=1}^d \frac{\eta^2 g_{t,i}^2}{b_{t-1,i}^2 + g_{t,i}^2}} \leq \sum_{s=t_1}^{t_2-1} \eta \sqrt{d} L = \eta \sqrt{d} L (t_2 - t_1)$$

Hence we have the stated result. \square

Lemma A.2. Let $\{a_s\}_{s=1}^\infty$ be any sequence such that $a_s \geq 0$ for all s . Moreover, define $A_t = A_{t-1} + a_t$, where $A_0 = 0$. Then we have

$$\sum_{t=1}^T \frac{a_t}{A_t + \delta^2} \leq \log \left(1 + \frac{A_T}{\delta^2} \right) \quad (21)$$

Proof. The proof is similar to [FTCMSW22, Lemma 15] and we repeat here for completeness. Note that for any $t \geq 1$, we have

$$\frac{a_t}{A_t + \delta^2} = 1 - \frac{A_{t-1} + \delta^2}{A_t + \delta^2} \leq \log \left(\frac{A_t + \delta^2}{A_{t-1} + \delta^2} \right).$$

The last step follows from $x \leq -\log(1-x)$. Summing the above inequalities from $t = 1$ to $t = T$, we obtain that

$$\sum_{t=1}^T \frac{a_t}{A_t + \delta^2} \leq \log \left(\frac{A_T + \delta^2}{A_0 + \delta^2} \right) = \log \left(1 + \frac{A_T}{\delta^2} \right).$$

\square

Now we are ready to prove (5) in Lemma 3.2. From the definition of the step size for (AdaGrad) we have the following equation,

$$\mathbb{E} \left[\sum_{t=1}^T \eta_{t,i}^2 g_{t,i}^2 \right] \leq \eta^2 \mathbb{E} \left[\sum_{t=1}^T \frac{g_{t,i}^2}{b_{t,i}^2 + \delta^2} \right]$$

Using Lemma A.2, we can bound the summation with a log term as follows,

$$\eta^2 \mathbb{E} \left[\sum_{t=1}^T \frac{g_{t,i}^2}{b_{t-1,i}^2 + g_{t,i}^2 + \delta^2} \right] \leq \eta^2 \mathbb{E} \left[\log \left(1 + \frac{b_{T,i}^2}{\delta^2} \right) \right] \leq \eta^2 \log \left(1 + \frac{\mathbb{E} [b_{T,i}^2]}{\delta^2} \right),$$

where we apply Jensen's Inequality to the concave log function in the last inequality. Moreover, since $b_{T,i}^2 = \sum_{t=1}^T g_{t,i}^2$, by using Assumptions 2.1 and 2.2 we have

$$\mathbb{E} [b_{T,i}^2] = \sum_{t=1}^T \mathbb{E} [g_{t,i}^2] \leq \sum_{t=1}^T (\sigma_i^2 + \nabla_i F(\mathbf{w}_t)^2) \leq T\sigma_i^2 + \sum_{t=1}^T \nabla_i F(\mathbf{w}_t)^2.$$

Using the result from Lemma A.1, for any $t \in [T]$, we further have

$$\nabla_i F(\mathbf{w}_t)^2 \leq \left(\|\nabla F(\mathbf{w}_1)\| + \eta\sqrt{d}Lt \right)^2 \leq \left(\|\nabla F(\mathbf{w}_1)\| + \eta\sqrt{d}LT \right)^2$$

Combining all the inequalities above, we obtain that

$$\eta^2 \mathbb{E} \left[\sum_{t=1}^T \frac{g_{t,i}^2}{b_{t-1,i}^2 + g_{t,i}^2 + \delta^2} \right] \leq \eta^2 \log \left(1 + \frac{T\sigma_i^2}{\delta^2} + \frac{T(\|\nabla F(\mathbf{w}_1)\| + \eta\sqrt{d}LT)^2}{\delta^2} \right)$$

Hence we have proved the bound in (5) of Lemma 3.2. This completes the proof of the results in Lemma 3.2.

A.3 Proof of Corollary 3.3

We begin with the result from Lemma 3.1, rearranging we have,

$$\begin{aligned} \mathbb{E} [F(\mathbf{w}_{t+1}) | \mathcal{F}_{t-1}] - F(\mathbf{w}_t) &\leq \sum_{i=1}^d \left[-\hat{\eta}_{t,i} \nabla_i F(\mathbf{w}_t)^2 - \mathbb{E} [(\eta_{t,i} - \hat{\eta}_{t,i}) \langle \nabla_i F(\mathbf{w}_t), g_{t,i} \rangle | \mathcal{F}_{t-1}] \right. \\ &\quad \left. + \frac{L_i}{2} \mathbb{E} [\eta_{t,i}^2 g_{t,i}^2 | \mathcal{F}_{t-1}] \right]. \end{aligned}$$

From the result in Lemma 3.2 we can bound the bias term,

$$\begin{aligned} \mathbb{E} [F(\mathbf{w}_{t+1}) - F(\mathbf{w}_t) | \mathcal{F}_{t-1}] &\leq \sum_{i=1}^d -\hat{\eta}_{t,i} \nabla_i F(\mathbf{w}_t)^2 + \frac{\hat{\eta}_{t,i}}{2} \nabla_i F(\mathbf{w}_t)^2 + 2\eta\sigma_i \mathbb{E} \left[\frac{g_{t,i}^2}{b_{t-1,i}^2 + g_{t,i}^2} | \mathcal{F}_{t-1} \right] \\ &\quad + \frac{L_i}{2} \mathbb{E} [\eta_{t,i}^2 g_{t,i}^2 | \mathcal{F}_{t-1}] \\ &\leq \sum_{i=1}^d -\frac{\hat{\eta}_{t,i}}{2} \nabla_i F(\mathbf{w}_t)^2 + \left(2\eta\sigma_i + \frac{L_i\eta^2}{2} \right) \mathbb{E} \left[\frac{g_{t,i}^2}{b_{t-1,i}^2 + g_{t,i}^2} | \mathcal{F}_{t-1} \right]. \end{aligned}$$

Rearranging the terms and taking the sum over the iterations we have,

$$\sum_{t=1}^T \sum_{i=1}^d \frac{\hat{\eta}_{t,i}}{2} \nabla_i F(\mathbf{w}_t)^2 \leq F(\mathbf{w}_1) - \mathbb{E} [F(\mathbf{w}_T) | \mathcal{F}_{T-1}] + \sum_{t=1}^T \sum_{i=1}^d \left(2\eta\sigma_i + \frac{L_i\eta^2}{2} \right) \mathbb{E} \left[\frac{g_{t,i}^2}{b_{t-1,i}^2 + g_{t,i}^2} | \mathcal{F}_{t-1} \right]$$

We can now take the expectation and use (5) from Lemma 3.2:

$$\begin{aligned}\mathbb{E} \left[\sum_{t=1}^T \sum_{i=1}^d \frac{\hat{\eta}_{t,i}}{2} \nabla_i F(\mathbf{w}_t)^2 \right] &\leq F(\mathbf{w}_1) - \mathbb{E} [F(\mathbf{w}_T)] + \sum_{i=1}^d \left(2\eta\sigma_i + \frac{L_i\eta^2}{2} \right) \mathbb{E} \left[\sum_{t=1}^T \frac{g_{t,i}^2}{b_{t-1,i}^2 + g_{t,i}^2} \right] \\ \mathbb{E} \left[\sum_{t=1}^T \sum_{i=1}^d \frac{\hat{\eta}_{t,i}}{2} \nabla_i F(\mathbf{w}_t)^2 \right] &\leq F(\mathbf{w}_1) - F^* + \sum_{i=1}^d \left(2\eta\sigma_i + \frac{L_i\eta^2}{2} \right) \log h_i(T)\end{aligned}$$

Using $h(T) = 1 + \frac{T\|\boldsymbol{\sigma}\|_\infty^2}{\delta^2} + \frac{T(\|\nabla F(\mathbf{w}_1)\| + \eta\sqrt{d}\|\mathbf{L}\|_\infty T)^2}{\delta^2}$ we have,

$$\begin{aligned}\mathbb{E} \left[\sum_{t=1}^T \sum_{i=1}^d \frac{\hat{\eta}_{t,i}}{2} \nabla_i F(\mathbf{w}_t)^2 \right] &\leq F(\mathbf{w}_1) - F^* + \left(\sum_{i=1}^d 2\eta\sigma_i + \sum_{i=1}^d \frac{L_i\eta^2}{2} \right) \log h(T) \\ &\leq F(\mathbf{w}_1) - F^* + \left(2\eta\|\boldsymbol{\sigma}\|_1 + \frac{\eta^2\|\mathbf{L}\|_1}{2} \right) \log h(T)\end{aligned}$$

Hence we have proved the lemma.

A.4 Proof of Lemma 3.4

From the definition of $\tilde{\eta}_{T,i}$ and using $b_{t-1,i}^2 = \sum_{s=1}^{t-1} g_{s,i}^2 \leq \sum_{t=1}^{T-1} g_{t,i}^2$,

$$\begin{aligned}\mathbb{E} \left[\frac{\eta}{\tilde{\eta}_{T,i}} \right] &= \mathbb{E} \left[\sqrt{b_{t-1,i}^2 + \sigma_i^2 + \nabla_i F(\mathbf{w}_t)^2} + \delta \right] \\ &\leq \mathbb{E} \left[\sqrt{\sum_{t=1}^T g_{t,i}^2 + \sigma_i^2 + \sum_{t=1}^T \nabla_i F(\mathbf{w}_t)^2} + \delta \right]\end{aligned}$$

We can use the upper bound of $g_{t,i}^2 \leq 2((g_{t,i} - \nabla_i F(\mathbf{w}_t))^2 + \nabla_i F(\mathbf{w}_t)^2)$,

$$\begin{aligned}\mathbb{E} \left[\frac{\eta}{\tilde{\eta}_{T,i}} \right] &\leq \mathbb{E} \left[\sqrt{\sum_{t=1}^T 2((g_{t,i} - \nabla_i F(\mathbf{w}_t))^2 + \nabla_i F(\mathbf{w}_t)^2) + \sigma_i^2 + \sum_{t=1}^T \nabla_i F(\mathbf{w}_t)^2} + \delta \right] \\ &\leq \mathbb{E} \left[\sqrt{2 \sum_{t=1}^{T-1} (g_{t,i} - \nabla_i F(\mathbf{w}_t))^2 + 3 \sum_{t=1}^T \nabla_i F(\mathbf{w}_t)^2 + \sigma_i^2} + \delta \right] \\ &\leq \mathbb{E} \left[\sqrt{2 \sum_{t=1}^{T-1} (g_{t,i} - \nabla_i F(\mathbf{w}_t))^2 + \sigma_i^2} \right] + \mathbb{E} \left[\sqrt{3 \sum_{t=1}^T \nabla_i F(\mathbf{w}_t)^2} \right] + \delta\end{aligned}$$

Applying Jensen's inequality and bounded variance from Assumption 2.2,

$$\begin{aligned}\mathbb{E} \left[\frac{\eta}{\tilde{\eta}_{T,i}} \right] &\leq \sqrt{2 \sum_{t=1}^{T-1} \mathbb{E} [(g_{t,i} - \nabla_i F(\mathbf{w}_t))^2] + \sigma_i^2} + \mathbb{E} \left[\sqrt{3 \sum_{t=1}^T \nabla_i F(\mathbf{w}_t)^2} \right] + \delta \\ &\leq \sqrt{2T\sigma_i^2} + \sqrt{3}\mathbb{E} \left[\sqrt{\sum_{t=1}^T \nabla_i F(\mathbf{w}_t)^2} \right] + \delta\end{aligned}$$

Rearranging the terms we have,

$$\mathbb{E} \left[\frac{1}{\tilde{\eta}_{T,i}} \right] \leq \frac{\sigma_i \sqrt{2T} + \delta}{\eta} + \frac{\sqrt{3}}{\eta} \mathbb{E} \left[\sqrt{\sum_{t=1}^T \nabla_i F(\mathbf{w}_t)^2} \right]$$

Hence we have proved the stated lemma.

A.5 Proof of Lemma 3.5

From equation 8 we have,

$$\mathbb{E} \left[\sqrt{\sum_{t=1}^T \nabla_i F(\mathbf{w}_t)^2} \right]^2 \leq \mathbb{E} \left[\sum_{t=1}^T \frac{\hat{\eta}_{t,i}}{2} \nabla_i F(\mathbf{w}_t)^2 \right] \mathbb{E} \left[\frac{2}{\tilde{\eta}_{T,i}} \right]$$

Using the result from Lemma 3.4 we get a quadratic inequality as follows,

$$\begin{aligned} \mathbb{E} \left[\sqrt{\sum_{t=1}^T \nabla_i F(\mathbf{w}_t)^2} \right] &\leq \sqrt{\mathbb{E} \left[\sum_{t=1}^T \frac{\hat{\eta}_{t,i}}{2} \nabla_i F(\mathbf{w}_t)^2 \right]} \sqrt{\mathbb{E} \left[\frac{2}{\tilde{\eta}_{T,i}} \right]} \\ &\leq \sqrt{\frac{2}{\eta}} \sqrt{\mathbb{E} \left[\sum_{t=1}^T \frac{\hat{\eta}_{t,i}}{2} \nabla_i F(\mathbf{w}_t)^2 \right]} \sqrt{(\sigma_i \sqrt{2T} + \delta) + \sqrt{3} \mathbb{E} \left[\sqrt{\sum_{t=1}^T \nabla_i F(\mathbf{w}_t)^2} \right]} \end{aligned}$$

Solving the quadratic we have the following bound,

$$\mathbb{E} \left[\sqrt{\sum_{t=1}^T \nabla_i F(\mathbf{w}_t)^2} \right] \leq \frac{2\sqrt{3}}{\eta} \mathbb{E} \left[\sum_{t=1}^T \frac{\hat{\eta}_{t,i}}{2} \nabla_i F(\mathbf{w}_t)^2 \right] + \sqrt{\frac{2}{\eta}} \sqrt{(\sigma_i \sqrt{2T} + \delta)} \sqrt{\mathbb{E} \left[\sum_{t=1}^T \frac{\hat{\eta}_{t,i}}{2} \nabla_i F(\mathbf{w}_t)^2 \right]}$$

Combining the bounds from all the coordinates and using Cauchy-Schwartz for the second term,

$$\begin{aligned} \mathbb{E} \left[\sum_{i=1}^d \sqrt{\sum_{t=1}^T \nabla_i F(\mathbf{w}_t)^2} \right] &\leq \frac{2\sqrt{3}}{\eta} \mathbb{E} \left[\sum_{i=1}^d \sum_{t=1}^T \frac{\hat{\eta}_{t,i}}{2} \nabla_i F(\mathbf{w}_t)^2 \right] \\ &\quad + \sqrt{\frac{2}{\eta}} \sqrt{\left(\sum_{i=1}^d \sigma_i \sqrt{2T} + d\delta \right)} \sqrt{\mathbb{E} \left[\sum_{i=1}^d \sum_{t=1}^T \frac{\hat{\eta}_{t,i}}{2} \nabla_i F(\mathbf{w}_t)^2 \right]} \end{aligned} \quad (22)$$

We can further bound the term using the result from Corollary 3.3,

$$\begin{aligned} \mathbb{E} \left[\sum_{i=1}^d \sqrt{\sum_{t=1}^T \nabla_i F(\mathbf{w}_t)^2} \right] &\leq \frac{2\sqrt{3}}{\eta} \left(F(\mathbf{w}_1) - F^* + \left(2\eta \|\boldsymbol{\sigma}\|_1 + \frac{\eta^2 \|\mathbf{L}\|_1}{2} \right) \log h(T) \right) \\ &\quad + \sqrt{\frac{2}{\eta T}} \sqrt{(\|\boldsymbol{\sigma}\|_1 \sqrt{2T} + d\delta)} \sqrt{F(\mathbf{w}_1) - F^* + \left(2\eta \|\boldsymbol{\sigma}\|_1 + \frac{\eta^2 \|\mathbf{L}\|_1}{2} \right) \log h(T)} \end{aligned}$$

where $h(T) = 1 + \frac{T \|\boldsymbol{\sigma}\|_\infty^2}{\delta^2} + \frac{T(\|\nabla F(\mathbf{w}_1)\| + \eta \sqrt{d} \|\mathbf{L}\|_\infty T)^2}{\delta^2}$. Hence the result in the lemma follows. This completes the proof.

B Lower Bound Result

In this section we provide details of the lower bound function. The function $p : \mathbb{R}^d \mapsto \mathbb{R}$ to prove the lower bound result is stated below,

$$p(\mathbf{x}) = \frac{1}{d} \sum_{i=1}^d f_i(\sqrt{d}x_i)$$

From the result in Theorem 4 in [CBS23], the explicit definition of $f_i : \mathbb{R} \mapsto \mathbb{R}$ can be constructed as follows. Consider the queries made as $\mathbf{y}_t, \dots, \mathbf{y}_N$ and for some coordinate i the queries are then $y_{1,i}, \dots, y_{N,i}$ such that $y_{t,i} \in (0, 1/\epsilon) \forall t \in [N]$. Let $l_{t,i} := y_{t+1,i} - y_{t,i}$. Further there exists a function $\Phi_{t,i}$ that satisfies the following properties,

1. $\Phi_{t,i}$ is continuously differentiable and 1-smooth on $[y_{t,i}, y_{t+1,i}]$
2. $\Phi_{t,i}(y_{t,i}) = 0$ and $\Phi_{t,i}(y_{t+1,i}) = l_{t,i}(\frac{l_{t,i}}{4} - \epsilon)$
3. $\Phi'_{t,i}(y_{t,i}) = \Phi'_{t,i}(y_{t+1,i}) = -\epsilon$

The definition of the main function for each coordinate is as below,

$$f_i(z_i) := \begin{cases} 1 - \epsilon z_i & z_i \in (-\infty, 0] \\ f_i(y_{t,i}) - \epsilon(z_i - y_{t,i}) & z \in [y_{t,i}, y_{t+1,i}] \text{ and } l_{t,i} < 8\epsilon(0 \leq t \leq N) \\ f_i(y_{t,i}) + \Phi_{t,i}(z_i) & z \in [y_{t,i}, y_{t+1,i}] \text{ and } l_{t,i} < 8\epsilon(0 \leq t \leq N) \\ f_i(z_i - 1/\epsilon) + f_i(1/\epsilon) - f_i(0) & z \in [1/\epsilon, \infty) \end{cases}$$

The function is 1-smooth and satisfies $f_i(0) - \inf f_i \leq 1$ when $N \leq O(1/\epsilon^2)$, this completes the construction.