

PRESCRIBED MEAN CURVATURE FLOW FOR NONCOMPACT HYPERSURFACES IN LORENTZ MANIFOLDS

LUEN-FAI TAM

ABSTRACT. Motivated by previous study on mean curvature flow and prescribed mean curvature flow on spatially compact space or asymptotically flat spacetime, in this work we will find sufficient conditions for the short time existence of prescribed mean curvature flow on a Lorentz manifold with a smooth time function starting from a complete noncompact spacelike hypersurface. Long time existence and convergence will also be discussed. Results will be applied to study some prescribed mean curvature flows inside the future of the origin in the Minkowski spacetime. Examples of spacetime related to the existence and convergence results near the future null infinity of the Schwarzschild spacetime are also discussed.

1. INTRODUCTION

There are well-developed theories on mean curvature flow and prescribed mean curvature flow in Euclidean space and in Riemannian manifolds. Classical mean curvature flow was developed by of Huisken [20] and others. One may consult the book [2] by Andrews-Chow-Guenther-Langford and the references therein. On the other hand, people are interested in constructing maximal or constant mean curvature spacelike hypersurfaces in Lorentz manifolds. Using elliptic method important results have been obtained by Bartnik [3, 4], Bartnik-Simon [5], Gerhard [15], Treibergs [29] and Andersson-Iriondo [1], to name a few. It is natural to try to use mean curvature flow or prescribed mean curvature flow to construct such kind of hypersurfaces. In this direction, works have been done by Ecker-Huisken [10], Gerhard [13] in the spatially compact case, and by Ecker [9, 8] in the noncompact case, for example. In particular, using the constructed maximal surfaces in [3] as barriers, in [8] Ecker was able to use the mean curvature flow to construct spacelike maximal hypersurfaces in asymptotically flat spacetime. Other results on mean curvature flow and prescribed mean curvature flow for noncompact hypersurfaces in Minkowski spacetime have been studied thoroughly in [9]. Recently there are also interesting works by Kröcke et al. [22] and Gentile-Vertman [12] on graphical prescribed mean curvature flows on some product or warped product

Date: June 10, 2024.

2010 Mathematics Subject Classification. Primary 53C44, Secondary 83C30.

Key words and phrases. prescribed mean curvature flow, Lorentz manifolds, spacelike hypersurfaces.

spacetimes. These motivate us to study prescribed mean curvature flows for complete noncompact spacelike hypersurface in general Lorentz manifolds. In this work, we want to find sufficient conditions for short time existence, long time existence and convergence of prescribed mean curvature flows in Lorentz manifolds. The results will be obtained by using well developed techniques, especially those by Bartnik [3], Ecker-Husiken [10] and Ecker [8].

To state our results, let (N^{n+1}, G) be a Lorentz manifold with connection $\overline{\nabla}$ and curvature tensor $\overline{\text{Rm}}$ and Ricci tensor $\overline{\text{Ric}}$. Assume there is a time function τ on N so that $\overline{\nabla}\tau$ is time like past directed with *lapse function* α given by $\alpha^{-2} = -\langle \overline{\nabla}\tau, \overline{\nabla}\tau \rangle$ and let $\mathbf{T} = -\alpha\overline{\nabla}\tau$ be the future pointing unit time like vector field. Suppose M is spacelike hypersurface in N with future pointing unit normal \mathbf{n} , the *tilt factor* of M with respect to \mathbf{T} is given by $-\langle \mathbf{n}, \mathbf{T} \rangle_G$ and the *height function* u of M is the function $u = \tau|_M$. To measure the size of a tensor, as in [3, 19], introduce the reference Riemannian metric G_E with respect to \mathbf{T} by

$$G_E = G + 2\omega \otimes \omega$$

where ω is the one form dual to \mathbf{T} . For a tensor B in N , the square of its norm at each point with respect to G_E is given by $\langle B, B \rangle_{G_E}$ and its norm is denoted by $|||B|||_{G_E}$. For any $\Omega \subset N$, we define for $k \geq 0$:

$$(1.1) \quad |||B|||_{G_E; k, \Omega} =: \sup_{0 \leq l \leq k} \sup_{\Omega} |||\overline{\nabla}^l B|||_{G_E}.$$

We will omit the subscript G_E if there is no confusion.

Let $\mathbf{X}_0 : M \rightarrow N$ be an immersed spacelike hypersurface and \mathcal{H} be a smooth function in N , then the prescribed mean curvature flow starting from \mathbf{X}_0 is a map $\mathbf{F} : M \times [0, s_0) \rightarrow N$ satisfying

$$(1.2) \quad \begin{cases} \frac{\partial}{\partial s} \mathbf{F}(p, s) = [(H - \mathcal{H})\vec{\nu}](p, s) & \text{in } M \times [0, s_0), \\ \mathbf{F}|_{s=0} = \mathbf{X}_0, \end{cases}$$

such that $\mathbf{F}^s(\cdot) = \mathbf{F}(\cdot, s)$ is a spacelike immersion, with mean curvature vector $\mathbf{H} = \vec{\nu}H$ where H is the mean curvature, $\vec{\nu}$ is the future timelike unit normal. $H, \mathcal{H}, \vec{\nu}$ are evaluated at the point $\mathbf{F}(p, s)$. We also denote M_s to be the immersed surface given by \mathbf{F}^s . In this work, we will study the questions of short time existence, longtime existence and long time behavior of solutions of (1.2) with M_0 being noncompact. For the short time existence, we have the following:

Theorem 1.1. *Let (N^{n+1}, G) , $n \geq 3$, and τ be as above. Let $\mathbf{X}_0 : M^n \rightarrow N$ be an immersed noncompact spacelike hypersurface with future time like unit normal \mathbf{n} and second fundamental form A so that $g_0 = \mathbf{X}_0^*(G)$ is a complete Riemannian metric and let \mathcal{H} be a smooth function on N . Assume the following:*

- (i) $\mathbf{X} : (-a, a) \times M \rightarrow N$ given by $\mathbf{X}(p, t) = \exp_{\mathbf{X}_0(p)}(\mathbf{tn}(\mathbf{X}_0(p)))$ is defined for some $a > 0$;
- (ii) $|||\overline{\nabla} \mathbf{T}|||_{0, \Omega} \leq c$ for some constant c , where $\Omega = \mathbf{X}((-a, a) \times M)$;
- (iii) $|||\overline{\text{Rm}}|||_{k, \Omega} \leq c_k$ for some constant c_k for all $k \geq 0$;
- (iv) $||\nabla^k A|| \leq a_k$ for some constant a_k on $\mathbf{X}_0(M)$ for all $k \geq 0$, where ∇ is the covariant derivative of the induced metric g_0 and the norm is with respect to this metric;
- (v) the tilt factor $-\langle \mathbf{n}(\mathbf{X}_0(p)), \mathbf{T} \rangle \leq v_0$ for some $v_0 > 0$ for all $p \in M$; and
- (vi) For $k \geq 0$, $|||\mathcal{H}|||_{k, \Omega} < \infty$.

Then the prescribed mean curvature flow (1.2) has a solution \mathbf{F} on $M \times [0, \epsilon]$ for some $\epsilon > 0$. Moreover, the immersed hypersurface M_s is complete, the second fundamental forms of M_s together with all covariant derivatives are uniformly bounded in space and time and the tilt factor $-\langle \vec{\nu}(s), \mathbf{T} \rangle$ of M_s is also uniformly bounded in space and time.

To study the question of long time existence, let us fix some notation. Let

$$\tau_- = \inf_N \tau, \quad \tau_+ = \sup_N \tau$$

which may be infinite. For any $\tau_- < \tau_1 < \tau_2 < \tau_+$, let

$$\Omega_{\tau_1, \tau_2} = \{p \in N \mid \tau_1 < \tau(p) < \tau_2\}.$$

Consider the following conditions:

$$(1.3) \quad \begin{cases} |||\alpha|||_{0, \Omega_{\tau_1, \tau_2}}, |||\overline{\nabla} \log \alpha|||_{0, \Omega_{\tau_1, \tau_2}}, |||\overline{\nabla} \mathbf{T}|||_{1, \Omega_{\tau_1, \tau_2}} \text{ are finite;} \\ |||\overline{\text{Rm}}|||_{k, \Omega_{\tau_1, \tau_2}}, \text{ is finite for all } k \geq 0; \\ |||\mathcal{H}|||_{k, \Omega_{\tau_1, \tau_2}} \text{ is finite all } k \geq 0. \end{cases}$$

Since G_E is not complete in general, we introduce the condition: There is a smooth function $\rho > 0$ defined on N such that for any $\tau_- < \tau_1 < \tau_2 < \tau_+$,

$$(1.4) \quad \begin{cases} |||\overline{\nabla} \rho|||_{\Omega_{\tau_1, \tau_2}} \text{ is finite;} \\ \text{for any } \rho_0 > 0 \text{ the set } \overline{\Omega}_{\tau_1, \tau_2} \cap \{p \in N \mid \rho(p) \leq \rho_0\} \text{ is compact.} \end{cases}$$

We have the following characterization of maximum time of existence:

Theorem 1.2. *Let (N^{n+1}, G) be a Lorentz manifold with a time function τ and a smooth function $\rho > 0$ and let \mathcal{H} be a smooth function on N so that they satisfy (1.3) and (1.4). In addition, assume $\overline{\text{Ric}}(w, w) \geq -c^2$ for all unit timelike vector w for some c . Suppose $\mathbf{X}_0 : M \rightarrow N$ is a noncompact immersed spacelike hypersurface with $\mathbf{X}_0(M) \subset \Omega_{\tau_1, \tau_2}$ for some $\tau_- < \tau_1 < \tau_2 < \tau_+$, such that*

- (i) the induced metric on $\mathbf{X}_0(M)$ is complete so that its second fundamental form A satisfies $\sup_{\mathbf{X}_0(M)} |\nabla^k A| < \infty$ for all $k \geq 0$;
- (ii) its tilt factor κ_0 is uniformly bounded;
- (iii) $\inf_{M_0} \alpha > 0$, where α is the lapse function of τ .

Let $s_{\max} > 0$ be the supremum of s_0 so that the prescribed mean curvature flow (1.2) is defined on $M \times [0, s_0]$ with $\sup_{M \times [0, s_0]} |\nabla^k A| < \infty$ for all $k \geq 0$, and $\sup_{M \times [0, s_0]} \kappa < \infty$, where A is the second fundamental form of M_s and κ is its tilt factor. If $s_{\max} < \infty$ then

$$\sup_{M \times [0, s_{\max})} \tau(\mathbf{F}) = \tau_+, \text{ or } \inf_{M \times [0, s_{\max})} \tau(\mathbf{F}) = \tau_-.$$

Finally, we want to study the long time behavior of solution to (1.2). We state two results.

Theorem 1.3. *Let (N^{n+1}, G) , \mathbf{X}_0 , \mathcal{H} and ρ be as in Theorem 1.2. Suppose (1.2) has a long time solution \mathbf{F} defined on $M \times [0, \infty)$. In addition, assume the following:*

- (i) $\tau_- < \tau_1 \leq \tau(\mathbf{F}) \leq \tau_2 < \tau_+$ on $M \times [0, \infty)$ for some τ_1, τ_2 .
- (ii) For any $0 < s < \infty$ and any $m \geq 0$,

$$\sup_{M \times [0, s)} (|\nabla^m A| + \kappa) < \infty,$$

where A is the second fundamental form of M_s and κ is its tilt factor.

- (iii) $\mathcal{H} \geq a > 0$ for some constant a , \mathcal{H} is monotone in the sense that $\langle \overline{\nabla} \mathcal{H}, w \rangle \geq 0$ for all future directed timelike vector. $H - \mathcal{H} \geq -a + \varepsilon_0$ for some $0 < \varepsilon_0 < a$ on M_0 .
- (iv) $\overline{\text{Ric}}(w, w) \geq -c^2$ for any timelike unit vector w with $\frac{c_0^2}{n} - c^2 > 0$.

Then as $s \rightarrow \infty$, $\mathbf{F}(\cdot, s)$ converges in C^∞ norm in compact sets in M to \mathbf{F}_∞ so that $\mathbf{F}_\infty : M \rightarrow N$ is an immersed spacelike hypersurface which is complete with the induced metric and such that the mean curvature H_∞ is equal to \mathcal{H} . Moreover, if α^{-1} is uniformly bounded in Ω_{τ_1, τ_2} and if there is a smooth function $\phi \geq 1$ on M_0 with $\sup_{M_0} (|\nabla \phi| + |\nabla^2 \phi|) < \infty$ and $\phi \rightarrow \infty$ as $\mathbf{x} \rightarrow \infty$ in M_0 so that $\sup_{M_0} \phi |H - \mathcal{H}| < \infty$, then the height functions u_∞ of M_∞ and u_0 of M_0 satisfy $\lim_{\mathbf{x} \rightarrow \infty} (u_\infty(\mathbf{x}) - u_0(\mathbf{x})) = 0$.

We may relax the condition that $\mathcal{H} \geq a > 0$ in the above.

Theorem 1.4. *With the same assumptions and notation as in Theorem 1.3, with conditions (iii) and (iv) to be replaced by the following:*

- (iii') $\mathcal{H} \geq 0$ and \mathcal{H} is monotone in the sense that $\langle \overline{\nabla} \mathcal{H}, w \rangle \geq 0$ for all future directed timelike vector. $H - \mathcal{H} \geq 0$ on M_0 .
- (iv') $\overline{\text{Ric}}(w, w) \geq 0$ for any timelike unit vector w .

Suppose $\mathbf{F}(\cdot, s)$ is an embedding for all s . Then there exist $s_k \rightarrow \infty$ and a spacelike hypersurface M_∞ with mean curvature \mathcal{H} such that $M_{s_k} \rightarrow M_\infty$ in the sense that for any compact $W \subset \overline{\Omega}_{\tau_1, \tau_2}$ set $d_h(M_{s_k} \cap W, M \cap W) \rightarrow 0$ as $k \rightarrow \infty$, where d_h is the Hausdorff distance defined by G_E .

Let us discuss some of the assumptions in the above theorems. In Theorem 1.1, unlike the case of complete Riemannian manifold, condition (i) in general may not be true, unless, for example, the spacetime is timelike geodesically

complete. Condition (ii) is to ensure that the level sets $t = \text{constant}$ have uniformly bounded tilt factors, at least for short time under the assumption (v). Together with (iii), (iv) one can construct suitable Banach spaces in order to apply inverse function theorem. From the proof, one can see that it is sufficient to assume (iii), (iv), (vi) are true up to $k = 3$. In case M_0 is smooth and compact without boundary, then (iv)–(v) will be satisfied automatically, and the proof can be applied. We would like to point out that the method of proof of Theorem 1.1, in a more simple way, can be used to obtain short time existence of prescribed mean curvature flow starting from a complete noncompact hypersurface in a complete Riemannian manifold under suitable assumptions. In Theorem 1.2, the condition on $\overline{\text{Ric}}$ is used to obtain estimate of the tilt factor which has been derived by Ecker [8]. The assumption on the existence of the function ρ is to ensure limit exists. Condition (iii) is related to condition (i) of Theorem 1.1, so that one can use that theorem to extend the prescribed mean curvature flow. Roughly speaking, Theorem 1.2 means that longtime existence is true if we have height estimates, which can be obtained if there exist suitable barrier surfaces. In Theorem 1.3, conditions (iii) and (iv) are used to assure the flow \mathbf{F} will converge exponentially fast as in [12]. (iii) can be replaced by the conditions that \mathcal{H} is monotone and $\overline{\text{Ric}}(w, w) \geq c^2 > 0$ for all unit timelike vector. In Theorem 1.4, the result is weaker. We do not claim that $\mathbf{F}(\cdot, s)$ is convergent as maps. We only consider the convergence of hypersurfaces in terms of Hausdorff distance as in [4, Theorem 3.8]. Moreover, it is unclear if the limit surface is complete. In case, $\mathcal{H} \equiv 0$, then we do not need to assume $H \geq 0$. In this case, this is the result in [8]. We should remark by changing the time orientation, dual results are true for Theorems 1.3 and 1.4. See Remark 4.1 for more details.

By the above results, in order to use prescribed mean curvature flow to obtain prescribed mean curvature hypersurfaces, under suitable assumptions, it is sufficient to obtain height estimates. This can be done if suitable barrier surfaces exist. See Corollary 4.1 and Proposition 5.1 for examples. Also, under additional assumptions on the foliation in [1], result similar to the result for maximal surface in asymptotically flat spacetime in [8] might also true for positive constant mean curvature cut in asymptotically Schwarzschild spacetime which was introduced and studied in [1].

The organization of this work is as follows: In section 2 we will prove the short time existence result. We need some preliminary results which will be discussed in subsection 2.1. In section 3, we will prove the long time existence result Theorem 1.2 and in section 4, we will prove the convergence results Theorems 1.3 and 1.4. In subsection 5.1, we will give some applications on Minkowski spacetime. In subsection 5.2, we will do some computations in Schwarzschild spacetime near the future null infinity, which are related to the conditions in theorems mentioned above and might be related to the work [1]. In the appendix, we recall the definitions of certain Hölder spaces, which are

used in the proof of short time existence, and we prove a convergence lemma which is used in the proof of Theorem 1.4.

Acknowledgement: The author would like to thank Albert Chau for useful discussions.

2. SHORT TIME EXISTENCE

2.1. Setup and preliminary results. Before we prove the short time existence Theorem 1.1, we need some preparation. Let (N, G) , $\overline{\nabla}$, $\overline{\text{Rm}}$, τ , α , \mathbf{T} , G_E be as in section 1. First, it is obvious that G_E depends on τ and will be called the reference Riemannian metric with respect to τ . If τ' is another time function with corresponding future pointing unit normal \mathbf{T}' , and let G'_E be the reference Riemannian metric with respect to τ' , then G_E, G'_E are different. However, if $-\langle \mathbf{T}, \mathbf{T}' \rangle$ (which is bounded from below by 1) is uniformly bounded from above then they will be equivalent by the following result which is well-known and has been used in [3], see [10] for example:

Lemma 2.1. *Let (V^{n+1}, G) be a Lorentz vector space. Choose a fixed time orientation. Let \mathbf{T}, \mathbf{T}' be two future unit time like unit vectors with dual one forms ω, ω' . Consider the Euclidean metrics $G_E = G + 2\omega \otimes \omega$, $G'_E = G + 2\omega' \otimes \omega'$. Let B be a tensor on V . Then the norms $\| \| B \| \|_{G_E}, \| \| B \| \|_{G'_E}$ are equivalent. In fact, if $v = -\langle \mathbf{T}, \mathbf{T}' \rangle$, and B is a (k, l) tensor, then*

$$\| \| B \| \|_{G'_E}^2 \leq (n+1)^{k+l} (4v^2)^{k+l} \| \| B \| \|_{G_E}^2.$$

Proof. We sketch the proof. Let $e_0 = \mathbf{T}, e'_0 = \mathbf{T}'$. By choosing an orthonormal basis for the intersection of e_0^\perp and $(e'_0)^\perp$, which are the subspaces perpendicular to e_0, e'_0 respectively, we may find e_1, e'_1 and e_2, \dots, e_n so that $e_0, e_1, e_2, \dots, e_n$ and $e'_0, e'_1, e_2, \dots, e_n$ are orthonormal. It is easy to see that

$$e_0 = \cosh \lambda e'_0 + \sinh \lambda e'_1; \quad e_1 = \sinh \lambda e'_0 + \cosh \lambda e'_1$$

where $v = \cosh \lambda$. Let w be a vector so that

$$w = \sum_{\alpha=0}^n a_\alpha e_\alpha = (a_0 \cosh \lambda + a_1 \sinh \lambda) e'_0 + (a_0 \sinh \lambda + a_1 \cosh \lambda) e'_1 + \sum_{i=2}^n a_i e_i.$$

Hence

$$\begin{aligned} \| \| w \| \|_{G'_E}^2 &= (a_0 \cosh \lambda + a_1 \sinh \lambda)^2 + (a_0 \sinh \lambda + a_1 \cosh \lambda)^2 + \sum_{i=2}^n a_i^2 \\ &\leq 2(\cosh^2 \lambda + \sinh^2 \lambda) \sum_{\alpha=0}^n a_\alpha^2 \\ &\leq 4v^2 \| \| w \| \|_{G_E}^2. \end{aligned}$$

Similarly, $|||w|||_{G_E}^2 \leq 4v^2 |||w|||_{G'_E}^2$. On the other hand, take a $(0, 2)$ tensor B for example. Let e'_α be orthonormal with $e'_0 = \mathbf{T}'$.

$$\begin{aligned} |||B|||_{G'_E}^2 &= \sum_{\alpha, \beta} B^2(e'_\alpha, e'_\beta) \\ &\leq \sum_{\alpha, \beta} |||B|||_{G_E}^2 |||e'_\alpha|||_{G_E}^2 |||e'_\beta|||_{G_E}^2 \\ &\leq 16v^4(n+1)^2 |||B|||_{G_E}^2. \end{aligned}$$

The general case can be proved similarly. \square

Let $\mathbf{X}_0 : M \rightarrow N$ be an immersion as a spacelike hypersurface. We assume that the induced metric $g_0 = \mathbf{X}_0^*(G)$ on M is complete. Define: $\mathbf{X} : (-a, a) \times M \rightarrow N$ by

$$(2.1) \quad \mathbf{X}(t, p) = \exp_{\mathbf{X}_0(p)}(\mathbf{tn}(\mathbf{X}_0(p)))$$

We assume that there is $a > 0$ so that \mathbf{X} is defined for all $p \in M$ and $t \in (-a, a)$. Let $\Omega = \mathbf{X}((-a, a) \times M)$. Note that

$$\mathbf{X}^*(G) = -dt^2 + g(t)$$

where $g(t)$ is the symmetric tensor on the slice $M_t = \{t\} \times M$: $g(t) = (\mathbf{X}^t)^*(\cdot, \cdot)(G)$ where $\mathbf{X}^t(p) = \mathbf{X}(t, p)$. In local coordinates x^1, \dots, x^n of M , then $t = x^0, x^1, \dots, x^n$ will be local coordinates of $(-a, a) \times M$. In the following α, β, \dots run from 0 to n and i, j, \dots run from 1 to n . We want to find prescribed mean curvature flow which can be expressed as a function $t = w(p, s)$. Namely, the flow up to diffeomorphisms on M is given by $\mathbf{F}(p, s) = \exp_{\mathbf{X}_0(p)}(w(p, s)\mathbf{n}(\mathbf{X}_0(p)))$. We try to solve for w . We need some preparations.

Lemma 2.2. *Assume $|||\bar{\nabla} \mathbf{T}|||_{G_E} \leq c$ in Ω . Let $v = -\langle \frac{\partial}{\partial t}, \mathbf{T} \rangle_G$, where we identify $\frac{\partial}{\partial t}$ in $(-a, a) \times M$ and $\mathbf{X}_*(\frac{\partial}{\partial t})$. Suppose $v(p, 0) \leq v_0$ for all $p \in M$. Then*

$$v(p, t) \leq 2v_0$$

for all for $p \in M$ and $|t| \leq \frac{1}{4cv_0}$.

Proof. In the following, we will denote $|||\cdot|||_{G_E}$ with respect to \mathbf{T} simply by $|||\cdot|||$. $\frac{\partial}{\partial t}$ is just the tangent vector of the geodesic $t \rightarrow \exp_{\mathbf{X}_0(p)}(\mathbf{tn}(\mathbf{X}_0(p)))$ which is a future pointing unit vector.

$$|||\bar{\nabla}_{\frac{\partial}{\partial t}} \mathbf{T}||| \leq |||\frac{\partial}{\partial t}||| |||\bar{\nabla} \mathbf{T}||| \leq c |||\frac{\partial}{\partial t}|||.$$

Since $\langle \frac{\partial}{\partial t}, \frac{\partial}{\partial t} \rangle = -1$, by Lemma 2.1, we have

$$|||\frac{\partial}{\partial t}|||^2 \leq 2v^2.$$

Hence

$$\left| \frac{\partial}{\partial t} v \right| = \left| \left\langle \bar{\nabla}_{\frac{\partial}{\partial t}} \mathbf{T}, \frac{\partial}{\partial t} \right\rangle \right| \leq \| \bar{\nabla}_{\frac{\partial}{\partial t}} \mathbf{T} \| \| \frac{\partial}{\partial t} \| \leq 2cv^2.$$

So if $|t| \leq \frac{1}{4cv_0}$, then $v(t) \leq 2v_0$. □

From now on, in the above setting, we always assume the following:

- (i) $\| \bar{\nabla} \mathbf{T} \| \leq c$;
- (ii) $-\langle \frac{\partial}{\partial t}, \mathbf{T} \rangle \leq v_0$ at $t = 0$;
- (iii) $a < \frac{1}{4cv_0}$ so that $-\langle \frac{\partial}{\partial t}, \mathbf{T} \rangle \leq 2v_0$ for $|t| < a$.

Lemma 2.3. *Assume $c_0 = \sup_{\Omega} \| \bar{\text{Rm}} \|_{G_E} < \infty$ and $a_0 = \sup_{M_0} |A| < \infty$ where $M_0 = \{0\} \times M$ and A is the second fundamental form of the immersed surface given by \mathbf{X}_0 , which will also be denoted by M_0 . Then \mathbf{X}^t is an immersion for $|t| < \min\{a, \frac{1}{2}C(n)(a_0^2 + c_0v_0^2)^{-\frac{1}{2}}\}$, for some constant $C(n)$ depending only on n . Moreover,*

$$e^{-2b_0t} g_0 \leq g \leq e^{2b_0t} g_0; \quad |A(t)| \leq b_0.$$

where $b_0 = 2(a_0^2 + c_0v_0^2)^{\frac{1}{2}}$. Here $A(t)$ is the second fundamental form of the hypersurface given by the immersion \mathbf{X}^t .

Proof. Let $p \in M$ be fixed. Let x^1, \dots, x^n be local coordinates of p and let $t = x^0$. Denote $\mathbf{X}_*(\frac{\partial}{\partial x^i})$ simply by ∂_i and $\mathbf{X}_*(\frac{\partial}{\partial t})$ by ∂_t or ∂_0 . Since ∂_t is the unit tangent vector of the time like geodesic $t \rightarrow \exp_{\mathbf{X}_0(q)}(t\mathbf{n}(\mathbf{X}_0(q)))$,

$$\frac{\partial}{\partial t} \langle \partial_t, \partial_i \rangle = \langle \bar{\nabla}_{\frac{\partial}{\partial t}} \frac{\partial}{\partial t}, \partial_i \rangle + \langle \frac{\partial}{\partial t}, \bar{\nabla}_{\frac{\partial}{\partial t}} \partial_i \rangle = 0.$$

because $\bar{\nabla}_{\frac{\partial}{\partial t}} \frac{\partial}{\partial t} = 0$, $\bar{\nabla}_{\frac{\partial}{\partial t}} \partial_i = \bar{\nabla}_{\partial_i} \frac{\partial}{\partial t}$ and $\langle \frac{\partial}{\partial t}, \frac{\partial}{\partial t} \rangle = -1$. Let $g_{ij} = \langle \partial_i, \partial_j \rangle$. Then

$$(2.2) \quad \frac{\partial}{\partial t} g_{ij} = \langle \bar{\nabla}_{\partial_t} \partial_i, \partial_j \rangle + \langle \partial_i, \bar{\nabla}_{\partial_t} \partial_j \rangle = 2A_{ij}$$

where $A_{ij} = \langle \bar{\nabla}_{\partial_t} \partial_i, \partial_j \rangle$ which is equal to $\langle \partial_i, \bar{\nabla}_{\partial_t} \partial_j \rangle$ by the above. Suppose $a \geq a_1 > 0$ is the largest value so that \mathbf{X}^t is an immersion near p for all $|t| < a_1$. Then A_{ij} is the second fundamental form of the immersed hypersurface defined by \mathbf{X}^t . For such t

$$\begin{aligned} \frac{\partial}{\partial t} A_{ij} &= \langle \bar{\nabla}_{\partial_t} \bar{\nabla}_{\partial_t} \partial_i, \partial_j \rangle + \langle \bar{\nabla}_{\partial_t} \partial_i, \bar{\nabla}_{\partial_t} \partial_j \rangle \\ &= \langle \bar{\nabla}_{\partial_t} \bar{\nabla}_{\partial_i} \partial_t, \partial_j \rangle + \langle \bar{\nabla}_{\partial_t} \partial_i, \bar{\nabla}_{\partial_t} \partial_j \rangle \\ &= \bar{\text{R}}_{0i0j} + g^{kl} A_{ik} A_{jl}. \end{aligned}$$

because $\bar{\nabla}_{\partial_t} \partial_t = 0$. Here $\bar{\text{R}}_{0i0j} = \bar{\text{R}}(\partial_0, \partial_i, \partial_0, \partial_j)$. Then by Lemma 2.1,

$$\begin{aligned} \partial_t |A|^2 &= -2g^{ip} g^{lq} g^{kj} A_{pq} A_{ij} A_{kl} + 2g^{il} g^{kj} \bar{\text{R}}_{0i0l} A_{kj} \\ &\leq 2|A|^3 + C(n)c_0v_0^2|A| \\ &\leq C(n)(|A|^2 + c_0v_0^2)^{\frac{3}{2}}. \end{aligned}$$

Hence

$$(|A|^2 + c_1 v_0^2)^{-\frac{1}{2}}(0) - (|A|^2 + c_0 v_0^2)^{-\frac{1}{2}}(t) \leq C(n)t.$$

and

$$(|A|^2 + c_1 v_0^2)^{-\frac{1}{2}}(t) \geq (k^2 + c_0 v_0^2)^{-\frac{1}{2}} - C(n)t.$$

Hence if $t \leq \frac{1}{2}C(n)(a_0^2 + c_0 v_0^2)^{-\frac{1}{2}}$, then

$$|A|(t) \leq 2(a_0^2 + c_1 v_0^2)^{\frac{1}{2}} =: b_0$$

By (2.2), we conclude that

$$e^{-2b_0 t} g_0 \leq g \leq e^{2b_0 t} g_0.$$

From this it is easy to see that $a \geq a_1 \geq \frac{1}{2}C(n)(a_0^2 + c_0 v_0^2)^{-\frac{1}{2}}$. This completes the proof of the lemma. \square

From now on we further assume:

(iv) a also satisfies $a \leq \frac{1}{2}C(n)(a_0^2 + c_0 v_0^2)^{-\frac{1}{2}}$.

Hence $\mathbf{X} : (-a, a) \times M \rightarrow N$ is an immersion. We also denote the immersed surface given by \mathbf{X}^t by M_t .

In the following, in order to study short time existence, we will identify N as $(-a, a) \times M$ and identify G is with $\mathbf{X}^(G)$. This will simplify the exposition.*

To study the prescribed mean curvature flow, let us first compute the mean curvature of a spacelike surface S given by a function w on M : Namely, the surface is the level zero set $\{f(t, \mathbf{x}) = 0\}$ where $f(t, \mathbf{x}) = t - w(\mathbf{x})$, $\mathbf{x} \in M$. Here the ambient manifold is $(-a, a) \times M$ with metric $G = -dt^2 + g(t)$.

Lemma 2.4. *The mean curvature of S with respect to the future pointing unit normal is given by*

$$H = (1 - |\nabla w|^2)^{-\frac{1}{2}} H^o + \operatorname{div}^o \left(\frac{\nabla w}{(1 - |\nabla w|^2)^{\frac{1}{2}}} \right) + \frac{\partial}{\partial t} ((1 - |\nabla w|^2)^{-\frac{1}{2}})$$

evaluated at $t = w$. Here H^o is the mean curvature of the reference slice $M_t = \{t\} \times M$, div^o is divergence on this slice, ∇ is the covariant derivative with respect to $g(t)$ on the slice M_t .

Proof. This follows from [3, p.160]. We sketch the proof. Let $f(t, \mathbf{x}) = t - w(\mathbf{x})$. Then $\bar{\nabla} f = \bar{\nabla} t - \bar{\nabla} w = \bar{\nabla} t - \nabla w$. So the mean curvature of the surface $\{f = 0\}$ with respect to the future pointing unit normal is given by

$$H = \operatorname{div}_G \left(-\frac{\bar{\nabla} f}{(1 - |\nabla w|^2)^{\frac{1}{2}}} \right) \Big|_{t=w}.$$

Let $T = \frac{\partial}{\partial t}$, $U = \nabla w$, $\lambda = (1 - |\nabla w|^2)^{-\frac{1}{2}}$. Then $H = \operatorname{div}_G(\lambda(T + U))|_{t=w}$. Now

$$\begin{aligned} \operatorname{div}_G(\lambda(T + U)) &= \operatorname{div}_G(\lambda U) + \operatorname{div}_G(\lambda T) \\ &= \operatorname{div}^o(\lambda U) + \langle \overline{\nabla}_T(\lambda U), T \rangle + \lambda \operatorname{div}_G(T) + T(\lambda) \\ &= \operatorname{div}^o(\lambda U) + \lambda H^o + T(\lambda) \end{aligned}$$

because $U \perp T$ and $\overline{\nabla}_T T = 0$. This completes the proof of the lemma. \square

Let x^i be local coordinates of M so that t, x^i are local coordinates of $(-a, a) \times M$. Then

$$\begin{aligned} H^o &= g^{ij} A_{ij}^o = \frac{1}{2} g^{ij} \partial_t g_{ij} \\ T(\lambda) &= \lambda^2 \partial_t g^{ij} w_i w_j \end{aligned}$$

$$\operatorname{div}^o \left(\frac{\nabla w}{(1 - |\nabla w|^2)^{\frac{1}{2}}} \right) = \lambda (g^{ij} + \lambda^2 w^i w^j) w_{;ij}$$

where $w_{;ij}$ is the Hessian of w in M_t . Hence we have the following remark.

Remark 2.1. The mean curvature is of the form

$$\begin{aligned} (2.3) \quad H &= (1 - |\nabla w|^2)^{-\frac{1}{2}} \left(g^{ij} + \frac{w^i w^j}{1 - |\nabla w|^2} \right) w_{;ij} + \frac{w^i w^j}{1 - |\nabla w|^2} \partial_t g^{ij} \\ &\quad + \frac{1}{2} (1 - |\nabla w|^2)^{-\frac{1}{2}} g^{ij} \partial_t g_{ij} \end{aligned}$$

evaluated at $t = w$.

From the remark, in order to study prescribed mean curvature flow, we need to study the equation involving H , which is a second order differential operator with coefficients of the form $\phi(t, \mathbf{x})|_{t=w}$ where $\phi(\mathbf{x}, t)$ can be expressed in terms of $g(t, \mathbf{x})$, $g^{-1}(t, \mathbf{x})$ and their derivatives. So we need to estimate g, g^{-1} and their derivatives as functions of t, \mathbf{x} .

In the above setting, let us make the following assumptions. Recall that $\Omega = (-a, a) \times M$ in this setting.

Assumption 2.1. *Assume:*

- (i) $\|\overline{\nabla} \mathbf{T}\|_{\Omega} \leq c$;
- (ii) $-\langle \mathbf{X}_* \left(\frac{\partial}{\partial t} \right), \mathbf{T} \rangle \leq v_0$ at $t = 0$;
- (iii) $\|\overline{\nabla}^k \overline{\mathbf{R}}\|_{\Omega} \leq c_k$ for all $k \geq 0$;
- (iv) $|\nabla^k A| \leq a_k$ on $M_0 = \{0\} \times M$ for all $k \geq 0$, where the ∇ is the covariant derivative of the metric g_0 induced by G and the norm is with respect to this metric.
- (v) $a \leq \min \left\{ \frac{1}{2} C(n) (k_0^2 + c_0 v_0^2)^{-\frac{1}{2}}, \frac{1}{2} C(n) (a_0^2 + c_0 v_0^2)^{-\frac{1}{2}} \right\}$ which are given in Lemmas 2.2 and 2.3.

Under this assumption, we identify N with $(-a, a) \times M$ with metric $G = -dt^2 + g(t)$.

Lemma 2.5. *Under the Assumption 2.1, there is $r > 0$ such that for all $p \in M_0$, $\xi_p =: \exp_p : D(r) \rightarrow B_p(r)$ is a surjective local diffeomorphism where $D(r) = \{\mathbf{x} \in \mathbb{R}^n \mid |\mathbf{x}| < r\}$ and $B_p(r)$ is the geodesic ball of M_0 with respect to the metric g_0 . Moreover, $C^{-1}g_e \leq \xi_p^*(g_0) \leq Cg_e$ for some constant $C > 0$ independent of p , where g_e is the Euclidean metric on $D(r)$, and in the standard coordinates x^i of $D(r)$, $\partial^\beta(\xi_p^*(g_0)_{ij})$ are uniformly bounded in $D(r)$ independent of p for all multi-index β and i, j .*

Proof. By [18], it is sufficient to show that the curvature together with its covariant derivatives of g_0 are uniformly bounded. But this follows from (ii), (iii), (iv) in Assumption 2.1, the Gauss equation and Lemma 2.1. \square

In general, the Hölder norm of a function in $B_p(r)$ dominates the Hölder norm of its pull back under ξ_p . On the other hand, if ξ_p is bijective, then these two norms are equivalent. We will call $\xi_p : D(r) \rightarrow B_p(r)$ a *quasi-coordinate neighborhood* and the standard coordinates x^i 's of \mathbb{R}^n will be called *quasi-coordinates*. Suppose Assumption 2.1 holds, and let $\xi_p : D(r) \rightarrow B_p(r)$, $p \in M$ be quasi-coordinate neighborhoods. By Lemma 2.5, one can define the following Hölder spaces for functions on $M \times [0, S_0]$ for $S_0 > 0$:

Definition 2.1. *Let f be a function on $M \times [0, S_0]$ and $k \geq 0$ be an integer. f is said to be in $C^{2k+\sigma, k+\frac{\sigma}{2}}(M \times [0, S_0])$ for $0 < \sigma < 1$ if $\xi_p^*(f) \in C^{2k+\sigma, k+\frac{\sigma}{2}}(D(r))$ so that its Hölder norm $\|\xi_p^*(f)\|_{2k+\sigma, k+\frac{\sigma}{2}}$ in $D(r) \times [0, S_0]$ is uniformly bounded independent of p .*

For the definition of Hölder space, see appendix. $C^{2k+\sigma, k+\frac{\sigma}{2}}(M \times [0, S])$ is a Banach space. We have the following, see [7, Appendix]:

Lemma 2.6. *Suppose $\xi'_p : D(r') \rightarrow B_p(r')$ is another family of quasi local coordinates satisfying the same conditions as in Lemma 2.5. Then the Hölder norms defined by $\{\xi_p \mid p \in M\}$ and the Hölder norms defined by $\{\xi'_p \mid p \in M\}$ are equivalent.*

Since ξ_p is a local diffeomorphism, we may consider the standard coordinates of $D(r)$ as local coordinates of points in $B_p(r)$. In the following lemma, since the computations are local, for simplicity, we still denote $\xi_p^*(g(t))$ by $g(t)$.

Lemma 2.7. *In the setting of Lemma 2.3. Under Assumption 2.1, in each quasi-coordinate neighborhood ξ_p , any order of derivatives of $g_{ij}(t)$ with respect to quasi-coordinates and with respect to t are uniformly bounded independent of p and $|t| < a$. Moreover, $C^{-1}g_e \leq \xi_p^*(g(t)) \leq Cg_e$ for some constant $C > 0$ independent of p, t . Hence any order of derivatives of $g^{ij}(t)$ with respect to quasi-coordinates and with respect to t are uniformly bounded independent of p and $|t| < a$.*

Proof. As mentioned above, we will not distinguish between $\xi_p^*(g(t))$ and $g(t)$. By Lemmas 2.5 and 2.3, it is easy to see that

$$(2.4) \quad C^{-1}g_e \leq \xi_p^*(g(t)) \leq Cg_e$$

in $D(r)$ for some C independent of p, t . Hence $g_{ij}(t), g^{ij}(t)$ are uniformly bounded, where g_{ij} are the components of g with respect to the Euclidean coordinates x^i . By Lemma 2.3 and its proof we have

$$(2.5) \quad \partial_t g_{ij} = 2A_{ij}$$

and $|A(t)|$ is uniformly bounded. Hence A_{ij} and $|\partial_t g_{ij}|$ are uniformly bounded.

Since all order of derivatives of $g_{ij}(0)$ are uniformly bounded, by (2.5), to prove the lemma it is sufficient to prove that $\partial_t^k \partial^\beta A_{ij}$ are uniformly bounded for all $k \geq 0$ and for all multi-index $\beta = (\beta_1, \beta_2, \dots, \beta_n)$ with $\beta_i \geq 0$. Here $\partial^\beta = \partial_1^{\beta_1} \dots \partial_n^{\beta_n}$ and $\partial_i = \frac{\partial}{\partial x^i}$. Let $|\beta| = \sum_{i=1}^n \beta_i$.

We first prove that $\partial^\beta A_{ij}$ are uniformly bounded for all β . Let $\bar{\Gamma}(t) = \Gamma(t) - \Gamma(0)$ where $\Gamma(t)$ are the Christoffel symbols of $g(t)$ with respect to the coordinates x^i . If $B_{ijk\dots}$ is a smooth tensor on $(-a, a) \times M$ which are tangential to $\{t\} \times M$ for all t , for tangential ∂_u :

$$\begin{aligned} (\partial_t \nabla_u B)_{ijk\dots} &= \partial_t \partial_u B_{ijk\dots} - \partial_t (\Gamma_{ui}^p B_{pjkd\dots}) - \partial_t (\Gamma_{uj}^p B_{ipkd\dots}) - \partial_t (\Gamma_{uk}^p B_{ijpd\dots}) - \dots \\ &= \nabla_u (\partial_t B)_{ijk\dots} + \partial_t \bar{\Gamma} * B. \end{aligned}$$

because $\partial_t \bar{\Gamma} = \partial_t \Gamma$. The number of terms depends only on n and the degree of B . We write this simply as:

$$(\partial_t \nabla B) = \nabla \partial_t B + \partial_t \bar{\Gamma} * B.$$

So

$$\begin{aligned} \partial_t (\nabla^k B) &= \nabla \partial_t (\nabla^{k-1} B) + \partial_t \bar{\Gamma} * \nabla^{k-1} B \\ &= \nabla (\nabla \partial_t (\nabla^{k-2} B) + \partial_t \bar{\Gamma} * \nabla^{k-2} B) + \partial_t \bar{\Gamma} * \nabla^{k-1} B \\ &= \nabla^2 (\partial_t (\nabla^{k-2} B)) + \nabla \partial_t \bar{\Gamma} * \nabla^{k-2} B + \partial_t \bar{\Gamma} * \nabla^{k-1} B \\ (2.6) \quad &= \dots \\ &= \nabla^k \partial_t B + \sum_{i=0}^{k-1} \nabla^i \partial_t \bar{\Gamma} * \nabla^{k-i-1} B. \end{aligned}$$

Now

$$(2.7) \quad \frac{\partial}{\partial t} A_{ij} = \bar{R}_{0i0j} + g^{kl} A_{ik} A_{jl}$$

and

$$(2.8) \quad \frac{\partial}{\partial t} \bar{\Gamma}_{ij}^k = \frac{\partial}{\partial t} \Gamma_{ij}^k = g^{kl} (A_{lj;i} + A_{il;j} - A_{ij;l})$$

where ; is the covariant derivative with respect to $g(t)$ and 0 is the index corresponding to $\frac{\partial}{\partial t}$. So $\bar{R}_{0i0j} = \bar{R}(\frac{\partial}{\partial t}, \frac{\partial}{\partial x^i}, \frac{\partial}{\partial t}, \frac{\partial}{\partial x^j})$. Let $B_{ij} = \bar{R}_{0i0j}$. By Lemma 2.3 and the choice of a , the tilt factor of $M \times \{t\}$ for $|t| < a$ is

bounded by $2v_0$. Let c_k be such that $||\overline{\nabla}^k \overline{\text{Rm}}|| \leq c_k$. By (2.4) and Lemma 2.1 we have $|B| \leq C(n, v_0, c_0)$. Suppose $|\nabla^k A| \leq b_k$. Since

$$\nabla_u \overline{\text{R}}_{0i0j} = \overline{\nabla}_u \overline{\text{R}}_{0i0j} + A_u^k (\overline{\text{R}}_{ki0j} + \overline{\text{R}}_{0ikj})$$

we have $||\nabla B|| \leq C(n, v_0, c_0, c_1, b_0)$. Here the norm is taken with respect to $g(t)$. In general, one can show that

$$|\nabla^k B| \leq C(n, v_0, c_0, \dots, c_k; b_0, \dots, b_{k-1}).$$

By (2.8),

$$\nabla^k \partial_t \overline{\Gamma} = g^{-1} * \nabla^{k+1} A.$$

Hence by (2.6) and (2.7), we have

$$\begin{aligned} \partial_t \nabla^k A &= \nabla^k \partial_t A + \sum_{i=0}^{k-1} \nabla^i \partial_t \overline{\Gamma} * \nabla^{k-i-1} A \\ &= \nabla^k (B + g^{-1} * A * A) + \sum_{i=0}^{k-1} \nabla^i \partial_t \overline{\Gamma} * \nabla^{k-i-1} A. \end{aligned}$$

$$|\partial_t \nabla^k A| \leq C(n, v_0, c_0, c_1, \dots, c_k; b_0, b_1, \dots, b_{k-1})(1 + ||\nabla^k A||)$$

Hence

$$\partial_t |\nabla^k A|^2 \leq C(1 + ||\nabla^k A||^2)$$

for some $C = C(n, v_0, c_0, c_1, \dots, c_k; b_0, b_1, \dots, b_{k-1})$. Since $|\nabla^k A|(0)$ is bounded, we conclude that $|\nabla^k A|$ is bounded provided $|\nabla^p A|$ is bounded for $0 \leq p \leq k-1$. Since $|A|$ is bounded, inductively, we have $|\nabla^k A|$ is bounded for all k . By (2.8), and that $|\nabla^k A|$ is bounded for all k , we have $\nabla^k \partial_t \overline{\Gamma}$ is bounded. Using (2.6) and the fact that $\overline{\Gamma} = 0$ at $t = 0$, we can argue as before to show that $\nabla^k \overline{\Gamma}$ is bounded for all k . From these we conclude that $\partial^\beta A_{ij}$ are uniformly bounded.

Next, we want to prove that $\partial_t^k \partial^\beta A_{ij}$ are uniformly bounded. Suppose $\partial_t^k \partial^\beta A_{ij}$ are uniformly bounded for all β , for all $0 \leq k \leq \ell$. Then $\partial_t^{k+1} \partial^\beta g_{ij}$ are uniformly bounded by (2.5) and hence $\partial_t^{k+1} \partial^\beta g^{ij}$ for all β and for all $0 \leq k \leq \ell$. By (2.7)

$$(2.9) \quad \partial_t^{k+1} \partial^\beta A_{ij} = \partial_t^k \partial^\beta (\overline{\text{R}}_{0i0j} + g^{pq} A_{pi} A_{qj}) = \partial_t^k \partial^\beta \overline{\text{R}}_{0i0j} + \partial_t^k \partial^\alpha (g^{pq} A_{pi} A_{qj}).$$

Let $k = \ell$ in the above. Then second term is uniformly bounded. To estimate the first term, by (2.9), $\partial_t^k \partial^\beta \overline{\text{R}}_{0i0j}$ is uniformly bounded for all α and $0 \leq k \leq \ell-1$.

let $\partial_0 = \frac{\partial}{\partial t}$ and $\partial_i = \frac{\partial}{\partial x^i}$ for $i = 1, \dots, n$. Since

$$\begin{cases} \overline{\nabla}_{\partial_i} \partial_j = \Gamma_{ij}^k \partial_k + A_{ij} \frac{\partial}{\partial t} \\ \overline{\nabla}_{\frac{\partial}{\partial t}} \partial_i = \overline{\nabla}_{\partial_i} \frac{\partial}{\partial t} = g^{kl} A_{il} \partial_k \\ \overline{\nabla}_{\frac{\partial}{\partial t}} \frac{\partial}{\partial t} = 0, \end{cases}$$

$$\bar{\nabla} \overline{\text{Rm}} = \partial \overline{\text{Rm}} + \overline{\text{Rm}} * f(A, g, g^{-1}, \partial g).$$

All terms are evaluated using the basis $\partial_0, \partial_1, \dots, \partial_n$. Here and in the following f is a polynomial of its arguments which may change from line to line and ∂ means ∂_γ , $0 \leq \gamma \leq n$. Differentiate the above relation, we have

$$\begin{aligned} \bar{\nabla}^2 \overline{\text{Rm}} &= \partial \bar{\nabla} \overline{\text{Rm}} - \bar{\nabla} \overline{\text{Rm}} * f(A, g, \partial g, g^{-1}) \\ &= \partial^2 \overline{\text{Rm}} + (\partial \overline{\text{Rm}}) * f(A, g, \partial g, g^{-1}) + \overline{\text{Rm}} * f(A, \partial A, g, \partial g, \partial^2 g, g^{-1}). \end{aligned}$$

Continue in this way, we have:

$$\bar{\nabla}^k \overline{\text{Rm}} = \partial^k \overline{\text{Rm}} + \sum_{i=1}^k \partial^{k-i} \overline{\text{Rm}} * f_i$$

where f_i denote polynomials in $\partial^\beta A$ with $|\beta| = s$ for $0 \leq s \leq i-1$, $\partial^s g$ for $0 \leq s \leq i$ and g^{-1} . First note that $\bar{\nabla}^k \overline{\text{Rm}}$ are uniformly bounded for all k by the facts that $\|\bar{\nabla}^k \overline{\text{Rm}}\|_\Omega$ are uniformly bounded, the tilt factors of M_t are uniformly bounded in space time, and by Lemma 2.1 and (2.4). Since $\partial_t^k \partial^\alpha A, \partial^{k+1} \partial^\alpha g, \partial^k \partial^\alpha \overline{\text{Rm}}$ are uniformly bounded for $0 \leq k \leq \ell$ and for all α , $\partial_t^k \partial^\alpha \overline{\text{Rm}}$ are uniformly bounded for $0 \leq k \leq \ell-1$, and g^{-1} is uniformly bounded, we conclude that $\partial_t^\ell \partial^\beta \overline{\text{Rm}}$ are uniformly bounded for all α . This implies that $\partial_t^\ell \partial^\alpha A$ are uniformly bounded for all β . By induction, the lemma follows. \square

2.2. Proof of short time existence. Let $N, G, \tau, M, \mathbf{X}_0$ be as in Theorem 1.1. Then by Lemmas 2.3 and 2.7, we may assume $N = (-a, a) \times M$ with metric $G = -dt^2 + g(t)$ with $a > 0$ small enough. With this setting, $\mathbf{F} : M \times [0, S_0] \rightarrow N$, $S_0 > 0$, is a prescribed mean curvature flow starting from $\mathbf{X}_0 : M \rightarrow N$ if it satisfies (1.2) such that $\mathbf{F}(\cdot, s)$ is a spacelike immersion, with mean curvature vector $\mathbf{H} = \vec{\nu}H$ where H is the mean curvature, $\vec{\nu}$ is the future timelike unit normal, and \mathcal{H} is a function on N . The following fact is well-known.

Lemma 2.8. *Let \mathbf{F} be a solution of the prescribed mean curvature flow (1.2). Suppose in the setting of Assumption 2.1 there is a smooth family of diffeomorphisms $\phi(p, s)$, $0 \leq s \leq S_0$ of M with $\phi(\cdot, 0)$ being the identity map, and a smooth function $w(p, s)$ on $M \times [0, S_0]$ so that in this setting*

$$\mathbf{F}(p, s) = \mathbf{X}(w(\phi(p, s), s), \phi(p, s)) = (w(\phi(p, s), s), \phi(p, s))$$

with $|w| < a$, where a is as in Assumption 2.1 and \mathbf{X} is as in (2.1). Then w satisfies:

$$(2.10) \quad \frac{\partial w}{\partial s} = -\left\langle \frac{\partial}{\partial t}, \vec{\nu} \right\rangle^{-1} (H - \mathcal{H})$$

in $M \times [0, S_0]$ and $w(p, 0) = 0$.

Proof. Since $|w| < a$ so that $\mathbf{X}(w(\phi(p, s), s), \phi(p, s))$ is well-defined. In local coordinates x^i of M in the domain and local coordinates y^i of the image of ϕ ,

$$\begin{aligned}\partial_s \mathbf{F} &= \frac{\partial y^j}{\partial s} \frac{\partial}{\partial y^j} + \left(\frac{\partial w}{\partial y^j} \frac{\partial y^j}{\partial s} + w_s \right) \frac{\partial}{\partial t} \\ &= \frac{\partial y^j}{\partial s} \left(\frac{\partial}{\partial y^j} + \frac{\partial w}{\partial y^j} \frac{\partial}{\partial t} \right) + w_s \frac{\partial}{\partial t}.\end{aligned}$$

Here $w_s = \frac{\partial w}{\partial s}$ as a function of in the form $w(p, s)$. For fixed s , $\phi(p, s)$ is a diffeomorphism of M and so $\mathbf{F}_* \left(\frac{\partial}{\partial x^i} \right)$ is tangential for all i . Since

$$\mathbf{F}_* \left(\frac{\partial}{\partial x^i} \right) = \frac{\partial y^k}{\partial x^i} \left(\frac{\partial}{\partial y^k} + \frac{\partial w}{\partial y^k} \frac{\partial}{\partial t} \right),$$

and $\mathbf{y} = \phi(\mathbf{x}, s)$ is a diffeomorphism, we have

$$(\partial_s \mathbf{F})^\perp = \left(w_s \frac{\partial}{\partial t} \right)^\perp$$

where \perp is the projection to the normal of immersion given by $\mathbf{F}^s = \mathbf{F}(\cdot, s)$.

So the first term in the above is tangential. Hence

$$w_s \left\langle \frac{\partial}{\partial t}, \vec{\nu} \right\rangle = \langle \partial_s \mathbf{F}, \vec{\nu} \rangle = -(H - \mathcal{H}).$$

From this the result follows because $-\langle \vec{\nu}, \frac{\partial}{\partial t} \rangle \geq 1$. \square

The following is also well-known.

Lemma 2.9. *Under Assumption 2.1, suppose w is a smooth function on $M \times [0, S_0]$ such that $|w| < a$, $w(p, 0) = 0$, and $w \in C^{2+\sigma, 1+\frac{\sigma}{2}}(M \times [0, S_0])$ such that $1 - |\nabla w|^2 \geq \lambda^{-2}$ uniformly for some constant $\lambda > 0$. Here ∇ is the gradient on the slice $\{t\} \times M$. Let $\tilde{\mathbf{F}}(p, s) = \mathbf{X}(p, w(p, s))$. Then the $\tilde{\mathbf{F}}(\cdot, s)$ is an immersion for all s . Suppose w satisfies (2.10) with mean curvature vector $\mathbf{H} = \vec{\nu}H$ with \mathcal{H} being bounded. Then one can find a smooth family of diffeomorphisms $\phi(p, s)$, $0 \leq s \leq S_0$ of M with $\phi(\cdot, 0)$ being the identity map such that*

$$\mathbf{F}(p, s) = \tilde{\mathbf{F}}(\phi(p, s), w(\phi(p, s), s))$$

is a smooth solution to (1.2).

Proof. Denote $\tilde{\mathbf{F}}(\cdot, s)$ by $\tilde{\mathbf{F}}^s(\cdot)$. Let y^i be local coordinates of M so that y^i, t are local coordinates of $N = (-a, a) \times M$. We may assume that y^i are quasi local coordinates. Then $\tilde{\mathbf{F}}$ is of the form

$$\tilde{\mathbf{F}}(\mathbf{y}, s) = (\mathbf{y}, w(\mathbf{y}, s)).$$

Let $e_i =: \tilde{\mathbf{F}}_*^s \left(\frac{\partial}{\partial y^i} \right) = \frac{\partial}{\partial y^i} + w_i \frac{\partial}{\partial t}$ where $w_i = \frac{\partial w}{\partial y^i}$. So $\langle e_i, e_j \rangle = g_{ij}(\mathbf{y}, t)|_{t=w} - w_i w_j$. Let $\xi_i, 1 \leq i \leq n$ be real numbers and let $\xi^i = g^{ik} \xi_k$. Then

$$\langle e_i, e_j \rangle \xi^i \xi^j = \xi^i \xi_i - \left(\sum_{i=1}^n w_i \xi^i \right)^2.$$

So

$$\xi^i \xi_i \geq \langle e_i, e_j \rangle \xi^i \xi^j \geq \xi^i \xi_i (1 - |\nabla w|^2) \geq \lambda^{-2} \xi^i \xi_i.$$

Hence the graph $\tilde{\mathbf{F}}^s$ is spacelike with induced metric uniformly equivalent to $g(\mathbf{y}, t)|_{t=w}$ at $\tilde{\mathbf{F}}^s(\mathbf{y})$, and hence is equivalent to $g(\mathbf{y}, 0)$ by Lemma 2.3. From this one can see that there is a smooth family of vector fields $V(\mathbf{y}, s)$ on M $s \in [0, S_0]$ such that $\tilde{\mathbf{F}}_*^s(V) = \left(\partial_s \tilde{\mathbf{F}}\right)^\top$ which is the tangential component of $\partial_s \tilde{\mathbf{F}}$ to the graph of $\tilde{\mathbf{F}}^s$. To be more precise, let $\vec{\nu}$ be the future timelike unit normal of the graph of $\tilde{\mathbf{F}}^s$,

$$\partial_s \tilde{\mathbf{F}} = w_s \frac{\partial}{\partial t} = \left(w_s \frac{\partial}{\partial t}\right)^\top - w_s \langle \vec{\nu}, \frac{\partial}{\partial t} \rangle \vec{\nu} = \left(w_s \frac{\partial}{\partial t}\right)^\top + (H - \mathcal{H}) \vec{\nu}$$

because w satisfies (2.10). Now $(w_s \frac{\partial}{\partial t})^\top(\mathbf{y}, s) = v^i(\mathbf{y}, s) e_i$ where e_i are tangent to the graph of $\tilde{\mathbf{F}}^s$ as before. Then $V = v^i \frac{\partial}{\partial y^i}$. It is easy to see that V is smooth in \mathbf{y}, s . Let $\phi(p, s)$ be the integral curve of $-V$ so that $\phi(\cdot, 0)$ is the identity map. We claim that $|V|_{g(0)}$ is uniformly bounded. If the claim is true, then ϕ can be defined on $M \times [0, S_0]$. In this case, let

$$\mathbf{F}(p, s) = \tilde{\mathbf{F}}(q, w(q, s))|_{q=\phi(p, s)}.$$

Then \mathbf{F} is smooth, $\mathbf{F}(p, 0) = \mathbf{X}_0(p)$ because $w(p, 0) = 0$. Moreover,

$$\begin{aligned} \partial_s \mathbf{F} &= \tilde{\mathbf{F}}_*^s(\partial_s \phi(p, s)) + (\partial_s \tilde{\mathbf{F}})^\top + (H - \mathcal{H}) \vec{\nu} \\ &= (H - \mathcal{H}) \vec{\nu}. \end{aligned}$$

The lemma follows provided we can prove the claim that $\|V\|_{g(0)}$ is bounded. To prove the claim, let e_i be as before,

$$\langle (\partial_s \tilde{\mathbf{F}})^\top, e_i \rangle = w_i w_s$$

which is uniformly bounded because $w \in C^{2+\sigma, 1+\frac{\sigma}{2}}(M \times [0, S_0])$, H is given by (2.3), $1 - |\nabla w|^2 \geq \lambda^{-2}$, and \mathcal{H} is uniformly bounded. Here we have also used Lemma 2.7. Since the metric $\langle e_i, e_j \rangle$ is uniformly equivalent to $g_{ij}(0)$, this implies the claim is true. \square

Hence in order to find short time existence of the prescribed mean curvature flow (1.2), it is sufficient to find w satisfying the conditions in the above lemma on $M \times [0, S_0]$ for some $S_0 > 0$. We will use the inverse theorem on C^1 maps on Banach spaces. We proceed as in [7] (see also [6]) using the argument outlined in [17]. Consider the Banach space

$$C_0^{2+\sigma, 1+\frac{\sigma}{2}}(M \times [0, S_0]) = \{f \in C^{2+\sigma, 1+\frac{\sigma}{2}}(M \times [0, S_0]) \mid f(\cdot, 0) = 0\}$$

for some $0 < \sigma < 1$ and S_0 will be chosen later. If there is no confusion, we will drop $M \times [0, S_0]$ in the definition of Hölder spaces.

Let \mathcal{B} be the open set in $C_0^{2+\sigma, 1+\frac{\sigma}{2}}(M \times [0, S_0])$ consisting functions f such that

$$(2.11) \quad \sup_{M \times [0, S_0]} |f| < a \text{ and } \inf_{M \times [0, S_0]} (1 - |\nabla f|^2) > \frac{1}{2}$$

where $|\nabla f|^2$ is the norm of the gradient with respect to $g(t)$. More precisely in local coordinates x^i of M ,

$$|\nabla f|^2(\mathbf{x}, s) = f^i f_i$$

where $f_i = \partial_i f = \frac{\partial}{\partial x^i} f$ and $f^i(\mathbf{x}, s) = g^{ij}(\mathbf{x}, t)|_{t=f(\mathbf{x}, s)} f_j(\mathbf{x}, s)$.

We want to define a suitable map \mathcal{F} from \mathcal{B} to $C^{\sigma, \frac{\sigma}{2}}(M \times [0, S_0])$. Let $w \in \mathcal{B}$ consider the graph of w in N which is identified with $(-a, a) \times M$ with metric $G = -dt^2 + g(t)$. The graph of $w(\cdot, s)$ is given by $(\mathbf{x}, w(\mathbf{x}, s)) \in (-a, a) \times M$, $\mathbf{x} \in M$. This is well defined because $|w| < a$. The graph is spacelike because $1 - |\nabla w|^2 > \frac{1}{2}$. For fixed s , the future pointing unit normal is given by

$$\vec{\nu}(w) = \frac{\frac{\partial}{\partial t} + \nabla w}{(1 - |\nabla w|^2)^{\frac{1}{2}}}.$$

The mean curvature vector is $\mathbf{H} = H\vec{\nu}$. Let \mathcal{H} be a smooth function in $(-a, a) \times M$. Define

$$(2.12) \quad \mathcal{F}(w) = \frac{\partial w}{\partial s} + \left\langle \frac{\partial}{\partial t}, \vec{\nu}(w) \right\rangle^{-1} (H - \mathcal{H}) = \frac{\partial w}{\partial s} - (1 - |\nabla w|^2)^{\frac{1}{2}} (H - \mathcal{H})$$

for $w \in \mathcal{B}$. In the following, it is more convenient to use the quasi local coordinates to express the map. By (2.3), we have

$$(2.13) \quad \begin{aligned} \mathcal{F}(w) &= \frac{\partial w}{\partial s} - \left(g^{ij} + \frac{w^i w^j}{1 - |\nabla w|^2} \right) w_{;ij} - \frac{w_i w_j}{(1 - |\nabla w|^2)^{\frac{1}{2}}} \partial_t g^{ij} \\ &\quad - \frac{1}{2} g^{ij} \partial_t g_{ij} + (1 - |\nabla w|^2)^{\frac{1}{2}} \mathcal{H} \\ &= \frac{\partial w}{\partial s} - \left(g^{ij} + \frac{w^i w^j}{1 - |\nabla w|^2} \right) (w_{ij} - \Gamma_{ij}^k w_k) - \frac{w_i w_j}{(1 - |\nabla w|^2)^{\frac{1}{2}}} \partial_t g^{ij} \\ &\quad - \frac{1}{2} g^{ij} \partial_t g_{ij} + (1 - |\nabla w|^2)^{\frac{1}{2}} \mathcal{H}. \end{aligned}$$

For any function $f(t, \mathbf{x})$ in the above it is understood the function will be evaluated at $t = w$. Here Γ_{ij}^k is the Christoffel symbols of g with respect to the quasi local coordinates.

Lemma 2.10. *Suppose $\|\overline{\nabla}^k \mathcal{H}\|_{\Omega}$ is uniformly bounded for $k = 0, 1$, then $\mathcal{F}(w) \in C^{\sigma, \frac{\sigma}{2}}$ for $w \in \mathcal{B}$.*

Proof. Let $\xi_p : D(r) \rightarrow B_p(r)$ be quasi local coordinate neighborhood. By Lemma 2.7 all the derivatives of the components of $\xi_p^*(g(t))$ and its inverse with respect to t and the quasi local coordinates x^i are uniformly bounded independent of p and for $|t| < a$. Moreover, the $C^{2+\sigma, 1+\frac{\sigma}{2}}$ norm of $\xi^*(w)$ in $D(r) \times [0, S_0]$ is uniformly bounded independent of p . In order to prove

the lemma, it is sufficient to prove that the $C^{\sigma, \frac{\sigma}{2}}$ norm in $D(r) \times [0, S_0]$ of each term of the last expression in (2.13) is uniformly bounded, where w is understood to be $\xi_p^*(w)$ and g is understood to be $\xi_p^*(g)$ etc. By the assumption on w , the $C^{2\sigma, \frac{\sigma}{2}}$ norms of w_i, w_{ij} are uniformly bounded. On the other hand, since the first derivative of g_{ij} in x^i, t are uniformly bounded and since w_i, w_s are uniformly bounded, we conclude that the $C^{\sigma, \frac{\sigma}{2}}$ norms of $g_{ij}(w(\mathbf{x}, s), \mathbf{x})$ are uniformly bounded. Similarly, the $C^{\sigma, \frac{\sigma}{2}}$ norms of $g^{ij}, \partial_t g_{ij}, \partial_t g^{ij}, \Gamma_{ij}^k, g$ are uniformly bounded evaluated at $t = w(\mathbf{x}, s)$. By the assumption on w and Lemma 2.7, $w^i = g^{ik} w_k$ has uniformly bounded $C^{\sigma, \frac{\sigma}{2}}$ norm. Since $1 - |\nabla w|^2 > \frac{1}{2}$, one can see that for any c , the $C^{2\sigma, \frac{\sigma}{2}}$ norm of $(1 - |\nabla w|^2)^c = (1 - w^i w_i)^c$ are uniformly bounded. Since $\|\overline{\nabla}^k \mathcal{H}\|_{\Omega}$ is uniformly bounded for $k = 0, 1$, $\mathcal{H}(w(\mathbf{x}, s), \mathbf{x})$ is in $C^{\sigma, \frac{\sigma}{2}}$ by Lemmas 2.3 and 2.7. From these, we conclude that the lemma is true. \square

Next we want to compute the differential of \mathcal{F} . Let $v(\mathbf{x}, s)$ be a function on $M \times [0, S_0]$ with $|v| < a$. For any family of tensors $T(t, \mathbf{x})$ in \mathbf{x} for $(t, \mathbf{x}) \in (-a, a) \times M$, denote

$${}^v T(\mathbf{x}, s) =: T(v(\mathbf{x}, s), \mathbf{x}).$$

For examples:

$$\begin{cases} v^i = {}^v g^{ij} v_j \\ v_{;ij} = v_{ij} - {}^v \Gamma_{ij}^k v_k. \end{cases}$$

We will compute $D\mathcal{F}$ at v in quasi local coordinates x^i, s in $M \times [0, S_0]$.

Lemma 2.11. *In addition to Assumption 2.1, suppose $\|\overline{\nabla}^k \mathcal{H}\|_{\Omega}$ are uniformly bounded for all $k \geq 0$. Then in the quasi local coordinates, for any $v \in \mathcal{B}$, the differential $D\mathcal{F}_v$ of \mathcal{F} at v is given by:*

$$(2.14) \quad D\mathcal{F}_v(\phi) = \phi_s - {}^v a^{ij} \phi_{ij} + {}^v b^k \phi_k + {}^v c \phi$$

where

$$(2.15) \quad \begin{cases} {}^v a^{ij} = {}^v g^{ij} + (1 - |\nabla v|^2)^{-1} v^i v^j; \\ {}^v b^k = -\frac{2({}^v g^{ki} v^j (1 - |\nabla v|^2) + v^i v^j v^k)(v_{ij} - {}^v \Gamma_{ij}^m v_m)}{(1 - |\nabla v|^2)^2} + \left({}^v g^{ij} + \frac{v^i v^j}{1 - |\nabla v|^2} \right) {}^v \Gamma_{ij}^k \\ 2 - \left[\frac{{}^v \partial_t g^{ij} v_i v_j v^k + (1 - |\nabla v|^2) {}^v \partial_t g^{ik} v_i + v^k ((1 - |\nabla v|^2) {}^v \mathcal{H})}{(1 - |\nabla v|^2)^{\frac{3}{2}}} \right] \\ {}^v c = - \left[{}^v \partial_t g^{ij} + \frac{2 {}^v \partial_t g^{ik} v_k}{1 - |\nabla v|^2} + \frac{v^i v^j {}^v \partial_t g^{lm} v_l v_m}{(1 - |\nabla v|^2)^2} \right] (v_{ij} - {}^v \Gamma_{ij}^k v_k) \\ + \left({}^v g^{ij} + \frac{v^i v^j}{1 - |\nabla v|^2} \right) {}^v \partial_t \Gamma_{ij}^k v_k - \left[\frac{{}^v \partial_t g^{lm} v_l v_m {}^v \partial_t g^{ij}}{(1 - |\nabla v|^2)^{\frac{3}{2}}} + \frac{{}^v \partial_t^2 g^{ij}}{(1 - |\nabla v|^2)^{\frac{1}{2}}} \right] v_i v_j \\ - \frac{1}{2} ({}^v \partial_t g^{ij} {}^v \partial_t g_{ij} + {}^v g^{ij} {}^v \partial_t^2 g_{ij}) - \frac{{}^v \partial_t g^{lm} v_l v_m {}^v \mathcal{H}}{(1 - |\nabla v|^2)^{\frac{1}{2}}} + (1 - |\nabla v|^2)^{\frac{1}{2}} {}^v \partial_t \mathcal{H} \end{cases}$$

Moreover, \mathcal{F} is C^1 in \mathcal{B} .

Proof. Fix $v \in \mathcal{B}$. Let $\phi \in C_0^{2+\alpha, 1+\frac{\alpha}{2}}$. Here and below the Höder spaces are defined on $M \times [0, S_0]$. Then for $|\epsilon|$ small, $v + \epsilon\phi \in C_0^{2+\alpha, 1+\frac{\alpha}{2}}$ and $\mathcal{F}(v + \epsilon\phi)$ is in $C^{\alpha, \frac{\alpha}{2}}$. In fact, for $|\epsilon|$ small enough, $v_\epsilon =: v + \epsilon\phi \in \mathcal{B}$. Then at $\epsilon = 0$,

$$\left\{ \begin{array}{l} \frac{\partial}{\partial \epsilon} (v + \epsilon\phi)_s = \phi_s \\ \frac{\partial}{\partial \epsilon} (v_\epsilon g^{ij}) = \phi^v \partial_t g^{ij} \\ \frac{\partial}{\partial \epsilon} (v_\epsilon \partial_t g^{ij}) = \phi^v \partial_t^2 g^{ij} \\ \frac{\partial}{\partial \epsilon} v_\epsilon^i = \phi^v \partial_t g^{ik} v_k + \phi_k^v g^{ki} \\ \frac{\partial}{\partial \epsilon} v_\epsilon \mathcal{H} = \phi^v \partial_t \mathcal{H} \\ \frac{\partial}{\partial \epsilon} v_\epsilon \Gamma_{ij}^k = \phi^v \partial_t \Gamma_{ij}^k. \end{array} \right.$$

In the following f' means $\frac{\partial f}{\partial \epsilon}|_{\epsilon=0}$. We have:

$$\begin{aligned} & \left[\left(g^{ij} + \frac{v^i v^j}{1 - |\nabla v|^2} \right) (v_{ij} - \Gamma_{ij}^k v_k) \right]' \\ &= \left[\phi^v \partial_t g^{ij} + \frac{2(\phi^v \partial_t g^{ik} v_k + \phi_k^v g^{ki} v_j)}{1 - |\nabla v|^2} + \frac{v^i v^j (\phi^v \partial_t g^{lm} v_l v_m + 2\phi_k v^k)}{(1 - |\nabla v|^2)^2} \right] (v_{ij} - v \Gamma_{ij}^k v_k) \\ &+ \left(v g^{ij} + \frac{v^i v^j}{1 - |\nabla v|^2} \right) (\phi_{ij} - v \Gamma_{ij}^k \phi_k - \phi^v \partial_t \Gamma_{ij}^k v_k) \\ & \left(\frac{1}{(1 - |\nabla v|^2)^{\frac{1}{2}}} \partial_t g^{ij} v_i v_j \right)' = \frac{\phi^v \partial_t g^{lm} v_l v_m + 2\phi_k v^k}{(1 - |\nabla v|^2)^{\frac{3}{2}}} v \partial_t g^{ij} v_i v_j \\ & \quad + \frac{1}{(1 - |\nabla v|^2)^{\frac{1}{2}}} (\phi^v \partial_t^2 g^{ij} v_i v_j + 2^v \partial_t g^{ik} v_i \phi_k) \\ & \left(\frac{1}{2} g^{ij} \partial_t g_{ij} \right)' = \frac{1}{2} \phi^v (\partial_t g^{ij} v \partial_t g_{ij} + v g^{ij} v \partial_t^2 g_{ij}) \\ & \left((1 - |\nabla v|^2)^{\frac{1}{2}} \mathcal{H} \right)' = - \frac{\phi^v \partial_t g^{lm} v_l v_m + 2\phi_k v^k}{(1 - |\nabla v|^2)^{\frac{1}{2}}} v \mathcal{H} + \phi (1 - |\nabla v|^2)^{\frac{1}{2}} v \partial_t \mathcal{H}. \end{aligned}$$

Let $D\mathcal{F}_v(\phi) = \frac{\partial}{\partial \epsilon} \mathcal{F}(v + \epsilon\phi)|_{\epsilon=0}$. Then

$$D\mathcal{F}_v(\phi) = \phi_s - v a^{ij} \phi_{ij} + v b^k \phi_k + v c \phi,$$

where $v a^{ij}$, $v b^k$, $v c$ are as in (2.15). Since $\| |\overline{\nabla}^k \mathcal{H} \|_{\Omega}$ is uniformly bounded for all k , together with Lemma 2.7, all the derivatives of g_{ij} and \mathcal{H} with respect to t and the quasi local coordinates x^i are uniformly bounded. Here we identify g , \mathcal{H} etc to their pull back to $D(r)$. From this and the fact that $v \in C_0^{2+\sigma, 1+\frac{\sigma}{2}}$, $D\mathcal{F}_v$ is a bounded linear operator from $C_0^{2+\sigma, 1+\frac{\sigma}{2}}$ to $C^{\sigma, \frac{\sigma}{2}}$. Moreover, by the structure of $v a^{ij}$, $v b^k$, $v c$, we have for $u, v \in \mathcal{B}$, the $C^{\sigma, \frac{\sigma}{2}}(D(r) \times [0, S_0])$ norms

of ${}^u a^{ij} - {}^v a^{ij}$, ${}^u b^k - {}^v b^k$, and ${}^u c - {}^v c$ are bounded by $C\|u - v\|_{C_0^{2+\sigma, 1+\frac{\sigma}{2}}(M \times [0, S_0])}$ for some constant C independent of the coordinate neighborhoods. Here C may depend on the upper bound of the norms of u, v in $C_0^{2+\sigma, 1+\frac{\sigma}{2}}(M \times [0, S_0])$. Hence the norms of the operators $D\mathcal{F}_u, D\mathcal{F}_v$ satisfy:

$$\|D\mathcal{F}_u - D\mathcal{F}_v\| \leq C\|u - v\|_{C_0^{2+\sigma, 1+\frac{\sigma}{2}}(M \times [0, S_0])}.$$

This implies $D\mathcal{F}_v$ is continuous in v . Here we have used the fact that $(1 - |\nabla w|^2) > \frac{1}{2}$ for all $w \in \mathcal{B}$. We claim that $D\mathcal{F}_v$ is the differential of \mathcal{F} at v . Let $v \in \mathcal{B}$ be fixed and $h \in \mathcal{B}$ so that $\|h\|_{C_0^{2+\sigma, 1+\frac{\sigma}{2}}(M \times [0, S_0])}$ (which will be denoted simply by $\|h\|$ in the following) is small enough. Then

$$\begin{aligned} \mathcal{F}(v+h) - \mathcal{F}(v) - D\mathcal{F}_v(h) &= \int_0^1 \frac{d}{d\epsilon} \mathcal{F}(v + \epsilon h) d\epsilon - D\mathcal{F}_v(h) \\ &= \int_0^1 (D\mathcal{F}_{v+\epsilon h}(h) - D\mathcal{F}_v(h)) d\epsilon \end{aligned}$$

Hence

$$\|\mathcal{F}(v+h) - \mathcal{F}(v) - D\mathcal{F}_v(h)\|_{C^{\sigma, \frac{\sigma}{2}}} \leq C\|h\|^2.$$

So $D\mathcal{F}_v$ is the differential of \mathcal{F} at v . This completes the proof of the lemma. \square

Lemma 2.12. *In addition to Assumption 2.1, suppose $\|\overline{\nabla}^k \mathcal{H}\|_{\Omega}$ are uniformly bounded. For any $v \in \mathcal{B}$, $D\mathcal{F}_v$ is a bijection from $C_0^{2+\sigma, 1+\frac{\sigma}{2}}(M \times [0, s_0])$ onto $C^{\sigma, \frac{\sigma}{2}}(M \times [0, s_0])$.*

Proof. Again, denote $C_0^{2+\sigma, 1+\frac{\sigma}{2}}(M \times [0, s_0])$ by $C_0^{2+\sigma, 1+\frac{\sigma}{2}}$ etc. To prove that $D\mathcal{F}_v$ is injective, let $\phi \in C_0^{2+\sigma, 1+\frac{\sigma}{2}}$ such that $D\mathcal{F}(\phi) = 0$. Since $(M, g(0))$ has bounded curvature, we can find a smooth function η on M such that α is uniformly equivalent to the distance function of $g(0)$ with respect to a fixed point such that η has bounded gradient and bounded Hessian with respect to $g(0)$, see [26, 27]. Hence η_i, η_{ij} with respect to any quasi local coordinates are uniformly bounded by a constant C_1 independent of the coordinate neighborhood. Here we denote the pull back of a function f by f again. Using η one can proceed in a standard way. Namely in any quasi local coordinate neighborhood, we have

$$|D\mathcal{F}_v(\exp(\alpha))| \leq C_2 \exp(\alpha)$$

where C_2 is independent of the quasi local coordinate neighborhood. Let $\Theta = \exp(C_3 s + \alpha)$ for some C_3 to be determined later. Then for $\epsilon > 0$, $\phi - \epsilon\Theta \leq 0$ at $s = 0$. Suppose it is positive somewhere, then there is $p \in M$ and $0 < s \leq s_0$ such that $\phi - \epsilon\Theta$ attains maximum at (p, s) . In the quasi

coordinate neighborhood centered at p , we have, $\phi_i = \epsilon\Theta_i$ and

$$\begin{aligned} 0 &\leq (\phi_s - \epsilon\Theta)_s - {}^v a^{ij}(\phi - \epsilon\Theta)_{ij} \\ &= - {}^v c(\phi - \epsilon\Theta) - D\mathcal{F}_v(\epsilon\Theta) \\ &\leq \epsilon\Theta(-C_3 + C_2) \\ &= -\epsilon C_2\Theta \end{aligned}$$

if $C_3 = 2C_2$. This is a contradiction. Hence $\phi \leq \epsilon\Theta$. Let $\epsilon \rightarrow 0$ we have $\phi \leq 0$. Similarly, $-\phi \leq 0$. So $\phi = 0$ and $D\mathcal{F}_v$ is injective.

To prove that \mathcal{F}_v is surjective, let $f \in C^{\sigma, \frac{\sigma}{2}}$ and let Ω_ℓ be compact domains in M with smooth boundary such that $\Omega_\ell \Subset \Omega_{\ell+1}$ which exhaust M . Let η_ℓ be smooth function so that $0 \leq \eta_\ell \leq 1$, $\eta_\ell = 0$ outside $\Omega_{\ell+1}$ and $\eta_\ell = 1$ on Ω_ℓ . Then one can find a solution ϕ_ℓ of the following initial-boundary value problem:

$$\begin{cases} D\mathcal{F}_v(\phi_\ell) = \eta_\ell f, & \text{in } \Omega_{\ell+1} \times [0, s_0]; \\ \phi_\ell = 0, & \text{at } s = 0 \text{ and } \partial\Omega_{\ell+1} \times [0, s_0]. \end{cases}$$

The solution ϕ_ℓ exists by [11, Theorem 4, Chapter 3], because $\eta_\ell f = 0$ on $\partial\Omega_{\ell+1}$, so that the compatibility condition $D\mathcal{F}_v(\psi) = \eta_\ell f$ at $\partial\Omega_{\ell+1} \times \{0\}$, where $\psi \equiv 0$, is satisfied. Moreover, the pull back of ϕ_ℓ is in $C^{2+\sigma, 1+\frac{\sigma}{2}}(D(r) \times [0, s_0])$ in any quasi local coordinate neighborhood contained in Ω_ℓ . Since ${}^v c$ is uniformly bounded independent of quasi local coordinates, by the maximum principle [24, Theorem 2.10] we can see ϕ_ℓ is uniformly bounded by a constant independent of ℓ . Standard Schauder estimates (see [11, Theorem 4, Chapter 4]) show that we can find ϕ_ℓ satisfying $D\mathcal{F}_v(\phi_\ell) = f$ in $\Omega_\ell \times [0, s_0]$ such that for any $p \in \Omega_\ell$ with $B_p(r) \Subset \Omega_\ell$, for the quasi local coordinates neighborhood $\xi_p : D(r) \rightarrow B_p(r)$, $\|\phi_\ell\|_{C^{2+\sigma, 1+\frac{\sigma}{2}}(D(\frac{r}{2}) \times [0, s_0])}$ is bounded by a constant independent of p and ℓ . Here we denote the pull back of ϕ_ℓ also by ϕ_ℓ . Passing to a subsequence, we can find ϕ such that $D\mathcal{F}_v(\phi) = f$ on $M \times [0, s_0]$ with $\phi = 0$ at $s = 0$. Moreover, $\|\phi\|_{C^{2+\sigma, 1+\frac{\sigma}{2}}(D(\frac{r}{2}) \times [0, s_0])}$ uniformly bounded by a constant in any quasi local coordinate neighborhood $\xi_p : D(r) \rightarrow B_p(r)$. From this we see that $\phi \in C^{2+\sigma, 1+\frac{\sigma}{2}}(M \times [0, s_0])$ by Lemma 2.6. \square

Now we are ready to prove the following short time existence result.

Proof of Theorem 1.1. By Lemmas 2.2, 2.3, we may assume $a > 0$ is small enough so that \mathbf{X} is an immersion. In order to construct the flow, without loss of generality we may identify N with $(-a, a) \times M$ with metric $G = -dt^2 + g(t)$. Moreover, we can introduce a family of quasi local coordinate neighborhoods as described by Lemma 2.5 so that $g(t)$ satisfy the conclusions of Lemma 2.7.

Let $v_0(\mathbf{x}, s) = sH^o(\mathbf{x}, 0) - s\mathcal{H}(\mathbf{x}, 0)$ defined on $M \times [0, s_0]$, where H^o is the mean curvature of the slice $t = 0$ in $(-a, a) \times M$. By Lemmas 2.7, 2.2, 2.1 and the fact that $\|\overline{\nabla}^k \mathcal{H}\|_N$ is finite for $k = 0, 1$, we can choose $s_0 > 0$ small enough so that $\sup_{M \times [0, s_0]} |v_0| < a$ and $\inf_{M \times [0, s_0]} (1 - |\nabla v_0|^2) > \frac{1}{2}$. On the other hand,

fixed $0 < \sigma < 1$, one can define $C_0^{2\sigma, \frac{\sigma}{2}}(M \times [0, s_0])$ and $C^{\sigma, \frac{\sigma}{2}}(M \times [0, s_0])$. As before let

$$\mathcal{B} = \{v \in C_0^{2+\sigma, 1+\frac{\sigma}{2}}(M \times [0, s_0]) \mid \sup_{M \times [0, s_0]} |v| < a, \inf_{M \times [0, s_0]} (1 - |\nabla v|^2) > \frac{1}{2}\}.$$

Hence $v_0 \in \mathcal{B}$. In fact, by Lemma 2.7 and the fact that $\|\overline{\nabla}^k \mathcal{H}\|_N$ are finite for any k , we have

$$(2.16) \quad |\xi_p^*(v_0)(\mathbf{x}, s) - \xi_p^*(v_0)(\mathbf{x}', s')| \leq C_1 (|\mathbf{x} - \mathbf{x}'| + |s - s'|)$$

in $D(r) \times [0, s_0]$, for any quasi local coordinate neighborhood $\xi_p^* : D(r) \rightarrow B_p(r)$. Here C_1 is a constant independent of p .

Let

$$\mathcal{F}(v) = \frac{\partial v}{\partial s} + \langle \frac{\partial}{\partial t}, \vec{\nu} \rangle^{-1} (H - \mathcal{H})$$

where $\vec{\nu}$ is the future direct unit normal of the graph of v : $(\mathbf{x}, v(\mathbf{x}, s)) \in (-a, a) \times M$ and H is the mean curvature. Then \mathcal{F} maps \mathcal{B} to $C^{\sigma, \frac{\sigma}{2}}(M \times [0, s_0])$ which is C^1 and $D\mathcal{F}_v$ is bijective by Lemmas 2.10 and 2.12. Hence \mathcal{F} maps a neighborhood of v_0 onto a neighborhood of $f_0 =: \mathcal{F}(v_0)$. By the definition of v_0 and (2.13), $f_0(\mathbf{x}, 0) = 0$ because $H^o = \frac{1}{2}g^{ij}\partial_t g_{ij}$.

For $0 < \lambda < s_0$, define

$$f_\lambda(\mathbf{x}, s) = \begin{cases} 0 & \text{if } 0 \leq s \leq \lambda. \\ f_0(\mathbf{x}, s - \lambda) & \text{if } \lambda < s \leq s_0. \end{cases}$$

Claim 2.1. $\|f_0 - f_\lambda\|_{C^{\sigma, \frac{\sigma}{2}}(M \times [0, s_0])} \leq C\lambda^{1-\sigma}$ for some C independent of λ .

If the claim is true, then $f_\lambda \in C^{\sigma, \frac{\sigma}{2}}(M \times [0, s_0])$ and by the inverse function theorem, for $\lambda > 0$ small enough, we can find w such that

$$\mathcal{F}(w) = f_\lambda.$$

In particular, w is a solution of (2.10) on $M_0 \times [0, \lambda]$, $w \in \mathcal{B}$.

To prove **Claim 2.1**, let $\xi_p : D(r) \rightarrow B_p(r)$ be a quasi local coordinate neighborhood. Since $f_0 = 0$, at $s = 0$, by (2.16), that there is $C_2 > 0$ independent of p , λ such that if $x, x' \in D(r)$ and $s, s' \in [0, s_0]$, we have

$$(2.17) \quad |\xi_p^* f_\lambda(\mathbf{x}, s) - \xi_p^* f_\lambda(\mathbf{x}', s')| \leq C_2 (|\mathbf{x} - \mathbf{x}'| + |s - s'|).$$

Let $\eta = f_0 - f_\lambda$. Then

$$\eta(\mathbf{x}, s) = \begin{cases} f_0(\mathbf{x}, s) & \text{if } 0 \leq s \leq \lambda. \\ f_0(\mathbf{x}, s) - f_0(\mathbf{x}, s - \lambda) & \text{if } \lambda < s \leq s_0. \end{cases}$$

There is a constant $C_3 > C_2$ independent of p such that $|\eta| \leq C_3\lambda$ by (2.16) and the fact that $f_0 = 0$ at $s = 0$. For $\mathbf{x}, \mathbf{x}' \in D(r)$, $s, s' \in [0, s_0]$.

Case 1: $s, s' \leq \lambda$. Then

$$|\xi_p^* \eta(\mathbf{x}, s) - \xi_p^* \eta(\mathbf{x}', s')| = |\xi_p^* f_0(\mathbf{x}, s) - \xi_p^* f_0(\mathbf{x}', s')| \leq C_3 \min\{\lambda, |x - x'| + |s - s'|\}$$

because $f_0(\mathbf{x}, 0) = 0$ and $s, s' \leq \lambda$. If $|\mathbf{x} - \mathbf{x}'| \geq \lambda$, then

$$|\xi_p^* \eta(\mathbf{x}, s) - \xi_p^* \eta(\mathbf{x}', s')| \leq C_3 \lambda^{1-\sigma} \lambda^\sigma \leq C_3 \lambda^{1-\sigma} (|\mathbf{x} - \mathbf{x}'|^\sigma + |s - s'|^{\frac{\sigma}{2}}).$$

If $|\mathbf{x} - \mathbf{x}'| \leq \lambda$, then by (2.16)

$$\begin{aligned} |\xi_p^* \eta(\mathbf{x}, s) - \xi_p^* \eta(\mathbf{x}', s')| &\leq C_2 (|\mathbf{x} - \mathbf{x}'| + |s - s'|) \\ &= C_2 (|\mathbf{x} - \mathbf{x}'|^{1-\sigma} |\mathbf{x} - \mathbf{x}'|^\sigma + |s - s'|^{1-\frac{\sigma}{2}} |s - s'|^{\frac{\sigma}{2}}) \\ &\leq C_4 \lambda^{1-\sigma} (|\mathbf{x} - \mathbf{x}'|^\sigma + |s - s'|^{\frac{\sigma}{2}}). \end{aligned}$$

for some C_4 independent of p .

Case 2: $s, s' \geq \lambda$.

$$\begin{aligned} |\xi_p^* \eta(\mathbf{x}, s) - \xi_p^* \eta(\mathbf{x}', s')| &= |\xi_p^* f_0(\mathbf{x}, s) - \xi_p^* f_0(\mathbf{x}, s - \lambda) - \xi_p^* f_0(\mathbf{x}', s') + \xi_p^* f_0(\mathbf{x}', s' - \lambda)| \\ &\leq C_3 \min\{\lambda, |\mathbf{x} - \mathbf{x}'| + |s - s'|\} \end{aligned}$$

As before, we have

$$|\xi_p^* \eta(\mathbf{x}, s) - \xi_p^* \eta(\mathbf{x}', s')| \leq C_4 \lambda^{1-\sigma} (|\mathbf{x} - \mathbf{x}'|^\sigma + |s - s'|^{\frac{\sigma}{2}}).$$

Case 3: $s \geq \lambda \geq s'$, then

$$\begin{aligned} |\xi_p^* \eta(\mathbf{x}, s) - \xi_p^* \eta(\mathbf{x}', s')| &\leq |\xi_p^* \eta(\mathbf{x}, s) - \xi_p^* \eta(\mathbf{x}, \lambda)| + |\xi_p^* \eta(\mathbf{x}, \lambda) - \xi_p^* \eta(\mathbf{x}', s')| \\ &\leq 2C_4 \lambda^{1-\sigma} (|s - \lambda|^{\frac{\sigma}{2}} + |\mathbf{x} - \mathbf{x}'|^\sigma + |s' - \lambda|^{\frac{\sigma}{2}}) \\ &\leq 3C_4 \lambda^{1-\sigma} (|\mathbf{x} - \mathbf{x}'|^\sigma + |s - s'|^{\frac{\sigma}{2}}) \end{aligned}$$

because $s \geq \lambda \geq s'$. Hence the claim is true.

To conclude, we can find a solution w to (2.10) on $M \times [0, \epsilon]$ for some $\epsilon > 0$, so that $w \in \mathcal{B}$. By differentiate the equation (2.13) with respect to x^k , by Schauder estimates, one can conclude that $\xi_p^*(w)_i$ the $C^{2+\sigma, 1+\frac{\sigma}{2}}(D(r') \times [0, \epsilon])$ are uniformly bounded for $r' < r$, in any quasi local coordinate neighborhood. Here we have used the fact that all derivatives of $g_{ij}(t)$ and all derivatives of \mathcal{H} are bounded and the fact that $1 - |\nabla w|^2 > \frac{1}{2}$. Continue in this way, we see that all the derivatives of $\xi_p^*(w)$ with respect to \mathbf{x} are bounded. Using the equation (2.13) again, we conclude that all the derivatives of $\xi^*(w_s)$ with respect to \mathbf{x} are bounded. Differentiate with respect to s , one can finally conclude that all the derivatives of $\xi_p^*(w)$ in $D(\frac{r}{2})$ are uniformly bounded independent of p . By Lemma 2.9, we conclude that the short time existence of the prescribed mean curvature flow. Moreover, \mathbf{F} is smooth and the second fundamental form of $\mathcal{F}(\cdot, s)(M)$ together with its covariant derivatives are uniformly bounded by the bounds of derivatives of w and the bounds of derivatives of $g(t)$. Since $1 - |\nabla w|^2 > \frac{1}{2}$, we have

$$-\left\langle \frac{\partial}{\partial t}, \vec{v} \right\rangle \leq C$$

for some constant C in $M \times [0, \lambda]$. Hence by [3]

$$-\langle \mathbf{T}, \vec{v} \rangle \leq 2 \langle \mathbf{T}, \frac{\partial}{\partial t} \rangle \langle \frac{\partial}{\partial t}, \vec{v} \rangle \leq C'$$

for some constant C' in $M \times [0, \lambda]$. Since $1 - |\nabla w|^2 > \frac{1}{2}$, by Lemma 2.7, the metric pulled back by $\mathcal{F}(\cdot, s)$ is equivalent to $g(0)$ and is complete. Hence the theorem is proved. \square

We would like to discuss the condition (i) in Theorem 1.1. If N is time like geodesically complete, then (i) is satisfied automatically. But this is too strong. On the other hand, we have the following.

Lemma 2.13. *Let (N, G) be a Lorentz manifold with time function τ as before. Suppose there is a smooth function $\rho > 0$ satisfying condition (1.4). Let $p \in \Omega_{\tau_1, \tau_2} \subset N$ and let v be a future time like unit vector at p . Assume $\|\|\bar{\nabla} \mathbf{T}\|\| \leq c$ in Ω_{τ_1, τ_2} . Suppose either $\|\|1/\alpha\|\|_{0, \Omega_{\tau_1, \tau_2}} \leq b$, or $\|\|\bar{\nabla} \log \alpha\|\|_{0, \Omega_{\tau_1, \tau_2}} \leq b$. Then the time like geodesic $\gamma(t)$ with $\gamma(0) = p, \gamma'(0) = v$ can be defined on $(-t_0, t_0)$ for some t_0 depending only on $b, -\langle v, \mathbf{T} \rangle, \tau_2 - \tau(p), \tau(p) - \tau_1$ and upper bound of $1/(\alpha(p))$ so that $\gamma(t) \in \Omega_{\tau_1, \tau_2}$.*

Proof. Let $a > 0$ be given by Lemma 2.2 depending only on $c, \kappa_0 = -\langle v, \mathbf{T} \rangle$. Suppose we can prove the following a priori bound. There is a constant C depending only on the quantities mentioned in the lemma such that whenever $\gamma(t) \subset \Omega_{\tau_1, \tau_2}, t \in (-t', t')$ with $t' \leq a$, then $\alpha(\gamma(t)) \geq C^{-1}$. Then the lemma is true with

$$t_0 = \min\{a, (\tau(p) - \tau_1)/(2C\kappa_0), (\tau_2 - \tau(p))/(2C\kappa_0)\}.$$

In fact if t_1 is the supremum of t so that $\gamma(t)$ is defined and $\gamma(-t, t) \subset \Omega_{\tau_1, \tau_2}$. If $t_1 < t_0$, by Lemma 2.2, $\kappa =: -\langle \gamma'(t), \mathbf{T} \rangle \leq 2\kappa_0$.

$$\tau(\gamma(t)) - \tau(p) = \int_0^t \langle \gamma', \bar{\nabla} \tau \rangle d\sigma = \int_0^t -\frac{1}{\alpha} \langle \gamma', \mathbf{T} \rangle d\sigma,$$

and

$$|\tau(\gamma(t)) - \tau(p)| \leq 2C\kappa_0|t|.$$

Hence

(2.18)

$$\tau_1 + \epsilon \leq (\tau(p) - \tau_1) - 2C\kappa_0|t| + \tau_1 \leq \tau(\gamma(t)) \leq -(\tau_2 - \tau(p)) + 2C\kappa_0|t| + \tau_2 \leq \tau_2 - \epsilon$$

for all t for some $\epsilon > 0$. On the other hand,

$$|\rho(\gamma(t)) - \rho(p)| = \left| \int_0^t \langle \gamma', \bar{\nabla} \rho \rangle d\sigma \right| \leq C_1|t|.$$

for some constant C_1 depending only on $\kappa_0, \|\|\bar{\nabla} \rho\|\|_{0, \Omega_{\tau_1, \tau_2}}$. In particular,

(2.19)

$$\rho(\gamma(t)) \leq C_2$$

on $(-t_1, t_1)$. For $t, t' \uparrow t_1$,

$$d_{G_E}(\gamma(t), \gamma(t')) \leq \left| \int_t^{t'} \|\|\gamma'\|\| d\sigma \right| \leq 2\kappa_0|t - t'|.$$

By the compactness of the set $\overline{\Omega}_{\tau_1, \tau_2} \cap \{\rho \leq C_2\}$, we conclude that $\gamma(t)$ can be extended up to $-t_1, t_1$. By (2.18), we must have $t_1 \geq t_0$.

It remains to obtain the a priori estimate mentioned above. In case $|||1/\alpha|||_{0, \Omega_{\tau_1, \tau_2}}$ this is obvious. Suppose $|||\overline{\nabla} \log \alpha|||_{0, \Omega_{\tau_1, \tau_2}}$, then as long as $\gamma \subset \Omega_{\tau_1, \tau_2}$, then

$$\log(\alpha(\gamma(t))) - \log \alpha(p) = \int_0^t \langle \gamma', \overline{\nabla} \log \alpha \rangle d\sigma.$$

From this one can obtain the required estimate. \square

Remark 2.2. Hence if in Theorem 1.1, condition (i) is replaced by $\tau_1 < \inf_M \tau \circ \mathbf{X}_0 \leq \sup_M \tau \circ \mathbf{X}_0 < \tau_2$, and either $|||1/\alpha|||_{0, \Omega_{\tau_1, \tau_2}} < \infty$ or $\sup_M 1/(\alpha \circ \mathbf{X}_0) < \infty$ and $|||\overline{\nabla} \log \alpha|||_{0, \Omega_{\tau_1, \tau_2}} < \infty$, then the conclusion of the theorem is still true. Here conditions involving Ω will be replaced by Ω_{τ_1, τ_2} .

3. LONG TIME EXISTENCE

In this section, we will prove the long time existence result: Theorem 1.2. Let us begin with the following lemma.

Lemma 3.1. *Let (N, G) , \mathbf{X}_0 , τ , ρ be as in Theorem 1.2. Suppose $\mathbf{F} : M \times [0, s_0)$, $s_0 < \infty$, is a solution of (1.2) so that the following are true:*

- (i) *For all $s < s_0$, $\sup_{M \times [0, s]} \kappa < \infty$ and $\sup_{M \times [0, s]} |\nabla^k A| < \infty$ for all $k \geq 0$.*
- (ii) *The height function $u(\mathbf{x}, s) =: \tau(\mathbf{F}(\mathbf{x}, s))$ satisfies*

$$\tau_1 < \inf_{M \times [0, s_0)} u \leq \sup_{M \times [0, s_0)} u < \tau_2$$

for some $\tau_- < \tau_1 < \tau_2 < \tau_+$.

Then \mathbf{F} can be extended smoothly up to $s = s_0$ as a solution of the prescribed mean curvature flow so that the tilt factor and all the covariant derivatives of the second fundamental form of $\mathbf{F}^{s_0}(M)$ are uniformly bounded.

Assume the lemma is true, let us prove Theorem 1.2.

Proof. By Theorem 1.1 and by Remark 2.2, there is $\epsilon > 0$ such that the prescribed mean curvature flow (1.2) has a solution \mathbf{F} on $M \times [0, \epsilon]$ such that $\sup_{M \times [0, \epsilon]} |\nabla^k A| < \infty$ for all $k \geq 0$, and $\sup_{M \times [0, \epsilon]} \kappa < \infty$. Let s_{\max} be as in the theorem and $s_{\max} < \infty$. Suppose

$$\tau_- < \tau_1 < \inf_{M \times [0, s_{\max})} \tau(\mathbf{F}) \leq \sup_{M \times [0, s_{\max})} \tau(\mathbf{F}) < \tau_2 < \tau_+.$$

for some τ_1, τ_2 . By Lemma 3.1, \mathbf{F} can be extended smoothly as a solution to the prescribed mean curvature flow up to s_{\max} so that all the derivatives of the second fundamental form and the tilt factor of $\mathbf{F}(\cdot, s_{\max})$ are uniformly

bounded. In particular, $\tau(\mathbf{F}(\mathbf{x}, s)) \rightarrow \tau(\mathbf{F}(\mathbf{x}, s_{\max}))$ as $s \uparrow s_{\max}$. Hence $\tau_1 \leq \tau(\mathbf{F}(\mathbf{x}, s_{\max})) \leq \tau_2$ for all $\mathbf{x} \in M$. On the other hand,

$$\begin{aligned} \log \alpha(\mathbf{F}(\mathbf{x}, s_{\max})) - \log \alpha(\mathbf{F}(\mathbf{x}, 0)) &= \int_0^{s_{\max}} \langle \partial_s \mathbf{F}, \bar{\nabla} \log \alpha \rangle d\sigma \\ &= \int_0^{s_{\max}} \langle (H - \mathcal{H})\vec{\nu}, \bar{\nabla} \log \alpha \rangle d\sigma. \end{aligned}$$

Since $\|\bar{\nabla} \log \alpha\|$, $H - \mathcal{H}$, and the tilt factor κ are all uniformly bounded on Ω_{τ_1, τ_2} , we have

$$\log \alpha(\mathbf{F}(\mathbf{x}, s_{\max})) \geq C \log \alpha(\mathbf{F}(\mathbf{x}, 0))$$

for some $C > 0$ independent of \mathbf{x} . Since $\inf_M \alpha(\mathbf{F}(\cdot, 0)) > 0$, we conclude that $\inf_M \alpha(\mathbf{F}(\cdot, s_{\max})) > 0$. By Theorem 1.1 and by Remark 2.2, \mathbf{F} can be extended beyond s_{\max} satisfying the conditions in Theorem 1.2. This contradicts the definition of s_{\max} . This completes the proof of the theorem. \square

It remains to prove Lemma 3.1. This follows from the basic estimates obtained by Ecker and Husiken [10, 8] and standard arguments as in [2, Chapter 6]. Here we use the existence of ρ to obtain C^0 convergence. For completeness, we sketch the proof. First recall the evolution equations of geometric quantities along the flow, see [10, 8]: Let \mathbf{F} satisfies (1.2).

Lemma 3.2. *With the notation as in Theorem 1.2, let $g(s)$ be the induced metric on M_s with unit future pointing unit normal $\vec{\nu}$, second fundamental form A , mean curvature H , height function u , tilt factor κ .*

- (i) $\frac{\partial}{\partial s} g_{ij} = 2(H - \mathcal{H})A_{ij}$.
- (ii) $\frac{\partial}{\partial s} g^{ij} = -2(H - \mathcal{H})A^{ij}$.
- (iii)

$$\begin{aligned} \frac{\partial}{\partial s} \Gamma_{ij}^k &= -g^{kl} \left[(H - \mathcal{H})(A_{jl;i} + A_{il;j} - A_{ij;l}) \right. \\ &\quad \left. + (H - \mathcal{H})_i A_{jl} + (H - \mathcal{H})_j A_{il} - (H - \mathcal{H})_l A_{ij} \right]. \end{aligned}$$

Here $;$ is the covariant derivative of $g(s)$ and Γ_{ij}^k is the connection.

- (iv) $\frac{\partial}{\partial s} \vec{\nu} = \nabla(H - \mathcal{H})$.
- (v)

$$\begin{aligned} \left(\frac{\partial}{\partial s} - \Delta \right) A_{ij} &= -(|A|^2 + \overline{\text{Ric}}(\vec{\nu}, \vec{\nu}))A_{ij} + 2HA_{ri}A_j^r \\ &\quad + 2g^{kl}A_k^q \overline{\text{R}}_{qijl} - g^{kl}A_j^q \overline{\text{R}}_{qli} - g^{kl}A_l^q \overline{\text{R}}_{qikj} \\ &\quad + g^{kl} \overline{\nabla}_l \overline{\text{R}}_{0ijk} - g^{kl} \overline{\nabla}_j \overline{\text{R}}_{0kli} \\ &\quad - \mathcal{H}(g^{kl}A_{ik}A_{jl} - \overline{\text{R}}_{0ij0}) - \mathcal{H}_{;ij}. \end{aligned}$$

Here the index 0 in $\overline{\mathbf{R}}$ corresponds to $\vec{\nu}$.

(vi)

$$\left(\frac{\partial}{\partial s} - \Delta\right)(H - \mathcal{H}) = -(|A|^2 + \overline{\text{Ric}}(\vec{\nu}, \vec{\nu}) + \langle \overline{\nabla} \mathcal{H}, \vec{\nu} \rangle)(H - \mathcal{H}).$$

(vii) $\frac{\partial}{\partial s} u = \alpha^{-1} \kappa (H - \mathcal{H})$, $\Delta u = \text{div}(\overline{\nabla} \tau) + \alpha^{-1} \kappa H$ and

$$\left(\frac{\partial}{\partial s} - \Delta\right)u = -\mathcal{H} \alpha^{-1} \kappa - \text{div} \overline{\nabla} \tau.$$

Here the divergence is on the immersed surface M_s .

(viii)

$$\left(\frac{\partial}{\partial s} - \Delta\right)\kappa = -\kappa(|A|^2 + \text{Ric}(\vec{\nu}, \vec{\nu})) - \mathbf{T}(H_{\mathbf{T}}) - (H - \mathcal{H})\langle \overline{\nabla}_{\vec{\nu}} \mathbf{T}, \vec{\nu} \rangle + \langle \mathbf{T}, \nabla^M \mathcal{H} \rangle.$$

Here $\mathbf{T}(H_{\mathbf{T}})$ is the variation of the mean curvature under the deformation of \mathbf{T} . (See [3, 2.10]).

By [3], we have the following facts:

Lemma 3.3. *Let M be a spacelike hypersurface with tilt factor κ and let $u = \tau|_M$. Then*

$$|\nabla^M u|^2 = \alpha^{-2}(\kappa^2 - 1).$$

For any $1 > \delta > 0$,

$$|\nabla_M \kappa|^2 \leq \left(1 + \frac{\delta}{n}\right)^{-1} \kappa^2 |A|^2 + C(\kappa^4 + \kappa^2 H^2),$$

for some $C = C(\|\overline{\nabla} \mathbf{T}\|, n, (1 - \delta)^{-1})$.

Since all the covariant derivatives of the curvature of (M, g_0) are bounded, we can find a smooth function η with bounded gradient and bounded Hessian, such that $d(\mathbf{x}) + C \leq \eta(\mathbf{x}) \leq 2d(\mathbf{x}) + C'$ where $d(\mathbf{x})$ is the distance function with respect to g_0 from a fixed point \mathbf{x}_0 , see [26, 27].

By Lemma 3.2 (i)–(iii) we have:

Lemma 3.4. *Let \mathbf{F} be a solution of (1.2) on $M \times [0, s_0]$. Suppose $|A|, |\nabla A|$ are uniformly bounded on $M \times [0, s_0]$, then gradient and the Hessian of η with respect to $g(s)$ are uniformly bounded.*

Using η , we have the following maximum principle.

Lemma 3.5. *Let \mathbf{F} and η be as in Lemma 3.4. Suppose f is a smooth function on $M \times [0, s_0]$ such that $f \leq C_1 e^{C_2 \eta}$ for some $C_1, C_2 > 0$ and $\left(\frac{\partial}{\partial s} - \Delta\right)f \leq 0$ whenever $f > 0$. Assume $\sup_{M_0} f < \infty$. Then $\sup_{M \times [0, s_0]} f \leq \max\{0, \sup_{M_0} f\}$.*

Proof. Let $a = \sup_{M_0} f$ and let $\phi = e^{\sigma s + 2C_2 \eta}$ where σ is to be determined. By Lemma 3.4, we have

$$\left(\frac{\partial}{\partial s} - \Delta\right)\phi \geq (\sigma - C_3)\phi$$

for some $C_3 > 0$ in $M \times [0, s_0]$. Choose $\sigma = 2C_3$, we have $(\frac{\partial}{\partial s} - \Delta)\phi > 0$. For any $\epsilon > 0$,

$$\left(\frac{\partial}{\partial s} - \Delta\right)(f - a - \epsilon\phi) < 0$$

whenever $f > 0$. $\sup_{M \times [0, s_0]}(f - a - \epsilon\phi)$ will be attained at some point in $(\mathbf{x}, s) \in M \times [0, s_0]$. If $s > 0$, we must have $f \leq 0$ at this point. So $\sup_{M \times [0, s_0]}(f - a - \epsilon\phi) \leq \max\{-a, 0\}$. Let $\epsilon \rightarrow 0$, the result follows. \square

The following is a basic estimate obtained by Ecker [8]. We modify the proof a little bit using the idea in [3]. The assumptions are slightly different from those in [8]. We give details of the proof in order to understand what conditions are sufficient.

Lemma 3.6. *[Ecker's basic estimate] Let \mathbf{F} be a solution to the prescribed mean curvature flow (1.2) such that $\tau_- < \tau_1 < \tau(\mathbf{F}) < \tau_2 < \tau_+$.*

- (i) $\sup_{M \times [0, s_0]}(\kappa + |A| + |\nabla A|) < \infty$.
- (ii) $\|\|\nabla \mathbf{T}\|\|_{1, \Omega_{\tau_1, \tau_2}}, \|\|\mathcal{H}\|\|_{1, \Omega}, \|\|\overline{\text{Rm}}\|\|_{1, \Omega_{\tau_1, \tau_2}}, \|\|\overline{\nabla} \log \alpha\|\|_{1, \Omega_{\tau_1, \tau_2}}, \|\|\alpha\|\|_{1, \Omega_{\tau_1, \tau_2}}$ are uniformly bounded by c_1 .
- (iii) $\overline{\text{Ric}}(w, w) \geq -c_2$ for some $c_2 \geq 0$ for all time like unit vector w in Ω_{τ_1, τ_2} .

Then there exist $\lambda > 0, \mu > 0, C$ depending only on $c_1, c_2, \tau_1, \tau_2, n$ with C also depending on $\sup_{M_0}(\kappa + |H - \mathcal{H}|)$ such that

$$\sup_{M \times [0, s_0]} (e^{\lambda u} \kappa^2 + \mu(H - \mathcal{H})^2) \leq C.$$

Proof. Let $u = \tau(\mathbf{F})$. Then $\tau_1 < u < \tau_2$. By Lemma 3.2(viii)

$$\begin{aligned} \left(\frac{\partial}{\partial s} - \Delta\right)e^{\lambda u} &= \lambda e^{\lambda u} (-\mathcal{H}\alpha^{-1}\kappa - \text{div}_M \overline{\nabla} \tau - \lambda|\nabla u|^2) \\ (3.1) \quad &\leq \lambda e^{\lambda u} (-\mathcal{H}\alpha^{-1}\kappa + \kappa^2 \|\|\overline{\nabla}(\alpha^{-1}\mathbf{T})\|\| - \lambda|\nabla u|^2) \\ &\leq \lambda e^{\lambda u} (C_1 \alpha^{-1} \kappa^2 - \lambda|\nabla u|^2) \end{aligned}$$

for some $C_1 = C_1(c_1, n)$ because $\overline{\nabla} \tau = -\alpha^{-1}\mathbf{T}$, $\overline{\nabla} \alpha^{-1} = -\alpha^{-1}\overline{\nabla} \log \alpha$ and $\kappa \geq 1$. Using the facts that

$$|\langle \overline{\nabla}_{\vec{\nu}} \mathbf{T}, \vec{\nu} \rangle| \leq 2\kappa^2 \|\|\overline{\nabla} \mathbf{T}\|\|; \quad |\langle \mathbf{T}, \nabla \mathcal{H} \rangle| \leq 2\kappa \|\|\overline{\nabla} \mathcal{H}\|\|,$$

and

$$|\mathbf{T}(H_{\mathbf{T}})| \leq C(n) \left(\|\|\overline{\nabla}^2 \mathbf{T}\|\| \kappa^3 + \|\|\overline{\nabla} \mathbf{T}\|\| \kappa^2 |A| \right)$$

by Bartnik [3, 2.10], using Lemma 3.2 we have for any $\epsilon > 0$:

$$\begin{aligned} \left(\frac{\partial}{\partial s} - \Delta\right)\kappa^2 &= 2\kappa \left(\frac{\partial}{\partial s} - \Delta\right)\kappa - 2|\nabla \kappa|^2 \\ &\leq -2(1 - \epsilon)\kappa^2 |A|^2 + 2C_2 \epsilon^{-1} \kappa^4 - 2|\nabla \kappa|^2 \end{aligned}$$

for some $C_2 = C_2(c_1, n)$. On the other hand, by Lemma 3.3, we have for $1 > \delta > 0$,

$$|\nabla\kappa|^2 \leq \left(1 + \frac{\delta}{n}\right)^{-1} \kappa^2 |A|^2 + C(\kappa^4 + \kappa^2 H^2)$$

for some $C = C(\|\overline{\nabla}T\|_{0,\Omega}, n, (1 - \delta)^{-1})$. Hence by choosing suitable δ, ϵ , we have

$$(3.2) \quad \left(\frac{\partial}{\partial s} - \Delta\right)\kappa^2 \leq -4\left(1 + \frac{1}{4n}\right)|\nabla\kappa|^2 + C_3(\kappa^4 + \kappa^2 H^2)$$

for some $C_3 = C_3(c_1, n)$. Since $\overline{\text{Ric}}(\vec{\nu}, \vec{\nu}) \geq -c_2$, and

$$(3.3) \quad |A|^2 \geq \frac{1}{n}H^2 \geq \frac{1}{2n}(H - \mathcal{H})^2 - \frac{1}{n}\mathcal{H}^2,$$

by Lemma 3.2(vi), we have

$$(3.4) \quad \begin{aligned} \left(\frac{\partial}{\partial s} - \Delta\right)(H - \mathcal{H})^2 &\leq -2(|A|^2 - C_4\kappa)(H - \mathcal{H})^2 - 2|\nabla(H - \mathcal{H})|^2 \\ &\leq -\frac{1}{n}(H - \mathcal{H})^4 + C_5\kappa(H - \mathcal{H})^2 + \frac{2}{n}\mathcal{H}^2 - 2|\nabla(H - \mathcal{H})|^2 \\ &\leq -\frac{1}{2n}(H - \mathcal{H})^4 - 2|\nabla(H - \mathcal{H})|^2 + C_6\kappa^2 \end{aligned}$$

for some $C_4, C_5, C_6 > 0$ depending only on c_1, c_2, n . Let $\mu > 0$ to be determined, and let

$$\Phi = e^{\lambda u} \kappa^2 + \mu(H - \mathcal{H})^2.$$

We may assume $C_3 \geq C_1$. Then by combining (3.1), (3.2), (3.4), we have:

$$(3.5) \quad \begin{aligned} \left(\frac{\partial}{\partial s} - \Delta\right)\Phi &\leq e^{\lambda u} \left[-4\left(1 + \frac{1}{4n}\right)|\nabla\kappa|^2 - \lambda^2 \kappa^2 |\nabla u|^2 \right] + C_3 e^{\lambda u} \kappa^2 ((\lambda\alpha^{-1} + 1)\kappa^2 + H^2) \\ &\quad - 2\langle \nabla e^{\lambda u}, \nabla \kappa^2 \rangle - \frac{\mu}{2n}(H - \mathcal{H})^4 - 2\mu|\nabla(H - \mathcal{H})|^2 + C_6\mu\kappa^2 \end{aligned}$$

Now

$$\begin{aligned} &e^{\lambda u} \left[-4\left(1 + \frac{1}{4n}\right)|\nabla\kappa|^2 - \lambda^2 \kappa^2 |\nabla u|^2 \right] - 2\langle \nabla e^{\lambda u}, \nabla \kappa^2 \rangle \\ &\leq e^{\lambda u} \left[-\frac{1 + 4n}{4n} \kappa^{-2} |\nabla \kappa^2|^2 - \lambda^2 \kappa^2 |\nabla u|^2 + 2\lambda |\nabla u| |\nabla \kappa^2| \right] \\ &= e^{\lambda u} \left[-\frac{1 + 4n}{4n} \kappa^{-2} |\nabla \kappa^2|^2 - \frac{4n}{1 + 4n} \lambda^2 \kappa^2 |\nabla u|^2 + 2\lambda |\nabla u| |\nabla \kappa^2| \right] - \frac{1}{1 + 4n} \lambda^2 e^{\lambda u} \kappa^2 |\nabla u|^2 \\ &\leq -\frac{1}{1 + 4n} \lambda^2 e^{\lambda u} \alpha^{-2} \kappa^2 (\kappa^2 - 1) \end{aligned}$$

by Lemma 3.3.

$$\begin{aligned} \left(\frac{\partial}{\partial s} - \Delta\right)\Phi &\leq -\frac{1}{4n+1}\lambda^2\kappa^2e^{\lambda u}\alpha^{-2}(\kappa^2-1) + C_3e^{\lambda u}\kappa^2((\lambda\alpha^{-1}+1)\kappa^2+H^2) \\ &\quad -\frac{\mu}{2n}(H-\mathcal{H})^4 - 2\mu|\nabla(H-\mathcal{H})|^2 + C_6\mu\kappa^2. \end{aligned}$$

Since $H^2 \leq 2(H-\mathcal{H})^2 + 2\mathcal{H}^2$,

$$e^{\lambda u}\kappa^2H^2 \leq e^{\lambda u}(\kappa^4 + (H-\mathcal{H})^4 + 2\mathcal{H}^2\kappa^2).$$

Hence

$$\begin{aligned} \left(\frac{\partial}{\partial s} - \Delta\right)\Phi &\leq -\frac{1}{4n+1}\lambda^2\kappa^2e^{\lambda u}\alpha^{-2}(\kappa^2-1) + C_7e^{\lambda u}(\lambda\alpha^{-1}+1)\kappa^4 \\ &\quad -\left(\frac{\mu}{2n} - C_3e^{\lambda u}\right)(H-\mathcal{H})^4 - 2\mu|\nabla(H-\mathcal{H})|^2 + C_8e^{-\lambda u}(\mu + e^{\lambda u})^2 \end{aligned}$$

for some positive constants C_7, C_8 depending only on c_1, c_2, n . Choose μ such that

$$\frac{\mu}{2n} - C_3e^{\lambda\tau_2} = 1.$$

Note that $\mu = \mu(c_1, c_2, n, \lambda, \tau_2)$. We have

$$(3.6) \quad \begin{aligned} \left(\frac{\partial}{\partial s} - \Delta\right)\Phi &\leq -\frac{1}{4n+1}\lambda^2\kappa^2e^{\lambda u}\alpha^{-2}(\kappa^2-1) + C_7e^{\lambda u}(\lambda\alpha^{-1}+1)\kappa^4 \\ &\quad - (H-\mathcal{H})^4 - 2\mu|\nabla(H-\mathcal{H})|^2 + C_8e^{-\lambda u}(\mu + e^{\lambda u})^2. \end{aligned}$$

Choose λ large enough depending only on $\|\alpha\|_{0,\Omega_{\tau_1,\tau_2}}$ and C_7 , so that if $\kappa \geq 2$, then

$$-\frac{1}{4n+1}\lambda^2\alpha^{-2}(\kappa^2-1) + C_7(\lambda\alpha^{-1}+1)\kappa^2 \leq -\kappa^2.$$

Note that $\lambda = \lambda(c_1, c_2, n)$ and $\mu = \mu(c_1, c_2, n, \tau_2)$. So for $\kappa \geq 2$, we have

$$(3.7) \quad \left(\frac{\partial}{\partial s} - \Delta\right)\Phi \leq -e^{\lambda u}\kappa^4 - (H-\mathcal{H})^4 + C_8e^{-\lambda u}(\mu + e^{\lambda u})^2$$

As in the proof of Lemma 3.5, let $\phi = e^{\sigma s + \eta}$ for some $\sigma > 0$ where η is as in Lemma 3.4 so that

$$\left(\frac{\partial}{\partial s} - \Delta\right)\phi > 0.$$

Let $m_0 = \sup_{M_0}\Phi$. For any $\epsilon > 0$. Then $\Phi - m_0 - \epsilon\phi < 0$ at $s = 0$ and at infinity. If $\Phi - m_0 - \epsilon\phi > 0$ somewhere, then there is $\mathbf{x}_1 \in M, 0 < s_1 \leq s_0$ so that $(\Phi - m_0 - \epsilon\phi)(\mathbf{x}_1, s_1) = \sup_{M \times [0, s_0]}(\Phi - m_0 - \epsilon\phi)$. At this point we have:

$$0 \leq \left(\frac{\partial}{\partial s} - \Delta\right)(\Phi - m_0 - \epsilon\phi) < \left(\frac{\partial}{\partial s} - \Delta\right)\Phi.$$

Suppose at this point $\kappa \geq 2$, then by (3.7)

$$0 \leq -e^{\lambda u}\kappa^4 - (H-\mathcal{H})^4 + C_9e^{-\lambda u}(\mu + e^{\lambda u})^2.$$

Hence $e^{\lambda u}\kappa^2 + (H-\mathcal{H})^2 \leq C(c_1, c_2, n, \tau_1, \tau_2)$. This implies $\Phi - \epsilon\phi \leq C$ for some $C = C(c_1, c_2, n, \tau_1, \tau_2)$.

Suppose at this point $\kappa \leq 2$, then by (3.6)

$$(H - \mathcal{H})^4 \leq C(c_1, c_2, n, \tau_1, \tau_2).$$

and we also have $\Phi - \epsilon\phi \leq C(c_1, c_2, n, \tau_1, \tau_2)$. In any case,

$$\Phi - \epsilon\phi \leq C(c_1, c_2, n, \tau_1, \tau_2) + m_0.$$

Let $\epsilon \rightarrow 0$, the result follows. \square

By [10, p.604] and the proof of [21, Th. 4.1], we have the following estimate for $|\nabla^k A|$.

Lemma 3.7. *Let $\mathbf{F} : M \times [0, s_0]$ be a solution to the prescribed mean curvature flow (1.2) so that the initial metric on M_0 is complete with bounded second fundamental form and so that the tilt factor of the immersed surface $\mathbf{F}^s : M \rightarrow N$ is uniformly bounded by κ_0 . Suppose $\mathbf{F}(M \times [0, s_0]) \subset \Omega$ and $|||\overline{\text{Rm}}|||_{k,\Omega}$, $|||\mathcal{H}|||_{k,\Omega}$ are finite for all $k \geq 0$ and suppose $\sup_{M \times [0, s_0]} |\nabla^k A| < \infty$ for all k . Then for all $m \geq 0$,*

$$\sup_{M \times [0, s_0]} |\nabla^m A|^2 \leq C_m$$

for some C_m depending only on $n, m, \sup_{M \times [0, s_0]} \kappa, |||\overline{\text{Rm}}|||_{m+1,\Omega}, |||\mathcal{H}|||_{m+2,\Omega}$, and $\sum_{j=0}^{m-1} \sup_{M_0} |\nabla^j A|$.

Now we are ready to prove Lemma 3.1.

Proof of Lemma 3.1. By Lemma 3.6, there is a constant κ_1 such that

$$(3.8) \quad \sup_{M \times [0, s_0]} \kappa \leq \kappa_1.$$

Using this, by Lemma 3.7, for any m , there is a constant c_m such that

$$(3.9) \quad \sup_{M \times [0, s_0]} |\nabla^m A| \leq c_m.$$

Let $\mathbf{x}_0 \in M$ and let $B_{\mathbf{x}_0}(r_0)$ be a geodesic ball with respect to g_0 with center \mathbf{x}_0 and radius r_0 , which is inside a coordinate chart with $\mathbf{x} = (x^1, \dots, x^n)$. First we want to obtain C^0 limit of $\mathbf{F}(\cdot, s)$ as $s \rightarrow s_0$. Let ρ be the function satisfying condition (1.4). Since $\mathbf{F}(\mathbf{x}, s) \in \Omega_{\tau_1, \tau_2}$, by the assumption on ρ , and (1.2) we have for $\mathbf{x} \in B_{\mathbf{x}_0}(r_0)$,

$$\rho(\mathbf{F}(\mathbf{x}, s)) = \int_0^s \langle \overline{\nabla} \rho, \partial_s \mathbf{F} \rangle d\sigma \leq C_1 s$$

for some constant C_1 depending only on $\kappa_0, \sup_{M \times [0, s_0]} (|H - \mathcal{H}|)$ and $|||\overline{\nabla} \rho|||_{0, \Omega_{\tau_1, \tau_2}}$. Hence $\mathbf{F}(B_{\mathbf{x}_0}(r_0) \times [0, s_0]) \subset \overline{\Omega}_{\tau_1, \tau_2} \cap \{\rho \leq \rho_0\} = K$ which is compact, where

$$\rho_0 = C_1 s_0 + \sup_{B_{\mathbf{x}_0}(r_0)} \rho + 1.$$

Hence K is compact with respect to the reference Riemannian metric G_E with respect to τ . Let $d(p, q)$ be the distance function defined by G_E . Then for $\mathbf{x} \in B_{\mathbf{x}_0}(r_0)$, for $s < s' < s_0$ by (1.2) we have:

$$d(\mathbf{F}(\mathbf{x}, s), \mathbf{F}(\mathbf{x}, s')) \leq \int_s^{s'} \|\partial_s \mathbf{F}\| d\sigma \leq C_2(s' - s)$$

for some constant C_2 depending only on $\kappa_0, \sup_{M \times [0, s_0]} (|H - \mathcal{H}|)$. Since K is compact, we conclude that $\mathbf{F}(\mathbf{x}, s)$ converges uniformly on $B_{\mathbf{x}_0}(r_0)$ as $s \rightarrow s_0$ to a continuous map \mathbf{F}^{s_0} from $B_{\mathbf{x}_0}(r_0)$ to N with image inside K . Let $p = \mathbf{F}^{s_0}(\mathbf{x}_0)$. Then there is $\sigma_0 > 0$ such that $\mathbf{F}^s(B_{\mathbf{x}_0}, r_0) \Subset D_p(\sigma_0)$ for all $s_1 < s < s_0$ for some $s_1 > 0$, for a possibly smaller r_0 . Here $D_p(\sigma_0)$ is the geodesic ball with respect to G_E with center at p , and radius σ_0 . We may also assume that $D_p(\sigma_0)$ is inside a coordinate neighborhood with coordinates y^0, y^1, \dots, y^n so that G_E is uniformly equivalent to the Euclidean metric and the components of the metric tensor and its derivatives with respect to y^a are uniformly bounded. Then \mathbf{F} is of the form:

$$\mathbf{F} = (f^0, f^1, \dots, f^n).$$

In order to prove the lemma, it is sufficient to prove that $\partial_s^k \partial^\beta f^a$ are uniformly bounded, for any $k, \ell \geq 0$ for any multi-index β for x^i with $|\beta| \leq \ell$. This is rather standard because of (3.8), (3.9) and Lemma 3.2. We sketch the proof as follows. Let $\partial_i = \frac{\partial}{\partial x^i}$ be the coordinate frame in $B_{\mathbf{x}_0}(r_0)$. Then

$$(3.10) \quad \begin{cases} \mathbf{F}_*(\partial_i) = f_i^a \frac{\partial}{\partial y^a}; \\ (H - \mathcal{H})\vec{\nu} = \partial_s \mathbf{F} = f_s^a \frac{\partial}{\partial y^a} \\ A_{ij}\vec{\nu} = (f_{ij}^a - \Gamma_{ij}^k f_k^a + f_i^b f_j^c \bar{\Gamma}_{bc}^a) \frac{\partial}{\partial y^a}. \end{cases}$$

here f_i^a, f_{ij}^a are partial derivatives with respect to x^i , $\bar{\Gamma}$ is the connection of the target.

By (3.8), (3.9) and Lemma 3.2(i)–(iii), we can conclude that $g_{ij}(s)$ is uniformly equivalent to $g_{ij}(0)$, and $\Gamma_{ij}^k(s)$ is uniformly bounded. Hence f_i^a, f_s^a are uniformly bounded. By differentiate the equation of $\partial_s \Gamma_{ij}^k$ with respect to x^l and integrate, we can conclude that all the partial derivatives of Γ_{ij}^k with respect to x^l are uniformly bounded. Hence all the partial derivatives of A_{ij} and H are bounded. Since $\|\mathcal{H}\|_{k, \Omega_{\tau_1, \tau_2}} < \infty$ for all $k \geq 0$, we conclude that all the partial derivatives of \mathcal{H} are uniformly bounded using (3.8). Using (3.10), we may conclude that f_{ij}^a are uniformly bounded. Take derivatives of the third equation with respect to x^l in (3.10), we conclude that all partial derivatives of f^a with respect to x^i are uniformly bounded. Differentiate the third equation with respect to x^l , then we conclude inductively that $\partial^\beta f^a$ are uniformly bounded for all multi-index β for x^i . In particular, $\mathbf{F}(\mathbf{x}, s)$ converges in C^∞ in $B_{\mathbf{x}_0}(r_0)$ to $\mathbf{F}^{s_0}(\mathbf{x})$. Next, we want to investigate $\partial_s^k \partial^\beta f^a$. By Lemma 3.2(v), (vi) one can conclude that $\partial_s^k \partial^\beta (H - \mathcal{H}), \partial_s^k \partial^\beta A_{ij}$ are uniformly bounded. For

example

$$\partial_s(H - \mathcal{H}) = \Delta(H - \mathcal{H}) - (|A|^2 + \overline{\text{Ric}}(\vec{\nu}, \vec{\nu}) + \langle \overline{\nabla} \mathcal{H}, \vec{\nu} \rangle)(H - \mathcal{H}).$$

Hence $\partial_s(H - \mathcal{H})$ is uniformly bounded. Note that

$$\partial_s \vec{\nu} = \nabla(H - \mathcal{H})$$

and

$$\overline{\nabla}_{\mathbf{F}_*(\partial_i)} \vec{\nu} = g^{kl} A_{il} \mathbf{F}_*(\partial_k) = g^{kl} A_{il} f_l^a \frac{\partial}{\partial y^a}.$$

Differentiate with respect to x^l , we conclude that $\partial_s \partial^\beta(H - \mathcal{H})$ are uniformly bounded. Similarly, $\partial_s \partial^\beta A_{ij}$ are uniformly bounded. By differentiate $\partial_s \partial^\beta(H - \mathcal{H})$ with respect to s one can conclude that $\partial_s^2 \partial^\beta(H - \mathcal{H})$ are uniformly bounded. Here we also use the fact that $\partial_s g_{ij}$ are uniformly bounded. Inductively, one can prove that $\partial_s^k \partial^\beta(H - \mathcal{H})$, $\partial_s^k \partial^\beta A_{ij}$ are uniformly bounded. Differentiate the second equation in (3.10) with respect to x^l, s , we conclude inductively that $\partial_s^k \partial^\beta f^a$ are uniformly bounded. Hence the lemma is true. \square

4. CONVERGENCE

We will prove the results on convergence. First we prove Theorem 1.3.

Proof of Theorem 1.3. As before, we will denote the immersed surface $\mathbf{F}(\cdot, s) : M \rightarrow N$ simply by M_s . By Lemmas 3.6, 3.7 and the assumptions of the theorem, there exists κ_1 and for any $m \geq 0$, there exists c_m such that

$$(4.1) \quad \sup_{M \times [0, \infty)} \kappa \leq \kappa_1; \quad \sup_{M \times [0, \infty)} |\nabla^m A| \leq c_m.$$

At $s = 0$, $H \geq \mathcal{H} - a + \varepsilon_0 \geq \varepsilon_0$ and

$$(4.2) \quad |A|^2 + \overline{\text{Ric}}(\vec{\nu}, \vec{\nu}) + \langle \overline{\nabla} \mathcal{H}, \vec{\nu} \rangle \geq \frac{H^2}{n} - c^2 \geq \frac{\varepsilon_0^2}{n} - c^2 > 0.$$

We want to prove that this is still true for $s > 0$. Since $\tau_- < \tau_1 \leq \tau(\mathbf{F}) \leq \tau_2 < \tau_+$, using (4.1), Lemma 3.2(vi) and the assumptions on $\overline{\text{Rm}}, \mathcal{H}$, we conclude that for any $\epsilon > 0$, there is $s_0 > 0$ so that for $0 \leq s \leq s_0$,

$$|A|^2 + \overline{\text{Ric}}(\vec{\nu}, \vec{\nu}) + \langle \overline{\nabla} \mathcal{H}, \vec{\nu} \rangle > 0.$$

By Lemma 3.2(vi), in $M \times [0, s_0]$,

$$\left(\frac{\partial}{\partial s} - \Delta \right) (\mathcal{H} - H) \leq -(|A|^2 + \overline{\text{Ric}}(\vec{\nu}, \vec{\nu}) + \langle \overline{\nabla} \mathcal{H}, \vec{\nu} \rangle)(\mathcal{H} - H)$$

which is negative at the points where $\mathcal{H} - H > 0$. By Lemma 3.5,

$$\sup_{M \times [0, s_0]} (\mathcal{H} - H) \leq \max\{0, \sup_{M_0} (\mathcal{H} - H)\} \leq \max\{0, a - \epsilon_0\} = a - \epsilon_0.$$

because $a > \epsilon_0$. So $H \geq \mathcal{H} - a + \epsilon_0 \geq \epsilon_0$ because $\mathcal{H} \geq a$. Hence (4.2) is true on $M \times [0, s_0]$. Because $H - \mathcal{H} \geq -a + \epsilon_0$ at $s = s_0$, iterating, we conclude that (4.2) is true for all s . So by Lemma 3.2(vi),

$$(4.3) \quad \left(\frac{\partial}{\partial s} - \Delta \right) (H - \mathcal{H})^2 \leq -\epsilon_1 (H - \mathcal{H})^2 - 2|\nabla(H - \mathcal{H})|^2.$$

Using Lemmas 3.4 and 3.5, we can conclude that there is a constant $C_0, b_0 > 0$ such that

$$(4.4) \quad \sup_{M_s} |H - \mathcal{H}|^2 \leq C_0 e^{-b_0 s}.$$

By Lemma 3.2(i), we have that

$$(4.5) \quad \lambda^{-1} g(0) \leq g(s) \leq \lambda g(0)$$

for some $\lambda > 0$ for all s , where $g(s)$ is the induced metric on M_s . Next, we want to show that for any $k \geq 1$, there are constants $C_k, b_k > 0$ such that

$$(4.6) \quad \sup_{M_s} |\nabla^k (H - \mathcal{H})|^2 \leq C_k e^{-b_k s}.$$

We will prove this inductively. To simplify the notation, we write the evolution equation Lemma 3.2(vi) as

$$(4.7) \quad \left(\frac{\partial}{\partial s} - \Delta \right) (H - \mathcal{H}) = \psi (H - \mathcal{H}).$$

where $\psi = -(|A|^2 + \overline{\text{Ric}}(\vec{\nu}, \vec{\nu}) + \langle \overline{\nabla} \mathcal{H}, \vec{\nu} \rangle)$. By (3.10) and the fact that (1.3) is true, $|\nabla^m \psi|$ is uniformly bounded in space time for all $m \geq 0$. Let us prove that the estimate (4.6) is true for $k = 1$. The general case can be proved similarly by induction.

By the Bochner's formula:

$$\begin{aligned} \left(\frac{\partial}{\partial s} - \Delta \right) |\nabla(H - \mathcal{H})|^2 &= -2|\nabla^2(H - \mathcal{H})|^2 - \langle \nabla(\psi(H - \mathcal{H})), \nabla(H - \mathcal{H}) \rangle \\ &\quad - 2\text{Ric}(\nabla(H - \mathcal{H}), (\nabla(H - \mathcal{H}))) \\ &\leq -2|\nabla^2(H - \mathcal{H})|^2 + K_1 (|\nabla(H - \mathcal{H})|^2 + (H - \mathcal{H})^2) \end{aligned}$$

in $M \times [0, \infty)$ for some $K_1 > 0$. Here Ric is the Ricci tensor of M_s which is uniformly bounded from below because of (3.10) and (1.3). Combining with (4.3), there exists $K_2 > 0$ and $b_2 > 0$ such that on $M \times [0, \infty)$:

$$\begin{aligned} \left(\frac{\partial}{\partial s} - \Delta \right) (|\nabla(H - \mathcal{H})|^2 + K_2 (H - \mathcal{H})^2) &\leq -2|\nabla^2(H - \mathcal{H})|^2 \\ &\quad - b_2 (|\nabla(H - \mathcal{H})|^2 + K_2 (H - \mathcal{H})^2). \end{aligned}$$

From this one can conclude that (4.6) is true for $k = 1$. Using (4.1), (1.3), we conclude that all the derivatives of Rm on M_s are uniformly bounded in space time. Hence by differentiating (4.7), argue as above we can conclude that (4.6) is true for all k by induction.

Using Lemma 3.2(iii), one can conclude that $\partial_s \Gamma_{ij}^k$ is of exponential decay and so $\Gamma(s) - \Gamma(0)$ are uniformly bounded in spacetime. Differentiate the equation, we have $\nabla^k(\Gamma(s) - \Gamma(0))$ are also uniformly bounded in space time. One can proceed as in the proof of Theorem 1.2 to conclude that as $s \rightarrow \infty$, $\mathbf{F}(\cdot, s)$ converges in C^∞ norm in compact sets in M to \mathbf{F}_∞ so that $\mathbf{F}_\infty : M \rightarrow N$ is an immersed spacelike hypersurface which is complete in the induced metric because of (4.5). By (4.4), the mean curvature H_∞ of M_∞ is equal to \mathcal{H} .

To prove the last assertion, we follow [8]. For any $s_0 > 0$, by (4.1) and Lemma 3.2, $\nabla\phi$ and $\nabla^2\phi$ are uniformly bounded with respect to $g(s)$ in $M \times [0, s_0]$. Using (4.3), one obtain

$$\left(\frac{\partial}{\partial s} - \Delta\right)(\phi^2(H - \mathcal{H})^2) \leq C\phi^2(H - \mathcal{H})^2$$

on $M \times [0, s_0]$ for some $C > 0$. Hence

$$\left(\frac{\partial}{\partial s} - \Delta\right)(e^{-Cs}\phi^2(H - \mathcal{H})^2) \leq 0.$$

By Lemmas 3.4 and 3.5, we have

$$\sup_{M \times [0, s_0]} \phi^2(H - \mathcal{H})^2 < \infty.$$

Combining this with Lemma 3.2(vii) and (4.1) and the fact that α^{-1} is uniformly bounded in $M \times [0, s_0]$, one can conclude that

$$\left|\frac{\partial}{\partial s}\phi^2(u(s) - u_0)^2\right| \leq C$$

in $M \times [0, s_0]$ for some $C > 0$. Here $u(s)$ is the height function of M_s . This implies that $u(\mathbf{x}, s) - u_0(\mathbf{x}) \rightarrow 0$ as $\mathbf{x} \rightarrow \infty$ uniformly in $s \in [0, s_0]$. On the other hand, by Lemma 3.2(vii), (4.4), (4.1) and the fact that α^{-1} is uniformly bounded because the image of \mathbf{F} is inside Ω_{τ_1, τ_2} , we have

$$|u_\infty(\mathbf{x}) - u(\mathbf{x}, s)| \leq C'e^{-b_0s}$$

for some $C' > 0$ for all \mathbf{x}, s . Combining these two facts, it is easy to see that the last assertion is true. □

The method of proof of the last assertion in the theorem can be used to obtain height estimate of the prescribed mean curvature flow if we have suitable barrier surface. In this case, one may construct spacelike hypersurface with prescribed mean curvature. One example is as follows. First recall that the *past domain of dependence* $D^-(S)$ of an achronal set S is the set consisting of points p in the spacetime so that every inextendible future causal curve from p will meet S . We have the following:

Corollary 4.1. *Let $(N^{n+1}, G), \mathbf{X}_0, \tau, \rho, \mathcal{H}$ be as in Theorem 1.2 so that $M_0 \subset \Omega_{\tau_1, \tau_2}$ for some $\tau_- < \tau_1 < \tau_2 < \tau_+$. In addition, assume the following:*

- (i) $\overline{\text{Ric}}(w, w) \geq 0$ for all time like vector;
- (ii) \mathcal{H} is monotone, $\mathcal{H} \geq a > 0$ for some constant a , $H - \mathcal{H} \geq 0$ on M_0 , and $\sup_{M_0} \phi(H - \mathcal{H}) < \infty$ for some smooth function $\phi \geq 1$ on M_0 with $\sup_{M_0} (|\nabla \phi| + |\nabla^2 \phi|) < \infty$ and $\phi \rightarrow \infty$ as $\mathbf{x} \rightarrow \infty$ in M_0 ;
- (iii) α^{-1} is uniformly bounded on $\Omega_{\tau', \tau''}$ for any $\tau_- < \tau' < \tau'' < \tau_+$;
- (iv) there exists an achronal spacelike hypersurface M^+ with mean curvature $H^+ < \mathcal{H}$ on M^+ , $\tau_+ > \sup \tau|_{M^+} \geq \inf \tau|_{M^+} > \tau_2$ such that $M_0 \subset D^-(M^+)$.

Then the prescribed mean curvature flow (1.2) has long time solution \mathbf{F} and \mathbf{F} converges in C^∞ norms on compact sets of M to a complete spacelike hypersurface with prescribed mean curvature \mathcal{H} .

Proof. Suppose $H \equiv \mathcal{H}$ at M_0 , then there is nothing to prove. So we assume that $H - \mathcal{H} > 0$ somewhere initially. By Theorems 1.2 and 1.3 it is sufficient to obtain height estimates for solution of (1.2). Let \mathbf{F} be a solution on $M \times [0, s_0]$ so that $\sup_{M \times [0, s_0]} (|\nabla^k A| + \kappa) < \infty$. As in the proof of the last assertion of Theorem 1.3, if we let $u = \tau(\mathbf{F})$ be the height function of M_s , then $u(\mathbf{x}, s) - u_0(\mathbf{x}) \rightarrow 0$ as $\mathbf{x} \rightarrow \infty$ uniformly on s , where $u_0 = \tau|_{M_0}$. Hence there is $\epsilon > 0$ and a compact set K , such that $u \leq \tau_{M^+} - \epsilon$ for all $\mathbf{x} \notin K$ and for all $0 \leq s \leq s_0$. By Lemma 3.2(vi) and assumptions (i) and the fact that \mathcal{H} is monotone, we have

$$\left(\frac{\partial}{\partial s} - \Delta \right) (H - \mathcal{H}) = \psi(H - \mathcal{H})$$

with $\psi \leq 0$. Hence Lemma 3.5 and the strong maximum principle [24, Theorem 2.7], $H - \mathcal{H} > 0$ if $s > 0$. By (1.2), $\partial_s \mathbf{F}$ is future pointing time like vector for $s > 0$. Hence for any $\mathbf{x} \in M$, $\mathbf{F}(\mathbf{x}, s)$, $0 \leq s \leq s_0$ is a future nonspacelike curve. Since τ is nondecreasing on such a curve, we have $\tau(\mathbf{F}(\mathbf{x}, s)) \geq \tau_1$ for all $\mathbf{x} \in M$, $s \in [0, s_0]$. To obtain an upper bound, we proceed as in [10]. Note that $\mathbf{F}(\mathbf{x}, s)$, $0 \leq s \leq s_0$ is part of an inextendible future nonspacelike curve. Suppose $\mathbf{F}(\mathbf{x}, s)$, $0 \leq s \leq s_0$ does not intersect M^+ , then $\tau(\mathbf{F}(\mathbf{x}, s)) \leq \sup \tau|_{M^+}$ because τ is nondecreasing on a future causal curve. Hence we may assume that $M_s \cap M^+ \neq \emptyset$ for some $0 < s \leq s_0$. Since $u \leq \tau_{M^+} - \epsilon$ for all $\mathbf{x} \notin K$ and for all $0 \leq s \leq s_0$, we conclude that there exists $0 < s_1 \leq s_0$ and $\mathbf{x}_1 \in K$ so that $\mathbf{F}(\mathbf{x}_1, s_1) \in M^+$ and for all $0 \leq s < s_1$, $\mathbf{F}(\mathbf{x}, s)$ is in the past of M^+ . So M_{s_1} is tangent to M^+ at $\mathbf{F}(x_1, s_1)$. Moreover, at (\mathbf{x}_1, s_1) , $H - \mathcal{H} \geq 0$. Hence $H > H^+$ at $\mathbf{F}(\mathbf{x}_1, s_1)$ this contradicts the strong maximum principle. Hence we must have $\tau(\mathbf{F}) \leq \sup \tau|_{M^+}$. One then apply Theorems 1.2, 1.3 to conclude the result. \square

Next, we want to prove a convergence result without assuming $\mathcal{H} > \delta > 0$.

Proof of Theorem 1.4. Since $H - \mathcal{H} \geq 0$ in M_0 , $\overline{\text{Ric}}(w, w) \geq 0$, \mathcal{H} is monotone, as in the proof of Theorem 1.3, one can conclude that $H - \mathcal{H} \geq 0$ for all s .

Since $\mathcal{H} \geq 0$, we have $H \geq H - \mathcal{H} \geq 0$. By Lemma 3.2(vi), we have:

$$\left(\frac{\partial}{\partial s} - \Delta \right) (H - \mathcal{H})^2 \leq -(H - \mathcal{H})^4.$$

Comparing the function s^{-1} and $(H - \mathcal{H})^2$, using Lemmas 3.4 and 3.5, we have

$$(4.8) \quad (H - \mathcal{H})^2(\mathbf{x}, s) \leq s^{-1}$$

for all $\mathbf{x} \in M$. Now we claim that for any compact set $K \subset M$, $\rho(\mathbf{F}(\mathbf{x}, s)) \leq \rho_0$ for some ρ_0 for all $(\mathbf{x}, s) \in K \times [0, \infty)$. Let $u(\mathbf{x}, s) = \tau(\mathbf{F}(\mathbf{x}, s))$ be the height function, then in $K \times [0, \infty)$, by Lemma 3.2(vii), we have

$$\frac{\partial}{\partial s} u = \alpha^{-1} \kappa (H - \mathcal{H}) \geq C_1 \kappa (H - \mathcal{H})$$

for some $C_1 > 0$ because α is bounded from above, and

$$\frac{\partial}{\partial s} \rho(\mathbf{F}(\mathbf{x}, s)) = \langle \bar{\nabla} \rho, \partial_s \mathbf{F} \rangle(\mathbf{x}, s) \leq C_2 \kappa (H - \mathcal{H})$$

for some $C_2 > 0$ by (1.2), the fact that $\|\bar{\nabla} \rho\|_{\Omega_{\tau_1, \tau_2}}$ is finite, and that $H - \mathcal{H} \geq 0$. Hence

$$\frac{\partial}{\partial s} u - \delta \rho \geq 0$$

for some $\delta > 0$ on $K \times [0, \infty)$. Hence

$$u(\mathbf{x}, s) - \delta \rho(\mathbf{F}(\mathbf{x}, s)) \geq u(\mathbf{x}, 0) - \delta \rho(\mathbf{F}(\mathbf{x}, 0)).$$

Since $\tau_1 \leq u \leq \tau_2$, it is easy to see the claim is true. From this one can conclude that $\mathbf{F}(K \times [0, \infty)) \subset W$ for some compact set W in N . As in the proof of Theorem 1.3, since the tilt factor κ and all the covariant derivatives of the second fundamental form of $\mathbf{F}(\cdot, s)$ are uniformly bounded in space and time, using (4.8) one may proceed as in [4, Theorem 3.8] and [8] to conclude that the theorem is true. The idea is to cover W with finitely many suitable simply connected coordinate neighborhoods so that one can apply Lemma B.1. Note that the limit is strictly spacelike because κ is uniformly bounded. \square

Remark 4.1. It is easy to see that Theorems 1.3, 1.4 are true under dual conditions. For example, in Theorem 1.4 if we assume $\mathcal{H} \leq 0$, $H - \mathcal{H} \leq 0$ in M_0 instead, then the result is still true. We can appeal to the theorem by changing the time orientation and replaced \mathcal{H} by $-\mathcal{H}$.

5. EXAMPLES

5.1. Application on Minkowski spacetime. We will apply the previous results to study some prescribed mean curvature flow in Minkowski spacetime. Consider the Minkowski spacetime $\mathbb{R}^{1,n}$ with metric

$$g_{\text{Mink}} = -dt^2 + \sum_{i=1}^n (dx^i)^2.$$

We consider $\frac{\partial}{\partial t}$ as future pointing. Consider the family of spacelike surfaces with constant mean curvature $1/\tau > 0$ given by

$$S_\tau = \{(t, \mathbf{x}) \mid t > 0, t^2 - r^2 = \tau^2\}$$

where $\mathbf{x} = (x^1, \dots, x^n)$ and $r = |\mathbf{x}|$. Then S_τ will foliate $N = I^+(0)$, the future of the origin. We will use τ as time function, see Lemma 5.1 below.

$$\tau_- = \inf_N \tau = 0, \quad \tau_+ = \sup_N \tau = +\infty.$$

As an application of previous results, we want to prove the following:

Proposition 5.1. (i) *A small perturbation of S_τ in a compact set will be deformed under a prescribed mean curvature flow to a constant mean curvature surface with mean curvature $1/\tau$. Namely, if M is a spacelike surface which coincide with S_τ outside a compact set of S_τ so that the mean curvature H of M satisfies $H \geq \varepsilon_0$ for some $\varepsilon_0 > 0$. Then the prescribed mean curvature flow with $\mathcal{H} = \frac{1}{\tau}$ has long time solution and will converge to a complete spacelike hypersurface M_∞ of constant mean curvature $1/\tau$. Moreover, the height function u_∞ will tend to τ at infinity.*

(ii) *Let \mathcal{H} be a smooth function in N which is monotone and satisfies the condition in (1.3). Moreover, suppose $|\mathcal{H} - \frac{1}{\tau_0}| < \frac{1}{2\tau_0}$ on S_{τ_0} and $\sup_{S_{\tau_0}} \phi |H - \mathcal{H}| < \infty$, where $\phi \geq 1$ is an exhaustion function with bounded gradient and Hessian on S_{τ_0} . Then the prescribed mean curvature flow with prescribed mean curvature \mathcal{H} has a long time solution starting from S_{τ_0} and will converge to a complete spacelike hypersurface M_∞ of prescribed mean curvature \mathcal{H} . Moreover, the height function u_∞ will tend to τ at infinity.*

We will give some examples of \mathcal{H} in (ii) after we prove the theorem. First, we need the following lemma.

Lemma 5.1. *We have the following:*

- (i) $\overline{\nabla}\tau = -\frac{1}{\tau}(t\frac{\partial}{\partial t} + r\frac{\partial}{\partial r})$, $\langle \overline{\nabla}\tau, \frac{\partial}{\partial t} \rangle = \frac{t}{\tau}$. Hence $\overline{\nabla}\tau$ is past directed time-like. The lapse function α of τ is 1.
- (ii) Let G_E be the reference Riemannian metric with respect to τ and let $\mathbf{T} = -\overline{\nabla}\tau$. Then $|||\overline{\nabla}\mathbf{T}|||_{\Omega_{\tau_1, \infty}} \leq C\tau_1^{-1}$ and $|||\overline{\nabla}^2\mathbf{T}|||_{\Omega_{\tau_1, \infty}} \leq C\tau_1^{-2}$ for some constant C for all $\tau_1 > 0$.
- (iii) Let $\rho = \log(r+2) > 0$. Then $|||\overline{\nabla}\rho|||_{\Omega_{\tau_1, \infty}} < \infty$ for all $\tau_1 > 0$. Moreover, for any $0 < \tau_1 < \tau_2 < \infty$ and any $\rho_0 > 0$, then set $\overline{\Omega}_{\tau_1, \tau_2} \cap \{\rho \leq \rho_0\}$ is compact.

Proof. (i) follows by direct computation.

(ii) Since $\overline{\nabla}\tau = -\frac{1}{\tau}(t\frac{\partial}{\partial t} + x^i\frac{\partial}{\partial x^i})$, at $(\tau_1, 0)$ the first derivatives and the second derivatives of the coefficients of $\frac{\partial}{\partial t}, \frac{\partial}{\partial x^i}$ are uniformly bounded by $C\tau_1^{-1}$ and $C\tau_1^{-2}$ for some constant C . Since $\overline{\nabla}\tau = -\frac{\partial}{\partial t}$, one can conclude that

$|||\bar{\nabla}\mathbf{T}|||, |||\bar{\nabla}^2\mathbf{T}|||$ are bounded by $C\tau_1^{-1}$ and $C\tau_1^{-2}$ at $(\tau_1, 0)$, for some constant C . Since for any point $p = (t, x^1, \dots, x^n) \in S_{\tau_1}$, there is a restricted Lorentz transformation Φ which is an isometry of N , and maps $(\tau_1, 0)$ to p . Moreover, $\tau \circ \Phi(q) = \tau(q)$ for all q . From this we can conclude that (ii) is true.

(iii) Let G'_E be the reference Riemannian metric with respect to $\frac{\partial}{\partial t}$. Then $|||\bar{\nabla}\rho|||_{G'_E} \leq \frac{1}{r+2}$. By (i) and Lemma 2.1, we have

$$|||\bar{\nabla}\rho||| \leq \frac{2t}{\tau} \cdot \frac{1}{r+2} = \frac{2(\tau^2 + r^2)^{\frac{1}{2}}}{\tau(r+2)} \leq 2\left(1 + \frac{1}{\tau_1}\right).$$

It is easy to see that (iii) is true. □

Proof of Proposition 5.1. (i) By Theorems 1.1, 1.2, 1.3, it is sufficient to show that if \mathbf{F} is a solution of the prescribed mean curvature flow (1.2) with $\mathcal{H} = 1/\tau_0$ starting from M , with $0 \leq s \leq s_0$, so that κ and $|\nabla^m A|$ are uniformly bounded on $M \times [0, s_0]$, then $0 < \tau_1 \leq \tau(\mathbf{F}) \leq \tau_2 < \infty$ for some τ_1, τ_2 independent of s . As in the proof of the last part of Theorem 1.3, the height function u of M_s will tends to τ_0 at infinity uniformly for $s \in [0, s_0]$. Choose $\tau_2 > \tau_0$ and $\tau_2 > \sup_{M_0} u_0$, where u_0 is the height function of $M_0 = M$. Since the mean curvature of $S_{\tau_2} = 1/\tau_2 < 1/\tau_0$, by the maximum principle as in [10, p.605], we can conclude that $\tau(\mathbf{F}) < \tau_2$. Note that τ_2 does not depend on s_0 . Similarly, one can prove that $\tau(\mathbf{F}) \geq \tau_1 > 0$ where τ_1 does not depend on s_0 . From this the result follows.

(ii) By the assumption, we have $\mathcal{H} > \frac{1}{2\tau_0}$ and $|H - \mathcal{H}| \leq \frac{1}{2\tau_0} - \varepsilon_0$ for some $\varepsilon_0 > 0$. Hence the proof is similar to the proof of (i). □

Example: In the Minkowski space, let $f(t, \mathbf{x}) = e^{-4t+(r^2+1)^{\frac{1}{2}}}$ where $r = |\mathbf{x}|$. We want to check that $\mathcal{H}(t, \mathbf{x}) = 2 - f(\mathbf{x}, t)$ satisfies the assumptions in Proposition 5.1(ii) with $\tau_0 = \frac{1}{2}$. We use $(r^2 + 1)^{\frac{1}{2}}$ because r is not a smooth function. Let us first check that $\sup_{S_{\frac{1}{2}}} |\mathcal{H} - 2| < 1$ on $S_{\frac{1}{2}}$. On $S_{\frac{1}{2}}$, $t^2 = r^2 + \frac{1}{4}$ and $t \geq r, t \geq \frac{1}{2}$ and so $f \leq e^{-\frac{1}{2}} < 1$. Hence $\sup_{S_{\frac{1}{2}}} |\mathcal{H} - 2| < 1$. Next we want to check that \mathcal{H} is monotone. Let w be a future directed timelike vector so that $w = a_0 \frac{\partial}{\partial t} + \sum_{i=1}^n a_i \frac{\partial}{\partial x^i}$. Then $a_0 \geq |\mathbf{a}|$ where $\mathbf{a} = (a_1, \dots, a_n)$. Now $\bar{\nabla}\mathcal{H} = f \left(-4 \frac{\partial}{\partial t} - (1 + r^2)^{-\frac{1}{2}} \sum_{i=1}^n x^i \frac{\partial}{\partial x^i} \right)$. Hence

$$\langle \bar{\nabla}\mathcal{H}, w \rangle = f \left(4a_0 - (1 + r^2)^{-\frac{1}{2}} \sum_{i=1}^n a_i x^i \right) \geq f \left(4a_0 - r(1 + r^2)^{-\frac{1}{2}} |\mathbf{a}| \right) \geq 0.$$

Next, we want to show that $|||\bar{\nabla}^k \mathcal{H}|||_{\Omega_{\tau_1, \tau_2}} < \infty$ for any $0 < \tau_1 < \tau_2 < \infty$. It is easy to see that \mathcal{H} is uniformly bounded in N . Let G_e be the Euclidean

metric $dt^2 + \sum_{i=1}^n (dx^i)^2$. For any $k \geq 1$

$$|\bar{\nabla}^k \mathcal{H}|_{G_e} \leq Cf$$

for some constant C depending only on k, n . By Lemmas 2.1 , 5.1 , we have

$$|||\bar{\nabla}^k \mathcal{H}||| \leq Cf \cdot \frac{t^k}{r^k} \leq C' \tau_1^{-k}$$

for some constant C' . Here we have used the fact that $t \geq r$ in Ω_{τ_1, τ_2} . Finally, we claim that $d(p)|H - \mathcal{H}|(p) \leq C''$ on $S_{\frac{1}{2}}$ for some constant C'' where $d(p)$ is the intrinsic distance of $S_{\frac{1}{2}}$ from a fixed point. At a point $p = (t, \mathbf{x})$ in $S_{\frac{1}{2}}$, one can check that $d(p) \sim \log r$, where $r = |\mathbf{x}|$. From this it is easy to see the claim is true. Since the distance function on $S_{\frac{1}{2}}$ has bounded gradient and Hessian away from $d = 0$, hence all the conditions in Proposition 5.1 are satisfied and we obtain a complete spacelike hypersurface in N with prescribed mean curvature given by \mathcal{H} .

5.2. Computation near the future null infinity of the Schwarzschild spacetime. We want to do some computations which might be related to the study of prescribed mean curvature flow in a asymptotically Schwarzschild spacetime introduced in [1]. Let us first recall the future null infinity of the Schwarzschild spacetime. The standard Schwarzschild metric in $r > 2m > 0$ is

$$(5.1) \quad g_{\text{Sch}} = - \left(1 - \frac{2m}{r}\right) dt^2 + \left(1 - \frac{2m}{r}\right)^{-1} dr^2 + r^2 \sigma.$$

defined on $\{r > 2m\}$, $-\infty < t < \infty$ where $r = \sum_{i=1}^3 (x^i)^2$ with $(x^1, x^2, x^3) \in \mathbb{R}^3$ and σ is the standard metric of the unit sphere \mathbb{S}^2 . Consider the retarded null coordinate

$$(5.2) \quad v = t - r_*$$

where $r_* = r + 2m \log\left(\frac{r}{2m} - 1\right)$ in the region $r > 2m$. Let $x = r^{-1}$,

$$(5.3) \quad \begin{aligned} g &= g_{\text{Sch}} = - (1 - 2mx) dv^2 + 2x^{-2} dv dx + x^{-2} \sigma \\ &=: x^{-2} \tilde{g}, \end{aligned}$$

with $0 < x < \frac{1}{2m}$, $-\infty < v < \infty$. Here the unphysical metric \tilde{g} is the product metric:

$$(5.4) \quad \tilde{g} = (\sigma_{AB}) \oplus \begin{pmatrix} 0 & 1 \\ 1 & -x^2(1 - 2mx) \end{pmatrix}.$$

Here (σ_{AB}) is the standard metric for \mathbb{S}^2 in local coordinates y^1, y^2 . \tilde{g} can be extended smoothly is Lorentz up to $x = 0$. Then future null infinity \mathcal{I}^+ can be identified as $x = 0$ in the compactification. In the following, let $h = 1 - \frac{2m}{r} =$

$1 - 2mx$. Given a smooth function $f(\mathbf{y})$ on \mathbb{S}^2 . Consider the cut \mathcal{C} given by $(\mathbf{y}, f(\mathbf{y})), \mathbf{y} \in \mathbb{S}^2$ in \mathcal{I}^+ . For $\tau > 0$, let

$$(5.5) \quad P(\mathbf{y}, x, \tau) = f(\mathbf{y}) + x\phi(\tau, \mathbf{y}) + \frac{1}{2!}x^2\psi(\tau, \mathbf{y}),$$

where $\phi = P_x, \psi = P_{xx}$ at $s = 0$ are smooth functions in τ, \mathbf{y} , given by

$$(5.6) \quad \begin{cases} \phi = -\frac{1}{2} \left(\tau^2 + |\tilde{\nabla} f|_{\mathbb{S}^2}^2 \right); \\ \psi = \frac{1}{2} \left(\tau^2 \tilde{\Delta} f + \langle \tilde{\nabla} |\tilde{\nabla} f|_{\mathbb{S}^2}^2, \tilde{\nabla} f \rangle_{\mathbb{S}^2} \right). \end{cases}$$

Here $\tilde{\nabla}, \tilde{\Delta}$ are the covariant derivative and the Laplacian of the standard \mathbb{S}^2 . By [23, Theorem 3.1], so that if Σ_τ is the surface given by $(\mathbf{y}, x) \rightarrow (\mathbf{y}, x, -P)$ in the \mathbf{y}, x, v coordinates, then Σ_τ is spacelike near $x = 0$. Let $0 < \tau_1 < \tau_2 < \infty$ be fixed. Let

$$M = \left\{ \mathbf{y} \in \mathbb{S}^2, x \in \left(0, \frac{1}{2m}\right), \tau \in (\tau_1, \tau_2) \right\} = \mathbb{S}^2 \times (0, s_0) \times (\tau_1, \tau_2).$$

Consider the map Φ from M to the Schwarzschild spacetime in \mathbf{y}, x, v coordinates defined by:

$$(5.7) \quad \Phi(\mathbf{y}, x, \tau) = (\mathbf{y}, x, v(\mathbf{y}, x, \tau))$$

with $v(\mathbf{y}, x, \tau) = -P(\mathbf{y}, x, \tau)$. Then there is $s_0 > 0$ such that Φ is diffeomorphic to its image. Hence τ can be considered a function in the image. We also have $\bar{\nabla}\tau$ is past directed time like vector field if $x_0 > 0$ is small enough. See [28]. Let $\mathbf{T} = -\alpha\bar{\nabla}\tau$, where α is the lapse function of τ . We have the following facts which are related to the assumptions in the results in previous sections.

Proposition 5.2. *In the above setting, let $N = \Phi(M)$ and g_{Sch} be the Schwarzschild metric and let Θ be the reference Riemannian metric with respect to τ . Denote the norm with respect to Θ by $||| \cdot |||$. Then there exists $x_0 > 0$ such that the following are true.*

- (i) $|||\bar{\nabla}\mathbf{T}|||_{1,N}$ is finite.
- (ii) α, α^{-1} and $|||\bar{\nabla}\alpha|||$ are uniformly bounded.
- (iii) $|||\bar{\nabla}\log r|||$ is uniformly bounded.
- (iv) The second fundamental form A and its covariant derivatives of the level set $\tau = \text{constant}$ are uniformly bounded.
- (v) $|||\overline{\text{Rm}}|||_{k,N}$ is finite for all $k \geq 0$.
- (vi) The mean curvature of the level set $\tau = \text{constant}$ satisfies: $H = \frac{1}{\tau} + O(x^2)$ and that $d(p)|H(p) - \frac{1}{\tau} \rightarrow 0$ as $p \rightarrow \infty$ on $\tau = \text{constant}$, where $d(p)$ is the intrinsic distance from a fixed point on the level surface.

Proof. (i) follows from [28, Lemmas 3.7, 3.8].

(ii) follows from [28, Lemma 3.2].

(iii) On the other hand, let $\tilde{\mathbf{T}} = h^{-\frac{1}{2}} \frac{\partial}{\partial t}$ in the standard coordinates t, x^i for the exterior of the Schwarzschild spacetime. Let G'_E be the reference Riemannian metric with respect to t . One can compute by (ii):

$$(5.8) \quad \langle \mathbf{T}, \tilde{\mathbf{T}} \rangle_G = \alpha h^{-1} \langle \bar{\nabla} \tau, \frac{\partial}{\partial t} \rangle_G = -\alpha h^{-1} P_\tau^{-1} = O(x^{-1})$$

near $x = 0$. From this it is easy to see that (iii) is true.

(iv) The bound of A follows from the [28, Lemma 3.4]. From the proof of this fact, one can also show that $\nabla^k A$ is also uniformly bounded for all $k \geq 1$.

(v) To prove $|||\overline{\text{Rm}}|||$ is bounded, we need to prove that $\overline{\text{Rm}}$ is bounded when evaluated at an orthonormal basis w_1, w_2, w_3, w_4 where $w_4 = \mathbf{T}$ and w_1, w_2, w_3 are tangential to the level surface of τ . Note that one cannot just appeal to Lemma 2.1 to conclude that $|||\overline{\text{Rm}}|||$ is bounded because $|||\overline{\text{Rm}}|||_{G'_E}$ decays like x^3 , $\overline{\text{Rm}}$ is a four tensor, and we only have (5.8). Let us choose w_i , $1 \leq i \leq 3$ as follows. Let y^1, y^2 be local coordinates of \mathbb{S}^2 , and let Φ be the diffeomorphism in (5.7):

$$(5.9) \quad \begin{cases} e_A =: \Phi_* \left(\frac{\partial}{\partial y^A} \right) = -P_A \partial_v + \partial_A, A = 1, 2; \\ e_3 =: \Phi_* \left(\frac{\partial}{\partial x} \right) = -P_x \partial_v + \partial_x; \\ e_4 =: \Phi_* \left(\frac{\partial}{\partial \tau} \right) = -P_\tau \partial_v. \end{cases}$$

Then e_1, e_2, e_3 are tangential to the level set of τ . We have

$$(5.10) \quad \begin{cases} \langle e_A, \tilde{\mathbf{T}} \rangle = -P_A h^{\frac{1}{2}} & A = 1, 2; \\ \langle e_3, \tilde{\mathbf{T}} \rangle = -P_x h^{\frac{1}{2}}. \end{cases}$$

Since $\langle e_i, e_i \rangle = P_i^2 + r^2 = P_i^2 + x^{-2}$, for $i = 1, 2, 3$ we have

$$|||e_i|||_{G'_E}^2 = \langle e_i, e_i \rangle + 2(\langle e_i, \tilde{\mathbf{T}} \rangle)^2 \leq Cx^{-2}.$$

We may obtain orthonormal basis w_i from e_i so that $w_i = xc_{ij}e_j$, with c_{ij} being bounded, see [28, Lemma 3.1]. $|||w_i|||_{G'_E} \leq C$ for some constant C . From (5.8), we also have $|||w_4|||_{G'_E} = |||\mathbf{T}|||_{G_E} \leq Cx^{-1}$. To summarise:

$$(5.11) \quad \begin{cases} |||w_i|||_{G'_E} \leq C, & \text{for } 1 \leq i \leq 3; \\ |||w_4|||_{G'_E} \leq Cx^{-1}, \end{cases}$$

for some constant $C > 0$, provided s_0 is small enough. Now we are ready to estimate $|||\overline{\text{Rm}}|||$. Since $\overline{\text{Rm}}(w_4, w_4, \cdot, \cdot) = 0$, by (5.11), we have

$$|\overline{\text{Rm}}(w_a, w_b, w_c, w_d)| \leq |||\overline{\text{Rm}}|||_{G'_E} |||w_a||| |||w_b||| |||w_c||| |||w_d||| \leq Cx.$$

because $|||\overline{\text{Rm}}|||_{G'_E} \leq Cx^3$. Therefore, $|||\overline{\text{Rm}}||| \leq Cx$. Similarly, one can conclude that $|||\overline{\nabla}^k \overline{\text{Rm}}||| \leq Cx$.

(vi) By [23, Lemma 2.1], The fact that $|H - \frac{1}{\tau}| = O(x^2) = O(r^{-2})$ follows from [23, Theorem 2.1] and the definition of \bar{P} . Since $\tau = \text{constant}$ can be written as a spacelike graph over \mathbf{x} in the standard coordinates t, x^i , as before one can see the last assertion is true. \square

APPENDIX A. HÖLDER SPACES

We begin by explicitly defining the local Hölder norms used in the definition parabolic Hölder spaces. Let Ω be an open set in \mathbb{R}^m . Let $k \geq 0$ be an integer and $0 < \sigma < 1$, for $s_0 > 0$, the $C^{2k+\sigma, k+\frac{\sigma}{2}}$ norm on $\Omega_{s_0} = \Omega \times [0, s_0]$ for functions $f(\mathbf{x}, s)$ is defined as:

$$\begin{aligned} \|f\|_{2k+\sigma, k+\frac{\sigma}{2}; \Omega_{s_0}} &= \sum_{|\alpha|+2r=0}^{2k} \sup_{\Omega} |\partial_s^r \partial^\alpha f| \\ &+ \sum_{|\alpha|+2r=2k} \sup_{(\mathbf{x}, s) \neq (\mathbf{x}', s') \in \Omega_{s_0}} \frac{|\partial_s^r \partial^\alpha f(\mathbf{x}, s) - \partial_s^r \partial^\alpha f(\mathbf{x}', s')|}{|\mathbf{x} - \mathbf{x}'|^\sigma + |s - s'|^{\frac{\sigma}{2}}} \end{aligned}$$

where

$$\partial_s^r \partial^\alpha u = \frac{\partial^{r+|\alpha|} u}{\partial s^r \partial x^\alpha}$$

and α is a multi-index. If there is no confusion on the domain of definition of f , we will simply write the norm as $\|f\|_{2k+\sigma, k+\frac{\sigma}{2}}$.

APPENDIX B. A CONVERGENCE LEMMA

Let U be a coordinate neighborhood of a Lorentz manifold with time function τ , $\overline{\nabla}\tau$ is past directed, so that U is given by

$$U = (-2, 2) \times B(2) = \{(x^0, x^1, \dots, x^n) \mid |x^a| < 2, 0 \leq a \leq n\}$$

where $\tau = x^0$. Let

$$V = (-1, 1) \times B(1) = \{(x^0, x^1, \dots, x^n) \mid |x^a| < 1, 0 \leq a \leq n\}$$

Moreover the metric is of the form $G = -\alpha^2 d\tau^2 + g_{ij} dx^i dx^j$, which is zero shift. Assume the lapse function α is uniformly bounded from above and below. We also assume that g_{ij} is uniformly equivalent to the Euclidean metric δ_{ij} .

Lemma B.1. *Let M_k be a sequence of spacelike surfaces so that*

- (i) $M_k \cap U$ is connected, $\partial M_k \cap U = \emptyset$;
- (ii) the mean curvature H_k of M_k together with all its derivatives are bounded; and
- (iii) the tilt factor κ_k of M_k with respect to τ are uniformly bounded.

If there is a limit point p of M_k inside V , then there is a subsequence of M_k , still denoted by M_k and a spacelike hypersurface M_∞ inside V so that $d_h(M_k, M_\infty) \rightarrow 0$ in every compact subsets of V . Moreover, the mean curvature H_∞ is such that $\lim_{k \rightarrow \infty} H_k = H_\infty$.

Proof. First we may assume that $H_k \rightarrow H_\infty$ in C^∞ norm in any compact set of U . Since U is simply connected, M_k is achronal [25, p.427]. Hence each M_k is a graph. Namely, for each k , there is a connected open set $W_k \subset U$ and a smooth function u_k such that

$$M_k = \{(u_k(\mathbf{x}), \mathbf{x}) \mid \mathbf{x} \in W_k\}$$

where $\mathbf{x} = (x^1, \dots, x^n)$. Since M_k is spacelike, there is $C_1 > 0$ such that $|Du_k| \leq C_1$ for all k , where Du_k is the gradient with respect to the Euclidean metric. Hence there is $\delta > 0$ independent of \mathbf{x} such that if $(u_k(\mathbf{x}), \mathbf{x}) \in M_k \cap V$, with $|u_k(\mathbf{x})| \leq 1 - \eta$ for some $\eta > 0$. Then $B_{\mathbf{x}}(\delta) \subset B_{\mathbf{x}}(2\delta) \subset W_k$ and $|u_k(\mathbf{y})| < 1$ for $\mathbf{y} \in B_{\mathbf{x}}(\delta)$. Here $B_{\mathbf{x}}(\delta) = \{\mathbf{y} \mid |y^i - x^i| < \delta, 1 \leq i \leq n\}$. Since the tilt factors of M_k are uniformly bounded and $H_k \rightarrow H_\infty$, the equation satisfied by u_k is uniformly elliptic in $B_{\mathbf{x}}(\delta)$. Hence if p is a limit point of M_k inside V , passing to a subsequence, u_k will converge in C^∞ norm on compact sets of $B_{\mathbf{x}}(\delta)$ where \mathbf{x} is the projection of p to $B(2)$. From this it is easy to see the lemma is true. \square

REFERENCES

- [1] Andersson, L.; Iriondo, M. S., *Existence of constant mean curvature hypersurfaces in asymptotically flat spacetimes*, Ann. Global Anal. Geom. **17** (1999), no. 6, 503–538.
- [2] Andrews, B.; Chow, B.; Guenther, C.; Langford, M., *Extrinsic geometric flows*. Graduate Studies in Mathematics, 206. American Mathematical Society, Providence, RI, [2020], ©2020. xxviii+759 pp. ISBN: 978-1-4704-5596-5
- [3] Bartnik, R., *Existence of maximal surfaces in asymptotically flat spacetimes*, Comm. Math. Phys. **94** (1984), no. 2, 155–175.
- [4] Bartnik, R., *Regularity of variational maximal surfaces*, Acta Math. **161** (1988), no. 3–4, 145–181.
- [5] Bartnik, R.; Simon, L., *Space-like hypersurfaces with prescribed boundary values and mean curvature*, Commun. Math. Phys. **87** (1982) 131–152.
- [6] Chau, A.; Chen, J.; He, W., *Lagrangian Mean Curvature flow for entire Lipschitz graphs*, Calc. Var. **44** (2012), 199–220.
- [7] Chau, A.; Tam, L.-F., *On a modified parabolic complex Monge-Ampère equation with applications*, Math. Z. **269** (2011), no. 3–4, 777–800.
- [8] Ecker, K., *On mean curvature flow of spacelike hypersurfaces in asymptotically flat spacetimes*, J. Austral. Math. Soc. Ser. A **55** (1993), no. 1, 41–59.
- [9] Ecker, K., *Interior estimates and longtime solutions for mean curvature flow of non-compact spacelike hypersurfaces in Minkowski space*, J. Differential Geom. **46** (1997), no. 3, 481–498.
- [10] Ecker, K.; Huisken, G., *Parabolic methods for the construction of spacelike slices of prescribed mean curvature in cosmological spacetimes* Comm. Math. Phys. **135** (1991), no. 3, 595–613.
- [11] A. Friedman, A., *Partial differential equations of parabolic type*, Englewood Cliffs, N.J., : Prentice-Hall (1964).
- [12] Gentile, G.; Vertman, B., *Prescribed mean curvature flow of non-compact space-like Cauchy hypersurfaces*, Ann. Global Anal. Geom. **64** (2023), no. 2, Paper No. 11, 45 pp.
- [13] Gerhardt, C., *Hypersurfaces of prescribed mean curvature in Lorentzian manifolds*, Math. Z. **235** (2000), no. 1, 83–97.
- [14] Gerhardt, C., *H-surfaces in Lorentzian manifolds*, Comm. Math. Phys. **89** (1983), no. 4, 523–553.
- [15] Gerhardt, C., *H-surfaces in Lorentzian manifolds*, Comm. Math. Phys. **89** (1983), no. 4, 523–553. 2
- [16] Gilbarg, D; Trudinger, N.S., *Elliptic partial differential equations of second order*, 2nd ed., rev. 3rd printing. Berlin ; New York : Springer c1998.

- [17] Hamilton, R.S., *Three-manifolds with positive Ricci curvature*, J. Differ. Geom. **17**(1982), 255–306.
- [18] Hamilton, R. S., *A compactness property for solutions of the Ricci flow*, Amer. J. Math. **117** (1995), no. 3, 545–572.
- [19] Hawking, S.; Ellis, G., *The large-scale structure of spacetime*, Cambridge: Cambridge University Press 1973.
- [20] Husiken, G., *Flow by mean curvature of convex surfaces into spheres*, J. Differential Geom. **20** (1984), no. 1, 237–266.
- [21] Husiken, G., *The volume preserving mean curvature flow*, J. Reine Angew. Math. **382** (1987), 35–48.
- [22] Kröcke, K.; Petersen, O.L.; Lubbe, F.; Marxen, T.; Maurer, W.; Meiser, W.; Schnürer, O.C.; Szabó, Á.; Vertman, B., *Mean curvature flow in asymptotically flat product spacetimes*, J. Geom. Anal. **31**(6), 5451–5479 (2021)
- [23] Li, C.; Shi, Y.-G.; Tam, L.-F., *Boundary behaviors of spacelike constant mean curvature surfaces in Schwarzschild spacetime*, arXiv:2202.00926.
- [24] Lieberman, G. M., *Second order parabolic differential equations*, World Scientific Publishing Co., Inc., River Edge, NJ, 1996.
- [25] O’Neill, B., *Semi-Riemannian geometry : with applications to relativity*, New York : Academic Press 1983.
- [26] Shi, W.-X., *Ricci Flow and the uniformization on complete non compact Kähler manifolds*, J. of Differential Geometry **45** (1997), 94-220.
- [27] Tam, L.-F., *Exhaustion functions on complete manifolds*, Recent advances in geometric analysis, 211–215, Adv. Lect. Math. (ALM), **11**, Int. Press, Somerville, MA, 2010.
- [28] Tam, L.-F., *Spacelike CMC surfaces near null infinity of the Schwarzschild spacetime*, arXiv:2306.03353
- [29] Treibergs, A. E., *Entire spacelike hypersurfaces of constant mean curvature in Minkowski space*, Invent. Math. **66** (1982), no. 1, 39–56.

(Luen-Fai Tam) THE INSTITUTE OF MATHEMATICAL SCIENCES AND DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN, HONG KONG, CHINA.

Email address: lftam@math.cuhk.edu.hk