On k-submodular subgroups of finite groups

T.I. Vasilyeva

Abstract

Let n be a natural number. A subgroup H of a group G will be called nmodularly embedded in G if either H is normal in G or $H \neq \operatorname{Core}_G(H)$, |G:H| = pand $|G/\operatorname{Core}_G(H)| = pq^n$, q^n divides p-1 for some primes p and q. Let k be a fixed natural number. A subgroup H of a group G will be called k-submodular in G if there exists a chain $H = H_0 \leq H_1 \leq \cdots \leq H_{m-1} \leq H_m = G$ of subgroups such that H_{i-1} is n-modularly embedded in H_i for some natural $n \leq k$ and every $i = 1, \ldots, m$. In this article, properties of k-submodular subgroups and classes of groups with given systems of k-submodular subgroups are obtained.

Keywords: *n*-modularly embedded subgroup, *k*-submodular subgroup, Schunck class, formation, supersoluble group

MSC2010 20D10, 20E15, 20F16

Introduction

All groups under consideration are finite. In a nilpotent group, every subgroup is subnormal. From Huppert's result [1, Chap. VI, Theorem 9.2 (b)] it follows that if the group G is supersoluble, then every its proper subgroup can be connected to G by a chain of subgroups of prime indices. Schmidt proved in [2] that the subgroup M of the group G is a maximal modular subgroup in G if and only if either M is maximal normal subgroup of G, or $G/\text{Core}_G(M)$ is a non-abelian group of order pq for some primes p and q. We introduce the following

Definition 1. Let n be a natural number. A subgroup H of a group G will be called n-modularly embedded in G if either $H \leq G$ or $H \neq \operatorname{Core}_G(H)$, |G:H| = p and $|G/\operatorname{Core}_G(H)| = pq^n$, q^n divides p-1 for some primes p and q.

Definition 2. Let k be a fixed natural number. A subgroup H of a group G will be called k-submodular in G if there exists a chain

$$H = H_0 \le H_1 \le \dots \le H_{m-1} \le H_m = G \tag{1.1}$$

of subgroups such that H_{i-1} is n-modularly embedded in H_i for some natural $n \leq k$ and every $i = 1, \ldots, m$.

Note if n = 1 and a subgroup in G is n-modularly embedded, then it is modular in G. If k = n = 1 and a subgroup in G is k-submodular, then it is submodular in G. Groups with given systems of submodular subgroups were studied in [3]–[8]. Groups with \mathfrak{U}_k -subnormal subgroups were studied in [9]. Here \mathfrak{U}_k is the class of all supersoluble groups in which exponents are not divided by the (k+1)th powers of primes. If G is a soluble group, then an k-submodular in G subgroup is \mathfrak{U}_k -subnormal in G (Lemma 2.7). As follows from Example 3.1, the converse statement does not hold in the general case.

In this article, properties of k-submodular subgroups of groups and classes of groups with given systems of k-submodular subgroups are obtained.

1. Preliminary results

We use the notation and terminology from [10, 11]. We recall some concepts significant in the paper.

Let G be a group. If H is a subgroup of G, we write $H \leq G$ and if $H \neq G$, we write H < G. We denote by |G| the order of G; by $\pi(G)$ the set of all distinct prime divisors of the order of G; by $\operatorname{Syl}_p(G)$ the set of all Sylow p-subgroups of G; by $\operatorname{Syl}(G)$ the set of all Sylow subgroups of G; by F(G) the Fitting subgroup of G; by |G : H| the index of H in G; by $\pi(G : H)$ the set of all different prime divisors of |G : H|; by Z_n the cyclic group of order n; by \mathbb{P} the set of all primes; by \mathfrak{S} the class of all soluble groups; by \mathfrak{A} the class of all supersoluble groups; by \mathfrak{N} the class of all nilpotent groups; by \mathfrak{N}_p the class of all p-groups for $p \in \mathbb{P}$; by $\mathfrak{A}(p-1)$ the class of all abelian groups of exponent dividing p-1.

A group G of order $p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_n^{\alpha_n}$ (where p_i is a prime, $i = 1, 2, \ldots, n$) is called Ore dispersive or Sylow tower group whenever $p_1 > p_2 > \cdots > p_n$ and G has a normal subgroup of order $p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_i^{\alpha_i}$ for every $i = 1, 2, \ldots, n$.

A class of groups \mathfrak{F} is called a *formation* if the following conditions hold: (a) every quotient group of a group lying in \mathfrak{F} also lies in \mathfrak{F} ; (b) if $G/N_i \in \mathfrak{F}$, $N_i \leq G$, i = 1, 2 then $G/N_1 \cap N_2 \in \mathfrak{F}$. A formation \mathfrak{F} is called *hereditary* whenever \mathfrak{F} together with every group contains all its subgroups, and *saturated*, if $G/\Phi(G) \in \mathfrak{F}$ implies that $G \in \mathfrak{F}$.

A subgroup H of a group G is said to be:

 \mathbb{P} -subnormal in G [12] if either H = G or there exists a chain of subgroups (1.1) such that $|H_i: H_{i-1}|$ is a prime for every $i = 1, \ldots, m$;

K- \mathbb{P} -subnormal in G [13] if there exists a chain of subgroups (1.1) such that either $H_{i-1} \leq H_i$ or $|H_i: H_{i-1}|$ is a prime for every $i = 1, \ldots, m$.

Let \mathfrak{F} be a non-empty formation. A subgroup H of a group G is said to be:

 \mathfrak{F} -subnormal in G [11] if either H = G or there exists a chain of subgroups (1.1) such that $H_i^{\mathfrak{F}} \leq H_{i-1}$ for every $i = 1, \ldots, m$;

K- \mathfrak{F} -subnormal in G [11] if there exists a chain of subgroups (1.1) such that either $H_{i-1} \leq H_i$ or $H_i^{\mathfrak{F}} \leq H_{i-1}$ for every $i = 1, \ldots, m$.

Here $G^{\mathfrak{F}}$ is the least normal subgroup of G for which $G/G^{\mathfrak{F}} \in \mathfrak{F}$.

A function $f : \mathbb{P} \to \{\text{formations}\}\$ is called a *formation function*. A formation \mathfrak{F} is called *local*, if there exists a formation function f such that $\mathfrak{F} = LF(f) = (G|G/C_G(H/K) \in f(p) \text{ for every chief factor } H/K \text{ of a group } G \text{ and for all primes } p \in \pi(H/K)).$

Lemma 1.1 [14, Chap. I, Theorem 1.4]. Let H/K be a chief p-factor of a group G. |H/K| = p if and only if $\operatorname{Aut}_G(H/K)$ is abelian group of exponent dividing p-1.

Lemma 1.2 [1, Chap. VI, Theorem 9.5]. A group G is supersoluble if and only if all its maximal subgroups in G have a prime index.

Lemma 1.3 [10, Chap. IV, Example 3.4(f), Chap. VII, Theorem 2.1]. (1) If $G \in \mathfrak{U}$, then G is Ore dispersive and has nilpotent commutator subgroup.

(2) The class \mathfrak{U} is a hereditary saturated formation and has a formation function f such that $f(p) = \mathfrak{A}(p-1)$ for every prime p.

Lemma 1.4 [15, Theorem 2]. A group G is supersoluble if and only if G can be represented as the product of two nilpotent \mathbb{P} -subnormal subgroups.

According to [12], w \mathfrak{U} denotes the class of all groups in which every Sylow subgroup is \mathbb{P} -subnormal.

Lemma 1.5. (1) If $G \in \mathfrak{wU}$, then G is Ore dispersive [12, Proposition 2.8].

(2) The class will is a hereditary saturated formation and it has a formation function f such that $f(p) = (G \in \mathfrak{S}|Syl(G) \subseteq \mathfrak{A}(p-1))$ for every prime p [12, Theorems 2.7, 2.10].

In [9] \mathfrak{U}_k denotes the class of all supersoluble groups in which exponents are not divided by the (k+1)th powers of primes, where k is a natural number; w \mathfrak{U}_k denotes the class of all groups in which every Sylow subgroup is \mathfrak{U}_k -subnormal.

Lemma 1.6. (1) The class \mathfrak{U}_k is a hereditary formation. [9, Lemma 3(2)].

(2) The class \mathfrak{wL}_k is a hereditary saturated formation. [9, Proposition 1].

(3) If $G \in \mathfrak{wU}_k$, then $G/F(G) \in \mathcal{A}_k$ [9, Corollary 1].

Here \mathcal{A}_k is the class of all groups with abelian Sylow subgroups in which exponents are not divided by the (k + 1)th powers of primes.

2. Properties of k-submodular subgroups of groups

Lemma 2.1. Let n be a natural number. Let H be a subgroup of a group G and $N \leq G$. If $N \leq H$ and H/N is n-modularly embedded in G/N, then H is n-modularly embedded in G.

Proof. Assume that H/N is n-modularly embedded in G/N and write $L = \operatorname{Core}_G(H)$. Then $N \leq L$. If H = L, then $H \leq G$. Let $H \neq L$. Since $L/N = \operatorname{Core}_{G/N}(H/N)$, we have $H/N \neq L/N$, |G:H| = |G/N:H/N| = p, $|G/L| = |G/N/L/N| = pq^n$, q^n divides p-1 for some primes p and q. Therefore H is n-modularly embedded in G. \Box

Lemma 2.2. Let n be a natural number and M be a maximal subgroup of a group G. Then the following statements are equivalent.

(1) M is n-modularly embedded in G.

(2) Either $M \leq G$ or $G/\operatorname{Core}_G(M)$ is non-nilpotent, $|G/\operatorname{Core}_G(M)| = pq^n$, q^n divides p-1 for prime p and q, and $G/\operatorname{Core}_G(M)$ is a group with a normal Sylow p-subgroup and a cyclic Sylow q-subgroup $M/\operatorname{Core}_G(M)$.

Proof. Proof. Assume that statement (1) holds. Let $D = \operatorname{Core}_G(M)$. If M = D, then $M \trianglelefteq G$. Let's assume that $M \neq D$. Then G/D is non-nilpotent and $|G/D| = pq^n$ by Statement (1). Since G/D is biprimary, by Burnside's theorem G/D is solvable. By [10, Chap. A, Theorem 15.6] $G/D = N/D \cdot M/D$, where N/D — minimal normal subgroup in G/D, $(N/D) \cap (M/D) = 1$, $N/D = C_{G/D}(N/D)$. It follows that |N/D| = |G:M| = p. Then

$$M/D \cong G/D/C_{G/D}(N/D)$$

is isomorphic to a subgroup of $\operatorname{Aut}(Z_p) \cong Z_{p-1}$. Therefore p does not divide |M/D|, M/D is a cyclic Sylow q-subgroup of G/D, $|M/D| = q^n$ divides p-1 and $N/D \in \operatorname{Syl}_p(G/D)$. So, statement (1) implies statement (2). If assertion (2) holds, then it is easy to verify that M is *n*-modularly embedded in G. Thus, statement (2) implies statement (1).

Лемма 2.3. Let k be a natural number.

(1) If every maximal subgroup of a group G is n-modularly embedded in G for some natural number $n, n \leq k$, then G is supersoluble.

(2) If M is a maximal subgroup of a group G and M is n-modularly embedded in G, then M is \mathbb{P} -subnormal and K- \mathbb{P} -subnormal in G, n is a natural number.

Proof. (1) Let W be any maximal subgroup of G. If $W = \operatorname{Core}_G(W)$, then $W \leq G$. Then G = WP for some $P \in \operatorname{Syl}_p(G)$ and $p \in \pi(G : W)$. From the maximality of W in G it follows that |G : W| = p. Let's assume that $W \neq \operatorname{Core}_G(W)$. Then by Definition 1 |G : W| is a prime number. By Lemma 1.2 G is supersoluble.

(2) The statement follows from proof (1) and Definition 1.

Lemma 2.4. Let k be a natural number and let H be a subgroup of a group G.

(1) If H is k-submodular in G, then H^x is k-submodular in G for all $x \in G$.

(2) If $H \leq R \leq G$, H is k-submodular in R and R is k-submodular in G, then H is k-submodular in G.

Proof. Statements (1) and (2) follow from the definition 2.

Lemma 2.5. Let k be a natural number. Let H and U be subgroups of a group G.

(1) If H is k-submodular in G, then $H \cap U$ is k-submodular in U.

(2) If H is k-submodular in G and U is k-submodular in G, then $H \cap U$ is k-submodular in G.

Proof. (1) Suppose that H is k-submodular in G. We can assume that $H \neq G$. There is a chain (1.1) of subgroups such that H_{i-1} is n-modularly embedded in H_i for some natural $n \leq k$ and every $i = 1, \ldots, m$. Consider the chain of subgroups

$$H \cap U = H_0 \cap U \le H_1 \cap U \le \dots \le H_{m-1} \cap U \le H_m \cap U = U.$$

If $H_{i-1} \cap U \leq H_i \cap U$ for i = 1, ..., m, then $H \cap U$ is k-submodular in U.

Let's assume that $H_{j-1} \cap U \not \trianglelefteq H_j \cap U$ for some $j \in \{1, \ldots, m\}$. In that case $H_{j-1} \cap U \neq H_j \cap U$ and $H_{j-1} \not \trianglelefteq H_j$. Let us denote $L = \operatorname{Core}_{H_j}(H_{j-1})$. Since H_{j-1} is *n*-modularly embedded in H_j for some natural $n \leq k$, we have $|H_j : H_{j-1}| = p$ and $|H_j/L| = pq^n$, q^n divides p-1 for some primes p and q. Let $P \in \operatorname{Syl}_p(H_j)$. Then $PL/L \in \operatorname{Syl}_p(H_j/L)$. From Lemma 2.2. it follows that $H_j/L = PL/L \cdot H_{j-1}/L$, $PL/L \trianglelefteq H_j/L$, $H_{j-1}/L \in \operatorname{Syl}_q(H_j/L)$ and H_{j-1}/L is a cyclic subgroup. We have two cases.

1. Let's assume that p does not divide $|(H_j \cap U)L/L|$. By Sylow's theorem $(H_j \cap U)L/L \leq (H_{j-1}/L)^{xL}$ for some $x \in H_j$. Then $(H_j \cap U)L/L$ is cyclic. From

$$(H_{j-1} \cap U)L/L \cong H_{j-1} \cap U/L \cap U \le H_j \cap U/L \cap U \cong (H_j \cap U)L/L,$$

we conclude that $H_{j-1} \cap U \leq H_j \cap U$. We have obtained a contradiction with $H_{j-1} \cap \cap U \not\leq H_j \cap U$.

2. Suppose that p divides $|(H_j \cap U)L/L|$. Since $PL/L \leq H_j/L$, we have $PL/L \leq (H_j \cap U)L/L$ by Sylow's theorem. Therefore

$$(H_j \cap U)L/L = ((H_j \cap U)L/L) \cap PL/L \cdot H_{j-1}/L = PL/L \cdot (H_{j-1} \cap U)L/L.$$

It follows that

$$p = |PL/L| = |(H_j \cap U)L/L : (H_{j-1} \cap U)L/L| = |(H_j \cap U) : (H_{j-1} \cap U)|.$$

Let us denote $D = \operatorname{Core}_{H_j \cap U}(H_{j-1} \cap U)$. From $H_{j-1} \cap U \not \leq H_j \cap U$ we have $D \neq H_{j-1} \cap U$ and $(H_j \cap U)/D = PD/D \cdot (H_{j-1} \cap U)/D$. Note that $H_{j-1} \cap U/L \cap U \cong (H_{j-1} \cap U)/L/L \leq H_{j-1}/L$ is a cyclic q-group. Since $L \cap U \leq H_j \cap U$ and $L \cap U \leq H_{j-1} \cap U$, we have $L \cap U \leq D$. It follows that $(H_{j-1} \cap U)/D \cong (H_{j-1} \cap U)/L \cap U/D/L \cap U$ is a cyclic q-group. Thus, $|(H_{j-1} \cap U)/D| = q^{n_1}$ for some natural $n_1 \leq n \leq k$, and q^{n_1} divides p-1. We have proved that $H_{j-1} \cap U$ is n_1 -modularly embedded in $H_j \cap U$ for natural $n_1 \leq k$. Thus, $H \cap U$ is k-submodular in U.

Assertion (2) follows from (1) and Lemma 2.1(2).

Lemma 2.6. Let k be a natural number. Let H be a subgroup of a group G and $N \trianglelefteq G$.

(1) If H is k-submodular in G, then HN/N is k-submodular in G/N.

(2) If $N \leq H$ and H/N is k-submodular in G/N, then H is k-submodular in G.

(3) If HN is k-submodular in G if and only if HN/N is k-submodular in G/N.

Proof. (1) By Definition 2 there is a chain of subgroups (1.1). If $H_{i-1}N/N \leq H_iN/N$ for i = 1, ..., m, then HN/N is k-submodular in HN/N.

Let's assume that $H_{i-1}N/N \not\leq H_iN/N$ for some $i \in \{1, \ldots, m\}$. Then $H_{i-1} \not\leq H_i$ and $H_{i-1}N \neq H_iN$. Let's denote $L = \operatorname{Core}_{H_i}(H_{i-1})$ and $T = \operatorname{Core}_{H_iN}(H_{i-1}N)$. Then $|H_i: H_{i-1}| = p$ and $|H_i/L| = pq^n$, q^n divides p-1 for some primes p and q and some natural $n \leq k$. From

$$1 \neq |H_i N/N : H_{i-1} N/N| = \frac{|H_i : H_{i-1}|}{|H_i \cap N : H_{i-1} \cap N|}$$

we have $H_i \cap N = H_{i-1} \cap N$ and $|H_iN/N : H_{i-1}N/N| = p$. Since $H_i \cap N \leq H_i$ and $H_i \cap N \leq H_{i-1}$ it follows that $H_i \cap N \leq L$ and $|H_iN/LN| = |H_i/L|$. Thus $|H_iN/T|$ divides pq^n because $LN \leq T$.

Note that $T/N = \operatorname{Core}_{H_iN/N}(H_{i-1}N/N) \neq H_{i-1}N/N$. Therefore $|H_iN/N/T/N| = pq^{n_1}$, $n_1 \leq n$ and q^{n_1} divides p-1. We have proved that $H_{i-1}N/N$ is n_1 -modularly embedded in H_iN/N for natural $n_1 \leq k$. Thus, HN/N is k-submodular in G/N.

(2) Let there be a chain of subgroups

$$H/N = H_0/N \le H_1/N \le \dots \le H_{m-1}/N \le H_m/N = G/N$$

such that H_{i-1}/N is *n*-modularly embedded in H_i/N for some natural $n \leq k$ and every $i = 1, \ldots, m$. Since $N \leq H_i$ and $N \leq H_{i-1}$, Assertion (2) follows from Lemma 2.1.

Statement (3) follows from Statements (1) and (2).

Lemma 2.7. Let k be a natural number and let G be a soluble group. If a subgroup H is k-submodular in G, then H is \mathfrak{U}_k -subnormal in G.

Proof. By Definition 2 there is a chain (1.1) of subgroups of subgroups such that H_{i-1} is *n*-modularly embedded in H_i for some natural $n \leq k$ and every $i = 1, \ldots, m$. If H = G then H is \mathfrak{U}_k -subnormal in G. Assume that $H \neq G$. Then $H_{i-1} \neq H_i$ for some $i \in \{1, \ldots, m\}$.

Assume that $H_{i-1} \leq H_i$. Since G is soluble, we draw a composition series through H_{i-1} and H_i :

$$H_{i-1} = R_0 < R_1 < \dots < R_{s-1} < R_s = H_i,$$

there R_{j-1} is normal in R_j and $|R_j : R_{j-1}|$ is some prime $q, j = 1, \ldots, s$. Since $R_{j-1} = \text{Core}_{R_j}(R_{j-1})$, we have $R_j/R_{j-1} \in \mathfrak{U}_k$.

Assume that $H_{i-1} \not \leq H_i$. Let's denote $L = \operatorname{Core}_{H_i}(H_{i-1})$. Then $L \neq H_{i-1}$. We have that $|H_i: H_{i-1}| = p$ and $|H_i/L| = pq^n$, q^n divides p-1 for some primes p and q. Thus, $H_i/L \in \mathfrak{U}_k$.

Therefore H is \mathfrak{U}_k -subnormal in G.

3. Classes of groups with systems of k-submodular subgroups

Theorem 3.1. Let k be a natural number, G a group. Then the following statements are equivalent.

(1) Every maximal subgroup of G is n-modularly embedded in G for some natural number $n, n \leq k$.

(2) G is supersoluble and $G/C_G(H/K)$ is either 1 or a cyclic group of order q^n for every complemented chief factor H/K of G, where q is some prime and n is some natural number, $n \leq k$.

Proof. Assume that (1) holds. By Lemma 2.3 G is supersoluble.

Let H/K be any complemented chief factor from G. Then G = HM for some subgroup M of G and $H \cap M = K$. Since G/K is supersoluble, it follows that |H/K| = pis a prime number. Then from |G/K : M/K| = |H/K| we conclude that M is maximal in G. Let $D = \text{Core}_G(M)$.

If M = D, then $G = HM \leq C_G(H/K)$ and $G/C_G(H/K) = 1$.

Let $M \neq D$. Then $G/D = HD/D \cdot M/D$, and $K = H \cap D$. From $HD/D \cong$ $\cong H/K$ it follows that |HD/D| = p and HD/D is the minimum normal subgroup of G/D. By [10, Chap. A, Theorem 15.6(1)] $HD/D = C_{G/D}(HD/D)$. Since M is *n*-modularly embedded in G and G/D is non-nilpotent, by Lemma 2.2 $|G/D| = pq^n$ for some prime $q, M/D \in \text{Syl}_q(G/D)$ and M/D are a cyclic group of order q^n, q^n divides p - 1. Note that $HD \leq C_G(HD/D)$. Then $G/C_G(HD/D) \cong G/D/C_{G/D}(HD/D) \cong M/D$. From [14, Appendix B, Theorems 1 and 3] it follows that $G/C_G(H/K) \cong \text{Aut}_G(HM_G/D) \cong G/C_G(HD/D)$. Thus, (1) \Rightarrow (2).

Assume that (2) holds. Take any maximal subgroup M in G. Let $K = \operatorname{Core}_G(M)$. If $M \leq G$ then M is n-modularly embedded in G. Let $M \neq K$. Since G/K is solvable, $G/K = N/K \cdot M/K$, where N/K is the minimum normal subgroup in G/K, $(N/K) \cap$ $\cap (M/K) = 1$, $N/K = C_{G/K}(N/K)$. Note that |N/K| = p is a prime number because G is supersoluble. We have $M/K \cong G/K/C_{G/K}(N/K) \cong G/C_G(N/K)$ — cyclic group of order q^n , q^n divides p-1. Thus, G/K is non-nilpotent and $|G/K| = pq^n$. By Lemma 2.2, M is n-modularly embedded in G. Thus, $(2) \Rightarrow (1)$.

For k = n = 1, Theorem 3.1 includes Theorem 5.3.10 from [16].

Theorem 3.2. Let k be a natural number, G a group. Then the following statements are equivalent.

(1) Every maximal subgroup of every subgroup A of G is n-modularly embedded in A for some natural number $n, n \leq k$.

(2) G is supersoluble and $G/C_G(H/K)$ is either 1 or a cyclic group of order q^n for every chief factor H/K of G, where q is some prime and n is some natural number, $n \leq k$.

Proof. Assume that (1) holds. Then every maximal subgroup of G is n-modularly embedded in G. By Lemma 2.3 G is supersoluble. Let H/K be the chief factor of G. Then |H/K| = p is a prime number. There is a Hall p'-subgroup R/K in G/K because

G/K is supersoluble. Then $G/K = P/K \cdot R/K$ for some $P/K \in \text{Syl}_p(G/K)$. Let us denote $A/K = H/K \cdot R/K$. Then A = HR, $H \cap R = K$ and H/K are the chief factor in A. By Theorem 3.1, $A/C_A(H/K)$ is either 1 or a cyclic group of order q^n , q^n divides p-1, $n \leq k$. Since $H \leq C_A(H/K)$, we have $A = C_A(H/K)R$ and

$$A/C_A(H/K) \cong R/R \cap C_A(H/K) = R/C_R(H/K)$$

On the other hand, from $H/K \leq Z(P/K)$ it follows that $P \leq C_G(H/K)$. Then $G = PR = C_G(H/K)R$ and

$$G/C_G(H/K) \cong R/R \cap C_G(H/K) = R/C_R(H/K).$$

Thus, it has been proven that $(1) \Rightarrow (2)$.

Assume that (2) holds. Take any subgroup $A \leq G$. Since G is supersoluble, there is a chief series

$$1 = G_0 < G_1 < \dots < G_{m-1} < G_m = G,$$

where $|G_i/G_{i-1}|$ is a prime number, i = 1, ..., m. Consider the series

$$1 = G_0 \cap A \le G_1 \cap A \le \dots \le G_{m-1} \cap A \le G_m \cap A = A.$$

$$(2.1)$$

If $G_{i-1} \cap A = G_i \cap A$ for any i = 1, ..., m, then 1 = A and for A the statement (1) of the theorem is satisfied.

Suppose that there exists $j \in \{1, \ldots, m\}$ for which $G_{j-1} \cap A \neq G_j \cap A$. Then $1 \neq A$. For m = 1 the theorem is true. Therefore let m > 1. From the maximality of G_{j-1} in G_j , $G_{j-1} \leq (G_j \cap A)G_{j-1} \leq G_j$ and $G_{j-1} \cap A \neq G_j \cap A$, we conclude that $G_j = (G_j \cap A)G_{j-1}$. Then $G_j/G_{j-1} \cong G_j \cap A/G_{j-1} \cap A$ is the chief factor of A. Omitting repetitions from the series (2.1), we obtain the chief series

$$1 = A_0 < A_1 < \dots < A_{l-1} < A_l = A,$$

where $A_i = G_t \cap A$ for some $t \in \{1, \ldots, m\}$ for which $G_{t-1} \cap A \neq G_t \cap A$.

Let T/S be an any chief factor of A. Then

$$S \leq (A_0 \cap T)S \leq (A_1 \cap T)S \leq \cdots \leq (A_{l-1} \cap T)S \leq (A_l \cap T)S = T.$$

From $S \neq T$ it follows that $(A_{i-1} \cap T)S \neq (A_i \cap T)S$ for some $i \in \{1, \ldots, l\}$. Since S is maximal in T, we have $S = (A_{i-1} \cap T)S$ and $T = (A_i \cap T)S$. Then

$$T/S \cong (A_i \cap T)/(A_{i-1} \cap T)(A_i \cap S).$$

On the other side

$$(A_i \cap T)A_{i-1}/(A_i \cap S)A_{i-1} \cong (A_i \cap T)/(A_{i-1} \cap T)(A_i \cap S).$$

Then $A_{i-1} \leq (A_i \cap S)A_{i-1} \neq (A_i \cap T)A_{i-1} \leq A_i$. Hence $A_{i-1} = (A_i \cap S)A_{i-1}$ and $A_i = (A_i \cap T)A_{i-1}$. This means $T/S \cong A_i/A_{i-1} = G_t \cap A/G_{t-1} \cap A$ for some $t \in \{1, \ldots, m\}$.

Let $|G_t/G_{t-1}| = p$ be a prime number. By assumption, $G/C_G(G_t/G_{t-1})$ is either 1 or a cyclic group of order q^n , q^n divides p-1, $n \leq k$.

Since $AC_G(G_t/G_{t-1})/C_G(G_t/G_{t-1})$ is a subgroup of $G/C_G(G_t/G_{t-1})$, we conclude that $A/C_G(G_t/G_{t-1}) \cap A \cong AC_G(G_t/G_{t-1})/C_G(G_t/G_{t-1})$ is either 1 or a cyclic group of order q^{n_1} , q^{n_1} divides p-1 for some natural number $n_1 \leq n$. From $C_G(G_t/G_{t-1}) \cap$ $\cap A \leq C_A(G_t \cap A/G_{t-1} \cap A) = C_A(A_i/A_{i-1})$ we conclude that $A/C_A(A_i/A_{i-1})$ is a homomorphic image of $A/C_G(G_t/G_{t-1}) \cap A$. In view of [14, Appendix B, Theorems 1 and 3] $A/C_A(T/S) \cong \operatorname{Aut}_A(T/S) \cong \operatorname{Aut}_A(A_i/A_{i-1}) \cong A/C_A(A_i/A_{i-1})$ is either 1 or a cyclic group of order q^{n_2} , q^{n_2} divides p-1 for some natural number $n_2 \leq n_1 \leq n$. By Theorem 3.1, any maximal subgroup in A is n-modularly embedded. Thus, (2) \Rightarrow (1). \Box For k = n = 1, Theorem 3.2 includes Theorem 4.4 from [14, Chap. 4].

Theorem 3.3. Let k be a natural number. The class \mathfrak{X} of all groups G in which any maximal subgroup is n-modularly embedded for some natural number n, $n \leq k$, is a Schunk class consisting of supersoluble groups.

Proof. By Lemma 2.3(1), \mathfrak{X} consists of supersoluble groups.

Let us show that \mathfrak{X} is a homomorph. Let $G \in \mathfrak{X}$ and $N \leq G$. For any maximal subgroup M/N in G/N, the subgroup M is maximal in G. Since $N \leq \operatorname{Core}_G(M)$, we have $\operatorname{Core}_{G/N}(M/N) = \operatorname{Core}_G(M)/N$. The subgroup M is *n*-modularly embedded in G. If $M \leq C$ then $M/N \leq C/N$

If $M \leq G$ then $M/N \leq G/N$.

Let $M/N \neq \operatorname{Core}_{G/N}(M/N)$. From $M \neq \operatorname{Core}_G(M)$ by Definition 1 we have |G : M| = p and $|G/\operatorname{Core}_G(M)| = pq^n$, q^n divides p-1. Therefore |G/N : M/N| = |G : M| = p and $|G/N/\operatorname{Core}_{G/N}(M/N)| = pq^n$, q^n divides p-1. Thus, M/N is n-modularly embedded in G/N and $G/N \in \mathfrak{X}$. This means that \mathfrak{X} is a homomorph.

Let us prove that \mathfrak{X} is primitively closed. Let $G/\operatorname{Core}_G(M) \in \mathfrak{X}$ for any maximal subgroup M in G. If $M = \operatorname{Core}_G(M)$, then $M \trianglelefteq G$. Let $M \neq \operatorname{Core}_G(M)$. From $G/\operatorname{Core}_G(M) \in \mathfrak{X}$ it follows that $M/\operatorname{Core}_G(M)$ is n-modularly embedded in $G/\operatorname{Core}_G(M)$. Since $\operatorname{Core}_{G/\operatorname{Core}_G(M)}(M/\operatorname{Core}_G(M)) \cong 1$ we have that |G : M| = $= |G/\operatorname{Core}_G(M) : M/\operatorname{Core}_G(M)| = p$ and $|G/\operatorname{Core}_G(M)| = pq^n$, q^n divides p - 1, i.e. M is n-modularly embedded in G. Thus, $G \in \mathfrak{X}$ and \mathfrak{X} are primitively closed. \Box

Theorem 3.4. Let k be a natural number. The class \mathfrak{Y} of all groups G such that every maximal subgroup of any subgroup A of G is n-modularly embedded in A for some natural number $n, n \leq k$, is a hereditary homomorph that has the following property: if $G = G_1 \times G_2$ and $G_i \in \mathfrak{Y}$, i = 1, 2, then $G \in \mathfrak{Y}$.

Proof. It is obvious that \mathfrak{Y} is a hereditary class of groups.

Let us prove that \mathfrak{Y} is a homomorph. Take $G \in \mathfrak{Y}$ and $N \leq G$. Let A/N be an any subgroup of G/N and M/N be a maximal subgroup in A/N. From the maximality of M in $A \leq G \in \mathfrak{Y}$ it follows that M is n-modularly embedded in A for some natural number $n, n \leq k$. If $M = \operatorname{Core}_A(M)$, then $M \leq A$ and $M/N \leq A/N$. Let $M \neq \operatorname{Core}_A(M)$. Then $M/N \leq A/N$. Since M is n-modularly embedded in A, we have |A : M| = p and $|A/\operatorname{Core}_A(M)| = pq^n, q^n$ divides p - 1. From this we obtain that |A/N : M/N| = pand $|A/N/(M/N)_{A/N}| = pq^n$. This means that M/N is n-modularly embedded in A/N. Thus, \mathfrak{Y} is a homomorph.

Suppose that $G = G_1 \times G_2$, $G_i \in \mathfrak{Y}$, i = 1, 2. Since $\mathfrak{Y} \subseteq \mathfrak{U}$ and \mathfrak{U} are a formation, we have $G \in \mathfrak{U}$. Let A be a subgroup of G and H/K be its chief factor. Since \mathfrak{Y} is hereditary and $G/G_i \in \mathfrak{Y}$, it follows that $A/A \cap G_i \cong AG_i/G_i \in \mathfrak{Y}$, i = 1, 2. In the case when $A \cap G_i = 1$ or $A \leq G_i$, i = 1, 2, we have $A \in \mathfrak{Y}$.

Let $1 \neq A \cap G_i \neq A$. By [14, Appendix B, Theorem 4] $A \cap G_1$ either covers or avoids H/K.

1. If $A \cap G_1$ covers H/K, then $H(A \cap G_1) = K(A \cap G_1)$. Therefore $H = K(H \cap G_1)$ and H/K is isomorphic to the chief factor $(H \cap G_1)(A \cap G_2)/A \cap G_2/(K \cap G_1)(A \cap G_2)/A \cap G_2$ of $A/A \cap G_2 \in \mathfrak{Y}$. From this and Theorem 3.2 we conclude that $A \in \mathfrak{Y}$.

2. If $A \cap G_1$ avoids H/K, then $H \cap G_1 = H \cap A \cap G_1 = K \cap A \cap G_1 = K \cap G_1$. Then H/K is isomorphic to the chief factor $H(A \cap G_1)/A \cap G_1/K(A \cap G_1)/A \cap G_1$ of $A/A \cap G_1 \in \mathfrak{Y}$. From this and Theorem 3.2 it follows that $A \in \mathfrak{Y}$.

Note that $\mathfrak{Y} \subseteq \mathfrak{X}$ and $\mathfrak{Y} \neq \mathfrak{X}$ in the general case.

Example 3.1. Let n = 1, let $P = \langle a, b, c | a^7 = b^7 = c^7 = 1$, ab = ba, $ac = ca, b^c = ba \rangle$ be the non-abelian group of order 7³. Define $g \in \operatorname{Aut}(P)$ by $(c^i b^j a^k)^g = c^{i6} b^{j2} a^{k12}$ for any $i, j, k \in \{0, 1, \dots, 6\}$. The group P has exponent 7 and $|\langle g \rangle| = 6$. In the group $G = P \setminus \langle g \rangle$, the subgroup $\langle a \rangle = \Phi(P) = \Phi(G)$ and $\langle a \rangle \langle g \rangle \notin \mathfrak{Y}$, but $G \in \mathfrak{X}$.

In $\langle a \rangle \langle g \rangle$ for k = n = 1 the maximal subgroup $\langle g \rangle$ is \mathfrak{U}_k -subnormal, since $\langle a \rangle \langle g \rangle \in \mathfrak{U}_k$. But $\langle g \rangle$ is not k-submodular, since $\operatorname{Core}_{\langle a \rangle \langle g \rangle}(\langle g \rangle) = 1$.

Lemma 3.1. Let \mathfrak{F} be the class of all groups in which every Sylow subgroup is k-submodular. Let G be a group. Then the following assertions hold.

(1) $\mathfrak{N} \subseteq \mathfrak{F} \subseteq \mathfrak{wU}.$

(2) If $G \in \mathfrak{F}$ and $N \leq G$, then $G/N \in \mathfrak{F}$.

(3) If $G/N_1 \in \mathfrak{F}$ and $G/N_2 \in \mathfrak{F}$ for any $N_i \leq G$, i = 1, 2, then $G/N_1 \cap N_2 \in \mathfrak{F}$.

(4) A direct product of groups from \mathfrak{F} lies in \mathfrak{F} .

(5) If $G \in \mathfrak{F}$ and U is a subgroup of G, then $U \in \mathfrak{F}$.

Proof. Statement (1) follows from Definitions 1, 2.

Statement (2) follows from Lemmas 2.4, 2.6 and Sylow's theorem.

(3) Let G be a group of the least order such that $G/N_i \in \mathfrak{F}$, $N_i \leq G$, i = 1, 2, but $G/N_1 \cap N_2 \notin \mathfrak{F}$. We can assume that $N_1 \cap N_2 = 1$. Let $Q \in \operatorname{Syl}_q(G)$. Then $QN_i/N_i \in \operatorname{Syl}_q(G/N_i)$ and QN_i/N_i is k-submodular in G/N_i for i = 1, 2. By Lemma 2.6(2) QN_i is k-submodular in G for i = 1, 2. By Lemma 2.5(2) $QN_1 \cap QN_2$ is k-submodular in G. By [10, Chap. A, Theorem 6.4(b)] $QN_1 \cap QN_2 = Q(N_1 \cap N_2) = Q$ is k-submodular in G. Therefore $G = G/N_1 \cap N_2 \in \mathfrak{F}$. The contradiction thus obtained completes the proof of Statement (3).

Statement (4) follows from (3).

Statement (5) follows from Lemma 2.5(1) and Sylow's theorem.

Further $\mathfrak{A}(p-1)_k$ means the class of all abelian groups of exponent dividing p-1 and not dividing by the (k+1)th powers of primes.

Theorem 3.5. Let \mathfrak{K} be the class of all supersoluble groups in which every Sylow subgroup is k-submodular. Then \mathfrak{K} is a hereditary saturated formation and \mathfrak{K} is defined by a formation function h such that $h(p) = (G \mid G \in \mathfrak{A}(p-1)_k \text{ and all Sylow subgroups}$ of G are cyclic).

Proof. Since $\mathfrak{K} = \mathfrak{U} \cap \mathfrak{F}$ we have that \mathfrak{K} is a hereditary formation.

Let us prove that \mathfrak{K} is saturated. Let G be a group of the least order for which $G/\Phi(G) \in \mathfrak{K}$ and $G \notin \mathfrak{K}$. Then there exists a Sylow q-subgroup Q of G that is not k-submodular in G. From $G/\Phi(G) \in \mathfrak{U}$ it follows that $G \in \mathfrak{U}$. Then a Sylow p-subgroup is normal in G for the largest prime $p \in \pi(G)$. Let N be a minimal normal subgroup of G. By [10, Chap. A, Theorem 9.2(e)] $\Phi(G)N/N \leq \Phi(G/N)$. Since $G/\Phi(G)N \in \mathfrak{K}$, we have that $(G/N)/\Phi(G/N) \in \mathfrak{K}$. From |G/N| < |G|, it follows $G/N \in \mathfrak{K}$. Then N is a unique minimal normal subgroup of G, $N \leq \Phi(G) \leq F(G)$ and $F(G) \in \text{Syl}_p(G)$. We note that |N| = p. By [10, Chap. A, Theorem 10.6(c)] $\Phi(G) < F(G)$. Since $QN/N \in \text{Syl}_q(G/N)$ and $QN/N \notin \Phi(G/N)$, we have that $QN/N \neq N/N$ and QN/N is k-submodular in G/N. By Lemma 2.6(2) QN is k-submodular in G. Then $Q \not \leq QN$ and $q \neq p$ by choice of Q.

Let B = QN. Then $T = \text{Core}_B(Q) \neq Q$. We have that $B/T = NT/T \cdot Q/T$ and $C_{B/T}(NT/T) = NT/T$. Therefore $Q/T \cong B/T/C_{B/T}(NT/T)$ is isomorphic to a subgroup from Z_{p-1} and $|Q/T| = q^n$, *n* is a natural number, q^n делит p-1. 1. Assume that G/N = QF(G)/N. By Lemma 2.7 $G/N \in \mathfrak{wU}_k$. By Lemma 1.6(3) $G/N/F(G/N) \in \mathcal{A}_k$. Since F(G/N) = F(G)/N. We have that $Q \cong G/F(G) \in \mathcal{A}_k$. So $n \leq k$. By Lemmas 2.2 and 2.4(2) Q is k-submodular in G; a contradiction.

2. Assume that $G/N \neq QF(G)/N = H/N$. From hereditary \mathfrak{K} it follows $H/N \in \mathfrak{K}$. By Lemma 2.7 $H/N \in \mathfrak{wU}_k$. By Lemma 1.6(2) \mathfrak{wU}_k is a hereditary saturated formation. Since H is a Hall subgroup in G, by [17, Corollary 16.2.3] from $H/H \cap \Phi(G) \in \mathfrak{wU}_k$ it follows that $H \in \mathfrak{wU}_k$. Then $B = QN \in \mathfrak{wU}_k$. By Lemma 1.6(3) $B/F(B) \in \mathcal{A}_k$. Then $Q/F(B) \cap Q \cong QF(B)/F(B) \in \mathcal{A}_k$. We note that $F(B) = N(F(B) \cap Q)$ and $(F(B) \cap Q) \in \mathcal{A}_k$ and $n \leq k$. By Lemmas 2.2 and 2.4(2) Q is k-submodular in G; a contradiction. This \mathfrak{K} is saturated.

Let us prove that $LF(h) = \mathfrak{K}$. By [10, Chap. IV, Proposition 3.14 and Theorem 4.6] LF(h) is a hereditary saturated formation.

Show that $LF(h) \subseteq \mathfrak{K}$. Let G be a group of least order in $LF(h) \setminus \mathfrak{K}$. Let N be a minimal normal subgroup of G. By choice of G we have that $G/N \in \mathfrak{K}$, $\Phi(G) = 1$ and N is a unique minimal normal subgroup of G, since LF(h) and \mathfrak{K} are saturated formation. From $h(p) \subseteq \mathfrak{A}(p-1)$ it follows that $G \in \mathfrak{U}$, so |N| = p for some prime p. From $G \notin \mathfrak{K}$ it follows that G has a Sylow q-subgroup Q that is not k-submodular in G. Then $QN/N \in \mathfrak{Syl}_q(G/N)$. From $G/N \in \mathfrak{K}$ by Lemma 2.6(2) we have that QN is k-submodular in G. So Q is not normal in QN and $q \neq p$.

If $G \neq QN$, then $QN \in \mathfrak{K}$ since LF(h) and \mathfrak{K} are hereditary. Thus Q is k-submodular in QN and Q is k-submodular in G; a contradiction.

Let G = QN. We have that $Q \cong G/N = G/C_G(N) \in h(p)$ because $G \in LF(h)$. Therefore $Q \in \mathfrak{A}(p-1)_k$ and Q is cyclic. Then Q is k-submodular in G; a contradiction. Thus $LF(h) \subseteq \mathfrak{K}$.

Prove that $\mathfrak{K} \subseteq LF(h)$. Let G be a group of least order in $\mathfrak{K} \setminus LF(h)$. We denote by N a minimal normal subgroup of G. Then $G/N \in \mathfrak{K}$. Since \mathfrak{K} and LF(h) are saturated formations, we have that $\Phi(G) = 1$ and N is a unique minimal normal subgroup of G. Then G = NM, where M is a maximal subgroup of G, $\operatorname{Core}_G(M) = 1$, $M \cap N = 1$ and $N = C_G(N)$. From $G \in \mathfrak{K} \subseteq \mathfrak{U}$ it follows that |N| = p for some prime p. Then $M \cong G/C_G(N)$ is isomorphic to a subgroup from Z_{p-1} . Let $R \in \operatorname{Syl}_q(M)$.

If $G \neq RN = D$ then $D \in \mathfrak{K}$, since \mathfrak{K} is hereditary. By choose of G, $D \in LF(h)$. So $R \cong D/C_D(N) \in h(p)$, since $N = C_D(N)$. Thus $R \in \mathfrak{A}(p-1)_k$. So $G/C_G(N) \cong M \in h(p)$ and $G \in LF(h)$; a contradiction.

Let G = RN. Then $M^x = R$ for some $x \in G$. Since $G \in \mathfrak{K}$, R is k-submodular in G. From maximality of R in G it follows that R is n-modularly embedded in G for some natural $n \leq k$. So $G/C_G(N) \in h(p)$ and $G \in LF(h)$; a contradiction. Thus $\mathfrak{K} \subseteq LF(h)$. This means that the equality $\mathfrak{K} = LF(h)$ is proven. \Box

Theorem 3.6. The class \mathfrak{F} of all groups in which every Sylow subgroup is k-submodular is a hereditary saturated formation and \mathfrak{F} is defined by a formation function f such that $f(p) = (G \mid \text{Syl}(G) \subseteq \mathfrak{A}(p-1)_k$ and all Sylow subgroups of G are cyclic).

Proof. By Lemma 3.1 \mathfrak{F} is a hereditary formation.

Prove that \mathfrak{F} is saturated. Let G be a group of least order such that $G/\Phi(G) \in \mathfrak{F}$ and $G \notin \mathfrak{F}$. Since $\mathfrak{F} \subseteq \mathfrak{wU}$ and by Lemma 1.5(2) \mathfrak{wU} is saturated formation, we have that $G \in \mathfrak{wU}$. So G is soluble. Let N be a minimal normal subgroup of G. By choose of G, it follows that $G/N \in \mathfrak{F}$. Then N is a unique minimal normal subgroup of G. Hence, $N \leq \Phi(G) < F(G)$ and F(G) is p-group for some prime p. Since G is Ore dispersive, we have that p is the greatest prime divisor of |G| and $F(G) \in \operatorname{Syl}_p(G)$. From $G \notin \mathfrak{F}$ it follows that G has a Sylow q-subgroup Q which is not k-submodular in G. Since $G/N \in \mathfrak{F}$ we have that QN/N is k-submodular in G/N. By Lemma 2.6(3) QN is k-submodular in G. So $q \neq p$ and $\operatorname{Core}_{QN}(Q) \neq Q$.

1. $|\pi(G)| = 2$. Then $|\pi(G/N)| = 2$ and $G/N = QN/N \cdot F(G)/N$. By [15, Theorem 2], $G/N \in \mathfrak{U}$. So $G/N \in \mathfrak{K}$. By Theorem 3.5 \mathfrak{K} is saturated. Then $G \in \mathfrak{K} \subseteq \mathfrak{F}$; a contradiction.

2. $|\pi(G)| > 2$. Then QN/N is contained in a Hall $\{p,q\}$ -subgroup H/N of G/Nand $H/N = QN/N \cdot F(G)/N$. By Theorem 3.5 \mathfrak{K} is hereditary, so $H/N \in \mathfrak{K}$. Therefore, QN/N and F(G)/N are k-submodular in H/N. By Lemma 1.4 $H/N \in \mathfrak{U}$, so $H/N \in \mathfrak{K}$. Since by Theorem 3.5 \mathfrak{K} a local formation and H = QF(G) is a Hall $\{p,q\}$ -subgroup of G, we can apply Corollary 16.2.3 of [17]. From $H\Phi(G)/\Phi(G) \cong H/H \cap \Phi(G) \cong H/N/H \cap$ $\cap \Phi(G)/N \in \mathfrak{K}$ it follows that $H \in \mathfrak{K}$. By Theorem 3.5, $QN \in \mathfrak{K}$. Then Q is k-submodular in QN and so Q is k-submodular in G. This contradiction completes the proof that \mathfrak{F} is saturated.

Let us prove that $LF(f) = \mathfrak{F}$.

It is easy to verify that f(p) is a hereditary formation. By [10, Chap. IV, Proposition 3.14 and Theorem 4.6] LF(f) is hereditary saturated formation. By Lemma 1.5 $LF(f) \subseteq \subseteq$ w \mathfrak{U} and LF(f) consists of Ore dispersive groups.

Show that $LF(f) \subseteq \mathfrak{F}$. Let G be a group of least order in $LF(f) \setminus \mathfrak{F}$. Since LF(f) and \mathfrak{F} are saturated formations, we conclude that G has a unique minimal normal subgroup N and $\Phi(G) = 1$. From $G \in \mathfrak{S}$ it follows that $N = C_G(N)$ is an elementary abelian p-group for some prime p. The Ore dispersiveness of G implies that $N \in \text{Syl}_p(G)$ and p is the greatest prime divisor of |G|. Sice $G \notin \mathfrak{N}$, $|\pi(G)| \geq 2$.

If $QN \neq G$ for all $Q \in \text{Syl}_q(G)$, $q \neq p$, then from $QN \in LF(f)$ it follows that $QN \in \mathfrak{F}$. So Q is k-submodular in QN. Since $QN/N \in \text{Syl}_q(G/N)$ and $G/N \in \mathfrak{F}$, we conclude that QN/N is k-submodular in G/N. By Lemma 2.6(2) QN is k-submodular in G. So Q is k-submodular in G. Hence $G \in \mathfrak{F}$; a contradiction.

Let QN = G for some $Q \in \text{Syl}_q(G)$. Then Q is a maximal subgroup of G and $\text{Core}_G(Q) = 1$. From $G \in LF(f)$ it follows that $Q \cong G/C_G(N) \in f(p)$. So $Q \in \mathfrak{A}(p-1)_k$ and Q is cyclic.

Since $G \in LF(f) \subseteq \mathfrak{wU}$ it follows that Q is \mathbb{P} -subnormal in G, we have that |G:Q| = p. By Lemma 2.2 Q is k-submodular in G. Hence $G \in \mathfrak{F}$ is a contradiction to the choice of G. Thus, $LF(f) \subseteq \mathfrak{F}$.

Prove that $\mathfrak{F} \subseteq LF(f)$. Let G be a group of least order in $\mathfrak{F} \setminus LF(f)$. Since \mathfrak{F} and LF(f) are saturated formations, we have that $G/N \in LF(f)$, N is a unique minimal normal subgroup of G, $\Phi(G) = 1$. Then G = NM, where M is a maximal subgroup of G, $N \cap M = 1$, $N = C_G(N)$ is an p-group for some prime p, because $G \in \mathfrak{S}$. From $G \in \mathfrak{wU}$ we conclude that G is Ore dispersive. Then p is the greatest prime divisor of |G| and $N \in \mathrm{Syl}_p(G)$. From $G \notin LF(f)$ it follows that $G/C_G(H/K) \notin f(p)$ for some chief factor H/K of G. Then H = N and K = 1 because $G/N \in LF(f)$. So $G/C_G(N) \notin f(p)$.

Let $R \in \text{Syl}_q(M)$. Then $R \in \text{Syl}_q(G)$.

Suppose that $D = RN \neq G$. Since $N \leq F(D)$, $F(D) = N(Q \cap F(D))$. So $Q \cap \cap F(D) \in \operatorname{Syl}_q(F(D))$. Then $Q \cap F(D) \leq C_D(N) \leq C_G(N) = N$ and F(D) = N. From $D \in \mathfrak{F} \subseteq \mathfrak{wl}$ and Lemma 1.4, we have that $D \in \mathfrak{K}$. By Theorem 3.5 $D/C_D(L/T) \in h(p)$ for every chief factor L/T of D. By [17, Corollary 4.1.1] $F(D) = \cap C_D(L/T)$ for every chief factor L/T of D. So $R \cong D/N \in h(p)$. Then $G/C_G(N) = G/N \cong M \in f(p)$; a

contradiction.

Suppose that RN = G. By Lemma 1.4, $G \in \mathfrak{U}$. Then |N| = p and $R \cong G/N = G/C_G(N) \cong L \leq Z_{p-1}$. From $G \in \mathfrak{F}$ and R = M it follows that R is *n*-modularly embedded in G, $n \leq k$. Therefore $G/C_G(N) \cong R \in \mathfrak{A}(p-1)_k$ and $G/C_G(N) \in f(p)$; a contradiction.

Hence $\mathfrak{F} \subseteq LF(f)$. It has been proven that $\mathfrak{F} = LF(f)$.

Proposition 3.1. $\mathfrak{K} = \mathfrak{U} \cap \mathfrak{wU}_k$, *i.e.* the class \mathfrak{K} of all supersoluble groups in which every Sylow subgroup is k-submodular coincides with the class of all supersoluble groups in which every Sylow subgroup is \mathfrak{U}_k -subnormal.

Proof. (1) Since $\mathfrak{K} \subseteq \mathfrak{U}$, by Lemma 2.7 $\mathfrak{K} \subseteq \mathfrak{U} \cap \mathfrak{w}\mathfrak{U}_k$.

Show that $\mathfrak{U} \cap \mathfrak{wl}_k \subseteq \mathfrak{K}$. Let G be a group of least order in $\mathfrak{U} \cap \mathfrak{wl}_k \setminus \mathfrak{K} \neq \emptyset$. Let N be a minimal normal subgroup of G. Then |N| = p for some prime p. From $G \notin \mathfrak{K}$ it follows that there exists a Sylow q-subgroup Q of G that is not k-submodular in G. Since \mathfrak{U}_k is hereditary, we have that $QN \in \mathfrak{wl}_k$ and Q is \mathfrak{U}_k -subnormal in QN. Then G = QN by choose of G. So $q \neq p$ and Q is a maximal subgroup in G. Since $\mathfrak{U} \cap \mathfrak{wl}_k$ and \mathfrak{K} are hereditary formations, we have that N is a unique minimal normal subgroup of G, $\operatorname{Core}_G(Q) = 1$ and $N = C_G(N) = F(G)$. Since $G/C_G(N)$ is isomorphic to a subgroup from Z_{p-1} we have that $Q \cong G/N$ is cyclic, $|Q| = q^n$ divides p - 1. By Lemma 1.6(3) $Q \cong G/F(G) \in \mathcal{A}_k$. So $n \leq k$ and Q is k-submodular in G; a contradiction. Thus $\mathfrak{U} \cap \mathfrak{wl}_k \subseteq \mathfrak{K}$. This means that the equality $\mathfrak{K} = \mathfrak{U} \cap \mathfrak{wl}_k$ is proven.

This work was supported by the Ministry of Education of the Republic of Belarus (Grant no. 20211750"Convergence-2025").

References

- [1] Huppert B. Endliche Gruppen. I. / B. Huppert. Berlin: Springer, 1967.
- [2] Schmidt R. Modulare Untergruppen endlicher Gruppen / R. Schmidt // J. Ill. Math. 1969. Vol. 13. P. 358–377.
- [3] Zimmermann I. Submodular subgroups in finite groups // Math. Z. 1989. Vol. 202. P. 545–557.
- [4] Vasilyev V.A. Finite groups with submodular Sylow subgroups // Siberian Math. J. 2015. Vol. 56, No. 6. P. 1019–1027.
- [5] Vasil'ev A.F., Vasil'eva T.I., Parfenkov K.L. Finite groups with three given subgroups // Siberian Math. J. 2018. Vol. 59, No. 1. P. 50–58.
- [6] Murashka V.I. Finite Groups With Given Sets of *S*-Subnormal Subgroups // Asian-Eur. J. Math. 2020. Vol. 13, No. 4. P. 2050073.
- [7] Monakhov V.S., Sokhor I.L. Finite groups with submodular primapy subgroups // Arch. Math. 2023. Vol. 121, P. 1–10.
- [8] Murashka V.I. Formations of finite groups in polynomial time: *S*-residuals and *S*-subnormality // Journal of Symbolic Computation. 2024. Vol. 122. P. 102271.

- [9] Monakhov V.S., Sokhor I.L. Finite groups with formational subnormal primapy subgroups of bounded exponent // Siberian Electronic Mathematical News. 2023. Vol. 20, No. 2. P. 785–796.
- [10] *Doerk K., Hawkes T.* Finite Soluble Groups. Berlin-New York: Walter de Gruyter, 1992.
- [11] Ballester-Bolinches A., Ezquerro L.M. Classes of Finite Groups. Springer-Verlag, 2006.
- [12] Vasil'ev A.F., Vasil'eva T.I., Tyutyanov V.N. On the finite groups of supersoluble type // Siberian Math. J. 2010. Vol. 51, No. 6. P. 1004–1012.
- [13] Vasil'ev A.F., Vasil'eva T.I., Tyutyanov V.N. On K-P-Subnormal Subgroups of Finite Groups // Math. Notes. 2014. Vol. 95, No. 4. P. 471–480.
- [14] Between Nilpotent and Solvable / H.G. Bray [and others]; edited by M. Weinstein. Passaic: Polugonal Publishing House, 1982.
- [15] Vasil'ev A.F. New properties of finite dinilpotent groups // Vestsi Nats. Akad. Navuk Belarusi. Ser. Fiz.-Mat. Navuk. 2004. No. 2. P. 29–33 (In Russian).
- [16] Schmidt R. Subgroup Lattices of Groups. Berlin etc: Walter de Gruyter, 1994.
- [17] Shemetkov L.A. Formations of finite groups. Moscow: Nauka, 1987. (In Russian)

T. I. Vasil'eva

Belarusian State University of Transport, Kirov str., 34, Gomel 246653, Belarus. E-mail address: tivasilyeva@mail.ru