

# On Quantifiers for Quantitative Reasoning

Matteo Capucci  
MSP Group, University of Strathclyde  
[matteo.capucci@gmail.com](mailto:matteo.capucci@gmail.com)

June 10, 2024

## Abstract

We explore a kind of first-order predicate logic with intended semantics in the reals. Compared to other approaches in the literature, we work predominantly in the multiplicative reals  $[0, \infty]$ , showing they support three generations of connectives, that we call *non-linear*, *linear additive* and *linear multiplicative*. Means and harmonic means emerge as natural candidates for bounded existential and universal quantifiers, and in fact we see they behave as expected in relation to the other logical connectives. We explain this fact through the well-known fact that min/max and arithmetic mean/harmonic mean sit at opposite ends of a spectrum, that of *p-means*. We give syntax and semantics for this quantitative predicate logic, and as example applications, we show how softmax is the quantitative semantics of argmax, and Rényi entropy/Hill numbers are additive/multiplicative semantics of the same formula. Indeed, the additive reals fit into the story by exploiting the *Napierian duality*  $-\log \dashv 1/\exp$ , which highlights a formal distinction between ‘additive’ and ‘multiplicative’ quantities. Finally, we describe two attempts at a categorical semantics via enriched hyperdoctrines. We discuss why hyperdoctrines are in fact probably inadequate for this kind of logic.

## 1. INTRODUCTION

In 1847, Boole [Boo47] published his treatise on the algebra of logic, where he gave algebraic axioms to manipulate truth values as quantities. Ever since, mathematicians have been trying to go back a certain measure and formulate formal languages which unite the *symbolic* and the *quantitative* in a single system. On the one hand, such a system should enable clean symbolic reasoning, like for traditional formal languages; on the other, it should have the nuance of the continuously-valued real number.

There’s more to it, however, than just these two aspects: after all, mathematicians started investigating many-valued logic a long time ago, with the work of Łukasiewicz [Łuk20] on the one hand, and Scott [Sco67] on the other. Recently, applying that technology to the case of the reals, seen as a poset of truth values, Figueroa and van der Berg studied the topos of real-valued sets in [FB22].

However, the mix of symbolic and quantitative that mathematicians *intue* to be needed is different than just having continuum-many truth values. It also involves giving arithmetic operations, like  $+$  and  $\times$ , a logical status. Probability theory gets very close to such a logic, with probability measures assigning belief degrees to propositions which are then summed or multiplied in the ways every mathematician is familiar with, and there are indeed axiomatizations of probability theory which brings to light their *rational*<sup>1</sup> content [Cox46]. However, probability theory lacks a formal or syntactic aspect to it, so that the probabilist must reason *with the theory* instead of *within the theory*.

Something different is the *real-valued generalised logic* of Lawvere [Law73], which tries to take seriously the idea that operations on the reals are logical connectives which can be manipulated symbolically while targeting a quantitative semantics. Lawvere’s ideas have been a great source of inspiration in category-theoretic circles, though only very recently Bacci, Mardare, Panangaden and Plotkin picked up where Lawvere left and started developing its logic [BMPP23; BMPP24].

Parallel to these coherent attempts, we must mention the vast development on fuzzy logic and fuzzy set theory [Zad65; Zad88; BMW85] as well as the growing literature on differential logics with continuous semantics [TMP24; SKD+23]. Also various works dealing with probability have foreshadowed links between the symbolic and the quantitative, such as Kozen’s probabilistic PDL [Koz83] where predicates are measurable functions and satisfaction is measured by integration (not unlike what we are going to do in this work). Other works in this space that we would like to mention are [Gra07; DP22a; DP22b].

This work started as an attempt to understand the semiotics<sup>2</sup> of softmax, which is used and sometimes advertised as a ‘quantitative’ (alternatively: ‘smooth’, ‘continuous’, ‘fuzzy’) version of argmax. Trying to understand in what sense is this intuition true led us to analyze the structure of the extended positive reals and we discovered a logical structure of *degrees of belief* which partially recovers known observations (such as [BMPP24; Shu22; Gra07; Koz83]), but are also original to some degree.

Nonetheless, this work is the proverbial tip of an iceberg, which remains largely submerged under the water. We only got glimpses of the logic we are about to describe, and this concerns syntax and semantics alike. Specifically, we lack a proof system, which limits the reach of the justifications we can give formally to symbolic manipulations.

**1.1. Outline of the paper.** In Section 2 we analyze the extended positive real numbers  $[0, \infty]_{\otimes}$  and single out the algebraic structure it naturally hosts, in the form of three generations of logical connectives—a non-linear one and two linear ones (additive and multiplicative). We single out positive reals as a *multiplicative* world, dual to the *additive* world which is instead

---

<sup>1</sup>As in: pertaining reasoning.

<sup>2</sup>Meaning not just the crude mathematical definition but the set of latent meaning and informal intuitions that a mathematical idea bears and is communicated with.

given by the full extended real line  $[-\infty, \infty]_{\oplus}$ . Taking seriously this duality is an important conceptual idea of this work.

In the multiplicative reals, the linear additive connectives given by *sum* and its de Morgan dual, *harmonic sum* [Gra07], form the basis for Section 3, where we notice two things. First, both kinds of sums are actually part of a spectrum, well known in real analysis, of  $p$ -sums, given by conjugating  $+$  with the non-trivial automorphisms of  $[0, \infty]_{\otimes}$ . Second, by normalizing these sums we get  $p$ -means, readily generalised to  $p$ -integrals over measure spaces (since means are integrals over probability spaces) and these behave like **soft quantifiers** for predicates bounded by a measure space. In Section 3.3 we show our main result, which is that softmax and argmax are semantics of the same sentence.

In Section 4 we try to frame integrals as quantifiers in the sense of Lawvere [Law69], thus as adjoints to reindexing. We first attempt to do so naively, and then by deploying the theory of enriched indexed categories of Shulman [Shu13] to come up with an enriched hyperdoctrine of quantitative predicates. We fail both times in telling ways, which is why we describe our failure: idempotency at the meta-level, manifesting in different forms, ends up ruining our plans in both cases.

**1.2. Acknowledgments.** I am deeply indebted to Ekaterina Komendatskaya, Bob Atkey, and Radu Mardare for their encouragement, support and advice regarding this work. We also thank Toby St Clere Smithe, Owen Lynch, David Jaz Myers, Kevin Carlson, Paolo Perrone, Frederik Nordvall-Forsberg and Neil Ghani for many conversations which helped shape this investigation.

## 2. THE EXTENDED POSITIVE REAL NUMBERS

The algebraic structure of the real numbers seems a fairly simple matter, yet it took us quite some time to reach the definition below. The reason is, as always in these cases, that many different structures are at play and one has to distillate the various tendrils to reach a satisfying state of affairs.

For the reals, there are two major confounding factors: the mix of additive and multiplicative structure and the role played by 0 and  $\pm\infty$ . In high-school, and throughout college, we are taught multiplication and addition (or, more accurately, subtraction and division) don't play nice with these extremal values. Specifically,  $0\infty$  could be both 0 and  $\infty$ , and  $\infty - \infty$  could be both  $+\infty$  or  $-\infty$ .

We make sense of this situation by first separating additive and multiplicative algebraic structures, realizing they are two faces of the same coin, related by *Napierian duality*  $-\log + 1/\exp : [-\infty, \infty]_{\oplus} \cong [0, \infty]_{\otimes}$ . Then, focusing on the multiplicative reals, we resolve the conflict of defining  $0\infty$  by weakening the structure of  $[0, \infty]_{\otimes}$  from 'compact closed' (thus admitting multiplicative inverses) to '\*-autonomous' (admitting an inverse-like duality). In practice, this doesn't weaken the structure but perfectly captures the sense in which  $[0, \infty]_{\otimes}$  allows taking inverses.

**Definition 2.1.** The poset  $([0, \infty]_{\otimes}, \leq)$  is a  $*$ -autonomous quantale, meaning:

1. it has all small joins  $\bigvee_i a_i$ , given by suprema;
2. it is equipped with a tensor product  $\otimes : [0, \infty]_{\otimes} \times [0, \infty]_{\otimes} \rightarrow [0, \infty]_{\otimes}$ , defined by multiplication extended with the rules

$$\forall a \in (0, \infty], a \otimes \infty = \infty, \quad 0 \otimes \infty = 0, \quad (2.1)$$

and satisfying the properties

- (a)  $\otimes$  distributes over joins:  $k \otimes \bigvee_i a_i = \bigvee_i k \otimes a_i$ ,
  - (b)  $\otimes$  is commutative, associative and unital, with unit  $1 \in [0, \infty]_{\otimes}$ .
3. it is equipped with a duality  $(-)^* : [0, \infty]_{\otimes} \rightarrow [0, \infty]_{\otimes}^{\text{op}}$ , defined as

$$\forall a \in (0, \infty), a^* := 1/a, \quad 0^* = \infty, \quad \infty^* = 0, \quad (2.2)$$

satisfying the property

$$\forall a \in [0, \infty]_{\otimes}, a \otimes b \leq c^* \iff a \leq (b \otimes c)^*. \quad (2.3)$$

*Remark 2.2.* The definitions of  $a \otimes \infty$  and  $0 \otimes \infty$  are dictated by the requirement  $\otimes$  preserves joins.

We remind the reader that the law  $(a \otimes b)^* = a^* \otimes b^*$  isn't required to hold. In fact, it doesn't hold in  $[0, \infty]_{\otimes}$ , since  $0 = 0 \otimes \infty = \infty^* \otimes 0^* \neq (\infty \otimes 0)^* = \infty$ . It does hold for all  $0 < a, b < \infty$  though.

The reason this doesn't hold is that a  $*$ -autonomous category (see [Bar79; BZ20]) like  $[0, \infty]_{\otimes}$  really features two distinct monoidal products, where the second is de Morgan dual to the first:  $a \otimes^* b := (a^* \otimes b^*)^*$ . In the case of  $[0, \infty]_{\otimes}$ ,  $\otimes^*$  is again given by multiplication on all finite values but, due to the fact  $\otimes^*$  is a monoidal product on the dual order  $[0, \infty]_{\otimes}^{\text{op}}$ , preservation of joins now forces the definition  $0 \otimes^* \infty = \infty$ .

The two structures lax-linearly distribute [CS97] over each other, meaning  $a \otimes (b \otimes^* c) \leq (a \otimes b) \otimes^* c$ , though equality holds for all finite values of  $a, b, c$  and only breaks for e.g.  $(a, b, c) = (0, 0, \infty)$ .

One can then define for  $a, b \in [0, \infty]_{\otimes}$ , the **internal hom**

$$a \multimap b := a^* \otimes b = \begin{cases} 0 & b = 0 \text{ or } a = \infty \\ \infty & a = 0 \text{ and } b \neq \infty \\ b/a & \text{else} \end{cases} \quad (2.4)$$

Observe that on  $[0, \infty]_{\otimes}$  we can define a second join-preserving monoidal operation, together with its induced dual:

polarity		non-linear	linear	
			additive	multiplicative
duality $a^* := 1/a$	positive	<b>false</b> := 0 $a \vee b := \max\{a, b\}$	<b>0</b> := 0 $a \oplus b := a + b$	<b>1</b> := 1 $a \otimes b := ab, 0\infty = 0$
	negative	<b>true</b> := $\infty$ $a \wedge b := \min\{a, b\}$	$\top$ := $\infty$ $a \oplus^* b := a +^* b$	$\perp$ := 1 $a \otimes^* b := ab, 0\infty = \infty$

**Table 1:** Semantics of very linear logic in  $[0, \infty]_{\otimes}$ .

**Definition 2.3.** Let  $a, b \in [0, \infty]_{\otimes}$ , their **sum** is

$$a \oplus b := \begin{cases} \infty & a = \infty \text{ or } b = \infty \\ a + b & \text{else} \end{cases} \quad (2.5)$$

while their **harmonic sum** is the de Morgan dual of their sum:

$$a \oplus^* b := (a^* \oplus b^*)^* = \begin{cases} 0 & a = 0 \text{ or } b = 0 \\ a & b = \infty \\ b & a = \infty \\ \frac{1}{1/a + 1/b} & \text{else} \end{cases} \quad (2.6)$$

If  $(a_i)_{i \in I} \in [0, \infty]_{\otimes}$  is a finite set of numbers, we write  $\bigoplus_{i \in I}^* a_i$  for their harmonic sum.

Harmonic sum is evidently associative, commutative, and unital with unit  $\infty$ .

**2.1. Very linear logic.** Hence  $[0, \infty]_{\otimes}$  supports a very rich structure: not only a tensor  $\otimes$  and a sum  $\oplus$ , but also joins  $\vee$  and a duality  $(-)^*$  which doubles them all. We summarise it in Table 1.

The various monoidal products distribute over each other in distinct ways:

$$\begin{aligned} \text{lax-linear} \quad a \otimes^* (b \otimes c) &\leq (a \otimes^* b) \otimes c & \mathbf{1} &\leq \top \\ \text{nonlinear} \quad a \otimes (b \oplus c) &= (a \otimes b) \oplus (a \otimes c) & a \otimes 0 &= 0 \\ \text{nonlinear} \quad a \otimes (b \vee c) &= (a \otimes b) \vee (a \otimes c) & a \otimes \text{false} &= \text{false} \\ & a \oplus (b \vee c) = (a \oplus b) \vee (a \oplus c) & a \oplus \text{true} &= \text{true} \end{aligned}$$

If we interpret these as logical connectives, we are faced with a version of linear logic with *three* generations of connectives: **non-linear**, **linear additive** and **linear multiplicative**. Linear ones behave like the ‘multiplicatives’ in classical linear logic, and the non-linear ones like the ‘additives’. But also linear multiplicatives and linear additives interact in the same way they do in linear logic, hence their names. Like in classical linear logic, all connectives come in two polarities which are exchanged by a formal duality, which also functions as negation.

For lack of a better name, we refer this extended system **very linear logic**.

Unfortunately, it’s not clear to us in which way the rules for linear additive connectives should distinguish them from linear multiplicative ones. In fact we are not aware of other

polarity		non-linear	linear	
			additive	multiplicative
duality $a^* := -a$	positive	$\text{false} := -\infty$ $a \vee b := \max\{a, b\}$	$\mathbf{0} := 0$ $a \oplus b := -\log(e^{-a} + e^{-b})$	$\mathbf{1} := 0$ $a \otimes b := a + b$
	negative	$\text{true} := +\infty$ $a \wedge b := \min\{a, b\}$	$\top := \infty$ $a \oplus^* b := \log(e^a + e^b)$	$\perp := 0$ $a \otimes^* b := a +^* b$

**Table 2:** Semantics of very linear logic in  $[-\infty, \infty]_{\oplus}$ .

occurrences of this kind of (likely substructural) logic. In some regards, the relation between additive and multiplicative is remindful of the logic of bunched implication [OP99], where contexts have the form of trees; but very linear logic has a further generation of linear connectives, as well as a different kind of distributivity between the connectives.

Additionally, while we don't know whether it is legitimate to call this logic linear: we don't have a proof system that justifies calling  $\oplus$  and  $\otimes$  'linear connectives'. However, it seems very likely this will be the case given what we know about Lawvere logic [BMPP24].

**2.2. Napierian duality.** Indeed, last year [BMPP23] introduced  $\mathbb{A}L$ , a propositional logic valued in  $([0, \infty], \geq, \oplus)$ , and recently [BMPP24] introduced  $\mathbb{P}L$  which extends  $\mathbb{A}L$  with  $\otimes$  (using our notation). The link with the structure we are contemplating is given by the following well-known diagram of quantales:

$$\begin{array}{ccc}
 & \text{Napierian} & \\
 & \text{duality} & \\
 \text{multiplicative} & \leftarrow \quad \quad \quad \rightarrow & \text{additive} \\
 & \begin{array}{ccc}
 & \xrightarrow{1/\text{exp}} & \\
 [0, 1]_{\otimes} & \xleftrightarrow{\sim} & [-\infty, 0]_{\oplus}^{\text{op}} \\
 & \xleftarrow{-\log} & \\
 \downarrow & & \downarrow \\
 [0, \infty]_{\otimes} & \xleftrightarrow{1/\text{exp}} & [-\infty, \infty]_{\oplus}^{\text{op}} \\
 & \xleftarrow{-\log} & \\
 \uparrow & & \uparrow \\
 [1, \infty]_{\otimes} & \xleftrightarrow{1/\text{exp}} & [0, \infty]_{\oplus}^{\text{op}} \\
 & \xleftarrow{-\log} & 
 \end{array} & (2.7) \\
 & & \\
 \text{negative} & & \text{positive} \\
 \uparrow & & \downarrow \\
 \text{de Morgan} & & \\
 \text{duality} & & 
 \end{array}$$

The logics  $\mathbb{A}L/\mathbb{P}L$  deal with the positive additive reals, while the very linear logic above is interpreted in the multiplicative reals. The reason we favour working multiplicatively rather than additively is that the three generations of structure we expounded above are more explicit in this setting. Indeed, the linear additive structure on  $[-\infty, \infty]_{\oplus}$  is awkward and it exists only because Napierian duality can be used to transport the structure from  $[0, \infty]_{\otimes}$ . In Table 2 we give the result of doing so: the linear multiplicative fragment of the additive reals is given by addition, so that the linear additive operation has to be something in-between, an operation known as log-sum-exp [NS16].

Since we are ultimately interesting in understanding quantification, and  $\sum$  is a clear candidate for the role of  $\exists$ , we are particularly interested in working in a setting where the linear

additive structure is natural. Moreover, we prefer using both negative and positive values than limiting ourselves to positives as to have available not only degrees of truth but also of falsehood.

*Remark 2.4.* The very existence of two ‘worlds’ of quantitative operations, related by a perfect duality, is conceptually and mathematically stimulating. One can observe this duality manifesting in various occurrences, where multiplicative quantities (e.g. probabilities, likelihood, perplexity, diversity, relative errors, ratios) and additive ones (e.g. logits, information, entropy, energy, absolute errors, distance) interact in a controlled way. In particular this dichotomy makes one pay attention to the different meanings attached to real numbers and their operation. This is particularly intriguing in the twofold nature of sum: sum of additive quantities is different, conceptually, from the sum of multiplicative ones.

**2.3. Degrees of truth.** A few lines above we noted we can think of the numbers in  $[0, \infty]_{\otimes}$  as truth values. To convert these values back to classical ones, we can borrow a technique from multivalued logic and use maps into the quantale of (meta-level) truth values  $\mathbf{Prop}_{\wedge}$ :

**Definition 2.5.** A **separator** in a quantale  $Q_{\otimes}$  is a map of quantales  $[-] : Q_{\otimes} \rightarrow \mathbf{Prop}_{\wedge}$ .

A separator casts each  $q \in Q$  to an actual truth value  $[q] \in \mathbf{Prop}$ . This defines a subset  $S = \{q \mid [q] = \top\}$  of **qualitatively true** elements which is upper-closed, closed under  $\otimes$ , and contains the top element, i.e. an **ideal** in  $Q_{\otimes}$  (which is the quantalic analogue to a *filter*). *Vice versa*, any ideal  $F$  defines a separator by the map  $_{-} \in F$ .

It’s easy to classify the ideals of  $[0, \infty]_{\otimes}$ : there is one big family given by choosing a ‘threshold for truth’  $t \geq 1$  (possibly  $t = \infty$ ), the one according to which every non-zero value is true, and the inconsistent one for which everything is true:

$$[t, \infty], t \geq 0 \quad (0, \infty] \quad [0, \infty] \quad (2.8)$$

We call the ideal  $[1, \infty]$  **unitary**. In general, is the ideal defined by the upperset of the unit of  $\otimes$ . It corresponds to ‘positive numbers’: indeed this (after passing through  $-\log$ ) is the choice (implicitly) made for  $\mathcal{AL}$  and  $\mathcal{PL}$ .

*Remark 2.6.* Enriched category theory [Kel82] is often done over a quantalic base, or the categorified variant, Bénabou cosmoi. In that context, the unitary ideal has a special role, e.g. that’s what’s used to define the ‘underlying category’ of an enriched category, but notice that any ideal would do: given a separator  $S : \mathcal{V} \rightarrow \mathbf{Set}$ , the associated notion of underlying category is given by change of base along  $S$ .

### 3. PREDICATES AND QUANTIFICATION

As anticipated, sums emerge as natural candidates for quantifiers: they seem to aggregate evidence by summing up the values of a ‘quantitative predicate’  $\varphi : I \rightarrow [0, \infty]_{\otimes}$ . This is hardly an original observation: the correspondence between the symbolic and the quantitative has

long lived in the collective subconscious of mathematicians, but also surfaced consciously in places like e.g. [Koz83; PT21].

Let us highlight some easy-to-verify properties of  $\bigoplus$  and its dual,  $\bigoplus^*$ , which hint at their role as quantifiers:

**Lemma 3.1.** *Harmonic sum*

1. *satisfies the identity:*

$$\bigoplus_{i \in I}^* \frac{a}{b_i} = \frac{a}{\bigoplus_{i \in I} b_i} \quad (3.1)$$

2. *satisfies the Fubini property:*

$$\bigoplus_{i \in I}^* \bigoplus_{j \in J}^* a_{ij} = \bigoplus_{(i,j) \in I \times J}^* a_{ij} = \bigoplus_{j \in J}^* \bigoplus_{i \in I}^* a_{ij}, \quad (3.2)$$

3. *is homogeneous (i.e. multiplication distributes over it):*

$$k \otimes \bigoplus_{i \in I}^* a_i = \bigoplus_{i \in I}^* k \otimes a_i, \quad k \otimes \bigoplus_{i \in I}^* a_i = \bigoplus_{i \in I}^* k \otimes a_i, \quad (3.3)$$

implying moreover

$$\left( \bigoplus_{i \in I}^* a_i \right) \left( \bigoplus_{j \in J}^* b_j \right) = \bigoplus_{i \in I}^* \bigoplus_{j \in J}^* a_i b_j, \quad (3.4)$$

4. *is monotonic in the argument: if, for each  $i \in I$ ,  $a_i \leq b_i$ , then*

$$\bigoplus_{i \in I}^* a_i \leq \bigoplus_{i \in I}^* b_i, \quad (3.5)$$

5. *is antitonic in the index: when  $J \subseteq I$ , one has*

$$\bigoplus_{i \in I}^* a_i \leq \bigoplus_{j \in J}^* a_j. \quad (3.6)$$

The identity (3.1), which we call the **fundamental property of harmonic sums**, can be written more suggestively as

$$\bigoplus_i^* (b_i \multimap a) = \left( \bigoplus_i b_i \right) \multimap a, \quad (3.7)$$

which is remindful of the usual  $\forall i (b_i \rightarrow a) = (\exists i b_i) \rightarrow a$ . Homogeneity can also be recast as

$$\bigoplus_i^* (b \multimap a_i) = b \multimap \left( \bigoplus_i^* a_i \right), \quad (3.8)$$

which is analogous to  $\forall i (b \rightarrow a_i) = b \rightarrow (\forall i a_i)$ .

In fact,  $\bigoplus / \bigoplus^*$  and  $\exists / \forall$  sit together in a spectrum:



**Definition 3.2** (*p*-sum). For any  $p \in (-\infty, \infty)$ ,  $p \neq 0$ , the *p*-**sum** of a finite set of numbers  $(a_i)_{i \in I}$  is

$$\bigoplus_{i \in I}^p a_i := \left( \bigoplus_{i \in I} a_i^p \right)^{1/p}. \quad (3.9)$$

To refer explicitly to negative *p* case, we talk of **harmonic p-sums** instead of  $(-p)$ -sums.

*Remark 3.3.* Like harmonic sum is the conjugate of arithmetic sum under taking reciprocals (i.e. under the automorphism  $(-)^{-1}$  of  $[0, \infty]_{\otimes}$ ), *p*-sums are the conjugate of arithmetic sum under the automorphisms  $(-)^p$ . These are the only continuous automorphisms of  $[0, \infty]_{\otimes}$ .

One extends the above definition to  $p = \pm\infty$  (but not  $p = 0$ ) by taking suitable limits (see [MV70, p. 75]). Then varying *p* we recover sum ( $p = 1$ ), harmonic sum ( $p = -1$ ),  $\bigwedge$  ( $p = -\infty$ ) and  $\bigvee$  ( $p = +\infty$ ), the latter two being the semantic counterpart of  $\forall$  and  $\exists$ :

$$\begin{array}{ccccccccccc} \bigwedge & \dots & \bigoplus^{-p} & \dots & \bigoplus^* & \not\exists & \bigoplus & \dots & \bigoplus^p & \dots & \bigvee \\ -\infty & \xrightarrow{\hspace{2em}} & -p & \xrightarrow{\hspace{2em}} & -1 & \xrightarrow{\hspace{1em}} & 0 & \xrightarrow{\hspace{1em}} & 1 & \xrightarrow{\hspace{2em}} & p & \xrightarrow{\hspace{2em}} & \infty \end{array} \quad (3.10)$$

*Remark 3.4.* We observe that all of the items of Lemma 3.1 still hold for general *p*-sums, with the exception of monotonicity in the indexing which reverses for positive *p*. In particular, the fundamental properties (3.7) and (3.8) still put in relation  $\bigoplus^{-p}$  with  $\bigoplus^p$  for all  $p \in [-\infty, \infty]$ .

For all  $p \neq 0$ , we have

$$\bigoplus^{-p} \leq \bigwedge \leq \bigvee \leq \bigoplus^p, \quad (3.11)$$

with the gap between linear additive and non-linear narrowing as *p* grows (see Fig. 1).


Intuitively, compared to  $\exists$ , an harmonic *p*-sum looks up existence in a set by allowing ‘non-local’ effects to compensate (similarly to how existential quantification in a topos might be satisfied locally, even if not globally). Dually,  $\forall$  checks uniform satisfaction of a predicate while  $\bigoplus^{-p}$  imposes extra ‘regularity’ conditions. For instance,  $\bigoplus_{i \in I}^* a_i \geq 1$  implies that each  $a_i \geq 1$ , but that’s not sufficient: they have to be sufficiently larger than 1 as to also satisfy  $\bigoplus_{i \in I} 1/a_i \leq 1$ ! Increasing *p* makes the latter condition weaker, until we reach that of  $\bigwedge$ .

**3.1. Justifying the means.** These characteristics are why *p*-sums are so useful, however if left unbridled they make *p*-sums depend strongly on the ‘size’ of the quantification domain. Specifically, quantifying with a *p*-sum can yield arbitrarily high truth values depending only on how many ‘false’ values we sum. This is apparent in Fig. 1, where you can see the *p*-sum going literally off the charts for small values of *p*.

This isn’t really a drawback *per se*, but since there is an alternative worth exploring, we consider it. Indeed, we can compensate for size effects by taking means instead of sums.

For  $p \in (-\infty, \infty)$  but  $p \neq 0$ , let the *p*-mean of a finite set of numbers  $(a_i)_{i \in I}$  to be

$$\int_{i \in I}^p a_i := \left( \bigoplus_{i \in I}^p \frac{a_i^p}{|I|} \right)^{1/p}. \quad (3.12)$$



**Figure 1:** A comparison of  $p$ -sums (solid lines) and  $p$ -means (dashed lines) in their behaviour as  $p$  changes. One can clearly see the tendency towards max/min (dotted) as  $p \rightarrow +\infty$ , and the convergence of the  $p$ -means to the harmonic one as  $p \rightarrow 0$ .

Again, this definition can be extended to  $p = \pm\infty$  by taking suitable limits. Unlike for sums, this time the limit exists even for  $p = 0$ , and it's given by the *geometric mean* (which we denote by  $\bigoplus^0$ ):

$$\begin{array}{cccccccccccc} \wedge & \dots & \int^{-p} & \dots & \int^* & \bigoplus^0 & \int & \dots & \int^p & \dots & \vee \\ -\infty & \text{---} & -p & \text{---} & -1 & \text{---} & 0 & \text{---} & 1 & \text{---} & p & \text{---} & \infty \end{array} \quad (3.13)$$

The extraordinary fact is that  $p$ -means satisfy again the properties of Lemma 3.1, including Fubini, monotonicity in the argument and index, the fundamental relation and homogeneity.

Unlike  $p$ -sums however, their values sit in between those of the traditional, non-linear quantifiers, with the gap closing as  $p$  grows:

$$\bigwedge \leq \bigoplus^{-p} \leq \bigoplus^p \leq \bigvee, \quad (3.14)$$

Thus, unlike  $p$ -sums,  $p$ -means compensate for size effects.

The intuition we propose is that sums and means are, respectively, unbounded and bounded quantifiers.

$$\begin{array}{l} \exists i. a(i) \quad \leftarrow \rightsquigarrow \quad \bigoplus_{i \in I}^p a(i) \\ \exists (i \in I). a(i) \quad \leftarrow \rightsquigarrow \quad \int_{i \in I}^p a(i) \equiv \left( \bigoplus_{i \in I}^p \frac{1}{|I|} \otimes a(i) \right)^{1/p} \end{array} \quad (3.15)$$

In fact a mean is a particular instance of a weighted mean, which is an integral over a probability space  $(I, di)$ :

$$\exists (i \in I). a(i) \quad \leftarrow \rightsquigarrow \quad \int_{i \in I}^p a(i) di. \quad (3.16)$$

The mass element  $di$  behaves as the bound  $i \in I$  in a quantification. The fact that this is part of the quantifier can be seen in two different ways. First, on the syntactic side, we observe that  $di$  doesn't participate in the duality relating  $\bigoplus_{i \in I}^p$  and  $\bigoplus_{i \in I'}^{-p}$ , like  $i \in I$  doesn't for  $\exists(i \in I)/\forall(i \in I)$ :

$$\exists(i \in I). a(i) = \neg \forall(i \in I) \neg a(i) \quad \longleftrightarrow \quad \int_{i \in I}^p a(i) di = \left( \int_{i \in I}^{-p} a(i)^* \right)^*. \quad (3.17)$$

Secondly, on the semantic side, integrating with respect to  $di$  is a more general operation than summing expressions such as  $a(i) \otimes di$  since  $di$  might not admit a density, witnessing the fact that, generally speaking, it is part of the quantification itself and not of the argument.

Notice, moreover, that sums are particular kinds of integral too: they are integrals over non-unitary measure spaces. Therefore, we propose real-valued quantification should correspond to integration.

**3.2. A sketch of syntax and semantics.** To formalise the analogies sketched above, we need first to formalise what predicates are.

We start by saying that each measure space  $(I, di)$  is a **context** (or **domain of discourse**). **Multiplicative predicates** over each context are defined by the following inductive definition:

$$\begin{aligned} \varphi(i) := & \text{const} \mid a \\ & \mid \varphi(i) \text{op} \varphi(i) \mid \varphi(i) \multimap \varphi(i) \mid \varphi(i)^* \\ & \mid k \cdot \varphi(i) \\ & \mid \exists^p(j \in J). \psi(i, j) \mid \forall^p(j \in J). \psi(i, j) \end{aligned} \quad (3.18)$$

where

- (i)  $\text{const}$  stands for any of the constants **false**, **true**, **0**, **1**,  $\top$ ,  $\perp$  introduced in Table 1,
- (ii)  $a$  ranges in measurable functions  $I \rightarrow [0, \infty]_{\otimes}$ ,  $\text{op} \in \{\vee, \wedge, \oplus, \oplus^*, \otimes, \otimes^*\}$ ,
- (iii)  $k \in [0, \infty]$ ,
- (iv)  $p \in [0, \infty]$ , and  $\psi$  is another predicate over context  $I \times J$ .

We thus obtain the sets  $\text{LT}(I)_{\otimes}$ . Ideally, these would be further quotiented by syntactic rewrites or, even better, equipped with an entailment relation but it's not yet clear how this entailment would look like. We discuss this in Section 4.

Still, the **multiplicative semantics** bracket  $\llbracket - \rrbracket_{\otimes}$  can be defined inductively on  $\text{LT}(I)_{\otimes}$ . It takes values in measurable functions  $I \rightarrow [0, \infty]_{\otimes}$ , and since every measurable function determines a predicate too we sometimes terminologically conflate the two notions.

The propositional fragment is covered by Table 1 where we gave semantics for constants, negation (the duality  $(-)^*$ ), and binary operators; except for  $k \cdot \varphi(i)$  which is interpreted as

exponentiation<sup>3</sup>:

$$\llbracket k \cdot \varphi(i) \rrbracket_{\otimes} := \begin{cases} (\llbracket \varphi(i) \rrbracket_{\otimes})^k & 0 < k, \llbracket \varphi(i) \rrbracket_{\otimes} < \infty, \\ \infty & \llbracket \varphi(i) \rrbracket_{\otimes} = \infty, 0 < k \leq \infty, \\ 1 & \llbracket \varphi(i) \rrbracket_{\otimes} = \infty, k = 0, \\ 0 & \llbracket \varphi(i) \rrbracket_{\otimes} = 0. \end{cases} \quad (3.19)$$

Finally, the  $p$ -soft quantifiers are interpreted as

$$\begin{aligned} \llbracket \exists^p(j \in J). \varphi(i, j) \rrbracket_{\otimes} &:= \int_{j \in J} \llbracket \varphi(i, j) \rrbracket_{\otimes} dj, \\ \llbracket \forall^p(j \in J). \varphi(i, j) \rrbracket_{\otimes} &:= \int_{j \in J} \llbracket \varphi(i, j) \rrbracket_{\otimes} dj. \end{aligned} \quad (3.20)$$

As noted above, in the limit  $p = \pm\infty$  the right hand side *means*  $\vee$  and  $\wedge$ , and for  $p = 0$  we get  $\llbracket \forall^0 \rrbracket = \llbracket \exists^0 \rrbracket =$  the geometric mean.

*Remark 3.5.* It's clear the syntax presented in (3.18) is not minimal. For instance, we have:

$$\llbracket \exists^p(i \in I). \varphi(i) \rrbracket_{\otimes} = \llbracket \frac{1}{p} \cdot \exists^1(i \in I). p \cdot \varphi(i) \rrbracket_{\otimes}. \quad (3.21)$$

We can define **additive predicates** by the same inductive definition as in (3.18), except now constants range in  $[-\infty, \infty]_{\oplus}$ . This defines a set  $\mathbf{LT}(I)_{\oplus}$ . There is a syntactic translation  $-\log : \mathbf{LT}(I)_{\oplus} \rightarrow \mathbf{LT}(I)_{\otimes}$  given by  $-\log$  on constants and defined to commute with all the other formers (operations and quantifiers alike). The inverse syntactic translation  $1/\exp : \mathbf{LT}(I)_{\otimes} \rightarrow \mathbf{LT}(I)_{\oplus}$  is defined analogously.

Then **additive semantics**  $\llbracket - \rrbracket_{\oplus}$  takes values in measurable functions  $I \rightarrow [-\infty, \infty]_{\oplus}$ . It can be defined from the multiplicative one by Napierian duality:

$$\llbracket \varphi(i) \rrbracket_{\oplus} := -\log \llbracket 1/\exp(\varphi(i)) \rrbracket_{\otimes} \quad (3.22)$$

On the propositional fragment, the effect is described by Table 2, then we have (note  $k \geq 0$ ):

$$\llbracket k \cdot \varphi(i) \rrbracket_{\oplus} := k \llbracket \varphi(i) \rrbracket_{\oplus}. \quad (3.23)$$

and  $p$ -soft quantifiers become:

$$\begin{aligned} \llbracket \exists^p(i \in I). \varphi(i) \rrbracket_{\oplus} &:= -\frac{1}{p} \log \int_{i \in I} e^{-p \llbracket \varphi(i) \rrbracket_{\oplus}} di, \\ \llbracket \forall^p(i \in I). \varphi(i) \rrbracket_{\oplus} &:= +\frac{1}{p} \log \int_{i \in I} e^{+p \llbracket \varphi(i) \rrbracket_{\oplus}} di. \end{aligned} \quad (3.24)$$

Again, in the limit  $p = \pm\infty$  these recover  $\vee$  and  $\wedge$ , and for  $p = 0$  we get  $\llbracket \forall^0 \rrbracket = \llbracket \exists^0 \rrbracket = \int_{i \in I}$ .

As far as we know, these expressions haven't been considered *as quantifiers* before, though, as we are going to see now, they are not new to mathematics.

---

<sup>3</sup>Notice that this definition defines an action of  $[0, \infty]_{\otimes}$  onto itself, since  $a^1 = a$  and  $(a^h)^k = a^{h \otimes k}$ .

**3.3. Quantifiers in the wild.** We have one important reason to believe the quantifiers of (3.24) (and by extension the original multiplicative ones of (3.20)) are interesting, which is their appearance in statistics and machine learning.

**3.3.1. Argmax and softmax.** The softmax operator is used in machine learning to turn a vector of real numbers into a probability distribution, and in statistical mechanics, where it's known as *Gibbs* or *Boltzmann distribution* to get a probability distribution out of a positive energy functional.

**Definition 3.6.** Let  $f : X \rightarrow [0, \infty]_{\otimes}$  be a function. Its **softmax** is the function  $X \rightarrow [0, 1]_{\otimes}$

$$(\text{softmax } f)(x^*) = \frac{f(x^*)}{\int_{x \in X} f(x)}. \quad (3.25)$$

On the other hand, in the Boolean world we have a similar construction:

$$(\text{argmax } f)(x^*) = \forall x \in X, f(x) \leq f(x^*). \quad (3.26)$$

We get a 'soft version' by replacing hard quantifiers with soft one. Define:

$$(p\text{-softmax } f)(x^*) = \llbracket \forall^p(x \in X). f(x) \multimap f(x^*) \rrbracket_{\otimes} = \int_{x \in X}^{-p} \frac{f(x^*)}{f(x)}. \quad (3.27)$$

Then it's easy to see that  $\text{softmax} = 1\text{-softmax}$ , whereas  $\text{argmax} = \lfloor \infty\text{-softmax} \rfloor$ :

$$\lfloor (\infty\text{-softmax } f)(x^*) \rfloor = \lfloor \bigwedge_{x \in X} f(x) \multimap f(x^*) \rfloor = \lfloor \bigwedge_{x \in X} f(x^*)/f(x) \rfloor = (\text{argmax } f)(x^*). \quad (3.28)$$

where the map  $\lfloor - \rfloor : [0, \infty]_{\otimes} \rightarrow 2$  is the unitary separator of Section 2.3. Indeed, the ratio  $f(x^*)/f(x)$  is greater or equal than 1 iff  $x^*$  is a maximum point, thus yielding  $\top$  for those points. Else, it is always strictly below 1 and yields  $\perp$ .

More commonly, softmax is applied to functions of arbitrary sign by first passing them through Napierian duality, i.e. usually  $f = e^{-u}$  in (3.25). The decaying exponential is antitone which means that softmax, for functions so constructed, puts more mass on small values of  $u$  than large ones, making it more like a 'softmin'. In any case, we still want to 'extract back' the distribution obtained by softmax, thus it stands to reason to apply  $-\log$  to the result:

$$L(x^*) = -\log \text{softmax } e^{-u(x^*)} = u(x^*) - \log \bigoplus_{x \in X} e^{-u(x)} = \llbracket \forall^1(x \in X). u(x^*) \multimap u(x) \rrbracket_{\oplus}. \quad (3.29)$$

The quantity  $L$  is called **log-likelihood**, and we just reconstructed it by purely formal considerations.

*Remark 3.7.* Log-likelihood, softmax and, generally speaking, the operations defining additive  $p$ -soft existential quantification (3.24) are all known in the machine learning community under the self-descriptive name of "log-sum-exponential" (LSE) [NS16], also *softplus* [GBB11]. As expounded in [BKBP22], while  $\text{ReLU}(x) = 0 \vee x$  is a popular and effective choice of activation function, it is plagued by various numerical problems which can be mitigated by replacing it with a smooth approximant. LSE seems a well-motivated candidate for such a smoothing.

3.3.2. *Entropy.* Consider a predicate  $\varphi : I \rightarrow [0, \infty]_{\otimes}$ . For simplicity, assume  $\varphi$  is negative and unitary, meaning it lands in  $[0, 1]_{\otimes}$  and  $\int_i \varphi(i) = 1$ . The **Rényi entropy** of order  $p \in [0, \infty]$  [Lei21, Definition 4.3.1] is the quantity

$$H_p(\varphi) = \frac{1}{1-p} \log \int_{i \in I} \varphi(i)^p. \quad (3.30)$$

When  $p = 0, 1, \infty$  (3.30) must be interpreted in the limit. Notably, for  $p = 1$ ,  $H_p(\varphi)$  recovers **Shannon entropy**.

The quantity  $H_p(\varphi)$  is the additive semantics of  $p$ -universally quantifying the negation of  $-\log \varphi$ , up to a multiplicative constant:

$$H_p(\varphi) = \llbracket \frac{p}{1-p} \cdot (\forall^p(i \in I). -\log \varphi(i)^*) \rrbracket_{\oplus} \quad (3.31)$$

This expression showcases some of the ideas regarding additive/multiplicative type discipline we talked about in Remark 2.4. Indeed, the logarithm appearing in (3.31) is very natural and can formally be justified as the necessary casting from the multiplicative world of  $\varphi$  to the additive world—indeed, entropy is an additive quantity.

In fact, one could define a purely multiplicative quantity:

$$D_p(\varphi) = \llbracket \frac{p}{1-p} \cdot (\forall^p(i \in I). \varphi(i)^*) \rrbracket_{\otimes}. \quad (3.32)$$

This is known as  **$p$ -diversity** (or  **$p$ -th Hill number**) [Lei21, Definition 4.3.4].

#### 4. TOWARDS A CATEGORICAL SEMANTICS

The definition of quantifiers we gave above seems to be directly in the groove of an algebraic account of first-order predicate logic, where contexts are provided by measure spaces and algebras of predicates are given by the (wannabe) quantales  $\mathbf{LT}(I)_{\otimes}$  and take semantics in analogous quantale of real-valued random variables.

We focus now on the latter, since that's a setting we can test our guesses on. Define the  $[0, \infty]_{\otimes}$ -enriched *quantale of functionals*  $\mathcal{M}^p(I, [0, \infty]_{\otimes})$  with entailment relation tentatively given by

$$\varphi \vdash_I \psi := \llbracket \forall^p(i \in I). \varphi(i) \multimap \psi(i) \rrbracket_{\otimes} = \left( \int_{i \in I} \frac{\varphi(i)^p}{\psi(i)^p} di \right)^{-1/p}. \quad (4.1)$$

This seems to define the action on objects as a functor  $\mathcal{M}^p : \mathbf{Meas}^{\text{op}} \rightarrow [0, \infty]_{\otimes}\text{-Cat}$ . The action on measure non-increasing maps  $f : J \rightarrow I$  is readily given by precomposition, which is indeed an enriched functor  $f^* : \mathcal{M}^p(I) \rightarrow \mathcal{M}^p(J)$ , since, for every  $\varphi, \psi \in \mathcal{M}^p(I)$ , we have

$$\varphi \vdash_I \psi \leq f^* \varphi \vdash_J f^* \psi \rightsquigarrow \left( \int_{i \in I} \frac{\varphi(i)^p}{\psi(i)^p} di \right)^{-1/p} \leq \left( \int_{j \in J} \frac{\varphi(f(j))^p}{\psi(f(j))^p} dj \right)^{-1/p}. \quad (4.2)$$

The proof hinges on the fact that, if  $u : I \rightarrow [0, \infty]_{\otimes}$  is a measurable function, then  $\int_j u(f(j)) dj = \int_i u(i) d(f_* dj) \leq \int_i u(i) di$  (and the inequality reverses when we take reciprocals, as we do above).

Moreover, we see enriched left and right adjoints to reindexing are quantification, a property that famously characterises quantifiers [Law69]. For simplicity, let  $f = \pi_I : I \times K \rightarrow I$ , then we have  $\int_k^P \dashv \pi_I^* \dashv \int_I^P$  since for all  $\rho \in \mathcal{M}^p(I \times K)$ ,  $\psi \in \mathcal{M}^p(I)$ , we have

$$\int_{k \in K}^P \rho(i, k) dk \dashv_I \psi(i) = \rho(i, k) \dashv_{I \times K} \pi_I^* \psi \quad (4.3)$$

which unpacks to

$$\int_{i \in I}^{-p} \left( \int_{k \in K}^p \rho(i, k) dk \right) \multimap \psi(i) di = \int_{i \in I}^{-p} \rho(i, k) \multimap \psi(i) dk di \quad (4.4)$$

and this identity is true by the fundamental property of harmonic  $p$ -means. Similarly, we can conclude  $\pi_I^* \dashv \int_k^P$ . Thus it would seem that  $\mathcal{M}^p$  could be used a semantics for first-order very linear logic with  $\exists^p / \forall^p$  sent to  $\int^p / \int^{-p}$ .

Unfortunately, this approach is broken: the entailment relation we proposed isn't a proper enriched preorder since it doesn't satisfy reflexivity nor transitivity. Reflexivity says

$$\varphi \dashv_I \varphi = \left( \int_{i \in I} \frac{\varphi(i)^p}{\varphi(i)^p} di \right)^{-1/p} = |I|^{-1/p} \geq 1 \quad (4.5)$$

and it is satisfied only when  $|I| \leq 1$ , thus when  $I$  is a probability spaces. But even if restricted to those, transitivity still breaks. Fixing  $p = 1$  for simplicity, notice first that

$$(\varphi \dashv_I \psi) \otimes (\psi \dashv_I \sigma) \leq (\varphi \dashv_I \sigma) \iff (\varphi \dashv_I \psi)^{-1} \otimes (\psi \dashv_I \sigma)^{-1} \geq (\varphi \dashv_I \sigma)^{-1}. \quad (4.6)$$

However, unpacking definitions, we see that:

$$\int_{i \in I} \int_{i' \in I} \frac{\varphi(i) \psi(i')}{\psi(i) \sigma(i')} di di' \geq \int_{i \in \Delta_I} \frac{\varphi(i) \psi(i)}{\psi(i) \sigma(i)} di \not\geq \int_{i \in I} \frac{\varphi(i)}{\sigma(i)} di. \quad (4.7)$$

From a logical standpoint, we are witnessing the failure of  $\dashv_I$  to satisfy the identity axiom and the cut rule, at least if expressed naively.

**4.1. The indexed way.** One might try to fix the previous attempt by varying the base of enrichment along with the indexing—the rationale being that prematurely taking integrals when defining entailment relation is what breaks the approach.

This kind of variable enrichment has been invented by Shulman in [Shu13], where he describes enrichment of an indexed category in an indexed monoidal category.

In our case the basis of enrichment is the following (we stick to  $p = 1$ ):

**Definition 4.1.** For each  $\sigma$ -finite<sup>4</sup> measure space  $(I, di)$ , the **quantale of functionals** on  $I$  is the set  $\mathcal{M}^1(I)$  of measurable functions  $I \rightarrow [0, \infty]_{\otimes}$  with essential pointwise ordering, meaning  $\varphi \leq \psi$  iff  $\varphi$  bounds  $\psi$  from below pointwise almost-everywhere, and pointwise multiplication (using  $\otimes$ ).

<sup>4</sup>Meaning it can be covered by a countable family of finite measure spaces: it is the technical requirement for the existence of Radon–Nikodym derivatives which we employ below.

Evidently, each quantale  $\mathcal{M}(I)$  is closed and  $*$ -autonomous, and in fact has the same structure as  $[0, \infty]_{\otimes}$ , inherited pointwise. The fact that  $\mathcal{M}(I)$  is complete and cocomplete is [Mey91, Lemma 2.6.1]. Notice, in  $\mathcal{M}(I)$ ,  $\varphi \cong \psi$  when  $\varphi$  equals  $\psi$  almost everywhere on  $I$ .

We would like to define the **indexed quantale of functionals** as the pseudofunctor

$$\mathcal{M} : \mathbf{Meas}^{\text{op}} \longrightarrow \mathbf{Qnt} \quad (4.8)$$

defined as above on objects and acting on measure non-increasing functions  $f : I \rightarrow J$  by precomposition. Thus  $f^* : \mathcal{M}(J) \rightarrow \mathcal{M}(I)$  sends  $\psi$  to  $\psi(f)$ .

This definition is clearly well-posed, since  $f^*$  is strict functorial as well as strict monoidal:

$$((f^* \varphi) \otimes (f^* \psi))(i) = \varphi(f(i)) \otimes \psi(f(i)) = f^*(\varphi \otimes \psi)(i). \quad (4.9)$$

Now, still following [Shu13, Definition 4.1], we self-enrich  $\mathcal{M}$  to make it into an indexed  $\mathcal{M}$ -category. The data of such an enrichment, which we denote  $\mathcal{M}_{\mathcal{M}}$ , consists of

1. An  $\mathcal{M}(I)$ -enriched category  $\mathcal{M}(I)_{\mathcal{M}(I)}$  for each  $I \in \mathbf{Meas}$ , and this enrichment is given by the fact  $\mathcal{M}(I)$  is closed monoidal:

$$[\varphi, \psi]_{\mathcal{M}(I)_{\mathcal{M}(I)}} := \lambda(i \in I). \varphi(i) \multimap \psi(i). \quad (4.10)$$

2. For each  $f : I \rightarrow J$ , a fully faithful  $\mathcal{M}(I)$ -functor  $\overline{f^*} : \mathcal{M}(J)_{f^*} \rightarrow \mathcal{M}(I)_{\mathcal{M}(I)}$ , where the domain is change of enrichment for  $\mathcal{M}(J)$  along  $f^*$ . This means the hom-objects of this  $\mathcal{M}(I)$ -category are

$$[\varphi, \psi]_{\mathcal{M}(J)_{f^*}} := \lambda(i \in I). \varphi(f(i)) \multimap \psi(f(i)). \quad (4.11)$$

Then  $\overline{f^*}$  is defined on objects by precomposition with  $f$ , and on hom-objects is given by the identity

$$[\varphi, \psi]_{\mathcal{M}(J)_{f^*}} = [f^* \varphi, f^* \psi]_{\mathcal{M}(I)_{\mathcal{M}(I)}}. \quad (4.12)$$

This data is also subject to coherence conditions which mostly trivialise for  $\mathcal{M}_{\mathcal{M}}$ , and thus we have a well-defined indexed quantale.

Instead, one could try defining a covariant indexing

$$\mathcal{M} : \mathbf{Meas} \longrightarrow \mathbf{Qnt} \quad (4.13)$$

defined as above on objects and on measure non-increasing functions  $f : I \rightarrow J$  defined as the monoidal monotone function  $f_! : \mathcal{M}(I) \rightarrow \mathcal{M}(J)$  given by Radon–Nikodym derivative of the pushforward of measures:

$$f_! \varphi := \frac{df_*(\varphi \cdot di)}{dj} \quad (4.14)$$

where  $\varphi \cdot di$  denotes the measure obtained by integrating  $\varphi$  against  $di$ .

Thus  $f_! \varphi$  is the density of the function  $\int_{i \in f^{-1}(-)} \varphi(i) di$ .



Pseudofunctoriality of  $\mathcal{M}$  follows from applying the definition of Radon–Nikodym derivative and functoriality of pushforward of measures:

$$g_! f_! \varphi = \frac{dg_* \left( \frac{df_*(\varphi \cdot di)}{dj} \cdot dj \right)}{dk} = \frac{dg_*(f_*(\varphi \cdot di))}{dk} = \frac{d(fg)_*(\varphi \cdot di)}{dk} = (fg)_! \varphi. \quad (4.15)$$

and the lack of strictness comes from the definition of Radon–Nikodym derivative only up to almost-everywhere equality.

Lax monoidality is where the definition fails, in fact we have:

$$\begin{aligned} (f_! \varphi) \otimes (f_! \psi) \cdot dj &= (f_! \varphi) \cdot f_*(\psi \cdot di) \\ &= \int_{j \in (-)} \frac{df_*(\varphi \cdot di)}{dj}(j) f_*(\psi \cdot di) \\ &= \int_{i \in f^{-1}(-)} \frac{df_*(\varphi \cdot di)}{dj}(f(i)) \psi(i) di \\ &\not\leq \int_{i \in f^{-1}(-)} \varphi(i) \psi(i) di \\ &= f_!(\varphi \otimes \psi) \cdot dj \end{aligned} \quad (4.16)$$

We would like to use the inequality  $\frac{df_*(\varphi \cdot di)}{dj}(f(\bar{i})) \leq \varphi(\bar{i})$ , i.e. that  $\int_{i \in f^{-1}(f(\bar{i}))} \varphi(i) di \leq \varphi(\bar{i})$ . But the fact  $f^{-1}f(\bar{i}) \supseteq \{\bar{i}\}$  can't be used to prove either direction of such an inequality since  $\int_{i \in \{\bar{i}\}} \varphi(i) di \neq \varphi(\bar{i})$  in general.

Thus the desired pseudofunctor  $\mathcal{M} : \mathbf{Meas} \rightarrow \mathbf{Qnt}$  doesn't exist.

*Remark 4.2.* One might object that  $f_!$  is supposed to be left adjoint to  $f^*$ , thus colax monoidal, and indeed the fact we are comparing integrals over  $f^{-1}f(\bar{i}) \supseteq \{\bar{i}\}$  seems to suggest this is the case. However, that direction also fails since we can't compare evaluation at a point with integrals on a neighborhood in general.

## 5. CONCLUSIONS

We described the syntax and semantics of a *quantitative predicate logic* (QPIL), which naturally arises from the algebraic structure of the reals. We made our case that a logical interpretation can be given to some objects, like softmax and entropy, which are widely employed in non-symbolic quantitative reasoning in statistics, machine learning, physics, etc.

We exhibited a syntax and a semantics for QPIL, but we couldn't come up with a convincing 'proof theory' for it. In fact, the usual Boole–Gentzen–Lawvere algebraic framework for sequent calculus turned out to be inadequate, as testified by the failure of hyperdoctrines to capture QPIL. We are in need of a different algebraic structure corresponding to the rules of QPIL, capable of accomodating the lack of idempotency in its interpretation of identity and cut rules for entailment.

Indeed, the influence a metatheory exerts on the theories it hosts is a fundamental aspect which is often underplayed. Hosting an unapologetically quantitative theory within a traditional qualitative metatheory seems very problematic, and we are thus left with a hard

bootstrapping problem: that of formulating both a theory and a metatheory apt to host it at the same time.

This obstacle is the only thing preventing us from declaring QPIL a theory of quantitative *reasoning*. As the present work stands, this is a theory for quantitative syntax.

Finally, we haven't covered an interesting use we can make of QPIL, which is to do category theory 'enriched' in it. The result is to be able to formulate universal properties as equations some quantitative object has to satisfy.

We had a taste of this idea in (3.7), the fundamental property of harmonic sum: the universal characterization of a quantitative object becomes an *equation* that can be *solved* to yield the desired object.<sup>5</sup> This enables to characterise some constructions: for instance, argmax can be characterised as a certain right Kan lift in the bicategory **Rel**. The same universal property, written in QPIL, yields softmax as its unique solution.

We are particularly interested in clarifying the universal properties of conditionals [Fri20] and updates [Jac19; DR23] in statistics: can we frame them as universal reasoning rules, in a logical sense?

#### REFERENCES

- [Bar79] M. Barr. *\*-Autonomous Categories*. Vol. 752. Lecture Notes in Mathematics. Berlin, Heidelberg: Springer, 1979. ISBN: 978-3-540-09563-7 978-3-540-34850-4. DOI: [10.1007/BFb0064579](https://doi.org/10.1007/BFb0064579). URL: <http://link.springer.com/10.1007/BFb0064579> (visited on 03/20/2024).
- [BKBP22] K. Biswas et al. *SMU: smooth activation function for deep networks using smoothing maximum technique*. en. arXiv:2111.04682 [cs]. Apr. 2022. URL: <http://arxiv.org/abs/2111.04682> (visited on 04/16/2024).
- [BMPP23] G. Bacci et al. *Propositional Logics for the Lawvere Quantale*. arXiv:2302.01224 [cs]. Feb. 2023. DOI: [10.48550/arXiv.2302.01224](https://doi.org/10.48550/arXiv.2302.01224). URL: <http://arxiv.org/abs/2302.01224> (visited on 03/01/2023).
- [BMPP24] G. Bacci et al. *Polynomial Lawvere Logic*. en. arXiv:2402.03543 [cs]. Feb. 2024. URL: <http://arxiv.org/abs/2402.03543> (visited on 02/12/2024).
- [BMW85] M. Barr, C. McLarty, and C. Wells. "Variable set theory". In: *Unpublished manuscript* (1985). URL: <https://www.math.mcgill.ca/barr/papers/vst.pdf> (visited on 05/10/2024).
- [Boo47] G. Boole. *The Mathematical Analysis of Logic*. English. Barclay, Macmillan and Co., 1847. URL: <https://www.gutenberg.org/ebooks/36884> (visited on 06/03/2024).

---

<sup>5</sup>In truth, a similar phenomenon holds in traditional settings, e.g. a product is made of tuples because it is universally equipped with projections.

- [BZ20] N. Blanco and N. Zeilberger. “Bifibrations of Polycategories and Classical Linear Logic”. In: *Electronic Notes in Theoretical Computer Science*. The 36th Mathematical Foundations of Programming Semantics Conference, 2020 352 (Oct. 2020), pp. 29–52. ISSN: 1571-0661. DOI: [10.1016/j.entcs.2020.09.003](https://doi.org/10.1016/j.entcs.2020.09.003). URL: <https://www.sciencedirect.com/> (visited on 03/20/2024).
- [Cox46] R. T. Cox. “Probability, Frequency and Reasonable Expectation”. In: *American Journal of Physics* 14.1 (Jan. 1946), pp. 1–13. ISSN: 0002-9505. DOI: [10.1119/1.1990764](https://doi.org/10.1119/1.1990764). URL: <https://doi.org/10.1119/1.1990764> (visited on 06/03/2024).
- [CS97] R. Cockett and R. Seely. “Weakly distributive categories”. In: *Journal of Pure and Applied Algebra* 114 (Nov. 1997), pp. 133–173. DOI: [10.1016/0022-4049\(95\)00160-3](https://doi.org/10.1016/0022-4049(95)00160-3).
- [DP22a] F. Dagnino and F. Pasquali. “Logical Foundations of Quantitative Equality”. In: *Proceedings of the 37th Annual ACM/IEEE Symposium on Logic in Computer Science*. LICS ’22. New York, NY, USA: Association for Computing Machinery, Aug. 2022, pp. 1–13. ISBN: 978-1-4503-9351-5. DOI: [10.1145/3531130.3533337](https://doi.org/10.1145/3531130.3533337). URL: <https://doi.org/10.1145/3531130> (visited on 03/03/2023).
- [DP22b] F. Dagnino and F. Pasquali. *Quantitative Equality in Substructural Logic via Lipschitz Doctrines*. arXiv:2110.05388 [cs, math]. Nov. 2022. DOI: [10.48550/arXiv.2110.05388](https://doi.org/10.48550/arXiv.2110.05388). URL: <http://arxiv.org/abs/2110.05388> (visited on 03/01/2023).
- [DR23] E. Di Loreto and M. Román. *Evidential Decision Theory via Partial Markov Categories*. en. arXiv:2301.12989 [cs, math]. Apr. 2023. URL: <http://arxiv.org/abs/2301.12989> (visited on 03/13/2024).
- [FB22] D. Figueroa and B. van den Berg. “A topos for continuous logic”. In: *Theory and Applications of Categories* 38.28 (2022). Published 2022-09-07, pp. 1108–1135. URL: <http://www.tac.mta.ca/tac/volumes/38/28/38-28.pdf>.
- [Fri20] T. Fritz. “A synthetic approach to Markov kernels, conditional independence and theorems on sufficient statistics”. In: *Advances in Mathematics* 370 (Aug. 2020). arXiv:1908.07021 [cs, math, stat], p. 107239. ISSN: 00018708. DOI: [10.1016/j.aim.2020.107239](https://doi.org/10.1016/j.aim.2020.107239). URL: <http://arxiv.org/abs/1908.07021> (visited on 09/27/2023).
- [GBB11] X. Glorot, A. Bordes, and Y. Bengio. “Deep sparse rectifier neural networks”. In: *Proceedings of the fourteenth international conference on artificial intelligence and statistics*. JMLR Workshop and Conference Proceedings, 2011, pp. 315–323. URL: <http://proceedings.mlr.press/v15/glorot11a> (visited on 04/16/2024).
- [Gra07] M. Grandis. “Categories, norms and weights”. en. In: *Journal of Homotopy and Related Structures* 2 (2007).
- [Jac19] B. Jacobs. “The mathematics of changing one’s mind, via Jeffrey’s or via Pearl’s update rule”. In: *Journal of Artificial Intelligence Research* 65.1 (May 2019), pp. 783–806. ISSN: 1076-9757. DOI: [10.1613/jair.1.11349](https://doi.org/10.1613/jair.1.11349). URL: <https://doi.org/10.1613/jair.1.11349> (visited on 04/16/2023).

- [Kel82] M. Kelly. *Basic concepts of enriched category theory*. Vol. 64. CUP Archive, 1982.
- [Koz83] D. Kozen. “A probabilistic PDL”. en. In: *Proceedings of the fifteenth annual ACM symposium on Theory of computing - STOC '83*. Not Known: ACM Press, 1983, pp. 291–297. ISBN: 978-0-89791-099-6. DOI: [10.1145/800061.808758](https://doi.org/10.1145/800061.808758). URL: <http://portal.acm.org/citation.cfm?id=291297> (visited on 06/07/2024).
- [Law69] F. W. Lawvere. “Adjointness in foundations”. In: *Dialectica* (1969). Publisher: JSTOR, pp. 281–296. URL: [https://www.jstor.org/stable/42969800?casa\\_token=GL45s\\_dCKqQAAAAA:SA](https://www.jstor.org/stable/42969800?casa_token=GL45s_dCKqQAAAAA:SA) (visited on 03/22/2024).
- [Law73] F. W. Lawvere. “Metric spaces, generalized logic, and closed categories”. In: *Rendiconti del seminario matematico e fisico di Milano* 43 (1973). Publisher: Springer, pp. 135–166.
- [Lei21] T. Leinster. *Entropy and diversity: the axiomatic approach*. Cambridge university press, 2021. ISBN: 978-1-108-96355-8. URL: <https://books.google.com/books?hl=en&lr=&id=pI4qEAAAQ> (visited on 06/03/2024).
- [Łuk20] J. Łukasiewicz. “O logice trójwartościowej”. In: *Ruch Filozoficzny* 5 (1920), pp. 170–171.
- [Mey91] P. Meyer-Nieberg. *Banach lattices*. Universitext. Springer-Verlag, 1991. ISBN: 3-540-54201-9. URL: <https://books.google.com/books?hl=en&lr=&id=2TPvCAAQBAJ&oi=fnd&pg=PA1> (visited on 03/26/2024).
- [MV70] D. S. Mitrinovic and P. M. Vasic. *Analytic inequalities*. Vol. 61. Springer, 1970.
- [NS16] F. Nielsen and K. Sun. “Guaranteed bounds on the Kullback-Leibler divergence of univariate mixtures using piecewise log-sum-exp inequalities”. en. In: *Entropy* 18.12 (Dec. 2016). arXiv:1606.05850 [cs, math, stat], p. 442. ISSN: 1099-4300. DOI: [10.3390/e18120442](https://doi.org/10.3390/e18120442). URL: <http://arxiv.org/abs/1606.05850> (visited on 04/16/2024).
- [OP99] P. W. O’Hearn and D. J. Pym. “The Logic of Bunched Implications”. en. In: *Bulletin of Symbolic Logic* 5.2 (June 1999), pp. 215–244. ISSN: 1079-8986, 1943-5894. DOI: [10.2307/421090](https://doi.org/10.2307/421090). URL: <https://www.cambridge.org/core/journals/bulletin-of-symbolic-logic> (visited on 03/27/2024).
- [PT21] P. Perrone and W. Tholen. *Kan extensions are partial colimits*. en. arXiv:2101.04531 [math]. Feb. 2021. URL: <http://arxiv.org/abs/2101.04531> (visited on 03/22/2024).
- [Sco67] D. Scott. “A Proof of the Independence of the Continuum Hypothesis”. In: *Mathematical systems theory* 1.2 (1967), pp. 89–111. URL: <https://www.karlin.mff.cuni.cz/~krajicek/s>
- [Shu13] M. Shulman. “Enriched Indexed Categories”. In: *Theory & Applications of Categories* 28.21 (2013). URL: <https://search.ebscohost.com/login.aspx?direct=true&profile=ehost&> (visited on 03/26/2024).

- [Shu22] M. Shulman. “Affine logic for constructive mathematics”. en. In: *Bulletin of Symbolic Logic* 28.3 (Sept. 2022). Publisher: Cambridge University Press, pp. 327–386. ISSN: 1079-8986, 1943-5894. DOI: [10.1017/bsl.2022.28](https://doi.org/10.1017/bsl.2022.28). URL: <https://www.cambridge.org/core/journal> (visited on 07/25/2023).
- [SKD+23] N. Slusarz et al. “Logic of Differentiable Logics: Towards a Uniform Semantics of DL”. In: *LPAR 2023: Proceedings of 24th International Conference on Logic for Programming, Artificial Intelligence and Reasoning, Manizales, Colombia, 4-9th June 2023*. Ed. by R. Piskac and A. Voronkov. Vol. 94. EPiC Series in Computing. EasyChair, 2023, pp. 473–493. DOI: [10.29007/C1NT](https://doi.org/10.29007/C1NT). URL: <https://doi.org/10.29007/c1nt>.
- [TMP24] S. Teuber, S. Mitsch, and A. Platzer. “Provably Safe Neural Network Controllers via Differential Dynamic Logic”. In: *CoRR* abs/2402.10998 (2024). DOI: [10.48550/ARXIV.2402.10998](https://doi.org/10.48550/ARXIV.2402.10998). arXiv: [2402.10998](https://arxiv.org/abs/2402.10998). URL: <https://doi.org/10.48550/arXiv.2402.10998>.
- [Zad65] L. A. Zadeh. “Fuzzy sets”. In: *Information and control* 8.3 (1965). Publisher: Elsevier, pp. 338–353. URL: <https://www.sciencedirect.com/science/article/pii/S001999586590241X> (visited on 05/16/2024).
- [Zad88] L. A. Zadeh. “Fuzzy logic”. In: *Computer* 21.4 (1988). Publisher: IEEE, pp. 83–93. URL: <https://ieeexplore.ieee.org/abstract/document/53/> (visited on 05/16/2024).