

MEETING TIMES OF MARKOV CHAINS VIA SINGULAR VALUE DECOMPOSITION

THOMAS VAN BELLE AND ANTON KLIMOVSKY

ABSTRACT. We suggest a non-asymptotic matrix perturbation-theoretic approach to get sharp bounds on the expected meeting time of random walks on large (possibly random) graphs. We provide a formula for the expected meeting time in terms of the singular value decomposition of the diagonally killed generator of a pair of independent random walks, which we view as a perturbation of the generator. Employing a rank-one approximation of the diagonally killed generator as the proof of concept, we work out sharp bounds on the expected meeting time of simple random walks on sufficiently dense Erdős–Rényi random graphs.

1. INTRODUCTION

Let $G_n = (V_n := [n], E_n)$, $n \in \mathbb{N}$ be a sequence of (possibly random) undirected graphs. Consider a pair of independent discrete time random walks on G_n : $(X_t^{(n)}, Y_t^{(n)})_{t \in \mathbb{Z}_+}$ started at vertex $i \in V_n$ resp. vertex $j \in V_n$. How long does it take them to meet? The *meeting time of two independent random walks* on G_n is a stopping time defined as

$$\tau_{\text{meet}}^{(n)}(i, j) = \inf\{t \in \mathbb{Z}_{\geq 0} : X_t^{(n)} = Y_t^{(n)}, X_0^{(n)} = i, Y_0^{(n)} = j\}, \quad i, j \in V_n \quad (1.1)$$

(with the natural convention that $\tau_{\text{meet}}^{(n)}(i, j) = \infty$ if the random walks never meet.) Of interest is also the *expected meeting time from stationarity*

$$t_{\text{meet}}^\pi = \mathbf{E}_{(i,j) \sim \pi \otimes \pi}[\tau_{\text{meet}}^{(n)}(i, j)] = \sum_{i,j \in V_n} \pi_i \pi_j \mathbf{E}[\tau_{\text{meet}}^{(n)}(i, j)], \quad (1.2)$$

where π is the invariant distribution of the random walk transition matrix P and \mathbf{E} is the expectation w.r.t. the law of the random walks on G_n . Both (1.1) and (1.2) are key quantities characterizing the behavior of random walks on graphs, see, e.g., [AF02].

Outline of the article. In Sections 1.1-1.3, we state our main results. In Section 1.4, we point to related literature. In Section 2, we give proof of the spectral formula for the expected meeting time, and we prove the rank- k approximation. In Section 3, we apply a result from matrix perturbation theory to investigate properties of the diagonally killed generator of a pair of independent Markov chains. In Section 4, we analyze the spectral formula applied to sufficiently dense Erdős–Rényi random graphs to deduce sharp non-asymptotic bounds on the expected meeting time from stationarity. In Section 5, we close with a discussion.

Date: June 10, 2024.

2010 Mathematics Subject Classification. 60K35.

Key words and phrases. meeting time, normalized Laplacian, generator, Markov chain, random walk, random graph, voter model, Erdős–Rényi, singular value decomposition, spectral theory, random matrix theory, non-asymptotic matrix perturbation theory.

This work was partly supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project-IDs 412848929; 443891315.

1.1. Meeting time formula via singular value decomposition. The first main result is a formula for the expected meeting times of two independent Markov chains both with transition matrix P from any pair of initial states in terms of the singular values decomposition (SVD) of what one might view as the “*diagonally killed*” generator of a pair of independent Markov chains

$$L_{\text{kill}} := I_{n^2} - (P \otimes P)E \in [-1, 1]^{n^2 \times n^2}, \quad (1.3)$$

$I_{n^2} \in \{0, 1\}^{n^2 \times n^2}$ is the unit matrix and $E \in \{0, 1\}^{n^2 \times n^2}$ is a matrix with

$$E_{ij} := \mathbb{1}\{i = j\} \mathbb{1}\{i \in \mathcal{D}\}, \quad i, j \in [n^2], \quad (1.4)$$

where $\mathcal{D} = \{1, n + 2, 2n + 3, \dots, n^2\}$ and $\mathbb{1}$ denotes the indicator function.

Remark 1.1 (Meaning of E). Let $C \in \mathbb{R}^{n^2 \times n^2}$ be a matrix. Then, the multiplication of C by E , e.g., from the right zeros out n columns of the matrix C , which exactly correspond to the diagonal $\{(i, i) \in V \times V : i \in V\}$ in the product state space. Thus, E encodes the meeting event. Note that $(P \otimes P)E$ is therefore a sub-stochastic matrix.

In order to better understand the Kronecker product structure of matrices, we define an “array flattening” mapping $f: [n]^2 \rightarrow [n^2]$ with

$$[n]^2 \ni (k, \ell) \xrightarrow{f} (k - 1)n + \ell \in [n^2]. \quad (1.5)$$

We will slightly abuse the notation and write as an index (k, ℓ) instead of $f(k, \ell)$. We choose this notation since it allows us to concisely write

$$(P \otimes P)_{(i,j),(k,\ell)} = P_{ik}P_{j\ell}. \quad (1.6)$$

Furthermore, the matrix E can be written as

$$(E)_{(i,j),(k,\ell)} := \mathbb{1}\{(i, j) = (k, \ell)\} \mathbb{1}\{k \neq \ell\}, \quad i, j, k, \ell \in [n]. \quad (1.7)$$

For a matrix $C \in \mathbb{R}^{n^2 \times n^2}$, we denote by $\text{vec}(C) \in \mathbb{R}^{n^2}$ the vector such that

$$\text{vec}(C)_{(i,j)} = C_{ij}. \quad (1.8)$$

In what follows we equip \mathbb{R}^n with the Euclidean (ℓ^2 -norm) $\|\cdot\|$ which induces the 2-2 operator norm on the space of the matrices, which we also denote by $\|\cdot\|$. Now we are in position to state the first main technical result.

Proposition 1.1 (Expected meeting times via SVD of the diagonally killed generator). *Let P the transition matrix of an irreducible Markov chain. Let $\tilde{\sigma}_1, \tilde{\sigma}_2, \dots, \tilde{\sigma}_{n^2}$ be the singular values (with multiplicities) of L_{kill} ordered non-increasingly:*

$$\tilde{\sigma}_1 \geq \tilde{\sigma}_2 \geq \dots \geq \tilde{\sigma}_{n^2} > 0. \quad (1.9)$$

Let \tilde{u}_i and \tilde{v}_i be the corresponding left and right singular vectors, $i \in [n^2]$. Then,

$$\text{vec}(\mathbf{E}[\tau_{\text{meet}}^{(n)}(i, j)]_{i,j \in [n]}) = E(I_{n^2} - (P \otimes P)E)^{-1} \mathbf{1}_{n^2}, \quad (1.10)$$

and in particular

$$t_{\text{meet}}^\pi = (\pi \otimes \pi)^t E(I_{n^2} - (P \otimes P)E)^{-1} \mathbf{1}_{n^2} \quad (1.11)$$

$$= -1 + \sum_{i=1}^{n^2} \frac{1}{\tilde{\sigma}_i} (\pi \otimes \pi)^t \tilde{v}_i \tilde{u}_i^t \mathbf{1}_{n^2}, \quad (1.12)$$

where $\mathbf{1}_{n^2}$ is the n^2 -dimensional vector with 1 at all coordinates, π is the stationary distribution of P .

The second main technical result is the following non-asymptotic rank- k approximation bound for the expected meeting time from stationarity.

Proposition 1.2 (Rank- k approximation of the meeting time). *Let $k \in [n^2 - 1]$, and define the rank- k approximation of the expected meeting time as*

$$\hat{t}_{\text{meet}}^{\pi, (k)} := -1 + \sum_{i=n^2-k}^{n^2} \frac{1}{\tilde{\sigma}_i} (\pi \otimes \pi)^t \tilde{v}_i \tilde{u}_i^t \mathbf{1}_{n^2}. \quad (1.13)$$

Then, the expected meeting time can be approximated by $\hat{t}_{\text{meet}}^{\pi, (k)}$:

$$\left| \hat{t}_{\text{meet}}^{\pi, (k)} - t_{\text{meet}}^{\pi} \right| \leq \frac{n \|\pi\|^2}{\tilde{\sigma}_{n^2-k-1}}. \quad (1.14)$$

Remark 1.2. In “mean-field-like” graphs, in which π is close to the uniform distribution, we get $\|\pi\|^2 = \Theta\left(\frac{1}{n}\right)$. So the l.h.s. of (1.14) reduces to $\Theta\left(\frac{1}{\tilde{\sigma}_{n^2-k-1}}\right)$.

1.2. A naive matrix perturbation theoretic approach to computations with diagonally killed generator. To apply Propositions 1.1 and 1.2, it can be useful to view L_{kill} as a perturbation of the generator of a pair of independent random walks

$$L := I_{n^2} - (P \otimes P) \quad (1.15)$$

to control the r.h.s. of (1.12). We denote the singular values (with multiplicities) of L by

$$\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_{n^2-1} > \sigma_{n^2} = 0. \quad (1.16)$$

The corresponding left and right singular vectors are denoted by u_i and v_i respectively. A naive application of the non-asymptotic second order perturbation for the least singular value [Ste84] yields

$$\tilde{\sigma}_{n^2}^2 = \sigma_{n^2}^2(L_{\text{kill}}) = \gamma_{11}^2 + O(\|(P \otimes P)(E - I)\|^3), \quad (1.17)$$

where $\gamma_{11} = u_{n^2}^t (P \otimes P)(E - I)v_{n^2}$. Note that the vectors u_{n^2} and v_{n^2} are given by

$$u_{n^2} = \frac{\pi \otimes \pi}{\|\pi \otimes \pi\|} \quad \text{and} \quad v_{n^2} = \frac{\mathbf{1}_{n^2}}{\|\mathbf{1}_{n^2}\|}. \quad (1.18)$$

The expansion (1.17) can be applied by noting that universally, i.e., for any graph,

$$\gamma_{11} = \frac{(\pi \otimes \pi)^t (P \otimes P)(E - I)\mathbf{1}_{n^2}}{\|\pi \otimes \pi\| \|\mathbf{1}_{n^2}\|} = \frac{(\pi \otimes \pi)^t (E - I)\mathbf{1}_{n^2}}{\|\pi \otimes \pi\| \|\mathbf{1}_{n^2}\|} = -\frac{1}{n}. \quad (1.19)$$

This naive application already fails even for the complete graph: a simple computation shows that the norm of the perturbation $\|(P \otimes P)(E - I)\|$ is of order $n^{-1/2}$. Even though (1.17) cannot be applied directly, it does show that the least singular value of L_{kill} is shifted by $\frac{1}{n}$ plus some graph-dependent terms. As we show below, see Section 3, it is possible to upgrade our understanding of the graph-dependent terms to make the method applicable to sufficiently dense Erdős–Rényi random graphs. This yields rather sharp non-asymptotic bounds on (1.12), see Theorem 1.3. In particular, the approximation

$$t_{\text{meet}}^{\pi} \approx \frac{1}{|\gamma_{11}|} = n \quad (1.20)$$

turns out to be valid with high probability.

Remark 1.3. A word of caution: It is known that the meeting time is *not* universally asymptotically equivalent to n (i.e., with proportionality constant 1), cf., e.g., the random regular graph [Che21] for another proportionality constant than 1 (but the same order $\Theta(n)$). Moreover, the meeting time must not be $\Theta(n)$ at all. It can be, e.g., $\Theta(1)$ like it is on star graphs, or $\Theta(n^2)$ like it is on the cycle graph, cf. [KMS23, Table 1].

1.3. Meeting times of random walks on sufficiently dense Erdős–Rényi random graphs.

In this section, we report on a rigorous proof-of-concept implementation of the idea for sufficiently dense Erdős–Rényi random graphs. Our result is non-asymptotic and identifies the exact proportionality constant. This constitutes our third main result.

Recall that the Erdős–Rényi random graph $\mathcal{G}(n, p)$ where $p \in (0, 1)$, $n \in \mathbb{N}$ is a random graph on $V := [n]$ such that the symmetric adjacency matrix $A = (a_{i,j})_{i,j \in [n]}$ has independent Bernoulli(p) distributed entries $(a_{i,j})_{i < j}$ and $a_{i,i} = 0$, $i \in V$. In what follows, we denote by \mathbb{P} the law of the Erdős–Rényi random graph.

Theorem 1.3 (Meeting time for sufficiently dense Erdős–Rényi graphs). *Consider two independent simple random walks on Erdős–Rényi random graph $\mathcal{G}(n, p_n)$, where the edge probability $p_n = cn^{\beta-1} \wedge 1$ with $c > 0$ is arbitrary but fixed. Let $\beta > \frac{1}{2}$. Then, for all $\epsilon > 0$, there exist $N \in \mathbb{N}$ and constants $\nu_1, \nu_2, \theta > 0$ such that for all $n \geq N$,*

$$\mathbb{P} \left(\left| \frac{1}{n} t_{\text{meet}}^\pi - 1 \right| > \epsilon \right) \leq 2n \exp \left(-\frac{\nu_1^2 d}{3} \right) + 2 \binom{n}{2} \exp \left(-\frac{\nu_2^2 d^2}{3n} \right) + e^{-\theta (\log n)^2}, \quad (1.21)$$

where $d := np_n$ is the mean degree. In particular, the r.h.s. (1.21) converges to zero, as $n \rightarrow \infty$.

See Section 5 for a discussion including possible extensions.

1.4. Related literature. There is a sizeable literature on the meeting times of random walks, see, e.g., [AF02; Ald13; KMS23]. Meeting times have applications to the long-run behavior of interacting particle systems such as voter model, see, e.g., [Ald13; Oli13; MT17; MPL17; Che21; Her+22; FO23], and some related algorithms on networks, see, e.g., [CFR10; KMS23].

Recently, there has been an increased interest in meeting times on random graphs, see, e.g., [Dur10; Che21; FO23; Ave+24; Ave+23]. For a sufficiently dense Erdős–Rényi random graph, i.e., $\beta > 0.5$, to the best of our knowledge, the sharpest bound on t_{meet} can be derived from [KMS23, Theorem B.1]. Specifically, $t_{\text{meet}} = \Theta(n)$, which gives no information about the proportionality constant. For $\beta \in (0, 0.5]$, the result gives $t_{\text{meet}} = O(nt_{\text{mix}})$ w.h.p.

Spectral characteristics (eigenvalues and -vectors) of the random walk transition matrix P are widely used to analyze the behavior of random walks on graphs (and more generally, Markov processes). For example, mixing times can be bounded using the spectral gap and the Perron-Frobenius eigenvector of P [LP17]. Hitting times also can be exactly expressed in terms of the eigenvalues and -vectors of P [Lov96]. The latter expression has been applied in [LT14; LT23] to analyze hitting times for random walks on Erdős–Rényi random graphs in the connected regime. Metastability also has spectral characterizations [BH15].

Singular values of Markov chain generators were (independently of this research) employed to define the notion of a spectral gap for nonreversible Markov chains [Cha23].

A formula resembling (1.10) appeared in [GPB18] (However, no link to singular values (1.12) and matrix perturbation theory was made therein.)

Matrix perturbation theory has a relatively long history. Asymptotic expansions were developed in physics literature, e.g., [Sch26]. It became a key topic in functional, numerical analyses and statistics, see, e.g., [Kat95; SS90]. Non-asymptotic matrix perturbation theory [DK70; Wed72] especially for the modern random matrix models remains a current topic of research; see, e.g., [BBP05; BN12; OVW18; Ver18; Che+21; Abb+20] with numerous contemporary applications, e.g., in statistics and machine learning.

2. MEETING TIMES: A MATRIX REPRESENTATION AND A SPECTRAL FORMULA

In this section, we derive the spectral formula (1.12). Furthermore, we prove the rank- k approximation (1.14) bound.

Proof of Proposition 1.1. First, we follow [GPB18] to write the pairwise meeting time in vector form. That is, consider two simple random walks X_n and Y_n both evolving according to P , and recall that $\tau_{\text{meet}}^{(n)}(i, j)$ denotes the meeting time of the two walks when started from states i and j respectively. The pairwise meeting time can be found recursively through

$$\tau_{\text{meet}}^{(n)}(i, j) = \begin{cases} 1 & \text{with prob. } \sum_{k \in [n]} P_{ik} P_{jk} \\ \tau_{\text{meet}}^{(n)}(k, \ell) + 1 & \text{with prob. } P_{ik} P_{j\ell}. \end{cases} \quad (2.1)$$

For $i \neq j$, let $M_{ij} = \mathbf{E} \left(\tau_{\text{meet}}^{(n)}(i, j) \right)$, then it can be seen that,

$$\begin{aligned} M_{ij} &= 1 + \sum_{k \in [n]} \sum_{\ell \in [n] \setminus \{k\}} P_{ik} P_{j\ell} M_{k\ell} \\ &= 1 + \sum_{k \in [n]} P_{ik} \sum_{\ell \in [n]} P_{j\ell} M_{k\ell} - \sum_{k \in [n]} P_{ik} P_{jk} M_{kk}. \end{aligned} \quad (2.2)$$

Written in matrix form, we obtain,

$$M = \mathbf{1}_n \mathbf{1}_n^t + P (M - M_d) P^t, \quad (2.3)$$

where M_d is a diagonal matrix with the same diagonal elements as M . Rewriting in vector form, yields

$$\begin{aligned} \text{vec}(M) &= \mathbf{1}_{n^2} + (P \otimes P) (\text{vec}(M) - \text{vec}(M_d)) \\ &= \mathbf{1}_{n^2} + (P \otimes P) (I_{n^2} - \text{vec}(I_n)) \text{vec}(M) \\ &= \mathbf{1}_{n^2} + (P \otimes P) E \text{vec}(M). \end{aligned} \quad (2.4)$$

By rearranging terms it can be seen that,

$$\text{vec}(M) = (I_{n^2} - (P \otimes P) E)^{-1} \mathbf{1}_{n^2}. \quad (2.5)$$

Note that $\mathbf{E}[\tau_{\text{meet}}^{(n)}(i, i)] = 0$ for all $i \in [n]$, so that

$$\mathbf{E}[\tau_{\text{meet}}^{(n)}(i, j)] = M_{ij} \mathbb{1}\{i \neq j\} = (E \text{vec}(M))_{(i,j)} \quad (2.6)$$

hence Equation (1.10) follows. We observe that

$$t_{\text{meet}}^\pi = \sum_{i=1}^n \sum_{\substack{j=1 \\ j \neq i}}^n \pi_i \pi_j M_{ij} = (\pi \otimes \pi)^t E \text{vec}(M), \quad (2.7)$$

so that (1.11) easily follows. The expected meeting time can also be written as

$$t_{\text{meet}}^{\pi} = \sum_{i=1}^n \sum_{\substack{j=1 \\ j \neq i}}^n \pi_i \pi_j M_{ij} = \sum_{i=1}^n \sum_{j=1}^n \pi_i \pi_j M_{ij} - \sum_{i=1}^n \pi_i^2 M_{ii}. \quad (2.8)$$

First we compute the second term. Using (2.3) it can be seen that

$$\pi^t M \pi = \pi^t \mathbf{1}_n \mathbf{1}_n^t \pi + \pi^t P (M - M_d) P^t \pi = 1 + \pi^t (M - M_d) \pi. \quad (2.9)$$

So that,

$$\sum_{i=1}^n \pi_i^2 M_{ii} = \pi^t M_d \pi = 1. \quad (2.10)$$

The first term of (2.8) can be written as

$$\sum_{i=1}^n \sum_{j=1}^n \pi_i \pi_j M_{ij} = (\pi \otimes \pi)^t (I_{n^2} - (P \otimes P) E)^{-1} \mathbf{1}_{n^2}. \quad (2.11)$$

To find the inverse in the equation above, we use the singular value decomposition of L_{kill} , i.e., $I - (P \otimes P)E = \tilde{U} \tilde{\Sigma} \tilde{V}^t$. Then,

$$(I - (P \otimes P)E)^{-1} = \tilde{V} \tilde{\Sigma}^{-1} \tilde{U}^t = \sum_{i=1}^{n^2} \frac{1}{\tilde{\sigma}_i} \tilde{v}_i \tilde{u}_i^t. \quad (2.12)$$

Equation (1.12) follows by plugging these results into (2.8). \square

Proof of Proposition 1.2. From Proposition 1.1, it follows that

$$t_{\text{meet}}^{\pi} = -1 + \sum_{i=1}^{n^2} \frac{1}{\tilde{\sigma}_i} (\pi \otimes \pi)^t \tilde{v}_i \tilde{u}_i^t \mathbf{1}_{n^2}. \quad (2.13)$$

The result then follows from the observation that

$$\begin{aligned} |\hat{t}_{\text{meet}}^{\pi} - t_{\text{meet}}^{\pi}| &= \left| (\pi \otimes \pi)^t \left(\sum_{i=n^2-k}^{n^2} \frac{\tilde{v}_i \tilde{u}_i^t}{\tilde{\sigma}_i} - \sum_{i=1}^{n^2} \frac{\tilde{v}_i \tilde{u}_i^t}{\tilde{\sigma}_i} \right) \mathbf{1}_{n^2} \right| \\ &\leq \|\pi \otimes \pi\| \left\| \sum_{i=1}^{n^2-k-1} \frac{\tilde{v}_i \tilde{u}_i^t}{\tilde{\sigma}_i} \right\| \|\mathbf{1}_{n^2}\| \\ &= \frac{1}{\tilde{\sigma}_{n^2-k-1}} n \|\pi\|^2, \end{aligned} \quad (2.14)$$

where the last equality follows from the Eckhart-Young(-Mirsky) Theorem [EY36] and the observation that

$$\|\pi \otimes \pi\|^2 = \sum_{i=1}^{n^2} (\pi \otimes \pi)_i^2 = \sum_{(i,j)=(1,1)}^{(n,n)} (\pi \otimes \pi)_{(i,j)}^2 = \sum_{i=1}^n \sum_{j=1}^n \pi_i^2 \pi_j^2 = \|\pi\|^4. \quad (2.15)$$

\square

3. MATRIX PERTURBATION THEORY

In order to usefully apply Proposition 1.2, we need to obtain a sufficiently sharp estimates for the $k + 1$ least singular values of the diagonally killed generator L_{kill} . We use a result from non-asymptotic matrix perturbation theory [Ste73, Theorems 4.11 and 4.12] to obtain useful bounds on the least singular value. Recall the definitions of v_{n^2} and u_{n^2} in (1.18) and σ_{n^2} in (1.16), and denote the singular value decomposition of the unperturbed matrix L in block form by

$$L = I_{n^2} - (P \otimes P) =: [u_{n^2} \quad U_2]^t \begin{bmatrix} \sigma_{n^2} & 0 \\ 0 & \Sigma_2 \end{bmatrix} [v_{n^2} \quad V_2], \quad (3.1)$$

where U_2 has the columns $(u_i)_{i < n^2}$ and V_2 has the columns $(v_i)_{i < n^2}$, and $\Sigma_2 := \text{diag}(\sigma_i)_{i < n^2}$ is a diagonal matrix with the remaining singular values. We apply Theorem A.1 with

$$\begin{aligned} B &= L^t L \\ \Delta &= L_{\text{kill}}^t L_{\text{kill}} - L^t L \\ X &= [v_{n^2} \quad V_2]. \end{aligned} \quad (3.2)$$

Note that in this application the block matrices defined in (A.1) are given by $B_{11} = 0$, $B_{22} = \Sigma_2^2$, $B_{12} = 0$. We define the parameters γ_{11} , g_{12} , g_{21} , and G_{22} through

$$[u_{n^2} \quad U_2]^t (P \otimes P)(E - I) [v_{n^2} \quad V_2] = \begin{bmatrix} \gamma_{11} & g_{12}^t \\ g_{21} & G_{22} \end{bmatrix}. \quad (3.3)$$

By observing that

$$[u_{n^2} \quad U_2]^t (I - (P \otimes P)E) [v_{n^2} \quad V_2] = \begin{bmatrix} \gamma_{11} & g_{12}^t \\ g_{21} & \Sigma_2 + G_{22} \end{bmatrix}, \quad (3.4)$$

and that $B_{11} = \sigma_{n^2} = 0$, it can be seen that

$$X^t (B + \Delta) X = \begin{bmatrix} \gamma_{11}^2 + g_{21}^t g_{21} & g_{21}^t \Sigma_2 + \gamma_{11} g_{12}^t + g_{21}^t G_{22} \\ \Sigma_2 g_{21} + \gamma_{11} g_{12} + G_{22}^t g_{21} & (\Sigma_2 + G_{22})^t (\Sigma_2 + G_{22}) + g_{12} g_{12}^t \end{bmatrix}. \quad (3.5)$$

Therefore, the blocks in (A.2) are given by,

$$\begin{aligned} \Delta_{11} &= \gamma_{11}^2 + g_{21}^t g_{21}, \\ \Delta_{12} &= g_{21}^t \Sigma_2 + \gamma_{11} g_{12}^t + g_{21}^t G_{22}, \\ \Delta_{21} &= \Sigma_2 g_{21} + \gamma_{11} g_{12} + G_{22}^t g_{21}, \\ \Delta_{22} &= G_{22}^t \Sigma_2 + \Sigma_2 G_{22} + G_{22}^t G_{22} + g_{12} g_{12}^t. \end{aligned} \quad (3.6)$$

Let $B \in \mathbb{R}^{\ell \times \ell}$ and $C \in \mathbb{R}^{m \times m}$. Define the operator $T_{B,C}: \mathbb{R}^{m \times \ell} \rightarrow \mathbb{R}^{m \times \ell}$ through,

$$T_{B,C} P = PB - CP. \quad (3.7)$$

The separation of B and C is the number $\text{sep}(B, C)$ defined by

$$\text{sep}(B, C) := \begin{cases} \|T_{B,C}^{-1}\|^{-1}, & \text{if } 0 \notin \lambda(T), \\ 0 & \text{otherwise.} \end{cases} \quad (3.8)$$

Let

$$\delta = \text{sep}(B_{11}, B_{22}) - \|\Delta_{11}\| - \|\Delta_{22}\|. \quad (3.9)$$

Note that since $B_{11} = 0$, and $B_{22} = \Sigma_2^2$, we get

$$\text{sep}(B_{11}, B_{22}) = \|\Sigma_2^{-2}\|^{-1}. \quad (3.10)$$

If

$$\frac{4}{\delta^2} \|\Delta_{21}\| (\|B_{12}\| + \|\Delta_{12}\|) \leq 1, \quad (3.11)$$

there exists a matrix Q satisfying $\|Q\| \leq \frac{2}{\delta} \|\Delta_{12}\|$ such that

$$\tilde{\sigma}_{n^2} = \lambda_{\min}(B + \Delta) = \gamma_{11}^2 + g_{21}^t g_{21} + (g_{21}^t \Sigma_2 + \gamma_{11} g_{12}^t + g_{21}^t G_{22})Q. \quad (3.12)$$

Furthermore, the columns of $X'_1 = (1 + \|Q\|^2)^{-\frac{1}{2}}(v_{n^2} + V_2 Q)$ span an invariant subspace of $B + \Delta$. Note that (3.12), gives us an exact expression for the least singular value of the diagonally killed generator L_{kill} . As we have seen in (1.19), $\gamma_{11} = -\frac{1}{n}$ for any transition matrix P . The remaining terms depend on the structure of the graph and therefore need to be bounded on a case by case basis. In Section 4, we will bound the terms g_{12} , g_{21} , G_{22} , Σ_2 , and Q by their operator norms, for the specific case where the graph is an Erdős-Rényi random graph. In order to obtain useful upper- and lower-bounds on the least singular value, we define

$$\begin{aligned} \tilde{\gamma}_{11}^2 &:= \|g_{21}\|^2 + \gamma_{11}^2 = v_{n^2}^t (E - I)^t (P \otimes P)^t U_2 U_2^t (P \otimes P) (E - I) v_{n^2} + \gamma_{11}^2 \\ &= v_{n^2}^t (E - I)^t (P \otimes P)^t (I - u_{n^2} u_{n^2}^t) (P \otimes P) (E - I) v_{n^2} + \gamma_{11}^2 \\ &= \frac{1}{n^2} \mathbf{1}_{n^2} (E - I)^t (P \otimes P)^t (P \otimes P) (E - I) \mathbf{1}_{n^2}, \end{aligned} \quad (3.13)$$

to see that

$$\begin{aligned} \tilde{\sigma}_{(n,n)}^2 &\geq \tilde{\gamma}_{11}^2 - 2 \cdot \frac{\|\Delta_{12}\|^2}{\delta}, \\ \tilde{\sigma}_{(n,n)}^2 &\leq \tilde{\gamma}_{11}^2 + 2 \cdot \frac{\|\Delta_{12}\|^2}{\delta}. \end{aligned} \quad (3.14)$$

In Section 4, we provide an example of graphs in which the term $2 \cdot \frac{\|\Delta_{12}\|^2}{\delta}$ is of lower order than $\tilde{\gamma}_{11}^2$. If this latter condition is satisfied, we can estimate the least singular value and use Propositions 1.1 and 1.2 to conclude that w.h.p. the approximation in (1.20) is valid.

4. APPLICATION TO DENSE RANDOM GRAPHS

In this section, we will provide an application of Proposition 1.1 to dense Erdős-Rényi graphs. This will result in a proof of Theorem 1.3.

We start by proving some concentration results. It will be useful to define the following quantities,

$$\epsilon_i := \frac{\deg i}{d} - 1 \quad R_1 := \max_{i \in V} |\epsilon_i|, \quad (4.1)$$

where $d := cn^\beta$ is the average degree for each vertex. Note these definitions imply

$$\min_{i \in V} \deg i \geq d(1 - R_1) \quad \max_{i \in V} \deg i \leq d(1 + R_1). \quad (4.2)$$

Furthermore for $k \neq \ell$ define

$$\epsilon_{(k,\ell)} := \frac{n}{d^2} \left(\sum_i \mathbb{1}\{k \sim i\} \mathbb{1}\{\ell \sim i\} \right) - 1 \quad R_2 := \max_{(k,\ell) \in V \times V, k \neq \ell} |\epsilon_{(k,\ell)}|. \quad (4.3)$$

Finally, for $\nu_1, \nu_2 > 0$, we define the events

$$F_{\nu_1, \nu_2} := \{R_1 \leq \nu_1\} \cap \{R_2 \leq \nu_2\} \quad \text{and} \quad F_{\nu_1} := \{R_1 \leq \nu_1\}. \quad (4.4)$$

Lemma 4.1. For any $\nu_1, \nu_2 > 0$,

$$\mathbb{P}(F_{\nu_1}^c) \leq 2n \exp\left(-\frac{\nu_1^2 np}{3}\right), \quad (4.5)$$

and

$$\mathbb{P}(F_{\nu_1, \nu_2}^c) \leq 2n \exp\left(-\frac{\nu_1^2 np}{3}\right) + 2 \binom{n}{2} \exp\left(-\frac{\nu_2^2 np^2}{3}\right). \quad (4.6)$$

Remark 4.1. Note that the upper-bound in (4.5) goes to zero whenever np grows faster than $\log n$, which holds for any $\beta > 0$. The upper-bound in (4.6) however, goes to zero whenever np^2 grows faster than $\log \binom{n}{2}$. Since $np^2 = \frac{1}{c^2} n^{1-2\beta}$, this is the case for $\beta > \frac{1}{2}$.

Proof. We use the union bound to get that for any $\nu_1 > 0$,

$$\mathbb{P}(R_1 > \nu_1) = \mathbb{P}\left(\exists i \in V : \left|\frac{\deg_i}{np} - 1\right| > \nu_1\right) \leq n \mathbb{P}\left(\left|\frac{\deg_1}{np} - 1\right| > \nu_1\right). \quad (4.7)$$

The probability on the right-hand side can be bounded using a Chernoff bound, i.e.,

$$\mathbb{P}\left(\left|\frac{\deg_1}{np} - 1\right| > \nu_1\right) = \mathbb{P}(|\deg_1 - np| > np\nu_1) \leq 2 \exp\left(-\frac{\nu_1^2 np}{3}\right). \quad (4.8)$$

Equation (4.5) follows. Similarly, note that

$$\begin{aligned} \mathbb{P}(R_2 > \nu_2) &= \mathbb{P}\left(\exists k \neq \ell : \left|\frac{1}{np^2} \sum_i \mathbb{1}\{k \sim i\} \mathbb{1}\{\ell \sim i\} - 1\right| > \nu_2\right) \\ &\leq \binom{n}{2} \mathbb{P}\left(\left|\frac{1}{np^2} \sum_i \mathbb{1}\{1 \sim i\} \mathbb{1}\{2 \sim i\} - 1\right| > \nu_2\right) \\ &\leq 2 \binom{n}{2} \exp\left(-\frac{\nu_2^2 np^2}{3}\right). \end{aligned} \quad (4.9)$$

The standard bound

$$\mathbb{P}(F_{\nu_1, \nu_2}^c) \leq \mathbb{P}(R_1 > \nu_1) + \mathbb{P}(R_2 > \nu_2) \quad (4.10)$$

completes the proof of (4.6). \square

Another graph-dependent event we frequently use is the convergence of the second least singular value. Denote the adjacency matrix of $G = G_n$ by $A = A_n$. We denote by F the event that the second least singular value of $\frac{A}{\sqrt{d}}$ is at most 8:

$$F := \left\{ \sigma_2\left(\frac{A}{\sqrt{d}}\right) \leq 8 \right\}. \quad (4.11)$$

We need the following adaptation from [Erd+12; Erd+13] as follows,

Proposition 4.2. There exist a constant $\theta > 0$ and $N \in \mathbb{R}_+$, such that for all $n \geq N$,

$$\mathbb{P}(F^c) \leq \exp(-\theta (\log n)^2) \quad (4.12)$$

Remark 4.2. In fact, [Erd+12] shows that for $d \gg (\log n)^{2C\xi}$,

$$\sigma_2\left(\frac{A}{\sqrt{d}}\right) \rightarrow 2 \quad (4.13)$$

with high probability. For our purposes, it suffices that $\sigma_2 \left(\frac{A}{\sqrt{d}} \right)$ is bounded. The constant 8 is arbitrary and chosen for (subjective) aesthetic reasons. It could be replaced by any constant bigger than 2.

Proof. By [Erd+12, Equation (3.19)], there exist constants $\theta, C > 0$ such that

$$\sigma_2 \left(\frac{A}{\sqrt{d}} \right) \leq \sigma_2 \left(\frac{A}{\sqrt{np(1-p)}} \right) \leq 2 + (\log n)^{C\xi} \left(\frac{1}{d} + \frac{1}{n^{2/3}} \right) \quad (4.14)$$

holds with probability at least $1 - \exp \left(-\theta (\log n)^\xi \right)$, where

$$1 + a_0 \leq \xi \leq A_0 \log \log n \quad (4.15)$$

for constants $a_0 > 0$ and $A_0 \geq 10$. We pick $\xi = 2$ to see that

$$\sigma_2 \left(\frac{A}{\sqrt{d}} \right) \leq 2 + (\log n)^{2C} \left(\frac{1}{d} + \frac{1}{n^{2/3}} \right) \quad (4.16)$$

holds with probability at least $1 - \exp \left(-\theta (\log n)^2 \right)$. Let n be such that,

$$\frac{(\log n)^{2C}}{\sqrt{n}} \leq 3c \quad \text{and} \quad \frac{(\log n)^{2C}}{n^{2/3}} \leq 3. \quad (4.17)$$

Note that there exists $N \in \mathbb{R}_+$ such that (4.17) is satisfied for all $n \geq N$. Therefore, for all $n \geq N$,

$$\sigma_2 \left(\frac{A}{\sqrt{d}} \right) \leq 8 \quad (4.18)$$

holds with probability at least $1 - \exp \left(-\theta (\log n)^2 \right)$. \square

Another key quantity is the norm of the perturbation, i.e., $\| (P \otimes P)(E - I) \|$. It can be usefully bounded in terms of the previously defined quantities R_1 and R_2 as follows.

Lemma 4.3. *Let $\beta \in (0, 1)$ then,*

$$\| (P \otimes P)(E - I) \|^2 \leq \frac{(1 + R_1)^2}{d^2(1 - R_1)^4} + \frac{(1 + R_2)^2}{n(1 - R_1)^4} - \frac{(1 + R_2)^2}{n^2(1 - R_1)^4}. \quad (4.19)$$

For $\beta > \frac{1}{2}$,

$$\| (P \otimes P)(E - I) \|^2 \geq \frac{(1 - R_1)^2}{d^2(1 + R_1)^4} + \frac{(1 - R_2)^2}{n(1 + R_1)^4} - \frac{(1 - R_2)^2}{n^2(1 + R_1)^4}. \quad (4.20)$$

Remark 4.3. This lemma shows that for $\beta > \frac{1}{2}$ the norm of the perturbation behaves like

$$\| (P \otimes P)(E - I) \|^2 \sim \frac{1}{n}. \quad (4.21)$$

For $\beta < \frac{1}{2}$, R_2 grows. Therefore, we expect that the upper-bound in (4.19) is not sharp.

Proof. Define the matrix $S \in \mathbb{R}^{n \times n}$

$$S_{i,k} = \left(\sum_q A_{qi} A_{qk} \right)^2 = \begin{cases} d^2(1 + \epsilon_i)^2 & \text{if } i = k \\ \frac{d^4}{n^2}(1 + \epsilon_{(i,k)})^2 & \text{if } i \neq k, \end{cases} \quad (4.22)$$

so that it contains the non-zero elements of $(E - I)^t (A \otimes A)^t (A \otimes A) (E - I)$. Then

$$\| (P \otimes P)(E - I) \|^2 \leq \frac{\|S\|}{d^4(1 - R_1)^4} \quad \text{and} \quad \| (P \otimes P)(E - I) \|^2 \geq \frac{\|S\|}{d^4(1 + R_1)^4}. \quad (4.23)$$

Note that by the Gershgorin Circle Theorem,

$$\|S\| \leq d^2 (1 + R_1)^2 + \frac{d^4}{n} (1 + R_2)^2 - \frac{d^4}{n^2} (1 + R_2)^2. \quad (4.24)$$

Equation (4.19) follows from (4.23) and (4.24). To prove the second statement (4.20), we note that,

$$\|(P \otimes P)(E - I)\|^2 \geq \frac{\|S\|}{d^4(1 + R_1)^4}. \quad (4.25)$$

Furthermore,

$$\|S\| = \max_{\|x\|=1} x^t S x \geq \frac{1}{n} \sum_{(i,k)} S_{i,k} \geq d^2 (1 - R_1)^2 + \left(1 - \frac{1}{n}\right) \frac{d^4}{n} (1 - R_2)^2, \quad (4.26)$$

where the first inequality follows from taking x to be the uniform vector. By combining (4.25) and (4.26), we obtain (4.20). \square

Next we show that some key quantities in the application of Theorem A.1 are bounded. These bounds will provide the backbone of the proof of Theorem 1.3.

Lemma 4.4. *Let $\beta > \frac{1}{2}$. For every $\varepsilon_1 > 0$, there exists $N_1 \in \mathbb{R}_+$ and constants $\nu_1, \nu_2 > 0$ such that, for all $n \geq N_1$,*

$$\mathbb{P}(\|\Sigma_2\| \leq 2 + \varepsilon_1 \mid F_{\nu_1}) = 1, \quad (4.27)$$

$$\mathbb{P}(n\|g_{12}\|^2 \leq 1 + \varepsilon_1 \mid F_{\nu_1}) = 1, \quad (4.28)$$

$$\mathbb{P}(n\|\pi\|^2 \leq 1 + \varepsilon_1 \mid F_{\nu_1}) = 1, \quad (4.29)$$

$$\mathbb{P}(\sqrt{n}\|G_{22}\| \leq 1 + \varepsilon_1 \mid F_{\nu_1, \nu_2}) = 1, \quad (4.30)$$

$$\mathbb{P}(n^2\|g_{21}\|^2 \leq (1 + \varepsilon_1)^2 - 1 \mid F_{\nu_1, \nu_2}) = 1, \quad (4.31)$$

$$\mathbb{P}((1 - \varepsilon_1)^2 \leq \|\Sigma_2^{-2}\|^{-1} \leq (1 + \varepsilon_1)^2 \mid F_{\nu_1} \cap F) = 1. \quad (4.32)$$

Proof. Let $\varepsilon_1 > 0$, and define the ν_1 and ν_2 by

$$\nu_1 = \nu_2 = \frac{(\varepsilon_1 + 1)^{\frac{1}{4}} - 1}{(\varepsilon_1 + 1)^{\frac{1}{4}} + 1}. \quad (4.33)$$

and recall that conditioning on F_{ν_1, ν_2} and F gives the conditions that $R_1 \leq \nu_1$, $R_2 \leq \nu_2$, and $\sigma_2(A) \leq 8\sqrt{d}$. This choice for ν_1 and ν_2 is not optimal, but it ensures that ν_1 and ν_2 are positive and bounded by 1. Furthermore,

$$\max \left\{ \frac{(1 + \nu_1)^2}{(1 - \nu_1)^4}, \frac{(1 + \nu_2)^2}{(1 - \nu_1)^4}, \frac{(1 + \nu_1)^4}{(1 - \nu_1)^4} \right\} \leq 1 + \varepsilon_1. \quad (4.34)$$

Finally let

$$N_1 = \max \left\{ c^{\frac{1}{1-\beta}}, \left(\frac{1}{c^2 \varepsilon_1} \right)^{\frac{1}{2\beta-1}}, \left(\frac{64(1 + \varepsilon_1)^2}{c \varepsilon_1^2} \right)^{\frac{1}{\beta}} \right\}, \quad (4.35)$$

and assume that $n \geq N_1$. Note that the condition $n \geq c^{\frac{1}{1-\beta}}$ ensures that

$$p = cn^{\beta-1} \leq 1. \quad (4.36)$$

We start by proving (4.27). The norm of the matrix Σ_2 can be bounded using a standard inequality for singular values,

$$\|\Sigma_2\| = \sigma_1(L) \leq \sigma_1(I) + \sigma_1(P \otimes P) \leq 1 + \frac{\sigma_1(A)^2}{d^2(1 - R_1)^2}. \quad (4.37)$$

Note that the maximal singular value of the adjacency matrix A is at most the maximal degree of the graph, so that by (4.2),

$$\|\Sigma_2\| \leq 1 + \left(\frac{1+R_1}{1-R_1}\right)^2 < 1 + \left(\frac{1+\nu_1}{1-\nu_1}\right)^2 \leq 2 + \varepsilon_1. \quad (4.38)$$

Next we prove (4.28). The norm of g_{12} can be bounded by observing that

$$\begin{aligned} n\|g_{12}\|^2 &= nu_1^t(P \otimes P)(E - I)V_2V_2^t(E - I)^t(P \otimes P)^tu_1 \\ &= nu_1^t(P \otimes P)(E - I)(I - v_1v_1^t)(E - I)^t(P \otimes P)^tu_1 \\ &= n \frac{(\pi \otimes \pi)^t(I - E)(\pi \otimes \pi)}{\|\pi \otimes \pi\|^2} - \gamma_{11}^2 \\ &= n \frac{\sum_i (1 + \epsilon_i)^4}{\sum_{(i,j)} (1 + \epsilon_i)^2(1 + \epsilon_j)^2} - \frac{1}{n} \\ &\leq \left(\frac{1+R_1}{1-R_1}\right)^4 < \left(\frac{1+\nu_1}{1-\nu_1}\right)^4 \leq 1 + \varepsilon_1. \end{aligned} \quad (4.39)$$

To prove (4.29), note that

$$n\|\pi\|^2 = n \sum_i \pi_i^2 \leq \frac{(1+R_1)^2}{(1-R_1)^2} \leq 1 + \varepsilon_1. \quad (4.40)$$

To prove (4.30), we observe that

$$\|G_{22}\| \leq \|U_2\| \|(P \otimes P)(E - I)\| \|V_2\| \leq \|(P \otimes P)(E - I)\|. \quad (4.41)$$

Using Lemma 4.3, it can be seen that therefore,

$$\begin{aligned} n\|G_{22}\|^2 &\leq \frac{n(1+R_1)^2}{d^2(1-R_1)^4} + \frac{(1+R_2)^2}{(1-R_1)^4} \\ &< \frac{n(1+\nu_1)^2}{d^2(1-\nu_1)^4} + \frac{(1+\nu_2)^2}{(1-\nu_1)^4} \\ &\leq \left(\frac{n}{d^2} + 1\right)(1 + \varepsilon_1). \end{aligned} \quad (4.42)$$

For $n \geq N_1$, we have in particular that $n \geq \left(\frac{1}{c^2\varepsilon_1}\right)^{\frac{1}{2\beta-1}}$, so that

$$\frac{n}{d^2} = \frac{1}{c^2} n^{1-2\beta} \leq \varepsilon_1. \quad (4.43)$$

Hence we can conclude that

$$\sqrt{n}\|G_{22}\| \leq 1 + \varepsilon_1. \quad (4.44)$$

Next we prove (4.31). Recall that

$$n^2\|g_{21}\|^2 = n^2\tilde{\gamma}_{11}^2 - 1. \quad (4.45)$$

We can upper-bound the expression above by observing that

$$\begin{aligned}
n^2 \tilde{\gamma}_{11}^2 &\leq \frac{1}{(1-R_1)^4} \frac{1}{d^4} \sum_{(k,\ell)} \left(\sum_i \mathbb{1}\{i \sim k\} \mathbb{1}\{i \sim \ell\} \right)^2 \\
&\leq \frac{1}{(1-R_1)^4} \left(2 \binom{n}{2} \frac{1}{n^2} (1+R_2)^2 + \frac{1}{np^2} (1+R_1)^2 \right) \\
&< \frac{(1+\nu_2)^2}{(1-\nu_1)^4} + \frac{1}{np^2} \frac{(1+\nu_1)^2}{(1-\nu_1)^4} \leq (1+\varepsilon_1)^2,
\end{aligned} \tag{4.46}$$

where the last inequality follows again from (4.43). Finally we prove (4.32). It holds that

$$\|\Sigma_2^{-2}\|^{-1} = \sigma_{\min}(\Sigma_2)^2 \geq (1 - \sigma_2(P \otimes P))^2, \tag{4.47}$$

and

$$\|\Sigma_2^{-2}\|^{-1} \leq (1 + \sigma_2(P \otimes P))^2. \tag{4.48}$$

Note that the second least singular value of $P \otimes P$ can be bounded using

$$\sigma_2(P \otimes P) \leq \frac{1}{d^2(1-R_1)^2} \sigma_2(A \otimes A) \leq \frac{(1+R_1)}{\sqrt{d}} \sigma_2\left(\frac{A}{\sqrt{d}}\right) \leq \frac{8}{\sqrt{d}}(1+\varepsilon_1). \tag{4.49}$$

For $n \geq N_1$, we have in particular that $n \geq \left(\frac{64(1+\varepsilon_1)^2}{c\varepsilon^2}\right)^{\frac{1}{\beta}}$, so that

$$\frac{8}{\sqrt{d}}(1+\varepsilon_1) \leq \frac{8}{\sqrt{c\left(\frac{64(1+\varepsilon_1)^2}{c\varepsilon^2}\right)}} \leq \varepsilon_1. \tag{4.50}$$

Therefore the result follows. \square

We use Lemma 4.4 to bound the terms of the rank-one approximation $\hat{t}_{\text{meet}}^{\pi, (1)}$ and the error approximation in Proposition 1.2.

Lemma 4.5. *Let $\beta > \frac{1}{2}$. For every $\varepsilon_2 > 0$, there exists $N_2 \in \mathbb{N}$ and constants $\nu_1, \nu_2 > 0$ such that, for all $n \geq N_2$,*

$$\mathbb{P}\left(\left|\frac{1}{n\tilde{\sigma}_{n^2}} - 1\right| \leq \varepsilon_2 \mid F_{\nu_1, \nu_2} \cap F\right) = 1 \tag{4.51}$$

$$\mathbb{P}\left(\left|(\pi \otimes \pi)^t \tilde{v}_{n^2} \tilde{u}_{n^2}^t \mathbf{1}_{n^2} - 1\right| \leq \varepsilon_2 \mid F_{\nu_1, \nu_2} \cap F\right) = 1 \tag{4.52}$$

$$\mathbb{P}\left(\frac{n\|\pi\|^2}{n\tilde{\sigma}_{n^2-1}} \leq \varepsilon_2 \mid F_{\nu_1} \cap F\right) = 1 \tag{4.53}$$

Proof. Let $\varepsilon_1 \in (0, \frac{1}{20})$, ν_1, ν_2 be defined as in (4.33), recall that conditioning on F_{ν_1, ν_2} and F gives the conditions that $R_1 \leq \nu_1$, $R_2 \leq \nu_2$, and $\sigma_2(A) \leq 8\sqrt{d}$. These conditions allow us to use the estimates in Lemma 4.4. First we note that

$$n\tilde{\gamma}_{11}^2 = 1 + n^2\|g_{21}\|^2 \geq 1. \tag{4.54}$$

Therefore, the perturbed least singular value can be upper-bounded by

$$\frac{1}{n\tilde{\sigma}_{n^2}} \leq \left(n^2\tilde{\gamma}_{11}^2 - 2n^2\frac{\|\Delta_{21}\|^2}{\delta}\right)^{-\frac{1}{2}} \leq \left(1 - 2n^2\frac{\|\Delta_{21}\|^2}{\delta}\right)^{-\frac{1}{2}}, \tag{4.55}$$

and lower-bounded by

$$\frac{1}{n\tilde{\sigma}_{n^2}} \geq \left(1 + n^2\|g_{21}\|^2 + 2n^2\frac{\|\Delta_{21}\|^2}{\delta}\right)^{-\frac{1}{2}}. \quad (4.56)$$

We will use the estimates from Lemma 4.4 to conclude that for every $\varepsilon_1 \in (0, \frac{1}{20})$, there exist constants $\nu_1, \nu_2 > 0$ and $N_1 \in \mathbb{R}_+$ such that for $n \geq N_1$, conditioned on F_{ν_1, ν_2} we have,

$$\begin{aligned} n\|\Delta_{21}\| &\leq n\|g_{21}\|\|\Sigma_2\| + \|g_{12}\| + n\|g_{21}\|\|G_{22}\| \\ &\leq 5\sqrt{\varepsilon_1}. \end{aligned} \quad (4.57)$$

and

$$n^2\|g_{21}\|^2 \leq \varepsilon_1(2 + \varepsilon_1). \quad (4.58)$$

Furthermore, conditioned on F_{ν_1, ν_2} and F we have,

$$\begin{aligned} \delta &= \text{sep}(B_{11}, B_{22}) - \|\Delta_{11}\| - \|\Delta_{22}\| \\ &= \|\Sigma_2^{-2}\|^{-1} - \tilde{\gamma}_{11}^2 - \|G_{22}^t\Sigma_2 + \Sigma_2G_{22} + G_{22}^tG_{22} + g_{12}g_{12}^t\| \\ &\geq \|\Sigma_2^{-2}\|^{-1} - \tilde{\gamma}_{11}^2 - 2\|G_{22}\|\|\Sigma_2\| - \|G_{22}\|^2 - \|g_{12}\|^2 \\ &\geq 1 - 5\varepsilon_1. \end{aligned} \quad (4.59)$$

Therefore,

$$\frac{1}{n\tilde{\sigma}_{n^2}} \leq \left(1 - \frac{50\varepsilon_1}{1 - 5\varepsilon_1}\right)^{-\frac{1}{2}} \quad \text{and} \quad \frac{1}{n\tilde{\sigma}_{n^2}} \geq \left(1 + \frac{50\varepsilon_1}{1 - 5\varepsilon_1} + \varepsilon_1(2 + \varepsilon_1)\right)^{-\frac{1}{2}}. \quad (4.60)$$

First, note that by Cauchy-Schwarz,

$$(\pi \otimes \pi)^t \tilde{v}_{n^2} \tilde{u}_{n^2}^t \mathbf{1}_{n^2} - 1 \leq n\|\pi\|^2 - 1 \leq \varepsilon_1. \quad (4.61)$$

By the matrix perturbation Theorem A.1, we can write

$$\tilde{v}_{(n,n)} = (v_{(n,n)} + V_2Q) (1 + \|Q\|^2)^{-\frac{1}{2}}, \quad (4.62)$$

where $\|Q\| \leq \frac{2\|\Delta_{21}\|}{\delta}$. Applying Theorem A.1 to $A = LL^t$, we find that there exists a matrix W with

$$\|W\| \leq \frac{2}{\delta_W} \|\gamma_{11}g_{21}^t + g_{12}^t\Sigma_2 + g_{12}^tG_{22}\| \quad (4.63)$$

such that

$$\tilde{u}_{(n,n)} = (u_{(n,n)} + U_2W) (1 + \|W\|^2)^{-\frac{1}{2}}. \quad (4.64)$$

The parameter δ_W is given by

$$\delta_W = \|\Sigma_2^{-2}\|^{-1} - \gamma_{11}^2 - \|g_{12}\|^2 - \|G_{22}^t\Sigma_2 + \Sigma_2G_{22} + G_{22}^tG_{22} + g_{21}g_{21}^t\|. \quad (4.65)$$

Note that we can directly compute the unperturbed projector through

$$|(\pi \otimes \pi)^t v_{(n,n)} u_{(n,n)}^t \mathbf{1}_{n^2}| = \frac{1}{n\|\pi\|}. \quad (4.66)$$

Furthermore, we note that

$$|(\pi \otimes \pi)^t v_{(n,n)} W^t U_2^t \mathbf{1}_{n^2}| \leq \frac{1}{n} \|W\| \|U_2\| n \leq \|W\|, \quad (4.67)$$

$$|(\pi \otimes \pi)^t V_2 Q u_{(n,n)}^t \mathbf{1}_{n^2}| \leq \frac{1}{\|\pi\|^2} \|\pi \otimes \pi\| \|V_2\| \|Q\| \leq \|Q\|, \quad (4.68)$$

$$|(\pi \otimes \pi)^t V_2 Q W^t U_2^t \mathbf{1}_{n^2}| \leq n\|\pi\|^2 \|V_2\| \|Q\| \|W\| \|U_2\| \leq n\|\pi\|^2 \|Q\| \|W\|. \quad (4.69)$$

So that in total we obtain,

$$(\pi \otimes \pi)^t \tilde{v}_{(n,n)} \tilde{u}_{(n,n)}^t \mathbf{1}_{n^2} \geq \frac{\left(\frac{1}{n\|\pi\|^2} - \|W\| - \|Q\| - n\|\pi\|^2 \|W\| \|Q\| \right)}{(1 + \|Q\|^2)^{\frac{1}{2}} (1 + \|W\|^2)^{\frac{1}{2}}}. \quad (4.70)$$

The norm of Q can be bounded using (4.57),

$$\|Q\| \leq \frac{2\|\Delta_{21}\|}{\delta} \leq \frac{10\sqrt{\varepsilon_1}}{(1 - 5\varepsilon_1)n}. \quad (4.71)$$

The norm of W is bounded from above by

$$\|W\| \leq \frac{2\|\gamma_{11}g_{21}^t + g_{12}\Sigma_2 + g_{12}^tG_{22}\|}{\delta_W} \leq \frac{8(1 + \varepsilon_1)^2}{(1 - 5\varepsilon_1)\sqrt{n}}. \quad (4.72)$$

Note that for

$$n \geq \max \left\{ \frac{10\sqrt{\varepsilon_1}}{(1 - 5\varepsilon_1)\varepsilon_1}, \left(\frac{8(1 + \varepsilon_1)}{\varepsilon_1} \right)^2 \right\}, \quad (4.73)$$

we have that $\|Q\| \leq \varepsilon_1$ and $\|W\| \leq \varepsilon_1$. Therefore,

$$(\pi \otimes \pi)^t \tilde{v}_{(n,n)} \tilde{u}_{(n,n)}^t \mathbf{1}_{n^2} \geq \frac{\left(\frac{1}{(1+\varepsilon_1)} - 2\varepsilon_1 - (1 + \varepsilon_1)\varepsilon_1^2 \right)}{(1 + \varepsilon_1^2)}. \quad (4.74)$$

Finally, we see that

$$\frac{n\|\pi\|^2}{n\tilde{\sigma}_{n^2-1}} \leq \frac{n\|\pi\|^2}{n(1 - \sigma_2(P \otimes P))} \leq \frac{1 + \varepsilon_1}{n(1 - \varepsilon_1)}. \quad (4.75)$$

Therefore, for $n \geq \frac{1+\varepsilon_1}{\varepsilon_1(1-\varepsilon_1)}$,

$$\frac{n\|\pi\|^2}{n\tilde{\sigma}_{n^2-1}} \leq \varepsilon_1. \quad (4.76)$$

The result follows from observing that for every $\varepsilon_2 > 0$, there exists $\varepsilon_1 \in (0, \frac{1}{20})$ such that

$$\begin{aligned} \frac{\left(\frac{1}{(1+\varepsilon_1)} - 2\varepsilon_1 - (1 + \varepsilon_1)\varepsilon_1^2 \right)}{(1 + \varepsilon_1^2)} &\geq 1 - \varepsilon_2, & \varepsilon_1 &\leq \varepsilon_2 \\ \left(1 - \frac{50\varepsilon_1}{1 - 5\varepsilon_1} \right)^{-\frac{1}{2}} &\leq 1 + \varepsilon_2 & \text{and} & \left(1 + \frac{50\varepsilon_1}{1 - 5\varepsilon_1} + \varepsilon_1(2 + \varepsilon_1) \right)^{-\frac{1}{2}} &\geq 1 - \varepsilon_2. \end{aligned} \quad (4.77)$$

Setting

$$N_2 = \max \left\{ N_1, \frac{10\sqrt{\varepsilon_1}}{(1 - 5\varepsilon_1)\varepsilon_1}, \left(\frac{8(1 + \varepsilon_1)}{\varepsilon_1} \right)^2, \frac{1 + \varepsilon_1}{\varepsilon_1(1 - \varepsilon_1)} \right\}, \quad (4.78)$$

where N_1 is the constant in Lemma 4.4, we obtain the result. \square

Finally, we are ready to prove Theorem 1.3.

Proof of Theorem 1.3. Note that for any $\varepsilon, \nu_1, \nu_2 > 0$ and $n \in \mathbb{N}$,

$$\mathbb{P} \left(\left| \frac{1}{n} t_{\text{meet}}^\pi - 1 \right| \leq \varepsilon \right) \geq \mathbb{P} \left(\left| \frac{1}{n} t_{\text{meet}}^\pi - 1 \right| \leq \varepsilon \mid F_{\nu_1, \nu_2} \cap F \right) - \mathbb{P} \left((F_{\nu_1, \nu_2} \cap F)^c \right). \quad (4.79)$$

From Lemma 4.1 and Proposition 4.2, it follows that there exists $\theta > 0$, such that for any $\nu_1, \nu_2 > 0$,

$$\mathbb{P}((F_{\nu_1, \nu_2} \cap F)^c) \leq 2n \exp\left(-\frac{\nu_1^2 np}{3}\right) + 2 \binom{n}{2} \exp\left(-\frac{\nu_2^2 np^2}{3}\right) + \exp(-\theta (\log n)^2). \quad (4.80)$$

Therefore, it remains to show that for every $\epsilon > 0$, there exist $\nu_1, \nu_2 > 0$ and $N \in \mathbb{R}_+$ such that for all $n \geq N$,

$$\mathbb{P}\left(\left|\frac{1}{n}t_{\text{meet}}^\pi - 1\right| \leq \epsilon \mid F_{\nu_1, \nu_2} \cap F\right) = 1. \quad (4.81)$$

We use the rank- k perturbation result 1.2, with $k = 1$, to deduce that,

$$\frac{1}{n}t_{\text{meet}}^\pi \leq \frac{1}{n}\hat{t}_{\text{meet}}^{\pi, (1)} + \frac{n\|\pi\|^2}{n\tilde{\sigma}_{n^2-1}} \leq \frac{1}{n\tilde{\sigma}_{n^2}}(\pi \otimes \pi)^{t\tilde{v}_{n^2}\tilde{u}_{n^2}\mathbf{1}_{n^2}} + \frac{n\|\pi\|^2}{n\tilde{\sigma}_{n^2-1}}. \quad (4.82)$$

$$\frac{1}{n}t_{\text{meet}}^\pi \geq \frac{1}{n}\hat{t}_{\text{meet}}^{\pi, (1)} - \frac{n\|\pi\|^2}{n\tilde{\sigma}_{n^2-1}} \geq \frac{1}{n\tilde{\sigma}_{n^2}}(\pi \otimes \pi)^{t\tilde{v}_{n^2}\tilde{u}_{n^2}\mathbf{1}_{n^2}} - \frac{1}{n} - \frac{n\|\pi\|^2}{n\tilde{\sigma}_{n^2-1}}. \quad (4.83)$$

By Lemma 4.5, for every $\varepsilon_2 > 0$, there exist constants $\nu_1, \nu_2 > 0$ and $N_2 \in \mathbb{R}_+$ such that for all $n \geq N_2$, conditioned on F_{ν_1, ν_2} and F we have,

$$\frac{1}{n}t_{\text{meet}}^\pi \leq (1 + \varepsilon_2)^2 + \varepsilon_2. \quad (4.84)$$

if additionally $n \geq \frac{1}{\varepsilon_2}$,

$$\frac{1}{n}t_{\text{meet}}^\pi \geq (1 - \varepsilon_2)^2 - 2\varepsilon_2. \quad (4.85)$$

Equation (4.81) follows from noting that for every $\epsilon > 0$, there exists $\varepsilon_2 > 0$ such that

$$(1 + \varepsilon_2)^2 + \varepsilon_2 - 1 \leq \epsilon \quad \text{and} \quad 1 - (1 - \varepsilon_2)^2 + 2\varepsilon_2 \leq \epsilon. \quad (4.86)$$

□

5. DISCUSSION

We proposed an approach to estimate the meeting time of two independent random walks on graphs via SVD of the diagonally killed generator of the pair. Specifically, we viewed the diagonally killed generator as a perturbation of the (plain) generator. We devised non-asymptotic perturbation bounds on the expected meeting time via a rank-one approximation. Next, we conducted non-asymptotic perturbation analysis of the rank one approximation to the diagonally killed generator. As a proof of the concept, we specialized to a sufficiently dense Erdős–Rényi random graph and obtained sharp bounds on the expected meeting time, cf. Theorem 1.3. In Section 4, we defined two graph characteristics R_1 and R_2 (see (4.1), (4.3)) which we used to control the meeting-time. While our main Theorem 1.3 is stated for sufficiently dense Erdős–Rényi graphs, this dependence can be factored out to some extent. By inspecting the proof of Theorem 1.3, one can convince oneself that the theorem can be generalized as follows.

Theorem 5.1 (Informal). *Let $(G_n)_{n \in \mathbb{N}}$ be a sequence of (random) graphs such that each G_n has n vertices. Suppose there exists a real sequence $(d_n)_{n \in \mathbb{N}}$ such that (w.h.p.), as $n \rightarrow \infty$, the conditions*

- (1) $\max_{i \in V_n} \left| \frac{\deg i}{d_n} - 1 \right| \rightarrow 0$,
- (2) $\max_{(k, \ell) \in V_n \times V_n} \left| \frac{n}{d_n^2} \left(\sum_i A_{ik} A_{i\ell} \right) - 1 \right| \rightarrow 0$,
- (3) $\sigma_2(A) = O(\sqrt{d_n})$,

hold, where $\sigma_2(A)$ is the second-largest eigenvalue of the adjacency matrix $A = A(G_n)$. Then,

$$\frac{1}{n}t_{\text{meet}}^\pi \approx 1 \text{ (w.h.p.)}. \quad (5.1)$$

When it comes to applying Theorem 5.1 to the Erdős–Rényi random graph, it is the second condition which is the most restrictive. Specifically, it restricts the statement of Theorem 1.3 to sufficiently dense Erdős–Rényi random graphs ($\beta > 0.5$). Indeed, Items (1) and (3) hold with high probability for $p_n \gg (\log n)/n$, cf. Lemma 4.1 and [ADK21], whereas the probability of (2) is only high exactly in the regime, where the average degree grows faster than \sqrt{n} . We conjecture that the perturbation method can be adapted to go through under Conditions (1), (3) and a substantially milder condition than (2). We will explore this elsewhere.

6. LITERATURE

- [Abb+20] E. Abbe, J. Fan, K. Wang, and Y. Zhong. “Entrywise eigenvector analysis of random matrices with low expected rank”. *Ann. Statist.* 48.3 (2020), pp. 1452–1474. DOI: 10.1214/19-AOS1854.
- [ADK21] J. Alt, R. Ducatez, and A. Knowles. “Extremal eigenvalues of critical Erdős–Rényi graphs”. *Ann. Probab.* 49.3 (2021), pp. 1347–1401. DOI: 10.1214/20-aop1483.
- [AF02] D. Aldous and J. A. Fill. *Reversible Markov Chains and Random Walks on Graphs*. Unfinished monograph, recompiled 2014, available at <http://www.stat.berkeley.edu> 2002.
- [Ald13] D. Aldous. “Interacting particle systems as stochastic social dynamics”. *Bernoulli* 19.4 (2013), pp. 1122–1149. DOI: 10.3150/12-BEJSP04.
- [Ave+23] L. Avena, F. Capannoli, R. S. Hazra, and M. Quattropani. “Meeting, coalescence and consensus time on random directed graphs”. *arXiv preprint* (2023). DOI: 10.48550/arXiv.2308.01832.
- [Ave+24] L. Avena, R. Baldasso, R. S. Hazra, F. den Hollander, and M. Quattropani. “Discordant edges for the voter model on regular random graphs”. *ALEA Lat. Am. J. Probab. Math. Stat.* 21.1 (2024), pp. 431–464. DOI: 10.30757/alea.v21-18.
- [BBP05] J. Baik, G. Ben Arous, and S. Péché. “Phase transition of the largest eigenvalue for nonnull complex sample covariance matrices”. *Ann. Probab.* 33.5 (2005), pp. 1643–1697. DOI: 10.1214/009117905000000233.
- [BH15] A. Bovier and F. den Hollander. *Metastability. A potential-theoretic approach*. Springer, Cham, 2015, pp. xxi+581. DOI: 10.1007/978-3-319-24777-9.
- [BN12] F. Benaych-Georges and R. R. Nadakuditi. “The singular values and vectors of low rank perturbations of large rectangular random matrices”. *J. Multivariate Anal.* 111 (2012), pp. 120–135. DOI: 10.1016/j.jmva.2012.04.019.
- [CFR10] C. Cooper, A. Frieze, and T. Radzik. “Multiple random walks in random regular graphs”. *SIAM J. Discrete Math.* 23.4 (2010), pp. 1738–1761. DOI: 10.1137/080729542.
- [Cha23] S. Chatterjee. “Spectral gap of nonreversible Markov chains”. *Preprint* (2023). DOI: 10.48550/arXiv.2310.10876.
- [Che+21] Y. Chen, Y. Chi, J. Fan, C. Ma, et al. “Spectral methods for data science: A statistical perspective”. *Found. Trends Mach. Learn.* 14.5 (2021), pp. 566–806. DOI: 10.1561/22000000079.

- [Che21] Y.-T. Chen. “Precise asymptotics of some meeting times arising from the voter model on large random regular graphs”. *Electron. Commun. Probab.* 26 (2021), Paper No. 5, 13. DOI: 10.1214/21-ECP373.
- [DK70] C. Davis and W. M. Kahan. “The rotation of eigenvectors by a perturbation. III”. *SIAM J. Numer. Anal.* 7 (1970), pp. 1–46. DOI: 10.1137/0707001.
- [Dur10] R. Durrett. “Some features of the spread of epidemics and information on a random graph”. *Proc. Natl. Acad. Sci. USA* 107.10 (2010), pp. 4491–4498. DOI: 10.1073/pnas.0914402107.
- [Erd+12] L. Erdős, A. Knowles, H.-T. Yau, and J. Yin. “Spectral statistics of Erdős-Rényi Graphs II: Eigenvalue spacing and the extreme eigenvalues”. *Comm. Math. Phys.* 314.3 (2012), pp. 587–640. DOI: 10.1007/s00220-012-1527-7.
- [Erd+13] L. Erdős, A. Knowles, H.-T. Yau, and J. Yin. “Spectral statistics of Erdős-Rényi graphs I: Local semicircle law”. *Ann. Probab.* 41.3B (2013), pp. 2279–2375. DOI: 10.1214/11-AOP734.
- [EY36] C. Eckart and G. Young. “The approximation of one matrix by another of lower rank”. *Psychometrika* 1.3 (1936), pp. 211–218.
- [FO23] J. Fernley and M. Ortgiese. “Voter models on subcritical scale-free random graphs”. *Random Structures Algorithms* 62.2 (2023), pp. 376–429. DOI: 10.1002/rsa.21107.
- [GPB18] M. George, R. Patel, and F. Bullo. “The meeting time of multiple random walks”. *Preprint* (2018). DOI: 10.48550/arXiv.1806.08843.
- [Her+22] J. Hermon, S. Li, D. Yao, and L. Zhang. “Mean field behavior during the big bang regime for coalescing random walks”. *Ann. Probab.* 50.5 (2022), pp. 1813–1884. DOI: 10.1214/22-aop1571.
- [Kat95] T. Kato. *Perturbation theory for linear operators*. Classics in Mathematics. Reprint of the 1980 edition. Springer-Verlag, Berlin, 1995, pp. xxii+619. DOI: 10.1007/978-3-642-66282-9.
- [KMS23] V. Kanade, F. Mallmann-Trenn, and T. Sauerwald. “On coalescence time in graphs: when is coalescing as fast as meeting?” *ACM Trans. Algorithms* 19.2 (2023), Art. 18, 46. DOI: 10.1145/3576900.
- [Lov96] L. Lovász. “Random walks on graphs: a survey”. In: *Combinatorics, Paul Erdős is eighty, Vol. 2 (Keszthely, 1993)*. Vol. 2. Bolyai Soc. Math. Stud. János Bolyai Math. Soc., Budapest, 1996, pp. 353–397.
- [LP17] D. A. Levin and Y. Peres. *Markov chains and mixing times*. American Mathematical Society, Providence, RI, 2017, pp. xvi+447. DOI: 10.1090/mbk/107.
- [LT14] M. Löwe and F. Torres. “On hitting times for a simple random walk on dense Erdős-Rényi random graphs”. *Statist. Probab. Lett.* 89 (2014), pp. 81–88. DOI: 10.1016/j.spl.2014.02.017.
- [LT23] M. Löwe and S. Terveer. “A central limit theorem for the mean starting hitting time for a random walk on a random graph”. *J. Theoret. Probab.* 36.2 (2023), pp. 779–810. DOI: 10.1007/s10959-022-01195-9.
- [MPL17] N. Masuda, M. A. Porter, and R. Lambiotte. “Random walks and diffusion on networks”. *Phys. Rep.* 716/717 (2017), pp. 1–58. DOI: 10.1016/j.physrep.2017.07.007.
- [MT17] E. Mossel and O. Tamuz. “Opinion exchange dynamics”. *Probab. Surv.* 14 (2017), pp. 155–204. DOI: 10.1214/14-PS230.
- [Oli13] R. I. Oliveira. “Mean field conditions for coalescing random walks”. *Ann. Probab.* 41.5 (2013), pp. 3420–3461. DOI: 10.1214/12-AOP813.

- [OVW18] S. O’Rourke, V. Vu, and K. Wang. “Random perturbation of low rank matrices: improving classical bounds”. *Linear Algebra Appl.* 540 (2018), pp. 26–59. DOI: 10.1016/j.laa.2017.11.014.
- [Sch26] E. Schrödinger. “Quantisierung als Eigenwertproblem”. *Ann. Physik* 385.13 (1926), pp. 437–490.
- [SS90] G. W. Stewart and J. G. Sun. *Matrix perturbation theory*. Computer Science and Scientific Computing. Academic Press, Inc., Boston, MA, 1990, pp. xvi+365.
- [Ste73] G. W. Stewart. “Error and perturbation bounds for subspaces associated with certain eigenvalue problems”. *SIAM Rev.* 15 (1973), pp. 727–764. DOI: 10.1137/1015095.
- [Ste84] G. W. Stewart. “A second order perturbation expansion for small singular values”. *Linear Algebra Appl.* 56 (1984), pp. 231–235. DOI: 10.1016/0024-3795(84)90128-9.
- [Ver18] R. Vershynin. *High-dimensional probability*. Vol. 47. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, Cambridge, 2018, pp. xiv+284. DOI: 10.1017/9781108231596.
- [Wed72] P.-A. Wedin. “Perturbation bounds in connection with singular value decomposition”. *Nordisk Tidskr. Informationsbehandling (BIT)* 12 (1972), pp. 99–111. DOI: 10.1007/bf01932678.

APPENDIX A.

Theorem A.1. *Let $B, \Delta \in \mathbb{R}^{n \times n}$. Let $X = (X_1, X_2)$ be unitary with $X_1 \in \mathbb{R}^{n \times 1}$ and suppose $\mathcal{R}(X_1)$ is an invariant subspace of B . Let $X^t B X$ and $X^t \Delta X$ be partitioned conformally with X in the forms*

$$X^t B X = \begin{bmatrix} B_{11} & B_{12} \\ 0 & B_{22} \end{bmatrix} \quad (\text{A.1})$$

and

$$X^t \Delta X = \begin{bmatrix} \Delta_{11} & \Delta_{12} \\ \Delta_{21} & \Delta_{22} \end{bmatrix} \quad (\text{A.2})$$

Let $\delta = \text{sep}(B_{11}, B_{22}) - \|\Delta_{11}\| - \|\Delta_{22}\|$. If

$$\frac{\|\Delta_{21}\|(\|A_{12}\| + \|\Delta_{12}\|)}{\delta^2} \leq \frac{1}{4}, \quad (\text{A.3})$$

there is a matrix Q with $\|Q\| \leq 2 \frac{\|\Delta_{21}\|}{\delta}$ such that the columns of

$$X'_1 = (X_1 + X_2 Q)(I + Q^t Q)^{-1/2} \quad (\text{A.4})$$

span an invariant subspace of $B + \Delta$. Furthermore, the matrices B'_{11} and B'_{22} are given by

$$B'_{11} = B_{11} + \Delta_{11} + (B_{12} + \Delta_{12})Q \quad (\text{A.5})$$

and

$$B'_{22} = (I + Q Q^t)^{1/2} (B_{22} + \Delta_{22} + Q(B_{12} + \Delta_{12})) (I + Q Q^t)^{-1/2}. \quad (\text{A.6})$$

The set $\lambda(B + \Delta)$ is the disjoint union of $\lambda(B'_{11})$ and $\lambda(B'_{22})$.

T.M. VAN BELLE, UNIVERSITÄT DUISBURG-ESSEN, FAKULTÄT FÜR MATHEMATIK, THEA-LEYMANN-STR. 9, 45127 ESSEN, GERMANY

Email address: thomas.vanbelle@uni-due.de

A. KLIMOVSKY, JULIUS-MAXIMILIANS-UNIVERSITÄT WÜRZBURG, INSTITUT FÜR MATHEMATIK, LEHRSTUHL FÜR MATHEMATIK VIII: ANGEWANDTE STOCHASTIK, EMIL-FISCHER-STR. 30, 97074 WÜRZBURG, GERMANY

Email address: anton.klimovsky@mathematik.uni-wuerzburg.de