

Experimental Bounds on Deformed Muon Lifetime Dilation

Iarley P. Lobo^{1,*} and Christian Pfeifer^{2,†}

¹*Department of Chemistry and Physics, Federal University of Paraíba,
Rodovia BR 079 - km 12, 58397-000 Areia-PB, Brazil*

²*ZARM, University of Bremen, 28359 Bremen, Germany.*

(Dated: June 11, 2024)

We analyze Planck scale induced modifications of the relativistic time dilation using data from the Muon Storage Ring experiment at CERN. By examining the lifetimes of muons, we establish, for the first time, a constraint on such quantum gravity-inspired deformations using this channel.

I. INTRODUCTION

The debate over whether Poincaré symmetry is preserved at a fundamental level has been ongoing for decades, particularly in discussions about extensions of the Standard Model [1] and quantum gravity phenomenology [2, 3]. In general, the models discussed in the literature can be categorized into two scenarios: a) Lorentz Invariance Violation (LIV), where the relativity principle is broken and inertial frames – those in which particle behavior is invariant – would no longer exist; b) Deformed (or Doubly) Special Relativity (DSR), where a deformation or modification of Poincaré symmetry maps inertial frames onto each other.

In the LIV case, one still has that the map between frames that are related by relative velocities, translations and rotations as the standard Poincaré transformations however, these are no longer symmetry transformations of the physical system described and thus preferred frames exist. In the DSR case, the transformations between inertial frames are modified with corrections suppressed by the energy scale that indicates deviations from Poincaré symmetry [4–7]. In general, both models predict some effects such as time delays in simultaneously emitted massless particles, with corrections that are similar enough to make it challenging to distinguish between LIV and DSR models.

A series of papers [8–15] have shown that deformations of boosts in DSR present a promising avenue for investigating quantum gravity effects through the time dilation of particle lifetimes. In particular, the authors of [12–15] (which we are included) identified an amplification factor proportional to the square of the ratio between the energy and mass of the unstable particle $(E/m)^2$. If confirmed, this effect could serve as strong evidence that Poincaré symmetry needs to be extended at a fundamental level.

To place bounds on the modified time dilation effect and identify better ways to analyze this scenario with higher sensitivity, we examine one of the few cases where the dilated lifetime of a fundamental particle is reported with a specific energy: the $g-2$ experiment performed at the Muon Storage Ring at CERN in [16, 17]. This paper represents the first instance of analyzing a boosted lifetime report from the perspective of a deformation of Lorentz symmetry by Planck scale effects, incorporating such a generic range of corrections.

This paper is organized as follows. In section II, we describe how the deformed time dilation is computed in an effective spacetime. In section III, we analyze the data from CERN to set bounds on the deformed time dilation. In section IV we discuss our results. We use units in which $\hbar = c = 1$.

II. REVISITING TIME DILATION COMPUTATION

Our starting point is a generic class of modified dispersion relations that describe propagation through a quantum spacetime, expressed as:

$$E^2 - |p|^2 = m^2 + s\xi^{(n)} \frac{|p|^{n+2}}{E_{\text{Pl}}^n}, \quad (1)$$

where $s = +1$ (-1) and determines if the cases of superluminal (subluminal) propagation, E is the energy of the test particle, $|p| = |\vec{p}|$ is its spatial momentum and m is its mass. We refer to $\xi^{(n)}$ as a dimensionless parameter that controls the kinematical deformations suppressed by the n -th power of the Planck energy $E_{\text{Pl}} = \sqrt{\hbar c^5/G} \approx 1.22 \times 10^{19}$ GeV.

* lobofisica@gmail.com

† christian.pfeifer@zarm.uni-bremen.de

A suitable geometry to describe the motion of such a particle through a quantum spacetime is Finsler geometry [18, 19]. The arc-length in Finsler geometry, derived from a Legendre transformation [20] on the Helmholtz action of a free particle with the above Modified Dispersion Relation (MDR) (1), is shown in to be [12, 15, 19, 21–24]:

$$S = \int d\lambda F(x, \dot{x}) = \int d\lambda \left[\sqrt{\dot{t}^2 - \dot{x}^2} + s \frac{\xi^{(n)}}{2} \left(\frac{m}{E_{\text{Pl}}} \right)^n \frac{|\dot{x}|^{n+2}}{(\dot{t}^2 - \dot{x}^2)^{\frac{n+1}{2}}} \right], \quad (2)$$

where F is the Finsler function and “dot” means derivative with respect to the parameter λ . The resulting particle trajectories extremising these length functions are Finsler geodesics, and the deformed Poincaré transformations are isometries of the Finsler function. From these results, it is shown in [15] that the deformed Lorentz transformation connecting the rest frame of the particle and the laboratory (lab) frame can be derived from an extension of the clock postulate to Finsler geometry [12]. Specifically, the proper time τ measured by a comoving observer is given by the arc-length S of its trajectory in spacetime divided by the speed of light c .

Using this observation, along with the expression for the energy and momentum of the particle in the lab frame, $p_\mu = m\partial F/\partial \dot{x}^\mu$, we derive the following formula for the dilation of the proper time τ measured by the comoving frame of a fundamental particle of mass m and the time t measured by lab frame for which it has energy and momentum $(E, |p|)$

$$t_{\text{DSR}} = \gamma_{\text{DSR}}\tau = \frac{E}{m} \left[1 + s \frac{n\xi^{(n)}}{2} \left(\frac{|p|}{m} \right)^2 \left(\frac{|p|}{E_{\text{Pl}}} \right)^n \right] \tau \quad (3)$$

$$= \frac{\sqrt{|p|^2 + m^2}}{m} \left\{ 1 + s \frac{n\xi^{(n)}}{E_{\text{Pl}}^n} \frac{|p|^{2+n} [m^2(n+1) + np^2]}{2m^2(p^2 + m^2)} \right\} \tau, \quad (4)$$

where in the last step, we used the MDR (1). For an unstable particle, this expression gives the dilation of its proper lifetime τ by a deformed Lorentz factor, γ_{DSR} , which is consistent with DSR principles [15].

In the next section, we will set bounds on the parameter $\xi^{(n)}$ using data from experiments measuring the anomalous magnetic moment of the muon $g - 2$ at CERN. These experiments provide explicit values for the Lorentz factor and the dilated lifetime of the muon, allowing us to constrain $\xi^{(n)}$.

III. BOUNDS FROM THE MUON STORAGE RING EXPERIMENT AT CERN

At the Muon Storage Ring at CERN, researchers measured the anomalous magnetic moment of both muons and antimuons, as detailed in the final report [17]. This experiment scrutinized the orbital and spin motion of highly polarized muons within a magnetic storage ring. The process involved proton collisions within a synchrotron accelerator, yielding pions that decayed into muons alongside neutrinos. Subsequently, these muons were injected into a region featuring a uniform magnetic field, where they were accelerated to traverse circular paths, thereby experiencing time dilation.

Notably, the same procedure was employed in the recent $g - 2$ experiment at the Fermilab [25]. This endeavor significantly enhanced the precision of CERN’s experiment. However, while both experiments aimed to measure the anomalous magnetic moment, the CERN setup also delved into quantifying the dilation of muon lifetimes—a facet not explored in Fermilab $g - 2$, which primarily focused on measuring the muon’s magnetic moment. Therefore, we have opted for the CERN experiment as it is more conducive to establishing constraints on a modified time dilation effect.

In [16], the reported momentum of the muon and the weighted average of the measured lifetime for both muons and antimuons, along with its mass and mean lifetime according to the Particle Data Group (PDG) [26], are

$$|p| = 3.094 \text{ GeV}, \quad t_{\text{exp}} = 64.378 \pm 0.026 \mu\text{s}, \quad m_\mu = 105.658 \text{ MeV}, \quad \tau_\mu = 2.197 \mu\text{s}. \quad (5)$$

Using these values, we can calculate the bounds on the quantum gravity parameter $\xi^{(n)}$ by considering the maximum and minimum values of t_{exp} . For $s = \pm 1$, the bounds for $n = 1$ and $n = 2$ are

$$\xi^{(1)} \leq 9.191 \times 10^{21}, \quad \xi^{(2)} \leq 1.813 \times 10^{40} \quad (6)$$

These bounds are far from the Planck scale, a consequence of the measurements being conducted with particles at GeV energy levels. This is typical of experiments aimed at measuring the anomalous magnetic moment of the muon. This value of the Lorentz factor effectively removes the contribution of the stabilizing quadrupole electrostatic field

from the muon's relation between the angular frequency and the electromagnetic field, described by the Thomas-Bargmann-Michel-Telegdi equation [27].

However, since this effect has an amplifier of the order $(|p|/m)^2$ (4), improvements could be achieved by examining observables that depend on the dilated lifetime of more energetic particles, such as the spectra of particles produced by cosmic rays [13].

IV. DISCUSSION

We have calculated the first bound on a modification of time dilation due to DSR effects from the lifetime of elementary particles. This effect can be derived by extending the clock postulate to an effective spacetime, a Finsler manifold, probed by particles whose kinematics are modified by quantum gravity effects. This result is compatible with a deformation of special relativity, as the Finsler norm is preserved by such transformations, as shown in [15].

We compared the lifetimes of particles in the comoving frame with those in the lab frame. To directly set a bound, we used an experiment where the momentum of a massive particle is known with high accuracy and the corresponding dilated lifetime was measured and published. The appropriate experiment was conducted at the CERN Muon Storage Ring [17], which provided data on measurements of relativistic time dilation in a controlled environment [16].

We found that the bound for the first-order correction lies 21 orders of magnitude above the Planck scale, and for the second-order correction, it lies 40 orders of magnitude above. The smallness of the effect is due to the relatively low energies of the muons, approximately in the GeV range. To improve these constraints, it would be necessary to either consider experiments or observations with particles at higher energies or to enhance the precision of the measurements. Observables that include the dilated lifetimes of particles are candidates to probe this effect. We discuss this matter in regarding prospects in accelerators [14] and for cosmic rays [13]. In the future, we expect to consider these cases following appropriate data analysis.

ACKNOWLEDGMENTS

I. P. L. was partially supported by the National Council for Scientific and Technological Development - CNPq grant 312547/2023-4. CP acknowledges support by the excellence cluster QuantumFrontiers of the German Research Foundation (Deutsche Forschungsgemeinschaft, DFG) under Germany's Excellence Strategy – EXC-2123 Quantum-Frontiers – 390837967 and was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) - Project Number 420243324.

-
- [1] Kostelecky, V. Alan, “Gravity, Lorentz violation, and the standard model,” *Phys. Rev. D* **69** (2004) 105009, [arXiv:hep-th/0312310](#).
 - [2] G. Amelino-Camelia, “Quantum-Spacetime Phenomenology,” *Living Rev. Rel.* **16** (2013) 5, [arXiv:0806.0339 \[gr-qc\]](#).
 - [3] Addazi, A. and others, “Quantum gravity phenomenology at the dawn of the multi-messenger era—A review,” *Prog. Part. Nucl. Phys.* **125** (2022) 103948, [arXiv:2111.05659 \[hep-ph\]](#).
 - [4] Lukierski, Jerzy and Ruegg, Henri and Nowicki, Anatol and Tolstoi, Valerii N., “Q deformation of Poincare algebra,” *Phys. Lett. B* **264** (1991) 331–338.
 - [5] Lukierski, Jerzy and Nowicki, Anatol and Ruegg, Henri, “New quantum Poincare algebra and k deformed field theory,” *Phys. Lett. B* **293** (1992) 344–352.
 - [6] Majid, S. and Ruegg, H., “Bicrossproduct structure of kappa Poincare group and noncommutative geometry,” *Phys. Lett. B* **334** (1994) 348–354, [arXiv:hep-th/9405107](#).
 - [7] G. Amelino-Camelia, “Relativity in space-times with short distance structure governed by an observer independent (Planckian) length scale,” *Int. J. Mod. Phys. D* **11** (2002) 35–60, [arXiv:gr-qc/0012051 \[gr-qc\]](#).
 - [8] Arzano, Michele and Kowalski-Glikman, Jerzy and Wislicki, Wojciech, “A bound on Planck-scale deformations of CPT from muon lifetime,” *Phys. Lett. B* **794** (2019) 41–44, [arXiv:1904.06754 \[hep-ph\]](#).
 - [9] Arzano, Michele and Kowalski-Glikman, Jerzy and Wislicki, Wojciech, “Planck-scale deformation of CPT and particle lifetimes,” [arXiv:2009.03135 \[hep-ph\]](#).
 - [10] Bevilacqua, Andrea and Kowalski-Glikman, Jerzy and Wislicki, Wojciech, “ κ -deformed complex scalar field: Conserved charges, symmetries, and their impact on physical observables,” *Phys. Rev. D* **105** no. 10, (2022) 105004, [arXiv:2201.10191 \[hep-th\]](#).
 - [11] Bevilacqua, Andrea and Kowalski-Glikman, Jerzy and Wislicki, Wojciech, “Deformation-induced CPT violation in entangled pairs of neutral kaons,” [arXiv:2404.03600 \[hep-ph\]](#).

- [12] Lobo, Iarley P. and Pfeifer, Christian, “Reaching the Planck scale with muon lifetime measurements,” *Phys. Rev. D* **103** no. 10, (2021) 106025, [arXiv:2011.10069 \[hep-ph\]](#).
- [13] Lobo, Iarley P. and Pfeifer, Christian and Morais, Pedro H. and Batista, Rafael Alves and Bezerra, Valdir B., “Two-body decays in deformed relativity,” *JHEP* **09** (2022) 003, [arXiv:2112.12172 \[hep-ph\]](#).
- [14] Lobo, Iarley P. and Pfeifer, Christian, “Muon accelerators—muon lifetime measurements as window to Planck scale physics,” *Class. Quant. Grav.* **41** no. 1, (2024) 015008, [arXiv:2306.07210 \[hep-ph\]](#).
- [15] Morais, Pedro H. and Lobo, Iarley P. and Pfeifer, Christian and Alves Batista, Rafael and Bezerra, Valdir B., “Modified particle lifetimes as a signature of deformed relativity,” *Phys. Lett. B* **848** (2024) 138380, [arXiv:2308.03716 \[gr-qc\]](#).
- [16] Bailey, J. and others, “Measurements of Relativistic Time Dilatation for Positive and Negative Muons in a Circular Orbit,” *Nature* **268** (1977) 301–305.
- [17] **CERN-Mainz-Daresbury** Collaboration, Bailey, J. and others, “Final Report on the CERN Muon Storage Ring Including the Anomalous Magnetic Moment and the Electric Dipole Moment of the Muon, and a Direct Test of Relativistic Time Dilation,” *Nucl. Phys. B* **150** (1979) 1–75.
- [18] Girelli, Florian and Liberati, Stefano and Sindoni, Lorenzo, “Planck-scale modified dispersion relations and Finsler geometry,” *Phys. Rev. D* **75** (2007) 064015, [arXiv:gr-qc/0611024](#).
- [19] Pfeifer, Christian, “Finsler spacetime geometry in Physics,” *Int. J. Geom. Meth. Mod. Phys.* **16** no. supp02, (2019) 1941004, [arXiv:1903.10185 \[gr-qc\]](#).
- [20] Rodrigues, Ernesto and Lobo, Iarley P., “Revisiting legendre transformations in finsler geometry,” *International Journal of Geometric Methods in Modern Physics* (Mar., 2024) . <http://dx.doi.org/10.1142/S021988782450155X>.
- [21] Amelino-Camelia, Giovanni and Barcaroli, Leonardo and Gubitosi, Giulia and Liberati, Stefano and Loret, Niccoló, “Realization of doubly special relativistic symmetries in Finsler geometries,” *Phys. Rev. D* **90** no. 12, (2014) 125030, [arXiv:1407.8143 \[gr-qc\]](#).
- [22] Raetzel, Dennis and Rivera, Sergio and Schuller, Frederic P., “Geometry of physical dispersion relations,” *Phys. Rev. D* **83** (2011) 044047, [arXiv:1010.1369 \[hep-th\]](#).
- [23] Letizia, Marco and Liberati, Stefano, “Deformed relativity symmetries and the local structure of spacetime,” *Phys. Rev. D* **95** no. 4, (2017) 046007, [arXiv:1612.03065 \[gr-qc\]](#).
- [24] Lobo, Iarley P. and Loret, Niccoló and Nettel, Francisco, “Investigation of Finsler geometry as a generalization to curved spacetime of Planck-scale-deformed relativity in the de Sitter case,” *Phys. Rev. D* **95** no. 4, (2017) 046015, [arXiv:1611.04995 \[gr-qc\]](#).
- [25] **Muon g-2** Collaboration, Abi, B. and others, “Measurement of the Positive Muon Anomalous Magnetic Moment to 0.46 ppm,” *Phys. Rev. Lett.* **126** no. 14, (2021) 141801, [arXiv:2104.03281 \[hep-ex\]](#).
- [26] **Particle Data Group** Collaboration, Workman, R. L. and others, “Review of Particle Physics,” *PTEP* **2022** (2022) 083C01.
- [27] Jegerlehner, F., *The Anomalous Magnetic Moment of the Muon*. Springer Tracts in Modern Physics. Springer, 2008.