SPECTRAL PROPERTIES OF THE KRAMERS-FOKKER-PLANCK OPERATOR WITH A LONG-RANGE POTENTIAL

XUE PING WANG

ABSTRACT. We study real resonances and embedded eigenvalues of the Kramers-Fokker–Planck operator with a long-range potential. We prove that thresholds are only possible accumulation points of eigenvalues and that the limiting absorption principle holds true for energies outside an exceptional set. We also prove that the eigenfunctions associated with discrete eigenvalues decay exponentially and those associated with embedded non-threshold ones decay polynomially.

1. INTRODUCTION

In this work, we consider the Kramers–Fokker–Planck (KFP, for short) operator, also called the Kramers operator by physicists ([19]), given by

$$P = -\Delta_v + \frac{1}{4}|v|^2 - \frac{n}{2} + v \cdot \nabla_x - \nabla V(x) \cdot \nabla_v, \quad (x,v) \in \mathbb{R}^{2n}.$$
 (1.1)

V(x) is supposed to be a real-valued C^1 potential satisfying, for some constants $\rho > -1, C > 0$ the estimate

$$|V(x)| + \langle x \rangle |\nabla V(x)| \le C \langle x \rangle^{-\rho}, \quad x \in \mathbb{R}^n.$$
(1.2)

Here $\langle x \rangle = (1 + |x|^2)^{1/2}$ and $n \ge 1$. In most part of this work, we assume $\rho > 0$, i.e., the potential is long-range, unless in the analysis of decay of eigenfunctions associated with the discrete spectrum, the condition $\rho > -1$ is sufficient.

Let P_0 be the free KFP operator:

$$P_0 = -\Delta_v + \frac{1}{4}|v|^2 - \frac{n}{2} + v \cdot \nabla_x, \quad (x, v) \in \mathbb{R}^{2n}.$$

 P_0 defined on $\mathcal{C}_0^{\infty}(\mathbb{R}^{2n})$ is essentially maximal accretive (see Proposition 5.5, [9]). We still denote by P_0 its closed extension with maximal domain $D(P_0) = \{f \in L^2(\mathbb{R}^{2n}); P_0 f \in L^2(\mathbb{R}^{2n})\}$. One has: $\sigma(P_0) = [0, \infty[$. The following subellitptic estimate holds true for P_0 ([9]):

$$\|\Delta_v f\| + \|v^2 f\| + \|\langle D_x \rangle^{\frac{2}{3}} f\| \le C(\|P_0 f\| + \|f\|) \quad f \in D(P_0).$$
(1.3)

Date: June 11, 2024.

²⁰⁰⁰ Mathematics Subject Classification. 35J10, 35P15, 47A55.

Key words and phrases. Limiting absorption principle, eigenfunction decay, Kramers-Fokker-Planck operator, kinetic equation, embedded eigenvalues.

For $r \geq 0, s \in \mathbb{R}$, we introduce accordingly the weighted Sobolev space:

$$\mathcal{H}^{r,s} = \{ u \in \mathcal{S}'(\mathbb{R}^{2n}); (1 + \langle D_v \rangle^2 + |v|^2 + \langle D_x \rangle^{\frac{2}{3}})^{\frac{r}{2}} \langle x \rangle^s u \in L^2 \}.$$

For r < 0 and $s \in \mathbb{R}$, $\mathcal{H}^{r,s}$ is taken to be the dual space of $H^{-r,-s}$. Let $\mathcal{L}(r,s;r',s')$ be the space of bounded linear operators from $\mathcal{H}^{r,s}$ to $\mathcal{H}^{r',s'}$. To simplify notation we denote $\mathcal{H}^s = \mathcal{H}^{0,s}$, $\mathcal{H} = \mathcal{H}^0$ and $\mathcal{L}(s,s') = \mathcal{L}(0,s;0,s')$.

We write the full KFP operator P as

$$P = P_0 + W$$
 with $W = -\nabla_x V(x) \cdot \nabla_v$.

Under the condition (1.2) with $\rho > -1$, $\nabla_x V(x)$ tends to 0 as $|x| \to \infty$. (1.3) implies that W is relatively compact with respect to P_0 . Therefore P defined on $D(P) = D(P_0)$ is closed and maximally accretive. Let $\sigma(P)$ (resp., $\sigma_d(P)$, $\sigma_p(P)$) denote the spectrum (resp., the discrete spectrum, the point spectrum) of P. $\sigma_d(P)$ is the set of isolate eigenvalues with finite (algebraic) multiplicity. Let $\sigma_{\text{ess}}(P) = \sigma(P) \setminus \sigma_d(P)$ be the essential spectrum of P. When $\rho > -1$, zero may be an embedded eigenvalue and under some additional conditions, the return to equilibrium is proven in [14] with subexpoential convergence rate.

The general theory on relatively compact perturbation ([5]) for closed non-selfadjoint operators affirms that

$$\sigma_{\rm ess}(P) = \sigma(P_0) = [0, \infty[. \tag{1.4})$$

and $\sigma(P) \setminus [0, \infty] = \sigma_d(P)$ and that the discrete spectrum $\sigma_d(P)$ may accumulate towards any point in $[0, \infty]$. One of the goals of this work is to improve this statement.

The set of thresholds, \mathcal{T} , of the KFP operators is composed of eigenvalues of the harmonic oscillator $-\Delta_v + \frac{1}{4}|v|^2 - \frac{n}{2}$ which are non-negative integers:

$$\mathcal{T} = \mathbb{N}.$$

Motivated by spectral analysis of non-self-adjoint Schrödinger operators, we introduce real (non-threshold) resonances of the KFP operator using the boundary values of the free resolvent $R_0(z) = (P_0 - z)^{-1}$

$$R_0(\lambda \pm i0) = \lim_{z \to \lambda, \pm \Im z > 0} R_0(z), \quad \text{in} \quad \mathcal{L}(s, -s)$$

for $s > \frac{1}{2}$. (See Proposition 2.3 below.) The notion of threshold resonance is dimension-dependent and requires faster decay of potentials. For the KFP operator with a fastly decaying potential, see [17, 24] when n = 1 and 3.

Definition 1.1. Assume condition (1.2) with $\rho > 0$. Let $1/2 < s < (1 + \rho)/2$. We define the set \mathcal{E}_+ (resp;, \mathcal{E}_-) of outgoing (resp., incoming) resonances of P by

$$\mathcal{E}_{\pm} = \{\lambda \in]0, \infty[\backslash \mathcal{T}; \exists f \in \mathcal{H}^{-s} \setminus \mathcal{H}, (1 + R_0(\lambda \pm i0)W)f = 0\}$$

For $\lambda \in \mathcal{E}_+$, a solution to the equation $(1 + R_0(\lambda + i0)W)f = 0$ with $f \in \mathcal{H}^{-s} \setminus \mathcal{H}$ is called an outgoing resonant state associated with λ . Incoming resonant states are defined similarly. $\mathcal{E}_+ \cup \mathcal{E}_-$ is the set of non-threshold resonances of P.

The restriction on s in the interval $]\frac{1}{2}, \frac{1+\rho}{2}[$ is required by the use of Fredholm theory for compact operators in \mathcal{H}^{-s} . The above definition seems to be s-dependent, but using the properties of $R_0(\lambda \pm i0)$, one sees that if $f \in \mathcal{H}^{-s}$ is a solution to the equation $(1 + R_0(\lambda \pm i0)W)f = 0$ for some $s \in]\frac{1}{2}, \frac{1+\rho}{2}[$, then $f \in \bigcap_{r>\frac{1}{2}} \mathcal{H}^{-r}$. Therefore, real resonances and resonant states given in Definition 1.1 are s-independent. For real resonances of non-self-adjoint Schrödinger operators and their role in spectral and scattering theories, the interested reader can see [1, 6, 7, 12, 20, 21, 22, 24]. For dissipative Schrödinger operators, outgoing resonances are absent, but for every positive number $\lambda > 0$, one can construct a smooth compactly-supported dissipative potential such that λ is an incoming resonance ([22]). For this reason, we do not expect $\mathcal{E}_+ = \mathcal{E}_-$ for the KFP operator in general.

Denote $R(z) = (P - z)^{-1}$ for $z \notin \sigma(P)$. The main result of this work is the limiting absorption principle for the KFP operator with a long-range potential.

Theorem 1.2. Assume condition 1.2 with $\rho > 0$. Then

(a). The sets of accumulation points of $\sigma_p(P)$ and \mathcal{E}_{\pm} are included in \mathcal{T} .

(b). Let s > 1/2. For $\lambda \in \mathbb{R}_+ \setminus (\sigma_p(P) \cup \mathcal{T} \cup \mathcal{E}_{\pm})$, the boundary values of the resolvent

$$R(\lambda \pm i0) = \lim_{\pm \Im z > 0, z \to \lambda} R(z)$$
(1.5)

exist in $\mathcal{L}(s, -s)$ and are continuous in $\lambda \in \mathbb{R}_+ \setminus (\sigma_p(P) \cup \mathcal{T} \cup \mathcal{E}_{\pm})$.

We also study decay properties of the eigenfunctions of P.

Theorem 1.3. Let $\lambda \in \sigma_p(P)$ and $u \in D(P)$ such that $Pu = \lambda u$.

(a). If $\lambda \in \sigma_d(P)$ and (1.2) is satisfied with $\rho > -1$, then there exists some constant c > 0 such that $e^{c(\langle x \rangle + |v|^2)} u \in \mathcal{H}$.

(b). If $\lambda \in \mathbb{R}_+ \setminus \mathcal{T}$ and (1.2) is satisfied with $\rho > 0$, then u decays polynomially: for any $s \ge 0$, $(\langle x \rangle + \langle v \rangle)^s u \in \mathcal{H}$.

Remark 1.4. Most of works in the literature on spectral analysis of the KFP operator concern the case of compact resolvent. In this connection, let us mention that Helffer-Nier [9] conjectured that the KFP operator is of compact resolvent if and only if the associated Witten Laplacian is of compact resolvent. See [4, 15, 16] for some partial results. Under condition (1.2) with $\rho > -1$, the resolvent of P is never compact. To prove Theorem 1.2 and 1.3, we use the scattering approach initiated in [23], considering the potential term W as perturbation of the free KFP operator P₀. See also [17, 25]. In [23], the resolvent of the KFP operator is studied near the first threshold 0 for short-range potentials ($\rho > 1$) and n = 3. It is proven that under these conditions, 0 is not an accumulation point of eigenvalues.

It remains an open question to study whether or not eigenvalues may accumulate towards a nonzero threshold for short-range potentials in dimension three.

The remaining part of this paper is organized as follows. In Section 2 we recall from [23] some results on the free KFP operator and improve some of them for later use. In Section 3 we analyze non-threshold resonances and embedded eigenvalues of P and prove that these sets do not have an accumulation point outside the thresholds \mathcal{T} . Once these preparations done, Theorem 1.2 can be easily inferred (see Section 4). In Section 5 we discuss decay properties of eigenfunctions. Appendix A. contains the proof for a technical lemma.

2. LAP FOR THE FREE KFP OPERATOR

We first recall from [23] some results needed in this work. Let

$$P_0 = -\Delta_v + \frac{1}{4}|v|^2 - \frac{n}{2} + v \cdot \nabla_x, \quad (x, y) \in \mathbb{R}^{2n},$$

and $R_0(z) = (P_0 - z)^{-1}$. By the partial Fourier transform $\mathcal{F}_{x \to \xi}$ in x-variables, the free KFP P_0 is a direct integral of the family $\{\widehat{P}_0(\xi); \xi \in \mathbb{R}^n\}$, where

$$\widehat{P}_0(\xi) = -\Delta_v + \frac{v^2}{4} - \frac{n}{2} + iv \cdot \xi.$$
(2.1)

The operator $\widehat{P}_0(\xi)$ is given by

$$\widehat{P}_0(\xi) = \mathcal{F}_{x \to \xi} P_0 \mathcal{F}_{\xi \to x}^{-1} = -\Delta_v + \frac{1}{4} \sum_{j=1}^n (v_j + 2i\xi_j)^2 - \frac{n}{2} + |\xi|^2.$$

 $\{\widehat{P}_0(\xi), \xi \in \mathbb{R}^n\}$ is a holomorphic family of type (A) in sense of Kato with constant domain $D = D(-\Delta_v + \frac{v^2}{4})$ in $L^2(\mathbb{R}^n_v)$. Let $F_j(s) = (-1)^j e^{\frac{s^2}{2}} \frac{d^j}{ds^j} e^{-\frac{s^2}{2}}, j \in \mathbb{N}$, be Hermite polynomials and

$$\varphi_j(s) = (j!\sqrt{2\pi})^{-\frac{1}{2}}e^{-\frac{s^2}{4}}F_j(s)$$

the normalized Hermite functions. For $\xi \in \mathbb{R}^n$ and $\alpha = (\alpha_1, \alpha_2, \cdots, \alpha_n) \in \mathbb{N}^n$, define

$$\psi_{\alpha}(v) = \prod_{j=1}^{n} \varphi_{\alpha_j}(v_j) \text{ and } \psi_{\alpha}^{\xi}(v) = \psi_{\alpha}(v+2i\xi).$$
(2.2)

One can check ([23]) that each eigenvalue $E_l(\xi) = l + |\xi|^2$ of $\widehat{P}_0(\xi)$ is semi-simple and the associated Riesz projection is given by

$$\Pi_l^{\xi} \phi = \sum_{\alpha, |\alpha|=l} \langle \psi_{\alpha}^{-\xi}, \phi \rangle \psi_{\alpha}^{\xi}, \quad \phi \in L^2.$$
(2.3)

Proposition 2.1 ([23]). Let $l \in \mathbb{N}$ and l < a < l + 1 be fixed. Take $\chi \ge 0$ and $\chi \in C_0^{\infty}(\mathbb{R}^n_{\xi})$ with supp $\chi \subset \{\xi, |\xi|^2 \le a + 4\}$, $\chi(\xi) = 1$ when $|\xi|^2 \le a + 3$ and $0 \le \chi(\xi) \le 1$. Then one has

$$\widehat{R}_0(z,\xi) = \sum_{k=0}^l \chi(\xi) \frac{\Pi_k^{\xi}}{\xi^2 + k - z} + r_l(z,\xi), \qquad (2.4)$$

for any $\xi \in \mathbb{R}^n$ and $z \in \mathbb{C}$ with $\Re z < a$ and $\Im z \neq 0$. Here $r_l(z,\xi)$ is holomorphic in z with $\Re z < a$ verifying the estimate

$$\sup_{\Re z < a, \xi \in \mathbb{R}^n} \|r_l(z,\xi)\|_{\mathcal{L}(L^2(\mathbb{R}^n_v))} < \infty.$$
(2.5)

Here and in the following, for two normed spaces \mathcal{F} and \mathcal{G} , $\mathcal{L}(\mathcal{H}, \mathcal{G})$ denotes the space of bounded linear operators from \mathcal{H} to \mathcal{G} and $\mathcal{L}(\mathcal{H}) = \mathcal{L}(\mathcal{H}, \mathcal{H})$. (2.4) is a kind of spectral decomposition for $\widehat{R}_0(z,\xi)$. The proof of the assertions on $r_l(z,\xi)$ is technical and uses time-dependent method (cf. Proposition 2.7 and Appendix A. in [23]).

For a temperate symbol $a(x, v; \xi, \eta)$ ([10]), denote by $a^w(x, v, D_x, D_v)$ the associated Weyl pseudo-differential operator defined by

$$a^{w}(x,v,D_{x},D_{v})u(x,v)$$

$$= \frac{1}{(2\pi)^{2n}} \int \int e^{i(x-x')\cdot\xi+i(v-v')\cdot\eta} a(\frac{x+x'}{2},\frac{v+v'}{2},\xi,\eta)u(x',v')dx'dv'd\xi d\eta$$
(2.6)

for $u \in \mathcal{S}(\mathbb{R}^{2n}_{x,v})$.

Remark that $\chi(D_x)\Pi_k^{D_x}$ is a Weyl pseudodifferential operator with a nice symbol b_k independent of x:

$$\chi(D_x)\Pi_k^{D_x} = b_k^w(v, D_x, D_v) \tag{2.7}$$

where $b_k(v,\xi,\eta)$ is given by

$$b_k(v,\xi,\eta) = \int_{\mathbb{R}^n} e^{-iv'\cdot\eta/2} \left(\sum_{|\alpha|=k} \chi(\xi)\psi_\alpha(v+v'+2i\xi)\psi_\alpha(v-v'+2i\xi) \right) dv'.$$
(2.8)

In particular,

$$b_0(v,\xi,\eta) = 2^{\frac{n}{2}}\chi(\xi)e^{-v^2 - \eta^2 + 2iv\cdot\xi + 2\xi^2}.$$
(2.9)

These pseudodifferential operators belong to $\mathcal{L}(r, s; r', s)$ for any $r, r', s \in \mathbb{R}$ (Ch. 18, [10]). Set

$$r_l(z) = \mathcal{F}_{\xi \to x}^{-1} r_l(z,\xi) \mathcal{F}_{x \to \xi}$$

Proposition 2.1 implies the following representation formula for the free resolvent

$$R_0(z) = \sum_{k=0}^{l} b_k^w(v, D_x, D_v) (-\Delta_x + k - z)^{-1} + r_l(z), \qquad (2.10)$$

for $z \in \mathbb{C}$ with $\Re z < a$ and $\Im z \neq 0$ and that $r_l(z)$ is bounded on L^2 and holomorphic in z with $\Re z < a, a \in]l, l+1[$.

Lemma 2.2. Let $z \in \mathbb{C} \setminus \mathbb{R}_+$. For any $m \in \mathbb{N}$, $s \in \mathbb{R}$ and $r \in [0, 2m]$, one has $R_0(z)^m \in \mathcal{L}(-r, s; 2m - r, s)$.

Proof. Let m = 1. Consider first the case r = 0 and $s \in \mathbb{N}$. For s = 1, one has $\langle x \rangle R_0(z) \langle x \rangle^{-1} = R_0(z) \left(1 + [-v \cdot \nabla_x, \langle x \rangle] R_0(z) \langle x \rangle^{-1} \right).$

The subelliptic estimate (1.3) shows that $R_0(z) \in \mathcal{L}(0,2)$ and $[-v \cdot \nabla_x, \langle x \rangle] R_0(z) \in \mathcal{L}(0,0)$. Therefore, $\langle x \rangle R_0(z) \langle x \rangle^{-1} \in \mathcal{L}(0,2)$. Similarly, using the formula

$$\langle x \rangle^k R_0(z) \langle x \rangle^{-k} = \langle x \rangle^{k-1} R_0(z) \left(\langle x \rangle^{-k+1} + \left[-v \cdot \nabla_x, \langle x \rangle \right] R_0(z) \langle x \rangle^{-k} \right),$$

we can show by an induction that $\langle x \rangle^k R_0(z) \langle x \rangle^{-k} \in \mathcal{L}(0,2)$ for all $k \in \mathbb{N}$. This proves $R_0(z) \in \mathcal{L}(0,k;2,k)$. By an argument of duality, one has $R_0(z)^* \in \mathcal{L}(-2,-k;0,-k)$ for $k \in \mathbb{N}$. Note that $P_0^* = JP_0J$ where Jf(x,v) = f(x,-v) and J conserves any $\mathcal{H}^{r,s}$ norm. From $R_0(\overline{z})^* = JR_0(z)J$, we deduce that $R_0(z) \in \mathcal{L}(-2,-k;0;-k)$ for $k \in \mathbb{N}$. A complex interpolation between $(r_1,s_1) = (-2,-k)$ and $(r_2,s_2) = (2,k)$ gives that $R_0(z) \in \mathcal{L}(-r,s;2-r,s)$ for $r \in [0,2]$ and $|s| \leq k$, for any $k \in \mathbb{N}$. The lemma is proven for m = 1.

When $m \geq 2$, one can use the result obtained for $R_0(z)$ and the above commutator method to show that $R_0(z)^m \in \mathcal{L}(0,s;2m,s)$ for any $s \in \mathbb{R}$. The details are omitted here. The arguments of duality and complex interpolation used above allow us to conclude that $R_0(z)^m \in \mathcal{L}(-r,s;2m-r,s)$ for $s \in \mathbb{R}$ and $r \in [0,2m]$. \Box

Proposition 2.3. Let $n \ge 1$. For any $s > \frac{1}{2}$, the boundary values of the resolvent $R_0(\lambda \pm i\epsilon) = \lim_{\pm \Im z > 0, z \to \lambda} R_0(z)$

exist in $\mathcal{L}(0, s; 2, -s)$ for $\lambda \in \mathbb{R}_+ \setminus \mathcal{T}$ and is continuous in λ . More generally, let $k \in \mathbb{N}$ and $s > k + \frac{1}{2}$ and $r, r' \geq 0$ with $r + r' \leq 2(k+1)$, $R_0(\lambda \pm i0)$ is of class C^k in $\mathcal{L}(-r, s; r', -s)$ for $\lambda \notin \mathcal{T}$.

Proof. When k = 0, r = 0 and r' = 2, the result has been proven in [23] by using (2.10) and the known results for $(-\Delta_x - (\mu \pm i0))^{-1}$ in weighted L^2 -spaces(cf. [2]). In the general case, the existence and the continuity of $\frac{d^k}{d\lambda^k}R_0(\lambda \pm i0)$ in $\mathcal{L}(0,s;0,-s)$, $s > k + \frac{1}{2}$, can be deduced by the same argument. To see its continuity in weighted Sobolev spaces, we use Lemma 2.2 to conclude $R_0(-1)^{k+1} \in$ $\mathcal{L}(0,s;2k+2,-s)$ and the equation

$$R_0(\lambda \pm i0)^{k+1} = R_0(-1)^{k+1} \left(1 + (\lambda + 1)R_0(\lambda \pm i0)\right)^{k+1}$$

to show that $\frac{d^k}{d\lambda^k}R_0(\lambda \pm i0)$ is continuous in λ in topology of $\mathcal{L}(0,s;2k+2,-s)$, if $s > k + \frac{1}{2}$. The general statement of Proposition 2.3 follows from the arguments of duality and complex interpolation already used in Lemma 2.2.

3. Real resonances and embedded eigenvalues

Let us now consider the full KFP operator P with a long-range potential satisfying the condition 1.2 with $\rho > 0$.

Lemma 3.1. Let $\lambda_0 \in \mathbb{R}_+ \setminus \mathcal{T}$ be an embedded eigenvalue of P and u an associated eigenfunction: $Pu = \lambda_0 u$. Then one has

$$u = -R_0(\lambda_0 \pm i0)Wu. \tag{3.1}$$

Proof. Proposition 2.3 shows that $R_0(\lambda_0 \pm i0)Wu \in \mathcal{H}^{1,-s}$, $1/2 < s < (1+\rho)/2$. To show (3.1) for sign +, we set $z = \lambda_0 + i\epsilon$ with $\epsilon > 0$ and write u as

$$u = -R_0(z)Wu - i\epsilon R_0(z)u.$$

To see that $\lim_{\epsilon \to 0_+} \epsilon R_0(z)u = 0$ in \mathcal{H} , we use (2.10) with $l > \lambda_0$. Clearly

$$\lim_{\epsilon \to 0_+} \epsilon r_l(z) u = 0,$$

because $r_l(w)$ is holomorphic in w with $\Re w < l + a$ for some a > 0. The symbol of $\epsilon(-\Delta + k - z)^{-1}$ is equal to $\epsilon(|\xi|^2 + k - \lambda_0 - i\epsilon)^{-1}$ which converges pointwise to 0 and is uniformly bounded by 1. The Dominated Convergence Theorem shows that

$$\lim_{\epsilon \to 0_+} \sum_{k=0}^{\iota} \epsilon b_k^w(v, D_x, D_v) (-\Delta_x + k - z)^{-1} u = 0$$

in \mathcal{H} . This proves that $\lim_{\epsilon \to 0_+} \epsilon R_0(z)u = 0$ in \mathcal{H} , hence $u = -R_0(\lambda_0 + i0)Wu$ in \mathcal{H} . Similarly one can show that $u = -R_0(\lambda_0 - i0)Wu$.

Lemma 3.2. Let $\lambda_0 \in \mathbb{R}_+ \setminus \mathcal{T}$ and $1/2 < s < 1 + \rho/2$. If $u \in \mathcal{H}^{-s} \setminus \{0\}$ satisfies both equations

$$u = -R_0(\lambda_0 \pm i0)Wu,$$

then $u \in \mathcal{H}$ and $\lambda_0 \in \sigma_p(P)$.

Proof. Making use of (2.10) with $l > \lambda_0$, one has

$$u = -\sum_{k=0}^{l} b_k^w(v, D_x, D_v) (-\Delta_x + k - (\lambda_0 \pm i0))^{-1} W u - r_l(\lambda_0) W u.$$
(3.2)

Since $Wu \in \mathcal{H}^{-1,1+\rho-s}$ and $1+\rho-s>s$, $r_l(\lambda_0)Wu \in \mathcal{H}$. Let $d = d(\lambda_0, \mathcal{T})$. Take $\chi_1 \in C_0^{\infty}(\mathbb{R}^n_{\mathcal{E}} \setminus 0)$ such that

$$\chi_1(\xi) = 1$$
 for $\frac{d}{2} \le |\xi|^2 \le l+1$.

Let $\chi_2(\xi) = 1 - \chi_1(\xi)$ and decompose

$$u = u_1 + u_2$$
 with $u_j = \chi_j(D_x)u$.

From (3.2) and the fact that $\chi_2(\xi)(|\xi|^2 + k - (\lambda_0 \pm i0))^{-1}$, $k = 0, \dots, l$, are bounded for $\xi \in \mathbb{R}^n$, we deduce that $u_2 \in \mathcal{H}$. To show that $u_1 \in \mathcal{H}$, we decompose $\chi_1(D_x)$ as

$$\chi_1(D_x) = B_+ + B_-$$

where $B_{\pm} = b_{\pm}^w(x, D_x)$ are Weyl pseudo-differential operators with bounded symbols $b_{\pm}(x, \xi)$ such that

$$\operatorname{supp} b_{\pm} \subset \{(x,\xi); |x| \le R\} \cup \{(x,\xi); \pm \widehat{x} \cdot \widehat{\xi} > (1-\delta_{\pm})\}$$

for some R > 1 and $0 < \delta_{\pm} < 1$. Here $\hat{x} = \frac{x}{|x|}$, $\hat{\xi} = \frac{\xi}{\xi}$ for $x, \xi \neq 0$. One has the microlocal resolvent estimates for $-\Delta_x$ in $L^2(\mathbb{R}^n_x)$ ([11]): for $\mu > 0$ and s > 1/2,

$$\|\langle x \rangle^{s-1} B_{\mp}(-\Delta_x - (\mu \pm i0))^{-1} \langle x \rangle^{-s} \| < \infty.$$
 (3.3)

It follows that

$$B_{\pm}b_k^w(v, D_x, D_v)(-\Delta_x + k - (\lambda_0 \pm i0))^{-1} \in \mathcal{L}(r, s; r', 1 - s)$$
(3.4)

for all $r, r' \in \mathbb{R}$ and s > 1/2. Write u_1 as

$$u_{1} = B_{-}u + B_{+}u$$

$$= -B_{-}\left(\sum_{k=0}^{l} b_{k}^{w}(v, D_{x}, D_{v})(-\Delta_{x} + k - (\lambda_{0} + i0))^{-1}Wu + r_{l}(\lambda_{0})Wu\right)(3.5)$$

$$-B_{+}\left(\sum_{k=0}^{l} b_{k}^{w}(v, D_{x}, D_{v})(-\Delta_{x} + k - (\lambda_{0} - i0))^{-1}Wu + r_{l}(\lambda_{0})Wu\right).$$

Applying (3.3) with $r = 1 + \rho - s$ and $\mu = \lambda - k$, $k = 0, \dots, l$, one obtains that $u_1 \in \mathcal{H}^{-(s-\rho)}$. If $s-\rho \leq 0$, then $u = u_1+u_2 \in \mathcal{H}$. The proof is finished. If $s-\rho > 0$, then $u \in \mathcal{H}^{-(s-\rho)}$. We apply (3.3) once more with $r = 1 + \rho - (s-\rho)$ and deduce from (3.5) that $u_1 \in \mathcal{H}^{-(s-2\rho)}$ which implies $u \in \mathcal{H}^{-\max\{0,s-2\rho\}}$. Repeating these arguments for at most a finite number of times, we obtain $u \in \mathcal{H}$.

Remark 3.3. We can also give large-x expansions of resonant states. If u is an outgoing resonant state associated with a resonance $\lambda \notin \mathcal{T}$, then the argument used above shows that

$$u - \sum_{k=0}^{l_0} b_k^w(v, D_x, D_v) (-\Delta_x + k - (\lambda + i0))^{-1} W u \in \mathcal{H},$$

where l_0 is the integer such that $l_0 < \lambda < l_0 + 1$. We can study the large |x| behavior of u, by making use of the distributional kernel of $(-\Delta_x - z)^{-1}$ given by

$$K(x,y;z) = \frac{i}{4} \left(\frac{\sqrt{z}}{2|x-y|} \right)^{\frac{n}{2}-1} H_{n/2-1}^{(1)}(\sqrt{z}|x-y|), \quad x,y \in \mathbb{R}^n,$$

where $\Im\sqrt{z} \geq 0$ and $H_{\nu}^{(1)}$ is the Bessel function of the third kind ([18]). In particular, for n = 3, one has simply

$$K(x,y;z) = \frac{e^{i\sqrt{z}|x-y|}}{4\pi|x-y|}.$$

For $\rho > 1$ and n = 3, one can check that

$$(-\Delta_x + k - (\lambda + i0))^{-1}Wu - \phi_k \in \mathcal{H},$$

where

$$\psi_k(x,v) = \frac{e^{i\sqrt{\lambda-k}|x|}}{4\pi|x|} \int_{\mathbb{R}^3_y} (Wu)(y,v)dy.$$
 (3.6)

This proves

$$u = \sum_{k=0}^{l_0} b_k^w(v, D_x, D_v)\psi_k + h, \qquad (3.7)$$

with $h \in \mathcal{H}$. In particular, if

$$\int_{\mathbb{R}^3} \nabla V(y) u(y, v) dy = 0, \quad a.e. \text{ in } v, \qquad (3.8)$$

then u is an eigenfunction and λ an embedded eigenvalue of P. Similar results are also true for incoming resonant states.

Let $K \subset \overline{\mathbb{C}}_{\pm}$ and $1/2 < s < (1+\rho)/2$. Let E_K be the subspace of \mathcal{H}^{-s} spanned by eigenfunctions and outgoing or incoming resonant states (according to sign + or -) associated with eigenvalues and resonances in K.

Proposition 3.4. If $K \subset \overline{\mathbb{C}}_{\pm}$ is compact and $K \cap \mathcal{T} = \emptyset$, then E_K is of finite dimension. In particular, the sets of accumulation points of $\sigma_p(P)$ and \mathcal{E}_{\pm} are included in \mathcal{T} .

Proof. To fix the idea, let $K \subset \overline{\mathbb{C}}_+$ and 1/2 < s' < s. Since $1 + \rho - s > 1/2$, Proposition 2.3 shows that

$$||R_0(z)W||_{\mathcal{L}(0,-s;1,-s')} \le C_K$$

uniformly for $z \in K$. Here R(z) is understood as $R(\lambda + i0)$ if $z = \lambda \in K \cap \mathbb{R}$. Let B be the unit ball in E_K and $u \in B$. Using the equation $u = -R_0(z_0)Wu$ for some $z_0 \in K$, one has

$$\|u\|_{\mathcal{H}^{1,-s'}} \le C_K \|u\|_{\mathcal{H}^{-s}} = C_K,$$

This means that B is bounded in $\mathcal{H}^{1,-s'}$. Since the canonical injection $\mathcal{H}^{1,-s'} \hookrightarrow \mathcal{H}^{-s}$ is compact for 1/2 < s' < s, B is compact in \mathcal{H}^{-s} . This proves that dim $E_K < \infty$ by F. Riesz's criteria on the finiteness of dimension for normed vector spaces. \Box

Proposition 3.4 implies that the geometric multiplicities of embedded eigenvalues of P are finite, but we have no information on their algebraic multiplicities.

4. LAP for the perturbed KFP operator

Proof of Theorem 1.2. Part (a) of Theorem 1.2 is a consequence of Proposition 3.4. For Part (b), we only study $R(\lambda + i0)$. Let $1/2 < s < (1 + \rho)/2$. For $\lambda \in \mathbb{R}_+ \setminus (\mathcal{T} \cup \sigma_p(P) \cup \mathcal{E}_+), 1 + R_0(\lambda + i0)W$ is injective in \mathcal{H}^{-s} and $R_0(\lambda)W$ is a compact operator in \mathcal{H}^{-s} . Therefore $1 + R_0(\lambda + i0)W$ is invertible in $\mathcal{L}(-s, -s)$. By the continuity of $R_0(\lambda + i0)$ (Proposition 2.3), $(1 + R_0(\mu + i0)W)^{-1}$ also exists

in $\mathcal{L}(-s, -s)$ for μ near λ and is continuous in μ . Taking the limit $z \to \mu + i0$ in the equation

$$R(z) = (1 + R_0(z)W)^{-1}R_0(z)$$

we see that for μ near λ , $R(\mu + i0) = \lim_{z \to \mu + i0} R(z)$ exists in $\mathcal{L}(s, -s)$ and is given by

$$R(\mu + i0) = (1 + R_0(\mu + i0)W)^{-1}R_0(\mu + i0).$$
(4.1)

As $\mathcal{L}(s, -s)$ -valued function, $R(\mu + i0)$ is continuous in μ , because $R_0(\mu + i0) \in \mathcal{L}(s, -s)$ is continuous in μ and the same is true for $((1 + R_0(\mu + i0)W)^{-1})$ as $\mathcal{L}(-s, -s)$ -valued function. This proves Part (b) for $R(\lambda + i0)$. The proof for $R(\lambda - i0)$ is the same.

Remark 4.1. If (1.2) is satisfied with $\rho > 2k$, $k \in \mathbb{N}^*$, then one can use Proposition 2.3 to show that $R(\lambda \pm i0)$ is of class C^k in $\mathcal{L}(0,s;2k+2,-s)$, $s > k + \frac{1}{2}$, for $\lambda \in \mathbb{R} \setminus (\mathcal{T} \cup \sigma_p(P) \cup \mathcal{E}_{\pm})$.

5. Decay properties of eigenfunctions

We prove Theorem 1.3 by studying separately discrete and embedded eigenvalues. The following result is a cosmetic improvement of Theorem 1.3 (a).

Theorem 5.1. Suppose that condition (1.2) is satisfied for some $\rho > -1$. Then there exist some positive constant c_0 and positive-valued function $\tau(\zeta)$ (defined by (5.2) below) for $\zeta \in \mathbb{C} \setminus \mathbb{R}$ such that if u is an eigenfunction of P associated with an eigenvalue $z \in \sigma_d(P)$, then $e^{c_0 \tau(z)(\langle x \rangle + |v|^2)} u \in \mathcal{H}$.

Proof. Let $u \in D(P)$ and $z \in \sigma_d(P)$ such that that Pu = zu. Then $z \notin \mathbb{R}_+$ and one has

$$u = -R_0(z)Wu. (5.1)$$

Let $\chi(s)$ be a smooth real-valued function on \mathbb{R} such that

$$\chi(s) = \begin{cases} s, & \text{if } s \in [0, 1], \\ \frac{3}{2}, & \text{if } s \ge 2. \end{cases}$$

Let $\chi_r(s) = r\chi(\frac{s}{r})$ for $r \ge 1$. Then $\chi_r(s) \to s$ pointwise as $r \to \infty$ and $|\chi'_r(s)| \le C$, uniformly for $r \ge 1$ and $s \ge 0$. Denote

$$\phi_r(x,v) = \chi_r(\langle x \rangle) + \chi_r(|v|^2).$$

Set $P_{0,r} = e^{a\phi_r} P_0 e^{-a\phi_r}$ with a > 0 which is to be chosen later. One can compute $P_{0,r} = P_0 + Q_r$,

where

$$Q_r = a\left(4\chi_r'(|v|^2)v \cdot \nabla_v + \Delta_v(\chi_r(|v|^2)) - a|\nabla_v\chi_r(|v|^2)|^2 - \chi_r'(\langle x \rangle)v \cdot \frac{x}{\langle x \rangle}\right)$$

By the choice of χ_r , there exists some constant C such that

$$||Q_r R_0(z)|| \le Ca(||v \cdot \nabla_v R_0(z)|| + ||v \cdot \frac{x}{\langle x \rangle} R_0(z)||) + (1+a)||R_0(z)||).$$

Define $\tau(\zeta)$ by

$$\tau(\zeta) = \min\{(\|v \cdot \nabla_v R_0(\zeta)\| + \|v \cdot \frac{x}{\langle x \rangle} R_0(\zeta)\| + \|R_0(z)\|)^{-1}, \|R_0(\zeta)\|^{-\frac{1}{2}}\}$$
(5.2)

for $\zeta \in \mathbb{C} \setminus \mathbb{R}_+$. For $a = c_0 \tau(z)$, one has

$$||Q_r R_0(z)|| \le C(c_0 + c_0^2) < 1$$

if $c_0 > 0$ is chosen appropriately small. With such a choice of a, the resolvent $R_{0,r}(z) = (P_{0,r} - z)^{-1}$ exists for all $r \ge 1$ and is given by

$$R_{0,r}(z) = R_0(z)(1 + Q_r R_0(z))^{-1}.$$

It follows that there exists some constant C depending on z such that

$$||R_{0,r}(z)|| \le C \tag{5.3}$$

uniformly for $r \ge 1$. From the subellipticity of P_0 , one deduces that $R_{0,r}(z) \in \mathcal{L}(-s,0;2-s,0), s \in [0,2]$, is uniformly bounded for $r \ge 1$. In particular, $\|R_{0,r}(z)(\langle v \rangle + \langle D_v \rangle)\| \le C$ uniformly for $r \ge 1$.

Let $u_r = e^{c_0 \tau(z)\phi_r} u$. Equation (5.1) gives

$$u_r = -R_{0,r}(z)(\nabla_v \chi_r(|v|^2) - \nabla_v) \cdot \nabla_x V u_r$$
(5.4)

Since $\nabla_x V = O(\langle x \rangle^{-1-\rho})$ and $1+\rho > 0$, there exists some constant C independent of R_0 and r such that for any $R_0 > 1$,

$$\|\nabla_x V u_r\| \le C_{R_0} \|u\| + C \langle R_0 \rangle^{-1-\rho} \|u_r\|.$$

Seeing that there exists some constant M > 0 such that

$$||R_{0,r}(z)(\nabla_v \chi_r(|v|^2) - \nabla_v)|| \le C' ||R_{0,r}(z)||_{\mathcal{L}(-1,0;0,0)} \le M,$$

uniformly for $r \ge 1$, one has

$$\begin{aligned} \|u_r\| &\leq \|R_{0,r}(z)(\nabla_v \chi_r(|v|^2) - \nabla_v)\| \|\nabla_x V u_r\| \\ &\leq M(C_{R_0} \|u\| + C \langle R_0 \rangle^{-1-\rho} \|u_r\|), \quad \forall r \geq 1. \end{aligned}$$

Taking R_0 appropriately large so that $MC\langle R_0\rangle^{-1-\rho} < \frac{1}{2}$, one obtains

$$||u_r|| \le 2MC_{R_0}||u||, \quad r \ge 1.$$

Since $u_r(x,v) \to e^{c_0\tau(z)(\langle x \rangle + v^2)}u(x,v)$ pointwise as $r \to \infty$, Fatou's Lemma shows that $e^{c_0\tau(z)(\langle x \rangle + v^2)}u \in \mathcal{H}$.

Remark 5.2. Theorem 5.1 gives some quantitative information about decay properties of eigenfunctions. Comparing with the Green function of P_0 , we believe that the phase function $\langle x \rangle + |v|^2$ is the appropriate one to use in this context, but the numerical factor $c_0\tau(z)$ before it can very likely be improved. If $z = b + i\mu$ with b > 0 fixed and $\mu \in \mathbb{R}$, one has uppers bounds for $||v \cdot \nabla_v R_0(z)||$, $||v \cdot \frac{x}{\langle x \rangle} R_0(z)||$) and $||R_0(z)||$ for $|\mu| \neq 0$. However to calculate $\tau(z)$, one also needs to evaluate their lower bounds, which seems not to be an easy task.

Theorem 1.3 (b) is proven by applying the boost-up argument of Lemma 3.2 to

$$u = -R_0(\lambda \pm i0)Wu$$

to improve the decay of u. To do this, we need to show that the remainder $r_l(\lambda)$ given by (2.10) with $l > \lambda$ and $z = \lambda$ conserves polynomial decay. Since $\nabla_x V(x)$ does not operate in $\mathcal{F}^{r,s}$ for r large, we modify the definition of $\mathcal{F}^{r,s}$ and introduce $\mathcal{G}^{r,s}$ by

$$\mathcal{G}^{r,s} = \{ f \in \mathcal{S}'(\mathbb{R}^{2n}; \langle 1 - \Delta_v + v^2 \rangle^{\frac{r}{2}} \langle x \rangle^s f \in L^2 \}.$$

 $\mathcal{G}^{r,s}$ is equipped with its natural norm. Then, $\nabla_x V \in \mathcal{L}(\mathcal{G}^{r,s}; \mathcal{G}^{r,s+1+\rho})$ for all r, s. The proof of the following technical lemma is given in Appendix A.

Lemma 5.3. For any $r, s \in \mathbb{R}$, one has $r_l(\lambda) \in \mathcal{L}(\mathcal{G}^{r,s}; \mathcal{G}^{2+r,s})$.

Proof of Theorem 1.3 (b). As in the proof of Lemma 3.2, we apply (2.10) with $l > \lambda$ to $u = -R_0(\lambda \pm i0)Wu$ and write

$$u = -\sum_{k=0}^{l} b_{k}^{w}(v, D_{x}, D_{v})(-\Delta_{x} + k - (\lambda \pm i0))^{-1}Wu - r_{l}(\lambda)Wu$$

Since $u \in \mathcal{H}$ and $Wu \in \mathcal{G}^{-1,1+\rho}$, Lemma A.2 shows that

$$r_l(\lambda)Wu \in \mathcal{G}^{1,1+\rho}$$

Introduce the microlocal partition of unity $\chi_2(D_x) + B_+ + B_- = 1$ as constructed in Lemma 3.2. Then an argument of support in ξ gives

$$\chi_2(D_x)b_k^w(v, D_x, D_v)(-\Delta_x + k - (\lambda \pm i0))^{-1}Wu \in \mathcal{G}^{r, 1+\rho}$$

and (3.3) with $r = 1 + \rho_0$ shows that

$$B_{\pm}b_k^w(v, D_x, D_v)(-\Delta_x + k - (\lambda \pm i0))^{-1}Wu \in \mathcal{G}^{r,\rho}$$

for all $r \in \mathbb{R}$. Putting these pieces together, one obtains $u \in \mathcal{G}^{1,\rho}$. Now $Wu \in \mathcal{G}^{0,1+2\rho}$. Repeating the above arguments with this improved property on Wu, we obtain $u \in \mathcal{G}^{2,2\rho}$. By an induction on m, we can show that $u \in \mathcal{G}^{m,m\rho}$ for any $m \in \mathbb{N}$. Consequently, u decays polynomially.

Remark 5.4. For N-body self-adjoint Schrödinger operator H with long-range interactions, it is known ([3, 8]) that an eigenfunction associated with a nonthreshold eigenvalue λ of H decays like $O(e^{-(1-\epsilon)\sqrt{d(\lambda)}|x|})$ for any $\epsilon > 0$ and small, where $d(\lambda)$ is the distance between λ and the thresholds of H located on the right-hand side of λ . For the KFP operator P with a long-range potential, one may naturally ask the questions whether there exists an exponential decay of eigenfunctions associated with an embedded eigenvalue $\lambda \in \mathbb{R}_+ \setminus \mathcal{T}$ and whether their decay rate can be described in terms of the distance between λ and \mathcal{T} .

Appendix A. Proof of Lemma 5.3

We first give two results needed for the proof of Lemma 5.3. The first one concerns the evolution of elementary observables by the semigroup e^{-tP_0} .

Lemma A.1 (Lemma 2.12, [23]). Let $n \ge 1$. For $t \ge 0$ and $0 \le s \le t$, one has the following equalities as operators from $\mathcal{S}(\mathbb{R}^{2n}_{x,v})$ to $L^2(\mathbb{R}^{2n}_{x,v})$:

$$e^{-(t-s)P_{0}}v_{j}e^{-sP_{0}} = e^{-tP_{0}}(v_{j}\cosh s - 2\partial_{v_{j}}\sinh s + 2(\cosh s - 1)\partial_{x_{j}}) (A.1)$$

$$e^{-(t-s)P_{0}}\partial_{v_{j}}e^{-sP_{0}} = -\frac{1}{2}e^{-tP_{0}}((v_{j}\sinh s - 2\partial_{v_{j}}\cosh s + 2\partial_{x_{j}}\sinh s)) (A.2)$$

$$e^{-(t-s)P_{0}}x_{j}e^{-sP_{0}} = e^{-tP_{0}}(x_{j} + v_{j}\sinh s - 2(\cosh s - 1)\partial_{v_{j}}$$

$$+2(\sinh s - s)\partial_{x_{j}}) (A.3)$$

The second one is about the smoothing properties of some integrals of e^{-tP_0} .

Lemma A.2. Let $k \in \mathbb{N}$. If $\theta(t)$ is a smooth function with $\theta^{(j)}(0) = 0$ for $j = 0, \dots, k$, then for any fixed T > 3, the operator

$$(P_0+1)^{k+1} \int_0^T e^{-tP_0} \theta(t) \, dt \in \mathcal{L}(\mathcal{H}). \tag{A.4}$$

Proof. For an abstract C_0 -contraction semigroup generated by a maximally accretive operator, results similar to (A.4) hold true for k = 0, but in general they fail for $k \ge 1$. To prove (A.4) for all $k \in \mathbb{N}$, we use the following formula (cf. (A.10) in [23]):

$$\sum_{l=0}^{\infty} e^{-t(l+|\xi|^2)} \|\Pi_l^{\xi}\| \le \frac{e^{-|\xi|^2(t-2-\frac{4}{e^{t-1}})}}{(1-e^{-t})^n},\tag{A.5}$$

for t > 0 and $\xi \in \mathbb{R}$. Here Π_l^{ξ} is the Riesz projection of $\widehat{P}_0(\xi)$ with eigenvalue $l + |\xi|^2$. For any $k \in \mathbb{N}$, one has

$$\begin{split} \| (\widehat{P}_{0}(\xi) + 1)^{k} e^{-t\widehat{P}_{0}(\xi)} \| \\ &= \| \sum_{l=0}^{\infty} (l + |\xi|^{2})^{k} e^{-t(l+\xi^{2})} \Pi_{l}^{\xi} \| \\ &\leq \sum_{l=0}^{\infty} \| (l + |\xi|^{2})^{k} e^{-t(l+\xi^{2})} \Pi_{l}^{\xi} \| \\ &\leq C_{k,\delta} \sum_{l=0}^{\infty} e^{-(t-\delta)(l+|\xi|^{2})} \| \Pi_{l}^{\xi} \| \quad (0 < \delta < t) \\ &\leq C_{k} e^{-(t-T)|\xi|^{2}}, \end{split}$$

for $\xi \in \mathbb{R}^n$ and $t \ge T > 3$ (with $\delta > 0$ chosen appropriately). Consequently for any $k \in \mathbb{N}$, one has

$$\|(P_0+1)^k e^{-tP_0}\| \le C_k,\tag{A.6}$$

uniformly for $t \ge T > 3$. Lemma A.2 follows from integration by parts and the above estimate with t = T > 3.

Lemma A.2 shows that if $\theta^{(j)}(0) = 0$ for $j = 0, \dots, k$, then

$$\int_0^T e^{-tP_0}\theta(t)dt \in \mathcal{L}(\mathcal{F}^{r,0}, \mathcal{F}^{r+2(k+1),0})$$
(A.7)

for $r \in [-2(k+1), 0]$. In particular, $\int_0^T e^{-tP_0} \langle D_x \rangle^{\frac{2(k+1)}{3}} \theta(t) dt \in \mathcal{L}(\mathcal{H}).$

Proof of Lemma 5.3. Let $\chi_1 = 1 - \chi$, where χ is the cut-off used in Proposition 2.1. By the partial Fourier transform in x, $r_l(\lambda)$ is the multiplication by $r_l(\lambda, \xi)$ given by (2.40) in [23]. We decompose it as

$$r_l(\lambda,\xi) = S_1(\lambda,\xi) + S_2(\lambda,\xi) + S_3(\lambda,\xi),$$
(A.8)

where for some constant T > 3,

$$S_{1}(\lambda,\xi) = \int_{T}^{\infty} \left(\chi_{1}(\xi)e^{-t(\widehat{P}_{0}(\xi)-\lambda)} + \chi(\xi)(\sum_{k=l+1}^{\infty}e^{-(k+|\xi|^{2})t}\Pi_{k}^{\xi}) \right) dt,$$

$$S_{2}(\lambda,\xi) = \int_{0}^{T} \chi_{1}(\xi)e^{-t(\widehat{P}_{0}(\xi)-\lambda)} dt,$$

$$S_{3}(\lambda,\xi) = \int_{0}^{T} \chi(\xi)e^{t\lambda} \left(e^{-t\widehat{P}_{0}(\xi)} - \sum_{k=0}^{l}e^{-(k+|\xi|^{2})t}\Pi_{k}^{\xi}\right) dt.$$

Denote

$$S_j = S_j(\lambda, D_x) = \mathcal{F}_{\xi \to x}^{-1} S_j(\lambda, \xi) \mathcal{F}_{x \to \xi},$$

for j = 1, 2, 3.

 $S_1(\lambda,\xi)$ is smooth and rapidly decreasing in ξ . Therefore $S_1 \in \mathcal{L}(\mathcal{G}^{0,s};\mathcal{G}^{0,s})$ for any $s \in \mathbb{R}$. From (A.6) and the subelliptic estimate for P_0 , it follows that $S_1 \in \mathcal{L}(\mathcal{G}^{r,s};\mathcal{G}^{r',s})$ for any $r, r', s \in \mathbb{R}$.

To study S_2 , we use the method of commutators. (A.3) with s = 0 gives

$$[x_j, e^{-tP_0}] = e^{-tP_0} (v_j \sinh t - 2(\cosh t - 1)\partial_{v_j} + 2(\sinh t - t)\partial_{x_j})$$
(A.9)
for $j = 1, \dots, n$. Hence

$$[S_{2}, x_{j}] = -i(\partial_{\xi_{j}}\chi_{1})(D_{x})\int_{0}^{T} e^{-t(P_{0}-\lambda)}dt$$

$$-\chi_{1}(D_{x})\int_{0}^{T} e^{-t(P_{0}-\lambda)}(v_{j}\sinh t - 2(\cosh t - 1)\partial_{v_{j}} + 2(\sinh t - t)\partial_{x_{j}})dt$$
(A.10)

Seeing that

$$\sinh t = O(t), \quad \cosh t - 1 = O(t^2), \quad \sinh t - t = O(t^3),$$

as $t \to 0$, we can apply Lemma A.2 to the three terms in the last integral in (A.10) with, respectively, k = 0, 1 and 2. It follows that

$$[S_2, x_j] \in \mathcal{L}(\mathcal{G}^{0,0}; \mathcal{G}^{2,0}),$$

for $j = 1, \dots, n$. This proves

$$x_j S_2 \langle x \rangle^{-1} = (S_2 x_j - [S_2, x_j]) \langle x \rangle^{-1} \in \mathcal{L}(\mathcal{G}^{0,0}; \mathcal{G}^{2,0}).$$

Consequently, $S_2 \in \mathcal{L}(\mathcal{G}^{0,1}; \mathcal{G}^{2,1})$. By an induction on k and using the commutator method, one can show

$$S_2 \in \mathcal{L}(\mathcal{G}^{0,k}; \mathcal{G}^{2,k}), \quad \forall k \in \mathbb{N}.$$

By the argument of duality and complex interpolation, we deduce that

$$S_2 \in \mathcal{L}(\mathcal{G}^{0,s}; \mathcal{G}^{2,s}), \quad s \in \mathbb{R}.$$

Similarly, using (A.1) and (A.2), we can evaluate the commutators $[S_2, v_j]$ and $[S_2, \partial_{v_j}]$. For $[S_2, v_j]$, one has

$$S_{2}, v_{j}]$$

$$= \chi_{1}(D_{x}) \int_{0}^{T} e^{-t(P_{0}-\lambda)} (v_{j}(\cosh t - 1) - 2\partial_{v_{j}} \sinh t + 2(\cosh t - 1)\partial_{x_{j}}) dt$$
(A.11)

Applying Lemma A.2 and noticing that $(P_0 + 1)^{-1}v_j$, $(P_0 + 1)^{-1}\partial_{v_j}$ and $(P_0 + 1)^{-2}\partial_{x_j}$ belong to $\mathcal{L}(s,s)$ for any s (cf. Lemma 2.2), we infer that $[S_2, v_j] \in \mathcal{L}(\mathcal{G}^{0,s}, \mathcal{G}^{2,s})$. Similarly, we can prove $[S_2, \partial_{v_j}] \in \mathcal{L}(\mathcal{G}^{0,s}, \mathcal{G}^{2,s})$, using (A.2). We deduce that $S_2 \in \mathcal{L}(\mathcal{G}^{1,s}, \mathcal{G}^{3,s})$. By an induction, we can show that $S_2 \in \mathcal{L}(\mathcal{G}^{k,s}, \mathcal{G}^{k+2,s})$ and conclude that $S_2 \in \mathcal{L}(\mathcal{G}^{r,s}, \mathcal{G}^{r+2,s})$ for $r, s \in \mathbb{R}$.

The term S_3 is easier to study. Since $\chi(\xi)$ is of compact support in ξ , S_3 is a convolution with a rapidly decaying function in x variable, which gives that $S_3 \in \mathcal{L}(\mathcal{G}^{0,s}, \mathcal{G}^{2,s})$ for $s \in \mathbb{R}$. In addition, for any $k \in \mathbb{Z}$, $\chi(D_x)(P_0+1)^k(1-\Delta_v+|v|^2)^{-k}$ is bounded. We conclude from Lemma 2.2 and the equation

$$(1 - \Delta_v + |v|^2)^{-k} S_3 (1 - \Delta_v + |v|^2)^k = \left((1 - \Delta_v + |v|^2)^{-k} (P_0 + 1)^k \right) S_3 ((P_0 + 1)^{-k} \left(1 - \Delta_v + |v|^2)^k \right)$$

that $S_3 \in \mathcal{L}(\mathcal{G}^{k,s}, \mathcal{G}^{k+2,s})$ for any $k \in \mathbb{Z}$ and $s \in \mathbb{R}$. A complex interpolation gives that $S_3 \in \mathcal{L}(\mathcal{G}^{r,s}, \mathcal{G}^{r+2,s})$ for $r, s \in \mathbb{R}$.

References

- M. Aafarani, X.P. Wang, Gevrey estimates of the resolvent and sub-exponential time-decay for the heat and Schrödinger semigroups. II. J. Differential Equations 316 (2022), 387-424.
- [2] S. Agmon, Spectral properties of Schrödinger operators and scattering theory. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) 2 (1975), no. 2, 151-218
- [3] S. Agmon, Lectures on Exponential Decay of Solutions to Second Order Elliptic Equations, Princeton Univ. Press, Princeton NJ, 1982.
- M. Ben Said, Kramers-Fokker-Planck operators with homogeneous potentials, Math. Methods in the Appl. Sci., 45 (2) (2022), 914-927
- [5] E. B. Davies, *Linear Operators and Their Spectra*, Cambridge Univ. Press, London, 2007.
- [6] J. Faupin, N. Frantz, Spectral decomposition of some non-self-adjoint operators, Annales Henri Lebesgue, 6 (2023), 1115-1167.
- J. Faupin and J. Fröhlich, Asymptotic completeness in dissipative scattering theory, Adv. Math., 340 (2018), 300-362.

- [8] R. Froese and I. Herbst, Exponential bounds and absence of positive eigenvalues for N-body Schrödinger operators, Commun. Math. Phys. 87 (1982), 429–447.
- B. Helffer, F. Nier, Hypoelliptic estimates and spectral theory for Fokker-Planck operators and Witten Laplacians. Lecture Notes in Mathematics, 1862. Springer-Verlag, Berlin, 2005. x+209 pp. ISBN: 3-540-24200-7
- [10] L. Hörmander, The analysis of linear partial differential operators. III. Pseudo-differential operators. Classics in Mathematics. Springer, Berlin, 2007.
- H. Isozaki, H. Kitada, Micro-local resolvent estimates for 2-body Schrödinger operators, J. Funct. Anal., 57 (1984), 270-300.
- [12] T. Kako, K. Yajima, Spectral and scattering theory for a class of non-self-adjoint operators. Sci. Papers College Gen. Ed. Univ. Tokyo, 26 (1976), no. 2, 73-89
- [13] T. Kato, Perturbation Theory for Linear Operators, Springer, Berlin, 1984.
- [14] T. Li, Z. Zhang, Large time behaviour for the Fokker-Planck equation with general potential, Sci. China, Mathematics, 61 (2018), Iss. 1, pp 137-150.
- [15] W.X. Li, Global hypoellipticity and compactness of resolvent for Fokker-Planck operator, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) 11 (2012), no. 4, 789-815.
- [16] W.X. Li, Compactness criteria for the resolvent of the Fokker-Planck operator, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) 18 (2018), no. 1, 119-143.
- [17] R. Novak, X.P. Wang, On the Kramers-Fokker-Planck equation with decreasing potentials in dimension one, J. of Spectral Theory, 10(1) (2020), 1-32.
- [18] F. W. J. Olver, Asymptotics and Special Functions, A. K. Peters Classics, Massachusetts, 1997.
- [19] H. Risken, The Fokker-Planck equation, Methods of solutions and applications. Springer, Berlin, 1989.
- [20] Y. Saito, The principle of limiting absorption for the nonself-adjoint Schrödinger operator in $\mathbb{R}^{N}(N \neq 2)$. Publ. Res. Inst. Math. Sci. 9 (1973/74), 397-428.
- [21] J. T. Schwartz, Some non-self-adjoint operators, Comm. Pure and Appl. Math., 13(1960), 609-639.
- [22] X. P. Wang, Time-decay of semigroups generated by dissipative Schrödinger operators. J. Differential Equations 253 (2012), no. 12, 3523-3542.
- [23] X. P. Wang, Large-time asymptotics of solutions to the Kramers-Fokker-Planck equation with a short-range potential. Comm. Math. Phys. 336 (2015), no. 3, 1435-1471.
- [24] X.P. Wang, Gevrey estimates of the resolvent and sub-exponential time-decay for the heat and Schrödinger semigroups, J. Math. Pures Appl. 135 no. 9 (2020), 284-338
- [25] X.P. Wang, L. Zhu, Global-in-time $L^p L^q$ estimates for solutions of the Kramers-Fokker-Planck equation, Commun. Math. Res., **38**(4) (2022), 560-578.

LABORATOIRE DE MATHÉMATIQUES JEAN LERAY, UMR CNRS 6629, NANTES UNIVERSITÉ, 44322 NANTES CEDEX 3 FRANCE, E-MAIL: XUE-PING.WANG@UNIV-NANTES.FR