Quantum Algorithms and Applications for Open Quantum Systems

Luis H. Delgado-Granados,[†] Timothy J. Krogmeier,^{‡,¶} LeeAnn M.

Sager-Smith,[§] Irma Avdic,[†] Zixuan Hu,^{∥,⊥} Manas Sajjan,[∥] Maryam Abbasi,^{‡,¶}

Scott E. Smart,[#] Prineha Narang,[#] Sabre Kais,[∥] Anthony W. Schlimgen,^{‡,¶}

Kade Head-Marsden,*,^{‡,¶} and David A. Mazziotti^{*,†}

†Department of Chemistry and The James Franck Institute, The University of Chicago, Chicago, IL 60637 USA

[‡]Department of Chemistry, Washington University in St. Louis, St. Louis, MO 61630 USA ¶Department of Chemistry, University of Minnesota, Minneapolis, MN 55455 USA

§Department of Chemistry and Physics, Saint Mary's College, Notre Dame, IN 46556 USA ||Department of Chemistry, Department of Electrical and Computer Engineering, and Purdue Quantum Science and Engineering Institute, Purdue University, West Lafayette, IN 47907, USA *LKey Laboratory of Precision and Intelligent Chemistry, University of Science and Technology of China, Hefei 230026, China*

#Division of Physical Sciences, College of Letters and Science, University of California, Los Angeles, CA 90095, USA

E-mail: khm@umn.edu; damazz@uchicago.edu

Abstract

Accurate models for open quantum systems—quantum states that have non-trivial interactions with their environment—may aid in the advancement of a diverse array of fields, including quantum computation, informatics, and the prediction of static and dynamic molecular properties. In recent years, quantum algorithms have been leveraged for the computation of open quantum systems as the predicted quantum advantage of quantum devices over classical ones may allow previously inaccessible applications. Accomplishing this goal will require input and expertise from different research perspectives, as well as the training of a diverse quantum workforce, making a compilation of current quantum methods for treating open quantum systems both useful and timely. In this Review, we first provide a succinct summary of the fundamental theory of open quantum systems and then delve into a discussion on recent quantum algorithms. We conclude with a discussion of pertinent applications, demonstrating the applicability of this field to realistic chemical, biological, and material systems.

1 Introduction

Open quantum systems (OQS) arise when a quantum system has non-negligible interactions with its environment. Theoretical treatment of an open quantum system entails separating the degrees of freedom into a *system* and the remaining degrees of freedom as an *environment* or *bath*.^{1,2} The dynamics of OQS are often divided into two primary categories. The first category, which we call *Markovian*, is when the relaxation time of the system is significantly slower than that of the environment, and second, which we call *non-Markovian*, is when the time scales of relaxation within the system and the environment are comparable.^{1–6} OQS have potentially significant applications in advancing quantum computing,^{7–11} cryptography,¹² metrology,^{13,14} simulation,¹⁵ thermodynamics,¹⁶ control,^{17–19} and information processing,²⁰ as well as in understanding and optimizing chemical reactions in solvents, thereby bridging fundamental science with practical applications across technology, biology,^{21,22} and materials science.²³

Modeling OQS on quantum computers, in particular, has the potential to revolutionize various classes of computation by harnessing the unique information processing available in quantum mechanics.^{24–26} While classical computers perform their calculations through combinations of bits, with each bit being a Boolean choice of 0 or 1, quantum computers perform their calculations through combinations of quantum bits, or qubits, in which each qubit can be any linear superposition of 0 and 1. A crucial application area for quantum computing is the prediction of the static

and dynamic properties of many-electron atoms, molecules, and materials. Although a plethora of algorithms have been proposed,^{27–33} most algorithms have focused on treating electronic structure in closed quantum systems. In this review article, we summarize a new, emerging frontier of quantum algorithms for open quantum systems. Because these systems require significant computational resources on classical devices, their simulation on quantum devices has the potential for realizing significant computational savings and reaching important, potentially hitherto inaccessible applications.



Figure 1: Schematic representation of the translation of OQS dynamics into a quantum computing framework for potential applications beyond classical capabilities.

In recent years, with the rapid progress in different quantum computing platforms such as superconducting qubits, ³⁴ ion-traps, ³⁵ and cold atoms, ³⁶ it has become desirable to test and benchmark new algorithms on quantum devices. To this end, a variety of approaches have been proposed, benchmarked, and applied to model OQS using quantum computers. ^{37–48} Because quantum computers work within the language of unitary transformations, ^{24,25} a fundamental challenge to simulating OQS on quantum devices is to treat their *non-unitary* dynamics. Efforts have been made to address this challenge through embedding non-unitary operators into larger unitary operators in a process referred to as dilation or block encoding, ^{42,46,48–53} as well as using imaginary time evolution, ⁴⁷ variational approaches, ^{45,54–67} tensor train approaches, ⁶⁸ and many more. ^{41,43,44,69–92} This process is depicted schematically in Figure 1, and with more step-by-step detail in Figure 2.

Existing quantum algorithms have been used to explore a variety of problems relevant to chem-



Figure 2: Step-by-step representation of the mapping from a classical open system approach into a quantum open system framework.

istry, ^{93,94} biology, ^{95–100} condensed matter, ^{47,101–105} electrodynamics, ¹⁰⁶ and materials science. ¹⁰⁷ The majority of these algorithms have shared a few common benchmarking systems including twolevel systems with amplitude damping or dephasing channels, ^{42,44,46,49,108} the Jaynes-Cummings model with weak, strong, and detuned couplings, ^{44,108,109} and the spin-boson model. ¹¹⁰ Beyond benchmarking, these algorithms have focused primarily on systems where the dynamics involve two types of processes, unitary or reversible dynamics often captured in the Hamiltonian, and non-unitary or irreversible dynamics often captured by additional terms. These initial efforts form a foundation for future applications in the treatment of open quantum systems that are classically challenging or intractable. In this Review, we begin by providing a brief overview of the fundamental theory of open quantum systems in Section 2. We then discuss current quantum algorithms and methodologies to consider the dynamics of OQS on quantum computers in Section 3. Applications to fundamental open quantum system models and specific problems including photosynthetic light harvesting, the avian compass, and other systems are presented in Section 4. Finally, we leave the reader with some concluding thoughts and an outlook towards future advances and applications in this field in Section 5.

2 Theoretical Foundations

In this section, we present an overview of the foundations of the theory of open quantum systems, including their definition, quantum maps, Markovian and non-Markovian dynamics, and quantum master equations.

2.1 Defining an Open Quantum System



Figure 3: A quantum system partitioned into the system of interest (S, \hat{H}_S, ρ_S) and the environment (E, \hat{H}_E, ρ_E) interacting through \hat{H}_{SE} .

Starting from an overall closed system, we can express an open quantum system as a system S embedded in an environment E. Here S is coupled to E to generate a system-environment supersystem SE, described for time t by a density matrix $\rho_{SE}(t)$ that belongs to the Hilbert space

 $\mathcal{H}_{SE} = \mathcal{H}_S \otimes \mathcal{H}_E$, where dim $(\mathcal{H}_E) \gg \dim(\mathcal{H}_S)$. This is shown schematically in Figure 3.^{1,2,111}

The time evolution of the system-environment density matrix can be expressed as a unitary operator $\hat{U}(t, t_0)$ acting on the initial density matrix $\rho_{SE}(t_0)$,

$$\rho_{SE}(t) = \hat{U}(t, t_0) \rho_{SE}(t_0) \hat{U}^{\dagger}(t, t_0), \qquad (1)$$

with $t > t_0$ and $\hat{U}(t, t_0)$ defined as, ^{1,2}

$$\hat{U}(t,t_0) = \hat{T} \exp\left\{-i \int_{t_0}^t \hat{H}_T(\tau) d\tau\right\},$$
(2)

in which \hat{T} is the time-ordering operator and \hat{H}_T is the total Hamiltonian of the system and the environment,

$$\hat{H}_T(\tau) = \hat{H}_S(\tau) \otimes \hat{I}_E + \hat{I}_S \otimes \hat{H}_E(\tau) + \hat{H}_{SE}(\tau),$$
(3)

where $\hat{H}_S(\tau)$, $\hat{H}_E(\tau)$, and $\hat{H}_{SE}(\tau)$ are the Hamiltonians of the system, effective environment, and the interaction term between the two, respectively.^{1,2,111} Formally, we can obtain the system by tracing the environment,

$$\rho_S(t) = \operatorname{Tr}_E\left(\hat{U}(t,t_0)\rho_{SE}(t_0)\hat{U}^{\dagger}(t,t_0)\right).$$
(4)

Moreover, assuming that the system and the environment are not initially correlated, we can represent the embedding of the system S into its environment E by an extension or assignment map E_{v} ,^{112–114}

$$E_v: \rho_S(t_0) \to \rho_S(t_0) \otimes \rho_E(t_0), \tag{5}$$

which leads to,

$$\rho_S(t) = \operatorname{Tr}_E\left(\hat{U}(t, t_0)(\rho_S(t_0) \otimes \rho_E(t_0))\hat{U}^{\dagger}(t, t_0)\right).$$
(6)

The above equation provides a density matrix representing our open quantum system with the only

assumption being the initial separability of the system and environment.^{1,111,113,114}

2.2 Quantum Maps

The time evolution of the system in the previous subsection defines a quantum map $\mathcal{E}_{(t,t_0)}$,

$$\mathcal{E}_{(t,t_0)}: \rho_S(t_0) \to \rho_S(t), \quad (t > t_0), \tag{7}$$

which is Hermitian, trace-preserving, and positive. ^{1–3,111–113,115} Unless specifically indicated in this review, we consider quantum maps to be not only positive but also completely positive (CP). A CP map places additional restrictions from the requirement that the system-environment density matrix remain positive semidefinite for all time. ^{111,112,115–117}

By rewriting the environment in Eq. (6) as sums over the environmental states $|m\rangle$ and $|n\rangle$ with associated weights $\omega_n(t_0)$, we obtain the operator-sum representation of the system density matrix,

$$\rho_{S}(t) = \sum_{m} \langle m | \hat{U}(t, t_{0}) \rho_{S}(t_{0}) \otimes \rho_{E}(t_{0}) \hat{U}^{\dagger}(t, t_{0}) | m \rangle$$

$$= \sum_{m,n} \omega_{n}(t_{0}) \langle m | \hat{U}(t, t_{0}) | n \rangle \rho_{S}(t_{0}) \langle n | \hat{U}^{\dagger}(t, t_{0}) | m \rangle$$

$$= \sum_{m,n} \hat{M}_{m,n}(t, t_{0}) \rho_{S}(t_{0}) \hat{M}^{\dagger}_{m,n}(t, t_{0})$$

$$= \sum_{k} \hat{M}_{k}(t, t_{0}) \rho_{S}(t_{0}) \hat{M}^{\dagger}_{k}(t, t_{0}) \qquad (8)$$

where $\hat{M}_k(t, t_0) = \hat{M}_{m,n}(t, t_0) = \sqrt{\omega_n(t_0)} \langle m | \hat{U}(t, t_0) | n \rangle$ are the Kraus operators. These operators obey the contraction mapping,

$$\sum_{k} \hat{M}_{k}(t, t_{0}) \hat{M}_{k}^{\dagger}(t, t_{0}) = \hat{I}_{S}.$$
(9)

Importantly, this representation is not unique, as it is dependent on the basis one chooses to represent the environment.^{1,111} Additionally, the number of Kraus maps needed to describe fully the dynamics

of the system is upper bounded by the square of the dimension of the system, $\dim(\mathcal{H}_S)^2$.^{48,111}

In general, quantum maps are not unitary. To obtain a unitary treatment, we can dilate S to a larger Hilbert space according to the *the Stinespring dilation theorem*.^{48,118,119} The theorem states that if one has a well-defined quantum map, then there exists a state $\rho_A \in \mathcal{H}_A$ where A is an ancillary environment such that,

$$\rho_S(t) = \operatorname{Tr}_A\left(\hat{U}_{SA}(t, t_0)\rho_S(t_0) \otimes \rho_A \hat{U}_{SA}^{\dagger}(t, t_0)\right),\tag{10}$$

where $\hat{U}_{SA}(t,t_0) \in \mathcal{H}_S \otimes \mathcal{H}_A$ and $\dim(\mathcal{H}_A) \leq \dim(\mathcal{H}_S)^2$. The unitary $\hat{U}_{SA}(t,t_0)$, is built by stacking the Kraus maps to obtain the first block column,

$$\hat{U}_{SA}(t,t_0) = \begin{bmatrix} \hat{M}_1 & | & | & | \\ \vdots & \mathbf{U}_2 & \mathbf{U}_3 & \cdots & \mathbf{U}_k \\ \hat{M}_k & | & | & | \end{bmatrix},$$
(11)

where $k \leq \dim(\mathcal{H}_S)$ and the remaining block columns \mathbf{U}_k are chosen such that $\hat{U}_{SA}(t, t_0)\hat{U}_{SA}^{\dagger}(t, t_0) = \hat{I}_{SA}$.

In addition to the Kraus maps and unitary representation, there are two other representations known as dynamical matrices defined as^{111,120,121}

$$\mathfrak{A} = \sum_{k} \hat{M}_{k} \otimes \hat{M}_{k}^{*} \tag{12}$$

$$\mathfrak{B} = \sum_{k} \left| \hat{M}_{k} \right\rangle \left\langle \hat{M}_{k} \right|. \tag{13}$$

where $|\hat{M}_k\rangle$ is the vectorized form of \hat{M}_k . The properties of these representations and their relationship with Kraus maps are fully described in Refs. 111 and 120. An advantage of using these representations is that they provide a direct way to determine if a map is CP; if at least one of the eigenvalues of \mathfrak{B} is negative, then the quantum map is not CP.¹²²

2.3 Quantum Master Equations

2.3.1 Markovian Master Equations

A quantum map, $\mathcal{E}_{(t,t_0)}$, that describes Markovian dynamics can be represented as,

$$\mathcal{E}_{(t,t_0)} = \exp\{\mathcal{L}(t-t_0)\},\tag{14}$$

where the superoperator \mathcal{L} is the dynamics generator that acts on $\rho_S(t)$.^{1,3,115,123} This leads to the time-local differential equation,

$$\frac{d}{dt}\rho_S(t) = \mathcal{L}\rho_S(t),\tag{15}$$

which is referred to as the Markovian master equation.¹²⁴ We can derive \mathcal{L} by projecting the Kraus maps onto an orthogonal set of operators in the Fock-Liouville space, ^{125–127}

$$\frac{d}{dt}\rho_{S}(t) = -i\left[\hat{H},\rho_{S}(t)\right] + \sum_{k=1}\gamma_{k}\left(\hat{L}_{k}\rho_{S}(t)\hat{L}_{k}^{\dagger} + \frac{1}{2}\{\hat{L}_{k}^{\dagger}\hat{L}_{k},\rho_{S}(t)\}\right)$$

$$= -i\left[\hat{H},\rho_{S}(t)\right] + \hat{\mathcal{D}}(\rho_{S}(t)),$$
(16)

where the first term is inherited from the quantum Liouville equation, and the second term is called the dissipator in which \hat{L}_k are the Lindblad operators with their associated decay rates γ_k . The notations $\{\cdot\}$ and $[\cdot]$ represent the anti-commutator and commutator, respectively. Provided the decay rates are all of the same sign, this master equation preserves the trace and complete positivity of the density matrix.

The Gorini-Kossakowski-Sudarshan-Lindblad (GKSL) equation can be derived from the Liouville equation of the combined system and environment. In this derivation, several key assumptions are at play. The system and environment are weakly coupled, known as the Born approximation, and the bath correlations decay quicker than the time scale of system relaxation, known as the Markov approximation. Often both the rotating wave and secular approximations are also invoked.^{1,115,125,126} Following a similar derivation, one can arrive at the Redfield master equation through a perturbative approximation.^{1,128}

2.3.2 Non-Markovian Master Equations

The general non-Markovian master equations can be derived by projection operator techniques introduced simultaneously by Nakajima¹²⁹ and Zwanzig.¹³⁰ The projection operator approach is based on two projection superoperators, \mathcal{P} and \mathcal{Q} , both of which act in the composite Hilbert space \mathcal{H}_{SE} .^{1,110,123,131} \mathcal{P} projects $\rho_{SE}(t)$ to a state in which S and E are not coupled,

$$\mathcal{P}\rho_{SE}(t) = \operatorname{Tr}_{E}\left(\rho_{SE}(t)\right) \otimes \rho_{E} = \rho_{S}(t) \otimes \rho_{E}, \tag{17}$$

where $\mathcal{P}\rho_{SE}(t)$ is referred to as the *relevant* part of $\rho_{SE}(t)$ and ρ_E represents an arbitrary state in \mathcal{H}_E , which is usually chosen to be a stationary state of the environment. The state resulting from $\mathcal{Q}\rho_{SE}(t)$,^{1,123}

$$\mathcal{Q}\rho_{SE}(t) = \rho_{SE}(t) - \mathcal{P}\rho_{SE}(t) \tag{18}$$

is called the *irrelevant* part of $\rho_{SE}(t)$. Both projection superoperators follow the same properties as one expects from a projection operator: $\mathcal{P} + \mathcal{Q} = \hat{I}_{SE}$, $\mathcal{P}^2 = \mathcal{P}$, $\mathcal{Q}^2 = \mathcal{Q}$, and $\mathcal{P}\mathcal{Q} = \mathcal{Q}\mathcal{P} = \hat{0}_{SE}$.

Using these projection operators, we can derive the generalized master equation in both nonlocal and local forms, often referred to as the Nakajima-Zwanzig^{129,130} equation and the timeconvolutionless (TCL) equation, respectively.^{132,133} The Nakajima-Zwanzig equation can be written as,¹

$$\frac{d}{dt}\rho_{S}(t) = \alpha \operatorname{Tr}_{E}\left(\mathcal{PL}(t)\mathcal{G}(t,t_{0})\mathcal{Q}\rho_{SE}(t_{0})\right) + \int_{t_{0}}^{t} ds \operatorname{Tr}_{E}\left(\mathcal{K}(t,s)\mathcal{P}\rho_{SE}(s)\right),$$
(19)

where $\mathcal{G}(t, t_0)$ is a time non-unitary propagator of the form,

$$\mathcal{G}(t,t_0) = \hat{T} \exp\left\{\alpha \int_{t_0}^t ds \mathcal{QL}(s)\right\},\tag{20}$$

and $\mathcal{K}(t, s)$, which is referred to as the memory kernel, is,

$$\mathcal{K}(t,s) = \alpha^2 \mathcal{PL}(t) \mathcal{G}(t,s) \mathcal{QL}(s) \mathcal{P}.$$
(21)

Alternatively, the TCL equation can be expressed as,

$$\frac{d}{dt}\rho_{S}(t) = \hat{A}(t)\rho_{S}(t) + \rho_{S}(t)\hat{B}^{\dagger}(t) + \sum_{i}\hat{C}_{i}(t)\rho_{S}(t)\hat{D}_{i}^{\dagger}(t),$$
(22)

where $\hat{A}(t)$, $\hat{B}(t)$, $\hat{C}(t)$, and $\hat{D}(t)$ are time-dependent linear operators.

While the above approaches provide generalizations to the Markovian methods, they are challenging to solve due to computational scaling and complexity. While much progress has been made in terms of method development,^{3,134} simulating non-Markovianity efficiently and accurately remains a challenge.

3 Algorithms for Quantum Simulation



Figure 4: Simulations to open quantum systems, in comparison to closed quantum systems, introduces additional challenges such as the need for non-unitary quantum dynamics.

Quantum simulation for closed quantum systems has significant challenges, including efficient mapping of orbitals onto qubits and optimization of circuits. The extension of simulations to open quantum systems introduces additional challenges, such as the need for non-unitary quantum dynamics, as depicted schematically in Figure 4. One of the most common approaches to this need is to translate the non-unitary dynamics into purely unitary dynamics. In this section, we review algorithms for the quantum simulation of open quantum systems, a sampling of which is depicted

schematically in Figure 5.



Figure 5: A selection of quantum algorithms for the simulation of open quantum systems.

3.1 Probabilistic Methods or Block-Encoding Techniques

The general process of block encoding, or dilation, is where a non-unitary system is mapped into a unitary framework in a larger Hilbert space, as shown on the left of Figure 6. This unitary can be encoded into a qubit framework, where the dilation is performed with ancillary qubits. For the noisy intermediate-scale quantum (NISQ) era, in which high gate count and number of qubits can limit the usefulness of an algorithm, it is preferable to keep dilation to a minimum such that the qubit space stays as small as possible. We will discuss several different algorithms that fall within this category, including the Sz.-Nagy dilation,^{42,97,98,108,110} the unitary operator decomposition,^{46,102} and the classical singular value decomposition (SVD),^{49,99} of which the latter two are depicted schematically on the right of Figure 6.



Figure 6: (Left) Block-encoding or dilation techniques where the original system Hilbert space is dilated into a larger space such that the dilated system evolves in a unitary fashion. (Right) Two examples of dilation-based approaches with the singular value decomposition approach⁴⁹ depicted schematically on the top and the operator decomposition approach⁴⁶ depicted schematically on the bottom.

3.1.1 Markovian Semigroups and Linear Combination

Some early work in understanding quantum algorithms for open quantum systems focused on finding universal sets of qubit gates for Markovian dissipative dynamics. Bacon and co-workers demonstrated that for one qubit, a single non-unitary operator parametrized by one variable is sufficient to generate all Markovian dynamics.³⁷ They did so by decomposing the GKSL generator into simple components through a linear combination of semigroups and unitary conjugation. Building on this work, Sweke and co-workers described a universal set of one- and two-qubit gates to simulate arbitrary one-qubit Markovian dynamics. The resulting minimal dilation utilized techniques from Hamiltonian simulation to show complexity bounds for implementing Markovian channels with controlled accuracy. In particular, using the Suzuki-Trotter-Lie formula, the authors draw an analogy to Hamiltonian simulation to derive complexity bounds that are polynomial in the norm of the Markovian generators and the desired error tolerance.^{39,135} This was also generalized to treat non-Markovian channels.⁴⁰ A thorough tutorial on these approaches can be found in Ref.

136.

3.1.2 Sz.-Nagy Dilation-Based Algorithm

The Stinespring dilation theorem states that any non-unitary quantum operator can be converted into a unitary one;^{118,137} however, naïve application of the Stinespring dilation increases the operator dimension dramatically and thus incurs high computational cost. Alternatively, the Sz.-Nagy dilation can be used to map Kraus operators into a larger unitary form that was then implementable on a quantum device.⁴² This dilation works with the density matrix in operator-sum form,

$$\rho(t) = \sum_{k} \hat{M}_{k} \rho(0) \hat{M}_{k}^{\dagger}, \qquad (23)$$

where $\rho(t)$ is the density matrix, \hat{M}_k are the Kraus operators, and the summation is over all possible channels of interaction with the environment. The non-unitary operators \hat{M} are dilated through the Sz.-Nagy approach such that the unitary representation is given by,

$$\hat{U}_k = \begin{pmatrix} \hat{M}_k & \hat{D}_{\hat{M}_k^{\dagger}} \\ \hat{D}_{\hat{M}_k} & -\hat{M}_k^{\dagger} \end{pmatrix}, \qquad (24)$$

where $\hat{D}_{\hat{M}_k}$ is the defect operator given by $\sqrt{I - \hat{M}_k^{\dagger} \hat{M}_k}$. This dilated unitary acts on each wavefunction of a decomposed density matrix,

$$\rho(t) = \sum_{k} \sum_{j} |c_j|^2 \hat{U}_k |\psi_j\rangle \langle \psi_j | \hat{U}_k^{\dagger}, \qquad (25)$$

in which the wavefunctions are padded with zeroes to match the dimensionality of the dilated unitary operator.

We can relate the integrated Kraus representation to any differential master equation such as the Lindblad equation by defining the Kraus operators as $\hat{M}_k = \sqrt{\gamma_k \delta t} L_k$, where L_k are the Lindbladian operators, with decay rates γ_k , and time step δt .^{97,120} This mapping recasts the basic Sz.-Nagy dilation to apply to the Lindblad master equation. One challenge of the Sz.-Nagy dilation approach is that it incurs an undesirable classical cost in taking the matrix square root while computing the dilated unitary operator.

This approach has been effective to consider basic Kraus dynamics,⁴² non-Markovian dynamics,^{108,110} and the dynamics in a few biological systems.^{97,98} This algorithm has also been applied in NMR quantum processing.⁵⁰

3.1.3 Unitary Decomposition Algorithm

One alternative dilation approach referred to as the unitary decomposition algorithm,⁴⁶ considers the decomposition of an operator into the sum of its Hermitian and anti-Hermitian components,

$$\hat{M} = \hat{H} + \hat{A}.$$
(26)

This can be recast as a sum of unitary operators

$$\hat{M} = \lim_{\epsilon \to 0} \frac{1}{2\epsilon} (ie^{-i\epsilon\hat{H}} - ie^{i\epsilon\hat{H}} + e^{\epsilon\hat{A}} - e^{-\epsilon\hat{A}}).$$
(27)

Because each of the exponentials above is a unitary operator, the equation can be implemented on a quantum computer. The summation can be performed with the linear combination of unitaries (LCU) technique.¹³⁸ While this unitary decompoition algorithm requires two ancillary qubits in the dilation in contrast to the Sz.-Nagy dilation, it overcomes the potentially significant classical cost of computing the matrix square root of \hat{M} . This method has been applied using the operator-sum and the vectorized Lindblad equation.^{46,102} A related dilation has been employed in electronic structure in the context of the contracted quantum eigensolvers, ^{139–142} which use the residual of a contraction of the Schrödinger equation to define an exact, iterative *Ansatz* for the wave function.

3.1.4 Classical Singular Value Decomposition Algorithm

Another recent approach to dilating non-unitary operators efficiently relies on computing the classical singular value decomposition (SVD). This involves a classical calculation of the SVD, dilating only the singular value matrix, and implementing this new form on a quantum computer to obtain the desired quantum dynamics. The SVD is written as,

$$\hat{M} = \hat{U}\hat{\Sigma}\hat{V}^{\dagger},\tag{28}$$

in which \hat{U} and \hat{V}^{\dagger} are unitary operators, while the $\hat{\Sigma}$ operator is non-unitary. This can be cast into unitary form via the dilation,

$$\hat{U}_{\hat{\Sigma}} = \begin{pmatrix} \hat{\Sigma}_{+} & 0\\ 0 & \hat{\Sigma}_{-} \end{pmatrix}, \qquad (29)$$

where $\hat{\Sigma}_+$ and $\hat{\Sigma}_-$ are generated using the singular values of \hat{M} .

This approach has been applied for unnormalized state preparation, basic dynamics, and systems in quantum biology.^{49,99} A recent study proposed a quantum singular value transformation algorithm to calculate the SVD on a quantum device, thus avoiding the classical complexity.⁴⁸

3.1.5 Monte Carlo and Minimal Dilation Approaches

For certain classes of channels, efficient dilations can be realized, which are more amenable for quantum computers. In particular, given a quantum map of the form Eq. (8), the channel is unital if it preserves the identity state, $\rho = I_d/d$. A channel is mixed unitary, or random unitary or a random external field, if it has the form,

$$\mathcal{E}[\rho] = \sum_{i} p_i U_i \rho U_i^{\dagger}, \tag{30}$$

where the collection of p_i is convex and U_i are unitary operators. Mixed unitary channels represent a subset of unital operators and are relevant for quantum information and quantum processes where unitary controls have some probabilistic error.

By preparing an ancilla state with probability amplitudes proportional to the square root of p_i , one can efficiently prepare a LCU circuit that affects the desired channel on some state.⁴⁸ However, when the number of p_i is large, this can become challenging, particularly in the near-term era of quantum computation, and recent works have instead utilized Monte Carlo approaches to realize these channels.^{143–145} Explicitly, an estimator of a quantum channel can be derived that samples from a multinomial distribution representing a collection of unitary channels as outcomes $\{(p_i, U_i)\}$.¹⁴⁶ The state estimator

$$\hat{\rho}_{\mathcal{E}} = \sum_{k} \hat{s}_{k} U_{k} \rho U_{k}^{\dagger} \tag{31}$$

can be implemented with no dilation, and represents an aggregate construction of the channel by probabilistically sampling quantum circuits against a measurement outcome with a given frequency. This approach was shown to be scalable to very large systems and completely bypasses the costs associated with dilation. Additionally, successive probabilistic channels can be sampled in a Markovian fashion, allowing for time evolution of channels to be performed.

These techniques have also been applied in the context of simulating spin systems.⁹³ Certain non-unitary operators, such as the amplitude damping channel, can be realized as a random unitary channel on a dilated space, and thus, the above procedure can be used to avoid further dilation. Approximate random unitary channels can also be constructed for given channels.¹⁴⁷ While inherently probabilistic, the above methods have approximate success probabilities of unity, i.e., they are not post-selected and hence, scale similarly to deterministic approaches.

3.2 Deterministic Algorithms

In addition to block-encoding methods, a variety of deterministic methods have been developed to simulate the non-unitary evolution of quantum systems. In particular, several algorithms are based on purification techniques, ^{148,149} quantum imaginary time evolution (QITE), ⁴⁷ and variational approaches. ^{54,150} Purification relies on doubling the size of the system by representing the

mixed state as a wavefunction, or column vector.^{151–153} QITE is a measurement-based approach in which a unitary operator is generated that approximates the non-unitary evolution, typically by sampling a known decomposition of the operator.^{45,47,154–157} Finally, variational approaches utilize a parametrized circuit where the parameters are variationally optimized to simulate the time evolution.

3.2.1 Purification

The density-matrix purification 24,25,148,151,152,158 introduces an effective bath, *B*, whose dimension equals the dimension of the system. The openness of the system is explicitly created through the entanglement of the system with the effective bath, such that the system density matrix takes the form,

$$\rho_S(t) = \operatorname{Tr}_B\left(\left|\Psi^{SB}(t)\right\rangle \left\langle \Psi^{SB}(t)\right|\right),\tag{32}$$

where $|\Psi^{SB}(t)\rangle$ is a pure state of the composite system SB,

$$\left|\Psi^{SB}(t)\right\rangle = \sum_{i} \sqrt{\omega_{i}(t)} \left|\Psi_{i}^{S}(t)\right\rangle \left|\Psi_{i}^{B}(t)\right\rangle.$$
(33)

This definition of the composite system allows a unitary description of its dynamics,

$$\left|\Psi^{SB}(t)\right\rangle = \hat{U}_{SB}(t,t_0) \left|\Psi^{SB}(t_0)\right\rangle \tag{34}$$

$$= \hat{T} \exp\left\{-i \int_{t_0}^t \hat{H}_T(\tau) d\tau\right\} \left|\Psi^{SB}(t_0)\right\rangle,\tag{35}$$

where \hat{T} is the time-ordering operator, and $\hat{H}_T(\tau)$ is the total Hamiltonian,

$$\hat{H}_T(\tau) = \hat{H}_S(\tau) \otimes \hat{I}_B + \hat{I}_S \otimes \hat{H}_B(\tau) + \hat{H}_{SB}(\tau).$$
(36)

This theory results in a deterministic algorithm because the effective bath qubits are not used as postselected ancillas. Instead, the effective environment's degrees of freedom are averaged after the time propagation. Although purification doubles the number of qubits for simulation, this approach allows for treating initially entangled system-environment dynamics in Markovian and non-Markovian regimes in a common framework. It also can treat cases in which the assumption of complete positivity breaks down.¹⁴⁹

3.2.2 Quantum Imaginary Time Evolution

Quantum imaginary time evolution (QITE) is a general algorithm that effectively solves an eigenvalue problem by a unitary least-squares approximation of the non-unitary dynamics.^{154,159} It can also be used in open quantum systems when a propagator form of the operator is available; for example, in the vectorized Lindblad equation.⁴⁷ In this case, we can write the time propagation in terms of the Lindbladian \mathcal{L} ,

$$|\rho(t)\rangle = e^{\mathcal{L}t}|\rho(0)\rangle. \tag{37}$$

Here, we have used the vectorized form of the density matrix, $|\rho\rangle$. The QITE algorithm seeks a Hermitian matrix Q, which approximately satisfies,

$$\frac{e^{\mathcal{L}t}|\rho(0)\rangle}{\|e^{\mathcal{L}t}|\rho(0)\rangle\|} = e^{-iQt}|\rho(0)\rangle.$$
(38)

If Q is written as a linear combination of Pauli operators, σ_i ,

$$Q = \sum_{i} x_i \sigma_i,\tag{39}$$

then we can solve a least squares equation Ax = b where,

$$A_{ij} = \langle \rho | \sigma_i^{\dagger} \sigma_j | \rho \rangle,$$

$$b_i = \frac{-i}{\|e^{\mathcal{L}t}|\rho\rangle\|} \langle \rho | \sigma_i H | \rho \rangle,$$
(40)

is measured with a quantum circuit. After finding Q, the vectorized time-evolved state is prepared with the unitary e^{-iQt} , which effectively acts as an *Ansatz* for the target state.

QITE has been used to simulate open quantum systems in a variety of contexts, including dissipative two-level systems and Ising models.⁴⁷ Other work has focused on reducing the circuit depth, an important development for any algorithm to be implementable on NISQ devices.^{160–162}

3.2.3 Variational Approaches

Time-dependent variational methods for closed or open systems rely on using a parameterized quantum circuit as an *Ansatz*,

$$|\psi(\vec{\theta}(t))\rangle = U(\vec{\theta}(t))|0\rangle^{\otimes N},\tag{41}$$

where $U(\vec{\theta_i}(t))$ corresponds to a unitary operator parameterized by a set of angles $\{\theta_i\}$ and N is the number of qubits. The variational approach is applicable to any differential equation of the following form,

$$\eta(t)\frac{d|\psi(\vec{\theta}(t))\rangle}{dt} = |d\psi(\vec{\theta}(t))\rangle = \hat{\chi}(t)|\psi(\vec{\theta}(t))\rangle, \tag{42}$$

where $\eta(t) \in \mathbb{C}$ and $\hat{\chi}(t) \in \mathcal{L}(\mathbb{C}^{2^N})$ can be chosen to represent either closed or open system quantum dynamics.⁴⁵

Using the traditional time-dependent variational principle, we construct the following equation of motion (EOM), ^{163–169}

$$\sum_{j} \operatorname{Im}(M_{ij}(t))\dot{\theta}_{j}(t) = \operatorname{Im}\left(\frac{1}{\eta(t)}V_{i}(t)\right),\tag{43}$$

where $M_{ij} = \langle \frac{\partial \psi(\vec{\theta}(t))}{\partial \theta_i} | \frac{\partial \psi(\vec{\theta}(t))}{\partial \theta_j} \rangle$, $\dot{\theta}_j(t) = \frac{d\theta_j}{dt}$, and $V_i = \langle \frac{\partial \psi(\vec{\theta}(t))}{\partial \theta_i} | \hat{\chi}(t) | \psi(\vec{\theta}(t)) \rangle$. For certain choices of the *Ansatz*, Im(M(t)) can become singular or nearly singular, leading to numerical instability.¹⁶⁷ Alternative EOMs are derived using two approaches, the Dirac-Frenkel^{170–172} and MacLaghlan

variational principles, 173-178

$$\sum_{j} M_{ij}(t)\dot{\theta}_{j}(t) = \frac{V_{i}(t)}{\eta(t)} \implies M(t)\dot{\vec{\theta}}(t) = \frac{V(t)}{\eta(t)},$$
(44)

and

$$\sum_{j} \operatorname{Re}(M_{ij}(t))\dot{\theta}_{j}(t) = \frac{V_{i}(t)}{\eta(t)},$$
(45)

respectively.

Whichever scheme is selected for generating the EOM for the parameters $\{\theta_j(t)\}_{j=0}^k$, the elements of matrix M(t) and of the vector V(t) can be estimated according to,

$$M_{ij}(t) = \langle 0|^{\otimes N} \mathcal{U}_i^{\dagger} \mathcal{U}_j | 0 \rangle^{\otimes N}, \qquad (46)$$

and

$$V_i(t) = \langle 0|^{\otimes N} \mathcal{U}_i^{\dagger} \hat{\chi}(t) U(\vec{\theta}(t)) | 0 \rangle^{\otimes N}.$$
(47)

where $\mathcal{U}_i = \frac{\partial U(\vec{\theta}(t))}{\partial \theta_i}$. Both quantities can be estimated from the quantum circuit directly either via indirect measurements or the modified Hadamard test.^{45,167,179} For the treatment of open quantum systems, the generators $\hat{\chi}(t)$ are non-Hermitian, requiring their decomposition as $\hat{\chi}_1(t) + i\hat{\chi}_2(t)$. The generator $\hat{\chi}_1(t)$ yields a modified set of *Ansatz* parameters corresponding to the nearest unitarily-evolved, parametrized state.^{54,180,181}

Following the computation of M(t) and V(t) via quantities extracted from the quantum circuits, the EOM is solved on a classical device.^{180,182} Thereafter, a new set of evolved variational parameters is suggested by the classical solver, which updates the variational *Ansatz* in the quantum circuit. This process is repeated to simulate the time evolution. The variational approach requires an explicit choice of the *Ansatz*, but also has the potential for compact and efficient circuits.

4 Applications



Figure 7: Quantum algorithms for the treatment of dynamics. (a) The excited-state populations (blue) and non-Markovianity witness (orange) corresponding to Markovian dynamics of a two-level system in a zero-temperature amplitude damping channel. Reprinted with permission from Ref. 44. (b) Average magnetization over time of a dissipative Transverse-Field Ising Model using a quantum algorithm based on Imaginary Time Evolution. Reprinted with permission from Ref. 47. (c) Excitonic population in the Fenna-Matthews-Olson complex over time using the generalized Sz.-Nagy dilation algorithm. Reprinted with permission from Ref. 97. (d) The singlet yield of a radical pair mechanism of avian navigation undergoing different rates of dissipation using an SVD-based algorithm. Reprinted with permission from Ref. 99. (e) Ground-state populations of a parity-time symmetric system in the unbroken (top) and broken (bottom) PT-symmetric regimes using a dilation method. Reprinted with permission from Ref. 183. (f) Electronic population difference between the donor and acceptor over time in the spin-boson model using a generalized Sz.-Nagy dilation algorithm. Reprinted with permission from Ref. 110. Copyright 2023 American Chemical Society.

In this section, we briefly review several applications of open quantum system algorithms on quantum devices, starting with a few benchmark systems. Figure 7(a) and (b) show a two-level system in an amplitude damping channel and a two-side transverse-field Ising model, respectively.

One application is the Fenna-Matthews-Olson (FMO) complex, a widely studied biochemical complex found in green-sulfur bacteria.^{184–193} The FMO complex is responsible for the efficient

exciton energy transfer from the antenna structure to the reaction center in the photosynthetic light harvesting process.^{194,195} Studied as a model system, the detailed understanding of this trimericpigment protein complex can become valuable for understanding other light-harvesting complexes and for designing artificial photovoltaic systems.^{196,197} In the first quantum simulation of the FMO dynamics, a dilation-based algorithm was used to study the FMO complex, as shown in Figure 7(c).⁹⁷ In the study, a Lindblad master equation with five states and seven elementary physical processes was constructed using a simplified but sufficient model provided by an earlier study showing the existence of multiple redundant pathways in the chromophore systems for transferring exciton energy to the reaction center.^{97,198–200} Recently, the entire seven-site system was also simulated for 2000 fs using the singular value decomposition (SVD) of the Lindblad superoperator for the system.⁹⁹ Another study examined this complex, calculating the SVD of the Kraus operators using a Walsh series representation.^{201,202}

To explain the unusual ability of birds to detect their orientation with respect to the Earth's magnetic field—an avian compass, the radical pair mechanism (RPM) has been proposed, which involves the reaction of a pair of radicals being correlated with each other as they are affected by magnetic fields.^{203,204} In the RPM, molecules in the bird's eyes are excited by photons with certain wavelengths to generate a radical pair in the singlet state. The radical pair then converts between the singlet and triplet states as affected by both the external magnetic field and the internal nuclear spin couplings. Finally, the singlet and triplet states generate different chemical products that are sensed by the bird for directional perception. By modeling two electron spins, one nuclear spin, and two "shelving states" representing the chemical yields of singlet and triplet states, the Lindblad master equation for the RPM of the avian compass' dynamics over time, in addition to the difference in the dynamics due to varying levels of environmental noise, with sample results shown using the SVD-based algorithm in Fig. 7(d).^{98,99}

A third application is a unique example of an open quantum system that simultaneously preserves parity and time (PT) symmetry.²⁰⁶ Despite their non-Hermiticity, these systems can exhibit real

eigenspectra within certain parameter regimes. In other regimes, these systems can possess complex spectra which results in intriguing topological features, ^{207,208} such as trifold knots and braiding of the eigenstates. ²⁰⁹ While analog simulation has been used to investigate these systems, ^{210,211} dilation-based algorithms have also been successful in simulating all regimes of these dynamics on quantum processors, as shown in Figure 7(e). ^{183,212}

The algorithms discussed in this review have been generalized to capture dynamics beyond the Markovian approximation, although these approaches are still in their infancy. The first conceptual approach relied on the Suzuki-Lie-Trotter decomposition of the system propagator, followed by a dilated implementation of the necessary unitaries.⁴⁰ Other approaches include the use of the non-Markovian witness formulation and the Sz.-Nagy dilation applied to the Jaynes-Cummings model,^{44,108,109} the latter of which was also applied with the generalized master equation to consider the dynamics in a spin-boson model, as depicted in Figure 7(f).¹¹⁰

5 Conclusions and Outlook

Quantum computing holds the promise to transform our ability to solve computational problems in science and engineering that are essential to pivotal advances in society. Areas that have been targeted for quantum computing applications include optimization problems, financial modeling, artificial intelligence and machine learning, cryptography and cybersecurity, and quantum chemistry. Quantum computing in quantum chemistry may lead to markedly more efficient approaches to simulating the structure and dynamics of many-electron molecules and materials. Such improvements to the current limitations of classical computers in molecular modeling would lead to significant advances in drug discovery and design, climate modeling, energy, and material science. The vast majority of quantum computing algorithms for quantum chemistry have focused on the description and prediction of closed quantum systems, and yet most real-world systems in the applications above involve open quantum systems. This review article provides an overview of the first steps towards realizing efficient quantum computing algorithms for treating open quantum systems.

This review begins with an overview of the theory of open quantum systems in Section 2 that is applicable to both classical and quantum algorithms. The overview provides a terse but in-depth perspective on different theoretical frameworks for open quantum systems, including both Markovian and non-Markovian dynamics. Second, Section 3 discusses pioneering work in the quantum simulation of open quantum systems. Several general methodological frameworks are covered, including block-encoding techniques, density matrix purification, imaginary time evolution, and variational approaches. Third, demonstrations of these quantum algorithms in problems ranging from quantum biology to quantum materials are summarized in Section 4, laying the groundwork for applications to a range of important problems in science and society. Despite advances made to date, this article describes only the beginning of a new frontier in the quantum simulation of open quantum systems. We hope that it will serve as a resource and inspiration for future advances that will take us closer to realizing the potential of quantum computing in chemistry.

6 Acknowledgments

D.A.M., P.N., and S.K. gratefully acknowledge funding from the Department of Energy, Office of Basic Energy Sciences, Grant No. DE-SC0019215. D.A.M. and P.N. acknowledge the U.S. National Science Foundation (NSF) Grant No. DMR-2037783, and D.A.M. acknowledges the U.S. NSF Grant No. CHE-2155082 and the NSF QuBBE Quantum Leap Challenge Institute (NSF OMA-2121044). S.K. acknowledges the financial support of the National Science Foundation under award number 2124511, CCI Phase I: NSF Center for Quantum Dynamics on Modular Quantum Devices (CQD-MQD) and financial support from the Quantum Science Center, a quantum partnership funded by the U.S. Department of Energy (DOE). K.H.M. acknowledges start-up funding from Washington University in St. Louis and the University of Minnesota. L.H.D.G. acknowledges that this material is based upon work supported by the U.S. Department of Energy, Office of Science,

Office of Advanced Scientific Computing Research, Department of Energy Computational Science Graduate Fellowship under Award Number DE-SC0024386. I.A. acknowledges support from the NSF Graduate Research Fellowship Program under Grant No. 2140001.

References

- Breuer, H.-P.; Petruccione, F. *The Theory of Open Quantum Systems*; Oxford University Press, 2007.
- (2) Rivas, A.; Huelga, S. F. Open quantum systems; Springer, 2012; Vol. 10.
- (3) Breuer, H.-P.; Laine, E.-M.; Piilo, J.; Vacchini, B. Colloquium: Non-Markovian dynamics in open quantum systems. *Rev. Mod. Phys.* **2016**, *88*, 021002.
- (4) Ángel Rivas,; Huelga, S. F.; Plenio, M. B. Quantum non-Markovianity: characterization, quantification and detection. *Reports on Progress in Physics* **2014**, *77*, 094001.
- (5) Li, C.-F.; Guo, G.-C.; Piilo, J. Non-Markovian quantum dynamics: What does it mean? *Europhysics Letters* 2019, 127, 50001.
- (6) Li, C.-F.; Guo, G.-C.; Piilo, J. Non-Markovian quantum dynamics: What is it good for? *Europhysics Letters* 2020, 128, 30001.
- (7) Kraus, B.; Büchler, H. P.; Diehl, S.; Kantian, A.; Micheli, A.; Zoller, P. Preparation of entangled states by quantum Markov processes. *Physical Review A* **2008**, *78*, 042307.
- (8) Diehl, S.; Micheli, A.; Kantian, A.; Kraus, B.; Büchler, H. P.; Zoller, P. Quantum states and phases in driven open quantum systems with cold atoms. *Nature Physics* **2008**, *4*, 878–883.
- (9) Verstraete, F.; Wolf, M. M.; Ignacio Cirac, J. Quantum computation and quantum-state engineering driven by dissipation. *Nature Physics* **2009**, *5*, 633–636.

- (10) Head-Marsden, K.; Flick, J.; Ciccarino, C. J.; Narang, P. Quantum information and algorithms for correlated quantum matter. *Chemical Reviews* **2021**, *121*, 3061–3120.
- (11) Olivera-Atencio, M. L.; Lamata, L.; Casado-Pascual, J. Benefits of open quantum systems for quantum machine learning. *Advanced Quantum Technologies* **2023**, *n/a*, 2300247.
- (12) Gisin, N.; Ribordy, G.; Tittel, W.; Zbinden, H. Quantum cryptography. *Reviews of Modern Physics* 2002, 74, 145–195.
- (13) Kolkowitz, S.; Bromley, S. L.; Bothwell, T.; Wall, M. L.; Marti, G. E.; Koller, A. P.; Zhang, X.; Rey, A. M.; Ye, J. Spin–orbit-coupled fermions in an optical lattice clock. *Nature* 2016, *542*, 66–70.
- (14) Martin, M. J.; Bishof, M.; Swallows, M. D.; Zhang, X.; Benko, C.; von Stecher, J.; Gorshkov, A. V.; Rey, A. M.; Ye, J. A Quantum Many-Body Spin System in an Optical Lattice Clock. *Science* **2013**, *341*, 632–636.
- (15) Georgescu, I.; Ashhab, S.; Nori, F. Quantum simulation. *Reviews of Modern Physics* 2014, 86, 153–185.
- (16) Yunger Halpern, N.; Faist, P.; Oppenheim, J.; Winter, A. Microcanonical and resourcetheoretic derivations of the thermal state of a quantum system with noncommuting charges. *Nature Communications* 2016, 7.
- (17) Wiseman, H. M.; Milburn, G. J. *Quantum Measurement and Control*; Cambridge University Press, 2009.
- (18) Pechen, A.; Rabitz, H. Incoherent Control of Open Quantum Systems. *Journal of Mathematical Sciences* 2014, 199, 695–701.
- (19) Wu, R.; Pechen, A.; Brif, C.; Rabitz, H. Controllability of open quantum systems with Krausmap dynamics. *Journal of Physics A: Mathematical and Theoretical* **2007**, *40*, 5681–5693.

- (20) Avdic, I.; Sager-Smith, L. M.; Mazziotti, D. A. Open quantum system violates generalized Pauli constraints on quantum device. *Communications Physics* **2023**, *6*, 180.
- (21) Mohseni, M.; Omar, Y.; Engel, G. S.; Plenio, M. B. *Quantum effects in biology*; Cambridge University Press, 2014.
- (22) Mohseni, M.; Shabani, A.; Lloyd, S.; Omar, Y.; Rabitz, H. Geometrical effects on energy transfer in disordered open quantum systems. *The Journal of Chemical Physics* 2013, *138*, 204309.
- (23) Kitai, K.; Guo, J.; Ju, S.; Tanaka, S.; Tsuda, K.; Shiomi, J.; Tamura, R. Designing metamaterials with quantum annealing and factorization machines. *Physical Review Research* 2020, 2.
- (24) Nielsen, M. A.; Chuang, I. L. *Quantum computation and quantum information*; Cambridge university press, 2010.
- (25) Wilde, M. M. Quantum information theory; Cambridge university press, 2013.
- (26) Kassal, I.; Whitfield, J. D.; Perdomo-Ortiz, A.; Yung, M.-H.; Aspuru-Guzik, A. Simulating chemistry using quantum computers. *Annual review of physical chemistry* 2011, 62, 185–207.
- (27) Cao, Y., et al. Quantum Chemistry in the Age of Quantum Computing. *Chemical Reviews* 2019, *119*, 10856–10915.
- (28) Tilly, J.; Chen, H.; Cao, S.; Picozzi, D.; Setia, K.; Li, Y.; Grant, E.; Wossnig, L.; Rungger, I.; Booth, G. H.; Tennyson, J. The Variational Quantum Eigensolver: A review of methods and best practices. *Physics Reports* 2022, *986*, 1–128.
- (29) Miessen, A.; Ollitrault, P. J.; Tacchino, F.; Tavernelli, I. Quantum algorithms for quantum dynamics. *Nature Computational Science* **2023**, *3*, 25–37.

- (30) Fauseweh, B. Quantum many-body simulations on digital quantum computers: State-of-theart and future challenges. *Nature Communications* **2024**, *15*.
- (31) Cerezo, M.; Arrasmith, A.; Babbush, R.; Benjamin, S. C.; Endo, S.; Fujii, K.; McClean, J. R.; Mitarai, K.; Yuan, X.; Cincio, L., et al. Variational quantum algorithms. *Nature Reviews Physics* 2021, *3*, 625–644.
- (32) Bharti, K.; Cervera-Lierta, A.; Kyaw, T. H.; Haug, T.; Alperin-Lea, S.; Anand, A.; Degroote, M.; Heimonen, H.; Kottmann, J. S.; Menke, T., et al. Noisy intermediate-scale quantum algorithms. *Reviews of Modern Physics* **2022**, *94*, 015004.
- (33) Wecker, D.; Hastings, M. B.; Troyer, M. Progress towards practical quantum variational algorithms. *Phys. Rev. A* **2015**, *92*, 042303.
- (34) Huang, H.-L.; Wu, D.; Fan, D.; Zhu, X. Superconducting quantum computing: a review. *Science China Information Sciences* **2020**, *63*.
- (35) Bruzewicz, C. D.; Chiaverini, J.; McConnell, R.; Sage, J. M. Trapped-ion quantum computing: Progress and challenges. *Applied Physics Reviews* **2019**, *6*, 021314.
- (36) Wintersperger, K.; Dommert, F.; Ehmer, T.; Hoursanov, A.; Klepsch, J.; Mauerer, W.; Reuber, G.; Strohm, T.; Yin, M.; Luber, S. Neutral atom quantum computing hardware: performance and end-user perspective. *EPJ Quantum Technology* **2023**, *10*, 32.
- (37) Bacon, D.; Childs, A. M.; Chuang, I. L.; Kempe, J.; Leung, D. W.; Zhou, X. Universal simulation of Markovian quantum dynamics. *Phys. Rev. A* **2001**, *64*, 062302.
- (38) Wang, H.; Ashhab, S.; Nori, F. Quantum algorithm for simulating the dynamics of an open quantum system. *Physical Review A* **2011**, *83*.
- (39) Sweke, R.; Sinayskiy, I.; Bernard, D.; Petruccione, F. Universal simulation of Markovian open quantum systems. *Phys. Rev. A* **2015**, *91*, 062308.

- (40) Sweke, R.; Sanz, M.; Sinayskiy, I.; Petruccione, F.; Solano, E. Digital quantum simulation of many-body non-Markovian dynamics. *Phys. Rev. A* **2016**, *94*, 022317.
- (41) Wei, S.-J.; Ruan, D.; Long, G.-L. Duality quantum algorithm efficiently simulates open quantum systems. *Scientific Reports* **2016**, *6*, 30727.
- (42) Hu, Z.; Xia, R.; Kais, S. A quantum algorithm for evolving open quantum dynamics on quantum computing devices. *Sci. Rep.* **2020**, *10*, 3301.
- (43) Patsch, S.; Maniscalco, S.; Koch, C. P. Simulation of open-quantum-system dynamics using the quantum Zeno effect. *Physical Review Research* 2020, *2*, 023133.
- (44) Garcia-Perez, G.; Rossi, M. A. C.; Maniscalco, S. IBM Q Experience as a versatile experimental testbed for simulating open quantum systems. *NPJ Quantum Inf.* **2020**, *6*, 1.
- (45) Endo, S.; Sun, J.; Li, Y.; Benjamin, S. C.; Yuan, X. Variational Quantum Simulation of General Processes. *Phys. Rev. Lett.* **2020**, *125*, 010501.
- (46) Schlimgen, A. W.; Head-Marsden, K.; Sager, L. M.; Narang, P.; Mazziotti, D. A. Quantum Simulation of Open Quantum Systems Using a Unitary Decomposition of Operators. *Phys. Rev. Lett.* 2021, *127*, 270503.
- (47) Kamakari, H.; Sun, S.-N.; Motta, M.; Minnich, A. J. Digital Quantum Simulation of Open Quantum Systems Using Quantum Imaginary–Time Evolution. *PRX Quantum* 2022, *3*, 010320.
- (48) Suri, N.; Barreto, J.; Hadfield, S.; Wiebe, N.; Wudarski, F.; Marshall, J. Two-Unitary Decomposition Algorithm and Open Quantum System Simulation. *Quantum* **2023**, *7*, 1002.
- (49) Schlimgen, A. W.; Head-Marsden, K.; Sager-Smith, L. M.; Narang, P.; Mazziotti, D. A. Quantum state preparation and nonunitary evolution with diagonal operators. *Physical Review A* 2022, 106.

- (50) Gaikwad, A.; Arvind,; Dorai, K. Simulating open quantum dynamics on an NMR quantum processor using the Sz.-Nagy dilation algorithm. *Phys. Rev. A* **2022**, *106*, 022424.
- (51) Ding, Z.; Li, X.; Lin, L. Simulating Open Quantum Systems Using Hamiltonian Simulations. *PRX Quantum* 2024, 5, 020332.
- (52) Basile, T.; Pineda, C. Quantum simulation of Pauli channels and dynamical maps: Algorithm and implementation. *PLOS One* **2024**, *19*, e0297210.
- (53) Xuereb, J.; Campbell, S.; Goold, J.; Xuereb, A. Deterministic quantum computation with one-clean-qubit model as an open quantum system. *Physical Review A* **2023**, *107*.
- (54) Shivpuje, S.; Sajjan, M.; Wang, Y.; Hu, Z.; Kais, S. Designing Variational Ansatz for Quantum-Enabled Simulation of Non-Unitary Dynamical Evolution - An Excursion into Dicke Supperradiance. *Advanced Quantum Technologies* 2024,
- (55) Watad, T. M.; Lindner, N. H. Variational quantum algorithms for simulation of Lindblad dynamics. *Quantum Science and Technology* **2024**, *9*.
- (56) Luo, J.; Lin, K.; Gao, X. Variational Quantum Simulation of Lindblad Dynamics via Quantum State Diffusion. *Journal OF Physical Chemistry Letters* 2024, 15, 3516–3522.
- (57) Mahdian, M.; Yeganeh, H. D. Hybrid quantum variational algorithm for simulating open quantum systems with near-term devices. *Journal of Physics A Mathematical and Theoret-ical* **2020**, *53*.
- (58) Liu, H.-Y.; Sun, T.-P.; Wu, Y.-C.; Guo, G.-P. Variational Quantum Algorithms for the Steady States of Open Quantum Systems. *Chinese Physics Letters* **2021**, *38*.
- (59) Lau, J. W. Z.; Lim, K. H.; Bharti, K.; Kwek, L.-C.; Vinjanampathy, S. Convex Optimization for Nonequilibrium Steady States on a Hybrid Quantum Processor. *Physical Review Letters* 2023, 130, 240601.

- (60) Joo, J.; Spiller, T. P. Commutation simulator for open quantum dynamics. *New Journal of Physics* 2023, 25.
- (61) Suri, N.; Binder, F. C.; Muralidharan, B.; Vinjanampathy, S. Speeding up thermalisation via open quantum system variational optimisation. *The European Physical Journal Special Topics* 2018, 227, 203–216.
- (62) Santos, S.; Song, X.; Savona, V. Low-Rank Variational Quantum Algorithm for the Dynamics of Open Quantum Systems. *arXiv* 2024,
- (63) Lee, C. K.; Patil, P.; Zhang, S.; Hsieh, C. Y. Neural-network variational quantum algorithm for simulating many-body dynamics. *Physical Review Research* **2021**, *3*, 023095.
- (64) Ollitrault, P. J.; Jandura, S.; Miessen, A.; Burghardt, I.; Martinazzo, R.; Tacchino, F.; Tavernelli, I. Quantum algorithms for grid-based variational time evolution. *Quantum* 2023, 7.
- (65) Gravina, L.; Savona, V. Adaptive variational low-rank dynamics for open quantum systems. *Phys. Rev. Res.* 2024, *6*, 023072.
- (66) Schlegel, D. S.; Minganti, F.; Savona, V. Coherent-State Ladder Time-Dependent Variational Principle for Open Quantum Systems. *arXiv* 2023,
- (67) Zhou, H.; Mao, R.; Sun, X. Hybrid algorithm simulating non-equilibrium steady states of an open quantum system. *arXiv* **2023**,
- (68) Lyu, N.; Mulvihill, E.; Soley, M. B.; Geva, E.; Batista, V. S. Tensor-Train Thermo-Field Memory Kernels for Generalized Quantum Master Equations. *Journal of Chemical Theory and Computation* **2023**, *19*, 1111–1129.
- (69) Gupta, P.; Chandrashekar, C. M. Optimal quantum simulation of open quantum systems. *arXiv* **2020**,

- (70) Barreiro, J. T.; Mueller, M.; Schindler, P.; Nigg, D.; Monz, T.; Chwalla, M.; Hennrich, M.; Roos, C. F.; Zoller, P.; Blatt, R. An open-system quantum simulator with trapped ions. *Nature* 2011, *470*, 486–491.
- (71) Mostame, S.; Huh, J.; Kreisbeck, C.; Kerman, A. J.; Fujita, T.; Eisfeld, A.; Aspuru-Guzik, A. Emulation of complex open quantum systems using superconducting qubits. *Quantum Information Processing* **2017**, *16*.
- (72) Su, H.-Y.; Li, Y. Quantum algorithm for the simulation of open-system dynamics and thermalization. *Phys. Rev. A* 2020, *101*, 012328.
- (73) Sun, S.; Shih, L.-C.; Cheng, Y.-C. Efficient Quantum Simulation of Open Quantum System Dynamics on Noisy Quantum Computers. 2021; https://arxiv.org/abs/2106.12882.
- (74) Burger, A.; Kwek, L. C.; Poletti, D. Digital Quantum Simulation of the Spin-Boson Model under Markovian Open-System Dynamics. *Entropy* 2022, 24.
- (75) Wang, A.; Zhang, J.; Li, Y. Error-mitigated deep-circuit quantum simulation of open systems: Steady state and relaxation rate problems. *Physical Review Research* 2022, *4*.
- (76) Cattaneo, M.; Rossi, M. A. C.; Garcia-Perez, G.; Zambrini, R.; Maniscalco, S. Quantum Simulation of Dissipative Collective Effects on Noisy Quantum Computers. *PRX Quantum* 2023, *4*.
- (77) Zanetti, M. S.; Pinto, D. F.; Basso, M. L. W.; Maziero, J. Simulating noisy quantum channels via quantum state preparation algorithms. *Journal of Physics B-Atomic Molecular and Optical Physics* **2023**, *56*.
- (78) Leppaekangas, J.; Vogt, N.; Fratus, K. R.; Bark, K.; Vaitkus, J. A.; Stadler, P.; Reiner, J.-M.; Zanker, S.; Marthaler, M. Quantum algorithm for solving open-system dynamics on quantum computers using noise. *Physical Review A* 2023, *108*.

- (79) Guimaraes, J. D.; Lim, J.; Vasilevskiy, M. I.; Huelga, S. F.; Plenio, M. B. Noise-Assisted Digital Quantum Simulation of Open Systems Using Partial Probabilistic Error Cancellation. *PRX Quantum* 2023, 4.
- (80) Guimaraes, J. D.; Vasilevskiy, M. I.; Barbosa, L. S. Digital quantum simulation of nonperturbative dynamics of open systems with orthogonal polynomials. *Quantum* **2024**, *8*.
- (81) Del Re, L.; Rost, B.; Foss-Feig, M.; Kemper, A. F.; Freericks, J. K. Robust Measurements of n-Point Correlation Functions of Driven-Dissipative Quantum Systems on a Digital Quantum Computer. *Physical Review Letters* 2024, 132.
- (82) Childs, A. M.; Li, T. Efficient Simulation of Sparse Markovian Quantum Dynamics. *Quantum Info. Comput.* 2017, *17*, 901–947.
- (83) Li, X.; Wang, C. Succinct Description and Efficient Simulation of Non-Markovian Open Quantum Systems. *Communications in Mathematical Physics* 2023, 401, 147–183.
- (84) Li, X.; Wang, C. Simulating Markovian open quantum systems using higher-order series expansion. **2023**,
- (85) Cosacchi, M.; Cygorek, M.; Ungar, F.; Barth, A. M.; Vagov, A.; Axt, V. M. Path-integral approach for nonequilibrium multitime correlation functions of open quantum systems coupled to Markovian and non-Markovian environments. *Physical Review B* 2018, 98.
- (86) Liu, J.; Segal, D.; Hanna, G. Hybrid quantum-classical simulation of quantum speed limits in open quantum systems. *Journal of Physics A-Mathematical and Theoretical* **2019**, *52*.
- (87) Cygorek, M.; Cosacchi, M.; Vagov, A.; Axt, V. M.; Lovett, B. W.; Keeling, J.; Gauger, E. M. Simulation of open quantum systems by automated compression of arbitrary environments. *Nature Physics* 2022, 18, 662.
- (88) Andreadakis, F.; Anand, N.; Zanardi, P. Scrambling of algebras in open quantum systems. *Physical Review A* 2023, 107.

- (89) Régent, F.-M. L.; Rouchon, P. Adiabatic elimination for composite open quantum systems: reduced model formulation and numerical simulations. *arXiv* 2023,
- (90) Müller, M.; Hammerer, K.; Zhou, Y. L.; Roos, C. F.; Zoller, P. Simulating open quantum systems: from many-body interactions to stabilizer pumping. *New Journal of Physics* 2011, *13*, 085007.
- (91) Ramusat, N.; Savona, V. A quantum algorithm for the direct estimation of the steady state of open quantum systems. *Quantum* **2021**, *5*.
- (92) Mahdian, M.; Yeganeh, H. D. Incoherent quantum algorithm dynamics of an open system with near-term devices. *Quantum Information Processing* **2020**, *19*.
- (93) Rost, B.; Jones, B.; Vyushkova, M.; Ali, A.; Cullip, C.; Vyushkov, A.; Nabrzyski, J. Simulation of Thermal Relaxation in Spin Chemistry Systems on a Quantum Computer Using Inherent Qubit Decoherence. 2020; https://arxiv.org/abs/2001.00794.
- (94) Tolunay, M.; Liepuoniute, I.; Vyushkova, M.; Jones, B. A. Hamiltonian simulation of quantum beats in radical pairs undergoing thermal relaxation on near-term quantum computers. *Phys. Chem. Chem. Phys.* 2023, 25, 15115–15134.
- (95) Mostame, S.; Rebentrost, P.; Eisfeld, A.; Kerman, A. J.; Tsomokos, D. I.; Aspuru-Guzik, A. Quantum simulator of an open quantum system using superconducting qubits: exciton transport in photosynthetic complexes. *New Journal of Physics* 2012, *14*.
- (96) Gupta, P.; Chandrashekar, C. M. Digital quantum simulation framework for energy transport in an open quantum system. *New Journal of Physics* **2020**, 22.
- (97) Hu, Z.; Head-Marsden, K.; Mazziotti, D. A.; Narang, P.; Kais, S. A general quantum algorithm for open quantum dynamics demonstrated with the Fenna-Matthews-Olson complex. *Quantum* 2022, *6*, 726.

- (98) Zhang, Y.; Hu, Z.; Wang, Y.; Kais, S. Quantum Simulation of the Radical Pair Dynamics of the Avian Compass. *The Journal of Physical Chemistry Letters* **2023**, *14*, 832–837.
- (99) Oh, E. K.; Krogmeier, T. J.; Schlimgen, A. W.; Head-Marsden, K. Singular Value Decomposition Quantum Algorithm for Quantum Biology. ACS Physical Chemistry Au 0, 0, null.
- (100) Sun, S.; Shih, L.-C.; Cheng, Y.-C. Efficient quantum simulation of open quantum system dynamics on noisy quantum computers. *Physica Scripta* **2024**, *99*.
- (101) Rost, B.; Del Re, L.; Earnest, N.; Kemper, A. F.; Jones, B.; Freericks, J. K. Demonstrating robust simulation of driven-dissipative problems on near-term quantum computers. 2021; https://arxiv.org/abs/2108.01183.
- (102) Schlimgen, A. W.; Head-Marsden, K.; Sager, L. M.; Narang, P.; Mazziotti, D. A. Quantum simulation of the Lindblad equation using a unitary decomposition of operators. *Phys. Rev. Res.* 2022, *4*, 023216.
- (103) Tornow, S.; Gehrke, W.; Helmbrecht, U. Non-equilibrium dynamics of a dissipative two-site Hubbard model simulated on IBM quantum computers. *Journal of Physics A: Mathematical and Theoretical* 2022, 55, 245302.
- (104) Tan, A. T. K.; Sun, S.-N.; Tazhigulov, R. N.; Chan, G. K.-L.; Minnich, A. J. Realizing symmetry-protected topological phases in a spin-1/2 chain with next-nearest-neighbor hopping on superconducting qubits. *Phycial Review A* 2023, 107.
- (105) Liang, J.-M.; Lv, Q.-Q.; Wang, Z.-X.; Fei, S.-M. Assisted quantum simulation of open quantum systems. *IScience* **2023**, *26*.
- (106) Jong, W. A. d.; Lee, K.; Mulligan, J.; P loskoń, M.; Ringer, F.; Yao, X. Quantum simulation of nonequilibrium dynamics and thermalization in the Schwinger model. *Physical Review D* 2022, *106*, 054508.

- (107) Del Re, L.; Rost, B.; Kemper, A. F.; Freericks, J. K. Driven-dissipative quantum mechanics on a lattice: Simulating a fermionic reservoir on a quantum computer. *Phys. Rev. B* 2020, *102*, 125112.
- (108) Head-Marsden, K.; Krastanov, S.; Mazziotti, D. A.; Narang, P. Capturing non-Markovian dynamics on near-term quantum computers. *Phys. Rev. Res.* **2021**, *3*, 013182.
- (109) Warren, S.; Wang, Y.; Benavides-Riveros, C. L.; Mazziotti, D. A. Exact Ansatz of Fermion-Boson Systems for a Quantum Device. *arXiv* 2024,
- (110) Wang, Y.; Mulvihill, E.; Hu, Z.; Lyu, N.; Shivpuje, S.; Liu, Y.; Soley, M. B.; Geva, E.; Batista, V. S.; Kais, S. Simulating Open Quantum System Dynamics on NISQ Computers with Generalized Quantum Master Equations. *Journal of Chemical Theory and Computation* 2023, *19*, 4851–4862.
- (111) Jagadish, V.; Petruccione, F. An Invitation to Quantum Channels. Quanta 2018, 7, 54-67.
- (112) Pechukas, P. Reduced Dynamics Need Not Be Completely Positive. *Phys. Rev. Lett.* 1994, 73, 1060–1062.
- (113) Carteret, H. A.; Terno, D. R.; Życzkowski, K. Dynamics beyond completely positive maps: Some properties and applications. *Phys. Rev. A* 2008, 77, 042113.
- (114) Sargolzahi, I. Positivity of the assignment map implies complete positivity of the reduced dynamics. *Quantum Information Processing* **2020**, *19*, 310.
- (115) Manzano, D. A short introduction to the Lindblad master equation. *AIP Advances* 2020, *10*, 025106.
- (116) Alicki, R. Comment on "Reduced Dynamics Need Not Be Completely Positive". *Phys. Rev. Lett.* 1995, 75, 3020–3020.
- (117) Pechukas, P. Pechukas Replies. Phys. Rev. Lett. 1995, 75, 3021–3021.

- (118) Levy, E.; Shalit, O. M. Dilation theory in finite dimensions: The possible, the impossible and the unknown. *Rocky Mountain Journal of Mathematics* **2014**, *44*, 203 221.
- (119) Ticozzi, F.; Viola, L. Quantum and classical resources for unitary design of open-system evolutions. *Quantum Science and Technology* **2017**, *2*, 034001.
- (120) Havel, T. F. Robust procedures for converting among Lindblad, Kraus and matrix representations of quantum dynamical semigroups. *J. Math. Phys.* **2003**, *44*, 534–557.
- (121) Kunold, A. Vectorization of the density matrix and quantum simulation of the von Neumann equation of time-dependent Hamiltonians. *Physica Scripta* **2024**, *99*.
- (122) Byrd, M.; Ceballos, R.; Chitambar, E. Open system quantum evolution and the assumption of complete positivity (A Tutorial). *International Journal of Quantum Information* **2016**, *14*.
- (123) Alicki, R.; Lendi, K. *Quantum dynamical semigroups and applications*; Springer, 2007; Vol. 717.
- (124) Cubitt, T. S.; Eisert, J.; Wolf, M. M. The Complexity of Relating Quantum Channels to Master Equations. *Communcations in Mathematical Physics* **2012**, *310*, 383–418.
- (125) Gorini, V.; Kossakowski, A.; Sudarshan, E. C. G. Completely positive dynamical semigroups of N-level systems. *Journal of Mathematical Physics* **1976**, *17*, 821–825.
- (126) Lindblad, G. On the generators of quantum dynamical semigroups. Communications in Mathematical Physics 1976, 48, 119–130.
- (127) Gorini, V.; Kossakowski, A.; Sudarshan, E. C. G. Completely positive dynamical semigroups of N-level systems. *Journal of Mathematical Physics* **2008**, *17*, 821–825.
- (128) Redfield, A. G. In Advances in Magnetic Resonance; Waugh, J. S., Ed.; Advances in Magnetic and Optical Resonance; Academic Press, 1965; Vol. 1; pp 1–32.

- (129) Nakajima, S. On quantum theory of transport phenomena: steady diffusion. Progress of Theoretical Physics 1958, 20, 948–959.
- (130) Zwanzig, R. Ensemble method in the theory of irreversibility. *The Journal of Chemical Physics* **1960**, *33*, 1338–1341.
- (131) Haake, F. Springer Tracts in Modern Physics: Ergebnisse der exakten Naturwissenschaftenc;
 Volume 66; Springer, 1973; pp 98–168.
- (132) Tokuyama, M.; Mori, H. Statistical-Mechanical Theory of Random Frequency Modulations and Generalized Brownian Motions*). *Progress of Theoretical Physics* **1976**, *55*, 411–429.
- (133) Shibata, F.; Takahashi, Y.; Hashitsume, N. A generalized stochastic liouville equation. Non-Markovian versus memoryless master equations. *Journal of Statistical Physics* 1977, *17*, 171–187.
- (134) de Vega, I.; Alonso, D. Dynamics of non-Markovian open quantum systems. *Reviews of Modern Physics* 2017, 89.
- (135) Sweke, R.; Sinayskiy, I.; Petruccione, F. Simulation of single-qubit open quantum systems.*Phys. Rev. A* 2014, *90*, 022331.
- (136) David, I.; Sinayskiy, I.; Petruccione, F. Digital Simulation of Single Qubit Markovian Open Quantum Systems: A Tutorial. *Quanta* 2023, *12*, 131–163.
- (137) Buscemi, F.; D'Ariano, G. M.; Sacchi, M. F. Physical realizations of quantum operations. *Phys. Rev. A* 2003, 68, 042113.
- (138) Childs, A. M.; Wiebe, N. Hamiltonian simulation using linear combinations of unitary operations. *Quantum Info. Comput.* **2012**, *12*, 901–924.
- (139) Smart, S. E.; Mazziotti, D. A. Verifiably exact solution of the electronic Schrödinger equation on quantum devices. *Physical Review A* 2024, *109*, 022802.

- (140) Wang, Y.; Mazziotti, D. A. Electronic excited states from a variance-based contracted quantum eigensolver. *Physical Review A* 2023, *108*, 022814.
- (141) Smart, S. E.; Boyn, J.-N.; Mazziotti, D. A. Resolving correlated states of benzyne with an error-mitigated contracted quantum eigensolver. *Physical Review A* **2022**, *105*.
- (142) Smart, S. E.; Mazziotti, D. A. Quantum solver of contracted eigenvalue equations for scalable molecular simulations on quantum computing devices. *Physical Review Letters* 2021, *126*, 070504.
- (143) Mazzola, G. Quantum computing for chemistry and physics applications from a Monte Carlo perspective. *Journal of Chemical Physics* 2024, *160*.
- (144) Nagy, A.; Savona, V. Driven-dissipative quantum Monte Carlo method for open quantum systems. *Physical Review A* **2018**, *97*.
- (145) Kadowaki, T. Dynamics of open quantum systems by interpolation of von Neumann and classical master equations, and its application to quantum annealing. *Physical Review A* 2018, 97.
- (146) Peetz, J.; Smart, S. E.; Tserkis, S.; Narang, P. Simulation of Open Quantum Systems via Low-Depth Convex Unitary Evolutions. https://arxiv.org/abs/2307.14325v2, 2023.
- (147) Rosgen, B. Additivity and Distinguishability of Random Unitary Channels. *Journal of Mathematical Physics* **2008**, *49*, 102107.
- (148) Schlimgen, A. W.; Head-Marsden, K.; Sager-Smith, L. M.; Narang, P.; Mazziotti, D. A. Quantum Simulation of Open Quantum Systems Using Density-Matrix Purification. 2022.
- (149) Delgado-Granados, L. H.; Warren, S.; Mazziotti, D. A. Unitary Dynamics for Open Quantum Systems with Density-Matrix Purification. 2024.

- (150) Zong, Z.; Huai, S.; Cai, T.; Jin, W.; Zhan, Z.; Zhang, Z.; Bu, K.; Sui, L.; Fei, Y.; Zheng, Y.; Zhang, S.; Wu, J.; Yin, Y. Determination of molecular energies via variational-based quantum imaginary time evolution in a superconducting qubit system. *Science China-Physics Mechanics & Astronomy* **2024**, 67.
- (151) Kleinmann, M.; Kampermann, H.; Meyer, T.; Bruß, D. Physical purification of quantum states. *Phys. Rev. A* **2006**, *73*, 062309.
- (152) Bassi, A.; Ghirardi, G. A general scheme for ensemble purification. *Physics Letters A* 2003, 309, 24–28.
- (153) Hughston, L. P.; Jozsa, R.; Wootters, W. K. A complete classification of quantum ensembles having a given density matrix. *Physics Letters A* **1993**, *183*, 14–18.
- (154) Motta, M.; Sun, C.; Tan, A. T. K.; O'Rourke, M. J.; Ye, E.; Minnich, A. J.; Brandão, F. G. S. L.; Chan, G. K.-L. Determining eigenstates and thermal states on a quantum computer using quantum imaginary time evolution. *Nature Physics* 2020, *16*, 205–210.
- (155) Cao, C.; An, Z.; Hou, S.-Y.; Zhou, D. L.; Zeng, B. Quantum imaginary time evolution steered by reinforcement learning. *Communications Physics* **2022**, *5*.
- (156) Mao, Y.; Chaudhary, M.; Kondappan, M.; Shi, J.; Ilo-Okeke, E. O.; Ivannikov, V.; Byrnes, T. Measurement-Based Deterministic Imaginary Time Evolution. *Physical Review Letters* 2023, 131.
- (157) Lin, S.-H.; Dilip, R.; Green, A. G.; Smith, A.; Pollmann, F. Real- and Imaginary-Time Evolution with Compressed Quantum Circuits. *PRX Quantum* 2021, 2.
- (158) Inoue, K.; Fukumoto, Y. Typical Purification Reproducing the Time Evolution of an Open Quantum System. *arXiv* **2018**,
- (159) McArdle, S.; Jones, T.; Endo, S.; Li, Y.; Benjamin, S. C.; Yuan, X. Variational ansatz-based quantum simulation of imaginary time evolution. *npj Quantum Information* **2019**, *5*, 75.

- (160) Shirakawa, T.; Seki, K.; Yunoki, S. Discretized quantum adiabatic process for free fermions and comparison with the imaginary-time evolution. *Phys. Rev. Res.* **2021**, *3*, 013004.
- (161) Nishi, H.; Kosugi, T.; Matsushita, Y.-i. Implementation of quantum imaginary-time evolution method on NISQ devices by introducing nonlocal approximation. *npj Quantum Information* 2021, 7, 85.
- (162) Lin, S.-H.; Dilip, R.; Green, A. G.; Smith, A.; Pollmann, F. Real- and Imaginary-Time Evolution with Compressed Quantum Circuits. *PRX Quantum* 2021, 2, 010342.
- (163) Kramer, P.; Saraceno, M. Geometry of the time-dependent variational principle in quantum mechanics; Springer, 1981.
- (164) Broeckhove, J.; Lathouwers, L.; Kesteloot, E.; Van Leuven, P. On the equivalence of timedependent variational principles. *Chemical physics letters* **1988**, *149*, 547–550.
- (165) Pedersen, T. B.; Koch, H. On the time-dependent Lagrangian approach in quantum chemistry. *The Journal of chemical physics* **1998**, *108*, 5194–5204.
- (166) Kerman, A. K.; Koonin, S. E. Hamiltonian formulation of time-dependent variational principles for the many-body system. *Annals of Physics* **1976**, *100*, 332–358.
- (167) Yuan, X.; Endo, S.; Zhao, Q.; Li, Y.; Benjamin, S. C. Theory of variational quantum simulation. *Quantum* **2019**, *3*, 191.
- (168) Hackl, L.; Guaita, T.; Shi, T.; Haegeman, J.; Demler, E.; Cirac, J. I. Geometry of variational methods: dynamics of closed quantum systems. *SciPost Physics* **2020**, *9*, 048.
- (169) Haegeman, J.; Osborne, T. J.; Verstraete, F. Post-matrix product state methods: To tangent space and beyond. *Phys. Rev. B* 2013, 88, 075133.
- (170) Dirac, P. A. M. Note on exchange phenomena in the Thomas atom. *Mathematical Proceedings* of the Cambridge Philosophical Society **1930**, 26, 376–385.

- (171) Frenkel, J. Wave Mechanics: Advanced General Theory; 1934.
- (172) Langhoff, P.; Epstein, S.; Karplus, M. Aspects of time-dependent perturbation theory. *Reviews of Modern Physics* **1972**, *44*, 602.
- (173) McLachlan, A. A variational solution of the time-dependent Schrodinger equation. *Molecular Physics* 1964, 8, 39–44.
- (174) Messina, M.; Garrett, B. C.; Schenter, G. K. Variational solutions for the thermal and real time propagator using the McLachlan variational principle. *The Journal of chemical physics* **1994**, *100*, 6570–6577.
- (175) Yao, Y.-X.; Gomes, N.; Zhang, F.; Wang, C.-Z.; Ho, K.-M.; Iadecola, T.; Orth, P. P. Adaptive variational quantum dynamics simulations. *PRX Quantum* **2021**, *2*, 030307.
- (176) Lee, C.-K.; Hsieh, C.-Y.; Zhang, S.; Shi, L. Variational quantum simulation of chemical dynamics with quantum computers. *Journal of Chemical Theory and Computation* 2022, 18, 2105–2113.
- (177) Miessen, A.; Ollitrault, P. J.; Tavernelli, I. Quantum algorithms for quantum dynamics: A performance study on the spin-boson model. *Physical Review Research* **2021**, *3*, 043212.
- (178) Alghassi, H.; Deshmukh, A.; Ibrahim, N.; Robles, N.; Woerner, S.; Zoufal, C. A variational quantum algorithm for the Feynman-Kac formula. *Quantum* **2022**, *6*, 730.
- (179) Nakaji, K.; Endo, S.; Matsuzaki, Y.; Hakoshima, H. Measurement optimization of variational quantum simulation by classical shadow and derandomization. *Quantum* **2023**, *7*, 995.
- (180) Chen, H.; Gomes, N.; Niu, S.; de Jong, W. A. Adaptive variational simulation for open quantum system dynamics. *Quantum* **2024**, *8*.
- (181) Linteau, D.; Barison, S.; Lindner, N. H.; Carleo, G. Adaptive projected variational quantum dynamics. *Phys. Rev. Res.* **2024**, *6*, 023130.

- (182) Gulliksson, M.; Wedin, P.-Å.; Wei, Y. Perturbation identities for regularized Tikhonov inverses and weighted pseudoinverses. *BIT Numerical Mathematics* **2000**, *40*, 513–523.
- (183) Dogra, S.; Melnikov, A. A.; Paraoanu, G. S. Quantum simulation of parity-time symmetry breaking with a superconducting quantum processor. *Communications Physics* **2021**, *4*.
- (184) Fenna, R. E.; Matthews, B. W. Chlorophyll arrangement in a bacteriochlorophyll protein from Chlorohium limicola. *Nature* 1975, 258, 1476–4687.
- (185) Ishizaki, A.; Fleming, G. Unified treatment of quantum coherent and incoherent hopping dynamics in electronic energy transfer: Reduced hierarchy equation approach. *The Journal* of Chemical Physics 2009, 130, 234111.
- (186) Zhu, J., et al. Multipartite quantum entanglement evolution in photosynthetic complexes. *The Journal of Chemical Physics* **2012**, *137*, 074112.
- (187) Thyrhaug, E., et al. Identification and characterization of diverse coherences in the Fenna–Matthews–Olson complex. *Nature Chemistry* **2018**, *10*, 780–786.
- (188) Irgen-Gioro, S., et al. Electronic coherence lifetimes of the Fenna–Matthews–Olson complex and light harvesting complex II. *Chemical Science* **2019**, *10*, 10503–10509.
- (189) Oh, S.; Coker, D.; Hutchinson, D. Optimization of energy transport in the Fenna-Matthews-Olson complex via site-varying pigment-protein interactions. *The Journal of Chemical Physics* 2019, 150, 085102.
- (190) Kim, Y., et al. Predictive First-Principles Modeling of a Photosynthetic Antenna Protein: The Fenna–Matthews–Olson Complex. *The Journal of Physical Chemistry Letters* 2020, *11*, 1636–1643.
- (191) Suzuki, Y., et al. Comparative study on model parameter evaluations for the energy transfer dynamics in Fenna–Matthews–Olson complex. *Chemical Physics* **2020**, *539*, 110903.

- (192) Engel, G. S.; Calhoun, T. R.; Read, E. L.; Ahn, T.-K.; Mančal, T.; Cheng, Y.-C.; Blankenship, R. E.; Fleming, G. R. Evidence for wavelike energy transfer through quantum coherence in photosynthetic systems. *Nature* 2007, 446.
- (193) Lee, H.; Cheng, Y.-C.; Fleming, G. R. Coherence Dynamics in Photosynthesis: Protein Protection of Excitonic Coherence. *Science* 2007, *316*, 1462–1465.
- (194) Sension, R. Quantum path to photosynthesis. *Nature* 2007, 446, 740–741.
- (195) Barroso-Flores, J. Evolution of the Fenna–Matthews–Olson Complex and Its Quantum Coherence Features. Which Led the Way? ACS Central Science 2017, 3, 1061–1062.
- (196) Hu, Z.; amd Fahhad H. Alharbi, G. S. E.; Kais, S. Dark states and delocalization: Competing effects of quantum coherence on the efficiency of light harvesting systems. *The Journal of Chemical Physics* 2018, *148*, 064304.
- (197) Hu, Z.; Engel, G.; Kais, S. Double-excitation manifold's effect on exciton transfer dynamics and the efficiency of coherent light harvesting. *Physical Chemistry Chemical Physics* 2018, 20, 30032–30040.
- (198) Skochdopole, N.; Mazziotti, D. Functional Subsystems and Quantum Redundancy in Photosynthetic Light Harvesting. *The Journal of Physical Chemistry Letters* **2011**, *2*, 2989–2993.
- (199) Avdic, I.; Sager-Smith, L. M.; Ghosh, I.; Wedig, O. C.; Higgins, J. S.; Engel, G. S.; Mazziotti, D. A. Quantum sensing using multiqubit quantum systems and the Pauli polytope. *Phys. Rev. Res.* 2023, *5*, 043097.
- (200) Schouten, A. O.; Sager-Smith, L. M.; Mazziotti, D. A. Exciton-Condensate-Like Amplification of Energy Transport in Light Harvesting. *PRX Energy* 2023, 2, 023002.
- (201) Seneviratne, A.; Walters, P. L.; Wang, F. Exact Non-Markovian Quantum Dynamics on the NISQ Device Using Kraus Operators. ACS Omega 2024, 9, 9666–9675.

- (202) Welch, J.; Greenbaum, D.; Mostame, S.; Aspuru-Guzik, A. Efficient quantum circuits for diagonal unitaries without ancillas. *New Journal of Physics* **2014**, *16*, 033040.
- (203) Ritz, T., et al. Resonance effects indicate a radical-pair mechanism for avian magnetic compass. *Nature* **2004**, *429*, 177–180.
- (204) Rodgers, C.; Hore, P. Chemical magnetoreception in birds: The radical pair mechanism. *Proceedings of the National Academy of Sciences* **2009**, *106*, 353–360.
- (205) Gauger, E., et al. Sustained Quantum Coherence and Entanglement in the Avian Compass. *Physical Review Letters* 2011, *106*, 040503.
- (206) Bender, C. M.; Boettcher, S. Real Spectra in Non-Hermitian Hamiltonians Having PT Symmetry. *Physical Review Letters* 1998, 80, 5243–5246.
- (207) El-Ganainy, R.; Makris, K. G.; Khajavikhan, M.; Musslimani, Z. H.; Rotter, S.; Christodoulides, D. N. Non-Hermitian physics and PT symmetry. *Nature Physics* 2018, 14, 11–19.
- (208) Wang, Z.; Chong, Y.; Joannopoulos, J. D.; Soljačić, M. Observation of unidirectional backscattering-immune topological electromagnetic states. *Nature* **2009**, *461*, 772–775.
- (209) Patil, Y. S. S.; Höller, J.; Henry, P. A.; Guria, C.; Zhang, Y.; Jiang, L.; Kralj, N.; Read, N.; Harris, J. G. E. Measuring the knot of non-Hermitian degeneracies and non-commuting braids. *Nature* **2022**, *607*, 271–275.
- (210) Wu, Y.; Liu, W.; Geng, J.; Song, X.; Ye, X.; Duan, C.-K.; Rong, X.; Du, J. Observation of parity-time symmetry breaking in a single-spin system. *Science* **2019**, *364*, 878–880.
- (211) Naghiloo, M.; Abbasi, M.; Joglekar, Y. N.; Murch, K. W. Quantum state tomography across the exceptional point in a single dissipative qubit. *Nature Physics* **2019**, *15*, 1232–1236.
- (212) Kazmina, A. S. et al. Demonstration of a parity-time-symmetry-breaking phase transition using superconducting and trapped-ion qutrits. *Physical Review A* **2024**, *109*.



Figure 8: TOC Graphics