

ON SOME RINGS OF DIFFERENTIABLE TYPE

SAYED SADIQUL ISLAM AND TONY J. PUTHENPURAKAL

ABSTRACT. Let K be a field of characteristic 0 and $S = K[x_1, \dots, x_m]/I$ be an affine domain. Consider $R = S_P$ where $P \in \text{Spec}(S)$ such that R is regular. In this paper we construct a field F which is contained in R such that

- (1) The residue field of R is a finite extension of F .
- (2) $D_F(R)$, the ring of F -linear differential operators on R is left and right Noetherian with finite global dimension.
- (3) The Bernstein class of $D_F(R)$ is closed under localization at one element of R .

We also prove a similar result for R^h , the Henselization of R . As an application we prove that $\frac{D_F(R)}{D_F(R)_P} \cong E(\kappa(P))$ where $E(\kappa(P))$ is the injective hull of the residue field of R .

1. INTRODUCTION

Let K be a field of characteristic zero, and R be a K -algebra. let $D(R) = D_K(R)$ be the ring of K -linear differential operators on R . There has been a lot of research on the finiteness properties of rings of differential operators. The following are example of R related to $D(R)$ as a left and right Noetherian with finite global dimension:

- (1) Polynomial rings over K in n variables. In this case $D(R) = A_n(K)$, the n^{th} -Weyl algebra over K . We have that global dimension of $D(R)$ is equal to n ; see [3, Chapter 2, Theorem 3.15].
- (2) Power series rings over K in n variables. In this case global dimension is n ; see [3, Chapter 3, Proposition 1.8].
- (3) The ring of convergent power series over \mathbb{C} in n variables; see [3, Page 197].
- (4) Let $(V, \pi V, K)$ be a DVR of mixed characteristic $p > 0$, and Let F denote its fraction field. Let $S = V[[x_1, \dots, x_{n+1}]] \otimes_V F$ be the tensor product of the power series ring with coefficients in V and F . Let $R = S/(f)S$ be a regular ring where $f = \pi - h$ for an element h in the square of maximal ideal of $V[[x_1, \dots, x_{n+1}]]$. In this case also $D(R)$ is Noetherian with finite global dimension; see [2, Page 4].

Date: June 11, 2024.

2020 Mathematics Subject Classification. Primary 13N10; Secondary 13N15, 13D45 .

Key words and phrases. Rings of differential operators, D-modules, local cohomology.

- (5) Localization of complete local domains of equal characteristic zero at certain elements; see [10].

However $D(R)$ is not always Noetherian. Consider in the space \mathbb{C}^3 with the coordinates x_1, x_2, x_3 the surface X defined by the equation $x_1^3 + x_2^3 + x_3^3 = 0$, then the ring of differential operator on X is not Noetherian. This example is due to Bernstein- Gelfand[1, Theorem 1]. In this article we provide a new infinite class of Noetherian domains R with $D(R)$ is both left and right Noetherian and with finite global dimension.

We now state the following results of this paper.

1.1. Setup: Let K be a field of characteristic 0 and $S = K[x_1, \dots, x_m]/I$ be an affine domain. Consider $R = S_P$ where $P \in \text{Spec}(S)$ such that R is regular. We shall show that R contains a field F with $\kappa(P) = S_P/PS_P$ a finite extension of F . Let $\text{Der}_F(R)$ be the set of F -linear derivations of R and $D_F(R)$ be the ring of F -linear differential operators on R . Our result is as follows:

Theorem 1.2. *Let R and F be as in Setup 1.1. Then,*

- (1) *The residue field of R is a finite extension of F .*
- (2) *$\text{Der}_F(R)$ is a finitely generated projective R -module of rank equals to $\dim R$.*
- (3) *The ring $D_F(R)$ is a left and right Noetherian and the Bernstein class of $D_F(R)$ is closed under localization at one element of R .*

1.3. Setup: Let R and S be as in Setup 1.1. Denote by R^h , the Henselization of R . Since $R \hookrightarrow R^h$, the field F in 1.1 is contained in R^h . We prove a similar result.

Theorem 1.4. *For the setup 1.3, we have the following;*

- (1) *The residue field of R^h is a finite extension of F .*
- (2) *$\text{Der}_F(R^h)$ is a finitely generated projective R^h -module of rank same as dimension of R^h .*
- (3) *The ring $D_F(R^h)$ is a left and right Noetherian and the Bernstein class of $D_F(R^h)$ is closed under localization at one element of R^h .*

We also prove the following result as an application of 1.2;

Theorem 1.5. *Let R and F be as in Setup 1.1. Then,*

- (1) *$H_P^{\dim R}(R) \cong E(\kappa(P))$ is a $D_F(R)$ -module in the Bernstein class.*
- (2) *$\frac{D_F(R)}{D_F(R)P} \cong E(\kappa(P))$, where $E(\kappa(P))$ is the injective hull of the residue field of R .*

Our paper comprises six sections. Section 2 provides an overview of D -modules. In section 3, we present crucial theorems essential for proving our results. Sections 4, 5, 6 and 7 are where we explain our proofs.

2. BACKGROUND ON D-MODULES

Let R be a commutative A -algebra. The ring of A -linear differential operators of R is a subring $D_A(R) \subseteq \text{Hom}_A(R, R)$ whose elements are defined inductively as follows: the differential operators of order zero are defined by the multiplication by elements of R , i.e. $D^0(R) \cong R$. Suppose we have defined operators of order $< n$. An operator $\delta \in \text{Hom}_A(R, R)$ is of order less than or equal to n if $[\delta, r] = \delta r - r\delta$ is an operator of order less than or equal to $n - 1$ for all $r \in R$. We have a filtration $D_A^0(R) \subset D_A^1(R) \subset \dots$ and ring of differential operators is defined as

$$D_A(R) = \bigcup_{m \geq 0} D_A^m(R).$$

Example 2.1. If R is either a polynomial ring $A[x_1, \dots, x_n]$ or the formal power series ring $A[[x_1, \dots, x_n]]$ with coefficients in A , then ring of A -linear differential operators is :

$$D_A(R) = R \left\langle \frac{1}{t!} \frac{d^t}{dx_i^t} \mid i = 1, \dots, n; t \in \mathbb{N} \right\rangle.$$

2.2. Let M be a $D_A(R)$ -module and $f \in R$. Then M_f is also a $D_A(R)$ -module, where the action of a differential operator $\partial \in D_A(R)$ on $\frac{v}{f^t} \in M_f$ is defined inductively as follows: If $\partial \in D_A^0(R)$ has order 0 then $\partial \cdot \frac{v}{f^t} = \frac{\partial \cdot v}{f^t}$. Now suppose that action of every element in $D_A^n(R)$ has been defined. Let $\partial \in D_A^{n+1}(R)$. Then,

$$\partial \cdot \frac{v}{f^t} = \frac{\partial \cdot v - [\partial, f^t] \cdot \frac{v}{f^t}}{f^t}$$

With this $D_A(R)$ -module structure on M_f , the map $M \rightarrow M_f$ is a morphism of $D_A(R)$ -modules.

2.3. Let $I \subseteq R$ be an ideal of R generated by $\underline{f} = f_1, f_2, \dots, f_r \in R$, and let M be an R -module. Then the Čech complex of M with respect to \underline{f} is defined by

$$\check{C}^\bullet(\underline{f}, M) : 0 \rightarrow M \rightarrow \bigoplus_i M_{f_i} \rightarrow \bigoplus_{i,j} M_{f_i f_j} \rightarrow \dots \rightarrow M_{f_1 \dots f_r} \rightarrow 0$$

where the maps on every summand are localization map upto a sign. The local cohomology module of M with support on I is defined by

$$H_I^i(M) = H^i(\check{C}^\bullet(\underline{f}, M))$$

It follows from this construction that every local cohomology module over a $D_A(R)$ -module is again a $D_A(R)$ -module.

Definition 2.4. We say that a associative ring R is filtered if there exists an ascending filtration $\Sigma_0 \subset \Sigma_1 \subset \Sigma_2 \dots$ of additive subgroups such that $1 \in \Sigma_0$, $\bigcup \Sigma_i = R$ and $\Sigma_i \Sigma_j \subset \Sigma_{i+j}$ for every i and j . We denote by $gr^\Sigma(R)$ the associated graded ring $\Sigma_0 \oplus \Sigma_1/\Sigma_0 \oplus \Sigma_2/\Sigma_1 \oplus \dots$

We have the following definition of a ring of differentiable type from [7];

Definition 2.5. A filtered ring R is a ring of differentiable type if its associated graded ring is commutative Noetherian regular with unity and pure graded dimension.

2.6. Let R be a filtered ring of differentiable type with filtration $\{\Sigma_i\}$. A filtration $\{\Gamma_i\}$ compatible with $\{\Sigma_i\}$ on a left R -module M is called good if the associated graded $gr^\Sigma(R)$ -module $gr^\Gamma(M)$ is finitely generated. It is well known that a left R -module M can be equipped with good filtration iff it is finitely generated. If M is a finitely generated left R -module we can choose some good filtration Γ on M . Now the associated graded $gr^\Sigma(R)$ -module $gr^\Gamma(M)$ is finitely generated. Then we define:

$$d_\Gamma(M) = \dim \text{Supp}(gr^\Gamma(M))$$

In general, M can be equipped with many good filtrations. But it turns out that the integer $d_\Gamma(M)$ do not depend on the choice of Γ , i.e. we have the equalities $d_\Gamma(M) = d_{\Gamma'}(M)$ for all pairs of good filtrations Γ and Γ' on M . This common integer is therefore denoted by $d(M)$ and called the dimension of R -module M .

Definition 2.7. A ring R has finite weak global dimension w if $\text{Tor}_v^R(N, M) = 0$ for all $v > w$ and every right R -module N and left R -module M , while $\text{Tor}_w^R(N, M) \neq 0$ for some pair.

Let R be a ring of differentiable type with pure graded dimension μ and assume that weak global dimension of R is w . We recall the following corollary from [3, Chapter 2, 7.2],

Corollary 2.8. $d(M) \geq \mu - w$ for every non-zero finitely generated R -module M .

Definition 2.9. The Bernstein class consists of all finitely generated R -modules M for which equality holds in the above corollary i.e. $d(M) = \mu - w$ (or if $M = 0$).

3. PRELIMINARIES

If A is a k -algebra, $\Omega_{A/k}$ denotes the module of Kähler differentials and let the universal derivation be $d_{A/k}: A \rightarrow \Omega_{A/k}$. For A -module M we have that $\text{Der}_k(A, M) \cong \text{Hom}_A(\Omega_{A/k}, M)$.

The following two results are useful:

Theorem 3.1. (First fundamental exact sequence)[6, Theorem 25.1] A composite $k \xrightarrow{f} A \xrightarrow{g} B$ of ring homomorphisms leads to an exact sequence of B -modules

$$\Omega_{A/k} \otimes_A B \xrightarrow{\alpha} \Omega_{B/k} \xrightarrow{\beta} \Omega_{B/A} \rightarrow 0$$

where the maps are given by $\alpha(d_{A/k}a \otimes b) = b d_{B/k}g(a)$ and $\beta(d_{B/k}b) = d_{B/A}b$ for $a \in A$ and $b \in B$.

Theorem 3.2. (Second fundamental exact sequence)[6, Theorem 25.2] Consider $k \xrightarrow{f} A \xrightarrow{g} B$ when g is surjective; set $\ker g = \mathfrak{m}$, so $B = A/\mathfrak{m}$. Then we have an exact sequence as follows:

$$\mathfrak{m}/\mathfrak{m}^2 \xrightarrow{\delta} \Omega_{A/k} \otimes_A B \xrightarrow{\alpha} \Omega_{B/k} \rightarrow 0$$

where δ is the B -linear map defined by $\delta(\bar{x}) = d_{A/k}x \otimes 1$.

In general $\Omega_{A/F}$ need not be finitely generated A -module. However we have the following result for $\text{Der}_F(A)$.

Theorem 3.3. [6, Theorem 30.7] (R, \mathfrak{m}) be an n -dimensional Noetherian local integral domain containing \mathbb{Q} , and $F \subset R$ be a subfield such that $\text{tr.deg}_F(R/\mathfrak{m}) = r < \infty$. Then $\text{Der}_F(R)$ is isomorphic to a submodule of R^{n+r} , and is therefore a finite R -module, with

$$\text{rank Der}_F(R) \leq \dim R + \text{tr.deg}_F(R/\mathfrak{m})$$

The following result gives a sufficient condition for $\text{Der}_F(R)$ to be free.

Theorem 3.4. [6, Theorem 30.6] Let (R, \mathfrak{m}, F) be a regular local commutative Noetherian ring with unity of dimension n containing a field F_0 . Suppose that F is an algebraic separable extension F_0 . Let \widehat{R} denote the \mathfrak{m} -adic completion of R . Let x_1, \dots, x_n be a regular system of parameters of R . Then, $\widehat{R} = F[[x_1, \dots, x_n]]$ is the power series ring with coefficients in F , and $\text{Der}_F \widehat{R}$ is a free \widehat{R} -module with basis $\partial/\partial x_i$ for $1 \leq i \leq n$. Moreover, the following conditions are equivalent:

- (1) $\partial/\partial x_i$ maps R into R for $1 \leq i \leq n$, so that they can be considered as elements of $\text{Der}_{F_0}(R)$;
- (2) $\text{Der}_{F_0}(R)$ is a free R -module of rank n ;
- (3) $\text{rank Der}_{F_0}(R) = n$.

Corollary 3.5. [8, Corollary 1.2.2.3] Under the hypotheses above, assume that any of the equivalent conditions in theorem 3.4 are satisfied. Then the following holds:

$$\widehat{R} \otimes_R \text{Der}_{F_0}(R) \cong \text{Der}_{F_0}(\widehat{R}) \cong \text{Der}_{F_0}(R) \otimes_R \widehat{R}$$

The following result is due to Lyubeznik;

Theorem 3.6. [4, Proposition 2.3] Let K be a field of characteristic 0, let $R = K[[x_1, \dots, x_n]]$ be the ring of formal power series ring in n variables over K and let \mathfrak{m} be maximal ideal of R . Then, as an R -module $\frac{D_K(R)}{D_K(R)\mathfrak{m}}$ is isomorphic to $E_R(K)$, the injective hull of the residue field of R .

Remark 3.7. Let R be an integral domain and L be quotient field of R .

- (1) Let M be R -module of rank r . Then $M \otimes L \cong L^r$. But $\text{Hom}_R(M, R) \otimes L \cong \text{Hom}_L(L^r, L) \cong L^r$. Hence $\text{Hom}_R(M, R)$ and M have the same rank.

- (2) Let M is R -module and $P \in \text{Spec}(R)$. We note that R and R_P have same quotient field, denoted as L . Since $M_P \otimes_{R_P} L \cong M \otimes_R R_P \otimes_{R_P} L \cong M \otimes_R L$, so M and M_P have the same rank.

Definition 3.8. A local homomorphism $\phi : (R, \mathfrak{m}, k) \rightarrow (S, \mathfrak{n}, l)$ of local rings is unramified provided S is essentially of finite type over R (that is, S is localization of some finitely generated R -algebra) and the following properties hold.

- (1) $\mathfrak{m}S = \mathfrak{n}$, and
(2) $S/\mathfrak{m}S$ is a finite separable field extension of R/\mathfrak{m} .

If, in addition, ϕ is flat, then we say ϕ is étale. (We say also that S is an unramified, respectively, étale extension of R .) A pointed étale neighborhood is an étale extension $(R, \mathfrak{m}, k) \rightarrow (S, \mathfrak{n}, l)$ inducing an isomorphism on residue fields.

The isomorphism classes of pointed étale neighborhoods form a direct system. The reason for this is that if $R \rightarrow S$ and $R \rightarrow T$ are pointed étale neighborhoods then there is at most one homomorphism $S \rightarrow T$ making the obvious diagram commute. We refer [5, Chapter 3] for details.

Definition 3.9. The Henselization R^h of R is the direct limit of a set of representatives of isomorphism classes of pointed étale neighborhoods of R .

Theorem 3.10. [9, Theorem 43.9] *Let (R, \mathfrak{m}) be a local ring and let R^h be its Henselization. For a finite number of elements b_i of R^h , there is an element a of $\mathfrak{m}R^h$ which is a root of a monic polynomial $f(x) = x^r + c_1x^{r-1} + \dots + c_r$ over R with $c_{r-1} \notin \mathfrak{m}$, $c_r \in \mathfrak{m}$, such that the ring $R' = R[a]_{(\mathfrak{m} + aR[a])}$ contains b_i . Furthermore R^h contains R' .*

We recall the following two results from [2].

Theorem 3.11. *Let R be a regular commutative Noetherian ring with unity that contains a field F of characteristic 0 satisfying the following conditions:*

- (1) R is equidimensional of dimension n ;
(2) every residual field with respect to a maximal ideal is an algebraic extension of F ;
(3) $\text{Der}_F(R)$ is a finitely generated projective R -module of rank n such that $R_{\mathfrak{m}} \otimes_R \text{Der}_F(R) = \text{Der}_F(R_{\mathfrak{m}})$ for each maximal ideal \mathfrak{m} of R .

Then, the ring of F -linear differential operators $D_F(R)$ is a ring of differentiable type of weak global dimension equal to $\dim(R)$. Moreover, the Bernstein class of $D_F(R)$ is closed under localization at one element of R .

Let $D_F(R)$ be as in the last theorem. Then,

Theorem 3.12. *If M is a left $D_F(R)$ -module in the Bernstein class. Then, each local cohomology module has a natural structure of $D_F(R)$ -module and it belongs to Bernstein class of $D_F(R)$ -module.*

4. EXISTENCE OF THE FIELD OF 1.1

Let R be as in setup 1.1. We first show the existence of a field F contained in R with $\kappa(P)$ a finite extension of F . As S/P is a finitely generated K -algebra, there exist variables y_1, \dots, y_r such that $T = K[y_1, \dots, y_r] \hookrightarrow S/P$ and S/P is integral over $K[y_1, \dots, y_r]$ (By Noether normalization). So quotient field of S/P is a finite extension of quotient field of $K[y_1, \dots, y_r]$. Set F to be the quotient field of $K[y_1, \dots, y_r]$. Therefore S_P/PS_P is a finite extension of F . We claim that F is contained in $R = S_P$. Let ϕ denotes the inclusion map between T and S/P . If $\phi(y_i) = \bar{a}_i$ then the map $\psi : T \rightarrow S$ given by $\psi(y_i) = a_i$ is injective. Composition of ψ with the natural map from S to S_P is injective. Let this map be $h : T \rightarrow S_P$.

$$\begin{array}{ccc} T & \xrightarrow{h} & S_P \\ \downarrow & \nearrow g & \\ T_{(0)} & & \end{array}$$

Every non zero element in T is non zero in S/P and hence unit in S_P . So by universal property of localization there exists a map $g : T_{(0)} \rightarrow S_P$ such that the above diagram is commutative. Hence $T_{(0)} = F$ is contained in S_P as g is injective.

5. PROOF OF THEOREM 1.2

From the discussions above we can see that S_P/PS_P is finite extension of F and this proves first part of 1.2. To prove second part of 1.2, it is enough to prove that $\text{Der}_F(R)$ is free of rank $n = \dim R$ because R is local ring.

Proof. If S is not regular, let $\text{Sing}(S)$, the singular locus of S , be defined by an ideal J in S . Note that $J \neq 0$ and $J \not\subset P$. So there exists $\alpha \in J$ such that $\alpha \notin P$. Then $S_\alpha = V$ is a regular affine domain and $R = S_P = V_{PV}$. From this observation we may assume that S is regular affine domain.

It suffices to show $\text{rank Der}_F(R) = n$ by 3.4. Using 3.3, $\text{rank Der}_F(R) \leq n$. We have that $T = K[y_1, \dots, y_r] \subset S$ and $K \hookrightarrow T \hookrightarrow S$. By 3.1 we get the following exact sequence of S -modules

$$\Omega_{T/K} \otimes_T S \rightarrow \Omega_{S/K} \rightarrow \Omega_{S/T} \rightarrow 0$$

$\Omega_{T/K} = T^r$ and $\Omega_{S/K}$ is a projective S -module of rank equals to $\dim S = m$. So the exact sequence becomes

$$S^r = T^r \otimes_T S \rightarrow \Omega_{S/K} \rightarrow \Omega_{S/T} \rightarrow 0$$

Let L be quotient field of S and $t = \text{rank } \Omega_{S/T}$. Tensoring the above sequence with L we get the exact sequence

$$L^r \rightarrow L^m \rightarrow L^t \rightarrow 0$$

Hence $t \geq m - r = \dim S - \dim S/P = \text{ht } P = \dim R = n$. Hence $\text{rank } \Omega_{S/T} \geq n$. Since $\text{Der}_T(S) \cong \text{Hom}_S(\Omega_{S/T}, S)$, remark 3.7 implies that $\text{rank } \text{Der}_T(S) = \text{rank } \Omega_{S/T} \geq n$. Therefore using 3.7 we get $\text{rank}_R \text{Der}_T(S)_P \geq n$. This implies that $\text{rank } \text{Der}_T(S_P) \geq n$ as $\text{Der}_T(S)_P \subset \text{Der}_T(S_P)$. Clearly $\text{Der}_F(S_P) \subset \text{Der}_T(S_P)$. We claim that this is an equality. Let $D \in \text{Der}_T(S_P)$ and $\frac{a}{s} \in F$, $u \in S_P$. If

$$v = \frac{a}{s} u \implies sv = au \implies D(sv) = D(au) \implies sD(v) = aD(u)$$

Hence we get that $D(v) = \frac{a}{s} D(u)$. Thus $D \in \text{Der}_F(S_P)$. So $\text{Der}_F(S_P) = \text{Der}_T(S_P)$. Therefore $\text{rank } \text{Der}_F(R) \geq n$. Hence $\text{rank } \text{Der}_F(R) = n$ and this proves that $\text{Der}_F(R)$ is a free R module by 3.4. \square

As a corollary of Theorem 3.11 we get 3 of 1.2 that is $D(R)$ is a left and right Noetherian with global dimension n . Moreover, the Bernstein class of $D(R)$ is closed under localization at one element of R .

6. PROOF OF THEOREM 1.4

Since the residue fields of R^h and R are same, the residue field of R^h is finite extension of F . Only thing is to prove that $\text{Der}_F(R^h)$ is a finitely generated free R^h -module of rank same as $\dim R^h = \dim R = n$.

Proof. Using 3.10, if $a \in R^h$, then $a \in \frac{R[x]_{(\mathfrak{m}_R, x)}}{(f(x))} = T \subseteq R^h$ for some $f(x) = x^n + a_{n-1}x^{n-1} + \dots + a_1x + a_0$ with $a_0 \in \mathfrak{m}$, $a_1 \notin \mathfrak{m}$.

Let $L = R[x]_{(\mathfrak{m}_R, x)} = S[x]_{(P, x)}$ and $T = \frac{R[x]_{(\mathfrak{m}_R, x)}}{(f(x))} = L/(f)$. Since $F \hookrightarrow R$, we have that $F \hookrightarrow L$ and $L/\mathfrak{m}_L = R/\mathfrak{m}$ is a finite extension of F .

Using 3.2 for $F \rightarrow L \rightarrow L/(f)$, we get the exact sequence

$$(f)/(f)^2 \rightarrow \Omega_{L/F} \otimes_L T \rightarrow \Omega_{T/F} \rightarrow 0$$

Applying $\text{Hom}_T(-, T)$, the exact sequence becomes

$$0 \rightarrow \text{Hom}_T(\Omega_{T/F}, T) \rightarrow \text{Hom}_T(\Omega_{L/F} \otimes_L T, T) \rightarrow \text{Hom}_T((f)/(f)^2, T)$$

Since $\text{Hom}_T(\Omega_{T/F}, T) \cong \text{Der}_F(T, T)$, $\text{Hom}_T(\Omega_{L/F} \otimes_L T, T) \cong \text{Hom}_L(\Omega_{L/F}, T) \cong \text{Der}_F(L, T)$ and $(f)/(f)^2 \cong T$, we have the exact sequence

$$0 \rightarrow \text{Der}_F(T, T) \rightarrow \text{Der}_F(L, T) \rightarrow \text{Hom}_T(T, T) = T$$

We claim that $\text{rank}_T \text{Der}_F(L, T) \geq n + 1$. Assume the claim for the moment. The last exact sequence implies $\text{rank}_T \text{Der}_F(T) \geq n$. Hence $\text{rank } \text{Der}_F(T) = n$ as 3.3 implies $\text{rank } \text{Der}_F(T) \leq n$. If $a \in T$, then $\frac{\partial}{\partial x_i}(a) \in T$ using 3.4 for all i . Therefore, $\frac{\partial}{\partial x_i}(a) \in R^h$ because $T \subseteq R^h$. Since this is true for all $a \in R^h$, we get $\frac{\partial}{\partial x_i} \in \text{Der}_F(R^h)$ for each i . Again using 3.4, $\text{Der}_F(R^h)$ is free R^h -module of rank n .

Now we prove our claim that $\text{rank}_T \text{Der}_F(L, T) \geq n + 1$. Consider the sequence $0 \rightarrow L \xrightarrow{f} L \rightarrow T \rightarrow 0$. This gives the following exact sequence

$$0 \rightarrow \text{Hom}_L(\Omega_{L/F}, L) \xrightarrow{f} \text{Hom}_L(\Omega_{L/F}, L) \rightarrow \text{Hom}_L(\Omega_{L/F}, T)$$

That is,

$$0 \rightarrow \text{Der}_F(L, L) \xrightarrow{f} \text{Der}_F(L, L) \rightarrow \text{Der}_F(L, T)$$

By our result in 5, $\text{Der}_F(L)$ is a free L -module of rank equals to $(\dim R + 1)$. Therefore, the last exact sequence becomes

$$0 \rightarrow L^{n+1} \xrightarrow{f} L^{n+1} \rightarrow \text{Der}_F(L, T)$$

Since cokernal of f is T^{n+1} , $\text{rank}_T \text{Der}_F(L, T) \geq n + 1$. \square

As a corollary of Theorem 3.11 we get that $D(R^h)$ is left and right Noetherian ring with global dimension n . Moreover, the Bernstein class of $D(R^h)$ is closed under localization at one element of R^h .

7. PROOF OF THEOREM 1.5

In this section, we present the proof of Theorem 1.5. Since R is regular of dimension n , $E_R(\kappa(P)) \cong H_P^n(R)$. We note that R is a $D_F(R)$ -module in the Bernstein class, so is $H_P^n(R)$ by 3.12 and this prove first part of 1.5. We now prove the second part.

Proof. Let $\kappa(P) = K$, $D = D_F(R)$ and $\widehat{D} = D_F(\widehat{R})$. We note that $E_R(K) \cong H_P^n(R)$ has socle dimension 1. Let the socle be spanned by t . Let ϕ be the map between D and $E_R(K)$ defined by $\phi(1) = t$. Since $\phi(DP) = 0$, so it induces a map $\bar{\phi} : \frac{D}{DP} \rightarrow E_R(K)$.

Now

$$\begin{aligned} E_R(K) \otimes_R \widehat{R} &\cong H_P^n(R) \otimes_R \widehat{R} \\ &\cong H_{P\widehat{R}}^n(\widehat{R}) \\ &\cong E_{\widehat{R}}\left(\frac{\widehat{R}}{P\widehat{R}}\right) \end{aligned}$$

And

$$\begin{aligned} \frac{D}{DP} \otimes_R \widehat{R} &\cong (D \otimes_R \frac{R}{P}) \otimes_R \widehat{R} \\ &\cong (D \otimes_R \widehat{R}) \otimes_R \frac{R}{P} \end{aligned}$$

Using 3.5, $(D \otimes_R \widehat{R}) \cong \widehat{D}$. Hence we get the following

$$\begin{aligned} \frac{D}{DP} \otimes_R \widehat{R} &\cong \widehat{D} \otimes_R \frac{R}{P} \\ &\cong \widehat{D} \otimes_{\widehat{R}} (\widehat{R} \otimes_R \frac{R}{P}) \\ &\cong \widehat{D} \otimes_{\widehat{R}} \frac{\widehat{R}}{P\widehat{R}} \\ &\cong \frac{\widehat{D}}{\widehat{DP}} \end{aligned}$$

3.6 implies $\frac{\widehat{D}}{\widehat{DP}} \cong E_{\widehat{R}}(\frac{\widehat{R}}{\widehat{PR}})$. Therefore, $E_R(K) \otimes_R \widehat{R} \cong \frac{D}{DP} \otimes_R \widehat{R}$. Since $R \rightarrow \widehat{R}$ is faithfully flat map we get $\frac{D}{DP} \cong E_R(K)$ and this proves our result. \square

ACKNOWLEDGEMENT

The first named author expresses sincere thanks for the invaluable support received from the Government of India through the Prime Minister's Research Fellowship (PMRF ID 1303161), which significantly aided in the completion of this work.

REFERENCES

- [1] I. N. Bernšteĭn, I. M. Gel'fand, and S. I. Gel'fand, *Differential operators on a cubic cone*, Uspehi Mat. Nauk **27** (1972), no. 1(163), 185–190. MR 385159
- [2] Luis Núñez Betancourt, *On certain rings of differentiable type and finiteness properties of local cohomology*, J. Algebra **379** (2013), 1–10. MR 3019242
- [3] J.-E. Björk, *Rings of differential operators*, North-Holland Mathematical Library, vol. 21, North-Holland Publishing Co., Amsterdam-New York, 1979. MR 549189
- [4] Gennady Lyubeznik, *Finiteness properties of local cohomology modules (an application of D-modules to commutative algebra)*, Invent. Math. **113** (1993), no. 1, 41–55. MR 1223223
- [5] B. Iversen, *Generic local structure of the morphisms in commutative algebra*, Lecture Notes in Math., Vol. 310 Springer-Verlag, Berlin-New York, 1973.
- [6] Hideyuki Matsumura, *Commutative ring theory*, Cambridge Studies in Advanced Mathematics, vol. 8, Cambridge University Press, Cambridge, 1986, Translated from the Japanese by M. Reid. MR 879273
- [7] Z. Mebkhout and L. Narváez-Macarro, *La théorie du polynôme de Bernstein-Sato pour les algèbres de Tate et de Dwork-Monsky-Washnitzer*, Ann. Sci. École Norm. Sup. (4) **24** (1991), no. 2, 227–256. MR 1097693
- [8] L. Narváez-Macarro, *Differential structures in commutative algebra*, Mini-course at the XXIII Brazilian Algebra Meeting, July 27–August 1, 2014, Maringá, Brazil.
- [9] Masayoshi Nagata, *Local rings*, Interscience Tracts in Pure and Applied Mathematics, vol. No. 13, Interscience Publishers (a division of John Wiley & Sons, Inc.), New York-London, 1962. MR 155856
- [10] Tony J. Puthenpurakal, *On the ring of differential operators of certain regular domains*, Proc. Amer. Math. Soc. **146** (2018), no. 8, 3333–3343. MR 3803659

Email address: 22d0786@iitb.ac.in

Email address: tputhen@math.iitb.ac.in

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY BOMBAY, POWAI, MUMBAI 400076, INDIA