

Schottky Anomaly of Reissner-Nordström-de Sitter spacetime

Hai-Long Zhen,^{1,2} Yu-Bo Ma,^{1,2} Huai-Fan Li,^{1,2} Li-Chun Zhang,^{1,2} Yun-Zhi Du,^{1,2*}

¹*Department of Physics, Shanxi Datong University,
Datong 037009, China*

²*Institute of Theoretical Physics,
Shanxi Datong University, Datong, 037009, China*

In the extended thermodynamics of black holes, there exists a thermodynamical pressure and its conjugate volume. Extensive studies have been conducted on the phase structure of numerous black holes, which have demonstrated striking similarities to the phase structure of various ordinary matter systems. A comparison of the thermodynamic properties of spherically symmetric AdS black holes with those of ordinary thermodynamic systems reveals that the isovolumetric heat capacity of the former is zero, whereas that of the latter is non-zero. It is a subject of interest for the intrinsic reason for this discrepancy. The effective thermodynamic quantities, based on the context of the boundary between the black hole horizon and the cosmological horizon in dS spacetime, as well as the interaction between the two horizons, are presented. The heat capacity in the Reissner-Nördstrom-de Sitter (RN-dS) spacetime is then investigated, and it is demonstrated that the behavior of the heat capacity in the RN-dS spacetime is analogous to that of Schottky specific heat. Treating the two horizon interfaces in the RN-dS spacetime as two distinct energy levels in a two-energy system, and the thermodynamic properties in the RN-dS spacetime are discussed with the method of studying the thermodynamic properties in an ordinary two-energy system, to elucidate the intrinsic reasons for the occurrence of Schottky specific heat in the RN-dS spacetime. The heat capacity observed in the RN-dS spacetime is not only consistent with that of the Schottky specific heat described by the effective thermodynamic quantities in the RN-dS spacetime, but also with that of the heat capacity in an ordinary two-energy-level system. The quantum properties in the RN-dS spacetime are reflected, thereby providing a new avenue for further in-depth study of the quantum properties of black holes and dS spacetime.

Keywords: de Sitter spacetime, effective thermodynamic quantities, heat capacity, Schottky specific heat.

1. INTRODUCTION

On 10 April 2019, the Event Horizon Telescope (EHT) research team released the first photograph of the supermassive black hole at the centre of the galaxy M87 [1, 2]. The morphology of the photograph is consistent with the Kerr black hole model predicted by general relativity. This observation provides compelling evidence for the presence of a supermassive black hole in the central engine of active galactic nucleus. Black hole is not only a strong gravitational system, but also a thermodynamic system. As early as the 1970s, physicists such as Hawking and Bekenstein established the four laws of black hole thermodynamics [3, 4]. The investigation of the thermodynamic properties of black holes not only contributes to a deeper understanding of their nature, but also provides insights into their internal microscopic state.

In order to investigate the microstructure of black holes as well as the formation and evolution of black holes in the Universe, the study of the thermodynamic properties of AdS and dS spacetime has attracted significant attention. Recently, in the context of the (n)-dimensional AdS spacetime, a potential correlation between the cosmological constant, Λ , and thermodynamical pressure, P , has been proposed. This hypothesis, which can be found in [5, 6], has attracted significant attention. This relation is defined as follows $P = \frac{n(n-1)}{16\pi l^2}$, $\Lambda = \frac{n(n-1)}{2l^2}$, in which the thermodynamical volume, V , is the conjugate quantity to pressure as $V = \left(\frac{\partial M}{\partial P}\right)_{S, Q_i, J_k}$. Based on these considerations, the thermodynamic properties of AdS black holes have been studied from different perspectives. Firstly, treating AdS black hole as an ordinary thermodynamic system, compared to the phase transition of ordinary system, the phase transition of AdS black hole has been studied. It has been demonstrated that the phase transition of the AdS black hole is analogous to that of the ordinary van der Waals system [7–31]. Secondly, the John-Thomson coefficients of AdS black holes have been investigated. The effects of different parameters and dimensions of AdS black holes on

* e-mail: duyzh22@sxdtu.edu.cn (corresponding author)

the John-Thomson coefficient have been explored [32–36]. Recently, the study of the topological properties of AdS black holes has attracted interest. The properties of black holes at the phase transition point are classified through the study of topological numbers. [37–41]. The microstructure of black holes has been one of the most significant theoretical problems. Notable advances have been made in exploring the microstructure of black holes by various methods [42–50]. Nevertheless, the microstructure of black holes remains a mystery. In order to gain insight into the quantum properties and microstructure of black holes, it is necessary to have a comprehensive understanding of the thermodynamic properties of black holes. Consequently, the discussion of various thermodynamic properties of black holes remains a significant topic in contemporary theoretical physics.

During the early inflationary epoch, our Universe is regarded as a quasi-de Sitter spacetime. By corresponding the cosmological constant to dark energy, our Universe will evolve to a new de Sitter phase [51]. To construct the entire history of the evolution of the Universe and identify the intrinsic cause of the accelerated expansion of the Universe, it is necessary to have a clear understanding of the classical, quantum and thermodynamic properties of the de Sitter spacetime. Consequently, the investigation of the thermodynamic characteristics of de Sitter spacetime has attracted considerable attention [52–71]. It is well established that in de Sitter spacetime, the black hole horizon and the cosmological horizon coexist if the spacetime parameters satisfy certain conditions. In this region, the thermodynamic equilibrium stability requirement in spacetime is not satisfied due to the existence of two horizon interfaces with different radiation temperatures. Consequently, in order to discuss the thermodynamic properties of this region, it is necessary for establishing a thermodynamic system to satisfy the requirement of thermodynamic equilibrium stability. Because the thermodynamic system must satisfy the universal relation for thermodynamic systems, which is the first law of thermodynamics. Firstly, the state parameters of the effective thermodynamic system established in dS spacetime satisfy the first law of thermodynamics. Secondly, the effective thermodynamic quantities satisfy the boundary conditions when the values of the state parameter in spacetime are that of the boundary of the coexistence region between the two horizon interfaces. Consequently, the effective thermodynamic system, based on the aforementioned conditions, provides a more comprehensive representation for the thermodynamic properties in spacetime.

A differential equation for the entropy of an effective thermodynamic system, based on the consideration of the interaction between the two horizon interfaces in dS spacetime, is given for the state parameter satisfying the universal first law of thermodynamics. The solution for the entropy of the effective thermodynamic system is obtained by the condition satisfied by the effective thermodynamic system at the boundary. Subsequently, the effective temperature and effective pressure of the effective thermodynamic system are then determined, and the equation of state of the effective thermodynamic system satisfying the boundary conditions is found. This provides a foundation for further in-depth study of the thermodynamic effect in dS spacetime. The study reveals that when the thermodynamic properties are described by the effective thermodynamic quantities in RN-dS spacetime, the behavior of the heat capacity in RN-dS spacetime is analogous to that of the Schottky specific heat, which is highly consistent with that of the heat capacity treating the two horizons in RN-dS spacetime as a two-energy-level system. This conclusion contributes to a deeper understanding of dS spacetime, which can be used to employed to investigate the state of motion of microscopic particles within black holes and to simulate the evolution of the Universe. Furthermore, this discovery offers a novel avenue for investigating the intrinsic causes of the accelerated expansion of the Universe.

The paper is arranged as follows: in Sec. 2, the conditions of in the presence of the black hole horizon and the cosmological horizon in RN-dS spacetime, along with the effects of parameters in spacetime on the existence of spacetime with the black hole horizon and the cosmological horizon are discussed. Furthermore, the interval of the value of the position ratio, x , between the two horizons in spacetime is determined. In Sec. 3, the effective thermodynamic quantities in the RN-dS spacetime satisfying the boundary conditions are presented. The behavior of the effective temperature, T_{eff} , the radiation temperature of the black hole horizon, T_+ , and the radiation temperature of the cosmological horizon, T_c , with respect to the position ratio between the two horizons, x , in spacetime is analysed. The Smarr relation treating effective quantities as state parameters is expressed. In Sec. 4, the behavior of the heat capacity, $C_{Q,l}$, described by the effective state parameter in the RN-dS spacetime, under both the charge, Q , and the cosmological constant, l , being fixed, is investigated. The results demonstrate that the behavior of the heat capacity, $C_{Q,l}$, in the RN-dS spacetime with the effective temperature, T_{eff} , and the position ratio, x , between the two horizons, exhibits Schottky peaks. It is similar to that of the heat capacity, $\hat{C}_{Q,l}$, with the effective temperature, T_{eff} , and the position ratio between the two horizons, x , in a two-level system treating the black hole horizon and the cosmological horizon as two distinct energy levels within an effective thermodynamic system. It indicates that the Schottky peaks in RN-dS spacetime is determined by a two-energy level system comprising two distinct horizons. Finally, a brief summary is presented in Sec. 5

2. RN-DS SPACETIME

The RN-dS black hole represents a static solution of the Einstein-Maxwell equations, incorporating an additional term for electric charge in comparison to the SdS case. [67]

$$ds^2 = -g(r)dt^2 + g^{-1}dr^2 + r^2d\Omega_2^2, \quad (2.1)$$

with the horizon function

$$g(r) = 1 - \frac{2M}{r} + \frac{Q^2}{r^2} - \frac{r^2}{l^2}, \quad (2.2)$$

M and Q represent the mass and charge of the black hole, respectively. The curvature radius of dS space is denoted by l . The positions of the black hole and cosmological horizons, r_+ and r_c respectively, are satisfied by the equation $g(r_{+,c}) = 0$. Substitution of eq. (2.2) into this relation yields

$$M = \frac{r}{2} \left(1 + \frac{Q^2}{r^2} - \frac{r^2}{l^2} \right), \quad (2.3)$$

The $M - r$ curve is then drawn from eq. (2.3).




FIG. 2.1: The $M - r$ curve

Fig. 2.1 of the $M - r$ curve demonstrates that the RN-dS spacetime possesses an inner black hole horizon r_- , a black hole horizon r_+ and a cosmological horizon r_c when $Q=1, l^2=60$ as well as the RN-dS spacetime energy is constrained by the condition $M_A \leq M \leq M_C$. At the point of local maxima, designated as C , the black hole horizon and cosmological horizon coincide. This point corresponds to the maximum energy, $M = M_C$, when the black hole and cosmological horizon exist in spacetime. At the point of the local minimum, designated as A , the inner and outer horizons of the black hole coincide. This point corresponds to the minimum energy, $M = M_A$, when the black hole and cosmological horizon exist in spacetime. In the RN-dS spacetime, a black hole does not exist when $M > M_C$ or $M < M_A$ [72]. The Nariai radius and mass [72]

$$r_N^2 = \frac{l^2}{6} \left(1 + \sqrt{1 - \frac{12Q^2}{l^2}} \right),$$

$$M_N = \frac{l}{\sqrt{6} \left(1 + \sqrt{1 - \frac{12Q^2}{l^2}} \right)} \left(\frac{1}{3} \left(1 + \sqrt{1 - \frac{12Q^2}{l^2}} \right) + \frac{4Q^2}{l^2} \right). \quad (2.4)$$

when $Q^2 \ll l^2$, the mass of spacetime corresponding to point A , $M_A \rightarrow 0$, and the mass of spacetime corresponding to point C , $M_C \rightarrow \frac{1}{3\sqrt{3}}$. It can be shown from Eq. (2.4) that the charge Q^2 and the cosmological constant l^2 in spacetime must satisfy $\frac{12Q^2}{l^2} \leq 1$. When the local minimum and local maximum points of the black hole mass curve merge into an inflection point, the three horizons coincide. In this case, the black hole is referred to as the ultracold black hole, whose horizon radius and mass are, with given magnetic charge Q , given as

$$r_{AC}^4 = \frac{Q^2 l^2}{3}, \quad r_{AC}^4 = 4Q^4, \quad M_{AC} = \frac{4Q}{3\sqrt{2}}. \quad (2.5)$$

From Eq. (2.4), it can be seen that when $Q=0$, $r_{A,C}^2 = \frac{l^2}{3}$, which indicates that there is no region of coexistence between the two horizons in spacetime. By treating the black hole and the cosmological horizon as independent thermodynamic systems, it can be demonstrated that the thermodynamic quantities on the two horizons each satisfy the first law of thermodynamics [51, 60, 72].

For the black hole horizon, the associated thermodynamic quantities are

$$dM = T_+ dS_+ + V_+ dP + \phi_+ dQ, \quad (2.6)$$

with

$$T_+ = \frac{g'(r_+)}{4\pi} = \frac{1}{4\pi r_+} \left(1 - \frac{3r_+^2}{l^2} - \frac{Q^2}{r_+^2} \right), \quad S_+ = \pi r_+^2, \quad \phi_+ = \frac{Q}{r_+}. \quad (2.7)$$

Some thermodynamic quantities associated with the cosmological horizon are

$$dM = -T_c dS_c + V_c dP + \phi_c dQ, \quad (2.8)$$

$$T_c = -\frac{g'(r_c)}{4\pi} = -\frac{1}{4\pi r_c} \left(1 - \frac{3r_c^2}{l^2} - \frac{Q^2}{r_c^2} \right), \quad S_c = \pi r_c^2, \quad \phi_c = \frac{Q}{r_c}. \quad (2.9)$$

T_c , S_c and ϕ_c denote the Hawking temperature, the entropy and the electric potential, respectively.

From the relationship $g(r_{+,c}) = 0$, which is satisfied by the position of the black hole horizon, r_+ , and the position of the cosmological horizon, r_c , we can derive

$$M = \frac{r_c x(1+x)}{2(1+x+x^2)} + \frac{Q^2(1+x)(1+x^2)}{2r_c x(1+x+x^2)}, \quad \frac{1}{l^2} = \frac{1}{r_c^2(1+x+x^2)} - \frac{Q^2}{r_c^4(1+x+x^2)}, \quad (2.10)$$

with $x = r_+/r_c$, which denotes the ratio of the position of the black hole horizon to that of the cosmological horizon. From Eq. (2.10), it can be seen that this corresponds to the point C when $x \rightarrow 1$. Consequently, only the '+' sign will be considered. namely,

$$r_c^2 = \frac{l^2}{6} \left(1 + \sqrt{1 - \frac{12Q^2}{l^2}} \right), \quad (2.11)$$

upon taking $l^2 = \alpha 12Q^2$ (with $\alpha \geq 1$ being constant), from Eq. (2.10), we can get

$$r_c^2 = \frac{6\alpha Q^2}{(1+x+x^2)} \left(1 \pm \sqrt{1 - \frac{(1+x+x^2)}{3x\alpha}} \right), \quad \frac{1}{r_c^2} = \frac{x}{2Q^2} \left(1 - \sqrt{1 - \frac{(1+x+x^2)}{3x\alpha}} \right), \quad (2.12)$$

the position of the ultracold black hole horizon at $\alpha = 1$ and $x \rightarrow 1$ is given by Eq. (2.12).

Substituting Eq. (2.10) into Eqs. (2.7) and (2.9) yields the radiation temperatures of the black hole horizon and the cosmological horizon, respectively, as

$$T_+ = \frac{(1-x)}{4\pi r_c x} \left(\frac{(1+2x)}{(1+x+x^2)} - \frac{Q^2(1+2x+3x^2)}{r_c^2 x^2(1+x+x^2)} \right), \quad (2.13)$$

$$T_c = \frac{(1-x)}{4\pi r_c} \left(\frac{(2+x)}{(1+x+x^2)} - \frac{Q^2(3+2x+x^2)}{r_c^2 x(1+x+x^2)} \right). \quad (2.14)$$

From Fig. 2.1, it can be observed that the black hole horizon and the cosmological horizon appear simultaneously in spacetime in the case of the electric charge, Q , the cosmological constant, l , fixed, as well as the energy in spacetime meets the constraints $M_A \leq M \leq M_C$. The energy, M , the electric charge, Q , and the cosmological constant, l , corresponding to the two points B and D are identical. Nevertheless, in general, the radiation temperature of the black hole horizon (B point), T_+ , is not equal to the radiation temperature of the cosmological horizon (D point), T_c . In this region, two thermodynamic subsystems with different temperatures emerge in spacetime, corresponding to the thermodynamic system on the black hole horizon and the cosmological horizon, respectively. Consequently, under the same parameters, the spacetime in this region is in a state of thermodynamic equilibrium that is unstable. At point A , the radiation temperature of the black hole is zero, which also marks the end of the coexistence region of two different

radiation temperatures in spacetime. Point E , which corresponds to point A , represents the endpoint of spacetime from two thermodynamic subsystems to the thermodynamic system only including the cosmological horizon, as well as the endpoint of spacetime with the two thermodynamic subsystems. The RN-dS spacetime exhibits disparate thermodynamic characteristics across distinct regions. Of particular interest is the thermodynamic behavior in the coexistence region, where two disparate temperatures, T_+ and T_c , represent a topic of great theoretical interest [51, 53–72].

In recent years, a search has been conducted for the state quantities of thermodynamic systems in spacetime in the region of coexistence of two subthermodynamic systems under various conditions. In [53, 62], the sum of the entropies corresponding to the two horizons is defined as the total entropy of spacetime. The effective temperature of the thermodynamic system in spacetime is then obtained by satisfying the first law of thermodynamics, which requires that the system in question be in equilibrium. This concept paves the way for the investigation of the thermodynamic properties of non-equilibrium states in spacetime. However, the interaction between two horizons is not considered, and the conclusions obtained are therefore not comprehensive. In [64, 66], the effective temperature and interaction entropy of spacetime are presented based on the interaction between the two horizons. This provides a foundation for the investigation of the thermodynamic properties of dS spacetime and the entropic force between the two horizons. It is crucial to acknowledge that this approach does not account for the condition that when the spacetime is at the endpoint of the two subthermodynamic systems, E , only the temperature of the cosmological horizon, T_c , exists in spacetime, with the radiation temperature of the black hole horizon, T_+ , being zero. This paper presents an extension of the study of the thermodynamic properties of dS spacetime, as previously outlined in [64, 66], to encompass the effective thermodynamic quantities of the RN-dS spacetime under various known boundary conditions.

3. THE EFFECTIVE THERMODYNAMIC QUANTITIES IN RN-DS SPACETIME

The analyses presented in Sec. 2 demonstrate that in order to establish a global thermodynamic system in a region with two subsystems in spacetime, certain boundary conditions must be satisfied by the system and be universal. The thermodynamic properties in RN-dS spacetime, which exhibits a black hole horizon and a cosmological horizon, are being considered. It is therefore of interest to consider the interval $M_A \leq M \leq M_C$. The exact values of M_A and M_C are related to the spacetime parameters, such as charge, Q , and the cosmological constant, l , which are given by Eq. (2.10). Considering the connection between the black hole horizon and the cosmological horizon, the effective thermodynamic quantities and the corresponding first law of black hole thermodynamics are derived [64, 66].

$$dM = T_{eff}dS - P_{eff}dV + \phi_{eff}dQ. \quad (3.1)$$

Where the thermodynamic volume is that between the black hole horizon and the cosmological horizon, namely [53, 60, 62].

$$V = V_c - V_+. \quad (3.2)$$

From Eq. (2.13), it can be seen when the potential, denoted by ϕ_A , on the black hole horizon at point A satisfies

$$\frac{Q^2}{r_A^2} = \frac{1 + 2x}{1 + 2x + 3x^2}. \quad (3.3)$$

The radiation temperature of the black hole horizon, $T_+=0$. The radiation temperature of the cosmological horizon, corresponding to the point E , is

$$T_E = \frac{(1-x)^2(1+x)}{2\pi r_c(1+2x+3x^2)}. \quad (3.4)$$

Given that the radiation temperature at point A is zero, it can be demonstrated that the effective temperature in spacetime should be T_E when the potential on the black hole horizon in the RN-dS spacetime satisfies Eq. (3.3). In this context, the effective temperature in spacetime, T_{eff} , can be derived from Eqs. (3.1), (3.2) and the boundary condition (3.4)

$$T_{eff} = \frac{1-x}{4\pi r_c x^5} \left\{ [(1+x)(1+x^3) - 2x^2] - \frac{Q^2}{r_c^2 x^2} [(1+x+x^2)(1+x^4) - 2x^3] \right\}, \quad (3.5)$$

the effective pressure in spacetime, P_{eff} , is

$$\begin{aligned}
P_{eff} &= -\frac{(1-x)}{16\pi r_c^2 x^5} \left[F'(x) \left(x(1+x) - \frac{Q^2(1+x+x^2+x^3)}{r_c^2 x} \right) - \frac{2F(x)}{(1+x+x^2)} \left((1+2x) - \frac{Q^2(1+2x+3x^2)}{r_c^2 x^2} \right) \right] \\
&= -\frac{(1-x)}{8\pi r_c^2 x^5 (1-x^3)} \left[\frac{x^6}{(1-x^3)} \left((1+x) - \frac{Q^2(1+x+x^2+x^3)}{r_c^2 x^2} \right) \right. \\
&\quad \left. - F(x) \left([(1+x)(1+x^3) - 2x^2] - \frac{Q^2[(1+x+x^2)(1+x^4) - 2x^3]}{r_c^2 x^2} \right) \right], \tag{3.6}
\end{aligned}$$

with

$$\begin{aligned}
F(x) &= \frac{8}{5}(1-x^3)^{2/3} + \frac{2}{5(1-x^3)} - 1 = 1+x^2 + f_0(x), \tag{3.7} \\
f_0(x) &= \frac{8}{5}(1-x^3)^{2/3} - \frac{8+5x^2-10x^3-5x^5}{5(1-x^3)}.
\end{aligned}$$

The thermodynamic volume, V , entropy, S and potential, ϕ_{eff} , of the effective thermodynamic system are, respectively

$$V = \frac{4\pi}{c} r_c^3 (1-x^3), \quad S = \pi r_c^2 F(x), \quad \phi_{eff} = \frac{Q(1+x)(1+x^2)}{r_c x(1+x+x^2)}. \tag{3.8}$$

From Eqs. (2.10) and (3.3), the minimum value, $x = x_{min}$, of the ratio, x , between the two horizon positions in spacetime, given the charge, Q , and the cosmological constant, l , in spacetime, is given by

$$\frac{6\alpha}{(1+x+x^2)} \left(1 \pm \sqrt{1 - \frac{(1+x+x^2)}{3x\alpha}} \right) = \frac{1+2x+3x^2}{x^2(1+2x)}, \tag{3.9}$$

i.e. $(1+2x=3x^2)^2 - 12\alpha x^2(1+2x) = 0$.




FIG. 3.1: The $x_{min} - \alpha$ curve

As illustrated in Fig. 3.1, it can be observed that x_{min} towards zero with an increase in α . This is demonstrated by the values of x_{min} , which are 0.0560513957; 0.042749089; 0.029789688, respectively, when the values of α are 30; 50; 100, respectively. It can be concluded that the position ratio x between the two horizons assumes a value within the interval $x_{min} \leq x \leq 1$ when the coexistence region of the two horizons exists in spacetime.

When the potential on the black hole horizon in spacetime satisfies [73]

$$\frac{Q^2}{r_+^2} = \frac{1}{(1+x)^2}, \tag{3.10}$$

the radiation temperature of the black hole horizon, \bar{T}_+ , is equal to that of the cosmological horizon, \bar{T}_c , i.e.

$$\bar{T}_+ = \bar{T}_c = \frac{1-x}{2\pi r_c(1+x)^2}. \tag{3.11}$$

Substituting Eq. (3.10) into Eq. (3.5), we obtain

$$T_{eff} = \bar{T}_{eff} = \bar{T}_+ + \frac{(1-x)}{2\pi r_c x^4(1+x)^2} = \bar{T}_c + \frac{(1-x)}{2\pi r_c x^4(1+x)^2}. \tag{3.12}$$

From Eq. (3.12), it can be observed that treating the thermodynamic system corresponding to the two horizons with the same radiation temperature in RN-dS spacetime as an independent thermodynamic system leads to a discrepancy in the effective radiation temperature which takes into account the conditions of interaction between the two horizons. It can be demonstrated that the effective temperature, \bar{T}_{eff} , of an effective thermodynamic system consisting of two subsystems with the same temperature in RN-dS spacetime, is not equal to the temperature of the subsystems, i.e. $\bar{T}_{eff} > \bar{T}_+ = \bar{T}_c$. This phenomenon can be attributed to the interaction between the two horizons. This phenomenon can be attributed to the existence of an interaction between the two subsystems. Despite the temperatures of the two subsystems being identical, namely, $\bar{T}_+ = \bar{T}_c$, the interaction between them results the effective temperature, \bar{T}_{eff} , of the effective thermodynamic system being distinct from that of the respective subsystems. This illustrates the pivotal role of the interaction between the two horizons in RN-dS spacetime in the effective thermodynamic system. This demonstrates that the thermodynamic system in the RN-dS spacetime is distinct from the ordinary thermodynamic system. The discovery of this phenomenon suggests that it is necessary to take into account the behavior of thermodynamic quantities due to the interaction between black holes and the external environment when the thermodynamic properties of black holes are studied by analogy with the method used to study ordinary thermodynamic systems. Consequently, it is of the importance to investigate the intrinsic causes of the interaction between the two horizons in greater depth.

The behavior of T_+ , T_c and T_{eff} can be plotted against the ratio between the two horizon positions, x , for different values of the parameter α by substituting Eq. (2.12) into Eqs. (2.13), (2.14) and (3.5) (taking a + sign in Eq. (2.12)).




FIG. 3.2: The $T_+ - x$ curve for different α , $Q=1$






FIG. 3.3: The $T_c - x$ curve for different α , $Q=1$

Figs. 3.2, 3.3 and 3.4 illustrate that T_+ , T_c and T_{eff} decrease as α increases at a fixed value of x . The behavior of T_+ , T_c and T_{eff} as a function of x is plotted in Fig. 3.5

Fig. 3.5 demonstrates that the temperature in question, which is a consequence of the interaction between the two horizons, is constrained to satisfy $T_{eff} \geq T_+ \geq T_c$ under a fixed x . The boundary condition of the effective thermodynamic system is satisfied when the effective temperature, T_{eff} , and the radiation temperature of the cosmological horizon, T_c , are equal, as well as the radiation temperature of the black hole horizon, T_+ , is zero. Furthermore, the Smarr relation of the effective thermodynamic system is as follows:

$$M = 2T_{eff}S + \varphi_{eff}Q - 3P_{eff}V \quad (3.13)$$

FIG. 3.4: The $T_{eff} - x$ curve for different α , $Q=1$ FIG. 3.5: The $T_+, T_c, T_{eff} - x$ curves for $\alpha=50$, $Q=1$

4. SCHOTTKY ANOMALY OF RN-DS BLACK HOLES

The heat capacity of a thermal system is typically an increasing function of temperature. However, a peak in the heat capacity, known as a Schottky anomaly, can occur in a system that has a maximum energy. It is well established that for a paramagnetic mass system with $j = 1/2$, which is a two-level system, the partition function

$$z = e^{\beta\varepsilon} + e^{-\beta\varepsilon}, \quad (4.1)$$

from Eq. (4.1), the internal energy of the system as follows:

$$U = F + TS = -N\varepsilon \tanh(\beta\varepsilon), \quad (4.2)$$

with $kT = 1/\beta$. The specific heat, when the interaction between particles is not taken into account, can be expressed in the following general form

$$C = Nk \left(\frac{\Delta}{T} \right)^2 \frac{e^{\frac{\Delta}{T}}}{(1 + e^{\frac{\Delta}{T}})^2}. \quad (4.3)$$

is a characteristic of a two-energy-level system with an energy gap $\Delta = 2\varepsilon$, which is the celebrated Schorrky specific heat. It is of great importance for the treatment of a multitude of systems as two-energy systems. The energy and specific heat capacity curves plotted versus temperature are presented in Fig. 4.1 [68, 70].

The effective thermodynamic system, comprising with two thermodynamic subsystems, was obtained through the consideration of the interaction between the two horizons with disparate temperatures in dS spacetime. The interaction between the two subsystems occurs through the two horizons. It is therefore necessary to determine whether the thermodynamic properties between the two horizons can be described as those of a two-energy level system. This is one of the key issues to be addressed in the study of the thermodynamic properties in dS spacetime. In recent years, the Schorrky specific heat with some promising results emerging in dS spacetime has been investigated [68, 70, 74]. Nevertheless, the conclusions obtained are not exhaustive in the absence of the interaction between the black hole and cosmological horizons. The Schorrky specific heat in RN-dS spacetime based on the interaction between the two horizons is investigated in this section.




FIG. 4.1: The curves of $U/N\varepsilon - kT/\varepsilon$ and $\hat{C} - \hat{T}$

When the charge, Q , and the cosmological constant, l , in spacetime are held constant,

$$C_{Q,l} = T_{\text{eff}} \left(\frac{\partial S}{\partial T_{\text{eff}}} \right)_{Q,l} = T_{\text{eff}} \left(\frac{dS/dx}{dT_{\text{eff}}/dx} \right)_{Q,l}, \quad (4.4)$$

from Eq. (4.4), the behavior of $C_{Q,l}$ with respect to the ratio of the two horizon positions, x , and the effective temperature, T_{eff} , can be plotted when different parameters, α , are taken into account.




FIG. 4.2: The $C_{Q,l} - x$ curve for different α






FIG. 4.3: The $C_{Q,l} - T_{\text{eff}}$ curve for different α

As illustrated in Figs. 4.2 and 4.3, the energy and heat capacity curves in RN-dS spacetime exhibit similarities to those of the two-energy-level curves depicted in Fig. 4.1. The highest energy and heat capacity exhibit an extreme value within certain range of values. The effective thermodynamic system in RN-dS spacetime is constituted by thermodynamic subsystems with the interaction between the two horizons. Consequently, the thermodynamic properties between the two horizons are of significant importance in the context of the effective thermodynamic system. The specific heat, $C_{Q,l}$, of the effective thermodynamic system in RN-dS spacetime exhibits a pattern of behavior consistent with that of a two-energy-level thermodynamic system. The effective thermodynamic system

FIG. 4.4: The $M - x$ curve for different α FIG. 4.5: The $M - T_{eff}$ curve for different α

in RN-dS spacetime is constituted by the subsystems that correspond to the two horizons. Consequently, the two horizons in RN-dS spacetime can be regarded as two distinct energy levels of a two-energy system. When the particles are situated on different horizons, they exhibit distinct energy levels. According to this hypothesis, the two horizons in RN-dS spacetime are regarded as two different energy levels. The heat capacity generated by the transitions of particles between energy levels is in accordance with Eq. (4.2). It is proposed that the energies of the particles on the black hole and cosmological horizons are, respectively,

$$\varepsilon_+ = \varepsilon T_+, \quad \varepsilon_c = \varepsilon T_c, \quad (4.5)$$

where ε is a positive constant. The two horizons in an effective thermodynamic system. Thus Δ in Eq. (4.2) as follows

$$\Delta = \varepsilon_+ - \varepsilon_c = \varepsilon(T_+ - T_c), \quad T = T_{eff}/\varepsilon. \quad (4.6)$$

when $\varepsilon = 1$ (the behavior of the curve is independent of the value of ε). Substituting Eq. (4.6) into Eq. (4.2) yields

$$\frac{\Delta}{T_{eff}} = \frac{\varepsilon x^4 (1-x)^2 (1+x) \left(1 - \frac{1}{2} \left(1 - \sqrt{1 - \frac{(1+x+x^2)}{3x\alpha}}\right) \frac{(1+x)^2}{x}\right)}{(1-x^3) \left\{ [(1+x)(1+x^3) - 2x^2] - \frac{1}{2x} \left(1 - \sqrt{1 - \frac{(1+x+x^2)}{3x\alpha}}\right) [(1+x+x^2)(1+x^4) - 2x^3] \right\}} \quad (4.7)$$

In order to differentiate it from the two-energy thermodynamic system, we designate the heat capacity of the two energy levels in the RN-dS spacetime as

$$\hat{C}_{Q,i} = Nk \left(\frac{\varepsilon \Delta}{T_{eff}} \right)^2 \frac{e^{\frac{\varepsilon \Delta}{T_{eff}}}}{\left(1 + e^{\frac{\varepsilon \Delta}{T_{eff}}}\right)^2}. \quad (4.8)$$

A comparison of Figs. 4.2 and 4.6, and Figs. 4.3 and 4.7, reveals that the two horizon interfaces in the RN-dS spacetime are regarded as two distinct energy levels of the effective thermodynamic system. The two-level system, $\hat{C}_{Q,i}$, exhibits the fundamental properties of the effective thermodynamic system, $C_{Q,i}$. As illustrated in Fig. 4.7, a quantum system with a split spectrum (two energy levels) in RN-dS spacetime is considered. This system exhibits




FIG. 4.6: The $\hat{C}_{Q,l}/Nk - x$ curve for different α , $\varepsilon=1$




FIG. 4.7: The $\hat{C}_{Q,l}/Nk - T_{eff}/\varepsilon$ curve for different α

the behavior of $C_{Q,l} \rightarrow 0$ when $T_{eff} \rightarrow 0$. The specific heat capacity reaches an extreme value. The specific heat capacity in RN-dS spacetime is in accordance with the universal Eq. (4.2) for two energy levels, which is known as the Schottky specific heat. It is significant that numerous systems can be treated as two-energy systems. This result demonstrates a novel quantum property in the RN-dS spacetime.

5. CONCLUSION

The effective thermodynamic quantities, based on the interaction between the black hole horizon and the cosmological horizon under the boundary condition in dS spacetime, as well as the Smarr relation and the system free energy expressed in the effective thermodynamic quantities, are presented in the RN-dS spacetime. This lays the foundation for further study of the thermodynamic properties in RN-dS spacetime. The effective thermodynamic quantities obtained in the RN-dS spacetime were then employed to investigate the heat capacity in the RN-dS spacetime with the region of coexistence between the black hole horizon and the cosmological horizon. It was found that the behavior of the heat capacity in the RN-dS spacetime described by the effective thermodynamic quantity was analogous to that of Schottky specific heat. In order to elucidate the underlying causes of this phenomenon, we posit that the black hole horizon interface and cosmological horizon interface in the RN-dS spacetime can be regarded as two distinct energy levels of an effective thermodynamic system, i.e. a two-energy-level system. The heat capacity of this two-energy-level system is discussed by a comparison between Figs. 4.2 and 4.6, and Figs. 4.3 and 4.7. It is found that the behavior of the heat capacity of this two-energy-level system is highly similar to that of the effective thermodynamic system. Consequently, it can be concluded that the behavior of the heat capacity, $C_{Q,l}$, in the RN-dS spacetime is primarily determined by the heat capacity, $\hat{C}_{Q,l}$, of the two-energy-level system.

From Fig. 4.7 it can be observed that the behavior of $\hat{C} \rightarrow 0$ when $T_{eff} \rightarrow 0$ is obeyed by the quantum system with a split spectrum (two energy levels) in the RN-dS spacetime. The specific heat capacity in the RN-dS spacetime shows an extreme value, and is in accordance with the universal relation Eq. (4.2) for the two energy levels, i.e. the well-known Schottky specific heat. The significance of ordinary two-level systems is that many systems are treated as two-level systems. In general, the two thermodynamic horizons with different radiation temperatures exist in dS spacetime. Therefore, the two horizons may be treated as two energy levels in an effective thermodynamic system, with the effective thermodynamic quantities employed to describe the behavior of heat capacity of the effective system. This leads to the question of whether the heat capacity of this two-energy-level quantum system has a behavior of

RN-dS spacetime with the heat capacity described by the effective thermodynamic quantity. Furthermore, it would be beneficial to ascertain whether the behavior of heat capacity of thermodynamic systems in RN-dS spacetime is universal to dS spacetime. Additionally, it would be advantageous to determine the effect of different parameters in different dS spacetimes on the aforementioned behavior. A comprehensive investigation of this topic would provide a deeper understanding of the quantum properties of black holes. At the same time, it would also suggest a new avenue for research into the thermodynamic properties of dS spacetime. This result encapsulates the quantum properties of RN-dS spacetime and offers a novel avenue for further in-depth investigation into the quantum properties of black holes and dS spacetime.

Acknowledgments

We would like to thank Prof. Ren Zhao and Meng-Sen Ma for their indispensable discussions and comments. This work was supported by the Natural Science Foundation of China (Grant No. 12375050, Grant No. 11705106, Grant No. 12075143), the Scientific Innovation Foundation of the Higher Education Institutions of Shanxi Province (Grant Nos. 2023L269), the Science Foundation of Shanxi Datong University (2022Q1, 2020Q2), the Natural Science Foundation of Shanxi Province (Grant No. 202203021221209, Grant No. 202303021211180) and the Teaching Reform Project of Shanxi Datong University (XJG2022234).

-
- [1] The Event Horizon Telescope Collaboration et al, First M87 Event Horizon Telescope Results. I. The Shadow of the Supermassive Black Hole, *ApJL* 875 L1 (2019)
 - [2] The Event Horizon Telescope Collaboration et al, First M87 Event Horizon Telescope Results. II. Array and Instrumentation, *ApJL* 875 L2 (2019)
 - [3] Hawking S. W, *Commun. Math. Phys* 43(3), 199-220 (1975)
 - [4] Bekenstein J. D, *Black Holes and Entropy*, *Phys. Rev. D* 7, 2333-2346 (1973)
 - [5] D. Kubiznak, R. B. Mann, P-V criticality of charged AdS black holes, *JHEP* 1207, 033 (2012), arXiv:1205.0559
 - [6] R.-G. Cai, L.-M. Cao, L. Li, R.-Q. Yang, P-V criticality in the extended phase space of Gauss-Bonnet black holes in AdS space, *JHEP* 09, 005 (2013), arXiv:1306.6233[gr-qc]
 - [7] R.-G. Cai, S.-M. Ruan, S.-J. Wang, R.-Q. Yang, R.-H. Peng, Complexity Growth for AdS Black Holes, *JHEP* 09, 161 (2016), arXiv:1606.08307[gr-qc]
 - [8] J.-L. Zhang, R.-G. Cai, H.-W. Yu, Phase transition and Thermodynamical geometry of Reissner-Nordstrom-AdS Black Holes in Extended Phase Space, *Phys. Rev. D* 91, 044028 (2015), arXiv:1502.01428[hep-th]
 - [9] S. Gunasekaran, D. Kubiznak, R. B. Mann, Extended phase space thermodynamics for charged and rotating black holes and Born-Infeld vacuum polarization, *JHEP* 11, 110 (2012), arXiv:1208.6251
 - [10] A. M. Frassino, D. Kubiznak, R. B. Mann, F. Simovic, Multiple Reentrant Phase Transitions and Triple Points in Lovelock Thermodynamics, *JHEP* 09, 080 (2014), arXiv:1406.7015[hep-th]
 - [11] N. Altamirano, D. Kubiznak, R. B. Mann, Z. Sherkatghanad, Kerr-AdS analogue of triple point and solid/liquid/gas phase transition, *Class. Quant. Grav* 31, 042001 (2014), arXiv:1308.2672 [hep-th]
 - [12] J.-Y. Yang, R. B. Mann, Dynamic behaviours of black hole phase transitions near quadruple points, *JHEP* 08, 028 (2023), arXiv:2304.08969[gr-qc]
 - [13] J. Wu, R. B. Mann, Multicritical Phase Transitions in Lovelock AdS Black Holes, *Phys. Rev. D* 107, 084035 (2023), arXiv:2212.08087[hep-th]
 - [14] S.-W. Wei, Y.-P. Zhang, Y.-X. Liu, R. B. Mann, Static spheres around spherically symmetric black hole spacetime, *Phys. Rev. Res.* 5, 043050 (2023), arXiv:2303.06814[gr-qc]
 - [15] T. Durgut, R. A. Hennigar, H. K. Kunduri, R. B. Mann, Phase Transitions and Stability of Eguchi-Hanson-AdS Solitons, *JHEP* 03, 114 (2023), arXiv:2212.12685[gr-qc]
 - [16] J. Wu, R. B. Mann, Multicritical Phase Transitions in Multiply Rotating Black Holes, *Class. Quan. Grav* 40, 06LT01 (2023), arXiv:2208.00012[gr-qc]
 - [17] Y. Xiao, Y. Tian, Y.-X. Liu, Extended Black Hole Thermodynamics from Extended Iyer-Wald Formalism, *Phys. Rev. L* 132, 021401 (2024), arXiv:2308.12630
 - [18] M.-T. Wang, H.-Y. Jiang, Y.-X. Liu, Generalized Volume-Complexity for RN-AdS Black Hole, *JHEP* 07, 178 (2023), arXiv:2304.05751[hep-th]
 - [19] R. Zhao, H.-H. Zhao, M.-S. Ma, L.-C. Zhang, On the critical phenomena and thermodynamics of charged topological dilaton AdS black holes. *Eur. Phys. J. C* 73, 2645 (2013), arXiv:1305.3725
 - [20] S. H. Hendi, R. B. Mann, S. Panahiyan, B. E. Panah, van der Waals like behavior of topological AdS black holes in massive gravity. *Phys. Rev. D* 95, 021501(R) (2017), arXiv:1702.00432 [gr-qc]
 - [21] S. H. Hendi, S. Hajkhalili, M. Jamil, M. Momennia, Stability and phase transition of rotating Kaluza-Klein black holes, *Fort. der. Phys* 71, 8-9 (2023), arXiv:2204.11558[gr-qc]

- [22] M. Momennia, S. H. Hendi, Critical Phenomena and Reentrant Phase Transition of Asymptotically Reissner-Nordstrom Black Holes, *Phys. Lett. B* 822, 136692 (2021), arXiv:2101.12039[gr-qc]
- [23] S. H. Hendi, K. Jafarzade, Critical behavior of charged AdS black holes surrounded by quintessence via an alternative phase space, *Phys. Rev. D* 103, 104011 (2021), arXiv:2012.13271[gr-qc]
- [24] H. Laassiri, A. Daassou, R. Benbrik, Analytical Critical Phenomena of Rotating Bardeen AdS Black Holes with Dark Energy, arXiv:2401.00522[hep-th]
- [25] X.-Q. Li, H.-P. Yan, L.-L. Xing, S.-W. Zhou, Critical behavior of AdS black holes surrounded by dark fluid with Chaplygin-like equation of state, *Phys. Rev. D* 107, 104055 (2023), arXiv:2305.03028[gr-qc]
- [26] S. I. Kruglov, Rational nonlinear electrostatics of AdS black holes and extended phase space thermodynamics, *Eur. Phys. J. C* 82, 4, 292 (2022), arXiv:2210.06344[physics.gen-ph]
- [27] S. I. Kruglov, Nonlinearly charged AdS black holes, extended phase space thermodynamics and Joule-Thomson expansion, *Annals Phys.* 441, 168894 (2022), arXiv:2208.13662[physics.gen-ph]
- [28] Y.-Z. Du, H.-F. Li, Ren Zhao, Overview of thermodynamic properties for Reissner-Nordstrom-de Sitter spacetime in induced phase space, *Eur. Phys. J. C* 82, 859 (2022), arXiv:2207.03126[hep-th]
- [29] Y.-Z. Du, H.-F. Li, F. Liu, L.-C. Zhang, Photon orbits and phase transition for Non-Linear charged Anti-de Sitter black holes, *JHEP* 01, 137 (2023), arXiv:2204.01007[hep-th]
- [30] Z.-Y. Gao, X.-Q. Kong, L. Zhao, Thermodynamics of Kerr-AdS black holes in the restricted phase space, *Eur. Phys. J. C* 82, 112 (2022), arXiv:2112.08672[hep-th]
- [31] S. Ghaffari, G. G. Luciano, A. Sheykhi, Nonextensive entropies impact onto thermodynamics and phase structure of Kerr-Newman black holes, *Phys. Dark Univ.* 44 101447 (2024), arXiv:2312.13407[hep-th]
- [32] M.-Y. Zhang, H. Chen, H. Hassanabadi, Z.-W. Long, H. Yang, Critical behavior and Joule-Thomson expansion of charged AdS black holes surrounded by exotic fluid with modified Chaplygin equation of state, arXiv:2401.17589 [gr-qc]
- [33] D. J. Gogoi, Y. Sekhmani, D. Kalita, N. J. Gogoi, J. Bora, Joule-Thomson expansion and Optical behaviour of Reissner-Nordstrom-Anti-de Sitter black holes in Rastall gravity surrounded by a Quintessence field, *Fort. Der. Phy* 71, 2300010 (2023), arXiv:2306.02881[gr-qc]
- [34] M. Yasir, X. Tiecheng, F. Javed, G. Mustafa, Thermal analysis and Joule-Thomson expansion of black hole exhibiting metric-affine gravity, *Chin. Phys. C* 48, 015103 (2024), arXiv:2305.13709[gr-qc]
- [35] J. Barrientos, J. Mena, Joule-Thomson expansion of AdS black holes in quasi-topological electromagnetism, *Phys. Rev. D* 106, 044064 (2022), arXiv:2206.06018[gr-qc]
- [36] J. Sadeghi, M. R. Alipour, S. N. Gashti, M. Ali, S. Afshar, Cooling and heating regions of Joule-Thomson expansion for AdS black holes: Maxwell-power-Yang-Mills and Kerr Sen black holes, arXiv:2402.02257 [hep-th]
- [37] T. N. Hunga, C. H. Nam, Topology in thermodynamics of regular black strings with Kaluza-Klein reduction, *Eur. Phys. J. C* 83, 582 (2023), arXiv:2305.15910 [gr-qc]
- [38] N. J. Gogoi, P. Phukon, Thermodynamic topology of 4D dyonic AdS black holes in different ensembles, *Phys. Rev. D* 108, 066016 (2023), arXiv:2304.05695 [hep-th]
- [39] C. Fairros, and T. Sharqui, Topological Nature of Black Hole Solutions in dRGT Massive Gravity, *Int. J. Mod. Phys. A* 38, 2350133 (2023), arXiv:2304.028895 [hep-th]
- [40] D. Wu, S.-Y. Gu, X.-D. Zhu, Q.-Q. Jiang, S.-Z. Yang, Topological classes of thermodynamics of the static multi-charge AdS black holes in gauged supergravities, arXiv:2402.00106 [hep-th]
- [41] P. K. Yerra, C. Bhamidipati, S. Mukherji, Topology of critical points in boundary matrix duals, *JHEP* 03, 138 (2024), arXiv:2304.14988 [hep-th]
- [42] S.-W. Wei, Y.-X. Liu, Clapeyron equations and fitting formula of the coexistence curve in the extended phase space of charged AdS black holes, *Phys. Rev. D* 91, 044018 (2015), arXiv:1411.5749 [hep-th]
- [43] S.-W. Wei, Y.-X. Liu, Insight into the Microscopic Structure of an AdS Black Hole from Thermodynamical Phase Transition, *Phys. Rev. Lett.* 115, 111302 (2015), arXiv:1502.00386 [gr-qc]
- [44] S.-W. Wei, Y.-X. Liu, R. B. Mann, Repulsive Interactions and Universal Properties of Charged Anti-de Sitter Black Hole Microstructures, *Phys. Rev. Lett.* 123, 071103 (2019), arXiv:1906.10840[gr-qc]
- [45] D.-Y. Chen, J. Tao, X.-T. Yang, Attractive interactions in the microstructures of asymptotically flat black holes, *Phys. Dark. Universe* 42, 101379 (2023), arXiv:2311.02145[gr-qc]
- [46] O. Sokoliuk, S. Pradhan, A. Baransky, P. K. Sahoo, AdS black hole thermodynamics and microstructures from $f(Q)$ gravitation, *Fort. Phys* 72, 2300043 (2024), arXiv:2311.02145[gr-qc]
- [47] G. Ruppeiner, A. M. Sturzu, Black hole microstructures in the extremal limit, *Physical Review D* 108, 086004 (2023), arXiv:2304.06187[gr-qc]
- [48] S. Masood A. S. Bukhari, B. Pourhassan, H. Aounallah, L.-G. Wang, On the microstructure of higher-dimensional Reissner-Nordstrom black holes in quantum regime, *Class. Quant. Grav* 40, 225007 (2023), arXiv:2304.00940[gr-qc]
- [49] G.-R. Li, G.-P. Li, S. Guo, Phase transition grade and microstructure of AdS black holes in massive gravity, *Class. Quant. Grav* 39, 195011 (2022), arXiv:2304.00842[gr-qc]
- [50] X.-Y. Guo, H.-F. Li, L.-C. Zhang, R. Zhao, Microstructure and continuous phase transition of RN-AdS black hole. *Phys. Rev. D* 100, 064036 (2019), arXiv:1901.04703[gr-qc]
- [51] R.-G. Cai, Cardy-Verlinde formula and thermodynamics of black holes in de Sitter spaces. *Nucl. Phys. B* 628, 375-386 (2002), arXiv:hep-th/0112253
- [52] G. A. Marks, F. Simovic, R. B. Mann, Phase Transitions in 4D Gauss-Bonnet-de Sitter Black Holes, *Phys.Rev.D* 104, 104056 (2021), arXiv:2107.11352v1 [gr-qc]
- [53] Y. Sekiwa. Thermodynamics of de Sitter black holes: Thermal cosmological constant, *Phys. Rev. D* 73, 084009 (2006),

arXiv:hep-th/0602269

- [54] M. Urano, A. Tomimatsu, H. Saida, Mechanical First Law of Black Hole Spacetimes with Cosmological Constant and Its Application to Schwarzschild-de Sitter Spacetime. *Class. Quant. Grav* 26, 105010 (2009),
- [55] S. Mbarek, R. B. Mann, Reverse Hawking-Page Phase Transition in de Sitter Black Holes, *JHEP* 02, 103 (2019), arXiv:1808.03349[hep-th]
- [56] D. Kubiznak, F. Simovic, Thermodynamics of horizons: de Sitter black holes and reentrant phase transitions, *Class. Quant. Grav* 33, 245001 (2016), arXiv:1507.08630[hep-th]
- [57] F. Simovic, R. B. Mann, Critical Phenomena of Born-Infeld-de Sitter Black Holes in Cavities, *JHEP* 05, 136 (2019), arXiv:1904.04871 [gr-qc]
- [58] S. Haroon, R. A. Hennigar, R. B. Mann, F. Simovic, Thermodynamics of Gauss-Bonnet-de Sitter Black Holes, *Phys. Rev. D* 101, 084051 (2020), arXiv:2002.01567 [gr-qc]
- [59] F. Simovic, D. Fusco, R. B. Mann, Thermodynamics of de Sitter Black Holes with Conformally Coupled Scalar Fields, *JHEP* 02, 219 (2021), arXiv:2008.07593[gr-qc]
- [60] B. P. Dolan, D. Kastor, D. Kubiznak, R. B. Mann, J. Traschen, Thermodynamic Volumes and Isoperimetric Inequalities for de Sitter Black Holes, *Phys. Rev. D* 87, 104017 (2013), arXiv:1301.5926
- [61] S. Bhattacharya, A note on entropy of de Sitter black holes, *Eur. Phys. J. C* 76, 112 (2016), [arXiv:1506.07809]
- [62] J. McInerney, G. Satishchandana, J. Traschena, Cosmography of KNdS Black Holes and Isentropic Phase Transitions, *Class. Quantum Grav* 33, 105007 (2016), arXiv:1509.02343 [hep-th]
- [63] P. Kanti, T. Pappas, Effective Temperatures and Radiation Spectra for a Higher-Dimensional Schwarzschild-de-Sitter Black-Hole, *Phys. Rev. D* 96, 024038 (2017), arXiv:1705.09108 [hep-th]
- [64] L.-C. Zhang, R. Zhao, M.-S. Ma, Entropy of Reissner-Nordstrom-de Sitter black hole, *Phys. Lett. B* 761, 74-76 (2016),
- [65] L.-C. Zhang, R. Zhao, The entropic force in Reissner-Nordstrom-de Sitter spacetime. *Phys. Lett. B* 797, 134798 (2019),
- [66] Y.-B. Ma, Y. Zhang, L.-C. Zhang, L. Wu, Y.-M. Huang, Y. Pan, Thermodynamic properties of higher-dimensional dS black holes in dRGT massive gravity, *Euro. Phys. J. C* 80, 213 (2020), arXiv:2003.05483 [hep-th]
- [67] J. Ko, B. Gwak, Universality on thermodynamic relation with corrections in de Sitter black holes, *JHEP* 03, 072 (2024), arXiv:2312.17014 [gr-qc]
- [68] V. Chakrabhavi, M. Etheredge, Y. Qiu, J. Traschen, Constrained Spin Systems and KNdS Black Holes, *JHEP* 02, 231 (2024), arXiv:2311.07777 [hep-th]
- [69] P. R. Anderson, J. Traschen, Horizons and Correlation Functions in 2D Schwarzschild-de Sitter Spacetime, *JHEP* 192, (2022), arXiv:2012.08494[hep-th]
- [70] J. Dinsmore, P. Draper, D. Kastor, Y. Qiu, J. Traschen, Schottky Anomaly of de Sitter Black Holes, *Class. Quant. Grav* 37, 054001 (2020), arXiv:1907.00248[hep-th]
- [71] Y.-B. Du, X.-D. Zhang, Topological classes of black holes in de-Sitter spacetime, *Eur. Phys. J. C* 83, 927 (2023), arXiv:2303.13105 [gr-qc]
- [72] C. H. Nam, Non-linear charged dS black hole and its thermodynamics and phase transitions, *Eur. Phys. J. C* 78, 418 (2018),
- [73] F. Mellor, I. Moss, Black holes and gravitational instantons, *Class. Quant. Grav* 6, 1379 (1989),
- [74] C. V. Johnson, Specific Heats and Schottky Peaks for Black Holes in Extended Thermodynamics, *Class. Quant. Grav.* 37, 054003 (2020), arXiv:1905.00539 [hep-th]