

A Generalized Version of Chung’s Lemma and its Applications

Li Jiang

LIJIANG@LINK.CUHK.EDU.CN

Xiao Li

LIXIAO@CUHK.EDU.CN

Andre Milzarek

ANDREMILZAREK@CUHK.EDU.CN

Junwen Qiu

JUNWENQIU@LINK.CUHK.EDU.CN

School of Data Science

The Chinese University of Hong Kong, Shenzhen (CUHK-Shenzhen)

Guangdong 518172, P.R. China

Abstract

Chung’s lemma is a classical tool for establishing asymptotic convergence rates of (stochastic) optimization methods under strong convexity-type assumptions and appropriate polynomial diminishing step sizes. In this work, we develop a generalized version of Chung’s lemma, which provides a simple non-asymptotic convergence framework for a more general family of step size rules. We demonstrate broad applicability of the proposed generalized Chung’s lemma by deriving tight non-asymptotic convergence rates for a large variety of stochastic methods. In particular, we obtain partially new non-asymptotic complexity results for stochastic optimization methods, such as Stochastic Gradient Descent (SGD) and Random Reshuffling (RR), under a general (θ, μ) -Polyak-Lojasiewicz (PL) condition and for various step sizes strategies, including polynomial, constant, exponential, and cosine step sizes rules. Notably, as a by-product of our analysis, we observe that exponential step sizes can adapt to the objective function’s geometry, achieving the optimal convergence rate without requiring exact knowledge of the underlying landscape. Our results demonstrate that the developed variant of Chung’s lemma offers a versatile, systematic, and streamlined approach to establish non-asymptotic convergence rates under general step size rules.

Keywords: Chung’s lemma, non-asymptotic convergence rates, stochastic methods, exponential and cosine step sizes, (θ, μ) -Polyak-Lojasiewicz condition

1 Introduction

Chung’s lemma (Chung (1954)) is widely utilized to derive asymptotic convergence rates of stochastic optimization methods; see, e.g., Bottou et al. (2018); Gürbüzbalaban et al. (2019); Nguyen et al. (2021); Gürbüzbalaban et al. (2021); Li et al. (2023). In the pioneering paper (Chung (1954)), Chung analyzed the stochastic gradient method (SGD, Robbins and Monro (1951)) using the polynomial step size rule $\alpha_k \sim 1/k^p$. Let $\{\mathbf{x}^k\}_k$ and \mathbf{x}^* denote the sequence of iterates generated by SGD and the corresponding optimal solution, respectively. The recursion for $a_k := \mathbb{E}[\|\mathbf{x}^k - \mathbf{x}^*\|^2]$ analyzed in Chung (1954) is then given by

$$a_{k+1} \leq \left(1 - \frac{c}{k^p}\right) a_k + \frac{d}{k^{p+q}}, \quad \text{where } p \in (0, 1], \quad c, d, q > 0. \quad (1)$$

A crucial task in the convergence analysis is to study how fast $\{a_k\}_k$ converges to 0. Chung provided a useful tool as follows (see Lemma 23 for a full version).

Chung’s Lemma (Informal)

Let $\{a_k\}_k \subseteq \mathbb{R}_{++}$ be given as in (1). If $0 < p < 1$ or $p = 1, c > q$, then $a_k = \mathcal{O}(k^{-q})$.

The classical version of Chung’s lemma only applies to polynomial step sizes. In order to go beyond such a specific step size rule and to encompass a broader class of strategies (e.g., exponential and cosine step size rules), we consider the following general recursion:

$$a_{k+1} \leq \left(1 - \frac{1}{s(b_k)}\right) a_k + \frac{1}{t(b_k)}, \tag{2}$$

where $\{b_k\}_k \subseteq \mathbb{R}_{++}$ is given and $s, t : \mathbb{R}_{++} \rightarrow \mathbb{R}_{++}$ are continuously differentiable functions. Clearly, $s(b_k)$ and $t(b_k)$ generalize the terms k^p/c and k^{p+q}/d in (1). Existing strategies for solving recursions of the form (2) mainly rely on explicit expansions:

$$a_{K+1} \leq a_0 \prod_{k=0}^K \left(1 - \frac{1}{s(b_k)}\right) + \sum_{k=0}^K \left[\prod_{i=k+1}^K \left(1 - \frac{1}{s(b_i)}\right) \right] \frac{1}{t(b_k)}, \tag{3}$$

where $\prod_{i=K+1}^K (1 - \frac{1}{s(b_i)}) := 1$. While this methodology can seemingly deal with any choice of s and t , it requires to derive tight bounds of the different appearing product and sum terms, which can be highly nontrivial and case-dependent; see, e.g., Li et al. (2021). Based on these observations, our work is motivated by the following core question:

Can we provide a *simple, non-asymptotic* upper bound for the sequence $\{a_k\}_k$ defined in (2) (as in the classic Chung’s lemma) for more *general step size rules*, i.e., for general $s(b_k), t(b_k)$?

1.1 Contributions

In this work, we address the latter core question affirmatively by imposing an additional (mild) structure on s and t . We now list our main contributions:

Generalized Chung’s Lemma. We derive a novel generalized variant of Chung’s Lemma to characterize the growth behavior of $\{a_k\}_k$. An informal version is shown below:

Generalized Chung’s Lemma (Informal Theorem 1)

Suppose that $\{a_k\}_k \subseteq \mathbb{R}_{++}$ follows the recursion (2) with $r := s/t$ being convex on some interval I and $\{b_k\}_k \subseteq I$. Then, under certain conditions, we have for all $k \geq 0$

$$a_{k+1} \leq \underbrace{C_1 \cdot r(b_{k+1})}_{\mathcal{T}\text{-induced rate}} + \underbrace{C_2 a_0 \cdot \prod_{i=0}^k \left[1 - \frac{1}{s(b_i)}\right]}_{\mathcal{S}\text{-induced rate}}, \quad \text{for some } C_1, C_2 > 0. \tag{4}$$

As indicated in the informal theorem, our generalized variant of Chung’s lemma naturally decomposes the rate bound into two components, a \mathcal{T} - and an \mathcal{S} -induced rate. The

\mathcal{T} -induced (often “slower”) rate is linked with the “error term” $1/t(b_k)$ which, in turn, is typically associated with stochastic gradient errors or other types of errors. The \mathcal{S} -induced (often “faster”) part represents the convergence rate derived from recursion (2) without the error term $1/t(b_k)$ and is solely connected to the mapping s . In Corollary 6, we present evidence demonstrating that the \mathcal{S} -induced term usually converges at a faster rate compared to the \mathcal{T} -induced part. Thus, the convergence rate of $\{a_k\}_k$ in (2) is $\mathcal{O}(s(b_k)/t(b_k)) = \mathcal{O}(r(b_k))$, which generalizes Chung’s classical result $\mathcal{O}(k^p/k^{p+q}) = \mathcal{O}(k^{-q})$ for recursion (1). Overall, our generalized Chung’s lemma provides a simple tool to obtain *non-asymptotic convergence rates* based on the recursion (2). As a direct application, we utilize Theorem 1 to derive a non-asymptotic extension of the classical asymptotic Chung’s Lemma (cf. Theorem 24).

(θ, μ) -PL Condition, Exponential and Cosine Step Sizes. Building on the generalized Chung’s lemma, we provide non-asymptotic analyses for SGD and the random reshuffling method (RR, Algorithm 2) with exponential and cosine step size rules under a general (θ, μ) -PL condition; see Theorems 13 to 16. Note that the existing analyses for these step size rules primarily focus on SGD with $\theta = \frac{1}{2}$. We recover existing complexity bounds and improve the convergence rates for cosine step sizes. Moreover, we establish new results for general PL-exponents $\theta \in (\frac{1}{2}, 1]$. A detailed comparison of our findings to the existing ones is displayed in Table 1. From a technical standpoint, there are two main challenges. The first challenge is that under the general (θ, μ) -PL condition, we need to analyze a more intricate recursion,

$$y_{k+1} \leq (1 + \ell_1 \alpha_k^\tau) y_k - \ell_2 \alpha_k y_k^{2\theta} + \ell_3 \alpha_k^\tau, \tag{5}$$

which renders the non-asymptotic Chung’s Lemma inapplicable when $\theta \in (\frac{1}{2}, 1]$. We tackle this issue by constructing auxiliary sequences, for which our generalized Chung’s lemma offers important instructional guidance, to simplify (5) to a recursion of the form (2). We refer to Section 3.2 for more details. The analysis of exponential and cosine step size rules presents additional challenges, i.e., the required assumptions for applying the generalized Chung’s lemma only hold for part of the iterates, but not all of them. We introduce a novel splitting technique by combining the generalized Chung’s lemma with an extension lemma (see Lemma 5) to resolve this problem. See Section 3.3 for more details.

Polynomial and Constant Step Size Rules. For completeness, we also provide non-asymptotic analyses for SGD and RR with polynomial and constant step size rules under the (θ, μ) -PL condition based on our generalized Chung’s lemma; see Theorems 17, 18, 20 and 21. Our findings not only corroborate the established results in the literature but also introduce several new ones, particularly for scenarios where $\theta \in (\frac{1}{2}, 1]$. A comparison between our results and the existing body of work is provided in Table 1.

In summary, our generalized Chung’s lemma serves as a versatile tool, providing a systematic way to establish convergence rates and complexity bounds for various optimization methods with different step sizes rules.

1.2 Related Works

Recursions of the form (2) are abundant in stochastic optimization; see, e.g., Mishchenko et al.; Nguyen et al. (2021); Bottou et al. (2018); Shamir (2016); Gürbüzbalaban et al. (2019, 2021); Ahn et al. (2020); Huang et al. (2021). As mentioned, the functions s, t in (2) are often related to the step size dynamics. Our overview focuses on Chung’s lemma and its

Algorithms		SGD		RR	
θ	step size	existing	ours	existing	ours
$\frac{1}{2}$	Exp	$\mathcal{O}(\frac{\log(K)}{K})$ (a)	$\mathcal{O}(\frac{\log(K)}{K})$	–	$\mathcal{O}(\frac{\log^2(K)}{NK^2})$
	Cos (b)	$\mathcal{O}(\frac{1}{K^{2/3}})$ (c)	$\mathcal{O}(\frac{\log^{1/(2\beta+1)}(K)}{K})$	–	$\mathcal{O}(\frac{\log^{2/(2\beta+1)}(NK)}{NK^2})$
	Const	$\mathcal{O}(\frac{\log(K)}{K})$ (d)	$\mathcal{O}(\frac{\log(K)}{K})$	$\mathcal{O}(\frac{\log^2(NK)}{NK^2})$ (e)	$\mathcal{O}(\frac{\log^2(NK)}{NK^2})$
	Poly	$\mathcal{O}(K^{-1})$ (f)	$\mathcal{O}(K^{-1})$	$\mathcal{O}(\frac{\log(K)}{NK^2})$ (g)	$\mathcal{O}(\frac{\log^2(N)}{NK^2})$
$(\frac{1}{2}, 1]$ (h)	Exp	–	$\mathcal{O}([\frac{\log(K/\beta)}{K}]^{\omega_1})$	X	$\mathcal{O}([\frac{\log(K/\beta)}{\sqrt{NK}}]^{\omega_2})$
	Cos	–	$\mathcal{O}([\frac{\log^{v_1}(K)}{K}]^{\omega_1})$		$\mathcal{O}([\frac{\log^{v_2}(NK)}{\sqrt{NK}}]^{\omega_2})$
	Const	–	$\mathcal{O}([\frac{\log(K)}{K}]^{\omega_1})$		$\mathcal{O}([\frac{\log(NK)}{\sqrt{NK}}]^{\omega_2})$
	Poly	$\mathcal{O}(K^{-\omega_1})$ (i)	$\mathcal{O}(K^{-\omega_1})$		$\mathcal{O}([\frac{\log(N)}{\sqrt{NK}}]^{\omega_2})$

Table 1: Comparison of our results with existing literature.

(a) See (Wang et al., 2021, Theorem 4.1). The result in Wang et al. (2021) is achieved under stricter assumptions compared to ours, necessitating f to be strongly convex and L -smooth, along with a bounded stochastic gradient condition.

(b) We use cosine step sizes of the form $\alpha_k = \alpha[\frac{1+\cos(\frac{k\pi}{K})}{2}]^\beta$, $\beta > 0$; Existing literature only covers the case when $\beta = 1$.

(c) See (Li et al., 2021, Theorem 2). We improve this rate by suggesting a particular method for selecting α in cosine step sizes, i.e., setting $\alpha \sim \frac{\log(K)}{K}$.

(d) See Shamir and Zhang (2013); Needell et al. (2014); Bottou et al. (2018); Gower et al. (2019).

(e) See (Nguyen et al., 2021, Theorem 1).

(f) See (Gower et al., 2021, Theorem 4.8).

(g) See (Nguyen et al., 2021, Theorem 7). Here, a direct comparison of our result is not possible due to some minor mistakes in the proof of (Nguyen et al., 2021, Theorem 7), which cannot be easily remedied. In Nguyen et al.’s proof of Theorem 7, the step size $\eta_t = \frac{2}{t+\beta}$ is selected, where $\beta = 1 + \frac{1}{N}$. The choice of β is critical as $\beta - 1$ is required to yield the factor $1/N$ in the convergence rate. However, we identified a missing parameter, ρ , in the selection of η_t . According to (Nguyen et al., 2021, Lemma 1), η_t should satisfy $\frac{q}{\rho(t+\beta)}$, where $q = 2$ and $\rho = \frac{1}{4\tau}$, i.e., $\rho = \frac{\mu}{2}$ in our notation. Hence, it holds that $\eta_t = 4(\mu(t + 1 + \frac{1}{N}))^{-1}$. Meanwhile, the proof of (Nguyen et al., 2021, Theorem 7) requires η_t to satisfy $\eta_t \leq (L\sqrt{2(\Theta/N + 1)})^{-1}$ for all $t \geq 1$. However, this leads to a contradiction requiring the condition number to satisfy $\frac{1}{\mu} \leq \frac{1}{4\sqrt{2}}(2 + \frac{1}{N}) < 1$. It is important to note that the rescaling argument in Nguyen et al. (2021) is not applicable in this context because the condition number is invariant under rescaling. One straightforward solution is to increase β ; however, this adjustment would eliminate the factor $\frac{1}{N}$ in the convergence rate in their proof.

(h) Here, we use the notation $\omega_1 = \frac{1}{4\theta-1}$, $\omega_2 = \frac{1}{3\theta-1}$, $\varrho_1 = \frac{2\theta}{4\theta-1}$, $\varrho_2 = \frac{\theta}{3\theta-1}$, $v_1 = \frac{\varrho_1}{2\beta+\varrho_1}$ and $v_2 = \frac{\varrho_2}{2\beta+\varrho_2}$.

(i) See (Fontaine et al., 2021, Theorem 12); (Fatkhullin et al., 2022, Corollary 1).

extensions, different step size rules, and convergence analyses based on PL-type conditions.

► **Chung’s Lemma and Extensions.** Chung’s lemma (Chung (1954)) is a fundamental tool extensively used in the literature to analyze the convergence rate of stochastic approximation and optimization algorithms, (Chung (1954); Derman (1956); Sacks (1958); Fabian

(1968); Nemirovski et al. (2009); Moulines and Bach (2011); Gürbüzbalaban et al. (2019); Karimi et al. (2016); Bottou et al. (2018); Gürbüzbalaban et al. (2021); Ahn et al. (2020); Nguyen et al. (2021); Fatkhullin et al. (2022)). Non-asymptotic extensions of Chung’s lemma are established in (Moulines and Bach (2011); Ahn et al. (2020); Nguyen et al. (2021)), which are tailored to polynomial step size rules. In addition, some studies have considered the recursion (2) with specific non-polynomial forms of $s(b_k)$ and $t(b_k)$, see (Li et al. (2021); Wang et al. (2021)). To the best of our knowledge, the literature appears to lack non-asymptotic variants of Chung’s lemma that accommodate fully general terms $s(b_k)$ and $t(b_k)$.

► **Exponential and Cosine Step Sizes.** Exponential step sizes (Li et al. (2021)), also known as geometrically decaying step sizes (Goffin (1977); Ge et al. (2019)), were utilized by Goffin (1977) for subgradient methods. This step size strategy is also widely used in machine learning, and is currently one of the learning rate decay schedules in Tensorflow (Abadi et al. (2015)) and PyTorch (Paszke et al. (2019)). The theoretical analysis of exponential step size is limited. In Ge et al. (2019), the authors provide derivations for least-squares regression problems. Theoretical and broader guarantees for a more general class of functions are shown in Li et al. (2021); Wang et al. (2021). Cosine step sizes, initially proposed by Loshchilov and Hutter (2016), are used as a popular warm restart scheme in performing stochastic algorithms. Recently, cosine step sizes have been widely employed in the optimization and training of large language models. For instance, the Llama-2 model family (Touvron et al. (2023)) is pre-trained using an AdamW optimizer with a cosine step size rule. Its fine-tuning procedure uses the same step size scheme. Therefore, understanding the convergence properties of representative stochastic optimization methods with this step size rule is crucial. Yet, the convergence behavior of stochastic methods under cosine step sizes has not been well understood. The first complexity bounds for SGD with cosine step sizes are presented in (Li et al., 2021, Theorem 2). In the general nonconvex setting, extensions to an RR-type method (with momentum) are considered in (Tran et al., 2021, Corollary 4).

► **Stochastic Optimization under PL-type Conditions.** The Polyak-Łojasiewicz (PL) condition (Polyak (1963); Łojasiewicz (1959, 1963)) provides a measure of the “strong convexity-like” behavior of a function without requiring it to be convex. The recent convergence frameworks discussed in Fontaine et al. (2021); Fatkhullin et al. (2022) rely on certain generalizations of the PL condition. In this paper, we consider a (θ, μ) -PL condition (Definition 9), which encompasses the well-known Polyak-Łojasiewicz condition when $\theta = \frac{1}{2}$. The additional PL exponent, $\theta \in [\frac{1}{2}, 1]$, enables the PL condition to hold for a wider range of functions, particularly those relevant to contemporary machine learning applications (see (Fatkhullin et al., 2022, Appendix A) for pertinent examples). Specifically, in the field of reinforcement learning, the objective function may satisfy the $(1, \mu)$ -PL condition for a certain $\mu > 0$, (Wang and Zou (2022); Agarwal et al. (2021); Mei et al. (2020)). These evidences indicate that analyzing the cases beyond the standard setting $\theta = \frac{1}{2}$ holds significance in real applications.

2 Generalized Chung's Lemma

Throughout this work, we concentrate on the non-asymptotic behavior of sequence $\{a_k\}_k$ satisfying

$$a_{k+1} \leq \left(1 - \frac{1}{s(b_k)}\right) a_k + \frac{1}{t(b_k)}, \quad \forall k \in [K] := \{0, 1, \dots, K\}. \quad (6)$$

The following generalized Chung's lemma provides a non-asymptotic upper bound for $\{a_k\}_k$.

Theorem 1 (Generalized Chung's Lemma) *Let $\{a_k\}_k \subseteq \mathbb{R}_{++}$, $\{b_k\}_k \subseteq \mathbb{R}$ and the continuous differentiable functions $s, t : \mathbb{R} \rightarrow \mathbb{R}_{++}$ be given. Assume that $\{a_k\}_k$ follows (6) and there is an interval I such that $b_k \in I$ for all $k \in [K]$, $s(x) \geq 1$ for all $x \in I$, and $r := s/t$ is convex on I . Define $u := s' - t'r$; if either of the following statements holds:*

(a) *There exists a sequence $\{\lambda_k\}_k \subseteq \mathbb{R}_+$ such that for all $k \in [K-1]$,*

$$\lambda_{k+1}(s(b_k) + (b_{k+1} - b_k)u(b_k)) \geq \lambda_k(s(b_k) - 1) + 1, \quad (7)$$

(b) *There exists a non-decreasing sequence $\{\lambda_k\}_k \subseteq \mathbb{R}_{++}$ such that for all $k \in [K-1]$,*

$$(b_{k+1} - b_k)u(b_k) \geq -1 + \frac{1}{\lambda_{k+1}}, \quad (8)$$

then we have

$$a_{k+1} \leq \lambda_{k+1}r(b_{k+1}) + (a_0 - \lambda_0r(b_0)) \prod_{i=0}^k \left(1 - \frac{1}{s(b_i)}\right), \quad \forall k \in [K-1]. \quad (9)$$

Proof Let us set $\beta_k := b_{k+1} - b_k$. By the convexity of the mapping r , we have

$$r(b_{k+1}) \geq r(b_k) + r'(b_k)\beta_k = r(b_k) + \beta_k \frac{s'(b_k)t(b_k) - s(b_k)t'(b_k)}{t(b_k)^2} = \frac{\beta_k u(b_k) + s(b_k)}{t(b_k)}. \quad (10)$$

If statement (a) holds, substituting the latter estimate into (7), this yields

$$\begin{aligned} \lambda_{k+1}r(b_{k+1}) &\geq \frac{1}{t(b_k)} \cdot \lambda_{k+1} (\beta_k u(b_k) + s(b_k)) \geq \lambda_k \left(\frac{s(b_k)}{t(b_k)} - \frac{1}{t(b_k)} \right) + \frac{1}{t(b_k)} \\ &= \lambda_k \left(r(b_k) - \frac{r(b_k)}{s(b_k)} \right) + \frac{1}{t(b_k)} = \lambda_k r(b_k) \left(1 - \frac{1}{s(b_k)} \right) + \frac{1}{t(b_k)}. \end{aligned}$$

Hence, invoking the recursion (6), we obtain $a_{k+1} - \lambda_{k+1}r(b_{k+1}) \leq (1 - \frac{1}{s(b_k)})(a_k - \lambda_k r(b_k))$. This verifies (9).

When statement (b) is valid, we notice that the condition (8) readily implies (7), then (9) follows. \blacksquare

We now provide a simple example to briefly illustrate the application of Theorem 1.

Example 1 *Let us consider the recursion $a_{k+1} \leq (1 - \alpha)a_k + \beta$, where $\alpha \in (0, 1)$, $\beta > 0$. Setting $I = \mathbb{R}$, $s \equiv \frac{1}{\alpha}$, $t \equiv \frac{1}{\beta}$, $b_k = k$, $\lambda = 1$, the rate mapping $r \equiv \frac{\beta}{\alpha}$ is convex and we have $(b_{k+1} - b_k)u(b_k) = 0 = -1 + \frac{1}{\lambda}$ for all k . Consequently, Theorem 1 is applicable, then $a_k \leq \frac{\beta}{\alpha} + (1 - \alpha)^k(a_0 - \frac{\beta}{\alpha})$ holds for all k .*

Remark 2 *The most significant and useful setting for Theorem 1 arises when the sequence $\{\lambda_k\}_k$ in condition (b) can be chosen to be constant. A majority of our subsequent analysis will be built based on this particular scenario.*

Remark 3 *The convexity requirement on $r := s/t$ in Theorem 1 is not strict since there is flexibility in choosing $\{b_k\}_k$. Consider the example $r(b_k) = 1 + \cos(\frac{k\pi}{K})$, $k \in [K]$. A direct choice $b_k := k$, $r(x) := 1 + \cos(\frac{x\pi}{K})$, and $I = [0, K]$ does not meet the convexity condition. However, we can ensure the convexity of r by instead selecting $b_k := 1 + \cos(\frac{x\pi}{K})$, $r(x) := x$, and $I = \mathbb{R}$. This strategy proves valuable in the analysis of cosine step sizes, see Section 3.3.*

Non-asymptotic Extension of Chung's Lemma. As a quick application, we use Theorem 1 to obtain a non-asymptotic extension of the classical Chung's lemma.

Lemma 4 (Non-asymptotic Chung's Lemma) *Let $\{a_k\}_k \subseteq \mathbb{R}_{++}$ be given satisfying*

$$a_{k+1} \leq \left(1 - \frac{c}{(k+\gamma)^p}\right) a_k + \frac{d}{(k+\gamma)^{p+q}}, \quad p \in (0, 1], \quad q > 0, \quad c, d > 0, \quad \gamma \geq c^{\frac{1}{p}}.$$

Then, the following statements hold:

(a) *If $p \in (0, 1)$, $\gamma \geq c^{\frac{1}{p}}$ and $\gamma > (\frac{q}{c})^{\frac{1}{1-p}}$, then*

$$a_{k+1} \leq \frac{\lambda d}{c} (k+1+\gamma)^{-q} + \left(a_0 - \frac{\lambda d}{c\gamma^q}\right)^+ \exp\left(\frac{c\gamma^{1-p}}{1-p}\right) \exp\left[-\frac{c(k+1+\gamma)^{1-p}}{1-p}\right],$$

holds for all $k \in [K]$, where $\lambda = \frac{c\gamma^{1-p}}{c\gamma^{1-p}-q}$ and $(x)^+ := \max\{x, 0\}$.

(b) *If $p = 1$, $c > q$, and $\gamma \geq c$, then*

$$a_{k+1} \leq \frac{d}{c-q} (k+1+\gamma)^{-q} + \gamma^c \left(a_0 - \frac{d}{(c-q)\gamma^q}\right)^+ (k+1+\gamma)^{-c}.$$

Here, we only discuss the cases $0 < p < 1$ and $p = 1$, $c > q$, as these are most relevant to the convergence analysis of stochastic optimization methods. A full derivation is provided in Appendix B.

Proof We now apply our main theorem (especially Theorem 1 (b)) to prove the non-asymptotic version of the classical Chung's lemma. To make use of Theorem 1 (b), we first specify the following sequences and functions:

$$b_k := k + \gamma, \quad s(x) := x^p/c, \quad \text{and} \quad t(x) := x^{p+q}/d.$$

We only need to check if the conditions in Theorem 1 (b) are fulfilled. Let us note that the mapping $r(x) := s(x)/t(x) = \frac{d}{c}x^{-q}$ is convex on \mathbb{R}_{++} . In addition, it holds that

$$u(x) := s'(x) - r(x)t'(x) = \frac{p}{c}x^{p-1} - \frac{d}{c}x^{-q} \cdot \frac{p+q}{d}x^{p+q-1} = -\frac{q}{c}x^{p-1}$$

and $s(x) \geq 1$ for all $x \geq c^{1/p}$. By assumption, we have $\gamma \geq c^{1/p}$ implying $b_k \in I := [c^{1/p}, \infty) \subseteq \mathbb{R}_{++}$ for all $k \in [K]$. Now, it only remains to verify the condition (8). Since $b_{k+1} - b_k = 1$, we need to find a non-decreasing sequence $\{\lambda_k\}_k \subseteq \mathbb{R}_+$ such that

$$-\frac{q}{c}(k+\gamma)^{p-1} \geq -1 + \frac{1}{\lambda_{k+1}}, \quad \forall k \in [K]. \quad (11)$$

Based on the parameter $p \in (0, 1]$, we now discuss different cases.

Case I: $p \in (0, 1)$. Let us set $\lambda_k \equiv \lambda = \frac{c\gamma^{1-p}}{c\gamma^{1-p}-q}$. Per assumption, we further have $\gamma > (\frac{q}{c})^{\frac{1}{1-p}}$ and thus, we can infer $\lambda > 0$ and

$$-\frac{q}{c}(k+\gamma)^{p-1} \geq -\frac{q}{c}\gamma^{p-1} = -1 + \frac{c\gamma^{1-p}-q}{c\gamma^{1-p}} = -1 + \frac{1}{\lambda} \quad \forall k \in [K]. \quad (12)$$

Hence, the condition (11) is satisfied for this choice of λ_k and λ . Utilizing Theorem 1 (b), the inequality $1-x \leq \exp(-x)$, and the integral test (Proposition 22), we obtain

$$\begin{aligned} a_{k+1} &\leq \lambda r(b_{k+1}) + (a_0 - \lambda r(\gamma)) \cdot \prod_{i=0}^k \left[1 - \frac{1}{s(b_i)} \right] \\ &= \frac{\lambda d}{c}(k+1+\gamma)^{-q} + \left(a_0 - \frac{\lambda d}{c\gamma^q} \right) \prod_{i=0}^k \left[1 - \frac{c}{(i+\gamma)^p} \right] \\ &\leq \frac{\lambda d}{c}(k+1+\gamma)^{-q} + \left(a_0 - \frac{\lambda d}{c\gamma^q} \right)^+ \exp \left(-\sum_{i=0}^k \frac{c}{(i+\gamma)^p} \right) \\ &\leq \frac{\lambda d}{c}(k+1+\gamma)^{-q} + \left(a_0 - \frac{\lambda d}{c\gamma^q} \right)^+ \exp \left(\frac{c\gamma^{1-p}}{1-p} \right) \exp \left(-\frac{c(k+1+\gamma)^{1-p}}{1-p} \right), \end{aligned}$$

where the last line is due to $\sum_{i=0}^k (i+\gamma)^{-p} \geq \int_0^{k+1} (x+\gamma)^{-p} dx = \frac{1}{1-p} [(k+1+\gamma)^{1-p} - \gamma^{1-p}]$.

Case II: $p = 1$, $c > q$, and $\gamma \geq c$. Setting $\lambda_k \equiv \lambda = \frac{c}{c-q}$, the condition (11) is obviously satisfied (cf. (12)). Repeating the core steps in **Case I**, we obtain

$$\begin{aligned} a_{k+1} &\leq \frac{\lambda d}{c}(k+1+\gamma)^{-q} + \left(a_0 - \frac{\lambda d}{c\gamma^q} \right) \prod_{i=0}^k \left[1 - \frac{c}{i+\gamma} \right] \\ &\leq \frac{d}{c-q}(k+1+\gamma)^{-q} + \gamma^c \left(a_0 - \frac{d}{(c-q)\gamma^q} \right)^+ (k+1+\gamma)^{-c}. \end{aligned}$$

Notice that the difference in the analysis only stems from the integral test (Proposition 22): $\sum_{i=0}^k \frac{1}{i+\gamma} \geq \int_0^{k+1} \frac{1}{x+\gamma} dx = \log(k+1+\gamma) - \log(\gamma)$. \blacksquare

Unfortunately, many of the following applications are not as simple as Example 1 or Lemma 4. In general, condition (7) in Theorem 1 (a) is less restrictive compared to (8). However, determining λ_k in (7) can be challenging as it requires the non-trivial construction of a new recursive sequence $\{\lambda_k\}_k$. It is worth noting that, in most cases, Theorem 1 (b) is easier to apply. Nevertheless, there may be situations where it is not possible to construct the sequence $\{\lambda_k\}_k$ in (8) for all $k \in [K]$, although it is feasible for a subset of $[K]$. This issue becomes evident when analyzing exponential and cosine step sizes (as discussed in Section 3.3). To address this potential limitation, we introduce the following important lemma, which significantly enhances the applicability of Theorem 1 (b).

Lemma 5 (Extension Lemma) *Let $\{a_k\}_k \subseteq \mathbb{R}_{++}$, $\{b_k\}_k \subseteq \mathbb{R}_+$ and the C^1 -functions $s, t : \mathbb{R} \rightarrow \mathbb{R}_{++}$ be given. Suppose that $\{a_k\}_k$ follows the recursion (6). Assume that $s(b_k) \geq$*

1, and $\{r(b_k)\}_k$ is non-increasing w.r.t. k for all $k \in [K]$. If there exist $k_0, \bar{K} \in [K-1]$, $k_0 \leq \bar{K}$, $C \in \mathbb{R}$, and $\bar{\lambda} \geq 1$ such that

$$a_{\bar{K}+1} \leq \bar{\lambda}r(b_{\bar{K}+1}) + C \prod_{i=k_0}^{\bar{K}} \left(1 - \frac{1}{s(b_i)}\right), \quad (13)$$

then, we have $a_K \leq \bar{\lambda}r(b_{\bar{K}+1}) + C \prod_{i=k_0}^{K-1} \left(1 - \frac{1}{s(b_i)}\right)$.

Proof There is nothing to show in the case $\bar{K} = K-1$. We can verify the result for $\bar{K} < K-1$ via induction. For any $k = \bar{K}+1, \dots, K-1$, if $a_k \leq \bar{\lambda}r(b_{\bar{K}+1}) + C \prod_{i=k_0}^{k-1} \left(1 - \frac{1}{s(b_i)}\right)$, and by applying (6), we obtain

$$\begin{aligned} a_{k+1} &\leq \left[\frac{1}{t(b_k)} + \left(1 - \frac{1}{s(b_k)}\right) \bar{\lambda}r(b_{\bar{K}+1}) \right] + C \prod_{i=k_0}^k \left(1 - \frac{1}{s(b_i)}\right) \\ &= \left[\frac{1}{s(b_k)}r(b_k) + \left(1 - \frac{1}{s(b_k)}\right) \bar{\lambda}r(b_{\bar{K}+1}) \right] + C \prod_{i=k_0}^k \left(1 - \frac{1}{s(b_i)}\right) \\ &\leq \bar{\lambda}r(b_{\bar{K}+1}) + C \prod_{i=k_0}^k \left(1 - \frac{1}{s(b_i)}\right), \end{aligned}$$

where we used $\bar{\lambda} \geq 1$ and $r(b_k) \leq r(b_{\bar{K}+1})$ for any $k \geq \bar{K}+1$ in the last step. Hence, by induction, it follows $a_K \leq \bar{\lambda}r(b_{\bar{K}+1}) + C \prod_{i=k_0}^{K-1} \left(1 - \frac{1}{s(b_i)}\right)$. \blacksquare

The combination of Theorem 1 and Lemma 5 is crucial in the subsequent applications. The key idea is to partition the index set $[K]$ into two subsets. Initially, we determine the largest value of \bar{K} that satisfies condition (8) for all $k \in [\bar{K}]$. By applying Theorem 1 (b), we can establish an upper bound for $a_{\bar{K}+1}$ resembling (13). Lemma 5 then enables us to extend this estimate from $a_{\bar{K}+1}$ to the final iterate a_K . This technique will be important when analyzing exponential and cosine step size rules (cf. Section 3.3).

Forgetting Initial Conditions. The following corollary offers insight into the relationship between the \mathcal{S} - and \mathcal{T} -induced components in (4).

Corollary 6 *Suppose that the assumptions in Theorem 1 (b) hold with $\lambda_k \equiv \lambda \geq 1$. Then, we have*

$$a_{k+1} \leq \lambda r(b_{k+1}) + \max \left\{ \frac{a_0}{r(b_0)} - \lambda, 0 \right\} \prod_{i=0}^k \left(1 - \frac{1/\lambda}{s(b_i) - 1 + 1/\lambda}\right) r(b_{k+1}), \quad \forall k \in [K].$$

Proof From Theorem 1 (b), we have for all $k \in [K]$,

$$a_{k+1} \leq \lambda r(b_{k+1}) + \max \left\{ \frac{a_0}{r(b_0)} - \lambda, 0 \right\} \prod_{i=0}^k \left(1 - \frac{1}{s(b_i)}\right) r(b_0)$$

$$= \lambda r(b_{k+1}) + \max \left\{ \frac{a_0}{r(b_0)} - \lambda, 0 \right\} \prod_{i=0}^k \left[\left(1 - \frac{1}{s(b_i)} \right) \frac{r(b_i)}{r(b_{i+1})} \right] r(b_{k+1}).$$

Combining equation (10) and the condition (8), we obtain

$$r(b_{i+1}) \geq \frac{(b_{i+1} - b_i)u(b_i) + s(b_i)}{t(b_i)} \geq \frac{s(b_i) - 1 + 1/\lambda}{t(b_i)} > 0.$$

Thus, we can establish the upper bound

$$\left(1 - \frac{1}{s(b_i)} \right) \frac{r(b_i)}{r(b_{i+1})} \leq \left(1 - \frac{1}{s(b_i)} \right) \frac{s(b_i)}{s(b_i) - 1 + 1/\lambda} = 1 - \frac{1/\lambda}{s(b_i) - 1 + 1/\lambda}.$$

Substituting this bound into the first estimate finishes the proof. ■

Corollary 6 shows that if the assumptions in Theorem 1 (b) are satisfied, then the \mathcal{S} -induced rate will generally be faster than the \mathcal{T} -induced rate, which can be quantified by the term $\prod_{i=0}^k \left(1 - \frac{1/\lambda}{s(b_i) - 1 + 1/\lambda} \right)$. This phenomenon is known as the “forgetting initial conditions” in the literature; but it is usually discussed for specific step size rules. For instance, in (Moulines and Bach (2011)), this phenomenon is studied for polynomial step sizes. Corollary 6 reveals that this property holds in a more general context.

3 Convergence Rates of Stochastic Methods under (θ, μ) -PL Condition

In this section, we establish non-asymptotic rates for stochastic methods under various step size strategies through the application of the generalized Chung’s lemma—Theorem 1. We consider the optimization problem

$$\min_{x \in \mathbb{R}^n} f(x), \tag{14}$$

where $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is a continuously differentiable mapping.

3.1 Basic Assumptions and Descent Properties

Below, in Algorithms 1 and 2, we present the main algorithmic steps of SGD and the random reshuffling (RR) method to solve the problem (14). We note that, in RR, the objective function f is assumed to have the following finite-sum structure $f(x) := \frac{1}{N} \sum_{i=1}^N f_i(x)$.

Algorithm 1 Stochastic Gradient Descent (SGD)

- Require:** Choose an initial point $x^0 \in \mathbb{R}^n$ and step sizes $\{\alpha_k\}_k \subset \mathbb{R}_{++}$;
- 1: **while** stopping criterion is not met **do**
 - 2: Generate a new stochastic gradient $g^k \approx \nabla f(x^k)$;
 - 3: Perform the update $x^{k+1} = x^k - \alpha_k g^k$;
 - 4: **end while**
-

Algorithm 2 Random Reshuffling (RR)

Require: Choose an initial point $x^0 \in \mathbb{R}^n$ and step sizes $\{\alpha_k\}_k \subset \mathbb{R}_{++}$;

- 1: **while** stopping criterion not met **do**
 - 2: Generate a uniformly random permutation π^k of $\{1, \dots, N\}$ and set $\tilde{x}_0^k = x^k$;
 - 3: **for** $i = 1, 2, \dots, N$ **do**
 - 4: Update the inner iterate via $\tilde{x}_i^k = \tilde{x}_{i-1}^k - \frac{\alpha_k}{N} \nabla f_{\pi_i^k}(\tilde{x}_{i-1}^k)$;
 - 5: **end for**
 - 6: Perform the update $x^{k+1} = \tilde{x}_N^k$;
 - 7: **end while**
-

Our analysis will be based on a filtered probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_k\}_k, \mathbb{P})$ to which the stochastic processes $\{\mathbf{x}^k\}_k$ generated by these two algorithms are adapted. We make the following standard assumptions for SGD and RR, respectively.

Assumption 7 (Conditions for SGD) *We consider the assumptions:*

- (A.1) *The objective f is L -smooth, i.e., $\nabla f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is L -Lipschitz continuous.*
- (A.2) *The stochastic gradient $\mathbf{g}^k : \Omega \rightarrow \mathbb{R}^n$ is an unbiased estimator of $\nabla f(\mathbf{x}^k)$, and there exist $A, \sigma \geq 0, \bar{f} \in \mathbb{R}$ such that $\mathbb{E}[\|\mathbf{g}^k - \nabla f(\mathbf{x}^k)\|^2 \mid \mathcal{F}_{k-1}] \leq A(f(\mathbf{x}^k) - \bar{f}) + \sigma^2$ a.s..*

Assumption 8 (Conditions for RR) *We work with the conditions:*

- (B.1) *The individual component functions f_1, f_2, \dots, f_N are all L -smooth.*
- (B.2) *There exist constants $A, \sigma > 0, \bar{f} \in \mathbb{R}$ such that for all $x \in \mathbb{R}^n$, it holds that*

$$\frac{1}{N} \sum_{i=1}^N \|\nabla f_i(x) - \nabla f(x)\|^2 \leq A(f(x) - \bar{f}) + \sigma^2.$$

- (B.3) *The permutations $\{\pi^k\}_k$ are sampled independently (for each k) and uniformly without replacement from $\{1, \dots, N\}$.*

The Lipschitz smoothness in (A.1) and (B.1) is ubiquitous in the analysis of optimization methods, see, e.g., Bolte et al. (2014); Gürbüzbalaban et al. (2019); HaoChen and Sra (2019). The assumptions (A.2) and (B.2) have been extensively employed in the analysis of stochastic gradient-based methods (Nguyen et al. (2021); Mishchenko et al.; Shamir (2016); Sohl-Dickstein et al. (2014)). In the case $A = 0$, (A.2) and (B.2) reduce to bounded variance conditions. Furthermore, (B.2) is applicable in a wide range of scenarios. For instance, (B.2) is satisfied when each individual component function f_i is Lipschitz smooth and bounded from below, see (Mishchenko et al., Proposition 2).

For both SGD and RR, we will work with the following core property that allows characterizing the geometric landscape of f .

Definition 9 ((θ, μ)-PL Condition) *Given $\theta \in [\frac{1}{2}, 1]$ and $\mu > 0$, the function f is said to satisfy the (θ, μ)-PL condition if f is bounded from below by $f^* := \inf_{x \in \mathbb{R}^n} f(x) > -\infty$ and*

$$\|\nabla f(x)\| \geq \sqrt{2\mu}(f(x) - f^*)^\theta, \quad \forall x \in \mathbb{R}^n.$$

Definition 9 covers the well-known, classical Polyak-Lojasiewicz condition (Polyak (1963)) as a special case when $\theta = \frac{1}{2}$.

Assumption 10 (PL Condition for f) *We consider the following assumption:*

(P) *For some $\theta \in [\frac{1}{2}, 1]$, $\mu > 0$, the objective function f satisfies (θ, μ) -PL condition.*

In the following sections, we will employ Theorem 1 to establish the convergence rates for various sampling schemes, under the (θ, μ) -PL condition where $\theta \in [\frac{1}{2}, 1]$, with different step size schemes in stochastic gradient-based methods. Under assumption (P) and without loss generality, we set $\bar{f} = f^*$ in (A.2) and (B.2). We start with (approximate) descent-type properties for SGD and RR.

Lemma 11 (Descent-type Properties) *Suppose assumption (P) holds.*

(a) *Let $\{\mathbf{x}^k\}_k$ be generated by Algorithm 1 with step sizes $\alpha_k \in (0, \frac{1}{L}]$ and let the assumptions (A.1)–(A.2) be satisfied. Then, it holds that*

$$\mathbb{E}[f(\mathbf{x}^{k+1}) - f^*] \leq \left(1 + \frac{AL\alpha_k^2}{2}\right) \mathbb{E}[f(\mathbf{x}^k) - f^*] - \mu\alpha_k (\mathbb{E}[f(\mathbf{x}^k) - f^*])^{2\theta} + \frac{L\sigma^2\alpha_k^2}{2}. \quad (15)$$

(b) *Let the sequence $\{\mathbf{x}^k\}_k$ be generated by Algorithm 2 with step sizes $\alpha_k \in (0, \frac{1}{2L}]$ and let (B.1)–(B.3) hold. Then, we have*

$$\mathbb{E}[f(\mathbf{x}^{k+1}) - f^*] \leq \left(1 + \frac{AL^2\alpha_k^3}{2N}\right) \mathbb{E}[f(\mathbf{x}^k) - f^*] - \frac{\mu\alpha_k}{2} (\mathbb{E}[f(\mathbf{x}^k) - f^*])^{2\theta} + \frac{L^2\sigma^2\alpha_k^3}{2N}. \quad (16)$$

Proof For Algorithm 1, with assumptions (A.1)–(A.2), and $\alpha_k \in (0, \frac{1}{L}]$, we have

$$\mathbb{E}[f(\mathbf{x}^{k+1}) - f^*] \leq \left(1 + \frac{AL\alpha_k^2}{2}\right) \mathbb{E}[f(\mathbf{x}^k) - f^*] - \frac{\alpha_k}{2} \mathbb{E}[\|\nabla f(\mathbf{x}^k)\|^2] + \frac{L\sigma^2\alpha_k^2}{2}.$$

For Algorithm 2, we refer to the result in (Mishchenko et al., Appendix 9.5, Proof of Theorem 4, eq. (47)). With (B.1)–(B.3), for $\alpha_k \in (0, \frac{1}{2L}]$, we have

$$\mathbb{E}[f(\mathbf{x}^{k+1}) - f^*] \leq \left(1 + \frac{AL^2\alpha_k^3}{2N}\right) \mathbb{E}[f(\mathbf{x}^k) - f^*] - \frac{\alpha_k}{4} \mathbb{E}[\|\nabla f(\mathbf{x}^k)\|^2] + \frac{L^2\sigma^2\alpha_k^3}{2N}.$$

Combining $\|\nabla f(x)\|^2 \geq 2\mu(f(x) - f^*)^{2\theta}$ in (P) and Hölder's inequality, we obtain

$$\mathbb{E}[\|\nabla f(x)\|^2] \geq 2\mu\mathbb{E}[(f(x) - f^*)^{2\theta}] \geq 2\mu\mathbb{E}[(f(x) - f^*)]^{2\theta},$$

which finalizes the proof. ■

3.2 Main Recursion and Analysis Roadmap

From Lemma 11, it is evident that under the (θ, μ) -PL framework, the problem reduces to analyzing the recursion with the given form:

$$y_{k+1} \leq (1 + \ell_1 \alpha_k^\tau) y_k - \ell_2 \alpha_k y_k^{2\theta} + \ell_3 \alpha_k^\tau, \quad k \in [K-1], \quad (17)$$

where $y_k = \mathbb{E}[f(\mathbf{x}^k) - f^*]$. For SGD, we have $\ell_1 = \frac{A\mathbf{L}}{2}$, $\ell_2 = \mu$, $\ell_3 = \frac{\mathbf{L}\sigma^2}{2}$, $\tau = 2$, and for RR, it holds that $\ell_1 = \frac{A\mathbf{L}^2}{2N}$, $\ell_2 = \frac{\mu}{2}$, $\ell_3 = \frac{\mathbf{L}^2\sigma^2}{2N}$, $\tau = 3$. Recursion (17) generally lies beyond the scope of the generalized Chung's lemma in the case $\theta \neq \frac{1}{2}$. To address this discrepancy, we propose a solution that involves constructing an auxiliary sequence $\{u_k\}_k$, for which our generalized Chung's lemma offers valuable instructional guidance. Here, we provide a general roadmap for the analysis.

Step 1: Linearization and Relaxation. In the case $\theta \neq \frac{1}{2}$, the term " $\ell_2 \alpha_k y_k^{2\theta}$ " depends on y_k in a nonlinear fashion and hence, Chung's lemma is not immediately applicable to the recursion (17). This can be resolved by substituting " $y_k^{2\theta}$ " with a suitable expression that only depends linearly on y_k . Since the mapping $y \mapsto h_\theta(y) := y^{2\theta}$ is convex (on \mathbb{R}_+), this can be essentially accomplished by utilizing the estimate

$$y^{2\theta} \geq x^{2\theta} + 2\theta x^{2\theta-1}(y-x) \quad \forall x, y \geq 0.$$

Upon setting $y = y_k$ and $x = u_k$, this allows relaxing (17) to

$$\begin{aligned} y_{k+1} &\leq (1 + \ell_1 \alpha_k^\tau) y_k - \ell_2 \alpha_k [h_\theta(u_k) + 2\theta u_k^{2\theta-1} y_k - 2\theta h_\theta(u_k)] + \ell_3 \alpha_k^\tau \\ &= [1 + \ell_1 \alpha_k^\tau - 2\theta \ell_2 \alpha_k u_k^{2\theta-1}] y_k + [(2\theta - 1) \ell_2 \alpha_k u_k^{2\theta} + \ell_3 \alpha_k^\tau], \end{aligned} \quad (18)$$

where $\{u_k\}_{k \geq 0}$ is a (free) non-negative auxiliary sequence.

Step 2: Optimal Choice of $\{u_k\}_k$. The sequence $\{u_k\}_k$ should be chosen such that

1. The term " $\theta \ell_2 \alpha_k u_k^{2\theta-1}$ " is not smaller than " $\ell_1 \alpha_k^\tau$ " (for all k), then (18) can be further relaxed to

$$y_{k+1} \leq [1 - \theta \ell_2 \alpha_k u_k^{2\theta-1}] y_k + [(2\theta - 1) \ell_2 \alpha_k u_k^{2\theta} + \ell_3 \alpha_k^\tau]. \quad (19)$$

2. For recursion (19), the expected rate $r_k \sim \frac{(2\theta-1)\ell_2\alpha_k u_k^{2\theta} + \ell_3\alpha_k^\tau}{\alpha_k u_k^{2\theta-1}} = (2\theta - 1)\ell_2 u_k + \ell_3 \frac{\alpha_k^{\tau-1}}{u_k}$ is optimal (as fast as possible), cf. Theorem 1.

Optimizing r_k (as a function of u_k), we can obtain the following candidate for $\{u_k\}_k$:

$$u_k = (\ell_3/\ell_2)^{\frac{1}{2\theta}} \alpha_k^{\frac{\tau-1}{2\theta}}. \quad (20)$$

To satisfy the first requirement, we need $\theta \ell_2 \alpha_k u_k^{2\theta-1} \geq \ell_1 \alpha_k^\tau$, i.e., $\alpha_k \leq \left(\frac{\theta \ell_2^{\frac{1}{2\theta}} \ell_3^{\frac{2\theta-1}{2\theta}}}{\ell_1}\right)^{\frac{2\theta}{\tau-1}}$. Substituting (19) into (18), this yields

$$y_{k+1} \leq \left[1 - \theta \ell_2^{\frac{1}{2\theta}} \ell_3^{\frac{2\theta-1}{2\theta}} \alpha_k^{\frac{(2\theta-1)\tau+1}{2\theta}}\right] y_k + 2\theta \ell_3 \alpha_k^\tau.$$

To simplify the notations, let us define $\zeta := \theta \ell_2^{\frac{1}{2\theta}} \ell_3^{\frac{2\theta-1}{2\theta}}$, $\varrho := \frac{2\theta}{(2\theta-1)\tau+1}$, and $\omega := \frac{\tau-1}{(2\theta-1)\tau+1}$. Then, we conclude that for all

$$\alpha_k \leq \left(\frac{\zeta}{\ell_1} \right)^{\frac{\varrho}{\omega}}, \quad (21)$$

we have

$$y_{k+1} \leq [1 - \zeta \alpha_k^{1/\varrho}] y_k + 2\theta \ell_3 \alpha_k^\tau. \quad (22)$$

The simplified and relaxed recursion (22) is the basis of our final step and of the application of the generalized Chung's lemma.


Step 3: Application of Chung's Lemma to (22). To facilitate the analysis, we need to assume $\zeta \alpha_k^{1/\varrho} \leq 1$, i.e.,

$$\alpha_k \leq \zeta^{-\varrho}. \quad (23)$$

Moreover, suppose that the step sizes α_k can be represented as a function $\eta : \mathbb{R} \rightarrow \mathbb{R}_+$ evaluated at b_k , i.e., $\alpha_k = \eta(b_k)$. Then, in Theorem 1, we can set

$$s(x) = \frac{1}{\zeta} \eta(x)^{-\frac{1}{\varrho}}, \quad t(x) = \frac{1}{2\theta \ell_3} \eta(x)^{-\tau}, \quad r(x) = 2 \left(\frac{\ell_3}{\ell_2} \right)^{\frac{1}{2\theta}} \eta(x)^{\frac{\omega}{\varrho}}, \quad u(x) = \frac{\omega}{\varrho \zeta} \eta(x)^{-(1+\frac{1}{\varrho})} \eta'(x), \quad (24)$$

which satisfy $u(x) = s'(x) - r(x)t'(x)$. The rest of derivation involves determining the appropriate function η and sequence $\{b_k\}_k$ to use for different types of step sizes and verifying the conditions stated in Theorem 1 for such η and $\{b_k\}_k$.



Notation. Throughout this paper, we use K to represent the total iterations in SGD (or epochs in RR). For all $k \in [K]$, we denote

$$y_k := \mathbb{E}[f(\mathbf{x}^k) - f^*].$$

We will use the following two sets of notations in the analysis of SGD (Algorithm 1) and RR (Algorithm 2), respectively. For SGD, we use

$$\varrho_1 = \frac{2\theta}{4\theta-1}, \quad \omega_1 = \frac{1}{4\theta-1}, \quad \zeta_1 = \theta \mu^{\frac{1}{2\theta}} \left(\frac{\mathsf{L}\sigma^2}{2} \right)^{1-\frac{1}{2\theta}}, \quad C_1 = \frac{\mathsf{A}^{4\theta-1} \mathsf{L}}{2\theta^{4\theta} \sigma^{8\theta-4} \mu^2}, \quad C_2 = \frac{2^{1-\frac{1}{2\theta}} \mathsf{L}}{\theta \mu^{\frac{1}{2\theta}} (\sigma^2)^{1-\frac{1}{2\theta}}}.$$

For RR, we use

$$\varrho_2 = \frac{\theta}{3\theta-1}, \quad \omega_2 = \frac{1}{3\theta-1}, \quad \zeta_2 = \frac{\theta}{2} \mu^{\frac{1}{2\theta}} \left(\frac{\mathsf{L}^2 \sigma^2}{N} \right)^{1-\frac{1}{2\theta}}, \quad C_3 = \frac{2\mathsf{A}^{3\theta-1} \mathsf{L}}{\theta^{3\theta} \sigma^{6\theta-3} \mu^{\frac{3}{2}}}, \quad C_4 = \frac{2^{4-\frac{1}{\theta}} \mathsf{L}}{\theta \mu^{\frac{1}{2\theta}} (\sigma^2)^{1-\frac{1}{2\theta}}}.$$

3.3 The Analysis of Exponential and Cosine Step Sizes

Exponential and Cosine Step Size. We will use the following formats for exponential step sizes:

$$\alpha_k = \alpha \gamma^k, \quad \text{where } \gamma = \left(\frac{\beta}{K}\right)^{\frac{p}{K}}, \beta, p > 0, K \geq 2^{\frac{p}{\beta}} \beta,$$

and for cosine step sizes:

$$\alpha_k = \alpha \left[\frac{1 + \cos\left(\frac{k\pi}{K}\right)}{2} \right]^\beta, \quad \text{where } \beta > 0.$$

Selection of b_k and η . For exponential step sizes, the naïve choice $b_k = k$, $\eta(x) = \alpha \gamma^x$ already ensures the convexity of r . However, for cosine step sizes, the choice $b_k = k$, $\eta(x) = \alpha \left(\frac{1 + \cos\left(\frac{x\pi}{K}\right)}{2}\right)^\beta$ does not work. As we will see later in the proof of Lemma 12, the alternative choice $b_k = \left[\frac{1 + \cos\left(\frac{k\pi}{K}\right)}{2}\right]^{\frac{\beta\omega}{\varrho}}$ and $\eta(x) = \alpha x^{\frac{\varrho}{\omega}}$ can resolve this problem. Here, we set $\varrho = \varrho_1 = \frac{2\theta}{4\theta-1}$, $\omega = \omega_1 = \frac{1}{4\theta-1}$ in SGD, and $\varrho = \varrho_2 = \frac{\theta}{3\theta-1}$, $\omega = \omega_2 = \frac{1}{3\theta-1}$ in RR.

An Important Splitting Technique. A direct application of Theorem 1 to exponential and cosine step sizes may lead to failure. To illustrate this, let us consider SGD with exponential step sizes. For simplicity, we set $\theta = \frac{1}{2}$, $\beta = 1$, and $p = 1$. Following the previous roadmap, we only need to analyze recursion (22). Using the same notations as in (24) with $b_k = k$ and $\eta(x) = \alpha K^{-\frac{x}{K}}$, we obtain

$$s(x) = \frac{2}{\ell_2 \alpha} K^{\frac{x}{K}}, \quad t(x) = \frac{1}{\ell_3 \alpha^2} K^{\frac{2x}{K}}, \quad r(x) = \frac{2\ell_3 \alpha}{\ell_2} K^{-\frac{x}{K}}, \quad u(x) = -\frac{1}{\zeta \alpha} K^{\frac{x}{K}} \frac{\log(K)}{K}.$$

To verify (8), we need to find a non-decreasing positive sequence $\{\lambda_k\}_{1 \leq k \leq K}$ such that for all $k \in [K-1]$, it holds that

$$(b_{k+1} - b_k)u(b_k) = -\frac{1}{\zeta \alpha} K^{\frac{k}{K}} \frac{\log(K)}{K} \geq -1 + \frac{1}{\lambda_{k+1}}. \quad (25)$$

By simply setting $k = K-1$, we can observe that this is infeasible. When $k = K-1$, equation (25) implies

$$(b_K - b_{K-1})u(b_{K-1}) \geq -1 + \frac{1}{\lambda_K} > -1. \quad (26)$$

However, by (25), we have

$$(b_K - b_{K-1})u(b_{K-1}) = -\frac{\log(K)}{\zeta \alpha} K^{-\frac{1}{K}} \leq -C \log(K), \quad \text{for some } C > 0. \quad (27)$$

Consequently, (27) contradicts (26) whenever $K \geq \exp(-1/C)$. Therefore, for large K , condition (25) cannot be satisfied for the entire sequence. Based on the diminishing property of the function u , this problem arises when k approaches the upper limit $K-1$.

Nonetheless, it is possible to fulfill (25) when k is not close to $K-1$. This observation inspires the subdivision of the index set $[K]$ into two distinct sets. During the first iterations,

(25) can be satisfied, and subsequently, a non-asymptotic estimation for the endpoint can be derived by applying Theorem 1. Subsequently, using Lemma 5, this estimation can be extended to the final iterate y_k . We proceed by demonstrating the execution of this idea in a simple example. Let us set $\lambda_k \equiv 2$. Then, condition (25) is satisfied if and only if

$$k \leq \log \left(\frac{\zeta \alpha K}{2 \log(K)} \right) \frac{K}{\log(K)},$$

which implies that (25) holds for all $0 \leq k \leq \bar{K}$, where $\bar{K} := \lfloor \log \left(\frac{\zeta \alpha K}{2 \log(K)} \right) \frac{K}{\log(K)} \rfloor$. Applying Theorem 1 (b) to the sequence $\{y_k\}_{0 \leq k \leq \bar{K}+1}$, we obtain:

$$y_{\bar{K}+1} \leq 2r(b_{\bar{K}+1}) + (y_0 - 2r(b_0)) \prod_{i=1}^{\bar{K}} \left(1 - \frac{1}{s(b_k)} \right).$$

By utilizing Lemma 5, we can extend this estimate to y_K :

$$y_K \leq 2r(b_{\bar{K}+1}) + (y_0 - 2r(b_0)) \prod_{i=1}^{K-1} \left(1 - \frac{1}{s(b_k)} \right),$$

where

$$r(b_{\bar{K}+1}) = \frac{2\ell_3 \alpha}{\ell_2} K^{-\frac{\bar{K}+1}{K}} \leq \frac{2\ell_3 \alpha}{\ell_2} \exp \left(-\log(K) \cdot \frac{\log \left(\frac{\zeta \alpha K}{2 \log(K)} \right)}{\log(K)} \right) = \frac{4\ell_3}{\ell_2 \zeta} \frac{\log(K)}{K}.$$

According to Corollary 6, the leading rate is $\mathcal{O}(r(b_{\bar{K}+1}))$, i.e., $\mathcal{O}\left(\frac{\log(K)}{K}\right)$.

Results on Exponential and Cosine Step Sizes. Following the general roadmap and with the help of the above splitting technique, we are able to show the convergence rate of both SGD and RR under general (θ, μ) -PL condition with exponential and cosine step sizes. We first introduce an important technical lemma.

Lemma 12 *Consider a non-negative sequence $\{y_k\}_{0 \leq k \leq K}$ satisfying*

$$y_{k+1} \leq (1 + \ell_1 \alpha_k^\tau) y_k - \ell_2 \alpha_k y_k^{2\theta} + \ell_3 \alpha_k^\tau \quad (28)$$

for all $k \in [K-1]$, where $\ell_1, \ell_2, \ell_3 > 0$, $K \in \mathbb{N}$ are given. Let us define

$$\varrho := \frac{2\theta}{(2\theta-1)\tau+1}, \quad \omega := \frac{\tau-1}{(2\theta-1)\tau+1}, \quad \zeta := \theta \ell_2^{\frac{1}{2\theta}} \ell_3^{\frac{2\theta-1}{2\theta}}.$$

Then, the following statements hold.

- (a) Let $\alpha_k = \alpha \gamma^k$ and $\gamma = \left(\frac{\beta}{K}\right)^{\frac{p}{K}}$ be given and assume $0 < \alpha \leq \min\left\{\left(\frac{\zeta}{\ell_1}\right)^{\frac{\varrho}{\omega}}, \zeta^{-\varrho}\right\}$, $p > 0$, $K \geq 2^{\frac{\varrho}{p}} \beta > 0$ and $\frac{\log(K/\beta)}{K} \leq \frac{\varrho \zeta \alpha^{1/\varrho}}{2p\omega}$. Then, it holds that

$$y_K \leq 4 \left(\frac{2p\omega}{\varrho} \right)^\omega \zeta^{-\omega} \left(\frac{\ell_3}{\ell_2} \right)^{\frac{1}{2\theta}} \left[\frac{\log(K/\beta)}{K} \right]^\omega + y_0 \exp \left(-\frac{\varrho \zeta \alpha^{1/\varrho}}{2p} \frac{K}{\log(K/\beta)} \right).$$

(b) Let $\alpha_k = \alpha \left[\frac{1 + \cos(\frac{k\pi}{K})}{2} \right]^\beta$ be given with $0 < \alpha \leq \min\{(\frac{\zeta}{\ell_1})^{\frac{\rho}{\omega}}, \zeta^{-\rho}\}$, then:

- if $\beta \geq \frac{\rho}{\omega}$, and $K \geq \frac{\pi^2 \beta \omega}{\alpha^{1/\rho} \rho \zeta}$, we have

$$y_K \leq 4 \left(\frac{\pi}{2} \right)^{\frac{2\beta\omega}{\rho}} \left(\frac{\ell_3}{\ell_2} \right)^{\frac{1}{2\theta}} \alpha^{\frac{\omega}{\rho}} \left[\frac{\pi^2 \beta \omega}{\alpha^{1/\rho} \rho \zeta} \frac{1}{K} \right]^{\frac{2\beta\omega}{2\beta+\rho}} + y_0 \exp \left(- \frac{\zeta \alpha^{1/\rho}}{2^{\max\{1, \beta/\rho\}}} K \right).$$

- if $0 < \beta < \frac{\rho}{\omega}$, and $K \geq \frac{4\pi^2 \beta \omega}{\alpha^{1/\rho} \rho \zeta}$, we have

$$y_K \leq 4 \left(\frac{\pi}{2} \right)^{\frac{2\beta\omega}{\rho}} \left(\frac{\ell_3}{\ell_2} \right)^{\frac{1}{2\theta}} \alpha^{\frac{\omega}{\rho}} \cdot \max \left\{ K^{-\frac{2\beta\omega}{\rho}}, \left(\frac{4\pi^2 \beta \omega}{\alpha^{1/\rho} \rho \zeta} \frac{1}{K} \right)^{\frac{2\beta\omega}{2\beta+\rho}} \right\} + y_0 \exp \left(- \frac{\zeta \alpha^{1/\rho} K}{2^{\max\{1, \beta/\rho\}}} \right).$$

Proof We prove the different parts of Lemma 12 in a step-by-step fashion and following the previous derivations. Recall that the roadmap outlined in Section 3.2 is applicable provided that $\alpha_k \leq \min\{(\frac{\zeta}{\ell_1})^{\frac{\rho}{\omega}}, \zeta^{-\rho}\}$, see (21) and (23).

Proof of Part (a). We consider exponential step sizes of the form $\alpha_k = \alpha \gamma^k$ where $\gamma = (\frac{\beta}{K})^{\frac{\rho}{K}}$, $\alpha \leq \min\{(\frac{\zeta}{\ell_1})^{\frac{\rho}{\omega}}, \zeta^{-\rho}\}$ and we have $p > 0$, $K \geq 2^{\frac{\rho}{p}} \beta > 0$ and $\frac{\log(K/\beta)}{K} \leq \frac{\rho \zeta \alpha^{1/\rho}}{2p\omega}$. Let us set $b_k = k$ and $\eta(x) = \alpha \gamma^x$. The explicit formula of s, t, r, u can be obtained by substituting $\eta(x) = \alpha \gamma^x$ into (24), i.e.,

$$s(x) = \frac{1}{\zeta \alpha^{1/\rho}} \gamma^{-\frac{x}{\rho}}, \quad t(x) = \frac{1}{2\theta \ell_3 \alpha^\tau} \gamma^{-\tau x}, \quad r(x) = 2 \left(\frac{\ell_3}{\ell_2} \right)^{\frac{1}{2\theta}} \alpha^{\frac{\omega}{\rho}} \gamma^{\frac{\omega x}{\rho}}, \quad u(x) = \frac{\omega \log(\gamma)}{\rho \zeta \alpha^{1/\rho}} \gamma^{-\frac{x}{\rho}}.$$

Since $0 < \gamma < 1$, we have r is convex over the real line and $r(b_k)$ is decreasing w.r.t. k , which meets the requirements for r in Theorem 1 and Lemma 5. Meanwhile, u is also decreasing on \mathbb{R} . Then for all

$$k \leq \rho \log \left(- \frac{2\omega \log(\gamma)}{\rho \zeta \alpha^{1/\rho}} \right) / \log(\gamma) = \frac{\rho}{p} \log \left(\frac{\rho \zeta \alpha^{1/\rho} K}{2p\omega \log(K/\beta)} \right) \frac{K}{\log(K/\beta)},$$

we have

$$(b_{k+1} - b_k)u(b_k) \geq -\frac{1}{2}.$$

We can now apply Theorem 1 (b) and Lemma 5 to (22) in tandem by setting

$$k_0 = 0, \quad \bar{K} = \min \left\{ K - 1, \left\lfloor \frac{\rho}{p} \log \left(\frac{\rho \zeta \alpha^{1/\rho} K}{2p\omega \log(K/\beta)} \right) \frac{K}{\log(K/\beta)} \right\rfloor \right\} \quad \text{and} \quad \lambda_k \equiv 2.$$

This yields

$$y_K \leq 2r(\bar{K} + 1) + y_0 \prod_{k=0}^{\bar{K}-1} \left(1 - \frac{1}{s(b_k)} \right) \leq 4 \left(\frac{\ell_3}{\ell_2} \right)^{\frac{1}{2\theta}} \alpha^{\frac{\omega}{\rho}} \left(\frac{\beta}{K} \right)^{\frac{\bar{K}+1}{K} \frac{p\omega}{\rho}} + y_0 \prod_{k=0}^{\bar{K}-1} \left(1 - \zeta \alpha^{\frac{1}{\rho}} \gamma^{\frac{k}{\rho}} \right).$$

In addition, we can establish the following relation

$$\left(\frac{\beta}{K} \right)^{\frac{\bar{K}+1}{K}} = \exp \left(- \frac{\bar{K}+1}{K} \log(K/\beta) \right) \leq \exp \left(- \frac{\rho}{p} \log \left(\frac{\rho \zeta \alpha^{1/\rho} K}{2p\omega \log(K/\beta)} \right) \right) = \left[\frac{2p\omega \log(K/\beta)}{\rho \zeta \alpha^{1/\rho} K} \right]^{\frac{\rho}{p}}.$$

Noticing $\gamma^{\frac{K}{e}} = \left(\frac{\beta}{K}\right)^{\frac{p}{e}} \leq \frac{1}{2}$ and using the geometric series $\sum_{k=0}^{K-1} x^k = \frac{1-x^K}{1-x}$, $x \neq 1$, we have

$$\begin{aligned} \prod_{k=0}^{K-1} (1 - \zeta \alpha^{\frac{1}{e}} \gamma^{\frac{k}{e}}) &\leq \exp\left(-\zeta \alpha^{\frac{1}{e}} \sum_{k=0}^{K-1} \gamma^{\frac{k}{e}}\right) \\ &= \exp\left(-\zeta \alpha^{\frac{1}{e}} \frac{1 - \gamma^{K/e}}{1 - \gamma^{1/e}}\right) \leq \exp\left(-\frac{\zeta \alpha^{1/e}}{2} \frac{1}{1 - \gamma^{1/e}}\right). \end{aligned}$$

Applying the inequality $\exp(x) \geq 1 + x$, we further obtain

$$(1 - \gamma^{\frac{1}{e}})^{-1} = (1 - \exp(\varrho^{-1} \log(\gamma)))^{-1} \geq (-\varrho^{-1} \log(\gamma))^{-1} = \frac{\varrho}{p} \frac{K}{\log(K/\beta)}.$$

Hence, we can conclude

$$\begin{aligned} y_K &\leq 4 \left(\frac{\ell_3}{\ell_2}\right)^{\frac{1}{2\theta}} \alpha^{\frac{\omega}{e}} \left[\frac{2p\omega \log(K/\beta)}{\varrho \zeta \alpha^{1/e} K}\right]^{\omega} + y_0 \prod_{k=0}^{K-1} (1 - \zeta \alpha^{\frac{1}{e}} \gamma^{\frac{k}{e}}) \\ &\leq 4 \left(\frac{2p\omega}{\varrho}\right)^{\omega} \zeta^{-\omega} \left(\frac{\ell_3}{\ell_2}\right)^{\frac{1}{2\theta}} \left[\frac{\log(K/\beta)}{K}\right]^{\omega} + y_0 \exp\left(-\frac{\varrho \zeta \alpha^{1/e}}{2p} \frac{K}{\log(K/\beta)}\right). \end{aligned}$$

Proof of Part (b). Now we consider cosine step sizes $\alpha_k = \alpha \left[\frac{1 + \cos(\frac{k\pi}{K})}{2}\right]^{\beta}$ where $\alpha \leq \min\left\{\left(\frac{\zeta}{\ell_1}\right)^{\frac{\varrho}{\omega}}, \zeta^{-\varrho}\right\}$, $\beta > 0$. Let us define $b_k := \left[\frac{1 + \cos(\frac{k\pi}{K})}{2}\right]^{\frac{\beta\omega}{e}}$ and $\eta(x) := \alpha x^{\frac{\varrho}{\omega}}$, then

$$s(x) = \frac{1}{\zeta \alpha^{1/e}} x^{-\frac{1}{\omega}}, \quad t(x) = \frac{1}{2\theta \ell_3 \alpha^{\tau}} x^{-\frac{\varrho\tau}{\omega}}, \quad r(x) = 2 \left(\frac{\ell_3}{\ell_2}\right)^{\frac{1}{2\theta}} \alpha^{\frac{\omega}{e}} x, \quad u(x) = \frac{1}{\alpha^{1/e} \zeta} x^{-(1+\frac{1}{\omega})}.$$

Firstly, r is linear, thus convex, over the real line, satisfying the requirement for r in Theorem 1. Moreover, as r is increasing and b_k is decreasing, $r(b_k)$ decreases with respect to k , making Lemma 5 applicable. To prepare, we present an explicit formula for $u(b_k)$:

$$u(b_k) = \frac{1}{\alpha^{1/e} \zeta} \left[\frac{1 + \cos(\frac{k\pi}{K})}{2}\right]^{-\left(\frac{\beta\omega}{e} + \frac{\beta}{e}\right)}.$$

Case I: $\beta \geq \frac{\varrho}{\omega}$. Since the function $x \mapsto x^{\frac{\beta\omega}{e}}$ is convex on $[0, +\infty)$, we have

$$b_{k+1} - b_k \geq \frac{\beta\omega}{\varrho} \left[\frac{1 + \cos(\frac{k\pi}{K})}{2}\right]^{\frac{\beta\omega}{e}-1} \left[\frac{\cos(\frac{(k+1)\pi}{K}) - \cos(\frac{k\pi}{K})}{2}\right].$$

Thus, it follows

$$(b_{k+1} - b_k)u(b_k) \geq \frac{\beta\omega}{\alpha^{1/e} \varrho \zeta} \left[\frac{1 + \cos(\frac{k\pi}{K})}{2}\right]^{-\left(\frac{\beta}{e} + 1\right)} \left[\frac{\cos(\frac{(k+1)\pi}{K}) - \cos(\frac{k\pi}{K})}{2}\right].$$

For $k \in [K]$, by double angle formula $1 - \cos(2a) = 2 \sin^2(a)$, it holds that $1 + \cos(\frac{k\pi}{K}) = 1 - \cos(\frac{(K-k)\pi}{K}) = 2 \sin^2(\frac{(K-k)\pi}{2K})$. The function $x \mapsto \frac{\sin(x)}{x}$ is monotonically decreasing on $[0, \frac{\pi}{2}]$ and hence, we have $\sin(x) \geq \frac{2}{\pi}x$ for all $x \in [0, \frac{\pi}{2}]$. Hence, we can infer $1 + \cos(\frac{k\pi}{K}) \geq 2(1 - \frac{k}{K})^2$ and

$$\left[\frac{1 + \cos(\frac{k\pi}{K})}{2} \right]^{-(1+\frac{\beta}{e})} \leq \left(1 - \frac{k}{K} \right)^{-2(1+\frac{\beta}{e})}.$$

In addition, utilizing the sum-to-product identity $\cos(a) - \cos(b) = -2 \sin(\frac{1}{2}(a+b)) \sin(\frac{1}{2}(a-b))$, $\sin(x) = \sin(\pi - x)$, and $\sin(x) \leq x$, it holds that

$$\cos\left(\frac{(k+1)\pi}{K}\right) - \cos\left(\frac{k\pi}{K}\right) = -2 \sin\left(\frac{(2k+1)\pi}{2K}\right) \sin\left(\frac{\pi}{2K}\right) \geq -\frac{\pi^2}{K} \left(1 - \frac{k}{K}\right). \quad (29)$$

Then, we obtain

$$(b_{k+1} - b_k)u(b_k) \geq -\frac{\pi^2 \beta \omega}{2\alpha^{1/e} \rho \zeta} \frac{1}{K} \left(1 - \frac{k}{K}\right)^{-(1+2\frac{\beta}{e})}.$$

Thus, for all $0 \leq k \leq K(1 - [\frac{\pi^2 \beta \omega}{\alpha^{1/e} \rho \zeta} \frac{1}{K}]^{\frac{1}{1+2\beta/e}})$, we have $(b_{k+1} - b_k)u(b_k) \geq -\frac{1}{2}$. Moreover, the rate function r is convex on the whole real line and $r(b_k)$ is non-increasing w.r.t. k for all $k \in [K]$. Invoking Theorem 1 (b) and Lemma 5 in tandem for (22) with

$$k_0 = 0, \quad \bar{K} = \left\lfloor K - K \left(\frac{\pi^2 \beta \omega}{\alpha^{1/e} \rho \zeta} \frac{1}{K} \right)^{\frac{1}{2\beta/e+1}} \right\rfloor \quad \text{and} \quad \lambda_k \equiv 2,$$

it follows

$$y_K \leq 4 \left(\frac{\ell_3}{\ell_2} \right)^{\frac{1}{2\theta}} \alpha^{\frac{\omega}{e}} \left[\frac{1 + \cos(\frac{(\bar{K}+1)\pi}{K})}{2} \right]^{\frac{\beta\omega}{e}} + y_0 \prod_{k=0}^{K-1} \left(1 - \zeta \alpha^{\frac{1}{e}} \left[\frac{1 + \cos(\frac{k\pi}{K})}{2} \right]^{\frac{\beta}{e}} \right).$$

Here, the condition $\bar{K} \geq 0$ is guaranteed by $K \geq \frac{\pi^2 \beta \omega}{\alpha^{1/e} \rho \zeta}$. Using $\bar{K}+1 \geq K(1 - [\frac{\pi^2 \beta \omega}{\alpha^{1/e} \rho \zeta} \frac{1}{K}]^{\frac{1}{2\beta/e+1}})$ and $1 + \cos(\frac{k\pi}{K}) \leq \frac{\pi^2}{2}(1 - \frac{k}{K})^2$, we further have

$$1 + \cos\left(\frac{(\bar{K}+1)\pi}{K}\right) \leq \frac{\pi^2}{2} \left(1 - \frac{\bar{K}+1}{K}\right)^2 \leq \frac{\pi^2}{2} \left[\frac{\pi^2 \beta \omega}{\alpha^{1/e} \rho \zeta} \frac{1}{K} \right]^{\frac{2}{2\beta/e+1}}.$$

To handle the product term, we notice

$$\prod_{k=0}^{K-1} \left(1 - \zeta \alpha^{\frac{1}{e}} \left[\frac{1 + \cos(\frac{k\pi}{K})}{2} \right]^{\frac{\beta}{e}} \right) \leq \exp \left(-\zeta \alpha^{\frac{1}{e}} \sum_{k=0}^{K-1} \left[\frac{1 + \cos(\frac{k\pi}{K})}{2} \right]^{\frac{\beta}{e}} \right).$$

Using Lagrange's trigonometric identity $\sum_{k=0}^n \cos(ka) = \frac{1}{2} + \frac{\sin(\frac{(2n+1)a}{2})}{2 \sin(\frac{a}{2})}$, we have $\sum_{k=0}^{K-1} [1 + \cos(\frac{k\pi}{K})] = K + \frac{1}{2} + \frac{\sin((K-\frac{1}{2})\frac{\pi}{K})}{2 \sin(\frac{\pi}{2K})} = K + 1$. Next, we further consider the following cases:

- If $\beta > \varrho$, applying Jensen's inequality, it follows

$$\sum_{k=0}^{K-1} \left[\frac{1 + \cos(\frac{k\pi}{K})}{2} \right]^{\frac{\beta}{\varrho}} \geq K \left[\frac{1}{K} \sum_{k=0}^{K-1} \frac{1 + \cos(\frac{k\pi}{K})}{2} \right]^{\frac{\beta}{\varrho}} \geq \frac{K}{2^{\beta/\varrho}}.$$

- If $\beta \leq \varrho$, we then have $\sum_{k=0}^{K-1} \left[\frac{1 + \cos(\frac{k\pi}{K})}{2} \right]^{\frac{\beta}{\varrho}} \geq \sum_{k=0}^{K-1} \left[\frac{1 + \cos(\frac{k\pi}{K})}{2} \right] \geq \frac{K}{2}$.

This shows $\sum_{k=0}^{K-1} \left[\frac{1 + \cos(\frac{k\pi}{K})}{2} \right]^{\frac{\beta}{\varrho}} \geq \frac{K}{2^{\max\{1, \beta/\varrho\}}}$ and we can conclude

$$y_K \leq 4 \left(\frac{\pi}{2} \right)^{\frac{2\beta\omega}{\varrho}} \left(\frac{\ell_3}{\ell_2} \right)^{\frac{1}{2\theta}} \alpha^{\frac{\omega}{\varrho}} \left[\frac{\pi^2 \beta \omega}{\alpha^{1/\varrho} \varrho \zeta} \frac{1}{K} \right]^{\frac{2\beta\omega}{2\beta+\varrho}} + y_0 \exp \left(-\frac{\zeta \alpha^{1/\varrho}}{2^{\max\{1, \beta/\varrho\}}} K \right).$$

Case II: $0 < \beta < \frac{\varrho}{\omega}$. In this case, the function $x \mapsto x^{\frac{\beta\omega}{\varrho}}$ is concave on $[0, +\infty)$, and we obtain

$$b_{k+1} - b_k \geq \frac{\beta\omega}{\varrho} \left[\frac{1 + \cos(\frac{(k+1)\pi}{K})}{2} \right]^{\frac{\beta\omega}{\varrho}-1} \left[\frac{\cos(\frac{(k+1)\pi}{K}) - \cos(\frac{k\pi}{K})}{2} \right].$$

Thus, it follows

$$\begin{aligned} & (b_{k+1} - b_k)u(b_k) \\ & \geq \frac{\beta\omega}{\alpha^{1/\varrho} \varrho \zeta} \left[\frac{1 + \cos(\frac{k\pi}{K})}{2} \right]^{-\left(\frac{\beta\omega}{\varrho} + \frac{\beta}{\varrho}\right)} \left[\frac{1 + \cos(\frac{(k+1)\pi}{K})}{2} \right]^{\frac{\beta\omega}{\varrho}-1} \left[\frac{\cos(\frac{(k+1)\pi}{K}) - \cos(\frac{k\pi}{K})}{2} \right] \\ & = \frac{\beta\omega}{\alpha^{1/\varrho} \varrho \zeta} \left[\frac{1 + \cos(\frac{(k+1)\pi}{K})}{1 + \cos(\frac{k\pi}{K})} \right]^{\frac{\beta\omega}{\varrho}} \left[\frac{1 + \cos(\frac{k\pi}{K})}{2} \right]^{-\frac{\beta}{\varrho}} \left[\frac{1 + \cos(\frac{(k+1)\pi}{K})}{2} \right]^{-1} \left[\frac{\cos(\frac{(k+1)\pi}{K}) - \cos(\frac{k\pi}{K})}{2} \right] \\ & \geq \frac{\beta\omega}{\alpha^{1/\varrho} \varrho \zeta} \left[\frac{1 + \cos(\frac{k\pi}{K})}{2} \right]^{-\frac{\beta}{\varrho}} \left[\frac{1 + \cos(\frac{(k+1)\pi}{K})}{2} \right]^{-1} \left[\frac{\cos(\frac{(k+1)\pi}{K}) - \cos(\frac{k\pi}{K})}{2} \right]. \end{aligned}$$

Utilizing $1 + \cos(\frac{k\pi}{K}) \geq 2(1 - \frac{k}{K})^2$ as shown in **Case I**, we have for all $k \in [K-1]$,

$$\left[\frac{1 + \cos(\frac{k\pi}{K})}{2} \right]^{-\frac{\beta}{\varrho}} \leq \left(1 - \frac{k}{K} \right)^{-\frac{2\beta}{\varrho}},$$

and for all $k \in [K-2]$,

$$\left[\frac{1 + \cos(\frac{(k+1)\pi}{K})}{2} \right]^{-1} \leq \left(1 - \frac{k+1}{K} \right)^{-2} \leq 4 \left(1 - \frac{k}{K} \right)^{-2}.$$

For the last inequality, we use $1 - \frac{k+1}{K} \geq \frac{1}{2} \left(1 - \frac{k}{K} \right)$. Combining the previous estimates and (29), we can infer:

$$(b_{k+1} - b_k)u(b_k) \geq -\frac{2\pi^2 \beta \omega}{\alpha^{1/\varrho} \varrho \zeta} \frac{1}{K} \left(1 - \frac{k}{K} \right)^{-(1+2\frac{\beta}{\varrho})}.$$

Then, for all $0 \leq k \leq \min\{K-2, K(1 - [\frac{4\pi^2\beta\omega}{\alpha^{1/e}\rho\zeta} \frac{1}{K}]^{\frac{1}{1+2\beta/e}})\}$, we have $(b_{k+1} - b_k)u(b_k) \geq -\frac{1}{2}$. Invoking Theorem 1 (b) and Lemma 5 in tandem for (22) with

$$k_0 = 0, \quad \bar{K} = \min\{K-2, \left\lceil K(1 - [\frac{4\pi^2\beta\omega}{\alpha^{1/e}\rho\zeta} \frac{1}{K}]^{\frac{1}{2\beta/e+1}}) \right\rceil\}, \quad \text{and } \lambda_k \equiv 2,$$

it follows

$$y_K \leq 4 \left(\frac{\ell_3}{\ell_2}\right)^{\frac{1}{2\theta}} \alpha^{\frac{\omega}{e}} \left[\frac{1 + \cos\left(\frac{(\bar{K}+1)\pi}{K}\right)}{2} \right]^{\frac{\beta\omega}{e}} + y_0 \prod_{k=0}^{K-1} \left(1 - \zeta \alpha^{\frac{1}{e}} \left[\frac{1 + \cos\left(\frac{k\pi}{K}\right)}{2} \right]^{\frac{\beta}{e}} \right).$$

Here, the condition $\bar{K} \geq 0$ is guaranteed by $K \geq \frac{4\pi^2\beta\omega}{\alpha^{1/e}\rho\zeta}$. Using $\bar{K} + 1 \geq \min\{K-1, K(1 - [\frac{4\pi^2\beta\omega}{\alpha^{1/e}\rho\zeta} \frac{1}{K}]^{\frac{1}{2\beta/e+1}})\}$ and $1 + \cos\left(\frac{k\pi}{K}\right) \leq \frac{\pi^2}{2} \left(1 - \frac{k}{K}\right)^2$, we have

$$1 + \cos\left(\frac{(\bar{K}+1)\pi}{K}\right) \leq \frac{\pi^2}{2} \left(1 - \frac{\bar{K}+1}{K}\right)^2 \leq \frac{\pi^2}{2} \max\left\{ \frac{1}{K^2}, \left[\frac{4\pi^2\beta\omega}{\alpha^{1/e}\rho\zeta} \frac{1}{K} \right]^{\frac{2}{2\beta/e+1}} \right\}.$$

Using the same strategy in **Case I** to handle the product term, we can conclude

$$y_K \leq 4 \left(\frac{\pi}{2}\right)^{\frac{2\beta\omega}{e}} \left(\frac{\ell_3}{\ell_2}\right)^{\frac{1}{2\theta}} \alpha^{\frac{\omega}{e}} \max\left\{ K^{-\frac{2\beta\omega}{e}}, \left[\frac{4\pi^2\beta\omega}{\alpha^{1/e}\rho\zeta} \frac{1}{K} \right]^{\frac{2\beta\omega}{2\beta/e+1}} \right\} + y_0 \exp\left(-\frac{\zeta \alpha^{1/e}}{2^{\max\{1, \beta/e\}}} K\right).$$

This finishes the proof of Lemma 12. \blacksquare

The following results on exponential and cosine step sizes can be obtained directly from Lemma 12.

Theorem 13 (Convergence Rate of SGD with Exponential Step Sizes) *Let (A.1)–(A.2) and (P) hold, and let the sequence $\{\mathbf{x}^k\}_{0 \leq k \leq K}$, $K \geq 1$, be generated by SGD with $\alpha_k = \alpha\gamma^k$ and $\gamma = \left(\frac{\beta}{K}\right)^{\frac{p}{K}}$. If $0 < \alpha \leq \min\left\{\left(\frac{2\zeta_1}{\text{AL}}\right)^{2\theta}, \zeta_1^{-e_1}, \text{L}^{-1}\right\}$ and K is sufficiently large such that $K \geq 2^{\frac{e_1}{p}} \beta > 0$ and $\frac{\log(K/\beta)}{K} \leq \frac{\varrho_1 \zeta_1 \alpha^{1/e_1}}{2p\omega_1}$, then we have*

$$y_K \leq 4 \left(\frac{p}{2\theta^2}\right)^{\omega_1} \left(\frac{\text{L}\sigma^2}{\mu^2}\right)^{\omega_1} \left[\frac{\log(K/\beta)}{K}\right]^{\omega_1} + y_0 \exp\left(-\frac{\varrho_1 \zeta_1 \alpha^{1/e_1}}{2p} \frac{K}{\log(K/\beta)}\right). \quad (30)$$

The full proof can be found in Appendix C.1. In the case $\alpha = \min\left\{\left(\frac{2\zeta_1}{\text{AL}}\right)^{2\theta}, \zeta_1^{-e_1}, \text{L}^{-1}\right\}$, the term $\zeta_1 \alpha^{1/e_1}$ in (30) further simplifies to $\zeta_1 \alpha^{1/e_1} = \min\{C_1^{-1}, C_2^{-1}, 1\}$. When $\theta = \frac{1}{2}$, (Li et al., 2021, Theorem 1) shows a similar rate to (30), with the difference being that they have a worse $\log^2(K)$ dependence in the dominant term. The work (Wang et al., 2021, Theorem 4.1) derives the same type of results under strong convexity and a more stringent bounded variance assumption. It is worth noting that our result also covers the range $\theta \in (\frac{1}{2}, 1]$, which is not present in these previous works.

Theorem 14 (Convergence Rate of RR with Exponential Step Sizes) *Let (B.1)–(B.3) and (P) hold, and let $\{\mathbf{x}^k\}_{0 \leq k \leq K}$, $K \geq 1$, be generated by RR with $\alpha_k = \alpha\gamma^k$ and $\gamma = \left(\frac{\beta}{K}\right)^{\frac{p}{K}}$.*

If $\alpha \leq \min\{(\frac{2N\zeta_2}{\mathbf{AL}^2})^\theta, \zeta_2^{-\varrho_2}, \frac{1}{2\mathbf{L}}\}$ and K is large enough such that $K \geq 2^{\frac{\varrho_2}{p}}\beta > 0$ and $\frac{\log(K/\beta)}{K} \leq \frac{\varrho_2\zeta_2\alpha^{1/\varrho_2}}{2p\omega_2}$, then it holds that

$$y_K \leq 4 \left(\frac{4p}{\theta^2}\right)^{\omega_2} \left(\frac{\mathbf{L}\sigma}{\mu^{3/2}}\right)^{\omega_2} \left[\frac{\log(K/\beta)}{\sqrt{NK}}\right]^{\omega_2} + y_0 \exp\left(-\frac{\varrho_2\zeta_2\alpha^{1/\varrho_2}}{2p} \frac{K}{\log(K/\beta)}\right). \quad (31)$$

Moreover, if $\alpha = \min\{(\frac{2N\zeta_2}{\mathbf{AL}^2})^\theta, \zeta_2^{-\varrho_2}, \frac{1}{2\mathbf{L}}\}$, $\frac{K}{\log(K/\beta)} \geq \frac{4p}{\theta} \max\{\frac{C_3}{\sqrt{N}}, C_4N^{1-\frac{1}{2\theta}}, 1\} \log(NK)$, and $K \geq 2^{\frac{\varrho_2}{p}}\beta > 0$, then

$$y_K \leq 4 \left(\frac{4p}{\theta^2}\right)^{\omega_2} \left(\frac{\mathbf{L}\sigma}{\mu^{3/2}}\right)^{\omega_2} \left[\frac{\log(K/\beta)}{\sqrt{NK}}\right]^{\omega_2} + y_0(NK)^{-\omega_2}. \quad (32)$$

The proof of Theorem 14 is provided in Appendix C.1. Theorem 14 extends the theoretical results for RR by including exponential step sizes case under the general (θ, μ) -PL condition.

Theorem 15 (Convergence Rate for SGD with cosine step sizes) Suppose (A.1)–(A.2) and (P) hold. Let the sequence $\{\mathbf{x}^k\}_{0 \leq k \leq K}$ where $K \geq 2$ be generated by Algorithm 1 with $\alpha_k = \alpha \left[\frac{1+\cos(\frac{k\pi}{K})}{2}\right]^\beta$. Let $\alpha \leq \min\{(\frac{2\zeta_1}{\mathbf{AL}})^{2\theta}, \zeta_1^{-\varrho_1}, \mathbf{L}^{-1}\}$, then

- if $\beta \geq 2\theta$ and $K \geq \frac{\pi^2\beta}{2\theta\alpha^{1/\varrho_1}\zeta_1}$, we have

$$y_K \leq 4 \left(\frac{\pi}{2}\right)^{\frac{2\beta\omega_1}{\varrho_1}} \left(\frac{\mathbf{L}\sigma}{2\mu}\right)^{\frac{1}{2\theta}} \alpha^{\frac{\omega_1}{\varrho_1}} \left[\frac{\pi^2\beta}{2\theta\alpha^{1/\varrho_1}\zeta_1} \frac{1}{K}\right]^{\frac{2\beta\omega_1}{2\beta+\varrho_1}} + y_0 \exp\left(-\frac{\zeta_1\alpha^{1/\varrho_1}}{2^{\max\{1, \beta/\varrho_1\}}} K\right). \quad (33)$$

- if $\beta < 2\theta$ and $K \geq \frac{2\pi^2\beta}{\theta\alpha^{1/\varrho_1}\zeta_1}$, we have

$$y_K \leq 4 \left(\frac{\pi}{2}\right)^{\frac{2\beta\omega_1}{\varrho_1}} \left(\frac{\mathbf{L}\sigma}{2\mu}\right)^{\frac{1}{2\theta}} \alpha^{\frac{\omega_1}{\varrho_1}} \max\left\{K^{-\frac{2\beta\omega_1}{\varrho_1}}, \left[\frac{\pi^2\beta}{2\theta\alpha^{1/\varrho_1}\zeta_1} \frac{1}{K}\right]^{\frac{2\beta\omega_1}{2\beta+\varrho_1}}\right\} + y_0 \exp\left(-\frac{\zeta_1\alpha^{1/\varrho_1}}{2^{\max\{1, \beta/\varrho_1\}}} K\right). \quad (34)$$

Moreover, if $\alpha = 2^{\max\{\beta, \varrho_1\}} \left(\frac{2\omega_1 \log(K)}{\zeta_1 K}\right)^{\varrho_1}$, $K \geq \Theta(\max\{C_1, C_2, 1\}) \log(K)$, then

$$y_K = \mathcal{O}\left(\left[\frac{(\log(K))^{\frac{\varrho_1}{2\beta+\varrho_1}}}{K}\right]^{\omega_1}\right). \quad (35)$$

The proof is deferred to Appendix C.2. To the best of our knowledge, the only existing theoretical analysis on the convergence of cosine step sizes under PL-type condition is provided in Li et al. (2021). In (Li et al., 2021, Theorem 2), the authors obtained the complexity bound $\mathcal{O}(K^{-\frac{2}{3}})$ for SGD with $\beta = 1$ under the $(\frac{1}{2}, \mu)$ -PL condition. Our Theorem 15 can recover their result by setting $\beta = 1$ and $\theta = \frac{1}{2}$ in (33). Furthermore, we can improve this rate to $\mathcal{O}(\log^{1/3}(K)/K)$ by carefully selecting the step size parameter α ; see (35). It is worth mentioning that we also derive the result for $\theta \in (\frac{1}{2}, 1]$, which are new.

Theorem 16 (Convergence Rate for RR with Cosine Step Sizes) Suppose (B.1)–(B.3) and (P) hold. Let the sequence $\{\mathbf{x}^k\}_{0 \leq k \leq K}$, $K \geq 2$, be generated by SGD with $\alpha_k = \alpha \left[\frac{1+\cos(\frac{k\pi}{K})}{2}\right]^\beta$. Let $\alpha \leq \min\{(\frac{2N\zeta_2}{\mathbf{AL}^2})^\theta, \zeta_2^{-\varrho_2}, (2\mathbf{L})^{-1}\}$, then

- if $\beta \geq \theta$ and $K \geq \frac{\pi^2 \beta}{\theta \alpha^{1/\varrho_2} \zeta_2}$, we have

$$y_K \leq 4 \left(\frac{\pi}{2} \right)^{\frac{2\beta\omega_2}{\varrho_2}} \left(\frac{\mathbb{L}^2 \sigma^2}{\mu N} \right)^{\frac{1}{2\theta}} \alpha^{\frac{\omega_2}{\varrho_2}} \left[\frac{\pi^2 \beta}{\theta \alpha^{1/\varrho_2} \zeta_2} \frac{1}{K} \right]^{\frac{2\beta\omega_2}{2\beta+\varrho_2}} + y_0 \exp \left(- \frac{\zeta_2 \alpha^{1/\varrho_2}}{2^{\max\{1, \beta/\varrho_2\}}} K \right). \quad (36)$$

- if $\beta < \theta$ and $K \geq \frac{4\pi^2 \beta}{\theta \alpha^{1/\varrho_2} \zeta_2}$, we have

$$y_K \leq 4 \left(\frac{\pi}{2} \right)^{\frac{2\beta\omega_2}{\varrho_2}} \left(\frac{\mathbb{L}^2 \sigma^2}{\mu N} \right)^{\frac{1}{2\theta}} \alpha^{\frac{\omega_2}{\varrho_2}} \max \left\{ K^{-\frac{2\beta\omega_2}{\varrho_2}}, \left[\frac{\pi^2 \beta}{\theta \alpha^{1/\varrho_2} \zeta_2} \frac{1}{K} \right]^{\frac{2\beta\omega_2}{2\beta+\varrho_2}} \right\} + y_0 \exp \left(- \frac{\zeta_2 \alpha^{1/\varrho_2}}{2^{\max\{1, \beta/\varrho_2\}}} K \right). \quad (37)$$

Moreover, if $\alpha = 2^{\max\{\varrho_2, \beta\}} \left(\frac{2\omega_2 \log(NK)}{\zeta_2 K} \right)^{\varrho_2}$ and $K \geq \Theta(\max\{\frac{C_3}{\sqrt{N}}, C_4 N^{1-\frac{1}{2\theta}}, 1\}) \log(NK)$, then

$$y_K = \mathcal{O} \left(\left[\frac{(\log(NK))^{\frac{1}{2\beta/\varrho_2+1}}}{\sqrt{NK}} \right]^{\omega_2} \right). \quad (38)$$

The proof is deferred to Appendix C.2. The results in Theorem 16 are new for RR.

3.4 The Analysis of Constant and Polynomial Step Sizes

For comprehensiveness, we conduct the aforementioned analysis for polynomial and constant step size rules. The findings are presented in Theorems 17, 18, 20 and 21, accompanied by an in-depth comparison to prior research.

3.4.1 CONSTANT STEP SIZES

Based on the roadmap in Section 3.2, we can obtain the following convergence results for SGD and RR utilizing constant step sizes.

Theorem 17 (Convergence Rate for SGD with Constant Step Size) *Let (A.1)–(A.2) and (P) hold. Let the sequence $\{\mathbf{x}^k\}_{0 \leq k \leq K}$, $K \geq 2$, be generated by SGD with $\alpha_k = \left[\frac{\alpha \log(K)}{K} \right]^{\varrho_1}$. If $\alpha > 0$ and K satisfies $K \geq 1$, $\frac{\alpha \log(K)}{K} \leq \min\left\{ \left(\frac{2\zeta_1}{\mathbb{A}\mathbb{L}} \right)^{\frac{1}{\omega_1}}, \zeta_1^{-1}, \mathbb{L}^{-\frac{1}{\varrho_1}} \right\}$, then*

$$y_K \leq 2 \left(\frac{\mathbb{L}\sigma^2}{2\mu} \right)^{\frac{1}{2\theta}} \left[\frac{\alpha \log(K)}{K} \right]^{\omega_1} + y_0 K^{-\zeta_1 \alpha}. \quad (39)$$

Moreover, if $\alpha = \frac{2\omega_1}{\zeta_1}$ and $K \geq (2\omega_1 \max\{C_1, C_2, 1\}) \log(K)$, then

$$y_K \leq 2 \left(\frac{\omega_1}{\theta} \right)^{\omega_1} \left(\frac{\mathbb{L}\sigma^2}{\mu^2} \right)^{\omega_1} \left[\frac{\log(K)}{K} \right]^{\omega_1} + y_0 K^{-2\omega_1}. \quad (40)$$

The proof is provided in Appendix D.2. SGD under constant step sizes has been extensively studied, but mostly limited to the case where $\theta = \frac{1}{2}$ (or under stronger assumptions); see, e.g., Gower et al. (2019); Bottou et al. (2018); Needell et al. (2014); Shamir and Zhang (2013). We can recover the existing results and provide new complexity bounds when $\theta \in (\frac{1}{2}, 1]$, which have not been studied in the literature to our knowledge.

Theorem 18 (Convergence Rate for RR with Constant Step Size) *Let (B.1)–(B.3) and (P) hold. Let the sequence $\{\mathbf{x}^k\}_{0 \leq k \leq K}$, $K \geq 2$, be generated by RR with $\alpha_k = \lceil \frac{\alpha \log(K)}{K} \rceil e^2$. If $\alpha > 0$ and K satisfies $\frac{\alpha \log(K)}{K} \leq \min\{(\frac{2N\zeta_2}{\text{AL}^2})^{\frac{1}{\omega_2}}, \zeta_2^{-1}, (2L)^{-\frac{1}{e_2}}\}$, then*

$$y_K \leq 2 \left(\frac{L^2 \sigma^2}{\mu N} \right)^{\frac{1}{2\theta}} \left[\frac{\alpha \log(K)}{K} \right]^{\omega_2} + y_0 K^{-\zeta_2 \alpha}. \quad (41)$$

If we set $\alpha = \frac{2\omega_2}{\zeta_2} [1 + \frac{\log(N)}{\log(K)}]$ and K satisfies $K \geq (2\omega_2 \max\{\frac{C_3}{\sqrt{N}}, C_4 N^{1-\frac{1}{2\theta}}, 1\}) \log(NK)$, then we have

$$y_K \leq 2 \left(\frac{4\omega_2}{\theta} \right)^{\omega_2} \left(\frac{L\sigma}{\mu^{3/2}} \right)^{\omega_2} \left[\frac{\log(NK)}{\sqrt{NK}} \right]^{\omega_2} + y_0 (NK)^{-2\omega_2}. \quad (42)$$

The proof is deferred to Appendix D.2. To our knowledge, Theorem 18 is the first non-asymptotic result under the general (θ, μ) -PL condition for random shuffling (Algorithm 2) with constant step size. The existing literature on RR with constant step size mainly assumes the $(\frac{1}{2}, \mu)$ -PL condition (or even strong convexity); see, e.g., HaoChen and Sra (2019); Ahn et al. (2020); Mishchenko et al.; Nguyen et al. (2021). For instance, the work (Nguyen et al., 2021, Theorem 1) obtains the complexity bound $\mathcal{O}(\frac{L^2 \sigma^2 \log^2(NK)}{\mu^3 NK^2} + y_0 \frac{1}{NK^2})$ under strong convexity, which matches our result for $\theta = \frac{1}{2}$.

3.4.2 RESULTS ON POLYNOMIAL STEP SIZES.

We now study the convergence of SGD and RR under the general (θ, μ) -PL condition when utilizing polynomial step sizes. We first proceed with a technical lemma to provide the convergence analysis for sequences $\{y_k\}_k$ that follow the key algorithmic recursion (17).

Lemma 19 *Let $\{y_k\}_{k \geq 0}$ be a given sequence satisfying the following recursion:*

$$y_{k+1} \leq (1 + \ell_1 \alpha_k^\tau) y_k - \ell_2 \alpha_k y_k^{2\theta} + \ell_3 \alpha_k^\tau, \quad (43)$$

where $\ell_1, \ell_2, \ell_3, p > 0$, $\tau > 1$, $\theta \in [\frac{1}{2}, 1]$ and $\alpha_k = \frac{\alpha}{(k+\gamma)^p}$, $\alpha, \gamma > 0$. Let us set

$$\varrho := \frac{2\theta}{(2\theta-1)\tau+1}, \quad \zeta = \theta \ell_2^{\frac{1}{2\theta}} \ell_3^{\frac{2\theta-1}{2\theta}}, \quad u_1 = \frac{p(\tau-1)}{2\theta}, \quad u_2 = \frac{\tau-1}{(2\theta-1)\tau+1}, \quad u_3 = \frac{1-p}{2\theta-1}.$$

The following statements hold:

(a) *If $p \in (0, \varrho)$ and $\gamma \geq \max\{(\frac{\ell_1}{\zeta})^{\frac{1}{u_1}} \alpha^{\frac{1}{p}}, \zeta^{\frac{\varrho}{p}} \alpha^{\frac{1}{p}}, (\frac{2u_1}{\zeta \alpha^{1/\varrho}})^{\frac{1}{1-p/\varrho}}\}$, then we have*

$$y_{k+1} \leq 4 \left(\frac{\ell_3}{\ell_2} \right)^{\frac{1}{2\theta}} \alpha^{\frac{\tau-1}{2\theta}} (k+1+\gamma)^{-u_1} + y_0 \exp\left(-\frac{\alpha^{1/\varrho} \zeta (k+1)}{(k+1+\gamma)^{p/\varrho}}\right).$$

(b) *If $p = \varrho$ and $\alpha \geq (\frac{2u_2}{\zeta})^\varrho$, $\gamma \geq \max\{(\frac{\ell_1}{\zeta})^{\frac{1}{u_2}} \alpha^{\frac{1}{\varrho}}, \zeta \alpha^{\frac{1}{\varrho}}\}$, then it holds that*

$$y_{k+1} \leq 4 \left(\frac{\ell_3}{\ell_2} \right)^{\frac{1}{2\theta}} \alpha^{\frac{\tau-1}{2\theta}} (k+1+\gamma)^{-u_2} + y_0 (\gamma^{-1}(k+1+\gamma))^{-\zeta \alpha^{1/\varrho}}.$$

(c) If $\theta \in (\frac{1}{2}, 1]$, $p \in (0, 1)$, $\alpha \geq \frac{2u_3}{\theta\ell_2}$, and $\gamma \geq \max\{(\frac{\alpha^{\tau-1}\ell_1}{\theta\ell_2})^{\frac{1}{\tau p-1}}, (\frac{\alpha^{\tau-1}\ell_3}{\ell_2})^{\frac{1}{\tau p-u_3-1}}, \alpha\theta\ell_2\}$, then:

$$y_{k+1} \leq 4(k+1+\gamma)^{-u_3} + y_0 (\gamma^{-1}(k+1+\gamma))^{-\theta\ell_2\alpha}.$$

(d) Suppose $\theta \in (\frac{1}{2}, 1]$, $p = 1$. Then, for every $\alpha \geq \frac{2}{\theta(2\theta-1)\ell_2}$, there is $\gamma_0 > 0$, which only depends on $\alpha, \ell_1, \ell_2, \ell_3, \tau$, such that for all $\gamma \geq \gamma_0$, we have

$$y_{k+1} \leq 4 \log(k+1+\gamma)^{-\frac{1}{2\theta-1}} + y_0 \left(\frac{\log(k+1+\gamma)}{\log(\gamma)} \right)^{-\alpha\theta\ell_2}.$$

The proof mainly follows the roadmap in Section 3.2, but requires certain refinements. We provide a detailed discussion in Appendix E.1.

Theorem 20 (Convergence Rate for SGD with Polynomial Step Sizes) Suppose (A.1)–(A.2) and (P) hold. Let the sequence $\{\mathbf{x}^k\}_{0 \leq k \leq K}$ be generated by SGD with $\alpha_k = \frac{\alpha}{(k+\gamma)^{e_1}}$.

Then for any $\alpha \geq (\frac{2\omega_1}{\zeta_1})^{e_1}$, $\gamma \geq \max\{(\frac{L}{2\zeta_1})^{\frac{1}{\omega_1}} \alpha^{\frac{1}{e_1}}, \zeta_1 \alpha^{\frac{1}{e_1}}, (L\alpha)^{\frac{1}{e_1}}\}$, we have


$$y_K \leq 4 \left(\frac{L\sigma^2}{2\mu} \right)^{\frac{1}{2\theta}} \alpha^{\frac{1}{2\theta}} (K+\gamma)^{-\omega_1} + y_0 [\gamma^{-1}(K+\gamma)]^{-\zeta_1 \alpha^{\frac{1}{e_1}}}. \quad (44)$$

If we set $\alpha = (\frac{2\omega_1}{\zeta_1})^{e_1}$ and $\gamma = 2\omega_1 \max\{C_1, C_2, 1\}$, then

$$y_K \leq 4 \left(\frac{\omega_1}{\theta} \right)^{\omega_1} \left(\frac{L\sigma^2}{\mu^2} \right)^{\omega_1} (K+\gamma)^{-\omega_1} + y_0 \left[\frac{2\omega_1 \max\{C_1, C_2, 1\}}{K+\gamma} \right]^{2\omega_1}. \quad (45)$$

The proof is presented in Appendix E. For simplicity, Theorem 20 exclusively addresses the convergence concerning the optimal polynomial step sizes. However, it is apparent that the convergence for any polynomial step sizes $\frac{\alpha}{(k+\gamma)^p}$ with $p \in (0, 1]$ can be directly inferred from Lemma 19. In (Fontaine et al., 2021, Theorem 12), the authors studied polynomial step sizes $\frac{\alpha}{(k+\gamma)^p}$ with $p \in (0, 1)$ under a PL-type condition and a bounded variance condition. By setting $\ell_1 = 0$, $\ell_2 = \mu$, and $\ell_3 = \frac{L\sigma^2}{2}$ in Lemma 19, we can reproduce their findings. Furthermore, Lemma 19 allows us to cover the case $p = 1$ and $\theta \neq \frac{1}{2}$; this scenario, not addressed in (Fontaine et al., 2021, Theorem 12), leads to a rate of $\mathcal{O}((\log(K+\gamma))^{-\frac{1}{2\theta-1}})$ for SGD. In (Fatkhullin et al., 2022, Corollary 1), the convergence rate of SGD is studied under the same (θ, μ) -PL conditions, along with a more general assumption on the variance of the stochastic gradients, but only for the optimal polynomial step sizes. Theorem 20 can recover the most typical cases, i.e., for $h(t) = t$, $b_k = 1$ in (Fatkhullin et al., 2022, Assumption 4), or equivalently, setting $\beta = 1$ and $\tau = 0$ in (Fatkhullin et al., 2022, Corollary 1).

We can also derive some other interesting non-asymptotic results for specific θ in Theorem 20. If $\theta = \frac{1}{2}$, we can obtain the rate $\mathcal{O}(\frac{L\sigma^2}{\mu^2 K} + y_0 [\frac{L}{\mu^2 K}]^2)$. This result matches exactly that of (Gower et al., 2021, Theorem 4.8). For the case when $\theta = 1$ and $p \in [\frac{1}{2}, 1]$, the convergence rate resulting in Lemma 19 matches the results for the convex functions class contained in (Moulines and Bach, 2011, Theorem 4).


 Figure 1: 2D Heatmap of convergence rate with respect to p, θ ; left: SGD, right: RR

Theorem 21 (Convergence Rate for RR with Polynomial Step Sizes) *Suppose (B.1)–(B.3) and (P) hold. Let the sequence $\{\mathbf{x}^k\}_{0 \leq k \leq K}$ be generated by RR with $\alpha_k = \frac{\alpha}{(k+\gamma)^{2\theta}}$. Then for any $\gamma \geq \max\{(\frac{L\sigma^2}{2N\zeta_2})^{\frac{1}{\omega_2}} \alpha^{\frac{1}{2\theta}}, \zeta_2 \alpha^{\frac{1}{2\theta}}, (2L\alpha)^{\frac{1}{2\theta}}\}$, and $\alpha \geq (\frac{2\omega_2}{\zeta_2})^{2\theta}$, we obtain*

$$y_K \leq 4 \left(\frac{L\sigma^2}{\mu N} \right)^{\frac{1}{2\theta}} \alpha^{\frac{1}{2\theta}} (K + \gamma)^{-\omega_2} + y_0 [\gamma^{-1}(K + \gamma)]^{-\zeta_2 \alpha^{\frac{1}{2\theta}}}. \quad (46)$$

Moreover, if $\alpha = \lceil \frac{2\omega_2(1+\log(N))}{\zeta_2} \rceil^{2\theta}$, $\gamma = 2\omega_2(1 + \log(N)) \max\{\frac{C_3}{\sqrt{N}}, C_4 N^{1-\frac{1}{2\theta}}, 1\}$ and $K \geq 2\gamma$, then

$$y_K \leq 4 \left(\frac{4\omega_2}{\theta} \right)^{\omega_2} \left(\frac{L\sigma}{\mu^{3/2}} \right)^{\omega_2} \left[\frac{1 + \log(N)}{\sqrt{N}(K + \gamma)} \right]^{\omega_2} + y_0 \left[\frac{2\omega_2(1 + \log(N)) \max\{\frac{C_3}{\sqrt{N}}, C_4 N^{1-\frac{1}{2\theta}}, 1\}}{N(K + \gamma)} \right]^{2\omega_2}. \quad (47)$$

To the best of our knowledge, Theorem 21 is the first non-asymptotic result for RR under the general (θ, μ) -PL condition with polynomial step sizes.

The Dependence of Convergence Rate on p and θ . From Lemma 19, it is evident that the convergence rate of polynomial step sizes depends significantly on the values of p and θ . When $\theta = \frac{1}{2}$, the upper bounds for SGD and RR are $\mathcal{O}(K^{-\frac{p}{2\theta}})$ and $\mathcal{O}(K^{-\frac{p}{\theta}})$, respectively. For $\theta \in (1/2, 1]$, when $p = 1$, both SGD and RR share a common upper bound of $\mathcal{O}((\log(K + \gamma))^{-\frac{1}{2\theta-1}})$. In the case where $\theta \in (1/2, 1]$ and $p \in (0, 1)$, SGD converges with an upper bound of $\mathcal{O}(K^{-w_1(p, \theta)})$, while RR exhibits a convergence rate of the form $\mathcal{O}(K^{-w_2(p, \theta)})$, where $w_1(p, \theta)$ and $w_2(p, \theta)$ are defined as follows:

$$w_1(p, \theta) = \begin{cases} \frac{p}{2\theta}, & \text{if } p \in (0, \varrho_1], \\ \frac{1-p}{2\theta-1}, & \text{if } p \in (\varrho_1, 1), \end{cases} \quad w_2(p, \theta) = \begin{cases} \frac{p}{\theta}, & \text{if } p \in (0, \varrho_2], \\ \frac{1-p}{2\theta-1}, & \text{if } p \in (\varrho_2, 1). \end{cases}$$

A visualization of the rate mappings w_1 and w_2 is provided in Figure 1. Notably, the full characterization of the rate over all instances of $p \in (0, 1]$ and $\theta \in [\frac{1}{2}, 1]$ for RR seems new.

3.5 Exponential Step Sizes Adapt to the Geometry of Objective Functions

The convergence rates of SGD or RR employing constant, polynomial, and cosine step size rules rely significantly on the choice of parameters. Achieving the optimal convergence rate necessitates a careful selection of these parameters in consideration of θ . In the case of a constant step size, $\alpha_k = \lceil \frac{\alpha \log(K)}{K} \rceil^p$, and a polynomial step size, $\alpha_k = \frac{\alpha}{(k+\gamma)^p}$, optimal performance is attained when $p = \varrho = \frac{2\theta}{(2\theta-1)\tau+1}$. Similarly, for a cosine step size, $\alpha_k = \alpha \lceil \frac{1+\cos(\frac{k\pi}{K})}{2} \rceil^\beta$, the optimal rate is achieved when $\alpha \sim (\frac{\log(K)}{K})^e$. Intriguingly, the exponential step sizes rule, $\alpha_k = \alpha (\frac{\beta}{K})^{\frac{pk}{K}}$, exhibits insensitivity to the parameters α , β , and p , consistently matching the optimal convergence rate of other step size rules for any $\theta \in [\frac{1}{2}, 1]$ (up to a logarithmic term).

We conduct a numerical simulation to visually demonstrate this property. The simulation involves the following recursion, representing the worst-case scenario of equation (17) with $\ell_1 = \ell_2 = \ell_3 = 1$, $\tau = 2$:

$$y_{k+1} = (1 + \alpha_k^2)y_k - \alpha_k y_k^{2\theta} + \alpha_k^2, \quad k \in [K].$$

We calculate y_K using polynomial step sizes $\alpha_k = \frac{\alpha}{(k+\gamma)^p}$ with $p = 1, \frac{4}{5}, \frac{2}{3}, \frac{1}{2}$, and an exponential step size $\alpha_k = \alpha (\frac{1}{K})^{\frac{k}{K}}$, for $\theta = \frac{1}{2}, \frac{2}{3}, 1$, considering various total numbers of iterations K . In Figure 2, the relationship between $\log_2(y_K)$ and $\log_2(K)$ is depicted for large K . The slope of each line represents the sublinear convergence rate corresponding to each step size rule, where a steeper slope indicates a faster sublinear convergence rate. Two important facts are illustrated in this figure:

- Polynomial step sizes achieve the optimal rate with different values of p for different θ . The optimal choice in our simulation is $p = 1, \frac{4}{5}, \frac{2}{3}$ for $\theta = \frac{1}{2}, \frac{2}{3}, 1$ respectively, which matches the optimal choice $p = \varrho_1 = \frac{2\theta}{4\theta-1}$ in Lemma 19.
- For different values of θ , exponential step sizes automatically attain the optimal rate that can be achieved by the polynomial step size with optimal p .




Figure 2: The relation between $\log_2(y_K)$ and $\log_2(K)$ for different θ with polynomial and exponential step sizes. A steeper slope corresponds to a faster sublinear rate.

4 Conclusion

In this work, we propose a generalized version of Chung’s lemma, which can serve as a simple yet powerful tool to obtain non-asymptotic convergence results for sequences $\{a_k\}_k$ following the recursion

$$a_{k+1} \leq \left(1 - \frac{1}{s(b_k)}\right)a_k + \frac{1}{t(b_k)},$$

where $s(b_k)$ and $t(b_k)$ are general terms controlling and capturing the growth behavior of $\{a_k\}_k$. In the special setting $s(b_k) = \mathcal{O}(k^p)$ and $t(b_k) = \mathcal{O}(k^{p+q})$, the classical Chung’s lemma is applicable yielding $a_k = \mathcal{O}(k^p/k^{p+q}) = \mathcal{O}(k^{-q})$. Our results expand upon this classical case by providing mild and verifiable conditions on $s(b_k)$ and $t(b_k)$ to ensure $a_k = \mathcal{O}(s(b_k)/t(b_k))$. We illustrate the broad applicability of the generalized version of Chung’s lemma by developing a unified framework for the convergence analysis of stochastic algorithms under the (θ, μ) -PL condition and utilizing various step size rules and sampling schemes.

Our generalized Chung’s lemma plays an important role in two aspects: (a) it guides the construction of the auxiliary sequence $\{u_k\}_k$ when addressing general (θ, μ) -PL conditions, as illustrated in Step 2 of our roadmap detailed in Section 3.2; (b) by combining the generalized Chung’s lemma with the extension lemma (refer to Lemma 5), we provide a unified analysis path for exponential and cosine step size rules, which is summarized as a splitting technique in Section 3.3. Intriguingly, we observe that exponential step sizes can adapt to the geometry of the objective function (determined by the PL parameters (θ, μ)). More specifically, exponential step sizes can automatically achieve an optimal convergence rate without explicitly knowing the PL exponent θ in the (θ, μ) -PL condition. Further investigation of this geometric feature is left as a future research direction.

Acknowledgments and Disclosure of Funding

Xiao Li was partly supported by the National Natural Science Foundation of China under grant No. 12201534 and by the Shenzhen Science and Technology Program under Grant No. RCYX20221008093033010. Andre Milzarek was partly supported by the National Natural Science Foundation of China under grant No. 12150410304, by the Shenzhen Science and Technology Program under grant No. RCYX20210609103124047 and RCYX20221008093033010, and by the Guangdong Provincial Key Laboratory of Mathematical Foundations for Artificial Intelligence (2023B1212010001).

Appendix A. Preparatory Tools

In this first section, we list several well-known mathematical facts and tools that will be used repeatedly in the later analysis.

Proposition 22 (Integral Test) *Let $I = [\iota, \infty)$, $\iota \in \mathbb{R}$, be a given interval and let $f : I \rightarrow \mathbb{R}$ be continuous and non-increasing. Then, for all $a, b \in \mathbb{N} \cap I$ with $a \leq b$, we have*

$$\int_a^{b+1} f(x) dx \leq \sum_{k=a}^b f(k) \leq f(a) + \int_a^b f(x) dx.$$

Appendix B. Non-asymptotic Extension of Classic Chung's Lemma

Classic Chung's lemma focuses on the sequence $\{a_k\}_k \subseteq \mathbb{R}_{++}$ satisfying the following recursion:

$$a_{k+1} \leq \left(1 - \frac{c}{(k+\gamma)^p}\right) a_k + \frac{d}{(k+\gamma)^{p+q}}, \quad p \in (0, 1], \quad q > 0, \quad c, d > 0, \quad \gamma \geq c^{\frac{1}{p}}. \quad (48)$$

In [Chung \(1954\)](#), Chung proposed several lemmas to obtain asymptotic estimates for a_k , which are summarized as follows:

Lemma 23 [*Asymptotic Chung's Lemma* [Chung \(1954\)](#); [Polyak \(1987\)](#)] *Let the sequence $\{a_k\}_k \subseteq \mathbb{R}_{++}$ be given as in (48). Then, the following statements hold:*

- (a) *When $p \in (0, 1)$, we have $a_k = \mathcal{O}(k^{-q})$.*
- (b) *When $p = 1$, if $c = q$ we have $a_k = \mathcal{O}(k^{-q} \log(k))$; otherwise, $a_k = \mathcal{O}(k^{-\min\{c, q\}})$.*

There already have been several extensions to non-asymptotic versions of certain subcases of [Theorem 23](#) in previous literature, see [Moulines and Bach \(2011\)](#); [Ahn et al. \(2020\)](#); [Nguyen et al. \(2021\)](#). In this paper, we utilize [Theorem 1](#) to derive a complete non-asymptotic extension of the classic Chung's lemma, which encompasses the previous extensions.

Theorem 24 (Non-asymptotic Chung's Lemma) *Let the sequence $\{a_k\}_k \subseteq \mathbb{R}_{++}$ be given and satisfy (48). Then, the following statements hold:*

- (a) *If $p \in (0, 1)$, $\gamma \geq c^{\frac{1}{p}}$ and $\gamma > (\frac{q}{c})^{\frac{1}{1-p}}$, then*

$$a_{k+1} \leq \frac{\lambda d}{c} (k+1+\gamma)^{-q} + \left(a_0 - \frac{\lambda d}{c\gamma^q}\right)^+ \exp\left(\frac{c\gamma^{1-p}}{1-p}\right) \exp\left[-\frac{c(k+1+\gamma)^{1-p}}{1-p}\right],$$

holds for all $k \in [K]$, where $\lambda = \frac{c\gamma^{1-p}}{c\gamma^{1-p}-q}$ and $(x)^+ := \max\{x, 0\}$. Additionally, if we further assume $\gamma \geq \left[\frac{(1+\varsigma)q}{c}\right]^{\frac{1}{1-p}}$ for some $\varsigma > 0$, then

$$a_{k+1} \leq \frac{(1+\varsigma)d}{\varsigma c} (k+1+\gamma)^{-q} + \left(a_0 - \frac{\lambda d}{c\gamma^q}\right)^+ \exp\left[-\frac{c(k+1)}{(k+1+\gamma)^p}\right],$$

holds for all $k \in [K]$.

- (b) *If $p = 1$, then*

$$a_{k+1} \leq \begin{cases} \frac{d}{c-q} (k+1+\gamma)^{-q} + \gamma^c \left(a_0 - \frac{d}{(c-q)\gamma^q}\right)^+ (k+1+\gamma)^{-c}, & \text{for } c > q, \gamma \geq c, \\ \left[\frac{d}{\gamma-c} + d \log\left(\frac{k+\gamma-c}{\gamma-c}\right) + a_0 \gamma^c\right] (k+1+\gamma)^{-c}, & \text{for } c = q, \gamma > c, \\ \left[\frac{d}{(q-c)(\gamma-q-1)^{q-c}} + a_0 \gamma^c\right] (k+\gamma+1)^{-c} - \frac{d}{q-c} (k+\gamma+1)^{-q}, & \text{for } c < q, \gamma > q+1. \end{cases}$$

Proof We now apply our main theorem (Theorem 1) to prove the non-asymptotic version of the classical Chung's lemma. To make use of Theorem 1, we first specify the following sequences and functions:

$$b_k := k + \gamma, \quad s(x) := \frac{x^p}{c}, \quad \text{and} \quad t(x) := \frac{x^{p+q}}{d}.$$

We now only need to check whether the conditions in Theorem 1 are fulfilled in this setting. Let us notice that the rate mapping $r(x) := s(x)/t(x) = \frac{d}{c}x^{-q}$ is convex on \mathbb{R}_{++} . In addition, it holds that

$$u(x) := s'(x) - r(x)t'(x) = \frac{p}{c}x^{p-1} - \frac{d}{c}x^{-q} \cdot \frac{p+q}{d}x^{p+q-1} = -\frac{q}{c}x^{p-1}.$$

and $s(x) \geq 1$ for all $x \geq c^{\frac{1}{p}}$. Per assumption, we have $\gamma \geq c^{\frac{1}{p}}$ which implies $b_k \in I := [c^{\frac{1}{p}}, \infty) \subseteq \mathbb{R}_{++}$ for all $k \in [K]$. Hence, we only need to verify the condition (7) or (8) in Theorem 1. To that end and using $b_{k+1} - b_k = 1$, we need to find a sequence $\{\lambda_k\}_k \subseteq \mathbb{R}_+$ such that

$$\lambda_{k+1} \left[\frac{(k+\gamma)^p}{c} - \frac{q(k+\gamma)^{p-1}}{c} \right] \geq \lambda_k \left[\frac{(k+\gamma)^p}{c} - 1 \right] + 1, \quad \forall k \in [K] \quad (49)$$

or a non-decreasing sequence $\{\lambda_k\}_k \subseteq \mathbb{R}_+$ such that

$$-\frac{q}{c}(k+\gamma)^{p-1} \geq -1 + \frac{1}{\lambda_{k+1}}, \quad \forall k \in [K]. \quad (50)$$

Based on the parameter $p \in (0, 1]$, we now discuss different cases.

Case I: $p \in (0, 1)$. This has already been shown in Lemma 4 (a). Furthermore, if we additionally have $\gamma \geq [\frac{(1+\varsigma)q}{c}]^{\frac{1}{1-p}}$ for some $\varsigma > 0$, then it follows $\gamma^{1-p} \geq \frac{(1+\varsigma)q}{c}$ and $\lambda = \frac{c\gamma^{1-p}}{c\gamma^{1-p-q}} \leq \frac{1+\varsigma}{\varsigma}$. Using the concavity of the mapping $x \mapsto x^{1-p}$, it holds that

$$(k+1+\gamma)^{1-p} - \gamma^{1-p} \geq \frac{1-p}{(k+1+\gamma)^p} [(k+1+\gamma) - \gamma] = (1-p) \frac{k+1}{(k+1+\gamma)^p}.$$

Combining these last estimate, we obtain

$$a_{k+1} \leq \frac{(1+\varsigma)d}{\varsigma c} (k+1+\gamma)^{-q} + \left(a_0 - \frac{\lambda d}{c\gamma^q} \right)^+ \exp \left(-\frac{c(k+1)}{(k+1+\gamma)^p} \right).$$

Case II: $p = 1$. Since the analysis technique depends on the relation between the constants t and c , we will split the following discussion into three sub-cases.

Sub-Case 2-1: $c > q$ and $\gamma \geq c$. This has already been shown in Lemma 4 (b).

Sub-Case 2-2: $c = q$ and $\gamma > c$. In this case, we can no longer ensure existence of a sequence $\{\lambda_k\}_k \subseteq \mathbb{R}_+$ satisfying the (simpler) condition (50). To that end, we want to find $\{\lambda_k\}_k \subseteq \mathbb{R}_+$ such that the (more general) condition (49) is valid, i.e., the following inequality should hold

$$\lambda_{k+1} \left[\frac{k+\gamma}{c} - 1 \right] \geq \lambda_k \left[\frac{k+\gamma}{c} - 1 \right] + 1, \quad \forall k \in [K].$$

A feasible choice of $\{\lambda_k\}_k$ to guarantee the above condition is $\lambda_0 = 0$ and $\lambda_{k+1} = \sum_{i=0}^k \frac{c}{i+\gamma-c}$ for all $k \in [K]$. In fact, this choice satisfies

$$\lambda_{k+1} \left[\frac{k+\gamma}{c} - 1 \right] = \left[\lambda_k + \frac{c}{k+\gamma-c} \right] \left[\frac{k+\gamma}{c} - 1 \right] = \lambda_k \left[\frac{k+\gamma}{c} - 1 \right] + 1 \quad \forall k \in [K].$$

Invoking the integral test, we further have $\lambda_{k+1} \leq \frac{c}{\gamma-c} + c \int_0^k \frac{1}{x+\gamma-c} dx = \frac{c}{\gamma-c} + c \log\left(\frac{k+\gamma-c}{\gamma-c}\right)$, for all $k \in [K]$. Hence, utilizing Theorem 1 (a) and this upper bound for $\{\lambda_k\}_k$, we can conclude

$$\begin{aligned} a_{k+1} &\leq \lambda_{k+1} r(b_{k+1}) + (a_0 - \lambda_0 r(b_0)) \prod_{i=0}^k \left(1 - \frac{1}{s(b_i)}\right) \\ &\leq \left[\frac{d}{\gamma-c} + d \log\left(\frac{k+\gamma-c}{\gamma-c}\right) \right] (k+1+\gamma)^{-c} + a_0 \prod_{i=0}^k \left(1 - \frac{c}{i+\gamma}\right) \\ &\leq \left[\frac{d}{\gamma-c} + d \log\left(\frac{k+\gamma-c}{\gamma-c}\right) + \gamma^c a_0 \right] (k+1+\gamma)^{-c}, \end{aligned}$$

where the last inequality follows again from the inequality $1-x \leq \exp(-x)$ and the integral test (Proposition 22).

Sub-Case 2-3: $c < q$ and $\gamma > q+1$. Similar to **Sub-Case 2-2**, we need to find a sequence $\{\lambda_k\}_k \subseteq \mathbb{R}_+$ such that the condition (49) holds and to ensure applicability of Theorem 1 (a). We first reformulate (49) as follows:

$$\lambda_{k+1} \left[\frac{k+\gamma}{c} - \frac{q}{c} \right] \geq \lambda_k \left[\frac{k+\gamma}{c} - 1 \right] + 1 \iff \lambda_{k+1}(k+\gamma-q) \geq \lambda_k(k+\gamma-c) + c.$$

Defining $\lambda_k := \zeta(k-1+\gamma-q)^{q-c} - \frac{c}{q-c}$ and $\zeta := \frac{c}{(q-c)(\gamma-q-1)^{q-c}}$, we then have

$$\begin{aligned} \lambda_{k+1}(k+\gamma-q) &\geq \lambda_k(k+\gamma-c) + c \\ \iff \zeta(k+\gamma-q)^{q-c+1} &\geq \lambda_k(k+\gamma-c) + c + \frac{c}{q-c}(k+\gamma-q) \\ \iff \zeta(k+\gamma-q)^{q-c+1} &\geq \zeta(k-1+\gamma-q)^{q-c}(k+\gamma-c) + \frac{c^2}{q-c} + c - \frac{cq}{q-c} \\ \iff (k+\gamma-q)^{q-c+1} &\geq (k-1+\gamma-q)^{q-c}(k+\gamma-c). \end{aligned} \tag{51}$$

Let us now verify (51). For any $x \geq 0, \eta > 0$, by the mean value theorem, there exists $\xi \in (0, 1)$ such that $(x+1)^\eta - x^\eta = \eta(x+\xi)^{\eta-1} \geq \frac{\eta x^\eta}{x+1}$. Rearranging the terms in this inequality, it follows

$$(x+1)^\eta - x^\eta \geq \frac{\eta x^\eta}{x+1} \iff (x+1)^{\eta+1} \geq x^\eta(x+1+\eta).$$

Choosing $\eta = q-c$ and $x = k-1+\gamma-q$ in the above estimate (and recalling $\gamma > q+1$), this proves the condition (51). Finally, invoking Theorem 1 (a) with such choice of $\{\lambda_k\}_k$,

we obtain

$$\begin{aligned}
 a_{k+1} &\leq \lambda_{k+1} r(b_{k+1}) + (a_0 - \lambda_0 r(b_0)) \prod_{i=0}^k \left(1 - \frac{c}{i + \gamma}\right) \\
 &\leq \frac{\zeta d}{c} (k + \gamma + 1)^{-c} - \frac{d}{q - c} (k + \gamma + 1)^{-q} + a_0 \prod_{i=0}^k \left(1 - \frac{c}{i + \gamma}\right) \\
 &\leq \left(\frac{c}{(q - c)(\gamma - q - 1)^{q-c}} + a_0 \gamma^c\right) (k + \gamma + 1)^{-c} - \frac{d}{q - c} (k + \gamma + 1)^{-q},
 \end{aligned}$$

where the second line holds owing to $(k + \gamma - q) \leq (k + \gamma + 1)$ and $\lambda_0 = 0$, and the last inequality utilizes the inequality $1 - x \leq \exp(-x)$ and the integral test (Proposition 22). ■

Appendix C. Additional Derivations for Exponential and Cosine Step Sizes

We now study the convergence of SGD and RR under the general (θ, μ) -PL condition when utilizing exponential, and cosine step size schemes. We provide the details of the verification for Theorem 13 to 16 using Lemma 12.

C.1 Exponential Step Sizes: Verifying Theorem 13 and 14

Proof of Theorem 13 By assumption, we have $\alpha_k = \alpha \gamma^k \leq \alpha \left(\frac{1}{2}\right)^{\frac{e_1 k}{K}} \leq \alpha \leq \frac{1}{L}$ for all $k \in [K]$. Hence, the descent property (15) is applicable for all $k \in [K]$.

The remaining conditions on α , β , and K ensure that Lemma 12 (b) is applicable which establishes (30) by setting $\ell_1 = \frac{AL}{2}$, $\ell_2 = \mu$, and $\ell_3 = \frac{L\sigma^2}{2}$ and using

$$\left(\frac{2p\omega_1}{\rho_1}\right)^{\omega_1} \zeta^{-\omega_1} \left(\frac{\ell_3}{\ell_2}\right)^{\frac{1}{2\theta}} = \left(\frac{p}{\theta^2}\right)^{\omega_1} \mu^{-\frac{\omega_1+1}{2\theta}} \left(\frac{L\sigma^2}{2}\right)^{\frac{1-(2\theta-1)\omega_1}{2\theta}} = \left(\frac{p}{2\theta^2}\right)^{\omega_1} \mu^{-2\omega_1} (L\sigma^2)^{\omega_1},$$

where the last equation is due to $\frac{\omega_1+1}{2\theta} = 2\omega_1$ and $\frac{1-(2\theta-1)\omega_1}{2\theta} = \omega_1$. ■

Proof of Theorem 14 The inequality $\alpha \leq (2L)^{-\frac{1}{e_2}}$ ensures that $\alpha_k \leq \frac{1}{2L}$, which means that (16) is applicable. The first inequality (31) can be obtained by applying Lemma 19 to (16) with $\ell_1 = \frac{AL^2}{2N}$, $\ell_2 = \frac{\mu}{2}$, and $\ell_3 = \frac{L^2\sigma^2}{2N}$. Furthermore, if we have $K \geq \left(\frac{4p}{\theta^2}\right) \max\left\{\frac{C_3}{\sqrt{N}}, C_4 N^{1-\frac{1}{2\theta}}, 1\right\} \log\left(\frac{K}{\beta}\right) \log(NK)$, then

$$y_K \leq 4 \left(\frac{4p}{\theta^2}\right)^{\omega_2} \left(\frac{L\sigma}{\mu^{3/2}}\right)^{\omega_2} \left[\frac{\log\left(\frac{K}{\beta}\right)}{\sqrt{NK}}\right]^{\omega_2} + y_0 (NK)^{-\omega_2}.$$

■

C.2 Cosine Step Sizes: Verifying Theorem 15 and 16

Proof of Theorem 15 The inequality $\alpha \leq L^{-\frac{1}{e_1}}$ ensures that $\alpha_k \leq \frac{1}{L}$, which means that (15) is applicable. The first two inequalities (33) and (34) can be obtained by ap-

plying Lemma 19 to (15) with $\ell_1 = \frac{\text{AL}}{2}$, $\ell_2 = \mu$, $\ell_3 = \frac{\text{L}\sigma^2}{2}$, and $\frac{\varrho_1}{\omega_1} = 2\theta$. To establish (35), we substitute $\alpha = 2^{\max\{\beta, \varrho_1\}} \left(\frac{\omega_1 \log(K)}{\zeta_1 K}\right)^{\varrho_1}$ into (33) and (34). To make sure that (33) and (34) available, we need to guarantee that $\alpha \leq \min\left\{\left(\frac{2\zeta_1}{\text{AL}}\right)^{2\theta}, \zeta_1^{-\varrho_1}, \text{L}^{-1}\right\}$, i.e., $K \geq 2^{\max\{2, 1 + \frac{\beta}{\varrho_1}\}} \omega_1 (\max\{C_1, C_2, 1\} \log(K))$, and $K \geq \frac{2\pi^2\beta}{\theta\alpha^{1/\varrho_1}\zeta_1}$, i.e., $K \geq \exp\left[\frac{\pi^2\beta}{2^{\max\{1, \beta/\varrho_1\}}\omega_1}\right]$. We summarize these requirements on K as $K \geq \Theta(\max\{C_1, C_2, 1\} \log(K))$. Then,

$$y_K \leq \mathcal{O}\left(\alpha^{\frac{\omega_1}{\varrho_1} - \frac{2\beta\omega_1/\varrho_1}{2\beta+\varrho_1}} \left[\frac{1}{K}\right]^{\frac{2\beta\omega_1}{2\beta+\varrho_1}}\right) + \mathcal{O}(y_0 K^{-2\omega_1}) = \mathcal{O}\left(\left[\frac{(\log(K))^{\frac{\varrho_1}{2\beta+\varrho_1}}}{K}\right]^{\omega_1}\right).$$

■

Proof of Theorem 16 The inequality $\alpha \leq (2\text{L})^{-\frac{1}{\varrho_2}}$ ensures that $\alpha_k \leq \frac{1}{2\text{L}}$, which means that (16) is applicable. The first two inequalities (36) and (37) can be obtained by applying Lemma 12 to (16) with $\ell_1 = \frac{\text{AL}^2}{2N}$, $\ell_2 = \frac{\mu}{2}$, $\ell_3 = \frac{\text{L}^2\sigma^2}{2N}$, and $\frac{\varrho_2}{\omega_2} = \theta$. To establish (38), we substitute $\alpha = 2^{\max\{\varrho_2, \beta\}} \left(\frac{2\omega_2 \log(NK)}{\zeta_2 K}\right)^{\varrho_2}$ into (36) and (37). To make sure (36) and (37) available, we need to guarantee that $\alpha \leq \min\left\{\left(\frac{2N\zeta_2}{\text{AL}^2}\right)^\theta, \zeta_2^{-\varrho_2}, (2\text{L})^{-1}\right\}$, i.e.,

$$K \geq \left(2^{\max\{2, 1 + \frac{\beta}{\varrho_2}\}} \cdot \omega_2 \max\left\{\frac{C_3}{\sqrt{N}}, C_4 N^{1-\frac{1}{2\theta}}, 1\right\}\right) \log(NK),$$

and $K \geq \frac{4\pi^2\beta}{\theta\alpha^{1/\varrho_2}\zeta_2}$, i.e., $K \geq \exp\left(\frac{2\pi^2\beta}{2^{\max\{1, \beta/\varrho_2\}}\theta\omega_2 \log(NK)}\right)$. We summarize these requirements for K as $K \geq \Theta(\max\{\frac{C_3}{\sqrt{N}}, C_4 N^{1-\frac{1}{2\theta}}, 1\} \log(NK))$. Recall that $\zeta_2 = \frac{\theta}{2}\mu^{\frac{1}{2\theta}} \left(\frac{\text{L}^2\sigma^2}{N}\right)^{1-\frac{1}{2\theta}}$, then,

$$\begin{aligned} y_K &\leq \mathcal{O}\left(N^{-\frac{1}{2\theta}} \alpha^{\frac{\omega_2}{\varrho_2}} \left[\frac{1}{\alpha^{1/\varrho_2}\zeta_2}\right]^{\frac{2\beta\omega_2}{2\beta+\varrho_2}}\right) + \mathcal{O}(y_0 (NK)^{-2\omega_2}) \\ &= \mathcal{O}\left(N^{-\frac{1}{2\theta}} \left[\frac{\log(NK)}{\zeta_2 K}\right]^{\omega_2} \left[\frac{1}{\log(NK)}\right]^{\frac{2\beta\omega_2}{2\beta+\varrho_2}}\right) + \mathcal{O}(y_0 (NK)^{-2\omega_2}) \\ &= \mathcal{O}\left(N^{-\frac{1}{2\theta} - (1-\frac{1}{2\theta})\omega_2} \left[\frac{\log(NK)^{\frac{\varrho_2}{2\beta+\varrho_2}}}{K}\right]^{\omega_2}\right) + \mathcal{O}(y_0 (NK)^{-2\omega_2}) \\ &= \mathcal{O}\left(\left[\frac{\log(NK)^{\frac{\varrho_2}{2\beta+\varrho_2}}}{\sqrt{N}K}\right]^{\omega_2}\right), \end{aligned}$$

where in the last step, we utilize $\frac{1}{2\theta} + (1 - \frac{1}{2\theta})\omega_2 = \frac{\omega_2}{2}$. ■

Appendix D. Derivations for Constant Step Size

We now study convergence of SGD and RR under the general (θ, μ) -PL condition when utilizing constant step size schemes. We first proceed with a technical lemma and apply Chung's lemma to sequences $\{y_k\}_k$ that follow the key algorithmic recursion

$$y_{k+1} \leq (1 + \ell_1 \alpha_k^\tau) y_k - \ell_2 \alpha_k y_k^{2\theta} + \ell_3 \alpha_k^\tau$$

from equation (17). In subsection E.2, we then utilize the obtained results to provide proofs for Theorem 20 and Theorem 21.

D.1 A Technical Lemma for Constant Step Size

Lemma 25 Consider a non-negative sequence $\{y_k\}_{0 \leq k \leq K}$ satisfying the recursion

$$y_{k+1} \leq (1 + \ell_1 \alpha_k^\tau) y_k - \ell_2 \alpha_k y_k^{2\theta} + \ell_3 \alpha_k^\tau \quad (52)$$

for all $k \in [K-1]$, where $\ell_1, \ell_2, \ell_3 > 0$, $K \in \mathbb{N}$ and $K \geq 1$. Let us define

$$\varrho := \frac{2\theta}{(2\theta-1)\tau+1}, \quad \omega := \frac{\tau-1}{(2\theta-1)\tau+1}, \quad \zeta := \theta \ell_2^{\frac{1}{2\theta}} \ell_3^{\frac{2\theta-1}{2\theta}}.$$

Suppose $\alpha_k = \lceil \frac{\alpha \log(K)}{K} \rceil^\varrho$ where $\alpha > 0$ and K satisfy $\frac{\alpha \log(K)}{K} \leq \min\{(\frac{\zeta}{\ell_1})^{\frac{1}{\omega}}, \zeta^{-1}\}$; then:

$$y_K \leq 2 \left(\frac{\ell_3}{\ell_2} \right)^{\frac{1}{2\theta}} \left[\frac{\alpha \log(K)}{K} \right]^\omega + y_0 K^{-\zeta \alpha}.$$

Proof Recall that the preceding analysis outlined in the roadmap in Section 3.2 is applicable provided that $\alpha_k \leq \min\{(\frac{\zeta}{\ell_1})^{\frac{1}{\omega}}, \zeta^{-\varrho}\}$. As stated in the assumptions, suppose that the step sizes $\{\alpha_k\}_k$ satisfy $\alpha_k = \lceil \frac{\alpha \log(K)}{K} \rceil^\varrho$ and we have $\frac{\alpha \log(K)}{K} \leq \min\{(\frac{\zeta}{\ell_1})^{\frac{1}{\omega}}, \zeta^{-1}\}$. Let us set $b_k = k$ and $\eta(x) = \lceil \frac{\alpha \log(K)}{K} \rceil^\varrho$, where b_k and η follow the same definition in Section 3.2. Then, the rate function r is obviously non-increasing and convex and it holds that

$$(b_{k+1} - b_k)u(b_k) = 0 \quad \forall k \in [K-1].$$

Hence, Theorem 1 (b) is applicable to (22) with $\lambda_k \equiv 1$ and we obtain

$$\begin{aligned} y_K &\leq r(K) + (y_0 - r(0)) \prod_{i=0}^{K-1} \left(1 - \frac{1}{s(i)} \right) \\ &= 2 \left(\frac{\ell_3}{\ell_2} \right)^{\frac{1}{2\theta}} \left[\frac{\alpha \log(K)}{K} \right]^\omega + y_0 \prod_{k=0}^{K-1} \left(1 - \zeta \frac{\alpha \log(K)}{K} \right) \\ &\leq 2 \left(\frac{\ell_3}{\ell_2} \right)^{\frac{1}{2\theta}} \left[\frac{\alpha \log(K)}{K} \right]^\omega + y_0 \exp \left(-K \zeta \frac{\alpha \log(K)}{K} \right) \\ &= 2 \left(\frac{\ell_3}{\ell_2} \right)^{\frac{1}{2\theta}} \left[\frac{\alpha \log(K)}{K} \right]^\omega + y_0 K^{-\zeta \alpha}, \end{aligned}$$

where we utilized $1 - x \leq \exp(-x)$. ■

D.2 Verifying Theorem 17 and 18

Proof of Theorem 17 The inequality $\frac{\alpha \log(K)}{K} \leq L^{-\frac{1}{\varepsilon_1}}$ ensures that $\alpha_k \leq \frac{1}{L}$, and thus, the descent property (15) for SGD is valid. The rate in (39) can be obtained by applying Lemma 19 in (15) with $\ell_1 = \frac{A}{2}$, $\ell_2 = \mu$, and $\ell_3 = \frac{L\sigma^2}{2}$. To show (40), we substitute $\alpha = \frac{2\omega_1}{\zeta_1}$ into (39). Then, we have

$$y_K \leq 2 \left(\frac{\ell_3}{\ell_2} \right)^{\frac{1}{2\theta}} \left(\frac{2\omega_1}{\zeta} \right)^{\omega_1} \left[\frac{\log(K)}{K} \right]^{\omega_1} + \frac{y_0}{K^{2\omega_1}} = 2 \left(\frac{\omega_1}{\theta} \right)^{\omega_1} \left(\frac{L\sigma^2}{\mu^2} \right)^{\omega_1} \left[\frac{\log(K)}{K} \right]^{\omega_1} + \frac{y_0}{K^{2\omega_1}},$$

when $\frac{\log(K)}{K} \leq \frac{\zeta_1}{2\omega_1} \min\{(\frac{2\zeta_1}{\mathbf{AL}})^{\frac{1}{\omega_1}}, \zeta_1^{-1}, \mathbf{L}^{-\frac{1}{e_1}}\}$, i.e., for all $K \geq (2\omega_1 \max\{C_1, C_2, 1\}) \log(K)$. ■

Proof of Theorem 18 The inequality $\frac{\alpha \log(K)}{K} \leq (2\mathbf{L})^{-\frac{1}{e_2}}$ ensures that $\alpha_k \leq \frac{1}{2\mathbf{L}}$, which means that (16) is applicable. The first inequality (41) can be obtained by applying Lemma 19 to (16) with $\ell_1 = \frac{\mathbf{AL}^2}{2N}$, $\ell_2 = \frac{\mu}{2}$, and $\ell_3 = \frac{\mathbf{L}^2 \sigma^2}{2N}$. To establish (42), we substitute $\alpha = \frac{2\omega_2}{\zeta_2} [1 + \frac{\log(N)}{\log(K)}]$ into (41). Then, we have

$$\begin{aligned} y_K &\leq 2 \left(\frac{\ell_3}{\ell_2}\right)^{\frac{1}{2\theta}} \left(\frac{2\omega_2}{\zeta}\right)^{\omega_2} \left[\frac{\log(NK)}{K}\right]^{\omega_2} + y_0 K^{-2\omega_2(1+\frac{\log(N)}{\log(K)})} \\ &= 2 \left(\frac{4\omega_2}{\theta}\right)^{\omega_2} \left(\frac{\mathbf{L}\sigma}{\mu^{3/2}}\right)^{\omega_2} \left[\frac{\log(NK)}{\sqrt{NK}}\right]^{\omega_2} + y_0 (NK)^{-2\omega_2}, \end{aligned}$$

when $\frac{\alpha \log(K)}{K} \leq \min\{(\frac{2N\zeta_2}{\mathbf{AL}^2})^{\frac{1}{\omega_2}}, \zeta_2^{-1}, (2\mathbf{L})^{-\frac{1}{e_2}}\}$, which can be equivalently expressed as $K \geq (2\omega_2 \max\{\frac{C_3}{\sqrt{N}}, C_4 N^{1-\frac{1}{2\theta}}, 1\}) \log(NK)$. ■

Appendix E. Derivations for Polynomial Step Sizes

We now study the convergence of SGD and RR under the general (θ, μ) -PL condition when utilizing polynomial step size schemes. We first provide proof for the key technical lemma, i.e., Lemma 19. In subsection E.2, we then utilize the obtained results to provide proofs for Theorem 20 and Theorem 21.

E.1 Proof of Lemma 19

We will divide the proof into two core cases: $\theta = \frac{1}{2}$ and $\theta \neq \frac{1}{2}$.

Case I: $\theta = \frac{1}{2}$. In this case, we have $\varrho = 1$, $\zeta = \frac{1}{2}\ell_2$, and $u_1 = p(\tau - 1)$ and the recursion (43) simplifies to

$$y_{k+1} \leq [1 + \ell_1 \alpha_k^\tau - \ell_2 \alpha_k] y_k + \ell_3 \alpha_k^\tau. \quad (53)$$

Sub-Case 1-1: $p \in (0, 1)$. Using $\alpha_k = \frac{\alpha}{(k+\gamma)^p}$ and $\gamma \geq (\frac{\ell_1}{\zeta})^{\frac{1}{u_1}} \alpha^{\frac{1}{p}} = (\frac{2\ell_1}{\ell_2})^{\frac{1}{(\tau-1)p}} \alpha^{\frac{1}{p}}$, (53) can be expressed as follows

$$\begin{aligned} y_{k+1} &\leq \left[1 + \frac{\ell_1 \alpha^\tau}{(k+\gamma)^{\tau p}} - \frac{\ell_2 \alpha}{(k+\gamma)^p}\right] y_k + \frac{\alpha^\tau \ell_3}{(k+\gamma)^{\tau p}} \\ &= \left[1 + \left(\frac{\ell_1 \alpha^{\tau-1}}{\ell_2 (k+\gamma)^{(\tau-1)p}} - 1\right) \frac{\ell_2 \alpha}{(k+\gamma)^p}\right] y_k + \frac{\alpha^\tau \ell_3}{(k+\gamma)^{\tau p}} \\ &\leq \left[1 - \frac{\ell_2 \alpha}{2(k+\gamma)^p}\right] y_k + \frac{\alpha^\tau \ell_3}{(k+\gamma)^{\tau p}}. \end{aligned} \quad (54)$$

Noticing $\gamma \geq \zeta^{\frac{p}{2}} \alpha^{\frac{1}{p}} = (\frac{\ell_2 \alpha}{2})^{\frac{1}{p}}$, we can ensure $\frac{\ell_2 \alpha / 2}{(k+\gamma)^p} \leq 1$ for all k . Invoking Theorem 24 (a) for (54) by setting $q = (\tau - 1)p$, $c = \frac{\ell_2 \alpha}{2}$, $d = \ell_3 \alpha^\tau$, and $\varsigma = 1$, we can infer

$$y_{k+1} \leq \frac{4\ell_3}{\ell_2} \alpha^{\tau-1} (k+1+\gamma)^{-(\tau-1)p} + y_0 \exp\left(-\frac{\alpha \ell_2 (k+1)}{2(k+1+\gamma)^p}\right)$$

provided that $\gamma \geq \left[\frac{(1+\varsigma)q}{c}\right]^{\frac{1}{1-p}} = \left[\frac{4(\tau-1)p}{\alpha\ell_2}\right]^{\frac{1}{1-p}} = \left(\frac{2u_1}{\zeta\alpha^{1/e}}\right)^{\frac{1}{1-p/e}}$ (which holds by assumption).

Sub-Case 1-2: $p = 1$. In this sub-case, we have $u_2 = \tau - 1$ and the recursion (43) reduces to

$$y_{k+1} \leq \left[1 + \frac{\ell_1\alpha^\tau}{(k+\gamma)^\tau} - \frac{\ell_2\alpha}{k+\gamma}\right] y_k + \frac{\alpha^\tau\ell_3}{(k+\gamma)^\tau}. \quad (55)$$

As stated in the assumptions in part (b), let α and γ be given such that $\alpha \geq \left(\frac{2u_2}{\zeta}\right)^e = \frac{4(\tau-1)}{\ell_2}$ and $\gamma \geq \max\left\{\left(\frac{\ell_1}{\zeta}\right)^{\frac{1}{u_2}}\alpha^{\frac{1}{e}}, \zeta\alpha^{\frac{1}{e}}\right\} = \max\left\{\alpha\left(\frac{2\ell_1}{\ell_2}\right)^{\frac{1}{\tau-1}}, \frac{\ell_2\alpha}{2}\right\}$. This implies

$$\frac{\ell_1\alpha^\tau}{(k+\gamma)^\tau} - \frac{\ell_2\alpha}{k+\gamma} \leq \left(\frac{\ell_1\alpha^{\tau-1}}{\ell_2\gamma^{\tau-1}} - 1\right) \frac{\ell_2\alpha}{k+\gamma} \leq -\frac{\ell_2\alpha}{2(k+\gamma)}, \quad \frac{\ell_2\alpha}{2(k+\gamma)} \leq 1,$$

and $\frac{\ell_2\alpha}{2} \geq 2(\tau-1)$ for all $k \geq 0$. Then, (55) further simplifies to

$$y_{k+1} \leq \left[1 - \frac{\ell_2\alpha}{2(k+\gamma)}\right] y_k + \frac{\alpha^\tau\ell_3}{(k+\gamma)^\tau}.$$

Applying Theorem 24 (b) (for the case $c > q$) with $c = \frac{\ell_2\alpha}{2}$, $d = \alpha^\tau\ell_3$, $q = \tau - 1$ and using the inequality $c - q \geq \frac{c}{2}$, we have

$$y_{k+1} \leq \frac{4\ell_3}{\ell_2}\alpha^{\tau-1}(k+1+\gamma)^{-(\tau-1)} + y_0(\gamma^{-1}(k+1+\gamma))^{-\frac{\ell_2\alpha}{2}}.$$

Case II: $\theta \neq \frac{1}{2}$. As a starting point, we consider the convex function $x \mapsto h_\theta(x) := x^{2\theta}$. Then, applying the convexity of h_θ , it follows

$$h_\theta(y) \geq h_\theta(x) + 2\theta x^{2\theta-1}(y-x) \quad \forall x, y \geq 0. \quad (56)$$

Introducing an auxiliary sequence $\{u_k\}_k$ (which we still need to determine) and setting $y = y_k$ and $x = u_k$ in (56), we can relax (43) to

$$\begin{aligned} y_{k+1} &\leq (1 + \ell_1\alpha_k^\tau)y_k - \ell_2\alpha_k h_\theta(y_k) + \ell_3\alpha_k^\tau \\ &\leq [1 + \ell_1\alpha_k^\tau - 2\theta\ell_2\alpha_k u_k^{2\theta-1}]y_k + [(2\theta - 1)\ell_2 u_k^{2\theta}\alpha_k + \ell_3\alpha_k^\tau]. \end{aligned} \quad (57)$$

We will now discuss (57) and the explicit definition of $\{u_k\}_k$ in several sub-cases depending on the choice of p .

Sub-Case 2-1: $p \in (0, 1)$. We use the ansatz $u_k = \nu(k+\gamma)^{-u}$, where u and ν are to be determined. Intuitively, the term $2\theta\ell_2\alpha_k u_k^{2\theta-1}$ should dominate $\ell_1\alpha_k^\tau$ and it should not decay faster than $\frac{1}{k}$ (to ensure applicability of Theorem 24), i.e., u should be chosen such that

$$\alpha_k u_k^{2\theta-1} \gtrsim \alpha_k^\tau \quad \text{and} \quad \alpha_k u_k^{2\theta-1} \gtrsim \frac{1}{k}. \quad (58)$$

In terms of the exponents τ , s , and θ , (58) is equivalent to $u \leq \bar{u} := \min\left\{\frac{p(\tau-1)}{2\theta-1}, \frac{1-p}{2\theta-1}\right\}$. According to Theorem 24, the estimated convergence rate is $O(k^{-\min\{u, p(\tau-1) - (2\theta-1)u\}})$. Note that

$$\min\{u, p(\tau-1) - (2\theta-1)u\} = \begin{cases} u & \text{if } u \leq \frac{p(\tau-1)}{2\theta}, \\ p(\tau-1) - (2\theta-1)u & \text{if } u > \frac{p(\tau-1)}{2\theta}, \end{cases}$$

then, $\min\{u, p(\tau - 1) - (2\theta - 1)u\} \leq \frac{p(\tau-1)}{2\theta}$. Moreover, due to $\frac{p(\tau-1)}{2\theta} \leq \frac{p(\tau-1)}{2\theta-1}$, it holds that $\frac{p(\tau-1)}{2\theta} \leq \bar{u}$ if and only if $s \leq \frac{2\theta}{(2\theta-1)\tau+1} = \varrho$. This shows that

$$\operatorname{argmax}_{u \in (0, \bar{u})} \min\{u, p(\tau - 1) - (2\theta - 1)u\} = \begin{cases} \frac{p(\tau-1)}{2\theta} & \text{if } p \in (0, \varrho], \\ \frac{1-p}{2\theta-1} & \text{if } p \in (\varrho, 1). \end{cases}$$

Hence, depending on p different optimal rates can be achieved and we continue to split this case into additional sub-cases.

- Case a): $p \in (0, \varrho)$. (The conditions in part (a) hold). We can set $u = u_1 := \frac{p(\tau-1)}{2\theta}$ to obtain the optimal rate. In this case, (57) can be expressed as

$$y_{k+1} \leq \left[1 + \frac{\ell_1 \alpha^\tau}{(k + \gamma)^{\tau p}} - \frac{2\theta \ell_2 \nu^{2\theta-1} \alpha}{(k + \gamma)^{p/e}} \right] y_k + \frac{(2\theta - 1)\ell_2 \nu^{2\theta} \alpha + \ell_3 \alpha^\tau}{(k + \gamma)^{\tau p}}.$$

Assuming $\gamma \geq \left(\frac{\ell_1 \alpha^{\tau-1}}{\theta \ell_2 \nu^{2\theta-1}}\right)^{\frac{1}{u_1}}$ and using $\tau p - \frac{p}{\varrho} = \frac{p(\tau-1)}{2\theta} = u_1$, this yields

$$\begin{aligned} y_{k+1} &\leq \left[1 + \left(\frac{\ell_1 \alpha^{\tau-1}}{2\theta \ell_2 \nu^{2\theta-1} (k + \gamma)^{u_1}} - 1 \right) \frac{2\theta \ell_2 \nu^{2\theta-1} \alpha}{(k + \gamma)^{p/e}} \right] y_k + \frac{(2\theta - 1)\ell_2 \nu^{2\theta} \alpha + \ell_3 \alpha^\tau}{(k + \gamma)^{\tau p}} \\ &\leq \left[1 - \frac{\theta \ell_2 \nu^{2\theta-1} \alpha}{(k + \gamma)^{p/e}} \right] y_k + \frac{(2\theta - 1)\ell_2 \nu^{2\theta} \alpha + \ell_3 \alpha^\tau}{(k + \gamma)^{\tau p}}. \end{aligned} \quad (59)$$

To ensure $\theta \ell_2 \nu^{2\theta-1} \alpha (k + \gamma)^{-(p/\varrho)} \leq 1$ for all $k \geq 0$, we further assume $\gamma \geq (\alpha \theta \ell_2 \nu^{2\theta-1})^{\frac{\varrho}{p}}$. Noticing $p/\varrho < 1$ and invoking Theorem 24 (a) for (59) with $p \leftarrow \frac{p}{\varrho}$, $q = u_1$, $c = \theta \ell_2 \nu^{2\theta-1} \alpha$, $d = (2\theta - 1)\ell_2 \nu^{2\theta} \alpha + \ell_3 \alpha^\tau$, and requiring $\gamma \geq \left[\frac{2u_1}{\alpha \theta \ell_2 \nu^{2\theta-1}}\right]^{\frac{1}{1-p/\varrho}}$, we obtain

$$y_{k+1} \leq \frac{2[(2\theta - 1)\ell_2 \nu^{2\theta} + \ell_3 \alpha^{\tau-1}]}{\theta \ell_2 \nu^{2\theta-1}} (k + 1 + \gamma)^{-u_1} + y_0 \exp\left(-\frac{\alpha \theta \ell_2 \nu^{2\theta-1} (k + 1)}{(k + 1 + \gamma)^{p/e}}\right).$$

We can choose $\nu = \left(\frac{\ell_3 \alpha^{\tau-1}}{\ell_2}\right)^{\frac{1}{2\theta}}$ to make the coefficient of the leading term to be optimal. Plugging this ν into the different conditions on γ and in the previous estimate, it follows

$$\gamma \geq \max\left\{\left(\frac{\ell_1}{\zeta}\right)^{\frac{1}{u_1}} \alpha^{\frac{1}{p}}, \zeta^{\frac{\varrho}{p}} \alpha^{\frac{1}{p}}, \left(\frac{2u_1}{\alpha^{1/\varrho} \zeta}\right)^{\frac{1}{1-p/\varrho}}\right\} \quad \text{where } \zeta = \theta \ell_2^{\frac{1}{2\theta}} \ell_3^{\frac{2\theta-1}{2\theta}}, \quad (60)$$

and

$$y_{k+1} \leq 4 \left(\frac{\ell_3}{\ell_2}\right)^{\frac{1}{2\theta}} \alpha^{\frac{\tau-1}{2\theta}} (k + 1 + \gamma)^{-u_1} + y_0 \exp\left(-\frac{\alpha^{1/\varrho} \zeta (k + 1)}{(k + 1 + \gamma)^{p/e}}\right).$$

The imposed conditions on γ in (60) hold due to the assumptions in part (a).

- Case b): $p = \varrho$. (The conditions in part (b) hold). We can set $u = u_2 := \frac{\varrho(\tau-1)}{2\theta} = \frac{\tau-1}{(2\theta-1)\tau+1}$ to achieve the optimal rate. In this case, (57) reduces to

$$y_{k+1} \leq \left[1 + \frac{\ell_1 \alpha^\tau}{(k + \gamma)^{\tau \varrho}} - \frac{2\theta \ell_2 \nu^{2\theta-1} \alpha}{k + \gamma} \right] y_k + \frac{(2\theta - 1)\ell_2 \nu^{2\theta} \alpha + \ell_3 \alpha^\tau}{(k + \gamma)^{\tau \varrho}}.$$

As before, if $\gamma \geq \{(\frac{\ell_1 \alpha^{\tau-1}}{\theta \ell_2 \nu^{2\theta-1}})^{\frac{1}{u_2}}, \alpha \theta \ell_2 \nu^{2\theta-1}\}$ and using $\tau \varrho - 1 = u_2$, we can infer

$$y_{k+1} \leq \left[1 - \frac{\theta \ell_2 \nu^{2\theta-1} \alpha}{k + \gamma} \right] y_k + \frac{(2\theta - 1) \ell_2 \nu^{2\theta} \alpha + \ell_3 \alpha^\tau}{(k + \gamma)^{\tau \varrho}} \quad (61)$$

and $\theta \ell_2 \nu^{2\theta-1} \alpha (k + \gamma)^{-1} \leq 1$ for all k . Invoking Theorem 24 (b) with $p = 1$, $q = \tau \varrho - 1 = u_2$, $c = \theta \ell_2 \nu^{2\theta-1} \alpha$, $d = (2\theta - 1) \ell_2 \nu^{2\theta} \alpha + \ell_3 \alpha^\tau$, and letting $\theta \ell_2 \nu^{2\theta-1} \alpha \geq 2u_2$ to guarantee $c - q \geq \frac{c}{2}$, we have

$$y_{k+1} \leq \frac{2[(2\theta - 1) \ell_2 \nu^{2\theta} + \ell_3 \alpha^{\tau-1}]}{\theta \ell_2 \nu^{2\theta-1}} (k + 1 + \gamma)^{-u_2} + y_0 (\gamma^{-1} (k + 1 + \gamma))^{-\theta \ell_2 \nu^{2\theta-1} \alpha}$$

Similar to the previous case, we can select $\nu = (\frac{\ell_3 \alpha^{\tau-1}}{\ell_2})^{\frac{1}{2\theta}}$ to optimize the coefficient of the leading term. Then, the conditions on α and γ reduce to $\alpha \geq (\frac{2u_2}{\zeta})^\varrho$, $\gamma \geq \{(\frac{\ell_1}{\zeta})^{\frac{1}{u_2}} \alpha^{\frac{1}{\varrho}}, \zeta \alpha^{\frac{1}{\varrho}}\}$ and we have

$$y_{k+1} \leq 4(\frac{\ell_3}{\ell_2})^{\frac{1}{2\theta}} \alpha^{\frac{\tau-1}{2\theta}} (k + 1 + \gamma)^{-\varrho} + y_0 (\gamma^{-1} (k + 1 + \gamma))^{-\zeta \alpha^{1/\varrho}}.$$

- Case c): $p \in (\varrho, 1)$. (This corresponds to part (c)). We can set $u = u_3 := \frac{1-p}{2\theta-1}$ to attain the optimal rate. Taking $\nu = 1$, (57) can be written as

$$y_{k+1} \leq \left[1 + \frac{\ell_1 \alpha^\tau}{(k + \gamma)^{\tau p}} - \frac{2\theta \ell_2 \alpha}{k + \gamma} \right] y_k + \left[\frac{(2\theta - 1) \ell_2 \alpha}{(k + \gamma)^{1+u_3}} + \frac{\ell_3 \alpha^\tau}{(k + \gamma)^{\tau p}} \right].$$

If $\alpha \geq \frac{2u_3}{\theta \ell_2}$ and $\gamma \geq \max\{(\frac{\alpha^{\tau-1} \ell_1}{\theta \ell_2})^{\frac{1}{\tau p-1}}, (\frac{\alpha^{\tau-1} \ell_3}{\ell_2})^{\frac{1}{\tau p-u_3-1}}, \alpha \theta \ell_2\}$, it follows

$$\frac{\ell_1 \alpha^\tau}{(k + \gamma)^{\tau p}} - \frac{2\theta \ell_2 \alpha}{k + \gamma} \leq -\frac{\theta \ell_2 \alpha}{k + \gamma}, \quad \frac{\ell_3 \alpha^\tau}{(k + \gamma)^{\tau p}} \leq \frac{\ell_2 \alpha}{(k + \gamma)^{1+u_3}}, \quad \frac{\theta \ell_2 \alpha}{k + \gamma} \leq 1,$$

$\alpha \theta \ell_2 \geq 2u_3$ for all $k \geq 0$. Consequently, we can infer

$$y_{k+1} \leq \left[1 - \frac{\theta \ell_2 \alpha}{k + \gamma} \right] y_k + \frac{2\theta \ell_2 \alpha}{(k + \gamma)^{1+u_3}}.$$

Thus, applying Theorem 24 (b) with $c = \theta \ell_2 \alpha$, $d = 2\theta \ell_2 \alpha$, $q = u_3$ and utilizing $c - q \geq \frac{c}{2}$, we obtain

$$y_{k+1} \leq 4(k + 1 + \gamma)^{-u_3} + y_0 [\gamma^{-1} (k + 1 + \gamma)]^{-\theta \ell_2 \alpha}.$$

Sub-Case 2-2: $p = 1$. Here, we introduce the auxiliary sequence $u_k := \log(k + \gamma)^{-\frac{1}{2\theta-1}}$. Mimicking the earlier cases, there exists $\gamma_0 > 0$, which solely depends on $\alpha, \ell_1, \ell_2, \ell_3, \tau$, such that for any $\gamma \geq \gamma_0$, we have $\gamma \log(\gamma) \geq \alpha \theta \ell_2$ and (57) can be simplified to

$$y_{k+1} \leq \left[1 - \frac{\alpha \theta \ell_2}{(k + \gamma) \log(k + \gamma)} \right] y_k + \frac{2\alpha \theta \ell_2}{(k + \gamma) \log(k + \gamma)^{\frac{2\theta}{2\theta-1}}}.$$

(We will omit an explicit construction of such γ_0 here). Let us now define $s(x) := \frac{1}{\alpha\theta\ell_2}x \log(x)$, $t(x) := \frac{1}{2\alpha\theta\ell_2}x \log(x)^{\frac{2\theta}{2\theta-1}}$, $r(x) := 2 \log(x)^{-\frac{1}{2\theta-1}}$, and $b_k = k + \gamma$. Then, it holds that

$$\begin{aligned} u(x) &:= s'(x) - r(x)t'(x) \\ &= \frac{1}{\alpha\theta\ell_2}(1 + \log(x)) - \frac{1}{\alpha\theta\ell_2} \cdot \frac{1}{\log(x)^{\frac{1}{2\theta-1}}} \left(\frac{2\theta}{2\theta-1} \log(x)^{\frac{1}{2\theta-1}} + \log(x)^{\frac{2\theta}{2\theta-1}} \right) \\ &= -\frac{1}{\alpha\theta(2\theta-1)\ell_2}. \end{aligned}$$

Thus, if $\alpha \geq \frac{2}{\theta(2\theta-1)\ell_2}$, it follows $(b_{k+1} - b_k)u(b_k) \geq -\frac{1}{2}$. This the rate function r is clearly convex on $I = [\gamma, \infty)$, all assumptions in Theorem 1 (b) are satisfied with $\lambda_k \equiv 2$. Hence, applying Theorem 1 (b), we can infer

$$\begin{aligned} y_{k+1} &\leq 4 \log(k+1+\gamma)^{-\frac{1}{2\theta-1}} + y_0 \exp\left(-\sum_{i=0}^k \frac{\alpha\theta\ell_2}{(i+\gamma)\log(i+\gamma)}\right) \\ &\leq 4 \log(k+1+\gamma)^{-\frac{1}{2\theta-1}} + y_0 \left(\frac{\log(k+1+\gamma)}{\log(\gamma)}\right)^{-\alpha\theta\ell_2}, \end{aligned}$$

where we utilize the integral test $\sum_{i=0}^k \frac{1}{(k+\gamma)\log(k+\gamma)} \geq \int_0^{k+1} \frac{1}{(x+\gamma)\log(x+\gamma)} dx = \log(\log(k+1+\gamma)) - \log(\log(\gamma))$ in the last line. \blacksquare

E.2 Results for Polynomial Step Sizes

Proof of Theorem 20 The condition $\gamma \geq (\mathsf{L}\alpha)^{1/\varrho_1}$ ensures $\alpha_k \leq \frac{1}{\mathsf{L}}$ for all k and hence, the descent property (15) is applicable. The first convergence rate (44) in Theorem 20 can be obtained by applying Lemma 19 (b) to (15) with $\ell_1 = \frac{\mathsf{A}\mathsf{L}}{2}$, $\ell_2 = \mu$, $\ell_3 = \frac{\mathsf{L}\sigma^2}{2}$, and $\tau = 2$.

To establish (45), we substitute $\alpha = (\frac{2\omega_1}{\zeta_1})^{\varrho_1}$ in $\gamma = \max\left\{\left(\frac{\mathsf{A}\mathsf{L}}{2\zeta_1}\right)^{\frac{1}{\omega_1}} \alpha^{\frac{1}{\varrho_1}}, \zeta_1 \alpha^{\frac{1}{\varrho_1}}, (\mathsf{L}\alpha)^{\frac{1}{\varrho_1}}\right\}$ and (44). Then, using $\frac{\varrho_1}{\omega_1} = 2 - \frac{1}{2\theta}$, $\frac{1}{\omega_1} = 4\theta - 1$, and $\zeta_1 = \theta\mu^{\frac{1}{2\theta}} \left(\frac{\mathsf{L}\sigma^2}{2}\right)^{1-\frac{1}{2\theta}}$, we have

$$\begin{aligned} \gamma &= 2\omega_1 \max\left\{\left(\frac{\mathsf{A}\mathsf{L}}{2\zeta_1}\right)^{\frac{1}{\omega_1}} \frac{1}{\zeta_1}, 1, \frac{\mathsf{L}^{1/\varrho_1}}{\zeta_1}\right\} = 2\omega_1 \max\left\{\frac{(\mathsf{A}\mathsf{L})^{4\theta-1}}{2^{4\theta-1}\zeta_1^{4\theta}}, 1, \frac{2^{1-\frac{1}{2\theta}}\mathsf{L}}{\theta\mu^{\frac{1}{2\theta}}\sigma^{2-\frac{1}{\theta}}}\right\} \\ &= 2\omega_1 \max\{1, C_1, C_2\} \end{aligned}$$

and $\left(\frac{\mathsf{L}\sigma^2}{2\mu}\right)^{\frac{1}{2\theta}} \alpha^{\frac{1}{2\theta}} = \left(\frac{\mathsf{L}\sigma^2}{2\mu}\right)^{\frac{1}{2\theta}} \left(\frac{2\omega_1}{\zeta_1}\right)^{\omega_1} = \left(\frac{\omega_1}{\theta}\right)^{\omega_1} \left(\frac{\mathsf{L}\sigma^2}{\mu^2}\right)^{\omega_1}$. \blacksquare

Proof of Theorem 21 The inequality $\gamma \geq (2\mathsf{L}\alpha)^{\frac{1}{\varrho_2}}$ ensures that $\alpha_k \leq \frac{1}{2\mathsf{L}}$, which means that (16) is applicable. The first inequality (46) can be obtained by applying Lemma 19 to (16) with $\ell_1 = \frac{\mathsf{A}\mathsf{L}^2}{2\mathsf{N}}$, $\ell_2 = \frac{\mu}{2}$, and $\ell_3 = \frac{\mathsf{L}^2\sigma^2}{2\mathsf{N}}$. To establish (47), we substitute

$$\begin{aligned} \gamma &= \max\left\{\left(\frac{\mathsf{A}\mathsf{L}^2}{2\mathsf{N}\zeta_2}\right)^{\frac{1}{\omega_2}} \alpha^{\frac{1}{\varrho_2}}, \zeta_2 \alpha^{\frac{1}{\varrho_2}}, (2\mathsf{L}\alpha)^{\frac{1}{\varrho_2}}\right\} = 2\omega_2(1 + \log(\mathsf{N})) \max\left\{\frac{C_3}{\sqrt{\mathsf{N}}}, C_4\mathsf{N}^{1-\frac{1}{2\theta}}, 1\right\} \\ \alpha &= \left[\frac{2\omega_2(1+\log(\mathsf{N}))}{\zeta_2}\right]^{\varrho_2} \quad \text{and} \quad K \geq 2\gamma \end{aligned}$$

into (46), then it holds that

$$\begin{aligned}
y_K &\leq 4 \left(\frac{\ell_3}{\ell_2}\right)^{\frac{1}{2\theta}} \left(\frac{2\omega_2}{\zeta}\right)^{\omega_2} \left[\frac{1 + \log(N)}{K + \gamma}\right]^{\omega_2} + y_0 [\gamma^{-1}(K + \gamma)]^{-2\omega_2(1 + \log(N))} \\
&\leq 4 \left(\frac{4\omega_2}{\theta}\right)^{\omega_2} \left(\frac{L\sigma}{\mu^{3/2}}\right)^{\omega_2} \left[\frac{1 + \log(N)}{\sqrt{N}(K + \gamma)}\right]^{\omega_2} + y_0 \left[\frac{\gamma}{K + \gamma}\right]^{2\omega_2} [\gamma^{-1}(K + \gamma)]^{\frac{-2\omega_2 \log(N)}{\log[(K + \gamma)/\gamma]}} \\
&= 4 \left(\frac{4\omega_2}{\theta}\right)^{\omega_2} \left(\frac{L\sigma}{\mu^{3/2}}\right)^{\omega_2} \left[\frac{1 + \log(N)}{\sqrt{N}(K + \gamma)}\right]^{\omega_2} + y_0 \left[\frac{2\omega_2(1 + \log(N)) \max\{\frac{C_3}{\sqrt{N}}, C_4 N^{1 - \frac{1}{2\theta}}, 1\}}{N(K + \gamma)}\right]^{2\omega_2}.
\end{aligned}$$

This completes the proof. ■

References

- M. Abadi, A. Agarwal, P. Barham, E. Brevdo, Z. Chen, C. Citro, G. S. Corrado, A. Davis, et al. TensorFlow: Large-scale machine learning on heterogeneous systems, 2015.
- A. Agarwal, S. M. Kakade, J. D. Lee, and G. Mahajan. On the theory of policy gradient methods: Optimality, approximation, and distribution shift. *The Journal of Machine Learning Research*, 22(1):4431–4506, 2021.
- K. Ahn, C. Yun, and S. Sra. SGD with shuffling: optimal rates without component convexity and large epoch requirements. *Advances in Neural Information Processing Systems*, 33: 17526–17535, 2020.
- J. Bolte, S. Sabach, and M. Teboulle. Proximal alternating linearized minimization for nonconvex and nonsmooth problems. *Mathematical Programming*, 146(1-2, Ser. A):459–494, 2014.
- L. Bottou, F. E. Curtis, and J. Nocedal. Optimization methods for large-scale machine learning. *SIAM Review*, 60(2):223–311, 2018.
- K. L. Chung. On a stochastic approximation method. *The Annals of Mathematical Statistics*, pages 463–483, 1954.
- C. Derman. An application of chung’s lemma to the Kiefer-Wolfowitz stochastic approximation procedure. *The Annals of Mathematical Statistics*, pages 532–536, 1956.
- V. Fabian. On asymptotic normality in stochastic approximation. *The Annals of Mathematical Statistics*, pages 1327–1332, 1968.
- I. Fatkhullin, J. Etesami, N. He, and N. Kiyavash. Sharp analysis of stochastic optimization under global Kurdyka-Łojasiewicz inequality. In *Advances in Neural Information Processing Systems*, volume 35, pages 15836–15848, 2022.
- X. Fontaine, V. De Bortoli, and A. Durmus. Convergence rates and approximation results for SGD and its continuous-time counterpart. In *Conference on Learning Theory*, pages 1965–2058. PMLR, 2021.

- R. Ge, S. M. Kakade, R. Kidambi, and P. Netrapalli. The step decay schedule: A near optimal, geometrically decaying learning rate procedure for least squares. *Advances in neural information processing systems*, 32, 2019.
- J.-L. Goffin. On convergence rates of subgradient optimization methods. *Mathematical programming*, 13:329–347, 1977.
- R. Gower, O. Sebbouh, and N. Loizou. SGD for structured nonconvex functions: Learning rates, minibatching and interpolation. In *International Conference on Artificial Intelligence and Statistics*, pages 1315–1323. PMLR, 2021.
- R. M. Gower, N. Loizou, X. Qian, A. Sailanbayev, E. Shulgin, and P. Richtárik. SGD: General analysis and improved rates. In *International conference on machine learning*, pages 5200–5209. PMLR, 2019.
- M. Gürbüzbalaban, A. Ozdaglar, and P. A. Parrilo. Convergence rate of incremental gradient and incremental Newton methods. *SIAM Journal on Optimization*, 29(4):2542–2565, 2019.
- M. Gürbüzbalaban, A. Ozdaglar, and P. Parrilo. Why random reshuffling beats stochastic gradient descent. *Mathematical Programming*, 186(1-2):49–84, 2021.
- J. Z. HaoChen and S. Sra. Random shuffling beats SGD after finite epochs. In *International Conference on Machine Learning*, pages 2624–2633, 2019.
- X. Huang, K. Yuan, X. Mao, and W. Yin. An improved analysis and rates for variance reduction under without-replacement sampling orders. *Advances in Neural Information Processing Systems*, 34, 2021.
- H. Karimi, J. Nutini, and M. Schmidt. Linear convergence of gradient and proximal-gradient methods under the Polyak-Lojasiewicz condition. In *Machine Learning and Knowledge Discovery in Databases: European Conference, ECML PKDD 2016, Riva del Garda, Italy, September 19-23, 2016, Proceedings, Part I 16*, pages 795–811. Springer, 2016.
- X. Li, Z. Zhuang, and F. Orabona. A second look at exponential and cosine step sizes: Simplicity, adaptivity, and performance. In *International Conference on Machine Learning*, pages 6553–6564. PMLR, 2021.
- X. Li, A. Milzarek, and J. Qiu. Convergence of random reshuffling under the Kurdyka–Lojasiewicz inequality. *SIAM Journal on Optimization*, 33(2):1092–1120, 2023.
- S. Lojasiewicz. Sur le problème de la division. *Studia Mathematica*, 18:87–136, 1959.
- S. Lojasiewicz. Une propriété topologique des sous-ensembles analytiques réels. *Les équations aux dérivées partielles*, 117:87–89, 1963.
- I. Loshchilov and F. Hutter. SGDR: Stochastic gradient descent with warm restarts. In *International Conference on Learning Representations*, 2016.

- J. Mei, C. Xiao, C. Szepesvari, and D. Schuurmans. On the global convergence rates of softmax policy gradient methods. In *International Conference on Machine Learning*, pages 6820–6829. PMLR, 2020.
- K. Mishchenko, A. Khaled, and P. Richtarik. Random reshuffling: Simple analysis with vast improvements. In *Advances in Neural Information Processing Systems*, pages 17309–17320.
- E. Moulines and F. Bach. Non-asymptotic analysis of stochastic approximation algorithms for machine learning. In *Advances in Neural Information Processing Systems*, volume 24. Curran Associates, Inc., 2011.
- D. Needell, R. Ward, and N. Srebro. Stochastic gradient descent, weighted sampling, and the randomized Kaczmarz algorithm. *Advances in neural information processing systems*, 27, 2014.
- A. Nemirovski, A. Juditsky, G. Lan, and A. Shapiro. Robust stochastic approximation approach to stochastic programming. *SIAM Journal on optimization*, 19(4):1574–1609, 2009.
- L. M. Nguyen, Q. Tran-Dinh, D. T. Phan, P. H. Nguyen, and M. van Dijk. A unified convergence analysis for shuffling-type gradient methods. *Journal of Machine Learning Research*, 2021.
- A. Paszke, S. Gross, F. Massa, A. Lerer, J. Bradbury, G. Chanan, T. Killeen, Z. Lin, N. Gimelshein, L. Antiga, et al. Pytorch: An imperative style, high-performance deep learning library. *Advances in neural information processing systems*, 32, 2019.
- B. T. Polyak. Gradient methods for the minimisation of functionals. *USSR Computational Mathematics and Mathematical Physics*, 3(4):864–878, 1963.
- B. T. Polyak. *Introduction to optimization*. Translations Series in Mathematics and Engineering. Optimization Software, Inc., Publications Division, New York, 1987. Translated from the Russian, With a foreword by Dimitri P. Bertsekas.
- H. Robbins and S. Monro. A stochastic approximation method. *The annals of mathematical statistics*, pages 400–407, 1951.
- J. Sacks. Asymptotic distribution of stochastic approximation procedures. *The Annals of Mathematical Statistics*, 29(2):373–405, 1958.
- O. Shamir. Without-replacement sampling for stochastic gradient methods. In *Advances in Neural Information Processing Systems*, pages 46–54, 2016.
- O. Shamir and T. Zhang. Stochastic gradient descent for non-smooth optimization: Convergence results and optimal averaging schemes. In *International conference on machine learning*, pages 71–79. PMLR, 2013.
- J. Sohl-Dickstein, B. Poole, and S. Ganguli. Fast large-scale optimization by unifying stochastic gradient and quasi-newton methods. In *International Conference on Machine Learning*, pages 604–612. PMLR, 2014.

- H. Touvron, L. Martin, K. Stone, P. Albert, A. Almahairi, Y. Babaei, N. Bashlykov, S. Batta, P. Bhargava, S. Bhosale, et al. Llama 2: Open foundation and fine-tuned chat models. *arXiv preprint arXiv:2307.09288*, 2023.
- T. H. Tran, L. M. Nguyen, and Q. Tran-Dinh. Sng: A shuffling gradient-based method with momentum. In *International Conference on Machine Learning*, pages 10379–10389. PMLR, 2021.
- X. Wang, S. Magnússon, and M. Johansson. On the convergence of step decay step-size for stochastic optimization. *Advances in Neural Information Processing Systems*, 34: 14226–14238, 2021.
- Y. Wang and S. Zou. Policy gradient method for robust reinforcement learning. In *International Conference on Machine Learning*, pages 23484–23526. PMLR, 2022.