

Relativistic Corrections to the CBF Effective Nuclear Hamiltonian

Andrea Sabatucci,^{1,2,*} Omar Benhar,² and Alessandro Lovato^{3,4}

¹*Dipartimento di Fisica, “Sapienza” University of Rome, 00185 Roma, Italy*

²*INFN, Sezione di Roma, 00185 Roma, Italy*

³*Physics Division, Argonne National Laboratory, Argonne, Illinois 60439, USA*

⁴*INFN, Trento Institute of Fundamental Physics and Applications, 38123 Trento, Italy*

We discuss the inclusion of relativistic boost corrections into the CBF effective nuclear Hamiltonian, derived from a realistic model of two- and three-nucleon interactions using the formalism of correlated basis functions and the cluster expansion technique. Different procedures to take into account the effects of boost interactions are compared on the basis of the ability to reproduce the nuclear matter equation of state obtained from accurate many-body calculations. The results of our study show that the repulsive contribution of the boost interaction significantly depends on the underlying model of the non relativistic potential. On the other hand, the dominant relativistic correction turns out to be the corresponding reduction of the strength of repulsive three-nucleon interactions, leading to a significant softening of the equation of state at supranuclear densities.

I. INTRODUCTION

According to the paradigm of many-body theory, all nuclear systems—from the deuteron to neutron stars—can be treated as collections of point-like protons and neutrons interacting via instantaneous potentials [1]. It should be kept in mind, however, that this description, while proving adequate to explain a wealth of nuclear properties, is inherently inconsistent with the requirement of causality, because it leads to predict a speed of sound larger than the speed of light in dense nuclear matter.

Pioneering studies of relativistic effects in both nuclear matter and the three-nucleon bound states were carried out by Coester *et al.* in the 1970s and 1980s; see Refs. [2, 3]. More recently, relativistic corrections to the properties of the three- and four-nucleon systems have been thoroughly analysed by the authors of Ref. [4] using a relativistic Hamiltonian—obtained from state-of-the-art phenomenological potentials following the procedure described in Ref. [5]—and Quantum Monte Carlo techniques. The results of this study indicate that, owing to the occurrence of large cancellations, the dominant relativistic effects, referred to as boost corrections, are driven by the total momentum of the interacting particles. Boost corrections to the ground-state energies turn out to be repulsive, and account for a sizeable fraction of the potential energy originating from irreducible repulsive interactions involving three nucleons.

Relativistic boost corrections to the energies of pure neutron matter (PNM) and isospin-symmetric nuclear matter (SNM), computed within the scheme described in Refs. [4, 5], are included in the equation of state (EOS) of charge-neutral β -stable matter derived by Akmal, Pandharipande and Ravenhall [6], widely employed in studies of neutron star properties. The results of Ref. [6]—obtained from accurate variational calculations

performed with a phenomenological nuclear Hamiltonian comprising the Argonne v_{18} (AV18) nucleon-nucleon (NN) potential [7] and the Urbana IX (UIX) three-nucleon (NNN) potential [8, 9]—show that the suppression of repulsive three-nucleon interactions associated with relativistic boost corrections leads to a conspicuous softening of the EOS at supranuclear density, which appreciably affects the mass and radius of stable neutron stars obtained from the solution of the equations of Tolman, Oppenheimer and Volkoff. The resulting maximum mass turns out to decrease from 2.3 to 2.2 M_{\odot} , with M_{\odot} being the solar mass, while the radius of a 1.4 M_{\odot} neutron star is reduced from 12.7 to 11.6 Km. The inclusion of boost corrections also alleviates the problem of violation of causality, pushing its occurrence towards higher density.

Over the past decade, the dynamical model underlying the work of Ref. [6] has been used, in conjunction with the formalism of Correlated Basis Function (CBF) [10] and the cluster expansion technique [11], to derive a density-dependent effective interaction suitable to carry out perturbative calculations of nuclear matter properties at both zero and non-zero temperature [12–15]. The resulting NN potential includes the contributions of two- and three-nucleon forces, and is properly renormalised to take into account the screening of non-perturbative interactions arising from strong short-range correlations among the nucleons.

The CBF effective Hamiltonian is designed to reproduce the EOSs of PNM and SNM obtained from accurate many-body calculations, carried out using a simplified version of the AV18+UIX Hamiltonian without taking into account the relativistic boost corrections considered in Ref [6]. In this article, we discuss an extension of the approach of Ref. [12] allowing to include these corrections, which are known to affect significantly the properties of nuclear matter in the high-density regime relevant to neutron stars.

The rest of this article is organised as follows. The phenomenological nuclear Hamiltonian providing the basis of the CBF effective interaction and the associated

* Present address: INFN, Sezione di Pisa, 56127 Pisa, Italy

relativistic boost corrections are discussed in Section II, while Section III is devoted to a short description of the procedure employed to derive the nuclear Hamiltonian of Ref. [12], which essentially amounts to a renormalisation of nuclear forces in matter. The impact of relativistic boost corrections on nuclear matter energy and the explicit expression of the underlying effective interaction are analysed in Sections IV and V, respectively. Finally, in Section VI we summarise our findings and state the conclusions.

II. DYNAMICAL MODEL

In this section, we outline the model of nuclear dynamics underlying our approach, as well as the formalism employed to include the contribution of relativistic boost interactions.

A. Nuclear Hamiltonian

In non relativistic many-body theory, nuclear dynamics are described by the Hamiltonian

$$H_{\text{NR}} = \sum_{i=1}^A \frac{\mathbf{p}_i^2}{2m} + \sum_{j>i=1}^A v_{ij} + \sum_{k>j>i=1}^A V_{ijk}, \quad (1)$$

where A is the number of particles, \mathbf{p}_i and m denote the momentum of the i -th nucleon and its mass, and the potentials v_{ij} and V_{ijk} describe two- and three-nucleon interactions, respectively.

Phenomenological nucleon-nucleon (NN) potentials are determined by fitting the observed properties of the two-nucleon system—including the deuteron binding energy, magnetic moment and electric quadrupole moment, as well as the data obtained from the measured NN scattering cross sections—and reduce to Yukawa's one-pion-exchange potential at large distance. They are usually written in the form

$$v_{ij} = \sum_p v^p(r_{ij}) O_{ij}^p, \quad (2)$$

where the functions v^p only depend upon the distance between the interacting particles, $r_{ij} = |\mathbf{r}_i - \mathbf{r}_j|$. The operators O_{ij}^p with $p = 1, \dots, 6$ account for the strong spin-isospin dependence of nuclear forces, as well as for the occurrence of non-central interactions. They are defined as

$$O_{ij}^{p \leq 6} = [1, (\boldsymbol{\sigma}_i \cdot \boldsymbol{\sigma}_j), S_{ij}] \otimes [1, (\boldsymbol{\tau}_i \cdot \boldsymbol{\tau}_j)], \quad (3)$$

where $\boldsymbol{\sigma}_i$ and $\boldsymbol{\tau}_i$ are Pauli matrices acting in spin and isospin space, respectively, and

$$S_{ij} = \frac{3}{r_{ij}^2} (\boldsymbol{\sigma}_i \cdot \mathbf{r}_{ij}) (\boldsymbol{\sigma}_j \cdot \mathbf{r}_{ij}) - (\boldsymbol{\sigma}_i \cdot \boldsymbol{\sigma}_j). \quad (4)$$

State-of-the-art models, such as the Argonne v_{18} (AV18) potential [7], include twelve additional terms. The contributions corresponding to $p = 7, \dots, 14$ are associated with non-static components of the potential, including spin-orbit and other angular momentum-dependent terms, while those corresponding to $p = 15, \dots, 18$ take into account small violations of charge symmetry and charge independence.

Three-body forces are long known to be required to model the interactions of extended composite bodies, such as protons and neutrons, without considering their internal structure explicitly; see, e.g., Ref. [16]. In nuclear many-body theory, the inclusion of *irreducible* NNN forces, described by the potential V_{ijk} , is needed to explain both the observed binding energies of the three-nucleon systems and saturation—that is, the occurrence of a minimum of the energy per particle $E(\rho)/A$ at non-vanishing density $\rho_0 = 0.16 \text{ fm}^{-3}$ —in SNM.

Phenomenological models of the NNN force such as the UIX potential [8, 9] are written in the form

$$V_{ijk} = V_{ijk}^{2\pi} + V_{ijk}^R, \quad (5)$$

where $V_{ijk}^{2\pi}$ is the attractive Fujita-Miyazawa potential [17], describing two-pion-exchange processes associated with the appearance of a Δ resonance in the intermediate state, while V_{ijk}^R is a purely phenomenological repulsive term. The UIX model involves two parameters, determined from fits to the empirical data performed using the AV18+UIX nuclear Hamiltonian. The strength of the two-pion exchange contribution is fixed in such a way as to reproduce the observed ground-state energies of ${}^3\text{He}$ and ${}^4\text{He}$, while that of the isoscalar repulsive contribution is adjusted to obtain the saturation density of SNM inferred from extrapolation of nuclear data.

The non relativistic Hamiltonian employed to obtain the effective interaction discussed in this work comprises the Argonne v'_6 (AV6P) NN potential [18], determined *projecting* the full AV18 potential onto the basis of the six operators appearing in the expression of the one-pion-exchange potential, defined by Eqs.(3) and (4), and the UIX NNN potential. The AV6P potential predicts the binding energy and electric quadrupole moment of the deuteron with accuracy of 1%, and 4%, respectively, and provides an excellent fit of the elastic NN scattering phase shifts in the ${}^1\text{S}_0$ channel[‡]—which is dominant in pure neutron matter—up to laboratory energies ~ 600 MeV, well above the pion production threshold.

It must be pointed out that the procedure employed to obtain the AV6P potential effectively takes into account interactions described by components of the AV18 potential having $p > 6$. This feature is clearly illustrated

[‡] We will adopt the spectroscopic notation, according to which the two-nucleon states labelled ${}^1\text{S}_0$ and ${}^1\text{P}_1$ correspond to spin $S = 0$ and orbital angular momentum $\ell = 0$ and 1, respectively.

in Fig. 1, providing a comparison between the neutron-proton scattering phase shifts in the 1S_0 and 1P_1 channels computed using the AV6P potential and those obtained from the AV18 potential neglecting the terms $v^{p>6}$. It appears that the AV6P model provides a description of the data comparable to that obtained from the full AV18 potential in both channels. On the other hand, the truncated AV18 potential, while being capable to predict the 1S_0 phase shifts, conspicuously fails to reproduce the data in the 1P_1 channel at energies larger than ~ 100 MeV. This pattern suggests that the contributions of angular




FIG. 1. Proton-neutron scattering phase shifts in the 1S_0 (A) and 1P_1 (B) channels, plotted as a function of the kinetic energy of the beam particle in the laboratory frame. The results obtained using the AV6P and the truncated AV18 potentials are represented by solid and dashed lines, respectively. The triangles correspond to the full AV18 potential. The data are taken from Refs. [19, 20] (circles) and [21] (squares).

momentum-dependent interactions, which turn out to be significant, are, in fact, largely included in the projected AV6P potential.

The results of a study performed using accurate many-body techniques also show that the EOSs of PNM com-

puted using the AV6P potential and the full AV18 potential remarkably agree with one another. At saturation density, the discrepancy between the energies per particle obtained from Auxiliary Field Diffusion Monte Carlo (AFDMC) calculations using the two potential models turns out to be ~ 10 -15% [22].

B. Relativistic corrections

The relativistic description of nuclear systems is a long-standing challenge, which has been confronted within different theoretical approaches. Here, we outline the formalism originally developed by Krajcik and Foldy [23], and further elaborated by Friar [24].

The authors of Ref. [23] derived the Relativistic Hamiltonian of a system with fixed number of particles from a covariant approach based on the well established conceptual framework of non-relativistic quantum mechanics. In this context, *relativistic covariance* refers to the requirement that the descriptions of physical phenomena observed in different inertial frames can be obtained from one another by performing a Poincaré transformation. Such a requirement entails the definition of a representation of the Poincaré group in the Hilbert space associated with the observed system, and the derivation of the relativistic Hamiltonian from the commutation rules of the Poincaré algebra. A clear and exhaustive discussion of the assumptions underlying this approach can be found in Ref. [25].

The relativistic Hamiltonian of an interacting many-particle system, H_R , is defined as the sum of the relativistic single-particle kinetic energies, the interaction potentials and the associated relativistic boost corrections. The explicit expression comprising interactions involving up to three particles can be written in the form

$$H_R = \sum_{i=1}^A \sqrt{m_i^2 + \mathbf{p}_i^2} + \sum_{j>i=1}^A [\tilde{v}_{ij} + \delta v_{ij}(\mathbf{P}_{ij})] + \sum_{k>j>i=1}^A [\tilde{V}_{ijk} + \delta V_{ijk}(\mathbf{P}_{ijk})], \quad (6)$$

to be compared to Eq. (1). In the above equation, \tilde{v}_{ij} and \tilde{V}_{ijk} denote the two- and three-body potentials in the rest frames of the interacting nucleons, defined by the conditions

$$\mathbf{P}_{ij} = \mathbf{p}_i + \mathbf{p}_j = 0, \quad (7)$$

and

$$\mathbf{P}_{ijk} = \mathbf{p}_i + \mathbf{p}_j + \mathbf{p}_k = 0, \quad (8)$$

with \mathbf{p}_i , \mathbf{p}_j and \mathbf{p}_k being the nucleon momenta. The boost corrections δv_{ij} and δV_{ijk} originate from the motion of the two- and three-nucleon center of mass in the

rest frame of the A -nucleon system. They satisfy the obvious requirements

$$\delta v_{ij}(\mathbf{P}_{ij} = 0) = \delta V_{ijk}(\mathbf{P}_{ijk} = 0) = 0. \quad (9)$$

Because, in principle, \tilde{v}_{ij} and the non relativistic v_{ij} of Eq. (1) can be both obtained from a fit to experimental data [26], one can reasonably assume that relativistic effects taken into account using \tilde{v}_{ij} in conjunction with square root kinetic energies are, in fact, implicitly included in v_{ij} .

In the 1970s, Krajcik and Foldy [23] and Friar [24] obtained an expression of the boost correction $\delta v(\mathbf{P})^\ddagger$ from an expansion in powers of $\mathbf{P}^2/4m^2$. On the other hand, relativistic corrections to the NNN potential, whose contribution to the energy is known to be much smaller than the one arising from NN interactions, have been generally neglected [5, 26]. The present discussion will be limited to the calculation of the leading order contribution to $\delta v(\mathbf{P})$ within the scheme of Forest *et al.* [5]. This approximation appears to be amply justified by the results of numerical calculations, showing that the probability of \mathbf{P}^2 being larger than m^2 is, in fact, quite small; see, e.g., Ref. [26].

The starting point is the decomposition

$$H = H_0 + H_I, \quad (10)$$

where H_0 is the Hamiltonian describing two non interacting nucleons and H_I , containing all interaction terms, can be conveniently written in the form

$$H_I = v + \delta v + \dots. \quad (11)$$

The potential v , independent of \mathbf{P} , can be assumed to be of order $1/m$, because in nuclear systems interaction energy and non-relativistic kinetic energy are known to be of comparable magnitude. The correction δv is of order $1/m^3$, and the ellipsis represents terms of order $1/m^4$ or higher. The presence of H_I entails the appearance of a corresponding term in the boost generator, that can be written in the form

$$\mathbf{K} = \mathbf{K}_0 + \mathbf{K}_I. \quad (12)$$

The leading contributions to H_I and \mathbf{K}_I have been derived in Refs. [23, 24]. Substitution of the resulting expressions into the relevant commutation relations of the Poincaré algebra yields the boost interaction at order $\mathbf{P}^2/4m^2$ [5]

$$\begin{aligned} \delta v(\mathbf{P}) = & -\frac{\mathbf{P}^2}{8m^2}\tilde{v} + \frac{1}{8m^2}[(\mathbf{P} \cdot \mathbf{r})(\mathbf{P} \cdot \nabla), \tilde{v}] \\ & + \frac{1}{8m^2}[(\boldsymbol{\sigma}_1 - \boldsymbol{\sigma}_2) \times (\mathbf{P} \cdot \nabla), \tilde{v}], \end{aligned} \quad (13)$$

where the potential in the NN center-of-mass frame, \tilde{v} , includes all relativistic corrections independent of \mathbf{P} , and the gradient operator acts on the relative coordinate. The first two terms in the right hand side of the above equation are associated with the relativistic energy-momentum relation and Lorentz contraction, respectively, whereas the last contribution—which has been shown to be negligibly small in light nuclei [27]—account for Thomas precession and quantum mechanical effects.

The results reported in this work have been obtained using the Hamiltonian

$$\begin{aligned} H_{\text{NR}}^* = & \sum_{i=1}^A \frac{\mathbf{P}_i^2}{2m} + \sum_{j>i=1}^A [v_{ij} + \delta v(\mathbf{P}_{ij})] \\ & + \sum_{k>j>i=1}^A V_{ijk}^*, \end{aligned} \quad (14)$$

in which V_{ijk}^* denotes the NNN potential modified to account for the presence of the boost interaction δv . The calculation of δv has been carried out following the approximations employed in Ref. [6], which amounts to neglecting the second line of Eq. (13) as well as non-static NN interactions, corresponding to the $p > 6$ components of the NN potential.

III. EFFECTIVE HAMILTONIAN

Within the approach developed by the authors of Ref. [12]—a simplified implementation of which was originally proposed in Ref. [28]—the effective interaction is defined by the equation

$$\langle \Psi_0 | H_{\text{NR}} | \Psi_0 \rangle = T_F + \langle \Phi_0 | V_{\text{eff}} | \Phi_0 \rangle \quad (15)$$

where H_{NR} is the non relativistic Hamiltonian of Eq. (1), T_F denotes the energy of the non interacting system, and

$$V_{\text{eff}} = \sum_{j>i=1}^A v_{ij}^{\text{eff}}, \quad (16)$$

with the effective NN potential v_{ij}^{eff} written as in Eq.(2) and including $p \leq 6$ contributions only.

The nuclear matter ground state $|\Psi_0\rangle$ is obtained from the Fermi gas ground state, $|\Phi_0\rangle$, through the transformation

$$|\Psi_0\rangle = \frac{\mathcal{F}|\Phi_0\rangle}{\sqrt{\langle \Phi_0 | \mathcal{F}^\dagger \mathcal{F} | \Phi_0 \rangle}}, \quad (17)$$

with

$$\mathcal{F} = \mathcal{S} \prod_{j>i} F_{ij}. \quad (18)$$

The two-nucleon correlation functions appearing in the right-hand side of the above equation are meant to embody all dynamical effects. As a consequence, they can

[‡] The subscripts specifying the labels of the interacting particles will be omitted whenever not necessary.

be conveniently written in the form

$$F_{ij} = \sum_{p=1}^6 f^p(r_{ij}) O_{ij}^p, \quad (19)$$

reminiscent of the NN potential of Eq. (2). Note that because, in general, $[F_{ij}, F_{jk}] \neq 0$, the product in Eq.(18) needs to be symmetrised through the action of the operator \mathcal{S} . The most prominent effect of the inclusion of correlations is the appearance of *screening*, that is, a strong suppression of the probability to find two nucleons within the range of the non perturbative repulsive core of the NN interaction.

Using the formalism described in Ref. [12], the effective Hamiltonian of Eq. (15) can be readily used to obtain the energy of nuclear matter with arbitrary neutron excess. In the case one-component Fermi liquids, such as PNM and SNM, the ground state expectation value of the effective interaction reduces to the form[‡]

$$\langle V_{\text{eff}} \rangle = A \frac{\rho}{2} \int d^3r_{12} \text{CTr} \left[v_{12}^{\text{eff}} \left(1 - \hat{P}_{12}^{\sigma\tau} \ell^2(k_F r_{12}) \right) \right], \quad (20)$$

where k_F is the Fermi momentum and $r = |\mathbf{r}_1 - \mathbf{r}_2|$. In the above equation, $\text{CTr}[\dots]$ denotes the trace operation in spin-isospin space, normalised such that $\text{CTr}[\mathbb{1}] = 1$, and $\hat{P}_{12}^{\sigma\tau}$ is the spin-isospin exchange operator, defined as

$$\hat{P}_{12}^{\sigma\tau} = \frac{1}{4} (1 + \boldsymbol{\sigma}_1 \cdot \boldsymbol{\sigma}_2) (1 + \boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_2). \quad (21)$$

The function $\ell(k_F r)$ —often referred to as Slater function—is trivially related to the density matrix of the non interacting system at density ρ through

$$\begin{aligned} \ell(k_F r) &= \frac{1}{A} \sum_{\mathbf{k}} \theta(k_F - |\mathbf{k}|) e^{i\mathbf{k}\cdot\mathbf{r}} \\ &= 3 \left[\frac{\sin(k_F r) - (k_F r) \cos(k_F r)}{(k_F r)^3} \right], \end{aligned} \quad (22)$$

with $\theta(x)$ being the Heaviside step function, respectively.

The effective interaction is obtained from Eq. (15) using the cluster expansion formalism [11], which amounts to rewrite the left-hand side in the form

$$\langle \Psi_0 | H_{\text{NR}} | \Psi_0 \rangle = T_F + (\Delta E)_2 + \dots, \quad (23)$$

where $(\Delta E)_n$ is the contribution arising from subsystems—or clusters—comprising n particles.

Keeping only two-body terms, one obtains the simple expression

$$v_{ij}^{\text{eff}} = -\frac{1}{m} (\nabla F_{ij})^2 + F_{ij} v_{ij} F_{ij}, \quad (24)$$

[‡] Hereafter, $\langle \dots \rangle$ will denote an expectation value in the Fermi gas ground state




FIG. 2. Radial and density dependence of the CBF effective interaction obtained from the AV6P+UIX Hamiltonian. For comparison, the thick solid line shows the behaviour of the bare AV6P potential.

which clearly illustrates how the the bare NN potential is *renormalised* by the action of the correlation functions.

The shape of the $f^p(r)$ is found by solving the set of Euler-Lagrange equations obtained from the minimisation of the ground state energy, evaluated at the two-body cluster level using the NN potential

$$\bar{v}_{ij} = \sum_p [\delta_{p1} + (1 - \delta_{p1})\alpha] v^p(r_{ij}) O_{ij}^p, \quad (25)$$

with α being a parameter meant to take into account the quenching of two-nucleon interactions due to the presence of the nuclear medium. The correlation functions satisfy the boundary conditions

$$f^p(r \geq d^p) = \delta_{1p}, \quad \left. \frac{df^p(r)}{dr} \right|_{d^p} = 0, \quad (26)$$

where, following Ref. [29], we have set

$$d^{p<5} = d_c, \quad d^{p=5,6} = d_t. \quad (27)$$

From the above discussion, it follows that, for any density ρ , the correlation functions associated with a given nuclear Hamiltonian depend on three parameters: the quenching factor α and the two relaxation distances d_c and d_t . In principle, they should be regarded as variational parameters, to be determined from the minimisation of the ground-state energy within an advanced computational method, such as the FHNC/SOC summation scheme or the Monte Carlo technique. According to the philosophy underlying the effective interaction formalism, on the other hand, they are adjusted in such a way as to reproduce a set of *target* results—obtained from accurate non perturbative calculations—using the potential V_{eff} defined by Eqs. (15)–(16) and first order

perturbation theory. This procedure has been successfully implemented to derive an effective interaction from the AV6P+UIX Hamiltonian, with the role of target results being played by the EOSs of SNM and PNM, obtained from the same Hamiltonian using the FHNC/SOC and the AFDMC computational techniques, respectively.

As pointed out previously, the determination of V_{eff} from Eq. (15) involves the calculation of the expectation value of the the nuclear Hamiltonian in the correlated ground state, performed using the cluster expansion method. The lowest order approximation allowing to take into account both the NN and NNN potentials involves two- and three-body cluster contributions. The resulting expression can be cast in the form

$$\langle V_{\text{eff}} \rangle = \langle \mathcal{F}^\dagger [T, \mathcal{F}] \rangle_{|2b} + \langle \mathcal{F}^\dagger V_{\text{NN}} \mathcal{F} \rangle_{|2b} + \langle \mathcal{F}^\dagger V_{\text{NNN}} \mathcal{F} \rangle_{|3b}, \quad (28)$$

where T denotes the A -body kinetic energy operator,

$$V_{\text{NN}} = \sum_{j>i=1}^A v_{ij}, \quad V_{\text{NNN}} = \sum_{k>j>i=1}^A V_{ijk}, \quad (29)$$

and the subscripts $2b$ and $3b$ label two- and three-nucleon cluster contributions, respectively.

It should be emphasised that the correlation operator \mathcal{F} appearing in Eq. (28) is *not* the same as the one determined performing a minimisation of the ground-state energy of nuclear matter. The values of the parameters α , d_c , and d_t entering the calculation of the right-hand side of Eq. (28) are instead adjusted to satisfy Eq. (15) using the effective interaction V_{eff} —obtained from the AV6P+UIX Hamiltonian—and the target results appearing in the the left-hand side. The radial and density dependence of CBF effective potential obtained from the AV6P+UIX Hamiltonian are illustrated in Fig. 2

IV. BOOST CORRECTION TO THE EOS OF NUCLEAR MATTER

As a first approximation, the impact of including relativistic boost corrections into the effective interaction of Ref. [12] has been investigated by treating the corresponding contribution to the ground-state energy as a perturbation. The resulting expression reads

$$\tilde{V}_{\text{eff}} = V_{\text{eff}} + \langle \mathcal{F}^\dagger \delta V \mathcal{F} \rangle_{|2b} = V_{\text{eff}} + \delta V_{\text{eff}}, \quad (30)$$

with

$$\delta V = \sum_{j>i} \delta v_{ij}, \quad (31)$$

where, following Ref. [6], the form of δv_{ij} is approximated by

$$\delta v_{ij} = -\frac{\mathbf{P}_{ij}^2}{8m^2} v_{ij} + \frac{1}{8m^2} [(\mathbf{P}_{ij} \cdot \mathbf{r})(\mathbf{P}_{ij} \cdot \nabla), v_{ij}]. \quad (32)$$

In the above equation, the gradient operates on the relative coordinate and v_{ij} is a NN potential whose operator structure is described by the $O^{p \leq 6}$ of Eqs. (3) and (4).




FIG. 3. Energy per particle of PNM, obtained from AFDMC calculations performed including NN interactions only. Squares and triangles correspond to the AV6P potential and to the truncated AV18 potential discussed in the text.

A. AV18 and AV6P Hamiltonians

In order to pin down the effects of δv alone, we have first considered Hamiltonian models including only the two-nucleon potential.

The EOSs of PNM and SNM computed by the authors of Refs. [6, 30] using the AV18 and AV18+ δv Hamiltonians have been compared to those obtained from the corresponding effective interactions, derived from the AV6P potential using the AV18 EOSs as target.

It must be kept in mind that, while having the same operator structure, the truncated AV18 potential—employed to evaluate the contribution of the relativistic boost interaction in Ref. [6]—differs significantly from the projected AV6P potential. The discrepancies observed in the NN scattering phase shifts, discussed in Section II A, also strikingly emerge in the density dependence of the energy per nucleon of PNM computed using the AFDMC technique; see Fig. 3.

Figure 4 shows a comparison between the variational ground-state energies of PNM and SNM determined by Akmal *et al.* using the AV18 and AV18+ δv Hamiltonians and the corresponding results obtained using the CBF effective interaction with and without inclusion of the relativistic correction defined by Eq. (30). The results of calculations carried out using the AV6P and the truncated AV18 potential—labelled $\delta V_{\text{eff}}(\text{AV6P})$ and $\delta V_{\text{eff}}(\text{AV18})$, respectively—turn out to be significantly different, with $\delta V_{\text{eff}}(\text{AV18})$ being appreciably larger than $\delta V_{\text{eff}}(\text{AV6P})$ over the whole density range. It is apparent that both the $V_{\text{eff}} + \delta V_{\text{eff}}(\text{AV6P})$ and $V_{\text{eff}} + \delta V_{\text{eff}}(\text{AV18})$ models fail to reproduce the results obtained using the AV18+ δv Hamiltonian.

We investigated the origin of the pattern emerging from Fig. 4 by studying the sensitivity of the boost interaction energy to modifications of the NN correlation functions $f^p(r)$. In Fig. 5, the lines labelled (X,Y)—with X,Y \equiv AV18 or AV6P—represent results obtained us-




FIG. 4. Energy dependence of the binding energy per particle of PNM and SNM obtained using Hamiltonians including only NN potentials. The meaning of the curves is explained in the text.

ing the potential model X for the solution of the Euler-Lagrange equations determining the shape of the $f^p(r)$, and model Y in the calculation of the boost interaction energy. It appears that the observed behaviour is primarily driven by the NN potential, although in the case $X \equiv \text{AV6P}$ appreciable differences between the energies corresponding to different correlation functions are visible at high densities.




FIG. 5. Density dependence of the relativistic boost correction δV_{eff} —defined by Eq. (30)—in PNM. The results have been obtained using different NN potentials and correlation functions. The meaning of the curves is explained in the text.

B. AV18+UIX and AV6P+UIX Hamiltonians

The formalism described in the previous sections has been generalised to allow the treatment of models of nuclear dynamics including a NNN potential, the corresponding effective interaction being defined by Eq.(28).

The parameters α , d_c and d_t entering the definition of the two-nucleon correlation functions, discussed in Section III, have been adjusted to reproduce the EOSs of both PNM and SNM reported by Akmal *et al.* in Ref. [6]. In the following, the EOSs obtained by these authors using the AV18+UIX and AV18+ δv +UIX* Hamiltonians will be referred to as APR1 and APR2, respectively. We recall the reader that the presence of the repulsive boost interaction brings about a significant modification of the UIX potential. The strength of the isoscalar repulsive contribution to the resulting NNN interaction, denoted UIX*, turns out to be reduced by a factor $\gamma = 0.63$ [4, 6].

The energies per nucleon corresponding to the APR2 model reported in Ref. [6] have been obtained by adding to the corresponding APR1 results the contribution of the boost interaction and the associated modification of the NNN potential, computed in first order perturbation theory. The resulting expression can be written in the form

$$\left(\frac{E}{A}\right)_{\text{APR2}} = \left(\frac{E}{A}\right)_{\text{APR1}} + \frac{1}{A} \langle \mathcal{F}^\dagger [\delta V + (\gamma - 1) V^R] \mathcal{F} \rangle, \quad (33)$$

with the correlation operator \mathcal{F} being determined from minimisation of the expectation value of the AV18+UIX Hamiltonian in the correlated ground state, and

$$V^R = \sum_{k>j>i} V_{ijk}^R. \quad (34)$$

To identify the effects of relativistic corrections, we have evaluated the contribution of the effective potential to the ground-state energy using different prescriptions. In the first scenario, relativistic boost corrections are neglected altogether, whereas the two additional definitions are

$$\delta V_{\text{eff}} = \langle \mathcal{F}^\dagger [\delta V + (\gamma - 1) V^R] \mathcal{F} \rangle, \quad (35)$$

and

$$\delta V_{\text{eff}}^* = (\gamma - 1) \langle \mathcal{F}^\dagger V^R \mathcal{F} \rangle. \quad (36)$$

In all three cases the correlation functions determining the effective interaction are the same, with the corresponding parameters being adjusted using as target results the APR1 EOSs of Ref. [6].

The solid lines of Fig. 6—which reproduce the APR1 EOSs by construction—represent the results obtained using V_{eff} defined by Eq. (28), while the dashed lines correspond to calculations carried out using δV_{eff}^* of Eq. (36). The results obtained including the full relativistic correction of Eq.(35) and using the AV6P and the truncated




FIG. 6. Energy dependence of the binding energy per particle of PNM and SNM obtained using Hamiltonians including both NN and NNN potentials. The meaning of the curves is explained in the text.

AV18 in the calculation of δV are displayed by the dot-dash and dotted lines, respectively.

It clearly appears that the differences originating from the use of the AV6P or the truncated AV18 potential in the calculation of the boost interaction energy, while being still visible, are largely obscured by the size of the associated modification of the repulsive NNN potential.

V. BOOST CORRECTION TO THE CBF EFFECTIVE INTERACTION

In the previous sections, the relativistic boost interaction has been treated in perturbation theory, which amounted to adding to the CBF effective potential of Eq. (28) the expectation value of δV in the correlated ground state, defined by Eq. (30). Here, we will discuss a procedure to take into account the boost interaction at operator level, which implies determining the relativistic effective potential, denoted V'_{eff} , from the equation

$$\langle \mathcal{F}^\dagger H' \mathcal{F} \rangle = T_F + \langle V'_{\text{eff}} \rangle, \quad (37)$$

where

$$H' = \sum_i \frac{\mathbf{p}_i^2}{2m} + \sum_{j>i} (v_{ij} + \delta v_{ij}) + \sum_{k>j>i} V_{ijk}^*, \quad (38)$$

the operator structure of V'_{eff} being the same as that of Eq.(2) with $p \leq 6$.

It clearly appears that the procedure based on Eqs. (37) and (38) allows to take into account the effects of boost interactions in the determination of the radial dependence of the correlation functions $f^p(r)$, which in turn drives the behaviour of the effective interaction.

The numerical results obtained from the above method are summarised in Fig. 7. In analogy to Sect. IV B we have considered two different definitions of the effective potential. The first one, which takes into account boost corrections only through the replacement $V_{\text{NNN}} \rightarrow V_{\text{NNN}}^*$, is written in the form

$$\begin{aligned} \langle V_{\text{eff}} \rangle = & \langle \mathcal{F}^\dagger [T, \mathcal{F}] \rangle|_{2b} + \langle \mathcal{F}^\dagger V_{\text{NN}} \mathcal{F} \rangle|_{2b} \\ & + \langle \mathcal{F}^\dagger V_{\text{NNN}}^* \mathcal{F} \rangle|_{3b}, \end{aligned} \quad (39)$$

reminiscent of the non relativistic expression of Eq. (28). The second prescription, on the other hand, is obtained from Eq. (37), and includes relativistic corrections to both the NN and NNN potentials. The corresponding expression reads

$$\begin{aligned} \langle V'_{\text{eff}} \rangle = & \langle \mathcal{F}^\dagger [T, \mathcal{F}] \rangle|_{2b} + \langle \mathcal{F}^\dagger (V_{\text{NN}} + \delta V) \mathcal{F} \rangle|_{2b} \\ & + \langle \mathcal{F}^\dagger V_{\text{NNN}}^* \mathcal{F} \rangle|_{3b}. \end{aligned} \quad (40)$$

The effective potentials defined by Eqs. (39) and (40) have been both determined in such a way as to reproduce the APR2 EOSs, obtained using the AV18+ δv +UIX* Hamiltonian. The details of the calculations can be found in the appendix.

The density dependence of the parameters α , d_c and d_t obtained by fitting the target EOSs with the effective potentials of Eqs.(39) and (40) are displayed in panels (a) and (b) of Fig. 7, respectively, while panels (c) and (d) show the corresponding energies per particle of PNM and SNM. The jump in the values of the parameters determining the shape of the correlation functions, clearly visible at $\rho \approx 0.2 \text{ fm}^{-3}$, is likely to reflect the appearance of the spin-isospin ordered phase—associated with the occurrence of neutral pion condensation—discussed in Ref.[6]. The small kink at $\rho \approx 0.32 \text{ fm}^{-3}$ may also be ascribed to this phase transition in SNM. It appears that the inclusion of boost interactions has only a marginal impact on the parameter values, and the target energies can be accurately reproduced with both choices of the effective potential.

Figure 8 illustrates the radial behaviour of the CBF effective potential acting in the two-nucleon channel of spin-isospin $S = 0$ and $T = 1$ at density $\rho = 0.16$ and 0.64 fm^{-3} . The solid line corresponds to V_{eff} of Eq. (28), while the dashed line represents the effective interaction V'_{eff} determined including relativistic boost interactions according to Eq. (40). The UIX* NNN potential has been used in both cases. Note that, because the effective interactions V_{eff} and V'_{eff} are optimised to reproduce the same APR2 EOSs, the values of the adjustable parameters α , d_c and d_t determining the shape of the correlation functions involved in their definitions are not the same. The modification arising from boost interactions are clearly visible.




FIG. 7. The top panels show the correlation ranges and quenching parameters, defined by Eqs. (25)-(27) obtained optimizing the effective interaction to reproduce APR2 EOSs without (a) and with (b) the inclusion of boost corrections in its definition. The corresponding energies per nucleon of both SNM and PNM are displayed in panels (c) and (d). Squares and circles represent the results of Akmal *et al.* [6].

VI. SUMMARY AND OUTLOOK

We have investigated the impact of including relativistic boost corrections into the density-dependent effective interaction of Ref.[12], derived from the AV6P+UIX model of the nuclear Hamiltonian using the CBF formalism and the cluster expansion technique. The contribution of the boost interaction δV has been considered both at average level, by treating its expectation value in the correlated ground state as a perturbation, and at microscopic level, by explicitly taking into account δV in the determination the effective potential.

The form of δV has been obtained following the procedure described in Refs. [4, 5], designed to be used with static NN potentials. We found significant differences between the contributions of the boost interaction to nuclear matter energy computed with either the

AV6P potential or the truncated version of the AV18 model employed by Akmal *et al.* [6], in which all angular-momentum-dependent terms are neglected. These discrepancies suggest that the projection method used by the authors of Ref.[18] to obtain the AV6P potential allows to effectively capture some important features of non static interactions. The contribution of the boost interaction computed using the truncated AV18 interaction turns out to be considerably larger than that obtained from the AV6P at all densities.

The inclusion of a repulsive correction to the NN potential obviously entails a corresponding decrease of the repulsive component of the NNN potential, V^R , needed to preserve the capability of the Hamiltonian to explain the measured binding energies of ${}^3\text{He}$ and ${}^4\text{He}$ and the empirical value of the saturation density of SNM. Our analysis shows that the resulting $\sim 30\%$ reduction of the strength of repulsive NNN interactions is, in fact, the




FIG. 8. Radial dependence of the CBF effective interaction in the $S = 0, T = 1$ channel at density $\rho = 0.16$ (a) and 0.64 fm^{-3} (b). The solid line represents the potential V_{eff} obtained from Eq. (39) using the UIX* NNN potential, while the dashed line corresponds to the effective interaction V'_{eff} , determined including the contribution of relativistic boost interactions according to Eq. (40).

dominant relativistic correction, leading to a significant softening of the nuclear matter EOS at high densities.

In this context, it is worth mentioning that additional constraints on NNN interactions in dense matter may be provided by multimessenger neutron star observations. The results of the analyses performed by the authors of Refs.[31, 32] suggest that the existing and forthcoming data have the potential to allow an accurate determination of the parameter γ of Eq. (33), providing a measure of the relativistic boost correction to the strength of V^R .

The analysis described in this article is meant to be a first exploratory step towards a fully consistent inclusion of relativistic boost corrections into the effective interaction of Ref. [12]. The achievement of this goal—which involves using correlation functions optimised to reproduce the EOSs of PNM and SNM obtained from the AV6P+UIX* Hamiltonian—will extend the density region in which the CBF effective Hamiltonian can be expected to provide a reliable description of nuclear matter properties relevant to neutron star structure and dynamics.

Relativistic modifications of the three-nucleon potential have been also recently discussed by the authors of Ref. [33], who derived the boost corrections to the lead-

ing order NNN contact interaction from the constraints imposed by Poincaré algebra on non relativistic dynamics. However, their impact on nuclear properties has not been studied yet.

As a final remark, it should be emphasised that, in assessing the role of relativistic corrections to the dynamics of dense matter, one has to keep in mind that the approach followed in our study is only applicable in the density regime in which nuclear matter can be described in terms of nucleon degrees of freedom. The data recently reported by the Neutron Star Internal Composition Explorer (NICER)[34, 35], showing that neutron stars of ~ 1.4 and ~ 2.1 solar masses have similar radii, provide convincing evidence that the limit of applicability of this description may be as high as $\sim 4\rho_0$.

ACKNOWLEDGMENTS

This research was funded by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, under contract DE-AC02-06CH11357 (A.L.), the NUCLEI SciDAC program (A.L.), and the Italian National Institute for Nuclear Research (INFN), under grant TEONGRAV (O.B. and A.S.).

Appendix A: Boost corrections to the CBF effective interaction

In this Appendix, we provide details on the calculation of the effective interaction defined by Eq. (40), in which the corrections arising from relativistic boost interactions are explicitly taken into account at operator level. In view of the fact that the contribution of δV is computed in the two-body cluster approximation, we can limit our discussion

to the case of a Hamiltonian comprising two-nucleon terms only, which can be written

$$H = T + V_{\text{NN}} + \delta V = \sum_i \frac{\mathbf{p}_i^2}{2m} + \sum_{j>i} (v_{ij} + \delta v_{ij}). \quad (\text{A1})$$

The effective interaction is obtained by expanding the expectation value of H in the correlated ground state, and retaining only the two-body cluster contribution $(\Delta E)_2$; see Eq. (23). In this case, the boost correction, which will be denoted $\delta v_{ij}^{\text{eff}}$, is implicitly defined by the equation

$$(\Delta E)_2 = \langle \mathcal{F}^\dagger [T, \mathcal{F}] \rangle_{2b} + \langle \mathcal{F}^\dagger \sum_{j>i} (v_{ij} + \delta v_{ij}) \mathcal{F} \rangle_{2b} = \langle \sum_{j>i} (v_{ij}^{\text{eff}} + \delta v_{ij}^{\text{eff}}) \rangle, \quad (\text{A2})$$

where

$$v_{ij}^{\text{eff}} = -\frac{1}{m} (\nabla F_{ij})^2 + F_{ij} v_{ij} F_{ij}, \quad (\text{A3})$$

with F_{ij} being the NN correlation function of Eq. (19). Because both v_{ij} and F_{ij} depend linearly on the operators $O^{p \leq 6}$ defined by Eqs. (3) and (4), one can exploit the algebra

$$O_{ij}^p O_{ij}^q = \sum_{r=1}^6 K^{pqr} O_{ij}^r, \quad (\text{A4})$$

with the coefficients K^{pqr} given in, e.g., Ref.[36], to cast the NN effective potential in the form

$$v_{ij}^{\text{eff}} = \sum_{p=1}^6 v^{\text{eff},p}(r_{ij}) O_{ij}^p. \quad (\text{A5})$$

The form of the boost interaction employed in this work, given by Eq.(32), has been derived using the approximations discussed in Refs. [5, 6]. It does not include the terms arising from Thomas Precession and the commutators involving the spin-dependent part of the NN potential present in Eq. (13), whose contributions to the binding energies of ${}^3\text{He}$ and ${}^4\text{He}$ have been found to be negligible [37]. For a static NN potential written as in Eq. (2) one obtains

$$\delta v_{12} = \sum_{p=1}^6 \left[-\frac{\mathbf{P}^2}{8m^2} v^p(r_{12}) + \frac{(\mathbf{P} \cdot \hat{\mathbf{r}}_{12})^2}{8m^2} r_{12} \frac{dv^p}{dr_{12}} \right] O_{12}^p + \sum_{p=1}^6 \frac{(\mathbf{P} \cdot \mathbf{r}_{12})}{8m^2} v^p(r_{12}) (\mathbf{P} \cdot \nabla) O_{12}^p, \quad (\text{A6})$$

where $\hat{\mathbf{r}}_{12} = \mathbf{r}_{12}/r_{12}$. Note that the last term in the right-hand side of the above equation involves the gradient of the tensor operator S_{12} , whose expectation value vanishes in unpolarised matter. For this reason, here we limit ourselves to consider the expression

$$\delta v_{12} = \sum_{p=1}^6 \left[-\frac{\mathbf{P}^2}{8m^2} v^p(r_{12}) + \frac{(\mathbf{P} \cdot \hat{\mathbf{r}}_{12})^2}{8m^2} r_{12} \frac{dv^p}{dr_{12}} \right] O_{12}^p = \sum_{i=1}^6 \delta v^p(r_{12}) O_{12}^p. \quad (\text{A7})$$

To obtain the expression of the relativistic boost correction we need to calculate the two-body cluster contribution to the expectation value of δV in the correlated ground state. As a first step, we consider the expectation value in the Fermi gas ground state, which can be readily evaluated and split into direct and exchange contributions according to

$$\frac{1}{A} \langle \delta V \rangle = \frac{1}{A} \langle \sum_{j>i} \delta v_{ij} \rangle = \frac{1}{A} \langle \delta V \rangle_D - \frac{1}{A} \langle \delta V \rangle_E, \quad (\text{A8})$$

where

$$\frac{1}{A} \langle \delta V \rangle_D = \frac{1}{5} \frac{\rho}{8m^2} k_F^2 \sum_p \int d^3 r_{12} \left[r_{12} \frac{dv^p(r_{12})}{dr_{12}} - 3v^p(r_{12}) \right] \text{CTr}[O_{12}^p], \quad (\text{A9})$$

and

$$\frac{1}{A} \langle \delta V \rangle_E = \frac{\rho}{16m^2} \sum_p \int d^3 r_{12} \left[r_{12} \frac{dv^p(r_{12})}{dr_{12}} I_2(k_F, r_{12}) - v^p(r) I_1(k_F, r_{12}) \right] \text{CTr}[O^p \hat{P}_{12}^{\sigma\tau}], \quad (\text{A10})$$

with

$$\begin{aligned} I_1(k_F, r) &= 2[\ell'(k_F r)]^2 - 2\ell(k_F r)\nabla_r^2\ell(k_F r) , \\ I_2(k_F, r) &= 2[\ell'(k_F r)]^2 - 2\ell(k_F r)\ell''(k_F r) . \end{aligned}$$

Here a prime indicates derivative with respect to r ,

$$\nabla_r^2 \equiv \frac{1}{r} \frac{\partial^2}{\partial r^2} r , \quad (\text{A11})$$

and k_F denotes the Fermi momentum. Note that in this work we only consider one-component Fermi liquids, such as PNM and SNM.

A more compact expression can be obtained collecting the direct and exchange terms in the right-hand side of Eq. (A8). The result can be written in the form

$$\frac{1}{A} \langle \delta V \rangle = \frac{\varrho}{2} \sum_{p=1}^6 \left\{ \int d^3 r_{12} A^p(k_F, r_{12}) \text{CTr}[O_{12}^p] - \int d^3 r_{12} B^p(k_F, r_{12}) \text{CTr}[O_{12}^p \widehat{P}_{12}^{\sigma\tau}] \right\} . \quad (\text{A12})$$

where

$$\begin{aligned} A^p(k_F, r) &\equiv \frac{1}{8m^2} \frac{2}{5} k_F^2 \left[r \frac{dv^p(r)}{dr} - 3v^p(r) \right] , \\ B^p(k_F, r) &\equiv \frac{1}{8m^2} \left[r \frac{dv^p(r)}{dr} I_2(k_F, r) - v^p(r) I_1(k_F, r) \right] . \end{aligned} \quad (\text{A13})$$

Finally, because the spin-isospin exchange operator can be written in terms of the operators of Eq. (3) as

$$\widehat{P}_{12}^{\sigma\tau} = \frac{1}{4} \sum_{q=1}^4 O_{12}^q , \quad (\text{A14})$$

we can use the algebra (A4) to obtain

$$\frac{1}{A} \langle \delta V \rangle = \frac{\varrho}{2} \sum_{p,m=1}^6 \int d^3 r_{12} \left[A^p(k_F, r_{12}) \delta_{pm} - \frac{1}{4} \sum_{q=1}^4 K^{pmq} B^p(k_F, r_{12}) \right] \text{CTr}[O_{12}^m] . \quad (\text{A15})$$

Let us now consider the two-body cluster approximation to the expectation value of δV in the correlated ground state, derived in, e.g., Ref. [38], whose expression reads

$$\begin{aligned} \frac{1}{A} \langle \mathcal{F}^\dagger \left(\sum_{j>i} \delta v_{ij} \right) \mathcal{F} \rangle_{2b} &= \frac{\varrho}{2} \int d^3 r_{12} \text{CTr} \left[\sum_m F_{12}^\dagger \delta v^m(r_{12}) F_{12} O_{12}^m \left(1 - \widehat{P}_{12}^{\sigma\tau} \ell^2(k_F r_{12}) \right) \right] \\ &= \frac{\varrho}{2} \int d^3 r_{12} \text{CTr} \left[\sum_{qmn} f^q(r_{12}) \delta v^m(r_{12}) f^n(r_{12}) O_{12}^q O_{12}^m O_{12}^n \left(1 - \widehat{P}_{12}^{\sigma\tau} \ell^2(k_F r_{12}) \right) \right] . \end{aligned} \quad (\text{A16})$$

It clearly appears that by rewriting δv^p in the form

$$\delta v^p(r_{12}) = -\frac{\mathbf{P}^2}{8m^2} w^p(r_{12}) + \frac{(\mathbf{P} \cdot \hat{\mathbf{r}}_{12})^2}{8m^2} g^p(r_{12}) \quad (\text{A17})$$

with

$$w^p(r_{12}) = v^p(r_{12}) \quad , \quad g^p(r_{12}) = r_{12} \frac{dv^p}{dr_{12}} , \quad (\text{A18})$$

and exploiting again the algebra of Eq. (A4), the right-hand side of Eq. (A16) can be simply computed replacing

$$\begin{aligned} w^p(r) &\rightarrow \sum_{mnlq} K^{mnl} K^{qlp} f^q(r) v^m(r) f^n(r) , \\ g^p(r) &\rightarrow \sum_{mnlq} K^{mnl} K^{qlp} f^q(r) r \frac{dv^m}{dr} f^n(r) , \end{aligned} \quad (\text{A19})$$

in the expectation value of δV in the Fermi gas ground state.

From now on, summation over repeated indices of the $O^{p \leq 6}$ operators of Eq. (A4) will be understood, as well as the subscripts labeling the interacting particles. By defining the quantities

$$\begin{aligned}\xi^m(k_F, r) &= \left\{ A^m(k_F, r) - \frac{1}{4} \sum_{q=1}^4 K^{pqm} B^p(k_F, r) \right\}, \\ M_{pm} &= \frac{1}{4} \sum_{q=1}^4 K^{pqm}.\end{aligned}\tag{A20}$$

we can rewrite the Fermi gas result in the form

$$\begin{aligned}\frac{1}{A} \langle \delta V \rangle &= \frac{\varrho}{2} \int d^3r \text{CTr}[O^m] \xi^m(k_F, r) \\ &= \frac{\varrho}{2} \int d^3r \text{CTr}[O^m] [\delta_{mn} - \ell^2(k_F r) M_{mn}] [\delta_{np} - \ell^2(k_F r) M_{np}]^{-1} \xi^p(k_F, r).\end{aligned}\tag{A21}$$

Since

$$[\delta_{mp} - \ell^2(k_F r) M_{mp}]^{-1} = \frac{1}{1 - \ell^4(k_F r)} [\delta_{mp} + \ell^2(k_F r) M_{mp}],\tag{A22}$$

we finally obtain

$$\frac{1}{A} \langle \delta V \rangle = \frac{\varrho}{2} \int d^3r \text{CTr}[O^m] [\delta_{mp} - \ell^2(k_F r) M_{mp}] \delta v^p(k_F, r)\tag{A23}$$

where we have defined

$$\delta v^p(k_F, r) \equiv \frac{1}{1 - \ell^4(k_F r)} [\delta_{pm} + \ell^2(k_F r) M_{pm}] \xi^m(k_F, r).\tag{A24}$$

By recalling that the operator M is linked to the spin-isospin exchange operator $\widehat{P}^{\sigma\tau}$, Eq. (A23) can be written as

$$\frac{1}{A} \langle \delta V \rangle = \frac{\varrho}{2} \int d^3r \text{CTr} \left[O^p \left(1 - \ell^2(k_F r) \widehat{P}^{\sigma\tau} \right) \right] \delta v^p(k_F, r)\tag{A25}$$

which is the expression we were looking for.

Finally we are going to derive the expression of Eq. (A22). Since the operator we want to invert has the form

$$\mathbb{1} - \ell^2 M\tag{A26}$$

we have that

$$(\mathbb{1} - \ell^2 M) (\mathbb{1} + \ell^2 M^{-1}) = (1 - \ell^4) \mathbb{1} - \ell^2 M + \ell^2 M^{-1}.\tag{A27}$$

Therefore if we show that

$$M = M^{-1}\tag{A28}$$

we have done. We start from the identity

$$O^m O^p O^q = K^{mpl} K^{lqs} O^s\tag{A29}$$

and summing over $p, q = 1, \dots, 4$ we have

$$\sum_{p,q=1}^4 O^m O^p O^q = O^m \sum_{p=1}^4 O^p \sum_{q=1}^4 O^q = O^m (4 \widehat{P}^{\sigma\tau}) (4 \widehat{P}^{\sigma\tau}) = 16 O^m (\widehat{P}^{\sigma\tau})^2 = 16 O^m.\tag{A30}$$

Therefore we can write

$$O^m = \frac{1}{16} \sum_{p,q=1}^4 O^m O^p O^q = \left(\frac{1}{4} \sum_{p=1}^4 K^{mpl} \right) \left(\frac{1}{4} \sum_{q=1}^4 K^{lqs} \right) O^s = M_{ml} M_{ls} O^s\tag{A31}$$

which entails

$$M_{ml}M_{ls} = \delta_{ms} \Rightarrow M = M^{-1}. \quad (\text{A32})$$

-
- [1] R. Wiringa, “From deuterons to neutron stars: variations in nuclear many-body theory,” *Rev. Mod. Phys.* **65** (Jan, 1993) 231–242. <https://link.aps.org/doi/10.1103/RevModPhys.65.231>.
- [2] F. Coester, S. Pieper, and F. Serduke, “Relativistic effects in phenomenological nucleon-nucleon potentials and nuclear matter,” *Phys. Rev. C* **11** (Jan, 1975) 1–18. <https://link.aps.org/doi/10.1103/PhysRevC.11.1>.
- [3] W. Glöckle, T.-S. H. Lee, and F. Coester, “Relativistic effects in three-body bound states,” *Phys. Rev. C* **33** (Feb, 1986) 709–716. <https://link.aps.org/doi/10.1103/PhysRevC.33.709>.
- [4] J. L. Forest, V. R. Pandharipande, and A. Arriaga, “Quantum monte carlo studies of relativistic effects in light nuclei,” *Phys. Rev. C* **60** (Jun, 1999) 014002. <https://link.aps.org/doi/10.1103/PhysRevC.60.014002>.
- [5] J. Forest, V. R. Pandharipande, and J. L. Friar *Phys. Rev. C* **52** (1995) 568.
- [6] A. Akmal, V.R. Pandharipande, and D.G. Ravenhall, “Equation of state of nucleon matter and neutron star structure,” *Phys. Rev. C* **58** (1998) 1804.
- [7] R. B. Wiringa, V. G. J. Stoks, and R. Schiavilla, “An Accurate nucleon-nucleon potential with charge independence breaking,” *Phys. Rev. C* **51** (1995) 38–51.
- [8] J. Carlson, V. R. Pandharipande, and R. B. Wiringa *Nucl. Phys. A* **401** (1983) 59.
- [9] B. S. Pudliner, V. R. Pandharipande, J. Carlson, and R. B. Wiringa, “Quantum Monte Carlo calculations of $A \leq 6$ nuclei,” *Phys. Rev. Lett.* **74** (1995) 4396–4399.
- [10] S. Fantoni and A. Fabrocini, “Correlated Basis Function Theory for Fermion Systems,” in *Microscopic Quantum Many-Body Theories and Their Applications*, J. Navarro and A. Polls, ed., pp. 119–186. Springer, Berlin, Heidelberg, 1998. <https://doi.org/10.1007/BFb0104526>.
- [11] J. W. Clark, “Variational theory of nuclear matter,” *Progress in Particle and Nuclear Physics* **2** (1979) 89–199. <https://www.sciencedirect.com/science/article/pii/0146641079900048>.
- [12] O. Benhar and A. Lovato, “Perturbation theory of nuclear matter with a microscopic effective interaction,” *Phys. Rev. C* **96** (2017) 054301.
- [13] O. Benhar, A. Lovato, and G. Camelió, “Modeling neutron star matter in the age of multimessenger astrophysics,” *The Astrophysical Journal* **939** no. 1, (Nov, 2022) 52. <https://dx.doi.org/10.3847/1538-4357/ac8e61>.
- [14] L. Tonetto and O. Benhar, “Thermal effects on nuclear matter properties,” *Phys. Rev. D* **106** (Nov, 2022) 103020. <https://link.aps.org/doi/10.1103/PhysRevD.106.103020>.
- [15] O. Benhar, A. Lovato, and L. Tonetto, “Properties of Hot Nuclear Matter,” *Universe* **9** no. 8, (2023) 345.
- [16] J. L. Friar, “Trinucleon bound states,” in *New Vistas in Electro-Nuclear Physics*, E. L. Tomusiak, H. S. Caplan, and E. T. Dressler, eds., p. 213. Springer US, Boston, MA, 1986. https://doi.org/10.1007/978-1-4684-5200-6_6.
- [17] J. Fujita and H. Miyazawa, “Pion Theory of Three-Body Forces,” *Prog. Theor. Phys.* **17** (1957) 360–365.
- [18] R. Wiringa and S. Pieper, “Evolution of nuclear spectra with nuclear forces,” *Phys. Rev. Lett.* **89** (2002) 182501.
- [19] J. R. Bergervoet, P. C. van Campen, R. A. M. Klomp, J.-L. de Kok, T. A. Rijken, V. G. J. Stoks, and J. J. de Swart, “Phase shift analysis of all proton-proton scattering data below $t_{lab}=350$ mev,” *Phys. Rev. C* **41** (Apr, 1990) 1435–1452.
- [20] V. G. J. Stoks, R. A. M. Klomp, M. C. M. Rentmeester, and J. J. de Swart, “Partial-wave analysis of all nucleon-nucleon scattering data below 350 mev,” *Phys. Rev. C* **48** (Aug, 1993) 792–815.
- [21] R. A. Arndt, W. J. Briscoe, I. I. Strakovsky, and R. L. Workman *Phys. Rev. C* **76** (2007) 025209.
- [22] M. Piarulli, I. Bombaci, D. Logoteta, A. Lovato, and R. B. Wiringa, “Benchmark calculations of pure neutron matter with realistic nucleon-nucleon interactions,” *Phys. Rev. C* **101** (Apr, 2020) 045801. <https://link.aps.org/doi/10.1103/PhysRevC.101.045801>.
- [23] R. A. Krafcik and L. L. Foldy, “Relativistic center-of-mass variables for composite systems with arbitrary internal interactions,” *Phys. Rev. D* **10** (Sep, 1974) 1777–1795. <https://link.aps.org/doi/10.1103/PhysRevD.10.1777>.
- [24] J. L. Friar, “Relativistic effects on the wave function of a moving system,” *Phys. Rev. C* **12** (1975) 695–698. <https://journals.aps.org/prc/abstract/10.1103/PhysRevC.12.695>.
- [25] L. L. Foldy, “Relativistic particle systems with interaction,” *Phys. Rev.* **122** (Apr, 1961) 275–288. <https://link.aps.org/doi/10.1103/PhysRev.122.275>.
- [26] J. Carlson, V. R. Pandharipande, and R. Schiavilla, “Variational monte carlo calculations of ${}^3\text{H}$ and ${}^4\text{He}$ with a relativistic hamiltonian,” *Phys. Rev. C* **47** (Feb, 1993) 484–497. <https://link.aps.org/doi/10.1103/PhysRevC.47.484>.
- [27] J. Carlson, V. R. Pandharipande, and R. Schiavilla, “Variational monte carlo calculations of ${}^3\text{H}$ and ${}^4\text{He}$ with a relativistic hamiltonian,” *Phys. Rev. C* **47** (Feb, 1993) 484–497. <https://link.aps.org/doi/10.1103/PhysRevC.47.484>.
- [28] S. T. Cowell and V. R. Pandharipande, “Quenching of weak interactions in nucleon matter,” *Phys. Rev. C* **67** (2003) 035504, [arXiv:nuc1-th/0211013](https://arxiv.org/abs/nuc1-th/0211013).

- [29] I. Lagaris and V. Pandharipande, “Phenomenological two-nucleon interaction operator,” *Nuclear Physics A* **359** no. 2, (1981) 331–348. <https://www.sciencedirect.com/science/article/pii/0375947481902402>.
- [30] A. Akmal and V. R. Pandharipande, “Spin-isospin structure and pion condensation in nucleon matter,” *Phys. Rev. C* **56** (Oct, 1997) 2261–2279. <https://link.aps.org/doi/10.1103/PhysRevC.56.2261>.
- [31] A. Maselli, A. Sabatucci, and O. Benhar, “Constraining three-nucleon forces with multimessenger data,” *Phys. Rev. C* **103** no. 6, (2021) 065804, [arXiv:2010.03581](https://arxiv.org/abs/2010.03581) [astro-ph.HE].
- [32] A. Sabatucci, O. Benhar, A. Maselli, and C. Pacilio, “Sensitivity of neutron star observations to three-nucleon forces,” *Phys. Rev. D* **106** (Oct, 2022) 083010. <https://link.aps.org/doi/10.1103/PhysRevD.106.083010>.
- [33] A. Nasoni, E. Filandri, and L. Girlanda, “Relativistic constraints on 3N contact interactions,” *Eur. Phys. J. A* **59** no. 12, (2023) 293, [arXiv:2308.13341](https://arxiv.org/abs/2308.13341) [nucl-th].
- [34] T. E. Riley *et al.*, “A NICER View of the Massive Pulsar PSR J0740+6620 Informed by Radio Timing and XMM-Newton Spectroscopy,” *Astrophys. J. Lett.* **918** no. 2, (2021) L27, [arXiv:2105.06980](https://arxiv.org/abs/2105.06980) [astro-ph.HE].
- [35] T. E. Riley *et al.*, “A NICER View of PSR J0030+0451: Millisecond Pulsar Parameter Estimation,” *Astrophys. J. Lett.* **887** no. 1, (2019) L21, [arXiv:1912.05702](https://arxiv.org/abs/1912.05702) [astro-ph.HE].
- [36] V. R. Pandharipande and R. B. Wiringa, “Variations on a theme of nuclear matter,” *Rev. Mod. Phys.* **51** (1979) 821–859.
- [37] J. L. Forest, V. R. Pandharipande, J. Carlson, and R. Schiavilla, “Variational monte carlo calculations of ^3H and ^4He with a relativistic hamiltonian,” *Phys. Rev. C* **52** (Aug, 1995) 576–577. <https://link.aps.org/doi/10.1103/PhysRevC.52.576>.
- [38] J. W. Clark, “Variational theory of nuclear matter,” *Prog. Part. Nucl. Phys.* **2** (1979) 89–199.