NIRJAN BISWAS¹[©] AND ROHIT KUMAR²[©]

¹Department of Mathematics, Indian Institute of Science Education and Research Pune, Dr. Homi Bhabha Road, Pune 411008, India

² Department of Mathematics, Indian Institute of Technology Jodhpur,

Rajasthan 342030, India

ABSTRACT. In this article, for $N \ge 2, s \in (1, 2), p \in (1, \frac{N}{s}), \sigma = s - 1$ and $a \in [0, \frac{N-sp}{2})$, we establish an isometric isomorphism between the higher order fractional weighted Beppo-Levi space

$$\mathcal{D}_{a}^{s,p}(\mathbb{R}^{N}) := \overline{\mathcal{C}_{c}^{\infty}(\mathbb{R}^{N})}^{[\cdot]_{s,p,a}} \text{ where } [u]_{s,p,a} := \left(\iint_{\mathbb{R}^{N} \times \mathbb{R}^{N}} \frac{\left| \nabla u(x) - \nabla u(y) \right|^{p}}{\left| x - y \right|^{N + \sigma p}} \frac{\mathrm{d}x}{\left| x \right|^{a}} \frac{\mathrm{d}y}{\left| y \right|^{a}} \right)^{\frac{1}{p}},$$

and higher order fractional weighted homogeneous space

$$\mathring{W}_{a}^{s,p}(\mathbb{R}^{N}) := \left\{ u \in L_{a}^{p_{s}^{*}}(\mathbb{R}^{N}) : \left\| \nabla u \right\|_{L_{a}^{p_{\sigma}^{*}}(\mathbb{R}^{N})} + [u]_{s,p,a} < \infty \right\}$$

with the weighted Lebesgue norm

$$\|u\|_{L^{p^*}_{a}(\mathbb{R}^N)} := \left(\int_{\mathbb{R}^N} \frac{|u(x)|^{p^*}_{\alpha}}{|x|^{\frac{2ap^*}{\alpha}}} \, \mathrm{d}x \right)^{\frac{1}{p^*_{\alpha}}}, \text{ where } p^*_{\alpha} = \frac{Np}{N - \alpha p} \text{ for } \alpha = s, \sigma$$

To achieve this, we prove that $\mathcal{C}^{\infty}_{c}(\mathbb{R}^{N})$ is dense in $\mathring{W}^{s,p}_{a}(\mathbb{R}^{N})$ with respect to $[\cdot]_{s,p,a}$, and $[\cdot]_{s,p,a}$ is an equivalent norm on $\mathring{W}^{s,p}_{a}(\mathbb{R}^{N})$. Further, we obtain a finer embedding of $\mathcal{D}^{s,p}_{a}(\mathbb{R}^{N})$ into the Lorentz space $L^{\frac{Np}{N-sp-2a},p}(\mathbb{R}^{N})$, where $L^{\frac{Np}{N-sp-2a},p}(\mathbb{R}^{N}) \subsetneq L^{p^{*}}_{a}(\mathbb{R}^{N})$.

1. INTRODUCTION

For $a \ge 0, s = 1 + \sigma, \sigma \in (0, 1), p \in (1, \infty)$ and $N \ge 2$, we define the higher order fractional weighted-p Laplace operator of order s, up to a normalizing constant as

$$\langle (-\Delta)_{p,a}^{s} u, \varphi \rangle = \iint_{\mathbb{R}^{N} \times \mathbb{R}^{N}} \frac{|\nabla u(x) - \nabla u(y)|^{p-2} \left(\nabla u(x) - \nabla u(y)\right) \cdot \left(\nabla \varphi(x) - \nabla \varphi(y)\right)}{|x - y|^{N + \sigma p}} \frac{\mathrm{d}x}{|x|^{a}} \frac{\mathrm{d}y}{|y|^{a}},$$

for $u, \varphi \in \mathcal{C}^{\infty}_{c}(\mathbb{R}^{N})$. We define the following Gagliardo seminorm

$$[u]_{s,p,a} := \left(\iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{|\nabla u(x) - \nabla u(y)|^p}{|x - y|^{N + \sigma p}} \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a} \right)^{\frac{1}{p}}$$

For a > 0, we consider the higher order fractional weighted Sobolev space $W_a^{s,p}(\mathbb{R}^N)$ as

$$W_{a}^{s,p}(\mathbb{R}^{N}) := \left\{ u \text{ is measurable} : \|u\|_{p,a} + \|\nabla u\|_{p,a} + [u]_{s,p,a} < \infty \right\},$$
(1.1)

where

$$||u||_{p,a} := \left(\int_{\mathbb{R}^N} |u(x)|^p \frac{\mathrm{d}x}{|x|^a}\right)^{\frac{1}{p}} \text{ and } ||\nabla u||_{p,a} := \left(\int_{\mathbb{R}^N} |\nabla u(x)|^p \frac{\mathrm{d}x}{|x|^a}\right)^{\frac{1}{p}}.$$

E-mail address: nirjaniitm@gmail.com, nirjan.biswas@acads.iiserpune.ac.in, kumar.174@iitj.ac.in. 2020 Mathematics Subject Classification. 46E35 · 46E30.

Key words and phrases. higher order fractional weighted Sobolev spaces, density of smooth functions with compact support, equivalent norms, Lorentz-Sobolev embeddings.

¹Corresponding author.

For a = 0, the corresponding higher order fractional Sobolev space $W_0^{s,p}(\mathbb{R}^N)$ is defined in [14, Chapter 11]. We refer to [14] for more details on this space. Now we consider the following non-local elliptic problem

$$(-\Delta)_{p,a}^{s} u = f, \quad \text{in } \mathbb{R}^{N}, \tag{1.2}$$

under suitable assumptions on the source term f. For the existence of a weak solution to (1.2), an obvious choice is to apply the Direct Method in the Calculus of Variations. Consequently, the problem reduces to minimizing the associated energy functional \mathcal{J} given by

$$\mathcal{J}(u) = \frac{1}{p} \iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{|\nabla u(x) - \nabla u(y)|^p}{|x - y|^{N + \sigma p}} \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a} - \int_{\mathbb{R}^N} f(x)u(x) \,\mathrm{d}x.$$
(1.3)

Suppose, we consider the solution space for (1.2) as $W_a^{s,p}(\mathbb{R}^N)$ where $f \neq 0$ and $f \in (W_a^{s,p}(\mathbb{R}^N))^*$ (dual space of $W_a^{s,p}(\mathbb{R}^N)$). Observe that one can not have any control over $\|\cdot\|_{p,a}$ of minimizing sequences because the following weighted Poincaré type inequality

$$\int_{\mathbb{R}^N} |u(x)|^p \frac{\mathrm{d}x}{|x|^a} \le C[u]_{s,p,a}^p, \ \forall u \in W_a^{s,p}(\mathbb{R}^N),$$
(1.4)

does not hold for any C > 0. Suppose (1.4) holds for some $C_1 > 0$. For $\lambda > 0$ taking $u_{\lambda}(x) = u(\lambda x)$, we get

$$[u_{\lambda}]_{s,p,a}^{p} = \lambda^{2a-N+(\sigma+1)p} [u]_{s,p,a}^{p} \text{ and } \|u_{\lambda}\|_{p,a}^{p} = \lambda^{a-N} \|u\|_{p,a}^{p},$$

which in turn implies

$$\int_{\mathbb{R}^N} |u(x)|^p \frac{\mathrm{d}x}{|x|^a} \le C_1 \lambda^{a+(\sigma+1)p} [u]_{s,p,a}^p, \text{ for every } \lambda > 0.$$
(1.5)

Dividing (1.5) by $\lambda^{a+(\sigma+1)p}$ and taking $\lambda \to 0$ we get $[u]_{s,p,a} = \infty$. In a similar way, one can verify that the following inequality

$$\int_{\mathbb{R}^N} \frac{|\nabla u(x)|^p}{|x|^a} \, \mathrm{d}x \le C[u]_{s,p,a}^p, \ \forall u \in W_a^{s,p}(\mathbb{R}^N),$$

does not hold for any C > 0. This phenomenon conveys the fact that $W_a^{s,p}(\mathbb{R}^N)$ is not an appropriate function space to look for the weak solution for (1.2) as \mathcal{J} is not coercive on $W_a^{s,p}(\mathbb{R}^N)$.

The natural spaces to look for the existence of weak solutions for (1.2) are the higher order fractional weighted Beppo-Levi spaces $\mathcal{D}_{a}^{s,p}(\mathbb{R}^{N})$, which are defined by

$$\mathcal{D}_a^{s,p}(\mathbb{R}^N) := \text{closure of } \mathcal{C}_c^{\infty}(\mathbb{R}^N) \text{ with respect to } [\cdot]_{s,p,a}$$

The notation $\mathcal{D}_{a}^{s,p}(\mathbb{R}^{N})$ is reminiscent of the historical one introduced by Deny and Lions in their seminal paper [8], where the authors first studied homogeneous spaces obtained by the completion of $\mathcal{C}_{c}^{\infty}(\mathbb{R}^{N})$ with respect to a given seminorm. Notice that for $f \in (\mathcal{D}_{a}^{s,p}(\mathbb{R}^{N}))^{*}$, \mathcal{J} is well defined and

$$\mathcal{J}(u) \ge \frac{[u]_{s,p,a}^p}{p} - \|f\|_{(\mathcal{D}_a^{s,p}(\mathbb{R}^N))^*}[u]_{s,p,a}, \ \forall u \in \mathcal{D}_a^{s,p}(\mathbb{R}^N).$$

The above inequality conveys that \mathcal{J} is coercive on $\mathcal{D}_{a}^{s,p}(\mathbb{R}^{N})$. Hence, the existence of a solution for (1.2) can be assured in $\mathcal{D}_{a}^{s,p}(\mathbb{R}^{N})$ if one verifies other qualitative properties of \mathcal{J} ; for example, the weak lower semicontinuity of \mathcal{J} in $\mathcal{D}_{a}^{s,p}(\mathbb{R}^{N})$. The main focus of this article is to characterize $\mathcal{D}_{a}^{s,p}(\mathbb{R}^{N})$ as a function space. Verifying that a function $u \in \mathcal{D}_{a}^{s,p}(\mathbb{R}^{N})$ can be challenging, as it is not always feasible to find a sequence $\{u_{n}\} \subset \mathcal{C}_{c}^{\infty}(\mathbb{R}^{N})$ that converges to u with respect to $[\cdot]_{s,p,a}$. To address this difficulty, we study an equivalent characterization of $\mathcal{D}_{a}^{s,p}(\mathbb{R}^{N})$. For $s = 1 + \sigma, \sigma \in (0, 1), a \in [0, \frac{N-sp}{2})$ and $p \in (1, \frac{N}{s})$, we define the higher order fractional weighted homogeneous space $\mathring{W}_{a}^{s,p}(\mathbb{R}^{N})$ as

$$\mathring{W}_{a}^{s,p}(\mathbb{R}^{N}) := \left\{ u \in L_{a}^{p_{s}^{*}}(\mathbb{R}^{N}) : \|u\|_{\mathring{W}_{a}^{s,p}(\mathbb{R}^{N})} < \infty \right\},$$
(1.6)

where

$$||u||_{\mathring{W}^{s,p}_{a}(\mathbb{R}^{N})} := ||\nabla u||_{L^{p^{*}}_{a}(\mathbb{R}^{N})} + [u]_{s,p,a},$$

where the weighted Lebesgue space $L_a^{p^*_{\alpha}}(\mathbb{R}^N)$, for $\alpha = s, \sigma$, is given by

$$L_{a}^{p_{\alpha}^{*}}(\mathbb{R}^{N}) := \left\{ u \text{ is measurable} : \|u\|_{L_{a}^{p_{\alpha}^{*}}(\mathbb{R}^{N})} < \infty \right\}, \text{ where } \|u\|_{L_{a}^{p_{\alpha}^{*}}(\mathbb{R}^{N})}^{p_{\alpha}^{*}} := \int_{\mathbb{R}^{N}} \frac{|u(x)|^{p_{\alpha}^{*}}}{|x|^{\frac{2ap_{\alpha}^{*}}{p}}} \,\mathrm{d}x$$

and the fractional critical exponent p_{α}^* is defined as $p_{\alpha}^* := \frac{Np}{N-\alpha p}$. The name 'homogeneous' is due to the fact that

$$\|u_{\lambda}\|_{\mathring{W}^{s,p}_{a}(\mathbb{R}^{N})} = \|u\|_{\mathring{W}^{s,p}_{a}(\mathbb{R}^{N})}, \text{ for } u_{\lambda}(x) = \lambda^{\frac{N-sp-2a}{p}} u(\lambda x), \text{ where } \lambda > 0.$$

In the first part of this article, we aim to establish the existence of an isometric isomorphism between $\mathring{W}^{s,p}_{a}(\mathbb{R}^{N})$ and $\mathcal{D}^{s,p}_{a}(\mathbb{R}^{N})$. For that we investigate certain qualitative properties of $\mathring{W}^{s,p}_{a}(\mathbb{R}^{N})$, such as the density of $\mathcal{C}^{\infty}_{c}(\mathbb{R}^{N})$ in $\mathring{W}^{s,p}_{a}(\mathbb{R}^{N})$ and the equivalence between $[\cdot]_{s,p,a}$ and $\|\cdot\|_{\mathring{W}^{s,p}_{a}(\mathbb{R}^{N})}$.

For $s \in (0, 1]$, a similar characterization for the fractional Beppo Levi space is recently studied in [5]. For $a = 0, s \in (0, 1]$ and $p \in (1, \frac{N}{s})$, the fractional Beppo-Levi space $\mathcal{D}^{s,p}(\mathbb{R}^N)$ is defined as the completion of $\mathcal{C}_c^{\infty}(\mathbb{R}^N)$ with respect to the seminorms

$$\|u\|_{s,p} := \begin{cases} \left(\iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{|u(x) - u(y)|^p}{|x - y|^{N + sp}} \, \mathrm{d}x \mathrm{d}y\right)^{\frac{1}{p}}, & \text{for } s \in (0, 1)\\ \left(\int_{\mathbb{R}^N} |\nabla u|^p \, \mathrm{d}x\right)^{\frac{1}{p}}, & \text{for } s = 1. \end{cases}$$

From [17, Theorem 10.2.1] and [6, Theorem 9.9], the following Sobolev inequality holds

$$\|u\|_{L^{p_s^*}(\mathbb{R}^N)} \le C(N, p, s) \|u\|_{s, p}, \ \forall u \in \mathcal{C}^{\infty}_c(\mathbb{R}^N).$$

$$(1.7)$$

In [5, Theorem 3.1], using (1.7), Brasco et al. characterize the fractional Beppo-Levi space $\mathcal{D}^{s,p}(\mathbb{R}^N)$ as

$$\mathcal{D}^{s,p}(\mathbb{R}^N) = \left\{ u \in L^{p_s^*}(\mathbb{R}^N) : \|u\|_{s,p} < \infty \right\}$$

For $a \in (0, \frac{N-sp}{2})$ the following weighted Sobolev inequality holds (see [1, Theorem 1.5]):

$$\|u\|_{L^{p_{s}^{*}}_{a}(\mathbb{R}^{N})} \leq C(N, p, s, a) \|u\|_{s, p, a}, \ \forall u \in \mathcal{C}^{\infty}_{c}(\mathbb{R}^{N}),$$
(1.8)

where $\|\cdot\|_{s,p,a}$ is defined as

$$\|u\|_{s,p,a} := \left(\iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{|u(x) - u(y)|^p}{|x - y|^{N + sp}} \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a}\right)^{\frac{1}{p}}$$

Motivated by (1.8), the authors in [10] define $\mathring{W}^{s,p}_{a}(\mathbb{R}^{N})$ endowed with the following norm

$$||u||_{\mathring{W}^{s,p}_{a}(\mathbb{R}^{N})} := ||u||_{s,p,a} + ||u||_{L^{p^{*}}_{a}(\mathbb{R}^{N})}.$$

In [10, Theorem 1.1], the authors proved that $\mathcal{C}_c^{\infty}(\mathbb{R}^N)$ is dense in $\mathring{W}_a^{s,p}(\mathbb{R}^N)$. Clearly, this density result and (1.8) infer that $\|\cdot\|_{s,p,a}$ is an equivalent norm in $\mathring{W}_a^{s,p}(\mathbb{R}^N)$. The appearance of the space $\mathring{W}_a^{s,p}(\mathbb{R}^N)$ is seen in some literature; for example, Hardy-Lieb-Thirring inequalities (see [12]) and the non-local critical problems involving the Hardy-Leray potential (see [9]).

For $s \in (0,1)$, $a \in [0, \frac{N-sp}{2})$ and $p \in (1, \frac{N}{s})$, from [1, Theorem 2.7] we now recall the following weighted Hardy inequality

$$\int_{\mathbb{R}^N} \frac{|u(x)|^p}{|x|^{sp+2a}} \,\mathrm{d}x \le C \iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} \,\frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a}, \,\forall u \in \mathcal{C}^\infty_c(\mathbb{R}^N),\tag{1.9}$$

where C = C(N, p, s, a) > 0. For a = 0, (1.9) reduces to the classical Hardy inequality derived by Frank-Seiringer in [13] with the best constant. Further, using this inequality the authors in [13, Lemma 4.3] established an embedding of $\mathcal{D}^{s,p}(\mathbb{R}^N)$ into the Lorentz space $L^{p_s^*,p}(\mathbb{R}^N)$, namely $\|u\|_{p_s^*,p} \leq C(N,p,s)[u]_{s,p}$, for all $u \in \mathcal{D}_a^{s,p}(\mathbb{R}^N)$. This embedding is finer than the fractional Sobolev embedding $\mathcal{D}^{s,p}(\mathbb{R}^N) \hookrightarrow L^{p_s^*}(\mathbb{R}^N)$ due to the fact that $L^{p_s^*,p}(\mathbb{R}^N) \subsetneq L^{p_s^*,p_s^*}(\mathbb{R}^N) = L^{p_s^*}(\mathbb{R}^N)$. In the local case s = 1, the Lorentz Sobolev embeddings are studied for $W^{1,p}(\mathbb{R}^N)$ and $\mathcal{D}^{1,p}(\mathbb{R}^N)$. For example, the authors in [2,19,20] independently demonstrated that $W^{1,p}(\mathbb{R}^N) \hookrightarrow L^{p^*,p}(\mathbb{R}^N)$ where $p^* = \frac{Np}{N-p}$ for N > p. In [3], the authors showed that $\mathcal{D}^{1,p}(\mathbb{R}^N) \hookrightarrow L^{p^*,p}(\mathbb{R}^N)$. For $s \in (1,2)$, the corresponding fractional Lorentz Sobolev embedding is not known. However, for s = 2, in [18] the authors showed that the Beppo Levi space $\mathcal{D}^{2,2}(\mathbb{R}^N)$ (which is a completion of $\mathcal{C}^{\infty}_c(\mathbb{R}^N)$ with respect to $\|\Delta u\|_{L^2(\mathbb{R}^N)}$) is continuously embedded into the Lorentz space $L^{2^{**},2}(\mathbb{R}^N)$ where $2^{**} = \frac{2N}{N-4}$.

In the second part of this article, for $s \in (1,2)$, we establish an embedding $\mathcal{D}_a^{s,p}(\mathbb{R}^N)$ into the Lorentz space $L^{p^*,p}(\mathbb{R}^N)$, where $p^* = \frac{Np}{N-sp-2a}$. This embedding is finer since $L^{p^*,p}(\mathbb{R}^N) \subsetneq L_a^{p^*_s}(\mathbb{R}^N)$ (see Proposition 5.3). Moreover, p^* is an appropriate exponent in this case, since for s = p = 2 and $a = 0, p^*$ becomes $\frac{2N}{N-4}$. For this finer embedding, we derive a fractional weighted Rellich inequality in Proposition 5.2. For a = 0 and p = 2, a similar inequality has been obtained in [11, Corollary 7.5].

The following theorem collects the main results of this article.

Theorem 1.1. Let $N \ge 2, s \in (1,2), \sigma = s - 1, p \in (1, \frac{N}{s})$, and $a \in [0, \frac{N-sp}{2})$. Then the following hold:

(i) (Density of smooth functions) For every $u \in \mathring{W}^{s,p}_{a}(\mathbb{R}^{N})$, there exists $u_{n} \in \mathcal{C}^{\infty}_{c}(\mathbb{R}^{N})$ such that

$$[u-u_n]_{s,p,a} \to 0, as n \to \infty$$

namely $\mathcal{C}^{\infty}_{c}(\mathbb{R}^{N})$ is dense in $\mathring{W}^{s,p}_{a}(\mathbb{R}^{N})$ with respect to the seminorm $[\cdot]_{s,p,a}$.

(ii) (*Equivalent norm*) There exists $C = C(N, p, \sigma, a) > 0$ such that

$$\|\nabla u\|_{L^{p^*}_{a^{\sigma}}(\mathbb{R}^N)} \le C[u]_{s,p,a}, \ \forall u \in \mathring{W}^{s,p}_{a}(\mathbb{R}^N).$$

(iii) (*Characterization*) $\mathring{W}^{s,p}_{a}(\mathbb{R}^{N})$ is a Banach space with respect to the norm $[\cdot]_{s,p,a}$. Further, there exists an isometric isomorphism

$$\mathcal{J}: \mathcal{D}^{s,p}_a(\mathbb{R}^N) \to \mathring{W}^{s,p}_a(\mathbb{R}^N),$$

i.e., the space $\mathring{W}_{a}^{s,p}(\mathbb{R}^{N})$ is identified with $\mathcal{D}_{a}^{s,p}(\mathbb{R}^{N})$. (iv) (*Finer embeddings*) Denote $p^{\star} = \frac{Np}{N-sp-2a}$. Then

$$|u|_{p^{\star},p} \leq C(N,s,p,a)[u]_{s,p,a}, \ \forall u \in \mathcal{D}_{a}^{s,p}(\mathbb{R}^{N}),$$

$$(1.10)$$

i.e., $\mathcal{D}^{s,p}_a(\mathbb{R}^N) \hookrightarrow L^{p^{\star},p}(\mathbb{R}^N).$

The rest of the article is organized as follows. In Section 2, we state mollifier and cut-off functions and define Lorentz spaces, weak Young's inequality. In Section 3, we prove the approximations with smooth functions and norm equivalence in $\mathring{W}_{a}^{s,p}(\mathbb{R}^{N})$. Section 4 is devoted to study the characterization between $\mathring{W}_{a}^{s,p}(\mathbb{R}^{N})$ and $\mathcal{D}_{a}^{s,p}(\mathbb{R}^{N})$. In the final section, we prove fractional weighted Rellich inequality and finer embeddings of $\mathcal{D}_{a}^{s,p}(\mathbb{R}^{N})$.

2. Preliminaries

We begin by proving several basic inequalities to be used frequently in the subsequent sections. Next, we introduce the mollifier and cut-off functions essential for establishing the density results. Subsequently, we define the Lorentz spaces and state the weak Young's inequality. The following notations will be used throughout this paper.

Notation:

- (i) For $p \in [1, \infty]$, $L^p(\mathbb{R}^N)$ -norm of f is denoted as $\|f\|_p$.
- (ii) $B_r := B_r(0) \subset \mathbb{R}^N$ denotes a ball of radius r with centre at the origin.
- (iii) The convolution $f \star g$ is defined as $(f \star g)(x) = \int_{\mathbb{R}^N} f(x-z)g(z) \, \mathrm{d}z$.
- (iv) For ρ_n defined in (2.5), we denote

$$u_n = u \star \rho_n$$
 and $\frac{\partial u_n}{\partial x_i} = \frac{\partial u}{\partial x_i} \star \rho_n$,

for $i = 1, \cdot, \cdot, \cdot, N$.

- (v) For $A \subset \mathbb{R}^N$, it's complement $\mathbb{R}^N \setminus A$ is denoted as A^c .
- (vi) C denotes generic positive constant.

Lemma 2.1. Let $a_i \geq 0$, for $i = 1, 2, \cdot, \cdot, \cdot, N$ and $0 < q < \infty$. Then

(i) $(a_1 + a_2 + \dots + a_N)^q \le A(N,q) \left(a_1^q + a_2^q + \dots + a_N^q\right),$ (ii) $(a_1 + a_2 + \dots + a_N)^q \ge B(N,q) \left(a_1^q + a_2^q + \dots + a_N^q\right),$

where

$$A(N,q) = 1, \ B(N,q) = \begin{cases} \left(\frac{q}{2}\right)^{\frac{N+1}{2}}, & \text{if } N \text{ is odd} \\ \left(\frac{q}{2}\right)^{\frac{N}{2}}, & \text{if } N \text{ is even} \end{cases}, \text{ for } 0 < q < 1.$$

and

$$B(N,q) = 1, \ A(N,q) = \begin{cases} \left(2^{q-1}\right)^{\frac{N+1}{2}}, & \text{if } N \text{ is odd} \\ \left(2^{q-1}\right)^{\frac{N}{2}}, & \text{if } N \text{ is even} \end{cases}, \text{ for } 1 \le q < \infty.$$

Proof. (i) For 0 < q < 1: In this case, $(a_1 + a_2)^q \leq a_1^q + a_2^q$ and generalising this inequality we get

$$(a_1 + a_2 + \dots + a_N)^q \le (a_1^q + a_2^q + \dots + a_N^q)$$

For $1 \leq q < \infty$: In this case, using the convexity of $x \mapsto |x|^q$, we have $(a_1 + a_2)^q \leq 2^{q-1}(a_1^q + a_2^q)$ for any $a_1, a_2 \ge 0$. Using it, we find

$$(a_1 + a_2 + a_3)^q \le 2^{q-1}(a_1 + a_2)^q + 2^{q-1}a_3^q \le (2^{q-1})^2(a_1^q + a_2^q + a_3^q),$$

and

$$(a_1 + a_2 + a_3 + a_4)^q \le 2^{q-1}(a_1 + a_2)^q + 2^{q-1}(a_3 + a_4)^q \le (2^{q-1})^2 (a_1^q + a_2^q + a_3^q + a_4^q)$$

Hence, upon generalization,

$$(a_1 + a_2 + \dots + a_N)^q \le A(N,q) \left(a_1^q + a_2^q + \dots + a_N^q\right)$$

(ii) For 0 < q < 1: We first prove that

$$(a_1 + a_2)^q \ge \frac{q}{2}(a_1^q + a_2^q).$$
(2.1)

We define $f_1: [1,\infty) \to \mathbb{R}$ by $f_1(t) = (t+1)^q - \frac{q}{2}(t^q+1)$. Then $f_1(t) \ge (1-\frac{q}{2})t^q - \frac{q}{2} \ge 1 - \frac{q}{2} - \frac{q}{2} > 0$, which implies

$$(t+1)^q > \frac{q}{2}(t^q+1), \text{ for all } t \ge 1.$$
 (2.2)

Now (2.1) is trivial if either $a_1 = 0$ or $a_2 = 0$. So, we assume $a_1, a_2 \neq 0$. If $a_1 \ge a_2$, then for $t = \frac{a_1}{a_2}$ in (2.2) we get (2.1). If $a_1 \le a_2$, then for $t = \frac{a_2}{a_1}$ in (2.2) we get (2.1). From (2.1) we deduce that

$$(a_1 + a_2 + a_3)^q \ge \frac{q}{2}(a_1 + a_2)^q + \frac{q}{2}a_3^q \ge \left(\frac{q}{2}\right)^2(a_1^q + a_2^q + a_3^q),$$

and

$$(a_1 + a_2 + a_3 + a_4)^q \ge \frac{q}{2}(a_1 + a_2)^q + \frac{q}{2}(a_3 + a_4)^q \ge \left(\frac{q}{2}\right)^2(a_1^q + a_2^q + a_3^q + a_4^q).$$

Upon generalization, we obtain

$$(a_1 + a_2 + \dots + a_N)^q \ge B(N,q) \left(a_1^q + a_2^q + \dots + a_N^q\right)$$

For $1 \leq q < \infty$: We consider a function $f_2 : [1, \infty) \to \mathbb{R}$ by $f_2(t) = (t+1)^q - t^q - 1$. Then $f'_2(t) = q[(t+1)^{q-1} - t^{q-1}] \geq 0$. Thus, $f(t) \geq f(1)$ for all $t \geq 1$, which implies $(t+1)^q - t^q - 1 \geq 0$. $2^q - 2 \ge 0$ as $q \ge 1$. Thus,

$$(t+1)^q \ge t^q + 1$$
, for all $t \ge 1$. (2.3)

By substituting $t = \frac{a_1}{a_2}$ if $a_1 \ge a_2$ or $t = \frac{a_2}{a_1}$ if $a_1 \le a_2$ into (2.3), we get

$$(a_1 + a_2)^q \ge a_1^q + a_2^q. \tag{2.4}$$

Generalizing (2.4), $(a_1 + a_2 + \dots + a_N)^q \ge a_1^q + a_2^q + \dots + a_N^q$. **Definition 2.2** (Mollifier functions). Consider a radially symmetric and decreasing function $\rho \in C_c^{\infty}(\mathbb{R}^N)$ such that $\rho \geq 0$ on \mathbb{R}^N , $supp(\rho) \subset \overline{B_1}$ and $\int_{\mathbb{R}^N} \rho(x) dx = 1$. For each $n \in \mathbb{N}$, define

$$\rho_n(x) = n^N \rho(nx), \text{ for } x \in \mathbb{R}^N.$$
(2.5)

Then ρ_n satisfies

$$\rho_n \in \mathcal{C}^{\infty}_c(\mathbb{R}^N), \ supp(\rho_n) \subset \overline{B_{\frac{1}{n}}}, \ \int_{\mathbb{R}^N} \rho_n(x) \, dx = 1, \ \rho_n \ge 0 \ in \ \mathbb{R}^N$$

The sequence $\{\rho_n\}$ is called a sequence of mollifiers.

Definition 2.3 (Cut-off functions). Fix a function $\zeta \in \mathcal{C}^{\infty}_{c}(\mathbb{R}^{N})$ such that $0 \leq \zeta \leq 1$ and

$$\zeta(x) = \begin{cases} 1 & \text{if } |x| < 1\\ 0 & \text{if } |x| \ge 2 \end{cases}$$

For each $n \in \mathbb{N}$, define

$$\zeta_n(x) := \zeta\left(\frac{x}{n}\right). \tag{2.6}$$

The sequence $\{\zeta_n\}$ is called the cut-off function. Observe that

$$\|\nabla\zeta_n\|_{\infty} \le \frac{C}{n}, \text{ and } \|\nabla^2\zeta_n\|_{\infty} \le \frac{C}{n^2}, \tag{2.7}$$

where C = C(N) and ∇^2 represents the Hessian matrix.

Now, we recall the definition of Lorentz space introduced by Lorentz in [16]. For an open set $\Omega \subset \mathbb{R}^N$, let $M(\Omega)$ be the set of all extended real-valued Lebesgue measurable functions that are finite a.e. in Ω . For t > 0 and $f \in M(\Omega)$, define $E_f(t) := \{x \in \Omega : |f(x)| > t\}$. The distribution function μ_f of f is defined as $\mu_f(t) := |E_f(t)|$. For $(p,q) \in (0,\infty) \times (0,\infty]$ we consider the following quantity

$$|f|_{p,q} := \begin{cases} \left(p \int_0^\infty t^{q-1} \mu_f(t)^{\frac{q}{p}} \, \mathrm{d}t \right)^{\frac{1}{q}}, & q < \infty; \\ \sup_{t > 0} t \mu_f(t)^{\frac{1}{p}}, & q = \infty. \end{cases}$$

The Lorentz space $L^{p,q}(\Omega)$ is defined as

$$L^{p,q}(\Omega) := \{ f \in M(\Omega) : |f|_{p,q} < \infty \}$$

where $|f|_{p,q}$ is a quasi norm on $L^{p,q}(\Omega)$. Notice that $L^{p,p}(\mathbb{R}^N) = L^p(\mathbb{R}^N)$ and $L^{p,q_1}(\mathbb{R}^N) \subsetneq L^{p,q_2}(\mathbb{R}^N)$ provided $q_1 < q_2$ (see [11, Proposition 3.4.3]). For details about this space, we refer to [11].

Example 2.4. For d < N, consider the function $g(x) = |x|^{-d}$ where $x \in \mathbb{R}^N$. Using [4, Example 2.2.5] we get $g \in L^{\frac{N}{d},\infty}(\mathbb{R}^N)$.

In the following proposition, we state the weak Young's inequality. For more details and proofs of this inequality, we refer [15].

Proposition 2.5. Let $1 < p, q, r < \infty$ be such that $1 + \frac{1}{r} = \frac{1}{p} + \frac{1}{q}$. Let $f \in L^{p,\infty}(\mathbb{R}^N)$ and $h \in L^q(\mathbb{R}^N)$. Then $f \star h \in L^r(\mathbb{R}^N)$, and

$$||f \star h||_{r} \le C(N, p, q)|f|_{p,\infty} ||h||_{q}$$

3. Approximations by smooth functions with compact support

This section is devoted to proving the density of smooth functions in $\mathring{W}^{s,p}_{a}(\mathbb{R}^{N})$ and the equivalence of $[\cdot]_{s,p,a}$ with the $\mathring{W}^{s,p}_{a}(\mathbb{R}^{N})$ norm. Given a measurable function $w : \mathbb{R}^{2N} \to \mathbb{R}$, define

$$w * \rho_n(x, y) := \int_{\mathbb{R}^N} w(x - z, y - z) \rho_n(z) \, \mathrm{d}z$$

Then the following hold.

Proposition 3.1. (i) For every measurable function $w_1 : \mathbb{R}^{2N} \to \mathbb{R}$,

$$\iint_{\mathbb{R}^N \times \mathbb{R}^N} |w_1 * \rho_n(x, y)|^p \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a} \le C \iint_{\mathbb{R}^N \times \mathbb{R}^N} |w_1(x, y)|^p \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a}$$

where C is independent of n. (ii) Let $\alpha = s, \sigma$. For every measurable function $w_2 : \mathbb{R}^N \to \mathbb{R}$,

$$\int_{\mathbb{R}^N} \frac{|w_2 \star \rho_n(x)|^{p_\alpha^*}}{|x|^{\frac{2ap_\alpha^*}{p}}} \,\mathrm{d}x \le C \int_{\mathbb{R}^N} \frac{|w_2(x)|^{p_\alpha^*}}{|x|^{\frac{2ap_\alpha^*}{p}}} \,\mathrm{d}x,$$

where C is independent of n.

Proof. Proof follows using [10, Proposition 4.4 and Proposition 4.5] and the fact that for each $n \in \mathbb{N}$, supp $(\rho_n) \subset B_1$.

For $u \in \mathcal{C}^{\infty}_{c}(\mathbb{R}^{N})$, it follows from [10, Lemma 2.1] with $\frac{\partial u}{\partial x_{i}}$ that $[u]_{s,p,a} < \infty$. Moreover, using $a \in [0, \frac{N-sp}{2})$ and $\sigma < s$,

$$\begin{aligned} \|u\|_{L^{p^*_s}_{a^*}(\mathbb{R}^N)}^{p^*_s} &= \int_{\mathbb{R}^N} \frac{|u(x)|^{p^*_s}}{|x|^{\frac{2ap^*_s}{p}}} \, \mathrm{d}x \le \|u\|_{\infty}^{p^*_s} \int_{\mathrm{supp}(u)} \frac{\mathrm{d}x}{|x|^{\frac{2ap^*_s}{p}}} < \infty, \\ \|\nabla u\|_{L^{p^*_\sigma}_{a}(\mathbb{R}^N)}^{p^*_\sigma} &= \int_{\mathbb{R}^N} \frac{|\nabla u(x)|^{p^*_\sigma}}{|x|^{\frac{2ap^*_\sigma}{p}}} \, \mathrm{d}x \le \|\nabla u\|_{\infty}^{p^*_\sigma} \int_{\mathrm{supp}(u)} \frac{\mathrm{d}x}{|x|^{\frac{2ap^*_\sigma}{p}}} < \infty. \end{aligned}$$

Thus we observe that $\mathcal{C}^{\infty}_{c}(\mathbb{R}^{N}) \subset \mathring{W}^{s,p}_{a}(\mathbb{R}^{N}).$

Proposition 3.2. Let $N \ge 2, s \in (1,2), \sigma = s - 1, p \in (1, \frac{N}{s})$, and $a \in [0, \frac{N-sp}{2})$. Let ρ_n and ζ_n be defined as in (2.5) and (2.6) respectively. Then for every $u \in \mathring{W}_a^{s,p}(\mathbb{R}^N)$,

$$[u - (u \star \rho_n) \zeta_n]_{s,p,a} \to 0, \text{ as } n \to \infty.$$

Proof. For $u \in \mathring{W}^{s,p}_{a}(\mathbb{R}^{N})$ and $\rho_{n} \in \mathcal{C}^{\infty}_{c}(\mathbb{R}^{N})$, we have $u \star \rho_{n} \in \mathcal{C}^{\infty}(\mathbb{R}^{N})$ and $(u \star \rho_{n}) \zeta_{n} \in \mathcal{C}^{\infty}_{c}(\mathbb{R}^{N})$ (see [6, Proposition 4.20]). Using the triangle inequality, we write

$$[u - (u \star \rho_n) \zeta_n]_{s,p,a} \le [u - u \star \rho_n]_{s,p,a} + [u \star \rho_n - (u \star \rho_n) \zeta_n]_{s,p,a}.$$
(3.1)

We now estimate the first term of (3.1). Since $|\nabla u| \in L_a^{p^*_{\sigma}}(\mathbb{R}^N)$, interchange of differentiation and integration gives $\nabla(u \star \rho_n) = \nabla u \star \rho_n \in (L_a^{p^*_{\sigma}}(\mathbb{R}^N))^N$ (applying Proposition 3.1-(ii) with $\frac{\partial u}{\partial x_i}$). Using Lemma 2.1-(i), we estimate

$$\begin{aligned} \left[u - u \star \rho_n\right]_{s,p,a}^p &= \iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{\left|\left(\nabla u(x) - \nabla u(y)\right) - \left(\left(\nabla u \star \rho_n\right)(x) - \left(\nabla u \star \rho_n\right)(y)\right)\right|^p}{|x - y|^{N + \sigma p}} \frac{dx}{|x|^a} \frac{dy}{|y|^a} \\ &= \iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{\left(\sum_{i=1}^N \left(\frac{\partial u}{\partial x_i}(x) - \frac{\partial u}{\partial x_i}(y) - \left(\frac{\partial u_n}{\partial x_i}(x) - \frac{\partial u_n}{\partial x_i}(y)\right)\right)^2\right)^{\frac{p}{2}}}{|x - y|^{N + \sigma p}} \frac{dx}{|x|^a} \frac{dy}{|y|^a} \\ &\leq C(N, p) \iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{\sum_{i=1}^N \left|\frac{\partial u}{\partial x_i}(x) - \frac{\partial u}{\partial x_i}(y) - \left(\frac{\partial u_n}{\partial x_i}(x) - \frac{\partial u_n}{\partial x_i}(y)\right)\right|^p}{|x - y|^{N + \sigma p}} \frac{dx}{|x|^a} \frac{dy}{|y|^a} \\ &= C(N, p) \sum_{i=1}^N \iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{\left|\frac{\partial u}{\partial x_i}(x) - \frac{\partial u}{\partial x_i}(y) - \left(\frac{\partial u_n}{\partial x_i}(x) - \frac{\partial u_n}{\partial x_i}(y)\right)\right|^p}{|x - y|^{N + \sigma p}} \frac{dx}{|x|^a} \frac{dy}{|y|^a}. \end{aligned}$$
(3.2)

Further, using Lemma 2.1-(ii), we find

$$[u]_{s,p,a}^{p} = \iint_{\mathbb{R}^{N} \times \mathbb{R}^{N}} \frac{\left|\nabla u(x) - \nabla u(y)\right|^{p}}{\left|x - y\right|^{N + \sigma p}} \frac{\mathrm{d}x}{\left|x\right|^{a}} \frac{\mathrm{d}y}{\left|y\right|^{a}} \ge C(N,p) \sum_{i=1}^{N} \iint_{\mathbb{R}^{N} \times \mathbb{R}^{N}} \frac{\left|\frac{\partial u}{\partial x_{i}}(x) - \frac{\partial u}{\partial x_{i}}(y)\right|^{p}}{\left|x - y\right|^{N + \sigma p}} \frac{\mathrm{d}x}{\left|x\right|^{a}} \frac{\mathrm{d}y}{\left|y\right|^{a}},$$
which implies that for each $i = 1, \dots, N$

which implies that for each $i = 1, \cdot, \cdot, \cdot, N$,

$$\iint_{\mathbb{R}^N \times \mathbb{R}^N} |w_i(x,y)|^p \, \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a} < \infty,\tag{3.3}$$

where

$$w_i(x,y) = \frac{\frac{\partial u}{\partial x_i}(x) - \frac{\partial u}{\partial x_i}(y)}{|x-y|^{\frac{N}{p}+\sigma}}, \text{ for } x, y \in \mathbb{R}^N.$$
(3.4)

For each $i = 1, \cdot, \cdot, \cdot, N$, now we apply [10, Lemma 6.1] with $\frac{\partial u}{\partial x_i}$ to obtain

$$\lim_{n \to \infty} \iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{\left| \frac{\partial u}{\partial x_i}(x) - \frac{\partial u}{\partial x_i}(y) - \left(\frac{\partial u_n}{\partial x_i}(x) - \frac{\partial u_n}{\partial x_i}(y) \right) \right|^p}{|x - y|^{N + \sigma p}} \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a} = 0.$$
(3.5)

Combining (3.2) and (3.5), we deduce that

$$\lim_{n \to \infty} \left[u - u \star \rho_n \right]_{s,p,a} = 0. \tag{3.6}$$

Next, we estimate the second term on the right-hand side of the inequality (3.1). Define $\varphi_n = 1 - \zeta_n$. Using $\nabla ((u \star \rho_n) \varphi_n) = (\nabla u \star \rho_n) \varphi_n + (u \star \rho_n) \nabla \varphi_n$ and Lemma 2.1-(i), we derive the following estimate:

$$\begin{split} &[(u \star \rho_{n}) \varphi_{n}]_{s,p,a}^{p} \\ \leq C \sum_{i=1}^{N} \iint_{\mathbb{R}^{N} \times \mathbb{R}^{N}} \frac{\left| \frac{\partial u_{n}}{\partial x_{i}}(x)\varphi_{n}(x) - \frac{\partial u_{n}}{\partial x_{i}}(y)\varphi_{n}(y) + u_{n}(x)\frac{\partial \varphi_{n}}{\partial x_{i}}(x) - u_{n}(y)\frac{\partial \varphi_{n}}{\partial x_{i}}(y) \right|^{p}}{|x|^{a}} \frac{dx}{|y|^{a}} \frac{dy}{|x|^{a}} \\ \leq C \left(\sum_{i=1}^{N} \iint_{\mathbb{R}^{N} \times \mathbb{R}^{N}} \frac{\left| \frac{\partial u_{n}}{\partial x_{i}}(x)\varphi_{n}(x) - \frac{\partial u_{n}}{\partial x_{i}}(y)\varphi_{n}(y) \right|^{p}}{|x-y|^{N+\sigma p}} \frac{dx}{|x|^{a}} \frac{dy}{|y|^{a}} \\ &+ \sum_{i=1}^{N} \iint_{\mathbb{R}^{N} \times \mathbb{R}^{N}} \frac{\left| u_{n}(x)\frac{\partial \varphi_{n}}{\partial x_{i}}(x) - u_{n}(y)\frac{\partial \varphi_{n}}{\partial x_{i}}(y) \right|^{p}}{|x-y|^{N+\sigma p}} \frac{dx}{|x|^{a}} \frac{dy}{|y|^{a}} \end{split}$$
(3.7)

where C = C(N, p). Further, using Proposition 3.1-(i) and (3.3), we have

$$\begin{bmatrix} \frac{\partial u_n}{\partial x_i} \end{bmatrix}_{\sigma,p,a}^p = \iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{\left| \left(\frac{\partial u}{\partial x_i} \star \rho_n \right) (x) - \left(\frac{\partial u}{\partial x_i} \star \rho_n \right) (y) \right|^p}{|x - y|^{N + \sigma p}} \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a}$$

$$= \iint_{\mathbb{R}^N \times \mathbb{R}^N} \left| \int_{\mathbb{R}^N} \frac{\left(\frac{\partial u}{\partial x_i} (x - z) - \frac{\partial u}{\partial x_i} (y - z) \right) \rho_n(z) \, \mathrm{d}z}{|(x - z) - (y - z)|^{\frac{N}{p} + \sigma}} \right|^p \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a}$$

$$= \iint_{\mathbb{R}^N \times \mathbb{R}^N} |w_i \star \rho_n(x, y)|^p \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a} \le C \iint_{\mathbb{R}^N \times \mathbb{R}^N} |w_i(x, y)|^p \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a} \le C, \qquad (3.8)$$

where C does not depend on n. Since $\varphi_n(x) = 0 = \varphi_n(y)$ for $x, y \in B_n$, we write

$$\iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{\left|\frac{\partial u_n}{\partial x_i}(x)\varphi_n(x) - \frac{\partial u_n}{\partial x_i}(y)\varphi_n(y)\right|^p}{|x-y|^{N+\sigma p}} \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a} \le C(J_{n,i} + K_{n,i}),\tag{3.9}$$

where

$$J_{n,i} := \iint_{\mathbb{R}^N \times B_n^c} \frac{\left|\frac{\partial u_n}{\partial x_i}(x) - \frac{\partial u_n}{\partial x_i}(y)\right|^p}{|x - y|^{N + \sigma p}} |\varphi_n(x)|^p \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a}$$

and $K_{n,i} := \iint_{\mathbb{R}^N \times B_n^c} \frac{|\varphi_n(x) - \varphi_n(y)|^p}{|x - y|^{N + \sigma p}} \left|\frac{\partial u_n}{\partial x_i}(y)\right|^p \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a}.$

Further, we define

$$D_{n,0} := \left\{ (x,y) \in \mathbb{R}^N \times B_n^c \text{ s.t. } |x| \le \frac{|y|}{2} \right\},$$

$$D_{n,1} := \left\{ (x,y) \in \mathbb{R}^N \times B_n^c \text{ s.t. } |x| > \frac{|y|}{2} \text{ and } |x-y| \ge n \right\},$$

$$D_{n,2} := \left\{ (x,y) \in \mathbb{R}^N \times B_n^c \text{ s.t. } |x| > \frac{|y|}{2} \text{ and } |x-y| < n \right\}.$$

For $j \in \{0, 1, 2\}$, we define

$$K_{n,i,j} := \iint_{D_{n,j}} \frac{|\varphi_n(x) - \varphi_n(y)|^p}{|x - y|^{N + \sigma p}} \left| \frac{\partial u_n}{\partial x_i}(y) \right|^p \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a}.$$

Observe that

$$K_{n,i} = K_{n,i,0} + K_{n,i,1} + K_{n,i,2}.$$
(3.10)

From the arguments as in the proof of [10, Lemma 3.1], for each $j \in \{0, 1, 2\}$ we obtain

$$K_{n,i,j} \le C \left\| \frac{\partial u_n}{\partial x_i} \right\|_{L^{p^*}_a(B^c_n)}^{\frac{N-\sigma p}{N}},$$

where C does not depend on n, and hence

$$K_{n,i,j} \leq C\left(\left\|\frac{\partial u_n}{\partial x_i} - \frac{\partial u}{\partial x_i}\right\|_{L_a^{p^{\sigma}}(B_n^c)}^{\frac{N-\sigma p}{N}} + \left\|\frac{\partial u}{\partial x_i}\right\|_{L_a^{p^{\sigma}}(B_n^c)}^{\frac{N-\sigma p}{N}}\right)$$
$$\leq C\left(\left\|\frac{\partial u_n}{\partial x_i} - \frac{\partial u}{\partial x_i}\right\|_{L_a^{p^{\sigma}}(\mathbb{R}^N)}^{\frac{N-\sigma p}{N}} + \left\|\frac{\partial u}{\partial x_i}\right\|_{L_a^{p^{\sigma}}(B_n^c)}^{\frac{N-\sigma p}{N}}\right).$$
(3.11)

Since $|\nabla u| \in L^{p^*_{\sigma}}_{a}(\mathbb{R}^N)$, using Proposition 3.1-(ii) we have $\frac{\partial u_n}{\partial x_i} \in L^{p^*_{\sigma}}_{a}(\mathbb{R}^N)$. As $n \to \infty$, the first term of (3.11) tends to zero using [10, Lemma 6.1] with $\frac{\partial u}{\partial x_i}$, and the second term of (3.11) tends to zero using $\frac{\partial u}{\partial x_i} \in L^{p^*_{\sigma}}_{a}(\mathbb{R}^N)$. Hence

$$\lim_{n \to \infty} K_{n,i} = 0. \tag{3.12}$$

Using $0 \leq \varphi_n \leq 1$ and the triangle inequality, we have

$$J_{n,i} \leq C(p) \bigg(\iint_{\mathbb{R}^N \times B_n^c} \frac{\left| \frac{\partial u_n}{\partial x_i}(x) - \frac{\partial u_n}{\partial x_i}(y) - \left(\frac{\partial u}{\partial x_i}(x) - \frac{\partial u}{\partial x_i}(y) \right) \right|^p}{|x - y|^{N + \sigma p}} \frac{dx}{|x|^a} \frac{dy}{|y|^a} \\ + \iint_{\mathbb{R}^N \times B_n^c} \frac{\left| \frac{\partial u}{\partial x_i}(x) - \frac{\partial u}{\partial x_i}(y) \right|^p}{|x - y|^{N + \sigma p}} \frac{dx}{|x|^a} \frac{dy}{|y|^a} \bigg) \\ \leq C(p) \bigg(\iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{\left| \frac{\partial u_n}{\partial x_i}(x) - \frac{\partial u_n}{\partial x_i}(y) - \left(\frac{\partial u}{\partial x_i}(x) - \frac{\partial u}{\partial x_i}(y) \right) \right|^p}{|x - y|^{N + \sigma p}} \frac{dx}{|x|^a} \frac{dy}{|y|^a} \\ + \iint_{\mathbb{R}^N \times B_n^c} \frac{\left| \frac{\partial u}{\partial x_i}(x) - \frac{\partial u}{\partial x_i}(y) \right|^p}{|x - y|^{N + \sigma p}} \frac{dx}{|x|^a} \frac{dy}{|y|^a} \bigg).$$
(3.13)

As $n \to \infty$, the first integral of (3.13) tends to zero using (3.5), and the second integral of (3.13) tends to zero using

$$\iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{\left| \frac{\partial u}{\partial x_i}(x) - \frac{\partial u}{\partial x_i}(y) \right|^p}{|x - y|^{N + \sigma p}} \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a} < \infty.$$

Therefore,

$$\lim_{n \to \infty} J_{n,i} = 0. \tag{3.14}$$

From (3.9), (3.12) and (3.14), we deduce that

$$\lim_{n \to \infty} \iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{\left| \frac{\partial u_n}{\partial x_i}(x)\varphi_n(x) - \frac{\partial u_n}{\partial x_i}(y)\varphi_n(y) \right|^p}{|x - y|^{N + \sigma p}} \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a} = 0.$$
(3.15)

We now turn our attention to estimate the final term of (3.7). Denote $\psi_{n,i} := \frac{\partial \varphi_n}{\partial x_i}$. By noticing that $\operatorname{supp}(\psi_{n,i}) \subset \overline{B_{2n} \setminus B_n}$ (since $\operatorname{supp}(\varphi_n) \subset B_n^c$ and $\varphi_n = 1$ in B_{2n}^c), for each *i*, we split the following integral

$$I_{n,i} := \iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{|u_n(x)\psi_{n,i}(x) - u_n(y)\psi_{n,i}(y)|^p}{|x - y|^{N + \sigma p}} \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a}$$
$$= 2I_{n,i,1} + 2I_{n,i,2} + 2I_{n,i,3} + 2I_{n,i,4} + I_{n,i,5}, \tag{3.16}$$

where

$$\begin{split} I_{n,i,1} &:= \iint_{B_n \times (B_{2n} \setminus B_n)} \frac{|u_n(x)\psi_{n,i}(x) - u_n(y)\psi_{n,i}(y)|^p}{|x - y|^{N + \sigma p}} \frac{dx}{|x|^a} \frac{dy}{|y|^a} \\ &= \iint_{B_n \times (B_{2n} \setminus B_n)} \frac{|u_n(y)\psi_{n,i}(y)|^p}{|x - y|^{N + \sigma p}} \frac{dx}{|x|^a} \frac{dy}{|y|^a}, \\ I_{n,i,2} &:= \iint_{B_n \times B_{2n}^c} \frac{|u_n(x)\psi_{n,i}(x) - u_n(y)\psi_{n,i}(y)|^p}{|x - y|^{N + \sigma p}} \frac{dx}{|x|^a} \frac{dy}{|y|^a} = 0, \\ I_{n,i,3} &:= \iint_{(B_{2n} \setminus B_n) \times (B_{2n} \setminus B_n)} \frac{|u_n(x)\psi_{n,i}(x) - u_n(y)\psi_{n,i}(y)|^p}{|x - y|^{N + \sigma p}} \frac{dx}{|x|^a} \frac{dy}{|y|^a}, \\ I_{n,i,4} &:= \iint_{(B_{2n} \setminus B_n) \times B_{2n}^c} \frac{|u_n(x)\psi_{n,i}(x) - u_n(y)\psi_{n,i}(y)|^p}{|x - y|^{N + \sigma p}} \frac{dx}{|x|^a} \frac{dy}{|y|^a}, \\ I_{n,i,5} &:= \iint_{B_{2n}^c \times B_{2n}^c} \frac{|u_n(x)\psi_{n,i}(x) - u_n(y)\psi_{n,i}(y)|^p}{|x - y|^{N + \sigma p}} \frac{dx}{|x|^a} \frac{dy}{|y|^a} = 0. \end{split}$$

Estimation of $I_{n,i,1}$: We infer from (2.7) that $\|\nabla^2 \varphi_n\|_{\infty} \leq \frac{C(N)}{n^2}$. Therefore, applying the mean value theorem,

$$I_{n,i,1} = \iint_{B_n \times (B_{2n} \setminus B_n)} \frac{|u_n(y)|^p |\psi_{n,i}(x) - \psi_{n,i}(y)|^p}{|x - y|^{N + \sigma p}} \frac{dx}{|x|^a} \frac{dy}{|y|^a}$$

$$\leq \frac{C(N,p)}{n^{2p}} \iint_{B_n \times (B_{2n} \setminus B_n)} \frac{|u_n(y)|^p}{|x - y|^{N + \sigma p - p}} \frac{dx}{|x|^a} \frac{dy}{|y|^a}$$

$$\leq \frac{C(N,p)}{n^{2p}} \left\{ \iint_{B_{\frac{n}{2}} \times (B_{2n} \setminus B_n)} \frac{|u_n(y)|^p}{|x - y|^{N + \sigma p - p}} \frac{dx}{|x|^a} \frac{dy}{|y|^a} + \iint_{(B_n \setminus B_{\frac{n}{2}}) \times (B_{2n} \setminus B_n)} \frac{|u_n(y)|^p}{|x - y|^{N + \sigma p - p}} \frac{dx}{|x|^a} \frac{dy}{|y|^a} \right\}.$$
(3.17)

Notice that for $x \in B_{\frac{n}{2}}$ and $y \in B_{2n} \setminus B_n$, we have |y| > |x| and

$$|x-y| \ge |y| - |x| = \frac{|y|}{2} + \frac{|y|}{2} - |x| \ge \frac{|y|}{2}.$$

Thus, we obtain

$$\begin{aligned} \iint_{B_{\frac{n}{2}} \times (B_{2n} \setminus B_{n})} \frac{|u_{n}(y)|^{p}}{|x-y|^{N+\sigma p-p}} \frac{dx}{|x|^{a}} \frac{dy}{|y|^{a}} &\leq C \iint_{B_{\frac{n}{2}} \times (B_{2n} \setminus B_{n})} \frac{|u_{n}(y)|^{p}}{|y|^{N+\sigma p-p}} \frac{dxdy}{|x|^{2a}} \\ &\leq C \left(\int_{B_{\frac{n}{2}}} \frac{dx}{|x|^{2a}} \right) \left(\int_{B_{2n} \setminus B_{n}} \frac{|u_{n}(y)|^{p}}{|y|^{N+\sigma p-p}} dy \right) \\ &\leq C n^{N-2a} \left(\int_{B_{2n} \setminus B_{n}} \frac{|u_{n}(y)|^{p}}{|y|^{\frac{2ap_{s}^{*}}{p}}} dy \right)^{\frac{p}{p_{s}^{*}}} \left(\int_{B_{2n} \setminus B_{n}} \left(\frac{|y|^{2a}}{|y|^{N+\sigma p-p}} \right)^{\frac{N}{sp}} dy \right)^{\frac{sp}{N}} \text{ (by Hölder's inequality)} \\ &\leq C n^{N-2a} \frac{n^{2a}}{n^{N+\sigma p-p}} \left(\int_{B_{2n} \setminus B_{n}} \frac{|u_{n}(y)|^{p_{s}^{*}}}{|y|^{\frac{2ap_{s}^{*}}{p}}} dy \right)^{\frac{p}{p_{s}^{*}}} \left(\int_{B_{2n} \setminus B_{n}} \frac{|u_{n}(y)|^{p_{s}^{*}}}{|y|^{\frac{2ap_{s}^{*}}{p}}} dy \right)^{\frac{p}{p_{s}^{*}}} \left(\int_{B_{2n} \setminus B_{n}} \frac{|u_{n}(y)|^{p_{s}^{*}}}{|y|^{\frac{2ap_{s}^{*}}{p}}} dy \right)^{\frac{p}{p_{s}^{*}}} (\int_{B_{2n} \setminus B_{n}} \frac{|u_{n}(y)|^{p_{s}^{*}}}{|y|^{\frac{2ap_{s}^{*}}{p}}} dy \right)^{\frac{p}{p_{s}^{*}}} (\int_{B_{2n} \setminus B_{n}} \frac{|u_{n}(y)|^{p_{s}^{*}}}{|y|^{\frac{2ap_{s}^{*}}{p}}} dy \right)^{\frac{p}{p_{s}^{*}}} (\int_{B_{2n} \setminus B_{n}} \frac{|u_{n}(y)|^{p_{s}^{*}}}{|y|^{\frac{2ap_{s}^{*}}{p}}} dy \right)^{\frac{p}{p_{s}^{*}}} (3.18) \end{aligned}$$

where $C = C(N, p, \sigma, a)$. Further, notice that for $x \in B_n \setminus B_{\frac{n}{2}}$ and $y \in B_{2n} \setminus B_n$, we have $|y| \ge |x|$ and $B_n \setminus B_{\frac{n}{2}} \subset B_{4n}(y)$. Hence we estimate

$$\begin{aligned} \iint_{(B_n \setminus B_{\frac{n}{2}}) \times (B_{2n} \setminus B_n)} \frac{|u_n(y)|^p}{|x - y|^{N + \sigma p - p}} \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a} &\leq \iint_{(B_n \setminus B_{\frac{n}{2}}) \times (B_{2n} \setminus B_n)} \frac{|u_n(y)|^p}{|x - y|^{N + \sigma p - p}} \frac{\mathrm{d}x\mathrm{d}y}{|x|^{2a}} \\ &\leq \frac{C}{n^{2a}} \int_{B_{2n} \setminus B_n} |u_n(y)|^p \left(\int_{B_n \setminus B_{\frac{n}{2}}} \frac{\mathrm{d}x}{|x - y|^{N + \sigma p - p}} \right) \mathrm{d}y \\ &\leq \frac{C}{n^{2a}} \int_{B_{2n} \setminus B_n} |u_n(y)|^p \left(\int_{B_{4n}(y)} \frac{\mathrm{d}x}{|x - y|^{N + \sigma p - p}} \right) \mathrm{d}y \\ &\leq C \frac{n^{p - \sigma p}}{n^{2a}} \int_{B_{2n} \setminus B_n} \frac{|u_n(y)|^{p^*}}{|y|^{2a}} \mathrm{d}y \\ &\leq C \frac{n^{p - \sigma p}}{n^{2a}} \left(\int_{B_{2n} \setminus B_n} \frac{|u_n(y)|^{p^*}}{|y|^{2m^*}} \mathrm{d}y \right)^{\frac{p}{p^*_s}} \left(\int_{B_{2n} \setminus B_n} |y|^{\frac{2an^*}{sp}} \mathrm{d}y \right)^{\frac{p}{p^*_s}} (by \text{ Hölder's inequality}) \\ &\leq C \frac{n^{p - \sigma p + 2a + sp}}{n^{2a}} \left(\int_{B_{2n} \setminus B_n} \frac{|u_n(y)|^{p^*_s}}{|y|^{\frac{2an^*_s}{p}}} \mathrm{d}y \right)^{\frac{p}{p^*_s}} = Cn^{2p} \left(\int_{B_{2n} \setminus B_n} \frac{|u_n(y)|^{p^*_s}}{|y|^{\frac{2an^*_s}{p}}} \mathrm{d}y \right)^{\frac{p}{p^*_s}}, \quad (3.19) \end{aligned}$$

where C = C(N, p, s, a). Hence

$$\iint_{(B_n \setminus B_{\frac{n}{2}}) \times (B_{2n} \setminus B_n)} \frac{|u_n(y)|^p}{|x-y|^{N+\sigma p-p}} \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a} \le C n^{2p} \left(\int_{B_{2n} \setminus B_n} \frac{|u_n(y)|^{p_s^*}}{|y|^{\frac{2ap_s^*}{p}}} \mathrm{d}y \right)^{\frac{p}{p_s^*}}.$$
 (3.20)

From (3.17), (3.18) and (3.20), we get the following inequality

$$I_{n,i,1} \le C(N, p, \sigma, a) \left(\int_{B_{2n} \setminus B_n} \frac{|u_n(y)|^{p_s^*}}{|y|^{\frac{2ap_s^*}{p}}} \,\mathrm{d}y \right)^{\frac{p}{p_s^*}}.$$

Using the arguments as in [10, Lemma 6.1], we have

$$\lim_{n \to \infty} \|u - u_n\|_{L^{p^*}_a(\mathbb{R}^N)} = 0.$$
(3.21)

Thus,

$$\begin{split} \int_{B_{2n}\setminus B_n} \frac{|u_n(y)|^{p_s^*}}{|y|^{\frac{2ap_s^*}{p}}} \,\mathrm{d}y &\leq C \int_{B_{2n}\setminus B_n} \frac{|u_n(y) - u(y)|^{p_s^*}}{|y|^{\frac{2ap_s^*}{p}}} \,\mathrm{d}y + C \int_{B_{2n}\setminus B_n} \frac{|u(y)|^{p_s^*}}{|y|^{\frac{2ap_s^*}{p}}} \,\mathrm{d}y \\ &\leq C \int_{\mathbb{R}^N} \frac{|u(y) - u_n(y)|^{p_s^*}}{|y|^{\frac{2ap_s^*}{p}}} \,\mathrm{d}y + C \int_{B_n^c} \frac{|u(y)|^{p_s^*}}{|y|^{\frac{2ap_s^*}{p}}} \,\mathrm{d}y. \end{split}$$

The first integral on the right-hand side of the above inequality converges to 0 as $n \to \infty$ due to (3.21) and the second integral also converges to 0 as $n \to \infty$ by dominated convergence theorem. Hence, we conclude that $\lim_{n\to\infty} I_{n,i,1} = 0$. **Estimation of** $I_{n,i,4}$: Using the fact that $\psi_{n,i} = 0$ in B_{2n}^c , we write

$$I_{n,i,4} = \iint_{(B_{2n} \setminus B_n) \times (B_{3n}^c)} \frac{|u_n(x)\psi_{n,i}(x)|^p}{|x-y|^{N+\sigma p}} \frac{dx}{|x|^a} \frac{dy}{|y|^a} \\ + \iint_{(B_{2n} \setminus B_n) \times (B_{3n} \setminus B_{2n})} \frac{|u_n(x)|^p}{|x-y|^{N+\sigma p}} \frac{|u_n(x)|^p}{|x-y|^{N+\sigma p}} \frac{dx}{|x|^a} \frac{dy}{|y|^a} \\ \leq \underbrace{\frac{C}{n^p} \iint_{(B_{2n} \setminus B_n) \times (B_{3n}^c)} \frac{|u_n(x)|^p}{|x-y|^{N+\sigma p}} \frac{dx}{|x|^a} \frac{dy}{|y|^a}}_{A_{n,i}} + \underbrace{\frac{C}{n^{2p}} \iint_{(B_{2n} \setminus B_n) \times (B_{3n} \setminus B_{2n})} \frac{|u_n(x)|^p}{|x-y|^{N+\sigma p-p}} \frac{dx}{|x|^a} \frac{dy}{|y|^a}}_{B_{n,i}},$$

where C = C(N, p). For $A_{n,i}$, we use $\|\nabla \varphi_n\|_{\infty} \leq \frac{C(N)}{n}$, and $B_{n,i}$ follows using the mean value theorem and $\|\nabla^2 \varphi_n\|_{\infty} \leq \frac{C(N)}{n^2}$. Notice that $B_n(x) \subset B_{3n}$ for $x \in B_{2n} \setminus B_n$. Thus, we get

$$A_{n,i} = \frac{C(N,p)}{n^p} \int_{B_{2n} \setminus B_n} |u_n(x)|^p \left(\int_{B_{3n}^c} \frac{\mathrm{d}y}{|x-y|^{N+\sigma p}|y|^a} \right) \frac{\mathrm{d}x}{|x|^a}$$

$$\leq \frac{C(N,p)}{n^{p+a}} \int_{B_{2n} \setminus B_n} |u_n(x)|^p \left(\int_{B_n^c(x)} \frac{\mathrm{d}y}{|x-y|^{N+\sigma p}} \right) \frac{\mathrm{d}x}{|x|^a}$$

$$\leq \frac{C(N,p,\sigma)}{n^{p+\sigma p+a}} \int_{B_{2n} \setminus B_n} |u_n(x)|^p \frac{\mathrm{d}x}{|x|^a}.$$
(3.22)

Next, we estimate the second integral, i.e.,

$$B_{n,i} = \frac{C(N,p)}{n^{2p}} \iint_{(B_{2n}\setminus B_n)\times(B_{3n}\setminus B_{2n})} \frac{|u_n(x)|^p}{|x-y|^{N+\sigma p-p}} \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a}$$
$$\leq \frac{C(N,p)}{n^{2p+a}} \int_{B_{2n}\setminus B_n} |u_n(x)|^p \left(\int_{B_{3n}\setminus B_{2n}} \frac{\mathrm{d}y}{|x-y|^{N+\sigma p-p}}\right) \frac{\mathrm{d}x}{|x|^a}.$$

Observe that $B_{3n} \subset B_{6n}(x)$ for $x \in B_{2n} \setminus B_n$, and hence

$$\int_{B_{3n}\setminus B_{2n}} \frac{\mathrm{d}y}{|x-y|^{N+\sigma p-p}} \le \int_{B_{6n}(x)} \frac{\mathrm{d}y}{|x-y|^{N+\sigma p-p}} \le C(N,p,\sigma) n^{p-\sigma p}$$

Thus, we obtain

$$B_{n,i} \le \frac{C(N, p, \sigma)}{n^{p+\sigma p+a}} \int_{B_{2n} \setminus B_n} |u_n(x)|^p \frac{\mathrm{d}x}{|x|^a}.$$
(3.23)

Further, using Hölder's inequality, we estimate

$$\int_{B_{2n}\setminus B_n} |u_n(x)|^p \frac{\mathrm{d}x}{|x|^a} = \int_{B_{2n}\setminus B_n} \frac{|u_n(x)|^p}{|x|^{2a}} |x|^a \,\mathrm{d}x$$

$$\leq Cn^{a+sp} \left(\int_{B_{2n}\setminus B_n} \frac{|u_n(x)|^{p_s^*}}{|x|^{\frac{2ap_s^*}{p}}} \,\mathrm{d}x \right)^{\frac{p}{p_s^*}} \leq Cn^{a+sp} \left(\int_{B_n^c} \frac{|u_n(x)|^{p_s^*}}{|x|^{\frac{2ap_s^*}{p}}} \,\mathrm{d}x \right)^{\frac{p}{p_s^*}}, \quad (3.24)$$

for some C = C(N, p, s). Therefore, in view of (3.22),(3.23), and using (3.21) and dominated convergence theorem we get

$$\lim_{n \to \infty} A_{n,i} = \lim_{n \to \infty} B_{n,i} = 0.$$

Thus, $\lim_{n\to\infty} I_{n,i,4} = 0$. Estimation of $I_{n,i,3}$: We break the integral $I_{n,i,3}$ into two parts as follows

$$I_{n,i,3} \leq C(p) \left\{ \underbrace{\iint_{(B_{2n} \setminus B_n) \times (B_{2n} \setminus B_n)} \frac{|\psi_{n,i}(x) - \psi_{n,i}(y)|^p}{|x - y|^{N + \sigma p}} |u_n(x)|^p \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a}}{E_{n,i}} + \underbrace{\iint_{(B_{2n} \setminus B_n) \times (B_{2n} \setminus B_n)} \frac{|u_n(x) - u_n(y)|^p}{|x - y|^{N + \sigma p}} |\psi_{n,i}(y)|^p}{\frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a}}_{F_{n,i}}} \right\}_{F_{n,i}}$$

Using the mean value theorem and $\|\nabla^2 \varphi_n\|_{\infty} \leq \frac{C(N)}{n^2}$, we estimate $E_{n,i}$ as

$$E_{n,i} \leq \frac{C(N,p)}{n^{2p}} \iint_{(B_{2n}\setminus B_n)\times(B_{2n}\setminus B_n)} \frac{|u_n(x)|^p}{|x-y|^{N+\sigma p-p}} \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a}$$
$$\leq \frac{C(N,p)}{n^{2p+a}} \int_{B_{2n}\setminus B_n} |u_n(x)|^p \left(\int_{B_{2n}\setminus B_n} \frac{\mathrm{d}y}{|x-y|^{N+\sigma p-p}}\right) \frac{\mathrm{d}x}{|x|^a}$$

Observe that $B_{2n} \subset B_{4n}(x)$, for $x \in B_{2n} \setminus B_n$. Thus using (3.24) we get

$$E_{n,i} \leq \frac{C(N,p)}{n^{2p+a}} \int_{B_{2n} \setminus B_n} |u_n(x)|^p \left(\int_{B_{4n}(x)} \frac{\mathrm{d}y}{|x-y|^{N+\sigma p-p}} \right) \frac{\mathrm{d}x}{|x|^a}$$
$$\leq \frac{C(N,p,\sigma)}{n^{p+\sigma p+a}} \int_{B_{2n} \setminus B_n} |u_n(x)|^p \frac{\mathrm{d}x}{|x|^a} \leq C(N,p,\sigma) \left(\int_{B_{2n} \setminus B_n} \frac{|u_n(x)|^{p_s^*}}{|x|^{\frac{2ap_s^*}{p}}} \,\mathrm{d}x \right)^{\frac{p}{p_s^*}}$$

Using (3.21) and subsequent arguments, it follows that $\lim_{n\to\infty} E_{n,i} = 0$. Let us now consider the second integral, i.e.,

$$F_{n,i} = \iint_{(B_{2n} \setminus B_n) \times (B_{2n} \setminus B_n)} \frac{|u_n(x) - u_n(y)|^p}{|x - y|^{N + \sigma p}} |\psi_{n,i}(y)|^p \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a} \\ \leq \frac{C(N,p)}{n^{p+2a}} \iint_{(B_{2n} \setminus B_n) \times (B_{2n} \setminus B_n)} \frac{|u_n(x) - u_n(y)|^p}{|x - y|^{N + \sigma p}} \,\mathrm{d}x \mathrm{d}y,$$
(3.25)

where the inequality holds since $|\psi_{n,i}(y)| \leq \frac{C(N)}{n}$. Applying Jensen's inequality $(\rho_n dz \text{ is a probability measure and } \sup (\rho_n) \subset \overline{B_{\frac{1}{n}}})$, we get

$$|u_{n}(x) - u_{n}(y)|^{p} = \left| \int_{\mathbb{R}^{N}} \rho_{n}(z)u(x-z) \, \mathrm{d}z - \int_{\mathbb{R}^{N}} \rho_{n}(z)u(y-z) \, \mathrm{d}z \right|^{p}$$
$$= \left| \int_{B_{\frac{1}{n}}} \left(u(x-z) - u(y-z) \right) \rho_{n}(z) \, \mathrm{d}z \right|^{p}$$
$$\leq \int_{B_{\frac{1}{n}}} |u(x-z) - u(y-z)|^{p} \rho_{n}(z) \, \mathrm{d}z.$$
(3.26)

From (3.25) and (3.26),

$$F_{n,i} \leq \frac{C(N,p)}{n^{p+2a}} \iint_{(B_{2n}\setminus B_n)\times(B_{2n}\setminus B_n)} \left(\int_{B_{\frac{1}{n}}} |u(x-z) - u(y-z)|^p \rho_n(z) \,\mathrm{d}z \right) \frac{\mathrm{d}x\mathrm{d}y}{|x-y|^{N+\sigma p}}$$

By the fundamental theorem of calculus and Jensen's inequality,

$$F_{n,i} \leq \frac{C(N,p)}{n^{p+2a}} \iint_{(B_{2n}\setminus B_n)\times(B_{2n}\setminus B_n)} \left(\int_{B_{\frac{1}{n}}} \left(\int_0^1 |\nabla u(\theta)| \, \mathrm{d}t \right)^p \rho_n(z) \, \mathrm{d}z \right) \frac{\mathrm{d}x\mathrm{d}y}{|x-y|^{N+\sigma p-p}} \\ \leq \frac{C(N,p)}{n^{p+2a}} \iint_{(B_{2n}\setminus B_n)\times(B_{2n}\setminus B_n)} \left(\int_{B_{\frac{1}{n}}} \left(\int_0^1 |\nabla u(\theta)|^p \, \mathrm{d}t \right) \rho_n(z) \, \mathrm{d}z \right) \frac{\mathrm{d}x\mathrm{d}y}{|x-y|^{N+\sigma p-p}}, \quad (3.27)$$

where $\theta = t(x-z) + (1-t)(y-z)$. Consider the following function

$$A(t,x,y,z) = \frac{|\nabla u(\theta)|^p}{|x-y|^{N+\sigma p-p}}\rho_n(z), \text{ where } x, y \in B_{2n} \setminus B_n, z \in B_{\frac{1}{n}} \text{ and } t \in (0,1).$$

Observe that $\theta = \theta(t, x, y, z)$ and

$$|\theta| \le t \left(|x| + |z|\right) + (1 - t) \left(|y| + |z|\right) \le 2n + \frac{1}{n}.$$
(3.28)

Now for fixed $t \in (0,1)$ and $z \in B_{\frac{1}{n}}$ we estimate the following integral

$$\iint_{(B_{2n}\setminus B_n)\times(B_{2n}\setminus B_n)} \frac{|\nabla u(\theta(t,\cdot,\cdot,z))|^p}{|x-y|^{N+\sigma p-p}} \,\mathrm{d}x\mathrm{d}y.$$

For fixed $x \in B_{2n} \setminus B_n$, we consider the following function

$$f_1(y) = \frac{|\nabla u(\theta(t, x, \cdot, z))|^p}{|\theta(t, x, \cdot, z)|^{2a}}, \text{ for } y \in \mathbb{R}^N.$$

Then using $\|\nabla u\|_{L^{p^*}_a(\mathbb{R}^N)} < \infty$ we see that

$$\int_{\mathbb{R}^N} f_1(y)^{\frac{p_{\sigma}^*}{p}} \, \mathrm{d}y \le (1-t)^{-N} \int_{\mathbb{R}^N} \frac{|\nabla u(\theta(t,x,\cdot,z))|^{p_{\sigma}^*}}{|\theta(t,x,\cdot,z)|^{\frac{2ap_{\sigma}^*}{p}}} \, \mathrm{d}\theta < \infty.$$

Also by Example 2.4, we have

$$g_1(\eta) = \frac{1}{|\eta|^{N+\sigma p-p}} \in L^{\frac{N}{N+\sigma p-p},\infty}(\mathbb{R}^N).$$

Therefore, by Proposition 2.5, $h_1(\eta) = \int_{\mathbb{R}^N} g_1(\eta - y) f_1(y) \, \mathrm{d}y < \infty$ for a.e. $\eta \in \mathbb{R}^N$ and moreover

$$h_1 \in L^r(\mathbb{R}^N)$$
, where $\frac{1}{r} = \frac{p}{p_\sigma^*} + \frac{N + \sigma p - p}{N} - 1 = \frac{N - p}{N}$

Hence for $x \in B_{2n} \setminus B_n$ using (3.28) we get

$$h_{2}(x) := \int_{B_{2n} \setminus B_{n}} |\nabla u(\theta(t, x, \cdot, z))|^{p} g_{1}(x - y) \, \mathrm{d}y \le c_{1}(n) \int_{B_{2n} \setminus B_{n}} g_{1}(x - y) f_{1}(y) \, \mathrm{d}y \le c_{1}(n) h_{1}(x),$$
(3.29)

where $c_1(n) = \left(2n + \frac{1}{n}\right)^{2a}$. Further, using the Hölder's inequality,

$$\int_{B_{2n} \setminus B_n} h_2(x) \, \mathrm{d}x \le C(N, p) c_1(n) \|h_1\|_r n^{\frac{N}{r'}}$$

Therefore, applying Tonelli's theorem, for a.e. $z\in B_{\frac{1}{n}}$ and $t\in(0,1)$ we get

$$B(t,z) := \iint_{(B_{2n}\setminus B_n)\times(B_{2n}\setminus B_n)} \frac{|\nabla u(\theta(t,\cdot,\cdot,z))|^p}{|x-y|^{N+\sigma p-p}} \,\mathrm{d}x \,\mathrm{d}y < \infty.$$
(3.30)

Further, Fubini's theorem yields

$$B(t,z) = \int_{B_{2n} \setminus B_n} \left(\int_{B_{2n} \setminus B_n} \frac{|\nabla u(\theta(t,\cdot,\cdot,z))|^p}{|x-y|^{N+\sigma p-p}} \, \mathrm{d}y \right) \, \mathrm{d}x$$

$$= \int_{B_{2n} \setminus B_n} h_2(x) \, \mathrm{d}x \le c_1(n) \int_{B_{2n} \setminus B_n} h_1(x) \, \mathrm{d}x \text{ (using (3.29))}$$

$$\le c_1(n) \left(\int_{B_{2n} \setminus B_n} h_1(x)^r \, \mathrm{d}x \right)^{\frac{1}{r}} \left(\int_{B_{2n} \setminus B_n} \, \mathrm{d}x \right)^{\frac{p}{N}}$$

$$\le C(N,p)c_1(n)n^p \left(\int_{B_n^c} h_1(x)^r \, \mathrm{d}x \right)^{\frac{1}{r}}.$$
(3.31)

Notice that $h_1(x) \equiv h_1(t, x, z)$, and for every $z \in B_{\frac{1}{n}}$ and $t \in (0, 1)$, since $h_1(t, \cdot, z) \in L^r(\mathbb{R}^N)$, we have

$$\left(\int_{B_n^c} h_1(t,\cdot,z)^r \,\mathrm{d}x\right)^{\frac{1}{r}} \to 0, \text{ as } n \to \infty.$$
(3.32)

Then there exists $n_0 \in \mathbb{N}$ such that for $n \ge n_0$, $\int_{B_n^c} h_1(t, \cdot, z)^r dx < 1$ for every $t \in (0, 1)$ and $z \in B_{\frac{1}{n}}$. Therefore,

$$f(t) := \int_{B_{\frac{1}{n}}} \rho_n(z) B(t,z) \, \mathrm{d}z \le C(N,p) c_1(n) n^p \int_{B_{\frac{1}{n}}} \rho_n(z) \left(\int_{B_n^c} h_1(t,\cdot,z)^r \, \mathrm{d}x \right)^{\frac{1}{r}} \, \mathrm{d}z$$
$$\le C(N,p) c_1(n) n^p \int_{B_{\frac{1}{n}}} \rho_n(z) \, \mathrm{d}z = C(N,p) c_1(n) n^p,$$

and $\int_0^1 f(t) dt \leq C(N,p)c_1(n)n^p$. Hence for $n \geq n_0$ using Tonneli's theorem,

$$A \in L^1\left((B_{2n} \setminus B_n) \times (B_{2n} \setminus B_n) \times B_{\frac{1}{n}} \times (0,1) \right),$$

and subsequently using Fubini's theorem, from (3.27) we obtain

$$F_{n,i} \leq \frac{C(N,p)}{n^{p+2a}} \int_0^1 \left(\int_{B_{\frac{1}{n}}} \left(\iint_{(B_{2n} \setminus B_n) \times (B_{2n} \setminus B_n)} \frac{|\nabla u(\theta)|^p}{|x-y|^{N+\sigma p-p}} \, \mathrm{d}x \mathrm{d}y \right) \rho_n(z) \, \mathrm{d}z \right) \, \mathrm{d}t.$$
(3.33)

Take $\epsilon > 0$. Then using the convergence in (3.32) there exists $n_1 (\geq n_0) \in \mathbb{N}$ such that for $n \geq n_1$, using (3.31) and (3.33), we conclude

$$F_{n,i} \leq C(N,p) \frac{c_1(n)n^p}{n^{p+2a}} \int_0^1 \left(\int_{B_{\frac{1}{n}}} \left(\int_{B_n^c} h_1(t,\cdot,z)^r \,\mathrm{d}x \right)^{\frac{1}{r}} \rho_n(z) \,\mathrm{d}z \right) \,\mathrm{d}t$$
$$\leq C(N,p) \int_0^1 \left(\int_{B_{\frac{1}{n}}} \left(\int_{B_n^c} h_1(t,\cdot,z)^r \,\mathrm{d}x \right)^{\frac{1}{r}} \rho_n(z) \,\mathrm{d}z \right) \,\mathrm{d}t < C(N,p)\epsilon.$$

Therefore, $\lim_{n \to \infty} F_{n,i} = 0$. In conclusion, we get $\lim_{n \to \infty} I_{n,i,3} = 0$.

Finally, taking limit as $n \to \infty$ in (3.16) and using the convergence of $I_{n,i,k}, k = 1, \cdot, \cdot, \cdot, 5$, we get $\lim_{n \to \infty} I_{n,i} = 0, \text{ for each } i = 1, 2, \cdot, \cdot, \cdot, N.$ (3.34)

Therefore, by (3.7), (3.15) and (3.34), we obtain

$$\lim_{n \to \infty} [u \star \rho_n - (u \star \rho_n) \zeta_n]_{s,p,a} = 0.$$
(3.35)

From (3.1), (3.6) and (3.35), we get the required result.

Remark 3.3. In view of Proposition 3.2 and the definition of $\mathcal{D}_a^{s,p}(\mathbb{R}^N)$, we infer that $\mathring{W}_a^{s,p}(\mathbb{R}^N) \subset \mathcal{D}_a^{s,p}(\mathbb{R}^N)$. More precisely, if $u \in \mathring{W}_a^{s,p}(\mathbb{R}^N)$, then using Proposition 3.2 there exists $\{u_n\} \subset \mathcal{C}_c^{\infty}(\mathbb{R}^N)$ such that $[u - u_n]_{s,p,a} \to 0$ as $n \to \infty$, which implies that $u \in \mathcal{D}_a^{s,p}(\mathbb{R}^N)$.

In the following proposition, we prove that the seminorm $[\cdot]_{s,p,a}$ is an equivalent norm in $\mathring{W}_a^{s,p}(\mathbb{R}^N)$. **Proposition 3.4.** Let $N \ge 2, s \in (1,2), \sigma = s - 1, p \in (1, \frac{N}{s})$, and $a \in [0, \frac{N-sp}{2})$. Then there exists $C = C(N, p, \sigma, a) > 0$ such that

$$\|\nabla u\|_{L^{p^*}_a(\mathbb{R}^N)} \le C[u]_{s,p,a}, \ \forall u \in \check{W}^{s,p}_a(\mathbb{R}^N).$$

Proof. For $u \in \mathring{W}_a^{s,p}(\mathbb{R}^N)$, we consider the same sequence $v_n = (u \star \rho_n) \zeta_n$ for $n \in \mathbb{N}$ as in Proposition 3.2. Then $v_n \in \mathcal{C}_c^{\infty}(\mathbb{R}^N)$. Using [1, Theorem 1.5] with $\frac{\partial v_n}{\partial x_i}$ and Lemma 2.1, we get

$$\left\|\nabla v_n\right\|_{L^{p^*}_{a\sigma}(\mathbb{R}^N)} \le C(N, p, \sigma, a) \left[v_n\right]_{s, p, a}, \ \forall n \in \mathbb{N}.$$
(3.36)

We claim

$$\|\nabla u - \nabla v_n\|_{L^{p^*}_a(\mathbb{R}^N)} \to 0, \text{ as } n \to \infty.$$
(3.37)

Using the triangle inequality, we obtain

$$\begin{aligned} \|\nabla u - \nabla v_n\|_{L^{p^*}_{a}(\mathbb{R}^N)} &= \|\nabla u - (\nabla u \star \rho_n) \zeta_n - (u \star \rho_n) \nabla \zeta_n\|_{L^{p^*}_{a}(\mathbb{R}^N)} \\ &\leq \underbrace{\|\nabla u \star \rho_n - (\nabla u \star \rho_n) \zeta_n\|_{L^{p^*}_{a}(\mathbb{R}^N)}}_{A_n} + \underbrace{\|\nabla u - \nabla u \star \rho_n\|_{L^{p^*}_{a}(\mathbb{R}^N)}}_{B_n} + \underbrace{\|(u \star \rho_n) \nabla \zeta_n\|_{L^{p^*}_{a}(\mathbb{R}^N)}}_{C_n}. \end{aligned}$$
(3.38)

As before, we define $\varphi_n = 1 - \zeta_n$. By noticing $0 \le \varphi_n \le 1$, we deduce

$$\begin{aligned} A_{n}^{p_{\sigma}^{*}} &= \left\| \left(\nabla u \star \rho_{n} \right) \varphi_{n} \right\|_{L_{a}^{p_{\sigma}^{*}}(\mathbb{R}^{N})}^{p_{\sigma}^{*}} = \int_{\mathbb{R}^{N}} \frac{\left| \left(\nabla u \star \rho_{n} \right) \varphi_{n} \right|^{p_{\sigma}^{*}}}{\left| x \right|^{\frac{2ap_{\sigma}^{*}}{p}}} \mathrm{d}x = \int_{\mathbb{R}^{N}} \left(\sum_{i=1}^{N} \left| \frac{\partial u_{n}}{\partial x_{i}} \varphi_{n} \right|^{2} \right)^{\frac{p_{\sigma}^{*}}{2}} \frac{\mathrm{d}x}{\left| x \right|^{\frac{2ap_{\sigma}^{*}}{p}}} \\ &\leq A(N, \frac{p_{\sigma}^{*}}{2}) \sum_{i=1}^{N} \int_{\mathbb{R}^{N}} \left| \frac{\partial u_{n}}{\partial x_{i}} \varphi_{n} \right|^{p_{\sigma}^{*}} \frac{\mathrm{d}x}{\left| x \right|^{\frac{2ap_{\sigma}^{*}}{p}}} \left(\text{using Lemma 2.1-(i)} \right) \\ &\leq 2^{p_{\sigma}^{*}-1} A(N, \frac{p_{\sigma}^{*}}{2}) \left(\sum_{i=1}^{N} \int_{\mathbb{R}^{N}} \left| \left(\frac{\partial u_{n}}{\partial x_{i}} - \frac{\partial u}{\partial x_{i}} \right) \varphi_{n} \right|^{p_{\sigma}^{*}} \frac{\mathrm{d}x}{\left| x \right|^{\frac{2ap_{\sigma}^{*}}{p}}} + \sum_{i=1}^{N} \int_{\mathbb{R}^{N}} \left| \frac{\partial u}{\partial x_{i}} \varphi_{n} \right|^{p_{\sigma}^{*}} \frac{\mathrm{d}x}{\left| x \right|^{\frac{2ap_{\sigma}^{*}}{p}}} \right) \\ &\leq 2^{p_{\sigma}^{*}-1} A(N, \frac{p_{\sigma}^{*}}{2}) \left(\sum_{i=1}^{N} \int_{\mathbb{R}^{N}} \left| \frac{\partial u}{\partial x_{i}} - \frac{\partial u_{n}}{\partial x_{i}} \right|^{p_{\sigma}^{*}} \frac{\mathrm{d}x}{\left| x \right|^{\frac{2ap_{\sigma}^{*}}{p}}} + \sum_{i=1}^{N} \int_{\mathbb{R}^{N}} \left| \frac{\partial u}{\partial x_{i}} \varphi_{n} \right|^{p_{\sigma}^{*}} \frac{\mathrm{d}x}{\left| x \right|^{\frac{2ap_{\sigma}^{*}}{p}}} \right). \end{aligned}$$
(3.39)

Since $|\nabla u| \in L^{p^*}_a(\mathbb{R}^N)$, we get $\frac{\partial u}{\partial x_i} \in L^{p^*}_a(\mathbb{R}^N)$. Thus, applying [10, Lemma 6.1] with $\frac{\partial u}{\partial x_i}$, we have

$$\lim_{n \to \infty} \int_{\mathbb{R}^N} \left| \frac{\partial u}{\partial x_i} - \frac{\partial u_n}{\partial x_i} \right|^{p_{\sigma}^*} \frac{\mathrm{d}x}{|x|^{\frac{2ap_{\sigma}^*}{p}}} = 0, \text{ for each } i = 1, 2, \cdot, \cdot, \cdot, N.$$
(3.40)

By noticing that $\operatorname{supp}(\zeta_n) \subset \overline{B_{2n}}$ and $\operatorname{supp}(\varphi_n) \subset B_n^c$, the dominated convergence theorem yields

$$\lim_{n \to \infty} \int_{\mathbb{R}^N} \left| \frac{\partial u}{\partial x_i} \varphi_n \right|^{p_{\sigma}^*} \frac{\mathrm{d}x}{|x|^{\frac{2ap_{\sigma}^*}{p}}} = 0, \text{ for each } i = 1, 2, \cdot, \cdot, \cdot, N.$$
(3.41)

Considering (3.39), (3.40), and (3.41), we can deduce that

$$\lim_{n \to \infty} A_n = 0. \tag{3.42}$$

Further applying Lemma 2.1-(i), we derive

$$B_n^{p^*_{\sigma}} = \|\nabla u - \nabla u \star \rho_n\|_{L^{p^*_{\sigma}}_a(\mathbb{R}^N)}^{p^*_{\sigma}} = \int_{\mathbb{R}^N} \left(\sum_{i=1}^N \left| \frac{\partial u}{\partial x_i} - \frac{\partial u_n}{\partial x_i} \right|^2 \right)^{\frac{p^*_{\sigma}}{2}} \frac{\mathrm{d}x}{|x|^{\frac{2ap^*_{\sigma}}{p}}} \\ \leq A(N, \frac{p^*_{\sigma}}{2}) \sum_{i=1}^N \int_{\mathbb{R}^N} \left| \frac{\partial u}{\partial x_i} - \frac{\partial u_n}{\partial x_i} \right|^{p^*_{\sigma}} \frac{\mathrm{d}x}{|x|^{\frac{2ap^*_{\sigma}}{p}}}.$$

We use (3.40) in the above inequality to get

$$\lim_{n \to \infty} B_n = 0. \tag{3.43}$$

Using $\|\nabla \zeta_n\|_{\infty} \leq \frac{C}{n}$ and $\operatorname{supp}(|\nabla \zeta_n|) \subset \overline{B_{2n} \setminus B_n}$, we estimate the last term of (3.38) as follows

$$C_n^{p^*_{\sigma}} = \int_{\mathbb{R}^N} \left| (u \star \rho_n) \nabla \zeta_n \right|^{p^*_{\sigma}} \frac{\mathrm{d}x}{|x|^{\frac{2ap^*_{\sigma}}{p}}} \\ \leq \frac{C(N, p, \sigma)}{n^{p^*_{\sigma}}} \int_{B_{2n} \setminus B_n} |u \star \rho_n|^{p^*_{\sigma}} \frac{\mathrm{d}x}{|x|^{\frac{2ap^*_{\sigma}}{p}}} \leq \frac{C(N, p, \sigma)}{n^{p^*_{\sigma} + \frac{2ap^*_{\sigma}}{p}}} \int_{B_{2n} \setminus B_n} \left| \int_{B_{\frac{1}{n}}} u(x - z)\rho_n(z) \, \mathrm{d}z \right|^{p^*_{\sigma}} \, \mathrm{d}x \\ \leq \frac{C(N, p, \sigma)}{n^{2m^*_{\sigma}}} \int_{\mathbb{R}^n} \left(\int_{\mathbb{R}^n} |u(x - z)|^{p^*_{\sigma}} \rho_n(z) \, \mathrm{d}z \right) \, \mathrm{d}x.$$

$$(3.44)$$

 $\leq \frac{O(N, p, \sigma)}{n^{p_{\sigma}^{*} + \frac{2ap_{\sigma}^{*}}{p}}} \int_{B_{2n} \setminus B_{n}} \left(\int_{B_{\frac{1}{n}}} |u(x-z)|^{p_{\sigma}^{*}} \rho_{n}(z) \, \mathrm{d}z \right) \, \mathrm{d}x. \tag{3.44}$ The last inequality follows using Jensen's inequality ($\rho_{n} \mathrm{d}z$ is a probability measure and $\mathrm{supp}(\rho_{n}) \subset$

The last inequality follows using Jensen's inequality ($\rho_n dz$ is a probability measure and $\sup(\rho_n) \subset \overline{B_{\frac{1}{n}}}$). For $x \in B_{2n} \setminus B_n$ and $z \in B_{\frac{1}{n}}$, we have

$$n - \frac{1}{n} \le |x| - |z| \le |x - z| \le |x| + |z| \le 2n + \frac{1}{n}.$$
(3.45)

Now consider $F(x,z) = |u(x-z)|^{p_{\sigma}^*} \rho_n(z)$. Then for a.e. $z \in B_{\frac{1}{n}}$, we have

$$\begin{split} \int_{B_{2n}\setminus B_n} F(x,z) \, \mathrm{d}x &= \rho_n(z) \int_{B_{2n}\setminus B_n} |u(x-z)|^{p_{\sigma}^*} \, \mathrm{d}x \\ &\leq c_2(n)\rho_n(z) \int_{B_{2n}\setminus B_n} \frac{|u(x-z)|^{p_{\sigma}^*}}{|x-z|^{\frac{2ap_{\sigma}^*}{p}}} \, \mathrm{d}x, \text{ where } c_2(n) = \left(2n + \frac{1}{n}\right)^{\frac{2ap_{\sigma}^*}{p}} \\ &\leq c_2(n)\rho_n(z) \left(\int_{B_{2n}\setminus B_n} \frac{|u(x-z)|^{p_s^*}}{|x-z|^{\frac{2ap_s^*}{p}}} \, \mathrm{d}x\right)^{\frac{p_{\sigma}^*}{p_s^*}} \left(\int_{B_{2n}\setminus B_n} \, \mathrm{d}x\right)^{\frac{p}{N-\sigma p}}. \end{split}$$

Thus we obtain

$$\int_{B_{2n}\setminus B_n} F(x,z) \, \mathrm{d}x \le C(N,p,\sigma)c_2(n)n^{p_{\sigma}^*}\rho_n(z) \left(\int_{B_n^c} \frac{|u(x-z)|^{p_s^*}}{|x-z|^{\frac{2ap_s^*}{p}}} \, \mathrm{d}x\right)^{\frac{p_{\sigma}^*}{p_s^*}}.$$

Since $u \in L_a^{p_s^*}(\mathbb{R}^N)$, for arbitrary $0 < \epsilon < 1$, there exists $n_{\epsilon} \in \mathbb{N}$ such that

$$\left(\int_{B_n^c} \frac{|u(x-z)|^{p_s^*}}{|x-z|^{\frac{2ap_s^*}{p}}} \,\mathrm{d}x\right)^{\frac{p_a^*}{p_s^*}} < \epsilon, \ \forall n \ge n_\epsilon.$$
(3.46)

Using (3.46), for $n \ge n_{\epsilon}$ we obtain

$$\int_{B_{2n}\setminus B_n} F(x,z) \,\mathrm{d}x \le C(N,p,\sigma)c_3(n)\rho_n(z)\epsilon,$$

where $c_3(n) = n^{\frac{2ap_{\sigma}^*}{p} + p_{\sigma}^*}$. Further, we see that for every $n \ge n_{\epsilon}$,

$$\int_{B_{\frac{1}{n}}} \left(\int_{B_{2n} \setminus B_n} F(x, z) \, \mathrm{d}x \right) \, \mathrm{d}z \le C(N, p, \sigma) c_3(n) \epsilon \int_{B_{\frac{1}{n}}} \rho_n(z) \, \mathrm{d}z = C(N, p, \sigma) c_3(n) \epsilon. \tag{3.47}$$

Therefore, Tonelli's theorem yields $F \in L^1\left((B_{2n} \setminus B_n) \times B_{\frac{1}{n}}\right)$. Subsequently, applying Fubini's theorem in (3.44) and then using (3.47) we get

$$C_n^{p^*_{\sigma}} \leq \frac{C(N, p, \sigma)}{n^{p^*_{\sigma} + \frac{2ap^*_{\sigma}}{p}}} \int_{B_{\frac{1}{n}}} \left(\int_{B_{2n} \setminus B_n} |u(x-z)|^{p^*_{\sigma}} \rho_n(z) \,\mathrm{d}x \right) \,\mathrm{d}z < C(N, p, \sigma)\epsilon, \; \forall n \geq n_{\epsilon}.$$

Therefore,

$$\lim_{n \to \infty} C_n = 0. \tag{3.48}$$

From (3.38), (3.42), (3.43) and (3.48), we obtain (3.37). Moreover, we know from Proposition 3.2 that

$$[u - v_n]_{s,p,a} \to 0, \text{ as } n \to \infty.$$
(3.49)

Next, passing the limit to the inequality (3.36) and using (3.37) and (3.49), we get

$$\|\nabla u\|_{L^{p^*}_a(\mathbb{R}^N)} \le C[u]_{s,p,a}, \ \forall \, u \in \tilde{W}^{s,p}_a(\mathbb{R}^N),$$

which completes the proof.

Proof of Theorem 1.1-((i) and (ii)): The proof of part-(i) and part-(ii) is followed from Proposition 3.2 and Proposition 3.4, respectively.

4. CHARACTERIZATION OF FRACTIONAL WEIGHTED HOMOGENEOUS SPACES

Now, we proceed to characterize the function space $\mathcal{D}_a^{s,p}(\mathbb{R}^N)$. In the following proposition, we show that every Cauchy sequence in $\mathcal{C}_c^{\infty}(\mathbb{R}^N)$ converges to a function in $\mathring{W}_a^{s,p}(\mathbb{R}^N)$ with respect to the Gagliardo seminorm $[\cdot]_{s,p,a}$.

Proposition 4.1. Let $N \ge 2, s \in (1,2), \sigma = s - 1, p \in (1, \frac{N}{s})$, and $a \in [0, \frac{N-sp}{2})$. Let $\{w_n\} \subset C_c^{\infty}(\mathbb{R}^N)$ be a Cauchy sequence with respect to $[\cdot]_{s,p,a}$. Then there exists $w \in \mathring{W}_a^{s,p}(\mathbb{R}^N)$ such that

$$[w - w_n]_{s,p,a} \to 0, \text{ as } n \to \infty,$$

Proof. Let $\epsilon > 0$ be given. By hypothesis, there exists $n_{\epsilon} \in \mathbb{N}$ such that

$$[w_n - w_m]_{s,p,a} < \epsilon, \ \forall m, n \ge n_\epsilon.$$

$$(4.1)$$

By [7, Theorem 1], we have

$$\|u\|_{L^{p^*}_a(\mathbb{R}^N)} \le C_1(N, p, \sigma, a) \|\nabla u\|_{L^{p^*}_a(\mathbb{R}^N)}, \ \forall \, u \in \mathcal{C}^\infty_c(\mathbb{R}^N).$$

$$(4.2)$$

Further, by Proposition 3.4,

$$\|\nabla u\|_{L^{p^*}_a(\mathbb{R}^N)} \le C_2(N, p, \sigma, a)[u]_{s, p, a}, \ \forall \, u \in \mathcal{C}^\infty_c(\mathbb{R}^N),$$

$$(4.3)$$

From (4.1), (4.2), (4.3), and the fact that $\{w_n\} \subset \mathcal{C}_c^{\infty}(\mathbb{R}^N)$, we get $\{w_n\}$ is Cauchy in $L_a^{p_s^*}(\mathbb{R}^N)$ and $\{\frac{\partial w_n}{\partial x_i}\}$ is Cauchy in $L_a^{p_\sigma^*}(\mathbb{R}^N)$. Consequently, $w_n \to w$ in $L_a^{p_s^*}(\mathbb{R}^N)$ and $\frac{\partial w_n}{\partial x_i} \to v_i$ in $L_a^{p_\sigma^*}(\mathbb{R}^N)$. Moreover, up to a subsequence, $w_n \to w$ and $\frac{\partial w_n}{\partial x_i} \to v_i$ a.e. in \mathbb{R}^N . To show $v_i = \frac{\partial w}{\partial x_i}$, we consider $\varphi \in \mathcal{C}_c^{\infty}(\mathbb{R}^N)$ with $\operatorname{supp}(\varphi) \subset K$, where K is a bounded open set of class $\mathcal{C}^{1,1}$. The integration by parts formula yields

$$\int_{\mathbb{R}^N} w_n \frac{\partial \varphi}{\partial x_i} \, \mathrm{d}x = \int_K w_n \frac{\partial \varphi}{\partial x_i} \, \mathrm{d}x = -\int_K \frac{\partial w_n}{\partial x_i} \varphi \, \mathrm{d}x = -\int_{\mathbb{R}^N} \frac{\partial w_n}{\partial x_i} \varphi \, \mathrm{d}x. \tag{4.4}$$

Using Hölder's inequality and $w_n \to w$ in $L_a^{p_s^*}(\mathbb{R}^N)$, we get

$$\begin{split} \left\| \int_{\mathbb{R}^{N}} \left(w - w_{n} \right) \frac{\partial \varphi}{\partial x_{i}} \, \mathrm{d}x \right\| &\leq \left\| \frac{\partial \varphi}{\partial x_{i}} \right\|_{\infty} \int_{K} \left| w - w_{n} \right| \, \mathrm{d}x \\ &= \left\| \frac{\partial \varphi}{\partial x_{i}} \right\|_{\infty} \int_{K} \frac{\left| w - w_{n} \right|}{\left| x \right|^{\frac{2a}{p}}} \cdot \left| x \right|^{\frac{2a}{p}} \, \mathrm{d}x \leq \left\| \frac{\partial \varphi}{\partial x_{i}} \right\|_{\infty} \left(\int_{K} \frac{\left| w - w_{n} \right|^{p_{s}^{*}}}{\left| x \right|^{\frac{2aps^{*}}{p}}} \, \mathrm{d}x \right)^{\frac{1}{p_{s}^{*}}} \left(\int_{K} \left(\left| x \right|^{\frac{2a}{p}} \right)^{\frac{p_{s}^{*}-1}}_{\frac{p_{s}^{*}-1}{p_{s}^{*}}} \, \mathrm{d}x \right)^{\frac{p_{s}^{*}-1}{p_{s}^{*}}} \\ &\leq C(N, p, \sigma, a, K) \left\| \frac{\partial \varphi}{\partial x_{i}} \right\|_{\infty} \left\| w - w_{n} \right\|_{L^{p_{s}^{*}}_{a}(\mathbb{R}^{N})} \to 0, \text{ as } n \to \infty. \end{split}$$

$$\tag{4.5}$$

Using $\frac{\partial w_n}{\partial x_i} \to v_i$ in $L_a^{p^*_{\sigma}}(\mathbb{R}^N)$ and similar calculations as in (4.5), we also get

$$\left| \int_{\mathbb{R}^N} \left(v_i - \frac{\partial w_n}{\partial x_i} \right) \varphi \, \mathrm{d}x \right| \to 0, \text{ as } n \to \infty.$$
(4.6)

Taking the limit as $n \to \infty$ in (4.4), and using (4.5) and (4.6),

$$\int_{\mathbb{R}^N} w \frac{\partial \varphi}{\partial x_i} \, \mathrm{d}x = -\int_{\mathbb{R}^N} v_i \varphi \, \mathrm{d}x, \; \forall \, \varphi \in \mathcal{C}^\infty_c(\mathbb{R}^N)$$

which implies $\frac{\partial w}{\partial x_i} = v_i \in L_a^{p^*_{\sigma}}(\mathbb{R}^N)$, and hence $|\nabla w| \in L_a^{p^*_{\sigma}}(\mathbb{R}^N)$ (by Lemma 2.1-(i)). For $n \ge n_{\epsilon}$, the triangle inequality and (4.1) give

$$[w_n]_{s,p,a} \le [w_n - w_{n_{\epsilon}}]_{s,p,a} + [w_{n_{\epsilon}}]_{s,p,a} < \epsilon + [w_{n_{\epsilon}}]_{s,p,a}.$$
(4.7)

For a.e. $(x, y) \in \mathbb{R}^N \times \mathbb{R}^N$, we define

$$V_n(x,y) = \frac{|\nabla w_n(x) - \nabla w_n(y)|^p}{|x-y|^{N+\sigma p} |x|^a |y|^a} \text{ and } V(x,y) = \frac{|\nabla w(x) - \nabla w(y)|^p}{|x-y|^{N+\sigma p} |x|^a |y|^a}.$$
(4.8)

Using (4.7), $\{V_n\}$ is bounded in $L^1(\mathbb{R}^N \times \mathbb{R}^N)$. Since $\frac{\partial w_n}{\partial x_i} \to \frac{\partial w}{\partial x_i}$ a.e. in \mathbb{R}^N , we get $V_n \to V$ a.e. in $\mathbb{R}^N \times \mathbb{R}^N$. Hence the Fatou's Lemma yields

$$[w]_{s,p,a}^p = \iint_{\mathbb{R}^N \times \mathbb{R}^N} V(x,y) \, \mathrm{d}x \mathrm{d}y \le \liminf_{n \to \infty} \iint_{\mathbb{R}^N \times \mathbb{R}^N} V_n(x,y) \, \mathrm{d}x \mathrm{d}y < \infty.$$

Therefore, $w \in \mathring{W}_{a}^{s,p}(\mathbb{R}^{N})$. Now we show that $w_n \to w$ in $\mathcal{D}_{a}^{s,p}(\mathbb{R}^{N})$. We define

$$V_{n,i}(x,y) = \frac{\frac{\partial w_n}{\partial x_i}(x) - \frac{\partial w_n}{\partial x_i}(y)}{|x-y|^{\frac{N}{p}+\sigma} |x|^{\frac{a}{p}} |y|^{\frac{a}{p}}}, \text{ for a.e. } (x,y) \in \mathbb{R}^N \times \mathbb{R}^N.$$

Using Lemma 2.1-(ii), we have

$$[w_n - w_m]_{s,p,a}^p = \iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{\left| (\nabla w_n(x) - \nabla w_n(y)) - (\nabla w_m(x) - \nabla w_m(y)) \right|^p}{|x - y|^{N + \sigma p}} \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a}$$

$$\geq B(N, \frac{p}{2}) \left(\sum_{i=1}^N \iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{\left| \left(\frac{\partial w_n}{\partial x_i}(x) - \frac{\partial w_n}{\partial x_i}(y) \right) - \left(\frac{\partial w_m}{\partial x_i}(x) - \frac{\partial w_m}{\partial x_i}(y) \right) \right|^p}{|x - y|^{N + \sigma p}} \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a} \right)$$

$$= B(N, \frac{p}{2}) \sum_{i=1}^N ||V_{n,i} - V_{m,i}||_{L^p(\mathbb{R}^N \times \mathbb{R}^N)}^p.$$

$$(4.9)$$

From (4.1) and (4.9), we deduce that for each *i*, the sequence $\{V_{n,i}\}$ is Cauchy in $L^p(\mathbb{R}^N \times \mathbb{R}^N)$. By completeness of $L^p(\mathbb{R}^N \times \mathbb{R}^N)$, there exists $V_i \in L^p(\mathbb{R}^N \times \mathbb{R}^N)$ such that $V_{n,i} \to V_i$ in $L^p(\mathbb{R}^N \times \mathbb{R}^N)$. Also, $\frac{\partial w_n}{\partial x_i} \to \frac{\partial w}{\partial x_i}$ a.e. in \mathbb{R}^N . Therefore, to the uniqueness of the limit,

$$V_i(x,y) = \frac{\frac{\partial w}{\partial x_i}(x) - \frac{\partial w}{\partial x_i}(y)}{|x-y|^{\frac{N}{p}+\sigma} |x|^{\frac{a}{p}} |y|^{\frac{a}{p}}}, \text{ for a.e. } (x,y) \in \mathbb{R}^N \times \mathbb{R}^N.$$

Therefore,

$$\begin{split} [w - w_n]_{s,p,a}^p &= \iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{\left| (\nabla w(x) - \nabla w(y)) - (\nabla w_n(x) - \nabla w_n(y)) \right|^p}{|x - y|^{N + \sigma p}} \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a} \\ &\leq A(N, \frac{p}{2}) \left(\sum_{i=1}^N \iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{\left| \left(\frac{\partial w}{\partial x_i}(x) - \frac{\partial w}{\partial x_i}(y) \right) - \left(\frac{\partial w_n}{\partial x_i}(x) - \frac{\partial w_n}{\partial x_i}(y) \right) \right|^p}{|x - y|^{N + \sigma p}} \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a} \right) \\ &= A(N, \frac{p}{2}) \sum_{i=1}^N \|V_i - V_{n,i}\|_{L^p(\mathbb{R}^N \times \mathbb{R}^N)}^p \to 0, \text{ as } n \to \infty. \end{split}$$

This completes the proof.

Completions. From Proposition 3.4, the seminorm $[\cdot]_{s,p,a}$ becomes a norm in the space $\mathcal{C}_c^{\infty}(\mathbb{R}^N)$. However, $\mathcal{C}_c^{\infty}(\mathbb{R}^N)$ is not complete with respect to this norm. This leads us to define the completion of $\mathcal{C}_c^{\infty}(\mathbb{R}^N)$ with respect to $[\cdot]_{s,p,a}$. Let V be the vector space of Cauchy sequences in $\mathcal{C}_c^{\infty}(\mathbb{R}^N)$ and V_0 be the subspace of V consisting of the sequences converging to zero. We define an equivalence relation on V by

$$\boldsymbol{u} \sim \boldsymbol{v} \iff \boldsymbol{u} - \boldsymbol{v} \in V_0,$$

i.e., two sequences $\boldsymbol{u} = \{u_n\}$ and $\boldsymbol{v} = \{v_n\}$ in V are said to be equivalent if and only if

$$\lim_{n \to \infty} [u_n - v_n]_{s,p,a} = 0.$$
(4.10)

The equivalence class containing a vector $\boldsymbol{u} \in V$ is called a coset, and it is denoted as $\boldsymbol{u} + V_0$ and defined by

$$oldsymbol{u}+V_0:=ig\{oldsymbol{u}+oldsymbol{w}:oldsymbol{w}\in V_0ig\}$$
 .

The set of all cosets forms a quotient space and is defined as

$$V/V_0 := \left\{ oldsymbol{u} + V_0 : oldsymbol{u} \in V
ight\},$$

and it becomes a vector space under the following addition and scalar multiplication:

$$(\boldsymbol{u}+V_0)+(\boldsymbol{v}+V_0)=(\boldsymbol{u}+\boldsymbol{v})+V_0$$
 and $\alpha(\boldsymbol{u}+V_0)=\alpha\boldsymbol{u}+V_0, \ \forall \, \boldsymbol{u}, \boldsymbol{v}\in V, \alpha\in\mathbb{R}.$

We define

$$\|\boldsymbol{u} + V_0\| := \lim_{n \to \infty} [u_n]_{s,p,a}.$$
(4.11)

Since $u \in V$, the sequence $u = \{u_n\}$ is Cauchy with respect to $[\cdot]_{s,p,a}$ and therefore the sequence $\{[u_n]_{s,p,a}\}$ is Cauchy in \mathbb{R} by the following inequality

$$|[u_n]_{s,p,a} - [u_m]_{s,p,a}| \le [u_n - u_m]_{s,p,a}.$$

By completeness of \mathbb{R} , the sequence $\{[u_n]_{s,p,a}\}$ converges in \mathbb{R} and hence the right hand quantity in (4.11) is well defined. Further, if $\boldsymbol{u} = \{u_n\}$ and $\boldsymbol{v} = \{v_n\}$ are Cauchy sequences such that $\{u_n - v_n\} \in V_0$, then by the following inequalities

$$[u_n]_{s,p,a} \le [u_n - v_n]_{s,p,a} + [v_n]_{s,p,a}$$
 and $[v_n]_{s,p,a} \le [u_n - v_n]_{s,p,a} + [u_n]_{s,p,a}$,

it follows that $\lim_{n\to\infty} [u_n]_{s,p,a} = \lim_{n\to\infty} [v_n]_{s,p,a}$. Thus, (4.11) is well defined, and it is independent of the choice of a representative of the coset.

Remark 4.2. Using [21, Section 10, Page 56], we can verify that $\|\cdot\|$ becomes a norm on V/V_0 and V/V_0 forms a Banach space with respect to $\|\cdot\|$ and this space is unique upto isometric isomorphism.

Since $\mathcal{D}_a^{s,p}(\mathbb{R}^N)$ is a completion of $(\mathcal{C}_c^{\infty}(\mathbb{R}^N), [\cdot]_{s,p,a})$, we can identify the spaces V/V_0 and $\mathcal{D}_a^{s,p}(\mathbb{R}^N)$. Each element of $\mathcal{D}_a^{s,p}(\mathbb{R}^N)$ is considered as an equivalence class $U = \langle \{u_n\} \rangle$, where the Cauchy sequence $\{u_n\}$ in $\mathcal{C}_c^{\infty}(\mathbb{R}^N)$ is a representative of the class. Due to the isometric isomorphism between $\mathcal{D}_a^{s,p}(\mathbb{R}^N)$ and V/V_0 , the norm on $\mathcal{D}_a^{s,p}(\mathbb{R}^N)$ in terms of its representative is given by

$$||U||_{\mathcal{D}_a^{s,p}(\mathbb{R}^N)} = \lim_{n \to \infty} [u_n]_{s,p,a}.$$

Proof of Theorem 1.1-(iii): Using Proposition 3.4, it follows that $[\cdot]_{s,p,a}$ defines a norm on $\mathring{W}_{a}^{s,p}(\mathbb{R}^{N})$. Now, we will show the existence of an isometric isomorphism between $\mathcal{D}_{a}^{s,p}(\mathbb{R}^{N})$ and $\mathring{W}_{a}^{s,p}(\mathbb{R}^{N})$, thereby establishing that $\mathring{W}_{a}^{s,p}(\mathbb{R}^{N})$ is a Banach space. Let $U = \langle \{u_n\} \rangle \in \mathcal{D}_{a}^{s,p}(\mathbb{R}^{N})$. By Proposition 4.1, $u_n \to u$ with respect to $[\cdot]_{s,p,a}$ for some $u \in \mathring{W}_{a}^{s,p}(\mathbb{R}^{N})$. We define a map $\mathcal{J}: \mathcal{D}_{a}^{s,p}(\mathbb{R}^{N}) \to \mathring{W}_{a}^{s,p}(\mathbb{R}^{N})$ as

$$\mathcal{J}(U) = u$$

If $\{v_n\}$ is another representative of the equivalence class U, then using the equivalence relation (4.10), we have

$$\lim_{n \to \infty} [u - v_n]_{s,p,a} \le \lim_{n \to \infty} [u - u_n]_{s,p,a} + \lim_{n \to \infty} [u_n - v_n]_{s,p,a} = 0.$$

Thus, the map \mathcal{J} is well-defined, and one can verify that it is linear. Further, the map \mathcal{J} is an isometry by Proposition 4.1 i.e.,

$$||U||_{\mathcal{D}_a^{s,p}(\mathbb{R}^N)} = \lim_{n \to \infty} [u_n]_{s,p,a} = [u]_{s,p,a}.$$

Now we claim that \mathcal{J} is surjective. It follows from Proposition 3.2 that for every $w \in \mathring{W}^{s,p}_a(\mathbb{R}^N)$, there exists $\{w_n\} \subset \mathcal{C}^{\infty}_c(\mathbb{R}^N)$ such that

$$\lim_{n \to \infty} \left[w - w_n \right]_{s,p,a} = 0$$

Since $\{w_n\}$ is Cauchy with respect to $[\cdot]_{s,p,a}$, it is a representative of some equivalence class W i.e., $W = \langle \{w_n\} \rangle$. Thus, we obtain

$$\mathcal{J}(W) = w,$$

which implies \mathcal{J} is surjective. Hence, \mathcal{J} is a linear surjective isometry.

5. Fractional weighted Rellich inequality and finer embeddings

First, we recall the Caffarelli-Kohn-Nirenberg inequality [7].

Theorem 5.1 (Caffarelli-Kohn-Nirenberg). Let $p, q, r, \alpha, \beta, \gamma$ and l be real numbers such that $p, q \ge 1, r > 0, 0 \le l \le 1$, and

$$\frac{1}{p}+\frac{\alpha}{N}, \frac{1}{q}+\frac{\beta}{N}, \frac{1}{r}+\frac{m}{N}>0,$$

where $m = l\gamma + (1 - l)\beta$. Then there exists a positive constant C such that

$$\left(\int_{\mathbb{R}^N} |x|^{mr} |u(x)|^r \,\mathrm{d}x\right)^{\frac{1}{r}} \le C \left(\int_{\mathbb{R}^N} |x|^{\alpha p} |\nabla u(x)|^p \,\mathrm{d}x\right)^{\frac{l}{p}} \left(\int_{\mathbb{R}^N} |x|^{\beta q} |u(x)|^q \,\mathrm{d}x\right)^{\frac{1-l}{q}}, \,\forall u \in \mathcal{C}_c^{\infty}(\mathbb{R}^N),$$

if and only if the following relations hold

$$\frac{1}{r} + \frac{m}{N} = l\left(\frac{1}{p} + \frac{\alpha - 1}{N}\right) + (1 - l)\left(\frac{1}{q} + \frac{\beta}{N}\right)$$

with

$$0 \le \alpha - \gamma \ \text{if } l > 0,$$

and

$$\alpha - \gamma \leq 1$$
 if $l > 0$ and $\frac{1}{r} + \frac{m}{N} = \frac{1}{p} + \frac{\alpha - 1}{N}$

For $s \in (1,2), \sigma = s - 1, p \in (1, \frac{N}{s})$ and $a \in [0, \frac{N-sp}{2})$, we consider

$$r = p, l = 1, \alpha = -\frac{\sigma p + 2a}{p}, \text{ and } \gamma = -\frac{sp + 2a}{p}.$$
 (5.1)

Then, we notice that $m = \gamma = \alpha - 1$ and $\frac{1}{r} + \frac{m}{N} = \frac{1}{p} - \frac{sp+2a}{Np} = \frac{1}{p} + \frac{\alpha - 1}{N}$. Moreover,

$$\frac{1}{p} + \frac{\alpha}{N} = \frac{N - \sigma p - 2a}{Np} > 0 \text{ and } \frac{1}{r} + \frac{m}{N} = \frac{N - sp - 2a}{Np} > 0.$$

Hence the assumptions of Theorem 5.1 are verified, and we get

$$C(N, p, \sigma, a) \int_{\mathbb{R}^N} \frac{|u(x)|^p}{|x|^{sp+2a}} \,\mathrm{d}x \le \int_{\mathbb{R}^N} \frac{|\nabla u(x)|^p}{|x|^{\sigma p+2a}} \,\mathrm{d}x.$$
(5.2)

Moreover, by [1, Theorem 2.7],

$$C(N, p, \sigma, a) \int_{\mathbb{R}^N} \frac{|u(x)|^p}{|x|^{\sigma p+2a}} \, \mathrm{d}x \le \iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{|u(x) - u(y)|^p}{|x - y|^{N + \sigma p}} \, \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a}, \, \forall \, u \in \mathcal{C}^\infty_c(\mathbb{R}^N).$$
(5.3)

Now observe that

$$\begin{split} &\iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{|\nabla u(x) - \nabla u(y)|^p}{|x - y|^{N + \sigma p}} \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a} \\ &\geq B(N, \frac{p}{2}) \left(\sum_{i=1}^N \iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{|\frac{\partial u}{\partial x_i}(x) - \frac{\partial u}{\partial x_i}(y)|^p}{|x - y|^{N + \sigma p}} \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a} \right) \text{ (using Lemma 2.1-(ii))} \\ &\geq B(N, \frac{p}{2}) C \left(\sum_{i=1}^N \int_{\mathbb{R}^N} \frac{|\frac{\partial u}{\partial x_i}(x)|^p}{|x|^{\sigma p + 2a}} \mathrm{d}x \right) \text{ (using (5.3) with } \frac{\partial u}{\partial x_i} \text{)} \\ &\geq \frac{B(N, \frac{p}{2})}{A(N, \frac{p}{2})} C \int_{\mathbb{R}^N} \frac{|\nabla u(x)|^p}{|x|^{\sigma p + 2a}} \mathrm{d}x \text{ (using Lemma 2.1-(i))} \\ &\geq \frac{B(N, \frac{p}{2})}{A(N, \frac{p}{2})} C \int_{\mathbb{R}^N} \frac{|u(x)|^p}{|x|^{sp + 2a}} \mathrm{d}x, \text{ (using (5.2)).} \end{split}$$

Therefore, we have

Proposition 5.2. Let $N \ge 2, s \in (1, 2), \sigma = s - 1, p \in (1, \frac{N}{s}), and a \in [0, \frac{N-sp}{2})$. Then

$$\iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{|\nabla u(x) - \nabla u(y)|^p}{|x - y|^{N + \sigma p}} \frac{\mathrm{d}x}{|x|^a} \frac{\mathrm{d}y}{|y|^a} \ge C \int_{\mathbb{R}^N} \frac{|u(x)|^p}{|x|^{sp + 2a}} \,\mathrm{d}x, \,\forall \, u \in \mathcal{C}^\infty_c(\mathbb{R}^N), \tag{5.4}$$

where $C = C(N, p, \sigma, a)$.

Next, using (5.4), we study the finer embeddings of $\mathcal{D}_{a}^{s,p}(\mathbb{R}^{N})$. Let Ω be an open set in \mathbb{R}^{N} . For $f \in M(\Omega)$ we define the one dimensional decreasing rearrangement f^{*} of f as $f^{*}(\tau) = \inf\{t > 0 : \mu_{f}(t) < \tau\}$ for $\tau > 0$. Then, the following identity holds (see [11])

$$\int_{\Omega} f(x) \, \mathrm{d}x = \int_{0}^{|\Omega|} f^{*}(t) \, \mathrm{d}t.$$
(5.5)

Proof of Theorem 1.1-(iv): For $u \in C_c^{\infty}(\mathbb{R}^N)$, consider the set $A_t = \{x \in \mathbb{R}^N : |u(x)|^p > t\}$ and we write

$$|u(x)|^{p} = \int_{0}^{|u(x)|^{p}} dt = \int_{0}^{\infty} \chi_{t}(x) dt, \text{ where } \chi_{t}(x) = 1, \text{ if } x \in A_{t}, \text{ and } \chi_{t}(x) = 0, \text{ if } x \in A_{t}^{c}.$$

Using Fubini's theorem, we write

$$\int_{\mathbb{R}^N} \frac{|u(x)|^p}{|x|^{sp+2a}} \,\mathrm{d}x = \int_{\mathbb{R}^N} \left(\int_0^\infty \chi_t(x) \,\mathrm{d}t \right) \frac{\mathrm{d}x}{|x|^{sp+2a}} = \int_0^\infty \left(\int_{\mathbb{R}^N} \frac{\chi_t(x)}{|x|^{sp+2a}} \,\mathrm{d}x \right) \,\mathrm{d}t$$
$$x) = |x|^{-(sp+2a)} \text{ for } x \in \mathbb{R}^N \text{ Using } (5.5)$$

Denote $f(x) = |x|^{-(sp+2a)}$ for $x \in \mathbb{R}^N$. Using (5.5),

$$\int_{\mathbb{R}^N} f(x)\chi_t(x) \, \mathrm{d}x = \int_{A_t} f(x) \, \mathrm{d}x = \int_0^{|A_t|} f^*(t) \, \mathrm{d}t$$

By [4, Example 2.1.2], we see that $f^*(t) = \left(\frac{\omega_N}{t}\right)^{\frac{sp+2a}{N}}$, were ω_N is the measure of a unit ball in \mathbb{R}^N . This implies

$$\int_{\mathbb{R}^N} f(x)\chi_t(x) \,\mathrm{d}x = \omega_N^{\frac{sp+2a}{N}} \int_0^{|A_t|} \frac{\mathrm{d}t}{t^{\frac{sp+2a}{N}}} = \frac{N\omega_N^{\frac{sp+2a}{N}}}{N-sp-2a} |A_t|^{\frac{N-sp-2a}{N}}.$$

Observe that $|A_t| = \mu_{|u|^p}(t)$ for t > 0. Therefore,

$$\int_{\mathbb{R}^N} \frac{|u(x)|^p}{|x|^{sp+2a}} \, \mathrm{d}x = \frac{N\omega_N^{\frac{sp+2a}{N}}}{N-sp-2a} \int_0^\infty \mu_{|u|^p}(t)^{\frac{p}{p^\star}} \, \mathrm{d}t = \omega_N^{\frac{sp+2a}{N}} ||u|^p|_{\frac{p^\star}{p},1} = \omega_N^{\frac{sp+2a}{N}} |u|_{p^\star,p}^p.$$

Hence using (5.4) we obtain

$$|u|_{p^{\star},p} \le C(N,s,p,a)[u]_{s,p,a}, \ \forall u \in \mathcal{C}^{\infty}_{c}(\mathbb{R}^{N}).$$
(5.6)

Thus, (5.6) with the definition of $\mathcal{D}_a^{s,p}(\mathbb{R}^N)$ yields (1.10).

In the following proposition, using a similar layer cake decomposition technique, we show that $L^{p^*,p}(\mathbb{R}^N)$ is strictly contained in $L^{p^*,p}_{a}(\mathbb{R}^N)$.

Proposition 5.3. Let $N \ge 2, s \in (1,2), \sigma = s-1, p \in (1,\frac{N}{s})$, and $a \in [0,\frac{N-sp}{2})$. Then $L^{p^{\star},p}(\mathbb{R}^N) \subsetneq L_a^{p_s^{\star}}(\mathbb{R}^N)$, where $p^{\star} = \frac{Np}{N-sp-2a}$.

Proof. For t > 0, we consider the set $A_t = \{x \in \mathbb{R}^N : |u(x)|^{p_s^*} > t\}$, and write

$$|u(x)|^{p_s^*} = \int_0^{|u(x)|^{p_s^*}} \mathrm{d}t = \int_0^\infty \chi_t(x) \,\mathrm{d}t, \text{ where } \chi_t(x) = 1, \text{ if } x \in A_t, \text{ and } \chi_t(x) = 0, \text{ if } x \in A_t^c.$$
Using Fubini's theorem,

 $\int_{\mathbb{R}^N} \frac{|u(x)|^{p_s^*}}{|x|^{\frac{2ap_s^*}{p}}} \,\mathrm{d}x = \int_{\mathbb{R}^N} \left(\int_0^\infty \chi_t(x) \,\mathrm{d}t \right) \frac{\mathrm{d}x}{|x|^{\frac{2ap_s^*}{p}}} = \int_0^\infty \left(\int_{\mathbb{R}^N} \frac{\chi_t(x)}{|x|^{\frac{2ap_s^*}{p}}} \,\mathrm{d}x \right) \,\mathrm{d}t.$

Denote $f(x) = |x|^{-\frac{2ap_s^*}{p}}$ for $x \in \mathbb{R}^N$. Using (5.5),

$$\int_{\mathbb{R}^N} f(x)\chi_t(x) \, \mathrm{d}x = \int_{A_t} f(x) \, \mathrm{d}x = \int_0^{|A_t|} f^*(t) \, \mathrm{d}t$$

By [4, Example 2.1.2], $f^*(t) = \left(\frac{\omega_N}{t}\right)^{\frac{2a}{N-sp}}$ and hence

$$\int_{\mathbb{R}^N} f(x)\chi_t(x) \,\mathrm{d}x = \omega_N^{\frac{2a}{N-sp}} \int_0^{|A_t|} \frac{\mathrm{d}t}{t^{\frac{2a}{N-sp}}} = \frac{\omega_N^{\frac{N-sp}{N-sp}}(N-sp)}{N-sp-2a} |A_t|^{\frac{N-sp-2a}{N-sp}}$$

where $|A_t| = \mu_{|u|^{p_s^*}}(t)$ for t > 0. Therefore,

$$\int_{\mathbb{R}^N} \frac{|u(x)|^{p_s^*}}{|x|^{\frac{2ap_s^*}{p}}} \,\mathrm{d}x = \frac{\omega_N^{\frac{N-sp}{N-sp}}(N-sp)}{N-sp-2a} \int_0^\infty \mu_{|u|^{p_s^*}}(t)^{\frac{p_s^*}{p^*}} \,\mathrm{d}t = \omega_N^{\frac{2a}{N-sp}} ||u|^{p_s^*}|_{p_s^*,1}^{\frac{p^*}{N-sp}} = \omega_N^{\frac{2a}{N-sp}} |u|^{p_s^*}|_{p_s^*,p_s^*}^{\frac{2a}{N-sp}}.$$

We know that $L^{q,r}(\mathbb{R}^N) \subsetneq L^{q,s}(\mathbb{R}^N)$ for r < s. Taking $q = p^*, s = p_s^*$ and r = p yields

$$L^{p^{\star},p}(\mathbb{R}^{N}) \subsetneq L^{p^{\star},p^{\star}_{s}}(\mathbb{R}^{N}) = L^{p^{\star}_{s}}_{a^{\star}}(\mathbb{R}^{N}).$$
s.
$$\Box$$
(5.7)

20

Hence the strict inclusion holds.

Remark 5.4. From Remark 3.3 and Theorem 1.1-(iv), we have

$$\mathring{W}^{s,p}_{a}(\mathbb{R}^{N}) \subset L^{p^{\star},p}(\mathbb{R}^{N}), \text{ for } s \in (1,2) \text{ and } p \in (1,\frac{N}{s}).$$

Acknowledgments. The first author acknowledges the Science and Engineering Research Board, Government of India, for national postdoctoral fellowship. The second author acknowledges the support of the CSIR fellowship, file no. 09/1125(0016)/2020–EMR–I.

References

- B. Abdellaoui and R. Bentifour. Caffarelli-Kohn-Nirenberg type inequalities of fractional order with applications. J. Funct. Anal., 272(10):3998–4029, 2017. 3, 16, 21
- [2] T. V. Anoop and N. Biswas. Neumann eigenvalue problems on the exterior domains. Nonlinear Anal., 187:339– 351, 2019. 3
- [3] T. V. Anoop, N. Biswas, and U. Das. Admissible function spaces for weighted Sobolev inequalities. Commun. Pure Appl. Anal., 20(9):3259–3297, 2021. 4
- [4] N. Biswas. The (p,q)-Hardy Inequalities and application to nonlinear eigenvalue problems. PhD thesis, Indian Institute of Technology Madras, Chennai, 2021. 6, 22, 23
- [5] L. Brasco, D. Gómez-Castro, and J. L. Vázquez. Characterisation of homogeneous fractional Sobolev spaces. Calc. Var. Partial Differential Equations, 60(2):Paper No. 60, 40, 2021. 3
- [6] H. Brézis. Functional analysis, Sobolev spaces and partial differential equations. Universitext. Springer, New York, 2011. 3, 7
- [7] L. Caffarelli, R. Kohn, and L. Nirenberg. First order interpolation inequalities with weights. Compositio Math., 53(3):259-275, 1984. 18, 21
- [8] J. Deny and J. L. Lions. Les espaces du type de Beppo Levi. Ann. Inst. Fourier (Grenoble), 21:305–370, 1945. 2
- [9] S. Dipierro, L. Montoro, I. Peral, and B. Sciunzi. Qualitative properties of positive solutions to nonlocal critical problems involving the Hardy-Leray potential. *Calc. Var. Partial Differential Equations*, 55(4):Art. 99, 29, 2016.

- [10] S. Dipierro and E. Valdinoci. A density property for fractional weighted Sobolev spaces. Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl., 26(4):397–422, 2015. 3, 7, 8, 9, 11, 16
- [11] D. E. Edmunds and W. D. Evans. Hardy operators, function spaces and embeddings. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2004. 4, 6, 22
- [12] R. L. Frank, E. H. Lieb, and R. Seiringer. Hardy-Lieb-Thirring inequalities for fractional Schrödinger operators. J. Amer. Math. Soc., 21(4):925–950, 2008. 3
- [13] R. L. Frank and R. Seiringer. Non-linear ground state representations and sharp Hardy inequalities. J. Funct. Anal., 255(12):3407–3430, 2008. 3
- [14] G. Leoni. A first course in fractional Sobolev spaces, volume 229 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, [2023] ©2023.
- [15] E. H. Lieb and M. Loss. Analysis, volume 14 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, second edition, 2001. 6
- [16] G. G. Lorentz. Some new functional spaces. Ann. of Math. (2), 51:37–55, 1950. 6
- [17] V. Maz'ya. Sobolev spaces with applications to elliptic partial differential equations, volume 342 of Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer, Heidelberg, augmented edition, 2011. 3
- [18] M. Milman and E. Pustylnik. On sharp higher order Sobolev embeddings. Commun. Contemp. Math., 6(3):495– 511, 2004. 4
- [19] J. Peetre. Espaces d'interpolation et théorème de Soboleff. Ann. Inst. Fourier (Grenoble), 16:279–317, 1966. 3
- [20] L. Tartar. Imbedding theorems of Sobolev spaces into Lorentz spaces. Boll. Unione Mat. Ital. Sez. B Artic. Ric. Mat. (8), 1(3):479–500, 1998. 3
- [21] K. Yosida. Functional analysis. Classics in Mathematics. Springer-Verlag, Berlin, 1995. Reprint of the sixth (1980) edition. 20