

QUASICONFORMAL MAPPINGS AND THE RANK OF $\frac{dDf}{d|Df|}$ FOR $f \in \text{BV}(\mathbb{R}^n; \mathbb{R}^n)$

PANU LAHTI

ABSTRACT. We define a relaxed version H_f^{fine} of the distortion number H_f that is used to define quasiconformal mappings. Then we show that for a BV function $f \in \text{BV}(\mathbb{R}^n; \mathbb{R}^n)$, for $|Df|$ -a.e. $x \in \mathbb{R}^n$ it holds that $H_f^{\text{fine}}(x) < \infty$ if and only if $\frac{dDf}{d|Df|}(x)$ has full rank.

1. INTRODUCTION

Quasiconformal mappings are defined as homeomorphisms $f \in W_{\text{loc}}^{1,n}(\mathbb{R}^n; \mathbb{R}^n)$, $n \geq 2$, for which $|\nabla f(x)|^n \leq K |\det \nabla f(x)|$ for a.e. $x \in \mathbb{R}^n$ and a constant $K < \infty$. In particular, ∇f has full rank a.e. Quasiconformal mappings can also be defined between two metric spaces (X, d_X) and (Y, d_Y) . For a mapping $f: X \rightarrow Y$, for every $x \in X$ and $r > 0$ one defines

$$L_f(x, r) := \sup\{d_Y(f(y), f(x)) : d_X(y, x) \leq r\}$$

and

$$l_f(x, r) := \inf\{d_Y(f(y), f(x)) : d_X(y, x) \geq r\},$$

and

$$H_f(x, r) := \frac{L_f(x, r)}{l_f(x, r)};$$

we interpret this to be ∞ if either the numerator is infinity or the denominator is zero. Then one defines

$$H_f(x) := \limsup_{r \rightarrow 0} H_f(x, r).$$

A homeomorphism $f: X \rightarrow Y$ is said to be (metric) *quasiconformal* if there is a number $1 \leq H < \infty$ such that $H_f(x) \leq H$ for all $x \in X$. In Euclidean spaces and sufficiently regular metric measure spaces, this is equivalent with the ‘‘analytic’’ definition mentioned above, see e.g. [6, Theorem 9.8].

It is natural to ask whether the analytic condition of ∇f having full rank is connected with the geometric condition of H_f being finite also for more general mappings such as $f \in W_{\text{loc}}^{1,1}(\mathbb{R}^n; \mathbb{R}^n)$, or even functions of bounded variation. The answer is in general negative, since $L_f(x, r)$ can easily be ∞ for every $x \in \mathbb{R}^n$ and $r > 0$ already for a mapping

Date: June 11, 2024.

2020 Mathematics Subject Classification. 46E35, 31C40, 30C65.

Key words and phrases. Function of bounded variation, quasiconformal mapping, finely open set, Alberti’s rank one theorem.

$f \in W_{\text{loc}}^{1,n}(\mathbb{R}^n; \mathbb{R}^n)$; see Example 3.6. With this in mind, given $U \subset \mathbb{R}^n$ containing $x \in \mathbb{R}^n$ and $r > 0$, we let

$$L_{f,U}(x, r) := \sup\{|f(x) - f(z)| : |z - x| \leq r, z \in U\}$$

and

$$l_{f,U}(x, r) := \inf\{|f(x) - f(z)| : |z - x| \geq r, z \in U\},$$

and then

$$H_{f,U}(x, r) := \frac{L_{f,U}(x, r)}{l_{f,U}(x, r)} \quad \text{and} \quad H_f^{\text{fine}}(x) := \inf_U \limsup_{r \rightarrow 0} H_{f,U}(x, r),$$

where the infimum is taken over *1-finely open sets* U containing x . We give definitions in Section 2.

Now it turns out that H_f^{fine} and the rank of the Jacobian are connected in the following way.

Theorem 1.1. *Let $f \in \text{BV}(\mathbb{R}^n; \mathbb{R}^n)$, $n \geq 2$. Then for $|Df|$ -a.e. $x \in \mathbb{R}^n$, we have $H_{f^*}^{\text{fine}}(x) < \infty$ if and only if $\frac{dDf}{d|Df|}(x)$ has full rank.*

Here f^* is the precise representative of f .

2. NOTATION AND DEFINITIONS

2.1. Basic notation and definitions. We will always work in the Euclidean space \mathbb{R}^n , $n \geq 1$. We denote the n -dimensional Lebesgue measure by \mathcal{L}^n . We denote the s -dimensional Hausdorff measure by \mathcal{H}^s , $s \geq 0$. If a property holds outside a set of Lebesgue measure zero, we say that it holds almost everywhere, or ‘‘a.e.’’. With other measures, we write more explicitly e.g. ‘‘ \mathcal{H}^{n-1} -a.e.’’. The Borel σ -algebra on a set $H \subset \mathbb{R}^n$ is denoted by $\mathcal{B}(H)$.

We write $B(x, r)$ for an open ball in \mathbb{R}^n with center x and radius r , that is, $\{y \in \mathbb{R}^n : |y - x| < r\}$. We denote the \mathcal{L}^n -measure of the unit ball by ω_n . We always work with the Euclidean norm $|\cdot|$ for vectors $v \in \mathbb{R}^n$ as well as for matrices $A \in \mathbb{R}^{k \times n}$, $k \geq 1$. For $a \in \mathbb{R}^k$ and $b \in \mathbb{R}^n$, we define the tensor product $a \otimes b := ab^T \in \mathbb{R}^{k \times n}$, where a, b are considered as column vectors. Given $x \in \mathbb{R}^n$, $r > 0$, and $\nu \in \mathbb{R}^n$ with $|\nu| = 1$, we denote by $Q_\nu(x, r)$ an open cube with center x and side length r , such that ν is perpendicular to one of the faces. Note that such a cube is not unique. We denote by $Q(x, r)$ an open cube with sides in the directions of the coordinate axes (such a cube is unique).

If a function f is in $L^1(H; \mathbb{R}^k)$, for $k \in \mathbb{N}$ and for some \mathcal{L}^n -measurable set $H \subset \mathbb{R}^n$ of nonzero and finite Lebesgue measure, we write

$$f_H := \int_H f(y) d\mathcal{L}^n(y) := \frac{1}{\mathcal{L}^n(H)} \int_H f(y) d\mathcal{L}^n(y)$$

for its mean value in H .

We will always denote by $\Omega \subset \mathbb{R}^n$ an open set. Let $k \in \mathbb{N}$. The Sobolev space $W^{1,1}(\Omega; \mathbb{R}^k)$ consists of functions $f \in L^1(\Omega; \mathbb{R}^k)$ whose first weak partial derivatives $D_l f_j$, $j = 1, \dots, k$, $l = 1, \dots, n$, belong to $L^1(\Omega)$. The Sobolev norm is

$$\|f\|_{W^{1,1}(\Omega; \mathbb{R}^k)} := \|f\|_{L^1(\Omega; \mathbb{R}^k)} + \|\nabla f\|_{L^1(\Omega; \mathbb{R}^{k \times n})}.$$

Let $S \subset \mathbb{R}^n$ be an \mathcal{H}^{n-1} -measurable set. We say that S is countably \mathcal{H}^{n-1} -rectifiable if there exist countably many Lipschitz functions $h_j: \mathbb{R}^{n-1} \rightarrow \mathbb{R}^n$ such that

$$\mathcal{H}^{n-1} \left(S \setminus \bigcup_{j=1}^{\infty} h_j(\mathbb{R}^{n-1}) \right) = 0.$$

Let $H \subset \mathbb{R}^n$ be an $n - 1$ -dimensional hyperplane, let π be the orthogonal projection onto H , and let π^\perp be the orthogonal projection onto the orthogonal complement H^\perp . Let $h: H \rightarrow H^\perp$ be an L -Lipschitz function, and let

$$S := \{x \in \mathbb{R}^n : h(\pi(x)) = \pi^\perp(x)\}$$

be the graph of h . We call this an L -Lipschitz $n - 1$ -graph, also if h is only defined on a subset of H . Every countably \mathcal{H}^{n-1} -rectifiable set can be presented, modulo an \mathcal{H}^{n-1} -negligible set, as an at most countable union of disjoint 1-Lipschitz $n - 1$ -graphs, see [2, Proposition 2.76]. We can also assume the 1-Lipschitz $n - 1$ -graphs to be \mathcal{H}^{n-1} -measurable, since a graph defined on an entire $n - 1$ -dimensional hyperplane is a closed set.

2.2. Radon measures. In this and the following subsections, we will follow the monograph [2]. Let $\Omega \subset \mathbb{R}^n$ be an open set and $\ell \in \mathbb{N}$. We denote by $\mathcal{M}(\Omega; \mathbb{R}^\ell)$ the Banach space of vector-valued Radon measures. For $\mu \in \mathcal{M}(\Omega; \mathbb{R}^\ell)$, the total variation $|\mu|(\Omega)$ is defined with respect to the Euclidean norm on \mathbb{R}^ℓ . We further denote the set of positive measures by $\mathcal{M}^+(\Omega)$.

For a vector-valued Radon measure $\gamma \in \mathcal{M}(\Omega; \mathbb{R}^\ell)$ and a positive Radon measure $\mu \in \mathcal{M}^+(\Omega)$, we can write the Lebesgue–Radon–Nikodym decomposition

$$\gamma = \gamma^a + \gamma^s = \frac{d\gamma}{d\mu} d\mu + \gamma^s$$

of γ with respect to μ , where $\frac{d\gamma}{d\mu} \in L^1(\Omega, \mu; \mathbb{R}^\ell)$.

For open sets $E \subset \mathbb{R}^{n-m}$, $F \subset \mathbb{R}^m$ with $m \in \{1, \dots, n - 1\}$, a *parametrized measure* $(\nu_y)_{y \in E}$ is a mapping from E to the set $\mathcal{M}(F; \mathbb{R}^\ell)$ of vector-valued Radon measures on F . It is said to be *weakly* μ -measurable*, for $\mu \in \mathcal{M}^+(E)$, if $y \mapsto \nu_y(B)$ is μ -measurable for all Borel sets $B \in \mathcal{B}(F)$ (it suffices to check this for open subsets). Equivalently, $(\nu_y)_{y \in E}$ is weakly* μ -measurable if the function $y \mapsto \int_F f(y, t) d\nu_x(t)$ is μ -measurable for every bounded $\mathcal{B}_\mu(E) \times \mathcal{B}(F)$ -measurable function $f: E \times F \rightarrow \mathbb{R}$ (see [2, Proposition 2.26]),

where $\mathcal{B}_\mu(E)$ denotes the μ -completion of $\mathcal{B}(E)$. Suppose that we additionally have

$$\int_E |\nu_y|(F) d\mu(y) < \infty.$$

In this case we denote by $\mu \otimes \nu_y$ the generalized product measure defined by

$$\mu \otimes \nu_y(A) := \int_E \left(\int_F \chi_A(y, t) d\nu_y(t) \right) d\mu(y) \quad (2.1)$$

for every $A \in \mathcal{B}_\mu(E) \times \mathcal{B}(F)$.

2.3. Functions of bounded variation. We follow the monograph [2, Section 3]. Let $k \in \mathbb{N}$, and as before let $\Omega \subset \mathbb{R}^n$ be an open set. A function $f \in L^1(\Omega; \mathbb{R}^k)$ is a function of bounded variation, denoted $f \in \text{BV}(\Omega; \mathbb{R}^k)$, if its weak derivative is an $\mathbb{R}^{k \times n}$ -valued Radon measure with finite total variation. This means that there exists a (unique) Radon measure Df such that for all $\varphi \in C_c^1(\Omega)$, the integration-by-parts formula

$$\int_\Omega f^j \frac{\partial \varphi}{\partial y_l} d\mathcal{L}^n = - \int_\Omega \varphi dD_l f^j, \quad j = 1, \dots, k, \quad l = 1, \dots, n,$$

holds. If f is the characteristic function of a set $E \subset \mathbb{R}^n$, we say that E is a set of finite perimeter.

Let $f \in L^1_{\text{loc}}(\Omega)$. The precise representative is defined by

$$f^*(x) := \limsup_{r \rightarrow 0} \int_{B(x,r)} f d\mathcal{L}^n, \quad x \in \Omega.$$

This is easily seen to be a Borel function. For $f \in L^1_{\text{loc}}(\Omega; \mathbb{R}^k)$, we also define $f^* := (f_1^*, \dots, f_k^*)$.

We say that $x \in \Omega$ is a Lebesgue point of f if

$$\lim_{r \rightarrow 0} \int_{B(x,r)} |f(y) - \tilde{f}(x)| d\mathcal{L}^n(y) = 0$$

for some $\tilde{f}(x) \in \mathbb{R}^k$. We denote by $S_f \subset \Omega$ the set where this condition fails and call it the approximate discontinuity set.

Given $x \in \mathbb{R}^n$ and $r > 0$, and a unit vector $\nu \in \mathbb{R}^n$, we define the half-balls

$$\begin{aligned} B_\nu^+(x, r) &:= \{y \in B(x, r) : \langle y - x, \nu \rangle > 0\}, \\ B_\nu^-(x, r) &:= \{y \in B(x, r) : \langle y - x, \nu \rangle < 0\}, \end{aligned}$$

where $\langle \cdot, \cdot \rangle$ denotes the inner product. We say that $x \in \Omega$ is an approximate jump point of f if there exist a unit vector $\nu \in \mathbb{R}^n$ and distinct vectors $f^+(x), f^-(x) \in \mathbb{R}^k$ such that

$$\lim_{r \rightarrow 0} \int_{B_\nu^+(x,r)} |f(y) - f^+(x)| d\mathcal{L}^n(y) = 0 \quad (2.2)$$

and

$$\lim_{r \rightarrow 0} \int_{B_\nu^-(x,r)} |f(y) - f^-(x)| d\mathcal{L}^n(y) = 0. \quad (2.3)$$

The set of all approximate jump points is denoted by J_f , and the functions f^-, f^+ can be taken to be Borel functions. For $f \in \text{BV}(\Omega; \mathbb{R}^k)$, we have that $\mathcal{H}^{n-1}(S_f \setminus J_f) = 0$, see [2, Theorem 3.78].

The lower and upper approximate limits of a function $f \in \text{BV}_{\text{loc}}(\Omega)$ are defined respectively by

$$f^\wedge(x) := \sup \left\{ t \in \mathbb{R} : \lim_{r \rightarrow 0} \frac{\mathcal{L}^n(B(x, r) \cap \{f < t\})}{\mathcal{L}^n(B(x, r))} = 0 \right\}$$

and

$$f^\vee(x) := \inf \left\{ t \in \mathbb{R} : \lim_{r \rightarrow 0} \frac{\mathcal{L}^n(B(x, r) \cap \{f > t\})}{\mathcal{L}^n(B(x, r))} = 0 \right\},$$

for all $x \in \Omega$. We interpret the supremum and infimum of an empty set to be $-\infty$ and ∞ , respectively. Note that

$$\tilde{f}(x) = f^\wedge(x) = f^\vee(x) \quad \text{for } x \in \Omega \setminus S_f \quad (2.4)$$

and

$$f^\wedge(x) = \min\{f^-(x), f^+(x)\} \quad \text{and} \quad f^\vee(x) = \max\{f^-(x), f^+(x)\} \quad \text{for } x \in J_f. \quad (2.5)$$

Given $f \in \text{BV}(\Omega; \mathbb{R}^k)$, for every $x \in J_f$ we moreover have

$$f^*(x) = \frac{f^-(x) + f^+(x)}{2}. \quad (2.6)$$

We write the Radon-Nikodym decomposition of the variation measure of f into the absolutely continuous and singular parts with respect to \mathcal{L}^n as

$$Df = D^a f + D^s f = \nabla f \mathcal{L}^n + D^s f.$$

Furthermore, we define the Cantor and jump parts of Df by

$$D^c f := D^s f \llcorner (\Omega \setminus S_f), \quad D^j f := D^s f \llcorner J_f.$$

Here

$$D^s f \llcorner J_f(A) := D^s f(J_f \cap A), \quad \text{for } D^s f\text{-measurable } A \subset \Omega.$$

Since $\mathcal{H}^{n-1}(S_f \setminus J_f) = 0$ and $|Df|$ vanishes on \mathcal{H}^{n-1} -negligible sets, we get the decomposition (see [2, Section 3.9])

$$Df = D^a f + D^c f + D^j f. \quad (2.7)$$

For the jump part, we know that

$$d|D^j f| = |f^+ - f^-| d\mathcal{H}^{n-1} \llcorner J_f. \quad (2.8)$$

We say that a sequence $\{f_i\}_{i=1}^\infty$ of functions in $\text{BV}(\Omega; \mathbb{R}^k)$ converges to $f \in \text{BV}(\Omega; \mathbb{R}^k)$ strictly in $\text{BV}(\Omega; \mathbb{R}^k)$ if

$$f_i \rightarrow f \quad \text{in } L^1(\Omega; \mathbb{R}^k) \quad \text{and} \quad |Df_i|(\Omega) \rightarrow |Df|(\Omega) \quad \text{as } i \rightarrow \infty.$$

2.4. One-dimensional sections of BV functions. The following notation and results on one-dimensional sections of BV functions are given in [2, Section 3.11].

Let $n = 1$. Suppose $f \in \text{BV}_{\text{loc}}(\mathbb{R})$. We have $J_f = S_f$, J_f is at most countable, and $Df(\{x\}) = 0$ for every $x \in \mathbb{R} \setminus J_f$. For every $x, \tilde{x} \in \mathbb{R} \setminus J_f$ with $x < \tilde{x}$ we have

$$f^*(\tilde{x}) - f^*(x) = Df((x, \tilde{x})). \quad (2.9)$$

For every $x, \tilde{x} \in \mathbb{R}$ with $x < \tilde{x}$ we have

$$|f^*(x) - f^*(\tilde{x})| \leq |Df|([x, \tilde{x}]), \quad (2.10)$$

and the same with f^* replaced by f^\wedge or f^\vee , or any pairing of these.

We say that $f^\vee(x) - f^\wedge(x) = |Df|(\{x\})$ is the *jump size* of f at point x . For a jump point $x \in J_f$ and a sequence $x_j \rightarrow x$, we have

$$\lim_{j \rightarrow \infty} \min\{|f^*(x_j) - f^\wedge(x)|, |f^*(x_j) - f^\vee(x)|\} = 0 \quad (2.11)$$

and the same with $f^*(x_j)$ replaced by $f^\wedge(x_j)$ or $f^\vee(x_j)$.

For a point $x = (x_1, \dots, x_n) \in \mathbb{R}^n$, denote by π_n the projection

$$\pi_n(x) := (x_1, \dots, x_{n-1}),$$

and analogously we define π_j for $j = 1, \dots, n-1$. We also denote balls and cubes in \mathbb{R}^{n-1} by $B_{n-1}(x, r)$ and $Q_{n-1}(x, r)$, respectively.

In \mathbb{R}^n , we denote the standard basis vectors by e_i , $i = 1, \dots, n$. For a set $A \subset \mathbb{R}^n$ we denote the slices of A by

$$A_z^n := \{t \in \mathbb{R} : (z, t) \in A\}, \quad z \in \pi_n(A).$$

Analogously, we define A_z^j for all $j = 1, \dots, n$. Usually we consider A_z^n ; denote it by A_z . For $f \in \text{BV}(\Omega; \mathbb{R}^k)$, we denote

$$f_{z,n}(t) := f(z, t), \quad t \in \Omega_z, \quad z \in \pi_n(\Omega).$$

Again to avoid too heavy notation, write f_z instead of $f_{z,n}$. We know that $f_z \in \text{BV}(\Omega_z)$ for \mathcal{L}^{n-1} -almost every $z \in \pi_n(\Omega)$ (see [2, Theorem 3.103]).

Analogously, we define $f_{z,l}$ for all $l = 1, \dots, n$.

Recall (2.1). Denoting $D_l f := \langle Df, e_l \rangle$, we further have

$$D_l f = \mathcal{L}^{n-1} \otimes Df_{z,l} \quad \text{and} \quad D_l^j f = \mathcal{L}^{n-1} \otimes D^j f_{z,l},$$

see [2, Theorem 3.107 & Theorem 3.108]. It follows that

$$|D_l f| = \mathcal{L}^{n-1} \otimes |Df_{z,l}| \quad \text{and} \quad |D_l^j f| = \mathcal{L}^{n-1} \otimes |D^j f_{z,l}|, \quad (2.12)$$

see [2, Corollary 2.29]. Moreover, for \mathcal{L}^{n-1} -almost every $z \in \pi_n(\Omega)$ we have

$$J_{f_z} = (J_f)_z \quad \text{and} \quad (f^*)_z(t) = (f_z)^*(t) \quad \text{for every } t \in \mathbb{R} \setminus J_{f_z}, \quad (2.13)$$

see [2, Theorem 3.108]. By [2, Theorem 3.108] we also know that for \mathcal{L}^{n-1} -almost every $z \in \pi_n(\Omega)$, we have

$$\{(f_z)^-(t), (f_z)^+(t)\} = \{(f^-)_z(t), (f^+)_z(t)\} \quad \text{for every } t \in (J_f)_z.$$

Assume $k = 1$ so that $f \in \text{BV}(\Omega)$; recalling (2.4) and (2.5), for \mathcal{L}^{n-1} -almost every $z \in \pi_n(\Omega)$ we have

$$(f_z)^\wedge(t) = (f^\wedge)_z(t) \quad \text{and} \quad (f_z)^\vee(t) = (f^\vee)_z(t) \quad \text{for every } t \in \Omega_z. \quad (2.14)$$

2.5. Capacities and fine topology. The (Sobolev) 1-capacity of a set $A \subset \mathbb{R}^n$ is defined by

$$\text{Cap}_1(A) := \inf \|u\|_{W^{1,1}(\mathbb{R}^n)},$$

where the infimum is taken over Sobolev functions $u \in W^{1,1}(\mathbb{R}^n)$ satisfying $u \geq 1$ in a neighborhood of A .

Given sets $A \subset W \subset \mathbb{R}^n$, where W is open, the relative 1-capacity is defined by

$$\text{cap}_1(A, W) := \inf \int_W |\nabla u| d\mathcal{L}^n,$$

where the infimum is taken over functions $u \in W_0^{1,1}(W)$ satisfying $u \geq 1$ in a neighborhood of A . The class $W_0^{1,1}(W)$ is the closure of $C_c^1(W)$ in the $W^{1,1}(\mathbb{R}^n)$ -norm.

By [3, Proposition 6.16], we know that for a ball $B(x, r)$ and $A \subset B(x, r)$, we have

$$\frac{\text{Cap}_1(A)}{1+r} \leq C \text{cap}_1(A, B(x, 2r)), \quad (2.15)$$

where C is a constant depending only on n .

For the proof of the following fact, see e.g. [10, Lemma 2.16].

Lemma 2.16. *Suppose $x \in \mathbb{R}^n$, $0 < r < 1$, and $A \subset B(x, r)$. Then we have*

$$\frac{\mathcal{L}^n(A)}{\mathcal{L}^n(B(x, r))} \leq C \frac{\text{Cap}_1(A)}{r^{n-1}} \quad \text{and} \quad \text{cap}_1(A, B(x, 2r)) \leq C \text{Cap}_1(A),$$

where C is a constant depending only on n .

Definition 2.17. We say that $A \subset \mathbb{R}^n$ is 1-thin at the point $x \in \mathbb{R}^n$ if

$$\lim_{r \rightarrow 0} \frac{\text{Cap}_1(A \cap B(x, r))}{r^{n-1}} = 0.$$

We say that a set $U \subset \mathbb{R}^n$ is 1-finely open if $\mathbb{R}^n \setminus U$ is 1-thin at every $x \in U$. Then we define the 1-fine topology as the collection of 1-finely open sets on \mathbb{R}^n .

See also [10, Section 4] for discussion on Definition 2.17. In fact, in [10] the criterion

$$\lim_{r \rightarrow 0} \frac{\text{cap}_1(A \cap B(x, r), B(x, 2r))}{r^{n-1}} = 0$$

for 1-thinness was used, in the context of more general metric measure spaces. By (2.15) and Lemma 2.16, this is equivalent with our current definition in the Euclidean setting.

3. PRELIMINARY RESULTS

In this section we collect some preliminary results.

We have the following quasi-semicontinuity result from [4, Theorem 2.5]. Alternatively, see [13, Theorem 1.1] and [11, Corollary 4.2] for a proof of this result in more general metric spaces.

Theorem 3.1. *Let $\Omega \subset \mathbb{R}^n$ be open, let $f \in \text{BV}_{\text{loc}}(\Omega)$, and $\varepsilon > 0$. Then there exists an open set $G \subset \Omega$ such that $\text{Cap}_1(G) < \varepsilon$ and $f^\wedge|_{\Omega \setminus G}$ is finite and lower semicontinuous, and $f^\vee|_{\Omega \setminus G}$ is finite and upper semicontinuous.*

Lemma 3.2. *Let $A \subset \mathbb{R}^n$. Then*

$$2\mathcal{H}^{n-1}(\pi_n(A)) \leq \text{Cap}_1(A).$$

Proof. Consider $u \in W^{1,1}(\mathbb{R}^n)$ with $u \geq 1$ in a neighborhood of A . Then for \mathcal{L}^{n-1} -a.e. $z \in \pi_n(A)$, we have

$$\int_{A_z} \left| \frac{\partial u}{\partial x_n} \right| ds \geq 2.$$

Integrating over \mathbb{R}^{n-1} , we get

$$\int_{\mathbb{R}^n} |\nabla u| dx \geq 2\mathcal{H}^{n-1}(\pi_n(A)).$$

Thus $\|u\|_{W^{1,1}(\mathbb{R}^n)} \geq 2\mathcal{H}^{n-1}(\pi_n(A))$ and we get the result by taking infimum over all such u . \square

Lemma 3.3. *Let $D \subset \mathbb{R}^{n-1}$, let $h: D \rightarrow \mathbb{R}$ be 1-Lipschitz, and let $S \subset \mathbb{R}^n$ be the graph of h , that is, $S = \{x = (z, t) \in D \times \mathbb{R} : t = h(z)\}$. Then*

$$\mathcal{H}^{n-1}(S) \leq (2\sqrt{2})^{n-1} \text{Cap}_1(S).$$

Proof. By Lemma 3.2, we have $2\mathcal{H}^{n-1}(D) \leq \text{Cap}_1(S)$. On the other hand, since h is 1-Lipschitz, it is easy to check from the definition of Hausdorff measures that $\mathcal{H}^{n-1}(S) \leq (2\sqrt{2})^{n-1}\mathcal{H}^{n-1}(D)$, and so we get the result. \square

Lemma 3.4. *Let S be a countably \mathcal{H}^{n-1} -rectifiable set with $\mathcal{H}^{n-1}(S) < \infty$. Then for \mathcal{H}^{n-1} -a.e. $x \in S$, given any 1-finely open set $U \ni x$ we have*

$$\lim_{r \rightarrow 0} \frac{\mathcal{H}^{n-1}(B(x, r) \cap S \cap U)}{\omega_{n-1}r^{n-1}} = 1.$$

Proof. We have $S = \bigcup_{j=1}^{\infty} S_j \cup N$, where each S_j is a \mathcal{H}^{n-1} -measurable 1-Lipschitz $n-1$ -graph and $\mathcal{H}^{n-1}(N) = 0$. Consider $x \in S_j$; we can assume that S_j is the graph of $h: D \rightarrow \mathbb{R}$ where $D \subset \mathbb{R}^{n-1}$ and h is 1-Lipschitz. Excluding \mathcal{H}^{n-1} -negligible sets, we can also assume that

$$\lim_{r \rightarrow 0} \frac{\mathcal{H}^{n-1}(B(x, r) \cap S_j)}{\omega_{n-1}r^{n-1}} = 1 \quad \text{and} \quad \lim_{r \rightarrow 0} \frac{\mathcal{H}^{n-1}(B(x, r) \cap S \setminus S_j)}{\omega_{n-1}r^{n-1}} = 0,$$

see [2, Theorem 2.56, Theorem 2.83]. By Lemma 3.3,

$$\begin{aligned} \limsup_{r \rightarrow 0} \frac{\mathcal{H}^{n-1}(B(x, r) \cap S_j \setminus U)}{\omega_{n-1} r^{n-1}} &\leq (2\sqrt{n})^{n-1} \limsup_{r \rightarrow 0} \frac{\text{Cap}_1(B(x, r) \cap S_j \setminus U)}{\omega_{n-1} r^{n-1}} \\ &= 0, \end{aligned}$$

since U is 1-finely open. Combining these facts, we get the result. \square

Recall Definition 2.17. The following theorem is proved in [10, Theorem 3.6].

Theorem 3.5. *Let $f \in \text{BV}_{\text{loc}}(\mathbb{R}^n; \mathbb{R}^n)$. Then for a.e. $x \in \mathbb{R}^n$ we find a 1-finely open set $U \ni x$ such that*

$$\lim_{U \setminus \{x\} \ni y \rightarrow x} \frac{|f^*(y) - f^*(x) - \nabla f(x)(y - x)|}{|y - x|} = 0.$$

Note that in general we understand ∇f to be the density of the absolutely continuous part of Df , but at a.e. $x \in \mathbb{R}^n$ we can understand $\nabla f(x)$ to be pointwise defined according to the above theorem.

Example 3.6. Let $\{q_j\}_{j=1}^\infty$ be an enumeration of points in \mathbb{R}^n with rational coordinates. Let $f \in W_{\text{loc}}^{1,n}(\mathbb{R}^n; \mathbb{R}^n)$ be such that the first component function is

$$f_1(x) := \sum_{j=1}^{\infty} 2^{-j} \log \max\{-\log|x - q_j|, 1\}, \quad x \in \mathbb{R}^n.$$

Now clearly $L_f(x, r) = \infty$ for every $x \in \mathbb{R}^n$ and $r > 0$. Thus $H_f = \infty$ everywhere, while with a suitable choice of the other component functions, we can have that ∇f has full rank in a set of nonzero Lebesgue measure. Thus these quantities are unrelated, and we are motivated to define a relaxed version of H_f .

Definition 3.7. Let f be a function in \mathbb{R}^n taking values in \mathbb{R}^n a.e. For an arbitrary set $U \subset \mathbb{R}^n$ containing x and $r > 0$, we let

$$L_{f,U}(x, r) := \sup\{|f(x) - f(z)| : |z - x| \leq r, z \in U\}$$

and

$$l_{f,U}(x, r) := \inf\{|f(x) - f(z)| : |z - x| \geq r, z \in U\},$$

and then

$$H_{f,U}(x, r) := \frac{L_{f,U}(x, r)}{l_{f,U}(x, r)} \quad \text{and} \quad H_f^{\text{fine}}(x) := \inf_U \limsup_{r \rightarrow 0} H_{f,U}(x, r),$$

where the infimum is taken over 1-finely open sets U containing x . We interpret $H_{f,U}(x, r)$ to be ∞ if the denominator is zero or the numerator is ∞ . If $f(x)$ or $f(z)$ is not in \mathbb{R}^n , we interpret $|f(x) - f(z)|$ to be ∞ .

The last sentence of the definition will be needed because the precise representative f^* of a function $f \in L_{\text{loc}}^1(\mathbb{R}^n; \mathbb{R}^n)$ may take values in \mathbb{R}^n only a.e.; we use the same interpretation with the previously defined quantities h_f, H_f .

We record the following ‘‘Borel regularity’’.

Lemma 3.8. *Let μ be a positive Radon measure and let $D \subset \mathbb{R}^n$. Then there exists a Borel set $D^* \supset D$ such that*

$$\mu(D \cap Q) = \mu(D^* \cap Q)$$

for all open cubes $Q \subset \mathbb{R}^n$ (of all sizes and orientations).

Proof. See the proof of [12, Lemma 4.3]. □

We have the following Morse’s covering theorem, see. e.g. [8, Theorem 1.144].

Theorem 3.9. *Let $A \subset \mathbb{R}^n$ be a bounded set, let μ be a positive Radon measure on \mathbb{R}^n , let $\kappa \geq 1$, and let*

$$\mathcal{D} \subset \{x + K : x \in A, K \text{ convex, compact}\}$$

such that for all $x \in A$, for arbitrarily small $r > 0$ there exists $x + K \in \mathcal{D}$ with $B(x, r) \subset x + K \subset B(x, \kappa r)$. Then there exists a disjoint family $\mathcal{D}' \subset \mathcal{D}$ with

$$\mu\left(A \setminus \bigcup_{D \in \mathcal{D}'} D\right) = 0.$$

Lemma 3.10. *Let μ be a positive Radon measure and let $D \subset \mathbb{R}^n$ be a set, not necessarily μ -measurable. Let $\alpha > 0$. Suppose that for μ -a.e. $x \in \mathbb{R}^n$ we find a unit vector $\nu \in \mathbb{R}^n$, a sequence $r_j \searrow 0$, $j \in \mathbb{N}$, and μ -measurable sets $A_j \subset Q_\nu(x, r_j) \setminus D$ such that $\mu(\partial Q_\nu(x, r_j)) = 0$ and*

$$\frac{\mu(A_j)}{\mu(Q_\nu(x, r_j))} \geq \alpha.$$

Then $\mu(D) = 0$.

Note that the cube $Q_\nu(x, r_j)$ is not unique but we can understand the notation to represent a particular choice of such a cube for each j .

Proof. From Lemma 3.8 we find a Borel set $D^* \supset D$ such that

$$\mu(D \cap Q) = \mu(D^* \cap Q)$$

for all cubes $Q \subset \mathbb{R}^n$. For μ -a.e. $x \in \mathbb{R}^n$, for all $j \in \mathbb{N}$ we have

$$\begin{aligned} \frac{\mu(Q_\nu(x, r_j) \cap D^*)}{\mu(Q_\nu(x, r_j))} &= \frac{\mu(Q_\nu(x, r_j) \cap D)}{\mu(Q_\nu(x, r_j))} \\ &\leq \frac{\mu(Q_\nu(x, r_j) \setminus A_j)}{\mu(Q_\nu(x, r_j))} \\ &\leq 1 - \alpha. \end{aligned}$$

From Theorem 3.9 combined with the fact that $\mu(\partial Q_\nu(x, r_j)) = 0$, it follows that $\mu(D^*) = 0$ and then also $\mu(D) = 0$. □

The next two lemmas concern strict and uniform convergence of BV functions.

Lemma 3.11. *Let $\Omega \subset \mathbb{R}$ be an open interval. Let $f \in \text{BV}(\Omega)$ with $|D^j f|(\Omega) = 0$ and suppose $\{f_i\}_{i=1}^\infty$ is a sequence converging to f strictly in $\text{BV}(\Omega)$, and also $f_i^*(x) \rightarrow f^*(x)$ for \mathcal{L}^1 -a.e. $x \in \Omega$. Let $K \subset \Omega$ be a compact interval. Then f_i^* converges to f^* uniformly in K .*

Proof. Fix $\varepsilon > 0$. Since $f_i^*(x) \rightarrow f^*(x)$ for a.e. $x \in \Omega$ and $|Df|$ does not charge singletons, we find a finite collection of points $x_1 < \dots < x_N$ such that $K \subset (x_1, x_N)$, $f_i^*(x_j) \rightarrow f^*(x_j)$ for every $j = 1, \dots, N$, and

$$|Df|([x_j, x_{j+1}]) < \varepsilon \quad \text{for all } j = 1, \dots, N-1.$$

Thus for sufficiently large i , we have $|f_i^*(x_j) - f^*(x_j)| < \varepsilon$ for every $j = 1, \dots, N$. Since $|Df|(W) \leq \liminf_{i \rightarrow \infty} |Df_i|(W)$ for every open set $W \subset \Omega$ and

$$|Df|(\Omega) = \lim_{i \rightarrow \infty} |Df_i|(\Omega),$$

we necessarily have

$$|Df|(H) \geq \limsup_{i \rightarrow \infty} |Df_i|(H)$$

for every compact $H \subset \Omega$. Thus for sufficiently large i , we have also

$$|Df_i|([x_j, x_{j+1}]) < \varepsilon \quad \text{for all } j = 1, \dots, N-1.$$

Let $j \in \{1, \dots, N-1\}$. Using (2.10), for all $x \in [x_j, x_{j+1}]$ we have

$$\begin{aligned} |f_i^*(x) - f^*(x)| &\leq |f_i^*(x) - f_i^*(x_j)| + |f_i^*(x_j) - f^*(x_j)| + |f^*(x_j) - f^*(x)| \\ &\leq |Df_i|([x_j, x_{j+1}]) + |f_i^*(x_j) - f^*(x_j)| + |Df|([x_j, x_{j+1}]) \\ &< \varepsilon + \varepsilon + \varepsilon \end{aligned}$$

for all sufficiently large i . In total, $|f_i^*(x) - f^*(x)| < 3\varepsilon$ for all $x \in K$, for sufficiently large i . \square

Lemma 3.12. *Let $\Omega = (a, b)$ be an open interval. Let $f \in \text{BV}(\Omega)$ be increasing and suppose $\{f_i\}_{i=1}^\infty$ is a sequence converging to f strictly in $\text{BV}(\Omega)$, and $f_i^*(x) \rightarrow f^*(x)$ for a.e. $x \in \Omega$. Then for a.e. $t \in \Omega$, we have*

$$\liminf_{i \rightarrow \infty} \inf_{s \in \Omega: s \geq t} f_i^*(s) \geq f^*(t).$$

Proof. The measures Df_i can be decomposed into their positive and negative parts as $Df_i = (Df_i)_+ - (Df_i)_-$. Due to the strict convergence, we have

$$\lim_{i \rightarrow \infty} (Df_i)_-(\Omega) = 0.$$

For a.e. $t \in \Omega$, we have $\lim_{i \rightarrow \infty} f_i^*(t) = f^*(t)$. Fix such t . Then by (2.10),

$$\begin{aligned} \liminf_{i \rightarrow \infty} \inf_{s \geq t} f_i^*(s) &\geq \liminf_{i \rightarrow \infty} f_i^*(t) - \limsup_{i \rightarrow \infty} (Df_i)_-([t, b]) \\ &= f^*(t). \end{aligned}$$

□

We will also need the following measurability results.

Lemma 3.13. *Let $f \in \text{BV}(\mathbb{R}^n)$. Let $P \subset \mathbb{R}^{n-1}$ be a \mathcal{L}^{n-1} -measurable set and let $h_1 \leq h_2$ be Borel functions on \mathbb{R}^{n-1} . Then*

$$\bigcup_{z \in P} \{z\} \times [h_1(z), h_2(z)]$$

is $\mathcal{L}^{n-1} \otimes |Df_z| = |D_n f|$ -measurable.

Proof. There is a Borel set $P' \supset P$ such that $\mathcal{L}^{n-1}(P' \setminus P) = 0$. The set

$$\bigcup_{z \in P'} \{z\} \times [h_1(z), h_2(z)]$$

can be seen to be a Borel set e.g. as in the proof of [7, Lemma 2.3], and the difference between the sets is contained in

$$\bigcup_{z \in P' \setminus P} \{z\} \times (-\infty, \infty)$$

which is of zero $\mathcal{L}^{n-1} \otimes |Df_z|$ -measure. □

Recall that for a BV function f on the real line, we call $f^\vee(x) - f^\wedge(x) = |Df|(\{x\})$ the jump (size) of f at point x .

Lemma 3.14. *Let $\Omega \subset \mathbb{R}^n$ be open, let $k \in \mathbb{N}$, and let $f \in \text{BV}(\Omega; \mathbb{R}^k)$. For every $z \in \pi_n(\Omega)$ such that $f_z \in \text{BV}(\Omega_z; \mathbb{R}^k)$ (which holds for \mathcal{L}^{n-1} -a.e. $z \in \pi_n(\Omega)$), let $\alpha(z)$ be the size of the maximal jump. Then the function α is \mathcal{L}^{n-1} -measurable.*

Proof. Note that $|Df_z|$ is weakly* \mathcal{L}^{n-1} -measurable; recall Section 2.2. We have

$$\alpha(z) = \lim_{k \rightarrow \infty} \max_{i \in \mathbb{Z}} |Df_z|([i2^{-k}, (i+1)2^{-k}] \cap \Omega_z)$$

for a.e. $z \in \pi_n(\Omega)$. □

4. PROOF OF THE MAIN THEOREM

In this section we prove our main theorem.

Proof of Theorem 1.1. Recall the decomposition $Df = D^a f + D^c f + D^j f$ from before (2.7). We will show that the claim holds for $|D^a f|$ -a.e., $|D^c f|$ -a.e., and $|D^j f|$ -a.e. $x \in \mathbb{R}^n$.

The absolutely continuous part. We prove the claim for $|D^a f|$ -a.e. $x \in \mathbb{R}^n$. Suppose $\frac{dDf}{d|Df|}(x)$ is of full rank. We need to show that $H_{f^*}^{\text{fine}}(x) < \infty$. Excluding a set of \mathcal{L}^n -measure zero, which is also a set of $|D^a f|$ -measure zero, we have $\frac{dDf}{d|Df|}(x) = \frac{\nabla f(x)}{|\nabla f(x)|}$ and the fine

differentiability of Theorem 3.5 holds at x . Consider the quantities

$$\|\nabla f\|_{\max} := \max_{|v|=1} |\nabla f(x)v| \quad \text{and} \quad \|\nabla f\|_{\min} := \min_{|v|=1} |\nabla f(x)v|,$$

which are both in $(0, \infty)$ since $\nabla f(x)$ has full rank. By Theorem 3.5 we find a 1-finely open set V containing x such that

$$\lim_{V \setminus \{x\} \ni y \rightarrow x} \frac{|f^*(y) - f^*(x) - \nabla f(x)(y-x)|}{|y-x|} = 0.$$

Now letting $U := V \cap B(x, r)$ for a sufficiently small $r > 0$, we still have

$$\lim_{U \setminus \{x\} \ni y \rightarrow x} \frac{|f^*(y) - f^*(x) - \nabla f(x)(y-x)|}{|y-x|} = 0, \quad (4.1)$$

and also

$$\frac{|f^*(y) - f^*(x) - \nabla f(x)(y-x)|}{|y-x|} < \frac{\|\nabla f\|_{\min}}{2} \quad \text{for all } y \in U \setminus \{x\}. \quad (4.2)$$

Choose an arbitrary sequence $r_j \searrow 0$. For every $j \in \mathbb{N}$, also consider $y_j \in U \cap \overline{B}(x, r_j)$. Then we have

$$\limsup_{j \rightarrow \infty} \frac{|f^*(y_j) - f^*(x)|}{r_j} \leq \|\nabla f\|_{\max} \quad \text{by (4.1).}$$

Thus

$$\limsup_{j \rightarrow \infty} \frac{L_{f^*, U}(x, r_j)}{r_j} = \limsup_{j \rightarrow \infty} \frac{\sup\{|f^*(z) - f^*(x)| : |z-x| \leq r_j, z \in U\}}{r_j} \leq \|\nabla f\|_{\max}.$$

On the other hand, for every $j \in \mathbb{N}$ consider $\tilde{z}_j \in U$ with $|\tilde{z}_j - x| \geq r_j$ (if it exists, as it does for all sufficiently large j). We have

$$\begin{aligned} \liminf_{j \rightarrow \infty} \frac{|f^*(\tilde{z}_j) - f^*(x)|}{r_j} &= \liminf_{j \rightarrow \infty} \frac{|f^*(\tilde{z}_j) - f^*(x)|}{|\tilde{z}_j - x|} \frac{|\tilde{z}_j - x|}{r_j} \quad \text{by (4.1)} \\ &\geq \liminf_{j \rightarrow \infty} \frac{|f^*(\tilde{z}_j) - f^*(x)|}{|\tilde{z}_j - x|} \\ &\geq \frac{\|\nabla f\|_{\min}}{2} \quad \text{by (4.2)}. \end{aligned}$$

Thus

$$\liminf_{j \rightarrow \infty} \frac{l_{f^*, U}(x, r_j)}{r_j} = \liminf_{j \rightarrow \infty} \frac{\inf\{|f^*(z) - f^*(x)| : |z-x| \geq r_j, z \in U\}}{r_j} \geq \frac{\|\nabla f\|_{\min}}{2}.$$

In total, we obtain

$$\limsup_{j \rightarrow \infty} H_{f^*, U}(x, r_j) = \limsup_{j \rightarrow \infty} \frac{L_{f^*, U}(x, r_j)}{l_{f^*, U}(x, r_j)} \leq 2 \frac{\|\nabla f\|_{\max}}{\|\nabla f\|_{\min}}.$$

We conclude

$$H_{f^*}^{\text{fine}}(x) \leq 2 \frac{\|\nabla f\|_{\max}}{\|\nabla f\|_{\min}} < \infty.$$

Then suppose $\frac{dDf}{d|Df|}(x)$ is not of full rank. We need to show that $H_{f^*}^{\text{fine}}(x) = \infty$. Fix an arbitrary 1-finely open set U containing x . Again excluding a set of \mathcal{L}^n -measure zero, which is also a set of $|D^a f|$ -measure zero, we have $\frac{dDf}{d|Df|}(x) = \frac{\nabla f(x)}{|\nabla f(x)|}$ and the fine differentiability of Theorem 3.5 holds at x . This means that by making the 1-finely open set U smaller, which only decreases $H_{f,U}(x, r)$, we have that

$$\lim_{U \setminus \{x\} \ni y \rightarrow x} \frac{|f^*(y) - f^*(x) - \nabla f(x)(y - x)|}{|y - x|} = 0. \quad (4.3)$$

But now the $\nabla f_j(x)$'s do not span \mathbb{R}^n ; we can assume that they do not span e_n . By Lemma 3.2, we have

$$\frac{\mathcal{H}^{n-1}(\pi_n(\partial B(x, r) \setminus U))}{r^{n-1}} \leq \frac{\text{Cap}_1(\partial B(x, r) \setminus U)}{r^{n-1}} \rightarrow 0 \quad \text{as } r \rightarrow 0,$$

since U is 1-finely open. Let $r_j \searrow 0$. Now we can choose points $z_j \in \partial B(x, r_j) \cap U$ such that the angle between $z_j - x$ and e_n goes to zero. Thus

$$\begin{aligned} \limsup_{j \rightarrow \infty} \frac{l_{f^*, U}(x, r_j)}{r_j} &= \limsup_{j \rightarrow \infty} \frac{\inf\{|f^*(x) - f^*(z)| : |z - x| \geq |z_j - x|, z \in U\}}{r_j} \\ &\leq \limsup_{j \rightarrow \infty} \frac{|f^*(x) - f^*(z_j)|}{r_j} \\ &= \limsup_{j \rightarrow \infty} \frac{|\nabla f(x)(z_j - x)|}{r_j} \quad \text{by (4.3)} \\ &= 0. \end{aligned}$$

Similarly, we can choose points $\tilde{z}_j \in \partial B(x, r_j) \cap U$ such that the angle between $\tilde{z}_j - x$ and $\nabla f_k(x)$ goes to zero, where we choose some nonzero $\nabla f_k(x)$. It follows that

$$\begin{aligned} \liminf_{j \rightarrow \infty} \frac{L_{f^*, U}(x, r_j)}{r_j} &= \liminf_{j \rightarrow \infty} \frac{\sup\{|f^*(x) - f^*(z)| : |z - x| \leq |\tilde{z}_j - x|, z \in U\}}{r_j} \\ &\geq \liminf_{j \rightarrow \infty} \frac{|f^*(x) - f^*(\tilde{z}_j)|}{r_j} \\ &= \liminf_{j \rightarrow \infty} \frac{|\nabla f(x)(\tilde{z}_j - x)|}{r_j} \quad \text{by (4.3)} \\ &\geq |\nabla f_k(x)|. \end{aligned}$$

Thus we get

$$\limsup_{r \rightarrow 0} H_{f^*, U}(x, r) \geq \limsup_{j \rightarrow \infty} \frac{L_{f^*, U}(x, r_j)}{l_{f^*, U}(x, r_j)} = \infty,$$

and so $H_{f^*}^{\text{fine}}(x) = \infty$.

The Cantor part. By Alberti's rank one theorem, see [1], $\frac{dDf}{d|Df|}(x)$ is of rank one for $|D^c f|$ -a.e. $x \in \mathbb{R}^n$, so we need to show for such x that $H_{f^*}^{\text{fine}}(x) = \infty$. For such points, we

have

$$\frac{dDf}{d|Df|}(x) = \lim_{r \rightarrow 0} \frac{Df(B(x, r))}{|Df|(B(x, r))} = \eta(x) \otimes \xi(x)$$

for unit vectors $\eta(x), \xi(x) \in \mathbb{R}^n$. For every such x and $r > 0$, we can choose a cube $Q_{\xi(x)}(x, r)$ such that for different values of r these are dilations of each other. Combining Morse's covering Theorem 3.9 with standard arguments, see e.g. [7, Section 1.6], we get

$$\lim_{r \rightarrow 0} \frac{|D^c f|(Q_{\xi(x)}(x, r))}{|Df|(Q_{\xi(x)}(x, r))} = 1 \quad (4.4)$$

for $|D^c f|$ -a.e. $x \in \mathbb{R}^n$.

We define the auxiliary quantities

$$H_{f^*}^{\text{fine}, k, k'}(x) := \inf_U \sup_{0 < r \leq 1/k} H_{f^*, U}(x, r), \quad k \in \mathbb{N}, k' \in \mathbb{N},$$

where the infimum is taken over 1-finely open sets U containing x for which

$$\frac{\text{Cap}_1(B(x, r) \setminus U)}{r^{n-1}} < \frac{1}{k'} \quad \text{for all } 0 < r \leq 1/k.$$

Note that for all $k' \in \mathbb{N}$, we have

$$H_{f^*}^{\text{fine}} = \inf_{k \in \mathbb{N}} H_{f^*}^{\text{fine}, k, k'} = \lim_{k \rightarrow \infty} H_{f^*}^{\text{fine}, k, k'}.$$

Thus it is enough to show for an arbitrary fixed $M \in \mathbb{N}$ that there exists $k' \in \mathbb{N}$ such that for every $k \in \mathbb{N}$, we have

$$H_{f^*}^{\text{fine}, k, k'}(x) > M \quad \text{for } |D^c f| \text{-a.e. } x \in \mathbb{R}^n. \quad (4.5)$$

Fix $M \in \mathbb{N}$, a natural number

$$k' \geq \frac{(10n)^{2n}}{\omega_{n-1}}, \quad (4.6)$$

and also $k \in \mathbb{N}$, and then fix $x \in \mathbb{R}^n$ where $\frac{dDf}{d|Df|}(x)$ has rank one and (4.4) holds. We can assume that $\xi(x) = e_n$, and then we can assume that $Q_{\xi(x)}(x, r)$ is $Q(x, r)$, that is, the cube with sides in the direction of the coordinate axes.

For $r > 0$, define the scalings

$$f_r(y) = \frac{f(x + ry) - f_{Q(x, r)}}{|Df|(Q(x, r))/r^{n-1}}, \quad y \in Q(0, 1). \quad (4.7)$$

Excluding another $|Df|$ -negligible set, the following blowup behavior is known, see e.g. [2, Theorem 3.95]. For a suitable sequence $r_i \rightarrow 0$, we get $f_{r_i} \rightarrow w$ strictly in $\text{BV}(Q(0, 1); \mathbb{R}^n)$, where

$$w(y) = \eta h(\langle y, e_n \rangle),$$

where h is an increasing, nonconstant function on $(-1/2, 1/2)$. Note that necessarily $|Dh|(((-1/2, 1/2))) \leq 1$ and $|h| \leq 1$. Now we have that

$$f_{r_i} \rightarrow w \quad \text{in } L^1(Q(0, 1)) \quad \text{and} \quad |Df_{r_i}|(Q(0, 1)) \rightarrow |Dw|(Q(0, 1)) = 1,$$

and we also have

$$|D_n f_{r_i}|(Q(0, 1)) \rightarrow |D_n w|(Q(0, 1)) = |Dw|(Q(0, 1)); \quad (4.8)$$

see e.g. [2, Proposition 3.15].

We can assume that $|Df|(\partial Q(x, r_i)) = 0$. Passing to further subsequences if necessary, we can also assume that $f_{r_i} \rightarrow w$ a.e. in $Q(0, 1)$, and that $(f_{r_i})_z \rightarrow w_z$ in $L^1((-1/2, 1/2))$ for \mathcal{L}^{n-1} -a.e. $z \in Q_{n-1}(0, 1)$. Then by lower semicontinuity, we also have

$$|Dw_z|((-1/2, 1/2)) \leq \liminf_{i \rightarrow \infty} |D(f_{r_i})_z|((-1/2, 1/2)). \quad (4.9)$$

On the other hand,

$$\begin{aligned} & \lim_{j \rightarrow \infty} \int_{Q_{n-1}(0, 1)} |D(f_{r_i})_z|((-1/2, 1/2)) d\mathcal{L}^{n-1}(z) \\ &= \lim_{j \rightarrow \infty} |D_n f_{r_i}|(Q(0, 1)) \quad \text{by (2.12)} \\ &= |D_n w|(Q(0, 1)) \quad \text{by (4.8)} \\ &= \int_{Q_{n-1}(0, 1)} |Dw_z|((-1/2, 1/2)) d\mathcal{L}^{n-1}(z) \quad \text{by (2.12)}. \end{aligned} \quad (4.10)$$

Combining (4.9) and (4.10), we necessarily have

$$|Dw_z|((-1/2, 1/2)) = \lim_{i \rightarrow \infty} |D(f_{r_i})_z|((-1/2, 1/2))$$

for \mathcal{L}^{n-1} -a.e. $z \in Q_{n-1}(0, 1)$, and so

$$(f_{r_i})_z \rightarrow w_z \quad \text{strictly in } \text{BV}((-1/2, 1/2)) \quad (4.11)$$

for \mathcal{L}^{n-1} -a.e. $z \in Q_{n-1}(0, 1)$.

For $U \subset \mathbb{R}^n$ and $y \in \mathbb{R}^n$, denote

$$r_i^{-1}(U - y) := \{r_i^{-1}(y' - y) : y' \in U\}.$$

Recalling that $k \in \mathbb{N}$ is fixed, we can assume that $r_i < 1/(nk)$ for all $i \in \mathbb{N}$. For all $y \in \mathbb{R}^n$ we can choose a 1-finely open set $U_y \ni y$ with

$$\frac{\text{Cap}_1(B(y, r) \setminus U_y)}{r^{n-1}} < \frac{1}{k'} \quad \text{for all } 0 < r \leq 1/k$$

and

$$\sup_{0 < r \leq 1/k} H_{f^*, U_y}(y, r) \leq H_{f^*}^{\text{fine}, k, k'}(y) + 1.$$

In particular, for $y \in Q(0, 1)$ we have

$$\frac{\text{Cap}_1(B(x + r_i y, r) \setminus U_{x+r_i y})}{r^{n-1}} < \frac{1}{k'} \quad \text{for all } 0 < r \leq 1/k,$$

and

$$\sup_{0 < r \leq 1/k} H_{f^*, U_{x+r_i y}}(x + r_i y, r) \leq H_{f^*}^{\text{fine}, k, k'}(x + r_i y) + 1, \quad (4.12)$$

so that

$$\frac{\text{Cap}_1(B(y, r) \setminus r_i^{-1}(U_{x+r_i y} - x))}{r^{n-1}} < \frac{1}{k'} \quad \text{for all } 0 < r \leq \sqrt{n}$$

(in fact, for all $0 < r \leq r_i^{-1}/k$), so applying this with the choice $r = \sqrt{n}$, using Lemma 3.2 we get

$$\mathcal{L}^{n-1}(\pi_n(Q(0, 1) \setminus r_i^{-1}(U_{x+r_i y} - x))) \leq \frac{1}{k'} n^n. \quad (4.13)$$

Using also (4.6), we get

$$\mathcal{L}^{n-1}(\pi_n(Q(0, 1) \setminus r_i^{-1}(U_{x+r_i y} - x))) \leq \frac{1}{4} \mathcal{L}^{n-1}(B_{n-1}(0, 1/10)). \quad (4.14)$$

We consider two cases.

Case 1: $|D^j w|(Q(0, 1)) = 0$.

From (2.12) we have $|D^j h|((-1/2, 1/2)) = 0$, and then from (2.10) we have that h^* is continuous, and then so is w^* . For simplicity, let us denote by h, w these representatives.

From (2.14), we know that $((f_{r_i})^*)_z(t) = ((f_{r_i})_z)^*(t)$ for \mathcal{L}^{n-1} -a.e. $z \in Q_{n-1}(0, 1)$ and for every $t \in (-1/2, 1/2)$. Thus we can simply use the notation $(f_{r_i})_z^*(t)$, and similarly $(f_{r_i}^l)_z^\wedge(t), (f_{r_i}^l)_z^\vee(t)$ for the component functions $f_{r_i}^l, l = 1, \dots, n$.

Since $|D^j h|((-1/2, 1/2)) = 0$, $|Dh|$ is not concentrated on one point. Since h is nonconstant, we can assume that $|Dh|((0, 1/2)) > 0$. Then we find $0 < T_1 < T_2 < T_3 < T_4 < 1/2$ such that $|Dh|((T_1, T_2)) > 0$ and $|Dh|((T_2, T_3)) > 0$, and $T_4 - T_1 < 1/10$.

Recall that we denote points $y \in \mathbb{R}^n$ by $y = (y_1, \dots, y_n)$. Also recall (2.9). For some fixed $0 < \varepsilon < 1/10$ and for all $y, z \in Q(0, 1)$, we have

$$|w(y) - w(z)| = |h(y_n) - h(z_n)| \geq \varepsilon \quad \text{whenever } z_n \in (T_1, T_2), y_n \in (T_3, T_4). \quad (4.15)$$

For \mathcal{L}^{n-1} -a.e. $z \in Q_{n-1}(0, 1)$, by (4.11) and Lemma 3.11 we also have that

$$\lim_{i \rightarrow \infty} \sup_{t \in [T_1, T_4]} |(f_{r_i})_z^*(t) - w_z(t)| = 0,$$

and so for sufficiently large i depending on z , we have

$$\sup_{t \in [T_1, T_4]} |(f_{r_i})_z^*(t) - w_z(t)| \leq \frac{\varepsilon}{6(M+1)} < \frac{\varepsilon}{3}. \quad (4.16)$$

Moreover by lower semicontinuity,

$$\liminf_{i \rightarrow \infty} |D(f_{r_i})_z|((T_1, T_2)) \geq |Dw_z|((T_1, T_2)) = |Dh|((T_1, T_2)),$$

so that

$$|D(f_{r_i})_z|((T_1, T_2)) \geq \frac{1}{2} |Dh|((T_1, T_2)) \quad (4.17)$$

for sufficiently large i (depending on z). Denote the set of those $z \in Q_{n-1}(0, 1)$ for which (4.16) and (4.17) hold by D_i . This set is \mathcal{L}^{n-1} -measurable, since $|D(f_{r_i})_z|$ is a parametrized

measure, and $(f_{r_i})^*$ and w are Borel functions with

$$\sup_{t \in [T_1, T_4]} |(f_{r_i})^*_z(t) - w_z(t)| = \sup_{t \in [T_1, T_4] \cap \mathbb{Q}} |(f_{r_i})^*_z(t) - w_z(t)| \quad \text{for } \mathcal{L}^{n-1}\text{-a.e. } z \in Q_{n-1}(0, 1).$$

Consider $i \in \mathbb{N}$ sufficiently large that

$$\mathcal{L}^{n-1}(Q_{n-1}(0, 1) \setminus D_i) \leq \frac{1}{4} \mathcal{L}^{n-1}(B_{n-1}(0, 1/10)). \quad (4.18)$$

For such i , we have

$$\begin{aligned} & |Df_{r_i}|((B_{n-1}(0, 1/10) \cap D_i) \times (T_1, T_2)) \\ & \geq |D_n f_{r_i}|((B_{n-1}(0, 1/10) \cap D_i) \times (T_1, T_2)) \\ & = \int_{B_{n-1}(0, 1/10) \cap D_i} |D(f_{r_i})_z|((T_1, T_2)) d\mathcal{L}^{n-1}(z) \quad \text{by (2.12)} \\ & \geq \frac{\mathcal{L}^{n-1}(B_{n-1}(0, 1/10) \cap D_i)}{2} |Dh|((T_1, T_2)) \quad \text{by (4.17)} \\ & \geq \frac{\mathcal{L}^{n-1}(B_{n-1}(0, 1/10))}{4} |Dh|((T_1, T_2)) \quad \text{by (4.18)}. \end{aligned} \quad (4.19)$$

Let $y \in (B_{n-1}(0, 1/10) \cap D_i) \times (T_1, T_2)$. By (4.18) and (4.14) we find a point

$$q_y \in [(B_{n-1}(0, 1/10) \cap D_i) \times (T_3, T_4)] \cap r_i^{-1}(U_{x+r_i y} - x).$$

Since $T_4 - T_1 < 1/10$, it follows that $|(q_y)_n - y_n| < 1/10$, and then $|q_y - y| < 3/10$. Moreover, we have

$$\begin{aligned} |f_{r_i}^*(q_y) - f_{r_i}^*(y)| & \geq \left[-|f_{r_i}^*(q_y) - w(q_y)| + |w(q_y) - w(y)| - |w(y) - f_{r_i}^*(y)| \right] \\ & > -\varepsilon/3 + \varepsilon - \varepsilon/3 \quad \text{by (4.15), (4.16)} \\ & = \varepsilon/3. \end{aligned} \quad (4.20)$$

On the other hand, there is a point $p \in Q_{n-1}(0, 1)$ such that $B_{n-1}(p, 1/10) \subset Q_{n-1}(0, 1) \setminus Q_{n-1}(0, 4/5)$. Using (4.18) and (4.14), we also find a point

$$q'_y \in ((B_{n-1}(p, 1/10) \cap D_i) \times \{y_n\}) \cap r_i^{-1}(U_{x+r_i y} - x),$$

and then by (4.16),

$$\begin{aligned} |f_{r_i}^*(q'_y) - f_{r_i}^*(y)| & \leq |f_{r_i}^*(q'_y) - w(q'_y)| + |w(q'_y) - w(y)| + |w(y) - f_{r_i}^*(y)| \\ & \leq \frac{\varepsilon}{6(M+1)} + 0 + \frac{\varepsilon}{6(M+1)} \\ & = \frac{\varepsilon}{3(M+1)}. \end{aligned} \quad (4.21)$$

Note that $|q'_y - y| \geq 3/10$. Let

$$A_i := (B_{n-1}(0, 1/10) \cap D_i) \times (T_1, T_2).$$

Recall the definition of the scalings (4.7). For the original function f , for all sufficiently large i we get

$$\begin{aligned} \frac{|Df|(x + r_i A_i)}{|Df|(Q(x, r_i))} &= |Df_{r_i}|(A_i) \\ &\geq \frac{\mathcal{L}^{n-1}(B_{n-1}(0, 1/10))}{4} |Dh|((T_1, T_2)) \quad \text{by (4.19)}. \end{aligned}$$

Then by (4.4), for sufficiently large i we also have

$$\frac{|D^c f|(x + r_i A_i)}{|D^c f|(Q(x, r_i))} \geq \frac{\mathcal{L}^{n-1}(B_{n-1}(0, 1/10))}{5} |Dh|((T_1, T_2)).$$

Recall that $|q_y - y| \leq 3/10$ and $|q'_y - y| \geq 3/10$. Combining (4.20) and (4.21), we get

$$H_{f_{r_i}^*, r_i^{-1}(U_{x+r_i y-x})}(y, 3/10) \geq \frac{|f_{r_i}^*(q_y) - f_{r_i}^*(y)|}{|f_{r_i}^*(q'_y) - f_{r_i}^*(y)|} > \frac{\varepsilon/3}{\varepsilon/(3(M+1))} = M + 1.$$

Now

$$H_{f^*, U_{x+r_i y}}(x + r_i y, 3r_i/10) = H_{f_{r_i}^*, r_i^{-1}(U_{x+r_i y-x})}(y, 3/10) > M + 1.$$

Then by (4.12) we have

$$\begin{aligned} H_{f^*}^{\text{fine}, k, k'}(x + r_i y) + 1 &\geq \sup_{0 < r \leq 1/k} H_{f^*, U_{x+r_i y}}(x + r_i y, r) \\ &\geq H_{f^*, U_{x+r_i y}}(x + r_i y, 3r_i/10) \\ &> M + 1. \end{aligned}$$

Apply Lemma 3.10 with the choices $\mu = |D^c f|$ and D is the set where $H_f^{\text{fine}, k, k'} \leq M$. By the lemma, this set has $|D^c f|$ -measure zero. Thus $H_f^{\text{fine}, k, k'}(x) > M$ for $|D^c f|$ -a.e. $x \in \mathbb{R}^n$, which is what we needed to show (recall (4.5)).

Case 2: $|D^j w|(Q(0, 1)) > 0$.

Using (2.12), we find that also $|D^j h|((-1/2, 1/2)) > 0$. Consider $s \in (-1, 1) \cap J_h$ such that h has its biggest jump at s . In particular, $h^\vee(s) - h^\wedge(s) = |Dh|(\{s\}) > 0$.

We can assume that $s \geq 0$. From (4.4), we have

$$\lim_{i \rightarrow \infty} |D^j f_{r_i}|(Q(0, 1)) = 0. \quad (4.22)$$

Recall the representation $w(y) = \eta h(y_n)$. Pick $l \in \{1, \dots, n\}$ with the largest value of $|\eta_l| > 0$; we can assume that $l = 1$ and that in fact $\eta_1 > 0$. Recall that $M \in \mathbb{N}$ is a (large) fixed number. Choose $0 < \delta < 1/100$ sufficiently small that

$$\delta < \eta_1 \frac{h^\vee(s) - h^\wedge(s)}{4(M+1)}. \quad (4.23)$$

Note that $|D^c f|$ -almost all of the set of points of the type considered in the current Case 2 can be represented as a countable union of points where the jump size $h^\vee(s) - h^\wedge(s)$

is at least, say, $1/l'$ with $l' \in \mathbb{N}$. Thus we can assume that $h^\vee(s) - h^\wedge(s)$ is at least a strictly positive number, and so δ in fact only depends on M . Then choose $0 < \varepsilon < 1/10$ sufficiently small that $s + \varepsilon < 1$ and

$$40^n \omega_{n-1} \omega_n^{-1} \delta^{-n} \varepsilon^{n-1} \leq \frac{1}{4} \mathcal{L}^{n-1}(B_{n-1}(0, 1/10)). \quad (4.24)$$

Apart from (4.6), assume

$$k' \geq \frac{2^{4n^2} \varepsilon^{-n}}{\omega_{n-1}}, \quad (4.25)$$

so that k' only depends on M (apart from the dimension n). Consider the set D_i of those $z \in Q_{n-1}(0, 1)$ for which

$$(f_{r_i})_z \in \text{BV}((-1/2, 1/2)) \text{ has jumps at most size } \delta \min\{1, \eta_1(h^\vee(s) - h^\wedge(s))\}. \quad (4.26)$$

By Lemma 3.14, D_i is \mathcal{L}^{n-1} -measurable. Due to (4.22) and (2.12), we have

$$\lim_{i \rightarrow \infty} \mathcal{L}^{n-1}(D_i) = \mathcal{L}^{n-1}(Q_{n-1}(0, 1)). \quad (4.27)$$

By Theorem 3.1 and Lemma 3.2, we can also assume that $(f_{r_i}^1)^\wedge$ and $(f_{r_i}^1)^\vee$ restricted to $D_i \times (-1/2, 1/2)$ are lower and upper semicontinuous, respectively. Moreover, by (4.11) we can assume that $(f_{r_i})_z \rightarrow w_z$ strictly in $\text{BV}((-1/2, 1/2))$ for all $z \in D_i$, and $(f_{r_i})_z^*(t_0) \rightarrow w_z^*(t_0)$ for some $t_0 \in [s - \varepsilon, s + \varepsilon]$. Thus by (2.10), we get

$$\begin{aligned} \limsup_{i \rightarrow \infty} \sup_{t \in [s - \varepsilon, s + \varepsilon]} |(f_{r_i})_z^*(t) - w_z^*(t_0)| &\leq \limsup_{i \rightarrow \infty} \sup_{t, t' \in [s - \varepsilon, s + \varepsilon]} |(f_{r_i})_z^*(t) - (f_{r_i})_z^*(t')| \\ &\leq \limsup_{i \rightarrow \infty} |D(f_{r_i})_z|([s - \varepsilon, s + \varepsilon]) \\ &\leq |Dw_z|([s - \varepsilon, s + \varepsilon]) \\ &\leq 1. \end{aligned}$$

Note that $w_z(\cdot) = \eta h(\cdot)$, where η is a unit vector and $|h| \leq 1$. Thus for large enough i (depending on z) we have

$$(f_{r_i})_z^*([s - \varepsilon, s + \varepsilon]) \subset Q(0, 5). \quad (4.28)$$

By Lemma 3.12 we also have for large enough i

$$(f_{r_i}^1)_z^\vee(t) \geq \eta_1 h^\vee(s) - \delta \eta_1 (h^\vee(s) - h^\wedge(s)) \quad \text{for all } t \in [s + \varepsilon/2, s + \varepsilon] \quad (4.29)$$

and

$$(f_{r_i}^1)_z^\wedge(t) \leq \eta_1 h^\wedge(s) + \delta \eta_1 (h^\vee(s) - h^\wedge(s)) \quad \text{for all } t \in [s - \varepsilon, s - \varepsilon/2]. \quad (4.30)$$

Denote by $E_i \subset D_i$ the set of points z for which (4.28), (4.29), and (4.30) hold. By Borel measurability, lower semicontinuity, and upper semicontinuity, respectively, the set E_i is \mathcal{L}^{n-1} -measurable. Using (4.26), note that

$$(f_{r_i}^1)_z^\vee(s - \varepsilon/2) \leq \eta_1 h^\wedge(s) + 2\delta \eta_1 (h^\vee(s) - h^\wedge(s)). \quad (4.31)$$

By upper semicontinuity of $t \mapsto (f_{r_i}^1)^\vee_z(t)$ for $z \in E_i$, we find a minimal number $t_{i,z} \in [s - \varepsilon/2, s + \varepsilon/2]$ for which

$$(f_{r_i}^1)^\vee_z(t_{i,z}) \geq \eta_1 \frac{h^\wedge(s) + 2h^\vee(s)}{3}.$$

Then by (2.11) and (4.26), in fact

$$\eta_1 \frac{h^\wedge(s) + 2h^\vee(s)}{3} + \delta\eta_1(h^\vee(s) - h^\wedge(s)) \geq (f_{r_i}^1)^\vee_z(t_{i,z}) \geq \eta_1 \frac{h^\wedge(s) + 2h^\vee(s)}{3},$$

and so by (4.31), necessarily $t_{i,z} > s - \varepsilon/2$. Then we also find a maximal number $t'_{i,z} \in (s - \varepsilon/2, t_{i,z})$ for which

$$(f_{r_i}^1)^\wedge_z(t'_{i,z}) \leq \eta_1 \frac{2h^\wedge(s) + h^\vee(s)}{3}.$$

Then in total

$$(f_{r_i}^1)^\vee_z(t_{i,z}) \geq \eta_1 \frac{h^\wedge(s) + 2h^\vee(s)}{3} \quad \text{and} \quad (f_{r_i}^1)^\wedge_z(t'_{i,z}) \leq \eta_1 \frac{2h^\wedge(s) + h^\vee(s)}{3}, \quad (4.32)$$

and also

$$\begin{aligned} \eta_1 \frac{2h^\wedge(s) + h^\vee(s)}{3} - \delta\eta_1(h^\vee(s) - h^\wedge(s)) &\leq (f_{r_i}^1)^\wedge_z(t) \\ &\leq (f_{r_i}^1)^\vee_z(t) \\ &\leq \eta_1 \frac{h^\wedge(s) + 2h^\vee(s)}{3} + \delta\eta_1(h^\vee(s) - h^\wedge(s)) \end{aligned} \quad (4.33)$$

for all $t'_{i,z} \leq t \leq t_{i,z}$. By (2.10) and (4.32), we have

$$\begin{aligned} |D(f_{r_i})_z|([t'_{i,z}, t_{i,z}]) &\geq |(f_{r_i}^1)^\wedge_z(t'_{i,z}) - (f_{r_i}^1)^\vee_z(t_{i,z})| \\ &\geq \eta_1 \frac{h^\vee(s) - h^\wedge(s)}{3}. \end{aligned} \quad (4.34)$$

Due to (4.27), we have

$$\lim_{i \rightarrow \infty} \mathcal{L}^{n-1}(E_i) = \mathcal{L}^{n-1}(Q_{n-1}(0, 1)).$$

Consider the set $F_i \subset E_i$ consisting of points z for which we have $|f_{r_i}^*(z, u_{i,z}) - f_{r_i}^*(\widehat{z}, \widehat{u})| \geq \delta$ for some $u_{i,z} \in [t'_{i,z}, t_{i,z}]$ and all

$$(\widehat{z}, \widehat{u}) \in Q(0, 1) \cap r_i^{-1}(U_{x+r_i(z, u_{i,z})} - x) \quad \text{with} \quad \widehat{z} \notin B_{n-1}(z, 2\varepsilon).$$

Recursively, for a fixed i and starting from $l = 1$, take points $(z_l, u_l) \in F_i \times [t'_{i,z_l}, t_{i,z_l}]$ with $u_l = u_{i,z_l}$ such that $z_l \in F_i$ is not in any of the sets

$$B_{n-1}(z_{\widehat{l}}, 2\varepsilon) \cup \pi_n[Q_{n-1}(0, 1) \setminus r_i^{-1}(U_{x+r_i(z_{\widehat{l}}, u_{\widehat{l}})} - x)].$$

for $\widehat{l} < l$. Call this collection of points I . By (4.13) and (4.25), we have

$$\#I \geq \frac{\mathcal{L}^{n-1}(F_i)}{\omega_{n-1}(2\varepsilon)^{n-1} + n^n/k'} \geq \frac{\mathcal{L}^{n-1}(F_i)}{2^n \omega_{n-1} \varepsilon^{n-1}}.$$

On the other hand, we have $|f_{r_i}^*(z_l, u_l) - f_{r_i}^*(z_{\widehat{l}}, u_{\widehat{l}})| \geq \delta$ for distinct points (z_l, u_l) and $(z_{\widehat{l}}, u_{\widehat{l}})$ in I , and so by (4.28), we have $\#I \leq 20^n / (\delta^n \omega_n)$. By the choice of ε , recall (4.24), we get

$$\mathcal{L}^{n-1}(F_i) \leq \frac{1}{4} \mathcal{L}^{n-1}(B_{n-1}(0, 1/10)).$$

Denote $S_i := E_i \setminus F_i$.

Thus for large enough i , we have

$$\mathcal{L}^{n-1}(B_{n-1}(0, 1) \setminus S_i) \leq \frac{1}{2} \mathcal{L}^{n-1}(B_{n-1}(0, 1/10)). \quad (4.35)$$

Define

$$A_i := \bigcup_{z \in S_i \cap B_{n-1}(0, 1/10)} \{z\} \times [t'_{i,z}, t_{i,z}].$$

By Lemma 4.39 (proved below) we know that $S_i \ni z \mapsto t_{i,z}$ and $S_i \ni z \mapsto t'_{i,z}$ are Borel functions, and so the set A_i is $|D_n f_{r_i}|$ -measurable by Lemma 3.13.

We also have for some $s - \varepsilon < \kappa < s - \varepsilon/2$ that

$$(f_{r_i})_z^*(\kappa) \rightarrow \eta h(\kappa) \quad \text{for } \mathcal{L}^{n-1}\text{-a.e. } z \in Q_{n-1}(0, 1).$$

Consider the set H_i of those $z \in Q_{n-1}(0, 1)$ for which

$$|(f_{r_i})_z^*(\kappa) - \eta h(\kappa)| < \delta \eta_1 (h^\vee(s) - h^\wedge(s)).$$

Using (4.30), we get

$$(f_{r_i}^1)_z^*(\kappa) < \eta_1 h^\wedge(s) + 2\delta \eta_1 (h^\vee(s) - h^\wedge(s)). \quad (4.36)$$

For sufficiently large i , we have

$$\mathcal{L}^{n-1}(Q_{n-1}(0, 1) \setminus H_i) < \frac{1}{4} \mathcal{L}^{n-1}(B(0, \varepsilon/2)). \quad (4.37)$$

For all sufficiently large i , we have

$$\begin{aligned} |Df_{r_i}|(A_i) &\geq |D_n f_{r_i}^1|(A_i) \\ &= \int_{S_i \cap B_{n-1}(0, 1/10)} |D(f_{r_i}^1)_z|([t'_{i,z}, t_{i,z}]) d\mathcal{L}^{n-1}(z) \quad \text{by (2.12)} \\ &\geq \mathcal{L}^{n-1}(S_i \cap B_{n-1}(0, 1/10)) \eta_1 \frac{h^\vee(s) - h^\wedge(s)}{3} \quad \text{by (4.34)} \\ &\geq \mathcal{L}^{n-1}(B_{n-1}(0, 1/10)) \eta_1 \frac{h^\vee(s) - h^\wedge(s)}{6} \quad \text{by (4.35)}. \end{aligned} \quad (4.38)$$

Recall the definition of the scalings (4.7). For the original function f , for all sufficiently large i we get

$$\begin{aligned} \frac{|Df|(x + r_i A_i)}{|Df|(Q(x, r_i))} &= |Df_{r_i}|(A_i) \\ &\geq \mathcal{L}^{n-1}(B_{n-1}(0, 1/10)) \eta_1 \frac{h^\vee(s) - h^\wedge(s)}{6} \quad \text{by (4.38)}. \end{aligned}$$

Then by (4.4), for sufficiently large i we also have

$$\frac{|D^c f|(x + r_i A_i)}{|D^c f|(Q(x, r_i))} \geq \mathcal{L}^{n-1}(B_{n-1}(0, 1/10)) \eta_1 \frac{h^\vee(s) - h^\wedge(s)}{7}.$$

Let $y \in A_i$. From (4.14), the set of points

$$q'_y \in r_i^{-1}(U_{x+r_i y} - x) \quad \text{with} \quad |\pi_n(q'_y) - \pi_n(y)| \geq 2\varepsilon$$

is nonempty. By the definition of F_i , in fact we can find such a q'_y with

$$|f_{r_i}^*(q'_y) - f_{r_i}^*(y)| < \delta,$$

and note that $|q'_y - y| \geq 2\varepsilon$. Combining (4.13) and (4.25), we have

$$\mathcal{L}^{n-1}(\pi_n(Q(0, 1) \setminus r_i^{-1}(U_{x+r_i y} - x))) \leq \frac{1}{4} \mathcal{L}^{n-1}(B(0, \varepsilon/2)),$$

and then combining with (4.37), we also find a point

$$q_y := (z, \kappa) \in r_i^{-1}(U_{x+r_i y} - x) \cap H_i$$

with $|z - \pi_n(y)| < \varepsilon/2$. Then $|q_y - y| < 2\varepsilon$. Using (4.33) and (4.36), we have

$$|f_{r_i}^*(q_y) - f_{r_i}^*(y)| \geq \eta_1 \frac{h^\vee(s) - h^\wedge(s)}{4}.$$

Thus

$$\begin{aligned} H_{f_{r_i}^*, r_i^{-1}(U_{x+r_i y} - x)}(y, 2\varepsilon) &\geq \frac{|f_{r_i}^*(q_y) - f_{r_i}^*(y)|}{|f_{r_i}^*(q'_y) - f_{r_i}^*(y)|} \\ &> \frac{\eta_1(h^\vee(s) - h^\wedge(s))/4}{\delta} \\ &> M + 1 \quad \text{by (4.23)}. \end{aligned}$$

Now

$$H_{f^*, U_{x+r_i y}}(x + r_i y, 2r_i \varepsilon) \geq H_{f_{r_i}^*, r_i^{-1}(U_{x+r_i y} - x)}(y, 2\varepsilon) > M + 1.$$

It follows that

$$\begin{aligned} H_f^{\text{fine}, k, k'}(x + r_i y) + 1 &\geq \sup_{0 < r \leq 1/k} H_{f^*, U_{x+r_i y}}(x + r_i y, r) \quad \text{by (4.12)} \\ &\geq H_{f^*, U_{x+r_i y}}(x + r_i y, 2\varepsilon r_i) \\ &> M + 1. \end{aligned}$$

Apply Lemma 3.10 with the choices $\mu = |D^c f|$ and D is the set where Df has rank one and $H_f^{\text{fine}, k, k'} \leq M$. By the lemma, this set has $|D^c f|$ -measure zero. Thus $H_f^{\text{fine}, k, k'}(x) > M$ for $|D^c f|$ -a.e. $x \in \mathbb{R}^n$, which is what we needed to show (recall (4.5)).

The jump part. By Alberti's rank one theorem, $\frac{dDf}{d|Df|}(x)$ is of rank one for $|D^j f|$ -a.e. $x \in \mathbb{R}^n$, and so we need to show that $H_{f^*}^{\text{fine}}(x) = \infty$. Recall definitions from Section 2.3.

Consider the sets

$$J_f \cap \{f^- \in B(q_j, r_k)\} \cap \{f^+ \in B(q_l, r_m)\}.$$

for $q_j, q_l \in \mathbb{Q}^n$ and $r_k, r_m \in \mathbb{Q}_+$, $|q_j - q_l| > r_k + r_m$. Each of these sets has finite \mathcal{H}^{n-1} -measure due to (2.8), and is countably \mathcal{H}^{n-1} -rectifiable (see [2, Theorem 3.78]). Thus for each of these sets, from Lemma 3.4 we obtain an exceptional set of \mathcal{H}^{n-1} -measure zero. Since there are countably many sets, corresponding to different choices of q_j, q_l, r_k, r_m , we get countably many exceptional sets of \mathcal{H}^{n-1} -measure zero.

Fix $x \in J_f$ which is outside all of the exceptional sets; this is true of \mathcal{H}^{n-1} -a.e. $x \in J_f$ and thus of $|D^j f|$ -a.e. $x \in \mathbb{R}^n$. Fix $0 < \varepsilon < |f^-(x) - f^+(x)|/2$. Choose q_j, q_l, r_k, r_m such that $r_k + r_m < \varepsilon$ and $r_k + r_m < |q_j - q_l|$, and

$$f^-(x) \in B(q_j, r_k) \quad \text{and} \quad f^+(x) \in B(q_l, r_m).$$

Now x is contained in

$$S := J_f \cap \{f^- \in B(q_j, r_k)\} \cap \{f^+ \in B(q_l, r_m)\}.$$

Let U be an arbitrary 1-finely open set containing x . From Lemma 3.4, we have

$$\lim_{r \rightarrow 0} \frac{\mathcal{H}^{n-1}(B(x, r) \cap S \cap U)}{\omega_{n-1} r^{n-1}} = 1.$$

In particular, we find a sequence of points $y_i \in S \cap U$ with $y_i \rightarrow x$, $y_i \neq x$. By the definition of S and by (2.6), we have

$$\begin{aligned} |f^*(y_i) - f^*(x)| &= \left| \frac{f^-(y_i) + f^+(y_i)}{2} - \frac{f^-(x) + f^+(x)}{2} \right| \\ &\leq \frac{1}{2} |f^-(y_i) - f^-(x)| + \frac{1}{2} |f^+(y_i) - f^+(x)| \\ &\leq r_k + r_m \leq \varepsilon. \end{aligned}$$

By Lemma 2.16 we know that

$$\lim_{r \rightarrow 0} \frac{\mathcal{L}^n(B(x, r) \setminus U)}{\mathcal{L}^n(B(x, r))} = 0.$$

Now from the asymptotic behavior given in (2.2) and (2.3), for any $r > 0$ we find a sequence of points $\hat{y}_i \in B_r^+(x, r) \cap U$ converging to x with $|f^*(\hat{y}_i) - f^+(x)| < \varepsilon$, and then

$$\begin{aligned} |f^*(\hat{y}_i) - f^*(x)| &= |f^*(\hat{y}_i) - (f^-(x) + f^+(x))/2| \\ &\geq \frac{1}{2} |f^+(x) - f^-(x)| - |f^*(\hat{y}_i) - f^+(x)| \\ &\geq \frac{1}{2} |f^+(x) - f^-(x)| - \varepsilon. \end{aligned}$$

Hence for all sufficiently small $r > 0$,

$$H_{f^*, U}(x, r) = \frac{L_{f^*, U}(x, r)}{l_{f^*, U}(x, r)} \geq \frac{\frac{1}{2} |f^+(x) - f^-(x)| - \varepsilon}{\varepsilon}.$$

We get

$$\limsup_{r \rightarrow 0} H_{f^*, U}(x, r) \geq \frac{\frac{1}{2}|f^+(x) - f^-(x)| - \varepsilon}{\varepsilon}.$$

Since $\varepsilon > 0$ was arbitrary and so was the 1-finely open set U , we get $H_{f^*}^{\text{fine}}(x) = \infty$. \square

Lemma 4.39. *The functions $S_i \ni z \mapsto t_{i,z}$ and $S_i \ni z \mapsto t'_{i,z}$ in Case 2 of the proof of the Cantor part of Theorem 1.1 are Borel functions.*

Proof. We have that $(f_{r_i}^1)^\vee$ restricted to $S_i \times (-1/2, 1/2)$ is upper semicontinuous, which implies that $S_i \ni z \mapsto t_{i,z}$ is lower semicontinuous. The subgraph

$$\{(z, t) : z \in S_i, s - \varepsilon/2 < t < t_{i,z}\}$$

is then relatively open in $S_i \times (s - \varepsilon/2, s + \varepsilon/2)$ and can be represented as a countable union of closed cylinders

$$C_k := \overline{B}(x_k, r_k) \times [a_k, b_k], \quad k \in \mathbb{N},$$

intersected with $S_i \times (s - \varepsilon/2, s + \varepsilon/2)$. Given $z \in \overline{B}(x_k, r_k) \cap S_i$, let $t'_{i,z,k}$ be the maximal number in $(s - \varepsilon/2, b_k]$ for which

$$(f_{r_i}^1)^\wedge_z(t'_{i,z,k}) \leq \eta_1 \frac{2h^\wedge(s) + h^\vee(s)}{3}.$$

Since $(f_{r_i}^1)^\wedge$ restricted to S_i is lower semicontinuous, the mapping $\overline{B}(x_k, r_k) \cap S_i \ni z \mapsto t'_{i,z,k}$ is upper semicontinuous. Then

$$S_i \ni z \mapsto t'_{i,z} = \sup_{k \in \mathbb{N}} t'_{i,z,k}$$

is Borel. \square

REFERENCES

- [1] G. Alberti, *Rank one property for derivatives of functions with bounded variation*, Proc. Roy. Soc. Edinburgh Sect. A 123 (1993), no. 2, 239–274. [14](#)
- [2] L. Ambrosio, N. Fusco, and D. Pallara, *Functions of bounded variation and free discontinuity problems*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 2000. [3](#), [4](#), [5](#), [6](#), [7](#), [9](#), [15](#), [16](#), [24](#)
- [3] A. Björn and J. Björn, *Nonlinear potential theory on metric spaces*, EMS Tracts in Mathematics, 17. European Mathematical Society (EMS), Zürich, 2011. xii+403 pp. [7](#)
- [4] M. Carrero, G. Dal Maso, A. Leaci, and E. Pascali, *Relaxation of the nonparametric plateau problem with an obstacle*, J. Math. Pures Appl. (9) 67 (1988), no. 4, 359–396. [8](#)
- [5] L. C. Evans and R. F. Gariepy, *Measure theory and fine properties of functions*. Revised edition. Textbooks in Mathematics. CRC Press, Boca Raton, FL, 2015. xiv+299 pp. [12](#), [15](#)
- [6] J. Heinonen, P. Koskela, N. Shanmugalingam, and J. Tyson, *Sobolev classes of Banach space-valued functions and quasiconformal mappings*, J. Anal. Math. 85 (2001), 87–139. [1](#)
- [7] L. C. Evans and R. F. Gariepy, *Measure theory and fine properties of functions*. Revised edition. Textbooks in Mathematics. CRC Press, Boca Raton, FL, 2015. xiv+299 pp. [12](#), [15](#)

- [8] I. Fonseca and G. Leoni, *Modern Methods in the Calculus of Variations, L^p Spaces*. Springer, Berlin (2007). [10](#)
- [9] P. Lahti, *A notion of fine continuity for BV functions on metric spaces*, Potential Anal. 46 (2017), no. 2, 279–294. [7](#), [9](#)
- [10] P. Lahti, *Finely quasiconformal mappings*, preprint 2024. [7](#), [9](#)
- [11] P. Lahti, *Strong approximation of sets of finite perimeter in metric spaces*, Manuscripta Math. 155 (2018), no. 3-4, 503–522. [8](#)
- [12] P. Lahti, *The Choquet and Kellogg properties for the fine topology when $p = 1$ in metric spaces*, J. Math. Pures Appl. (9) 126 (2019), 195–213. [10](#)
- [13] P. Lahti and N. Shanmugalingam, *Fine properties and a notion of quasicontinuity for BV functions on metric spaces*, J. Math. Pures Appl. (9) 107 (2017), no. 2, 150–182. [8](#)

PANU LAHTI, ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES,
BEIJING 100190, PR CHINA, panulahti@amss.ac.cn