A MODEL FOR SLOWING PARTICLES IN RANDOM MEDIA

FRANÇOIS GOLSE¹ ^(a), VALERIA RICCI^{2,*} ^(b), AND ANA JACINTA SOARES³ ^(b)

ABSTRACT. We present a simple model in dimension $d \geq 2$ for slowing particles in random media, where point particles move in straight lines among and inside spherical identical obstacles with Poisson distributed centres. When crossing an obstacle, a particle is slowed down according to the law $\dot{V} = -\frac{\kappa}{\epsilon}S(|V|)V$, where Vis the velocity of the point particle, κ is a positive constant, ϵ is the radius of the obstacle and S(|V|) is a given slowing profile. With this choice, the slowing rate in the obstacles is such that the variation of speed at each crossing is of order 1. We study the asymptotic limit of the particle system when ϵ vanishes and the mean free path of the point particles stays finite. We prove the convergence of the point particles density measure to the solution of a kinetic-like equation with a collision term which includes a contribution proportional to a δ function in v = 0; this contribution guarantees the conservation of mass for the limit equation.

Key-words: particle systems, kinetic limit, slowing particles, point measure component.

¹École polytechnique & IP Paris, CMLS, 91128 Palaiseau Cedex, France

²UNIVERSITÀ DEGLI STUDI DI PALERMO, DIPARTIMENTO DI MATEMATICA E INFORMATICA, VIA ARCHIRAFI 34, 90123 PALERMO, ITALY

³UNIVERSIDADE DO MINHO, CENTRO DE MATEMÁTICA, CAMPUS DE GUAL-TAR, 4710-057 Braga, Portugal,

^{*}Corresponding Author

E-mail addresses: francois.golse@polytechnique.edu,

valeria.ricci@unipa.it, ajsoares@math.uminho.pt.

Acknowledgments V. Ricci acknowledges the support of the Gruppo Nazionale per la Fisica Matematica-Istituto Nazionale di Alta Matematica (GNFM -

INdAM) and of the University of Palermo, Fondi Finalizzati alla Ricerca di Ateneo FFR2021 Valeria Ricci, FFR2023, FFR2018-2020.

A.J. Soares thanks the support of the Portuguese FCT (Fundação para a Ciência e a Tecnologia) Projects UIDB/00013/2020 and UIDP/00013/2020 of CMAT-UM (Centro de Matemática da Universidade do Minho).

F. Golse and A.J. Soares thank the support of the Bilateral Project Pessoa PT-FCT-Ref.2021.09255.CBM and FR-PHC-Ref.47871WF. .

MSC: 35B40, 35Q49, 35Q70, 82C22, 82C40, 82D10, 82D30.

1. INTRODUCTION

In this article, we present a simple particle model describing the slowing down of point (*light*) particles in a random inhomogeneous medium, where the inhomogeneities (or *inclusions*) consist in spherical obstacles with Poisson distributed centres. In this model, the light particles move freely in the space free from obstacles and slow down when crossing the obstacles. We study the asymptotic behaviour of the particle system when the radius ϵ of the obstacles vanishes and the slowing rate grows to infinity in such a way that the loss of kinetic energy of the particles at each crossing is of order 1. Choosing the density of the Poisson centres in such a way that the mean free path is finite, we derive a kinetic type equation for the probability measure of the light particles in the phase space where the conservation of mass is guaranteed by a term proportional to a Dirac delta measure in v = 0.

One motivation for studying this type of dynamics comes from the modelling of Inertial Confinement Fusion (ICF) targets. While a detailed description of the intricate processes at work in ICF targets [1, 2] exceeds by far the scope of the present study, the following features are of special interest to the problem addressed in the present paper.

Some ICF experiments use pure deuterium (D) targets as a mean of measuring the fuel areal density, known to be a key parameter in ICF research [3, 4]. In these targets, tritium (T) ions are created by the thermonuclear reaction

$$D + D \rightarrow T + \text{ proton } + 4.03 MeV.$$

These T ions are created with a very large kinetic energy (about 1 MeV, see Section 2.1.1 in [1] or Section 1.3.1 in [2]), and therefore are referred to as *suprathermal*. The T ions travel in the plasma and lose kinetic energy, to the point that they combine with the D ions, producing α particles, neutrons and energy according to the thermonuclear reaction

$$D + T \rightarrow \alpha + \text{ neutron } + 17.58 MeV.$$

The cross-section of the D-T reaction is much larger than that of the D-D reaction, and the D-T reaction is the main source of energy in ICF experiments. Thus, the yield of ICF implosions will be reduced by any obstruction to the D-T reaction.

One significant such obstruction comes from the ICF implosion itself. The fuel is highly compressed by ablation of the outer shells of the fusion pellet (see Section 3.8 in [1]). This process initiates a Richtmyer-Meshkov instability (see Section 3.6 in [1], Section 8.1.3 in [2] and [5]) at

the interface between the fuel and the various layers of materials around it. As a result, inclusions of dense, inert material with a high stopping power are driven in the fuel. T ions stopping in these inclusions are prevented from reacting with the surrounding D ions, with the negative effect on the yield of the thermonuclear burning of the fuel in the target already mentioned above.

Returning to the mathematical problem sketched at the beginning of this section, the point particles in the context of ICF experiments would be the T ions, and the "light" material would be the thermonuclear fuel, while the inclusion of "dense" material with high stopping power would be the debris of the various outer shells surrounding the fusion pellet before implosion driven into the fuel by the hydrodynamic instabilities.

The specific problem mentioned above has been addressed in part in earlier mathematical publications. For instance Levermore and Zimmerman have obtained a formula for the fraction of T ions lost to the D-T reaction (formula (22) in [6]). Their argument is based on an earlier model of particle transport in a very special kind of random media proposed by Levermore, Pomraning, Sanzo and Wong [7]. In such media, a particle in medium A (resp. B) at the position s has a probability of finding itself in medium B (resp. A) at the position s + ds equal to ds/λ_A (resp. ds/λ_B), where the lengths λ_A and λ_B are constant. The same problem is considered by Clouet, Puel, Sentis and the first author in [8] with a completely different random media, corresponding to the setting chosen in the present paper — spherical inclusions centred at realizations of a Poisson point process. The analvsis in that reference leads to another formula for the probability of stopping in the inclusions (see formula (13) in [8]). Numerical simulations lead to comparisons between the results obtained in [6] and [8] concerning the fraction of T ions lost to the D-T reaction.

The present paper differs from either [6] or [8] in several ways. The random medium considered here involves a Poisson point process as in [8], but corresponds to a different asymptotic regime. Specifically, the situation considered here involves

- (i) very small, identical spherical inclusions,
- (ii) very high stopping power in each spherical inclusion.

We shall see that the effective equation for the expected velocity distribution function of point particles (T ions), equation (4) in Section 2, is a kinetic model including a collision integral acting on the particles speed only, and not on their direction as in most classical kinetic models, together with a Lagrange multiplier, equation (5) in Section 2,

proportional to the Dirac mass at zero velocity, which maintains the local conservation of mass.

The outline of the paper is the following: in Section 2 we present the particle model, fix the notation and state the main theorem; in Section 3 we recall some a priori estimates needed to derive some useful properties of the limit points of the probability measure of light particles; in Section 4 we derive the limit equation for the probability measure of light particles and prove the main theorem.

2. The particle model and the main result

In this section we shall define the particle model and state the main theorem.

In what follows, we denote by

$$B(c,\epsilon) = \{x \in \mathbb{R}^d : |x - c| < \epsilon\}$$

the open ball centred in c and having radius ϵ (we shall add a superscript denoting the dimension when needed, so that $B^d(c, \epsilon)$ will denote the open ball centred in c and having radius ϵ in dimension d) and by $B^d = |B^d(0, 1)|$ the volume of the d-dimensional unit ball.

Moreover, given a locally compact Hausdorff space X, we denote by $\mathcal{M}(X)$ the space of Radon measures, by $\mathcal{M}^1(X)$ the space of finite mass signed Radon measures and by $\mathcal{M}^1_+(X)$ the space of positive finite mass Radon measures on X.

In this paper we shall use the following notions of convergence in Radon measures spaces.

Denoting as $C_c(X)$ and $C_b(X)$ resp. the set of continuous functions with compact support and the set of continuous bounded functions on X, we say that a sequence of measures on X, $\{\mu_n\}$, converges to μ in the

• weak-* (of $\mathcal{M}(X)$) or vague topology when, for all $\phi \in C_c(X)$, $\int_X \phi(x)\mu_n(dx) \to \int_X \phi(x)\mu(dx)$, and we write:

 $\mu_n \stackrel{*}{\rightharpoonup} \mu \quad \text{as} \quad n \to \infty;$

• weak topology (of $\mathcal{M}^1(X)$) when, for all $\phi \in C_b(X)$, $\int_X \phi(x)\mu_n(dx) \to \int_X \phi(x)\mu(dx)$, and we write:

$$\mu_n \rightharpoonup \mu$$
 as $n \to \infty$.

As a last remark on the notation, if m(dxdy) is a positive Radon measure with finite mass on $\mathbb{R}^p \times \mathbb{R}^q$, we denote by

$$\int_{\mathbb{R}^q} m dy$$

the push-forward of m by the first projection $\mathbb{R}^p \times \mathbb{R}^q \ni (x, y) \mapsto x \in \mathbb{R}^p$, i.e. the Radon measure defined on \mathbb{R}^p by the linear functional

$$\psi \mapsto \int_{\mathbb{R}^p \times \mathbb{R}^q} \psi(x) m(dxdy)$$

We use a similar notation for the push-forward of m by the second projection $\mathbb{R}^p \times \mathbb{R}^q \ni (x, y) \mapsto y \in \mathbb{R}^q$.

We study then the asymptotics of the following particle system.

We consider a particle with initial position $x \in \mathbb{R}^d$ and velocity $v \in \mathbb{R}^d$, $d \ge 2$, and configurations of obstacles (or inclusions) of radius ϵ with Poisson distributed centres $C = (c_1, c_2 \dots)$, where $c_i \in \mathbb{R}^d$.

The probability to find n centres in the set $A \subset \mathbb{R}^d$ is then given by:

$$\mathbf{P}\left(\{C \text{ s.t. } \#(A \cap C) = n\}\right) = \frac{(\lambda_{\epsilon}|A|)^n}{n!} e^{-\lambda_{\epsilon}|A|},$$

with $\lambda_{\epsilon} = \frac{\lambda}{\epsilon^{d-1}}$ and $\lambda = O(1)$ (|A| denotes the Lebesgue measure of the set A). Hence the particle mean free path is finite uniformly as $\epsilon \to 0^+$, since the mean free length to the first collision is proportional to $(\lambda_{\epsilon}\epsilon^{d-1})^{-1}$ and the particle is slowed down inside the obstacles.

We shall denote by \mathbb{E} the expectation under \mathbf{P} and by

$$Z_{\epsilon}[C] := \mathbb{R}^d \setminus \bigcup_{c \in C} \overline{B(c, \epsilon)}$$

the complement in \mathbb{R}^d of the set occupied by the obstacles.

The equation of motion of the particle for a given configuration of obstacles C is given by (see also [6, 8]):

(1)
$$\begin{cases} \dot{X} = V, \\ \dot{V} = -\frac{\kappa}{\epsilon} \mathbf{1}_{Z_{\epsilon}^{c}[C]}(x)S(|V|)V = -\kappa_{\epsilon}(x;C)S(|V|)V, \\ (X,V)\big|_{t=0} = (x,v), \end{cases}$$

where $S \in \text{Lip}((0, +\infty))$, satisfying $S \ge S_0 > 0$, is the slowing profile $(\text{Lip}((0, +\infty)))$ denotes the space of Lipschitz functions on $(0, +\infty)$).

Calling $(X_t^{\epsilon}, V_t^{\epsilon})(x, v; C)$ the solution to the differential system (1), the evolution of an initial profile both on the space of test functions and on the space of Borel measures is defined through the following prescriptions:

$$\phi_{\epsilon}(t, x, v; C) := \phi((X_t^{\epsilon}, V_t^{\epsilon})(x, v; C)),$$

and

(2)
$$\int f_{\epsilon}(t, dxdv; C)\phi(x, v) := \int f_{\epsilon}^{0}(dxdv)\phi_{\epsilon}(t, x, v; C),$$

for each $\phi \in C_b(\mathbb{R}^d \times \mathbb{R}^d)$ and for each positive Borel measure with finite mass $f_{\epsilon}(0, dxdv; C) = f_{\epsilon}^0(dxdv)$.

The time-dependent measure f_{ϵ} is the weak solution to the Cauchy problem for the advection equation

(3)
$$(\partial_t + v \cdot \nabla_x) f_{\epsilon}(t, dxdv; C) = \kappa_{\epsilon}(x; C) \operatorname{div}_v(S(|v|)vf_{\epsilon}(t, dxdv; C))$$

with initial datum $f_{\epsilon}(0, dxdv; C) = f_{\epsilon}^{0}(dxdv)$.

Observe that f_{ϵ} is continuous in t with values in the set of positive measures with finite mass on $\mathbb{R}^d \times \mathbb{R}^d$ equipped with the weak topology (see Lemma 8.1.6 in [9]).

We are interested in studying the behaviour in the limit $\epsilon \to 0$ of $\mathbb{E}[f_{\epsilon}(t, dxdv; C)]$, i.e. in studying the Boltzmann–Grad limit for this specific particle system.

In what follows we shall prove the following theorem:

Theorem 1. Assume $f_0 \in C(\mathbb{R}^d \times \mathbb{R}^d)$, with compact support and $f_0 \geq 0$, and let $f_{\epsilon}(t, x, v; C)$ be the solution to equation (3) with initial datum f_0 . Let $a(z) = \int_0^z du \frac{1}{S(u)}$, with $S \in \text{Lip}((0, +\infty))$, satisfying $S \geq S_0 > 0$.

Then $\mathbb{E}[f_{\epsilon}(t, dxdv; C)]dt \rightarrow F(t, dxdv)dt$ where F(t, dxdv) is the unique solution in $\mathcal{D}'((0, +\infty) \times \mathbb{R}^d \times \mathbb{R}^d)$ to the equation

$$\partial_t F + v \cdot \nabla_x F = \sigma \left[\int_0^1 dh^{d-1} \frac{|z|^d}{|v|^{d-1}} \frac{S(|z|)}{S(|v|)} F(t,x,z) \mathbf{1}_{|v|>0} - |v| F(t,x,v) \right] dx dv + \lambda_F(t,x) \delta_0(dv) dx$$

with $|z| = a^{-1}(a(|v|) + 2\kappa\sqrt{1-h^2}), \ z = |z|\hat{v}, \ \sigma = \lambda B^{d-1}$ and

(5)
$$\lambda_F(t,x) = \int_{\mathbb{R}^d} dz \int_0^1 dh^{d-1} |z| F(t,x,z) \mathbf{1}_{|z| \le a^{-1}(2\kappa\sqrt{1-h^2})}.$$

3. A priori estimates and considerations about limit points

Before analysing in detail the evolution induced on the density measure by (1), we shall recall some a priori estimates which can be directly derived from the definitions (1), (2) and the evolution equation (3) implied by them.

 $\mathbf{6}$

The a priori estimates worked out in this section will help us to obtain some useful properties of the limit points of $\mathbb{E}[f_{\epsilon}(t, x, v; C)]$ when $\epsilon \to 0^+$ in both these topologies.

Lemma 2. Assume that $f_{\epsilon}^{0}(\cdot; C)$ is supported in $\overline{B(0, R)} \times \overline{B(0, R)}$ for some R > 0 and all configuration C of inclusions. Then, for all $t \ge 0$,

(6)
$$\operatorname{supp}(f_{\epsilon}(t,\cdot;C)) \subset \overline{B(0,R+Rt)} \times \overline{B(0,R)}$$

Proof. Let $\phi \in C_b(\mathbb{R}^d \times \mathbb{R}^d)$ satisfy $\phi = 0$ on $\overline{B(0, R + RT)} \times \overline{B(0, R)}$ for some T > 0. Since

$$\begin{aligned} \frac{d}{dt}|V_t^{\epsilon}(x,v;C)| &= -\kappa_{\epsilon}(X_t^{\epsilon}(x,v;C),C)S(|V_t^{\epsilon}(x,v;C)|)|V_t^{\epsilon}(x,v;C)| \le 0\,,\\ \frac{d}{dt}|X_t^{\epsilon}(x,v;C)| \le |V_t^{\epsilon}(x,v;C)| \le |v|\,. \end{aligned}$$

Therefore $\phi((X_t^{\epsilon}, V_t^{\epsilon})(x, v; C)) = 0$ for all $(x, v) \in \overline{B(0, R)} \times \overline{B(0, R)}$ and for all $t \in [0, T]$, so that

$$\begin{split} \int_{\mathbb{R}^d \times \mathbb{R}^d} &\phi(x, v) f_{\epsilon}(t, dx dv; C) \\ &= \int_{\overline{B(0, R)} \times \overline{B(0, R)}} \phi((X_t^{\epsilon}, V_t^{\epsilon})(x, v; C)) f_{\epsilon}^0(dx dv; C) = 0 \,. \end{split}$$

Therefore $\operatorname{supp}(f_{\epsilon}(t, \cdot; C)) \subset \overline{B(0, R + RT)} \times \overline{B(0, R)}$ for all $t \in [0, T)$, which implies the inclusion in the lemma.

Proposition 3. Assume that $f^0_{\epsilon}(\cdot; C)$ is supported in $\overline{B(0, R)} \times \overline{B(0, R)}$ for some R > 0 and all configuration C of inclusions. Then f_{ϵ} satisfies

(a) the continuity equation

$$\partial_t \int_{\mathbb{R}^d} f_{\epsilon}(t,\cdot;C) dv + \nabla_x \cdot \int_{\mathbb{R}^d} f_{\epsilon}(t,\cdot;C) v dv = 0 \quad in \ \mathcal{D}'((0,+\infty) \times \mathbb{R}^d);$$

(b) the global conservation of mass

$$\frac{d}{dt} \int_{\mathbb{R}^d \times \mathbb{R}^d} f_{\epsilon}(t, dx dv; C) = 0 \quad in \ \mathcal{D}'((0, +\infty));$$

(c) the momentum balance

$$\frac{d}{dt} \int_{\mathbb{R}^d \times \mathbb{R}^d} v f_{\epsilon}(t, dx dv; C) = -\int_{\mathbb{R}^d \times \mathbb{R}^d} \kappa_{\epsilon}(x, C) S(|v|) v f_{\epsilon}(t, dx dv; C)$$

in $\mathcal{D}'((0, +\infty));$

(d) the mass flux inequality

$$\frac{d}{dt} \int_{\mathbb{R}^d \times \mathbb{R}^d} \phi(|v|^2) f_{\epsilon}(t, dx dv; C) \le 0 \quad in \ \mathcal{D}'((0, +\infty)) \,,$$

provided that $\phi \in C^1([0, +\infty))$ is nondecreasing.

Proof. Since $f_{\epsilon}(t, \cdot; C)$ is compactly supported by Lemma 2, both sides of the advection equation (3) for f_{ϵ} are compactly supported distributions for $t \in (0, T)$, and can therefore be evaluated on C^{∞} functions, not necessarily with compact support. Choosing the test function of the form $(t, x, v) \mapsto \psi(t, x)$ with ψ of class C^{∞} on $(0, +\infty) \times \mathbb{R}^d$ with support in $(1/n, n) \times \mathbb{R}^d$ for $n \geq 1$ shows that

$$-\int_{(0,+\infty)\times\mathbb{R}^d\times\mathbb{R}^d} (\partial_t + v\cdot\nabla_x)\psi(t,x)f_{\epsilon}(t,dxdv;C)dt = 0.$$

The validity of this equality for all such test function ψ and all $n \geq 1$ is exactly the formulation of (a) in the sense of distributions on $(0, +\infty) \times \mathbb{R}^d$. The same argument with ψ independent of x implies (b).

If we pick a test vector field of the form $(t, x, v) \mapsto \chi(t)v$ where $\chi \in C_c^{\infty}((0, +\infty))$, we find that

$$\int_{(0,+\infty)\times\mathbb{R}^d\times\mathbb{R}^d} \chi'(t) v f_{\epsilon}(t, dxdv; C) dt$$

=
$$\int_{(0,+\infty)\times\mathbb{R}^d\times\mathbb{R}^d} \chi(t) \kappa_{\epsilon}(x, C) S(|v|) v f_{\epsilon}(t, dxdv; C) dt,$$

which is the formulation of (c) in the sense of distributions.

Finally, let us choose a test vector field of the form $(t, x, v) \mapsto \chi(t)\phi(|v|^2)$ where $\chi \in C_c^{\infty}((0, +\infty))$ (since the left-hand side of the advection equation satisfied by f_{ϵ} is a compactly support distribution of order 1 when t is restricted to a bounded interval of $(0, +\infty)$, one can use a test function whose dependence in v is of class C^1 only: see Remark 8.1.1 in [9]). Arguing as above leads to the equality

$$\int_{(0,+\infty)\times\mathbb{R}^d\times\mathbb{R}^d} \chi'(t)\phi(|v|^2)f_{\epsilon}(t,dxdv;C)dt$$

= $2\int_{(0,+\infty)\times\mathbb{R}^d\times\mathbb{R}^d} \chi(t)\kappa_{\epsilon}(x,C)S(|v|)|v|^2\phi'(|v|^2)f_{\epsilon}(t,dxdv;C)dt.$

Hence

$$\frac{d}{dt} \int_{\mathbb{R}^d \times \mathbb{R}^d} \phi(|v|^2) f_{\epsilon}(t, dx dv; C) = -2 \int_{\mathbb{R}^d \times \mathbb{R}^d} \kappa_{\epsilon}(x, C) S(|v|) |v|^2 \phi'(|v|^2) f_{\epsilon}(t, dx dv; C) dt$$

in the sense of distributions on $(0, +\infty)$. Since ϕ is nondecreasing and of class C^1 on $[0, +\infty)$, its derivative ϕ' is nonnegative, which implies (d).

Remark 1. Since, as a consequence of Lemma 2, supp $(\mathbb{E}[f_{\epsilon}(t, \cdot; C)]) \subset \overline{B(0, R + Rt)} \times \overline{B(0, R)}$, whenever $f_{\epsilon}^{0}(\cdot; C) \in \mathcal{M}^{1}_{+}(\mathbb{R}^{d} \times \mathbb{R}^{d})$ each vague limit point of the Radon measure $\mathbb{E}[f_{\epsilon}(t, \cdot; C)]$ on $\mathbb{R}^{d} \times \mathbb{R}^{d}$ is also a weak limit point. The same remark is true for the Radon measure $\mathbb{E}[f_{\epsilon}(t, ;C)]dt$ on $[0, T] \times \mathbb{R}^{d} \times \mathbb{R}^{d}$.

We are now ready to establish some properties of the limit points of $\mathbb{E}[f_{\epsilon}(t, \cdot; C)]$ in the weak-* (and therefore, because of the previous Remark 1, weak) topology.

Proposition 4. Assume that $f^0_{\epsilon}(\cdot; C)$ is supported in $\overline{B(0, R)} \times \overline{B(0, R)}$ for some R > 0 and all configuration C of inclusions, and that

$$\sup_{\epsilon>0} \mathbb{E} \int_{\mathbb{R}^d \times \mathbb{R}^d} f^0_{\epsilon}(dxdv; C) < +\infty \,.$$

Then the family of positive measures $\mathbb{E}[f_{\epsilon}(t, \cdot; C)]$ on $\mathbb{R}^d \times \mathbb{R}^d$ is weakly relatively compact for each $t \geq 0$ and $\mathbb{E}[f_{\epsilon}]$ is weakly relatively compact in the set of positive measures with finite total mass on $[0, T] \times \mathbb{R}^d \times \mathbb{R}^d$ for each T > 0. Any of its limit points F as $\epsilon \to 0^+$ satisfies

(7)
$$\partial_t \int_{\mathbb{R}^d} F(t, \cdot) dv + \nabla_x \cdot \int_{\mathbb{R}^d} F(t, \cdot) v dv = 0$$

in $\mathcal{D}'((0, +\infty) \times \mathbb{R}^d)$, and

(8)
$$\frac{d}{dt} \int_{\mathbb{R}^d \times \mathbb{R}^d} F(t, dx dv) = 0$$

in $\mathcal{D}'((0,+\infty))$.

Proof. By construction, $t \mapsto \mathbb{E}[f_{\epsilon}(t, \cdot; C)]$ is a family of continuous functions on $[0, +\infty)$ in the set of positive measures on $\mathbb{R}^d \times \mathbb{R}^d$ equipped with the weak topology, and Proposition 3 (b) shows that

$$\sup_{\epsilon>0} \mathbb{E} \int_{\mathbb{R}^d \times \mathbb{R}^d} f_{\epsilon}(t, dxdv; C) = \sup_{\epsilon>0} \mathbb{E} \int_{\mathbb{R}^d \times \mathbb{R}^d} f_{\epsilon}^0(dxdv; C) < +\infty.$$

The weak-* relative compactness of $\mathbb{E}[f_{\epsilon}]$ on $[0,T] \times \mathbb{R}^d \times \mathbb{R}^d$ and of $\mathbb{E}[f_{\epsilon}(t,\cdot;C)]$ on $\mathbb{R}^d \times \mathbb{R}^d$ for each $t \geq 0$ follow from statement (13.4.2) in [10]. Since the weak-* closures of both these families are metrizable, any element of these closures is a weak-* limit of a subsequence of $\mathbb{E}[f_{\epsilon}(t,\cdot;C)]$ for fixed t > 0, or of $\mathbb{E}[f_{\epsilon}]$ as $\epsilon \to 0$. Finally, Lemma 2 implies that

$$\operatorname{supp}(\mathbb{E}[f_{\epsilon}(t,\cdot;C)] \subset \overline{B(0,R+Rt)} \times \overline{B(0,R)}$$

for all $\epsilon > 0$ and all $t \in [0, T]$, so that weakly-* converging subsequences of $\mathbb{E}[f_{\epsilon}(t, \cdot; C)]$ for fixed t > 0, or of $\mathbb{E}[f_{\epsilon}]$ as $\epsilon \to 0$ are in fact weakly convergent. Thus, if F is a limit point of $\mathbb{E}[f_{\epsilon}]$, there exists a subsequence $\mathbb{E}[f_{\epsilon_n}] \to F$ weakly in $[0,T] \times \mathbb{R}^d \times \mathbb{R}^d$, and in particular

$$\int_{\mathbb{R}^d \times \mathbb{R}^d} \mathbb{E}[f_{\epsilon_n}](t, dx dv) \to \int_{\mathbb{R}^d \times \mathbb{R}^d} F(t, dx dv)$$

weakly on [0, T] as $\epsilon_n \to 0$, so that

$$\frac{d}{dt} \int_{\mathbb{R}^d \times \mathbb{R}^d} F(t, dx dv) = 0$$

by passing to the limit in Proposition 3 (b) in the sense of distributions on $(0, +\infty)$. Similarly $|v|\mathbb{E}[f_{\epsilon_n}] \to |v|F$ weakly in $[0, T] \times \mathbb{R}^d \times \mathbb{R}^d$, and in particular

$$\int_{\mathbb{R}^d} (1, v) \mathbb{E}[f_{\epsilon_n}](t, \cdot) dv \to \int_{\mathbb{R}^d} (1, v) F(t, \cdot) dv$$

weakly on $[0,T] \times \mathbb{R}^d$ as $\epsilon_n \to 0$. Hence

$$\partial_t \int_{\mathbb{R}^d} F(t, \cdot) dv + \nabla_x \cdot \int_{\mathbb{R}^d} F(t, \cdot) v dv = 0$$

by passing to the limit in the sense of distributions in $(0, +\infty) \times \mathbb{R}^d$. \Box

The last fundamental properties of the limit points of $\mathbb{E}[f_{\epsilon}(t, \cdot; C)]$ concern directly the structure of the limit equation and the uniqueness of its solution.

For the sake of simplicity and clarity we shall choose, from now on, $f_{\epsilon}(0, \cdot; C) = f_0(\cdot) \in \mathcal{M}^1_+(\mathbb{R}^d \times \mathbb{R}^d)$: the results are valid also in the generic case, with suitable assumptions on the sequence $f_{\epsilon}(0, \cdot; C)$.

Lemma 5. Assume that $f_{\epsilon}^{0} = f^{0}$ is a positive measure of finite mass supported in $\overline{B(0,R)} \times \overline{B(0,R)}$ for some R > 0, independent of the configuration C of inclusions, and let F be a weak limit point of $\mathbb{E}[f_{\epsilon}]$ as $\epsilon \to 0$. Assume the existence of $\ell \in C(\mathbb{R}^{d})$ such that $\ell \geq 0$ on \mathbb{R}^{d} , and of a time-dependent positive Radon measure $m_{F}(t, \cdot)$ on $\mathbb{R}^{d} \times \mathbb{R}^{d}$ supported in $\mathbb{R}^{d} \times \overline{B(0,R)}$ for all $t \geq 0$ such that

(9)
$$(\partial_t + v \cdot \nabla_x + \ell(v))F = m_F$$
 in $\mathcal{D}'((0, +\infty) \times \mathbb{R}^d \times (\mathbb{R}^d \setminus \{0\}))$.

Then

$$(\partial_t + v \cdot \nabla_x + \ell(v))F - m_F = \lambda_F(t, dx) \otimes \delta_0(dv) \quad in \ \mathcal{D}'((0, +\infty) \times \mathbb{R}^d \times \mathbb{R}^d) ,$$

where $\lambda_F(t, \cdot)$ is a time-dependent Radon measure on \mathbb{R}^d .

Proof. Consider the distribution

$$S = (\partial_t + v \cdot \nabla_x + \ell(v))F - m_F \in \mathcal{D}'((0, +\infty) \times \mathbb{R}^d \times \mathbb{R}^d).$$

It is assumed in the statement of the lemma that

$$S\big|_{(0,+\infty)\times\mathbb{R}^d\times(\mathbb{R}^d\setminus\{0\})}=0$$

Hence supp $(S) \subset (0, +\infty) \times \mathbb{R}^d \times \{0\}$. By Theorems XXXV-XXXVI in chapter III of [11],

$$S = (\partial_t + v \cdot \nabla_x + \ell(v))F - m_F = \sum_{\text{locally finite}} A_\alpha \otimes \partial_v^\alpha \delta_0(dv)$$

where $A_{\alpha} \in \mathcal{D}'((0, +\infty) \times \mathbb{R}^d)$. Consider next, for each $\phi \in C_c^{\infty}((0, +\infty) \times \mathbb{R}^d)$, the linear functionals

$$S_1: \psi \mapsto -\int_{(0,+\infty)\times\mathbb{R}^d\times\mathbb{R}^d} (\partial_t + v\cdot\nabla_x)\phi(t,x)\psi(v)F(t,dxdv)dt$$

and

$$S_2: \psi \mapsto \int_{(0,+\infty)\times\mathbb{R}^d\times\mathbb{R}^d} \phi(t,x)\psi(v)(\ell(v)F(t,dxdv) - m_F(t,dxdv))dt.$$

These linear functionals are Radon measures on \mathbb{R}^d with finite total variation supported in $\overline{B(0,R)}$. Therefore

$$\langle S, \phi \otimes \psi \rangle = \langle S_1, \psi \rangle + \langle S_2, \psi \rangle,$$

and since S_1 and S_2 are distributions of order 0,

$$|\alpha| > 0 \implies A_{\alpha} = 0,$$

and hence

$$S = (\partial_t + v \cdot \nabla_x + \ell(v))F - m_F = A_0 \otimes \delta_0 \quad \text{in } \mathcal{D}'((0, +\infty) \times \mathbb{R}^d \times \mathbb{R}^d)$$

By Proposition 3 (a), one has $\langle S_1, 1 \rangle = 0$, so that
 $\langle S, \phi \otimes 1 \rangle = \langle S_2, 1 \rangle$
$$= \int_{(0, +\infty) \times \mathbb{R}^d \times \mathbb{R}^d} \phi(t, x)(\ell(v)F(t, dxdv) - m_F(t, dxdv))dt = \langle A_0, \phi \rangle$$

meaning that

$$\lambda_F(t,\cdot) = A_0(t,\cdot) = \int_{\mathbb{R}^d} \ell(v) F(t,\cdot) dv - \int_{\mathbb{R}^d} m_F(t,\cdot) dv \,.$$

In particular λ_F is a time-dependent Radon measure.

Remark 2. Notice that the fact that the distribution λ_F is of order 0 is a consequence of the mass conservation property valid for the limit point F and proved in Proposition 4.

Let $\ell \in C_b(\mathbb{R}^d)$ and let $\mathcal{K} : C_0(\mathbb{R}^d) \to C_0(\mathbb{R}^d)$ be a continuous linear map. The notation $C_0(\mathbb{R}^d)$ designates the Banach space of real-valued continuous functions on \mathbb{R}^d vanishing at infinity, equipped with the norm defining uniform convergence on \mathbb{R}^d . Thus the adjoint of \mathcal{K} is a continuous linear map $\mathcal{K}^* : \mathcal{M}^1(\mathbb{R}^d) \to \mathcal{M}^1(\mathbb{R}^d)$, since the topological dual of $C_0(\mathbb{R}^d)$ is $\mathcal{M}^1(\mathbb{R}^d)$ by Theorem 6.6 in chapter II of [12]. Henceforth we denote by $\tilde{\mathcal{K}}$ the operator defined on $C_0(\mathbb{R}^d \times \mathbb{R}^d)$ by the formula

$$(\mathcal{K}\psi)(x,v) := (\mathcal{K}\psi(x,\cdot))(v) \,.$$

Lemma 6. Let $F(t, \cdot)$ be a time-dependent Radon measure on $\mathbb{R}^d \times \mathbb{R}^d$ with finite total variation, continuous on finite time intervals for the weak topology of $\mathcal{M}^1(\mathbb{R}^d \times \mathbb{R}^d)$ and such that $F(t, \cdot)$ has compact support for all $t \ge 0$. Assume that F is a weak solution to

$$\begin{cases} (\partial_t + v \cdot \nabla_x)F + \ell F = \tilde{\mathcal{K}}^*(\ell F), \quad x, v \in \mathbb{R}^d, \ t > 0, \\ F(0, \cdot) = 0. \end{cases}$$

Then $F(t, \cdot) = 0$ for all $t \ge 0$.

Proof. Let T > 0, and let $\phi \in C_0([0,T] \times \mathbb{R}^d \times \mathbb{R}^d)$ such that $\partial_t \phi$ and $\partial_{x_j} \phi$ exist and belong to $C([0,T] \times \mathbb{R}^d \times \mathbb{R}^d)$ for $j = 1, \ldots, d$. Using ϕ as a test function in the weak formulation of the Cauchy problem verified by F shows that

$$\int_{\mathbb{R}^d \times \mathbb{R}^d} \phi(T, x, v) F(T, dxdv) - \int_0^T \int_{\mathbb{R}^d \times \mathbb{R}^d} ((\partial_t + v \cdot \nabla_x) \phi(t, x, v) F(t, dxdv) dt)$$
$$= \int_0^T \int_{\mathbb{R}^d \times \mathbb{R}^d} \ell(v) (\mathcal{K}(\phi(t, x, \cdot))(v) - \phi(t, x, v)) F(t, dxdv) dt.$$

As mentioned above, see Remark 8.1.1 in [9] for using a class of test functions wider than $C_c^{\infty}((0,T) \times \mathbb{R}^d)$. Besides, since it is known a priori that $F(t, \cdot)$ is weakly continuous in $t \in [0,T]$ with values in the space of compactly supported Radon measures, one uses $\phi(T, \cdot, \cdot)$ as a test function to identify $F(T, \cdot)$, which is the reason for dropping the compact support assumption on ϕ in all its variables).

This identity shows that

$$\int_{\mathbb{R}^d \times \mathbb{R}^d} \phi(T, x, v) F(T, dx dv) = 0$$

for all $\phi \in C_0([0,T] \times \mathbb{R}^d \times \mathbb{R}^d)$ such that $\partial_t \phi$ and $\partial_{x_j} \phi$ exist and belong to $C([0,T] \times \mathbb{R}^d \times \mathbb{R}^d)$ for all $j = 1, \ldots, d$, satisfying

$$-(\partial_t + v \cdot \nabla_x)\phi + \ell(v)\phi = \ell(v)\tilde{\mathcal{K}}\phi, \qquad x, v \in \mathbb{R}^d, \ t \in (0,T).$$

The next lemma shows that $\phi(T, x, v)$ can be chosen to be any function in the space $C_0(\mathbb{R}^d \times \mathbb{R}^d)$ with $\partial_{x_j} \phi \in C_0(\mathbb{R}^d \times \mathbb{R}^d)$ for all $j = 1, \ldots, d$. Hence $F(T, \cdot) = 0$, and since T > 0 is arbitrary, this implies that $F(t, \cdot) = 0$ for all $t \ge 0$.

Lemma 7. Let T > 0, let $\phi^T \in C_0(\mathbb{R}^d \times \mathbb{R}^d)$, such that $\partial_{x_j} \phi^T$ exist and belong to $C_0(\mathbb{R}^d \times \mathbb{R}^d)$ for j = 1, ..., d. There exists $\phi \in C_0([0, T] \times \mathbb{R}^d \times \mathbb{R}^d)$ such that $\partial_{x_j} \phi$ exist and belong to $C_0([0, T] \times \mathbb{R}^d \times \mathbb{R}^d)$ for j = 1, ..., d, and such that $\partial_t \phi \in C([0, T] \times \mathbb{R}^d \times \mathbb{R}^d)$, solution to the (backward) Cauchy problem

$$\begin{cases} -(\partial_t + v \cdot \nabla_x)\phi + \ell(v)\phi = \ell(v)\tilde{\mathcal{K}}\phi, \quad x, v \in \mathbf{R}^d, \ 0 < t < T, \\ \phi(T, \cdot) = \phi^T. \end{cases}$$

Proof. A mild solution to the backward Cauchy problem in Lemma 7 is a function $\phi \in C_0([0,T] \times \mathbb{R}^d \times \mathbb{R}^d)$ satisfying the integral equation

$$\begin{split} \phi(t,x,v) = & e^{-\ell(v)(T-t)} \phi^T(x + (T-t)v,v) \\ & - \int_t^T e^{-\ell(v)(s-t)} \ell(v) \tilde{\mathcal{K}} \phi(s,x + (s-t)v,v) ds \,, \end{split}$$

for all $x, v \in \mathbb{R}^d$ and $t \in [0, T]$.

The family of linear maps defined by the formula

$$S_0(t)\psi(x,v) = e^{-t\ell(v)}\psi(x-tv,v)$$

is a strongly continuous semigroup of bounded operators on $C_0([0, T] \times \mathbb{R}^d \times \mathbb{R}^d)$ with norm $||S_0(t)|| \leq \exp(T \sup_{v \in \mathbb{R}^d} \ell(v))$. By construction $\tilde{\mathcal{K}}$ is a bounded operator on $C_0(\mathbb{R}^d \times \mathbb{R}^d)$ with the same norm as that of the bounded operator \mathcal{K} defined on $C_0(\mathbb{R}^d)$. The existence (and uniqueness) of the mild solution ϕ follows from Proposition 1.2 in chapter 3 of [13].

The regularity in x follows from applying the same argument to each partial derivative $\partial_{x_j} \phi$ for $j = 1, \ldots, d$.

Finally, the regularity in t follows from deriving in t both sides of the integral equation above:

$$\begin{split} \partial_t \phi(t,x,v) &= e^{-\ell(v)(T-t)} \ell(v) \phi^T (x + (T-t)v,v) \\ &\quad -e^{-\ell(v)(T-t)} v \cdot \nabla_x \phi^T (x + (T-t)v,v) + \ell(v) \tilde{\mathcal{K}} \phi(t,x,v) \\ &\quad -\int_t^T e^{-\ell(v)(s-t)} (\ell(v))^2 \tilde{\mathcal{K}} \phi(s,x + (s-t)v,v) ds \\ &\quad + \ell(v) \sum_{j=1}^d \int_t^T e^{\ell(v)(s-t)} v_j \tilde{\mathcal{K}} \partial_{x_j} \phi(s,x + (s-t)v,v) ds \,. \end{split}$$

Notice that the 2nd and 5th terms on the right-hand side of this identity involve products by the components of v, so that $\partial_t \phi$ is only continuous, but may fail to be bounded, let alone converge to 0 as $|v| \to +\infty$. \Box

It remains to check that Lemmas 6 and 7 apply to the equation satisfied by the limit points of $\mathbb{E}[f_{\epsilon}]$ as $\epsilon \to 0$. We shall see that, in that equation, $\ell(v) = \sigma |v|$, where σ is a positive constant, so that ℓ is obviously continuous on \mathbb{R}^d , but not bounded. However, since it is known a priori that $F(t, \cdot)$ has compact support included in $\overline{B(0, R(1+t))} \times \overline{B(0, R)}$ for each $t \ge 0$, one can replace $\ell(v) = \sigma |v|$ with $\ell(v) := \sigma \min(|v|, R+1)$, which is both continuous and bounded on \mathbb{R}^d , without affecting F.

As for the operator \mathcal{K} , it will turn out to be of the form

$$\frac{1}{B^{d-1}} \int_{\nu \in B^{d-1}(0,1)} \psi\left(\mathfrak{V}(|\nu|,v)\right) d\nu + k(|v|)\psi(0)$$

where

$$\mathfrak{V}(\beta, v) := \mathcal{V}(\beta, |v|) \frac{v}{|v|}.$$

In the definition of \mathfrak{V} , the function $\mathcal{V} : [0,1] \times [0,+\infty) \to [0,+\infty)$ is such that

$$\begin{cases} \mathcal{V}(\beta, \cdot) : (c(\beta), +\infty) \to (0, +\infty) \text{ is a } C^1 \text{-diffeomorphism, and} \\ \mathcal{V}(\beta, \cdot)([0, c(\beta)]) = \{0\}, \text{ while } \mathcal{V}(\beta, r) \leq r \text{ for all } r \geq 0, \end{cases}$$

with $c \in C^1((0,1)) \cap C([0,1])$ nonincreasing. Finally

$$k(u) := |\{\nu \in B^{d-1}(0,1) \text{ s.t. } |c(\nu)| \ge u\}|/B^{d-1},.$$

One easily checks that $\mathcal{K} : C_0(\mathbb{R}^d) \to C_0(\mathbb{R}^d)$ is a continuous linear map satisfying

$$\psi \in C_0(\mathbb{R}^d) \text{ and } \psi \ge 0 \text{ on } \mathbb{R}^d \implies \mathcal{K}\psi \ge 0,$$

and

$$|\mathcal{K}\psi(v)| \le \tilde{\mathcal{K}}|\psi|(v) \le \sup_{v \in \mathbb{R}^d} |\psi(v)|$$

Finally, the adjoint of \mathcal{K} is the continuous linear map $\mathcal{K}^* : \mathcal{M}^1(\mathbb{R}^d) \to \mathcal{M}^1(\mathbb{R}^d)$ given by the formula

$$\mathcal{K}^*\mu(dw) = \left(\int_{\mathbb{R}^d} k(|v|)\mu(dv)\right) \delta_0(dw) + \frac{1}{B^{d-1}} \int_{|\nu| \le 1} \mathfrak{V}(|\nu|, \cdot) \# \left(\mu\big|_{\mathbb{R}^d \setminus \{0\}}\right).$$

In the latter formula, the notation $\Phi \# m$ designates the push-forward of the measure m by the (measurable) map Φ , i.e. $\Phi \# m(B) = m(\Phi^{-1}(B))$ (see for instance formula 1.4.3 in chapter IV of [12]), while the notation $m|_A$ designates the restriction of the measure m to the measurable set $A \subset \mathbb{R}^d$.

4. The limit equation

In this section we shall derive the limit equation for $\mathbb{E}[f_{\epsilon}(t, \cdot; C)]$ in $(0, +\infty) \times \mathbb{R}^d \times \mathbb{R}^d \setminus \{0\}$, i.e. an equation of the form (9), with the help of the technique adopted for instance in [14], which is based on the construction of a semiexplicit solution to (3) making use of the backward flow associated to (1). We shall then apply Lemma 5 to recover the limit equation in $(0, +\infty) \times \mathbb{R}^d \times \mathbb{R}^d$.

Again for the sake of simplicity, we shall assume

$$f_{\epsilon}(0, dxdv; C) = f_0(x, v)dxdv.$$

Although in the physical model the light particles should have $x \in Z_{\epsilon}[C]$, this requirement becomes unnecessary when $\epsilon \to 0^+$, since $P(x \in B(c, \epsilon)) < \epsilon$; in any case, all results can be generalised to a generic sequence of initial data $f_{\epsilon}(0, \cdot; C) \in \mathcal{M}^1_+(\mathbb{R}^d \times \mathbb{R}^d)$, with suitable straightforward assumptions on this sequence.

We define, from (2), the evolution of

$$F_{\epsilon}(t, dxdv) = \mathbb{E}[f_{\epsilon}(t, dxdv; C)],$$

and analyse its limit evolution equation in $(0, +\infty) \times \mathbb{R}^d \times \mathbb{R}^d \setminus \{0\}$, in order to apply Lemma 5.

We study then

(10)
$$\int F_{\epsilon}(t, dxdv)\phi(x, v) := \int dxdv f_0(x, v)\mathbb{E}[\phi_{\epsilon}(t, x, v; C)]$$

for $\phi \in C_c(\mathbb{R}^d \times \mathbb{R}^d \setminus \{0\})$ (where $F_{\epsilon}(0, dxdv) = f_0(x, v)dxdv$).

Call $(X_{-t}^{\epsilon}, V_{-t}^{\epsilon})(x, v; C)$ the inverse flow associated to the equations of motion (1). Then,

(11)
$$\int F_{\epsilon}(t, dxdv)\phi(x, v) = \int dxdvF_{\epsilon}(t, x, v)\phi(x, v)$$
$$= \int dxdv \mathbb{E}[f_{0}((X_{-t}^{\epsilon}, V_{-t}^{\epsilon})(x, v; C))J_{\epsilon, C}(t, x, v)\mathbf{1}_{|v|>0}]\phi(x, v)$$

where $J_{\epsilon,C}(t, x, v)$ is the Jacobian of the change of coordinates $(x, v) \rightarrow (X_t^{\epsilon}, V_t^{\epsilon})(x, v; C)$ (we shall see in Subsection 4.2 that the inverse flow is defined whenever |v| > 0).

An equivalent expression for $F_{\epsilon}(t, x, v)$ may be derived from (3), so that we may write

$$F_{\epsilon}(t, x, v)$$

$$(12) = \mathbb{E}[f_{0}((X_{-t}^{\epsilon}, V_{-t}^{\epsilon})(x, v; C))J_{\epsilon,C}(t, x, v)\mathbf{1}_{|v|>0}]$$

$$(13) = \mathbb{E}\left[f_{0}\left((X_{-t}^{\epsilon}, V_{-t}^{\epsilon})(x, v; C)\right)\right)$$

$$\times \mathbf{1}_{|v|>0}e^{d\int_{0}^{t}ds\kappa_{\epsilon}(X_{-s}^{\epsilon}(x, v; C); C)\left[S(|V_{-s}^{\epsilon}(x, v; C)|)+V_{-s}^{\epsilon}(x, v; C)\cdot\nabla_{V}S(|V_{-s}^{\epsilon}(x, v; C)|)\right]}\right].$$

In next subsections we shall define all the elements which will allow us to make explicit (12).

4.1. The backward flow. Since $\mathbf{P}(x \in B_{\epsilon}(c)) < \epsilon$, without loss of generality, we shall assume $x \in Z_{\epsilon}[C]$.

Define

$$\left\{ t \in [0, +\infty) \text{ s.t. } X_t^{\epsilon}(x, v; C) \notin Z_{\epsilon}[C] \right\}$$

= $[\tau_-^1(x, v), \tau_+^1(x, v)] \cup [\tau_-^2(x, v), \tau_+^2(x, v)] \cup \dots$

with

$$0 < \tau_{-}^{1}(x,v) < \tau_{+}^{1}(x,v) < \tau_{-}^{2}(x,v) < \tau_{+}^{2}(x,v) < \dots$$

and the assumption that $\tau^k_+(x,v) = +\infty$ implies that $[\tau^l_-(x,v), \tau^l_+(x,v)] = \emptyset$ for all l > k (here $\tau^j_\pm(x,v) = \tau^j_\pm(x,v;C)$), but we drop the dependence on C in order to simplify the notation).

The times $\tau_{-}^{j}(x,v)$ and $\tau_{+}^{j}(x,v)$ are respectively the entering and the exit time in the *j*-th connected region occupied by one or more obstacles along the forward flow (1).

The number of connected regions occupied by the obstacles crossed by $\{X_s^{\epsilon}(x,v;C)\}_{s\in(0,t)}$ is therefore given by

$$N(t, x, v; C) = \max_{l} \{ l \text{ s.t. } \tau_{-}^{l}(x, v) < t \}.$$

Consider $(x,v) = (X_t^{\epsilon}, V_t^{\epsilon})(y, z; C)$ for $(y, z) \in \mathbb{R}^d \times \mathbb{R}^d$ and denote N = N(t, y, z; C).

We may define

$$\begin{aligned} \tau_{i,+}^{l}(x,v) &= t - \tau_{+}^{N+1-l}(y,z) \\ \tau_{i,-}^{l}(x,v) &= t - \tau_{-}^{N+1-l}(y,z) \\ \tau_{i,-}^{0}(x,v) &= 0 \\ \tau_{i,+}^{N+1}(x,v) &= t \end{aligned}$$

with

. .

$$0 < \tau_{i,+}^1(x,v) < \tau_{i,-}^1(x,v) < \tau_{i,+}^2(x,v) < \tau_{i,-}^2(x,v) < \dots$$

The backward flow is then obtained as the solution $(Y_t^{\epsilon}, Z_t^{\epsilon})(x, -v; C)$ to the equations of motion

(14)
$$\begin{cases} Y = Z \\ \dot{Z} = \frac{\kappa}{\epsilon} \mathbf{1}_{Z_{\epsilon}^{c}[C]}(Y)S(|Z|)Z = \kappa_{\epsilon}(Y;C)S(|Z|)Z \\ (Y,Z)\big|_{t=0} = (x,-v), \end{cases}$$

 $\text{i.e. } X^\epsilon_{-t}(x,v;C) = Y^\epsilon_t(x,-v;C), \text{ while } V^\epsilon_{-t}(x,v;C) = -Z^\epsilon_t(x,-v;C).$

In particular, we may express the solution to (14) in semiexplicit form as

$$Y_{t}^{\epsilon}(x, -v; C) = X_{-t}^{\epsilon}(x, v; C).$$

$$= x + \sum_{j=0}^{N-1} Z(\tau_{i,+}^{j+1})(\tau_{i,+}^{j+1} - \tau_{i,-}^{j}) + Z(\tau_{i,+}^{N+1})(t - \tau_{i,-}^{N})$$

$$(15) \qquad -\sum_{j=1}^{N} |Y_{\tau_{i,-}^{j}}^{\epsilon}(x, -v; C) - Y_{\tau_{i,+}^{j}}^{\epsilon}(x, -v; C)|\hat{v}$$

$$\begin{aligned} Z_t^{\epsilon}(x, -v; C) &= -V_{-t}^{\epsilon}(x, v; C) \\ &= -\left(v + \hat{v}\sum_{j=1}^N \frac{\kappa}{\epsilon} \int_{\tau_{i,+}^j}^{\tau_{i,-}^j} S(|Z|(s)) |Z(s)| ds\right), \end{aligned}$$

where $Z(t) = Z_t^{\epsilon}(x, -v; C)$ and $\hat{v} = \frac{v}{|v|}$.

For ease of notation, from now on, we shall drop the dependence on the configuration of obstacles C in the flows.

4.2. The Jacobian $J_{\epsilon,C}(t, x, v)$. We may now compute the Jacobian $J_{\epsilon,C}(t, x, v)$, observing that, from (12)-(13), we deduce that:

$$J_{\epsilon,C}(t,x,v)$$

(16)
$$= e^{d\int_0^t ds \kappa_{\epsilon}(X_{-s}^{\epsilon}(x,v);C) \left[S(|V_{-s}^{\epsilon}(x,v)|) + V_{-s}^{\epsilon}(x,v) \cdot \nabla_V S(|V_{-s}^{\epsilon}(x,v)|)\right]}.$$

This expression can be made more explicit since, when $Y_s^{\epsilon}(x, -v) \in Z_{\epsilon}^{c}[C]$,

$$\frac{\kappa}{\epsilon}S(|Z|) = \frac{|Z|}{|Z|},$$

so that

$$\exp\left(d\int_0^t \kappa_\epsilon(Y_s^\epsilon(x,-v);C)S(|Z_s^\epsilon(x,-v)|)\right) = \left(\frac{|Z_{\tau_{i,-}^N}^\epsilon|}{|v|}\right)^d$$
$$= \left(\frac{|Z_t^\epsilon(x,-v)|}{|v|}\right)^d$$

and

$$\exp\left(\int_0^t ds \,\kappa_\epsilon(X_{-s}^\epsilon(x,v);C)V_{-s}^\epsilon(x,v)\cdot\nabla_V S(|V_{-s}^\epsilon(x,v)|)\right) = \frac{S(|Z_t^\epsilon(x,-v)|)}{S(|v|)}$$

We have then

(17)
$$J_{\epsilon,C}(t,x,v) = \left(\frac{|Z_t^{\epsilon}(x,-v)|}{|v|}\right)^d \frac{S(|Z_t^{\epsilon}(x,-v)|)}{S(|v|)},$$

while the Jacobian of the change of coordinates $(x, v) \to (Y_t, -Z_t)$ is $\frac{1}{J_{\epsilon,C}}$.

The change of coordinates $(x, v) \to X_t^{\epsilon}(x, v), V_t^{\epsilon}(x, v)$ is therefore well defined when |v| > 0.

4.3. The flow tube. In order to perform explicit computations in the expectations, it is useful to define the notion of *flow tube* associated to a space trajectory.

On specific configurations of obstacles, this notion allows to simplify remarkably the expression of \mathbb{E} , giving also an intuition on the behaviour of the particle system.

We define here the flow tube associated to $(Y_t^{\epsilon}(x, -v), Z_t^{\epsilon}(x, -v))$ as

(18)
$$\mathcal{T}_{\epsilon}(-t;x,v)(C) := \{ y \in \mathbb{R}^d : |y - Y_s^{\epsilon}(x,-v)| < \epsilon \ s \in [0,t] \}.$$

A given trajectory $Y_t^{\epsilon}(x, -v)$ will meet an obstacle c only if $c \in \mathcal{T}_{\epsilon}(-t; x, v)(C)$.

Whenever $|V_{-t}^{\epsilon}(x,v)| = |Z_t^{\epsilon}(x,-v)| < V_{\max}$, we have the following inclusion

(19)
$$\mathcal{T}_{\epsilon}(-t;x,v)(C) \subset \mathcal{T}_{\epsilon}(-t;x,\hat{v})(V_{\max})$$

with

$$\mathcal{T}_{\epsilon}(-t; x, \hat{v})(V_{\max}) := \left\{ y \in \mathbb{R}^d : |y - (x - V_{\max}\hat{v}s)| < \epsilon \ s \in [0, t] \right\}$$
$$\subset B(x, V_{\max}t).$$

In the same way we may define the flow tube associated to the direct flow to $(X_t^{\epsilon}(x,v), V_t^{\epsilon}(x,v))$ as

(20)
$$\mathcal{T}_{\epsilon}(t;x,v)(C) := \{ y \in \mathbb{R}^d : |y - X_s^{\epsilon}(x,v)| < \epsilon \ s \in [0,t] \}.$$

We have then:

(21)
$$\mathcal{T}_{\epsilon}(t; Y_t^{\epsilon}(x, -v), -Z_t^{\epsilon}(x, -v))(C) = \mathcal{T}_{\epsilon}(t; Y, -Z)(C) = \mathcal{T}_{\epsilon}(t; Y, -Z)(C) = \mathcal{T}_{\epsilon}(t; Y, -Z)(C)$$

 $\{y\in \mathbb{R}^d: |y-X^\epsilon_s(Y,-Z)|<\epsilon \ s\in [0,t]\},$

and obviously, whenever $|V^{\epsilon}_{-t}(x,v)| = |Z^{\epsilon}_{t}(x,-v)| < V_{\max}$,

$$\mathcal{T}_{\epsilon}(t;Y,-Z)(C) = \mathcal{T}_{\epsilon}(-t;x,v)(C) \subset B(x,V_{\max}t)$$

and

(22)
$$\mathcal{T}_{\epsilon}(t;Y,-Z)(C) \subset \mathcal{T}_{\epsilon}(t;Y,\hat{v})(V_{\max}) = \mathcal{T}_{\epsilon}(-t;Y,-\hat{v})(V_{\max})$$
$$\subset B(x,V_{\max}t).$$

These inclusions will be useful to simplify computations.

4.4. The Markovian set $A_1^{\epsilon}(x, v)$ and the limit dynamics. For each $(x, v) \in \mathbb{R}^d \times \mathbb{R}^d$ and $t \in (0, +\infty)$, call

(23)
$$A_1^{\epsilon}(x,v) = \{C | \forall c, c' \in Cs.t.(x - (0, +\infty)v) \cap B(c,\epsilon) \neq \emptyset \text{ and}$$

 $(x - (0, +\infty)v) \cap B(c',\epsilon) \neq \emptyset \implies B(c,\epsilon) \cap B(c',\epsilon) = \emptyset\}.$

Such configurations of obstacles are the ones where the obstacles crossed by any trajectory with initial datum (x, v), defined by (14), do not superimpose.

The set $A_1^{\epsilon}(x, v)$ enjoys some relevant properties for the particle system with dynamics (1).

4.4.1. Irrelevance of $(A_1^{\epsilon}(x,v))^c$ when $\epsilon \to 0^+$. The first relevant property of $A_1^{\epsilon}(x,v)$ is given by the following lemma.

Lemma 8. Assume that $f_0(x,v) \in L^1_+(\mathbb{R}^d \times \mathbb{R}^d)$ with $\operatorname{supp}(f_0) \subset \overline{B(0,R)} \times \overline{B(0,R)}$. Then for each $\phi \in C_b(\mathbb{R}^d \times \mathbb{R}^d), \forall t > 0$

(24)
$$\lim_{\epsilon \to 0^+} \int_{\mathbb{R}^d \times \mathbb{R}^d} dx dv \phi(x, v) \mathbb{E}[f_{\epsilon}(t, x, v) \mathbf{1}_{|v| > 0} \mathbf{1}_{(A_1^{\epsilon})^c}] = 0$$

Proof. We have a.e. in $\mathbb{R}^d \times \mathbb{R}^d$

$$\mathbb{E}[f_{\epsilon}(t, x, v)\mathbf{1}_{|v|>0}\mathbf{1}_{(A_{1}^{\epsilon})^{c}}]$$

$$\leq \sum_{M\geq 0} \frac{\lambda_{\epsilon}^{M}}{M!} \int dc_{1} \dots dc_{M} e^{-\lambda_{\epsilon}|B(x,Rt)|}$$

$$\times \prod_{j=1, j\neq 2}^{M} \mathbf{1}_{B(x,Rt)}(c_{j})\mathbf{1}_{B(c_{1},2\epsilon)}(c_{2})\mathbf{1}_{|v|>0}$$

$$\times f_{0}((Y_{t}^{\epsilon}(x, -v), -Z_{t}^{\epsilon}(x, -v)))J_{\epsilon,C}(t, x, v)$$

Therefore, because of definition (18) and inclusion (22),

$$\int dv dx \,\phi(x,v) \mathbb{E}[f_{\epsilon}(t,x,v)\mathbf{1}_{|v|>0}\mathbf{1}_{(A_{1}^{\epsilon})^{c}}]$$

$$\leq \sum_{M\geq 0} \frac{\lambda_{\epsilon}^{M}}{M!} \int dY dZ \int dc_{1} \dots dc_{M} e^{-\lambda_{\epsilon}|\mathcal{T}_{\epsilon}(t;Y,\hat{v})(R)|}$$

$$\times \prod_{j=1, j\neq 2}^{M} \mathbf{1}_{\mathcal{T}_{\epsilon}(t;Y,\hat{v})(R)}(c_{j})\mathbf{1}_{B(c_{1},2\epsilon)}(c_{2})\mathbf{1}_{|v|>0}$$

$$\times f_{0}(Y,Z)\phi\big(X_{t}^{\epsilon}(Y,Z),V_{t}^{\epsilon}(Y,Z)\big),$$

where we have performed the change of variable $(x, v) \rightarrow (Y, Z) = (Y_t^{\epsilon}(x, -v), -Z_t^{\epsilon}(x, -v))$, with Jacobian $J_{\epsilon,C}^{-1}$.

We have then

r

$$\int dv dx \,\mathbb{E}[f_{\epsilon}(t, x, v) \mathbf{1}_{|v|>0} \mathbf{1}_{(A_{1}^{\epsilon})^{c}}]$$

$$\leq \|\phi\|_{\infty} \lambda_{\epsilon} |B(0, 2\epsilon)| \int dY dZ f_{0}(Y, Z)$$

$$\leq \lambda \epsilon 2^{d} B^{d} \|\phi\|_{\infty} \int dY dZ f_{0}(Y, Z) \xrightarrow{\epsilon \to 0} 0$$

and the lemma is proved.

We have then that $\mathbb{E}[f_{\epsilon}(t, x, v)\mathbf{1}_{|v|>0}\mathbf{1}_{(A_{1}^{\epsilon})^{c}}] \rightarrow 0$, and therefore $A_{1}^{\epsilon}(x, v)$ is the only set of configurations of obstacles which play a role in the computation of the weak limit of $\mathbb{E}[f_{\epsilon}(t, x, v)\mathbf{1}_{|v|>0}]$ when $\epsilon \rightarrow 0^{+}$.

4.4.2. The dynamics on $A_1^{\epsilon}(x, v)$. A second property of $A_1^{\epsilon}(x, v)$ is that, when $C \in A_1^{\epsilon}(x, v)$, the motion described by (14) is quite simplified.

Within this set, the trajectory starting at (x, -v) will cross $N = N(t, Y_t^{\epsilon}(x, -v), -Z_t^{\epsilon}(x, -v); C)$ single obstacles, which can be ordered according to their collision times; therefore $\tau_{i,+}^j(x, v)$ and $\tau_{i,-}^j(x, v)$ are resp. the entry and the exit time of the trajectory in and from the *j*-th obstacle.

For $C \in A_1^{\epsilon}(x, v)$ such that $C \cap \mathcal{T}_{\epsilon}(-t; x, v)(C) = (c_1, \ldots, c_N)$, where c_1, \ldots, c_N are ordered according to their collision times, we call:

$$[C]_{N} = (c_{1}, \dots, c_{N})$$

$$(25)$$

$$= \{ C' \in A_{1}^{\epsilon}(x, v) : \exists i_{1}, \dots, i_{N} \in \mathbb{N} \ s.t. \ (c'_{i_{1}}, \dots, c'_{i_{N}}) = (c_{1}, \dots, c_{N}) \}.$$

We call

$$[A_1^{\epsilon}(t, x, v)]_N = \{ [C]_N \text{ defined by}(25) \}$$

and define the set of ordered obstacles possibly met by the trajectories (14) up to time t by

$$[A_1^{\epsilon}(t,x,v)] = \bigcup_{N=0}^{\infty} [A_1^{\epsilon}(t,x,v)]_N$$

Once again, since $\mathbf{P}(x \in B_{\epsilon}(c)) < \epsilon$, without loss of generality, we shall assume $x \in Z_{\epsilon}[C]$.

For a given configuration of ordered obstacles $(c_1, \ldots, c_N) \in [A_1^{\epsilon}(t, x, v)]_N$ we define $n_j(x, v) = (n_{j,x_1}, n_{j,x_2}, \ldots, n_{j,x_d})$ as the unit vector such that

$$c_j + \epsilon n_j = Y^{\epsilon}_{\tau^j_{i,+}}(x, -v)$$

Then, choosing \hat{v} as the unit vector of the first coordinate axis, we define the *impact parameter* of the trajectory with the *j*-th obstacle as

$$h_j = \sqrt{\sum_{i=2}^d n_{j,x_i}^2}.$$

The solution to (14) on the set of configurations of obstacles

$$A_1^{\epsilon}(t, x, v)]_N \subset [A_1^{\epsilon}(t, x, v)]$$

can be expressed as:

$$Y_{t}^{\epsilon}(x,-v) = x + \sum_{j=0}^{N-1} Z_{\tau_{i,+}^{j+1}}^{\epsilon}(x,-v) (\tau_{i,+}^{j+1} - \tau_{i,-}^{j}) + Z_{\tau_{i,+}^{k+1}}^{\epsilon}(x,-v) (t - \tau_{i,-}^{N})$$

$$(26) \qquad - 2\epsilon \sum_{j=1}^{N-1} \sqrt{1 - h_{j}^{2}} \hat{v} - |Y_{\tau_{i,-}^{N}}^{\epsilon}(x,-v) - Y_{\tau_{i,+}^{N}}^{\epsilon}(x,-v)| \hat{v}$$

$$Z_t^{\epsilon}(x, -v) = -\left(v + \hat{v}\sum_{j=1}^N \frac{\kappa}{\epsilon} \int_{\tau_{i,+}^j}^{\tau_{i,-}^j} S(|Z|(s))|Z(s)|ds\right)$$

with $|V^{\epsilon}_{i,+}(x-v)| = V^{\epsilon}_{i,+}(x-v)|<2\epsilon$

with $|Y_{\tau_{i,-}^N}^{\epsilon}(x,-v) - Y_{\tau_{i,+}^N}^{\epsilon}(x,-v)| < 2\epsilon$. Within $[A_1^{\epsilon}(t,x,v)]_N$, calling $n_j = (\sqrt{1-\nu_j^2},\nu_j)$ we may perform the change of variable

(27)
$$(c_1...,c_N) \to (\nu_1,\tau_{i,+}^1,\ldots,\nu_N,\tau_{i,+}^N),$$

where

$$c_j = x + \sum_{k=0}^{j-1} Z_{\tau_{i,+}^{j+1}}^{\epsilon}(\tau_{i,+}^{j+1} - \tau_{i,-}^k) - 2\epsilon \sum_{k=1}^{j-1} \sqrt{1 - h_k^2} \hat{v} - \epsilon n_j.$$

The Jacobian matrix of (27) is a block triangular matrix, with Jacobian

(28)
$$J_{ct}(\nu_1, \tau_{i,+}^1, \dots, \nu_N, \tau_{i,+}^N) = \prod_{j=1}^N \epsilon^{d-1} |Z_{\tau_{i,+}^j}^{\epsilon_j}|$$

and each set $[A_1^{\epsilon}(t, x, v)]_N$ is expressed, in the new coordinates, as

$$[A_{1}^{\epsilon}(t, x, v)]_{N} = \Big\{ (\nu_{1}, \tau_{i,+}^{1}, \dots, \nu_{N}, \tau_{i,+}^{N}) \in (\mathbb{R}^{d-1} \times (0, t])^{N} :$$

$$(29) \qquad 0 < \tau_{i,+}^{1} < \dots < \tau_{i,+}^{N} \le t, \quad |\nu_{j}| \le 1, \quad j = 1, \dots, N \Big\}.$$

The size of the flow tube associated to the trajectory (26) is

$$\begin{aligned} |\mathcal{T}_{\epsilon}(-t;x,v)(c_{1}\dots c_{N}) &= B^{d-1}\epsilon^{d-1} \bigg[\sum_{j=0}^{N} |Z|^{\epsilon}_{\tau^{j+1}_{i,+}}(\tau^{j+1}_{i,+} - \tau^{j}_{i,-}) \\ (30) &+ 2\epsilon \sum_{j=1}^{N-1} \sqrt{1 - h_{j}^{2}} + \left| Y^{\epsilon}_{\tau^{N}_{i,-}}(x,-v) - Y^{\epsilon}_{\tau^{N}_{i,+}}(x,-v) \right| \bigg] \end{aligned}$$

Finally, thanks to (14), we may make explicit, at each complete crossing, the change in velocity, Indeed,

(31)
$$\int_{|Z|_{\tau_{i,+}^{j}}}^{|Z|_{\tau_{i,+}^{j}}} \frac{du}{S(u)} = \frac{\kappa}{\epsilon} |Y_{\tau_{i,-}^{j}}^{\epsilon}(x,-v) - Y_{\tau_{i,+}^{j}}(x,-v)| = 2\kappa \sqrt{1-h_{j}^{2}}.$$

Defining, for $z \in (0, +\infty)$, the invertible (because S(u) > 0) function

$$a(z) = \int_0^z du \frac{1}{S(u)},$$

we may write (31) as

$$a(|Z|(\tau_{i,-}^{j})) - a(|Z|(\tau_{i,+}^{j})) = 2\kappa\sqrt{1 - h_{j}^{2}}$$

and express the exit velocity from the obstacle j as

(32)
$$|Z|_{\tau_{i,-}^{j}}^{\epsilon} = a^{-1} [a(|Z|_{\tau_{i,+}^{j}}^{\epsilon}) + 2\kappa \sqrt{1 - h_{j}^{2}}].$$

where, in particular,

(33)
$$|Z|_{\tau_{i,-}^1}^{\epsilon} = a^{-1} \left[a(|v|) + 2\kappa \sqrt{1 - h_j^2} \right]$$

From (32-33) we see that $|Z|_{\tau_{i,-}^j}^{\epsilon}$ is independent of ϵ .

Remark 3. The independence of the exit velocity of ϵ follows from the choice $\kappa_{\epsilon} = \frac{\kappa}{\epsilon}$, i.e. the increment in the velocity is inversely proportional to the size of the obstacles.

4.4.3. The limit dynamics on $[A_1^{\epsilon}(t, x, v)]$. From the formulas given in the previous subsection, it is easy to compute the limit dynamics on the relevant set of obstacles.

By (32-33) and $S(u) > S_0$ we may estimate $\tau_{i,-}^j - \tau_{i,+}^j$ as follows

(34)
$$\tau_{i,-}^{j} - \tau_{i,+}^{j} = \frac{\epsilon}{\kappa} \int_{|Z|(\tau_{i,+}^{j})}^{|Z|(\tau_{i,+}^{j})} \frac{du}{uS(u)} \le \frac{\epsilon}{\kappa S_{0}} \ln \left[\frac{|Z|_{\tau_{i,-}^{j}}^{\epsilon}}{|Z|_{\tau_{i,+}^{j}}^{\epsilon}} \right]$$

so that

(35)
$$\lim_{\epsilon \to 0^+} (\tau_{i,-}^j - \tau_{i,+}^j) = 0.$$

When $\epsilon \to 0^+$, we obtain therefore the limit dynamics

(36)
$$Y_t^{\epsilon}(t, -v) \rightarrow Y_t(x, -v) = x + \sum_{j=0}^N Z_{t_{i,+}^{j+1}}^{-}(t_{i,+}^{j+1} - t_{i,+}^j)$$

(37)
$$Z_t^{\epsilon}(x, -v) \rightarrow Z_t(x, -v) = -v + \sum_{j=1}^N \left[Z_{t_{i,+}^j}^+ - Z_{t_{i,+}^j}^- \right],$$

where

$$t_{i,+}^j = \lim_{\epsilon \to 0^+} \tau_{i,+}^j$$

and (for a.a. configurations of obstacles)

$$Z_{t_{i,+}^{j}}^{+} = -a^{-1} \left[a(|Z_{t_{i,+}^{j}}^{-}|) + 2k\sqrt{1 - h_{j}^{2}} \right] \hat{v}$$

 $(\text{we note that } Z^{\pm}_{t^j_{i,+}} = \underset{t \to (t^j_{i,+})^{\pm}}{\lim} Z^{\epsilon}_t, \text{ recalling that } \underset{\epsilon \to 0^+}{\lim} Z^{\epsilon}_{\tau^j_{i,\pm}}(x,-v)) =$ $\lim_{t \to (t_{i,+}^j)^{\mp}} Z_t^{\epsilon} \big).$

More precisely, we have

$$|Y_t^{\epsilon}(x, -v) - Y_t(x, -v)| \le \epsilon \left[\frac{|Z|_{\max}}{\kappa S_0} \ln\left(\frac{|Z|_{\max}}{|v|}\right) + 2N\right]$$

and

$$|Z_t^{\epsilon}(x, -v) - Z_t(x, -v)| = \left| \sum_{j=1}^N \left[Z_{\tau_{i,-}^j}^{\varepsilon} - Z_{\tau_{i,+}^j}^{\epsilon} - \left(Z_{t_{i,+}^j}^+ - Z_{t_{i,+}^j}^- \right) \right] \right| = 0$$

Thanks to Lemma 8 and the considerations given before, we may now study the behaviour of the limit points of $\mathbb{E}[f_{\epsilon}(t, x, v)\mathbf{1}_{A_{1}^{\epsilon}(x, v)}]$ when $\epsilon \to 0^+$ on $\mathbb{R}^d \times \mathbb{R}^d \setminus \{0\}$ (or equivalently the behaviour of $\mathbb{E}[f_{\epsilon}(t, x, v)\mathbf{1}_{|v|>0}\mathbf{1}_{A_{1}^{\epsilon}(x, v)}]).$

4.5. Limit of $\mathbb{E}[f_{\epsilon}(t, x, v)\mathbf{1}_{|v|>0}\mathbf{1}_{A_{1}^{\epsilon}}]$. We are now ready to prove Theorem 1, through the following auxiliary theorem.

Theorem 9. Assume $f_0 \in C(\mathbb{R}^d \times \mathbb{R}^d)$, with compact support and $f_0 \geq 0$, and let $f_{\epsilon}(t, x, v; C)$ be the solution to (3) with initial datum f_0 Then, for $t \in (0, +\infty)$ on $\mathbb{R}^d \times \mathbb{R}^d \setminus \{0\}$,

$$\mu_{\epsilon} = \mathbb{E}[f_{\epsilon}(t, x, v)] dx dv \stackrel{*}{\rightharpoonup} \mu = F(t, x, v) dx dv$$

where F solves the equation (38) $\partial_t F + v \cdot \nabla_x F = \sigma \int_0^1 dh^{d-1} \frac{|z|^d}{|v|^{d-1}} \frac{S(|z|)}{S(|v|)} F(t,x,z) \mathbf{1}_{|v|>0} - \sigma |v| F(t,x,v)$

in $\mathcal{D}'((0, +\infty) \times \mathbb{R}^d \times \mathbb{R}^d \setminus \{0\})$, with $z = a^{-1} \left(a(|v|) + 2\kappa\sqrt{1-h^2}\right) \hat{v}$ and $\sigma = \lambda B^{d-1}$

Proof. Because of Lemma 8,

$$\mathbb{E}[f_{\epsilon}(t,x,v)\mathbf{1}_{|v|>0}] = \mathbb{E}[f_{\epsilon}(t,x,v)\mathbf{1}_{|v|>0}\mathbf{1}_{A_{1}^{\epsilon}(x,v)}] + O(\epsilon).$$

We write then, for $x \in Z_{\epsilon}[C]$,

$$\mathbb{E}[f_{\epsilon}(t,x,v)\mathbf{1}_{|v|>0}\mathbf{1}_{A_{1}^{\epsilon}(x,v)}]$$

$$=\sum_{N\geq0}\frac{\lambda_{\epsilon}^{N}}{N!}\int dc_{1}\dots dc_{N}\,\mathbf{1}_{|v|>0}\mathbf{1}_{A_{1}^{\epsilon}(x,v)}\,e^{-\lambda_{\epsilon}|\mathcal{T}_{\epsilon}(-t;x,v)(C)|(c_{1},\dots,c_{N})}$$

$$(39)\quad\mathbf{1}_{C\cap\mathcal{T}_{\epsilon}(-t;x,v)(C)=(c_{1},\dots,c_{N})}f_{0}((Y_{t}^{\epsilon}(x,-v),-Z_{t}^{\epsilon}(x,-v)))J_{\epsilon,C}(t,x,v)$$

which, because of the considerations in Section 4.4, can be rewritten as

$$\mathbb{E}[f_{\epsilon}(t, x, v) \mathbf{1}_{|v|>0} \mathbf{1}_{A_{1}^{\epsilon}}] = \sum_{N\geq 0} \lambda_{\epsilon}^{N} \int dc_{1} \dots dc_{N} \mathbf{1}_{[A_{1}^{\epsilon}(t, x, v)]_{N}}([C]_{N})$$

$$\times \exp\left\{-\lambda B^{d-1}\left[\sum_{j=0}^{N} |Z_{\tau_{i,+}^{j+1}}^{\epsilon}|(\tau_{i,+}^{j+1} - \tau_{i,-}^{j}) + O(\epsilon)\right]\right\}$$

$$\times f_{0}\left(x + \sum_{j=0}^{N} Z(\tau_{i,+}^{j+1})(\tau_{i,+}^{j+1} - \tau_{i,-}^{j}) + O(\epsilon),$$

$$(40) \qquad v + \hat{v}\sum_{j=1}^{N} \frac{\kappa}{\epsilon} \int_{\tau_{i,+}^{j}}^{\tau_{i,-}^{j}} S(|Z|(s))|Z(s)|ds\right)$$

$$\times \left(\frac{|v| + \sum_{j=1}^{N} \frac{\kappa}{\epsilon} \int_{\tau_{i,+}^{j}}^{\tau_{i,-}^{j}} S(|Z|(s))|Z(s)|ds)}{|v|}\right)^{d}$$

$$\times \frac{S\left(|v| + \sum_{j=1}^{N} \frac{\kappa}{\epsilon} \int_{\tau_{i,+}^{j}}^{\tau_{i,-}^{j}} S(|Z|(s))|Z(s)|ds)\right)}{S(|v|)}.$$

By the change of variables (27), and using the relations

$$\frac{|Z_{\tau_{i,-}^{N}}^{\epsilon}|}{|v|} = \prod_{j=1}^{N} \frac{\left|Z_{\tau_{i,-}^{j}}^{\epsilon}\right|}{\left|Z_{\tau_{i,+}^{j}}^{\epsilon}\right|}$$

and

$$\frac{S(|Z_{\tau_{i,-}^{\kappa}}^{\epsilon}|)}{S(|v|)} = \prod_{j=1}^{N} \frac{S(|Z_{\tau_{i,-}^{j}}^{\epsilon}|)}{S(|Z_{\tau_{i,+}^{j}}^{\epsilon}|)},$$

we get (recall that $\lambda_{\epsilon} = \frac{\lambda}{\epsilon^{d-1}}$)

$$\mathbb{E}[f_{\epsilon}(t, x, v)\mathbf{1}_{|v|>0}\mathbf{1}_{A_{1}^{\epsilon}}] = \sum_{N\geq0} \lambda^{N} \int_{0}^{t} d\tau_{i,+}^{N} \int_{\nu_{N}^{2}\leq1}^{1} d\nu_{N} \dots \int_{0}^{\tau_{i,+}^{2}} d\tau_{i,+}^{1} \int_{\nu_{1}^{2}\leq1}^{1} d\nu_{1}$$

$$(41) \qquad \times \prod_{j=1}^{N} \left[|Z_{\tau_{j,+}^{j}}^{\epsilon}| \left(\frac{|Z_{\tau_{j,+}^{j}}^{\epsilon}|}{|Z_{\tau_{i,+}^{j}}^{\epsilon}|} \right)^{d} \frac{S(|Z_{\tau_{j,+}^{j}}^{\epsilon}|)}{S(|Z_{\tau_{i,+}^{j}}^{\epsilon}|)} \right]$$

$$\times \exp\left\{ -\lambda B^{d-1} \left[\sum_{j=0}^{N} |Z_{\tau_{i,+}^{j+1}}^{\epsilon}| (\tau_{i,+}^{j+1} - \tau_{i,-}^{j}) + O(\epsilon) \right] \right\}$$

$$\times f_{0} \left(x + \sum_{j=0}^{N} Z(\tau_{i,+}^{j+1}) (\tau_{i,+}^{j+1} - \tau_{i,-}^{j}) + O(\epsilon) ,$$

$$v + \hat{v} \sum_{j=1}^{N} \frac{\kappa}{\epsilon} \int_{\tau_{i,+}^{j}}^{\tau_{i,-}^{j}} S(|Z|(s)) |Z(s)| ds) \right)$$

or equivalently

$$\mathbb{E}[f_{\epsilon}(t, x, v) \mathbf{1}_{|v|>0} \mathbf{1}_{A_{1}^{\epsilon}}] = \sum_{N\geq 0} (\lambda B^{d-1})^{N} \int_{0}^{t} d\tau_{i,+}^{N} \int_{0}^{1} dh_{N}^{d-1} \dots \int_{0}^{\tau_{i,+}^{2}} d\tau_{i,+}^{1} \int_{0}^{1} dh_{1}^{d-1} (42) \qquad \times \prod_{j=1}^{N} \left[|Z_{\tau_{i,+}^{j}}^{\epsilon}| \left(\frac{|Z_{\tau_{i,+}^{j}}^{\epsilon}|}{|Z_{\tau_{i,+}^{j}}^{\epsilon}|} \right)^{d} \frac{S(|Z_{\tau_{i,+}^{j}}^{\epsilon}|)}{S(|Z_{\tau_{i,+}^{j}}^{\epsilon}|)} \right] \qquad \times \exp\left\{ -\lambda B^{d-1} \left[\sum_{j=0}^{N} |Z_{\tau_{i,+}^{j+1}}^{\epsilon}| (\tau_{i,+}^{j+1} - \tau_{i,-}^{j}) + O(\epsilon) \right] \right\} \qquad f_{0} \left(x - \sum_{j=0}^{N} Z(\tau_{i,+}^{j+1}) (\tau_{i,+}^{j+1} - \tau_{i,-}^{j}) + O(\epsilon) , \qquad v + \hat{v} \sum_{j=1}^{N} \frac{\kappa}{\epsilon} \int_{\tau_{i,+}^{j}}^{\tau_{i,-}^{j}} S(|Z|(s)) |Z(s)| ds \right).$$

We may now perform the limit, using (36)-(37) and recalling that $\mathbf{P}(x \in$ $B_{\epsilon}(c)$ < ϵ , so that, for all $\phi \in C_c(\mathbb{R}^d \times \mathbb{R}^d \setminus \{0\})$, we have

$$\begin{split} \lim_{\epsilon \to 0^+} \int_{\mathbb{R}^d \times \mathbb{R}^d} dx dv \, \phi(x, v) \, \mathbb{E}[f_{\epsilon}(t, x, v)] \\ &= \lim_{\epsilon \to 0^+} \int_{\mathbb{R}^d \times \mathbb{R}^d} dx dv \, \phi(x, v) \, \mathbb{E}[f_{\epsilon}(t, x, v) \mathbf{1}_{|v| > 0} \mathbf{1}_{A_1^{\epsilon}}] \\ &= \int_{\mathbb{R}^d \times \mathbb{R}^d} dx dv \phi(x, v) F(t, x, v), \end{split}$$

where

$$F(t, x, v) = \sum_{N \ge 0} (\lambda B^{d-1})^N \int_0^t d\tau_{i,+}^N \int_0^1 dh_N^{d-1} \dots \int_0^{\tau_{i,+}^2} d\tau_{i,+}^1 \int_0^1 dh_1^{d-1} (43) \times \prod_{j=1}^N \left[|Z_{\tau_{i,+}^j}^-| \left(\frac{|Z_{\tau_{i,+}^j}^+|}{|Z_{\tau_{i,+}^j}^-|} \right)^d \frac{S(|Z_{\tau_{i,+}^j}^+|)}{S(|Z_{\tau_{i,+}^j}^-|)} \right] \\ \times \exp\left\{ -\lambda B^{d-1} \left[\sum_{j=0}^N |Z_{\tau_{i,+}^{j+1}}^-| (\tau_{i,+}^{j+1} - \tau_{i,+}^j) \right] \right\} \\ \times f_0 \left(x + \sum_{j=0}^N Z_{\tau_{i,+}^{j+1}}^- (\tau_{i,+}^{j+1} - \tau_{i,+}^j), v + \sum_{j=1}^N \left[Z_{\tau_{i,+}^j}^+ - Z_{\tau_{i,+}^j}^- \right] \right) \right)$$

is the series solution to (38).

Proof of Theorem 1.

We proved in Theorem 9 that $\mathbb{E}[f_{\epsilon}(t, x, v; C)]dxdv \xrightarrow{*} F(t, x, v)dxdv$ on $\mathbb{R}^d \times \mathbb{R}^d \setminus \{0\}$, where F(t, x, v) solves (38) in $\mathcal{D}'((0, +\infty) \times \mathbb{R}^d \times \mathbb{R}^d \setminus \{0\})$. Therefore the limit point F(t, dxdv)dt s.t. $F(t, dxdv)dt|_{(0,+\infty)\times\mathbb{R}^d\times\mathbb{R}^d\setminus\{0\}} =$ F(t, x, v) dt dx dv solves equation (9) with

$$m_F = \sigma \int_0^1 dh^{d-1} \frac{|z|^d}{|v|^{d-1}} \frac{S(|z|)}{S(|v|)} F(t, x, z) \mathbf{1}_{|v|>0} dx dv$$

and

$$\ell(v) = \sigma |v|.$$

Since the hypotheses in Lemma 5 are verified, we have:

(44)
$$(\partial_t + v \cdot \nabla_x + \ell(v))F - m_F = \lambda_F(t, x)\delta_0(dv)dx$$
.
in $\mathcal{D}'((0, +\infty) \times \mathbb{R}^d \times \mathbb{R}^d)$.

Since the jacobian of the change of coordinate $z \to v$ is

$$J' = \left(\frac{|z|}{|v|}\right)^{d-1} \frac{S(|z|)}{S(|v|)}$$

and $|v| > 0 \implies |z| > a^{-1}(2\kappa\sqrt{1-h^2}), \lambda_F(t,x)$ is given by

$$\lambda_F(t, dx) = \int_{\mathbb{R}^d} \ell(v) F(t, dx dv) - \int_{\mathbb{R}^d} m_F(t, dx dv)$$
$$= \int_{\mathbb{R}^d} dz \int_0^1 dh^{d-1} |z| F(t, x, z) \mathbf{1}_{|z| \le a^{-1} (2\kappa\sqrt{1-h^2})} dx.$$

Since the solution to (4) is unique by Lemma 6 and the weak-* limit of $\mathbb{E}[f_{\epsilon}(t, dxdv; C)]dt$ is also a weak limit, Theorem 1 is proved.

This concludes our description of the Boltzmann–Grad asymptotics for the particle dynamics described by (1).

Finally, we have a few remarks.

The first one is that, although the dynamics (1) at $\epsilon > 0$ induces an evolution on the Radon measures which preserves mass, in the limit mesoscopic evolution equation for the number density function (in the phase space) of the light particles the collision integral does not preserve mass. The conservation of mass is nevertheless guaranteed by the presence in the limit kinetic equation of a point measure component of the form $\lambda(t, x)\delta_0(dv)dx$ proportional to the Dirac delta distribution in 0, where $\lambda(t, x)$ is a sort of "Lagrange multiplier" of the conservation of mass, which describes the particles stopping in the random medium.

The second remark is that, because the collision integral does not satisfy the conservation of mass and the flow (1) is irreversible, although the dynamics of the particle system under exam looks quite simple, the asymptotics behaviour of its number density profile is more tricky to analyse than for systems like the ones described, e.g. in [14] or [15]. In particular, it is not possible to reduce the analysis to the asymptotics of the Markovian component of the number density (for a review on the terminology, see [16]).

DECLARATIONS

Conflict of interest: the authors declare that there are no conflicts of interest related to this article.

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no datasets were generated or analyzed during the current study.

References

- Duderstadt, J.J., Moses, G.A.: Inertial Confinement Fusion. J. Wiley and Sons, Inc. (1982).
- [2] Atzeni, S., Meyer Ter-Vehn, J.: The Physics of Inertial Fusion. Clarendon Press-Oxford (2004).
- [3] Azechi, H., Cable, M.D., Stapf, R.O.: Review of secondary and tertiary reactions, and neutron scattering as diagnostic techniques for inertial confinement fusion targets. Laser and Particle Beams 9, 119–134 (1993). https://doi.org/10.1017/S0263034600002378
- [4] Cable, M.D., Hatchett, S.P.: Neutron spectra from inertial confinement fusion targets for measurement of fuel areal density and charged particle stopping powers. J. Appl. Phys. 62, 2233-2236 (1987). https://doi.org/10.1063/1.339850
- Richtmyer, R.D.: Taylor Instability in Shock Acceleration of Compressible Fluids. Comm. on Pure and Appl. Math. 13, 297–319 (1960). https://doi.org/10.1002/cpa.3160130207
- [6] Levermore, C.D., Zimmermann, G.B.: Modeling charged particle loss in a fuel/shell mixture. J. Math. Phys. 34(10), 4725-4729 (1993). https://doi.org/10.1063/1.530367
- [7] Levermore, C.D., Pomraning, G.C., Sanzo, D.L., Wong J.: Linear transport theory in a random medium. J. Math. Phys.27(10), 2526-2536 (1986). https://doi.org/10.1063/1.527320
- [8] Clouet, J.-F., Golse, F., Puel, М., Sentis, R.: On the slowinbinary ing down of charged particles \mathbf{a} stochastic mixture. Kinetic and Related Models 1 (3).387 - 404(2008).https://www.aimsciences.org/article/doi/10.3934/krm.2008.1.387
- [9] Ambrosio, L., Gigli, N., Savaré, G.: Gradient Flows in Metric Spaces and in the Space of Probability Measures. 2nd ed. Birkhäuser Verlag AG, Basel, Boston, Berlin (2008)
- [10] Dieudonné, J.: Treatise on Analysis. Vol. 2. Academic Press, Inc. (1976)
- [11] Schwartz, L.: Théorie des distributions. Hermann Paris (1966)
- [12] Malliavin, P.: Integration and Probability. Springer-Verlag New York, Inc. (1983)
- [13] Pazy, A.: Semigroups of Operators and Applications to Partial Differential Equations. Springer-Verlag New York, Inc. (1983)
- [14] Gallavotti, G.: Rigorous theory of the Boltzmann equation in the Lorentz gas. Nota Interna No. 358, Istituto di Fisica, Università di Roma (1972), reprint in Statistical Mechanics, 1999. https://ipparco.roma1.infn.it/pagine/deposito/1967-1979/041-Bz.pdf

- [15] Desvillettes, L., Ricci, V.: A Rigorous Derivation of a Linear Kinetic Equation of Fokker-Planck Type in the Limit of Grazing Collisions. Journal of STatistical Physics 104 (5/6) 1173-1189 (2001). https://doi.org/10.1023/A:1010461929872
- [16] Ricci, V.: Non Markovian Behavior of the Boltzmann-Grad Limit of Linear Stochastic Particle Systems. Communications in Mathematical Sciences, Suppl. 1, 95–105 (2007). https://dx.doi.org/10.4310/CMS.2007.v5.n5.a8