Embedding Network Autoregression for time series analysis and causal peer effect inference

Jae Ho Chang

Department of Statistics, The Ohio State University

and

Subhadeep Paul Department of Statistics, The Ohio State University

Abstract

We propose an Embedding Network Autoregressive Model (ENAR) for multivariate networked longitudinal data. We assume the network is generated from a latent variable model, and these unobserved variables are included in a structural peer effect model or a time series network autoregressive model as additive effects. This approach takes a unified view of two related problems, (1) modeling and predicting multivariate time series data and (2) causal peer influence estimation in the presence of homophily from finite time longitudinal data. Our estimation strategy comprises estimating latent factors from the observed network adjacency matrix either through spectral embedding or maximum likelihood estimation, followed by least squares estimation of the network autoregressive model. We show that the estimated momentum and peer effect parameters are consistent and asymptotically normal in asymptotic setups with a growing number of network vertices N while including a growing number of time points T and finite T cases. We allow the number of latent vectors K to grow at appropriate rates, which improves upon existing rates when such results are available for related models.

Keywords: Network time series, social influence, peer effect, social network, latent homophily, network embedding

1 Introduction

A network of relationships and longitudinal node-level responses commonly appear in research problems in multiple domains, including social sciences, economics, public health, and biomedical sciences. We consider two key statistical problems associated with such data that have been widely investigated in the literature. The first is to causally estimate peer effects or social influence propagating through an observed network when the node level outcome of interest is measured in at least two-time points (Shalizi and Thomas, 2011; VanderWeele, 2011; Goldsmith-Pinkham and Imbens, 2013; McFowland III and Shalizi, 2021; Nath et al., 2022). The second problem is to model and predict a high dimensional time series when a network information is also observed (Zhu et al., 2017, 2019; Knight et al., 2020; Zhu and Pan, 2020; Chen et al., 2023). We take a unified view of these two problems and propose to include latent homophily variables in both of these problems to aid causal identification in the former case and improved prediction and model estimation in the latter case.

It will be convenient to formally introduce the network autoregressive model (NAR or NAM) model, which has been historically used for both problems, to facilitate further discussion on its interpretation. We assume that we have measurements y_{it} for i = 1, ..., N, and t = 0, ..., T - 1 with $N, T \in \mathbb{N}$ for an univariate outcome measured at N vertices of a network over T time periods. We denote the undirected network adjacency matrix with **A** and define its normalized (symmetric) Laplacian matrix as $\mathcal{L}_{\mathbf{A}} = \mathcal{D}^{-1/2} \mathbf{A} \mathcal{D}^{-1/2}$ with ℓ_{ij} denoting its (i, j)-th entry and \mathcal{D} is a diagonal matrix containing its degrees. Then, our measurements y_{it} are assumed to be generated via

$$y_{i,t+1} = \alpha y_{it} + \theta \sum_{j \neq i} \ell_{ij} y_{jt} + \mathbf{z}_{it}^{\mathsf{T}} \gamma + \epsilon_{i,t+1}, \qquad (1)$$

where z_{it} is a vector of (possibly time varying) covariates and ϵ_{it} is the error term. We differentiate between the utility and the interpretation of this model in terms of whether T is finite or growing, with the former being useful as a linear structural model for peer influence estimation and the latter being useful for multivariate time series modeling. The above NAR or NAM model with longitudinally measured outcomes in Equation 1 has been widely employed for the causal identification of peer influence (Christakis and Fowler, 2007; Shalizi and Thomas, 2011; VanderWeele et al., 2012; Christakis and Fowler, 2013; O'Malley et al., 2014; McFowland III and Shalizi, 2021; Nath et al., 2022). Typical restrictions on the model would be an assumption of exogeneity of error term $E[\epsilon_{i,t+1}|l_{ij}, z_{it}, y_{it}] = 0$, along with the regressors not being linearly dependent. However, several authors have noted issues with identifying peer effects from observational data with such models, including confounding due to latent homophily in peer selection, and other unobserved omitted variables (Shalizi and Thomas, 2011; Goldsmith-Pinkham and Imbens, 2013; O'Malley et al., 2014; An et al., 2022). For the longitudinal peer effects models, Mc-Fowland III and Shalizi (2021); Nath et al. (2022) suggest augmenting the linear peer effects model with additive latent variables, which are responsible for network formation as a way of controlling for latent homophily. They proceed to show that the peer influence effect can be estimated in an asymptotically unbiased way.

In the context of modeling and predicting multivariate high-dimensional network-linked time series, a line of work including Zhu et al. (2017, 2019); Knight et al. (2020); Zhu and Pan (2020); Chen et al. (2023) termed the above model the network vector autoregressive model. Those papers then investigate the stationarity of the model along with consistency and asymptotic normality of the parameter estimates in an asymptotic setup where $T \to \infty$ under the assumptions $\epsilon_{it} \stackrel{iid}{\sim} \mathcal{N}(0, \sigma^2)$. Several extensions of the NAR model in Equation 1 has been proposed, including the Community NAR (CNAR) model in Chen et al. (2023) which introduces community-dependent heterogeneous network effects and Grouped NAR Zhu and Pan (2020) which allows the peer effect parameter to differ by groups.

Our proposal in this paper is to augment the NAR model with latent variables that are

related to both the outcome and the formation of the network. Accordingly, we assume that the network adjacency matrix is generated from either the Random Dot Product Graph (RDPG) model Athreya et al. (2017), or the additive and multiplicative effects latent space model Hoff (2021); Ma et al. (2020). The RDPG model is a general latent variable model for network which contains several popular latent variable models for networks namely, the Stochastic Block Model (SBM), Degree Corrected SBM (DCSBM), Mixed Membership SBM (MMSBM), and DCMMSBM as special cases (Athreva et al., 2017; Rubin-Delanchy et al., 2022). This model can also be thought of as related to the multiplicative part of the additive and multiplicative latent factor model (Ma et al., 2020; Hoff, 2021; Li et al., 2023). Specifically, we assume network adjacency matrix \mathbf{A} is generated from an RDPG model (defined later) with parameters $\rho_N, \mathbf{X} \in \mathbb{R}^{N \times K}$ such that the probability of connections $\mathbf{P} = \rho_N \mathbf{X} \mathbf{X}^{\intercal}$. We let $\mathbf{U}_{\mathbf{P}} \in \mathbb{R}^{N \times K}$ be the matrix of orthogonal eigenvectors of \mathbf{P} for K leading eigenvalues, which contains information about the associated latent positions. Then, our ENAR model augments the NAR model with these K dimensional unknown (latent) eigen vector variables. Formally, the ENAR model assumes that the measurements y_{it} for i = 1, ..., N, are assumed to be generated via

$$y_{i,t+1} = \alpha y_{it} + \theta \sum_{j \neq i} \ell_{ij} y_{jt} + \mathbf{u}_i^{\mathsf{T}} \beta + \mathbf{z}_{it}^{\mathsf{T}} \gamma + \epsilon_{it}$$

However, the vectors of latent variables \mathbf{u}_i s are not observed. Therefore our proposal is to estimate the latent variables from the observed network and replace \mathbf{u}_i with its estimated version $\hat{\mathbf{u}}_i$. This idea has been explored previously in the peer influence literature (McFowland III and Shalizi, 2021; Nath et al., 2022) as a way of removing the omitted variable bias due to homophily. However, a natural concern is whether the true peer influence parameter θ can be estimated consistently and the asymptotic variance can be characterized to enable inference when the \mathbf{u}_i is replaced with estimated $\hat{\mathbf{u}}_i$. McFowland III and Shalizi (2021) considered this issue as trading off omitted variable bias with measurement error bias and intuitively will only succeed if the measurement error bias is low. In this article, we show that the answer is affirmative and develop a theory for the consistency and asymptotic normality of the peer influence parameter. In addition, we propose to include estimated latent variables in the context of time series modeling as well to enable accurate inference on parameters of the model as well as improve predictive performance.

We summarize our theoretical results for both of these statistical goals. We always operate under a large N asymptotic i.e., assume $N \to \infty$. We define the set of parameters as $\mu^{H} \triangleq (\beta^{\mathsf{T}}\mathbf{H}, \alpha, \theta, \gamma^{\mathsf{T}})^{\mathsf{T}}$ with $\mathbf{H} \in \mathcal{O}_{K}$, is a $K \times K$ matrix with orthonormal columns. It is well known that the multiplicative latent variables can only be estimated from a network up to an ambiguity of such a matrix from the class \mathcal{O}_K (Hoff et al., 2002; Athreya et al., 2017). Therefore, the parameter β can only be recovered up to the ambiguity of $\beta^{\dagger} \mathbf{H}$. In the case of modeling time series, we assume $T \to \infty$. We show that under certain regularity conditions on the eigengap π_N of the network, namely, $\pi_N = \omega(\sqrt{KNT\rho_N})$, the estimated parameter vector $\hat{\mu}$, suitably normalized, converges to a multivariate normal distribution with finite variance around the true parameter vector μ^H as long as $K^2 = o(N)$ and log K = o(T). The rates of convergence are different for different set of parameters, namely, it is \sqrt{T} for the parameters $\beta^{\mathsf{T}}\mathbf{H}$, while it is \sqrt{NT} for the remaining model parameters. To compare with existing results in related literature, the asymptotic growth rates necessary for the CNAR model in Chen et al. (2023) (which albeit is a different but related model) is $K^4 = o(N)$ and $K^2 \log K = o(T)$ and the eigengap $\pi_N = \omega(K\sqrt{NT\rho_N})$. Therefore, our results on the ENAR model require strictly weaker assumptions on the eigengap and the necessary sample size. In simulation, we see that ENAR can consistently estimate the latent effects and network effects, yet it achieves comparable prediction performance to CNAR and NAR.

Second, for the case of finite T (with $T \ge 2$), we are interested in the accurate inference of the peer influence parameter θ . Our result shows if $\pi_N = \omega(\sqrt{KN\rho_N})$ and $N = \omega(K^2)$, while $\hat{\beta} - \mathbf{H}^{\mathsf{T}} \beta = O_{\mathbb{P}}(1)$, the rest of the parameters including the peer influence parameter is estimated with a \sqrt{N} convergence to a multivariate normal distribution with finite variance. This result can be compared with the results in McFowland III and Shalizi (2021); Nath et al. (2022) where the authors showed asymptotic unbiasedness of the peer influence parameter under the SBM and the RDPG models respectively. In contrast, our results hold for RDPG models with growing dimension of the latent space K and generalize and supplement those results to include consistency and asymptotic normality. Finally, in the case of T = 1, when the interest is estimating the effect of covariates controlling for homophily related latent variables (termed as Network Regression), our results show that the regression coefficients for the covariates are estimated at \sqrt{N} rate.

We also consider the additive and multiplicative effects latent space model (LSN) (Hoff, 2021; Ma et al., 2020; Li et al., 2023) for modeling the network data. We incorporate these latent variables from the network model with appropriate scaling into a model that we call the additive and multiplicative NAR (AMNAR) model. For estimation, the latent factors are estimated from a maximum likelihood estimator (Ma et al., 2020). We study the consistency and asymptotic normality properties of the resulting estimators and show that the momentum and peer effect parameters can be estimated at \sqrt{NT} rate.

2 Embedding Network Autoregressive Model

In this section, we describe the proposed Embedding Network Autoregressive (ENAR) model. We start by defining our notations, then describe the NAR and RDPG models, which the ENAR model builds upon. Then we describe our model with motivations from both multivariate time series analysis and causal peer influence estimation. Finally, we also propose an extension of ENAR based on the additive and multiplicative latent variables from the LSN model (AMNAR).

Let a, \mathbf{a} , and \mathbf{A} be generic notations for scalars, vectors, and matrices, respectively. Let \mathbb{I}_A denotes an indicator with a support set A. For a matrix $\mathbf{A} \in \mathbb{R}^{n \times m}$, write its element at (i, j)-th entry as a_{ij} . Let $\mathcal{O}_{N,K}$ and \mathcal{O}_K be collections of $N \times K$ and $K \times K$ matrices with real orthonormal columns, respectively. Write an m-dimensional vector and $m \times m$ matrix with zeros as \mathbf{o}_m and \mathbf{O}_m , and let \mathbf{I}_m denotes an $m \times m$ identity matrix.

For an *n*-row square matrix \mathbf{X} , let tr $(\mathbf{X}) \triangleq \sum_{i=1}^{n} x_{ii}$ denote its trace and $\lambda_i(\mathbf{X})$ denote the *i*-th leading eigenvalue of \mathbf{X} (hence $|\lambda_1(\mathbf{X})| \geq |\lambda_2(\mathbf{X})| \geq \cdots \geq |\lambda_n(\mathbf{X})|$). The *i*th leading eigenvector of \mathbf{X} will mean the eigenvector corresponding to this *i*th leading eigenvalue. Let $\rho(\mathbf{X}) \triangleq |\lambda_1(\mathbf{X})|$ denote its spectral radius. Following are some matrix norms: ℓ_{∞} -norm $\|\mathbf{X}\|_{\infty} \triangleq \max_{i} \sum_{j} |x_{ij}|, \ell_2$ -norm $\|\mathbf{X}\| \triangleq \lambda_1 (\mathbf{X}^{\mathsf{T}} \mathbf{X})^{1/2}$, and Frobenius norm $\|\mathbf{X}\|_F \triangleq \operatorname{tr} (\mathbf{X}^{\mathsf{T}} \mathbf{X})^{1/2}$.

For $a_n, b_n \geq 0$, we write $a_n \gtrsim b_n$ if there exists c > 0 independent with n such that $a_n \geq cb_n$ for all n. Also, write $a_n = \omega(b_n)$ if for all e > 0 there exists $n_0 > 0$ such that $a_n > eb_n$ for any $n > n_0$. $f_n = \Theta(g_n)$ means that both $f_n = O(g_n)$ and $g_n = O(f_n)$ hold. We say $X_n = O(Y_n)$ whp. (with high probability) if for any c > 0 there exist $C = C(c) > 0, n_0 > 0$ such that $\mathbb{P}(|X_n| > C|Y_n|) < n^{-c}$ for all $n > n_0$. Also, we write $X_n = O(Y_n)$ as. if $|X_n| \leq |Y_n|$ almost surely, and write $X_n = O(1)$ as. if there exists M > 0 such that $|X_n| \leq M$ for all n almost surely. Finally, we denote a weak convergence of a sequence of random variables by $X_n \Rightarrow X$, and for a sequence of random vectors $\mathbf{x}_n \in \mathbb{R}^p$, write $\mathbf{x}_n = O_{\mathbb{P}}(a_n)$ if $\eta^{\mathsf{T}} \mathbf{x}_n / a_n$ is bounded in probability and $\mathbf{x}_n = o_{\mathbb{P}}(a_n)$ if $\eta^{\mathsf{T}} \mathbf{x}_n / a_n$ converges to zero in probability for all $\eta \in \mathbb{R}^p$ such that $||\eta|| \leq 1$.

2.1 Network Vector Autoregression

As stated in the introduction, we assume a statistical problem where we have time series or longitudinal measurements y_{it} on an univariate outcome over N subjects at T time points. We let i = 1, ..., N and t = 0, ..., T - 1 with $(N, T \in \mathbb{N})$. We further assume that these Nindividuals are connected in a network with an (undirected) adjacency matrix \mathbf{A} , which is also observed. For each unit i, we further have measurements on p dimensional covariates \mathbf{z}_{it} , where the subscript t indicates that the covariates may vary over time. Recall the network vector autoregressive model (Zhu et al., 2017; McFowland III and Shalizi, 2021) in Equation 1, $y_{i,t+1} = \alpha y_{it} + \theta \sum_{j \neq i} \ell_{ij} y_{jt} + \mathbf{z}_{it}^{\mathsf{T}} \gamma + \epsilon_{i,t+1}$. We assume that $\mathbf{z}_{it} \in \mathbb{R}^p$ are i.i.d. sub-gaussian random vectors with independent coordinates with zero mean and finite fourth-order-moments, while $\Sigma_z \triangleq \operatorname{diag}(v_1, ..., v_p)$ denotes their common covariance with strictly positive diagonals. This assumption is similar to one described in Chen et al. (2023). For the model errors, we assume $\epsilon_{it} \stackrel{id}{\sim} \mathcal{N}(0, \sigma^2)$ for $\sigma > 0$. Among the parameters, $\alpha \in \mathbb{R}$ denotes the momentum effect, $\theta \in \mathbb{R}$ denotes the peer influence effect, and $\gamma \in \mathbb{R}^p$ denotes the time-invariant covariate effects (Zhu et al., 2017).

Let $\mathcal{E}_t \triangleq (\epsilon_{1t}, \ldots, \epsilon_{Nt})^{\mathsf{T}} \in \mathbb{R}^N$, and $\mathbf{y}_t \triangleq (y_{1t}, \ldots, y_{Nt})^{\mathsf{T}} \in \mathbb{R}^N$ be the vectorized forms of the error term and the response obtained by stacking the corresponding terms for the N individuals. Similarly, let $\mathbf{Z}_t \triangleq [\mathbf{z}_{1t}, \ldots, \mathbf{z}_{Nt}]^{\mathsf{T}} \in \mathbb{R}^{N \times p}$ be the matrix of covariates whose *i*th row is \mathbf{z}_{it} , the covariate for the *i*th subject. Then the above model can be expressed in the vector and matrix notations as $\mathbf{y}_{t+1} = \alpha \mathbf{y}_t + \theta \mathcal{L}_{\mathbf{A}} \mathbf{y}_t + \mathbf{Z}_t \gamma + \mathcal{E}_{t+1}$.

2.2 Random Dot Product Graph

Latent position random graph models assume that a network is created by random edges independently sampled over the Euclidean space constrained with respect to a kernel of latent positions associated with edges. In a K-dimensional random dot product graph (RDPG), this kernel is the dot product of two K-dimensional latent vectors. We outline the RDPG model below.

Definition 1. Let \mathcal{X} be a subset of \mathbb{R}^{K} such that $\mathbf{x}_{1}^{\mathsf{T}}\mathbf{x}_{2} \in [0, 1]$ for all $\mathbf{x}_{1}, \mathbf{x}_{2} \in \mathcal{X}$. Let $K \leq N$ and ρ_{N} be a sequence such that $\rho_{N} \in (0, 1]$ for all N. Then, \mathbf{A} is said to follow a random dot product graph with latent positions $\mathbf{X} = [\mathbf{x}_{1}, ..., \mathbf{x}_{N}]^{\mathsf{T}} \in \mathcal{X}^{N}$ and sparsity factor ρ_{N} , denoted by $\mathbf{A} \sim \text{RDPG}(\rho_{N}, \mathbf{X})$, if $a_{ij} \stackrel{ind.}{\sim} \text{Bernoulli}(\rho_{N}\mathbf{x}_{i}^{\mathsf{T}}\mathbf{x}_{j})\mathbb{I}_{\{i \neq j\}}, \quad 1 \leq i, j \leq N$.

Note that the indicator $\mathbb{I}_{\{i \neq j\}}$ in the above definition ensures that **A** is hollow without self-loops. Using matrix notations, we have $\mathbb{E}[\mathbf{A}] = \mathbf{P}$ for $\mathbf{P} \triangleq \rho_N \mathbf{X} \mathbf{X}^{\intercal}$. The role of ρ_N is controlling the sparsity of the network. For example, the expected degrees are $\sum_{j=1}^{N} p_{ij} \in$ $[0, N\rho_N]$ for every i = 1, ..., N. Hence when $\rho_N = 1 \forall N$, the resulting graph is dense in the sense that expected number of edges $\sum_{i < j} p_{ij} \sim N^2$. If $\rho_N \to 0$ as $N \to \infty$, the graph becomes sparse in the sense that $\sum_{i < j} p_{ij} = o(N^2)$ (Xie and Xu, 2023).

Under our asymptotic framework where the network size N grows, it's reasonable to expect that the dimension of the latent space, denoted as K, will also increase. Under moderate sparsity assumption, the difference in eigenvectors of \mathbf{P} and \mathbf{A} remains bounded in probability (Lei and Rinaldo, 2015). The estimation of latent positions for random graphs has received extensive attention in the literature (Tang and Priebe, 2018; Cape et al., 2019; Xie and Xu, 2023; Rubin-Delanchy et al., 2022), and the results often involve an assumption on the minimum growth rate of ρ_n .

2.3 Embedding Network Autoregression

Now we define our model which augments the NAR model with latent variables that are common for both the model of the univariate responses and the model for the network. Accordingly, in this section, we further assume the network is generated from a RDPG model, $\mathbf{A} \sim \text{RDPG}(\rho_N, \mathbf{X})$. We assume that every vertex in \mathbf{A} is connected to at least one vertex.

The spectral decomposition of \mathbf{P} can be expressed as $\mathbf{P} = \mathbf{U}_{\mathbf{P}} \mathbf{S}_{\mathbf{P}} \mathbf{U}_{\mathbf{P}}^{\mathsf{T}}$ where $\mathbf{U}_{\mathbf{P}} \in \mathbb{R}^{N \times K}$ contains orthogonal eigenvectors for K leading eigenvalues, and $\mathbf{S}_{\mathbf{P}}$ is a diagonal matrix containing those eigenvalues. Simply write $\mathbf{U} \triangleq \mathbf{U}_{\mathbf{P}}$ so that $\mathbf{U} = [\mathbf{u}_1, ..., \mathbf{u}_N]^{\mathsf{T}}$ contains information about the associated latent position of each vertex. Then, we define the ENAR model as a set of two models as follows,

$$y_{i,t+1} = \alpha y_{it} + \theta \sum_{j \neq i} \ell_{ij} y_{jt} + \mathbf{u}_i^{\mathsf{T}} \beta + \mathbf{z}_{it}^{\mathsf{T}} \gamma + \epsilon_{it}, \qquad (2)$$
$$a_{ij} \stackrel{ind.}{\sim} \text{Bernoulli}(\rho_N \mathbf{x}_i^{\mathsf{T}} \mathbf{x}_j) \mathbb{I}_{\{i \neq j\}}.$$

As before, we can write the model in vector and matrix notation as follows,

$$\mathbf{y}_{t+1} = \alpha \mathbf{y}_t + \theta \mathcal{L}_{\mathbf{A}} \mathbf{y}_t + \mathbf{U}\beta + \mathbf{Z}_t \gamma + \mathcal{E}_{t+1}, \tag{3}$$

where now the parameter β denotes the global effect of latent positions, and the other parameters have the same meaning as before. Finally, assume that \mathbf{Z}_t is independent with $\{\mathcal{E}_{t+1}, \mathcal{E}_t, \ldots\}$ and $\{\mathbf{y}_t, \mathbf{y}_{t-1}, \ldots\}$ for each t and \mathbf{A} is generated independently with the rest of random components across all t.

2.4 Finite time model for peer influence

The ENAR model is also motivated from the problem of estimating causal peer influence adjusting for latent homophily in the settings of longitudinal data but perhaps with finite time points (e.g., T = 2). The model is identical to the ENAR model described in the earlier section, except we do not have a time series, but only a finite number of time periods (perhaps just 2) and the asymptotic setup is with respect to $N \to \infty$. The problem can be illustrated with a causal diagram similar to McFowland III and Shalizi (2021); Nath et al. (2022). In the causal diagram Pearl (2009) in Figure 1, the observed variables are represented by rectangles while the unobserved or latent variables are represented by circles. Intuitively, the causal peer influence is the causal effect of outcome of a "peer" who is linked in the network on the outcome of an individual. The problem of estimating causal peer influence is then estimating the causal effect corresponding to the path $Y_j^t \to Y_i^{t+1}$, conditioning on the observed network links a_{ij} s.

Figure 1: Causal diagram for peer effects



However, as the causal diagram shows, there are already several backdoor paths open. Moreover, conditioning on a_{ij} opens several backdoor paths since a_{ij} is a collider variable in several of those paths. Below, we enumerate all the backdoor paths as follows. (1) $Y_j^t \leftarrow Y_i^{t-1} \rightarrow Y_i^t \rightarrow Y_i^{t+1}$, (2) $Y_j^t \leftarrow \mathbf{u}_j \rightarrow a_{ij} \rightarrow Y_i^{t+1}$, (3) $Y_j^t \leftarrow \mathbf{u}_j \rightarrow a_{ij} \leftarrow \mathbf{u}_i \rightarrow Y_i^{t+1}$, (4) $Y_j^t \leftarrow \mathbf{u}_j \rightarrow a_{ij} \leftarrow \mathbf{u}_i \rightarrow Y_i^t \rightarrow$

 Y_i^{t+1} , (5) $Y_j^t \leftarrow \mathbf{u}_j \to a_{ij} \leftarrow \mathbf{u}_i \to Y_i^{t-l} \dots \to Y_i^{t-1} \to Y_i^t \to Y_i^{t+1}$. The first backdoor path can be closed by conditioning on Y_i^t , while the backdoor paths involving Z_i and Z_j can be closed by conditioning on those observed covariates. However, as we can see from the other open backdoor paths, we need to condition on \mathbf{u}_i and \mathbf{u}_j to close all of those backdoor paths. Therefore the linear structural equation model that we want to estimate is

$$\mathbf{y}_{t+1} = \alpha \mathbf{y}_t + \theta \mathcal{L}_{\mathbf{A}} \mathbf{y}_t + \mathbf{U}\beta + \mathbf{Z}_t \gamma + \mathcal{E}_{t+1}.$$

Here θ is our target peer influence parameter that we want to estimate consistently. We emphasize that **U** is a latent variable which is not observed. We theoretically show that we *can* estimate the structural peer effect parameter θ consistently with methodology described below under our modeling assumptions.

Next we consider a regression model for data observed in just one time point (T = 1). Here our goal is to model a response observed only once as a function of several covariates or predictors while controlling for latent homophily variables that may be correlated with the covariates whose effects we want to estimate. Accordingly, we propose the Embedding Network Regression (ENR) model as, $\mathbf{y} = \alpha \mathbf{1}_N + \mathbf{U}\beta + \mathbf{Z}\gamma + \mathcal{E}$, with \mathcal{E}_i being iid from $N(0, \sigma^2)$ distribution. This model has appeared in various forms previously in the literature Fosdick and Hoff (2015); He and Hoff (2019); Le and Li (2022). However, we are not aware of a study of the theoretical properties of estimators of this model. In addition, this model is a network analogue of the popular spatial confounding regression model used in spatial data analysis Guan et al. (2023).

2.5 Additive and multiplicative latent variables

Finally we consider an additive and multiplicative latent space model for the network data and propose to include both the additive and multiplicative latent variables in the network autoregressive model. Let \mathbf{q}_i, v_i be *K*-dimensional real vector and real scalars representing the multiplicative homophily and additive degree or activity parameter of node *i*, respectively. Let $\sigma : \mathbb{R} \to [0, 1]$ be a known link function that maps latent factors to the connection probability between two vertices. Then, latent space network can be generated via the density $f_{ij} \triangleq f(\cdot; \sigma(\mathbf{q}_i^{\mathsf{T}}\mathbf{q}_j + v_i + v_j))$ Li et al. (2023); Ma et al. (2020). Let $\mathbf{x}_i \triangleq (\mathbf{q}_i^{\mathsf{T}}, v_i)^{\mathsf{T}}$ denote a parameter vector containing all latent factors associated with node *i*. Collecting all latent factors for the network, we obtain $\mathbf{Q} \triangleq [\mathbf{q}_1, ..., \mathbf{q}_N]^{\mathsf{T}}$ and $\mathbf{v} \triangleq (v_1, ..., v_N)^{\mathsf{T}}$ hence $\mathbf{X} \triangleq [\mathbf{Q} | \mathbf{v}] \in \mathbb{R}^{N \times (K+1)}$. Here, we treat \mathbf{X} as fixed parameters. If we let $\chi \triangleq \mathbf{Q}\mathbf{Q}^{\mathsf{T}} + \mathbf{v}\mathbf{1}_N^{\mathsf{T}} + \mathbf{1}_N\mathbf{v}^{\mathsf{T}} \in \mathbb{R}^{N \times N}$ denote a latent factor matrix, then we can define the network connectivity \mathbf{P} as $p_{ij} \triangleq \sigma(\chi_{ij})$. Then, the log-likelihood of χ becomes

$$l(\chi; \mathbf{A}) = \sum_{i=1}^{n} \sum_{j>i} \log f(a_{ij}; p_{ij}) = \sum_{i} \sum_{j>i} \log f(a_{ij}; \sigma(\chi_{ij})).$$

Equipped with this network model, we define additive and multiplicative effect network autoregressive model (AMNAR) as following. Assume that a hollow and undirected graph is generated by $a_{ij} \stackrel{ind}{\sim} f_{ij}$. Define global effects of latent positions and degree parameters on responses as $\beta_1 \in \mathbb{R}^K$ and $\beta_2 \in \mathbb{R}$. Then, our measurements y_{it} are now assumed to be generated via $y_{i,t+1} = \alpha y_{it} + \theta \sum_{j \neq i} \ell_{ij} y_{jt} + (r\mathbf{q}_i)^{\mathsf{T}} \beta_1 + (rv_i) \beta_2 + \mathbf{z}_{it}^{\mathsf{T}} \gamma + \epsilon_{it}$, for i = 1, ..., Nand $t = 0, ..., T - 1(N, T \in \mathbb{N})$. The multiplier $r = r_{N,T} > 0$ controls the growth rate of \mathbf{X} to ensure the consistent estimation of considered parameters of AMNAR. In this regard, we let $r \triangleq \frac{1}{N^s \sqrt{T}}$ for a fixed constant $s \in (0, 1/2)$. The above model can be expressed with matrices and vectors as (with $\beta \triangleq (\beta_1^{\mathsf{T}}, \beta_2)^{\mathsf{T}}$)

$$\mathbf{y}_{t+1} = \alpha \mathbf{y}_t + \theta \mathcal{L}_{\mathbf{A}} \mathbf{y}_t + r \mathbf{X} \beta + \mathbf{Z}_t \gamma + \mathcal{E}_{t+1}.$$
(4)

Our proposals for ENAR and AMNAR models are related to but different from the Community NAR model of Chen et al. (2023). Similar to CNAR our frameworks use latent variables which are part of network formation, but the use of those factors are quite distinct. As Chen et al. (2023) noted, the CNAR outcome model is different from the peer effect NAR model since it contains the term UBU^T in place of the observed network $\mathcal{L}_{\mathbf{A}}$, where *B* is a matrix of unknown parameters. Our framework allows for the interpretation of peer influence given the observed network, and consistent estimation of peer effects is an important goal for us. In addition, our framework allows us to include more general latent effects than multiplicative factors.

3 Estimation

3.1 Strict stationarity

We start describing our estimation methodology with a discussion on the stationary distribution of \mathbf{y}_t for ENAR. Given our aim to establish its asymptotic distribution under both finite and diverging T and growing network size N, we first derive a stationary solution for \mathbf{y}_t . We denote $\mathbb{P}^* \triangleq \mathbb{P}(\cdot|\mathbf{A})$, $\mathbb{E}^* \triangleq \mathbb{E}(\cdot|\mathbf{A})$, and $\operatorname{Cov}^* \triangleq \operatorname{Cov}(\cdot, \cdot|\mathbf{A})$, as the conditional probability, expectation, and covariance respectively conditioning on \mathbf{A} . Using notations similar to Zhu et al. (2017), define $\mathbf{G} \triangleq \alpha \mathbf{I}_N + \theta \mathcal{L}_{\mathbf{A}}$ and $\tilde{\mathcal{E}}_{t+1} \triangleq \mathbf{Z}_t \gamma + \mathcal{E}_{t+1}$. Then, we can rewrite the ENAR model equivalently as

$$\mathbf{y}_{t+1} = \mathbf{U}\boldsymbol{\beta} + \mathbf{G}\mathbf{y}_t + \tilde{\mathcal{E}}_{t+1}.$$
 (5)

When N is fixed, the following results hold.

Theorem 3.1. If $|\alpha| + |\theta| < 1$, then there is a unique strictly stationary solution to the ENAR model 5 with a finite first moment, and the solution is given by,

$$\mathbf{y}_t = (\mathbf{I}_N - \mathbf{G})^{-1} \mathbf{U}\beta + \sum_{j=0}^{\infty} \mathbf{G}^j \tilde{\mathcal{E}}_{t-j}.$$
 (6)

Lemma 3.1. Define $\Gamma(h) \triangleq \operatorname{Cov}^*(\mathbf{y}_t, \mathbf{y}_{t-h})$ for all t. Upon the conditions in Theorem 3.1 and conditional on \mathbf{A} , 6 follows a normal distribution with the mean $\varphi \triangleq (\mathbf{I}_N - \mathbf{G})^{-1} \mathbf{U}\beta$ and $\operatorname{vec} \Gamma(0) = (\mathbf{I}_{N^2} - \mathbf{G} \otimes \mathbf{G})^{-1} \operatorname{vec} \{(\sigma^2 + \gamma^{\mathsf{T}} \Sigma_z \gamma) \mathbf{I}_N\}$ and $\Gamma(h) = \begin{cases} \mathbf{G}^h \Gamma(0) & , h > 0 \\ \Gamma(0)(\mathbf{G}^{\mathsf{T}})^{-h} & , h < 0. \end{cases}$ The proof of Theorem 3.1, along with all other theorems and lemmas, is contained in the Appendix. These results closely resemble the stationarity results presented in Zhu et al. (2017) for the model without the latent effects. As evident from the form of stationary mean, the location of \mathbf{y}_t is dependent on both the peer influence and the latent positions of the network. To simplify expressions, we will denote $\Gamma(0)$ by Γ henceforth. Next we note that \mathbf{y}_t is also asymptotically stationary in the sense of Definition 2 in the Appendix when the network size N grows. The following theorem is also proved in Appendix.

Theorem 3.2. Upon the conditions in Theorem 3.1 with $N \to \infty$, 6 is a unique strictly stationary solution with a finite first moment. i.e., $\max_{1 \le i < \infty} \mathbb{E} |y_{it}| < \infty$.

3.2 Least Squares Estimation of Parameters

One challenge in estimating the parameters of the ENAR model is that the latent vectors **U** are unobservable. To address this, we can utilize the asymptotic properties governing the differences between population spectra and sample spectra. In the literature, the random adjacency matrix is widely recognized as an additive perturbation of **P** by stochastically controllable noise. Specifically, if **A** follows $\text{RDPG}(\rho_N, \mathbf{X})$, then we have $\|\mathbf{A} - \mathbf{P}\| = O(\sqrt{N\rho_N})$ whp (Athreya et al., 2017; Xie and Xu, 2023). Moreover, the low-dimensional subspace represented by the spectra of **A** and **P** are close with high probability up to an orthogonal rotation as well. The following is the straightforward consequence of applying Davis-Kahan theorem to an undirected random network (Bhatia, 2013; Lei and Rinaldo, 2015; Chen et al., 2023):

Proposition 3.1. Let $\mathbf{A} \sim \text{RDPG}(\rho_N, \mathbf{X})$ and assume that \mathbf{X} is of rank K and $N\rho_N = \omega(\log N)$. Let $\mathbf{U}_{\mathbf{A}}$ be a matrix with orthogonal columns containing K leading eigenvectors of \mathbf{A} corresponding to its K leading eigenvalues and π_N denote the smallest non-zero

eigenvalue of **P**. Then, there exists $\mathbf{H} \in \mathcal{O}_K$ such that

$$\|\mathbf{U}_{\mathbf{A}} - \mathbf{U}\mathbf{H}\| = O\left(\frac{\sqrt{KN\rho_N}}{\pi_N}\right) whp.$$

We use the above result to estimate the additive latent factors as $\hat{\mathbf{U}} \triangleq \mathbf{U}_{\mathbf{A}}$ for fitting the ENAR model with these estimated latent factors.

From the equation of the model 3, we obtain the linear regression representation $y_{i,t+1} = \mathbf{w}_{i,t}^{\mathsf{T}} \mu + \epsilon_{i,t+1}$, where $\mathbf{w}_{it}^{\mathsf{T}} \triangleq (\mathbf{u}_{i}^{\mathsf{T}}, y_{it}, \ell_{i}^{\mathsf{T}} \mathbf{y}_{t}, \mathbf{z}_{it}^{\mathsf{T}})$ and our parameters of interest, $\mu \triangleq (\beta^{\mathsf{T}}, \alpha, \theta, \gamma^{\mathsf{T}})^{\mathsf{T}} \in \mathbb{R}^{K+p+2}$. Thus, the auto-regression of the networked measurements at time t + 1 can be written as

$$\mathbf{y}_{t+1} = \mathbf{W}_t \boldsymbol{\mu} + \mathcal{E}_{t+1}$$

where $\mathbf{W}_t = [\mathbf{w}_{1t}, ..., \mathbf{w}_{Nt}]^{\mathsf{T}}$. We can further collect the entire time series as $\mathbf{y} \triangleq (\mathbf{y}_1^{\mathsf{T}}, ..., \mathbf{y}_T^{\mathsf{T}})^{\mathsf{T}}$, $\mathbf{W} \triangleq [\mathbf{W}_0^{\mathsf{T}}, ..., \mathbf{W}_{T-1}^{\mathsf{T}}]^{\mathsf{T}}$, and $\mathcal{E} \triangleq (\mathcal{E}_1^{\mathsf{T}}, ..., \mathcal{E}_T^{\mathsf{T}})^{\mathsf{T}}$, thereby obtaining the representation $\mathbf{y} = \mathbf{W}\mu + \mathcal{E}$ in \mathbb{R}^{NT} . Note that \mathbf{W} contains the population latent positions \mathbf{U} of the observed network. Therefore, it is interpreted as the population design matrix for the ENAR model (Chen et al., 2023). Utilizing the estimated latent factors from $\hat{\mathbf{U}}$ from \mathbf{A} , we can obtain the approximated version $\hat{\mathbf{W}}_t = [\hat{\mathbf{w}}_{1t}, ..., \hat{\mathbf{w}}_{Nt}]^{\mathsf{T}}$ for $\hat{\mathbf{w}}_{it}^{\mathsf{T}} \triangleq (\hat{\mathbf{u}}_i^{\mathsf{T}}, y_{it}, \ell_i .^{\mathsf{T}} \mathbf{y}_t, \mathbf{z}_{it}^{\mathsf{T}})$ and $\hat{\mathbf{W}}$ accordingly.

This expression naturally motivates the least squares estimation of μ , and the asymptotic distribution of the estimator can be established through the classical martingale central limit theorem (Hall and Heyde, 2014; Zhu et al., 2017; Chen et al., 2023). Therefore, we target the least squares estimator

$$\hat{\mu} = \left(\hat{\mathbf{W}}^{\mathsf{T}}\hat{\mathbf{W}}\right)^{-1}\hat{\mathbf{W}}^{\mathsf{T}}\mathbf{y} = \left(\hat{\beta}^{\mathsf{T}}, \hat{\alpha}, \hat{\theta}, \hat{\gamma}^{\mathsf{T}}\right)^{\mathsf{T}}$$

and study its asymptotic properties as an estimator of μ .

3.3 Estimation with latent space model

In the same manner as ENAR, we establish the stationarity and the asymptotic properties of the AMNAR parameter estimators. The discussion of model stationarity is very similar to that of the ENAR model, and their proofs are contained in the Appendix. Therefore, we briefly state the stationarity results here. Note the model can be written as $\mathbf{y}_{t+1} =$ $r\mathbf{X}\beta + \mathbf{G}\mathbf{y}_t + \tilde{\mathcal{E}}_{t+1}$, for $\mathbf{G} \triangleq \alpha \mathbf{I}_N + \theta \mathcal{L}_{\mathbf{A}}$ and $\tilde{\mathcal{E}}_{t+1} \triangleq \mathbf{Z}_t \gamma + \mathcal{E}_{t+1}$ and we assume that \mathbf{A} is generated independently with the rest of random components across all t.

Theorem 3.3. If $|\alpha| + |\theta| < 1$, then there is a unique strictly stationary solution to 4 with a finite first moment:

$$\mathbf{y}_{t} = (\mathbf{I}_{N} - \mathbf{G})^{-1} r \mathbf{X} \beta + \sum_{j=0}^{\infty} \mathbf{G}^{j} \tilde{\mathcal{E}}_{t-j}$$
(7)

Lemma 3.2. Define $\Gamma(h) \triangleq \operatorname{Cov}^*(\mathbf{y}_t, \mathbf{y}_{t-h})$ for all t. Upon the conditions in Theorem 3.3 and conditional on \mathbf{A} , 7 follows a normal distribution with the mean $\psi \triangleq (\mathbf{I}_N - \mathbf{G})^{-1} r \mathbf{X} \beta$ and the auto-covariance function in Lemma 3.1.

Theorem 3.4. Upon the conditions in Theorem 3.3 with $N \to \infty$, 7 is a unique strictly stationary solution with a finite first moment. i.e., $\max_i \mathbb{E}|y_{it}| < \infty$.

Our main interest is to estimate the parameters of AMNAR consistently, but similarly to ENAR, the latent factors **X** are not observable. We tackle this issue by introducing the maximum likelihood estimation with Lagrange adjustment given in Li et al. (2023) where the authors found its asymptotic properties as well. Let $\hat{\mathbf{X}} \triangleq \arg \max_{\mathbf{X} \in \Xi} l(\chi; \mathbf{A})$ denote the maximum likelihood estimator over the constrained parameter space $\Xi \triangleq$ $\{\mathbf{X}; \mathbf{Q}^{\mathsf{T}} \mathbf{1}_N = \mathbf{o}_K, \mathbf{Q}^{\mathsf{T}} \mathbf{Q} \text{ is diagonal}, \|\mathbf{X}\|_{2,\infty} = O(1) \text{ as } N \to \infty\}$. The following proposition is analogous to Theorem 3.3 of Li et al. (2023) with slightly modified assumptions stated in the Appendix section . **Proposition 3.2.** Under assumptions on latent spade model given in section 9 in Appendix, we have $\|\hat{\mathbf{X}} - \mathbf{X}\|_F = O_{\mathbb{P}}(1)$.

Following the same estimation strategy as in ENAR model, we begin with the expression $y_{i,t+1} = \mathbf{m}_{it}{}^{\mathsf{T}}\mu + \epsilon_{i,t+1}$, where $\mathbf{m}_{it}{}^{\mathsf{T}} \triangleq (r\mathbf{x}_i{}^{\mathsf{T}}, y_{it}, \ell_i \cdot {}^{\mathsf{T}}\mathbf{y}_t, \mathbf{z}_{it}{}^{\mathsf{T}})$ and our parameters of interest, $\mu_m \triangleq (\beta^{\mathsf{T}}, \alpha, \theta, \gamma^{\mathsf{T}})^{\mathsf{T}} \in \mathbb{R}^{K+p+3}$. Thus, the autoregression of the networked measurements at time t + 1 can be written as

$$\mathbf{y}_{t+1} = \mathbf{M}_t \boldsymbol{\mu}_m + \mathcal{E}_{t+1} \in \mathbb{R}^{\Lambda}$$

and obtain the representation for the total observed data as $\mathbf{y} = \mathbf{M}\mu_m + \mathcal{E}$ in \mathbb{R}^{NT} . Note that \mathbf{M} contains the population latent positions \mathbf{X} of the observed network. Therefore, the population design matrix \mathbf{M} can be approximated by utilizing Proposition 3.2 as $\hat{\mathbf{M}}_t =$ $[\hat{\mathbf{m}}_{1t}, ..., \hat{\mathbf{m}}_{Nt}]^{\mathsf{T}}$ for $\hat{\mathbf{m}}_{it}^{\mathsf{T}} \triangleq (r\hat{\mathbf{x}}_i^{\mathsf{T}}, y_{it}, \ell_i.^{\mathsf{T}}\mathbf{y}_t, \mathbf{z}_{it}^{\mathsf{T}})$ and $\hat{\mathbf{M}}$ accordingly. Then, the least squares estimator for μ_m can be easily found as $\hat{\mu}_m = (\hat{\mathbf{M}}^{\mathsf{T}}\hat{\mathbf{M}})^{-1}\hat{\mathbf{M}}^{\mathsf{T}}\mathbf{y} = (\hat{\beta}^{\mathsf{T}}, \hat{\alpha}, \hat{\theta}, \hat{\gamma}^{\mathsf{T}})^{\mathsf{T}}$ which minimizes the residual sum of squares.

4 Large Sample Results

We next develop theory on consistency and asymptotic normality of the least squares estimator of the ENAR model under an asymptotic setup where we always assume $N \to \infty$, but consider both a finite T case as well as the case of $T \to \infty$. Unless otherwise stated, the results are for the ENAR model with the RDPG network model.

4.1 Growing T results

The first two theorems stated below show the asymptotic normality of the least squares estimator when both N and T grows under two cases, first when the number of latent dimensions K is fixed and then second when K also grows along with N and possibly also T. Define $\mu^H \triangleq (\beta^{\intercal} \mathbf{H}, \alpha, \theta, \gamma^{\intercal})^{\intercal}$, where $\mathbf{H} \in \mathcal{O}_K$ is an arbitrary matrix with orthonormal columns from Proposition 3.1, and recall from Proposition 3.1 that π_N denotes the smallest non-zero eigenvalue of \mathbf{P} .

Theorem 4.1. Assume that $|\alpha| + |\theta| < 1$ and $\pi_N = \omega \sqrt{TN\rho_N}$ for $\rho_N = \omega (\log N/N)$.

$$\Sigma_{w} \triangleq \begin{bmatrix} \mathbf{I}_{K} & \mathbf{O}_{K \times (p+2)} \\ & \Sigma_{-u} \end{bmatrix}, \quad \Sigma_{-u} \triangleq \begin{bmatrix} \tau_{2} & \tau_{23} & \mathbf{o}_{p}^{\mathsf{T}} \\ & \tau_{3} & \mathbf{o}_{p}^{\mathsf{T}} \\ & & \Sigma_{z} \end{bmatrix}$$

where $\tau_2 \triangleq \lim_{N \to \infty} \frac{1}{N} \mathbb{E} \operatorname{tr}(\Gamma), \tau_{23} \triangleq \lim_{N \to \infty} \frac{1}{N} \mathbb{E} \operatorname{tr}(\mathcal{L}_{\mathbf{A}}\Gamma), \text{ and } \tau_3 \triangleq \lim_{N \to \infty} \frac{1}{N} \mathbb{E} \operatorname{tr}(\mathcal{L}_{\mathbf{A}}^2 \Gamma).$ Then, for $\mathbf{D}_{NT} \triangleq \operatorname{diag}\left(\sqrt{T}\mathbf{I}_K, \sqrt{NT}\mathbf{I}_{p+2}\right)$, we have, as $N, T \to \infty$,

$$\mathbf{D}_{NT}\left(\hat{\mu}-\mu^{H}\right) \Rightarrow \mathcal{N}\left(\mathbf{o}_{K+p+2},\sigma^{2}\Sigma_{w}^{-1}\right).$$

The condition $|\alpha| + |\theta| < 1$ is required to ensure that \mathbf{y}_t has the stationary distribution discussed in section 3.1 as $N, T \to \infty$. The asymptotic precision matrix Σ_w has zero off-diagonal block hence the estimators $\hat{\beta}$ and $\hat{\mu}_{-\beta}$ are asymptotically independent. It is noteworthy that to consistently estimate the parameters, the population eigen gap π_N should grow faster than $\sqrt{TN\rho_N}$ with $N\rho_N = \omega (\log N)$ in Theorem 3.1. This rate can be compared to the assumption of $\pi_N = \omega(N\rho_N)$ made in Cape et al. (2019) for eigenvector deviation results in signal plus noise matrix models. The growth rate needed in Theorem 3.1 matches the rate in Cape et al. (2019) as long as $T = O(N\rho_N)$. If T grows faster then, the Theorem 3.1 requires a better concentration of $\hat{\mathbf{U}}$ in order for ENAR estimation to remain accurate. The convergence rate of \sqrt{T} for the latent position effects $\hat{\beta}$ is slower than \sqrt{NT} for the rest, which matches the rates obtained in Chen et al. (2023) for their CNAR model. We conjecture that the rate of convergence for $\hat{\beta}$ is dependent on the growth rate of the eigenvectors, which in turn depends on the growth rate of ρ_N and π_N . We explore this issue more in the context of AMNAR in a later section.

Next, we consider the case where K grows along with N and T as well.

Theorem 4.2. Assume that $|\alpha| + |\theta| < 1$ and $||\beta|| = O(1)$ as $K \to \infty$. Furthermore, assume that $\pi_N == \omega(\sqrt{TKN\rho_N})$, $N = \omega(K^2)$, and $T = \omega(\log K)$. For a positive integer m, suppose we have an $m \times (K + p + 2)$ matrix \mathbf{A}_K such that $||\mathbf{A}_k|| = O(1)$ as $K \to \infty$. If we define $\mathbf{V} \triangleq \lim_{K\to\infty} \mathbf{A}_K \Sigma_w^{-1} \mathbf{A}_K^{\mathsf{T}} \in \mathbb{R}^{m \times m}$, then we have, as $N, T, K \to \infty$,

$$\mathbf{A}_{K}\mathbf{D}_{NT}\left(\hat{\mu}-\mu^{H}\right) \Rightarrow \mathcal{N}\left(\mathbf{o}_{m},\sigma^{2}\mathbf{V}\right).$$

Compared to Theorem 4.1, when the dimension of the latent space K also grows, an additional \sqrt{K} is required for the growth rate of π_N . The condition on the population eigen gap π_N is that $\pi_N = \omega(\sqrt{TKN\rho_N})$ which improves upon the rate $\pi_N = \omega(K\sqrt{TN\rho_N})$ presented in the context of CNAR model in Chen et al. (2023) by a factor of \sqrt{K} which is a meaningful difference when K is large. Further, to consistently estimate the parameters, we only require $N = \omega(K^2)$ and $T = \omega(\log K)$. This can be compared to the required rates of $N = \omega(K^4)$ and $T = \omega(K^2 \log K)$ in Chen et al. (2023). Therefore consistent estimation in the CNAR model Chen et al. (2023) requires $O(K^2)$ times more sample size both in terms of N and T. Therefore, this represents a substantial relaxation of conditions. These reductions are largely due to ENAR model having **U** entering the model in an additive form as opposed to a multiplicative form in the CNAR model, and consequently, requiring us to estimate fewer parameters attached to latent factors.

4.2 Finite T results

Now, we prove asymptotic results for the model in the finite T case which is appropriate for the problem of causal peer influence estimation. This finite T case was not studied in earlier works of Zhu et al. (2017); Chen et al. (2023). Note that we no longer require $|\alpha| + |\theta| < 1$ in the finite *T* case. Distinctively from the Theorem 4.1 and 4.2, the consistency of $\hat{\mu}$ given finite *T* is different as the estimation error for the latent position effects, $\hat{\beta} - \mathbf{H}^{\mathsf{T}}\beta$, will only be bounded in probability. However, our result still shows the peer influence parameter θ along with other parameters, namely, α and γ , which we are typically interested in inferring are all \sqrt{N} consistent.

Theorem 4.3. Assume that $\pi_N = \omega(\sqrt{N\rho_N})$. Partition $\mu = (\beta^{\intercal}, \mu_{-\beta}^{\intercal})^{\intercal}$ and $\hat{\mu}$ accordingly as well. Then, we have $\hat{\beta} - \mathbf{H}^{\intercal}\beta = O_{\mathbb{P}}(1)$ and

$$\sqrt{N}\left(\hat{\mu}_{-\beta} - \mu_{-\beta}^{H}\right) \Rightarrow \mathcal{N}\left(\mathbf{o}_{p+2}, \frac{\sigma^{2}}{T}\Sigma_{-u}^{-1}\right), \quad as \ N \to \infty.$$
(8)

The result for growing K is as follows.

Theorem 4.4. In Theorem 4.3, further assume that $\|\beta\| = O(1)$ as $K \to \infty$, $\pi_N = \omega(\sqrt{KN\rho_N})$, and $N = \omega(K^2)$. For a positive integer m, suppose we have an $m \times (K + p + 2)$ matrix \mathbf{A}_K such that $\|\mathbf{A}_k\| = O(1)$ as $K \to \infty$. Then, $\mathbf{A}_K \left(\hat{\beta} - \mathbf{H}^{\mathsf{T}}\beta\right) = O_{\mathbb{P}}(1)$ and we have $\sqrt{N} \left(\hat{\mu}_{-\beta} - \mu_{-\beta}^H\right) \Rightarrow \mathcal{N} \left(\mathbf{o}_{p+2}, \frac{\sigma^2}{T} \Sigma_{-u}^{-1}\right)$ as $N, K \to \infty$.

As we limit T to be finite, π_N is allowed to grow at rates that are less restrictive compared to the case where T diverges. However, we still require the number of dimensions to grow at the same rate with the network size N as described in Theorem 4.2.

Our next two results are related to the Embedding Network Regression (ENR) model. As aforementioned, ENR is a special case of ENAR with $T = 1 < \infty$. Specifically, it can be derived from ENAR with finite T: let $\mathbf{y}_0 \triangleq \mathbf{1}_N$, $\mathbf{Z} \triangleq \mathbf{Z}_0$, and assume that there is no peer influence effect, i.e., $\theta = 0$.

Without loss of generality, we may omit the grand mean effect, α . Therefore, we have reduced data model as $\mathbf{w}_i^{R^{\intercal}} \triangleq (\mathbf{u}_i^{\intercal}, \mathbf{z}_i^{\intercal})$ and $\mu^R \triangleq (\beta^{\intercal}, \gamma^{\intercal})^{\intercal} \in \mathbb{R}^{K+p}$ hence giving the representation $\mathbf{y}_1 = \mathbf{W}^R \boldsymbol{\mu}^R + \mathcal{E}_1$. With a usual least square estimator $\hat{\boldsymbol{\mu}}^R \triangleq (\mathbf{W}^{R^{\mathsf{T}}} \mathbf{W}^R)^{-1} \mathbf{W}^{R^{\mathsf{T}}} \mathbf{y}_1$, the following two results are corollaries that follow from Theorems 4.3 and 4.4 respectively.

Corollary 4.1. (Fixed K.) Assume that $\pi_N = \omega(\sqrt{N\rho_N})$. Then, we have $\hat{\beta} - \mathbf{H}^{\mathsf{T}}\beta = O_{\mathbb{P}}(1)$ and $\sqrt{N}(\hat{\gamma} - \gamma) \Rightarrow \mathcal{N}(\mathbf{o}_p, \sigma^2 \Sigma_z^{-1})$, as $N \to \infty$.

Corollary 4.2. (Growing K.) Further assume that $\|\beta\| = O(1)$ as $K \to \infty$, $\pi_N = \omega(\sqrt{KN\rho_N})$, and $N = \omega(K^2)$. For a positive integer m, suppose we have an $m \times (K + p + 2)$ matrix \mathbf{A}_K such that $\|\mathbf{A}_k\| = O(1)$ as $K \to \infty$. Then, we have $\mathbf{A}_K \left(\hat{\beta} - \mathbf{H}^{\mathsf{T}}\beta\right) = O_{\mathbb{P}}(1)$ and $\sqrt{N}(\hat{\gamma} - \gamma) \Rightarrow \mathcal{N}(\mathbf{o}_p, \sigma^2 \Sigma_z^{-1})$ as $N, K \to \infty$.

4.3 Results for additive and multiplicative latent effects model

Before stating asymptotic properties of $\hat{\mu}_m$, we first define its asymptotic precision matrix as $\Omega_x \triangleq \begin{bmatrix} \Omega_q & \mathbf{q_v} \\ \nu \end{bmatrix}$, $\Omega_m \triangleq \begin{bmatrix} \Omega_x & \mathbf{O}_{(K+1) \times (p+2)} \\ \Sigma_{-u} \end{bmatrix}$. The values in the limiting precision matrix are defined as $\mathbf{q_v} \triangleq \lim_{N \to \infty} \frac{\mathbf{Q^{\intercal v}}}{N}$ and $\nu \triangleq \lim_{N \to \infty} \frac{\|\mathbf{v}\|^2}{N}$ for Ω_x . Then, the following result holds for $\hat{\mu}_m$.

Theorem 4.5. Assume that $|\alpha| + |\theta| < 1$ and the assumptions of Proposition 3.2. Then, for $\tilde{\mathbf{D}}_{NT} \triangleq \operatorname{diag}\left(N^{1/2-s}\mathbf{I}_{K+1}, \sqrt{NT}\mathbf{I}_{p+2}\right)$ and as $N, T \to \infty$, we have

$$\tilde{\mathbf{D}}_{NT}(\hat{\mu}_m - \mu_m) \Rightarrow \mathcal{N}\left(\mathbf{o}_{K+p+3}, \sigma^2 {\Omega_m}^{-1}\right)$$

As Theorem 4.5 implies, we have \sqrt{NT} -consistency for α and θ under AMNAR model as well, and $\hat{\beta}$ converges at the rate of $N^{1/2-s}$ for AMNAR, where, $s \in (0, 1/2)$, is controlled by the multiplier r. In ENAR model, by the definition of RDPG, we have $\|\mathbf{U}_{\mathbf{P}}\|_{2,\infty} \leq \frac{\sqrt{N}\rho_N}{\pi_N}$ for all N. Recall that we assumed that $N\rho_N = \omega (\log N)$ and $\pi_N = \omega(\sqrt{TN\rho_N})$ in Theorem 4.1. Therefore, we have $\|\mathbf{U}_{\mathbf{P}}\|_{2,\infty} = o\left(\sqrt{\rho_N/T}\right)$. So if $\rho_N = \Theta(N^{\epsilon-1})$ for $\epsilon \in (0, 1)$, for example, we have $\|\mathbf{U}_{\mathbf{P}}\|_{2,\infty} = o(N^{(\epsilon-1)/2}T^{-1/2})$ when $\min(N,T) \to \infty$. This is the motivation for the multiplier r in the AMNAR model since $\|r\mathbf{X}\|_{2,\infty} = \Theta(N^{-s}T^{-1/2})$. Next, the asymptotic results for the model under finite time is given below.

Theorem 4.6. Assume the settings for Theorem 4.5 with finite T. Then define $\mathbf{\tilde{D}}_N \triangleq \operatorname{diag}\left(\frac{N^{1/2-s}}{\sqrt{T}}\mathbf{I}_{K+1}, \sqrt{N}\mathbf{I}_{p+2}\right)$. As $N \to \infty$, we have

$$\tilde{\mathbf{D}}_N(\hat{\mu}_m - \mu_m) \Rightarrow \mathcal{N}\left(\mathbf{o}_{K+p+3}, \frac{\sigma^2}{T} \Omega_m^{-1}\right).$$

5 Simulation

In this section, we use Monte Carlo simulations to illustrate the finite sample performance of the ENAR model estimator and compare it with NAR and CNAR. We examine the sensitivity of the considered models under model misspecification in terms of estimations of model parameters and one-step-ahead prediction of \mathbf{y}_{T+1} . In this regard, we consider the scenarios where $\{\mathbf{y}_0, ..., \mathbf{y}_T\}$ and \mathbf{y}_{T+1} follow each of ENAR, CNAR, and NAR. In the case where we generate \mathbf{y}_t with CNAR (Chen et al., 2023), we assumed that there is no latent factor structure in the model noise.

We consider the DCSBM and DCMMSBM for the population distributions of **A**, and generate the networks using **fastRG** package in R Rohe et al. (2018). First, we used the matrix $2q\mathbf{I}_K + q\mathbf{1}_K\mathbf{1$ with parameter vector $(1, \ldots, 1)$. For DCSBM, the block memberships were sampled from a categorical distribution with equal probabilities.



Figure 2: Boxplot of estimates of θ (left) and α (right) from ENAR and NAR model with increasing N when data is generated from ENAR model with DCMMSBM. The rows corresponds to K = 3 and K = 12, and the columns corresponds to T = 40, 160, 320.

For model parametrization, we set the parameters β^{enar} associated with the latent effects as $(1, -1/2, ..., (-1)^{K-1}/K)^{\mathsf{T}} \in \mathbb{R}^{K}$. For the latent peer effect measured by the community structure of CNAR, we used $\mathcal{B}_1 \triangleq \frac{1}{10} \operatorname{diag}(\beta^{\text{enar}})$ where the definition of \mathcal{B}_1 comes from Chen et al. (2023). For ENAR, and NAR we set $\alpha = \theta = \frac{1}{5}$, and set the covariate effects as $\gamma = (\frac{1}{3}, -\frac{1}{6}, 0)^{\mathsf{T}}$ for all of considered models.

In order for comparison of estimation performance, we computed relative root mean squarred errors (RMSE) as $\frac{\|\mathbf{B}-\hat{\mathbf{B}}\|}{\|\mathbf{B}\|}$ where \mathbf{B} and $\hat{\mathbf{B}}$ are arbitrary matrices. Also, we report one-step prediction errors as $\frac{\|\mathbf{W}_{T}(\hat{\mu}-\mu)\|}{\|\mathbf{W}_{T}\mu\|}$ i.e., root mean squared prediction errors (RMSP), as the systematic noise incurred by \mathcal{E}_{T+1} is difficult to predict. Throughout simulations we generate the covariates \mathbf{z}_{it} from $\mathcal{N}(\mathbf{o}_{3}, \text{diag}(3, 2, 1))$ and \mathcal{E}_{t} from $\mathcal{N}(\mathbf{o}_{N}, 0.25 \mathbf{I}_{N})$. To track the model performance as its dimension grows, we take $N \in \{40, 80, 160, 320\}$ and $K \in \{3, 12\}$. We also consider finite T case where T = 2 and growing T case where



Figure 3: Boxplot of prediction error from ENAR, CNAR, and NAR model when data is generated from ENAR model with DCMMSBM and DCSBM respectively.

 $T \in \{40, 160, 320\}$. In all cases, we run 200 replications.

5.1 Generating data from ENAR model

The RMSE boxplots of peer influence and momentum effects are shown in Figures 2 for data generated from ENAR model with DCMMSBM. Each column and row of the facet grid corresponds to a different value of T and K, respectively, while the figures within the grids are with increasing N. From Figure 2, we observe that when the true model is ENAR, the RMSE of its estimates of α and θ consistency decreases as N and T grows, which is the expected phenomenon from our asymptotic theories.

For estimates from NAR model, while the RMSE for α parameter still decreases with increasing N, the RMSE for estimation of θ continues to remain high even when N and T increases. This is due to the fact that since the NAR fit omits the latent variable effects, it incurs irreducible bias in parameter estimation,

The predictions of ENAR, as shown in Figure 3 for graphs generated from DCMMSBM and DCSBM also improve with both N and T. This figure shows that if ENAR is the



Figure 4: Model misspecification: Boxplot of prediction error from CNAR, ENAR, and NAR model when data is generated from CNAR and NAR models respectively.

true data-generating model, then omitting the latent variables from the fitted model (i.e., fitting the NAR model) not only leads to inaccurate parameter estimates but also to higher prediction error.

5.2 Comparison Under Model misspecification

Next, we generate \mathbf{y}_t from NAR and CNAR while the underlying networks were generated from the DCMMSBM model. This corresponds to model misspecification for the ENAR model and we can compare the accuracy of prediction as well as parameter estimating in this setting. In Figure 4 (left), we see a very good performance of both ENAR and NAR even when the data is generated from the CNAR model. This is especially true for smaller values of N and T (e.g., N = T = 50). When K is increased to 12, the performance of CNAR is worse than NAR and ENAR even when the data is generated from CNAR. This is because with increasing K the performance of CNAR model estimators become worse.

When we assumed the true model NAR, as expected, the predictive ability of NAR was the best overall for all settings of N, T, K (Figure 4 (right)). However, in each case the



Figure 5: Boxplot of estimates of θ and α from ENAR, and NAR model when data is generated from ENAR and NAR model with DCMMSBM.

ENAR model came close in terms of predictive ability while the CNAR model produced large errors, especially when K was larger and N and T were smaller.

Finally, we evaluate the estimates of the θ and α parameters under model misspeciciation when the data is generated from the NAR model. Then, we observe that the estimation of the peer influence effect becomes biased for ENAR especially when T and N are larger (Figure A1 in Appendix). However, their estimation bias is significantly smaller than that of using NAR when data is generated from ENAR as shown in Figure 2. Moreover, ENAR model was able to consistently estimate α at comparable rates to NAR, showing robustness under model misspecification. In contrast, the NAR model produced large errors even for estimating α when ENAR was the true data generating model in Figure 2.

5.3 Finite T case

Next we investigate the performance of ENAR and NAR estimators in terms of accuracy of parameter estimation in the fixed T case. We set T = 2, and K = 3 and increased N. In Figure 5, it is clear that ENAR is able to consistently estimate α and θ even under model misspecification of generating data from NAR. The estimation error for estimating both α and θ from ENAR are comparable to NAR and decreases with increasing N. In contrast, when the data is generated from ENAR, we see that the estimate of θ from NAR is biased and continues to show high error even when N increases, while the estimation error decreases for ENAR.

Combined with our previous observations in growing N and T cases, ENAR shows robust estimation performance under various model misspecifications, while achieving better prediction and parameter estimation performance when the data is generated from ENAR in both growing and fixed model dimension cases.

6 Real Data Example

In this section, we will analyze two datasets. The first one is a finite-time dataset where the primary goal is to infer causal peer effects and effects of covariates, and the second one is a time series dataset where the goal is both accurate prediction and parameter estimation.

Knecht dutch students delinquency and alcohol data: The first dataset we analyze with this new method is the longitudinal Dutch students friendship network and delinquency study by Andrea Knecht Knecht (2008); Knecht et al. (2010); He and Hoff (2019). The dataset is taken from the R package "xergm.common". This longitudinal data consists of friendship networks along with responses relating to delinquency and alcohol consumption and some demographic covariates measured at four-time points on 26 students in one classroom. The measurements are taken 3 months apart during the first year of their secondary school. The demographic information includes sex, age, religion and ethnicity. There are two response variables that we are interested in - alcohol consumption measured at waves 2,3,4 and delinquency, which is defined as a rounded average over four types of minor delinquency (stealing, vandalism, graffiti, and fighting) and measured at all 4 waves. We construct an average friendship network by taking average of the friendship networks in waves 1 and 2. For alcohol consumption, we fit the ENAR and NAR models using 2 time periods, waves 1 and 2, and then predict the responses for Wave 3. For the delinquency response, we fit two models, one using data from waves 1 and 2, and predict the response in wave 3, and the other using data from waves 1, 2 and 3, and predict the response in wave 4. In both cases, we also fit a linear regression model with only the demographic covariates and do not include the lagged own response and lagged peer effects, which we call the OLS model.

Model	ENAR	NAR	OLS
Alcohol	1.9133	1.9726	2.5628
Delinquency	0.6712	0.6942	0.7375

Table 1: Comparison of Mean square prediction error (MSPE) for alcohol use in 4th wave and delinquency data from 2nd and 3rd wave.

The Table A1 in the Appendix shows the parameter estimates from the three models fitted to the alcohol consumption and delinquency data. We note that both the lagged own response and lagged peer response parameters are insignificant in both NAR and ENAR models. The coefficients corresponding to most predictors are also

insignificant in all 3 models. We compare the model fit in terms of out-of-sample prediction in Table 1 for the 3 models. We observe that ENAR performs the best in terms of mean square prediction error for both responses. The boxplots in Figure A2 shows the predicted values from the 3 models for various levels of actual response. We see that for both alcohol and delinquency, predictions from ENAR model is higher for higher values of the actual response indicating a good model fit to the data with strong predictive ability.

Wind speed time series data: Next we apply the ENAR model to a multivariate time series data containing wind speed measurements over 721 time periods at 102 weather sta-



Figure 6: (left) 1 step ahead mean square prediction error, (center) AIC and (right) BIC of model with increasing time for the Wind speed data.

tions in England and Wales. We take this dataset from the R package GNAR Knight et al. (2020). This is a data with large T and large N. We assess the accuracy of model fits in terms of the ability to accurately predict responses in 1 time period ahead. To compare the models over a range of time periods, we perform 1 step ahead predictions 200 times. In the *i* th prediction task, we fit the NAR and ENAR models to the time series until time 519 + i and predict the response at 520 + i th time. We continue to increase *i* and slide the window of training data until i = 200. For the ENAR model we estimate the dimension of the latent factors in the RDPG model is d = 7.



The figure 6 (left) shows the 1 step ahead prediction error from the ENAR and NAR model. Out of these 200 test time windows, ENAR gives smaller MSPE in 68.5% of the cases. Therefore Both the AIC and BIC

Figure 7: Boxplots of (left) Peer effect parameter and (right) windows, ENAR gives lagged effect parameter of 200 estimates of ENAR and NAR smaller MSPE in 68.5% model with training sample size 520 to 720. of the cases. Therefore we conclude ENAR outperforms NAR in this prediction task. Both the AIC and BIC

criteria in Figure 6 also point to superior model fit by the ENAR model over the NAR model. The parameter estimates of the peer effect and lagged effect (momentum effect) parameters along with their standard errors for one representative model with T = 520 is shown in Table A2 in the Appendix. We can see that both effects are statistically significant, with the momentum effect being roughly 10 times that of the peer effect. The parameter estimates for both of these effects are significantly different in the ENAR model from those in the NAR model, as is evident from the size of the differences of these estimates corresponding to the peer effect and lagged effect parameters from the 200 ENAR and NAR models. The plot shows a clear difference in the parameter estimates from the two models.

7 Conclusion

ENAR can successfully address two major statistical problems. It directly embeds the multiplicative homophily latent effects (plus the additive effect for AMNAR) in the time series network auto-regression problem, tackling both the consistent estimation of causal peer effects and predictive performance enhancement. We proved that the estimators of AMNAR and ENAR have asymptotic normality in both long-term and finite time. Our numerical study also illustrated that the estimation accuracy of the key peer effect parameter and predictive performance of the ENAR model is comparable to or better than that of other considered competitors. Therefore, this work is an extension and improvement of currently available approaches for both causal effect estimation and prediction tasks.

However, many topics remain uncovered in this paper and are worth studying in the future. First, many low-rank space models like ENAR often face the challenge of consistently estimating the dimension of the latent space from the observed data. We selected the dimension K corresponding to the values which minimized cross-validation error on the adjacency matrix. It is unclear whether one should utilize model selection techniques using the outcome model for estimating K instead. Second, further theoretical analysis of the prediction error and hypothesis testing will be beneficial for inferential questions. Additionally, as many real-world data exhibit evolving network structures over time, extending the AMNAR or ENAR models beyond static networks and embedding latent variables beneath the regime of dynamic networks would also be very interesting.

References

- An, W., Beauvile, R., and Rosche, B. (2022). Causal network analysis. Annual Review of Sociology, 48:23–41.
- Athreya, A., Fishkind, D. E., Tang, M., Priebe, C. E., Park, Y., Vogelstein, J. T., Levin, K., Lyzinski, V., and Qin, Y. (2017). Statistical inference on random dot product graphs: a survey. *The Journal of Machine Learning Research*, 18(1):8393–8484.
- Basu, S. and Michailidis, G. (2015). Regularized estimation in sparse high-dimensional time series models. The Annals of Statistics, 43(4):1535 – 1567.
- Bhatia, R. (2013). Matrix analysis, volume 169. Springer Science & Business Media.
- Cape, J., Tang, M., and Priebe, C. E. (2019). Signal-plus-noise matrix models: eigenvector deviations and fluctuations. *Biometrika*, 106(1):243–250.
- Chen, E. Y., Fan, J., and Zhu, X. (2023). Community network auto-regression for highdimensional time series. *Journal of Econometrics*, 235(2):1239–1256.

- Christakis, N. A. and Fowler, J. H. (2007). The spread of obesity in a large social network over 32 years. *New England journal of medicine*, 357(4):370–379.
- Christakis, N. A. and Fowler, J. H. (2013). Social contagion theory: examining dynamic social networks and human behavior. *Statistics in medicine*, 32(4):556–577.
- Fosdick, B. K. and Hoff, P. D. (2015). Testing and modeling dependencies between a network and nodal attributes. *Journal of the American Statistical Association*, 110(511):1047–1056.
- Goldsmith-Pinkham, P. and Imbens, G. W. (2013). Social networks and the identification of peer effects. *Journal of Business & Economic Statistics*, 31(3):253–264.
- Guan, Y., Page, G. L., Reich, B. J., Ventrucci, M., and Yang, S. (2023). Spectral adjustment for spatial confounding. *Biometrika*, 110(3):699–719.
- Hall, P. and Heyde, C. C. (2014). *Martingale limit theory and its application*. Academic press.
- He, Y. and Hoff, P. D. (2019). Multiplicative coevolution regression models for longitudinal networks and nodal attributes. *Social Networks*, 57:54–62.
- Hoff, P. (2021). Additive and multiplicative effects network models. Statistical Science.
- Hoff, P. D., Raftery, A. E., and Handcock, M. S. (2002). Latent space approaches to social network analysis. *Journal of the american Statistical association*, 97(460):1090–1098.
- Knecht, A., Snijders, T. A., Baerveldt, C., Steglich, C. E., and Raub, W. (2010). Friendship and delinquency: Selection and influence processes in early adolescence. *Social Development*, 19(3):494–514.

- Knecht, A. B. (2008). Friendship selection and friends' influence: Dynamics of networks and actor attributes in early adolescence.
- Knight, M., Leeming, K., Nason, G., and Nunes, M. (2020). Generalized network autoregressive processes and the gnar package. *Journal of Statistical Software*, 96(5):1–36.
- Le, C. M. and Li, T. (2022). Linear regression and its inference on noisy networklinked data. Journal of the Royal Statistical Society Series B: Statistical Methodology, 84(5):1851–1885.
- Lei, J. and Rinaldo, A. (2015). Consistency of spectral clustering in stochastic block models. The Annals of Statistics, 43(1).
- Li, J., Xu, G., and Zhu, J. (2023). Statistical inference on latent space models for network data. *arXiv preprint arXiv:2312.06605*.
- Ma, Z., Ma, Z., and Yuan, H. (2020). Universal latent space model fitting for large networks with edge covariates. *Journal of Machine Learning Research*, 21(4):1–67.
- McFowland III, E. and Shalizi, C. R. (2021). Estimating causal peer influence in homophilous social networks by inferring latent locations. *Journal of the American Statistical Association*, pages 1–12.
- Nath, S., Warren, K., and Paul, S. (2022). Identifying peer influence in therapeutic communities. arXiv preprint arXiv:2203.14223.
- O'Malley, A. J., Elwert, F., Rosenquist, J. N., Zaslavsky, A. M., and Christakis, N. A. (2014). Estimating peer effects in longitudinal dyadic data using instrumental variables. *Biometrics*, 70(3):506–515.
- Pearl, J. (2009). Causality. Cambridge university press.

- Rohe, K., Tao, J., Han, X., and Binkiewicz, N. (2018). A note on quickly sampling a sparse matrix with low rank expectation. *Journal of Machine Learning Research*, 19(77):1–13.
- Rubin-Delanchy, P., Cape, J., Tang, M., and Priebe, C. E. (2022). A statistical interpretation of spectral embedding: The generalised random dot product graph. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 84(4):1446–1473.
- Rudelson, M. and Vershynin, R. (2013). Hanson-Wright inequality and sub-gaussian concentration. *Electronic Communications in Probability*, 18(none):1 – 9.
- Shalizi, C. R. and Thomas, A. C. (2011). Homophily and contagion are generically confounded in observational social network studies. *Sociological methods & research*, 40(2):211–239.
- Tang, M. and Priebe, C. E. (2018). Limit theorems for eigenvectors of the normalized laplacian for random graphs. *The Annals of Statistics*, 46(5):2360–2415.
- VanderWeele, T. J. (2011). Sensitivity analysis for contagion effects in social networks. Sociological Methods & Research, 40(2):240–255.
- VanderWeele, T. J., Ogburn, E. L., and Tchetgen, E. J. T. (2012). Why and when" flawed" social network analyses still yield valid tests of no contagion. *Statistics, Politics* and Policy, 3(1).
- Vershynin, R. (2018). High-dimensional probability: An introduction with applications in data science, volume 47. Cambridge university press.
- Xie, F. and Xu, Y. (2023). Efficient estimation for random dot product graphs via a one-step procedure. *Journal of the American Statistical Association*, 118:651–664.

- Zhu, X. and Pan, R. (2020). Grouped network vector autoregression. *Statistica Sinica*, 30(3):1437–1462.
- Zhu, X., Pan, R., Li, G., Liu, Y., and Wang, H. (2017). Network vector autoregression. *The Annals of Statistics*, 45(3):1096 – 1123.
- Zhu, X., Wang, W., Wang, H., and H\u00e4rdle, W. K. (2019). Network quantile autoregression. Journal of econometrics, 212(1):345–358.
Appendix

8 Proofs for ENAR model in Sections 3.1 & 4

8.1 Technical Results

Here, we list some technical results that are useful in proving the asymptotic properties of our estimators for ENAR and AMNAR.

Proposition 8.1. If $X_n = O(Y_n)$ whp. and $Y_n = O(Z_n)$ whp., then $X_n = O(Z_n)$ whp. (or as.) If $X_n = O(Y_n)$ whp., then we have $\mathbb{P}(\{|X_n| \leq C |Y_n|\} ev.) = 1$ by Borel-Cantelli lemma.

By the model assumptions, our responses of interest, \mathbf{y}_t , will have stationary subgaussian distributions. To show the concentration of our estimators, we employ the theories of sub-Gaussian concentration as discussed in Vershynin (2018). However, since we assumed that both the observed graph \mathbf{A} and covariates \mathbf{z}_{it} are random, it necessitates investigating the asymptotic behaviors of the inner products of multiple random components. Therefore, we cannot directly apply well-known sub-Gaussian concentration results like Bernstein's inequality or Hoeffding's inequality.

For example, in the case of the Hanson-Wright inequality (Rudelson and Vershynin, 2013), we do not know how the tail behavior of concentrations will differ when studying the quadratic forms associated with random matrices. Under our assumptions on the random graphs and random predictors, we achieve stochastic boundedness for these elements. This fact can provide concentration results analogous to known inequalities such as Hoeffding's inequality and the Hanson-Wright inequality as follows.

Lemma 8.1 (Hoeffding's Inequality). Let $\mathbf{y} \in \mathbb{R}^n$ be a sub-gaussian random vector such that $\max_i ||Y_i||_{\psi_2} < \infty$ with independent components and zero mean. Let $\mathbf{x} \in \mathbb{R}^n$ be a random vector such that $||\mathbf{x}|| = O(g)$ whp. for some g > 0. Then, there exist some $c > 0, n_0 > 0$ such that

$$\mathbb{P}(|\mathbf{x}^{\mathsf{T}}\mathbf{y}| \ge \upsilon) \le 2\exp\left(-\frac{c\upsilon^2}{g^2}\right) + \frac{1}{n}$$

for all $n > n_0$ and v > 0.

Proof. Note that by general Hoeffding's inequality, with probability one we have

$$\mathbb{P}\left(|\mathbf{x}^{\mathsf{T}}\mathbf{y}| \geq \upsilon | \mathbf{x}\right) \leq 2 \exp\left(-\frac{c\upsilon^2}{\|\mathbf{x}\|^2}\right).$$

Denote the LHS and RHS by $f_n(\mathbf{x})$ and $h_n(\mathbf{x})$, respectively. By assumption, there exist contants $M, n_0 > 0$ such that $\mathbb{P}(||\mathbf{x}|| > Mg) < 1/n$ for all $n > n_0$. Then, for a sequence of events $A_n \triangleq \{||\mathbf{x}|| \le Mg\}$ we have

$$f_n(\mathbf{x})\mathbb{I}_{A_n} \le h_n(\mathbf{x})\mathbb{I}_{A_n} \le 2\exp\left(-\frac{c'\upsilon^2}{g^2}\right)\mathbb{I}_{A_n}$$

for some constant c' > 0. Taking expectation, we get

$$\mathbb{E}\left[\mathbb{P}\left(A_{n} \cap \{|\mathbf{x}^{\mathsf{T}}\mathbf{y}| \geq \upsilon\}|\mathbf{x}\right)\right] = \mathbb{P}\left(A_{n} \cap \{|\mathbf{x}^{\mathsf{T}}\mathbf{y}| \geq \upsilon\}\right) \leq 2\exp\left(-\frac{c'\upsilon^{2}}{g^{2}}\right)\mathbb{P}(A_{n})$$

hence

$$\mathbb{P}\left(\left\{|\mathbf{x}^{\mathsf{T}}\mathbf{y}| \geq \upsilon\right\} \setminus A_{n}^{c}\right) \leq 2\exp\left(-\frac{c'\upsilon^{2}}{g^{2}}\right)$$

$$\therefore \mathbb{P}\left(|\mathbf{x}^{\mathsf{T}}\mathbf{y} \geq \upsilon|\right) \leq 2\exp\left(-\frac{c'\upsilon^{2}}{g^{2}}\right) + 1 - \mathbb{P}(A_{n})$$

For almost surely bounded case, we will have no $\frac{1}{n}$ in the uppper bound.

By the same logic, we have the following lemma as well.

Lemma 8.2. Suppose that $\mathbf{x} \in \mathbb{R}^n$ is a random vector such that $\|\mathbf{x}\| = O_{\mathbb{P}}(g)$ for some g > 0 in Lemma 8.1. Then, $\forall \epsilon, v > 0$ there exist some $c > 0, n_0 > 0$ such that

$$\mathbb{P}(|\mathbf{x}^{\mathsf{T}}\mathbf{y}| \ge \upsilon) \le 2\exp\left(-\frac{c\upsilon^2}{g^2}\right) + \epsilon$$

for all $n > n_0$.

The proof of Lemma 8.2 coincides with Lemma 8.1.

Lemma 8.3. (Hanson-Wright Inequality) Let $\mathbf{y} \in \mathbb{R}^n$ be a random vector with independent components and zero mean such that $\max_i ||Y_i||_{\psi_2} < \infty$. Let $\mathbf{S} \in \mathbb{R}^{n \times n}$ be a random matrix such that $||\mathbf{S}||_F^2 = O(h)$ whp. and $||\mathbf{S}|| = O(k)$ whp. for some h, k > 0. Then, there exists $n_0 > 0$ such that

$$\mathbb{P}\left(|\mathbf{y}^{\mathsf{T}}\mathbf{S}\mathbf{y} - \mathbb{E}(\mathbf{y}^{\mathsf{T}}\mathbf{S}\mathbf{y})| > \upsilon\right) \le 2\exp\left[-c\min\left(\frac{\upsilon^2}{h}, \frac{\upsilon}{k}\right)\right] + \frac{2}{n}$$

for all $n > n_0$ and v > 0. If $\|\mathbf{S}\|_F^2 = O(h)$ as. and $\|\mathbf{S}\| = O(k)$ as., then we have

$$\mathbb{P}\left(|\mathbf{y}^{\mathsf{T}}\mathbf{S}\mathbf{y} - \mathbb{E}(\mathbf{y}^{\mathsf{T}}\mathbf{S}\mathbf{y})| > \upsilon\right) \le 2\exp\left[-c\min\left(\frac{\upsilon^2}{h}, \frac{\upsilon}{k}\right)\right]$$

for all n > 0.

Proof. Let $M, n_0 > 0$ be constants such that $\mathbb{P}\left(\|\mathbf{S}\|_F^2 > Mh\right) < \frac{1}{n}$ and $\mathbb{P}\left(\|\mathbf{S}\| > Mk\right) < \frac{1}{n}$ for all $n > n_0$. By Hanson-Wright inequality (Rudelson and Vershynin, 2013), with probability one we have

$$\mathbb{P}\left(|\mathbf{y}^{\mathsf{T}}\mathbf{S}\mathbf{y} - \mathbb{E}(\mathbf{y}^{\mathsf{T}}\mathbf{S}\mathbf{y})| > \upsilon|\mathbf{S}\right) < 2\exp\left[-c\min\left(\frac{\upsilon^2}{\|\mathbf{S}\|_F^2}, \frac{\upsilon}{\|\mathbf{S}\|}\right)\right].$$

Denote the LHS and RHS by $f_n(\mathbf{S})$ and $g_n(\mathbf{S})$, respectively. Then we have

$$g_n(\mathbf{S}) \le 2 \exp\left[-c' \min\left(\frac{v^2}{h}, \frac{v}{k}\right)\right]$$

on $A_n \triangleq \left\{ \left\| \mathbf{S} \right\|_F^2 \le Mh \right\} \cap \left\{ \left\| \mathbf{S} \right\| \le Mk \right\}$. Since

$$\mathbb{E}\left[f_n(\mathbf{S})\mathbf{1}_{A_n}\right] = \mathbb{E}\left[\mathbb{P}\left(A_n \cap \{|\mathbf{y}^{\mathsf{T}}\mathbf{S}\mathbf{y} - \mathbb{E}(\mathbf{y}^{\mathsf{T}}\mathbf{S}\mathbf{y})| > \upsilon\} | \mathbf{S}\right)\right],$$

we can obtain

$$\mathbb{P}\left(\{|\mathbf{y}^{\mathsf{T}}\mathbf{S}\mathbf{y} - \mathbb{E}(\mathbf{y}^{\mathsf{T}}\mathbf{S}\mathbf{y})| > \upsilon\} \setminus A_n^{c}\right) < 2\exp\left[-c'\min\left(\frac{\upsilon^2}{h}, \frac{\upsilon}{k}\right)\right]\mathbb{P}(A_n),$$

implying

$$\mathbb{P}\left(|\mathbf{y}^{\mathsf{T}}\mathbf{S}\mathbf{y} - \mathbb{E}(\mathbf{y}^{\mathsf{T}}\mathbf{S}\mathbf{y})| > t\right) < 2\exp\left[-c'\min\left(\frac{t^2}{h}, \frac{t}{k}\right)\right] + 1 - \mathbb{P}(A_n).$$

Therefore, the conclusion follows. For almost surely bounded case, $\mathbb{P}(A_n) = 1$.

8.2 Stationarity

For showing stationarity, we adapt the proofs of Zhu et al. (2017) to the setup of ENAR.

Proof of Theorem 3.1. It is straightforward that the spectral radius of \mathbf{G} , denoted by g, satisfies

$$g \le |\alpha| + |\theta|\rho\left(\mathcal{L}_{\mathbf{A}}\right) < 1 \tag{9}$$

with probability one. Therefore, $\sum_{j=0}^{\infty} \mathbf{G}^{j} \tilde{\mathcal{E}}_{t-j}$ exists as., and \mathbf{y}_{t} in 6 is a strictly stationary process. It is straightforward that 6 satisfies 3. Next, assume that $\bar{\mathbf{y}}_{t}$ is another strictly stationary solution with $\mathbb{E}\|\bar{\mathbf{y}}_{t}\| < \infty$. Then,

$$\bar{\mathbf{y}}_t = \sum_{j=0}^{m-1} \mathbf{G}^j (\mathbf{U}\beta + \tilde{\mathcal{E}}_{t-j}) + \mathbf{G}^m \bar{\mathbf{y}}_{t-m}$$

for any positive integer m. Therefore,

$$\mathbb{E}^* \|\mathbf{y}_t - \bar{\mathbf{y}}_t\| = \mathbb{E}^* \left\| \sum_{j=m}^{\infty} \mathbf{G}^j (\mathbf{U}\beta + \tilde{\mathcal{E}}_{t-j}) - \mathbf{G}^m \bar{\mathbf{y}}_t \right\| \le Cg^m$$

for a constant C independent of t and m. Growing m to infinity, we get $\mathbb{E} \|\mathbf{y}_t - \bar{\mathbf{y}}_t\| = 0$ hence $\mathbf{y}_t = \bar{\mathbf{y}}_t$ almost surely.

Next, we prove Theorem 3.2 according to the following definition.

Definition 2. (Zhu et al., 2017) Let $\{\mathbf{y}_t \in \mathbb{R}^N\}$ be an N-dimensional with $N \to \infty$. Define $\mathcal{M} \triangleq \{\omega \in \mathbb{R}^\infty : \sum_{i=1}^\infty |w_i| < \infty\}$. For each $\omega \in \mathcal{M}$, let $\mathbf{w}_N = (w_1, ..., w_N)^{\mathsf{T}} \in \mathbb{R}^N$ be the truncated N-dimensional process. \mathbf{y}_t is said to be strictly stationary if $\forall \omega \in \mathcal{M}$

- 1. $\mathbf{y}_t^{\omega} = \lim_{N \to \infty} \mathbf{w}_N^{\mathsf{T}} \mathbf{y}_t$ exists almost surely.
- 2. \mathbf{y}_t^{ω} is strictly stationary.

Proof of Theorem 3.2. To prove the existence of a stationary solution, it suffices to show that 6 is strictly stationary according to above definition. Write $|\mathbf{A}|_e$ as a matrix of absolute elements of a matrix \mathbf{A} . Moreover, write $\mathbf{A} \preccurlyeq \mathbf{B}$ if \mathbf{B} is not less than \mathbf{A} elementwisely.

Recall that $\mathbf{y}_t = \sum_{j=0}^{m-1} \mathbf{G}^j (\mathbf{U}\beta + \tilde{\mathcal{E}}_{t-j}) + \mathbf{G}^m \mathbf{y}_{t-m}$ hence

$$\mathbf{y}_t = \lim_{m \to \infty} \mathbf{y}_t = \sum_{j=0}^{\infty} \mathbf{G}^j (\mathbf{U}\beta + \tilde{\mathcal{E}}_{t-j}).$$

For the columns of \mathbf{U} , say $U_1, ..., U_K$, We have $\mathbf{U}\beta = \sum_{j=1}^K U_j\beta_j$ and its sup vector norm is bounded as $\|\mathbf{U}\beta\|_{\infty} \leq \sum_{j=1}^K |\beta_j| \|U_j\|_{\infty} \leq \|\beta\|_1$ hence $|\mathbf{U}\beta|_e \preccurlyeq \|\beta\|_1 \mathbf{1}_N$. So, we have $\mathbb{E} \left| \mathbf{U}\beta + \tilde{\mathcal{E}}_{t-j} \right|_e \preccurlyeq C \cdot \mathbf{1}_N$ for $C = \|\beta\|_1 + \mathbb{E} |\mathbf{z}_1^{\mathsf{T}}\gamma| + \mathbb{E} |\epsilon_{11}|$. Since $\mathcal{L}_{\mathbf{A}}$ is normal and symmetric, its spectral decomposition can be given as $\mathcal{L}_{\mathbf{A}} = \mathbf{U}_{\mathcal{L}} \mathbf{S}_{\mathcal{L}} \mathbf{U}_{\mathcal{L}}^{\mathsf{T}}$ with an orthogonal eigenvectors $\mathbf{U}_{\mathcal{L}} \in \mathbb{R}^{N \times N}$ and $\mathbf{S}_{\mathcal{L}}$ containing corresponding eigenvalues of $\mathcal{L}_{\mathbf{A}}$. So, we have $\|\mathcal{L}_{\mathbf{A}}^{j}\|_{\infty} \leq$ $\|\mathbf{U}_{\mathcal{L}}\|_{\infty} \|\mathbf{U}_{\mathcal{L}}^{\mathsf{T}}\|_{\infty} \rho (\mathcal{L}_{\mathbf{A}})^j \leq C'$ for a constant C' > 0 independent of j. Therefore, $|\mathbf{G}|_e^j \mathbf{1}_N =$ $(|\theta|\mathcal{L}_{\mathbf{A}} + |\alpha|\mathbf{I}_N)^j \mathbf{1}_N \preccurlyeq C'(|\theta| + |\alpha|)^j \mathbf{1}_N$. Consequently,

$$\begin{split} \mathbb{E}^* \left| \mathbf{w}_N^{\mathsf{T}} \mathbf{y}_t \right| &\leq \left\| \mathbf{w}_N \right\|_1 \mathbb{E}^* \left\| \mathbf{y}_t \right\|_{\infty} \leq \sum_{i=1}^{\infty} \left| w_i \right| \sum_{j=0}^{\infty} \mathbb{E}^* \left\| \left| \mathbf{G} \right|_e^j \left| \mathbf{U}\beta + \tilde{\mathcal{E}}_{t-j} \right|_e \right\|_{\infty} \\ &\lesssim \sum_{i=1}^{\infty} \left| w_i \right| \sum_{j=0}^{\infty} (\left| \theta \right| + \left| \alpha \right|)^j \end{split}$$

implying that $\lim_{N\to\infty} \mathbf{w}_N^{\mathsf{T}} \mathbf{y}_t$ exists almost surely. Next, assume that $\bar{\mathbf{y}}_t$ is another strictly

stationary solution with a finite first moment. Then, $\mathbb{E} |\bar{\mathbf{y}}_t|_e \preceq \mathbf{1}_N$. We have

$$\mathbb{E}^* \left| \mathbf{w}_N^{\mathsf{T}} (\mathbf{y}_t - \bar{\mathbf{y}}_t) \right| = \mathbb{E}^* \left| \sum_{j=m}^{\infty} \mathbf{w}_N^{\mathsf{T}} \mathbf{G}^j (\mathbf{U}\beta + \tilde{\mathcal{E}}_{t-j}) - \mathbf{w}_N^{\mathsf{T}} \mathbf{G}^m \bar{\mathbf{y}}_{t-m} \right|$$

which is bounded above by the product of a constant and

$$\sum_{i=1}^{\infty} |w_i| \sum_{j=m}^{\infty} \left\{ (|\alpha| + |\theta|)^j + (|\alpha| + |\theta|)^m \right\}$$

for any $\omega \in \mathcal{M}$. Growing $m \to \infty$, we have $\mathbf{w}_N^{\mathsf{T}}(\mathbf{y}_t - \bar{\mathbf{y}}_t) = 0$ as. hence $\mathbf{y}_t = \bar{\mathbf{y}}_t$ as. \Box

8.3 Consistency

Here, we prove the asymptotic normality of $\hat{\mu}$ for both cases where K is fixed and growing. First, we clarify some notations here. Let $\Gamma_{\mathbf{y}} \triangleq \operatorname{Cov}^*(\mathbf{y})$. Write $\Phi \triangleq \mathbf{1}_T \otimes \varphi$ and $\tilde{\mathbf{y}} \triangleq \Gamma_{\mathbf{y}}^{-\frac{1}{2}}(\mathbf{y} - \Phi)$ so that $\mathbf{y} = \Gamma_{\mathbf{y}}^{1/2}\tilde{\mathbf{y}} + \Phi$. Note that the entries of $\tilde{\mathbf{y}}$ are independent by the property of multivariate normal distribution. Also, for neater expressions, let us denote $\frac{\sqrt{KN\rho_N}}{\pi_N}$ by κ .

By Proposition 2.2 and 2.3 of Basu and Michailidis (2015), with probability one, we have

$$\|\Gamma_{\mathbf{y}}\| \le \frac{\sigma^2}{m_{\min}(\mathbf{G})} \le \frac{\sigma^2 \|\mathbf{U}_{\mathbf{G}}\|^2 \|\mathbf{U}_{\mathbf{G}}^{-1}\|^2}{(1-g)^2}$$

where g is the spectral radius of **G** in 9, $\mathbf{U}_{\mathbf{G}}$ is an orthogonal matrix that contains eigenvectors of **G** and $m_{\min}(\mathbf{G}) \triangleq \min_{\{z \in \mathcal{C}; |z|=1\}} (\mathbf{I}_N - \mathbf{G}z)^* (\mathbf{I}_N - \mathbf{G}z)$. The last upper bound holds because **G** is diagonalizable. So, we have $\|\Gamma_{\mathbf{y}}\| = O(1)$ as.

Proof of Theorem 4.1. First, define $\tilde{\mathbf{U}} \triangleq \mathbf{U}\mathbf{H}$ and let $\mathbf{W}_t^H \triangleq \left[\tilde{\mathbf{U}} | \mathbf{y}_t, \mathcal{L}_{\mathbf{A}}\mathbf{y}_t | \mathbf{Z}_t\right]$, $\mu_w^H = \left(\frac{\beta^{\intercal}\mathbf{H}}{\sqrt{N}}, \alpha, \theta, \gamma^{\intercal}\right)^{\intercal}, \ \mathcal{W}_t \triangleq \left[\sqrt{N}\mathbf{U} | \mathbf{y}_t, \mathcal{L}_{\mathbf{A}}\mathbf{y}_t | \mathbf{Z}_t\right], \ \mathcal{W}_t^H \triangleq \left[\sqrt{N}\tilde{\mathbf{U}} | \mathbf{y}_t, \mathcal{L}_{\mathbf{A}}\mathbf{y}_t | \mathbf{Z}_t\right]$, and $\hat{\mathcal{W}}_t = \left[\sqrt{N}\hat{\mathbf{U}} | \mathbf{y}_t, \mathcal{L}_{\mathbf{A}}\mathbf{y}_t | \mathbf{Z}_t\right]$. Under this representation, we have $\hat{\mathcal{W}}_t\mathbf{D}_{NT} = \sqrt{NT}\hat{\mathbf{W}}_t$, $\sqrt{NT}\mathbf{D}_{NT}^{-1}\mu_w^H = \mu^H$, and $\mathbf{W}_t^H\mu^H = \mathcal{W}_t^H\mu_w^H$. Collect and bind them row-wise for t = 0, ..., T-1 to obtain $NT \times (K+p+2)$ matrices $\mathbf{W}^H, \mathcal{W}^H$, and define $\hat{\mathcal{W}}$ analogously. For $\hat{\Sigma}_w \triangleq \frac{1}{NT} \hat{\mathcal{W}}^{\intercal} \hat{\mathcal{W}}$ and $\mathcal{E}_{\mathcal{W}} \triangleq \frac{1}{NT} \hat{\mathcal{W}}^{\intercal} \mathcal{E}$, we have

$$\hat{\mu} = \left(\hat{\mathbf{W}}^{\mathsf{T}}\hat{\mathbf{W}}\right)^{-1}\hat{\mathbf{W}}^{\mathsf{T}}\left(\mathbf{W}^{H}\mu^{H} + \mathcal{E}\right)$$

$$= \left(\frac{\mathbf{D}_{NT}}{NT}\hat{\mathcal{W}}^{\mathsf{T}}\hat{\mathcal{W}}\mathbf{D}_{NT}\right)^{-1}\frac{\mathbf{D}_{NT}}{\sqrt{NT}}\hat{\mathcal{W}}^{\mathsf{T}}\left(\mathcal{W}^{H}\mu^{H}_{w} + \mathcal{E}\right)$$

$$= \left(\hat{\Sigma}_{w}\mathbf{D}_{NT}\right)^{-1}\frac{1}{\sqrt{NT}}\hat{\mathcal{W}}^{\mathsf{T}}\left\{\left(\mathcal{W}^{H} - \hat{\mathcal{W}}\right)\mu^{H}_{w} + \hat{\mathcal{W}}\mu^{H}_{w} + \mathcal{E}\right\}$$

$$= \mathbf{D}_{NT}^{-1}\hat{\Sigma}_{w}^{-1}\left\{\frac{1}{\sqrt{NT}}\hat{\mathcal{W}}^{\mathsf{T}}\left(\mathcal{W}^{H} - \hat{\mathcal{W}}\right)\mu^{H}_{w} + \sqrt{NT}\mathcal{E}_{w}\right\} + \sqrt{NT}\mathbf{D}_{NT}^{-1}\mu^{H}_{w}$$

$$\therefore \mathbf{D}_{NT}\left(\hat{\mu} - \mu^{H}\right) = \frac{\hat{\Sigma}_{w}^{-1}}{\sqrt{NT}}\hat{\mathcal{W}}^{\mathsf{T}}\left(\mathcal{W}^{H} - \hat{\mathcal{W}}\right)\mu^{H}_{w} + \sqrt{NT}\hat{\Sigma}_{w}^{-1}\mathcal{E}_{w}.$$

Therefore, we next show that the first term on the RHS is negligible and $\sqrt{NT}\hat{\Sigma}_w^{-1}\mathcal{E}_W$ is converging to a multivariate normal distribution. We start with claiming that $\hat{\Sigma}_w$ is converging to a matrix with finite entries as N and T tend to infinity. This will allow us to focus on the behaviors of $\frac{1}{\sqrt{NT}}\hat{W}^{\mathsf{T}}(\mathcal{W}^H - \hat{\mathcal{W}})\mu_w^H$ and $\sqrt{NT}\mathcal{E}_W$, and then apply Slutsky's Theorem.

Claim 1. $\hat{\Sigma}_w$ converges to Σ_w in probability, i.e., $\hat{\Sigma}_w \Rightarrow \Sigma_w$.

Proof. As consequences of Lemma 8.4, we have

$$\hat{\Sigma}_{w} = \frac{1}{NT} \sum_{t} \begin{bmatrix} N\mathbf{I}_{K} & \sqrt{N}\hat{\mathbf{U}}^{\mathsf{T}}\mathbf{y}_{t} & \sqrt{N}\hat{\mathbf{U}}^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}\mathbf{y}_{t} & \sqrt{N}\hat{\mathbf{U}}^{\mathsf{T}}\mathbf{Z}_{t} \\ & \mathbf{y}_{t}^{\mathsf{T}}\mathbf{y}_{t} & \mathbf{y}_{t}^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}\mathbf{y}_{t} & \mathbf{y}_{t}^{\mathsf{T}}\mathbf{Z}_{t} \\ & & \mathbf{y}_{t}^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}^{2}\mathbf{y}_{t} & \mathbf{y}_{t}^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}\mathbf{Z}_{t} \\ & & & \mathbf{Z}_{t}^{\mathsf{T}}\mathbf{Z}_{t} \end{bmatrix} \\ \rightarrow \lim_{N,T \to \infty} \begin{bmatrix} \mathbf{I}_{K} & \frac{1}{\sqrt{N}}\tilde{\mathbf{U}}^{\mathsf{T}}\varphi & \frac{1}{\sqrt{N}}\tilde{\mathbf{U}}^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}\varphi & \mathbf{O}_{K \times p} \\ & \frac{1}{N}\varphi^{\mathsf{T}}\varphi + \tau_{2} & \frac{1}{N}\varphi^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}\varphi + \tau_{23} & \mathbf{O}_{p}^{\mathsf{T}} \\ & & \frac{1}{N}\varphi^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}^{2}\varphi + \tau_{3} & \mathbf{O}_{p}^{\mathsf{T}} \\ & & & \Sigma_{z} \end{bmatrix}$$

for $\pi_N = \omega(\sqrt{TN\rho_N})$. Existence of τ_2, τ_{23} , and τ_3 come by dominated convergence theorem after noting that $\operatorname{tr}(\mathcal{L}_{\mathbf{A}}\Gamma) \leq ||\mathcal{L}_{\mathbf{A}}|| \operatorname{tr}(\Gamma) = O(N)$ as. and $\operatorname{tr}(\mathcal{L}_{\mathbf{A}}^2\Gamma) = O(N)$ as. Also from the assumptions and the asymptotic order of each term found in the proof of Lemma 8.4, we have

$$\lim_{N \to \infty} \frac{1}{\sqrt{N}} \mathbf{h}_1^{\mathsf{T}} \tilde{\mathbf{U}}^{\mathsf{T}} \varphi = 0, \frac{1}{\sqrt{N}} \mathbf{h}_1^{\mathsf{T}} \tilde{\mathbf{U}}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \varphi \Rightarrow 0, \lim_{N \to \infty} \frac{1}{N} \varphi^{\mathsf{T}} \varphi = 0,$$
$$\frac{1}{N} \varphi^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \varphi \Rightarrow 0, \frac{1}{N} \varphi^{\mathsf{T}} \mathcal{L}_{\mathbf{A}}^2 \varphi \Rightarrow 0$$

for all $\mathbf{h}_1 \in \mathbb{R}^K$ such that $\|\mathbf{h}_1\| \leq 1$.

Claim 2. $\frac{1}{\sqrt{NT}}\hat{\mathcal{W}}^{\mathsf{T}}(\mathcal{W}^H - \hat{\mathcal{W}})\mu_w^H = o_{\mathbb{P}}(1).$

Proof. Since $\hat{\mathcal{W}}$ is different from \mathcal{W}^H by $\sqrt{N}\hat{\mathbf{U}}$ only, we have

$$\frac{1}{\sqrt{NT}}\hat{\mathcal{W}}^{\mathsf{T}}\left(\mathcal{W}^{H}-\hat{\mathcal{W}}\right)\mu_{w}^{H} = \begin{bmatrix} \sqrt{T}\hat{\mathbf{U}}^{\mathsf{T}} \\ \frac{1}{\sqrt{NT}}\sum_{t}\mathbf{y}_{t}^{\mathsf{T}} \\ \frac{1}{\sqrt{NT}}\sum_{t}\mathbf{y}_{t}^{\mathsf{T}}\mathcal{L}_{\mathbf{A}} \\ \frac{1}{\sqrt{NT}}\sum_{t}\sum_{t}\mathbf{z}_{t}^{\mathsf{T}} \end{bmatrix} [\hat{\mathbf{U}}-\tilde{\mathbf{U}} \mid \mathbf{O}_{N\times(p+2)}]\mu^{H}$$

Then, by the statements 10.–13. of Lemma 8.4, we have the conclusion by Cramér–Wold.

Claim 3. $\sqrt{NT}\mathcal{E}_{\mathcal{W}} \Rightarrow \mathcal{N}(\mathbf{o}_{K+p+2}, \sigma^2 \Sigma_w).$

Proof. It is sufficient to show that for any $\eta \in \mathbb{R}^{K+p+2}$ such that $\|\eta\| \leq 1$, we have $\sqrt{NT}\eta^{\mathsf{T}}\mathcal{E}_{\mathcal{W}} \Rightarrow \mathcal{N}(0, \sigma^{2}\eta^{\mathsf{T}}\Sigma_{w}\eta)$. Denoting $\xi_{N,t+1} \triangleq (NT)^{-1/2} \eta^{\mathsf{T}} \hat{\mathcal{W}}_{t}^{\mathsf{T}} \mathcal{E}_{t+1}$ and $\mathcal{F}_{Nt} = \sigma(\mathbf{A}, \epsilon_{is}, \mathbf{Z}_{is}; i \leq N, -\infty < s \leq t), \{\sum_{s=1}^{t} \xi_{Ns}, \mathcal{F}_{Nt}\}$ constitutes a martingale array for each $N, t \leq T$. Then, we can apply Corollary 3.1 of Hall and Heyde (2014) to $\sqrt{NT}\eta^{\mathsf{T}}\mathcal{E}_{\mathcal{W}} = \sum_{t} \xi_{N,t+1}$ by checking following two conditions:

(1) $\sum_{t} \mathbb{E}\left(\xi_{N,t+1}^{2} \mathbb{1}_{\left\{|\xi_{N,t+1}| > v\right\}} | \mathcal{F}_{Nt}\right) = o_{\mathbb{P}}(1).$ (2) $\sum_{t} \mathbb{E}\left(\xi_{N,t+1}^{2} | \mathcal{F}_{Nt}\right) = \eta^{\mathsf{T}} \sigma^{2} \Sigma_{w} \eta + o_{\mathbb{P}}(1).$ **Proof of (1).** First, we have

$$\sum_{t=0}^{T-1} \mathbb{E}\left(\xi_{N,t+1}^2 \mathbb{1}_{\left\{\left|\xi_{N,t+1}\right| > \upsilon\right\}} \middle| \mathcal{F}_{Nt}\right) \le \upsilon^{-2} \sum_{t} \mathbb{E}\left(\xi_{N,t+1}^4 \middle| \mathcal{F}_{Nt}\right).$$

One can easily verify that

$$\mathbb{E}\left(\xi_{N,t+1}^{4}|\mathcal{F}_{Nt}\right) \lesssim \sigma^{4}\left(\frac{1}{NT}\eta^{\mathsf{T}}\hat{\mathcal{W}}_{t}^{\mathsf{T}}\hat{\mathcal{W}}_{t}\eta\right)^{2}.$$

So, we only need to show that $\sum_t \left(\frac{1}{NT}\eta^{\mathsf{T}}\hat{\mathcal{W}}_t^{\mathsf{T}}\hat{\mathcal{W}}_t\eta\right)^2 = o_{\mathbb{P}}(1)$. First note that

$$(NT)^{-1} \hat{\mathcal{W}}_{t}^{\mathsf{T}} \hat{\mathcal{W}}_{t} = \begin{bmatrix} \frac{1}{T} \mathbf{I}_{K} & \frac{1}{\sqrt{NT}} \hat{\mathbf{U}}^{\mathsf{T}} \mathbf{y}_{t} & \frac{1}{\sqrt{NT}} \hat{\mathbf{U}}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \mathbf{y}_{t} & \frac{1}{\sqrt{NT}} \hat{\mathbf{U}}^{\mathsf{T}} \mathbf{Z}_{t} \\ & \frac{1}{NT} \mathbf{y}_{t}^{\mathsf{T}} \mathbf{y}_{t} & \frac{1}{NT} \mathbf{y}_{t}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \mathbf{y}_{t} & \frac{1}{NT} \mathbf{y}_{t}^{\mathsf{T}} \mathbf{Z}_{t} \\ & & \frac{1}{NT} \mathbf{y}_{t}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}}^{2} \mathbf{y}_{t} & \frac{1}{NT} \mathbf{y}_{t}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \mathbf{Z}_{t} \\ & & & \frac{1}{NT} \mathbf{Z}_{t}^{\mathsf{T}} \mathbf{Z}_{t} \end{bmatrix}.$$

Since similar arguments can be used to show the convergence of each entry, take

$$\sum_{t} \left(\frac{1}{\sqrt{N}T} \eta_1^{\mathsf{T}} \hat{\mathbf{U}}^{\mathsf{T}} \mathbf{y}_t \right)^2 = \frac{1}{NT^2} \sum_{t} \eta_1^{\mathsf{T}} \hat{\mathbf{U}}^{\mathsf{T}} \mathbf{y}_t \mathbf{y}_t^{\mathsf{T}} \hat{\mathbf{U}} \eta_1$$

for example, where $\eta_1 \in \mathbb{R}^K$; $\|\eta_1\| \leq 1$. Then, we have

$$\sum_{t} \eta_1^{\mathsf{T}} \hat{\mathbf{U}}^{\mathsf{T}} \mathbf{y}_t \mathbf{y}_t^{\mathsf{T}} \hat{\mathbf{U}} \eta_1 = \mathbf{y}^{\mathsf{T}} \left\{ \mathbf{I}_T \otimes \left(\hat{\mathbf{U}} \eta_1 \eta_1^{\mathsf{T}} \hat{\mathbf{U}}^{\mathsf{T}} \right) \right\} \mathbf{y}.$$

Denoting $\hat{\mathbf{U}} - \tilde{\mathbf{U}}$ by Δ , we get

$$\tilde{\mathbf{y}}^{\mathsf{T}}\Gamma_{\mathbf{y}}^{1/2}\left\{\mathbf{I}_{T}\otimes\left(\Delta\eta_{1}\eta_{1}^{\mathsf{T}}\Delta^{\mathsf{T}}\right)\right\}\Gamma_{\mathbf{y}}^{1/2}\tilde{\mathbf{y}}=\tilde{\mathbf{y}}^{\mathsf{T}}\left\{\mathbf{I}_{T}\otimes\left(\Gamma^{1/2}\Delta\eta_{1}\eta_{1}^{\mathsf{T}}\Delta^{\mathsf{T}}\Gamma^{1/2}\right)\right\}\tilde{\mathbf{y}}.$$

Letting $\hat{\mathbf{S}} \triangleq \mathbf{I}_T \otimes (\Gamma^{1/2} \Delta \eta_1 \eta_1^{\mathsf{T}} \Delta^{\mathsf{T}} \Gamma^{1/2})$ and $\tilde{\mathbf{S}} \triangleq \mathbf{I}_T \otimes (\Gamma^{1/2} \tilde{\mathbf{U}} \eta_1 \eta_1^{\mathsf{T}} \tilde{\mathbf{U}}^{\mathsf{T}} \Gamma^{1/2})$, we have $\left\| \hat{\mathbf{S}} \right\|_F^2 = T \operatorname{tr} \left\{ (\Delta \eta_1 \eta_1^{\mathsf{T}} \Delta^{\mathsf{T}} \Gamma)^2 \right\} \leq T \left\| \Delta \eta_1 \eta_1^{\mathsf{T}} \Delta^{\mathsf{T}} \right\|^2 \operatorname{tr} (\Gamma^2) = O(NT\kappa^4)$ whp. and $\left\| \hat{\mathbf{S}} \right\| = \left\| \Gamma^{1/2} \Delta \eta_1 \eta_1^{\mathsf{T}} \Delta^{\mathsf{T}} \Gamma^{1/2} \right\| \leq \left\| \Delta \right\|^2 \|\Gamma\| = O(\kappa^2)$ whp. By Lemma 8.3,

$$\mathbb{P}\left[\frac{1}{\sqrt{NT}}\left|\tilde{\mathbf{y}}^{\dagger}\hat{\mathbf{S}}\tilde{\mathbf{y}} - \mathbb{E}\left(\tilde{\mathbf{y}}^{\dagger}\hat{\mathbf{S}}\tilde{\mathbf{y}}\right)\right| > \upsilon\right] < 2\exp\left(-c\min\left\{\frac{\upsilon^{2}}{\kappa^{4}}, \frac{\upsilon\sqrt{NT}}{\kappa^{2}}\right\}\right) + \frac{2}{N}.$$

Note that $\mathbb{E}\left(\tilde{\mathbf{y}}^{\mathsf{T}}\hat{\mathbf{S}}\tilde{\mathbf{y}}\right) = T\mathbb{E}\operatorname{tr}\left(\Delta\eta_{1}\eta_{1}^{\mathsf{T}}\Delta^{\mathsf{T}}\Gamma\right)$ and $T\operatorname{tr}\left(\Delta\eta_{1}\eta_{1}^{\mathsf{T}}\Delta^{\mathsf{T}}\Gamma\right) \leq T \|\Delta\eta_{1}\eta_{1}^{\mathsf{T}}\Delta^{\mathsf{T}}\|\operatorname{tr}\left(\Gamma\right)$ hence is $O\left(NT\kappa^{2}\right)$ whp. Therefore, $\lim_{N\to\infty}\frac{\tilde{\mathbf{y}}^{\mathsf{T}}\hat{\mathbf{S}}\tilde{\mathbf{y}}}{NT\kappa^{2}} < \infty$ as. hence $\tilde{\mathbf{y}}^{\mathsf{T}}\hat{\mathbf{S}}\tilde{\mathbf{y}} = o_{\mathbb{P}}(NT^{2})$. So we have $\mathbb{E}\left(\tilde{\mathbf{y}}^{\mathsf{T}}\hat{\mathbf{S}}\tilde{\mathbf{y}}\right) = o\left(NT^{2}\right)$ by dominated convergence theorem. Next, observe that $\|\mathbf{G}\| \leq |\alpha_{1}| + |\theta| < 1$ hence $\|(\mathbf{I}_{N} - \mathbf{G})^{-1}\| \leq (1 - \|\mathbf{G}\|)^{-1} < 1/(1 - |\alpha_{1}| - |\theta|)$. So, $\|\varphi\| = O(\|\mathbf{U}\beta\|) = O(1)$ as. Using Lemma 8.1, we have

$$\mathbb{P}\left(\frac{1}{\sqrt{NT}} \left| 2\tilde{\mathbf{y}}^{\mathsf{T}} \hat{\mathbf{S}} \Phi \right| > \upsilon\right) < 2 \exp\left(-\frac{cNT\upsilon^2}{T\kappa^4}\right) + \frac{1}{N} \\
\mathbb{P}\left(\frac{1}{\sqrt{NT}} \left| 2\tilde{\mathbf{y}}^{\mathsf{T}} \tilde{\mathbf{S}} \Phi \right| > \upsilon\right) < 2 \exp\left(-\frac{cNT\upsilon^2}{T}\right) + \frac{1}{N}$$

because $\|\hat{\mathbf{S}}\Phi\| \leq \|\hat{\mathbf{S}}\| \sqrt{T} \|\varphi\| = O(\sqrt{T}\kappa^2)$ whp. and $\|\tilde{\mathbf{S}}\Phi\| = O(\sqrt{T})$ as. Also, $|\Phi^{\intercal}\tilde{\mathbf{S}}\Phi| = O(T)$ as. Therefore, we have $\frac{1}{NT^2} \sum_t \left(\eta_1^{\intercal}\hat{\mathbf{U}}^{\intercal}\mathbf{y}_t\right)^2 = o_{\mathbb{P}}(1)$. Noting that both \mathbf{y} and \mathbf{Z}_t have finite fourth-order moments, one can show that the rest are also $o_{\mathbb{P}}(1)$ similarly.

Proof of (2). Since $\sum_{t} \mathbb{E} \left(\xi_{N,t+1}^{2} | \mathcal{F}_{Nt} \right) = \frac{\sigma^{2}}{NT} \eta^{\mathsf{T}} \hat{\mathcal{W}}^{\mathsf{T}} \hat{\mathcal{W}} \eta = \sigma^{2} \eta^{\mathsf{T}} \hat{\Sigma}_{w} \eta$, by Lemma 8.5, we have (2).

Therefore, by Claims 1–3, we have

$$\mathbf{D}_{NT}\left(\hat{\mu}-\mu^{H}\right) \Rightarrow \Sigma_{w}^{-1} \mathcal{N}\left(\mathbf{o}_{K+p+2}, \sigma^{2} \Sigma_{w}\right)$$

which leads to the desired asymptotic normality.

Proof of Theorem 4.2. It suffices to show that for any A_K in Theorem 4.2, we have

$$\frac{\mathbf{A}_K \hat{\Sigma}_w^{-1}}{\sqrt{NT}} \hat{\mathcal{W}}^{\mathsf{T}} (\mathcal{W}^H - \hat{\mathcal{W}}) \mu_w^H = o_{\mathbb{P}}(1).$$

First provided that $\|\beta\| = 1$ as $K \to \infty$, we have $\frac{1}{\sqrt{NT}} \hat{\mathcal{W}}^{\intercal} (\mathcal{W}^H - \hat{\mathcal{W}}) \mu_w^H = o_{\mathbb{P}}(1)$ by the statements **10-13**. of Lemma 8.4. Since $\|\mathbf{A}_K \hat{\Sigma}_w^{-1}\| = \|\mathbf{A}_K \left(\hat{\Sigma}_w^{-1} - \Sigma_w^{-1} + \Sigma_w^{-1}\right)\|$, we only need to check if $\|\hat{\Sigma}_w^{-1} - \Sigma_w^{-1}\| = O_{\mathbb{P}}(\|\Sigma_w^{-1}\|)$. Since $\|\Sigma_w\| = O(1)$ and $\|\hat{\Sigma}_w - \Sigma_w\| = o_{\mathbb{P}}(\Sigma_w)$ by Lemma 8.5, we have the first claim.

Next, It is sufficient to show that for any $\eta \in \mathbb{R}^m$ such that $\|\eta\| \leq 1$, we have

 $\eta^{\mathsf{T}} \mathbf{A}_K \hat{\Sigma}_w^{-1} \sqrt{NT} \mathcal{E}_{\mathcal{W}} \Rightarrow \mathcal{N}(0, \sigma^2 \eta^{\mathsf{T}} \mathbf{V} \eta).$ Since

$$\eta^{\mathsf{T}} \mathbf{A}_{K} \hat{\Sigma}_{w}^{-1} \sqrt{NT} \mathcal{E}_{\mathcal{W}} = \eta^{\mathsf{T}} \mathbf{A}_{K} \left(\hat{\Sigma}_{w}^{-1} - \Sigma_{w}^{-1} \right) \sqrt{NT} \mathcal{E}_{\mathcal{W}} + \eta^{\mathsf{T}} \mathbf{A}_{K} \Sigma_{w}^{-1} \sqrt{NT} \mathcal{E}_{\mathcal{W}}$$
$$= o_{\mathbb{P}}(1) + \eta^{\mathsf{T}} \mathbf{A}_{K} \Sigma_{w}^{-1} O_{\mathbb{P}}(1),$$

we focus on the latter. Define $\xi_{N,t+1} \triangleq (NT)^{-1/2} \eta^{\mathsf{T}} \mathbf{A}_K \Sigma_w^{-1} \hat{\mathcal{W}}_t^{\mathsf{T}} \mathcal{E}_{t+1}$ and $\mathcal{F}_{Nt} = \sigma(\mathbf{A}, \epsilon_{is}, \mathbf{Z}_{is}; i \leq N, -\infty < s \leq t)$. Then, a set of pairs $\{\sum_{s=1}^t \xi_{Ns}, \mathcal{F}_{Nt}\}$ constitutes a martingale array for each $N, t \leq T$. Then, we repeat the following arguments (Hall and Heyde, 2014):

(1) $\sum_{t} \mathbb{E}\left(\xi_{N,t+1}^{2} \mathbb{1}_{\left\{|\xi_{N,t+1}| > \upsilon\right\}} | \mathcal{F}_{Nt}\right) = o_{\mathbb{P}}(1).$ (2) $\sum_{t} \mathbb{E}\left(\xi_{N,t+1}^{2} | \mathcal{F}_{Nt}\right) = \sigma^{2} \eta^{\mathsf{T}} \mathbf{V} \eta + o_{\mathbb{P}}(1).$

Proof of (1). We only need to show that

$$\sum_{t} \left(\frac{1}{NT} \eta^{\mathsf{T}} \mathbf{A}_{K} \Sigma_{w}^{-1} \hat{\mathcal{W}}_{t}^{\mathsf{T}} \hat{\mathcal{W}}_{t} \Sigma_{w}^{-1} \mathbf{A}_{K}^{\mathsf{T}} \eta \right)^{2} = o_{\mathbb{P}}(1).$$

Since both \mathbf{A}_K and Σ_w have spectral norm of O(1) as $K \to \infty$, this can be shown in the same manner as in the Proof of (1) of Claim 3 in Theorem 4.1.

Proof of (2). Since

$$\sum_{t} \mathbb{E}\left(\xi_{N,t+1}^{2} | \mathcal{F}_{Nt}\right) = \eta^{\mathsf{T}} \mathbf{A}_{K} \Sigma_{w}^{-1} \frac{\sigma^{2}}{NT} \hat{\mathcal{W}}^{\mathsf{T}} \hat{\mathcal{W}} \Sigma_{w}^{-1} \mathbf{A}_{K}^{\mathsf{T}} \eta = \sigma^{2} \eta^{\mathsf{T}} \mathbf{A}_{K} \Sigma_{w}^{-1} \hat{\Sigma}_{w} \Sigma_{w}^{-1} \mathbf{A}_{K}^{\mathsf{T}} \eta,$$

and noting that $\hat{\Sigma}_w = \Sigma_w + o_{\mathbb{P}}(1)$ for each K > 0, we have (2) provided that $\lim_{K \to \infty} \mathbf{A}_K \Sigma_w^{-1} \mathbf{A}_K^{\mathsf{T}}$ exists.

8.4 Asymptotic results

In the following lemma, we state each item for $N > N_0$ with large enough $N_0 > 0$.

Lemma 8.4. Let $\mathbf{h}_i \in \mathbb{R}^{j_i}$ for $j_1 = K$, $j_2 = p$ be real vectors such that $\|\mathbf{h}_i\| \leq 1$. Then under the conditions of Theorem 4.1, there exist $N_0, T_0 > 0$ and $v_0(N_0, T_0) \in (0, 1)$ such that the statements **1.–13.** hold for all $v \in (0, v_0)$ and $N > N_0, T > T_0$:

$$\begin{split} &\mathbf{I}. \ \mathbb{P}\left(\frac{1}{\sqrt{NT}} \left| \mathbf{h}_{1}^{\mathsf{T}} \sum_{t} (\hat{\mathbf{U}}^{\mathsf{T}} \mathbf{y}_{t} - \tilde{\mathbf{U}}^{\mathsf{T}} \varphi) \right| > \upsilon \right) < 2e^{-d_{1}\upsilon^{2}} + 2e^{-d_{2}\upsilon^{2}} + b/N \\ &\mathbf{2}. \ \mathbb{P}\left[\frac{1}{\sqrt{NT}} \left| \mathbf{h}_{1}^{\mathsf{T}} \sum_{t} \left(\hat{\mathbf{U}}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \mathbf{y}_{t} - \tilde{\mathbf{U}}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \varphi \right) \right| > \upsilon \right] < 2e^{-d_{1}\upsilon^{2}} + 2e^{-d_{2}\upsilon^{2}} + b/N \\ &\mathbf{3}. \ \mathbb{P}\left(\frac{1}{\sqrt{NT}} \left| \sum_{t} \mathbf{h}_{1}^{\mathsf{T}} \hat{\mathbf{U}}^{\mathsf{T}} \mathbf{Z}_{t} \mathbf{h}_{2} \right| > \upsilon \right) < 2e^{-d_{1}\upsilon^{2}} + 2e^{-d_{2}\upsilon^{2}} + b/N \\ &\mathbf{4}. \ \mathbb{P}\left[\frac{1}{\sqrt{NT}} \left| \sum_{t} (\mathbf{y}_{t}^{\mathsf{T}} \mathbf{y}_{t} - \operatorname{tr} \mathbb{E}\Gamma - \varphi^{\mathsf{T}} \varphi) \right| > \upsilon \right] < 2e^{-d_{2}\upsilon^{2}} + 2e^{-d_{3}\upsilon^{2}} \\ &\mathbf{5}. \ \mathbb{P}\left[\frac{1}{NT} \left| \sum_{t} \mathbf{y}_{t}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \mathbf{y}_{t} - T\varphi^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \varphi - T \operatorname{tr} \mathbb{E}\left(\mathcal{L}_{\mathbf{A}} \Gamma\right) \right| > \upsilon \right] < 2e^{-d_{2}\upsilon^{2}} + 2e^{-d_{3}\upsilon^{2}} \\ &\mathbf{6}. \ \mathbb{P}\left(\frac{1}{NT} \left| \sum_{t} \mathbf{y}_{t}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \mathbf{y}_{t} - \varphi^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \varphi - \operatorname{tr} \mathbb{E}\left(\mathcal{L}_{\mathbf{A}} \Gamma\right) \right| > \upsilon \right] < 2e^{-d_{1}\upsilon^{2}} + 2e^{-d_{3}\upsilon^{2}} \\ &\mathbf{6}. \ \mathbb{P}\left[\frac{1}{NT} \left| \sum_{t} \mathbf{y}_{t}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}}^{2} \mathbf{y}_{t} - \varphi^{\mathsf{T}} \mathcal{L}_{\mathbf{A}}^{2} \varphi - \operatorname{tr} \mathbb{E}\left(\mathcal{L}_{\mathbf{A}}^{2} \Gamma\right) \right| > \upsilon \right] < 2e^{-d_{1}\upsilon^{2}} + 2e^{-d_{3}\upsilon^{2}} \\ &\mathbf{7}. \ \mathbb{P}\left[\frac{1}{NT} \left| \sum_{t} \mathbf{y}_{t}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}}^{2} \mathbf{y}_{t} - \varphi^{\mathsf{T}} \mathcal{L}_{\mathbf{A}}^{2} \varphi - \operatorname{tr} \mathbb{E}\left(\mathcal{L}_{\mathbf{A}}^{2} \Gamma\right) \right\} \right| > \upsilon \right] < 2e^{-d_{1}\upsilon^{2}} + 2e^{-d_{3}\upsilon^{2}} \\ &\mathbf{8}. \ \mathbb{P}\left(\frac{1}{NT} \left| \sum_{t} \mathbf{y}_{t}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \mathbf{Z}_{t} \mathbf{h}_{2} \right| > \upsilon \right) < 2e^{-d_{4}\upsilon^{2}} + 2e^{-d_{3}\upsilon^{2}} + b/N \\ &\mathbf{9}. \ \mathbb{P}\left[\frac{1}{NT} \left| \sum_{t} \mathbf{y}_{t}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \mathbf{Z}_{t} \mathbf{h}_{2} \right| > \upsilon \right] < 2e^{-d_{2}\upsilon^{2}} \\ &\mathbf{10}. \ \mathbb{P}\left[\sqrt{T} \left| \mathbf{h}_{1}^{\mathsf{T}} \hat{\mathbf{U}}_{t}^{\mathsf{T}} \left(\hat{\mathbf{U}} - \tilde{\mathbf{U}}\right) \beta \right| > \upsilon \right] < 2e^{-c\upsilon^{2}/\kappa^{2}} + \frac{1}{N} \\ &\mathbf{11}. \ \mathbb{P}\left[\frac{1}{\sqrt{T}} \left| \sum_{t} \mathbf{y}_{t}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \left(\hat{\mathbf{U}} - \tilde{\mathbf{U}}\right) \beta \right| > \upsilon \right] < 2e^{-c\upsilon^{2}/\kappa^{2}} + \frac{1}{N} \\ &\mathbf{13}. \ \mathbb{P}\left[\frac{1}{\sqrt{T}} \left| \sum_{t} \mathbf{y}_{t}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \left(\hat{\mathbf{U}} - \tilde{\mathbf{U}}\right) \beta \right| > \upsilon \right] < 2e^{-c\upsilon^{2}/\kappa^{2}} + \frac{1}{N} \\ &\text{where } d_{1} \ a \frac{c_{1}NT}{\kappa^{2}}, d_{2} \ a c_{2}NT, d_{3}$$

constants.

Proof of 1. Note that

$$\sum_{t=1}^{T} \mathbf{h}_{1}^{\mathsf{T}} \left(\hat{\mathbf{U}}^{\mathsf{T}} \mathbf{y}_{t} - \tilde{\mathbf{U}}^{\mathsf{T}} \varphi \right) = \mathbf{h}_{1}^{\mathsf{T}} \left(\mathbf{1}_{T} \otimes \Delta \right)^{\mathsf{T}} \Gamma_{\mathbf{y}}^{1/2} \tilde{\mathbf{y}} + \mathbf{h}_{1}^{\mathsf{T}} \left(\mathbf{1}_{T} \otimes \tilde{\mathbf{U}} \right)^{\mathsf{T}} \Gamma_{\mathbf{y}}^{1/2} \tilde{\mathbf{y}} + T \mathbf{h}_{1}^{\mathsf{T}} \Delta^{\mathsf{T}} \varphi = \mathbf{h}_{1}^{\mathsf{T}} \hat{\mathbf{S}}^{\mathsf{T}} \tilde{\mathbf{y}} + \mathbf{h}_{1}^{\mathsf{T}} \tilde{\mathbf{S}}^{\mathsf{T}} \tilde{\mathbf{y}} + T \mathbf{h}_{1}^{\mathsf{T}} \Delta^{\mathsf{T}} \varphi.$$

where $\hat{\mathbf{S}} \triangleq \Gamma_{\mathbf{y}}^{1/2} (\mathbf{1}_T \otimes \Delta)$ and $\tilde{\mathbf{S}} \triangleq \Gamma_{\mathbf{y}}^{1/2} (\mathbf{1}_T \otimes \tilde{\mathbf{U}})$. Since $\|\hat{\mathbf{S}}\mathbf{h}_1\| = O(\sqrt{T\kappa})$ whp., by Lemma 8.1 we have

$$\mathbb{P}\left(\left|\mathbf{h}_{1}^{\mathsf{T}}\hat{\mathbf{S}}^{\mathsf{T}}\tilde{\mathbf{y}}\right| > \sqrt{N}T\upsilon\right) < 2\exp\left(\frac{-cNT^{2}\upsilon^{2}}{T\kappa^{2}}\right) + \frac{1}{N}.$$

Likewise, since $\|\tilde{\mathbf{S}}\| = O(\sqrt{T})$ as., by Lemma 8.1 we have $\mathbb{P}\left(\left|\mathbf{h}_1^{\mathsf{T}}\tilde{\mathbf{S}}\tilde{\mathbf{y}}\right| > \sqrt{N}Tv\right) < 2\exp(-c_2NT^2v^2/T)$. Finally, $|T\mathbf{h}_1^{\mathsf{T}}\Delta^{\mathsf{T}}\varphi| = O(T\kappa)$ whp., i.e., $\mathbb{P}\left(|T\mathbf{h}_1^{\mathsf{T}}\Delta^{\mathsf{T}}\varphi| > CT\kappa\right) < CT\kappa$

1/N for all large enough N. By selecting v such that $\sqrt{N}Tv = \omega(T\kappa)$, i.e., $v = \omega(\kappa/\sqrt{N})$, and noting that $\pi_N \gg \sqrt{TN\rho_N}$ hence $\kappa/\sqrt{N} = o(N^{-1/2})$, we can reduce v sufficiently for large enough N.

Proof of 2. First note that

$$\mathbf{h}_{1}^{\mathsf{T}} \sum_{t} \left(\hat{\mathbf{U}}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \mathbf{y}_{t} - \tilde{\mathbf{U}}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \varphi \right) = \mathbf{h}_{1}^{\mathsf{T}} \left[\mathbf{1}_{T} \otimes \left\{ \mathcal{L}_{\mathbf{A}} \Delta \right\} \right]^{\mathsf{T}} \Gamma_{\mathbf{y}}^{-1/2} \tilde{\mathbf{y}} + \mathbf{h}_{1}^{\mathsf{T}} \left\{ \mathbf{1}_{T} \otimes \mathcal{L}_{\mathbf{A}} \tilde{\mathbf{U}} \right\}^{\mathsf{T}} \Gamma_{\mathbf{y}}^{-1/2} \tilde{\mathbf{y}} + T \mathbf{h}_{1}^{\mathsf{T}} \Delta^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \varphi.$$

For $\hat{\mathbf{S}} \triangleq \Gamma_{\mathbf{y}}^{1/2} \{ \mathbf{1}_T \otimes (\mathcal{L}_{\mathbf{A}} \Delta) \}$ and $\tilde{\mathbf{S}} \triangleq \Gamma_{\mathbf{y}}^{1/2} \{ \mathbf{1}_T \otimes (\mathcal{L}_{\mathbf{A}} \tilde{\mathbf{U}}) \}$, we have $\left\| \hat{\mathbf{S}} \mathbf{h}_1 \right\| = O(\sqrt{T}\kappa)$ whp. because $\rho(\mathcal{L}_{\mathbf{A}}) = O(1)$ as. hence $\mathbb{P}\left(\left| \mathbf{h}_1^{\mathsf{T}} \hat{\mathbf{S}}^{\mathsf{T}} \tilde{\mathbf{y}} \right| > \sqrt{N} T \upsilon \right) < 2 \exp\left(\frac{-cNT^2 \upsilon^2}{T\kappa^2} \right) + \frac{1}{N}$. Since $\left\| \tilde{\mathbf{S}} \mathbf{h}_1 \right\| = O\left(\sqrt{T}\right)$ as., we have $\mathbb{P}\left(\left| \mathbf{h}_1^{\mathsf{T}} \tilde{\mathbf{S}}^{\mathsf{T}} \tilde{\mathbf{y}} \right| > \sqrt{N} T \upsilon \right) < 2 \exp\left(-cNT^2 \upsilon^2/T \right)$. The tail probability of $|T\mathbf{h}_1^{\mathsf{T}} \Delta^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \varphi|$ can be bounded similarly to 1.

Proof of 3. Let $\mathbf{Z} \triangleq [\mathbf{Z}_0^{\mathsf{T}}, ..., \mathbf{Z}_{T-1}^{\mathsf{T}}]^{\mathsf{T}}$ and $\hat{\mathbf{S}} \triangleq \mathbf{1}_T \otimes \Delta$ and $\tilde{\mathbf{S}} \triangleq \mathbf{1}_T \otimes \tilde{\mathbf{U}}$. Then, we can express $\sum_t \mathbf{h}_1^{\mathsf{T}} \hat{\mathbf{U}}^{\mathsf{T}} \mathbf{Z}_t \mathbf{h}_2$ as tr $(\mathbf{Z}\mathbf{h}_2\mathbf{h}_1^{\mathsf{T}} \hat{\mathbf{S}}^{\mathsf{T}}) + \operatorname{tr} (\mathbf{Z}\mathbf{h}_2\mathbf{h}_1^{\mathsf{T}} \hat{\mathbf{S}}^{\mathsf{T}})$. This is equal to

$$\operatorname{vec}\left(\mathbf{h}_{2}\mathbf{h}_{1}^{\mathsf{T}}\right)^{\mathsf{T}}\left\{\mathbf{I}_{p}\otimes\left(\hat{\mathbf{S}}+\tilde{\mathbf{S}}\right)^{\mathsf{T}}\right\}\operatorname{vec}\left(\mathbf{Z}\right).$$

Note that $\left\|\mathbf{I}_p \otimes \hat{\mathbf{S}}^{\mathsf{T}}\right\| = O\left(\sqrt{T}\kappa\right)$ whp. and $\left\|\mathbf{I}_p \otimes \tilde{\mathbf{S}}^{\mathsf{T}}\right\| = O\left(\sqrt{T}\right)$. Therefore, by Lemma 8.1, we have

$$\mathbb{P}\left[\left|\sum_{t} \mathbf{h}_{1}^{\mathsf{T}} \hat{\mathbf{U}}^{\mathsf{T}} \mathbf{Z}_{t} \mathbf{h}_{2}\right| > \sqrt{N} T \upsilon\right] \leq 2 \exp\left(-\frac{cNT^{2}\upsilon^{2}}{T\kappa^{2}}\right) + 2 \exp\left(-\frac{c'NT^{2}\upsilon^{2}}{T}\right) + \frac{1}{N}.$$

Proof of 4. First we have

$$\sum_{t} \left(\mathbf{y}_{t}^{\mathsf{T}} \mathbf{y}_{t} - \varphi^{\mathsf{T}} \varphi \right) = \left\| \Gamma_{\mathbf{y}}^{1/2} \tilde{\mathbf{y}} + \Phi \right\|^{2} - T \varphi^{\mathsf{T}} \varphi = \tilde{\mathbf{y}}^{\mathsf{T}} \Gamma_{\mathbf{y}} \tilde{\mathbf{y}} + 2 \Phi^{\mathsf{T}} \Gamma_{\mathbf{y}}^{1/2} \tilde{\mathbf{y}}.$$

Note that $\|\Gamma_{\mathbf{y}}\|_F^2 = \sum_{i=1}^{NT} \sigma_i(\Gamma_{\mathbf{y}})^2 = O(NT)$ as. By Theorem 8.3, we have

$$\mathbb{P}\left(|\tilde{\mathbf{y}}^{\mathsf{T}}\Gamma_{\mathbf{y}}\tilde{\mathbf{y}} - T\operatorname{tr}\mathbb{E}\Gamma| > \upsilon\right) < 2\exp\left(-c\min\left\{\frac{\upsilon^2}{NT}, \upsilon\right\}\right).$$

By Lemma 8.1,

$$\mathbb{P}\left(2|\Phi^{\mathsf{T}}\Gamma_{\mathbf{y}}^{1/2}\tilde{\mathbf{y}}| > \upsilon\right) < 2\exp\left(-\frac{c'\upsilon^2}{T}\right)$$

since $\left\|\Gamma_{\mathbf{y}}^{1/2}\Phi\right\|^2 \leq \left\|\Gamma_{\mathbf{y}}\right\| T\varphi^{\mathsf{T}}\varphi = O(T)$ as.

Proof of 5. Let $\mathbf{S}_{\mathbf{A}} \triangleq \Gamma_{\mathbf{y}}^{1/2} \left(\mathbf{I}_T \otimes \mathcal{L}_{\mathbf{A}} \right) \Gamma_{\mathbf{y}}^{1/2}$. We have

$$\sum_{t} \mathbf{y}_{t}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \mathbf{y}_{t} - \varphi^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \varphi = \left(\Gamma_{\mathbf{y}}^{1/2} \tilde{\mathbf{y}} + \Phi \right)^{\mathsf{T}} \left(\mathbf{I}_{T} \otimes \mathcal{L}_{\mathbf{A}} \right) \left(\Gamma_{\mathbf{y}}^{1/2} \tilde{\mathbf{y}} + \Phi \right) - T \varphi^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \varphi$$
$$= \tilde{\mathbf{y}}^{\mathsf{T}} \mathbf{S}_{\mathbf{A}} \tilde{\mathbf{y}} + 2 \Phi^{\mathsf{T}} \Gamma_{\mathbf{y}}^{-\frac{1}{2}} \mathbf{S}_{\mathbf{A}} \tilde{\mathbf{y}}.$$

Also,

$$\|\mathbf{S}_{\mathbf{A}}\|_{F}^{2} = \operatorname{tr}\left[\left(\mathbf{I}_{T} \otimes \mathcal{L}_{\mathbf{A}}\right) \Gamma_{\mathbf{y}}\left(\mathbf{I}_{T} \otimes \mathcal{L}_{\mathbf{A}}\right) \Gamma_{\mathbf{y}}\right] \leq \operatorname{tr}\left(\Gamma_{\mathbf{y}}^{2}\right) \left\|\mathbf{I}_{T} \otimes \mathcal{L}_{\mathbf{A}}^{2}\right\| = O(NT) \ as.$$
$$\left\|\mathbf{S}_{\mathbf{A}}\Gamma_{\mathbf{y}}^{-\frac{1}{2}}\Phi\right\| = O\left(\sqrt{T}\right) \ as.$$

We can apply Lemmas 8.3 and 8.1 to directly obtain the following:

$$\mathbb{P}\left[\left|\tilde{\mathbf{y}}^{\mathsf{T}}\mathbf{S}_{\mathbf{A}}\tilde{\mathbf{y}} - T\mathbb{E}\operatorname{tr}\left(\mathcal{L}_{\mathbf{A}}\Gamma\right)\right| > \upsilon\right] < 2\exp\left(-c\min\left\{\frac{\upsilon^{2}}{NT},\upsilon\right\}\right)$$
$$\mathbb{P}\left[\left|2\Phi^{\mathsf{T}}\Gamma_{\mathbf{y}}^{-\frac{1}{2}}\mathbf{S}_{\mathbf{A}}\tilde{\mathbf{y}}\right| > \upsilon\right] < 2\exp\left(-c\upsilon^{2}/T\right).$$

-	-	1

Proof of 6. We have

$$\sum_{t} \mathbf{y}_{t}^{\mathsf{T}} \mathbf{Z}_{t} \mathbf{h}_{2} = \left(\Gamma_{\mathbf{y}}^{1/2} \tilde{\mathbf{y}} + \Phi \right)^{\mathsf{T}} \mathbf{Z} \mathbf{h}_{2} = \operatorname{tr} \left(\mathbf{h}_{2} \tilde{\mathbf{y}}^{\mathsf{T}} \Gamma_{\mathbf{y}}^{1/2} \mathbf{Z} \right) + \operatorname{tr} \left(\mathbf{h}_{2} \Phi^{\mathsf{T}} \mathbf{Z} \right)$$
$$= \operatorname{vec} \left(\tilde{\mathbf{y}} \mathbf{h}_{2}^{\mathsf{T}} \right)^{\mathsf{T}} \left(\mathbf{I}_{p} \otimes \Gamma_{\mathbf{y}}^{1/2} \right) \operatorname{vec} \left(\mathbf{Z} \right) + \operatorname{vec} \left(\Phi \mathbf{h}_{2}^{\mathsf{T}} \right)^{\mathsf{T}} \left(\mathbf{I}_{p} \otimes \mathbf{I}_{NT} \right) \operatorname{vec} \left(\mathbf{Z} \right).$$

Therefore, since $\|\Gamma_{\mathbf{y}}\| = O(1)$ as., $\left\| \left(\mathbf{I}_p \otimes \Gamma_{\mathbf{y}}^{1/2} \right) \operatorname{vec} \left(\tilde{\mathbf{y}} \mathbf{h}_2^{\mathsf{T}} \right) \right\| \lesssim \|\operatorname{vec} \left(\tilde{\mathbf{y}} \mathbf{h}_2^{\mathsf{T}} \right) \|$ which is concentrated around \sqrt{NTp} (Vershynin, 2018) hence is $O(\sqrt{NTp})$ whp. Also, we have $\| (\mathbf{I}_p \otimes \mathbf{I}_{NT}) \operatorname{vec} \left(\Phi \mathbf{h}_2^{\mathsf{T}} \right) \| = O(\sqrt{T})$ as. By Lemma 8.1,

$$\mathbb{P}\left(\left|\sum_{t} \mathbf{y}_{t}^{\mathsf{T}} \mathbf{Z}_{t} \mathbf{h}_{2}\right| > \upsilon\right) \leq 2 \exp\left(-\frac{c\upsilon^{2}}{NTp}\right) + 2 \exp\left(-\frac{c\upsilon^{2}}{T}\right) + \frac{1}{N}.$$

Therefore, the conclusion follows after letting $\sqrt{NTv} = \omega(\sqrt{NTp})$.

Proof of 7. Similarly to the proof of 5., let $\mathbf{S}_{\mathbf{A}} \triangleq \Gamma_{\mathbf{y}}^{-1/2} \left(\mathbf{I}_T \otimes \mathcal{L}_{\mathbf{A}}^2 \right) \Gamma_{\mathbf{y}}^{-1/2}$ and check that

$$\|\mathbf{S}_{\mathbf{A}}\|_{F}^{2} \leq \operatorname{tr}\left(\Gamma_{\mathbf{y}}^{2}\right) \|\mathbf{I}_{T} \otimes \mathcal{L}_{\mathbf{A}}^{4}\| = O(NT) \ as., \left\|\mathbf{S}_{\mathbf{A}}\Gamma_{\mathbf{y}}^{-\frac{1}{2}}\Phi\right\| = O\left(\sqrt{T}\right) \ as.$$

Again by Lemmas 8.3 and 8.1, we directly have the following:

$$\mathbb{P}\left[\left|\tilde{\mathbf{y}}^{\mathsf{T}}\mathbf{S}_{\mathbf{A}}\tilde{\mathbf{y}} - T\mathbb{E}\operatorname{tr}\left(\mathcal{L}_{\mathbf{A}}^{2}\Gamma\right)\right| > \upsilon\right] < 2\exp\left(-c\min\left\{\frac{\upsilon^{2}}{NT},\upsilon\right\}\right)$$
$$\mathbb{P}\left[\left|2\Phi^{\mathsf{T}}\Gamma_{\mathbf{y}}^{-\frac{1}{2}}\mathbf{S}_{\mathbf{A}}\tilde{\mathbf{y}}\right| > \upsilon\right] < 2\exp\left(-c\upsilon^{2}/T\right).$$

Proof of 8. The same logic as the proof in 6. applies here. Since $\|\mathcal{L}_{\mathbf{A}}\| \leq 1$, we get the same bound by letting $\sqrt{NTv} = \omega(\sqrt{NTp})$.

Proof of 9. Note that

$$\sum_{t} \mathbf{h}_{2}^{\mathsf{T}} \mathbf{Z}_{t}^{\mathsf{T}} \mathbf{Z}_{t} \mathbf{h}_{2} = \operatorname{tr} \left(\mathbf{Z} \mathbf{h}_{2} \mathbf{h}_{2}^{\mathsf{T}} \mathbf{Z}^{\mathsf{T}} \right) = \operatorname{vec} \left(\mathbf{Z}^{\mathsf{T}} \right)^{\mathsf{T}} \left\{ \mathbf{I}_{NT} \otimes \left(\mathbf{h}_{2} \mathbf{h}_{2}^{\mathsf{T}} \right) \right\} \operatorname{vec} \left(\mathbf{Z}^{\mathsf{T}} \right).$$

Note that $\mathbb{E}\left[\operatorname{vec}\left(\mathbf{Z}^{\mathsf{T}}\right)\operatorname{vec}\left(\mathbf{Z}^{\mathsf{T}}\right)^{\mathsf{T}}\right] = \mathbf{I}_{NT} \otimes \Sigma_{z}$. Also, by the definition of \mathbf{h}_{2} , we have $\|\mathbf{I}_{NT} \otimes (\mathbf{h}_{2}\mathbf{h}_{2}^{\mathsf{T}})\|_{F}^{2} \leq NT$ and $\|\mathbf{I}_{NT} \otimes (\mathbf{h}_{2}\mathbf{h}_{2}^{\mathsf{T}})\| \leq 1$. Therefore, by Hanson-Wright inequality (Rudelson and Vershynin, 2013),

$$\mathbb{P}\left[\left|\mathbf{h}_{2}^{\mathsf{T}}\sum_{t}\mathbf{Z}_{t}^{\mathsf{T}}\mathbf{Z}_{t}\mathbf{h}_{2}-NT\mathbf{h}_{2}^{\mathsf{T}}\Sigma_{z}\mathbf{h}_{2}\right|>\upsilon\right]<2\exp\left[-c\min\left\{\frac{\upsilon^{2}}{NT},\upsilon\right\}\right].$$

Proof of 10. Note that $\|\Delta\|^2 = O(\kappa^2)$ whp. and $\|\tilde{\mathbf{U}}\| = 1$. Also, $\|\beta\| = O(1)$. So, since $\kappa = o(1)$ by assumption and

$$\left|\mathbf{h}_{1}^{\mathsf{T}}\hat{\mathbf{U}}^{\mathsf{T}}\Delta\beta\right| \leq \left(\left\|\Delta\right\|^{2} + \left\|\Delta\right\|\right)\left\|\beta\right\| = O\left(\kappa\right)$$

whp., by letting $v = \omega(\sqrt{T}\kappa)$ and noting that $\sqrt{T}\kappa = o(1)$, we have the conclusion. \Box

Proof of 11. Similar to the proof of 1., taking $\hat{\mathbf{S}} \triangleq \Gamma_{\mathbf{y}}^{1/2} (\mathbf{1}_T \otimes \Delta)$ which has the spectral norm of $O(\sqrt{T\kappa})$ whp., we have

$$\sum_{t} \mathbf{y}_{t}^{\mathsf{T}} \Delta \beta = \left(\Gamma_{\mathbf{y}}^{1/2} \tilde{\mathbf{y}} + \Phi \right)^{\mathsf{T}} \left(\mathbf{1}_{T} \otimes \Delta \right) \beta = \tilde{\mathbf{y}}^{\mathsf{T}} \hat{\mathbf{S}} \beta + \Phi^{\mathsf{T}} \Gamma_{\mathbf{y}}^{-\frac{1}{2}} \hat{\mathbf{S}} \beta.$$

First noting that $\|\beta\| = O(1)$ as $K \to \infty$, Lemma 8.1 gives

$$\mathbb{P}\left(\left|\tilde{\mathbf{y}}^{\mathsf{T}}\hat{\mathbf{S}}\beta\right| \geq \sqrt{T}\upsilon\right) \leq 2\exp\left(\frac{-cT\upsilon^2}{T\kappa^2}\right) + \frac{1}{N}.$$

It is straightforward that $\left| \Phi^{\dagger} \Gamma_{\mathbf{y}}^{-\frac{1}{2}} \hat{\mathbf{S}} \beta \right| = O(T\kappa)$ whp. hence by letting $\sqrt{T} \upsilon = \omega(T\kappa)$, we obtain the conclusion.

Proof of 12. Let $\hat{\mathbf{S}} \triangleq \Gamma_{\mathbf{y}}^{1/2} \{ \mathbf{1}_T \otimes (\mathcal{L}_{\mathbf{A}} \Delta) \}$ which is of $O\left(\sqrt{T}\kappa\right)$ whp. The rest of the proof coincides with the proof of 11.

Proof of 13. Note that $\mathbf{h}_2^{\mathsf{T}} \sum_t \mathbf{Z}_t^{\mathsf{T}} \Delta \beta = \mathbf{h}_2^{\mathsf{T}} \mathbf{Z}^{\mathsf{T}} \left(\mathbf{1}_T \otimes \Delta \right) \beta$ which equals

tr {
$$\mathbf{Z}\mathbf{h}_{2}\beta^{\mathsf{T}}(\mathbf{1}_{T}\otimes\Delta)^{\mathsf{T}}$$
} = vec ($\mathbf{h}_{2}\beta^{\mathsf{T}}$) ^{T} { $\mathbf{I}_{p}\otimes(\mathbf{1}_{T}\otimes\Delta)^{\mathsf{T}}$ } vec (\mathbf{Z})

Since the norm of vec $(\mathbf{h}_2\beta^{\intercal})^{\intercal} \{ \mathbf{I}_p \otimes (\mathbf{1}_T \otimes \Delta)^{\intercal} \}$ is of $O(\sqrt{T}\kappa)$ whp., we obtain the bound after applying Lemma 8.1.

Lemma 8.5. There exist $N_0, T_0 > 0$ and $v_0(N_0, T_0) \in (0, 1)$ such that for all $v \in (0, v_0)$ and $N > N_0, T > T_0$,

$$\mathbb{P}\left[\sup_{\mathbf{h}\in\mathbb{R}^{K+p+2};\|\mathbf{h}\|\leq 1}\left|\mathbf{h}^{\mathsf{T}}\left(\hat{\Sigma}_{w}-\Sigma_{w}\right)\mathbf{h}\right|>\upsilon\right]\lesssim K^{2}\left[\frac{1}{N}+\exp\left\{-c\upsilon^{2}\frac{NT}{K^{2}\max\left(p,\kappa^{2}\right)}\right\}\right]$$

Proof. For $\mathbf{h}_1 \in \mathbb{R}^K$ and $\mathbf{h}_2 \in \mathbb{R}^p$, let $\mathbf{h} \triangleq (\mathbf{h}_1^{\mathsf{T}}, h_1, h_2, \mathbf{h}_2^{\mathsf{T}})^{\mathsf{T}} \in \mathbb{R}^{K+p+2}$ such that $\|\mathbf{h}\| \leq 1$. Let \mathbf{e}_i be i^{th} canonical basis of \mathbb{R}^{K+p+2} . By Lemma 8.4, we have

$$\max_{i,j} \mathbb{P}\left(\left| \mathbf{e}_i^{\mathsf{T}} \left(\hat{\Sigma}_w - \Sigma_w \right) \mathbf{e}_j \right| > \upsilon \right) \lesssim \frac{1}{N} + \exp\left(-c_1 \upsilon^2 N T / p \right) + \exp\left(-c_2 \upsilon^2 N T / \kappa^2 \right).$$

Since

$$\sup_{\mathbf{h};\|\mathbf{h}\|\leq 1} \left| \mathbf{h}^{\mathsf{T}} \left(\hat{\Sigma}_w - \Sigma_w \right) \mathbf{h} \right| \leq \sup_{\mathbf{h};\|\mathbf{h}\|\leq 1} \|\mathbf{h}\|_1^2 \max_{i,j} \left| \hat{\sigma}_{w,ij} - \sigma_{w,ij} \right| \lesssim K \max_{i,j} \left| \hat{\sigma}_{w,ij} - \sigma_{w,ij} \right|,$$

we have

$$\mathbb{P}\left[\sup_{\mathbf{h};\|\mathbf{h}\|\leq 1} \left| \mathbf{h}^{\mathsf{T}} \left(\hat{\Sigma}_{w} - \Sigma_{w} \right) \mathbf{h} \right| > v \right] \leq \mathbb{P}\left(\max_{i,j} \left| \hat{\sigma}_{w,ij} - \sigma_{w,ij} \right| > \frac{ce}{K} \right) \\
\lesssim K^{2} \left\{ \frac{1}{N} + \exp\left(-c_{1}' v^{2} \frac{NT}{K^{2} p} \right) + \exp\left(-c_{2}' v^{2} \frac{NT}{K^{2} \kappa^{2}} \right) \right\}.$$

Lemma 8.6. If $\pi_N \gg \sqrt{TN\rho_N}$, we have

$$\sqrt{T}\left(\hat{\mathbf{U}}\hat{\boldsymbol{\beta}}-\mathbf{U}\boldsymbol{\beta}\right)\Rightarrow\mathcal{N}\left(\mathbf{o}_{K},\sigma^{2}\mathbf{U}\mathbf{U}^{\intercal}\right)$$

as $\min(N, T) \to \infty$.

Proof. Recall that $\sqrt{T} \left(\hat{\beta} - \mathbf{H}^{\mathsf{T}} \beta \right) \Rightarrow \mathcal{N} \left(\mathbf{o}_{K}, \sigma^{2} \mathbf{I}_{K} \right)$ by Theorem 4.1. By simple algebra, we have

$$\sqrt{T} \left(\hat{\mathbf{U}} \hat{\beta} - \mathbf{U} \beta \right) = \sqrt{T} \left(\hat{\mathbf{U}} \hat{\beta} - \tilde{\mathbf{U}} \mathbf{H}^{\mathsf{T}} \beta \right)
= \Delta \sqrt{T} \left(\hat{\beta} - \mathbf{H}^{\mathsf{T}} \beta \right) + \sqrt{T} \Delta \mathbf{H}^{\mathsf{T}} \beta + \tilde{\mathbf{U}} \sqrt{T} \left(\hat{\beta} - \mathbf{H}^{\mathsf{T}} \beta \right).$$
(10)

By assumption, $\sqrt{T}\Delta = O\left(\frac{\sqrt{TN\rho_N}}{\pi_N}\right)$ whp. hence we have $\mathbb{P}\left[\left\{\sqrt{T}\Delta \le C\frac{\sqrt{TN\rho_N}}{\pi_N}\right\} ev.\right] = 1$

for some constant C > 0. Provided that $\pi_N \gg \sqrt{TN\rho_N}$, this implies $\sqrt{T}\Delta \to 0$ as. Therefore, we have $10 = o_{\mathbb{P}}(1) + \tilde{\mathbf{U}}\sqrt{T}\left(\hat{\beta} - \mathbf{H}^{\intercal}\beta\right)$ and the conclusion by Slutsky. \Box

8.5 Theories for finite time models

Here, we present theoretical results for the finite-time corollary models specified in section 2.4. Recall ENR:

$$\mathbf{y} = \alpha \mathbf{1}_N + \mathbf{U}\beta + \mathbf{Z}\gamma + \mathcal{E}$$

and ENAR (with finite T) models:

$$\mathbf{y}_{t+1} = \alpha \mathbf{y}_t + \theta \mathcal{L}_{\mathbf{A}} \mathbf{y}_t + \mathbf{U}\beta + \mathbf{Z}\gamma + \mathcal{E}_{t+1}.$$

Note that ENR can be treated as a subset model of ENAR under finite time. Therefore, we only verify the asymptotic properties of ENAR with finite T.

Proof of Theorem 4.3 Let $\mathbf{D}_N \triangleq \operatorname{diag}\left(\mathbf{I}_K, \sqrt{N}\mathbf{I}_{p+2}\right)$. Then, we have $\mathcal{W}_t \mathbf{D}_N = \sqrt{N}\mathbf{W}_t, \sqrt{N}\mathbf{D}_N^{-1}\mu_w^H = \mu^H$, and $\mathbf{W}_t \mu = \mathbf{W}_t^H \mu^H = \mathcal{W}_t^H \mu_w^H$. Next,

$$\hat{\mu} = \left(\frac{\mathbf{D}_N}{N}\hat{\mathcal{W}}^{\mathsf{T}}\hat{\mathcal{W}}\mathbf{D}_N\right)^{-1}\frac{\mathbf{D}_N}{\sqrt{N}}\hat{\mathcal{W}}^{\mathsf{T}}\left(\mathcal{W}^H\mu_w^H + \mathcal{E}\right)$$
$$= \left(T\hat{\Sigma}_w\mathbf{D}_N\right)^{-1}\frac{1}{\sqrt{N}}\hat{\mathcal{W}}^{\mathsf{T}}\left\{\left(\mathcal{W}^H - \hat{\mathcal{W}}\right)\mu_w^H + \hat{\mathcal{W}}\mu_w^H + \mathcal{E}\right\}$$
$$= \frac{1}{T}\mathbf{D}_N^{-1}\hat{\Sigma}_w^{-1}\left\{\frac{1}{\sqrt{N}}\hat{\mathcal{W}}^{\mathsf{T}}\left(\mathcal{W}^H - \hat{\mathcal{W}}\right)\mu_w^H + \sqrt{N}T\mathcal{E}_W\right\} + \sqrt{N}\mathbf{D}_N^{-1}\mu_w^H$$

hence

$$\mathbf{D}_{N}\left(\hat{\mu}-\mu^{H}\right)=\frac{\hat{\Sigma}_{w}^{-1}}{\sqrt{N}T}\hat{\mathcal{W}}^{\mathsf{T}}(\mathcal{W}^{H}-\hat{\mathcal{W}})\mu_{w}^{H}+\sqrt{N}\hat{\Sigma}_{w}^{-1}\mathcal{E}_{\mathcal{W}}.$$

Claim 1. $\hat{\Sigma}_w \Rightarrow \Sigma_w$.

Proof. Note that we are assuming $\pi_N \gg \sqrt{N\rho_N}$ here. Then, the claim follows from Lemma 8.4 and Lemma 8.5 with fixed $T = T_0 > 0$.

Claim 2.
$$\frac{1}{\sqrt{NT}}\hat{\mathcal{W}}^{\mathsf{T}}(\mathcal{W}^H - \hat{\mathcal{W}})\mu_w^H = o_{\mathbb{P}}(1).$$

Proof. We have

$$\frac{1}{\sqrt{N}T}\hat{\mathcal{W}}^{\mathsf{T}}\left(\mathcal{W}^{H}-\hat{\mathcal{W}}\right)\mu_{w}^{H} = \begin{bmatrix} \hat{\mathbf{U}}^{\mathsf{T}} \\ \frac{1}{\sqrt{N}T}\sum_{t} \mathbf{y}_{t}^{\mathsf{T}} \\ \frac{1}{\sqrt{N}T}\sum_{t} \mathbf{y}_{t}^{\mathsf{T}}\mathcal{L}_{\mathbf{A}} \\ \frac{1}{\sqrt{N}T}\sum_{t} \mathbf{Z}_{t}^{\mathsf{T}} \end{bmatrix} [\hat{\mathbf{U}}-\tilde{\mathbf{U}} \mid \mathbf{O}_{N\times(p+2)}]\mu^{H}.$$

Again, we are assuming $\pi_N \gg \sqrt{N\rho_N}$ hence $\kappa = o(1)$. Then the result follows by the statements **10.–13.** of Lemma 8.4 with fixed $T = T_0 > 0$.

Next, since Claims 1 and 2 are true, for $\sqrt{N}\hat{\Sigma}_w^{-1}\mathcal{E}_W = \sqrt{N}\left(\hat{\Sigma}_w^{-1} - {\Sigma_w}^{-1}\right)\mathcal{E}_W + \sqrt{N}{\Sigma_w}^{-1}\mathcal{E}_W$ we only need to consider the latter. By definition, we have

$$\hat{\beta} - \mathbf{H}^{\mathsf{T}}\beta = \mathbf{L}_{K}\left(\hat{\mu} - \mu^{H}\right), \quad \hat{\mu}_{-\beta} - \mu_{-\beta} = \mathbf{R}_{p}\left(\hat{\mu} - \mu^{H}\right)$$

where the matrices $\mathbf{L}_{K} \triangleq [\mathbf{I}_{K}, \mathbf{O}_{K \times (p+2)}] \in \mathbb{R}^{K \times (K+p+2)}$ and $\mathbf{R}_{p} \triangleq [\mathbf{O}_{(p+2) \times K}, \mathbf{I}_{p+2}] \in \mathbb{R}^{(p+2) \times (K+p+2)}$ take out first $K \times 1$ and last $(p+2) \times 1$ sub-vectors from a K + p + 2 vector, respectively. Therefore, we focus on $\sqrt{N}\mathbf{L}_{K}\Sigma_{w}^{-1}\mathcal{E}_{W}$ and $\sqrt{N}\mathbf{R}_{p}\Sigma_{w}^{-1}\mathcal{E}_{W}$. Note that $\mathbf{L}_{K}\Sigma_{w}^{-1} = \mathbf{L}_{K}$ and $\mathbf{R}_{p}\Sigma_{w}^{-1} = [\mathbf{O}_{(p+2) \times K} | \Sigma_{-u}^{-1}]$. Let $\mathbf{R} \triangleq [\mathbf{O}_{(p+2) \times K} | \Sigma_{-u}^{-1}]$ and let η be an appropriate-dimensional vector such that $\|\eta\| \leq 1$. First, since $\eta^{\mathsf{T}}\mathbf{L}_{K}\mathcal{E}_{W} = \frac{\sqrt{N}}{NT}\eta^{\mathsf{T}}(\mathbf{1}_{T} \otimes \hat{\mathbf{U}})^{\mathsf{T}}\mathcal{E}$ and $\|\mathbf{1}_{T} \otimes \hat{\mathbf{U}}\| = \sqrt{T}$, by similar arguments as in Lemma 8.4 we have

$$\mathbb{P}\left(\sqrt{N}\left|\eta^{\mathsf{T}}\mathbf{L}_{K}\mathcal{E}_{W}\right| > \upsilon\right) = \mathbb{P}\left[\left|\eta^{\mathsf{T}}\left(\mathbf{1}_{T}\otimes\hat{\mathbf{U}}\right)^{\mathsf{T}}\mathcal{E}\right| > T\upsilon\right] < 2\exp\left(-c\upsilon^{2}\right)$$

for any $\upsilon > 0$ hence $\hat{\beta} - \mathbf{H}^{\intercal}\beta = O_{\mathbb{P}}(1)$.

Next, let $\zeta_{N,t+1} \triangleq \frac{1}{\sqrt{NT}} \eta^{\mathsf{T}} \mathbf{R} \hat{\mathcal{W}}_t^{\mathsf{T}} \mathcal{E}_{t+1}$. If we define $\mathcal{F}_{Nt} = \sigma(\mathbf{A}, \epsilon_{is}, \mathbf{Z}_{is}; i \leq N, -\infty < s \leq t)$, then the pairs $\{\sum_{s=1}^t \zeta_{Ns}, \mathcal{F}_{Nt}\}$ constitute a martingale array for each $N, t \leq T$. Then, check:

(1) $\sum_{t} \mathbb{E}\left(\zeta_{N,t+1}^{2} \mathbb{1}_{\left\{\left|\zeta_{N,t+1}\right| > v\right\}} \middle| \mathcal{F}_{Nt}\right) = o_{\mathbb{P}}(1)$ (2) $\sum_{t} \mathbb{E}\left(\zeta_{N,t+1}^{2} \middle| \mathcal{F}_{Nt}\right) = \frac{\sigma^{2}}{T} \eta^{\mathsf{T}} \Sigma_{-u} \eta + o_{\mathbb{P}}(1)$

Proof of (1). We only need to check if $\sum_{t} \left(\frac{1}{N} \eta^{\mathsf{T}} \mathbf{R} \hat{\mathcal{W}}_{t}^{\mathsf{T}} \hat{\mathcal{W}}_{t} \mathbf{R}^{\mathsf{T}} \eta\right)^{2} = o_{\mathbb{P}}(1)$. First note that

$$\frac{1}{N}\mathbf{R}\hat{\mathcal{W}}_{t}^{\mathsf{T}}\hat{\mathcal{W}}_{t}\mathbf{R}^{\mathsf{T}} = \Sigma_{-u}^{-1} \begin{bmatrix} \frac{1}{N}\mathbf{y}_{t}^{\mathsf{T}}\mathbf{y}_{t} & \frac{1}{N}\mathbf{y}_{t}^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}\mathbf{y}_{t} & \frac{1}{N}\mathbf{y}_{t}^{\mathsf{T}}\mathcal{Z}_{t} \\ & \frac{1}{N}\mathbf{y}_{t}^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}^{2}\mathbf{y}_{t} & \frac{1}{N}\mathbf{y}_{t}^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}\mathbf{Z}_{t} \\ & & \frac{1}{N}\mathbf{Z}_{t}^{\mathsf{T}}\mathbf{Z}_{t} \end{bmatrix} \Sigma_{-u}^{-1}.$$

Since similar arguments can be used to show the convergence of each entry, take $\sum_{t} \left(\frac{1}{N} \mathbf{y}_t^{\mathsf{T}} \mathbf{y}_t\right)^2$

for example. Since \mathbf{y}_t follows $\mathcal{N}(\varphi, \Gamma)$, we have

$$\mathbb{E}^{*}\left[\left(\mathbf{y}_{t}^{\mathsf{T}}\mathbf{y}_{t}\right)^{2}\right] = \mathbb{E}^{*}\left[\left\|\Gamma^{1/2}\tilde{\mathbf{y}}_{t} + \varphi\right\|^{2}\right] \leq \left\|\Gamma\right\| \mathbb{E}^{*}\left(\left\|\tilde{\mathbf{y}}_{t} + \Gamma^{-1/2}\varphi\right\|^{2}\right) = O\left(N\right)$$

as. Therefore, $\frac{1}{N^2} \sum_t (\mathbf{y}_t^{\mathsf{T}} \mathbf{y}_t)^2 = o_{\mathbb{P}}(1)$. One can show that the rest are also $o_{\mathbb{P}}(1)$ similarly.

Proof of (2). Since

$$\sum_{t} \mathbb{E} \left(\zeta_{N,t+1}^{2} | \mathcal{F}_{Nt} \right) = \frac{\sigma^{2}}{NT^{2}} \eta^{\mathsf{T}} \mathbf{R} \hat{\mathcal{W}}^{\mathsf{T}} \hat{\mathcal{W}} \mathbf{R}^{\mathsf{T}} \eta = \frac{\sigma^{2}}{T} \eta^{\mathsf{T}} \Sigma_{-u}^{-1} \hat{\Sigma}_{-u} \Sigma_{-u}^{-1} \eta$$

and noting that $\hat{\Sigma}_{-u}^{-1} \Rightarrow {\Sigma_{-u}}^{-1}$ by Claim 1, we have (2).

Proof of Theorem 4.4 First recall that $\hat{\beta} - \mathbf{H}^{\mathsf{T}}\beta = \mathbf{L}_{K}\mathbf{D}_{N}(\hat{\mu} - \mu^{H})$. So, it suffices to show that for any \mathbf{A}_{K} in Theorem 4.4, we have

$$\frac{\mathbf{A}_{K}\mathbf{L}_{K}\hat{\Sigma}_{w}^{-1}}{\sqrt{N}}\hat{\mathcal{W}}^{\mathsf{T}}(\mathcal{W}^{H}-\hat{\mathcal{W}})\mu_{w}^{H}=o_{\mathbb{P}}(1)$$
(11)

$$\sqrt{N}\mathbf{A}_{K}\mathbf{L}_{K}\mathcal{E}_{W} = O_{\mathbb{P}}(1).$$
(12)

Again we can show $\frac{\mathbf{A}_{K}\mathbf{L}_{K}\Sigma_{w}^{-1}}{\sqrt{N}}\hat{W}^{\mathsf{T}}(\mathcal{W}^{H}-\hat{\mathcal{W}})\mu_{w}^{H}=o_{\mathbb{P}}(1)$ by the same logic of the proof of Theorem 4.2. Writing $\mathbf{f} \triangleq \frac{1}{\sqrt{N}}\hat{W}^{\mathsf{T}}(\mathcal{W}^{H}-\hat{\mathcal{W}})\mu_{w}^{H}$, for $\mathbf{u} \in \mathbb{R}^{m}$; $\|\mathbf{u}\| \leq 1$ we have $\mathbf{u}^{\mathsf{T}}\mathbf{A}_{K}\mathbf{L}_{K}\Sigma_{w}^{-1}\mathbf{f} = O(\|\mathbf{A}_{K}\mathbf{L}_{K}\mathbf{f}\|)$ as. Note that $\|\mathbf{A}_{K}\| = O(1)$, $\|\mathbf{L}_{K}\| = 1$, and $\|\mathbf{f}\| =$ $o_{\mathbb{P}}(1)$ provided that $\|\beta\| = O(1)$. We have $\sqrt{N}\eta^{\mathsf{T}}\mathbf{A}_{K}\mathbf{L}_{K}\Sigma_{w}^{-1}\mathcal{E}_{W} = \sqrt{N}\eta^{\mathsf{T}}\mathbf{A}_{K}\mathbf{L}_{K}\mathcal{E}_{W} =$ $\frac{1}{\sqrt{N}T}\eta^{\mathsf{T}}\mathbf{A}_{K}\left(\mathbf{1}_{T}\otimes\hat{\mathbf{U}}\right)^{\mathsf{T}}\mathcal{E}$ and $\|\left(\mathbf{1}_{T}\otimes\hat{\mathbf{U}}\right)\mathbf{A}_{K}^{\mathsf{T}}\| = \sqrt{T}\|\mathbf{A}_{K}^{\mathsf{T}}\| = O(1)$. Therefore, by the same logic in the proof of Theorem 4.3, we have $\sqrt{N}\eta^{\mathsf{T}}\mathbf{A}_{K}\mathbf{L}_{K}\mathcal{E}_{W} = O_{\mathbb{P}}(1)$.

9 Proofs for Sections 3.3 & 4.3

Here, we present the proof for the model stationarity and the asymptotic properties of estimators for AMNAR. First, we provide the theoretical background for the estimation of latent variables **X**. Let $\tilde{\mathbf{Q}} \triangleq [\mathbf{Q} \mid \mathbf{1}_N]$. For simplicity, $l'(\chi_{ij}), l''(\chi_{ij}), l'''(\chi_{ij})$ denote

the 1st, 2nd, 3rd-order derivatives of the log-likelihood function $l(\chi_{ij}; a_{ij})$ with regard to χ_{ij} , respectively. Fix a constant $s \in (0, 1/2)$. Following assumptions are analogous to the assumptions made in the section 2.2 of Li et al. (2023).

I. X is contained in the constrained parameter space

$$\Xi \triangleq \left\{ \mathbf{X} \, ; \, \mathbf{Q}^{\mathsf{T}} \mathbf{1}_{N} = \mathbf{o}_{K}, \, \mathbf{Q}^{\mathsf{T}} \mathbf{Q} \text{ is diagonal}, \, \left\| \mathbf{X} \right\|_{2,\infty} = O(1) \text{ as } N \to \infty \right\}.$$

- II. There exist a positive definite $K \times K$ matrix Ω_q and some constant $\nu > 0$ such that $\frac{1}{N} \mathbf{Q}^{\mathsf{T}} \mathbf{Q} \to \Omega_q$ as $N \to \infty$ where Ω_q is a diagonal matrix with unique eigenvalues and $\frac{1}{N} \mathbf{v}^{\mathsf{T}} \mathbf{v} \to \nu$.
- III. $l'''(\cdot)$ exists within Ξ . Furthermore, there exists $0 < b_L < b_U$ such that $b_L \leq -l''(\cdot) \leq b_U$ and $|l'''(\cdot)| \leq b_U$ within Ξ .
- IV. There exist t > 0 and u > 0 such that for all $x \ge 0$, $P(|l_{ij}'| > x) \le \exp(-(x/t)^u)$.
- V. For any $1 \leq i \leq N$, there exist Σ_i such that

$$\frac{-1}{N}\sum_{j:j\neq i}\left(l''\circ\sigma\right)\left(\chi_{ij}\right)\tilde{\mathbf{q}}_{j}\tilde{\mathbf{q}}_{j}\stackrel{\mathsf{T}}{\to}\Sigma_{i}.$$

For an integer m and any m node indices $\mathcal{I} = (i_1, i_2, ..., i_m)$, there exists $\mathbf{S}_{\mathcal{I}}$ such that $\frac{1}{\sqrt{N}} ([S(\mathbf{x})]_{i_1}, [S(\mathbf{x})]_{i_2}, ..., [S(\mathbf{x})]_{i_m}) \to \mathcal{N}(0, \mathbf{S}_{\mathcal{I}})$, where $S(\mathbf{x}) = \frac{\partial L}{\partial \mathbf{x}}|_{\mathbf{x}=\mathbf{x}}$ is the score vector evaluated at the true parameters $\mathbf{x} \triangleq \operatorname{vec}(\mathbf{X}^{\intercal})$, and $[S(\mathbf{x})]_i$ denotes the subvector of $S(\mathbf{x})$ corresponding to all latent parameters associated with node i, i.e., \mathbf{x}_i .

Assumption I is often posed for the sake of theoretical analysis (Ma et al., 2020). Conditions on $\mathbf{Q}^{\intercal}\mathbf{1}_{N}$ and $\mathbf{Q}^{\intercal}\mathbf{Q}$ in Assumptions I and II ensure the identifiability of \mathbf{v} and give regular conditions on the asymptotic behavior of the covariance structure of latent positions \mathbf{Q} . However, the diagonality assumption can be relaxed as Li et al. (2023) noted. Assumption III requires $\sigma(\cdot)$ to be a smooth function and l to be a concave log-likelihood function which constitutes a widely accepted class of link functions such as logit links. Assumptions IV and V ensure the fast decadence of tail density and the asymptotic distributions of maximum likelihood estimator (Li et al., 2023).

Remark 1. In assumption II, we put an additional assumption as $\|\mathbf{v}\|^2 / N = \nu + o(1)$ so that AMNAR estimators can obtain a non-singular asymptotic precision matrix. In ENAR model, by the definition of RDPG, we have $\|\mathbf{U}_{\mathbf{P}}\|_{2,\infty} \leq \frac{\sqrt{N}\rho_N}{\pi_N}$ for all N. Recall that we assumed that $N\rho_N = \omega (\log N)$ and $\pi_N \gg \sqrt{TN\rho_N}$ in Theorem 4.1. Therefore, we have $\|\mathbf{U}_{\mathbf{P}}\|_{2,\infty} = o\left(\sqrt{\rho_N/T}\right)$ so if $\rho_N = \Theta(N^{\epsilon-1})$ for $\epsilon \in (0,1)$, for example, we have $\|\mathbf{U}_{\mathbf{P}}\|_{2,\infty} = o(N^{(\epsilon-1)/2}T^{-1/2})$ when $\min(N,T) \to \infty$. Similarly, we have $\|\mathbf{r}\mathbf{X}\|_{2,\infty} = \Theta(N^{-s}T^{-1/2})$ by the multiplier $r = \frac{1}{N^s\sqrt{T}}$.

9.1 Stationarity

Proof of Theorem 3.4. We have $\mathbf{y}_t = \sum_{j=0}^{m-1} \mathbf{G}^j (r \mathbf{X} \beta + \tilde{\mathcal{E}}_{t-j}) + \mathbf{G}^m \mathbf{y}_{t-m}$ hence

$$\mathbf{y}_t = \lim_{m \to \infty} \mathbf{y}_t = \sum_{j=0}^{\infty} \mathbf{G}^j (r \mathbf{X} \beta + \tilde{\mathcal{E}}_{t-j}).$$

First we have $\|r\mathbf{X}\beta\|_{\infty} = \max_{i=1,\dots,N} r |\mathbf{x}_i^{\mathsf{T}}\beta| \leq r \|\mathbf{X}\|_{2,\infty} \|\beta\|$ hence $|r\mathbf{X}\beta|_e \precsim r\mathbf{1}_N$ independently with j. So, we have $\mathbb{E} \left| r\mathbf{X}\beta + \tilde{\mathcal{E}}_{t-j} \right|_e \precsim C \cdot \mathbf{1}_N$ for $C = o(1) + \mathbb{E} |\mathbf{z}_1^{\mathsf{T}}\gamma| + \mathbb{E} |\epsilon_{11}|$. Therefore, we have

$$\mathbb{E}^* |\mathbf{w}_N^{\mathsf{T}} \mathbf{y}_t| \le \|\mathbf{w}_N\|_1 \mathbb{E}^* \|\mathbf{y}_t\|_{\infty} \le \sum_{i=1}^{\infty} |w_i| \sum_{j=0}^{\infty} \mathbb{E}^* \left\| |\mathbf{G}|_e^j \left| r \mathbf{X}\beta + \tilde{\mathcal{E}}_{t-j} \right|_e \right\|_{\infty}$$
$$\lesssim \sum_{i=1}^{\infty} |w_i| \sum_{j=0}^{\infty} (|\theta| + |\alpha|)^j + o(1)$$

implying that $\lim_{N\to\infty} \mathbf{w}_N^{\mathsf{T}} \mathbf{y}_t$ exists almost surely. Next, assume that $\bar{\mathbf{y}}_t$ is another strictly stationary solution with a finite first moment. Then, $\mathbb{E} |\bar{\mathbf{y}}_t|_e \preceq \mathbf{1}_N$. We have

$$\mathbb{E}^* \left| \mathbf{w}_N^{\mathsf{T}} (\mathbf{y}_t - \bar{\mathbf{y}}_t) \right| = \mathbb{E}^* \left| \sum_{j=m}^{\infty} \mathbf{w}_N^{\mathsf{T}} \mathbf{G}^j (r \mathbf{X} \beta + \tilde{\mathcal{E}}_{t-j}) - \mathbf{w}_N^{\mathsf{T}} \mathbf{G}^m \bar{\mathbf{y}}_{t-m} \right|$$

which is less than

$$\sum_{i=1}^{\infty} |w_i| \sum_{j=m}^{\infty} \left\{ (|\alpha| + |\theta|)^j (1 + o(1)) + (|\alpha| + |\theta|)^m \right\} + o(1)$$

up to a constant multiplication for any $\omega \in \mathcal{M}$. By growing $m \to \infty$, we again conclude that $\mathbf{y}_t = \bar{\mathbf{y}}_t$ as.

9.2 Consistency

Proof of Theorem 4.5. Recalling $r = \frac{1}{N^s \sqrt{T}} = o(1)$ as $\max(N, T) \to \infty$, let $\tilde{\mathbf{D}}_{NT} \triangleq \sqrt{NT} \operatorname{diag}(r\mathbf{I}_{K+1}, \mathbf{I}_{p+2})$. For $\hat{\Omega}_m \triangleq \tilde{\mathbf{D}}_{NT}^{-1} \hat{\mathbf{M}}^{\dagger} \hat{\mathbf{M}} \tilde{\mathbf{D}}_{NT}^{-1}$ and $\mathcal{E}_{\mathbf{M}} \triangleq \tilde{\mathbf{D}}_{NT}^{-1} \hat{\mathbf{M}}^{\dagger} \mathcal{E}$, we have

$$\hat{\mu}_{m} = \tilde{\mathbf{D}}_{NT}^{-1} \left(\hat{\Omega}_{m} \right)^{-1} \tilde{\mathbf{D}}_{NT}^{-1} \hat{\mathbf{M}}^{\mathsf{T}} \left\{ \left(\mathbf{M} - \hat{\mathbf{M}} \right) \mu_{m} + \hat{\mathbf{M}} \mu_{m} + \mathcal{E} \right\}$$
$$= \tilde{\mathbf{D}}_{NT}^{-1} \hat{\Omega}_{m}^{-1} \left\{ \tilde{\mathbf{D}}_{NT}^{-1} \hat{\mathbf{M}}^{\mathsf{T}} \left(\mathbf{M} - \hat{\mathbf{M}} \right) \mu_{m} + \mathcal{E}_{\mathbf{M}} \right\} + \mu_{m}$$

hence

$$\hat{\Omega}_m \tilde{\mathbf{D}}_{NT} (\hat{\mu}_m - \mu_m) = \tilde{\mathbf{D}}_{NT}^{-1} \hat{\mathbf{M}}^{\mathsf{T}} (\mathbf{M} - \hat{\mathbf{M}}) \mu_m + \mathcal{E}_{\mathbf{M}}.$$

Claim 1. $\hat{\Omega}_m \Rightarrow \Omega_m$.

Proof. As consequences of Lemma 9.2, we have

$$\hat{\Omega}_{m} = \frac{1}{NT} \sum_{t} \begin{vmatrix} \hat{\mathbf{X}}^{\mathsf{T}} \hat{\mathbf{X}} & \hat{\mathbf{X}}^{\mathsf{T}} \mathbf{y}_{t} & \hat{\mathbf{X}}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \mathbf{y}_{t} & \hat{\mathbf{X}}^{\mathsf{T}} \mathbf{Z}_{t} \\ & \mathbf{y}_{t}^{\mathsf{T}} \mathbf{y}_{t} & \mathbf{y}_{t}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \mathbf{y}_{t} & \mathbf{y}_{t}^{\mathsf{T}} \mathbf{Z}_{t} \\ & \mathbf{y}_{t}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}}^{2} \mathbf{y}_{t} & \mathbf{y}_{t}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \mathbf{Z}_{t} \\ & & \mathbf{Z}_{t}^{\mathsf{T}} \mathbf{Z}_{t} \end{vmatrix} \\ \Rightarrow \lim_{N,T \to \infty} \begin{bmatrix} \frac{1}{N} \mathbf{X}^{\mathsf{T}} \mathbf{X} & \frac{1}{N} \mathbf{X}^{\mathsf{T}} \psi & \frac{1}{N} \mathbf{X}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \psi & \mathbf{O}_{(K+1) \times p} \\ & \frac{1}{N} \psi^{\mathsf{T}} \psi + \tau_{2} & \frac{1}{N} \psi^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \psi + \tau_{23} & \mathbf{O}_{p}^{\mathsf{T}} \\ & & \frac{1}{N} \psi^{\mathsf{T}} \mathcal{L}_{\mathbf{A}}^{2} \psi + \tau_{3} & \mathbf{O}_{p}^{\mathsf{T}} \\ & & & \Sigma_{z} \end{bmatrix}$$

.

By assumptions in section 9, we have $\|\mathbf{v}\|^2 = O(N)$ and $\frac{1}{N}\mathbf{Q}^{\mathsf{T}}\mathbf{Q} \to \Omega_q$ hence we can infer that $\|\mathbf{Q}^{\mathsf{T}}\mathbf{Q}\| = O(N)$. Therefore, $|\mathbf{h}^{\mathsf{T}}\mathbf{Q}^{\mathsf{T}}\mathbf{v}| \leq \|\mathbf{Q}\| \|\mathbf{v}\| = O(N)$ for $\|\mathbf{h}\| \leq 1$ and $\frac{1}{N}\mathbf{X}^{\mathsf{T}}\mathbf{X} \to \Omega_x$. Since $\|\mathbf{X}\|_{2,\infty} = O(1)$, we have $\|\mathbf{X}\| = O(\sqrt{N})$ hence $\|\psi\| \leq r \|(\mathbf{I}_N - \mathbf{G})^{-1}\| \|\mathbf{X}\beta\| \leq r \|\mathbf{X}\| \|\beta\| / (1-g) = O(r\sqrt{N})$ as. Therefore, $\frac{1}{N}\mathbf{h}_1^{\mathsf{T}}\mathbf{X}^{\mathsf{T}}\psi$ and $\frac{1}{N}\mathbf{h}_1^{\mathsf{T}}\mathbf{X}^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}\psi$ are O(r) and $\frac{1}{N}\psi^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}\psi$, and $\frac{1}{N}\psi^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}^{2}\psi$ are $O(r^2)$ as. for all $\mathbf{h}_1 \in \mathbb{R}^{K+1}$ such that $\|\mathbf{h}_1\| \leq 1$. This imply that they are all o(1).

Claim 2. $\tilde{\mathbf{D}}_{NT}^{-1}\hat{\mathbf{M}}^{\mathsf{T}}(\mathbf{M}-\hat{\mathbf{M}})\mu_m = o_{\mathbb{P}}(1).$

Proof.

 $\hat{\mathbf{M}}$ is different from \mathbf{M} by $\hat{\mathbf{X}}$ only, so we have

$$\mathbf{h}^{\mathsf{T}} \tilde{\mathbf{D}}_{NT}^{-1} \hat{\mathbf{M}}^{\mathsf{T}} \left(\hat{\mathbf{M}} - \mathbf{M} \right) \mu_{m} = \mathbf{h}^{\mathsf{T}} \begin{bmatrix} \frac{r}{\sqrt{NT}} \hat{\mathbf{X}}^{\mathsf{T}} \\ \frac{r}{\sqrt{NT}} \sum_{t} \mathbf{y}_{t}^{\mathsf{T}} \\ \frac{r}{\sqrt{NT}} \sum_{t} \mathbf{y}_{t}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \\ \frac{r}{\sqrt{NT}} \sum_{t} \mathbf{Z}_{t}^{\mathsf{T}} \end{bmatrix} [\hat{\mathbf{X}} - \mathbf{X} \mid \mathbf{O}_{N \times (p+2)}] \mu_{m}$$

Then, by the statements 10.–13. of Lemma 9.1, we have the conclusion by Cramér–Wold.

Claim 3. $\mathcal{E}_{\mathbf{M}} \Rightarrow \mathcal{N}(\mathbf{o}_{K+p+3}, \sigma^2 \Omega_m).$

Proof. We show that for any $\eta \in \mathbb{R}^{K+p+3}$ such that $\|\eta\| \leq 1$, it holds that $\eta^{\mathsf{T}} \mathcal{E}_{\mathbf{M}} \Rightarrow \mathcal{N}(0, \sigma^2 \eta^{\mathsf{T}} \Omega_m \eta)$. Denoting $\xi_{N,t+1} \triangleq \eta^{\mathsf{T}} \tilde{\mathbf{D}}_{NT}^{-1} \hat{\mathbf{M}}_t^{\mathsf{T}} \mathcal{E}_{t+1}$ and $\mathcal{F}_{Nt} = \sigma(\mathbf{A}, \epsilon_{is}, \mathbf{Z}_{is}; i \leq N, -\infty < s \leq t)$, $\{\sum_{s=1}^t \xi_{Ns}, \mathcal{F}_{Nt}\}$ constitutes a martingale array for each $N, t \leq T$. Applying Corollary 3.1 of Hall and Heyde (2014), we check:

(1)
$$\sum_{t} \mathbb{E}\left(\xi_{N,t+1}^{2} \mathbb{1}_{\left\{\left|\xi_{N,t+1}\right| > \upsilon\right\}} | \mathcal{F}_{Nt}\right) = o_{\mathbb{P}}(1)$$

(2) $\sum_{t} \mathbb{E}\left(\xi_{N,t+1}^{2} | \mathcal{F}_{Nt}\right) = \sigma^{2} \eta^{\intercal} \Omega_{m} \eta + o_{\mathbb{P}}(1).$

Proof of (1). First, we have

$$\sum_{t=0}^{T-1} \mathbb{E}\left(\xi_{N,t+1}^2 \mathbb{1}_{\left\{\left|\xi_{N,t+1}\right| > \upsilon\right\}} \left|\mathcal{F}_{Nt}\right) \le \upsilon^{-2} \sum_{t} \mathbb{E}\left(\xi_{N,t+1}^4 \left|\mathcal{F}_{Nt}\right)\right).$$

One can easily verify that

$$\mathbb{E}\left(\xi_{N,t+1}^{4}|\mathcal{F}_{Nt}\right) \lesssim \sigma^{4}\left(\eta^{\mathsf{T}}\tilde{\mathbf{D}}_{NT}^{-1}\hat{\mathbf{M}}_{t}^{\mathsf{T}}\hat{\mathbf{M}}_{t}\tilde{\mathbf{D}}_{NT}^{-1}\eta\right)^{2}.$$

First, we have

$$\begin{split} \tilde{\mathbf{D}}_{NT}^{-1} \hat{\mathbf{M}}_{t}^{\mathsf{T}} \hat{\mathbf{M}}_{t} \tilde{\mathbf{D}}_{NT}^{-1} \\ &= \frac{1}{NT} \begin{bmatrix} \hat{\mathbf{X}}^{\mathsf{T}} \hat{\mathbf{X}} & \hat{\mathbf{X}}^{\mathsf{T}} \mathbf{y}_{t} & \hat{\mathbf{X}}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \mathbf{y}_{t} & \hat{\mathbf{X}}^{\mathsf{T}} \mathbf{Z}_{t} \\ & \mathbf{y}_{t}^{\mathsf{T}} \mathbf{y}_{t} & \mathbf{y}_{t}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \mathbf{y}_{t} & \mathbf{y}_{t}^{\mathsf{T}} \mathbf{Z}_{t} \\ & & \mathbf{y}_{t}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}}^{2} \mathbf{y}_{t} & \mathbf{y}_{t}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \mathbf{Z}_{t} \\ & & & \mathbf{Z}_{t}^{\mathsf{T}} \mathbf{Z}_{t} \end{bmatrix} \end{split}$$

Since similar arguments can be used to show the convergence of each entry, take

$$\sum_{t} \left(\frac{1}{NT} \eta_1^{\mathsf{T}} \hat{\mathbf{X}}^{\mathsf{T}} \mathbf{y}_t \right)^2 = \frac{1}{N^2 T^2} \sum_{t} \eta_1^{\mathsf{T}} \hat{\mathbf{X}}^{\mathsf{T}} \mathbf{y}_t \mathbf{y}_t^{\mathsf{T}} \hat{\mathbf{X}} \eta_1$$

for example, where $\eta_1 \in \mathbb{R}^K$; $\|\eta_1\| \leq 1$. Then, we have

$$\begin{split} \sum_{t} \eta_{1}^{\mathsf{T}} \hat{\mathbf{X}}^{\mathsf{T}} \mathbf{y}_{t} \mathbf{y}_{t}^{\mathsf{T}} \hat{\mathbf{X}} \eta_{1} &= \mathbf{y}^{\mathsf{T}} \left\{ \mathbf{I}_{T} \otimes \left(\hat{\mathbf{X}} \eta_{1} \eta_{1}^{\mathsf{T}} \hat{\mathbf{X}}^{\mathsf{T}} \right) \right\} \mathbf{y} \\ &= \left(\Gamma_{\mathbf{y}}^{1/2} \tilde{\mathbf{y}} + \Psi \right)^{\mathsf{T}} \left[\mathbf{I}_{T} \otimes \left\{ \left(\Delta + \mathbf{X} \right) \eta_{1} \eta_{1}^{\mathsf{T}} \left(\Delta + \mathbf{X} \right)^{\mathsf{T}} \right\} \right] \left(\Gamma_{\mathbf{y}}^{1/2} \tilde{\mathbf{y}} + \Psi \right). \end{split}$$

Denoting $\hat{\mathbf{X}} - \mathbf{X}$ by Δ , recall that $\|\Delta\| = O(1)$ whp. and $\|\Psi\| = O(r\sqrt{NT})$. Letting $\tilde{\mathbf{S}} \triangleq \Gamma_{\mathbf{y}}^{1/2} \{ \mathbf{I}_T \otimes (\mathbf{X}\eta_1\eta_1^{\mathsf{T}}\mathbf{X}^{\mathsf{T}}) \} \Gamma_{\mathbf{y}}^{1/2}$, we have $\|\tilde{\mathbf{S}}\| = \|\Gamma^{1/2}\mathbf{X}\eta_1\eta_1^{\mathsf{T}}\mathbf{X}^{\mathsf{T}}\Gamma^{1/2}\| \le \|\Gamma\| \|\mathbf{X}\|^2$ hence is O(N) as. and $\|\tilde{\mathbf{S}}\|_F^2 = T \operatorname{tr} \{ (\mathbf{X}\eta_1\eta_1^{\mathsf{T}}\mathbf{X}^{\mathsf{T}}\Gamma)^2 \} \le T \operatorname{tr} (\mathbf{X}\eta_1\eta_1^{\mathsf{T}}\mathbf{X}^{\mathsf{T}})^2 \|\Gamma\|^2 \le T \|\mathbf{X}\|^4 \|\Gamma\| = O(N^2T)$ as. Note that

$$\mathbb{E}\left(\tilde{\mathbf{y}}^{\mathsf{T}}\tilde{\mathbf{S}}\tilde{\mathbf{y}}\right) = \mathbb{E}\operatorname{tr}\left(\tilde{\mathbf{S}}\tilde{\mathbf{y}}\tilde{\mathbf{y}}^{\mathsf{T}}\right) = \operatorname{tr}\left\{\mathbb{E}(\tilde{\mathbf{S}})\mathbb{E}\left(\tilde{\mathbf{y}}\tilde{\mathbf{y}}^{\mathsf{T}}\right)\right\} = T\operatorname{tr}\left(\mathbf{X}\eta_{1}\eta_{1}^{\mathsf{T}}\mathbf{X}^{\mathsf{T}}\mathbb{E}\Gamma\right)$$

which is bounded above by $T \operatorname{tr} (\mathbf{X}\eta_1\eta_1^{\mathsf{T}}\mathbf{X}^{\mathsf{T}}) \|\mathbb{E}\Gamma\| \leq T \|\mathbf{X}\|^2 \mathbb{E} \|\Gamma\| = O(T \|\mathbf{X}\|^2) = O(NT).$ Applying Theorem 8.3, we get

$$\mathbb{P}\left[\left|\tilde{\mathbf{y}}^{\mathsf{T}}\tilde{\mathbf{S}}\tilde{\mathbf{y}} - \mathbb{E}\left(\tilde{\mathbf{y}}^{\mathsf{T}}\tilde{\mathbf{S}}\tilde{\mathbf{y}}\right)\right| > N^{2}T^{2}\upsilon\right] < 2\exp\left(-c\min\left\{\frac{N^{4}T^{4}\upsilon^{2}}{N^{2}T}, \frac{N^{2}T^{2}\upsilon}{N}\right\}\right)$$

and note that $\mathbb{E}\left(\tilde{\mathbf{y}}^{\mathsf{T}}\tilde{\mathbf{S}}\tilde{\mathbf{y}}\right) = o\left(N^{2}T^{2}\right)$ by dominated convergence. Similarly, we note that $\Psi^{\mathsf{T}}\left\{\mathbf{I}_{T} \otimes \left(\mathbf{X}\eta_{1}\eta_{1}^{\mathsf{T}}\mathbf{X}^{\mathsf{T}}\right)\right\}\Psi = O(r^{2}N^{2}T)$ as. hence $\sum_{t}\eta_{1}^{\mathsf{T}}\hat{\mathbf{X}}^{\mathsf{T}}\mathbf{y}_{t}\mathbf{y}_{t}^{\mathsf{T}}\hat{\mathbf{X}}\eta_{1} = o_{\mathbb{P}}(N^{2}T^{2})$ in conclusion. The rest of the proof uses the same logic as well so is omitted here. Noting that both \mathbf{y} and \mathbf{Z}_{t} have finite fourth-order moments, one can show that the rest terms of $\sum_{t}\left(\eta^{\mathsf{T}}\tilde{\mathbf{D}}_{NT}^{-1}\hat{\mathbf{M}}_{t}^{\mathsf{T}}\hat{\mathbf{M}}_{t}\tilde{\mathbf{D}}_{NT}^{-1}\eta\right)^{2}$ are also $o_{\mathbb{P}}(N^{2}T^{2})$ similarly. **Proof of (2).** Since $\hat{\Omega}_{m} = \Omega_{m} + o_{\mathbb{P}}(1)$, we directly have (2).

9.3 Asymptotic results

Lemma 9.1. Let $\mathbf{h}_i \in \mathbb{R}^{j_i}$ for $j_1 = K + 1$, $j_2 = p$ be real vectors such that $\|\mathbf{h}_i\| \leq 1$. Then under the conditions of Theorem 4.5, there exist $N_0, T_0 > 0$ and $v_0(N_0, T_0) \in (0, 1)$ such that for all $v \in (0, v_0)$ and $N > N_0, T > T_0$, the statements **0.-13.** hold.

$$\begin{array}{l} \mathbf{0.} \ \mathbb{P}\left[\frac{1}{N}\left|\mathbf{h}_{1}^{\mathsf{T}}\left(\hat{\mathbf{X}}^{\mathsf{T}}\hat{\mathbf{X}}-\mathbf{X}^{\mathsf{T}}\mathbf{X}\right)\mathbf{h}_{1}\right| > \upsilon\right] < \epsilon \\ \mathbf{1.} \ \mathbb{P}\left(\frac{1}{NT}\left|\mathbf{h}_{1}^{\mathsf{T}}\sum_{l}\left(\hat{\mathbf{X}}^{\mathsf{T}}\mathbf{y}_{l}-\mathbf{X}^{\mathsf{T}}\psi\right)\right| > \upsilon\right) < 2e^{-c_{1}N^{2}T\upsilon^{2}} + 2e^{-c_{2}NT\upsilon^{2}} + \epsilon \\ \mathbf{2.} \ \mathbb{P}\left[\frac{1}{NT}\left|\mathbf{h}_{1}^{\mathsf{T}}\sum_{l}\left(\hat{\mathbf{X}}^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}\mathbf{y}_{l}-\mathbf{X}^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}\psi\right)\right| > \upsilon\right] < 2e^{-c_{1}N^{2}T\upsilon^{2}} + 2e^{-c_{2}NT\upsilon^{2}} + \epsilon \\ \mathbf{3.} \ \mathbb{P}\left(\frac{1}{NT}\left|\sum_{t}\mathbf{h}_{1}^{\mathsf{T}}\hat{\mathbf{X}}^{\mathsf{T}}\mathbf{Z}_{t}\mathbf{h}_{2}\right| > \upsilon\right) < 2e^{-c_{1}N^{2}T\upsilon^{2}} + 2e^{-c_{2}NT\upsilon^{2}} + \epsilon \\ \mathbf{4.} \ \mathbb{P}\left[\frac{1}{NT}\left|\sum_{t}\left(\mathbf{y}_{t}^{\mathsf{T}}\mathbf{y}_{t}-\operatorname{tr}\mathbb{E}\Gamma-\psi^{\mathsf{T}}\psi\right)\right| > \upsilon\right] < 2e^{-c_{1}NT\upsilon^{2}} + 2e^{-c_{2}NT\upsilon^{2}/r^{2}} \\ \mathbf{5.} \ \mathbb{P}\left[\frac{1}{NT}\left|\sum_{t}\left\{\mathbf{y}_{t}^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}\mathbf{y}_{t}-\psi^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}\psi-\operatorname{tr}\mathbb{E}\left(\mathcal{L}_{\mathbf{A}}\Gamma\right)\right\}\right| > \upsilon\right] < 2e^{-c_{1}NT\upsilon^{2}} + 2e^{-c_{2}NT\upsilon^{2}/r^{2}} \\ \mathbf{6.} \ \mathbb{P}\left(\frac{1}{NT}\left|\sum_{t}\mathbf{y}_{t}^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}\mathbf{y}_{t}-\psi^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}\psi-\operatorname{tr}\mathbb{E}\left(\mathcal{L}_{\mathbf{A}}\Gamma\right)\right\}\right| > \upsilon\right] < 2e^{-c_{1}NT\upsilon^{2}} + 2e^{-c_{2}NT\upsilon^{2}/r^{2}} \\ \mathbf{6.} \ \mathbb{P}\left(\frac{1}{NT}\left|\sum_{t}\mathbf{y}_{t}^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}\mathbf{y}_{t}-\psi^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}^{2}\psi-\operatorname{tr}\mathbb{E}\left(\mathcal{L}_{\mathbf{A}}\Gamma\right)\right\}\right| > \upsilon\right] < 2e^{-c_{1}NT\upsilon^{2}} + 2e^{-c_{2}NT\upsilon^{2}/r^{2}} \\ \mathbf{6.} \ \mathbb{P}\left[\frac{1}{NT}\left|\sum_{t}\mathbf{y}_{t}^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}^{2}\mathbf{y}_{t}-\psi^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}^{2}\psi-\operatorname{tr}\mathbb{E}\left(\mathcal{L}_{\mathbf{A}}^{2}\Gamma\right)\right\}\right| > \upsilon\right] < 2e^{-c_{1}NT\upsilon^{2}} + 2e^{-c_{2}NT\upsilon^{2}/r^{2}} \\ \mathbf{6.} \ \mathbb{P}\left[\frac{1}{NT}\left|\sum_{t}\mathbf{y}_{t}^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}\mathbf{Z}_{t}\mathbf{h}_{2}\right| > \upsilon\right) < 2e^{-c_{1}NT\upsilon^{2}/r^{2}} + \epsilon \\ \mathbf{7.} \ \mathbb{P}\left[\frac{1}{NT}\left|\sum_{t}\mathbf{y}_{t}\mathbf{y}_{t}^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}\mathbf{Z}_{t}\mathbf{h}_{2}\right| > \upsilon\right) < 2e^{-c_{1}NT\upsilon^{2}/r^{2}} + \epsilon \\ \mathbf{9.} \ \mathbb{P}\left[\frac{1}{NT}\left|\sum_{t}\mathbf{y}_{t}\mathbf{y}_{t}^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}\mathbf{Z}_{t}\mathbf{h}\right| > \upsilon\right] < 2e^{-c_{1}NT\upsilon^{2}} \\ \mathbf{10.} \ \mathbb{P}\left[\frac{r}{\sqrt{NT}}\left|\mathbf{h}_{1}^{\mathsf{T}}\hat{\mathbf{X}}\left(\hat{\mathbf{X}-\mathbf{X}\right)\beta\right| > \upsilon\right] < \epsilon \\ \mathbf{11.} \ \mathbb{P}\left[\frac{r}{\sqrt{NT}}\left|\sum_{t}\mathbf{y}_{t}\mathbf{y}_{t}^{\mathsf{T}}\hat{\mathbf{A}}\left(\hat{\mathbf{X}-\mathbf{X}\right)\beta\right| > \upsilon\right] < 2e^{-c_{1}N\upsilon^{2}/r^{2}} \\ \mathbf{12.} \ \mathbb{P}\left[\frac{r}{\sqrt{NT}}\left|\sum_{t}\mathbf{y}_{t}\mathbf{y}_{t}\mathcal{L}_{\mathbf{A}}\left(\hat{\mathbf{X}-\mathbf{X}\right)\beta\right| > \upsilon\right] < 2e^{-c_{1}N\upsilon^{2}/r^{2}} + \epsilon \\$$

13.
$$\mathbb{P}\left[\frac{r}{\sqrt{NT}}\left|\mathbf{h}_{2}^{\mathsf{T}}\sum_{t}\mathbf{Z}_{t}^{\mathsf{T}}\left(\hat{\mathbf{X}}-\mathbf{X}\right)\beta\right| > \upsilon\right] < 2e^{-cN\upsilon^{2}/r^{2}} + \epsilon$$

where $c_1, \ldots, c_3 > 0$ are constants.

Proof of 0. First note that $\hat{\mathbf{X}}^{\mathsf{T}}\hat{\mathbf{X}} - \mathbf{X}^{\mathsf{T}}\mathbf{X} = \Delta^{\mathsf{T}}(\Delta + \mathbf{X}) + \mathbf{X}^{\mathsf{T}}\Delta$ and $\mathbf{h}_{1}^{\mathsf{T}}\Delta^{\mathsf{T}}\Delta\mathbf{h}_{1} \leq \|\Delta\|_{F}^{2} \leq \|\Delta\|_{F}^{2} = O_{\mathbb{P}}(1)$. Also, $\|\Delta^{\mathsf{T}}\mathbf{X}\| = O_{\mathbb{P}}(\|\mathbf{X}\|)$. Therefore,

$$\mathbb{P}\left[\frac{1}{N}\left|\mathbf{h}_{1}^{\mathsf{T}}\left(\hat{\mathbf{X}}^{\mathsf{T}}\hat{\mathbf{X}}-\mathbf{X}^{\mathsf{T}}\mathbf{X}\right)\mathbf{h}_{1}\right|>\upsilon\right]<\epsilon$$

provided that $Nv = \omega(||\mathbf{X}||)$. Since $||\mathbf{X}|| / N = O(1/\sqrt{N})$ and $1/\sqrt{N} = o(1)$, we have the conclusion.

Proof of 1. Note that

$$\sum_{t=1}^{T} \mathbf{h}_{1}^{\mathsf{T}} \left(\hat{\mathbf{X}}^{\mathsf{T}} \mathbf{y}_{t} - \mathbf{X}^{\mathsf{T}} \psi \right) = \mathbf{h}_{1}^{\mathsf{T}} \left(\mathbf{1}_{T} \otimes \Delta \right)^{\mathsf{T}} \Gamma_{\mathbf{y}}^{1/2} \tilde{\mathbf{y}} + \mathbf{h}_{1}^{\mathsf{T}} \left(\mathbf{1}_{T} \otimes \mathbf{X} \right)^{\mathsf{T}} \Gamma_{\mathbf{y}}^{1/2} \tilde{\mathbf{y}} + T \mathbf{h}_{1}^{\mathsf{T}} \Delta^{\mathsf{T}} \psi = \mathbf{h}_{1}^{\mathsf{T}} \hat{\mathbf{S}}^{\mathsf{T}} \tilde{\mathbf{y}} + \mathbf{h}_{1}^{\mathsf{T}} \tilde{\mathbf{S}}^{\mathsf{T}} \tilde{\mathbf{y}} + T \mathbf{h}_{1}^{\mathsf{T}} \Delta^{\mathsf{T}} \psi.$$

where $\hat{\mathbf{S}} \triangleq \Gamma_{\mathbf{y}}^{1/2} (\mathbf{1}_T \otimes \Delta)$ and $\tilde{\mathbf{S}} \triangleq \Gamma_{\mathbf{y}}^{1/2} (\mathbf{1}_T \otimes \mathbf{X})$. Since $\|\hat{\mathbf{S}}\mathbf{h}_1\| = O_{\mathbb{P}} (\sqrt{T})$, by Theorem 8.2 we have

$$\mathbb{P}\left(\left|\mathbf{h}_{1}^{\mathsf{T}}\hat{\mathbf{S}}^{\mathsf{T}}\tilde{\mathbf{y}}\right| > NT\upsilon\right) < 2\exp\left(\frac{-cN^{2}T^{2}\upsilon^{2}}{T}\right) + \epsilon.$$

Likewise, since $\|\tilde{\mathbf{S}}\| = O(\sqrt{T} \|\mathbf{X}\|) = O(\sqrt{NT})$ as., by Lemma 8.1 again we have $\mathbb{P}\left(\left|\mathbf{h}_{1}^{\mathsf{T}}\tilde{\mathbf{S}}\tilde{\mathbf{y}}\right| > NTv\right) < 2\exp\left\{-c_{2}NTv^{2}\right\}$. Also, $T |\mathbf{h}_{1}^{\mathsf{T}}\Delta^{\mathsf{T}}\psi| = O_{\mathbb{P}}(rT \|\mathbf{X}\|)$. By selecting v such that $\sqrt{NT}v = \omega(rT \|\mathbf{X}\|)$, that is, $v = \omega\left(r\sqrt{T}\right)$ where $r\sqrt{T} = o(1)$, we obtain the conclusion for all large enough N and T.

Proof of 2. First note that

$$\mathbf{h}_{1}^{\mathsf{T}} \sum_{t} \left(\hat{\mathbf{X}}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \mathbf{y}_{t} - \mathbf{X}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \psi \right) = \mathbf{h}_{1}^{\mathsf{T}} \left[\mathbf{1}_{T} \otimes \{ \mathcal{L}_{\mathbf{A}} \Delta \} \right]^{\mathsf{T}} \Gamma_{\mathbf{y}}^{1/2} \tilde{\mathbf{y}} + \mathbf{h}_{1}^{\mathsf{T}} \{ \mathbf{1}_{T} \otimes \mathcal{L}_{\mathbf{A}} \mathbf{X} \}^{\mathsf{T}} \Gamma_{\mathbf{y}}^{1/2} \tilde{\mathbf{y}} + T \mathbf{h}_{1}^{\mathsf{T}} \Delta^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \psi.$$

For $\hat{\mathbf{S}} \triangleq \Gamma_{\mathbf{y}}^{1/2} \{ \mathbf{1}_T \otimes (\mathcal{L}_{\mathbf{A}} \Delta) \}$ and $\tilde{\mathbf{S}} \triangleq \Gamma_{\mathbf{y}}^{1/2} \{ \mathbf{1}_T \otimes (\mathcal{L}_{\mathbf{A}} \mathbf{X}) \}$, we have $\left\| \hat{\mathbf{S}} \mathbf{h}_1 \right\| = O_{\mathbb{P}}(\sqrt{T})$ because $\rho(\mathcal{L}_{\mathbf{A}}) = O(1)$ as. hence $\mathbb{P}\left(\left| \mathbf{h}_1^{\mathsf{T}} \hat{\mathbf{S}}^{\mathsf{T}} \tilde{\mathbf{y}} \right| > NTv \right) < 2 \exp\left(\frac{-cN^2 T^2 v^2}{T} \right) + \epsilon$. Since $\left\|\tilde{\mathbf{S}}\mathbf{h}_{1}\right\| = O\left(\sqrt{T} \|\mathbf{X}\|\right) \text{ as., we have } \mathbb{P}\left(|\mathbf{h}_{1}^{\mathsf{T}}\tilde{\mathbf{S}}^{\mathsf{T}}\tilde{\mathbf{y}}| > NT\upsilon\right) < 2\exp\left\{-cNT\upsilon^{2}\right\}.$ The tail probability of $T |\mathbf{h}_{1}^{\mathsf{T}}\Delta^{\mathsf{T}}\mathcal{L}_{\mathbf{A}}\psi|$ can be bounded similarly to the proof of statement 1. \Box

Proof of 3. Let $\mathbf{Z} \triangleq [\mathbf{Z}_0^{\mathsf{T}}, ..., \mathbf{Z}_{T-1}^{\mathsf{T}}]^{\mathsf{T}}$ and $\hat{\mathbf{S}} \triangleq \mathbf{1}_T \otimes \Delta$ and $\tilde{\mathbf{S}} \triangleq \mathbf{1}_T \otimes \mathbf{X}$. Then, we can express $\sum_t \mathbf{h}_1^{\mathsf{T}} \hat{\mathbf{X}}^{\mathsf{T}} \mathbf{Z}_t \mathbf{h}_2$ as tr $(\mathbf{Z}\mathbf{h}_2\mathbf{h}_1^{\mathsf{T}} \hat{\mathbf{S}}^{\mathsf{T}}) + \operatorname{tr}(\mathbf{Z}\mathbf{h}_2\mathbf{h}_1^{\mathsf{T}} \hat{\mathbf{S}}^{\mathsf{T}})$. This is equal to

$$\operatorname{vec}\left(\mathbf{h}_{2}\mathbf{h}_{1}^{\mathsf{T}}\right)^{\mathsf{T}}\left\{\mathbf{I}_{p}\otimes\left(\hat{\mathbf{S}}+\tilde{\mathbf{S}}\right)^{\mathsf{T}}\right\}\operatorname{vec}\left(\mathbf{Z}\right).$$

Note that $\left\|\mathbf{I}_p \otimes \hat{\mathbf{S}}^{\mathsf{T}}\right\| = O_{\mathbb{P}}\left(\sqrt{T}\right)$ and $\left\|\mathbf{I}_p \otimes \tilde{\mathbf{S}}^{\mathsf{T}}\right\| = O\left(\sqrt{T} \|\mathbf{X}\|\right)$. Therefore, by Theorem 8.2, we have

$$\mathbb{P}\left[\left|\sum_{t} \mathbf{h}_{1}^{\mathsf{T}} \hat{\mathbf{X}}^{\mathsf{T}} \mathbf{Z}_{t} \mathbf{h}_{2}\right| > NT\upsilon\right] \leq 2\exp\left(-\frac{cN^{2}T^{2}\upsilon^{2}}{T}\right) + 2\exp\left(-c'NT\upsilon^{2}\right) + \epsilon.$$

Proof of 4. First we have

$$\sum_{t} \left(\mathbf{y}_{t}^{\mathsf{T}} \mathbf{y}_{t} - \psi^{\mathsf{T}} \psi \right) = \left\| \Gamma_{\mathbf{y}}^{1/2} \tilde{\mathbf{y}} + \Psi \right\|^{2} - T \psi^{\mathsf{T}} \psi = \tilde{\mathbf{y}}^{\mathsf{T}} \Gamma_{\mathbf{y}} \tilde{\mathbf{y}} + 2 \Psi^{\mathsf{T}} \Gamma_{\mathbf{y}}^{1/2} \tilde{\mathbf{y}}.$$

Note that $\|\Gamma_{\mathbf{y}}\|_F^2 = \sum_{i=1}^{NT} \sigma_i (\Gamma_{\mathbf{y}})^2 = O(NT)$ as. By Theorem 8.3, we have

$$\mathbb{P}\left(|\tilde{\mathbf{y}}^{\mathsf{T}}\Gamma_{\mathbf{y}}\tilde{\mathbf{y}} - T\operatorname{tr}\mathbb{E}\Gamma| > NT\upsilon\right) < 2\exp\left(-c\min\left\{NT\upsilon^{2}, NT\upsilon\right\}\right).$$

By Lemma 8.1,

$$\mathbb{P}\left(2|\Psi^{\mathsf{T}}\Gamma_{\mathbf{y}}^{-1/2}\tilde{\mathbf{y}}| > NT\upsilon\right) < 2\exp\left(-c'NT\upsilon^{2}/r^{2}\right)$$

since $\left\|\Gamma_{\mathbf{y}}^{1/2}\Psi\right\|^2 \leq \left\|\Gamma_{\mathbf{y}}\right\| T\psi^{\dagger}\psi = O\left(r^2T \left\|\mathbf{X}\right\|^2\right) = O(r^2NT)$ as.

Proof of 5. Let $\mathbf{S}_{\mathbf{A}} \triangleq \Gamma_{\mathbf{y}}^{1/2} \left(\mathbf{I}_T \otimes \mathcal{L}_{\mathbf{A}} \right) \Gamma_{\mathbf{y}}^{1/2}$. We have

$$\sum_{t} \mathbf{y}_{t}^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \mathbf{y}_{t} - \psi^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \psi = \left(\Gamma_{\mathbf{y}}^{1/2} \tilde{\mathbf{y}} + \Psi \right)^{\mathsf{T}} \left(\mathbf{I}_{T} \otimes \mathcal{L}_{\mathbf{A}} \right) \left(\Gamma_{\mathbf{y}}^{1/2} \tilde{\mathbf{y}} + \Psi \right) - T \psi^{\mathsf{T}} \mathcal{L}_{\mathbf{A}} \psi$$
$$= \tilde{\mathbf{y}}^{\mathsf{T}} \mathbf{S}_{\mathbf{A}} \tilde{\mathbf{y}} + 2 \Psi^{\mathsf{T}} \Gamma_{\mathbf{y}}^{-\frac{1}{2}} \mathbf{S}_{\mathbf{A}} \tilde{\mathbf{y}}.$$

Also,

$$\|\mathbf{S}_{\mathbf{A}}\|_{F}^{2} = \operatorname{tr}\left[\left(\mathbf{I}_{T} \otimes \mathcal{L}_{\mathbf{A}}\right) \Gamma_{\mathbf{y}}\left(\mathbf{I}_{T} \otimes \mathcal{L}_{\mathbf{A}}\right) \Gamma_{\mathbf{y}}\right] \leq \operatorname{tr}\left(\Gamma_{\mathbf{y}}^{2}\right) \left\|\mathbf{I}_{T} \otimes \mathcal{L}_{\mathbf{A}}^{2}\right\| = O(NT) \ as.$$
$$\left\|\mathbf{S}_{\mathbf{A}}\Gamma_{\mathbf{y}}^{-\frac{1}{2}}\Psi\right\| = O\left(r\sqrt{T} \left\|\mathbf{X}\right\|\right) \ as.$$

We can apply Lemmas 8.3 and 8.1 to directly obtain the following:

$$\mathbb{P}\left[\left|\tilde{\mathbf{y}}^{\mathsf{T}}\mathbf{S}_{\mathbf{A}}\tilde{\mathbf{y}} - T\mathbb{E}\operatorname{tr}\left(\mathcal{L}_{\mathbf{A}}\Gamma\right)\right| > NT\upsilon\right] < 2\exp\left(-c\min\left\{NT\upsilon^{2}, NT\upsilon\right\}\right)$$
$$\mathbb{P}\left[\left|2\Psi^{\mathsf{T}}\Gamma_{\mathbf{y}}^{-\frac{1}{2}}\mathbf{S}_{\mathbf{A}}\tilde{\mathbf{y}}\right| > NT\upsilon\right] < 2\exp\left(-cNT\upsilon^{2}/r^{2}\right).$$

Proof of 6. We have

$$\sum_{t} \mathbf{y}_{t}^{\mathsf{T}} \mathbf{Z}_{t} \mathbf{h}_{2} = \left(\Gamma_{\mathbf{y}}^{-1/2} \tilde{\mathbf{y}} + \Psi \right)^{\mathsf{T}} \mathbf{Z} \mathbf{h}_{2} = \operatorname{tr} \left(\mathbf{h}_{2} \tilde{\mathbf{y}}^{\mathsf{T}} \Gamma_{\mathbf{y}}^{-1/2} \mathbf{Z} \right) + \operatorname{tr} \left(\mathbf{h}_{2} \Psi^{\mathsf{T}} \mathbf{Z} \right)$$
$$= \operatorname{vec} \left(\tilde{\mathbf{y}} \mathbf{h}_{2}^{\mathsf{T}} \right)^{\mathsf{T}} \left(\mathbf{I}_{p} \otimes \Gamma_{\mathbf{y}}^{-1/2} \right) \operatorname{vec} \left(\mathbf{Z} \right) + \operatorname{vec} \left(\Psi \mathbf{h}_{2}^{\mathsf{T}} \right)^{\mathsf{T}} \left(\mathbf{I}_{p} \otimes \mathbf{I}_{NT} \right) \operatorname{vec} \left(\mathbf{Z} \right).$$

Therefore, since $\|\Gamma_{\mathbf{y}}\| = O(1)$ as., $\left\| \left(\mathbf{I}_p \otimes \Gamma_{\mathbf{y}}^{1/2} \right) \operatorname{vec} \left(\tilde{\mathbf{y}} \mathbf{h}_2^{\mathsf{T}} \right) \right\| \lesssim \|\operatorname{vec} \left(\tilde{\mathbf{y}} \mathbf{h}_2^{\mathsf{T}} \right) \|$ which is concentrated around \sqrt{NTp} (Vershynin, 2018) hence is $O(\sqrt{NTp})$ whp. Also, we have $\| (\mathbf{I}_p \otimes \mathbf{I}_{NT}) \operatorname{vec} \left(\Psi \mathbf{h}_2^{\mathsf{T}} \right) \| = O\left(r \sqrt{T} \| \mathbf{X} \| \right)$ as. By Lemma 8.1,

$$\mathbb{P}\left(\left|\sum_{t} \mathbf{y}_{t}^{\mathsf{T}} \mathbf{Z}_{t} \mathbf{h}_{2}\right| > NT\upsilon\right) \leq 2\exp\left(-\frac{c_{1}NT\upsilon^{2}}{p}\right) + 2\exp\left(-c_{2}NT\upsilon^{2}/r^{2}\right) + \epsilon$$

provided that $NTv = \omega(\sqrt{NTp})$.

Proof of 7. Similarly to the proof of statement 5., let $\mathbf{S}_{\mathbf{A}} \triangleq \Gamma_{\mathbf{y}}^{1/2} \left(\mathbf{I}_{T} \otimes \mathcal{L}_{\mathbf{A}}^{2} \right) \Gamma_{\mathbf{y}}^{1/2}$ and check that

$$\|\mathbf{S}_{\mathbf{A}}\|_{F}^{2} \leq \operatorname{tr}\left(\Gamma_{\mathbf{y}}^{2}\right) \|\mathbf{I}_{T} \otimes \mathcal{L}_{\mathbf{A}}^{4}\| = O(NT) \ as., \left\|\mathbf{S}_{\mathbf{A}}\Gamma_{\mathbf{y}}^{-\frac{1}{2}}\Psi\right\| = O\left(\sqrt{T} \|\mathbf{X}\|\right) \ as.$$

Again by Lemmas 8.3 and 8.1, we directly have the result by the same technique used in the proof of statement 5. $\hfill \Box$

Proof of 8. Since $\|\mathcal{L}_{\mathbf{A}}\| \leq 1$, the proof coincides with the proof of statement 6. \Box **Proof of 10.** Note that $\|\Delta\|^2 = O_{\mathbb{P}}(1)$. Also, $\|\beta\| = O(1)$. So, since

$$\left|\mathbf{h}_{1}^{\mathsf{T}}\hat{\mathbf{X}}^{\mathsf{T}}\Delta\beta\right| \leq \left(\|\Delta\| + \|\mathbf{X}\|\right)\|\Delta\| \|\beta\| = O_{\mathbb{P}}(\|\mathbf{X}\|),$$

by letting $v = \omega\left(\frac{r}{\sqrt{NT}} \|\mathbf{X}\|\right)$ where we have $\frac{r\|\mathbf{X}\|}{\sqrt{NT}} = O(r/\sqrt{T})$ and $r = o(1/\sqrt{T})$, we have the conclusion.

Proof of 11. Similar to the proof of 1., taking $\hat{\mathbf{S}} \triangleq \Gamma_{\mathbf{y}}^{1/2} (\mathbf{1}_T \otimes \Delta)$ which has the spectral norm of $O_{\mathbb{P}}(\sqrt{T})$, we have

$$\sum_{t} \mathbf{y}_{t}^{\mathsf{T}} \Delta \beta = \left(\Gamma_{\mathbf{y}}^{1/2} \tilde{\mathbf{y}} + \Psi \right)^{\mathsf{T}} \left(\mathbf{1}_{T} \otimes \Delta \right) \beta = \tilde{\mathbf{y}}^{\mathsf{T}} \hat{\mathbf{S}} \beta + \Psi^{\mathsf{T}} \Gamma_{\mathbf{y}}^{-\frac{1}{2}} \hat{\mathbf{S}} \beta.$$

First noting that $\|\beta\| = O(1)$ as $K \to \infty$, Lemma 8.1 gives

$$\mathbb{P}\left(r\left|\tilde{\mathbf{y}}^{\mathsf{T}}\hat{\mathbf{S}}\beta\right| \geq \sqrt{NT}\upsilon\right) \leq 2\exp\left(\frac{-cNT\upsilon^{2}}{Tr^{2}}\right) + \epsilon.$$

It is straightforward that $r \left| \Psi^{\dagger} \Gamma_{\mathbf{y}}^{-\frac{1}{2}} \hat{\mathbf{S}} \beta \right| = O_{\mathbb{P}}(r^2 T \| \mathbf{X} \|)$ hence by letting $\sqrt{NT} \upsilon = \omega(r^2 \sqrt{NT})$, i.e., $\upsilon = \omega \left(r^2 \sqrt{T} \right)$, we obtain the conclusion.

Proof of 12. Let $\hat{\mathbf{S}} \triangleq \Gamma_{\mathbf{y}}^{1/2} \{ \mathbf{1}_T \otimes (\mathcal{L}_{\mathbf{A}} \Delta) \}$ which is of $O_{\mathbb{P}} \left(\sqrt{T} \right)$. The rest of the proof coincides with the proof of 11.

Proof of 13. This follows by noting that $\operatorname{vec}(\mathbf{h}_2\beta^{\intercal})^{\intercal} \{ \mathbf{I}_p \otimes (\mathbf{1}_T \otimes \Delta)^{\intercal} \}$ is of $O_{\mathbb{P}}(\sqrt{T})$.

Lemma 9.2. There exist $N_0, T_0 > 0$ and $v_0(N_0, T_0) \in (0, 1)$ such that for all $v \in (0, v_0)$ and $N > N_0, T > T_0$,

$$\mathbb{P}\left[\sup_{\mathbf{h}\in\mathbb{R}^{K+p+2};\|\mathbf{h}\|\leq 1}\left|\mathbf{h}^{\mathsf{T}}\left(\hat{\Omega}_{m}-\Omega_{m}\right)\mathbf{h}\right|>\upsilon\right]\lesssim K^{2}\left\{\epsilon+\exp\left(-c\upsilon^{2}\frac{NT}{pK^{2}}\right)\right\}.$$

9.4 Theories for Corollary Models

Similarly, we verify the asymptotic properties of AMNAR with finite T.

Proof of Theorem 4.6

Let $\tilde{\mathbf{D}}_N \triangleq \sqrt{N} \operatorname{diag}(r\mathbf{I}_{K+1}, \mathbf{I}_{p+2})$. Then, $\sqrt{T}\tilde{\mathbf{D}}_N = \tilde{\mathbf{D}}_{NT}$. We have

$$\hat{\mu}_{m} = \tilde{\mathbf{D}}_{NT}^{-1} \left(\hat{\Omega}_{m} \right)^{-1} \tilde{\mathbf{D}}_{NT}^{-1} \hat{\mathbf{M}}^{\mathsf{T}} \left\{ \left(\mathbf{M} - \hat{\mathbf{M}} \right) \mu_{m} + \hat{\mathbf{M}} \mu_{m} + \mathcal{E} \right\}$$
$$= \frac{\tilde{\mathbf{D}}_{N}^{-1}}{\sqrt{T}} \hat{\Omega}_{m}^{-1} \left\{ \tilde{\mathbf{D}}_{NT}^{-1} \hat{\mathbf{M}}^{\mathsf{T}} \left(\mathbf{M} - \hat{\mathbf{M}} \right) \mu_{m} + \mathcal{E}_{\mathbf{M}} \right\} + \mu_{m}$$

hence

$$\hat{\Omega}_m \tilde{\mathbf{D}}_N (\hat{\mu}_m - \mu_m) = \frac{\tilde{\mathbf{D}}_{NT}^{-1}}{\sqrt{T}} \hat{\mathbf{M}}^{\mathsf{T}} (\mathbf{M} - \hat{\mathbf{M}}) \mu_m + \frac{\mathcal{E}_{\mathbf{M}}}{\sqrt{T}}$$

Claim 1. $\hat{\Omega}_m \Rightarrow \Omega_m$.

Proof. This is true under finite T as well as Lemma 9.1 and 9.2 imply when we fix $T = T_0 > 0$.

Claim 2. $\tilde{\mathbf{D}}_{NT}^{-1} \hat{\mathbf{M}}^{\mathsf{T}} (\mathbf{M} - \hat{\mathbf{M}}) \mu_m = o_{\mathbb{P}}(1).$

Proof. By the statements 10.–13. of Lemma 9.1 with fixed $T = T_0 > 0$, we have the conclusion for $N \to \infty$.

Claim 3. $\mathcal{E}_{\mathbf{M}} \Rightarrow \mathcal{N}(\mathbf{o}_{K+p+3}, \sigma^2 \Omega_m).$

Proof. Recall that $r = \Theta(N^{-s})$. Then, this directly holds by noticing that the Claim 3 of the proof of Theorem 4.5 also holds for fixed $T = T_0 > 0$ and $N \to \infty$.

Therefore, by Claims 1–3, the asymptotic normality follows. \Box

10 Additional tables and figures

10.1 Simulation

The figure A1 displays the boxplots of estimates of θ and α parameters with growing N and T when the data is generated from NAR and ENAR models respectively. The simulation setup and the figure has been described in the main text.



Figure A1: Boxplot of estimates of θ and α from AMNAR, ENAR, and NAR model when data is generated from NAR model with DCMMSBM.

10.2 Real data analysis

The table A1 below displays parameter estimates along with standard errors for the Knecht dataset.

	Dependent variable:					
	Alcohol Consumption			Delinquency		
	OLS	NAR	ENAR	OLS	NAR	ENAR
Alc. previous		0.278	0.372		0.101	0.164
		(0.288)	(0.310)		(0.176)	(0.151)
Peer effect		0.708	1.788		-0.714	-0.375
		(0.725)	(1.253)		(0.583)	(0.321)
Sex	-0.533	-0.328	-0.669	-0.061	-0.369	-0.083
	(0.480)	(0.513)	(1.284)	(0.250)	(0.661)	(0.243)
Age	0.029	0.182	0.212	-0.003	-0.050	0.009
	(0.472)	(0.497)	(0.560)	(0.245)	(0.286)	(0.240)
Ethnicity	0.444	0.224	0.144	-0.595	-0.646	-0.591
	(0.815)	(0.853)	(0.891)	(0.424)	(0.459)	(0.415)
Religion	0.664**	0.540	0.525	0.277	0.310	0.294
	(0.315)	(0.336)	(0.356)	(0.164)	(0.189)	(0.176)

Table A1: Parameter estimates and standard errors for the OLS, NAR and ENAR models fitted to the alcohol consumption data and delinquency data.

The figure A2 displays the predicted responses against the actual ones to show the quality of predictions.

Table A2 provides estimates from ENAR and NAR for one representative model with T = 520.



Figure A2: Boxplots of predicted values from the 3 models for different labels of actual values for the response on (a) alcohol and (b) delinquency

Table A2: Estimates of ENAR and NAR model for the Wind speed data where the model is fit on data between T = [1, 520].

	Dependent variable: Wind speed			
	ENAR	NAR		
Ylagged	0.810***	0.823***		
	(0.003)	(0.002)		
LYlagged	0.096***	0.081***		
	(0.003)	(0.002)		
AIC	53243.98	53576.56		
BIC	53394.89	53612.07		
Note:	*p<0.1; **p<0.05; ***p<0.01			

References

- An, W., Beauvile, R., and Rosche, B. (2022). Causal network analysis. Annual Review of Sociology, 48:23–41.
- Athreya, A., Fishkind, D. E., Tang, M., Priebe, C. E., Park, Y., Vogelstein, J. T., Levin, K., Lyzinski, V., and Qin, Y. (2017). Statistical inference on random dot product graphs: a survey. *The Journal of Machine Learning Research*, 18(1):8393–8484.
- Basu, S. and Michailidis, G. (2015). Regularized estimation in sparse high-dimensional time series models. *The Annals of Statistics*, 43(4):1535 1567.
- Bhatia, R. (2013). Matrix analysis, volume 169. Springer Science & Business Media.
- Cape, J., Tang, M., and Priebe, C. E. (2019). Signal-plus-noise matrix models: eigenvector deviations and fluctuations. *Biometrika*, 106(1):243–250.
- Chen, E. Y., Fan, J., and Zhu, X. (2023). Community network auto-regression for highdimensional time series. *Journal of Econometrics*, 235(2):1239–1256.
- Christakis, N. A. and Fowler, J. H. (2007). The spread of obesity in a large social network over 32 years. *New England journal of medicine*, 357(4):370–379.
- Christakis, N. A. and Fowler, J. H. (2013). Social contagion theory: examining dynamic social networks and human behavior. *Statistics in medicine*, 32(4):556–577.
- Fosdick, B. K. and Hoff, P. D. (2015). Testing and modeling dependencies between a network and nodal attributes. *Journal of the American Statistical Association*, 110(511):1047–1056.

- Goldsmith-Pinkham, P. and Imbens, G. W. (2013). Social networks and the identification of peer effects. *Journal of Business & Economic Statistics*, 31(3):253–264.
- Guan, Y., Page, G. L., Reich, B. J., Ventrucci, M., and Yang, S. (2023). Spectral adjustment for spatial confounding. *Biometrika*, 110(3):699–719.
- Hall, P. and Heyde, C. C. (2014). *Martingale limit theory and its application*. Academic press.
- He, Y. and Hoff, P. D. (2019). Multiplicative coevolution regression models for longitudinal networks and nodal attributes. *Social Networks*, 57:54–62.
- Hoff, P. (2021). Additive and multiplicative effects network models. Statistical Science.
- Hoff, P. D., Raftery, A. E., and Handcock, M. S. (2002). Latent space approaches to social network analysis. *Journal of the american Statistical association*, 97(460):1090–1098.
- Knecht, A., Snijders, T. A., Baerveldt, C., Steglich, C. E., and Raub, W. (2010). Friendship and delinquency: Selection and influence processes in early adolescence. *Social Development*, 19(3):494–514.
- Knecht, A. B. (2008). Friendship selection and friends' influence: Dynamics of networks and actor attributes in early adolescence.
- Knight, M., Leeming, K., Nason, G., and Nunes, M. (2020). Generalized network autoregressive processes and the gnar package. *Journal of Statistical Software*, 96(5):1–36.
- Le, C. M. and Li, T. (2022). Linear regression and its inference on noisy networklinked data. Journal of the Royal Statistical Society Series B: Statistical Methodology, 84(5):1851–1885.
- Lei, J. and Rinaldo, A. (2015). Consistency of spectral clustering in stochastic block models. *The Annals of Statistics*, 43(1).
- Li, J., Xu, G., and Zhu, J. (2023). Statistical inference on latent space models for network data. *arXiv preprint arXiv:2312.06605*.
- Ma, Z., Ma, Z., and Yuan, H. (2020). Universal latent space model fitting for large networks with edge covariates. *Journal of Machine Learning Research*, 21(4):1–67.
- McFowland III, E. and Shalizi, C. R. (2021). Estimating causal peer influence in homophilous social networks by inferring latent locations. *Journal of the American Statistical Association*, pages 1–12.
- Nath, S., Warren, K., and Paul, S. (2022). Identifying peer influence in therapeutic communities. arXiv preprint arXiv:2203.14223.
- O'Malley, A. J., Elwert, F., Rosenquist, J. N., Zaslavsky, A. M., and Christakis, N. A. (2014). Estimating peer effects in longitudinal dyadic data using instrumental variables. *Biometrics*, 70(3):506–515.
- Pearl, J. (2009). Causality. Cambridge university press.
- Rohe, K., Tao, J., Han, X., and Binkiewicz, N. (2018). A note on quickly sampling a sparse matrix with low rank expectation. *Journal of Machine Learning Research*, 19(77):1–13.
- Rubin-Delanchy, P., Cape, J., Tang, M., and Priebe, C. E. (2022). A statistical interpretation of spectral embedding: The generalised random dot product graph. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 84(4):1446–1473.
- Rudelson, M. and Vershynin, R. (2013). Hanson-Wright inequality and sub-gaussian concentration. *Electronic Communications in Probability*, 18(none):1 – 9.

- Shalizi, C. R. and Thomas, A. C. (2011). Homophily and contagion are generically confounded in observational social network studies. *Sociological methods & research*, 40(2):211–239.
- Tang, M. and Priebe, C. E. (2018). Limit theorems for eigenvectors of the normalized laplacian for random graphs. The Annals of Statistics, 46(5):2360–2415.
- VanderWeele, T. J. (2011). Sensitivity analysis for contagion effects in social networks. Sociological Methods & Research, 40(2):240–255.
- VanderWeele, T. J., Ogburn, E. L., and Tchetgen, E. J. T. (2012). Why and when" flawed" social network analyses still yield valid tests of no contagion. *Statistics, Politics* and Policy, 3(1).
- Vershynin, R. (2018). High-dimensional probability: An introduction with applications in data science, volume 47. Cambridge university press.
- Xie, F. and Xu, Y. (2023). Efficient estimation for random dot product graphs via a one-step procedure. *Journal of the American Statistical Association*, 118:651–664.
- Zhu, X. and Pan, R. (2020). Grouped network vector autoregression. *Statistica Sinica*, 30(3):1437–1462.
- Zhu, X., Pan, R., Li, G., Liu, Y., and Wang, H. (2017). Network vector autoregression. *The Annals of Statistics*, 45(3):1096 – 1123.
- Zhu, X., Wang, W., Wang, H., and H\u00e4rdle, W. K. (2019). Network quantile autoregression. Journal of econometrics, 212(1):345–358.