

STABILITY AND FINITENESS OF WASSERSTEIN SPACES

MOHAMMAD ALATTAR

ABSTRACT. Under Gromov–Hausdorff convergence, and equivariant Gromov–Hausdorff convergence, we prove stability results of Wasserstein spaces over certain classes of singular and non-singular spaces. For example, we obtain an analogue of Perelman’s stability theorem on Wasserstein spaces.

1. INTRODUCTION

Finiteness theorems for finite dimensional spaces with curvature bounds can be traced back to 1967, with Weinstein [32] proving that given any even natural number n , and any $\delta > 0$, there are only finitely many homotopy types of n -dimensional, δ -pinched manifolds. Shortly after, in 1970, Cheeger obtained several finiteness theorems in his celebrated work [10]. For instance, he proved that given any natural number n , and real numbers $D, v, k > 0$, there are only finitely many diffeomorphism types of compact n -dimensional manifolds X admitting a Riemannian metric such that $\text{diameter}(X) \leq D$, $\text{vol}(X) \geq v$ and $|\text{sec}(X)| \leq k^2$, where $\text{sec}(X)$ denotes the sectional curvature. These results served as departure points for further seminal and important developments in Riemannian geometry (see for instance [3, 21, 22, 20, 29]).

In 1981, Gromov [19] introduced a powerful notion of convergence, now known as *Gromov–Hausdorff convergence*, that has been useful in establishing (among other things) finiteness theorems through the notion of “stability”. For instance, using Gromov–Hausdorff convergence, Grove, Petersen and Wu [22], in 1990, relaxed the curvature condition in Cheeger’s finiteness theorem and showed that for any $n \neq 3$, real number k , and any positive numbers D and v , the class consisting of compact n -dimensional Riemannian manifolds X with $\text{diameter}(X) \leq D$, $\text{sec}(X) \geq k$, and $\text{vol}(X) \geq v$ contains finitely many homeomorphism types.

In 1991, Perelman [29, 25], using Gromov–Hausdorff convergence, showed that given a non-collapsing convergent sequence of compact Alexandrov spaces, with a uniform lower curvature bound, uniform finite dimension and uniform upper diameter bound, the tail of the sequence *stabilizes* topologically. That is, all spaces in the tail of the sequence are mutually homeomorphic. As a corollary, one obtains homeomorphism finiteness (in all dimensions) for a more general class of spaces that need not be manifolds.

In all cases of finiteness and stability, attention has been restricted to finite dimensional spaces. Thus, the question of stability in infinite dimensions remains open. In this paper, we focus our attention on *Wasserstein spaces*, which are infinite-dimensional, and prove stability and finiteness results for these spaces in different settings. Our proofs are not technical, and these results are, to the best of our knowledge, the first stability and finiteness results for infinite-dimensional spaces.

Date: June 11, 2024.

2020 Mathematics Subject Classification. 53C23, 53C20, 51K10.

Key words and phrases. equivariant Gromov–Hausdorff convergence, Gromov–Hausdorff convergence, Wasserstein spaces, stability.

Let $\mathcal{A}^{\leq}(n, K, D)$ denote the class of compact Alexandrov spaces with dimension at most n , curvature bounded below by K , and diameter bounded above by D . We denote by $\mathcal{A}(n, K, D)$ all Alexandrov spaces in $\mathcal{A}^{\leq}(n, K, D)$ with dimension n . Given $v > 0$, we will denote by $\mathcal{A}(n, K, D, v)$ the subset of $\mathcal{A}(n, K, D)$ where the Alexandrov spaces under consideration have volume bounded below by v . Furthermore, given a sequence of compact metric spaces $\{X_i\}_{i \in \mathbb{N}}$ and a compact metric space X , we will denote by $X_i \rightarrow_{GH} X$ the Gromov–Hausdorff convergence of X_i to X . We will denote by $\mathbb{P}_2(X)$ the Wasserstein space over X equipped with the 2-Wasserstein metric (see Definition 2.5).

Theorem A. *Assume $\{X_i\}_{i \in \mathbb{N}}$ and X are in $\mathcal{A}^{\leq}(n, K, D)$. Then the following assertions are equivalent:*

- (1) $\mathbb{P}_2(X_i) \rightarrow_{GH} \mathbb{P}_2(X)$.
- (2) $X_i \rightarrow_{GH} X$.
- (3) $(\mathbb{P}_2(X_i), X_i) \rightarrow_{GH} (\mathbb{P}_2(X), X)$.

We point out a few remarks concerning the above theorem. First, the interesting implication is (1) \implies (2). The implication (2) \implies (1) is well known (see Corollary 4.3 in [28]) and holds for general compact metric spaces. Second, the notion of convergence in (3) is known as *Gromov–Hausdorff convergence of metric pairs*. It was introduced in [9], and later studied extensively in [1]. This notion of convergence generalizes Gromov–Hausdorff convergence and takes into account pairs of the form (X, A) , where X is a metric space, and A is a closed non-empty subset of X . In our theorem, we identify the base space X with the set of Dirac deltas in the Wasserstein space $\mathbb{P}_2(X)$. Thus, (2) \implies (3) becomes trivial. However, the new idea for this implication is that we give an equivalent characterization of convergence of metric pairs using the notion of ϵ -isometries (see Proposition 2.1). This equivalent characterization will be useful when we introduce the notion of relative equivariant Gromov–Hausdorff convergence (see Definition 2.2) which takes into account a triple (X, A, G) , where X is a compact metric space, A is a closed, non-empty G -invariant subset of X and G is a closed subgroup of $\text{Isom}(X)$, the group of isometries of X .

A consequence of the above theorem, is the following analogue of Perelman’s stability theorem for Wasserstein spaces.

Corollary B. *Let $\{X_i\}_{i \in \mathbb{N}}$ and X be spaces in $\mathcal{A}(n, K, D)$. If $\mathbb{P}_2(X_i) \rightarrow_{GH} \mathbb{P}_2(X)$ then for all sufficiently large i , $\mathbb{P}_2(X_i)$ and $\mathbb{P}_2(X)$ are homeomorphic, and the homeomorphisms can be taken to be Gromov–Hausdorff approximations.*

Note that in the preceding corollary we do not impose any other structure on the Wasserstein spaces other than that coming from the base spaces.

We further have the following finiteness result. For any $n \in \mathbb{N}$, $K \in \mathbb{R}$, $D > 0$ and $v > 0$, let $W_2(n, K, D, v)$ denote the class consisting of Wasserstein spaces with base spaces in $\mathcal{A}(n, K, D, v)$ and equipped with the 2-Wasserstein metric.

Corollary C. *There are only finitely many topological types in $W_2(n, K, D, v)$.*

We note that the technique in this paper can be applied whenever we have a stability result on the base spaces and essentially reduces to the following result.

Theorem D. *Let \mathfrak{X} denote a precompact class (with respect to the Gromov–Hausdorff topology) of compact non-branching metric measure spaces having good transport behavior (GTB) and their reference measures*

are qualitatively non-degenerate. Assume that if $\{X_i\}_{i=1}^\infty$ is a sequence in \mathfrak{X} , and $X_i \rightarrow_{GH} X$, then X is a non-branching compact metric space admitting a measure that is qualitatively non-degenerate and has good transport behavior (we will take such a measure as a reference measure on X). Then, the following assertions are equivalent:

- (1) $\mathbb{P}_2(X_i) \rightarrow_{GH} \mathbb{P}_2(X)$.
- (2) $X_i \rightarrow_{GH} X$.

The notion of good transport behavior was first introduced in the paper by Galaz-García, Kell, Mondino, and Sosa [18], and further studied by Kell [26]. The class of metric measure spaces satisfying such condition is large. For instance, according to [18], strong $CD^*(K, N)$ and essentially non-branching $MCP(K, N)$ spaces admit good transport behavior. The notion of a measure being qualitatively non-degenerate was introduced by Cavalletti and Huesmann [7] and later studied by Kell [26]. It is technical to state, and we refer the reader to [26, 7, 30] for more details, but we note a measure being qualitatively non-degenerate yields desirable information. For example, a qualitatively non-degenerate measure is doubling (Lemma 2.8 in [30]).

We note that the classes $\mathcal{A}^\leq(n, K, D)$, $\mathcal{A}(n, K, D)$ satisfy the hypotheses of Theorem D. However, the properties in Theorem D are not closed under Gromov–Hausdorff convergence in general. This is because a compact metric measure space endowed with a qualitatively non-degenerate measure is doubling and thus must have finite dimension.

Using almost verbatim the proofs of Theorems D and A, and noting that $RCD(K, N)$ spaces are non-branching [14], we obtain the following corollary.

Corollary E. *Assume each $(X_i, d_i, \mathfrak{m}_i)$ is an compact $RCD(K, N)$ space with uniform upper diameter bound. Assume (X, d) is an N -dimensional compact Riemannian manifold. If $\mathbb{P}_2(X_i) \rightarrow_{GH} \mathbb{P}_2(X)$ then for all large i , there exists a homeomorphism $G_i: \mathbb{P}_2(X_i) \rightarrow \mathbb{P}_2(X)$ that is Lipschitz and its inverse is Hölder.*

A consequence of our next main theorem is an equivariant stability result for Wasserstein spaces. We first discuss notation. Given a closed subgroup G of $\text{Isom}(X)$, where X is a compact metric space, we denote by $G^\#$ the induced subgroup of $\text{Isom}(\mathbb{P}_2(X))$ given by the push-forward of maps. Given a sequence of pairs $\{(X_i, G_i)\}_{i \in \mathbb{N}}$, where each X_i is a compact metric space and G_i is a closed subgroup of $\text{Isom}(X)$, we will denote, by $(X_i, G_i) \rightarrow_{eGH} (X, G)$ the equivariant Gromov–Hausdorff convergence of (X_i, G_i) to (X, G) . This convergence, originally introduced by Fukaya (see [16, 17, 15]), has been useful in establishing interesting results in both the singular and non-singular setting (see for instance [34, 12, 5, 23, 33, 13, 8]). Furthermore, it has been useful in obtaining finiteness and stability results in the singular setting. For instance, using equivariant Gromov–Hausdorff convergence, Zamora extended Anderson finiteness to the $RCD^*(K, N)$ setting [33].

Theorem F. *Let $\{X_i\}_{i \in \mathbb{N}}$ and X be closed Riemannian manifolds with uniform lower sectional curvature bound $K > 0$, uniform upper diameter bound and uniform upper dimension bound. Assume H_i and H are closed subgroups of $\text{Isom}(\mathbb{P}_2(X_i))$ and $\text{Isom}(\mathbb{P}_2(X))$ respectively. Then the following assertions are equivalent:*

- (1) $(\mathbb{P}_2(X_i), H_i) \rightarrow_{eGH} (\mathbb{P}_2(X), H)$.
- (2) *There exists unique closed subgroups G_i of $\text{Isom}(X)$ and G of $\text{Isom}(X)$ such that $G_i^\# = H_i$ and $G^\# = H$ and such that $(X_i, G_i) \rightarrow_{eGH} (X, G)$.*
- (3) $(\mathbb{P}_2(X_i), X_i, H_i) \rightarrow_{eGH} (\mathbb{P}_2(X), X, H)$.

We emphasize that we identify the base space with the (closed) subset of Dirac deltas in the Wasserstein space. In the presence of a uniform lower curvature bound, we have the following corollary, an equivariant stability result on Wasserstein spaces.

Corollary G. *Let $\{X_i\}_{i \in \mathbb{N}}$ and X be closed Riemannian manifolds with uniform lower sectional curvature bound $K > 0$, uniform upper diameter bound and uniform dimension. Assume H_i and H are closed subgroups of $\text{Isom}(\mathbb{P}_2(X_i))$ and $\text{Isom}(\mathbb{P}_2(X))$ respectively of the same dimension. If $(\mathbb{P}_2(X_i), H_i) \rightarrow_{eGH} (\mathbb{P}_2(X), H)$, then for all large i , there exists a homeomorphism $f_i: \mathbb{P}_2(X_i) \rightarrow \mathbb{P}_2(X)$ that is compatible with some group isomorphism $\chi_i: H_i \rightarrow H$.*

Our article is organized as follows. In section 2, we discuss the preliminaries that will be useful in this paper. Moreover, we will give a new characterization of convergence of metric pairs. In section 3 we give proofs of the theorems and corollaries.

Acknowledgements. It is a pleasure to thank Fernando Galaz–García, Martin Kerin, Kohei Suzuki, Jaime Santos-Rodríguez, and Mauricio Che for helpful conversations and comments during the Durham Metric Geometry Reading Seminar. I would also like to thank Sergio Zamora and Yanpeng Zhi for several interesting conversations and Aseel alnajjar for several comments on the first draft of this manuscript. Part of this work was completed at the Erwin Schrödinger International Institute for Mathematics and Physics (ESI), where the author received financial support to participate in the workshop *Synthetic Curvature Bounds for Non-Smooth Spaces: Beyond Finite Dimension*. The author would like to thank ESI and the conference organizers for the excellent atmosphere and working conditions.

2. PRELIMINARIES

In this section, we collect preliminary definitions and results we will use in the proof of our main results.

2.1. Convergence of Metric Pairs and Equivariant Convergence of Metric Pairs. In this section, we give a new equivalent characterization of the notion of convergence of metric pairs which will allow us to introduce, canonically, the notion of relative equivariant Gromov–Hausdorff convergence. The equivalent characterization is demonstrated in the following proposition. We refer the reader to the paper by Ahumada Gómez and Che [1] for results concerning the (non-equivariant) convergence of metric pairs.

Proposition 2.1. *Let $\{X_i\}_{i \in \mathbb{N}}$ be a sequence of compact metric spaces. Let $\{A_i\}_{i \in \mathbb{N}}$ be a collection of closed non-empty subspaces $A_i \subseteq X_i$. Let X be a compact metric space, and $A \subseteq X$ a closed non-empty subset. Then the following assertions are equivalent:*

- (1) $(X_i, A_i) \rightarrow_{GH} (X, A)$.
- (2) *There exists a sequence of positive numbers $\{\epsilon_i\}_{i \in \mathbb{N}}$ such that $\epsilon_i \rightarrow 0$ and ϵ_i -Gromov–Hausdorff approximations $f_i: X_i \rightarrow X$, $g_i: A_i \rightarrow A$ such that f_i and g_i are ϵ_i -close, i.e., $d_X(f_i(x_i), g_i(x_i)) \leq \epsilon_i$ for all $x_i \in A_i$.*

Proof. Assume $(X_i, A_i) \rightarrow_{GH} (X, A)$. Then there exists a sequence of numbers $\{\epsilon_i\}_{i \in \mathbb{N}}$ that tends to 0 and ϵ_i -Gromov–Hausdorff approximations $f_i: X_i \rightarrow X$ such that $d_H(f_i(A_i), A) < \epsilon_i$. Here, d_H denotes the Hausdorff distance. We define $g_i: A_i \rightarrow A$ as follows. Since $d_H(f_i(A_i), A) < \epsilon_i$, for each $a_i \in A_i$ choose

$g_i(a_i) \in A$ such that $d_X(f_i(a_i), g_i(a_i)) < \epsilon_i$. We will show that the map $g_i: A_i \rightarrow A$ is a $3\epsilon_i$ -Gromov–Hausdorff approximation. Indeed, for each $a \in A$, there exists an $a_i \in A_i$ such that $d_X(f_i(a_i), a) \leq \epsilon$. Therefore, by the triangle inequality, we get $d_X(g_i(a_i), a) \leq 3\epsilon_i$. Now we show that g_i is almost distance preserving up to an error $3\epsilon_i$. We only show one side of the inequality, for the other is similar. Let $a_i, a'_i \in A_i$. Then

$$d_X(g_i(a_i), g_i(a'_i)) \leq d_X(g_i(a_i), f_i(a_i)) + d_X(f_i(a_i), f_i(a'_i)) + d_X(f_i(a'_i), g_i(a'_i)).$$

Now note that $d_X(g_i(a_i), f_i(a_i)) \leq \epsilon_i$ and $d_X(f_i(a'_i), g_i(a'_i)) \leq \epsilon_i$. Thus, as f_i is an ϵ_i -Gromov–Hausdorff approximation, we get

$$d_X(g_i(a_i), g_i(a'_i)) \leq 3\epsilon_i + d_{X_i}(a_i, a'_i).$$

Now we show the reverse direction. Define $\xi_i: X_i \rightarrow X$ by

$$\xi_i(x_i) = \begin{cases} f_i(x_i) & x_i \in X_i \setminus A_i \\ g_i(x_i) & x_i \in A_i. \end{cases}$$

Since $g_i: A_i \rightarrow A$ is an ϵ_i -Gromov–Hausdorff approximation, it follows that $d_H(g_i(A_i), A) \leq \epsilon_i$. It remains to verify that ξ_i is a Gromov–Hausdorff approximation. It is clear that ξ_i is ϵ_i -surjective. Therefore, it remains to prove that ξ_i is almost distance preserving up to an error that tends to 0. Once again, we only verify one direction of the inequality, since the other is similar. Let $x_i, x'_i \in X_i$. It is clear that it suffices to assume that $x_i \in X_i \setminus A_i$ and that $x'_i \in A_i$. Then

$$d_X(\xi_i(x_i), \xi_i(x'_i)) = d_X(f_i(x_i), g_i(x'_i)) \leq d_X(f_i(x_i), f_i(x'_i)) + d_X(f_i(x'_i), g_i(x'_i)).$$

Since f_i is an ϵ_i -Gromov–Hausdorff approximation, $d_X(f_i(x_i), f_i(x'_i)) \leq \epsilon_i + d_{X_i}(x_i, x'_i)$. Since f_i and g_i are ϵ_i -close, $d_X(f_i(x'_i), g_i(x'_i)) \leq \epsilon_i$. \square

The proposition above suggests the following definition of relative equivariant Gromov–Hausdorff convergence.

Definition 2.2. Let X and Y be compact metric spaces. Suppose G_X and G_Y are closed subgroups of $\text{Isom}(X)$ and $\text{Isom}(Y)$ respectively. Assume A_X is a closed non-empty G_X -invariant subset of X , and A_Y is a closed non-empty G_Y -invariant subset of Y . For $\epsilon > 0$, we define an ϵ -equivariant Gromov–Hausdorff approximation between (X, A_X, G_X) and (Y, A_Y, G_Y) to be a quadruple of maps

$$(f: X \rightarrow Y, f': A_X \rightarrow A_Y, \theta: G_X \rightarrow G_Y, \psi: G_Y \rightarrow G_X)$$

subject to the following conditions:

- (1) The triple (f, θ, ψ) is an ϵ -equivariant Gromov–Hausdorff approximation (in the usual sense).
- (2) f' is an ϵ -Gromov–Hausdorff approximation.
- (3) The maps f and f' are ϵ -close.

Remark 2.3. We use the terms “equivariant Gromov–Hausdorff approximations” and “equivariant approximations” interchangeably.

Definition 2.4. Given a sequence of triples $\{(X_i, A_i, G_i)\}_{i \in \mathbb{N}}$, where X_i is a compact metric space, G_i is a closed subgroup of $\text{Isom}(X_i)$ and A_i is a closed non-empty G_i -invariant subset of X_i , we say that (X_i, A_i, G_i) Gromov–Hausdorff equivariantly converges to (X, A, G) , denoted $(X_i, A_i, G_i) \rightarrow_{eGH} (X, A, G)$, where X

is a compact metric space, G is a closed subgroup of $\text{Isom}(X)$ and A is a closed subset of X that is G -invariant, if there exists a sequence of positive real numbers $\{\epsilon_i\}_{i \in \mathbb{N}}$ such that $\epsilon_i \rightarrow 0$ and ϵ_i -equivariant Gromov–Hausdorff approximations $(f_i, f'_i, \theta_i, \psi_i): (X_i, A_i, G_i) \rightarrow (X, A, G)$.

2.2. Further definitions and auxiliary results. For the convenience of the reader, we include the definitions of *Wasserstein spaces*, *good transport behavior* and *isometric rigidity*.

Definition 2.5. Let (X, d) be a Polish metric space and let $p \in [1, \infty)$. Given any two probability measure μ and ν on X , we define the L^p -Wasserstein metric between μ and ν , denoted by W_p , to be

$$W_p^p(\mu, \nu) = \inf_{\pi} \int d(x, y)^p d\pi(x, y),$$

where the infimum is taken over all admissible measures π having marginals μ and ν . The L^p -Wasserstein metric is a metric on the space of probability measures with finite p -moments, which is denoted by $\mathbb{P}_p(X)$.

Definition 2.6. A metric measure space (X, d, \mathfrak{m}) is said to have *good transport behavior* if, for any two probability measures μ, ν in $\mathbb{P}_2(X)$, where $\mu \ll \mathfrak{m}$, any optimal transport map between them is induced by a map.

We motivate the concept of isometric rigidity by the following example.

Example 2.7. If X is a compact metric space and $f: X \rightarrow X$ is an isometry, then f induces an isometry $f_{\#}: \mathbb{P}_2(X) \rightarrow \mathbb{P}_2(X)$.

Thus, it is natural to formulate the following definition.

Definition 2.8. Let (X, d) be a metric space. Then, X is said to be *isometrically rigid* if given any isometry $\theta: \mathbb{P}_2(X) \rightarrow \mathbb{P}_2(X)$, there exists an isometry $f: X \rightarrow X$ such that $\theta = f_{\#}$. In particular, every isometry of $\mathbb{P}_2(X)$ is induced by an isometry of X ; the base space.

The concept of isometric rigidity was first introduced by Kloeckner in the context of euclidean space in [27], where he showed, among other things, that, $\mathbb{P}_2(\mathbb{R})$ is not isometrically rigid. Later, Bertrand and Kloeckner [4] studied the concept of isometric rigidity and showed that if X is a negatively curved geodesically complete Hadamard space, then $\mathbb{P}_2(X)$ is isometrically rigid. In 2021, Santos Rodriguez showed [30] that if X is a positively curved closed Riemannian manifold, then X is isometrically rigid and that, if X is a CROSS, then, in fact, the isometry groups of $\mathbb{P}_p(X)$ and X are the same for all $p > 1$. In the same paper, Santos Rodriguez showed the following result which we will use frequently in this paper (Corollary 3.8, [30]).

Proposition 2.9. *Let (X, d_X, \mathfrak{m}_X) and (Y, d_Y, \mathfrak{m}_Y) be two compact non-branching metric measure spaces equipped with qualitatively non-degenerate measures and such that they have good transport behavior. Suppose that there exists an isometry $\Phi: \mathbb{P}_2(X) \rightarrow \mathbb{P}_2(Y)$. Then (X, d_X) and (Y, d_Y) are isometric.*

In fact, Santos Rodriguez in [30] showed the above proposition for Wasserstein spaces with finite p -moments, where $p \in (1, \infty)$.

3. PROOFS

In what follows, we assume that the maps are sufficiently regular. Indeed, this is possible because if X is a compact metric space then $\text{Isom}(X)$ is compact with the compact-open topology. Moreover, the compact

open topology on $\text{Isom}(X)$ can be metrized with the uniform metric. One can always replace ϵ -isometries between compact metric spaces by almost-isometries that are measurable. In particular, we have the following simple lemma.

Lemma 3.1. *Assume X and Y are compact metric spaces and G_X and G_Y are closed subgroups of $\text{Isom}(X)$ and $\text{Isom}(Y)$ respectively. Equip G_X and G_Y with the uniform metrics d_{G_X} and d_{G_Y} respectively. If $(f: X \rightarrow Y, \theta: G_X \rightarrow G_Y, \psi: G_Y \rightarrow G_X)$ is an ϵ -equivariant approximation, then, f , θ and ψ can be chosen to be measurable.*

Proof. By Corollary 3.4 in [2], it follows that θ is an 5ϵ -approximation. Furthermore, observe that ψ is an almost inverse to θ . In particular, for $\lambda \in G_Y$, and $g \in G_X$, one has $d_{G_Y}(\theta(\psi(\lambda)), \lambda) \leq 4\epsilon$ and $d_{G_X}(\psi(\theta(g)), g) \leq 4\epsilon$. Hence, as shown in Lemma 4.1 in [11], one can obtain measurable approximations $f_1: X \rightarrow Y$, $\theta_1: G_X \rightarrow G_Y$ and $\psi_1: G_Y \rightarrow G_X$ that remain close to f , θ and ψ respectively. Hence the result follows. \square

Proof of Theorem D. The implication (2) \implies (1) is well known (see Corollary 4.3 in [28]). Let us show (1) \implies (2). Assume for the sake of obtaining a contradiction that $\mathbb{P}_2(X_i) \rightarrow_{GH} \mathbb{P}_2(X)$ holds but $X_i \not\rightarrow_{GH} X$ does not hold. Hence, there exists a $\delta > 0$ such that (up to a subsequence), $d_{GH}(X_i, X) \geq \delta > 0$ for all i . Since the class \mathfrak{X} is pre-compact, and by our assumptions, it follows that up to a further subsequence, $X_i \rightarrow_{GH} X'$, where X' is a compact non-branching metric space. Hence, $\mathbb{P}_2(X_i) \rightarrow_{GH} \mathbb{P}_2(X')$. By assumption, $\mathbb{P}_2(X)$ and $\mathbb{P}_2(X')$ are isometric. Now, by Proposition 2.9, X and X' are therefore isometric, which is a contradiction. \square

Proof of Theorem A. The proof is similar to the proof of Theorem D. However, we give details for the convenience of the reader. The implications (2) \implies (1) and (3) \implies (1) are clear. The implication (2) \implies (3) follows at once from the following observation. Let $f_i: X_i \rightarrow X$ be ϵ_i -approximations by measurable maps. Then each f_i induces an $\tilde{\epsilon}_i$ -approximation $(f_i)_\# : \mathbb{P}_2(X_i) \rightarrow \mathbb{P}_2(X)$, with $\tilde{\epsilon}_i \rightarrow 0$, and such that, on Dirac deltas, $(f_i)_\#$ is a Hausdorff approximation. Hence, one may use Proposition 2.1 to establish this implication. We shall now show (1) \implies (2).

We must prove that $X_i \rightarrow_{GH} X$. Assume otherwise. Hence, there exists $\delta > 0$ such that, up to a subsequence, $d_{GH}(X_i, X) \geq \delta > 0$ for all i . Here, d_{GH} denotes the Gromov–Hausdorff distance. Since $\mathcal{A}^\leq(n, K, D)$ is, with respect to d_{GH} , compact [6], there exists a compact Alexandrov space X' such that, up to a further subsequence, $X_i \rightarrow_{GH} X'$. Hence, $\mathbb{P}_2(X_i) \rightarrow_{GH} \mathbb{P}_2(X')$. By assumption, $\mathbb{P}_2(X_i) \rightarrow_{GH} \mathbb{P}_2(X)$. Therefore, $\mathbb{P}_2(X')$ and $\mathbb{P}_2(X)$ are isometric. Alexandrov spaces have good transport behavior, and on them, the Hausdorff measure is qualitatively non-degenerate, therefore X and X' are isometric. Hence we have $X_i \rightarrow_{GH} X$. \square

Proof of Corollary B. The result follows from Perelman’s stability Theorem. Note that if $f: X \rightarrow Y$ is a map, then $f_\#$ is continuous if and only if f is continuous (see Remark 4.14 in [31]). \square

Proof of Corollary C. We proceed by contradiction. Assume we have a sequence $(\mathbb{P}_2(X_i))_{i \in \mathbb{N}}$ of topologically pairwise inequivalent Wasserstein spaces in $W_2(n, K, D, v)$. Then, since $\mathcal{A}(n, K, D, v)$ is precompact and we have volume bounded below by $v > 0$, there exists a compact n -dimensional Alexandrov space X such that (up to a subsequence), $X_i \rightarrow_{GH} X$. Hence, $\mathbb{P}_2(X_i) \rightarrow_{GH} \mathbb{P}_2(X)$. Therefore, by Corollary B, for all large i , $\mathbb{P}_2(X_i)$ and $\mathbb{P}_2(X)$ are homeomorphic, which is a contradiction. \square

Proof of Corollary E. This follows in an analogous manner as in the proof of Corollary B. However, one uses the result in [24] instead of Perelman's stability theorem. \square

Now we will prove Theorem F. First, we need two lemmas.

Lemma 3.2. *Let X and Y be compact metric spaces and G_X and G_Y be closed subgroups of $\text{Isom}(X)$ and $\text{Isom}(Y)$ respectively. An ϵ -equivariant approximation, (f, θ, ψ) between (X, G_X) and (Y, G_Y) induces an $\tilde{\epsilon}$ -equivariant approximation $(f_\#, \theta_\#, \psi_\#)$ between $(\mathbb{P}_2(X), G_X^\#)$ and $(\mathbb{P}_2(Y), G_Y^\#)$, where $\tilde{\epsilon}$ tends to 0 as ϵ tends to 0.*

Proof. Define $\theta_\#$ and $\psi_\#$ by the rules $\theta_\#(g_\#) = (\theta(g))_\#$ and $\psi_\#(\lambda_\#) = (\psi(\lambda))_\#$. By Corollary 4.3 in [28], the map $f_\#$ is an $\tilde{\epsilon}$ -approximation, where

$$\tilde{\epsilon} = 6\epsilon + \sqrt{\epsilon(2 \text{diam}(Y) + \epsilon)}.$$

Let $\mu \in \mathbb{P}_2(X)$. Then,

$$W_2^2(\theta_\#(g_\#)(f_\#\mu), f_\#(g_\#\mu)) \leq \int d^2(\theta(g)(f(x)), f(gx))d\mu \leq \epsilon^2.$$

The last inequality is similar. \square

As a corollary, we have the following result.

Corollary 3.3. *Let X_i and X be compact metric spaces. If $(X_i, G_i) \rightarrow_{eGH} (X, G)$, then $(\mathbb{P}_2(X_i), G_i^\#) \rightarrow_{eGH} (\mathbb{P}_2(X), G^\#)$.*

We will also need the following lemma, an equivariant analogue of Proposition 2.9.

Lemma 3.4. *Let X and Y be positively curved closed Riemannian manifolds and let G_X and G_Y be closed subgroups of $\text{Isom}(X)$ and $\text{Isom}(Y)$, respectively. If $(\mathbb{P}_2(X), G_X^\#)$ and $(\mathbb{P}_2(Y), G_Y^\#)$ are equivariantly isometric then so are (X, G_X) and (Y, G_Y) .*

Proof. Let $\Phi: \mathbb{P}_2(X) \rightarrow \mathbb{P}_2(Y)$ be an isometry and let $\Theta: G_X^\# \rightarrow G_Y^\#$ be an isomorphism such that for any $g \in \text{Isom}(X)$, one has $\Phi \circ g_\# = \Theta(g_\#) \circ \Phi$. Then, from Corollary 3.8 in [30], it follows that Φ sends Dirac deltas onto Dirac deltas. Thus, there exists an isometry $F: X \rightarrow Y$ such that $F_\#$ and Φ agree on Dirac deltas. Moreover, there exists an isomorphism $\theta: G_X \rightarrow G_Y$ such that for $g \in G_X$, $\theta(g)_\# = \Theta(g_\#)$. Thus it follows that $F \circ g = \theta(g) \circ F$. \square

We are now ready to prove theorem F.

Proof of Theorem F. The implication (2) \implies (1) follows from Lemma 3.2. The implication (2) \implies (3) is trivial. Now we shall verify the implication (1) \implies (2). Since X_i and X are isometrically rigid, there exists unique closed subgroups G_i of $\text{Isom}(X_i)$ and G of $\text{Isom}(X)$ such that $G_i^\# = H_i$ and $G^\# = H$. It now remains to verify that $(X_i, G_i) \rightarrow_{eGH} (X, G)$. As usual, assume otherwise. So, there exists a $\delta > 0$ such that, up to a subsequence, $d_{eGH}((X_i, G_i), (X, G)) \geq \delta$ for all i . Here, d_{eGH} denotes the equivariant Gromov–Hausdorff distance. Since $\{X_i\}_i$ forms a family of compact Alexandrov spaces with curvature uniformly bounded below, dimension uniformly bounded above, and uniform upper diameter bound, it follows that there exists a compact Alexandrov space X' , such that, up to a subsequence, $X_i \rightarrow_{GH} X'$. Up to a further

subsequence, there exists a closed subgroup G' of $\text{Isom}(X')$ such that $(X_i, G_i) \rightarrow_{eGH} (X', G')$. Hence, by Lemma 3.2, it follows that $(\mathbb{P}_2(X_i), H_i) \rightarrow_{eGH} (\mathbb{P}_2(X'), (G')^\#)$. Therefore, $(\mathbb{P}_2(X'), (G')^\#)$ and $(\mathbb{P}_2(X), G^\#)$ are equivariantly isometric. The previous lemma shows that (X', G') and (X, G) are equivariantly isometric, which is a contradiction. \square

Proof of Corollary G. Theorem F ensures there exists closed subgroups G_i and G such that $G_i^\# = H_i$, $G^\# = H$ and such that $(X_i, G_i) \rightarrow_{eGH} (X, G)$. Note that the isometric rigidity of the X_i and X ensures that G_i and G have the same dimension as H_i and H . Hence the result follows from Theorem A in [2]. \square

REFERENCES

- [1] Andrés Ahumada Gómez and Mauricio Che. “Gromov–Hausdorff convergence of metric pairs and metric tuples”. In: *Differential Geom. Appl.* 94 (2024), Paper No. 102135. ISSN: 0926-2245. DOI: [10.1016/j.difgeo.2024.102135](https://doi.org/10.1016/j.difgeo.2024.102135). URL: <https://doi.org/10.1016/j.difgeo.2024.102135>.
- [2] Mohammad Alattar. “Stability and equivariant Gromov–Hausdorff convergence”. In: *Bulletin of the London Mathematical Society* 47 (2024). DOI: <https://doi.org/10.1112/blms.13073>. URL: <https://doi.org/10.1112/blms.13073>.
- [3] Michael T. Anderson and Jeff Cheeger. “Diffeomorphism finiteness for manifolds with Ricci curvature and $L^{n/2}$ -norm of curvature bounded”. In: *Geom. Funct. Anal.* 1.3 (1991), pp. 231–252. ISSN: 1016-443X. DOI: [10.1007/BF01896203](https://doi.org/10.1007/BF01896203). URL: <https://doi.org/10.1007/BF01896203>.
- [4] Jérôme Bertrand and Benoît R. Kloeckner. “A geometric study of Wasserstein spaces: isometric rigidity in negative curvature”. In: *Int. Math. Res. Not. IMRN* 5 (2016), pp. 1368–1386. ISSN: 1073-7928. DOI: [10.1093/imrn/rnv177](https://doi.org/10.1093/imrn/rnv177). URL: <https://doi.org/10.1093/imrn/rnv177>.
- [5] Elia Brue, Aaron Naber, and Daniele Semola. “Stability of Tori under Lower Sectional Curvature”. In: (2023). arXiv: [2307.03824](https://arxiv.org/abs/2307.03824) [math.DG].
- [6] Yuri. Burago, Misha Gromov, and G. Perelman. “A. D. Aleksandrov spaces with curvatures bounded below”. In: *Uspekhi Mat. Nauk* 47.2(284) (1992), pp. 3–51, 222. ISSN: 0042-1316,2305-2872. DOI: [10.1070/RM1992v047n02ABEH000877](https://doi.org/10.1070/RM1992v047n02ABEH000877). URL: <https://doi.org/10.1070/RM1992v047n02ABEH000877>.
- [7] Fabio Cavalletti and Martin Huesmann. “Existence and uniqueness of optimal transport maps”. In: *Ann. Inst. H. Poincaré C Anal. Non Linéaire* 32.6 (2015), pp. 1367–1377. ISSN: 0294-1449. DOI: [10.1016/j.anihpc.2014.09.006](https://doi.org/10.1016/j.anihpc.2014.09.006). URL: <https://doi.org/10.1016/j.anihpc.2014.09.006>.
- [8] Nicola Cavallucci. *A GH-compactification of CAT(0)-groups via totally disconnected, unimodular actions*. 2023. arXiv: [2307.05640](https://arxiv.org/abs/2307.05640) [math.MG].
- [9] Mauricio Che et al. *Metric Geometry of Spaces of Persistence Diagrams*. 2022. arXiv: [2109.14697](https://arxiv.org/abs/2109.14697) [math.MG].
- [10] Jeff Cheeger. “Finiteness theorems for Riemannian manifolds”. In: *Amer. J. Math.* 92 (1970), pp. 61–74. ISSN: 0002-9327. DOI: [10.2307/2373498](https://doi.org/10.2307/2373498). URL: <https://doi.org/10.2307/2373498>.
- [11] Nhan-Phu Chung. “Gromov-Hausdorff distances for dynamical systems”. In: *Discrete Contin. Dyn. Syst.* 40.11 (2020), pp. 6179–6200. ISSN: 1078-0947. DOI: [10.3934/dcds.2020275](https://doi.org/10.3934/dcds.2020275). URL: <https://doi.org/10.3934/dcds.2020275>.
- [12] Diego Corro. *Collapsing regular Riemannian foliations with flat leaves*. 2024. arXiv: [2403.11602](https://arxiv.org/abs/2403.11602) [math.DG].
- [13] Diego Corro, Jaime Santos-Rodríguez, and Jesús Núñez-Zimbrón. *Cohomogeneity one RCD-spaces*. 2024. arXiv: [2405.09448](https://arxiv.org/abs/2405.09448) [math.MG].
- [14] Qin Deng. *Hölder continuity of tangent cones in RCD(K,N) spaces and applications to non-branching*. 2020. arXiv: [2009.07956](https://arxiv.org/abs/2009.07956) [math.DG].

- [15] Kenji Fukaya. “Hausdorff convergence of Riemannian manifolds and its applications”. In: *Recent topics in differential and analytic geometry*. Vol. 18. Adv. Stud. Pure Math. Academic Press, Boston, MA, 1990, pp. 143–238. DOI: [10.2969/aspm/01810143](https://doi.org/10.2969/aspm/01810143). URL: <https://doi.org/10.2969/aspm/01810143>.
- [16] Kenji Fukaya. “Theory of convergence for Riemannian orbifolds”. In: *Japan. J. Math. (N.S.)* 12.1 (1986), pp. 121–160. ISSN: 0289-2316. DOI: [10.4099/math1924.12.121](https://doi.org/10.4099/math1924.12.121). URL: <https://doi.org/10.4099/math1924.12.121>.
- [17] Kenji Fukaya and Takao Yamaguchi. “The fundamental groups of almost non-negatively curved manifolds”. In: *Ann. of Math. (2)* 136.2 (1992), pp. 253–333. ISSN: 0003-486X. DOI: [10.2307/2946606](https://doi.org/10.2307/2946606). URL: <https://doi.org/10.2307/2946606>.
- [18] Fernando Galaz-García et al. “On quotients of spaces with Ricci curvature bounded below”. In: *J. Funct. Anal.* 275.6 (2018), pp. 1368–1446. ISSN: 0022-1236. DOI: [10.1016/j.jfa.2018.06.002](https://doi.org/10.1016/j.jfa.2018.06.002). URL: <https://doi.org/10.1016/j.jfa.2018.06.002>.
- [19] Mikhael Gromov. “Groups of polynomial growth and expanding maps”. In: *Inst. Hautes Études Sci. Publ. Math.* 53 (1981), pp. 53–73. ISSN: 0073-8301. URL: http://www.numdam.org/item?id=PMIHES_1981__53__53_0.
- [20] Karsten Grove. “Finiteness theorems in Riemannian geometry”. In: *Explorations in complex and Riemannian geometry*. Vol. 332. Contemp. Math. Amer. Math. Soc., Providence, RI, 2003, pp. 101–120. DOI: [10.1090/conm/332/05931](https://doi.org/10.1090/conm/332/05931). URL: <https://doi.org/10.1090/conm/332/05931>.
- [21] Karsten Grove and Peter Petersen V. “Bounding homotopy types by geometry”. In: *Ann. of Math. (2)* 128.1 (1988), pp. 195–206. ISSN: 0003-486X. DOI: [10.2307/1971439](https://doi.org/10.2307/1971439). URL: <https://doi.org/10.2307/1971439>.
- [22] Karsten Grove, Peter Petersen V, and Jyh Yang Wu. “Geometric finiteness theorems via controlled topology”. In: *Invent. Math.* 99.1 (1990), pp. 205–213. ISSN: 0020-9910. DOI: [10.1007/BF01234417](https://doi.org/10.1007/BF01234417). URL: <https://doi.org/10.1007/BF01234417>.
- [23] John Harvey. “Convergence of isometries, with semicontinuity of symmetry of Alexandrov spaces”. In: *Proceedings of the American Mathematical Society* 144.8 (2016), pp. 3507–3515.
- [24] Shouhei Honda and Yuanlin Peng. “A note on the topological stability theorem from RCD spaces to Riemannian manifolds”. In: *Manuscripta Math.* 172.3-4 (2023), pp. 971–1007. ISSN: 0025-2611. DOI: [10.1007/s00229-022-01418-7](https://doi.org/10.1007/s00229-022-01418-7). URL: <https://doi.org/10.1007/s00229-022-01418-7>.
- [25] Vitali Kapovitch. “Perelman’s stability theorem”. In: *Surveys in differential geometry. Vol. XI*. Vol. 11. Surv. Differ. Geom. Int. Press, Somerville, MA, 2007, pp. 103–136. DOI: [10.4310/SDG.2006.v11.n1.a5](https://doi.org/10.4310/SDG.2006.v11.n1.a5). URL: <https://doi.org/10.4310/SDG.2006.v11.n1.a5>.
- [26] Martin Kell. “Transport maps, non-branching sets of geodesics and measure rigidity”. In: *Adv. Math.* 320 (2017), pp. 520–573. ISSN: 0001-8708. DOI: [10.1016/j.aim.2017.09.003](https://doi.org/10.1016/j.aim.2017.09.003). URL: <https://doi.org/10.1016/j.aim.2017.09.003>.
- [27] Benoît Kloeckner. “A geometric study of Wasserstein spaces: Euclidean spaces”. In: *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* 9.2 (2010), pp. 297–323. ISSN: 0391-173X.
- [28] John Lott and Cédric Villani. “Ricci curvature for metric-measure spaces via optimal transport”. In: *Ann. of Math. (2)* 169.3 (2009), pp. 903–991. ISSN: 0003-486X. DOI: [10.4007/annals.2009.169.903](https://doi.org/10.4007/annals.2009.169.903). URL: <https://doi.org/10.4007/annals.2009.169.903>.
- [29] Grigori Perelman. “Alexandrov spaces with curvatures bounded from below II”. In: *preprint* (1991).
- [30] Jaime Santos-Rodríguez. “On isometries of compact L^p -Wasserstein spaces”. In: *Adv. Math.* 409 (2022), Paper No. 108632, 21. ISSN: 0001-8708,1090-2082. DOI: [10.1016/j.aim.2022.108632](https://doi.org/10.1016/j.aim.2022.108632). URL: <https://doi.org/10.1016/j.aim.2022.108632>.
- [31] Cédric Villani et al. *Optimal transport: old and new*. Vol. 338. Springer, 2009.

- [32] Alan Weinstein. “On the homotopy type of positively-pinched manifolds”. In: *Arch. Math. (Basel)* 18 (1967), pp. 523–524. ISSN: 0003-889X. DOI: [10.1007/BF01899493](https://doi.org/10.1007/BF01899493). URL: <https://doi.org/10.1007/BF01899493>.
- [33] Sergio Zamora. “Anderson finiteness for RCD spaces”. In: *MPIM Preprint Series* (4) (2023). URL: https://archive.mpim-bonn.mpg.de/id/eprint/5009/1/mpim-preprint_2023-4.pdf.
- [34] Sergio Zamora. “First Betti number and collapse”. In: (2022). arXiv: [2209.12628](https://arxiv.org/abs/2209.12628) [[math.DG](https://arxiv.org/abs/2209.12628)].

(Al-Attar) DEPARTMENT OF MATHEMATICAL SCIENCES, DURHAM UNIVERSITY, UNITED KINGDOM
Email address: mohammad.al-attar@durham.ac.uk