# Bayesian Parametric Methods for Deriving Distribution of Restricted Mean Survival Time

Keisuke Hanada<sup>1</sup> and Masahiro Kojima<sup>\*2</sup>

<sup>1</sup>Osaka University <sup>2</sup>The Institute of Statistical Mathematics

June 12, 2024

#### Abstract

We propose a Bayesian method for deriving the distribution of restricted mean survival time (RMST) using posterior samples, which accounts for covariates and heterogeneity among clusters based on a parametric model for survival time. We derive an explicit RMST equation by devising an integral of the survival function, allowing for the calculation of not only the mean and credible interval but also the mode, median, and probability of exceeding a certain value. Additionally, We propose two methods: one using random effects to account for heterogeneity among clusters and another utilizing frailty. We developed custom Stan code for the exponential, Weibull, log-normal frailty, and log-logistic models, as they cannot be processed using the brm functions in R. We evaluate our proposed methods through computer simulations and analyze real data from the eight Empowered Action Group states in India to confirm consistent results across states after adjusting for cluster differences.

In conclusion, we derived explicit RMST formulas for parametric models and their distributions, enabling the calculation of the mean, median, mode, and credible interval. Our simulations confirmed the robustness of the proposed methods, and using the shrinkage effect allowed for more accurate results for each cluster.

Keywords: restricted mean survival time, frailty model, Weibull model, log-logistic model, log-normal model.

# 1 Introduction

The restricted mean survival time (RMST) up to a pre-specified evaluation time point allows for the assessment of survival time using an intuitive measure of the mean [1]. RMST can serve

<sup>\*</sup>Address:10-3 Midori-cho, Tachikawa, Tokyo 190-8562, Japan. Tel: +81-(0)50-5533-8500 E-Mail: ko-jima.masahiro@ism.ac.jp

as a valuable endpoint in Phase II and III randomized controlled studies. Most RMST methodological research is based on frequentist theory [2–4], allowing for the adjustment of baseline covariates and heterogeneity among clusters. However, these methods do not support deriving the distribution of RMST, estimating the probability of RMST exceeding a specific threshold, or utilizing prior distributions. While frequentist theory offers differences in means and confidence intervals through normal approximation, we aim to derive more exact distributions using Bayesian analysis. Although a non-parametric Bayesian method for RMST estimation has been proposed [5], it cannot handle covariate adjustment or cluster heterogeneity.

In this paper, we propose a Bayesian method for deriving the distribution of RMST using posterior samples. Because it is difficult to fit RMST directly to a statistical model, we present a Bayesian representation that can account for covariates and heterogeneity among clusters based on a parametric model for survival time. The RMST requires integrating the survival function up to the evaluation point. Although integrating the survival function is challenging for most parametric distributions, we use a formula that facilitates calculating the integral for all survival functions by transforming the existing equation. No explicit equation for RMST has ever been proposed, and this equation allows us to calculate the distribution of RMST using posterior samples. By obtaining the RMST distribution, we can compute not only the mean and confidence interval but also the mode, median, and probability of exceeding a certain value. It is often necessary to account for heterogeneity among clusters or to use shrinkage effects to increase estimation accuracy for clusters with small sample sizes. Hence, we introduce two methods for RMST: one using a random effect to account for heterogeneity among clusters and another utilizing frailty. We evaluate the performance of the RMST distribution using posterior samples in simulations and present the results of an analysis based on real data from the eight Empowered Action Group (EAG) states in India. We develop custom Stan source code to analyze the exponential, Weibull, log-normal frailty models and all log-logistic models, as they cannot be processed using the brm functions.

This paper is organized as follows. Section 2 introduces the explicit RMST formula calculation, Bayesian sampling, and RMST distribution. Section 3 details the settings and results of computer simulations. Section 4 presents an analysis plan and the results of analyzing data from the eight EAG states in India. Finally, Section 5 discusses all the results.

# 2 Method

#### 2.1 Restricted mean survival time

The RMST is defined as the integral of any survival function S(t) from 0 to  $\tau$ , where  $\tau > 0$  is the pre-specified time point. We assume a parametric model for the survival time. The RMST requires integrating the survival function, but this integral of the parametric survival function can generally be complex. Therefore, we use the following formula to calculate the RMST.

$$RMST(\tau) = \int_0^\tau S(t)dt = \int_0^\tau tf(t)dt + \tau S(\tau), \tag{1}$$

where f(t) is a density function for a random survival time T. Equation (1) implies that the RMST can be calculated for any distribution for which the process for deriving the expectation

is known. The equation (1) is obtained from the following integral.

$$\int_0^\tau tf(t)dt = [-tS(t)]_0^\tau + \int_0^\tau S(t)dt = -\tau S(\tau) + \int_0^\tau S(t)dt.$$

This paper deals with four parametric models: exponential, Weibull, log-logistic, and lognormal. We demonstrate the RMSTs of these models. The RMSTs for models other than these four may be calculated using the formula (1).

[Exponential distribution] The RMST for an exponential distribution  $T \sim Exp(\lambda)$  can be easily obtained, where a rate parameter  $\lambda > 0$ .

$$RMST_E(\tau) = \int_0^{\tau} S(t)dt = \frac{1 - e^{-\lambda\tau}}{\lambda}.$$

When  $\tau$  approaches infinitely, it becomes apparent that the expected value of the exponential distribution is obtained. The derivation is shown in the Appendix ?? in the Supplemental material. Note that it may not be necessary to apply the equation 1 because the survival function of the exponential distribution can be easily integrated.

**[Weibull distribution]** Next, we consider the RMST of the Weibull distribution  $T \sim W(\lambda, k)$ , where  $\lambda$  is the scale parameter ( $\lambda > 0$ ) and k is the shape parameter (k > 0). The RMST is challenging to calculate without using formula (1).

$$RMST_W(\tau) = \int_0^\tau S(t)dt = \lambda^{-\frac{1}{k}}\gamma\left(\lambda\tau^k; \frac{1}{k} + 1\right) + \tau \exp(-\lambda\tau^k),$$

where  $\gamma(z; a) = \int_0^z t^{a-1} e^{-t} dt$  is the incomplete gamma function. The incomplete gamma function can be computed in software such as R or Python. The derivation of the RMST is found in the Appendix ?? in the Supplemental material. By making  $\tau$  sufficiently large, the RMST coincides with the expectation of the Weibull distribution. The RMST can be obtained by substituting the estimated values into the parameters. Therefore, the ability to calculate the RMST using an explicit formula, even for distributions where directly integrating the survival function is challenging, is an advantage. We can also see that in subsequent distributions, we can calculate the RMST without using approximations.

[Log-logistic distribution] We show the RMST for the log-logistic distribution  $T \sim LL(\mu, k)$ , where  $\mu$  is the scale parameter ( $\mu \in \mathbf{R}$ ) and k is the shape parameter (k > 0).

$$RMST_{LL}(\tau) = \int_0^{\tau} S(t)dt = e^{-\frac{\mu}{k}} B\left(\frac{e^{\mu}\tau^k}{1+e^{\mu}\tau^k}; 1+\frac{1}{k}, 1-\frac{1}{k}\right) + \tau \frac{1}{1+e^{\mu}\tau^k},$$

where  $B(z; a, b) = \int_0^z t^{a-1}(1-t)^{b-1}dt$  is the incomplete beta function. The incomplete gamma function can also be computed in software such as R or Python. The derivation can be found in the Appendix ?? in the Supplemental material. Based on the first moment condition, we restrict to k > 1. Taking the limit as  $\tau \to \infty$ , we obtain the expectation of the log-logistic distribution.

[Log-normal distribution] Finally, we show the RMST in a log-normal distribution  $T \sim LN(\mu, \sigma^2)$ , where the parameters are  $\mu \in \mathbf{R}$  and  $\sigma^2 > 0$ .

$$RMST_{LN}(\tau) = \int_0^{\tau} S(t)dt = \exp\left\{\mu + \frac{\sigma^2}{2}\right\} \Phi\left(\frac{\log(\tau) - \mu - \sigma^2}{\sigma}\right) + \tau\left(1 - \Phi\left(\frac{\log(\tau) - \mu}{\sigma}\right)\right).$$
(2)

The derivation is shown in the Appendix ?? in the Supplemental material. As  $\tau \to \infty$  on the right-hand side of the equation, equation (2) becomes the expectation of the log-normal distribution.

We also consider the RMST for each cluster with shrinkage effects via the hierarchical Bayesian method. The RMST, including random effects, is shown in Appendix ?? in the Supplemental material. Additionally, the RMST under the hazard with frailty effects is presented in Appendix ?? in the Supplemental material.

The Weibull and log-logistic distributions have different representations, and the RMSTs in the different representations are shown in Appendix ?? and ?? in the Supplemental material.

The Bayesian distribution of RMSTs is given by substituting the posterior samples into the parameters. In the next section, we will introduce the method for generating posterior samples.

#### 2.2 Posterior probability

We show a posterior probability for each model. We assume that there are M clusters and sample size of  $n_i$  for the *i*-th cluster. Additionally, to account for the heterogeneity, we apply the random effect or frailty. We assume that the random effect  $u_i$  follows a normal distribution  $\mathcal{N}(0, \phi^2)$  and the frailty  $v_i$  follows a gamma distribution  $Gamma\left(\frac{1}{\phi}, \frac{1}{\phi}\right)$ .

**[Exponential model]** We transform the parameter  $\lambda$  to account for the q covariates vector  $\boldsymbol{x}$ .

$$\lambda_{ij} = \exp\{\boldsymbol{x}_{ij}^T \boldsymbol{\beta}\} \tag{3}$$

where *i* represents the *i*-th cluster, *j* represents the *j*-th subject in *i*-th cluster,  $x_{ij}$  includes the coefficient corresponding to the intercept term and the treatment group, and baseline covariates, and  $\beta$  is a *q*-dimensional coefficient parameter vector. The random effect  $u_i$  for *i*-th cluster is added to  $x_{ij}^T \beta$ .

The posterior probability is

$$\pi_r(\boldsymbol{\beta}, \boldsymbol{u}, \boldsymbol{\phi} | \boldsymbol{t}, \boldsymbol{X}) \propto L_r(\boldsymbol{t} | \boldsymbol{X}, \boldsymbol{\beta}, \boldsymbol{u}, \boldsymbol{\phi}) g_r(\boldsymbol{u} | \boldsymbol{\phi}) \pi(\boldsymbol{\beta} | \boldsymbol{c}) \pi(\boldsymbol{\phi} | \boldsymbol{\xi}),$$

where

$$L_r(\boldsymbol{t}|\boldsymbol{X},\boldsymbol{\beta},\boldsymbol{u},\boldsymbol{\phi}) = \prod_{i=1}^M \prod_{j=1}^{n_i} \{f_r(t_{ij}|v_i)\}^{\delta_{ij}} \{S_r(t_{ij}|v_i)\}^{1-\delta_{ij}}$$
$$= \prod_{i=1}^M \prod_{j=1}^{n_i} \left[\exp\{\boldsymbol{x}_{ij}^T\boldsymbol{\beta} + u_i\}\exp\{-\exp(\boldsymbol{x}_{ij}^T\boldsymbol{\beta} + u_i)t_{ij}\}\right]^{\delta_{ij}}$$
$$\times \left[\exp\{-\exp(\boldsymbol{x}_{ij}^T\boldsymbol{\beta} + u_i)t_{ij}\}\right]^{1-\delta_{ij}},$$

the variable  $\delta_{ij}$  is a censoring indicator that takes the value 1 if an event occurs and 0 if censored. The details of the other functions are shown in the Appendix ?? in the Supplemental material.

The frailty model is defined as the product of the hazard function and the frailty term. The hazard function with the frailty term  $v_i$  is

$$h(t_{ij}|v_i) = v_i h(t_{ij}) = v_i \exp\{\boldsymbol{x}_{ij}^T \boldsymbol{\beta}\},\$$

where  $h(t_{ij})$  is the hazard function of the exponential distribution. The posterior probability is

$$\pi_f(\boldsymbol{eta}, \boldsymbol{v}, \phi | \boldsymbol{t}, \boldsymbol{X}) \propto L_f(\boldsymbol{t} | \boldsymbol{X}, \boldsymbol{eta}, \boldsymbol{v}, \phi) g_f(\boldsymbol{v} | \phi) \pi(\boldsymbol{eta} | \boldsymbol{c}) \pi(\phi | \xi),$$

where

$$\begin{split} L_{f}(\boldsymbol{t}|\boldsymbol{X},\boldsymbol{\beta},\boldsymbol{v},\phi) &= \prod_{i=1}^{M} \prod_{j=1}^{n_{i}} \{f_{f}(t_{ij}|v_{i})\}^{\delta_{ij}} \{S_{f}(t_{ij}|v_{i})\}^{1-\delta_{ij}} \\ &= \prod_{i=1}^{M} \prod_{j=1}^{n_{i}} \left[v_{j} \exp\{\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta}\} \exp\{-v_{j} \exp\{\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta}\}t\}\right]^{\delta_{ij}} \left[\exp\{-v_{j} \exp\{\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta}\}t\}\right]^{1-\delta_{ij}}. \end{split}$$

The details of the other functions are shown in the Appendix ?? in the Supplemental material.

**[Weibull model]** Next, we consider the Weibull model. We transform the parameter  $\lambda$  to the equation (3) to account for covariates similarly to an exponential distribution. For the mixed effects model, the posterior probability is

$$\pi_r(\boldsymbol{\beta}, \boldsymbol{u}, \boldsymbol{\phi}, k | \boldsymbol{t}, \boldsymbol{X}) \propto L_r(\boldsymbol{t} | \boldsymbol{X}, \boldsymbol{\beta}, \boldsymbol{u}, \boldsymbol{\phi}, k) g_r(\boldsymbol{u} | \boldsymbol{\phi}) \pi(\boldsymbol{\beta} | \boldsymbol{c}) \pi(\boldsymbol{\phi} | \boldsymbol{\xi}) \pi(k | \boldsymbol{a}, \boldsymbol{b}),$$

where

$$L_{r}(\boldsymbol{t}|\boldsymbol{X},\boldsymbol{\beta},\boldsymbol{u},\phi,k) = \prod_{i=1}^{M} \prod_{j=1}^{n_{i}} \{f_{r}(t_{ij}|u_{i})\}^{\delta_{ij}} \{S_{r}(t_{ij}|u_{i})\}^{1-\delta_{ij}}$$
$$= \prod_{i=1}^{M} \prod_{j=1}^{n_{i}} \left[ \frac{\exp\{\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta} + u_{i}\}kt_{ij}^{k-1}}{\left(1 + \exp(\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta} + u_{i})t^{k}\right)^{2}} \right]^{\delta_{ij}} \left[ \frac{1}{1 + \exp(\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta} + u_{i})t^{k}} \right]^{1-\delta_{ij}}.$$

The details of the other functions are shown in the Appendix ?? in the Supplemental material. The hazard function with frailty term  $v_i$  is

$$h(t_{ij}|v_i) = v_i \exp\{\boldsymbol{x}_{ij}^T \boldsymbol{\beta}\} k t^{k-1}.$$

The posterior probability is

$$\pi_f(\boldsymbol{\beta}, \boldsymbol{v}, \phi, k | \boldsymbol{t}, \boldsymbol{X}) \propto L_f(\boldsymbol{t} | \boldsymbol{X}, \boldsymbol{\beta}, \boldsymbol{v}, \phi) g_f(\boldsymbol{v} | \phi) \pi(\boldsymbol{\beta} | \boldsymbol{c}) \pi(\phi | \xi) \pi(k | a, b)$$

$$\begin{split} L_f(\boldsymbol{t}|\boldsymbol{X},\boldsymbol{\beta},\boldsymbol{v},\phi,k) &= \prod_{i=1}^M \prod_{j=1}^{n_i} \{f_f(t_{ij}|v_i)\}^{\delta_{ij}} \{S_f(t_{ij}|v_i)\}^{1-\delta_{ij}} \\ &= \prod_{i=1}^M \prod_{j=1}^{n_i} \left[ v_j \exp\{\boldsymbol{x}_{ij}^T \boldsymbol{\beta}\} k t^{k-1} \exp\left\{-v_j \exp\{\boldsymbol{x}_{ij}^T \boldsymbol{\beta}\} t^k\right\} \right]^{\delta_{ij}} \\ &\times \left[ \exp\left\{-v_j \exp\{\boldsymbol{x}_{ij}^T \boldsymbol{\beta}\} t^k\right\} \right]^{1-\delta_{ij}}. \end{split}$$

The details of the other functions are shown in the Appendix ?? in the Supplemental material.

[Log-logistic distribution] For the log-logistic distribution, we transform the parameter  $\mu$  to account for covariates.

$$\mu_{ij} = \boldsymbol{x}_{ij}^T \boldsymbol{\beta}.$$

For the random effects model, the posterior probability is

$$\pi_r(\boldsymbol{\beta}, \boldsymbol{u}, \phi, k | \boldsymbol{t}, \boldsymbol{X}) \propto L_r(\boldsymbol{t} | \boldsymbol{X}, \boldsymbol{\beta}, \boldsymbol{u}, \phi, k) g_r(\boldsymbol{u} | \phi) \pi(\boldsymbol{\beta} | \boldsymbol{c}) \pi(\phi | \xi) \pi(k | a, b),$$

$$L_{r}(\boldsymbol{t}|\boldsymbol{X},\boldsymbol{\beta},\boldsymbol{u},\phi,k) = \prod_{i=1}^{M} \prod_{j=1}^{n_{i}} \{f_{r}(t_{ij}|u_{i})\}^{\delta_{ij}} \{S_{r}(t_{ij}|u_{i})\}^{1-\delta_{ij}}$$
$$= \prod_{i=1}^{M} \prod_{j=1}^{n_{i}} \left[ \frac{\exp\{\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta} + u_{i}\}kt_{ij}^{k-1}}{\left(1 + \exp(\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta} + u_{i})t^{k}\right)^{2}} \right]^{\delta_{ij}} \left[ \frac{1}{1 + \exp(\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta} + u_{i})t^{k}} \right]^{1-\delta_{ij}}$$

The details of the other functions are shown in the Appendix ?? in the Supplemental material. For the frailty model, the hazard function with frailty term  $v_i$  is

$$h(t_{ij}|v_i) = v_i \frac{e^{\boldsymbol{x}_{ij}^T \boldsymbol{\beta}} k t^{k-1}}{1 + e^{\boldsymbol{x}_{ij}^T \boldsymbol{\beta}} t^k}$$

The posterior probability is

$$\pi_f(\boldsymbol{\beta}, \boldsymbol{v}, \phi, k | \boldsymbol{t}, \boldsymbol{X}) \propto L_f(\boldsymbol{t} | \boldsymbol{X}, \boldsymbol{\beta}, \boldsymbol{v}, \phi) g_f(\boldsymbol{v} | \phi) \pi(\boldsymbol{\beta} | \boldsymbol{c}) \pi(\phi | \xi) \pi(k | a, b),$$

where

$$L_{f}(\boldsymbol{t}|\boldsymbol{X},\boldsymbol{\beta},\boldsymbol{v},\phi,k) = \prod_{i=1}^{M} \prod_{j=1}^{n_{i}} \{f(t_{ij}|v_{i})\}^{\delta_{ij}} \{S(t_{ij}|v_{i})\}^{1-\delta_{ij}}$$
$$= \prod_{i=1}^{M} \prod_{j=1}^{n_{i}} \left[ v_{i} \exp\{\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta}\} k t_{ij}^{k-1} \left(\frac{1}{1+e^{\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta}} t^{k}}\right)^{v_{i}+1} \right]^{\delta_{ij}} \left[ \left(\frac{1}{1+e^{\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta}} t^{k}}\right)^{v_{i}} \right]^{1-\delta_{ij}}$$

The details of the other functions are shown in the Appendix ?? in the Supplemental material.

[Log-normal distribution] We transform the parameter  $\mu$  to account for covariates.

$$\mu_{ij} = \boldsymbol{x}_{ij}^T \boldsymbol{\beta}.$$

For the random effect model, the posterior probability is

$$\pi_r(\boldsymbol{\beta}, \boldsymbol{u}, \phi, \sigma^2 | \boldsymbol{t}, \boldsymbol{X}) \propto L_r(\boldsymbol{t} | \boldsymbol{X}, \boldsymbol{\beta}, \boldsymbol{u}, \phi, \sigma^2) g_r(\boldsymbol{u} | \phi) \pi(\boldsymbol{\beta} | \boldsymbol{c}) \pi(\phi | \boldsymbol{\xi}) \pi(\sigma^2 | \boldsymbol{a}, \boldsymbol{b}),$$

$$L_r(\boldsymbol{t}|\boldsymbol{X},\boldsymbol{\beta},\boldsymbol{u},\phi,\sigma^2) = \prod_{i=1}^M \prod_{j=1}^{n_i} \{f_r(t_{ij}|u_i)\}^{\delta_{ij}} \{S_r(t_{ij}|u_i)\}^{1-\delta_{ij}}$$
$$= \prod_{i=1}^M \prod_{j=1}^{n_i} \left[\frac{1}{t\sqrt{2\pi\sigma^2}} \exp\left[-\frac{\{\log(t) - (\boldsymbol{x}_{ij}^T\boldsymbol{\beta} + u_i)\}^2}{2\sigma^2}\right]\right]^{\delta_{ij}}$$
$$\times \left[1 - \Phi\left(\frac{\log(t) - (\boldsymbol{x}_{ij}^T\boldsymbol{\beta} + u_i)}{\sigma}\right)\right]^{1-\delta_{ij}}.$$

The details of the other functions are shown in the Appendix ?? in the Supplemental material. For the frailty model, the hazard function with frailty term  $v_i$  is

$$h(t_{ij}|v_i) = v_i \frac{\frac{1}{t\sqrt{2\pi\sigma^2}} \exp\left\{-\frac{(\log(t)-\mu)^2}{2\sigma^2}\right\}}{1 - \Phi\left(\frac{\log(t)-\mu}{\sigma}\right)}.$$
(4)

The posterior probability is

$$\pi_f(\boldsymbol{\beta}, \boldsymbol{v}, \phi, \sigma^2 | \boldsymbol{t}, \boldsymbol{X}) \propto L_f(\boldsymbol{t} | \boldsymbol{X}, \boldsymbol{\beta}, \boldsymbol{v}, \phi) g_f(\boldsymbol{v} | \phi) \pi(\boldsymbol{\beta} | \boldsymbol{c}) \pi(\phi | \xi) \pi(\sigma^2 | a, b),$$

where

$$\begin{split} L_{f}(\boldsymbol{t}|\boldsymbol{X},\boldsymbol{\beta},\boldsymbol{v},\phi,k) &= \prod_{i=1}^{M} \prod_{j=1}^{n_{i}} \{f(t_{ij}|v_{i})\}^{\delta_{ij}} \{S(t_{ij}|v_{i})\}^{1-\delta_{ij}} \\ &= \prod_{i=1}^{M} \prod_{j=1}^{n_{i}} \left[ v_{i} \frac{1}{t\sqrt{2\pi\sigma^{2}}} \exp\left\{-\frac{(\log(t)-\mu)^{2}}{2\sigma^{2}}\right\} \left(1 - \Phi\left(\frac{\log(t)-\mu}{\sigma}\right)\right)^{v_{i}-1} \right]^{\delta_{ij}} \\ &\times \left[ \left(1 - \Phi\left(\frac{\log(t)-\mu}{\sigma}\right)\right)^{v_{i}} \right]^{1-\delta_{ij}}. \end{split}$$

The details of the other functions are shown in the Appendix ?? in the Supplemental material.

If random or frailty effects are not included, a function with  $u_i$  set to 0 or  $v_i$  set to 1 can be used, and the posterior probabilities can be calculated by excluding the distribution of the effects.

#### 2.3 Distribution of RMST

The distribution of RMST is obtained using the posterior sample. We are interested in the RMST for each group and the difference between groups. For  $\beta$ , when the intercept is  $\beta_0$  and the coefficient parameter for the dose group is  $\beta_1$ , the coefficient parameters reflected in the RMST are  $\beta_0$  and  $\beta_1$ .

[Exponential model] The distribution of RMST for each group is obtained below

$$RMST_E(\tau, x_1, \beta_0^*, \beta_1^*) = \frac{1 - e^{-\exp(\beta_0^* + x_1\beta_1^*)\tau}}{\exp(\beta_0^* + x_1\beta_1^*)},$$
(5)

where \* denotes the posterior sample, and  $x_1$  is 0 if the group is the control and 1 if the group is the treatment. The difference between the RMSTs is

$$RMST_E(\tau, 1, \beta_0^*, \beta_1^*) - RMST_E(\tau, 0, \beta_0^*, \beta_1^*).$$

[Weibull model] The distribution of RMST for each group is obtained below

$$RMST_{W}(\tau, x_{1}, \beta_{0}^{*}, \beta_{1}^{*}) = \exp\left(-\frac{\beta_{0}^{*} + x_{1}\beta_{1}^{*}}{k^{*}}\right)\gamma\left(\exp(\beta_{0}^{*} + x_{1}\beta_{1}^{*})\tau^{k^{*}}; \frac{1}{k^{*}} + 1\right) + \tau\exp\left(-\exp(\beta_{0}^{*} + x_{1}\beta_{1}^{*})\tau^{k^{*}}\right).$$
(6)

The difference between the RMSTs is

$$RMST_W(\tau, 1, \beta_0^*, \beta_1^*) - RMST_W(\tau, 0, \beta_0^*, \beta_1^*).$$

[Log-logistic model] The distribution of RMST for each group is obtained below

$$RMST_{LL}(\tau, x_1, \beta_0^*, \beta_1^*) = e^{-\frac{\beta_0^* + x_1 \beta_1^*}{k^*}} B\left(\frac{e^{\beta_0^* + x_1 \beta_1^*} \tau^{k^*}}{1 + e^{\beta_0^* + x_1 \beta_1^*} \tau^{k^*}}; 1 + \frac{1}{k^*}, 1 - \frac{1}{k^*}\right) + \tau \frac{1}{1 + e^{\beta_0^* + x_1 \beta_1^*} \tau^{k^*}}.$$
(7)

The difference between the RMSTs is

$$RMST_{LL}(\tau, 1, \beta_0^*, \beta_1^*) - RMST_{LL}(\tau, 0, \beta_0^*, \beta_1^*).$$

[Log-normal model] The distribution of RMST for each group is obtained below

$$RMST_{LN}(\tau, x_1, \beta_0^*, \beta_1^*) = \exp\left\{\beta_0^* + x_1\beta_1^* + \frac{\sigma^{2*}}{2}\right\} \Phi\left(\frac{\log(\tau) - (\beta_0^* + x_1\beta_1^* - \sigma^{2*})}{\sigma^*}\right) + \tau\left(1 - \Phi\left(\frac{\log(\tau) - (\beta_0^* + x_1\beta_1^*)}{\sigma^*}\right)\right).$$
(8)

The difference between the RMSTs is

$$RMST_{LN}(\tau, 1, \beta_0^*, \beta_1^*) - RMST_{LN}(\tau, 0, \beta_0^*, \beta_1^*).$$

The RMST for each cluster, using shrinkage effects via random or frailty effects, is shown in the Appendices ?? and ?? in the Supplemental material, as it follows a similar RMST formula to the ones introduced so far.

#### 2.4 Theoretical aspect

We show the consistency of the RMST for any models when the estimated parameters are consistent. The consistency of the RMST implies that the RMST derived from the posterior distribution approaches the true value as the number of subjects increases. Furthermore, it is known that the posterior probability has consistency if the true parameter is in support of the prior and the consistent Bayes estimator exists with rich enough information set [6-8].

Let  $\hat{\boldsymbol{\beta}}_n = (\hat{\beta}_{n0}, \hat{\beta}_{n1})$  be the estimators by the posterior distribution with *n* subjects and  $\boldsymbol{\beta} = (\beta_0, \beta_1)$  be the coefficient parameters. If  $\hat{\boldsymbol{\beta}}_n \to_p \boldsymbol{\beta}$ , then  $\int_0^\tau S(t|\hat{\boldsymbol{\beta}}_n) dt \to_p \int_0^\tau S(t|\boldsymbol{\beta}) dt$ .

Here, we show the proof. We can calculate the below inequality.

$$\begin{aligned} \left| \int_0^\tau S(t|\hat{\boldsymbol{\beta}}_n) dt - \int_0^\tau S(t|\boldsymbol{\beta}) dt \right| &= \left| \int_0^\tau \left\{ S(t|\hat{\boldsymbol{\beta}}_n) - S(t|\boldsymbol{\beta}) \right\} dt \right| \\ &\leq \int_0^\tau \left| S(t|\hat{\boldsymbol{\beta}}_n) - S(t|\boldsymbol{\beta}) \right| dt \\ &= \int_0^\tau \left| F(t|\hat{\boldsymbol{\beta}}_n) - F(t|\boldsymbol{\beta}) \right| dt \end{aligned}$$

Because the maximum difference for distribution functions is 1, there exists a constant  $t_c \in [0, \tau]$  that satisfies

$$\int_0^\tau \left| F(t|\hat{\boldsymbol{\beta}}_n) - F(t|\boldsymbol{\beta}) \right| dt \le \tau \left| F(t_c|\hat{\boldsymbol{\beta}}_n) - F(t_c|\boldsymbol{\beta}) \right|$$

By the first degree Taylor approximating polynomial of  $F(t|\hat{\beta}_n)$  at  $\hat{\beta}_n = \beta$ , the inequality can be transformed as

$$\begin{aligned} \tau \left| F(t_c | \hat{\boldsymbol{\beta}}_n) - F(t_c | \boldsymbol{\beta}) \right| &= \tau \left| \left\{ \frac{\partial F(t_c | \boldsymbol{\beta}')}{\partial \beta_0} \right|_{\boldsymbol{\beta} = \boldsymbol{\beta}'} \right\} (\hat{\beta}_{n0} - \beta_0) + \left\{ \frac{\partial F(t_c | \boldsymbol{\beta}')}{\partial \beta_1} \right|_{\boldsymbol{\beta} = \boldsymbol{\beta}'} \right\} (\hat{\beta}_{n1} - \beta_1) \right| \\ &\leq \tau \left| \left\{ \frac{\partial F(t_c | \boldsymbol{\beta})}{\partial \beta_0} \right|_{\boldsymbol{\beta} = \boldsymbol{\beta}'} \right\} \left| \left| \hat{\beta}_{n0} - \beta_0 \right| + \tau \left| \left\{ \frac{\partial F(t_c | \boldsymbol{\beta})}{\partial \beta_1} \right|_{\boldsymbol{\beta} = \boldsymbol{\beta}'} \right\} \right| \left| \hat{\beta}_{n1} - \beta_1 \right| \\ &\to 0 \quad (n \to \infty), \end{aligned}$$

where  $\boldsymbol{\beta}'$  is a constant that satisfied  $|\beta'_i - \beta_i| < |\beta'_i - \hat{\beta}_i|$  (i = 0, 1). Thus,  $\left|\int_0^{\tau} S(t|\hat{\boldsymbol{\beta}}_n) dt - \int_0^{\tau} S(t|\boldsymbol{\beta}) dt\right| \rightarrow 0$   $(n \rightarrow \infty)$ , and then the RMST has consistent.

#### 2.5 Model selection

We utilize model selection methods to determine the distribution, covariates, and random and frailty effects that best describe the time-to-event data. Model selection is conducted using the posterior samples. Established information criteria such as widely applicable information criterion (WAIC) [9], widely applicable Bayesian information criterion (WBIC) [10], deviance information criterion (DIC) [11], and leave-one-out cross-validation (LOO) [12] are available. In particular, WAIC and LOO can be calculated using the **brms** package in R.

# 3 Simulation

#### **3.1** Simulation configuration

We investigated the performance of our proposed method through computer simulations based on two data scenarios. For comparison purposes, we computed the bias, mean squared error (MSE), the difference between mode and true RMST (Mode), and the difference between median and true RMST (Median). Specifically, our primary interest was the bias of the difference in RMST between groups across exponential, Weibull, log-logistic, and log-normal distribution. We assumed that a censoring probability is 0.1, and all survival times larger than 100 months were censored. We considered different sample sizes ranging from small to large (N = 64, 512, 2048). The Markov chain Monte Carlo (MCMC) specification included two chains, each with 2000 iterations and 1000 burn-ins. The number of simulations was set to 100 due to the extensive computation time involved in MCMC.

The simulation datasets for Scenario A were generated from a log-logistic model. The scale parameter was as follows:

$$\mu_{ij} = -(\beta_0 + \beta_1 x_{1ij} + \beta_2 x_{2ij} + u_i), \tag{9}$$

where  $\beta_0 = 5$ ,  $\beta_1 = -0.2$ ,  $\beta_2 = 1$ ,  $u_i$  represented the *i*-th cluster random effect drawn from a normal distribution with mean 0 and variance 0.1,  $x_{1ij}$  denoted the treatment arm (0 for the control group and 1 for the treatment group), and  $x_{2ij}$  was a continuous covariate distributed as standard normal. The number of clusters was four, and the shape parameter k was set to 2. The true model-based RMST value for each group in Scenario A was 82.69 and 87.99 by equation (7), respectively. Thus, the true difference between the two RMSTs was -5.30.

The simulation datasets of Scenario B were generated from a log-normal model. The parameters were given as like equation (9), where  $\beta_0 = 3$ ,  $\beta_1 = -0.5$  and  $\beta_2 = 1$ . The variance parameter  $\sigma^2$  was set to 1. The true model-based RMST values for each group in Scenario B were 19.14 and 29.51 by equation (8), respectively. Thus, the true difference between the RMSTs was -10.37.

A supplementary simulation, Scenario C, was also conducted. Refer to Appendix ?? in the Supplemental material for details on Scenario C. All survival curves were shown in Supplemental Figure ?? in the Supplemental material.

#### 3.2 Simulation results

The simulation results were in Table 1 for Scenarios A and B, and Supplemental Table ?? for scenario C in the Supplemental material. In Scenario A, where the mixed-effects log-logistic model was the true model, the bias and MSE of the log-logistic model were small regardless of the number of subjects. However, when the number of subjects was large, the Weibull and log-normal models had smaller bias and MSE than the log-logistic model. The exponential model, which is incorrect in Scenario A, exhibited some bias, but the MSE tended to be similar to the log-logistic model. Regarding the Mode and Median, the log-logistic model with frailty effects had the smallest values among the models.

In Scenario B, where the log-normal model with random effects is the true model, the lognormal model had a smaller bias and MSE than the Weibull and exponential models. The loglogistic model produced results similar to those in Scenario A, but the Weibull and exponential models had a systematic bias. Despite the log-normal being the true model, the log-logistic model showed a smaller Mode and Median.

		Ez	kponent	ial	Lo	g-Logis	tic	Lo	og-Norn	nal		Weibul	l
	n	Т	F	М	Т	F	М	Т	F	М	Т	$\mathbf{F}$	Μ
Scenario	A: Log	g-Logist	ic mode	l with r	andom	effect		1			1		
Bias	64	1.57	2.04	1.46	1.06	1.16	1.24	1.17	0.96	1.35	1.28	0.42	1.08
	512	0.94	1.28	0.85	0.30	-0.14	0.35	0.18	1.09	0.23	0.61	0.11	0.57
	2048	1.10	1.38	0.98	0.39	-0.25	0.44	0.39	2.83	0.40	0.77	0.22	0.64
MSE	64	30.81	24.66	31.18	31.44	36.05	31.13	32.13	38.69	32.44	34.42	49.65	35.02
	512	6.45	6.10	6.21	6.65	7.97	5.95	6.33	5.84	5.66	7.18	8.01	7.07
	2048	2.68	3.19	2.35	2.46	2.79	2.40	2.22	9.61	2.21	2.47	1.93	2.20
Mode	64	1.77	3.07	2.20	1.44	2.02	2.26	1.10	1.84	1.95	1.47	1.84	2.10
	512	0.93	1.89	1.31	0.27	0.04	0.88	0.15	1.68	0.61	0.69	0.84	1.03
	2048	1.14	1.62	1.30	0.35	-0.17	0.69	0.39	3.26	0.61	0.79	0.70	0.97
Median	64	1.64	2.37	1.73	1.14	1.41	1.54	1.21	1.30	1.57	1.32	0.85	1.44
	512	0.95	1.50	1.03	0.30	-0.10	0.54	0.18	1.31	0.38	0.61	0.38	0.78
	2048	1.11	1.49	1.12	0.39	-0.24	0.53	0.39	2.97	0.48	0.77	0.42	0.78
Scenario	B: Log	g-Normε	al model	l with ra	andom e	effect		1			1		
Bias	64	-1.38	-1.42	-0.91	-0.29	-0.61	-0.45	-0.49	0.02	-0.42	-1.28	-0.19	-0.73
	512	-1.48	-1.74	-1.30	-0.52	-0.37	-0.77	-0.88	-0.84	-0.86	-1.44	-0.87	-1.23
	2048	-1.19	-1.33	-0.91	-0.13	-0.13	-0.30	-0.52	-0.61	-0.47	-1.12	-0.53	-0.82
MSE	64	59.42	53.76	51.03	38.26	53.88	38.17	39.61	35.20	37.46	58.34	47.91	51.58
	512	10.56	11.41	9.63	6.27	6.71	6.48	6.46	6.47	6.33	10.50	8.70	9.95
	2048	4.75	4.40	3.69	2.33	2.78	2.31	2.61	3.10	2.49	4.66	3.24	3.64
Mode	64	-1.11	-0.38	0.00	0.11	0.16	0.53	-0.64	0.82	0.40	-1.27	0.98	-0.05
	512	-1.52	-1.82	-1.14	-0.57	-0.20	-0.41	-0.80	-0.86	-0.66	-1.36	-0.68	-1.14
	2048	-1.18	-1.54	-1.05	-0.13	-0.12	-0.13	-0.53	-0.58	-0.50	-1.12	-0.72	-0.97
Median	64	-1.28	-1.17	-0.68	-0.22	-0.45	-0.18	-0.48	0.26	-0.20	-1.18	0.23	-0.45
	512	-1.47	-1.76	-1.30	-0.52	-0.34	-0.65	-0.88	-0.82	-0.79	-1.42	-0.88	-1.21
	2048	-1.19	-1.46	-0.99	-0.13	-0.12	-0.23	-0.52	-0.62	-0.47	-1.11	-0.66	-0.89

Table 1: Results of simulation in each scenario

T: typical fixed model, F: frailty model, R: mixed effects model with random effects, Mean, Mode, and Median are the difference from true RMST.

# 4 Actual data re-analysis

We analyzed survival data of children in eight states of the Empowered Action Group (EAG) in India, collected from 2019 to 2021 through the Demographic and Health Surveys (DHS) program. Our primary interest was the difference in RMSTs of children under five among different maternal age groups at birth ("12-19 years old", "20-30 years old", "31+ years old"). The restricted time was set at 50 months, where the mortality rate exceeded 80%. Kaplan-Meier plots were presented in Supplemental Figures ?? and ?? in the Supplemental material.

The covariates included sex, place of delivery (Respondent's home, Other home, Public sector, Government hospital, CS Govt health professional, Other public sector, Private hos-

pital/clinic, CS private health facility, and Other), size of child at birth (Very large, Larger than average, Average, Smaller than average, Very small, and Don't know), and birth order. The clusters represented the eight states: Bihar, Jharkhand, Odisha, Chhattisgarh, Madhya Pradesh, Uttarakhand, Rajasthan, and Uttar Pradesh. The RMST differences between the two age groups across the eight states reveal that the trends varied by state (see Figures 1 and 3). We reanalyzed the data to determine if these differences were due to heterogeneity between the states or if they resulted from the sample size causing random variations in trends.

The MCMC specification included four chains, each with 2000 iterations and 1000 burn-ins. We used the brm functions from the **brms** package to analyze the exponential, Weibull, log-normal models, and mixed effects models. However, we developed custom Stan source code to analyze the exponential, Weibull, log-normal frailty models and all log-logistic models, as they could not be processed using the brm functions. These Stan files are included as supplemental material. Model selection was carried out using WAIC, with all WAIC results presented in Appendix ?? in the Supplemental material.

For the subgroup ('12-19 years old' vs. '20-30 years old' groups), the WAIC for the mixed effects Weibull model was the lowest. Hence, the following discussion was based on the results of the mixed effects model. Other estimation results were summarized in Appendix ?? in the Supplemental material. In the mixed-effects model, the RMST differences across states were similar due to the shrinkage effect, as the mean of the parameter  $\phi$ , representing the variance in the random effect distribution (indicating heterogeneity across clusters), was small, as shown in Figure 1. All estimation results were presented in Table 2, indicating that Place and Size affect the survival time of children under five. We obtained the RMST distribution, and its histogram is shown in Figure 2. From the RMST distribution, we calculated the probability of RMST under 0, -3, and -6 months. The probability of the RMST difference under 0 months was 0.507, while the probabilities under -3 and -6 months were both 0.000.

State	DiffMean [95%CI]	DiffMode	DiffMedian	Deaths_1219	Censors_1219	Deaths_2030	Censors_2030							
All states	-0.12 [-0.77, 0.51] -0.02 [-0.60, 0.53]	-0.13 -0.13	-0.12 -0.02	1920 (5.2%)	35113 (94.8%)	2889 (4.6%)	60491 (95.4%)			-	+			
Bihar	0.08 [-2.16, 2.88] -0.02 [-0.56, 0.48]	0.01 -0.07	0.01 -0.02	515 (5.3%)	9172 (94.7%)	463 (4.9%)	8929 (95.1%)			_	+			
Jharkhand	-0.77 [-2.97, 1.15] -0.02 [-0.56, 0.48]	-0.44 -0.16	-0.71 -0.02	165 (4.0%)	3912 (96.0%)	188 (4.2%)	4302 (95.8%)	_		•_	+	-		
Odisha	0.63 [-0.20, 1.79] -0.03 [-0.67, 0.58]	0.39 -0.08	0.57 -0.03	107 (4.6%)	2206 (95.4%)	210 (4.1%)	4874 (95.9%)			_	+			
Chhattisgarh	0.18 [-1.01, 1.41] -0.03 [-0.63, 0.55]	0.19 -0.08	0.17 -0.02	102 (4.8%)	2031 (95.2%)	250 (5.0%)	4707 (95.0%)			_	-	_		
Madhya Pradesh	-1.09 [-3.41, 0.26] -0.02 [-0.61, 0.52]	-0.02 -0.15	-0.87 -0.02	252 (5.0%)	4744 (95.0%)	362 (4.3%)	8033 (95.7%)			• -	+			
Uttarakhand	-0.63 [-3.02, 1.65] -0.02 [-0.60, 0.53]	-0.43 0.08	-0.58 -0.02	32 (4.8%)	638 (95.2%)	77 (3.2%)	2308 (96.8%)	_		-	+	—		
Rajasthan	-0.12 [-1.55, 0.00] -0.03 [-0.66, 0.57]	0.00 -0.08	0.00 -0.02	152 (3.7%)	3941 (96.3%)	267 (3.4%)	7650 (96.6%)		-	_	•			
Uttar Pradesh	-0.29 [-1.68, 0.72] -0.02 [-0.62, 0.54]	-0.01 -0.08	-0.17 -0.02	595 (6.6%)	8469 (93.4%)	1072 (5.2%)	19688 (94.8%)		-		+			
							-4	-3	-2	-1	0	1 2	3	4
							~	12-	19 gro	up bette	er 20-30	group be	ətter	$\rightarrow$
				No	random effect	Random effect								

Figure 1: Forest plot of the difference in restricted mean survival times (12–19 group vs 20–30 group)

	Parameter	Mode	Median	Mean	SE	95%CI	Rhat	ESS
	Intercept	3.91	3.99	4.00	0.14	[3.73, 4.28]	1.01	622
	20–30 group	0.00	0.00	0.00	0.02	[-0.04, 0.03]	1.00	2658
	Sex	0.07	0.08	0.08	0.02	[0.04, 0.11]	1.00	2274
Place	Respondents home	0.62	0.54	0.54	0.14	[0.26, 0.79]	1.01	551
Place	Other home	0.51	0.58	0.60	0.27	[0.10, 1.17]	1.00	1204
Place	Public sector	0.65	0.68	0.68	0.15	[0.36,  0.95]	1.00	646
Place	Government hospital	0.57	0.58	0.58	0.14	[0.30,  0.83]	1.00	551
Place	CS Govt health professional	0.63	0.57	0.57	0.13	[0.29, 0.81]	1.01	547
Place	Other public sector	0.58	0.63	0.62	0.15	[0.32, 0.89]	1.00	602
Place	Private hospital/clinic	0.45	0.41	0.40	0.14	[0.12,  0.65]	1.01	563
Place	CS private health facility	0.52	0.62	0.62	0.17	[0.29, 0.96]	1.00	681
Size	Very large	0.67	0.67	0.67	0.05	[0.58,  0.76]	1.01	974
Size	Larger than average	0.73	0.71	0.70	0.05	[0.61,  0.79]	1.01	926
Size	Average	0.78	0.77	0.76	0.04	[0.69, 0.84]	1.01	734
Size	Smaller than average	0.54	0.54	0.53	0.04	[0.45, 0.62]	1.00	963
Size	Very small	0.10	0.08	0.08	0.05	[-0.01, 0.17]	1.00	991
	Order	-0.01	-0.01	-0.01	0.01	[-0.02, 0.00]	1.00	2864
	k	1.71	1.71	1.71	0.02	[1.67, 1.75]	1.00	1979
	$\phi$	0.08	0.10	0.11	0.04	[0.05, 0.22]	1.00	483
	$RMST_{12-19group}$	37.99	37.35	37.33	2.27	[32.73, 41.66]		
	$\mathrm{RMST}_{20-30group}$	37.14	37.34	37.31	2.28	[32.78, 41.56]		
	RMST <sub>diff</sub>	-0.13	-0.02	-0.02	0.29	[-0.60, 0.53]		

Table 2: Results of mixed effects Weibull model (12–19 group vs 20–30 group)



Figure 2: Histogram of the difference in restricted mean survival times (12–19 group vs 20–30 group)

For the subgroup ('12-19 years old' vs. '31+ years old' groups), the WAIC for the Weibull frailty model was the lowest. Hence, the following discussion is based on the results of the frailty model. Other estimation results were summarized in Appendix ?? in the Supplemental material. In the frailty model, the RMST differences across states were similar due to the shrinkage effect, as the mean of the parameter  $\phi$ , representing the variance in the gamma frailty distribution (indicating heterogeneity across clusters), was small, as shown in Figure 3. All estimation results were presented in Table 3, which implied that Place and Size affect the survival time of children under five. We obtained the RMST distribution, and its histogram is shown in Figure 4. From the RMST distribution, we calculated the probability of RMST under 0, -3, and -6 months. The probability of the RMST difference under 0 months was 0.999, the probability under -3 months was 0.69375, and the probability under -6 months was 0.057.

State	DiffMean [95%CI]	DiffMode	DiffMedian	Deaths_1219	Censors_1219	Deaths_2030	Censors_2030						
All states	-3.96 [-6.74, -1.36] -3.77 [-6.55, -1.31]	-3.68 -3.44	-3.92 -3.71	1920 (5.2%)	35113 (94.8%)	2889 (4.6%)	60491 (95.4%)		=				
Bihar	-0.13 [-1.52, 0.19] -3.87 [-6.79, -1.34]	-0.01 -3.49	0.00 -3.82	515 (5.3%)	9172 (94.7%)	463 (4.9%)	8929 (95.1%)			•			
Jharkhand	1.89 [-8.08, 14.05] -4.08 [-7.07, -1.44]	-0.09 -3.61	1.54 -4.01	165 (4.0%)	3912 (96.0%)	188 (4.2%)	4302 (95.8%)	-		-			-
Odisha	-2.94 [-7.61, 0.18] -3.42 [-6.47, -1.00]	-2.52 -3.56	-2.66 -3.33	107 (4.6%)	2206 (95.4%)	210 (4.1%)	4874 (95.9%)	-	-				
Chhattisgarh	-5.30 [-11.50, 1.70] -3.65 [-6.51, -1.23]	-4.99 -3.29	-5.43 -3.58	102 (4.8%)	2031 (95.2%)	250 (5.0%)	4707 (95.0%)			+			
Madhya Pradesh	-0.16 [-2.12, 0.00] -3.94 [-6.84, -1.37]	0.01 -3.63	0.00 -3.88	252 (5.0%)	4744 (95.0%)	362 (4.3%)	8033 (95.7%)			•			
Uttarakhand	0.17 [0.00, 2.31] -3.85 [-6.76, -1.31]	-0.01 -3.40	0.00 -3.80	32 (4.8%)	638 (95.2%)	77 (3.2%)	2308 (96.8%)			-			
Rajasthan	-0.10 [-1.00, 0.00] -3.72 [-6.55, -1.24]	0.01 -3.52	0.00 -3.65	152 (3.7%)	3941 (96.3%)	267 (3.4%)	7650 (96.6%)			•			
Uttar Pradesh	-2.88 [-10.85, 1.91] -3.71 [-6.54, -1.25]	-0.27 -3.37	-1.92 -3.65	595 (6.6%)	8469 (93.4%)	1072 (5.2%)	19688 (94.8%)			+			
							-15 (	-10	-5 aroup bette	0 r 31+ c	5 aroup be	10 tter	
				•	lo frailty effect	railty effect		0	5				

Figure 3: Forest plot of the difference in restricted mean survival times (12–19 group vs 31+ group)

	Parameter	Mode	Median	Mean	SE	95%CI	Rhat	ESS
	Intercept	3.98	4.02	4.03	0.19	[3.68, 4.42]	1.00	978
	31+ group	-0.22	-0.21	-0.21	0.07	[-0.34, -0.08]	1.00	5178
	Sex	0.05	0.05	0.05	0.03	[0.00, 0.10]	1.00	4795
Place	Respondents home	0.55	0.49	0.48	0.18	[0.12, 0.82]	1.00	1008
Place	Other home	0.45	0.35	0.37	0.34	[-0.24, 1.10]	1.00	2087
Place	Public sector	0.80	0.77	0.77	0.20	[0.37, 1.17]	1.00	1173
Place	Government hospital	0.55	0.52	0.52	0.18	[0.16, 0.86]	1.00	1022
Place	CS Govt health professional	0.58	0.55	0.54	0.18	[0.19, 0.87]	1.00	999
Place	Other public sector	0.70	0.69	0.68	0.22	[0.25, 1.11]	1.00	1341
Place	Private hospital/clinic	0.37	0.33	0.32	0.18	[-0.04, 0.66]	1.00	1024
Place	CS private health facility	0.81	0.86	0.87	0.28	[0.34, 1.42]	1.00	1824
Size	Very large	0.45	0.44	0.44	0.07	[0.30,  0.57]	1.00	2108
Size	Larger than average	0.57	0.56	0.56	0.07	[0.43, 0.69]	1.00	2055
Size	Average	0.63	0.63	0.63	0.06	[0.52, 0.73]	1.00	1754
Size	Smaller than average	0.49	0.48	0.49	0.07	[0.36,  0.61]	1.00	2070
Size	Very small	0.03	0.02	0.02	0.07	[-0.11, 0.15]	1.00	2183
	Order	0.00	0.00	0.00	0.01	[-0.02, 0.01]	1.00	5814
	k	1.82	1.83	1.83	0.04	[1.76, 1.90]	1.00	3477
	$\phi$	0.02	0.04	0.06	0.06	[0.01, 0.22]	1.00	1634
	$RMST_{12-19group}$	38.70	38.54	38.38	3.03	[31.98, 43.70]		
	$RMST_{31+group}$	34.53	34.78	34.61	3.73	[27.03, 41.36]		
	$\mathrm{RMST}_{diff}$	-3.44	-3.71	-3.77	1.36	[-6.55, -1.31]		

Table 3: Results of Weibull frailty model (12–19 group vs 31+ group)



Figure 4: Histogram of the difference in restricted mean survival times (12–19 group vs 31+ group)

# 5 Discussion

In this paper, we proposed a Bayesian method for determining the distribution of RMST using posterior samples. We showed that the estimated RMST converged to the true value as the sufficient number of subjects when the estimated Bayesian parameters were consistent. We derived an explicit RMST equation by devising an integral of the survival function to obtain the distribution of RMST. Our proposed method allows to derive not only the mean and credible interval but also the mode, median, and probability of exceeding a certain value. In addition, we proposed two methods: one using random effects to account for heterogeneity among clusters, and the other using frailty. Because the frailty model cannot be analyzed with the existing brms package in R, we created a custom Stan file; see the supplemental material for the Stan files. Similarly, a log-logistic model could not be analyzed with brms, so a custom Stan code was created; see the supplemental material for the Stan files.

A limitation of our study is the necessity to assume a parametric model for survival time to obtain posterior samples. However, because there are several parametric models and many methods of model selection, selecting the model most appropriate for the data helps avoid incorrect analysis results. Additionally, a more flexible model can be built using the spline function proposed by Zhong and Schaubel (2022) [13]. The RMST for each period between knots can be obtained using the formulas we have used, and the overall RMST can be derived by summing the RMSTs.

The simulation study confirms that the bias decreases, and the MSE becomes smaller as the sample size increases, as theory suggests. In Scenario A, the log-normal frailty model did not perform well due to a poor fit to the model, but in Scenario B, it performed adequately. The simulation results showed a small bias and MSE for models close enough to the true model.

The importance of selecting an appropriate model by model selection for actual data analysis was suggested.

In the actual data analysis, we adjusted for differences between clusters by considering heterogeneity and obtained consistent results for each state. When using the exponential distribution from Scenario C, the estimation results were unstable, as with the log-normal frailty model in the real data analysis. The log-normal frailty model was found to increase frailty when the fit to the model was poor. A sensitivity analysis in Appendix ?? in the Supplemental material was performed to check the behavior of the log-normal frailty estimation and the behavior of the WAIC for log-logistic frailty, and no particular problems were found. The simulations for log-normal frailty showed that the estimation accuracy may be poor if the data were not generated from a log-normal base.

In conclusion, we derived explicit RMST formulas for the parametric model and the distribution of RMST. This allows us to calculate the mean, median, mode, and credible interval. We created the Stan code for the frailty and log-logistic models. Simulations confirmed that there were no performance problems, and we used the shrinkage effect to derive more accurate results for each cluster.

# References

- Hajime Uno, Brian Claggett, Lu Tian, Eisuke Inoue, Paul Gallo, Toshio Miyata, Deborah Schrag, Masahiro Takeuchi, Yoshiaki Uyama, Lihui Zhao, et al. Moving beyond the hazard ratio in quantifying the between-group difference in survival analysis. *Journal of clinical* Oncology, 32(22):2380, 2014.
- [2] Per Kragh Andersen, John P Klein, and Susanne Rosthoj. Generalised linear models for correlated pseudo-observations, with applications to multi-state models. *Biometrika*, 90(1):15–27, 2003.
- [3] Lu Tian, Lihui Zhao, and LJ Wei. Predicting the restricted mean event time with the subject's baseline covariates in survival analysis. *Biostatistics*, 15(2):222–233, 2014.
- [4] Keisuke Hanada and Masahiro Kojima. Random effect restricted mean survival time model. arXiv preprint arXiv:2401.02048, 2024.
- [5] Chenyang Zhang and Guosheng Yin. Bayesian nonparametric analysis of restricted mean survival time. *Biometrics*, 79(2):1383–1396, 2023.
- [6] Joseph L Doob. Application of the theory of martingales. Le calcul des probabilites et ses applications, pages 23–27, 1949.
- [7] Andrew Gelman and Cosma Rohilla Shalizi. Philosophy and the practice of bayesian statistics. *British Journal of Mathematical and Statistical Psychology*, 66(1):8–38, 2013.
- [8] Mark J Schervish. Theory of statistics. Springer Science & Business Media, 2012.
- [9] Sumio Watanabe and Manfred Opper. Asymptotic equivalence of bayes cross validation and widely applicable information criterion in singular learning theory. *Journal of machine learning research*, 11(12), 2010.
- [10] Sumio Watanabe. A widely applicable bayesian information criterion. The Journal of Machine Learning Research, 14(1):867–897, 2013.

- [11] David J Spiegelhalter, Nicola G Best, Bradley P Carlin, and Angelika Van Der Linde. Bayesian measures of model complexity and fit. Journal of the royal statistical society: Series b (statistical methodology), 64(4):583–639, 2002.
- [12] Aki Vehtari, Andrew Gelman, and Jonah Gabry. Practical bayesian model evaluation using leave-one-out cross-validation and waic. *Statistics and computing*, 27:1413–1432, 2017.
- [13] Yingchao Zhong and Douglas E Schaubel. Restricted mean survival time as a function of restriction time. *Biometrics*, 78(1):192–201, 2022.

# Supplemental material of Bayesian Parametric Methods for Deriving Distribution of Restricted Mean Survival Time

A Appendix

# A.1 Derivation of RMST

#### A.1.1 Exponential distribution

For the exponential distribution  $Exp(\lambda)$ , we define the density function, survival function, and hazard function.

$$f(t) = \lambda e^{-\lambda t}, S(t) = e^{-\lambda t}, h(t) = \lambda.$$

The RMST can be calculated as follows:

$$\int_0^{\tau} S(t)dt = \int_0^{\tau} tf(t)dt + \tau S(\tau)$$
$$= \left[-te^{-\lambda t}\right]_0^t + \int_0^{\tau} e^{-\lambda t}dt + \tau e^{-\lambda \tau}$$
$$= -\tau e^{-\lambda \tau} + \left[-\frac{e^{-\lambda t}}{\lambda}\right]_0^{\tau} + \tau e^{-\lambda \tau}$$
$$= \frac{1 - e^{-\lambda \tau}}{\lambda}.$$

In the exponential distribution, the survival function can be directly integrated as  $\int_0^{\tau} S(t)dt = \int_0^{\tau} e^{-\lambda t} dt = \frac{1-e^{-\lambda t}}{\lambda}$ .

#### A.1.2 Weibull distribution

For the weibull distribution  $W(\lambda, k)$  where the scale parameter  $\lambda > 0$  and the shape parameter k > 0, we define the density function, survival function, and hazard function.

$$f(t) = \lambda k t^{k-1} \exp(-\lambda t^k), S(t) = \exp(-\lambda t^k), h(t) = \lambda k t^{k-1}.$$

The RMST is

$$\begin{split} \int_0^\tau S(t)dt &= \int_0^\tau t\lambda kt^{k-1}\exp(-\lambda t^k)dt + \tau\exp(-\lambda \tau^k) \\ &(\lambda t^k = x \Leftrightarrow t = \lambda^{-\frac{1}{k}}x^{\frac{1}{k}}, \lambda kt^{k-1}dt = dx) \\ &= \int_0^{\lambda \tau^k} \lambda^{-\frac{1}{k}}x^{\frac{1}{k}}\exp(-x)dx + \tau\exp(-\lambda \tau^k) \\ &= \lambda^{-\frac{1}{k}}\gamma\left(\lambda \tau^k; \frac{1}{k} + 1\right) + \tau\exp(-\lambda \tau^k), \end{split}$$

where  $\gamma(z; a) = \int_0^z t^{a-1} e^{-t} dt$  is the incomplete gamma function.

#### A.1.3 Log-logistic distribution

For the log-logistic distribution  $LL(\mu, k)$  where the parameters  $\mu \in \mathbf{R}$ , k > 0, we define the density function, survival function, and hazard function.

$$f(t) = \frac{e^{\mu}kt^{k-1}}{(1+e^{\mu}t^k)^2}, S(t) = \frac{1}{1+e^{\mu}t^k}, h(t) = \frac{e^{\mu}kt^{k-1}}{1+e^{\mu}t^k}$$

The RMST is calculated as follows:

$$\int_{0}^{\tau} S(t)dt = \int_{0}^{\tau} t \frac{e^{\mu}kt^{k-1}}{(1+e^{\mu}t^{k})^{2}}dt + \tau \frac{1}{1+e^{\mu}\tau^{k}}$$

$$(e^{\mu}t^{k} = x \Leftrightarrow t = e^{-\frac{\mu}{k}}x^{\frac{1}{k}}, dt = \frac{1}{k}e^{-\frac{\mu}{k}}x^{\frac{1}{k}-1}dx)$$

$$= \int_{0}^{e^{\mu}\tau^{k}} \frac{e^{-\frac{\mu}{k}}x^{\frac{1}{k}}}{(1+x)^{2}}dx + \tau \frac{1}{1+e^{\mu}\tau^{k}}$$

$$= e^{-\frac{\mu}{k}}B\left(\frac{e^{\mu}\tau^{k}}{1+e^{\mu}\tau^{k}}; 1+\frac{1}{k}, 1-\frac{1}{k}\right) + \tau \frac{1}{1+e^{\mu}\tau^{k}}, \qquad (1)$$

where  $B(z; a, b) = \int_0^z t^{a-1}(1-t)^{b-1}dt$  is the incomplete beta function. The following formula was used to derive the incomplete beta function in the equation 1.

$$\begin{split} &\int_0^z \frac{x^{a-1}}{(1+x)^{a+b}} dx \\ &\left(\frac{x}{1+x} = y \Leftrightarrow x = \frac{y}{1-y}, dx = \frac{1}{(1-y)^2} dy\right) \\ &= \int_0^{\frac{z}{1+z}} y^{a-1} (1-y)^{b-1} \\ &= B\left(\frac{z}{1+z}; a, b\right). \end{split}$$

#### A.1.4 Log-normal distribution

For the log-normal distribution,  $LN(\mu, \sigma^2)$  where the parameters  $\mu \in \mathbf{R}$ ,  $\sigma^2 > 0$ , we define the density function, survival function, and hazard function.

$$\begin{split} f(t) &= \frac{1}{t\sqrt{2\pi\sigma^2}} \exp\left\{-\frac{(\log(t)-\mu)^2}{2\sigma^2}\right\}, S(t) = 1 - \Phi\left(\frac{\log(t)-\mu}{\sigma}\right),\\ h(t) &= \frac{\frac{1}{t\sqrt{2\pi\sigma^2}} \exp\left\{-\frac{(\log(t)-\mu)^2}{2\sigma^2}\right\}}{1 - \Phi\left(\frac{\log(t)-\mu}{\sigma}\right)}. \end{split}$$

The RMST is

$$\begin{split} \int_0^\tau S(t)dt &= \int_0^\tau t \frac{1}{t\sqrt{2\pi\sigma^2}} \exp\left\{-\frac{(\log(t)-\mu)^2}{2\sigma^2}\right\} dt + \tau \left(1 - \Phi\left(\frac{\log(\tau)-\mu}{\sigma}\right)\right) \\ &(\log(t) = x \sim \mathcal{N}(\mu, \sigma^2), dt = e^x dx) \\ &= \int_{-\infty}^{\log(\tau)} \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left\{-\frac{(x-\mu)^2}{2\sigma^2}\right\} \exp\{x\} dt + \tau \left(1 - \Phi\left(\frac{\log(\tau)-\mu}{\sigma}\right)\right) \\ &= \exp\left\{\mu + \frac{\sigma^2}{2}\right\} \int_{-\infty}^{\log(\tau)} \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left\{-\frac{(x-\mu-\sigma^2)^2}{2\sigma^2}\right\} dt + \tau \left(1 - \Phi\left(\frac{\log(\tau)-\mu}{\sigma}\right)\right) \\ &\left(\frac{x-\mu-\sigma^2}{\sigma} = y \sim \mathcal{N}(-\sigma, 1), dx = \sigma dy\right) \\ &= \exp\left\{\mu + \frac{\sigma^2}{2}\right\} \int_{-\infty}^{\frac{\log(\tau)-\mu-\sigma^2}{\sigma}} \frac{1}{\sqrt{2\pi}} \exp\left\{-\frac{y^2}{2}\right\} dt + \tau \left(1 - \Phi\left(\frac{\log(\tau)-\mu}{\sigma}\right)\right) \\ &= \exp\left\{\mu + \frac{\sigma^2}{2}\right\} \Phi\left(\frac{\log(\tau)-\mu-\sigma^2}{\sigma}\right) + \tau \left(1 - \Phi\left(\frac{\log(\tau)-\mu}{\sigma}\right)\right). \end{split}$$

# A.2 Distribution of RMST for each cluster in mixed effect models

We demonstrate the computation of the RMST for each cluster, considering the heterogeneity via random effects.

[Mixed effects exponential model] The distribution of the RMST for the *i*-th cluster in each group is obtained below.

$$RMST_{E,r}(\tau, x_1, \beta_0^*, \beta_1^*, u_i^*) = \frac{1 - e^{-\exp\{\beta_0^* + x_1\beta_1^* + u_i^*\}\tau}}{\exp\{\beta_0^* + x_1\beta_1^* + u_i^*\}},$$

where the \* denotes the posterior sample, and  $x_1$  is 0 for the control group and 1 for the treatment group. The difference between the RMSTs is

$$RMST_{E,r}(\tau, 1, \beta_0^*, \beta_1^*, u_i^*) - RMST_{E,r}(\tau, 0, \beta_0^*, \beta_1^*, u_i^*).$$

If the model includes covariates,  $\beta_0^*$  and  $\beta_1^*$  are adjusted for the covariates.

[Mixed effects Weibull model] The distribution of the RMST for the *i*-th cluster in each group is obtained below.

$$RMST_{W,r}(\tau, x_1, \beta_0^*, \beta_1^*, u_i^*) = \exp\left\{-\frac{\beta_0^* + x_1\beta_1^* + u_i^*}{k^*}\right\} \gamma\left(\exp\{\beta_0^* + x_1\beta_1^* + u_i^*\}\tau^{k^*}; \frac{1}{k^*} + 1\right) + \tau \exp\left(-\exp\{\beta_0^* + x_1\beta_1^* + u_i^*\}\tau^{k^*}\right).$$

The difference between the RMSTs is

$$RMST_{W,r}(\tau, 1, \beta_0^*, \beta_1^*, u_i^*) - RMST_{W,r}(\tau, 0, \beta_0^*, \beta_1^*, u_i^*).$$

[Mixed effects log-logistic distribution] The distribution of the RMST for the *i*-th cluster in each group is obtained below.

$$RMST_{LL,r}(\tau, x_1, \beta_0^*, \beta_1^*, u_i^*) = e^{-\frac{\beta_0^* + x_1 \beta_1^* + u_i^*}{k^*}} B\left(\frac{e^{\beta_0^* + x_1 \beta_1^* + u_i^*} \tau^{k^*}}{1 + e^{\beta_0^* + x_1 \beta_1^* + u_i^*} \tau^{k^*}}; 1 + \frac{1}{k^*}, 1 - \frac{1}{k^*}\right) + \tau \frac{1}{1 + e^{\beta_0^* + x_1 \beta_1^* + u_i^*} \tau^{k^*}}.$$

The difference between the RMSTs is

$$RMST_{LL,r}(\tau, 1, \beta_0^*, \beta_1^*, u_i^*) - RMST_{LL,r}(\tau, 0, \beta_0^*, \beta_1^*, u_i^*).$$

[Mixed effects log-normal distribution] The distribution of the RMST for the *i*-th cluster in each group is obtained below.

$$RMST_{LN,r}(\tau, x_1, \beta_0^*, \beta_1^*, u_i^*) = \exp\left\{\beta_0^* + x_1\beta_1^* + u_i^* + \frac{\sigma^{2*}}{2}\right\} \Phi\left(\frac{\log(\tau) - (\beta_0^* + x_1\beta_1^* + u_i^* - \sigma^{2*})}{\sigma^*}\right) + \tau\left(1 - \Phi\left(\frac{\log(\tau) - (\beta_0^* + x_1\beta_1^* + u_i^*)}{\sigma^*}\right)\right).$$

The difference between the RMSTs is

$$RMST_{LN,r}(\tau, 1, \beta_0^*, \beta_1^*, u_i^*) - RMST_{LN,r}(\tau, 0, \beta_0^*, \beta_1^*, u_i^*).$$

#### A.2.1 Derivations of RMST for mixed effects models

We consider adding random effects to the rate parameters or the scale parameters.

[Mixed effects exponential model] The density and survival functions are conditioned on a random effect  $u \sim \mathcal{N}(0, \phi)$ , denoted as

$$f(t|u) = \lambda e^{u} e^{-\lambda e^{u}t}, S(t) = e^{-\lambda e^{u}t}.$$

The RMST for the mixed effects exponential model is

$$\int_0^\tau S(t|u)dt = \frac{1 - e^{-\lambda e^u \tau}}{\lambda e^u}.$$

The mixed effects exponential model is equivalent to the exponential model with the log-normal frailty.

[Mixed effects Weibull model] The density and survival functions are conditioned on a random effect  $u \sim \mathcal{N}(0, \phi)$ , denoted as

$$f(t|u) = \lambda e^{u} k t^{k-1} \exp(-\lambda e^{u} t^{k}), S(t) = \exp(-\lambda e^{u} t^{k}).$$

The RMST is

$$\int_0^\tau S(t|u)dt = \lambda^{-\frac{1}{k}} e^{-\frac{u}{k}} \gamma\left(\lambda e^u \tau^k; \frac{1}{k} + 1\right) + \tau \exp(-\lambda e^u \tau^k),$$

where  $\gamma(z; a) = \int_0^z t^{a-1} e^{-t} dt$  is an incomplete gamma function. The mixed effects Weibull model is equivalent to the Weibull model with the log-normal frailty.

[Mixed effects log-logistic model] The density and survival functions are conditioned on a random effect  $u \sim \mathcal{N}(0, \phi)$ , denoted as

$$f(t|u) = \frac{e^{\mu+u}kt^{k-1}}{(1+e^{\mu+u}t^k)^2}, S(t|u) = \frac{1}{1+e^{\mu+u}t^k}.$$

The RMST is

$$\int_0^{\tau} S(t|u)dt = e^{-\frac{\mu+u}{k}} B\left(\frac{e^{\mu+u}\tau^k}{1+e^{\mu+u}\tau^k}; 1+\frac{1}{k}, 1-\frac{1}{k}\right) + \tau \frac{1}{1+e^{\mu+u}\tau^k},$$

where B(z; a, b) is an incomplete beta function.

[Mixed effects log-normal model] The density and survival functions are conditioned on a random effect  $u \sim \mathcal{N}(0, \phi)$ , denoted as

$$f(t|u) = \frac{1}{t\sqrt{2\pi\sigma^2}} \exp\left\{-\frac{(\log(t) - \mu - u)^2}{2\sigma^2}\right\}, S(t|u) = 1 - \Phi\left(\frac{\log(t) - \mu - u}{\sigma}\right).$$

The RMST is

$$\int_0^{\tau} S(t|u)dt = \exp\left\{\mu + \frac{\sigma^2}{2}\right\} \Phi\left(\frac{\log(\tau) - \mu - u - \sigma^2}{\sigma}\right) + \tau\left(1 - \Phi\left(\frac{\log(\tau) - \mu - u}{\sigma}\right)\right).$$

# A.3 Distribution of RMST for frailty models

The derivations of the RMSTs for the frailty models are shown in the Appendix A.3.1.

[Exponential frailty model] The distribution of the RMST with frailty term for the *i*-th cluster in each group is obtained below.

$$RMST_{E,f}(\tau, x_1, \beta_0^*, \beta_1^*, v_i^*) = \frac{1 - e^{-v_i^* \exp\{\beta_0^* + x_1\beta_1^*\}\tau}}{v_i^* \exp\{\beta_0^* + x_1\beta_1^*\}}.$$

The difference between the RMSTs is

$$RMST_{E,f}(\tau, 1, \beta_0^*, \beta_1^*, v_i^*) - RMST_{E,f}(\tau, 0, \beta_0^*, \beta_1^*, v_i^*).$$

[Weibull frailty model] The distribution of the RMST with frailty term for the *i*-th cluster in each group is obtained below.

$$RMST_{W,f}(\tau, x_1, \beta_0^*, \beta_1^*, v_i^*) = (v_i^*)^{-\frac{1}{k}} \exp\left\{-\frac{\beta_0^* + x_1\beta_1^*}{k^*}\right\} \gamma\left(v_i^* \exp\{\beta_0^* + x_1\beta_1^*\}\tau^{k^*}; \frac{1}{k^*} + 1\right) + \tau \exp\left(-v_i^* \exp\{\beta_0^* + x_1\beta_1^*\}\tau^{k^*}\right).$$

The difference between the RMSTs is

$$RMST_{W,f}(\tau, 1, \beta_0^*, \beta_1^*, v_i^*) - RMST_{W,f}(\tau, 0, \beta_0^*, \beta_1^*, v_i^*).$$

[Log-logistic frailty model] The distribution of the RMST with frailty term for the *i*-th cluster in each group is obtained below.

$$RMST_{LL,f}(\tau, x_1, \beta_0^*, \beta_1^*, v_i^*) = v_i^* e^{-\frac{\beta_0^* + x_1 \beta_1^*}{k^*}} B\left(\frac{e^{\beta_0^* + x_1 \beta_1^*} \tau^{k^*}}{1 + e^{\beta_0^* + x_1 \beta_1^*} \tau^{k^*}}; 1 + \frac{1}{k^*}, v_i^* - \frac{1}{k^*}\right) + \tau \left(\frac{1}{1 + e^{\beta_0^* + x_1 \beta_1^*} \tau^{k^*}}\right)^{v_i^*}.$$

The difference between the RMSTs is

$$RMST_{LL,f}(\tau, 1, \beta_0^*, \beta_1^*, v_i^*) - RMST_{LL,f}(\tau, 0, \beta_0^*, \beta_1^*, v_i^*).$$

[Log-normal frailty model] The distribution of the RMST with frailty term for the *i*-th cluster in each group is obtained below.

$$RMST_{LN,f}(\tau, x_1, \beta_0^*, \beta_1^*, v_i^*) = \exp\left\{\beta_0^* + x_1\beta_1^* + \frac{\sigma^{2*}}{2}\right\} \frac{1}{v_i^*} \times \left(1 - \left(1 - \Phi\left(\frac{\log(\tau) - (\beta_0^* + x_1\beta_1^* - \sigma^{2*})}{\sigma^*}\right)\right)^{v_i^*}\right) + \tau \left(1 - \Phi\left(\frac{\log(\tau) - (\beta_0^* + x_1\beta_1^*)}{\sigma^*}\right)\right)^{v_i^*}.$$

The difference between the RMSTs is

$$RMST_{LN,f}(\tau, 1, \beta_0^*, \beta_1^*, v_i^*) - RMST_{LN,f}(\tau, 0, \beta_0^*, \beta_1^*, v_i^*).$$

#### A.3.1 Derivations of RMST for frailty models

We derive the RMST using frailty models before incorporating covariates. [Exponential frailty model] The RMST with frailty term is

$$\int_0^{\tau} S_f(t|v) dt = \int_0^{\tau} t f_f(t|v) dt + \tau S_f(\tau|v)$$
$$= \left[-te^{-v\lambda t}\right]_0^t + \int_0^{\tau} e^{-v\lambda t} dt + \tau e^{-v\lambda \tau}$$
$$= -\tau e^{-v\lambda \tau} + \left[-\frac{e^{-v\lambda t}}{v\lambda}\right]_0^{\tau} + \tau e^{-v\lambda \tau}$$
$$= \frac{1 - e^{-v\lambda \tau}}{v\lambda}.$$

[Weibull frailty model] The RMST with frailty term is

$$\int_0^\tau S_f(t|v)dt = \int_0^\tau tv\lambda kt^{k-1}\exp(-v\lambda t^k)dt + \tau\exp(-v\lambda\tau^k)$$
$$(v\lambda t^k = x \Leftrightarrow t = v^{-\frac{1}{k}}\lambda^{-\frac{1}{k}}x^{\frac{1}{k}}, v\lambda kt^{k-1}dt = dx)$$
$$= \int_0^{v\lambda\tau^k} v^{-\frac{1}{k}}\lambda^{-\frac{1}{k}}x^{\frac{1}{k}}\exp(-x)dx + \tau\exp(-v\lambda\tau^k)$$
$$= v^{-\frac{1}{k}}\lambda^{-\frac{1}{k}}\gamma\left(v\lambda\tau^k; \frac{1}{k} + 1\right) + \tau\exp(-v\lambda\tau^k).$$

[Log-logistic frailty model] The RMST with frailty term is

$$\begin{split} \int_0^\tau S_f(t|v)dt &= \int_0^\tau tv \frac{e^\mu k t^{k-1}}{(1+e^\mu t^k)^{v+1}} dt + \tau \left(\frac{1}{1+e^\mu \tau^k}\right)^v \\ &(e^\mu t^k = x \Leftrightarrow t = e^{-\frac{\mu}{k}} x^{\frac{1}{k}}, dt = \frac{1}{k} e^{-\frac{\mu}{k}} x^{\frac{1}{k}-1} dx) \\ &= \int_0^{e^\mu \tau^k} v \frac{e^{-\frac{\mu}{k}} x^{\frac{1}{k}}}{(1+x)^{v+1}} dx + \tau \left(\frac{1}{1+e^\mu \tau^k}\right)^v \\ &= v e^{-\frac{\mu}{k}} B\left(\frac{e^\mu \tau^k}{1+e^\mu \tau^k}; 1+\frac{1}{k}, v-\frac{1}{k}\right) + \tau \left(\frac{1}{1+e^\mu \tau^k}\right)^v. \end{split}$$

[Log-normal frailty model] The RMST with frailty term is

$$\begin{split} \int_{0}^{\tau} S_{f}(t|v) dt &= \int_{0}^{\tau} tv \frac{1}{t\sqrt{2\pi\sigma^{2}}} \exp\left\{-\frac{(\log(t)-\mu)^{2}}{2\sigma^{2}}\right\} \left(1 - \Phi\left(\frac{\log(t)-\mu}{\sigma}\right)\right)^{v-1} dt \\ &+ \tau \left(1 - \Phi\left(\frac{\log(\tau)-\mu}{\sigma}\right)\right)^{v} \\ (\log(t) &= x \sim \mathcal{N}(\mu, \sigma^{2}), dt = e^{x} dx) \\ &= \int_{-\infty}^{\log(\tau)} \frac{1}{\sqrt{2\pi\sigma^{2}}} \exp\left\{-\frac{(x-\mu)^{2}}{2\sigma^{2}}\right\} \exp\{x\} \left(1 - \Phi\left(\frac{x-\mu}{\sigma}\right)\right)^{v-1} dx \\ &+ \tau \left(1 - \Phi\left(\frac{\log(\tau)-\mu}{\sigma}\right)\right)^{v} \\ &= \exp\left\{\mu + \frac{\sigma^{2}}{2}\right\} \int_{-\infty}^{\log(\tau)} \frac{1}{\sqrt{2\pi\sigma^{2}}} \exp\left\{-\frac{(x-\mu-\sigma^{2})^{2}}{2\sigma^{2}}\right\} \left(1 - \Phi\left(\frac{x-\mu}{\sigma}\right)\right)^{v-1} dx \\ &+ \tau \left(1 - \Phi\left(\frac{\log(\tau)-\mu}{\sigma}\right)\right)^{v} \\ \left(\frac{x-\mu-\sigma^{2}}{\sigma} &= y \sim \mathcal{N}(-\sigma, 1), dx = \sigma dy\right) \\ &= \exp\left\{\mu + \frac{\sigma^{2}}{2}\right\} \int_{-\infty}^{\frac{\log(\tau)-\mu-\sigma^{2}}{\sigma}} \frac{1}{\sqrt{2\pi}} \exp\left\{-\frac{y^{2}}{2}\right\} (1 - \Phi\left(y + \sigma\right))^{v-1} dy \\ &+ \tau \left(1 - \Phi\left(\frac{\log(\tau)-\mu}{\sigma}\right)\right)^{v} \\ &\approx \exp\left\{\mu + \frac{\sigma^{2}}{2}\right\} \int_{-\infty}^{\frac{\log(\tau)-\mu-\sigma^{2}}{\sigma}} \frac{1}{\sqrt{2\pi}} \exp\left\{-\frac{y^{2}}{2}\right\} (1 - \Phi\left(y\right))^{v-1} dy \\ &+ \tau \left(1 - \Phi\left(\frac{\log(\tau)-\mu}{\sigma}\right)\right)^{v} \end{aligned}$$
(2)   
 
$$&= \exp\left\{\mu + \frac{\sigma^{2}}{2}\right\} \frac{1}{v} \left(1 - \left(1 - \Phi\left(\frac{\log(\tau)-\mu-\sigma^{2}}{\sigma}\right)\right)^{v}\right) \\ &+ \tau \left(1 - \Phi\left(\frac{\log(\tau)-\mu}{\sigma}\right)\right)^{v}. \end{split}$$

The second-to-last approximate equation 2 is the result of modifying the integral because it is not solvable. To obtain a better approximation, the Monte Carlo integral can be applied, but this may be computationally time-consuming since it is applied to each posterior sample.

#### Another representation of the Weibull distribution **A.4**

We demonstrate the RMST in the weibull distribution  $W(\lambda, k)$ , where a scale parameter  $\lambda > 0$  and the shape parameter k > 0. We define the density function, survival function, and hazard function.

$$f(t) = \frac{k}{\lambda} \left(\frac{t}{\lambda}\right)^{k-1} \exp\left\{-\left(\frac{t}{\lambda}\right)^k\right\}, S(t) = \exp\left\{-\left(\frac{t}{\lambda}\right)^k\right\}, h(t) = \frac{k}{\lambda} \left(\frac{t}{\lambda}\right)^{k-1}$$

The RMST is calculated as follows:

$$\int_{0}^{\tau} S(t)dt = \int_{0}^{\tau} t \frac{k}{\lambda} \left(\frac{t}{\lambda}\right)^{k-1} \exp\left\{-\left(\frac{t}{\lambda}\right)^{k}\right\} dt + \tau \exp\left\{-\left(\frac{\tau}{\lambda}\right)^{k}\right\}$$
$$\left(\left(\frac{t}{\lambda}\right)^{k} = x \Leftrightarrow t = \lambda x^{\frac{1}{k}}, \frac{k}{\lambda} \left(\frac{t}{\lambda}\right)^{k-1} dt = dx\right)$$
$$= \int_{0}^{\left(\frac{\tau}{\lambda}\right)^{k}} \lambda x^{\frac{1}{k}} \exp(-x) dx + \tau \exp\left\{-\left(\frac{\tau}{\lambda}\right)^{k}\right\}$$
$$= \lambda \gamma \left(\left(\frac{\tau}{\lambda}\right)^{k}; \frac{1}{k} + 1\right) + \tau \exp\left\{-\left(\frac{\tau}{\lambda}\right)^{k}\right\},$$

where  $\gamma(z; a) = \int_0^z t^{a-1} e^{-t} dt$  is an incomplete gamma function. Next we consider the Weibull model. We transform the parameter  $\lambda$  to account for covariates.

$$\lambda_{ij} = \exp\{\boldsymbol{x}_{ij}^T \boldsymbol{\beta}\}.$$

The hazard function with frailty term is

$$h(t_{ij}|v_i) = v_i \frac{k}{\exp(\boldsymbol{x}_{ij}^T \boldsymbol{\beta})} \left(\frac{t}{\exp(\boldsymbol{x}_{ij}^T \boldsymbol{\beta})}\right)^{k-1}.$$

For the Weibull distribution, the distribution of the RMST for each group is obtained below.

$$RMST_{W}(\tau, x_{1}, \beta_{0}^{*}, \beta_{1}^{*}) = \exp\left\{\beta_{0}^{*} + x_{1}\beta_{1}^{*}\right\} \gamma\left(\left(\frac{\tau}{\exp\left\{\beta_{0}^{*} + x_{1}\beta_{1}^{*}\right\}}\right)^{k^{*}}; \frac{1}{k^{*}} + 1\right) + \tau \exp\left\{-\left(\frac{\tau}{\exp\left\{\beta_{0}^{*} + x_{1}\beta_{1}^{*}\right\}}\right)^{k^{*}}\right\}.$$

The difference between the RMSTs is

$$RMST_W(\tau, 1, \beta_0^*, \beta_1^*) - RMST_W(\tau, 0, \beta_0^*, \beta_1^*).$$

The RMSTs for the mixed effects Weibull model are

$$RMST_{W,r}(\tau, x_1, \beta_0^*, \beta_1^*, u_i^*) = \exp\left\{\beta_0^* + x_1\beta_1^* + u_i^*\right\} \gamma\left(\left(\frac{\tau}{\exp\left\{\beta_0^* + x_1\beta_1^* + u_i^*\right\}}\right)^{k^*}; \frac{1}{k^*} + 1\right) + \tau \exp\left\{-\left(\frac{\tau}{\exp\left\{\beta_0^* + x_1\beta_1^* + u_i^*\right\}}\right)^{k^*}\right\}.$$

For the frailty model, the hazard function is

$$h_f(t|v) = v \frac{k}{\lambda} \left(\frac{t}{\lambda}\right)^{k-1}.$$

The density function and survival function are

$$f_f(t|v) = v\frac{k}{\lambda} \left(\frac{t}{\lambda}\right)^{k-1} \exp\left\{-v\left(\frac{t}{\lambda}\right)^k\right\}, S(t|v) = \exp\left\{-v\left(\frac{t}{\lambda}\right)^k\right\}$$

The RMST is calculated as follows

$$\int_{0}^{\tau} S(t|v)dt = \int_{0}^{\tau} vt \frac{k}{\lambda} \left(\frac{t}{\lambda}\right)^{k-1} \exp\left\{-v\left(\frac{t}{\lambda}\right)^{k}\right\} dt + \tau \exp\left\{-v\left(\frac{\tau}{\lambda}\right)^{k}\right\}$$
$$\left(v\left(\frac{t}{\lambda}\right)^{k} = x \Leftrightarrow t = v^{\frac{1}{k}}\lambda x^{\frac{1}{k}}, v\frac{k}{\lambda} \left(\frac{t}{\lambda}\right)^{k-1} dt = dx\right)$$
$$= \int_{0}^{v\left(\frac{\tau}{\lambda}\right)^{k}} v^{\frac{1}{k}}\lambda x^{\frac{1}{k}} \exp(-x)dx + \tau \exp\left\{-v\left(\frac{\tau}{\lambda}\right)^{k}\right\}$$
$$= v^{\frac{1}{k}}\lambda \gamma \left(v\left(\frac{\tau}{\lambda}\right)^{k}; \frac{1}{k} + 1\right) + \tau \exp\left\{-v\left(\frac{\tau}{\lambda}\right)^{k}\right\}.$$

The RMST for the Weibull frailty model is

$$RMST_{W,f}(\tau, x_1, \beta_0^*, \beta_1^*, v_i^*) = (v_i^*)^{\frac{1}{k^*}} \exp\left\{\beta_0^* + x_1\beta_1^*\right\} \gamma\left(v_i^*\left(\frac{\tau}{\exp\left\{\beta_0^* + x_1\beta_1^*\right\}}\right)^{k^*}; \frac{1}{k^*} + 1\right) + \tau \exp\left\{-v_i^*\left(\frac{\tau}{\exp\left\{\beta_0^* + x_1\beta_1^*\right\}}\right)^{k^*}\right\}.$$

### A.5 Another representation of the log-logistic distribution

We demonstrate the RMST in a log-logistic distribution  $LL(\alpha, k)$ , where parameters  $\alpha > 0$  and k > 0. We define the density function, survival function, and hazard function.

$$f(t) = \frac{\frac{k}{\alpha} \left(\frac{t}{\alpha}\right)^{k-1}}{\left(1 + \left(\frac{t}{\alpha}\right)^k\right)^2}, S(t) = \frac{1}{1 + \left(\frac{t}{\alpha}\right)^k}, h(t) = \frac{\frac{k}{\alpha} \left(\frac{t}{\alpha}\right)^{k-1}}{1 + \left(\frac{t}{\alpha}\right)^k}.$$

The RMST is calculated as follows:

$$\int_0^\tau S(t)dt = \int_0^\tau t \frac{\frac{k}{\alpha} \left(\frac{t}{\alpha}\right)^{k-1}}{\left(1 + \left(\frac{t}{\alpha}\right)^k\right)^2} dt + \tau \frac{1}{1 + \left(\frac{\tau}{\alpha}\right)^k}$$
$$\left(\left(\frac{t}{\alpha}\right)^k = x \Leftrightarrow t = \alpha x^{\frac{1}{k}}, \frac{k}{\alpha} \left(\frac{t}{\alpha}\right)^{k-1} dt = dx\right)$$
$$= \int_0^{\left(\frac{\tau}{\alpha}\right)^k} \frac{\alpha x^{\frac{1}{k}}}{(1+x)^2} dx + \tau \frac{1}{1 + \left(\frac{\tau}{\alpha}\right)^k}$$
$$= \alpha B\left(\frac{\left(\frac{\tau}{\alpha}\right)^k}{1 + \left(\frac{\tau}{\alpha}\right)^k}; 1 + \frac{1}{k}, 1 - \frac{1}{k}\right) + \tau \frac{1}{1 + \left(\frac{\tau}{\alpha}\right)^k},$$

where B(z; a, b) is an incomplete beta function. We transform the parameter  $\alpha$  to account for covariates.

$$\alpha_{ij} = \exp(\boldsymbol{x}_{ij}^T \boldsymbol{\beta}).$$

The hazard function with frailty term is

$$h(t_{ij}|v_i) = v_i \frac{\frac{k}{\exp(\boldsymbol{x}_{ij}^T \boldsymbol{\beta})} \left(\frac{t}{\exp(\boldsymbol{x}_{ij}^T \boldsymbol{\beta})}\right)^{k-1}}{1 + \left(\frac{t}{\exp(\boldsymbol{x}_{ij}^T \boldsymbol{\beta})}\right)^k},$$

$$S(t_{ij}|v_i) = \left(\frac{1}{1 + \left(\frac{t}{\exp(\boldsymbol{x}_{ij}^T \boldsymbol{\beta})}\right)^k}\right)^{v_i},$$

$$f(t_{ij}|v_i) = v_i \frac{k}{\exp(\boldsymbol{x}_{ij}^T \boldsymbol{\beta})} \left(\frac{t}{\exp(\boldsymbol{x}_{ij}^T \boldsymbol{\beta})}\right)^{k-1} \left(\frac{1}{1 + \left(\frac{t}{\exp(\boldsymbol{x}_{ij}^T \boldsymbol{\beta})}\right)^k}\right)^{v_i+1}.$$

For log-logistic distribution, the distribution of RMST for each group is obtained below

$$RMST_{LL}(\tau, x_1, \beta_0^*, \beta_1^*) = \exp(\beta_0^* + x_1\beta_1^*)B\left(\frac{\left(\frac{\tau}{\exp(\beta_0^* + x_1\beta_1^*)}\right)^{k^*}}{1 + \left(\frac{\tau}{\exp(\beta_0^* + x_1\beta_1^*)}\right)^{k^*}}; 1 + \frac{1}{k^*}, 1 - \frac{1}{k^*}\right) + \tau \frac{1}{1 + \left(\frac{\tau}{\exp(\beta_0^* + x_1\beta_1^*)}\right)^{k^*}}.$$

The difference between the RMSTs is

$$RMST_{LL}(\tau, 1, \beta_0^*, \beta_1^*) - RMST_{LL}(\tau, 0, \beta_0^*, \beta_1^*).$$

The RMST for the mixed effects log-logistic model is

$$RMST_{LL,r}(\tau, x_1, \beta_0^*, \beta_1^*, u_i^*) = \exp(\beta_0^* + x_1\beta_1^* + u_i^*)B\left(\frac{\left(\frac{\tau}{\exp(\beta_0^* + x_1\beta_1^* + u_i^*)}\right)^{k^*}}{1 + \left(\frac{\tau}{\exp(\beta_0^* + x_1\beta_1^* + u_i^*)}\right)^{k^*}}; 1 + \frac{1}{k^*}, 1 - \frac{1}{k^*}\right) + \tau \frac{1}{1 + \left(\frac{\tau}{\exp(\beta_0^* + x_1\beta_1^* + u_i^*)}\right)^{k^*}}.$$

For frailty model, the hazard function is

$$h_f(t|v) = v \frac{\frac{k}{\alpha} \left(\frac{t}{\alpha}\right)^{k-1}}{1 + \left(\frac{t}{\alpha}\right)^k}.$$

The density function and survival function are

$$f_f(t|v) = v \frac{k}{\alpha} \left(\frac{t}{\alpha}\right)^{k-1} \left(\frac{1}{1+\left(\frac{t}{\alpha}\right)^k}\right)^{v+1}, S(t|v) = \left(\frac{1}{1+\left(\frac{t}{\alpha}\right)^k}\right)^v.$$

The RMST is calculated as follows

$$\int_0^\tau S(t|v)dt = \int_0^\tau vt \frac{k}{\alpha} \left(\frac{t}{\alpha}\right)^{k-1} \left(\frac{1}{1+\left(\frac{t}{\alpha}\right)^k}\right)^{v+1} dt + \tau \left(\frac{1}{1+\left(\frac{t}{\alpha}\right)^k}\right)^v$$
$$\left(\left(\frac{t}{\alpha}\right)^k = x \Leftrightarrow t = \alpha x^{\frac{1}{k}}, \frac{k}{\alpha} \left(\frac{t}{\alpha}\right)^{k-1} dt = dx\right)$$
$$= \int_0^{\left(\frac{\tau}{\alpha}\right)^k} \alpha v x^{\frac{1}{k}} \left(\frac{1}{1+x}\right)^{v+1} dx + \tau \left(\frac{1}{1+\left(\frac{t}{\alpha}\right)^k}\right)^v$$
$$= \alpha v B \left(\frac{\left(\frac{\tau}{\alpha}\right)^k}{1+\left(\frac{\tau}{\alpha}\right)^k}; 1+\frac{1}{k}, v-\frac{1}{k}\right) + \tau \left(\frac{1}{1+\left(\frac{t}{\alpha}\right)^k}\right)^v.$$

The RMST for the log-logistic frailty model is

$$RMST_{LL,f}(\tau, x_1, \beta_0^*, \beta_1^*, v_i^*) = v \left(\beta_0^* + x_1 \beta_1^*\right) B \left(\frac{\left(\frac{\tau}{(\beta_0^* + x_1 \beta_1^*)}\right)^k}{1 + \left(\frac{\tau}{(\beta_0^* + x_1 \beta_1^*)}\right)^k}; 1 + \frac{1}{k}, v_i^* - \frac{1}{k}\right) + \tau \left(\frac{1}{1 + \left(\frac{t}{(\beta_0^* + x_1 \beta_1^*)}\right)^k}\right)^{v_i^*}.$$

#### A.6 Posterior probability of mixed effect models

#### A.6.1 Mixed effects exponential model

The density function and survival function, conditional on the random effect  $u_i$  for *i*-th cluster, are

$$f_r(t_{ij}|u_i) = \exp\{\boldsymbol{x}_{ij}^T \boldsymbol{\beta} + u_i\} \exp\{-\exp(\boldsymbol{x}_{ij}^T \boldsymbol{\beta} + u_i)t_{ij}\},\$$
  
$$S_r(t_{ij}|u_i) = \exp\{-\exp(\boldsymbol{x}_{ij}^T \boldsymbol{\beta} + u_i)t_{ij}\}.$$

The posterior probability is

$$\pi_r(\boldsymbol{\beta}, \boldsymbol{u}, \phi | \boldsymbol{t}, \boldsymbol{X}) \propto L_r(\boldsymbol{t} | \boldsymbol{X}, \boldsymbol{\beta}, \boldsymbol{u}, \phi) g_r(\boldsymbol{u} | \phi) \pi(\boldsymbol{\beta} | \boldsymbol{c}) \pi(\phi | \xi),$$

where

$$L_r(\boldsymbol{t}|\boldsymbol{X},\boldsymbol{\beta},\boldsymbol{u},\boldsymbol{\phi}) = \prod_{i=1}^M \prod_{j=1}^{n_i} \{f_r(t_{ij}|v_i)\}^{\delta_{ij}} \{S_r(t_{ij}|v_i)\}^{1-\delta_{ij}}$$
$$= \prod_{i=1}^M \prod_{j=1}^{n_i} \left[\exp\{\boldsymbol{x}_{ij}^T\boldsymbol{\beta} + u_i\}\exp\{-\exp(\boldsymbol{x}_{ij}^T\boldsymbol{\beta} + u_i)t_{ij}\}\right]^{\delta_{ij}}$$
$$\times \left[\exp\{-\exp(\boldsymbol{x}_{ij}^T\boldsymbol{\beta} + u_i)t_{ij}\}\right]^{1-\delta_{ij}},$$

the variable  $\delta_{ij}$  is a censoring indicator that takes the value 1 if an event occurs and 0 if censored,  $g_r(\boldsymbol{u}|\phi)$  is the density function of  $\boldsymbol{u}, \boldsymbol{\beta} \sim \mathcal{N}(0, \boldsymbol{\Sigma}_c), \boldsymbol{\Sigma}_c = \text{diag}(c_1, c_2, \ldots, c_p), u_i \sim \mathcal{N}(0, \phi^2), \phi \sim \mathcal{U}(0, \xi)$ , and  $\xi$  is a hyperparameter.

#### A.6.2 Mixed effects Weibull model

The density function and survival function, conditional on the random effect  $u_i$  for *i*-th cluster, are

$$f_r(t_{ij}|u_i) = \exp(\boldsymbol{x}_{ij}^T \boldsymbol{\beta} + u_i) k t_{ij}^{k-1} \exp\left\{-\exp(\boldsymbol{x}_{ij}^T \boldsymbol{\beta} + u_i) t_{ij}^k\right\}, \quad S_r(t_{ij}|u_i) = \exp\left\{-\exp(\boldsymbol{x}_{ij}^T \boldsymbol{\beta} + u_i) t_{ij}^k\right\}.$$

The posterior probability is

$$\pi_r(\boldsymbol{\beta}, \boldsymbol{u}, \phi, k | \boldsymbol{t}, \boldsymbol{X}) \propto L_r(\boldsymbol{t} | \boldsymbol{X}, \boldsymbol{\beta}, \boldsymbol{u}, \phi, k) g_r(\boldsymbol{u} | \phi) \pi(\boldsymbol{\beta} | \boldsymbol{c}) \pi(\phi | \xi) \pi(k | a, b),$$

where

$$L_{r}(\boldsymbol{t}|\boldsymbol{X},\boldsymbol{\beta},\boldsymbol{u},\phi,k) = \prod_{i=1}^{M} \prod_{j=1}^{n_{i}} \{f_{r}(t_{ij}|u_{i})\}^{\delta_{ij}} \{S_{r}(t_{ij}|u_{i})\}^{1-\delta_{ij}}$$
$$= \prod_{i=1}^{M} \prod_{j=1}^{n_{i}} \left[\exp(\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta}+u_{i})kt_{ij}^{k-1}\exp\left\{-\exp(\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta}+u_{i})t_{ij}^{k}\right\}\right]^{\delta_{ij}}$$
$$\times \left[\exp\left\{-\exp(\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta}+u_{i})t_{ij}^{k}\right\}\right]^{1-\delta_{ij}},$$

 $\boldsymbol{\beta} \sim \mathcal{N}(0, \boldsymbol{\Sigma}_{c}), \, \boldsymbol{\Sigma}_{c} = \text{diag}(c_{1}, c_{2}, \dots, c_{p}), \, u_{i} \sim \mathcal{N}(0, \phi^{2}), \, \phi \sim \mathcal{U}(0, \xi), \, \text{and} \, k \sim Gamma(a, b).$ 

#### A.6.3 Mixed effects log-logistic model

The density function and survival function, conditional on the random effect  $u_i$  for *i*-th cluster, are

$$f_r(t_{ij}|u_i) = \frac{\exp\{\boldsymbol{x}_{ij}^T \boldsymbol{\beta} + u_i\} k t_{ij}^{k-1}}{\left(1 + \exp(\boldsymbol{x}_{ij}^T \boldsymbol{\beta} + u_i) t^k\right)^2}, S_r(t_{ij}|u_i) = \frac{1}{1 + \exp(\boldsymbol{x}_{ij}^T \boldsymbol{\beta} + u_i) t^k}.$$

The posterior probability is

$$\pi_r(\boldsymbol{\beta}, \boldsymbol{u}, \phi, k | \boldsymbol{t}, \boldsymbol{X}) \propto L_r(\boldsymbol{t} | \boldsymbol{X}, \boldsymbol{\beta}, \boldsymbol{u}, \phi, k) g_r(\boldsymbol{u} | \phi) \pi(\boldsymbol{\beta} | \boldsymbol{c}) \pi(\phi | \xi) \pi(k | a, b),$$

where

$$L_{r}(\boldsymbol{t}|\boldsymbol{X},\boldsymbol{\beta},\boldsymbol{u},\phi,k) = \prod_{i=1}^{M} \prod_{j=1}^{n_{i}} \{f_{r}(t_{ij}|u_{i})\}^{\delta_{ij}} \{S_{r}(t_{ij}|u_{i})\}^{1-\delta_{ij}}$$
$$= \prod_{i=1}^{M} \prod_{j=1}^{n_{i}} \left[ \frac{\exp\{\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta} + u_{i}\}kt_{ij}^{k-1}}{\left(1 + \exp(\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta} + u_{i})t^{k}\right)^{2}} \right]^{\delta_{ij}} \left[ \frac{1}{1 + \exp(\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta} + u_{i})t^{k}} \right]^{1-\delta_{ij}},$$

 $\boldsymbol{\beta} \sim \mathcal{N}(0, \boldsymbol{\Sigma}_{c}), \ \boldsymbol{\Sigma}_{c} = \operatorname{diag}(c_{1}, c_{2}, \dots, c_{p}), \ u_{i} \sim \mathcal{N}(0, \phi^{2}), \ \phi \sim \mathcal{U}(0, \xi), \ \mathrm{and} \ k \sim Gamma(a, b).$ 

#### A.6.4 Mixed effects log-normal model

The density function and survival function, conditional on the random effect  $u_i$  for *i*-th cluster, are

$$f_r(t_{ij}|u_i) = \frac{1}{t\sqrt{2\pi\sigma^2}} \exp\left[-\frac{\{\log(t) - (\boldsymbol{x}_{ij}^T\boldsymbol{\beta} + u_i)\}^2}{2\sigma^2}\right],$$
$$S_r(t_{ij}|u_i) = 1 - \Phi\left(\frac{\log(t) - (\boldsymbol{x}_{ij}^T\boldsymbol{\beta} + u_i)}{\sigma}\right).$$

The posterior probability is

$$\pi_r(\boldsymbol{\beta}, \boldsymbol{u}, \phi, \sigma^2 | \boldsymbol{t}, \boldsymbol{X}) \propto L_r(\boldsymbol{t} | \boldsymbol{X}, \boldsymbol{\beta}, \boldsymbol{u}, \phi, \sigma^2) g_r(\boldsymbol{u} | \phi) \pi(\boldsymbol{\beta} | \boldsymbol{c}) \pi(\phi | \xi) \pi(\sigma^2 | a, b),$$

where

 $\boldsymbol{\beta} \sim \mathcal{N}(0,$ 

$$L_{r}(\boldsymbol{t}|\boldsymbol{X},\boldsymbol{\beta},\boldsymbol{u},\phi,\sigma^{2}) = \prod_{i=1}^{M} \prod_{j=1}^{n_{i}} \{f_{r}(t_{ij}|u_{i})\}^{\delta_{ij}} \{S_{r}(t_{ij}|u_{i})\}^{1-\delta_{ij}}$$
$$= \prod_{i=1}^{M} \prod_{j=1}^{n_{i}} \left[\frac{1}{t\sqrt{2\pi\sigma^{2}}} \exp\left[-\frac{\{\log(t) - (\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta} + u_{i})\}^{2}}{2\sigma^{2}}\right]\right]^{\delta_{ij}}$$
$$\times \left[1 - \Phi\left(\frac{\log(t) - (\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta} + u_{i})}{\sigma}\right)\right]^{1-\delta_{ij}},$$
$$\boldsymbol{\Sigma_{c}}, \boldsymbol{\Sigma_{c}} = \operatorname{diag}\left(c_{1}, c_{2}, \dots, c_{p}\right), u_{i} \sim \mathcal{N}\left(0, \phi^{2}\right), \phi \sim \mathcal{U}(0, \xi), \text{ and } \sigma^{2} \sim \mathcal{U}(0, s).$$

# A.7 Posterior probability of frailty models

#### A.7.1 Exponential frailty model

The survival function  $S_f(t_{ij}|v_i)$  conditional on the frailty  $u_i$  for *i*-th cluster is expressed as

$$S_f(t_{ij}|v_i) = \exp\left\{-\int_0^{t_{ij}} h_f(s|v_i)ds\right\} = \exp\left\{-\int_0^{t_{ij}} v_i \exp\{\boldsymbol{x}_{ij}^T\boldsymbol{\beta}\}ds\right\} = \exp\left\{-v_i \exp\{\boldsymbol{x}_{ij}^T\boldsymbol{\beta}\}t_{ij}\right\}.$$

The density function  $f_f(t_{ij}|v_i)$  conditional on the frailty  $u_i$  for *i*-th cluster is

$$f_f(t_{ij}|v_i) = -\frac{d}{dt}S_f(t|v_i)\bigg|_{t=t_{ij}} = v_i \exp\{\boldsymbol{x}_{ij}^T \boldsymbol{\beta}\} \exp\{-v_i \exp\{\boldsymbol{x}_{ij}^T \boldsymbol{\beta}\} t_{ij}\}.$$

The posterior probability is

$$\pi_f(\boldsymbol{\beta}, \boldsymbol{v}, \phi | \boldsymbol{t}, \boldsymbol{X}) \propto L_f(\boldsymbol{t} | \boldsymbol{X}, \boldsymbol{\beta}, \boldsymbol{v}, \phi) g_f(\boldsymbol{v} | \phi) \pi(\boldsymbol{\beta} | \boldsymbol{c}) \pi(\phi | \xi),$$

where

$$L_{f}(\boldsymbol{t}|\boldsymbol{X},\boldsymbol{\beta},\boldsymbol{v},\boldsymbol{\phi}) = \prod_{i=1}^{M} \prod_{j=1}^{n_{i}} \{f_{f}(t_{ij}|v_{i})\}^{\delta_{ij}} \{S_{f}(t_{ij}|v_{i})\}^{1-\delta_{ij}}$$
  
$$= \prod_{i=1}^{M} \prod_{j=1}^{n_{i}} [v_{j} \exp\{\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta}\} \exp\{-v_{j} \exp\{\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta}\}t\}]^{\delta_{ij}} [\exp\{-v_{j} \exp\{\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta}\}t\}]^{1-\delta_{ij}},$$

 $\boldsymbol{\beta} \sim \mathcal{N}(0, \boldsymbol{\Sigma}_{\boldsymbol{c}}), \, \boldsymbol{\Sigma}_{\boldsymbol{c}} = \operatorname{diag}(c_1, c_2, \dots, c_p), \, v_i \sim \operatorname{Gamma}\left(\frac{1}{\phi}, \frac{1}{\phi}\right), \, \phi \sim \mathcal{U}(0, \xi).$ 

### A.7.2 Weibull frailty model

The survival function conditional on the frailty  $u_i$  for *i*-th cluster is

$$S_f(t_{ij}|v_i) = \exp\left\{-\int_0^{t_{ij}} v_i \exp\{\boldsymbol{x}_{ij}^T \boldsymbol{\beta}\} k s^{k-1} ds\right\} = \exp\left\{-v_i \exp\{\boldsymbol{x}_{ij}^T \boldsymbol{\beta}\} t_{ij}^k\right\}.$$

The density function conditional on the frailty  $u_i$  for *i*-th cluster is

$$f_f(t_{ij}|v_i) = -\frac{d}{dt}S_f(t|v_i)\bigg|_{t=t_{ij}} = v_i \exp\{\boldsymbol{x}_{ij}^T \boldsymbol{\beta}\} k t_{ij}^{k-1} \exp\{-v_i \exp\{\boldsymbol{x}_{ij}^T \boldsymbol{\beta}\} t_{ij}^k\}.$$

The posterior probability is

$$\pi_f(\boldsymbol{\beta}, \boldsymbol{v}, \phi, k | \boldsymbol{t}, \boldsymbol{X}) \propto L_f(\boldsymbol{t} | \boldsymbol{X}, \boldsymbol{\beta}, \boldsymbol{v}, \phi, k) g_f(\boldsymbol{v} | \phi) \pi(\boldsymbol{\beta} | \boldsymbol{c}) \pi(\phi | \xi) \pi(k | a, b),$$

where

$$L_{f}(\boldsymbol{t}|\boldsymbol{X},\boldsymbol{\beta},\boldsymbol{v},\phi,k) = \prod_{i=1}^{M} \prod_{j=1}^{n_{i}} \{f_{f}(t_{ij}|v_{i})\}^{\delta_{ij}} \{S_{f}(t_{ij}|v_{i})\}^{1-\delta_{ij}}$$
$$= \prod_{i=1}^{M} \prod_{j=1}^{n_{i}} [v_{j} \exp\{\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta}\}kt^{k-1} \exp\{-v_{j} \exp\{\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta}\}t^{k}\}]^{\delta_{ij}}$$
$$\times \left[\exp\{-v_{j} \exp\{\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta}\}t^{k}\}\right]^{1-\delta_{ij}},$$

$$\boldsymbol{\beta} \sim \mathcal{N}(0, \boldsymbol{\Sigma}_{\boldsymbol{c}}), \, \boldsymbol{\Sigma}_{\boldsymbol{c}} = \operatorname{diag}(c_1, c_2, \dots, c_p), \, v_i \sim \operatorname{Gamma}\left(\frac{1}{\phi}, \frac{1}{\phi}\right), \, \phi \sim \mathcal{U}(0, \xi), \, \operatorname{and} \, k \sim \operatorname{Gamma}(a, b).$$

#### A.7.3 Log-logistic frailty model

The following formula is used to calculate the survival function conditional on the frailty.

$$S(t) = e^{-\int_0^t h(s)ds} \Leftrightarrow -\int_0^t h(s)ds = \log\left(S(t)\right)$$

The survival function conditional on the frailty  $u_i$  for *i*-th cluster is

$$S_f(t_{ij}|v_i) = \exp\left\{-\int_0^{t_{ij}} h_f(s|v_i)ds\right\} = \exp\left\{-v_i \int_0^{t_{ij}} h(s)ds\right\} = \exp\left\{\log\left(S(t)\right)^{v_i}\right\} = \left(\frac{1}{1+e^{\boldsymbol{x}_{ij}^T\boldsymbol{\beta}}t^k}\right)^{v_i}$$

The density function conditional on the frailty  $u_i$  for *i*-th cluster is

$$f_f(t_{ij}|v_i) = -\frac{d}{dt}S(t|v_i)\Big|_{t=t_{ij}} = v_i \exp\{\boldsymbol{x}_{ij}^T \boldsymbol{\beta}\} k t_{ij}^{k-1} \left(\frac{1}{1+e^{\boldsymbol{x}_{ij}^T \boldsymbol{\beta}} t^k}\right)^{v_i+1}$$

The posterior probability is

$$\pi_f(\boldsymbol{\beta}, \boldsymbol{v}, \phi, k | \boldsymbol{t}, \boldsymbol{X}) \propto L_f(\boldsymbol{t} | \boldsymbol{X}, \boldsymbol{\beta}, \boldsymbol{v}, \phi, k) g_f(\boldsymbol{v} | \phi) \pi(\boldsymbol{\beta} | \boldsymbol{c}) \pi(\phi | \xi) \pi(k | a, b),$$

where

$$\begin{split} L_{f}(\boldsymbol{t}|\boldsymbol{X},\boldsymbol{\beta},\boldsymbol{v},\phi,k) &= \prod_{i=1}^{M} \prod_{j=1}^{n_{i}} \{f(t_{ij}|v_{i})\}^{\delta_{ij}} \{S(t_{ij}|v_{i})\}^{1-\delta_{ij}} \\ &= \prod_{i=1}^{M} \prod_{j=1}^{n_{i}} \left[ v_{i} \exp\{\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta}\} k t_{ij}^{k-1} \left(\frac{1}{1+e^{\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta}} t^{k}}\right)^{v_{i}+1} \right]^{\delta_{ij}} \left[ \left(\frac{1}{1+e^{\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta}} t^{k}}\right)^{v_{i}} \right]^{1-\delta_{ij}}, \end{split}$$

 $\boldsymbol{\beta} \sim \mathcal{N}(0, \boldsymbol{\Sigma}_{c}), \ \boldsymbol{\Sigma}_{c} = \operatorname{diag}(c_{1}, c_{2}, \dots, c_{p}), \ v_{i} \sim \operatorname{Gamma}\left(\frac{1}{\phi}, \frac{1}{\phi}\right), \ \mathrm{and} \ \phi \sim \mathcal{U}(0, \xi), \ k \sim \operatorname{Gamma}(a, b).$ 

#### A.7.4 Log-normal frailty model

The survival function conditional on the frailty  $u_i$  for *i*-th cluster is

$$S_f(t_{ij}|v_i) = (S(t))^{v_i} = \left(1 - \Phi\left(\frac{\log(t) - \boldsymbol{x}_{ij}^T \boldsymbol{\beta}}{\sigma}\right)\right)^{v_i}$$

The density function conditional on the frailty  $u_i$  for *i*-th cluster is

$$f_f(t_{ij}|v_i) = -\frac{d}{dt}S(t|v_i)\bigg|_{t=t_{ij}} = v_i \frac{1}{t\sqrt{2\pi\sigma^2}} \exp\left\{-\frac{(\log(t) - \boldsymbol{x}_{ij}^T\boldsymbol{\beta})^2}{2\sigma^2}\right\} \left(1 - \Phi\left(\frac{\log(t) - \boldsymbol{x}_{ij}^T\boldsymbol{\beta}}{\sigma}\right)\right)^{v_i - 1}$$

The posterior probability is

$$\pi_f(\boldsymbol{\beta}, \boldsymbol{v}, \phi, \sigma^2 | \boldsymbol{t}, \boldsymbol{X}) \propto L_f(\boldsymbol{t} | \boldsymbol{X}, \boldsymbol{\beta}, \boldsymbol{v}, \phi, \sigma^2) g_f(\boldsymbol{v} | \phi) \pi(\boldsymbol{\beta} | \boldsymbol{c}) \pi(\phi | \xi) \pi(\sigma^2 | a, b),$$

where

$$L_{f}(\boldsymbol{t}|\boldsymbol{X},\boldsymbol{\beta},\boldsymbol{v},\boldsymbol{\phi},\sigma^{2}) = \prod_{i=1}^{M} \prod_{j=1}^{n_{i}} \{f(t_{ij}|v_{i})\}^{\delta_{ij}} \{S(t_{ij}|v_{i})\}^{1-\delta_{ij}}$$

$$= \prod_{i=1}^{M} \prod_{j=1}^{n_{i}} \left[ v_{i} \frac{1}{t\sqrt{2\pi\sigma^{2}}} \exp\left\{-\frac{(\log(t)-\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta})^{2}}{2\sigma^{2}}\right\} \left(1-\Phi\left(\frac{\log(t)-\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta}}{\sigma}\right)\right)^{v_{i}-1}\right]^{\delta_{ij}}$$

$$\times \left[ \left(1-\Phi\left(\frac{\log(t)-\boldsymbol{x}_{ij}^{T}\boldsymbol{\beta}}{\sigma}\right)\right)^{v_{i}}\right]^{1-\delta_{ij}},$$

 $\boldsymbol{\beta} \sim \mathcal{N}(0, \boldsymbol{\Sigma}_{\boldsymbol{c}}), \, \boldsymbol{\Sigma}_{\boldsymbol{c}} = \operatorname{diag}(c_1, c_2, \dots, c_p), \, v_i \sim \operatorname{Gamma}\left(\frac{1}{\phi}, \frac{1}{\phi}\right), \, \phi \sim \mathcal{U}(0, \xi), \, \operatorname{and} \, \sigma^2 \sim \mathcal{U}(0, s).$ 

# A.8 Supplemental information for simulation

A.8.1 Survival probability in each scenario



Figure 1: Survival probabilities of simulation.

#### A.8.2 Simulation specifications and results of scenario C

Scenario C is a basic proportional hazard assumption based on exponential distribution. The parameter is given as

$$\lambda_{ij} = \exp(\beta_0 + \beta_1 x_{1ij} + \beta_2 x_{2ij} + v_i),$$

where  $\beta_0 = -4.5$  is a baseline effect,  $\beta_1 = 0.5$  is a treatment effect,  $\beta_2 = 1$  is an effect of the covariate, and other parameters are set as well as scenario A. We consider 4 clusters for the simulation. The theoretical RMSTs in Scenario C are 45.85 and 60.37 by equation (??), respectively. Thus, the true difference between the RMSTs is -14.52.

The simulation results were in Table 1 for scenario C in the Appendix. In scenario C, where the exponential model with random effect is true, the exponential and Weibull models had small biases and MSEs. The log-logistic and log-normal models had some biases in the scenario. The log-normal model with frailty effects included a significant bias, and consequently, the MSE was also significant. In summary, bias and MSE were larger when inferences were made using models that deviated from the true model. It is noted that the frailty effect may obtain extreme accuracy, like the log-normal model in scenario C. Therefore, an appropriate model with or without frailty or random effects should be selected for estimating RMST with small bias and MSE.

		Exponential		Lo	g-Logis	tic	Log-Normal			Weibull			
	n	Т	$\mathbf{F}$	Μ	Т	$\mathbf{F}$	Μ	Т	$\mathbf{F}$	Μ	Т	$\mathbf{F}$	М
Scenario	C: Exp	ponentia	al model	l with ra	andom e	effect							
Mean	64	0.94	1.82	0.88	1.64	1.37	1.72	2.52	2.97	2.52	1.15	0.97	1.08
(Bias)	512	0.50	0.92	0.26	1.39	2.60	1.28	2.19	10.46	2.11	0.71	0.37	0.28
	2048	0.94	1.59	1.00	1.86	3.71	1.99	2.53	14.25	2.60	1.08	1.08	1.00
MSE	64	81.75	73.34	83.53	87.73	92.44	87.10	86.94	89.72	88.25	80.19	83.11	80.80
	512	11.00	10.03	10.27	12.84	15.08	11.91	15.43	113.58	14.85	10.89	10.13	9.95
	2048	4.13	5.27	3.75	6.48	15.55	6.75	9.27	203.06	9.41	4.39	3.77	3.68
Mode	64	0.74	2.27	1.28	1.28	1.40	1.64	1.99	3.50	2.35	1.00	1.22	1.02
	512	0.40	0.48	0.19	1.37	2.70	1.22	2.06	11.84	2.03	0.57	0.35	0.35
	2048	0.94	1.12	0.73	1.83	3.71	1.90	2.48	14.40	2.55	1.04	0.85	0.78
Median	64	0.81	1.94	0.84	1.53	1.33	1.68	2.39	3.06	2.44	1.06	0.99	1.05
	512	0.49	0.77	0.21	1.38	2.62	1.25	2.17	10.96	2.09	0.69	0.29	0.24
	2048	0.94	1.31	0.91	1.86	3.71	1.92	2.53	14.30	2.56	1.07	0.93	0.91

Table 1: Simulation results of scenario C: exponential model with random effects

T: typical fixed model, F: frailty effects model, R: random effects model, Mean, Mode, and Median are the difference from true RMST.

# A.9 Supplemental information for DHS data analysis

# A.9.1 Kaplan-Meier plot of DHS data



Figure 2: Kaplan-Meier plot (12–19 group vs 20–30 group)



Figure 3: Kaplan-Meier plot (12–19 group vs 31+ group)

# A.9.2 Estimation results (12–19 group vs 20–30 group)

	Parameter	Mode	Median	Mean	SE	95%CI	Rhat	ESS
	Intercept	4.44	4.42	4.43	0.25	[3.97, 4.95]	1.00	788
	20–30 group	0.02	0.01	0.01	0.03	[-0.04, 0.07]	1.00	3653
	Sex	0.12	0.13	0.13	0.03	[0.07,  0.19]	1.00	4055
Place	Respondents home	0.89	0.80	0.79	0.24	[0.28,  1.23]	1.00	758
Place	Other home	0.92	0.88	0.89	0.46	[0.00,  1.83]	1.00	1780
Place	Public sector	1.09	1.06	1.06	0.27	[0.50,  1.57]	1.00	850
Place	Government hospital	0.99	0.94	0.93	0.24	[0.42, 1.37]	1.00	723
Place	CS Govt health professional	0.86	0.90	0.90	0.24	[0.39,  1.34]	1.00	738
Place	Other public sector	0.95	0.98	0.97	0.27	[0.41,  1.51]	1.00	854
Place	Private hospital/clinic	0.58	0.61	0.61	0.24	[0.10,  1.06]	1.00	741
Place	CS private health facility	0.96	0.95	0.94	0.29	[0.33, 1.49]	1.00	963
Size	Very large	1.13	1.15	1.15	0.09	[0.98,  1.32]	1.00	1643
Size	Larger than average	1.23	1.22	1.22	0.08	[1.07, 1.38]	1.00	1554
Size	Average	1.33	1.32	1.32	0.07	[1.19, 1.45]	1.00	1271
Size	Smaller than average	0.96	0.94	0.94	0.08	[0.78,  1.10]	1.00	1453
Size	Very small	0.15	0.16	0.16	0.08	[0.00,  0.32]	1.00	1581
	Order	-0.03	-0.03	-0.03	0.01	[-0.05, -0.01]	1.00	4215
	$RMST_{12-19group}$	37.76	37.58	37.45	2.46	[32.33, 42.11]		
	$\mathrm{RMST}_{20-30group}$	38.13	37.71	37.58	2.43	[32.59, 42.16]		
	$\mathrm{RMST}_{diff}$	0.05	0.13	0.13	0.30	[-0.43, 0.72]		

Table 2: Results of exponential model (12–19 group vs 20–30 group)

	Parameter	Mode	Median	Mean	SE	95%CI	Rhat	ESS
	Intercept	4.35	4.43	4.44	0.26	[3.96, 4.97]	1.00	1230
	20–30 group	0.01	0.03	0.03	0.03	[-0.04, 0.09]	1.00	3905
	$\mathbf{Sex}$	0.14	0.13	0.13	0.03	[0.08,  0.19]	1.00	3528
Place	Respondents home	0.85	0.84	0.82	0.24	[0.32, 1.28]	1.01	1083
Place	Other home	0.82	0.92	0.93	0.47	[0.06,  1.89]	1.00	2359
Place	Public sector	1.08	1.08	1.07	0.27	[0.52,  1.58]	1.00	1231
Place	Government hospital	0.93	0.95	0.94	0.24	[0.44,  1.39]	1.01	1085
Place	CS Govt health professional	0.89	0.92	0.91	0.24	[0.42,  1.35]	1.01	1105
Place	Other public sector	1.08	1.02	1.01	0.27	[0.45,  1.53]	1.00	1300
Place	Private hospital/clinic	0.59	0.66	0.65	0.24	[0.15, 1.10]	1.01	1080
Place	CS private health facility	1.20	1.03	1.02	0.29	[0.42,  1.58]	1.00	1385
Size	Very large	1.15	1.16	1.16	0.08	[1.00, 1.33]	1.00	2092
Size	Larger than average	1.22	1.22	1.22	0.08	[1.07,  1.38]	1.00	1949
Size	Average	1.34	1.32	1.32	0.07	[1.19, 1.45]	1.00	1720
Size	Smaller than average	0.97	0.94	0.94	0.08	[0.79,  1.09]	1.00	1944
Size	Very small	0.16	0.17	0.17	0.08	[0.00,  0.33]	1.00	2105
	Order	-0.02	-0.02	-0.02	0.01	[-0.04, 0.00]	1.00	4496
	$\phi$	0.17	0.17	0.18	0.07	[0.09,  0.36]	1.01	1099
	$RMST_{12-19group}$	36.86	37.62	37.55	2.58	[32.18, 42.23]		
	$RMST_{20-30group}$	38.70	37.89	37.80	2.55	[32.43, 42.44]		
	RMST <sub>diff</sub>	0.19	0.25	0.25	0.32	[-0.36, 0.89]		

Table 3: Results of mixed effects exponential model (12–19 group vs 20–30 group)

	Parameter	Mode	Median	Mean	SE	95%CI	Rhat	ESS
	Intercept	4.39	4.44	4.45	0.26	[3.96, 5.00]	1.00	888
	20–30 group	0.02	0.02	0.02	0.03	[-0.03, 0.08]	1.00	4368
	Sex	0.13	0.13	0.13	0.03	[0.08, 0.19]	1.00	4676
Place	Respondents home	0.82	0.84	0.83	0.25	[0.30, 1.27]	1.00	849
Place	Other home	0.76	0.93	0.94	0.47	[0.08, 1.91]	1.00	2227
Place	Public sector	1.12	1.08	1.07	0.27	[0.51,  1.59]	1.00	907
Place	Government hospital	0.96	0.95	0.94	0.25	[0.42, 1.39]	1.00	833
Place	CS Govt health professional	0.86	0.93	0.92	0.24	[0.40, 1.36]	1.00	838
Place	Other public sector	1.06	1.02	1.02	0.27	[0.47, 1.54]	1.00	991
Place	Private hospital/clinic	0.63	0.66	0.65	0.25	[0.14, 1.10]	1.00	835
Place	CS private health facility	1.04	1.03	1.03	0.30	[0.42, 1.60]	1.00	1165
Size	Very large	1.15	1.16	1.16	0.08	[1.00, 1.33]	1.00	1812
Size	Larger than average	1.22	1.22	1.22	0.08	[1.07, 1.37]	1.00	1824
Size	Average	1.33	1.31	1.31	0.07	[1.18, 1.44]	1.00	1576
Size	Smaller than average	0.91	0.93	0.93	0.08	[0.78,  1.09]	1.00	1797
Size	Very small	0.16	0.17	0.17	0.08	[0.00, 0.33]	1.00	1940
	Order	-0.02	-0.02	-0.02	0.01	[-0.04, 0.00]	1.00	4882
	$\phi$	0.02	0.03	0.05	0.07	[0.01, 0.19]	1.00	1595
	$RMST_{12-19qroup}$	38.25	37.80	37.66	2.61	[32.21, 42.43]		
	$RMST_{20-30group}$	38.60	38.06	37.91	2.57	[32.59, 42.59]		
	RMST <sub>diff</sub>	0.22	0.24	0.25	0.30	[-0.32, 0.88]		

Table 4: Results of exponential frailty model (12–19 group vs 20–30 group)

	Parameter	Mode	Median	Mean	SE	95%CI	Rhat	ESS
	Intercept	3.97	3.99	3.99	0.15	[3.71, 4.28]	1.00	415
	20–30 group	0.00	-0.01	-0.01	0.02	[-0.04, 0.03]	1.00	2078
	$\mathbf{Sex}$	0.08	0.08	0.08	0.02	[0.04,  0.11]	1.00	1939
Place	Respondents home	0.52	0.52	0.51	0.14	[0.23,  0.79]	1.00	377
Place	Other home	0.54	0.56	0.57	0.27	[0.09, 1.12]	1.00	906
Place	Public sector	0.67	0.67	0.66	0.16	[0.34,  0.98]	1.00	427
Place	Government hospital	0.58	0.58	0.57	0.14	[0.30,  0.85]	1.00	371
Place	CS Govt health professional	0.55	0.56	0.56	0.14	[0.29, 0.83]	1.00	374
Place	Other public sector	0.58	0.60	0.60	0.16	[0.29,  0.91]	1.00	465
Place	Private hospital/clinic	0.33	0.37	0.37	0.14	[0.09,  0.65]	1.00	377
Place	CS private health facility	0.62	0.58	0.57	0.17	[0.24,  0.92]	1.00	504
Size	Very large	0.65	0.66	0.66	0.05	[0.57,  0.75]	1.00	1019
Size	Larger than average	0.71	0.70	0.70	0.05	[0.61,  0.79]	1.00	907
Size	Average	0.77	0.77	0.76	0.04	[0.69, 0.84]	1.00	815
Size	Smaller than average	0.54	0.53	0.53	0.05	[0.44,  0.62]	1.00	957
Size	Very small	0.07	0.07	0.07	0.05	[-0.03, 0.16]	1.00	1106
	Order	-0.02	-0.02	-0.02	0.01	[-0.03, -0.01]	1.00	2833
	k	1.70	1.70	1.70	0.02	[1.66, 1.75]	1.00	1049
	$RMST_{12-19group}$	38.80	37.34	37.21	2.39	[32.15, 41.57]		
	$\mathrm{RMST}_{20-30group}$	37.34	37.26	37.09	2.41	[31.90, 41.44]		
	$\mathrm{RMST}_{diff}$	-0.13	-0.12	-0.12	0.31	[-0.77, 0.51]		

Table 5: Results of Weibull model (12–19 group vs 20–30 group)

	Parameter	Mode	Median	Mean	SE	95%CI	Rhat	ESS
	Intercept	4.00	3.99	4.00	0.15	[3.71, 4.30]	1.00	489
	20–30 group	0.00	0.00	0.00	0.02	[-0.04, 0.03]	1.00	2376
	Sex	0.08	0.08	0.08	0.02	[0.04,  0.11]	1.00	2132
Place	Respondents home	0.52	0.53	0.53	0.14	[0.24,  0.80]	1.00	476
Place	Other home	0.46	0.58	0.59	0.27	[0.08, 1.14]	1.00	1090
Place	Public sector	0.69	0.67	0.66	0.16	[0.34,  0.96]	1.00	534
Place	Government hospital	0.67	0.57	0.57	0.14	[0.28,  0.83]	1.00	482
Place	CS Govt health professional	0.51	0.56	0.56	0.14	[0.27,  0.82]	1.00	462
Place	Other public sector	0.69	0.62	0.61	0.16	[0.28,  0.92]	1.00	515
Place	Private hospital/clinic	0.38	0.39	0.39	0.14	[0.09,  0.66]	1.00	471
Place	CS private health facility	0.60	0.61	0.61	0.18	[0.27,  0.96]	1.00	638
Size	Very large	0.66	0.67	0.67	0.05	[0.57,  0.77]	1.00	878
Size	Larger than average	0.72	0.70	0.70	0.05	[0.61,  0.79]	1.01	868
Size	Average	0.77	0.76	0.76	0.04	[0.68, 0.84]	1.01	727
Size	Smaller than average	0.52	0.53	0.53	0.05	[0.44,  0.62]	1.01	788
Size	Very small	0.08	0.07	0.07	0.05	[-0.02, 0.17]	1.00	912
	Order	-0.01	-0.01	-0.01	0.01	[-0.02, 0.00]	1.00	3125
	k	1.70	1.71	1.71	0.02	[1.67,  1.75]	1.00	2106
	$\phi$	0.02	0.04	0.05	0.06	[0.01,  0.20]	1.00	511
	$RMST_{12-19group}$	37.33	37.37	37.33	2.51	[32.20, 41.80]		
	$\mathrm{RMST}_{20-30group}$	36.53	37.37	37.31	2.51	[32.10, 41.79]		
	$\mathrm{RMST}_{diff}$	0.03	-0.01	-0.02	0.30	[-0.62, 0.60]		

Table 6: Results of Weibull frailty model (12–19 group vs 20–30 group)

	Parameter	Mode	Median	Mean	SE	95%CI	Rhat	ESS
	Intercept	4.44	4.48	4.49	0.23	[4.04, 4.96]	1.00	824
	20–30 group	-0.01	-0.02	-0.02	0.02	[-0.06, 0.02]	1.00	3736
	$\mathbf{Sex}$	0.10	0.11	0.11	0.02	[0.06,  0.15]	1.00	3710
Place	Respondents home	0.71	0.62	0.62	0.23	[0.14,  1.06]	1.00	854
Place	Other home	0.54	0.60	0.61	0.38	[-0.12, 1.36]	1.00	1651
Place	Public sector	0.83	0.78	0.78	0.24	[0.28, 1.26]	1.00	915
Place	Government hospital	0.78	0.73	0.72	0.23	[0.25, 1.16]	1.00	857
Place	CS Govt health professional	0.79	0.70	0.70	0.23	[0.23, 1.14]	1.00	837
Place	Other public sector	0.80	0.78	0.77	0.25	[0.27, 1.24]	1.00	927
Place	Private hospital/clinic	0.54	0.47	0.46	0.23	[-0.01, 0.90]	1.00	842
Place	CS private health facility	0.72	0.69	0.69	0.26	[0.19, 1.20]	1.00	1005
Size	Very large	0.87	0.87	0.87	0.07	[0.73, 1.02]	1.01	1836
Size	Larger than average	0.93	0.93	0.93	0.07	[0.79,  1.07]	1.01	1767
Size	Average	1.02	1.02	1.02	0.06	[0.89, 1.14]	1.01	1577
Size	Smaller than average	0.73	0.71	0.71	0.07	[0.58,  0.85]	1.00	1711
Size	Very small	0.05	0.07	0.07	0.08	[-0.07, 0.22]	1.00	1896
	Order	-0.02	-0.02	-0.02	0.01	[-0.03, 0.00]	1.00	4360
	$\sigma$	1.54	1.54	1.54	0.02	[1.50, 1.57]	1.00	2287
	$RMST_{12-19group}$	40.63	40.39	40.28	1.88	[36.31, 43.79]		
	$RMST_{20-30group}$	40.57	40.18	40.11	1.90	[36.11, 43.71]		
	$\mathrm{RMST}_{diff}$	-0.16	-0.17	-0.17	0.19	[-0.55, 0.19]		

Table 7: Results of log-normal model (12–19 group vs 20–30 group)

	Parameter	Mode	Median	Mean	SE	95%CI	Rhat	ESS
	Intercept	4.51	4.48	4.49	0.23	[4.04,  4.97]	1.00	1119
	20–30 group	-0.02	-0.01	-0.01	0.02	[-0.06, 0.03]	1.00	3885
	Sex	0.12	0.11	0.11	0.02	[0.07,  0.15]	1.00	4243
Place	Respondents home	0.58	0.65	0.65	0.22	[0.22,  1.09]	1.00	1074
Place	Other home	0.53	0.64	0.66	0.37	[-0.04, 1.39]	1.00	2022
Place	Public sector	0.73	0.79	0.79	0.24	[0.33, 1.25]	1.00	1125
Place	Government hospital	0.70	0.73	0.73	0.22	[0.29,  1.16]	1.00	1083
Place	CS Govt health professional	0.76	0.72	0.71	0.22	[0.29,  1.15]	1.00	1089
Place	Other public sector	0.84	0.80	0.80	0.24	[0.32,  1.26]	1.00	1144
Place	Private hospital/clinic	0.53	0.50	0.50	0.22	[0.07,  0.92]	1.00	1100
Place	CS private health facility	0.73	0.75	0.75	0.25	[0.26, 1.24]	1.00	1299
Size	Very large	0.89	0.89	0.89	0.08	[0.74,  1.03]	1.00	1815
Size	Larger than average	0.93	0.93	0.93	0.07	[0.79,  1.07]	1.00	1830
Size	Average	1.01	1.02	1.02	0.06	[0.90, 1.14]	1.00	1624
Size	Smaller than average	0.74	0.72	0.72	0.07	[0.58, 0.85]	1.00	1914
Size	Very small	0.04	0.07	0.08	0.08	[-0.08, 0.23]	1.00	2100
'	Order	-0.02	-0.01	-0.01	0.01	[-0.03, 0.00]	1.00	4405
	$\sigma$	1.54	1.53	1.53	0.02	[1.50, 1.57]	1.00	3455
	$\phi$	0.10	0.12	0.13	0.05	[0.07, 0.25]	1.00	1151
	$RMST_{12-19qroup}$	40.62	40.40	40.31	1.88	[36.43, 43.83]		
	$RMST_{20-30group}$	40.24	40.29	40.21	1.88	[36.27, 43.79]		
	$\mathrm{RMST}_{diff}$	-0.09	-0.10	-0.10	0.19	[-0.48, 0.26]		

Table 8: Results of mixed effects log-normal model (12–19 group vs 20–30 group)

	Parameter	Mode	Median	Mean	SE	95%CI	Rhat	ESS
	Intercept	19.26	19.11	19.10	1.19	[16.79, 21.42]	1.00	790
	20–30 group	0.00	0.00	0.00	0.02	[-0.04, 0.03]	1.00	3775
	$\mathbf{Sex}$	0.08	0.08	0.08	0.02	[0.05,  0.12]	1.00	3934
Place	Respondents home	0.60	0.57	0.56	0.15	[0.25, 0.85]	1.00	1039
Place	Other home	0.63	0.61	0.62	0.28	[0.11,  1.18]	1.00	1938
Place	Public sector	0.80	0.71	0.71	0.17	[0.36,  1.02]	1.00	1171
Place	Government hospital	0.58	0.62	0.61	0.16	[0.30,  0.90]	1.00	1022
Place	CS Govt health professional	0.59	0.60	0.60	0.15	[0.28, 0.89]	1.00	1030
Place	Other public sector	0.70	0.67	0.66	0.17	[0.32,  0.99]	1.00	1141
Place	Private hospital/clinic	0.39	0.43	0.42	0.16	[0.10,  0.71]	1.00	1007
Place	CS private health facility	0.61	0.65	0.65	0.19	[0.27,  1.02]	1.00	1319
Size	Very large	0.72	0.72	0.72	0.05	[0.61, 0.83]	1.00	1505
Size	Larger than average	0.75	0.75	0.75	0.05	[0.65,  0.85]	1.00	1565
Size	Average	0.82	0.82	0.82	0.04	[0.73, 0.90]	1.00	1225
Size	Smaller than average	0.59	0.57	0.57	0.05	[0.47,  0.67]	1.00	1509
Size	Very small	0.07	0.08	0.08	0.06	[-0.03, 0.18]	1.00	1676
	Order	-0.01	-0.01	-0.01	0.01	[-0.02, 0.00]	1.00	3479
	$\sigma$	3.27	3.27	3.27	0.11	[3.05, 3.49]	1.00	849
	$\phi$	131492	446651	841037	1300743	[55426, 4005760]	1.00	1007
	$RMST_{12-19group}$	50.00	50.00	50.00	0.00	[50.00, 50.00]		
	$RMST_{20-30group}$	50.00	50.00	50.00	0.00	[50.00, 50.00]		
	$\mathrm{RMST}_{diff}$	0.00	0.00	0.00	0.00	[0.00, 0.00]		

Table 9: Results of log-normal frailty model (12–19 group vs 20–30 group)

The estimated results of the log-normal frailty model differ from the others; therefore, we examined the behavior using different data in Appendix A.10.

	Parameter	Mode	Median	Mean	SE	95%CI	Rhat	ESS
	Intercept	4.01	4.01	4.02	0.16	[3.73, 4.35]	1.01	1059
	20–30 group	-0.01	-0.01	-0.01	0.02	[-0.04, 0.03]	1.00	4416
	$\mathbf{Sex}$	0.08	0.08	0.08	0.02	[0.04,  0.11]	1.00	4233
Place	Respondents home	0.54	0.53	0.53	0.16	[0.21,  0.81]	1.00	997
Place	Other home	0.55	0.55	0.56	0.29	[0.01, 1.14]	1.00	1979
Place	Public sector	0.68	0.68	0.67	0.17	[0.32,  0.99]	1.00	1119
Place	Government hospital	0.62	0.60	0.59	0.16	[0.27,  0.88]	1.00	1001
Place	CS Govt health professional	0.56	0.58	0.58	0.16	[0.26,  0.86]	1.00	1003
Place	Other public sector	0.62	0.62	0.62	0.17	[0.26,  0.95]	1.01	1197
Place	Private hospital/clinic	0.45	0.40	0.39	0.16	[0.07,  0.68]	1.00	1009
Place	CS private health facility	0.63	0.59	0.59	0.19	[0.21,  0.93]	1.00	1274
Size	Very large	0.68	0.68	0.68	0.05	[0.58,  0.78]	1.00	1728
Size	Larger than average	0.71	0.72	0.72	0.05	[0.63,  0.82]	1.00	1679
Size	Average	0.79	0.78	0.78	0.04	[0.70,  0.87]	1.00	1451
Size	Smaller than average	0.54	0.55	0.55	0.05	[0.45,  0.65]	1.00	1604
Size	Very small	0.05	0.06	0.06	0.05	[-0.04, 0.16]	1.00	1938
	Order	-0.02	-0.02	-0.02	0.01	[-0.03, -0.01]	1.00	4461
	k	1.73	1.73	1.73	0.02	[1.69, 1.77]	1.00	2020
	$RMST_{12-19group}$	39.76	39.64	39.62	1.95	[35.80, 43.30]		
	$\mathrm{RMST}_{20-30group}$	39.85	39.54	39.53	1.96	[35.66, 43.23]		
	$\mathrm{RMST}_{diff}$	-0.02	-0.09	-0.09	0.22	[-0.53, 0.34]		

Table 10: Results of log-logistic model (12–19 group vs 20–30 group)

	Parameter	Mode	Median	Mean	SE	95%CI	Rhat	ESS
	Intercept	4.00	4.02	4.03	0.16	[3.73, 4.36]	1.01	1278
	20–30 group	0.00	0.00	0.00	0.02	[-0.04, 0.03]	1.00	3898
	$\mathbf{Sex}$	0.08	0.08	0.08	0.02	[0.04, 0.11]	1.00	4324
Place	Respondents home	0.61	0.56	0.55	0.15	[0.24, 0.83]	1.01	1214
Place	Other home	0.46	0.58	0.60	0.28	[0.04, 1.18]	1.00	2242
Place	Public sector	0.71	0.70	0.69	0.17	[0.34, 1.01]	1.00	1350
Place	Government hospital	0.62	0.61	0.60	0.15	[0.29, 0.89]	1.00	1231
Place	CS Govt health professional	0.57	0.60	0.59	0.15	[0.28, 0.87]	1.01	1220
Place	Other public sector	0.68	0.65	0.64	0.17	[0.30,  0.97]	1.00	1394
Place	Private hospital/clinic	0.45	0.43	0.42	0.15	[0.11, 0.70]	1.00	1228
Place	CS private health facility	0.59	0.64	0.64	0.18	[0.27, 0.99]	1.00	1543
Size	Very large	0.69	0.69	0.69	0.05	[0.58, 0.79]	1.00	1944
Size	Larger than average	0.72	0.72	0.72	0.05	[0.63, 0.82]	1.00	1785
Size	Average	0.77	0.78	0.78	0.04	[0.70, 0.87]	1.00	1599
Size	Smaller than average	0.55	0.55	0.55	0.05	[0.45, 0.65]	1.00	1749
Size	Very small	0.07	0.07	0.07	0.05	[-0.04, 0.17]	1.00	2246
	Order	-0.01	-0.01	-0.01	0.01	[-0.02, 0.00]	1.00	4957
	k	1.74	1.73	1.73	0.02	[1.69, 1.77]	1.00	3740
	$\phi$	0.08	0.10	0.11	0.05	[0.05, 0.23]	1.00	1239
	$RMST_{12-19group}$	39.48	39.74	39.70	1.96	[35.73, 43.31]		
	$RMST_{20-30group}$	39.67	39.71	39.68	1.95	[35.69, 43.33]		
	$\mathrm{RMST}_{diff}$	0.03	-0.02	-0.03	0.23	[-0.48, 0.42]		

Table 11: Results of mixed effects log-logistic model (12–19 group vs 20–30 group)

 $95\%\mathrm{CI}:\,95\%$  credible interval, ESS: Effective sample size

	Parameter	Mode	Median	Mean	SE	95%CI	Rhat	ESS
	Intercept	3.17	3.18	3.18	0.21	[2.76, 3.59]	1.00	671
	20–30 group	0.00	0.00	0.00	0.02	[-0.04, 0.03]	1.00	3295
	$\mathbf{Sex}$	0.07	0.08	0.08	0.02	[0.04,  0.11]	1.00	2520
Place	Respondents home	0.58	0.55	0.55	0.14	[0.27,  0.81]	1.00	880
Place	Other home	0.56	0.60	0.61	0.27	[0.10,  1.17]	1.00	1821
Place	Public sector	0.71	0.69	0.69	0.16	[0.39,  0.98]	1.00	991
Place	Government hospital	0.57	0.60	0.59	0.14	[0.31,  0.85]	1.00	864
Place	CS Govt health professional	0.59	0.58	0.58	0.14	[0.31,  0.84]	1.00	870
Place	Other public sector	0.62	0.64	0.63	0.16	[0.32,  0.94]	1.00	1024
Place	Private hospital/clinic	0.43	0.42	0.41	0.14	[0.13,  0.68]	1.00	886
Place	CS private health facility	0.70	0.64	0.63	0.18	[0.28,  0.96]	1.00	1112
Size	Very large	0.69	0.67	0.67	0.05	[0.57,  0.77]	1.00	1762
Size	Larger than average	0.70	0.71	0.71	0.05	[0.61,  0.80]	1.00	1586
Size	Average	0.76	0.77	0.77	0.04	[0.68,  0.84]	1.00	1360
Size	Smaller than average	0.52	0.54	0.54	0.05	[0.44,  0.63]	1.00	1516
Size	Very small	0.08	0.07	0.07	0.05	[-0.02, 0.17]	1.00	1700
	Order	-0.01	-0.01	-0.01	0.01	[-0.02, 0.00]	1.00	3336
	k	1.71	1.71	1.71	0.02	[1.66, 1.75]	1.00	2485
	$\phi$	7.22	9.51	11.58	8.24	[3.00, 32.40]	1.00	997
	$RMST_{12-19qroup}$	26.85	27.13	27.16	3.36	[20.60, 33.57]		
	$RMST_{20-30group}$	26.75	27.13	27.14	3.36	[20.47, 33.55]		
	$\mathrm{RMST}_{diff}$	0.07	-0.02	-0.02	0.28	[-0.57, 0.53]		

Table 12: Results of log-logistic frailty model (12–19 group vs 20–30 group)

### A.9.3 Estimation results (12–19 group vs 31+ group)

	Parameter	Mode	Median	Mean	SE	95%CI	Rhat	ESS
	Intercept	4.46	4.45	4.47	0.34	[3.86, 5.18]	1.00	908
31+ group		-0.37	-0.37	-0.37	0.12	[-0.60, -0.13]	1.00	4149
	$\mathbf{Sex}$	0.09	0.10	0.10	0.04	[0.01,  0.19]	1.00	4255
Place	Respondents home	0.89	0.84	0.83	0.33	[0.15,  1.45]	1.00	825
Place	Other home	0.75	0.60	0.61	0.61	[-0.48,  1.87]	1.00	1817
Place	Public sector	1.31	1.39	1.39	0.37	[0.63,  2.09]	1.00	970
Place	Government hospital	1.02	0.97	0.96	0.33	[0.27,  1.58]	1.00	801
Place	CS Govt health professional	1.05	1.01	1.00	0.33	[0.32,  1.61]	1.01	790
Place	Other public sector	1.26	1.30	1.29	0.40	[0.48,  2.05]	1.00	1003
Place	Private hospital/clinic	0.61	0.58	0.57	0.33	[-0.11, 1.18]	1.01	800
Place	CS private health facility	1.34	1.49	1.50	0.50	[0.51,  2.50]	1.00	1414
Size	Very large	0.82	0.83	0.83	0.12	[0.59,  1.06]	1.00	1750
Size	Larger than average	1.09	1.05	1.05	0.12	[0.82,  1.29]	1.00	1635
Size	Average	1.19	1.17	1.17	0.10	[0.98,  1.36]	1.00	1406
Size	Smaller than average	0.92	0.92	0.91	0.12	[0.68,  1.15]	1.00	1721
Size	Very small	0.10	0.07	0.07	0.13	[-0.18, 0.32]	1.00	1846
	Order	-0.01	-0.01	-0.01	0.01	[-0.04, 0.02]	1.00	4436
	$RMST_{12-19group}$	37.98	37.90	37.71	3.26	[30.91, 43.56]		
	$RMST_{31+group}$	34.19	33.73	33.60	4.13	[25.25, 41.10]		
	$\mathrm{RMST}_{diff}$	-3.90	-4.05	-4.11	1.58	[-7.38, -1.23]		

Table 13: Results of exponential model (12–19 group vs 31– group)

	Parameter	Mode	Median	Mean	SE	95%CI	Rhat	ESS
	Intercept	4.58	4.49	4.50	0.34	[3.85, 5.17]	1.00	1131
	31+ group	-0.38	-0.36	-0.36	0.12	[-0.59, -0.12]	1.00	5588
	$\mathbf{Sex}$	0.11	0.10	0.10	0.04	[0.01,  0.18]	1.00	5349
Place	Respondents home	0.94	0.86	0.85	0.32	[0.22, 1.45]	1.00	1057
Place	Other home	0.45	0.62	0.65	0.64	[-0.51, 1.96]	1.00	2017
Place	Public sector	1.56	1.39	1.38	0.38	[0.64, 2.09]	1.00	1260
Place	Government hospital	0.82	0.96	0.96	0.32	[0.32,  1.56]	1.00	1056
Place	CS Govt health professional	1.04	1.00	1.00	0.32	[0.36,  1.60]	1.00	1050
Place	Other public sector	1.37	1.27	1.27	0.40	[0.50,  2.06]	1.00	1390
Place	Private hospital/clinic	0.50	0.62	0.61	0.32	[-0.03, 1.20]	1.00	1044
Place	CS private health facility	1.44	1.56	1.57	0.52	[0.61, 2.63]	1.00	1807
Size	Very large	0.84	0.83	0.83	0.12	[0.59,  1.08]	1.00	2446
Size	Larger than average	1.09	1.06	1.05	0.12	[0.82,  1.28]	1.00	2195
Size	Average	1.18	1.16	1.16	0.10	[0.97,  1.34]	1.00	1851
Size	Smaller than average	0.88	0.91	0.91	0.12	[0.67, 1.14]	1.00	2265
Size	Very small	0.07	0.08	0.09	0.13	[-0.17, 0.33]	1.00	2395
	Order	0.00	0.00	0.00	0.01	[-0.03, 0.03]	1.00	5932
	$\phi$	0.15	0.17	0.19	0.08	[0.09,  0.40]	1.00	1223
	$RMST_{12-19group}$	39.08	38.28	37.97	3.32	[30.74, 43.51]		
	$RMST_{31+group}$	35.00	34.36	34.08	4.16	[25.26, 41.36]		
	RMST <sub>diff</sub>	-3.99	-3.83	-3.88	1.54	[-7.25, -1.08]		

Table 14: Results of mixed effects exponential model (12–19 group vs 31+ group)

	Parameter	Mode	Median	Mean	SE	95%CI	Rhat	ESS
	Intercept	4.45	4.49	4.51	0.35	[3.86, 5.27]	1.00	865
	31+ group	-0.37	-0.36	-0.36	0.12	[-0.58, -0.12]	1.00	3971
	Sex	0.09	0.10	0.10	0.04	[0.01,  0.19]	1.00	4744
Place	Respondents home	0.91	0.87	0.85	0.32	[0.15, 1.44]	1.01	752
Place	Other home	0.46	0.60	0.63	0.62	[-0.49, 1.94]	1.00	1789
Place	Public sector	1.37	1.40	1.39	0.37	[0.59, 2.09]	1.00	933
Place	Government hospital	0.99	0.98	0.96	0.32	[0.25,  1.56]	1.01	749
Place	CS Govt health professional	1.13	1.02	1.00	0.32	[0.29,  1.59]	1.01	751
Place	Other public sector	1.36	1.29	1.27	0.40	[0.45, 2.04]	1.00	976
Place	Private hospital/clinic	0.66	0.63	0.61	0.33	[-0.11, 1.20]	1.01	758
Place	CS private health facility	1.52	1.56	1.57	0.51	[0.61, 2.62]	1.00	1308
Size	Very large	0.85	0.84	0.83	0.12	[0.59,  1.08]	1.00	1921
Size	Larger than average	1.09	1.06	1.06	0.12	[0.81, 1.29]	1.00	1813
Size	Average	1.14	1.17	1.17	0.10	[0.97,  1.36]	1.00	1428
Size	Smaller than average	0.89	0.92	0.92	0.12	[0.68, 1.15]	1.00	1685
Size	Very small	0.07	0.09	0.09	0.13	[-0.16, 0.34]	1.00	1881
	Order	0.00	0.00	0.00	0.01	[-0.03, 0.03]	1.00	4536
	$\phi$	0.02	0.04	0.06	0.06	[0.01, 0.21]	1.00	1235
	$RMST_{12-19qroup}$	37.92	38.27	38.11	3.28	[30.90, 44.08]		
	$RMST_{31+group}$	34.31	34.37	34.27	4.12	[25.71, 42.05]		
	RMST <sub>diff</sub>	-4.11	-3.81	-3.84	1.52	[-7.06, -1.08]		

Table 15: Results of exponential frailty model (12–19 group vs 31+ group)

	Parameter	Mode	Median	Mean	SE	95%CI	Rhat	ESS
	Intercept	4.01	4.01	4.02	0.19	[3.68, 4.42]	1.00	1363
	31+ group	-0.22	-0.22	-0.22	0.07	[-0.35, -0.08]	1.00	2631
	$\mathbf{Sex}$	0.05	0.05	0.05	0.02	[0.00,  0.10]	1.00	2900
Place	Respondents home	0.44	0.47	0.47	0.18	[0.09,  0.79]	1.00	1341
Place	Other home	0.35	0.33	0.34	0.34	[-0.28, 1.04]	1.00	2649
Place	Public sector	0.75	0.77	0.77	0.21	[0.34,  1.15]	1.01	1898
Place	Government hospital	0.56	0.52	0.52	0.18	[0.15,  0.85]	1.00	1310
Place	CS Govt health professional	0.55	0.55	0.54	0.18	[0.17,  0.86]	1.00	1295
Place	Other public sector	0.70	0.70	0.70	0.22	[0.26, 1.11]	1.00	1774
Place	Private hospital/clinic	0.33	0.31	0.30	0.18	[-0.08, 0.63]	1.01	1378
Place	CS private health facility	0.76	0.83	0.83	0.28	[0.29,  1.38]	1.00	2328
Size	Very large	0.43	0.44	0.44	0.07	[0.30,  0.57]	1.00	2547
Size	Larger than average	0.56	0.56	0.56	0.07	[0.43,  0.69]	1.00	2530
Size	Average	0.63	0.63	0.63	0.05	[0.52,  0.74]	1.00	2229
Size	Smaller than average	0.49	0.48	0.48	0.07	[0.35,  0.61]	1.00	2266
Size	Very small	0.05	0.02	0.02	0.07	[-0.12, 0.15]	1.00	2751
	Order	-0.01	-0.01	-0.01	0.01	[-0.02, 0.01]	1.00	2858
	k	1.82	1.83	1.83	0.04	[1.76,  1.90]	1.00	2280
	$RMST_{12-19group}$	38.80	38.41	38.26	2.98	[32.08, 43.72]		
	$RMST_{31+group}$	34.90	34.44	34.31	3.72	[27.19, 41.21]		
	$\mathrm{RMST}_{diff}$	-3.68	-3.92	-3.96	1.40	[-6.74, -1.36]		

Table 16: Results of Weibull model (12–19 group vs 31+ group)

	Parameter	Mode	Median	Mean	SE	95%CI	Rhat	ESS
	Intercept	4.09	4.03	4.04	0.19	[3.69, 4.43]	1.00	1059
	31+ group	-0.22	-0.21	-0.21	0.07	[-0.34, -0.08]	1.00	4577
	$\mathbf{Sex}$	0.05	0.05	0.05	0.02	[0.00,  0.10]	1.00	4768
Place	Respondents home	0.48	0.49	0.48	0.18	[0.10,  0.81]	1.00	950
Place	Other home	0.38	0.34	0.35	0.34	[-0.25, 1.05]	1.00	2002
Place	Public sector	0.83	0.77	0.76	0.21	[0.34, 1.14]	1.00	1142
Place	Government hospital	0.53	0.53	0.52	0.18	[0.14,  0.85]	1.00	956
Place	CS Govt health professional	0.59	0.55	0.54	0.18	[0.16,  0.86]	1.00	940
Place	Other public sector	0.63	0.68	0.68	0.22	[0.24, 1.09]	1.00	1304
Place	Private hospital/clinic	0.29	0.33	0.32	0.18	[-0.06, 0.65]	1.00	961
Place	CS private health facility	0.86	0.86	0.87	0.28	[0.33, 1.43]	1.00	1719
Size	Very large	0.44	0.44	0.44	0.07	[0.30,  0.58]	1.00	1917
Size	Larger than average	0.56	0.56	0.56	0.07	[0.43,  0.69]	1.00	1870
Size	Average	0.62	0.63	0.63	0.06	[0.52,  0.73]	1.00	1552
Size	Smaller than average	0.50	0.48	0.48	0.07	[0.35, 0.62]	1.00	1855
Size	Very small	0.02	0.02	0.02	0.07	[-0.11, 0.16]	1.00	2102
	Order	0.00	0.00	0.00	0.01	[-0.02, 0.01]	1.00	5911
	k	1.82	1.82	1.83	0.03	[1.76, 1.89]	1.00	3812
	$\phi$	0.08	0.09	0.10	0.04	[0.05, 0.21]	1.00	1389
	$RMST_{12-19qroup}$	39.57	38.76	38.59	3.00	[32.22, 43.90]		
	$RMST_{31+group}$	35.80	34.97	34.88	3.75	[27.03, 41.86]		
	$\mathrm{RMST}_{diff}$	-3.88	-3.66	-3.71	1.40	[-6.55, -1.18]		

Table 17: Results of mixed effects Weibull model (12–19 group vs 31+ group)

	Parameter	Mode	Median	Mean	SE	95%CI	Rhat	ESS
	Intercept	4.41	4.44	4.45	0.29	[3.92, 5.04]	1.00	980
	31– group	-0.34	-0.33	-0.33	0.09	[-0.51, -0.15]	1.00	3741
	$\mathbf{Sex}$	0.07	0.07	0.07	0.03	[0.00,  0.13]	1.00	4148
Place	Respondents home	0.59	0.65	0.64	0.28	[0.08, 1.18]	1.00	978
Place	Other home	0.19	0.33	0.34	0.49	[-0.59, 1.32]	1.00	1937
Place	Public sector	1.07	1.04	1.04	0.31	[0.42, 1.64]	1.00	1086
Place	Government hospital	0.65	0.71	0.71	0.28	[0.13, 1.23]	1.00	984
Place	CS Govt health professional	0.81	0.75	0.75	0.28	[0.18, 1.27]	1.00	973
Place	Other public sector	0.97	0.90	0.90	0.32	[0.27, 1.49]	1.00	1154
Place	Private hospital/clinic	0.41	0.43	0.42	0.28	[-0.16, 0.95]	1.00	1001
Place	CS private health facility	1.01	0.94	0.94	0.37	[0.22, 1.65]	1.00	1415
Size	Very large	0.58	0.57	0.57	0.10	[0.37,  0.76]	1.00	1615
Size	Larger than average	0.73	0.76	0.76	0.10	[0.56, 0.95]	1.00	1598
Size	Average	0.87	0.84	0.84	0.09	[0.67,  1.01]	1.00	1409
Size	Smaller than average	0.69	0.66	0.66	0.10	[0.46,  0.85]	1.00	1596
Size	Very small	-0.02	0.00	0.00	0.11	[-0.21, 0.21]	1.00	1748
	Order	-0.01	-0.01	-0.01	0.01	[-0.03, 0.01]	1.00	4046
	$\sigma$	1.42	1.43	1.43	0.02	[1.38, 1.48]	1.00	2188
	$RMST_{12-19group}$	41.05	40.67	40.58	2.40	[35.60, 44.96]		
	$RMST_{31+group}$	37.32	37.64	37.57	2.91	[31.65, 43.06]		
	$\mathrm{RMST}_{diff}$	-2.81	-2.98	-3.01	0.97	[-4.98, -1.26]		

Table 18: Results of log-normal model (12–19 group vs 31+ group)

	Parameter	Mode	Median	Mean	SE	95%CI	Rhat	ESS
	Intercept	4.44	4.47	4.47	0.30	[3.90, 5.09]	1.00	1041
	31+ group	-0.31	-0.32	-0.32	0.09	[-0.49, -0.15]	1.00	4358
	Sex	0.07	0.07	0.07	0.03	[0.00,  0.13]	1.00	4494
Place	Respondents home	0.76	0.66	0.66	0.29	[0.06, 1.21]	1.00	978
Place	Other home	0.27	0.36	0.37	0.48	[-0.53, 1.32]	1.00	1830
Place	Public sector	1.00	1.03	1.03	0.32	[0.40,  1.64]	1.00	1038
Place	Government hospital	0.75	0.71	0.70	0.29	[0.11,  1.26]	1.00	985
Place	CS Govt health professional	0.84	0.75	0.75	0.29	[0.16,  1.29]	1.00	973
Place	Other public sector	0.87	0.86	0.86	0.33	[0.19,  1.49]	1.00	1160
Place	Private hospital/clinic	0.50	0.45	0.45	0.29	[-0.14, 1.00]	1.00	979
Place	CS private health facility	1.16	1.00	1.00	0.38	[0.28,  1.76]	1.00	1430
Size	Very large	0.55	0.57	0.57	0.10	[0.36,  0.76]	1.00	1879
Size	Larger than average	0.76	0.76	0.76	0.10	[0.56,  0.95]	1.00	1772
Size	Average	0.85	0.84	0.84	0.09	[0.67,  1.00]	1.00	1569
Size	Smaller than average	0.66	0.67	0.67	0.10	[0.47,  0.86]	1.00	1840
Size	Very small	0.03	0.00	0.00	0.11	[-0.20, 0.21]	1.00	2041
	Order	0.00	0.00	0.00	0.01	[-0.02, 0.02]	1.00	4585
	$\sigma$	1.44	1.43	1.43	0.02	[1.38, 1.48]	1.00	3286
	$\phi$	0.13	0.12	0.13	0.05	[0.06, 0.27]	1.00	1172
	$RMST_{12-19group}$	40.69	40.90	40.76	2.50	[35.51, 45.27]		
	$RMST_{31+group}$	39.25	38.00	37.89	3.02	[31.78, 43.53]		
	RMST <sub>diff</sub>	-2.66	-2.83	-2.87	0.95	[-4.81, -1.17]		

Table 19: Results of mixed effects log-normal model (12–19 group vs 31+ group)

 $95\%\mathrm{CI}:\,95\%$  credible interval, ESS: Effective sample size

	Parameter	Mode	Median	Mean	SE	95%CI	Rhat	ESS
	Intercept	12.37	12.49	12.53	0.90	[10.81, 14.43]	1.00	713
	31+ group	-0.25	-0.25	-0.25	0.07	[-0.38, -0.10]	1.00	3585
	$\mathbf{Sex}$	0.05	0.06	0.06	0.03	[0.00,  0.11]	1.00	3723
Place	Respondents home	0.59	0.55	0.54	0.20	[0.11,  0.91]	1.00	959
Place	Other home	0.32	0.37	0.39	0.39	[-0.33, 1.19]	1.00	2026
Place	Public sector	0.89	0.87	0.86	0.23	[0.40,  1.30]	1.00	1136
Place	Government hospital	0.55	0.59	0.58	0.20	[0.16,  0.96]	1.00	966
Place	CS Govt health professional	0.63	0.62	0.61	0.20	[0.19,  0.98]	1.00	964
Place	Other public sector	0.81	0.76	0.75	0.25	[0.25, 1.24]	1.00	1244
Place	Private hospital/clinic	0.37	0.37	0.37	0.20	[-0.07, 0.73]	1.00	975
Place	CS private health facility	0.94	0.91	0.92	0.30	[0.33, 1.53]	1.00	1467
Size	Very large	0.49	0.49	0.48	0.08	[0.33,  0.63]	1.00	1863
Size	Larger than average	0.64	0.62	0.62	0.08	[0.47, 0.77]	1.00	1858
Size	Average	0.68	0.69	0.69	0.06	[0.56, 0.82]	1.00	1645
Size	Smaller than average	0.56	0.54	0.54	0.08	[0.39, 0.69]	1.00	1807
Size	Very small	0.03	0.02	0.02	0.08	[-0.14, 0.18]	1.00	1745
	Order	0.00	0.00	0.00	0.01	[-0.02, 0.02]	1.00	3913
k		2.50	2.50	2.51	0.10	[2.31, 2.72]	1.00	835
σ		1187	2298	3696	4759	[433, 15736]	1.00	890
	$RMST_{12-19group}$	50.00	50.00	50.00	0.00	[50.00, 50.00]		
$RMST_{31+group}$		50.00	50.00	50.00	0.00	[50.00, 50.00]		
	$\mathrm{RMST}_{diff}$	0.00	0.00	0.00	0.00	[0.00, 0.00]		

Table 20: Results of log-normal frailty model (12–19 group vs 31+ group)

The estimated results of the log-normal frailty model differ from the others; therefore, we examined the behavior using different data in Appendix A.10.

	Parameter	Mode	Median	Mean	SE	95%CI	Rhat	ESS
	Intercept	4.05	4.03	4.04	0.21	[3.65, 4.48]	1.00	931
	31+ group	-0.22	-0.23	-0.23	0.07	[-0.37, -0.10]	1.00	4101
	$\mathbf{Sex}$	0.05	0.05	0.05	0.03	[0.00,  0.10]	1.00	3482
Place	Respondents home	0.48	0.52	0.51	0.20	[0.09,  0.88]	1.00	939
Place	Other home	0.33	0.36	0.37	0.37	[-0.34, 1.10]	1.00	1873
Place	Public sector	0.88	0.82	0.82	0.22	[0.35, 1.24]	1.00	1096
Place	Government hospital	0.66	0.57	0.57	0.20	[0.14,  0.94]	1.00	925
Place	CS Govt health professional	0.58	0.60	0.59	0.20	[0.17,  0.95]	1.00	921
Place	Other public sector	0.77	0.75	0.74	0.24	[0.26, 1.22]	1.00	1169
Place	Private hospital/clinic	0.29	0.35	0.34	0.20	[-0.09, 0.71]	1.00	926
Place	CS private health facility	0.85	0.87	0.87	0.30	[0.28, 1.49]	1.00	1528
Size	Very large	0.43	0.45	0.45	0.07	[0.30,  0.59]	1.00	1729
Size	Larger than average	0.59	0.58	0.58	0.07	[0.44,  0.72]	1.00	1620
Size	Average	0.63	0.64	0.65	0.06	[0.53,  0.77]	1.00	1446
Size	Smaller than average	0.49	0.50	0.50	0.07	[0.35,  0.64]	1.00	1616
Size	Very small	0.02	0.00	0.00	0.08	[-0.15, 0.16]	1.00	1756
	Order	-0.01	-0.01	-0.01	0.01	[-0.02, 0.01]	1.00	4171
k		1.86	1.86	1.86	0.04	[1.79,  1.93]	1.00	2427
$RMST_{12-19group}$		40.93	40.40	40.26	2.56	[34.88, 45.00]		
$RMST_{31+qroup}$		37.31	37.22	37.09	3.13	[30.89, 43.06]		
	$\mathrm{RMST}_{diff}$	-3.43	-3.15	-3.17	1.08	[-5.37, -1.18]		

Table 21: Results of log-logistic model (12–19 group vs 31+ group)

	Parameter	Mode	Median	Mean	SE	95%CI	Rhat	ESS
	Intercept	4.00	4.05	4.06	0.21	[3.66, 4.49]	1.00	1133
	31+ group	-0.24	-0.23	-0.22	0.07	[-0.36, -0.08]	1.00	4336
	$\mathbf{Sex}$	0.05	0.05	0.05	0.03	[0.00, 0.10]	1.00	4488
Place	Respondents home	0.55	0.54	0.53	0.20	[0.12, 0.89]	1.00	1042
Place	Other home	0.29	0.37	0.39	0.36	[-0.31, 1.11]	1.00	1932
Place	Public sector	0.83	0.83	0.83	0.23	[0.37, 1.25]	1.00	1204
Place	Government hospital	0.64	0.58	0.57	0.20	[0.16,  0.94]	1.00	1035
Place	CS Govt health professional	0.61	0.60	0.59	0.20	[0.18, 0.96]	1.00	1036
Place	Other public sector	0.73	0.73	0.73	0.24	[0.24, 1.19]	1.00	1264
Place	Private hospital/clinic	0.33	0.37	0.36	0.20	[-0.05, 0.74]	1.00	1062
Place	CS private health facility	0.84	0.90	0.90	0.29	[0.33, 1.49]	1.00	1789
Size	Very large	0.45	0.45	0.45	0.07	[0.30,  0.59]	1.00	1698
Size	Larger than average	0.58	0.58	0.58	0.07	[0.43, 0.72]	1.00	1601
Size	Average	0.63	0.64	0.64	0.06	[0.52, 0.76]	1.00	1400
Size	Smaller than average	0.51	0.50	0.50	0.07	[0.36, 0.64]	1.00	1621
Size	Very small	0.00	0.01	0.01	0.08	[-0.15, 0.15]	1.00	1751
	Order	0.00	0.00	0.00	0.01	[-0.02, 0.01]	1.00	5308
k		1.87	1.86	1.86	0.04	[1.79, 1.93]	1.00	3258
$\phi$		0.08	0.10	0.11	0.05	[0.05, 0.22]	1.00	1256
$RMST_{12-19aroup}$		40.13	40.58	40.44	2.57	[35.09, 45.01]		
$RMST_{31+aroup}$		36.17	37.56	37.43	3.13	[31.18, 43.19]		
	$\mathrm{RMST}_{diff}$	-2.64	-2.96	-3.01	1.10	[-5.25, -0.97]		

Table 22: Results of mixed effects log-logistic model (12–19 group vs 31+ group)

	Parameter	Mode	Median	Mean	SE	95%CI	Rhat	ESS
	Intercept	3.55	3.60	3.60	0.26	[3.10, 4.10]	1.00	1260
	31+ group	-0.25	-0.22	-0.22	0.07	[-0.35, -0.08]	1.00	4729
	$\mathbf{Sex}$	0.05	0.05	0.05	0.03	[0.00,  0.10]	1.00	4304
Place	Respondents home	0.51	0.52	0.51	0.19	[0.11,  0.85]	1.00	1273
Place	Other home	0.45	0.37	0.38	0.36	[-0.28, 1.11]	1.00	2561
Place	Public sector	0.80	0.81	0.80	0.22	[0.36,  1.19]	1.00	1467
Place	Government hospital	0.54	0.56	0.55	0.19	[0.15,  0.89]	1.00	1282
Place	CS Govt health professional	0.62	0.58	0.57	0.19	[0.17,  0.91]	1.00	1267
Place	Other public sector	0.70	0.71	0.71	0.23	[0.24, 1.14]	1.00	1558
Place	Private hospital/clinic	0.31	0.36	0.35	0.19	[-0.06, 0.69]	1.00	1258
Place	CS private health facility	0.81	0.90	0.90	0.29	[0.35, 1.46]	1.00	1746
Size	Very large	0.42	0.44	0.44	0.07	[0.31,  0.58]	1.00	1802
Size	Larger than average	0.60	0.57	0.57	0.07	[0.43,  0.71]	1.00	1761
Size	Average	0.63	0.64	0.64	0.06	[0.52,  0.75]	1.00	1537
Size	Smaller than average	0.48	0.50	0.50	0.07	[0.36,  0.63]	1.00	1847
Size	Very small	0.03	0.02	0.02	0.07	[-0.13, 0.16]	1.00	1914
	Order	0.00	0.00	0.00	0.01	[-0.02, 0.01]	1.00	5181
k		1.82	1.83	1.83	0.04	[1.76, 1.90]	1.00	3725
$\phi$		1.16	2.15	2.90	2.74	[0.20, 10.01]	1.00	1302
$RMST_{12-19aroup}$		33.30	34.02	33.86	3.98	[25.70, 41.10]		
$RMST_{31+aroup}$		31.44	30.50	30.40	4.27	[21.91, 38.46]		
	$\mathrm{RMST}_{diff}$	-3.79	-3.46	-3.46	1.13	[-5.65, -1.16]		

Table 23: Results of log-logistic frailty model (12–19 group vs 31+ group)

#### A.9.4 Values of widely applicable information criterion

We presented the calculated WAICs in Tables 24 and 25. For the comparison between the "12-19 group" and the "20-30 group," the mixed effects Weibull model yielded the minimum WAIC. For the comparison between the "12-19 group" and the "31+ group," the Weibull frailty model had the minimum WAIC. The WAIC for the log-logistic frailty model was larger than that for the other models. We further investigated the WAIC using different data in Appendix A.10.

	WAIC
exponential model	70797.8912255
mixed effects exponential model	70729.22940
exponential frailty model	70728.64805
Weibull model	69233.90573
mixed effects Weibull model	69160.98410
Weibull frailty model	69162.08765
log-normal model	70373.81205
mixed effects log-normal model	70302.03889
log-normal frailty model	69379.64799
log-logistic model	69349.66898
mixed effects log-logistic model	69277.20932
log-logistic frailty model	1006049.864

Table 24: Results of WAIC (12–19 group vs 20-30 group)

Table 25: Results of WAIC (12–19 group vs 31+ group)

	WAIC
exponential model	29204.82671
mixed effects exponential model	29169.61918
exponential frailty model	29169.12216
Weibull model	28398.84546
mixed effects Weibull model	28363.75397
Weibull frailty model	28363.7075
log-normal model	28902.55726
mixed effects log-normal model	28862.84461
log-normal frailty model	28512.54361
log-logistic model	28448.37898
mixed effects log-logistic model	28412.54262
log-logistic frailty model	64523.48299

# A.10 Sensitivity analysis for log-normal frailty model

We conducted a sensitivity analysis of the log-normal frailty model using the LeukSurv dataset from the spBayesSurv package in R. We compared the 'Greater than or equal to 65 years old' group with the 'Less than 65 years old' group. The covariates included sex, white blood cell count at diagnosis (Wbc), and the Townsend score, where higher values indicate less affluent areas (Tpi). The clusters were the twenty-four administrative districts of residence. The sample size was 1043 (540 in the 'Greater than or equal to 65 years old' group and 503 in the 'Less than 65 years old' group). The distribution of the number of subjects in each group is as follows:

10010 201	Pipp	00101		0110	mann	001 0	1 Duo	Jeece	111 0	aon (	1100111	
Distinct ID	1	2	3	4	5	6	7	8	9	10	11	12
# subjects	35	69	44	15	46	12	71	25	34	12	17	27
Distinct ID	13	14	15	16	17	18	19	20	21	22	23	24
# subjects	18	58	37	52	84	45	61	53	63	27	36	102

Table 26: Disposition of the number of subjects in each distinct

The Kaplan-Meier plot comparing the 'Greater than or equal to 65 years old' group with the 'Less than 65 years old' group is illustrated below.



Strata + factor(arm)=Greater than or equal to 65 years old + factor(arm)=Less than 65 years old

Figure 4: Kaplan Meier plot of LeukSurv ('Greater than or equal to 65 years old' group vs 'Less than 65 years old' group)

The estimation results are shown in Tables 27,28, and 29. The log-normal frailty model is consistent with the other estimated results.

Parameter	Mode	Median	Mean	SE	95%CI
Intercept	4.58	4.60	4.60	0.12	[4.37, 4.84]
Less than 65 years old	1.69	1.69	1.69	0.13	[1.45, 1.93]
Sex	-0.01	-0.03	-0.04	0.13	[-0.30, 0.21]
Wbc	-0.01	-0.01	-0.01	0.00	[-0.01, -0.01]
Tpi	-0.06	-0.06	-0.06	0.02	[-0.10, -0.03]
σ	2.05	2.05	2.05	0.05	[1.95, 2.15]

Table 27: Results of log-normal model

 $95\%\mathrm{CI}:\,95\%$  credible interval

Table 28: Results of mixed effects log-normal model

Parameter	Mode	Median	Mean	SE	95%CI
Intercept	4.61	4.60	4.60	0.12	[4.36, 4.84]
Less than 65 years old	1.69	1.70	1.70	0.13	[1.44, 1.96]
Sex	-0.05	-0.04	-0.03	0.13	[-0.28, 0.22]
Wbc	-0.01	-0.01	-0.01	0.00	[-0.01, -0.01]
Tpi	-0.07	-0.06	-0.06	0.02	[-0.10, -0.03]
σ	2.05	2.04	2.04	0.05	[1.94, 2.14]
$\phi$	0.14	0.16	0.16	0.10	[0.01, 0.37]

95%CI: 95% credible interval

Table 29: Results of log-normal frailty model

Parameter	Mode	Median	Mean	SE	95%CI	
Intercept	4.56	4.60	4.60	0.14	[4.33, 4.87]	
less than 65 years old	1.71	1.71	1.71	0.13	[1.46, 1.96]	
Sex	0.02	-0.03	-0.03	0.13	[-0.28, 0.21]	05% CI. 05% and this internal
Wbc	-0.01	-0.01	-0.01	0.00	[-0.01, -0.01]	95%CI: 95% credible interval
$\mathrm{Tpi}$	-0.06	-0.07	-0.07	0.02	[-0.10, -0.03]	
$\sigma$	2.02	2.03	2.03	0.06	[1.92, 2.14]	
$\phi$	0.02	0.03	0.03	0.02	[0.01, 0.09]	
	$\begin{array}{c} \hline Parameter \\ \hline Intercept \\ ess than 65 years old \\ Sex \\ Wbc \\ Tpi \\ \sigma \\ \phi \\ \end{array}$	$\begin{tabular}{ c c c c } \hline Parameter & Mode \\ \hline Intercept & 4.56 \\ \hline less than 65 years old & 1.71 \\ \hline Sex & 0.02 \\ \hline Wbc & -0.01 \\ \hline Tpi & -0.06 \\ \hline \sigma & 2.02 \\ \hline \phi & 0.02 \\ \hline \end{tabular}$	$\begin{array}{c c c c c c c c c c c c c c c c c c c $	$\begin{array}{c c c c c c c c c c c c c c c c c c c $	$\begin{array}{c c c c c c c c c c c c c c c c c c c $	$\begin{array}{c c c c c c c c c c c c c c c c c c c $

We displayed the WAICs in Table 30 and confirmed that the WAIC of the log-logistic frailty model is almost the same as the other WAICs.

Table 30: Results of WA	IC
	WAIC
log-normal model	11983.72
mixed effects log-normal model	11983.72
log-normal frailty model	11977.70
log-logistic frailty model	12068.60

Table 3	30:	Results	of	WAIC