

ABSOLUTE DILATIONS OF UCP SELF-ADJOINT FOURIER MULTIPLIERS: THE NON UNIMODULAR CASE

CHARLES DUQUET AND CHRISTIAN LE MERDY

ABSTRACT. Let φ be a normal semi-finite faithful weight on a von Neumann algebra A , let $(\sigma_r^\varphi)_{r \in \mathbb{R}}$ denote the modular automorphism group of φ , and let $T: A \rightarrow A$ be a linear map. We say that T admits an absolute dilation if there exist another von Neumann algebra M equipped with a normal semi-finite faithful weight ψ , a w^* -continuous, unital and weight-preserving $*$ -homomorphism $J: A \rightarrow M$ such that $\sigma^\psi \circ J = J \circ \sigma^\varphi$, as well as a weight-preserving $*$ -automorphism $U: M \rightarrow M$ such that $T^k = \mathbb{E}_J U^k J$ for all integer $k \geq 0$, where $\mathbb{E}_J: M \rightarrow A$ is the conditional expectation associated with J . Given any locally compact group G and any real valued function $u \in C_b(G)$, we prove that if u induces a unital completely positive Fourier multiplier $M_u: VN(G) \rightarrow VN(G)$, then M_u admits an absolute dilation. Here $VN(G)$ is equipped with its Plancherel weight φ_G . This result had been settled by the first named author in the case when G is unimodular so the salient point in this paper is that G may be non unimodular, and hence φ_G may not be a trace. The absolute dilation of M_u implies that for any $1 < p < \infty$, the L^p -realization of M_u can be dilated into an isometry acting on a non-commutative L^p -space. We further prove that if u is valued in $[0, 1]$, then the L^p -realization of M_u is a Ritt operator with a bounded H^∞ -functional calculus.

2000 Mathematics Subject Classification : 47A20, 46L51, 43A22.

Key words: Fourier multipliers. Dilations. Non-commutative L^p -spaces.

1. INTRODUCTION

Let (Ω, μ) be any measure space, let $1 < p < \infty$ and let $T: L^p(\Omega) \rightarrow L^p(\Omega)$ be a positive contraction. Akcoglu's dilation theorem [1, 2] asserts that T admits an isometric dilation, that is, there exist a second measure space (Ω', μ') , an isometry $J: L^p(\Omega) \rightarrow L^p(\Omega')$, an isometry $U: L^p(\Omega') \rightarrow L^p(\Omega')$ and a contraction $Q: L^p(\Omega') \rightarrow L^p(\Omega)$ such that $T^k = QU^k J$ for all integer $k \geq 0$. This L^p -analogue of Nagy's dilation theorem has been instrumental in various topics concerning L^p -operators, such as functional calculus, harmonic analysis and ergodic theory. We refer to [8, 28, 34, 36] and the references therein for specific results.

Turning to non-commutative analysis, we note that Akcoglu's theorem does not extend to non-commutative L^p -spaces [29]. This obstacle led to new, interesting difficulties in the development of operator theory and harmonic analysis on non-commutative L^p -spaces. Also, it became important to exhibit classes of completely positive contractions on non-commutative L^p -spaces which do admit a dilation into an isometry acting on a second non-commutative

Date: June 11, 2024.

L^p -space. We refer to [4, 5, 6, 16, 17, 27, 30] for various results of this type. In all these papers (except [27]), the construction of an isometric dilation of a map $T: L^p(A) \rightarrow L^p(A)$ relies on the existence of a trace (or weight) preserving dilation of T on the underlying von Neumann algebra A . This leads us to introduce the following definition, which extends [17, Definition 2.1]. Let φ be a normal semi-finite faithful weight on A (we say that (A, φ) is a weighted von Neumann algebra in this case). Let $(\sigma_r^\varphi)_{r \in \mathbb{R}}$ denote the modular automorphism group of φ . We say that a linear map $T: A \rightarrow A$ admits an absolute dilation if there exist another weighted von Neumann algebra (M, ψ) , a w^* -continuous, unital and weight-preserving $*$ -homomorphism $J: A \rightarrow M$ such that $\sigma^\psi \circ J = J \circ \sigma^\varphi$, as well as a weight-preserving $*$ -automorphism $U: M \rightarrow M$ such that $T^k = \mathbb{E}_J U^k J$ for all integer $k \geq 0$, where $\mathbb{E}_J: M \rightarrow A$ is the conditional expectation associated with J . This dilation property implies that T is unital, completely positive and weight-preserving, that T has a natural extension to $L^p(A)$ and that the latter admits a dilation into an isometry acting on another non-commutative L^p -space (see Proposition 2.7).

In [17], we gave a characterization of bounded Schur multipliers which admit an absolute dilation. In [16], the first named author showed that for any unimodular locally compact group G , every unital completely positive Fourier multiplier $VN(G) \rightarrow VN(G)$ (in the sense of [14]) whose symbol is real-valued admits an absolute dilation. The case when G is discrete had been settled in [4]. (The discrete case also follows from [38, Corollary] and [22, Theorem 4.4]). The objective of this paper is to extend the above mentioned result to the non unimodular case. Let G be any locally compact group, let $u: G \rightarrow \mathbb{R}$ be a continuous function and assume that u induces a unital completely positive Fourier multiplier $M_u: VN(G) \rightarrow VN(G)$. We show that M_u admits an absolute dilation, see Theorem 4.1.

In the previous paragraph, the group von Neumann algebra $A = VN(G)$ is equipped with its Plancherel weight $\varphi = \varphi_G$ (see Section 3 for background). This specific weight is a trace if and only if G is unimodular. This is the reason why passing from the unimodular case (considered in [16]) to the non unimodular case (considered in the present paper) requires new arguments. When φ_G is a non tracial weight, we need to consider the Haagerup non-commutative L^p -spaces associated with $(VN(G), \varphi_G)$ (see Subsection 2.2 for background). In particular, the space $L^1(VN(G), \varphi_G)$ plays a key role in the proof of Theorem 4.1.

We note that a dilation theorem for semigroups of unital completely positive Fourier multiplier with real symbols was established in [6]. However it cannot be applied to our case. Indeed there is no known way to pass from a dilation result for semigroups to a dilation result for single operators.

In the last part of the paper (Section 5), we give an application of our dilation result to functional calculus. We consider a unital completely positive Fourier multiplier $M_u: VN(G) \rightarrow VN(G)$ and we assume that u is valued in $[0, 1]$. Then we show that for any $1 < p < \infty$, the L^p -realization of T on $L^p(VN(G), \varphi_G)$ is a Ritt operator which admits a bounded H^∞ -functional calculus in the sense of [7, 35].

2. ABSOLUTE DILATIONS AND ISOMETRIC p -DILATIONS IN THE WEIGHTED CASE

We refer the reader to [41] for general information on von Neumann algebras. Given any von Neumann algebra A , we let A_* denote its predual. We let A^+ and A_*^+ denote the positive

cones of A and A_* , respectively. A map $V: A \rightarrow A'$ between two von Neumann algebras is called positive if $V(A^+) \subset A'^+$. We will use the stronger notion of completely positive map, for which we refer to [41, Section IV.3].

2.1. Absolute dilations. We refer to [39, 42] for general information on weights. Let A be a von Neumann algebra equipped with a normal semi-finite faithful weight φ . In this situation, we say that (A, φ) is a weighted von Neumann algebra. We let $(\sigma_r^\varphi)_{r \in \mathbb{R}}$ denote the modular automorphism group of φ . Then we set

$$\mathcal{N}_\varphi = \{x \in A : \varphi(x^*x) < \infty\}$$

and

$$\mathcal{M}_\varphi = \text{Span}\{y^*x : x, y \in \mathcal{N}_\varphi\}.$$

We recall that \mathcal{N}_φ is a w^* -dense left ideal of A and that \mathcal{M}_φ is a w^* -dense $*$ -subalgebra of A . Further $\mathcal{M}_\varphi = \text{Span}\{x^*x : x \in \mathcal{N}_\varphi\}$ and the restriction of φ to the set $\{x^*x : x \in \mathcal{N}_\varphi\}$ uniquely extends to a functional (still denoted by)

$$\varphi: \mathcal{M}_\varphi \longrightarrow \mathbb{C}.$$

Let (M, ψ) be a second weighted von Neumann algebra and let $T: A \rightarrow M$ be any positive map. We say that T is weight-preserving provided that

$$\psi \circ T = \varphi \quad \text{on } A^+.$$

Let $J: A \rightarrow M$ be a w^* -continuous, unital and weight-preserving $*$ -homomorphism such that

$$\sigma^\psi \circ J = J \circ \sigma^\varphi.$$

Here and thereafter, this identity means that $\sigma_r^\psi \circ J = J \circ \sigma_r^\varphi$ on A for all $r \in \mathbb{R}$. According to either [39, Theorem 10.1] or [42, Theorem IX.4.2], there exists a unique weight-preserving conditional expectation

$$(2.1) \quad \mathbb{E}_J: M \longrightarrow A.$$

That is, $\mathbb{E}_J: M \rightarrow A$ is the unique linear map satisfying $\mathbb{E}_J \circ J = I_A$, $\|\mathbb{E}_J\| = 1$ and $\varphi \circ \mathbb{E}_J = \mathbb{E}_J \circ \psi$. We note that we further have $\sigma^\varphi \circ \mathbb{E}_J = \mathbb{E}_J \circ \sigma^\psi$, see [39, Corollary 10.5], and that \mathbb{E}_J is w^* -continuous (see also Lemma 2.12).

Definition 2.1. *Let (A, φ) be a weighted von Neumann algebra and let $T: A \rightarrow A$ be a bounded map. We say that T admits an absolute dilation if there exist another weighted von Neumann algebra (M, ψ) , a w^* -continuous, unital and weight-preserving $*$ -homomorphism $J: A \rightarrow M$ such that $\sigma^\psi \circ J = J \circ \sigma^\varphi$, as well as a weight-preserving $*$ -automorphism $U: M \rightarrow M$ such that*

$$(2.2) \quad \forall k \geq 0, \quad T^k = \mathbb{E}_J U^k J.$$

Remark 2.2.

(1) Assume that $T: (A, \varphi) \rightarrow (A, \varphi)$ admits an absolute dilation and consider (M, ψ) , J and U as in Definition 2.1. In particular, we have $T = \mathbb{E}_J U J$. We observe that J, U and \mathbb{E}_J are both unital, completely positive and weight-preserving. Further $\sigma^\psi \circ U = U \circ \sigma^\psi$, by

[42, Corollary VIII.1.4]. Since $\sigma^\psi \circ J = J \circ \sigma^\varphi$ and $\sigma^\varphi \circ \mathbb{E}_J = \mathbb{E}_J \circ \sigma^\psi$, we deduce that T is unital, completely positive, weight-preserving and that $\sigma^\varphi \circ T = T \circ \sigma^\varphi$.

(2) Assume that φ is a state and consider $T: (A, \varphi) \rightarrow (A, \varphi)$. If T admits an absolute dilation and if (M, ψ) and $J: A \rightarrow M$ are given by Definition 2.1, then $\psi(1) = \psi(J(1)) = \varphi(1) = 1$, hence ψ is a state. It therefore follows from [22, Theorem 4.4], that T admits an absolute dilation if and only if T is factorizable in the sense of [3, 38]. Absolute dilations in the state case go back to [33]. We refer to [5] for various examples.

(3) Assume that φ is a trace and that $T: (A, \varphi) \rightarrow (A, \varphi)$ admits an absolute dilation. Then the weighted von Neumann algebra (M, ψ) from Definition 2.1 can be chosen in such a way that ψ is a trace.

Indeed, assume that T satisfies Definition 2.1 and let M_0 be the centralizer of ψ , that is, $M_0 = \{m \in M : \sigma_r^\psi(x) = x \text{ for all } r \in \mathbb{R}\}$. Let

$$\mathcal{B} = \text{Span} \left\{ \sum_{k=1}^m U^{k_1} J(x_1) U^{k_2} J(x_2) \cdots U^{k_m} J(x_m) : m \geq 1, k_1, \dots, k_m \in \mathbb{Z}, x_1, \dots, x_m \in A \right\},$$

and let $B = \overline{\mathcal{B}}^{w^*}$. Since σ^φ is trivial, $U^k J(x) \in M_0$ for all $x \in A$ and $k \in \mathbb{Z}$. Hence B is a sub-von Neumann algebra of M_0 . Therefore, the restriction ψ_B of ψ to B^+ is a normal faithful trace. Let us show its semi-finiteness. For any $x \in \mathcal{N}_\varphi$ and any $k \in \mathbb{Z}$, we have

$$\psi((U^k J(x))^*(U^k J(x))) = \psi(U^k J(x^*x)) = \varphi(x^*x) < \infty,$$

that is, $U^k J(x) \in \mathcal{N}_\psi$. Since \mathcal{N}_ψ is w^* -dense in A and J, U are w^* -continuous, $U^k J(\mathcal{N}_\varphi)$ is w^* -dense in $U^k J(A)$. Since \mathcal{N}_ψ is a left ideal, we deduce that $\mathcal{N}_\psi \cap \mathcal{B}$ is w^* -dense in \mathcal{B} . Consequently, $\mathcal{N}_\psi \cap B$ is w^* -dense in B . Thus, ψ_B is semi-finite. Consequently, (M, ψ) can be replaced by (B, ψ_B) in the dilation of T .

In the paper [17], we considered absolute dilations in the tracial case. The above paragraph shows that in the case when φ is a trace, $T: (A, \varphi) \rightarrow (A, \varphi)$ admits an absolute dilation in the sense of Definition 2.1 if and only if it admits an absolute dilation in the sense of [17].

(4) Combining [22, Section 3] with (3) above, we obtain examples of unital, completely positive, weight-preserving maps $T: (A, \varphi) \rightarrow (A, \varphi)$ such that $\sigma^\varphi \circ T = T \circ \sigma^\varphi$, without any absolute dilation (see [17, Remark 7.3] for details).

We record for further use the so-called Pedersen-Takesaki theorem (see [39, Theorem 6.2]).

Lemma 2.3. *Let (A, φ) be a weighted von Neumann algebra and let $B \subset \mathcal{N}_\varphi$ be a w^* -dense $*$ -algebra such that $\sigma_r^\varphi(B) \subset B$ for all $r \in \mathbb{R}$. Let γ be a normal semi-finite weight on A such that $\gamma \circ \sigma^\varphi = \gamma$ and $\gamma(x^*x) = \varphi(x^*x)$ for all $x \in B$. Then $\gamma = \varphi$.*

2.2. Haagerup non-commutative L^p -spaces. Let (A, φ) be a weighted von Neumann algebra. We recall the definition of the Haagerup non-commutative L^p -spaces $L^p(A, \varphi)$, as well as some of their properties. We refer the reader to [20], [23, Section 1], [26, Chapter 9], [37] and [44] for details and complements. Assume that A acts on some Hilbert space H and let

$$\mathcal{R} = A \rtimes_{\sigma^\varphi} \mathbb{R} \subset B(L^2(\mathbb{R})) \overline{\otimes} A \subset B(L^2(\mathbb{R}; H))$$

be the crossed product associated with $\sigma^\varphi: \mathbb{R} \rightarrow \text{Aut}(A)$, see e.g. [39, Chapter IV] or [42, Chapter X]. We regard A as a sub-von Neumann algebra of \mathcal{R} in the natural way. Let $\widehat{\sigma}^\varphi: \mathbb{R} \simeq \widehat{\mathbb{R}} \rightarrow \text{Aut}(\mathcal{R})$ be the dual action of σ^φ and note that $\widehat{\sigma}_r^\varphi(x) = x$ for all $r \in \mathbb{R}$ if and only if $x \in A$.

There exists a unique normal semi-finite faithful trace τ_0 on \mathcal{R} such that

$$\tau_0 \circ \widehat{\sigma}_r^\varphi = e^{-r} \tau_0, \quad r \in \mathbb{R}.$$

This trace gives rise to the $*$ -algebra $L^0(\mathcal{R}, \tau_0)$ of τ_0 -measurable operators [26, Chapter 4]. Then for any $1 \leq p \leq \infty$, the Haagerup L^p -space $L^p(A, \varphi)$ is defined as

$$L^p(A, \varphi) = \{y \in L^0(\mathcal{R}, \tau_0) : \widehat{\sigma}_r^\varphi(y) = e^{-\frac{r}{p}} y \text{ for all } r \in \mathbb{R}\}.$$

In particular, $L^\infty(A, \varphi) = A \subset \mathcal{R}$. For all $1 \leq p < \infty$, $L^p(A, \varphi)^+ = L^p(A, \varphi) \cap L^0(\mathcal{R}, \tau_0)^+$ is a positive cone on $L^p(A, \varphi)$.

For any normal semi-finite weight γ on A , let $\widehat{\gamma}$ denote the dual weight on \mathcal{R} , see e.g. [39, Section 19] or [42, Section 10.1]. Let h_γ denote the Radon-Nikodym derivative of γ with respect to τ_0 . This is the unique positive operator on $L^2(\mathbb{R}; H)$ affiliated with \mathcal{R} , such that $\widehat{\gamma}(y) = \tau_0(h_\gamma^{\frac{1}{2}} y h_\gamma^{\frac{1}{2}})$, for all $y \in \mathcal{R}^+$. It turns out that h_γ belongs to $L^1(A, \varphi)^+$ if and only if $\gamma \in A_*^+$. Further the mapping $\gamma \mapsto h_\gamma$ on A_*^+ extends to a linear isomorphism from A_* onto $L^1(A, \varphi)$, still denoted by $\gamma \mapsto h_\gamma$. With this isomorphism in hands, we equip $L^1(A, \varphi)$ with the norm $\|\cdot\|_1$ inherited from A_* , that is, $\|h_\gamma\|_1 = \|\gamma\|_{A_*}$ for all $\gamma \in A_*$. Next, for any $1 \leq p < \infty$ and any $y \in L^p(A, \varphi)$, the positive operator $|y|$ belongs to $L^p(A, \varphi)$ as well and hence $|y|^p$ belongs to $L^1(A, \varphi)$. This allows to define $\|y\|_p = \| |y|^p \|_1^{\frac{1}{p}}$ for all $y \in L^p(A, \varphi)$. Then $\|\cdot\|_p$ is a complete norm on $L^p(A, \varphi)$.

The Banach spaces $L^p(A, \varphi)$ satisfy the following version of Hölder's inequality (see e.g. [26, Proposition 9.17]). For any $1 \leq p, p' \leq \infty$ such that $p^{-1} + p'^{-1} = 1$ and for any $x \in L^p(A, \varphi)$ and $y \in L^{p'}(A, \varphi)$, the product xy belongs to $L^1(A, \varphi)$ and we have

$$\|xy\|_1 \leq \|x\|_p \|y\|_{p'}.$$

We let $\text{Tr}_\varphi: L^1(A, \varphi) \rightarrow \mathbb{C}$ be defined by $\text{Tr}_\varphi(h_\gamma) = \gamma(1)$ for all $\gamma \in A_*$. This functional has two remarkable properties. First, for any $x \in L^p(A, \varphi)$ and $y \in L^{p'}(A, \varphi)$ with $p^{-1} + p'^{-1} = 1$, we have $\text{Tr}_\varphi(xy) = \text{Tr}_\varphi(yx)$. Second,

$$\text{Tr}_\varphi(h_\gamma x) = \gamma(x),$$

for all $\gamma \in A_*$ and $x \in A$. It follows from this identity and the definition of $\|\cdot\|_1$ that the duality pairing $\langle x, y \rangle = \text{Tr}_\varphi(xy)$ for $x \in A$ and $y \in L^1(A, \varphi)$ yields an isometric isomorphism

$$(2.3) \quad L^1(A, \varphi)^* \simeq A.$$

We also note that $L^2(A, \varphi)$ is a Hilbert space for the inner product

$$(2.4) \quad (x|y)_{L^2(A)} = \text{Tr}_\varphi(y^*x), \quad x, y \in L^2(A, \varphi).$$

We let $D_\varphi = h_\varphi$ be the Radon-Nikodym derivative of φ with respect to τ_0 . This operator is called the density of φ . Note that it belongs to $L^1(A, \varphi)$ if and only if φ is finite. It

was shown in [18, Section 2] that for any $1 \leq p < \infty$, $D_\varphi^{\frac{1}{2p}} x D_\varphi^{\frac{1}{2p}}$ belongs to $L^p(A, \varphi)$ for all $x \in \mathcal{M}_\varphi$, and that $D_\varphi^{\frac{1}{2p}} \mathcal{M}_\varphi D_\varphi^{\frac{1}{2p}}$ is dense in $L^p(A, \varphi)$. In the case $p = 1$, we further have

$$(2.5) \quad \varphi(x) = \text{Tr}_\varphi(D_\varphi^{\frac{1}{2}} x D_\varphi^{\frac{1}{2}}), \quad x \in \mathcal{M}_\varphi,$$

see [18, Proposition 2.13]. We refer to [18, Section 1] for the precise meaning of the above products.

2.3. The Haagerup-Junge-Xu extension property and p -dilations. Consider two weighted von Neumann algebras (A, φ) and (A', φ') , and let $V: A \rightarrow A'$ be any weight-preserving positive operator. If $x \in \mathcal{N}_\varphi$, then $\varphi'(V(x^*x)) = \varphi(x^*x) < \infty$, hence $V(x^*x) \in \mathcal{M}_{\varphi'}$. Consequently, V maps \mathcal{M}_φ into $\mathcal{M}_{\varphi'}$. Hence for any $1 \leq p < \infty$, one may define

$$V_p: D_\varphi^{\frac{1}{2p}} \mathcal{M}_\varphi D_\varphi^{\frac{1}{2p}} \longrightarrow L^p(A', \varphi'), \quad V_p(D_\varphi^{\frac{1}{2p}} x D_\varphi^{\frac{1}{2p}}) = D_{\psi'}^{\frac{1}{2p}} V(x) D_{\psi'}^{\frac{1}{2p}}, \quad \text{for all } x \in \mathcal{M}_\varphi.$$

The following extension result is due to Haagerup-Junge-Xu.

Proposition 2.4. [23, Proposition 5.4 & Remark 5.6] *For any weight-preserving positive map $V: (A, \varphi) \rightarrow (A', \varphi')$ and any $1 \leq p < \infty$, V_p extends to a (necessarily unique) bounded map from $L^p(A, \varphi)$ into $L^p(A', \varphi')$. If further $\|V\| \leq 1$, then we have*

$$\|V_p: L^p(A, \varphi) \longrightarrow L^p(A', \varphi')\| \leq 1, \quad 1 \leq p < \infty.$$

Remark 2.5. Consider $V: (A, \varphi) \rightarrow (A', \varphi')$ as in Proposition 2.4 and assume that V is an isometric isomorphism. Then for all $1 \leq p < \infty$, $V_p: L^p(A, \varphi) \rightarrow L^p(A', \varphi')$ is an isometric isomorphism as well. To prove this, it suffices to apply Proposition 2.4 to V and to V^{-1} , and to observe that $\{V^{-1}\}_p = \{V_p\}^{-1}$.

Definition 2.6. *Let (A, φ) be a weighted von Neumann algebra and let $T: L^p(A, \varphi) \rightarrow L^p(A, \varphi)$ be a contraction. We say that T admits an isometric p -dilation if there exist another weighted von Neumann algebra (M, ψ) , an isometric isomorphism $U: L^p(M, \psi) \rightarrow L^p(M, \psi)$ and two contractions $J: L^p(A, \varphi) \rightarrow L^p(M, \psi)$ and $Q: L^p(M, \psi) \rightarrow L^p(A, \varphi)$ such that*

$$\forall k \geq 0, \quad T^k = QU^k J.$$

The following is a straightforward application of Proposition 2.4 and Remark 2.5. It extends [17, Proposition 2.3] which was concerned by the tracial case only (see also [5, p. 2281] for the state case).

Proposition 2.7. *Let $T: (A, \varphi) \rightarrow (A, \varphi)$ be a positive weight-preserving contraction. For any $1 \leq p < \infty$, let $T_p: L^p(A, \varphi) \rightarrow L^p(A, \varphi)$ be the contraction provided by Proposition 2.4. If T admits an absolute dilation, then T_p admits an isometric p -dilation.*

2.4. The use of analytic elements. Let (A, φ) be a weighted von Neumann algebra. We let $A_a \subset A$ be the subset of all $x \in A$ such that $r \mapsto \sigma_r^\varphi(x)$ extends to an entire function $z \in \mathbb{C} \mapsto \sigma_z^\varphi(x) \in A$. Such elements are called analytic. They play a key role in the study of absolute dilations. Following [18, p.46], we introduce

$$\mathcal{N}_\varphi^\infty = \{x \in A_a : \sigma_z^\varphi(x) \in \mathcal{N}_\varphi \cap \mathcal{N}_\varphi^* \text{ for all } z \in \mathbb{C}\}$$

and

$$\mathcal{M}_\varphi^\infty = \text{Span}\{y^*x : x, y \in \mathcal{N}_\varphi^\infty\}.$$

It is easy to check that $\mathcal{M}_\varphi^\infty \subset \mathcal{N}_\varphi^\infty$. According to [18, Theorem 2.4,(a)], $\mathcal{M}_\varphi^\infty$ is a w^* -dense $*$ -subalgebra of A and by [18, Theorem 2.4,(b)], we have the following.

Lemma 2.8. *The space $D_\varphi^{\frac{1}{2}}\mathcal{M}_\varphi^\infty D_\varphi^{\frac{1}{2}} \subset L^1(A, \varphi)$ is dense in $L^1(A, \varphi)$.*

The following commutation property is given by [18, Lemma 2.5].

Lemma 2.9. *For all $x \in \mathcal{M}_\varphi^\infty$, we have $x D_\varphi^{\frac{1}{4}} = D_\varphi^{\frac{1}{4}} \sigma_{-\frac{i}{4}}^\varphi(x)$.*

Next we record a variant of (2.5), based on Lemma 2.9, for which we refer to [18, Proposition 2.13].

Lemma 2.10. *For any $x \in \mathcal{N}_\varphi^*$ and any $y \in \mathcal{M}_\varphi^\infty$, $x \sigma_{-\frac{i}{2}}^\varphi(y)$ belongs to \mathcal{M}_φ and we have*

$$\text{Tr}_\varphi(x D_\varphi^{\frac{1}{2}} y D_\varphi^{\frac{1}{2}}) = \varphi(x \sigma_{-\frac{i}{2}}^\varphi(y)).$$

Remark 2.11. Consider $V: (A, \varphi) \rightarrow (A', \varphi')$ as in Proposition 2.4. Using (2.3) both for A and for A' , we may consider $V_1^*: A' \rightarrow A$. Then V_1^* is a weight-preserving positive map. This is proved in [23, Proposition 5.4] in the case when the weights φ and φ' are finite. This actually extends to the general case [47]. The proof uses the space $\mathcal{N}_\varphi^\infty$.

Lemma 2.12. *Let $J: (A, \varphi) \rightarrow (M, \psi)$ be a w^* -continuous, unital and weight-preserving $*$ -homomorphism such that $\sigma^\psi \circ J = J \circ \sigma^\varphi$. Let \mathbb{E}_J be the conditional expectation associated with J , see (2.1). Then*

$$\mathbb{E}_J = J_1^*.$$

Proof. We know that J_1^* is a contraction. Furthermore, J_1^* is a weight-preserving positive map, by Remark 2.11. Hence it remains to prove that $J_1^* \circ J = I_A$.

Let $x \in \mathcal{M}_\varphi$ and let $y \in \mathcal{M}_\varphi^\infty$. Then

$$\langle J_1^* \circ J(x), D_\varphi^{\frac{1}{2}} y D_\varphi^{\frac{1}{2}} \rangle_{A, L^1(A)} = \langle J(x), J_1(D_\varphi^{\frac{1}{2}} y D_\varphi^{\frac{1}{2}}) \rangle_{M, L^1(M)} = \langle J(x), D_\psi^{\frac{1}{2}} J(y) D_\psi^{\frac{1}{2}} \rangle_{M, L^1(M)}.$$

Since J is a $*$ -homomorphism, we have $J(\mathcal{N}_\varphi) \subset \mathcal{N}_\psi$. Using the commutation property $\sigma^\psi \circ J = J \circ \sigma^\varphi$, we deduce that $J(\mathcal{N}_\varphi^\infty) \subset \mathcal{N}_\psi^\infty$. Using again that J is a $*$ -homomorphism, we deduce that

$$J(\mathcal{M}_\varphi^\infty) \subset \mathcal{M}_\psi^\infty.$$

Likewise,

$$J(\mathcal{M}_\varphi) \subset \mathcal{M}_\psi.$$

Hence $J(y) \in \mathcal{M}_\psi^\infty$ and $J(x) \in \mathcal{M}_\psi \subset \mathcal{N}_\psi^*$. By Lemma 2.10 (applied to ψ), we deduce that

$$\langle J_1^* \circ J(x), D_\varphi^{\frac{1}{2}} y D_\varphi^{\frac{1}{2}} \rangle_{A, L^1(A)} = \text{Tr}_\psi(J(x) D_\psi^{\frac{1}{2}} J(y) D_\psi^{\frac{1}{2}}) = \psi(J(x) \sigma_{-\frac{i}{2}}^\psi(J(y))).$$

Moreover, $J(x) \sigma_{-\frac{i}{2}}^\psi(J(y)) = J(x \sigma_{-\frac{i}{2}}^\varphi(y))$ and $\psi \circ J = \varphi$, hence

$$\langle J_1^* \circ J(x), D_\varphi^{\frac{1}{2}} y D_\varphi^{\frac{1}{2}} \rangle_{A, L^1(A)} = \varphi(x \sigma_{-\frac{i}{2}}^\varphi(y)).$$

Thus, by Lemma 2.10 again, we obtain that

$$\langle J_1^* \circ J(x), D_{\varphi}^{\frac{1}{2}} y D_{\varphi}^{\frac{1}{2}} \rangle_{A, L^1(A)} = \langle x, D_{\varphi}^{\frac{1}{2}} y D_{\varphi}^{\frac{1}{2}} \rangle_{A, L^1(A)}.$$

Since J, J_1^* are w^* -continuous and \mathcal{M}_{φ} is w^* -dense, the result follows from Lemma 2.8. \square

3. BACKGROUND ON FOURIER MULTIPLIERS

Let G be a locally compact group equipped with a fixed left Haar measure dg . We let ϵ denote the unit of G . Let $\lambda: G \rightarrow B(L^2(G))$ be the left regular representation of G , i.e.

$$[\lambda(g)h](t) = h(g^{-1}t), \quad h \in L^2(G), \quad g, t \in G.$$

Let $VN(G) \subset B(L^2(G))$ be the von Neumann algebra generated by $\{\lambda(g) : g \in G\}$. For any $f \in L^1(G)$, we set

$$\lambda(f) = \int_G f(g)\lambda(g) dg,$$

where the integral is defined in the strong sense. The operators $\lambda(f)$ belong to $VN(G)$ and $\lambda(L^1(G))$ is actually a w^* -dense $*$ -subalgebra of $VN(G)$. Further for all $f_1, f_2 \in L^1(G)$, we have $\lambda(f_1 * f_2) = \lambda(f_1)\lambda(f_2)$.

Let $\Delta_G: G \rightarrow (0, \infty)$ be the modular function of G and recall that it is characterized by the identity

$$\int_G f(g^{-1}) dg = \int_G f(g)\Delta_G(g)^{-1} dg.$$

For any $f \in L^1(G)$, let $f^* \in L^1(G)$ be defined by

$$f^*(g) = \Delta_G(g)^{-1} \overline{f(g^{-1})}, \quad g \in G.$$

Then it is easy to check that $\lambda(f)^* = \lambda(f^*)$.

Let $K(G)$ be the space of all continuous functions $G \rightarrow \mathbb{C}$ with compact support. It follows from above that $\lambda(K(G))$ is a $*$ -subalgebra of $VN(G)$. Since $K(G)$ is dense in $L^1(G)$, the latter is w^* -dense.

We equip $VN(G)$ with the so-called Plancherel weight

$$\varphi_G: VN(G)^+ \longrightarrow [0, \infty],$$

for which we refer to [21, Section 2] (see also [42, Section VII.3]). For simplicity we let

$$\sigma^G := \sigma^{\varphi_G} \quad \text{and} \quad D_G := D_{\varphi_G}$$

denote the modular automorphism group and the density of φ_G , respectively.

We record for later use that for any $f \in L^1(G) \cap L^2(G)$, $\lambda(f) \in \mathcal{N}_{\varphi_G}$ and

$$(3.1) \quad \varphi_G(\lambda(f)^* \lambda(f)) = \|f\|_{L^2(G)}^2, \quad f \in L^1(G) \cap L^2(G).$$

Moreover if $f \in K(G)$ is such that $\lambda(f) \geq 0$, then $\lambda(f) \in \mathcal{M}_{\varphi_G}$ and

$$(3.2) \quad \varphi_G(\lambda(f)) = f(\epsilon).$$

We also note that by the last but one line of [21, p. 125], we have

$$(3.3) \quad \sigma_r^G(\lambda(g)) = \Delta_G(g)^{ir} \lambda(g), \quad g \in G, \quad r \in \mathbb{R}.$$

This readily implies

$$(3.4) \quad \sigma_r^G(\lambda(f)) = \lambda(\Delta_G^{ir} f), \quad f \in L^1(G), \quad r \in \mathbb{R}.$$

Consequently, $\lambda(K(G))$ is σ^G -invariant.

Let $C_b(G)$ be the space of all bounded continuous functions $u: G \rightarrow \mathbb{C}$. Following [14], we say that $u \in C_b(G)$ is a bounded Fourier multiplier on $VN(G)$ if there exists a w^* -continuous operator $M_u: VN(G) \rightarrow VN(G)$ such that

$$M_u(\lambda(g)) = u(g)\lambda(g), \quad g \in G.$$

In this case, the operator M_u is necessarily unique. Furthermore, we have

$$M_u(\lambda(f)) = \lambda(uf), \quad f \in L^1(G),$$

see [14, Proposition 1.2] and its proof.

Lemma 3.1. *Assume that u is a bounded Fourier multiplier on $VN(G)$.*

- (1) *For all $r \in \mathbb{R}$, we have $\sigma_r^G \circ M_u = M_u \circ \sigma_r^G$.*
- (2) *The operator M_u is unital if and only if $u(\epsilon) = 1$.*
- (3) *If M_u is both unital and positive, then $\varphi_G \circ M_u = \varphi_G$.*

Proof. According to (3.3), we have

$$(\sigma_r^G \circ M_u)(\lambda(g)) = \Delta_G(g)^{ir} u(g)\lambda(g) = (M_u \circ \sigma_r^G)(\lambda(g)),$$

for all $g \in G$ and all $r \in \mathbb{R}$. Since M_u is w^* -continuous, (1) follows at once.

Part (2) is clear, since $\lambda(\epsilon)$ is the unit of $VN(G)$.

To prove (3), assume that M_u is both unital and positive. Let $f \in K(G)$ such that $\lambda(f) \geq 0$. Then $uf \in K(G)$ and $\lambda(uf) = M_u(\lambda(f)) \geq 0$. Hence by (3.2),

$$\varphi_G(M_u(\lambda(f))) = (uf)(\epsilon) = f(\epsilon) = \varphi_G(\lambda(f)).$$

This implies that for all $f_1 \in K(G)$, we have $(\varphi_G \circ M_u)(\lambda(f_1)^*\lambda(f_1)) = \varphi_G(\lambda(f_1)^*\lambda(f_1))$. Using part (1), the σ^G -invariance of $\lambda(K(G))$, the w^* -density of $\lambda(K(G))$ and Lemma 2.3, we deduce that $\varphi_G \circ M_u = \varphi_G$. \square

It follows from (3.4) that

$$\lambda(K(G)) \subset VN(G)_a, \quad \text{with } \sigma_z^G(\lambda(f)) = \lambda(\Delta_G^{iz} f)$$

for all $f \in K(G)$ and all $z \in \mathbb{C}$. This identity further implies that $\lambda(f) \in \mathcal{N}_{\varphi_G}^\infty$. Let $K(G) * K(G)$ denote the linear span of all functions $f_1 * f_2$, with $f_1, f_2 \in K(G)$. We deduce from above that

$$(3.5) \quad \lambda(K(G) * K(G)) \subset \mathcal{M}_{\varphi_G}^\infty.$$

We will use the Haagerup non-commutative L^p -spaces $L^p(VN(G), \varphi_G)$ associated with the Plancherel weight of G . If u is a bounded Fourier multiplier on $VN(G)$ and M_u is both unital and positive, then $\|M_u\| \leq 1$, and hence

$$(3.6) \quad |u(g)| \leq 1, \quad g \in G.$$

It further follows from Lemma 3.1, (3) and Proposition 2.4 that for any $1 \leq p < \infty$, M_u gives rise to a contraction $M_{u,p}: L^p(VN(G), \varphi_G) \rightarrow L^p(VN(G), \varphi_G)$ taking $D_G^{\frac{1}{2p}} x D_G^{\frac{1}{2p}}$ to $D_G^{\frac{1}{2p}} M_u(x) D_G^{\frac{1}{2p}}$ for all $x \in \mathcal{M}_{\varphi_G}$.

Lemma 3.2. *Assume that u is a bounded Fourier multiplier on $VN(G)$ such that M_u is both unital and positive. The following assertions are equivalent.*

- (i) *The function u is real-valued.*
- (ii) *The operator $M_{u,2}: L^2(VN(G), \varphi_G) \rightarrow L^2(VN(G), \varphi_G)$ is self-adjoint.*

In this case, $M_{u,2}$ is positive in the Hilbertian sense, that is, $(M_{u,2}(h)|h)_{L^2(VN(G))} \geq 0$ for all $h \in L^2(VN(G), \varphi_G)$, if and only if u is valued in $[0, 1]$.

Proof. We first note that $\varphi_G(\lambda(f')^* \lambda(f)) = (f|f')_{L^2(VN(G))}$ for all $f, f' \in L^1(G) \cap L^2(G)$, by polarisation of the identity (3.1). We use this identity twice below.

Let $f, f' \in K(G) * K(G)$ and recall (3.5). By Lemma 2.9 and the fact that M_u commutes with σ^G , see Lemma 3.1, (1), we have

$$\begin{aligned} M_{u,2}(\lambda(f) D_G^{\frac{1}{2}}) &= M_{u,2} \left(D_G^{\frac{1}{4}} \sigma_{\frac{i}{4}}^G(\lambda(f)) D_G^{\frac{1}{4}} \right) \\ &= D_G^{\frac{1}{4}} M_u \left(\sigma_{\frac{i}{4}}^G(\lambda(f)) \right) D_G^{\frac{1}{4}} \\ &= D_G^{\frac{1}{4}} \sigma_{\frac{i}{4}}^G(\lambda(uf)) D_G^{\frac{1}{4}} \\ &= \lambda(uf) D_G^{\frac{1}{2}}. \end{aligned}$$

Hence by (2.5), we have

$$\left(M_{u,2}(\lambda(f) D_G^{\frac{1}{2}}) | \lambda(f') D_G^{\frac{1}{2}} \right)_{L^2(VN(G))} = \text{Tr}_{\varphi_G} \left(D_G^{\frac{1}{2}} \lambda(f')^* \lambda(uf) D_G^{\frac{1}{2}} \right) = \varphi_G(\lambda(f')^* \lambda(uf)).$$

Therefore,

$$(3.7) \quad \left(M_{u,2}(\lambda(f) D_G^{\frac{1}{2}}) | \lambda(f') D_G^{\frac{1}{2}} \right)_{L^2(VN(G))} = \int_G \overline{f'} u f.$$

Likewise,

$$\left(\lambda(f) D_G^{\frac{1}{2}} | M_{u,2}(\lambda(f') D_G^{\frac{1}{2}}) \right)_{L^2(VN(G))} = \int_G \overline{u f'} f.$$

We deduce that u is real-valued if and only if

$$\left(M_{u,2}(\lambda(f) D_G^{\frac{1}{2}}) | \lambda(f') D_G^{\frac{1}{2}} \right)_{L^2(VN(G))} = \left(\lambda(f) D_G^{\frac{1}{2}} | M_{u,2}(\lambda(f') D_G^{\frac{1}{2}}) \right)_{L^2(VN(G))}$$

for all $f, f' \in K(G) * K(G)$.

It follows from [18, Theorem 2.4,(b)] that $\mathcal{N}_{\varphi_G}^\infty D_G^{\frac{1}{2}}$ is dense in $L^2(VN(G), \varphi_G)$. The proof of this theorem shows as well that $\lambda(K(G) * K(G)) D_G^{\frac{1}{2}}$ is dense in $L^2(VN(G), \varphi_G)$. We therefore deduce the equivalence “(i) \Leftrightarrow (ii)” from above.

By (3.6), the function u is valued in $[-1, 1]$. Hence the last assertion is obtained by applying (3.7) with $f' = f$. \square

4. DILATION RESULT

The main result of this paper is the following theorem. In the case when G is discrete, this result was established in [4, Theorem 4.6], see also [38]. Next, it was proved in [16, Theorem (B)] for any unimodular group G . The approach for the non-unimodular case is similar, however significant difficulties arise from the fact that the Plancherel weight may be non tracial.

Theorem 4.1. *Let G be a locally compact group and let $u \in C_b(G)$ such that $M_u: VN(G) \rightarrow VN(G)$ is unital and completely positive. If u is real valued, then M_u admits an absolute dilation.*

Proof. We let $\ell_G^{1,\mathbb{R}}$ be the real ℓ^1 -space over G , with standard basis $(e_t)_{t \in G}$. Define a bilinear form $[\cdot, \cdot]_u: \ell_G^{1,\mathbb{R}} \times \ell_G^{1,\mathbb{R}} \rightarrow \mathbb{R}$ by

$$[\alpha, \beta]_u = \sum_{s,t \in G} u(t^{-1}s) \alpha_t \beta_s, \quad \alpha = (\alpha_t)_{t \in G}, \beta = (\beta_s)_{s \in G} \in \ell_G^{1,\mathbb{R}}.$$

According to [14, Proposition 4.2], the complete positivity of $M_u: VN(G) \rightarrow VN(G)$ implies that the function u is positive definite. Hence $[\cdot, \cdot]_u$ is positive semi-definite. Let $E_u = \{\alpha \in \ell_G^{1,\mathbb{R}} : [\alpha, \alpha]_u = 0\}$ and let H_u be the real Hilbert space obtained by completing the space $\ell_G^{1,\mathbb{R}}/E_u$ equipped with the norm induced by $[\alpha, \alpha]_u(\alpha, \alpha)^{\frac{1}{2}}$. In the sequel, we let $h_t \in H_u$ denote the class of $e_t \in \ell_G^1(\mathbb{R})$, for all $t \in G$. We let $(\cdot | \cdot)_{H_u}$ denote the inner product on H_u .

Let N be the Clifford von Neumann algebra over H_u and let τ denote its standard trace. Let N_{sa} be the self-adjoint part of N . We recall that there exists an \mathbb{R} -linear map

$$w: H_u \longrightarrow N_{sa}$$

such that $w(\xi)$ is a unitary for all $\xi \in H_u$ with $\|\xi\|_{H_u} = 1$,

$$(4.1) \quad \tau(w(h)w(h')) = (h|h')_{H_u}, \quad h, h' \in H_u,$$

and N is generated (as a von Neumann algebra) by the range of w . We refer to [11, 12, 38] for the construction of the Clifford von Neumann algebra and more information.

Fix some $g \in G$. For any $(\alpha_t)_{t \in G} \in \ell_G^{1,\mathbb{R}}$, we have

$$\begin{aligned} \left\| \sum_t \alpha_{g^{-1}t} h_t \right\|_{H_u}^2 &= \sum_{s,t \in G} u(t^{-1}s) \alpha_{g^{-1}t} \alpha_{g^{-1}s} = \sum_{s,t \in G} u((gt)^{-1}(gs)) \alpha_t \alpha_s \\ &= \sum_{s,t \in G} u(t^{-1}s) \alpha_t \alpha_s = \left\| \sum_t \alpha_t h_t \right\|_{H_u}^2. \end{aligned}$$

Hence there exists a (necessarily unique) orthogonal map $\theta_g \in B(H_u)$ such that

$$\theta_g(h_t) = h_{gt}, \quad t \in G.$$

Let $\tilde{\theta}_g: N \rightarrow N$ be the second quantization of θ_g . By [11, Theorem 2.11], $\tilde{\theta}_g$ is a trace-preserving $*$ -automorphism and we have

$$(4.2) \quad \tilde{\theta}_g(w(h)) = w(\theta_g(h)), \quad h \in H_u.$$

Recall that τ is normalized. Let

$$N^\otimes = \overline{\otimes}_{\mathbb{Z}} N,$$

be the infinite tensor product of (N, τ) over the index set \mathbb{Z} , as considered in [43, Definition XIV.1.6]. Note that the associated state $\tau_\otimes := \overline{\otimes}_{\mathbb{Z}} \tau_N$ on N^\otimes is a normal faithful tracial state. In the sequel, we let 1 denote the unit of N and we let 1_\otimes denote the unit of N^\otimes .

For any integer $m \geq 1$ and for any finite family $(x_k)_{k=-m}^m$ in N , set

$$\otimes_k x_k = \cdots 1 \otimes 1 \otimes x_{-m} \otimes \cdots \otimes x_0 \otimes x_1 \otimes \cdots \otimes x_m \otimes 1 \otimes 1 \cdots,$$

where each x_k is in position k . For any $g \in G$, we let $\rho(g): N^\otimes \rightarrow N^\otimes$ be the $*$ -automorphism such that

$$[\rho(g)](\otimes_k x_k) = \otimes_k \tilde{\theta}_g(x_k)$$

for all finite families $(x_k)_k$ in N . Since $\tilde{\theta}_g$ is trace-preserving, $\rho(g)$ is trace-preserving. The mapping $G \rightarrow \text{Aut}(N)$ taking any $g \in G$ to $\tilde{\theta}_g$ is a group homomorphism and it follows from [16, Lemma 5.4] that it is continuous in the sense of [42, Proposition X.1.2]. We easily deduce that $\rho: G \rightarrow \text{Aut}(N^\otimes)$ is a continuous group homomorphism as well.

We are going to use crossed products, for which we refer again to either [39, Chapter IV] or [42, Chapter X]. We let

$$M = N^\otimes \rtimes_\rho G \subset B(L^2(G)) \overline{\otimes} N^\otimes$$

be the crossed product associated with ρ . We regard $N^\otimes \subset M$ in the usual way. We recall that $VN(G) \otimes 1_\otimes \subset M$ and that M is generated (as a von Neumann algebra) by N^\otimes and $VN(G) \otimes 1_\otimes$. We also recall (see e.g. [39, 19.1.(2)]) that

$$(4.3) \quad \rho(g)x = (\lambda(g) \otimes 1_\otimes)x(\lambda(g^{-1}) \otimes 1_\otimes), \quad x \in N^\otimes, g \in G.$$

This implies that the linear span of $\{x(y \otimes 1_\otimes) : x \in N^\otimes, y \in VN(G)\}$ is w^* -dense in M . We will use it in the sequel without any further reference.

Let $K(G, N^\otimes)$ be the space of all σ -strong $*$ -continuous functions $G \rightarrow N^\otimes$ with compact support. For any $F \in K(G, N^\otimes)$, we define $T_F \in M$ by

$$T_F = \int_G F(g)(\lambda(g) \otimes 1_\otimes) dg,$$

where the integral is defined in the w^* -sense. Following [42, pp. 242-243], given any $F_1, F_2 \in K(G, N^\otimes)$, we may define $F_1 * F_2 \in K(G, N^\otimes)$ by

$$(F_1 * F_2)(g) = \int_G F_1(t)\rho(t)(F_2(t^{-1}g)) dt, \quad g \in G,$$

and we have

$$T_{F_1} * T_{F_2} = T_{F_1 * F_2}.$$

Similarly, given any $F \in K(G, N^\otimes)$, we may define $F^* \in K(G, N^\otimes)$ by

$$F^*(g) = \Delta_G(g)^{-1}\rho(g)(F(g^{-1})^*), \quad g \in G,$$

and we have

$$T_F^* = T_{F^*}.$$

Using these definitions, we see that

$$(4.4) \quad (F^* * F)(g) = \int_G \Delta_G(t)^{-1} \rho(t) (F(t^{-1})^* F(t^{-1}g)) dt, \quad g \in G.$$

It actually follows from [42, Lemma X.1.8] that

$$\mathcal{B} := \{T_F : F \in K(G, N^\otimes)\}$$

is a w^* -dense $*$ -subalgebra of M .

We let ψ be the dual weight of τ_\otimes on M . This weight is normal, semi-finite and faithful [42, Lemma X.1.18], so that (M, ψ) is a weighted von Neumann algebra. Moreover

$$\mathcal{B} \subset \mathcal{N}_\psi$$

and for any $F \in K(G, N^\otimes)$, we have

$$\psi(T_F^* T_F) = \tau_\otimes((F^* * F)(\epsilon)),$$

see [42, Theorem X.1.17,(i)]. This is an analogue of (3.2). According to [42, Theorem X.1.17,(ii)] and (3.3), the modular automorphism group σ^ψ satisfies:

$$(4.5) \quad \forall x \in N^\otimes, \quad \sigma_r^\psi(x) = x \quad \text{and} \quad \forall y \in VN(G), \quad \sigma_r^\psi(y \otimes 1_\otimes) = \sigma_r^G(y) \otimes 1_\otimes,$$

for all $r \in \mathbb{R}$.

We define a w^* -continuous unital $*$ -homomorphism

$$J: VN(G) \longrightarrow M, \quad J(y) = y \otimes 1_\otimes \quad \text{for all } y \in VN(G).$$

It follows from the second part of (4.5) that

$$(4.6) \quad \sigma^\psi \circ J = J \circ \sigma^G.$$

Let $f \in K(G)$ and let $F = f \otimes 1_\otimes \in K(G, N^\otimes)$. Then $J(\lambda(f)) = T_F$ and we both have $F^* * F = (f^* * f) \otimes 1_\otimes$ and $J(\lambda(f)^* \lambda(f)) = T_{F^* * F}$. Therefore,

$$\psi(J(\lambda(f)^* \lambda(f))) = \tau_\otimes((F^* * F)(\epsilon)) = (f^* * f)(\epsilon) = \varphi_G(\lambda(f)^* \lambda(f)),$$

by (3.2). We deduce, using (4.6) and Lemma 2.3, that J is weight-preserving.

Let us record two remarkable properties of J . First, we have

$$(4.7) \quad \psi(xJ(y)) = \tau_\otimes(x) \varphi_G(y), \quad x \in N^\otimes, \quad y \in \mathcal{M}_{\varphi_G}.$$

To prove this, consider $x \in N^\otimes \setminus \{0\}$ and define

$$\gamma: VN(G)^+ \longrightarrow [0, \infty], \quad \gamma(y) = \tau_\otimes(x^* x)^{-1} \psi(xJ(y)x^*).$$

This is a normal semi-finite weight.

Given any $f \in K(G)$, let $F \in K(G, N^\otimes)$ be defined by

$$F(g) = \Delta_G(g)^{-1} \overline{f(g^{-1})} \rho(g)(x^*), \quad g \in G.$$

This function is chosen in such a way that $F^* = f \otimes x$. Thus, we have $T_F^* = T_{F^*} = xJ(\lambda(f))$. Consequently,

$$\gamma(\lambda(f)\lambda(f)^*) = \tau_\otimes(x^* x)^{-1} \psi(T_F^* T_F) = \tau_\otimes(x^* x)^{-1} \tau_\otimes((F^* * F)(\epsilon)).$$

Using (4.4), we compute

$$(F^* * F)(\epsilon) = \|f\|_2 x x^*.$$

We deduce that

$$\gamma(\lambda(f)\lambda(f)^*) = \varphi_G(\lambda(f)\lambda(f)^*).$$

By the first part of (4.5), x belongs to the centralizer of ψ , hence

$$\gamma(\cdot) = \tau_\otimes(x^*x)^{-1}\psi(J(\cdot)x^*x),$$

by [42, Theorem VIII.2.6]. Applying (4.6), we deduce that $\gamma \circ \sigma^G = \gamma$. Then arguing as in the proof of Lemma 3.1, (3), we infer that $\gamma = \varphi_G$. Using the fact that x^* belongs to the centralizer of ψ , this yields $\psi(x^*xJ(y)) = \tau_\otimes(x^*x)\varphi_G(y)$ for all $y \in \mathcal{M}_{\varphi_G}$. The identity (4.7) follows at once.

Let us now show that

$$(4.8) \quad \mathbb{E}_J(xJ(y)) = \tau_\otimes(x)y, \quad x \in N^\otimes, \quad y \in VN(G).$$

Since J and \mathbb{E}_J are w^* -continuous, we may assume in the proof of (4.8) that $y \in \mathcal{M}_{\varphi_G}$. Let $m \in \mathcal{M}_{\varphi_G}^\infty$. Since J is a weight-preserving $*$ -homomorphism and J satisfies (4.6), we easily obtain, as in the proof of Lemma 2.12, that

$$J(y) \in \mathcal{M}_\psi \quad \text{and} \quad J(m) \in \mathcal{M}_\psi^\infty.$$

Since x belongs to the centralizer of ψ , the product $xJ(y)$ belongs to \mathcal{M}_ψ , by [42, Theorem VIII.2.6]. Using Lemmas 2.12 and 2.10, we deduce that

$$\begin{aligned} \langle \mathbb{E}_J(xJ(y)), D_G^{\frac{1}{2}}mD_G^{\frac{1}{2}} \rangle_{VN(G), L^1(VN(G))} &= \langle xJ(y), J_1(D_G^{\frac{1}{2}}mD_G^{\frac{1}{2}}) \rangle_{M, L^1(M)} \\ &= \langle xJ(y), D_\psi^{\frac{1}{2}}J(m)D_\psi^{\frac{1}{2}} \rangle_{M, L^1(M)} \\ &= \text{Tr}_\psi(xJ(y)D_\psi^{\frac{1}{2}}J(m)D_\psi^{\frac{1}{2}}) \\ &= \psi(xJ(y)\sigma_{-\frac{i}{2}}^\psi(J(m))). \end{aligned}$$

Using (4.6) again and (4.7), we have

$$\psi(xJ(y)\sigma_{-\frac{i}{2}}^\psi(J(m))) = \psi(xJ(y\sigma_{-\frac{i}{2}}^G(m))) = \tau_\otimes(x)\varphi_G(y\sigma_{-\frac{i}{2}}^G(m)).$$

By Lemmas 2.12 and 2.10 again,

$$\varphi_G(y\sigma_{-\frac{i}{2}}^\psi(m)) = \text{Tr}_{\varphi_G}(yD_G^{\frac{1}{2}}mD_G^{\frac{1}{2}}) = \langle \mathbb{E}_J(y), D_G^{\frac{1}{2}}mD_G^{\frac{1}{2}} \rangle_{VN(G), L^1(VN(G))}.$$

Altogether, we obtain that

$$\langle \mathbb{E}_J(xJ(y)), D_G^{\frac{1}{2}}mD_G^{\frac{1}{2}} \rangle_{VN(G), L^1(VN(G))} = \tau_\otimes(x) \langle \mathbb{E}_J(y), D_G^{\frac{1}{2}}mD_G^{\frac{1}{2}} \rangle_{VN(G), L^1(VN(G))}.$$

Since $D_G^{\frac{1}{2}}\mathcal{M}_{\varphi_G}^\infty D_G^{\frac{1}{2}}$ is dense in $L^1(VN(G))$, see Lemma 2.8, we deduce (4.8).

We now aim at constructing a suitable $U: M \rightarrow M$. We proceed in two steps.

We let $\mathfrak{s}: N^\otimes \rightarrow N^\otimes$ be the $*$ -automorphism such that

$$\mathfrak{s}\left(\bigotimes_k x_k\right) = \bigotimes_k x_{k-1}$$

for all finite families $(x_k)_k$ in N . Clearly \mathfrak{s} is trace-preserving, i.e.

$$(4.9) \quad \tau_{\otimes} \circ \mathfrak{s} = \tau_{\otimes}.$$

Moreover $\rho(g) \circ \mathfrak{s} = \mathfrak{s} \circ \rho(g)$ for all $g \in G$. This commutation property ensures, by [40, Theorem 4.4.4], that there exists a (necessarily unique) $*$ -automorphism $\tilde{\mathfrak{s}}: M \rightarrow M$ such that

$$\forall x \in N^{\otimes}, \quad \tilde{\mathfrak{s}}(x) = \mathfrak{s}(x) \quad \text{and} \quad \forall y \in VN(G), \quad \tilde{\mathfrak{s}}(y \otimes 1_{\otimes}) = y \otimes 1_{\otimes}.$$

It follows from these identities and (4.5) that

$$(4.10) \quad \sigma^{\psi} \circ \tilde{\mathfrak{s}} = \tilde{\mathfrak{s}} \circ \sigma^{\psi}.$$

Let us show that $\tilde{\mathfrak{s}}$ is weight-preserving. Let $F \in K(G, N^{\otimes})$ and let $r \in \mathbb{R}$. Then

$$\sigma_r^{\psi}(T_F) = \int_G \sigma_r^{\psi}(F(g)) \sigma_r^{\psi}(\lambda(g) \otimes 1_{\otimes}) dg = \int_G F(g) \Delta_G(g)^{ir} (\lambda(g) \otimes 1_{\otimes}) dg = T_{\Delta_G^{ir} F}.$$

Hence \mathcal{B} is σ^{ψ} -invariant. Using Lemma 2.3, (4.10), and the fact that \mathcal{B} is a w^* -dense $*$ -subalgebra of M , it therefore suffices to show that

$$(4.11) \quad (\psi \circ \tilde{\mathfrak{s}})(T_F^* T_F) = \psi(T_F^* T_F), \quad F \in K(G, N^{\otimes}),$$

to obtain that $\psi \circ \tilde{\mathfrak{s}} = \psi$. To prove this, consider $F \in K(G, N^{\otimes})$ and write

$$\tilde{\mathfrak{s}}(T_F^* T_F) = \tilde{\mathfrak{s}}\left(\int_G (F^* * F)(g) (\lambda(g) \otimes 1_{\otimes}) dg\right) = \int_G \mathfrak{s}((F^* * F)(g)) (\lambda(g) \otimes 1_{\otimes}) dg.$$

Since $\mathfrak{s} \circ \rho(t) = \rho(t) \circ \mathfrak{s}$, we deduce, using (4.4), that

$$\begin{aligned} \mathfrak{s}((F^* * F)(g)) &= \int_G \Delta_G(t)^{-1} \rho(t) (\mathfrak{s}(F(t^{-1})^*) \mathfrak{s}(F(t^{-1}g))) dt \\ &= [(\mathfrak{s} \circ F)^* * (\mathfrak{s} \circ F)](g), \quad g \in G. \end{aligned}$$

Consequently,

$$(4.12) \quad \tilde{\mathfrak{s}}(T_F^* T_F) = T_{\mathfrak{s} \circ F}^* T_{\mathfrak{s} \circ F}.$$

We deduce, using (4.9), that

$$\begin{aligned} (\psi \circ \tilde{\mathfrak{s}})(T_F^* T_F) &= \tau_{\otimes}([(\mathfrak{s} \circ F)^* * (\mathfrak{s} \circ F)](\epsilon)) \\ &= \tau_{\otimes}(\mathfrak{s}((F^* * F)(\epsilon))) \\ &= \tau_{\otimes}((F^* * F)(\epsilon)) \\ &= \psi(T_F^* T_F), \end{aligned}$$

which proves (4.11).

For any $g \in G$, let $\delta_g \in N^{\otimes}$ be defined by

$$\delta_g = \cdots 1 \otimes \cdots \otimes 1 \otimes w(h_g) \otimes 1 \otimes \cdots \otimes 1 \otimes \cdots,$$

where $w(h_g)$ is in position 0. By assumption, M_u is unital hence $\|h_{\epsilon}\|_{H_u} = u(\epsilon) = 1$, see Lemma 3.1, (2). Consequently, $w(h_{\epsilon})$ is a unitary of N hence δ_{ϵ} is a self-adjoint unitary of $N^{\otimes} \subset M$. Let $U: M \rightarrow M$ be defined by

$$U(z) = \delta_{\epsilon} \tilde{\mathfrak{s}}(z) \delta_{\epsilon}, \quad z \in M.$$

Then U is a $*$ -automorphism. By the first part of (4.5), δ_ϵ belongs to the centralizer of ψ , hence we may write $\psi(U(y)) = \psi(\tilde{\mathfrak{f}}(y)\delta_\epsilon^2) = \psi(\tilde{\mathfrak{f}}(y)) = \psi(y)$. Thus we have

$$\psi \circ U = \psi.$$

Let us now check that $T = M_u$ satisfies (2.2). We observe that by (4.2), we have

$$\rho(g)(\delta_t) = \delta_{gt}, \quad g, t \in G.$$

Applying (4.3), we deduce that

$$(4.13) \quad (\lambda(g) \otimes 1_\otimes)\delta_\epsilon = \delta_g(\lambda(g) \otimes 1_\otimes), \quad g \in G.$$

Let $g \in G$. Then $\tilde{\mathfrak{f}}(\lambda(g) \otimes 1_\otimes) = \lambda(g) \otimes 1_\otimes$, hence by (4.13), we have

$$UJ(\lambda(g)) = \delta_\epsilon \delta_g(\lambda(g) \otimes 1_\otimes).$$

Next,

$$\tilde{\mathfrak{f}}(\delta_\epsilon \delta_g) = \cdots 1 \otimes \cdots \otimes 1 \otimes w(h_\epsilon)w(h_g) \otimes 1 \otimes \cdots \otimes 1 \otimes \cdots,$$

where $w(h_\epsilon)w(h_g)$ is in position 1. Hence δ_ϵ commutes with $\tilde{\mathfrak{f}}(\delta_\epsilon \delta_g)$ and using (4.13) again, we obtain

$$U^2 J(\lambda(g)) = \delta_\epsilon \tilde{\mathfrak{f}}(\delta_\epsilon \delta_g)(\lambda(g) \otimes 1_\otimes)\delta_\epsilon = \tilde{\mathfrak{f}}(\delta_\epsilon \delta_g)\delta_\epsilon \delta_g(\lambda(g) \otimes 1_\otimes).$$

More explicitly,

$$U^2 J(\lambda(g)) = (\cdots \otimes 1 \otimes w(h_\epsilon)w(h_g) \otimes w(h_\epsilon)w(h_g) \otimes 1 \otimes \cdots)(\lambda(g) \otimes 1_\otimes),$$

with $w(h_\epsilon)w(e_g)$ in positions 0 and 1. Then we obtain by induction that for all integer $k \geq 0$, we have

$$U^k J(\lambda(g)) = (\cdots \otimes 1 \otimes w(h_\epsilon)w(e_g) \otimes \cdots \otimes w(h_\epsilon)w(e_g) \otimes 1 \otimes \cdots)(\lambda(g) \otimes 1_\otimes),$$

with $w(h_\epsilon)w(e_g)$ in positions $0, \dots, k-1$. According to (4.8) and (4.1), this implies that

$$\begin{aligned} \mathbb{E}_J U^k J(\lambda(g)) &= \tau_\otimes(\cdots \otimes 1 \otimes w(h_\epsilon)w(e_g) \otimes \cdots \otimes w(h_\epsilon)w(e_g) \otimes 1 \otimes \cdots)\lambda(g) \\ &= \tau(w(h_\epsilon)w(h_g))^k \lambda(g) \\ &= (h_\epsilon | h_g)_{H_u}^k \lambda(g) \\ &= u(g)^k \lambda(g). \end{aligned}$$

Thus, $\mathbb{E}_J U^k J(\lambda(g)) = M_u^k(\lambda(g))$ for all $k \geq 0$ and all $g \in G$. Since the operators M_u, J, U and \mathbb{E}_J are all w^* -continuous and $\text{Span}\{\lambda(g) : g \in G\}$ is w^* -dense in $VN(G)$, we deduce (2.2) for all $k \geq 0$. \square

If we restrict to the case when G is unimodular, the above proof is different and somewhat simpler than the one in [16] or [4]. Indeed, we do not use [16, Lemma 2.7].

Remark 4.2. We observe (as in [16, Theorem 6.3]) that we can simultaneously dilate any finite family of Fourier multipliers satisfying the assumptions of Theorem 4.1. More explicitly, let $u_1, \dots, u_n \in C_b(G)$ such that $M_{u_j} : VN(G) \rightarrow VN(G)$ is unital and completely positive and u_j is real valued, for all $j = 1, \dots, n$. Then there exist a weighted von Neumann algebra (M, ψ) , a w^* -continuous, unital and weight-preserving $*$ -homomorphism $J : VN(G) \rightarrow M$

such that $\sigma^\psi \circ J = J \circ \sigma^\varphi$, as well as a commuting family (U_1, \dots, U_n) of weight-preserving $*$ -automorphisms on M , such that

$$\forall k_1, \dots, k_n \geq 0, \quad M_{u_1}^{k_1} \dots M_{u_n}^{k_n} = \mathbb{E}_J U_1^{k_1} \dots U_n^{k_n} J.$$

5. AN APPLICATION TO FUNCTIONAL CALCULUS

The aim of this section is to establish a functional calculus property of the L^p -realization of certain Fourier multipliers $M_u: VN(G) \rightarrow VN(G)$.

Let $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ be the open unit disc. Let X be a Banach space and let $T \in B(X)$. We say that T is power bounded if there exists a constant $C_0 \geq 1$ such that $\|T^k\| \leq C_0$ for all integer $k \geq 0$. Let $\sigma(T)$ denote the spectrum of T . We say that T is a Ritt operator if $\sigma(T) \subset \overline{\mathbb{D}}$ and there exists a constant $C > 0$ such that

$$\|(z - T)^{-1}\| \leq \frac{C}{|z - 1|}, \quad z \in \mathbb{C} \setminus \overline{\mathbb{D}}.$$

A classical result (see e.g. [46]) asserts that T is a Ritt operator if and only if T is power bounded and there exists a constant $C_1 \geq 1$ such that

$$n\|T^n - T^{n-1}\| \leq C_1, \quad n \geq 1.$$

For any $\gamma \in (0, \frac{\pi}{2})$, let B_γ be the interior of the convex hull of 1 and the disc $\{|z| < \sin \gamma\}$, see Figure 1 below. It is well-known that the spectrum of any Ritt operator T is included in the closure of one of the B_γ .




FIGURE 1. Stolz domain

Let \mathcal{P} be the algebra of complex polynomials and let $T \in B(X)$ be a Ritt operator. For any $\gamma \in (0, \frac{\pi}{2})$, we say that T admits a bounded $H^\infty(B_\gamma)$ -functional calculus if there exists a constant $K \geq 1$ such that

$$\|F(T)\| \leq K \sup\{|F(z)| : z \in B_\gamma\}, \quad F \in \mathcal{P}.$$

Then we simply say that T admits a bounded H^∞ -functional calculus if it admits a bounded $H^\infty(B_\gamma)$ -functional calculus for some $\gamma \in (0, \frac{\pi}{2})$. This implies that T is polynomially bounded, that is, T satisfies an estimate $\|F(T)\| \leq K \sup\{|F(z)| : z \in \mathbb{D}\}$ for all $F \in \mathcal{P}$.

Ritt operator and their H^∞ -functional calculus have attracted a lot of attention in the last twenty years. We refer the reader to [7, 8, 10, 15, 19, 24, 25, 35, 46] and the reference therein for more information.

The main result of this section is the following theorem. Note that in this statement, the multiplier u is not only a real valued function but a nonnegative one.

Theorem 5.1. *Let G be a locally compact group, let $u: G \rightarrow [0, 1]$ be a continuous function, and assume that $M_u: VN(G) \rightarrow VN(G)$ is unital and completely positive. Then for all $1 < p < \infty$, $M_{u,p}: L^p(VN(G), \varphi_G) \rightarrow L^p(VN(G), \varphi_G)$ is a Ritt operator which admits a bounded H^∞ -functional calculus.*

We need sectorial operators and their H^∞ functional calculus. We refer to [28, Chapter 10] for background and undefined terminology on this topic. As in this reference, we set

$$\Sigma_\theta = \{z \in \mathbb{C} \setminus \{0\} : |\text{Arg}(z)| < \theta\},$$

for any $\theta \in (0, \pi)$. The following was proved in [35, Proposition 4.1].

Proposition 5.2. *Let $T \in B(X)$ be a Ritt operator. Then $I_X - T$ is a sectorial operator. If there exists $\theta \in (0, \frac{\pi}{2})$ such that $I_X - T$ admits a bounded $H^\infty(\Sigma_\theta)$ functional calculus, then T admits a bounded H^∞ -functional calculus.*

The proof of Theorem 5.1 will also rely on interpolation. Given any interpolation couple (X_0, X_1) of Banach spaces and $\alpha \in [0, 1]$, we let $[X_0, X_1]_\alpha$ denote the Banach space obtained by the complex interpolation method. We refer to [9, Chapter 4] for background.

Proposition 5.3. *Let (X_0, X_1) be an interpolation couple of Banach spaces and consider an operator $T: X_0 + X_1 \rightarrow X_0 + X_1$ such that*

$$T: X_0 \longrightarrow X_0 \quad \text{and} \quad T: X_1 \longrightarrow X_1$$

are bounded. If $T: X_0 \rightarrow X_0$ is a Ritt operator and $T: X_1 \rightarrow X_1$ is power bounded, then for all $\alpha \in [0, 1)$, $T: [X_0, X_1]_\alpha \rightarrow [X_0, X_1]_\alpha$ is a Ritt operator.

Proof. This result was proved by Blunck in the case when $X_0 = L^p(\Omega)$ and $X_1 = L^q(\Omega)$ for a measure space (Ω, μ) and $1 \leq p, q \leq \infty$, see [10, Theorem 1.1]. His arguments apply as well in this abstract setting. \square

Proof of Theorem 5.1. We set $A = VN(G)$, $\varphi = \varphi_G$ and $D = D_G$ for simplicity. We use complex interpolation for the spaces $L^p(A, \varphi)$. Following [45] (see also [13, Section 6], [32] and [26, Chapter 9]), we regard $(A, L^1(A, \varphi))$ as an interpolation couple of Banach spaces, using the embedding $\mathcal{M}_\varphi \rightarrow L^1(A, \varphi)$ which takes x to $D^{\frac{1}{2}}x D^{\frac{1}{2}}$ for any $x \in \mathcal{M}_\varphi$. Then for any $1 \leq p \leq \infty$, the mapping $D^{\frac{1}{2p}} \mathcal{M}_\varphi D^{\frac{1}{2p}} \rightarrow L^1(A, \varphi)$ which takes $D^{\frac{1}{2p}}x D^{\frac{1}{2p}}$ to $D^{\frac{1}{2}}x D^{\frac{1}{2}}$ for any $x \in \mathcal{M}_\varphi$ extends to an isometric isomorphism

$$(5.1) \quad L^p(A, \varphi) \simeq [A, L^1(A, \varphi)]_{\frac{1}{p}}.$$

Furthermore, if $V: A \rightarrow A$ is a weight-preserving positive map, we can define $T: A + L^1(A, \varphi) \rightarrow A + L^1(A, \varphi)$ by $T(x + y) = V(x) + V_1(y)$ for all $(x, y) \in A \times L^1(A, \varphi)$ and the interpolation operator on $[A, L^1(A, \varphi)]_{\frac{1}{p}}$ induced by T coincides with V_p through the identification (5.1).

Let $u: G \rightarrow [0, 1]$ be a continuous function such that $M_u: VN(G) \rightarrow VN(G)$ is unital and completely positive and let us apply the above properties with $V = M_u$. We fix some $p \in [2, \infty)$, the case when $p \in (1, 2)$ being similar.

According to Lemma 3.2, $M_{u,2}$ is positive in the Hilbertian sense. Hence $\sigma(M_{u,2}) \subset [0, 1]$. Then, by the functional calculus of self-adjoint operators, $M_{u,2}$ is a Ritt operator. Indeed, for any $z \in \mathbb{C} \setminus \overline{\mathbb{D}}$, we have an estimate

$$\|(z - M_{u,2})^{-1}\| \leq \sup\{|z - t|^{-1} : t \in [0, 1]\} \lesssim |z - 1|^{-1}.$$

By (5.1) and the reiteration theorem for complex interpolation (see [9, Theorem 4.6.1]), we have

$$L^p(A, \varphi) \simeq [A, L^2(A, \varphi)]_{\frac{2}{p}}.$$

Hence by Proposition 5.3, $M_{u,p}$ is a Ritt operator.

The rest of the proof is similar to the one of [5, Theorem 7.1]. Let $q \in (p, \infty)$ and let $\theta_0 \in (\frac{\pi}{2}, \pi)$. By Theorem 4.1 and Proposition 2.7, the operator $M_{u,q}$ admits an isometric q -dilation. Hence by [7, Proposition 6.2], the sectorial operator $I_{L^q} - M_{u,q}$ admits a bounded $H^\infty(\Sigma_{\theta_0})$ functional calculus. (The non-commutative L^q -spaces considered in [7, Section 6] are the tracial ones. However the results contained therein extend verbatim to the framework of Haagerup non-commutative L^q -spaces, using the fact that for $1 < q < \infty$, these spaces satisfy the UMD property, see [37, Corollary 7.7].)

Let $\theta_1 \in (0, \frac{\pi}{2})$. The operator $I_{L^2} - M_{u,2}$ is self-adjoint and $\sigma(I_{L^2} - M_{u,2}) \subset [0, 1]$. Hence using again the functional calculus of self-adjoint operators, we see that $I_{L^2} - M_{u,2}$ admits a bounded $H^\infty(\Sigma_{\theta_0})$ functional calculus. Applying again the reiteration theorem for complex interpolation, we obtain that

$$L^p(A, \varphi) \simeq [L^q(A, \varphi), L^2(A, \varphi)]_\alpha, \quad \text{with } \frac{1 - \alpha}{q} + \frac{\alpha}{2} = \frac{1}{p}.$$

Now applying the interpolation theorem for the H^∞ -functional calculus of sectorial operators [31, Proposition 4.9] (see also [30, Proposition 5.8]), we deduce that $I_{L^p} - M_{u,p}$ admits a bounded $H^\infty(\Sigma_\theta)$ functional calculus, with $\theta = (1 - \alpha)\theta_0 + \alpha\theta_1$. We can choose q, θ_0, θ_1 so that $\theta \in (0, \frac{\pi}{2})$. (We can actually obtain bounded $H^\infty(\Sigma_\theta)$ functional calculus for any $\theta > (\frac{1}{2} - \frac{1}{p})\pi$.) By Proposition 5.2, this implies that $M_{u,p}$ admits a bounded H^∞ -functional calculus. \square

Acknowledgements: The two authors were supported by the ANR project Noncommutative analysis on groups and quantum groups (No./ANR-19-CE40-0002).

REFERENCES

- [1] M. A. Akcoglu, *A pointwise ergodic theorem in L_p -spaces*, Canadian J. Math. 27 (1975), no. 5, 1075–1082.

- [2] M. A. Akcoglu and L. Sucheston, *Dilations of positive contractions on L_p -spaces*, *Canad. Math. Bull.* 20 (1977), no. 3, 285–292.
- [3] C. Anantharaman-Delaroche, *On ergodic theorems for free group actions on noncommutative spaces*, *Probab. Theory Related Fields* 135 (2006), no. 4, 520–546.
- [4] C. Arhancet, *On Matsaev’s conjecture for contractions on noncommutative L_p -spaces*, *J. Operator Theory* 69 (2013), no. 2, 387–421.
- [5] C. Arhancet, *Dilations of semigroups on von Neumann algebras and noncommutative L_p -spaces*, *J. Funct. Anal.* 276 (2019), no. 7, 2279–2314.
- [6] C. Arhancet, *Dilations of Markovian semigroups of Fourier multipliers on locally compact groups*, *Proc. Amer. Math. Soc.* 148 (2020), no. 6, 2551–2563.
- [7] C. Arhancet and C. Le Merdy, *Dilation of Ritt operators on L_p -spaces*, *Israel J. Math.* 201 (2014), no. 1, 373–414.
- [8] C. Arhancet, S. Fackler and C. Le Merdy, *Isometric dilations and H^∞ -calculus for bounded analytic semigroups and Ritt operators*, *Trans. Amer. Math. Soc.* 369 (2017), no. 10, 6899–6933.
- [9] J. Bergh and J. Löfström, *Interpolation spaces, an introduction*, *Grundlehren der Mathematischen Wissenschaften*, No. 223. Springer-Verlag, Berlin-New York, 1976, x+207 pp.
- [10] S. Blunck, *Analyticity and discrete maximal regularity on L_p -spaces*, *J. Funct. Anal.* 183 (2001), no. 1, 211–230.
- [11] M. Bozejko, B. Kümmerer and R. Speicher, *q -Gaussian processes: non-commutative and classical aspects*, *Comm. Math. Phys.* 185 (1997), no. 1, 129–154.
- [12] O. Bratelli and D. Robinson, *Operator algebras and quantum statistical mechanics II*, *Texts and Monographs in Physics*. Springer-Verlag, New York-Berlin, 1981. xiii+505 pp.
- [13] M. Caspers and M. de la Salle, *Schur and Fourier multipliers of an amenable group acting on non-commutative L_p -spaces*, *Trans. Amer. Math. Soc.* 367 (2015), no. 10, 6997–7013.
- [14] J. De Cannière and U. Haagerup, *Multipliers of the Fourier algebras of some simple Lie groups and their discrete subgroups*, *Amer. J. Math.* 107 (1985), 455–500.
- [15] N. Dungey, *Subordinated discrete semigroups of operators*, *Trans. Amer. Math. Soc.* 363 (2011), no. 4, 1721–1741.
- [16] C. Duquet, *Dilation properties of measurable Schur multipliers and Fourier multipliers*, *Positivity* 26 (2022), no. 4, Paper No. 69, 41 pp.
- [17] C. Duquet and C. Le Merdy, *A characterization of absolutely dilatable Schur multipliers*, *Adv. Math.* 439 (2024), Paper No. 109492, 33 pp.
- [18] S. Goldstein and J. M. Lindsay, *Markov semigroups KMS-symmetric for a weight*, *Math. Ann.* 313 (1999), no. 1, 39–67.
- [19] A. Gomilko and Y. Tomilov, *On discrete subordination of power bounded and Ritt operators*, *Indiana Univ. Math. J.* 67 (2018), no. 2, 781–829.
- [20] U. Haagerup, *L^p -spaces associated with an arbitrary von Neumann algebra*, In “Algèbres d’opérateurs et leurs applications en physique mathématique (Proc. Colloq., Marseille, 1977)”, pp. 175–184, *Colloq. Internat. CNRS*, 274, CNRS, Paris, 1979.
- [21] U. Haagerup, *On the dual weights for crossed products of von Neumann algebras II*, *Math. Scand.* 43 (1978) 119–140.
- [22] U. Haagerup and M. Musat, *Factorization and dilation problems for completely positive maps on von Neumann algebras*, *Comm. Math. Phys.* 303 (2011), no. 2, 555–594.
- [23] U. Haagerup, M. Junge and Q. Xu, *A reduction method for noncommutative L_p -spaces and applications*, *Trans. Amer. Math. Soc.* 362 (2010), no. 4, 2125–2165.
- [24] B. Haak, *Remarks on Ritt operators, their H^∞ -functional calculus and associated square function estimates*, Preprint, arXiv:2303.03022.
- [25] B. Haak and M. Haase, *Square function estimates and functional calculi*, Preprint, arXiv:1311.0453.
- [26] F. Hiai, *Lectures on selected topics in von Neumann algebras*, *EMS Series of Lectures in Mathematics*, EMS Press, Berlin, 2021, viii+241 pp.

- [27] G. Hong, S. Ray and S. Wang, *Maximal ergodic inequalities for some positive operators on noncommutative L_p -spaces*, Preprint 2019, arXiv:1907.12967.
- [28] T. Hytönen, J. van Neerven, M. Veraar and L. Weis, *Analysis in Banach spaces, Vol. II, Probabilistic methods and operator theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics, 67, Springer, Cham, 2017. xxi+616 pp.
- [29] M. Junge and C. Le Merdy, *Dilations and rigid factorisations on noncommutative L_p -spaces*, J. Funct. Anal. 249 (2007), no. 1, 220–252.
- [30] M. Junge, C. Le Merdy and Q. Xu, *H^∞ functional calculus and square functions on noncommutative L_p -spaces*, Soc. Math. France, Astérisque No. 305 (2006), vi+138 pp.
- [31] N. Kalton, P. Kunstmann and L. Weis, *Perturbation and interpolation theorems for the H^∞ -calculus with applications to differential operators*, Math. Ann. 336 (2006), no. 4, 747–801.
- [32] H. Kosaki, *Applications of the complex interpolation method to a von Neumann algebra: noncommutative L_p -spaces*, J. Funct. Anal. 56 (1984), 29–78.
- [33] B. Kümmerner, *Markov dilations on W^* -algebras*, J. Funct. Anal. 63 (1985), no. 2, 139–177.
- [34] P. C. Kunstmann and L. Weis, *Maximal L_p -regularity for parabolic equations, Fourier multiplier theorems and H^∞ -functional calculus*, Functional analytic methods for evolution equations, 65–311, Lecture Notes in Math., 1855, Springer, Berlin, 2004.
- [35] C. Le Merdy, *H^∞ -functional calculus and square function estimates for Ritt operators*, Rev. Mat. Iberoam. 30 (2014), no. 4, 1149–1190.
- [36] C. Le Merdy and Q. Xu, *Maximal theorems and square functions for analytic operators on L_p -spaces*, J. Lond. Math. Soc. (2) 86 (2012), no. 2, 343–365.
- [37] G. Pisier and Q. Xu, *Non-commutative L^p -spaces*, Handbook of the geometry of Banach spaces, Vol. 2, pp. 1459–1517, North-Holland, Amsterdam, 2003.
- [38] É. Ricard, *A Markov dilation for self-adjoint Schur multipliers*, Proc. Amer. Math. Soc. 136 (2008), no. 12, 4365–4372.
- [39] S. V. Sratila, *Modular theory in operator algebras*, Editura Academiei Republicii Socialiste România, Bucharest; Abacus Press, Tunbridge Wells, 1981, 492 pp.
- [40] V. S. Sunder, *An invitation to von Neumann algebras*, Universitext, Springer-Verlag, New York, 1987. xiv+171 pp.
- [41] M. Takesaki, *Theory of operator algebras. I*, Springer-Verlag, New York-Heidelberg, 1979, vii+415 pp.
- [42] M. Takesaki, *Theory of operator algebras. II*, Encyclopaedia of Mathematical Sciences, 125, Operator Algebras and Non-commutative Geometry, 6, Springer-Verlag, Berlin, 2003, xxii+518 pp.
- [43] M. Takesaki, *Theory of operator algebras. III*, Encyclopaedia of Mathematical Sciences, 127, Operator Algebras and Non-commutative Geometry, 8, Springer-Verlag, Berlin, 2003. xxii+548 pp.
- [44] M. Terp, *L_p -spaces associated with von Neumann algebras*, Notes, Math. Institute, Copenhagen University, 1981.
- [45] M. Terp, *Interpolation spaces between a von Neumann algebra and its predual*, J. Operator Theory 8 (1982), no. 2, 327–360.
- [46] P. Vitse, *A band limited and Besov class functional calculus for Tadmor-Ritt operators*, Arch. Math. (Basel) 85 (2005), no. 4, 374–385.
- [47] Q. Xu, Private communication.

Email address: charles.duquet@univ-fcomte.fr

LABORATOIRE DE MATHÉMATIQUES DE BESANÇON, UNIVERSITÉ DE FRANCHE-COMTÉ, 16 ROUTE DE GRAY 25030 BESANÇON CEDEX, FRANCE

Email address: clemerdy@univ-fcomte.fr

LABORATOIRE DE MATHÉMATIQUES DE BESANÇON, UNIVERSITÉ DE FRANCHE-COMTÉ, 16 ROUTE DE GRAY 25030 BESANÇON CEDEX, FRANCE