

Recursive algorithm for generating high-temperature expansions for spin systems and the chiral non-linear susceptibility

Andreas Rückriegel,¹ Dmytro Tarasevych,¹ Jan Krieg,² and Peter Kopietz¹

¹*Institut für Theoretische Physik, Universität Frankfurt,
Max-von-Laue Straße 1, 60438 Frankfurt, Germany*

²*Eurex Clearing, Deutsche Börse AG, 60485 Frankfurt, Germany*

(Dated: June 10, 2024)

We show that the high-temperature expansion of the free energy and arbitrary connected correlation functions of quantum spin systems can be recursively obtained from the exact renormalization group flow equation for the generating functional of connected spin correlation functions derived by Krieg and Kopietz [Phys. Rev. B **99**, 060403(R) (2019)]. Our recursive algorithm can be explicitly written down in closed form including all combinatorial factors. We use our method to estimate critical temperatures of Heisenberg magnets from low-order truncations of the *inverse* spin susceptibility in the static limit. We also calculate the connected correlation function involving three different spin components (chiral non-linear susceptibility) of quantum Heisenberg magnets up to second order in the exchange couplings.

I. INTRODUCTION

High-temperature series expansions are a powerful numerical method to investigate lattice models for quantum magnetism or strongly correlated electrons [1, 2]. By numerically implementing graph theoretical and combinatorial algorithms it is nowadays possible to evaluate many terms in the high-temperature series [3–7]. State-of-the-art high-temperature expansions of quantum Heisenberg models include terms up to 12th order for thermodynamic quantities and up to 10th order for static correlation functions [5]. For classical spin models such as the Ising model the high-temperature series expansion can be pushed to even higher orders; for example, in Ref. [8] the first 25 terms of the high-temperature expansion of the zero-field susceptibility of the nearest-neighbor Ising model on a cubic lattice have been calculated. From the mathematical analysis of the truncated series one can obtain quantitative estimates for critical phenomena, such as phase boundaries, critical exponents, and critical temperatures. High-temperature series expansions for static two-spin correlation functions of quantum Heisenberg models have also been obtained [5].

While sophisticated numerical algorithms for generating the high-temperature series expansions for spin models are available [5, 7], for the calculation of the first few terms in the high-temperature series expansion it would be useful to have a simple algorithm which can be readily implemented using symbolic manipulation software such as MATHEMATICA without invoking additional graph theoretical or combinatorial concepts. Moreover, it would be useful to have an algorithm for calculating the high-temperature expansion of arbitrary dynamic correlation functions in a magnetic field. In this work we show that such a general algorithm can be obtained from the formally exact functional renormalization group flow equations for the generating functional of connected time-ordered spin correlation functions of quantum spinsystems derived in Ref. [9]. We then use

our method to calculate the *inverse* momentum- and frequency dependent spin susceptibility of a general class of quantum Heisenberg models in a magnetic field up to third order in the exchange couplings and use the result to estimate the critical temperatures for specific models where benchmarks are available. Finally, we use our algorithm to calculate the high-frequency behavior of the connected time-ordered correlation function of three different spin components to second order in the exchange couplings. This so-called *chiral non-linear susceptibility* [10] determines the quadratic response of the spin system to a time-dependent external magnetic field and exhibits a rather non-trivial frequency dependence which can in principle be measured experimentally.

II. RECURSIVE ALGORITHM FOR HIGH-TEMPERATURE EXPANSIONS OF CONNECTED SPIN CORRELATIONS

In this section we derive a new recursive algorithm for generating the high-temperature expansion of the free energy and arbitrary connected spin correlation functions of a general class of anisotropic quantum Heisenberg models with Hamiltonian

$$\mathcal{H} = \frac{1}{2} \sum_{ij} \sum_{ab} J_{ij}^{ab} S_i^a S_j^b - \sum_i \mathbf{H}_i \cdot \mathbf{S}_i, \quad (2.1)$$

where J_{ij}^{ab} are arbitrary anisotropic exchange couplings, \mathbf{H}_i is site-dependent external magnetic field in units of energy, and the spin-operators \mathbf{S}_i are normalized such that $\mathbf{S}_i^2 = S(S+1)$. The subscripts i and j label the positions \mathbf{R}_i , and \mathbf{R}_j of the spins while the superscripts a and b label their Cartesian components. At this point the spin positions \mathbf{R}_i are arbitrary, so that our expressions given below are also valid for finite spin clusters and systems where the spins are not located on a periodic lattice. For simplicity, we only consider spin systems

without single-site anisotropy so that we can set

$$J_{ii}^{ab} = 0. \quad (2.2)$$

It is straightforward to generalize our method to include spin models with finite single-site anisotropy. We assume that the local magnetic fields are aligned with the z -axis so that the part of the Hamiltonian representing the Zeeman energy can be written as

$$\mathcal{H}_0 = - \sum_i \mathbf{H}_i \cdot \mathbf{S}_i = - \sum_i H_i S_i^z. \quad (2.3)$$

For arbitrary \mathbf{H}_i we can locally rotate the coordinate system in spin space such that the rotated z axis points into the direction of the local magnetic field. Note that such a rotation re-defines the exchange couplings, but since we consider arbitrary J_{ij}^{ab} we do not lose any generality.

Our algorithm is based on the formally exact functional renormalization group (FRG) flow equation for the generating functional of the connected spin correlation functions in imaginary time [9]. For completeness, let us briefly outline the derivation of this flow equation. Following Ref. [9], we replace the exchange couplings by a deformation $J_{ij}^{ab} \rightarrow J_{ij,\Lambda}^{ab}$ depending on a continuous parameter Λ such that $J_{ij,\Lambda=0}^{ab} = 0$ and $J_{ij,\Lambda=1}^{ab} = J_{ij}^{ab}$. In

this work we satisfy this boundary condition via an interaction switch [9]

$$J_{ij,\Lambda}^{ab} = \Lambda J_{ij}^{ab}, \quad \Lambda \in [0, 1]. \quad (2.4)$$

In spite of its simplicity this deformation scheme has been useful in several recent applications of the spin FRG approach [11–15]. Following Ref. [9] we write the deformed generating functional $\mathcal{G}_\Lambda[\mathbf{h}]$ of the connected time-ordered spin correlation functions in imaginary time τ as the trace of a time-ordered exponential,

$$e^{\mathcal{G}_\Lambda[\mathbf{h}]} = \text{Tr} \left\{ e^{-\beta \mathcal{H}_0} \mathcal{T} e^{\int_0^\beta d\tau \sum_i \mathbf{h}_i(\tau) \cdot \mathbf{S}_i(\tau)} \times e^{-\int_0^\beta d\tau \frac{1}{2} \sum_{ij} \sum_{ab} J_{ij,\Lambda}^{ab} S_i^a(\tau) S_j^b(\tau)} \right\}. \quad (2.5)$$

Here β is the inverse temperature, \mathcal{T} denotes time-ordering in imaginary time, $\mathbf{h}_i(\tau)$ are fluctuating source fields, and the time dependence of all operators is in the interaction picture with respect to \mathcal{H}_0 . By differentiating Eq. (2.5) with respect to the deformation parameter Λ we obtain the exact flow equation [9]

$$\partial_\Lambda \mathcal{G}_\Lambda[\mathbf{h}] = -\frac{1}{2} \int_0^\beta d\tau \sum_{ij,ab} (\partial_\Lambda J_{ij,\Lambda}^{ab}) \left[\frac{\delta^2 \mathcal{G}_\Lambda[\mathbf{h}]}{\delta h_i^a(\tau) \delta h_j^b(\tau)} + \frac{\delta \mathcal{G}_\Lambda[\mathbf{h}]}{\delta h_i^a(\tau)} \frac{\delta \mathcal{G}_\Lambda[\mathbf{h}]}{\delta h_j^b(\tau)} \right]. \quad (2.6)$$

This is equivalent to an infinite hierarchy of flow equations for the connected time-ordered n -spin correlation functions $G_{i_1 \dots i_n, \Lambda}^{a_1 \dots a_n}(\tau_1, \dots, \tau_n)$, which are defined via the functional Taylor expansion of $\mathcal{G}_\Lambda[\mathbf{h}]$,

$$\mathcal{G}_\Lambda[\mathbf{h}] = \mathcal{G}_\Lambda[0] + \sum_{n=1}^{\infty} \frac{1}{n!} \int_0^\beta d\tau_1 \dots \int_0^\beta d\tau_n \sum_{i_1 \dots i_n} \sum_{a_1 \dots a_n} G_{i_1 \dots i_n, \Lambda}^{a_1 \dots a_n}(\tau_1, \dots, \tau_n) h_{i_1}^{a_1}(\tau_1) \dots h_{i_n}^{a_n}(\tau_n). \quad (2.7)$$

Omitting for simplicity the deformation label Λ the hierarchy of flow equations implied by the functional differential equation (2.6) can be written as [9]

$$\begin{aligned} \partial_\Lambda G_{i_1 \dots i_n}^{a_1 \dots a_n}(\tau_1, \dots, \tau_n) &= -\frac{1}{2} \int_0^\beta d\tau \sum_{ij,ab} (\partial_\Lambda J_{ij,\Lambda}^{ab}) \left[G_{i_1 \dots i_n ij}^{a_1 \dots a_n ab}(\tau_1, \dots, \tau_n, \tau, \tau) \right. \\ &\quad \left. + \sum_{m=0}^n \mathcal{S}_{1, \dots, m; m+1, \dots, n} \left\{ G_{i_1 \dots i_m i}^{a_1 \dots a_m a}(\tau_1, \dots, \tau_m, \tau) G_{i_{m+1} \dots i_n j}^{a_{m+1} \dots a_n b}(\tau_{m+1}, \dots, \tau_n, \tau) \right\} \right]. \end{aligned} \quad (2.8)$$

Here the symmetrization operator $\mathcal{S}_{1, \dots, m; m+1, \dots, n} \{ \dots \}$ symmetrizes the expression in the curly braces with respect to the exchange of all labels [16]. Explicitly, the action of this operator on a function $f(1, \dots, n)$ which is already symmetric with respect to the first m labels

$1, \dots, m$ and the last $n - m$ labels $m + 1, \dots, n$ is

$$\begin{aligned} &\mathcal{S}_{1, \dots, m; m+1, \dots, n} \{ f(1, \dots, n) \} \\ &= \frac{1}{m!(n-m)!} \sum_P f(P_1, \dots, P_n), \end{aligned} \quad (2.9)$$

where \sum_P is the sum of all $n!$ permutations (P_1, \dots, P_n) of $(1, \dots, n)$. Note that the number of terms generated by

$\mathcal{S}_{1,\dots,m;m+1,\dots,n} \{ \dots \}$ is given by the binomial coefficient

$$\binom{n}{m} = \frac{n!}{m!(n-m)!}. \quad (2.10)$$

A graphical representation of Eq. (2.8) can be found in Fig. 1 of Ref. [9]. At this point it is useful to introduce multi-labels $\alpha = (i, \tau, a)$ representing all parameters which are necessary to specify the degrees of freedom, i.e. lattice site, imaginary time, and field component. Our FRG flow equation (2.8) can then be written as

$$\begin{aligned} \partial_\Lambda G_{\alpha_1 \dots \alpha_n}^{(n)} &= -\frac{1}{2} \int_\alpha \int_{\alpha'} [(\partial_\Lambda \mathbf{J}_\Lambda)]_{\alpha\alpha'} \left[G_{\alpha\alpha'\alpha_1 \dots \alpha_n}^{(n+2)} \right. \\ &+ \left. \sum_{m=0}^n \mathcal{S}_{1,\dots,m;m+1,\dots,n} \left\{ G_{\alpha\alpha_1 \dots \alpha_m}^{(m+1)} G_{\alpha'\alpha_{m+1} \dots \alpha_n}^{(n-m+1)} \right\} \right], \end{aligned} \quad (2.11)$$

where $\int_\alpha = \int_0^\beta d\tau \sum_{ia}$ and \mathbf{J}_Λ is a matrix in the multi-labels with matrix elements

$$[\mathbf{J}_\Lambda]_{\alpha=(i\tau a), \alpha'=(j\tau' b)} = \delta(\tau - \tau') J_{ij, \Lambda}^{ab}. \quad (2.12)$$

The hierarchy (2.11) of flow equations can now be used to generate a systematic expansion of the connected spin correlation functions in powers of the exchange couplings, which amounts to a high-temperature expansion. The crucial observation is that with the interaction switch deformation scheme (2.4) an expansion in powers of Λ amounts to an expansion in powers of J/T [15]. To generate the high-temperature expansion, we simply substitute the Taylor expansion

$$G_{\alpha_1 \dots \alpha_n, \Lambda}^{(n)} = \sum_{k=0}^{\infty} \Lambda^k G_{\alpha_1 \dots \alpha_n}^{(n,k)} \quad (2.13)$$

into the flow equation (2.11) and compare powers of Λ on both sides. This yields the recursion relation,

$$\begin{aligned} G_{\alpha_1 \dots \alpha_n}^{(n,k)} &= -\frac{1}{2k} \int_\alpha \int_{\alpha'} J_{\alpha\alpha'} \left[G_{\alpha\alpha'\alpha_1 \dots \alpha_n}^{(n+2,k-1)} + \right. \\ &\left. \sum_{m=0}^n \mathcal{S}_{1,\dots,m;m+1,\dots,n} \left\{ \sum_{l=0}^{k-1} G_{\alpha\alpha_1 \dots \alpha_m}^{(m+1,l)} G_{\alpha'\alpha_{m+1} \dots \alpha_n}^{(n-m+1,k-l-1)} \right\} \right], \end{aligned} \quad (2.14)$$

which is the central result of this work. Keeping in mind that for $\Lambda = 1$ we recover our original Heisenberg model, the high-temperature expansion of the physical connected n -spin correlation function is

$$G_{\alpha_1 \dots \alpha_n}^{(n)} = \sum_{k=0}^{\infty} G_{\alpha_1 \dots \alpha_n}^{(n,k)}. \quad (2.15)$$

To define a graphical representation of the recursion relation we introduce the graphical elements shown in Fig. 1. With this notation our recursion recursion relation (2.14)




FIG. 1. Elements of our graphical representation of the perturbation series generated by the iteration of Eq. (2.14). The exact connected n -spin correlation function $G_{\alpha_1 \dots \alpha_n}^{(n)}$ is represented by a large empty circle with n labeled external legs. A number k inside the circle denotes the contribution $G_{\alpha_1 \dots \alpha_n}^{(n,k)}$ of order J^k to $G_{\alpha_1 \dots \alpha_n}^{(n)}$. The exchange interaction is represented by a (red) wavy line. Circles of varying size denote the corresponding connected spin correlation functions for $J = 0$, where the dots represent the external labels. These quantities correspond precisely to the generalized blocks introduced by Izyumov and Skryabin [17] who used the same graphical notation.

can be represented graphically as shown in Fig. 2.

The exact recursion relation (2.14) expresses the connected n -point function $G^{(n,k)}$ at order $(J/T)^k$ in terms of the connected m -point functions $G^{(m,l)}$ with $1 \leq m \leq n+2$ and $0 \leq l \leq k-1$. Each iteration lowers the order k of the expansion by one but raises the number n of external legs by two. After k iterations all terms generated by this recursion can be expressed in terms of the connected correlation functions $G^{(m,0)}$ of an isolated spin, i.e., the *generalized blocks* introduced by Izyumov and Skryabin [17]. Following their notation, we represent the generalized blocks graphically by circles of varying size with black dots representing the external labels as shown in Fig. 1. Given the fact that for coinciding lattice sites $J_{ii}^{ab} = 0$, we see that the contribution from the term involving $n + 2k$ external legs vanishes,

$$\int_\alpha \int_{\alpha'} J_{\alpha\alpha'} G_{\alpha\alpha'\alpha_1 \dots \alpha_n}^{(n+2k,0)} = 0. \quad (2.16)$$

By iteration we can thus express an arbitrary connected n -point function $G^{(n,k)}$ at order $(J/T)^k$ in terms of the connected correlation functions $G^{(m,0)}$ involving up to




FIG. 2. (a) Graphical representation of the exact recursion formula (2.14) relating the connected n -spin correlation function of order k in terms of a sum of connected spin correlation functions involving up to $n+2$ external legs but at most $k-1$ powers of the exchange couplings. (b) Special case of (a) for the free energy which is represented by an empty circle without external legs. In this case we may omit the symmetrization operator because there are no external labels to be symmetrized. Our recursion formula (2.14) then relates the free energy at order k to the connected two-point function at order $k-1$ and a sum of products of one-point functions (magnetic moments) of order $k-l-1$ and l with $l=0, \dots, k-1$.

$m = n + 2k - 1$ components of an isolated spin. Keeping in mind that for fixed m the symmetrization operator $\mathcal{S}_{1, \dots, m; m+1, \dots, n}$ in the second line of Eq. (2.14) generates $n!/(m!(n-m)!)$ different terms corresponding to distinct permutations of the external labels, the total number of terms generated the nested sum in the second line of Eq. (2.14) is

$$\sum_{m=0}^n \sum_{l=0}^{k-1} \binom{n}{m} = 2^n k. \quad (2.17)$$

All combinatorial factors are automatically generated by the symmetrization operator $\mathcal{S}_{1, \dots, m; m+1, \dots, n}$ so that the lowest few terms in the high-temperature expansion of the free energy and arbitrary connected correlation functions can be easily obtained purely algebraically without additional graph theoretical or combinatorial considerations. Of course, the explicit calculation of high orders in this expansion requires a numerical implementation of the recursion relation (2.14) which is beyond the scope of this work. In fact, at this point it is not clear whether our recursive algorithm offers any computational advan-

tages over the established methods [1, 2] for generating the high-temperature expansions of Ising or Heisenberg models.

III. TRUNCATED HIGH-TEMPERATURE EXPANSIONS

The first few terms in the high temperature expansions of the free energy and the connected n -point correlation functions for a general spin Hamiltonian of the form (2.1) can be obtained analytically by straightforward iteration of our recursion relation (2.14). To that end, it is convenient to transform the imaginary-time correlation functions to frequency space. We normalize the Fourier transform of the connected n -spin correlation function as follows [11],

$$G_{i_1 \dots i_n}^{a_1 \dots a_n}(\tau_1, \dots, \tau_n) = \frac{1}{\beta^n} \sum_{\omega_1 \dots \omega_n} e^{-i(\omega_1 \tau_1 + \dots + \omega_n \tau_n)} \times \beta \delta_{\omega_1 + \dots + \omega_n, 0} G_{i_1 \dots i_n}^{a_1 \dots a_n}(\omega_1, \dots, \omega_n), \quad (3.1)$$

where we have used time-translational invariance to factorize a frequency-conserving Kronecker- δ . For two-spin correlation functions we suppress the redundant second frequency label. In the atomic limit $J \rightarrow 0$ the corresponding correlation functions are diagonal in the site index,

$$\lim_{J \rightarrow 0} G_{i_1 \dots i_n}^{a_1 \dots a_n}(\omega_1, \dots, \omega_n) = \delta_{i_1 i_2} \delta_{i_2 i_3} \dots \delta_{i_{n-1} i_n} g_{i_1}^{a_1 \dots a_n}(\omega_1, \dots, \omega_n), \quad (3.2)$$

where the site-dependence of $g_{i_1}^{a_1 \dots a_n}(\omega_1, \dots, \omega_n)$ is due to the inhomogeneous external magnetic field H_i . For a homogeneous field $H_i = H$ the connected single-spin correlation functions $g_{i_1}^{a_1 \dots a_n}(\omega_1, \dots, \omega_n) = g^{a_1 \dots a_n}(\omega_1, \dots, \omega_n)$ are independent of the site labels. Following Ref. [17] we refer to the functions $g^{a_1 \dots a_n}(\omega_1, \dots, \omega_n)$ as *generalized blocks* (or simply *blocks*) and represent them graphically as shown in the lower part of Fig. 1. Explicit expressions for the blocks involving up to $n=4$ spin components have already been derived long time ago by Vaks, Larkin, and Pikin [18, 19], and can also be found in the textbook [17]. A recursive algorithm for calculating the blocks for arbitrary n has been constructed in Ref. [11], see also Ref. [20]. In Appendix A we explicitly give the blocks involving up to $n=5$ spin components. As will become evident below, the five-spin blocks are necessary to obtain the two-point function to order J^3 and the three-point function to order J^2 .

A. Free energy and magnetization

Let us now calculate the high-temperature expansion of the negative free energy $G^{(0)} = -\beta F$ in units of temperature up to third order in J . To generate this expansion we set $n=0$ in Eq. (2.14) and iterate up to order

$k = 3$, which can be easily done analytically. A graphical representation of the terms in this expansion up to order J^3 is shown in Fig. 3. These graphs represent the




FIG. 3. Diagrammatic representation of the high-temperature series for the negative free energy $G^{(0)} = -\beta F$ in units of temperature up to order J^3 . The symbols are defined in Fig. 1.

following mathematical expression,

$$G^{(0)} = \sum_i B(\beta H_i) - \frac{\beta}{2} \sum_{ij} J_{ij}^{zz} m_i m_j + G^{(0,2)} + G^{(0,3)} + \mathcal{O}(J^4), \quad (3.3)$$

where

$$B(y) = \ln \left[\frac{\sinh[(S + 1/2)y]}{\sinh(y/2)} \right] \quad (3.4)$$

is the primitive integral of the spin- S Brillouin function

$$b(y) = \left(S + \frac{1}{2} \right) \coth \left[\left(S + \frac{1}{2} \right) y \right] - \frac{1}{2} \coth \left[\frac{y}{2} \right], \quad (3.5)$$

which gives the local magnetic moments for vanishing exchange interaction,

$$m_i = b(\beta H_i). \quad (3.6)$$

For later reference we note that for small $y = \beta H$ the Brillouin function has the expansion

$$b(y) = b_1 y + \frac{b_3}{3!} y^3 + \mathcal{O}(y^5), \quad (3.7)$$

with

$$b_1 = \frac{(2S + 1)^2 - 1}{12} = \frac{S(S + 1)}{3}, \quad (3.8a)$$

$$b_3 = -\frac{(2S + 1)^4 - 1}{120} = -\frac{6}{5} b_1 \left(b_1 + \frac{1}{6} \right). \quad (3.8b)$$

The Cartesian components of the connected two-spin correlation function of a isolated spin are in frequency space

$$g_i^{xx}(\omega) = g_i^{yy}(\omega) = \frac{m_i H_i}{H_i^2 + \omega^2}, \quad (3.9a)$$

$$g_i^{xy}(\omega) = -g_i^{yx}(\omega) = -\frac{m_i \omega}{H_i^2 + \omega^2}, \quad (3.9b)$$

$$g_i^{zz}(\omega) = \beta \delta_{\omega,0} b'(\beta H_i), \quad (3.9c)$$

where $b'(y)$ is the derivative of the Brillouin function. For finite magnetic field it is more convenient to work with the spherical spin components,

$$S_i^+ = \frac{S_i^x + i S_i^y}{\sqrt{2}}, \quad (3.10a)$$

$$S_i^- = \frac{S_i^x - i S_i^y}{\sqrt{2}}, \quad (3.10b)$$

$$S_i^0 = S_i^z. \quad (3.10c)$$

In this basis the Heisenberg equations of motion for the spin components decouple,

$$\partial_\tau S_i^p(\tau) = -p H_i S_i^p(\tau), \quad \text{where } p = +, -, 0, \quad (3.11)$$

implying

$$S_i^p(\tau) = e^{-p H_i \tau} S_i^p. \quad (3.12)$$

In frequency space the transverse connected two-spin correlation function in the spherical basis is therefore

$$g_i^{+-}(\omega) = g_i^{-+}(-\omega) = \frac{m_i}{H_i - i\omega}. \quad (3.13)$$

Note that for $H_i \rightarrow 0$ the functions $g_i^{zz}(\omega)$, $g_i^{+-}(\omega)$, and $g_i^{-+}(\omega)$ approach the same limit

$$\lim_{H_i \rightarrow 0} g_i^{p\bar{p}}(\omega) = g_0(\omega) = \beta \delta_{\omega,0} b_1, \quad (3.14)$$

where $\bar{p} = -p$. With the above notation the second-order correction to the negative free energy in units of temperature can be written as

$$G^{(0,2)} = \frac{1}{4} \sum_{ij} \sum_{abcd} J_{ij}^{ab} J_{ij}^{cd} \sum_{\omega} g_i^{ac}(\omega) g_j^{db}(\omega) + \frac{\beta}{2} \sum_{ijk} \sum_{ab} J_{ij}^{za} J_{jk}^{bz} g_j^{ab}(0) m_i m_k. \quad (3.15)$$

The third-order correction $G^{(0,3)}$ is rather lengthy and is explicitly given in Eq. (B1) of Appendix B.

In general, the contribution $G^{(0,k)}$ to the dimensionless free energy of order J^k depends on generalized blocks $g_i^{a_1 \dots a_l}(\omega_1, \dots, \omega_l)$ with up to $l = k$ external legs. From the high-temperature expansion of the free energy $F = -TG^{(0)}$ we can obtain the corresponding expansion of the local magnetization M_i to the same order in J by taking a derivative with respect to the magnetic field,

$$M_i \equiv -\frac{\partial F}{\partial H_i} = m_i - \frac{\partial m_i}{\partial H_i} \sum_j J_{ij}^{zz} m_j + \frac{1}{\beta} \frac{\partial G^{(0,2)}}{\partial H_i} + \frac{1}{\beta} \frac{\partial G^{(0,3)}}{\partial H_i} + \mathcal{O}(J^4). \quad (3.16)$$

This expression gives the magnetic equation of state $M_i = M_i(H, T)$ of spin models with Hamiltonian of the type (2.1) at high temperatures up to order J^3 . The explicit evaluation of Eq. (3.16) for specific models such as the Kitaev-Heisenberg- Γ model [21], effective spin models for altermagnets [22, 23] or other spin models of current interest is straightforward but tedious and is not the subject of this work.

B. Dynamic spin susceptibility

Next, we calculate the high-temperature expansion of the connected two-spin correlation function up to order J^3 ,

$$G_{ij}^{ab}(\omega) = \delta_{ij} g_i^{ab}(\omega) + G_{ij}^{ab(1)}(\omega) + G_{ij}^{ab(2)}(\omega) + G_{ij}^{ab(3)}(\omega) + \mathcal{O}(J^4). \quad (3.17)$$

The terms contributing to the expansion (3.17) up to order J^2 are shown diagrammatically in Fig. 4 (a). The mathematical expression for the first order correction is

$$G_{ij}^{ab(1)}(\omega) = -\sum_{cd} g_i^{ac}(\omega) J_{ij}^{cd} g_j^{db}(\omega) - \delta_{ij} \sum_k \sum_c g_i^{abc}(\omega, -\omega, 0) J_{ik}^{cz} m_k. \quad (3.18)$$

The expressions for the seven second-order diagrams shown in Fig. 4 (a) are given in Eq. (B4) of Appendix B. The explicit evaluation of these expressions is rather tedious. To reduce the complexity, let us focus here on the case of vanishing magnetic field where all tadpole diagrams vanish and the spherical components of the two-point function are given by $g_0(\omega) = \beta \delta_{\omega,0} b_1$, see Eq. (3.14). Moreover, in this limit only two of the diagrams in Fig. 4 (a) can be separated into two parts by cutting a single interaction line; these are the interaction-reducible diagrams. Formally, the interaction-irreducible part $\Pi_{\alpha\alpha'}$ of the two-spin correlation function $G_{\alpha\alpha'} \equiv G_{\alpha\alpha'}^{(2)}$ can be defined by setting

$$G_{\alpha\alpha'} = [\mathbf{\Pi}(1 + \mathbf{J}\mathbf{\Pi})^{-1}]_{\alpha\alpha'} = [\mathbf{\Pi}^{-1} + \mathbf{J}]_{\alpha\alpha'}^{-1}, \quad (3.19)$$

where all quantities on the right-hand side are matrices in the multi-labels. In Fig. 4 (b) we show the diagrams contributing to the interaction-irreducible part $\Pi_{\alpha\alpha'}$ of the susceptibility up to second order in J , while the third-order diagrams are shown in Fig. 5. For simplicity, let us focus now on an isotropic Heisenberg magnet where $J_{ij}^{ab} = \delta^{ab} J_{ij}$ on a Bravais lattice with N sites. Then the matrix equation (3.19) can be diagonalized in momentum space so that the Fourier transform of the two-spin correlation function can be written as

$$G(\mathbf{k}, \omega) = \frac{\mathbf{\Pi}(\mathbf{k}, \omega)}{1 + J_{\mathbf{k}} \mathbf{\Pi}(\mathbf{k}, \omega)} = \frac{1}{J_{\mathbf{k}} + \mathbf{\Pi}^{-1}(\mathbf{k}, \omega)}, \quad (3.20)$$

where

$$J_{\mathbf{k}} = \frac{1}{N} \sum_{ij} e^{-i\mathbf{k} \cdot (\mathbf{R}_i - \mathbf{R}_j)} J_{ij} \quad (3.21)$$

is the Fourier transform of the exchange couplings. Evaluating the diagrams in Figs. 4 and 5 for vanishing mag-




FIG. 4. (a) Expansion of the connected two-spin correlation function $G_{\alpha\alpha'}^{(2)}$ up to second order in the exchange interaction. (b) Expansion of the interaction-irreducible part $\Pi_{\alpha\alpha'}$ of the connected two-spin correlation function up to second order in the exchange interaction for vanishing magnetic field.

netic field we obtain




FIG. 5. Diagrammatic representation of the third-order contribution $\Pi_{\alpha\alpha'}^{(3)}$ to the interaction-irreducible susceptibility.

$$\Pi(\mathbf{k}, \omega) = \beta\delta_{\omega,0}b_1 + \Pi^{(2)}(\mathbf{k}, \omega) + \Pi^{(3)}(\mathbf{k}, \omega) + \mathcal{O}(J^4), \quad (3.22)$$

where the second-order term is

$$\begin{aligned} \Pi^{(2)}(\mathbf{k}, \omega) &= \frac{1}{2\beta N} \sum_{\mathbf{q}} J_{\mathbf{q}}^2 \sum_{\omega'} g_0(\omega') [2g_0^{+-zz}(\omega', -\omega', \omega, -\omega) + g_0^{zzzz}(\omega', -\omega', \omega, -\omega)] \\ &+ \frac{1}{\beta N} \sum_{\mathbf{q}} J_{\mathbf{q}} J_{\mathbf{q}+\mathbf{k}} \sum_{\omega'} g_0^{+-z}(\omega', -\omega - \omega', \omega) g_0^{+-z}(\omega + \omega', -\omega', -\omega). \end{aligned} \quad (3.23)$$

Here the zero-field limits of the generalized blocks are denoted by $g_0(\omega)$, $g_0^{+-z}(\omega_1, \omega_2, \omega_3)$ and similarly for the higher-order blocks. Recall that according to Eq. (3.14) the two-point function for vanishing magnetic field is $g_0(\omega) = \beta\delta_{\omega,0}b_1$. Explicit expressions for the relevant generalized blocks are given in Appendix A. The frequency sums can be evaluated analytically and we obtain for vanishing magnetic field

$$\Pi^{(2)}(\mathbf{k}, \omega) = \delta_{\omega,0} \frac{\beta^3}{N} \sum_{\mathbf{q}} \left[\frac{5b_1b_3}{6} J_{\mathbf{q}}^2 - \frac{b_1^2}{12} J_{\mathbf{q}} J_{\mathbf{q}+\mathbf{k}} \right] + (1 - \delta_{\omega,0}) \frac{\beta}{\omega^2} \frac{2b_1^2}{N} \sum_{\mathbf{q}} J_{\mathbf{q}} (J_{\mathbf{q}} - J_{\mathbf{q}+\mathbf{k}}), \quad (3.24)$$

where b_1 and b_3 are defined in Eq. (3.8). The third-order contribution to the irreducible susceptibility is represented by the four diagrams in Fig. 5. The explicit mathematical expression represented by these diagrams is given in Eq. (B7) of Appendix B. After performing the frequency sums we obtain for vanishing magnetic field

$$\begin{aligned} \Pi^{(3)}(\mathbf{k}, \omega) &= -\delta_{\omega,0} \beta^4 \left\{ \frac{1}{72N^2} \sum_{\mathbf{q}_1 \mathbf{q}_2} [(b_1^2 - 10b_1b_3) J_{\mathbf{q}_1} J_{\mathbf{q}_2} J_{\mathbf{q}_1+\mathbf{q}_2} + (b_1^2 + 20b_3^2) J_{\mathbf{q}_1} J_{\mathbf{q}_2} J_{\mathbf{q}_1+\mathbf{q}_2+\mathbf{k}}] + \frac{5b_1^2b_3}{6N} \sum_{\mathbf{q}} J_{\mathbf{q}}^3 \right\} \\ &+ (1 - \delta_{\omega,0}) \frac{\beta^2}{\omega^2} \left\{ \frac{b_1^2}{2N^2} \sum_{\mathbf{q}_1 \mathbf{q}_2} J_{\mathbf{q}_1} J_{\mathbf{q}_2} (J_{\mathbf{q}_1+\mathbf{q}_2} - J_{\mathbf{q}_1+\mathbf{q}_2+\mathbf{k}}) - \frac{2b_1^3}{N} \sum_{\mathbf{q}} J_{\mathbf{q}}^2 (J_{\mathbf{q}} - J_{\mathbf{q}+\mathbf{k}}) \right\}. \end{aligned} \quad (3.25)$$

Note that the exact identity $\Pi(\mathbf{k} = 0, \omega \neq 0) = 0$ implied by the conservation of the total spin is satisfied order by order in perturbation theory. It is also straightforward to show that the sum rule

$$\frac{1}{\beta N} \sum_{\mathbf{k}\omega} G(\mathbf{k}, \omega) = b_1 = \frac{S(S+1)}{3} \quad (3.26)$$

implied by the spin-length constraint $\mathbf{S}_i^2 = S(S+1)$ is likewise satisfied order by order in perturbation theory.

For a Heisenberg model with nearest-neighbor exchange J on a D -dimensional hypercubic lattice with

lattice spacing a the Fourier transform of the exchange couplings is $J_{\mathbf{q}} = J_0 \gamma_{\mathbf{q}}$ where $J_0 = 2DJ$ and the nearest-neighbor form factor is

$$\gamma_{\mathbf{q}} = \frac{1}{D} \sum_{\mu=1}^D \cos(q_{\mu}a). \quad (3.27)$$

Using

$$\frac{1}{N} \sum_{\mathbf{q}} \gamma_{\mathbf{q}} \gamma_{\mathbf{q}+\mathbf{k}} = \frac{\gamma_{\mathbf{k}}}{2D}, \quad (3.28a)$$

$$\frac{1}{N} \sum_{\mathbf{q}} \gamma_{\mathbf{q}}^2 \gamma_{\mathbf{q}+\mathbf{k}} = 0, \quad (3.28b)$$

$$\frac{1}{N^2} \sum_{\mathbf{q}_1 \mathbf{q}_2} \gamma_{\mathbf{q}_1} \gamma_{\mathbf{q}_2} \gamma_{\mathbf{q}_1+\mathbf{q}_2+\mathbf{k}} = \frac{\gamma_{\mathbf{k}}}{(2D)^2}, \quad (3.28c)$$

we obtain for the second-order term

$$\begin{aligned} \Pi^{(2)}(\mathbf{k}, \omega) = & \delta_{\omega,0} \frac{\beta^3 J_0^2}{2D} \left[\frac{5b_1 b_3}{6} - \frac{b_1^2 \gamma_{\mathbf{k}}}{12} \right] \\ & + (1 - \delta_{\omega,0}) \frac{\beta J_0^2}{2D} 2b_1^2 \frac{1 - \gamma_{\mathbf{k}}}{\omega^2}, \end{aligned} \quad (3.29)$$

and for the third-order term

$$\begin{aligned} \Pi^{(3)}(\mathbf{k}, \omega) = & -\delta_{\omega,0} \frac{\beta^4 J_0^3}{(2D)^2} \left[\frac{b_1^2 - 10b_1 b_3}{72} + \frac{(b_1^2 + 20b_3^2) \gamma_{\mathbf{k}}}{72} \right] \\ & + (1 - \delta_{\omega,0}) \frac{\beta^2 J_0^3}{(2D)^2} \frac{b_1^2}{2} \frac{1 - \gamma_{\mathbf{k}}}{\omega^2}. \end{aligned} \quad (3.30)$$

Comparing our perturbative results at finite frequency ω with high-temperature expansions for the short-time ($|J|t \ll 1$) evolution of the Kubo relaxation function, we find that our results for $\Pi^{(2)}(\mathbf{k}, \omega)$ and $\Pi^{(3)}(\mathbf{k}, \omega)$ are consistent with the first two terms in the corresponding second order moment [24, 25].

In the static limit $\omega \rightarrow 0$ we may use the above results to estimate the critical temperature T_c where the system develops long-range magnetic order. At $T = T_c$ the static susceptibility $G(\mathbf{Q}, 0)$ diverges at the ordering wave-vector \mathbf{Q} . At this temperature

$$1 + J_{\mathbf{Q}} \Pi(\mathbf{Q}, 0) = 0, \quad (3.31)$$

or equivalently

$$J_{\mathbf{Q}} + \Pi^{-1}(\mathbf{Q}, 0) = 0. \quad (3.32)$$

If $\Pi(\mathbf{Q}, 0)$ is only known up to some finite order in J , these conditions are not equivalent if we consistently expand Eqs. (3.31) and (3.32) up to this order. For example, if we expanding the irreducible static susceptibility up to third order in J ,

$$\Pi(\mathbf{k}, 0) = \beta b_1 + \Pi^{(2)}(\mathbf{k}, 0) + \Pi^{(3)}(\mathbf{k}, 0) + \mathcal{O}(J^4), \quad (3.33)$$

the corresponding expansion of the inverse susceptibility is

$$\Pi^{-1}(\mathbf{k}, 0) = \frac{1}{\beta b_1} - \frac{\Pi^{(2)}(\mathbf{k}, 0)}{(\beta b_1)^2} - \frac{\Pi^{(3)}(\mathbf{k}, 0)}{(\beta b_1)^2} + \mathcal{O}(J^4). \quad (3.34)$$

Using this truncated expansion to estimate the critical temperature T_c in various cases where controlled benchmarks are available, we obtain the results summarized in Table I and in Fig. 6. It turns out that the

S	J	T_c/T_{c0}			rel. error / %	
		J^2	J^3	benchmark	J^2	J^3
1/2	< 0	-	0.667	0.559	-	19.3
1/2	> 0	0.667	0.667	0.629	6	6
1	< 0	0.645	0.741	0.650	0.8	14
1	> 0	0.750	0.771	0.684	9.6	12.7
3/2	< 0	0.724	0.773	0.685	5.7	12.8
3/2	> 0	0.769	0.791	0.702	9.5	12.7
∞	$\neq 0$	0.788	0.810	0.722	9.1	12.2
$\infty(d=4)$	$\neq 0$	0.853	0.862	0.822	3.7	4.9

TABLE I. Critical temperatures T_c in units of the mean-field critical temperature $T_{c0} = b_1 |J_{\mathbf{Q}}|$ of spin- S quantum Heisenberg ferromagnets ($J < 0$, $\mathbf{Q} = 0$) and antiferromagnets ($J > 0$, $\mathbf{Q} = (\frac{\pi}{a}, \frac{\pi}{a}, \frac{\pi}{a})$) with nearest neighbor exchange J . With the exception of the last row all temperatures are for a three-dimensional simple cubic lattice. The values in the third and fourth columns are obtained by inserting the high-temperature expansion (3.34) of $\Pi^{-1}(\mathbf{Q}, 0)$ at order J^2 or J^3 into Eq. (3.32). The benchmarks are obtained using quantum Monte Carlo simulations [27, 28] ($S = 1/2$) and a more sophisticated high-temperature series expansion [29]. The benchmark in the classical limit ($S = \infty$) in dimension $d = 4$ has been obtained via high-temperature series expansion [30].




FIG. 6. Graphical representation of the results for the critical temperatures of nearest-neighbor ferromagnets or antiferromagnets in a simple cubic lattice listed in Table I. The solid lines are obtained from the condition (3.32) using the order- J^3 truncated high-temperature expansion for the inverse static spin susceptibility given in Eq. (3.34). The crosses and squares represent the benchmarks mentioned in the caption of Table I and additional values for $S \geq 2$ are taken from Ref. [31].

truncated expansion (3.32) of the inverse susceptibility yields more accurate estimates for T_c than Eq. (3.31), which has already been noticed by Krieg [26]. In fact, if we substitute the truncated high-temperature expansion (3.33) for $\Pi(\mathbf{Q}, 0)$ into Eq. (3.31), we find that for three-dimensional spin- S Heisenberg ferromagnets or an-

tiferromagnets with nearest-neighbor coupling the resulting equation does not have any solutions, so that at this level of approximation we miss the magnetic instability. On the other hand, if we use the condition (3.32) involving $\Pi^{-1}(\mathbf{Q}, 0)$ and substitute the corresponding truncated high-temperature expansion for $\Pi^{-1}(\mathbf{Q}, 0)$ given in Eq. (3.34) we obtain the expected magnetic instabilities for arbitrary S , although for a $S = 1/2$ ferromagnet the J^3 -term is crucial to reproduce the instability and T_c is roughly 20% larger than the established result. Note also that for $S \geq 1$ the J^2 -truncation produces better estimates for T_c than the J^3 -truncation, indicating the alternating nature of the high-temperature series of the inverse susceptibility. A plausible reason why estimates of T_c based on a low-order truncated expansion of $\Pi^{-1}(\mathbf{Q}, 0)$ produce better results than the corresponding expansion of $\Pi(\mathbf{Q}, 0)$ is that in the former case we expand the physical inverse spin susceptibility in powers of J/T , whereas $\Pi(\mathbf{k}, 0)$ is an auxiliary quantity that cannot be directly measured. Improved estimates for T_c can be obtained with moderate numerical effort by solving truncated spin-FRG flow equations [14]. In this case it likewise turns out to be crucial to consider the flow of the physical inverse static spin susceptibility $\Pi^{-1}(\mathbf{k}, 0)$, which entails a hybrid approach treating static and dynamic fluctuations differently [13–15]. Note finally that our perturbative approach correctly reproduces the fact that for any finite S the critical temperature of an antiferromagnet is always larger than the critical temperature of a ferromagnetic with the same value of $|J|$.

Another limit where the critical temperature can be calculated systematically is the limit of high dimensions [9, 26]. In fact, by comparing the second-order correction $\Pi^{(2)}(\mathbf{k}, \omega)$ in Eq. (3.29) with the third-order correction $\Pi^{(3)}(\mathbf{k}, \omega)$ in Eq. (3.30), we see that in high dimensions successive orders in the high-temperature expansion are additionally controlled by the small parameter $1/(2D)$, as already pointed out in Refs. [9 and 26]. However, to collect all contributions to $\Pi^{-1}(\mathbf{k}, 0)$ of order $1/(2D)^2$ we have to take into account contributions up to order J^4 in the high-temperature expansion of $\Pi(\mathbf{k}, 0)$ [26].

C. Chiral non-linear susceptibility

With our recursive algorithm it is straightforward to obtain the high-temperature expansion of arbitrary connected spin correlation functions. Of particular interest is the connected three-spin correlation function, which determines the quadratic response of the magnetization to a time-dependent external magnetic field [10]. In Fig. 7 we show all diagrams contributing to the connected three-spin correlation function up to order J^2 . For simplicity let us now assume that the spins are located on a Bravais lattice and are exposed to homogeneous magnetic field H in z -direction, so that it is again convenient to work in a spherical basis. In momentum-frequency space the high-temperature expansion of the chiral three-spin




FIG. 7. Diagrammatic representation of the connected three-spin correlation function up to order J^2 .

correlation function is

$$\begin{aligned}
 G^{+-z}(K_1, K_2, K_3) &= g^{+-z}(\omega_1, \omega_2, \omega_3) \\
 &+ G^{+-z(1)}(K_1, K_2, K_3) \\
 &+ G^{+-z(2)}(K_1, K_2, K_3) + \mathcal{O}(J^3),
 \end{aligned}
 \tag{3.35}$$

where $K_i = (\mathbf{k}_i, \omega_i)$ are multi-labels for momentum and Matsubara frequency and it is understood that both sides of Eq. (3.35) should be multiplied by a factor of $\delta_{\mathbf{k}_1 + \mathbf{k}_2 + \mathbf{k}_3, 0} \delta_{\omega_1 + \omega_2 + \omega_3, 0}$ implementing conservation of momentum and frequency. The zeroth-order term $g^{+-z}(\omega_1, \omega_2, \omega_3)$ is given by the generalized block involving three different spin components defined in Eq. (A1). In the limit $H \rightarrow 0$ the chiral three-point function can be written in the manifestly symmetric form [11, 32],

$$\begin{aligned}
 g_0^{+-z}(\omega_1, \omega_2, \omega_3) &= -\beta b_1 (1 - \delta_{\omega_1, 0} \delta_{\omega_2, 0} \delta_{\omega_3, 0}) \\
 &\times \left[\frac{\delta_{\omega_1, 0}}{i\omega_2} + \frac{\delta_{\omega_2, 0}}{i\omega_3} + \frac{\delta_{\omega_3, 0}}{i\omega_1} \right],
 \end{aligned}
 \tag{3.36}$$

where the subscript indicates the zero-field limit and $b_1 = S(S+1)/3$ has already been introduced in Eq. (3.8a).

Note that for $H \rightarrow 0$ the purely longitudinal three-point function g_0^{zzz} vanishes so that in a Cartesian basis all non-zero components of the three-point function can be written in terms of the three-dimensional Levi-Civita symbol $\epsilon^{a_1 a_2 a_3}$ as follows,

$$\begin{aligned} G^{a_1 a_2 a_3}(K_1, K_2, K_3) &= \epsilon^{a_1 a_2 a_3} G^{xyz}(K_1, K_2, K_3) \\ &= -i \epsilon^{a_1 a_2 a_3} G^{+-z}(K_1, K_2, K_3), \end{aligned} \quad (3.37)$$

where $\epsilon^{xyz} = 1$. In particular, for an isolated spin we obtain for vanishing magnetic field

$$\begin{aligned} g_0^{a_1 a_2 a_3}(\omega_1, \omega_2, \omega_3) &= \epsilon^{a_1 a_2 a_3} \beta b_1 (1 - \delta_{\omega_1, 0} \delta_{\omega_2, 0} \delta_{\omega_3, 0}) \\ &\quad \times \left[\frac{\delta_{\omega_1, 0}}{\omega_2} + \frac{\delta_{\omega_2, 0}}{\omega_3} + \frac{\delta_{\omega_3, 0}}{\omega_1} \right], \end{aligned} \quad (3.38)$$

which is manifestly symmetric under the exchange of any pair of labels $1 \leftrightarrow 2$, $1 \leftrightarrow 3$ or $2 \leftrightarrow 3$. The frequency dependence of $g_0^{xyz}(\omega_1, \omega_2, \omega_3)$ reflects the non-trivial on-site correlations implied by the spin commutation relations [11, 18, 32]. As a result, the chiral quadratic response is only finite if one of the frequencies in $g_0^{xyz}(\omega_1, \omega_2, \omega_3)$ vanishes and the other two frequencies are finite. This is displayed explicitly in the left panel of Fig 8. Obviously, the chiral quadratic response of an isolated spin does not exhibit second-harmonic generation, which is a hallmark of non-linear response. The fact that the spin dynamics generated by the spin algebra yields such a singular frequency dependence of the chiral quadratic response has also been pointed out in a recent work by Kappl *et al.* [10] in the context of the Anderson impurity model. The frequency dependence of the chiral nonlinear susceptibility of an isolated spin in a magnetic field has recently also been discussed in Ref [20].

Let us now explore the fate of this singular frequency dependence when we include the effect of the exchange interaction at high temperatures and frequencies. To first order in J the chiral three-point function is given by the four first-order diagrams in the first and second line of Fig. 7. For $H = 0$ the tadpole diagram vanishes and the contribution from the three diagrams in the second line of Fig. 7 is

$$\begin{aligned} G^{xyz(1)}(K_1, K_2, K_3) &= -\beta^2 b_1^2 (1 - \delta_{\omega_1, 0} \delta_{\omega_2, 0} \delta_{\omega_3, 0}) \\ &\quad \times \left[J_{\mathbf{k}_1} \frac{\delta_{\omega_1, 0}}{\omega_2} + J_{\mathbf{k}_2} \frac{\delta_{\omega_2, 0}}{\omega_3} + J_{\mathbf{k}_3} \frac{\delta_{\omega_3, 0}}{\omega_1} \right]. \end{aligned} \quad (3.39)$$

At this level of approximation the chiral susceptibility still exhibits the same singular frequency dependence as the generalized block $g_0^{xyz}(\omega_1, \omega_2, \omega_3)$. To obtain the generic frequency dependence of the chiral non-linear susceptibility allowing for second-harmonic generation we have to take into account the second order in the exchange couplings. For simplicity let us focus again on the limit $H \rightarrow 0$ where all tadpole diagrams involving the one-point functions in Fig. 7 vanish. Assuming that

all three Matsubara frequencies ω_1 , ω_2 and ω_3 are finite, only four diagrams contribute to the chiral three-point function: the diagram involving the five-spin block in the third row of Fig. 7, and the three diagrams where the four-spin blocks are connected to the three-spin blocks in the fourth row of Fig. 7. Technical details for the evaluation of these diagrams are given in Appendix B. If none of the frequencies ω_1, ω_2 and ω_3 is zero we obtain for vanishing magnetic field

$$\begin{aligned} G^{xyz(2)}(K_1, K_2, K_3) &= \beta b_1^2 \left(\frac{1}{N} \sum_{\mathbf{q}} J_{\mathbf{q}}^2 \right) \\ &\quad \times \left[\frac{1 - \lambda_{\mathbf{k}_1}}{\omega_1^2} \left(\frac{1}{\omega_2} - \frac{1}{\omega_3} \right) \right. \\ &\quad \left. + \frac{1 - \lambda_{\mathbf{k}_2}}{\omega_2^2} \left(\frac{1}{\omega_3} - \frac{1}{\omega_1} \right) \right. \\ &\quad \left. + \frac{1 - \lambda_{\mathbf{k}_3}}{\omega_3^2} \left(\frac{1}{\omega_1} - \frac{1}{\omega_2} \right) \right], \end{aligned} \quad (3.40)$$

where the momentum-dependent form factor $\lambda_{\mathbf{k}}$ is defined by the following ratio of Brillouin zone averages,

$$\lambda_{\mathbf{k}} = \frac{\sum_{\mathbf{q}} J_{\mathbf{q}} J_{\mathbf{q}+\mathbf{k}}}{\sum_{\mathbf{q}} J_{\mathbf{q}}^2}. \quad (3.41)$$

For nearest-neighbor exchange the form factor $\lambda_{\mathbf{k}} = \gamma_{\mathbf{k}}$ agrees with nearest-neighbor form factor defined in Eq. (3.27). For the special case $S = 1/2$ the frequency-structure in Eq. (3.40) has been obtained previously by Krieg [26], who has calculated the corresponding irreducible three-point vertex. This frequency dependence is shown in the center and right panels of Fig. 8 for two different sets of wave-vectors. Note that the single-spin result (3.38) and the first-order correction (3.39) to the chiral non-linear susceptibility are only finite if one of the frequencies ω_1 , ω_2 or ω_3 vanishes and the other two frequencies are non-zero; the corresponding quadratic response does not exhibit second-harmonic generation. In contrast, the second-order non-linear chiral spin susceptibility (3.40) is finite for all frequencies compatible with frequency conservation, as illustrated in Fig. 8. In particular, for $\omega_1 = \omega_2 = \omega \neq 0$ and $\omega_3 = -2\omega$ we obtain

$$\begin{aligned} G^{xyz(2)}(\mathbf{k}_1, \omega, \mathbf{k}_2, \omega, -\mathbf{k}_1 - \mathbf{k}_2, -2\omega) \\ = -3\beta b_1^2 \left(\frac{1}{N} \sum_{\mathbf{q}} J_{\mathbf{q}}^2 \right) \frac{\lambda_{\mathbf{k}_1} - \lambda_{\mathbf{k}_2}}{2\omega^3}. \end{aligned} \quad (3.42)$$

Note that the frequencies in the above expressions are bosonic Matsubara frequencies so that an analytic continuation to the real frequency axis is necessary to obtain the physical chiral response function. We conclude that a necessary condition for observing second-harmonic generation in the chiral non-linear response function of a quantum spin system at high temperatures and frequencies is $\lambda_{\mathbf{k}_1} \neq \lambda_{\mathbf{k}_2}$, which means that the external magnetic field must be inhomogeneous such that the spatial




FIG. 8. Matsubara-frequency dependence of the chiral quadratic response. Left: Response (3.38) of an isolated spin, which is only finite if one of the three frequencies ω_1 , ω_2 or $\omega_3 = -\omega_1 - \omega_2$ vanishes. Center: The order- J^2/T^2 contribution (3.40) to the chiral quadratic response that exhibits the generic frequency dependence, for $\lambda_{\mathbf{k}_1} = \lambda_{\mathbf{k}_2} = \lambda_{\mathbf{k}_3} = -1$. Right: The same for $\lambda_{\mathbf{k}_1} = \lambda_{\mathbf{k}_2} = 1$ and $\lambda_{\mathbf{k}_3} = -1$. All plots are normalized to their maximum value.

dependence of two orthogonal components of the field is characterized by different wave-vectors.

IV. SUMMARY AND CONCLUSIONS

In this work we have derived a new recursive algorithm for generating the high-temperature series expansion of the free energy and arbitrary connected correlation functions of quantum spin systems. Our algorithm is based on a formally exact functional renormalization group flow equation for the generating functional of imaginary time-ordered connected spin correlation function [9]. Using a specific interaction-switch deformation scheme, we have obtained a system of recursion relations given by Eq. (2.14) which can be iteratively solved to obtain the coefficients in the high-temperature expansion of arbitrary connected correlation functions. It is easy to implement our iterative algorithm using symbolic manipulation software such as MATHEMATICA to obtain the free energy, the magnetic equation of state, and the two-point function up to order J^4 . The main advantage of our method is that it does not require any additional combinatorial considerations; all combinatorial factors are explicitly taken into account via the symmetrization operator $\mathcal{S}_{1,\dots,m;m+1,\dots,n}$ defined in Eq. (2.9).

Whether or not a fully numerical implementation of our recursive algorithm based on Eq. (2.14) can be used to obtain competitive high-order results in the high-temperature series expansion remains an interesting open question. For Heisenberg models, one has to invest some effort into calculating the initial conditions for the iteration given by the generalized blocks describing the frequency-dependent correlations between different components of a single spin. These correlations can be cal-

culated iteratively using the recursive form of the generalized Wick-theorem for spin operators [11, 20]. On the other hand, for Ising models the initial conditions can be explicitly written down in closed form. Note that the spin- S Ising model can be obtained from our general spin Hamiltonian (2.1) by setting $J_{ij}^{ab} = \delta^{az}\delta^{bz}J_{ij}$. In this case all operators in our spin Hamiltonian commute so that their correlation functions are time-independent. For vanishing exchange couplings the connected n -point functions are then given by [11]

$$G_{i_1\dots i_n}^{(n,0)} = \delta_{i_1 i_2} \delta_{i_2 i_3} \dots \delta_{i_{n-1} i_n} b^{(n-1)}(\beta H), \quad (4.1)$$

where $b^{(n-1)}(y)$ the $(n-1)$ -st derivative of the spin- S Brillouin function $b(y)$ defined in Eq. (3.5). Given this initial condition, a fully numerical implementation of our recursive algorithm for Ising models seems to be straightforward.

We have used our method to calculate the interaction-irreducible spin susceptibility $\Pi(\mathbf{k}, \omega)$ to order J^3 and the chiral non-linear spin susceptibility $G^{xyz}(K_1, K_2, K_3)$ to order J^2 . From our perturbative result for the static part $\Pi^{-1}(\mathbf{k}, 0)$ of the *inverse* interaction-irreducible susceptibility we have obtained estimates for the critical temperature of three-dimensional nearest-neighbor Heisenberg ferro- and antiferromagnets which overestimate the known results by roughly 10-20%. Given the simplicity of our method it should be useful to estimate critical temperatures of more complicated spin systems where accurate benchmarks are not readily available [21–23].

The fact that the chiral non-linear susceptibility in strongly correlated systems exhibits an interesting frequency dependence has recently been pointed out by Kappl *et al.* [10] who presented numerical results for the chiral non-linear susceptibility of the Anderson im-

purity model. Surprisingly, a thorough investigation of this quantity for Heisenberg magnets at low temperatures seems not to exist in the literature. Although the perturbative result for the chiral non-linear susceptibility in Eq. (3.40) is only valid at high temperatures and frequencies, it exhibits a rather non-trivial momentum- and frequency dependence which can in principle be observed experimentally by measuring the strength of second-harmonic generation in the chiral quadratic magnetization response of Heisenberg magnets.

ACKNOWLEDGMENTS

We thank Björn Sbierski for useful discussions. This work was financially supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) through Project No. 431190042.

APPENDIX A: Connected correlations of a single spin in a magnetic field

In this Appendix we summarize the connected imaginary time-ordered correlation functions for the components of a single spin- S operator \mathbf{S} in a magnetic field. Following Ref. [17], we call these quantities *generalized blocks*. Choosing our coordinate system such that the z -axis is aligned with the direction of the magnetic field, the Hamiltonian is simply $\mathcal{H}_0 = -HS^z$, where the magnetic field H is measured in units of energy. Although for $H \rightarrow 0$ the Hamiltonian vanishes, the spin correlations remain non-trivial due to the $SU(2)$ -commutation relations between different spin components. The generalized blocks involving up to four spin components can be found in Refs. [11, 17–19]. For the calculation of the interaction-irreducible spin susceptibility $\Pi(\mathbf{k}, \omega)$ to order J^3 and the chiral non-linear susceptibility $G^{xyz}(K_1, K_2, K_3)$ to order J^2 in Sec. III we also need the generalized five-spin blocks which can be obtained from the recursive algorithm described in the Appendix B of Ref. [11]. A similar algorithm has recently been constructed in Ref. [20].

For finite magnetic field it is convenient to work with the spherical spin components S^p defined in Eq. (3.10), where $p = +, -$ labels the two circular components transverse to the magnetic field and $p = 0 = z$ labels the longitudinal component in the direction of the field. In this basis the connected two-point functions are given in Eqs. (3.9c) and (3.13). The connected three-point functions are more interesting. Of particular interest is the chiral three-point function $g^{+-z}(\omega_1, \omega_2, \omega_3)$ involving three different spin components which can be written as [11]

$$bg^{+-z}(\omega_1, \omega_2, \omega_3) = -g(\omega_1) [g(-\omega_2) - \beta \delta_{\omega_3, 0} b']. \quad (\text{A1})$$

Here

$$g(\omega) = \frac{b}{H - i\omega}, \quad (\text{A2})$$

where $b = b(\beta H)$ is the Brillouin function and $b' = b'(\beta H)$ is its first derivative. For finite magnetic field H the purely longitudinal three-point function is also finite,

$$g^{zzz}(\omega_1, \omega_2, \omega_3) = \beta^2 \delta_{\omega_1, 0} \delta_{\omega_2, 0} b''(\beta H), \quad (\text{A3})$$

where $b''(\beta H)$ is the second derivative of the Brillouin function. Recall that with our convention (3.1) the above three-point functions should be multiplied by an overall factor of $\beta \delta_{\omega_1 + \omega_2 + \omega_3, 0}$ so that we can use this constraint to write our correlation functions in a more symmetric form. For $H \rightarrow 0$ the longitudinal three-point function vanishes because $b''(0) = 0$ while the chiral three-point function (A1) has a non-trivial limit which can be written in the highly symmetric form given in Eq. (3.36).

Next, consider the connected four-point functions in a magnetic field. The purely longitudinal four-point function is

$$g^{zzzz}(\omega_1, \omega_2, \omega_3, \omega_4) = \beta^3 \delta_{\omega_1, 0} \delta_{\omega_2, 0} \delta_{\omega_3, 0} b'''(\beta H), \quad (\text{A4})$$

where for $H \rightarrow 0$ the third derivative $b'''(\beta H)$ of the Brillouin function has a finite limit b_3 given by Eq. (3.8b). The four-point functions involving transverse spin components are more interesting. The purely transverse four-point function can be written as [11]

$$b^2 g^{++++}(\omega_1, \omega_2, \omega_3, \omega_4) = -g(\omega_1) g(\omega_2) \left\{ g(-\omega_3) + g(-\omega_4) - \beta [\delta_{\omega_1, -\omega_3} + \delta_{\omega_1, -\omega_4}] b' \right\}, \quad (\text{A5})$$

while the mixed transverse-longitudinal four-point function is

$$b^2 g^{+-zz}(\omega_1, \omega_2, \omega_3, \omega_4) = g(\omega_1) g(-\omega_2) \left\{ g(\omega_1 + \omega_3) + g(\omega_1 + \omega_4) - \beta [\delta_{\omega_3, 0} + \delta_{\omega_4, 0}] b' \right\} + g(\omega_1) \beta^2 \delta_{\omega_3, 0} \delta_{\omega_4, 0} b b''. \quad (\text{A6})$$

For vanishing magnetic field only special frequency combinations of Eqs. (A5) and (A6) have a non-zero limit. Here we give only those contributions needed for the evaluation of the second-order contribution $\Pi^{(2)}(\mathbf{k}, \omega)$ to the irreducible susceptibility for vanishing magnetic field given in Eq. (3.23),

$$g_0^{zzzz}(0, 0, \omega, -\omega) = \beta^3 \delta_{\omega, 0} b_3, \quad (\text{A7})$$

$$g_0^{+-zz}(0, 0, \omega, -\omega) = \frac{\beta^3}{3} \delta_{\omega, 0} b_3 + 2\beta(1 - \delta_{\omega, 0}) \frac{b_1}{\omega^2}. \quad (\text{A8})$$

For the calculation of the third-order correction $\Pi^{(3)}(\mathbf{k}, \omega)$ to the irreducible susceptibility we also need the generalized five-spin blocks which can be obtained either diagrammatically following Vaks, Larkin, and Pikin [18], or purely algebraically using the recursive form the generalized Wick theorem for spin operators derived in Appendix B of Ref. [11]. We obtain

$$\begin{aligned}
b^3 g^{+-z}(\omega_1, \omega_2, \omega_3, \omega_4, \omega_5) &= g(-\omega_3)g(-\omega_4) \left\{ \right. \\
&g(\omega_1) \left[g(\omega_1 + \omega_5) + g(-\omega_3 - \omega_5) + g(-\omega_4 - \omega_5) - \beta[\delta_{\omega_5,0} + \delta_{\omega_2,-\omega_3} + \delta_{\omega_2,-\omega_4}]b' \right] \\
&+ g(\omega_2) \left[g(\omega_2 + \omega_5) + g(-\omega_3 - \omega_5) + g(-\omega_4 - \omega_5) - \beta[\delta_{\omega_5,0} + \delta_{\omega_1,-\omega_3} + \delta_{\omega_1,-\omega_4}]b' \right] \\
&\left. + \beta^2 \delta_{\omega_5,0} [\delta_{\omega_1,-\omega_3} + \delta_{\omega_2,-\omega_3}]bb'' \right\}, \tag{A9}
\end{aligned}$$

and for the other mixed five-spin block,

$$\begin{aligned}
b^3 g^{+zzz}(\omega_1, \omega_2, \omega_3, \omega_4, \omega_5) &= -g(\omega_1)g(-\omega_2) \left\{ \right. \\
&\left[g(\omega_1 + \omega_3) - \beta\delta_{\omega_3,0}b' \right] \left[g(-\omega_2 - \omega_4) + g(-\omega_2 - \omega_5) \right] + \beta^2 \delta_{\omega_4,0} \delta_{\omega_5,0} bb'' \\
&+ \left[g(\omega_1 + \omega_4) - \beta\delta_{\omega_4,0}b' \right] \left[g(-\omega_2 - \omega_5) + g(-\omega_2 - \omega_3) \right] + \beta^2 \delta_{\omega_5,0} \delta_{\omega_3,0} bb'' \\
&+ \left[g(\omega_1 + \omega_5) - \beta\delta_{\omega_5,0}b' \right] \left[g(-\omega_2 - \omega_3) + g(-\omega_2 - \omega_4) \right] + \beta^2 \delta_{\omega_3,0} \delta_{\omega_4,0} bb'' \left. \right\} \\
&+ g(-\omega_2) \beta^3 \delta_{\omega_3,0} \delta_{\omega_4,0} \delta_{\omega_5,0} b^2 b'''. \tag{A10}
\end{aligned}$$

APPENDIX B: Evaluation of high-temperature expansions

In this Appendix, we give some technical details for the evaluation of the perturbative high-temperature ex-

pansions in Sec. III. Consider first the high-temperature expansion of the negative free energy $-\beta F = G^{(0)}$ in units of temperature given in Eq. (3.3). The third-order coefficient $G^{(0,3)}$ represented by the five third-order diagrams in Fig. 3 is given by

$$\begin{aligned}
G^{(0,3)} &= -\frac{1}{12\beta} \sum_{ij} \sum_{a_1 b_1} \sum_{a_2 b_2} \sum_{a_3 b_3} J_{ij}^{a_1 b_1} J_{ij}^{a_2 b_2} J_{ij}^{a_3 b_3} \sum_{\omega \omega'} g_i^{a_1 a_2 a_3}(\omega, \omega', -\omega - \omega') g_j^{b_1 b_2 b_3}(-\omega, -\omega', \omega + \omega') \\
&- \frac{1}{6} \sum_{ijk} \sum_{a_1 b_1} \sum_{a_2 b_2} \sum_{a_3 b_3} J_{ij}^{a_1 b_1} J_{jk}^{a_2 b_2} J_{ki}^{a_3 b_3} \sum_{\omega} g_i^{a_1 b_3}(\omega) g_j^{a_2 b_1}(\omega) g_k^{a_3 b_2}(\omega) \\
&- \frac{\beta}{6} \sum_{ijkl} \sum_{abc} J_{ij}^{az} J_{ik}^{bz} J_{il}^{cz} g_i^{abc}(0, 0, 0) m_j m_k m_l \\
&- \frac{1}{2} \sum_{ijk} \sum_{a_1 b_1} \sum_{a_2 b_2} \sum_c J_{ij}^{a_1 b_1} J_{ij}^{a_2 b_2} J_{ik}^{cz} \sum_{\omega} g_i^{a_1 a_2 c}(\omega, -\omega, 0) g_j^{b_2 b_1}(\omega) m_k \\
&- \frac{\beta}{2} \sum_{ijkl} \sum_{abcd} J_{ij}^{ab} J_{ik}^{cz} J_{jl}^{dz} g_i^{ac}(0) g_j^{bd}(0) m_k m_l. \tag{B1}
\end{aligned}$$

For the special case of an isotropic Heisenberg magnet ($J_{ij}^{ab} = \delta^{ab} J_{ij}$) in a homogeneous magnetic field ($H_i = H$) along the z -axis the generalized blocks $g^{a_1 \dots a_n}(\omega_1, \dots, \omega_n)$ are independent of the site label. The resulting high-temperature

expansion of the free energy in a magnetic field is then

$$\begin{aligned}
F = & -\frac{NB(\beta H)}{\beta} + \frac{m^2}{2} \sum_{ij} J_{ij} - \frac{1}{4\beta} \sum_{ij} J_{ij}^2 \sum_{\omega} \sum_{ab} g^{ab}(\omega) g^{ba}(\omega) - \frac{m^2}{2} g^{zz}(0) \sum_{ijk} J_{ij} J_{jk} \\
& + \frac{1}{12\beta^2} \sum_{ij} J_{ij}^3 \sum_{\omega\omega'} \sum_{abc} g^{abc}(\omega, \omega', -\omega - \omega') g^{abc}(-\omega, -\omega', \omega + \omega') \\
& + \frac{1}{6\beta} \sum_{ijk} J_{ij} J_{jk} J_{ki} \sum_{\omega} \sum_{abc} g^{ab}(\omega) g^{bc}(\omega) g^{ca}(\omega) + \frac{m^3}{6} g^{zzz}(0, 0, 0) \sum_{ijkl} J_{ij} J_{ik} J_{il} \\
& + \frac{m}{2\beta} \sum_{ijk} J_{ij}^2 J_{ik} \sum_{\omega} \sum_{ab} g^{abz}(\omega, -\omega, 0) g^{ba}(\omega) + \frac{m^2}{2} \sum_{ijkl} J_{ij} J_{ik} J_{jl} \sum_a [g^{az}(0)]^2 + \mathcal{O}(J^4). \tag{B2}
\end{aligned}$$

Next, consider the high-temperature expansion of the two-spin correlation function,

$$G_{ij}^{ab}(\omega) = \delta_{ij} g_i^{ab}(\omega) + G_{ij}^{ab(1)}(\omega) + G_{ij}^{ab(2)}(\omega) + G_{ij}^{ab(3)}(\omega) + \mathcal{O}(J^4), \tag{B3}$$

where the zeroth-order term $g_i^{ab}(\omega)$ is given in Eq. (3.9), the first-order term $G_{ij}^{ab(1)}(\omega)$ is given in Eq. (3.18), and the second-order term is

$$\begin{aligned}
G_{ij}^{ab(2)}(\omega) = & \frac{1}{2\beta} \sum_{a_1 b_1} \sum_{a_2 b_2} J_{ij}^{a_1 b_1} J_{ij}^{a_2 b_2} \sum_{\omega'} g_i^{a_1 a_2}(\omega, \omega', -\omega - \omega') g_j^{b_1 b_2}(-\omega, -\omega', \omega + \omega') \\
& + \frac{\delta_{ij}}{2\beta} \sum_k \sum_{a_1 b_1} \sum_{a_2 b_2} J_{ik}^{a_1 b_1} J_{ik}^{a_2 b_2} \sum_{\omega'} g_i^{a_1 a_2}(\omega, -\omega, \omega', -\omega') g_k^{b_1 b_2}(\omega', -\omega') \\
& + \sum_k \sum_{a_1 b_1} \sum_{a_2 b_2} J_{ik}^{a_1 b_1} J_{kj}^{a_2 b_2} g_i^{a_1 a_2}(\omega) g_k^{b_1 a_2}(\omega) g_j^{b_2 b_1}(\omega) \\
& + \frac{\delta_{ij}}{2} \sum_{kl} \sum_{cd} J_{ik}^{cz} J_{il}^{dz} g_i^{abcd}(\omega, -\omega, 0) m_k m_l \\
& + \delta_{ij} \sum_{kl} \sum_{a_1 b_1 c} J_{ik}^{a_1 b_1} J_{kl}^{cz} g_i^{a_1 a_2}(\omega, -\omega, 0) g_k^{b_1 c}(0) m_l \\
& + \sum_k \sum_{a_1 b_1 c} J_{ij}^{a_1 b_1} \left[J_{ik}^{cz} g_i^{a_1 a_2}(\omega, -\omega, 0) g_j^{b_1 b_2}(\omega) + J_{jk}^{cz} g_i^{a_1 a_2}(\omega) g_j^{b_1 b_2}(\omega, -\omega, 0) \right] m_k. \tag{B4}
\end{aligned}$$

Using Eq. (A8) we can easily evaluate the frequency sums in our expression for $\Pi^{(2)}(\mathbf{k}, \omega)$ in Eq. (3.23),

$$\frac{1}{2\beta} \sum_{\omega'} g_0(\omega') \left[2g_0^{+-zz}(\omega', -\omega', \omega, -\omega) + g_0^{zzzz}(\omega', -\omega', \omega, -\omega) \right] = \delta_{\omega,0} \beta^3 \frac{5}{6} b_1 b_3 + (1 - \delta_{\omega,0}) \frac{\beta}{\omega^2} 2b_1^2, \tag{B5}$$

$$\frac{1}{\beta} \sum_{\omega'} g_0^{+-z}(\omega', -\omega - \omega', \omega) g_0^{+-z}(\omega + \omega', -\omega', -\omega) = -\delta_{\omega,0} \beta^3 \frac{b_1^2}{12} - (1 - \delta_{\omega,0}) \frac{\beta}{\omega^2} 2b_1^2. \tag{B6}$$

The third-order correction $\Pi^{(3)}(\mathbf{k}, \omega)$ to the irreducible spin susceptibility represented by the four diagrams in Fig. 5 is

$$\begin{aligned}
\Pi^{(3)}(\mathbf{k}, \omega) = & -\frac{1}{N^2 \beta^2} \sum_{\mathbf{q}_1 \mathbf{q}_2} J_{\mathbf{q}_1} J_{\mathbf{q}_2} J_{\mathbf{q}_1 + \mathbf{q}_2} \sum_{\omega_1 \omega_2} g^{+-zzz}(\omega_1, \omega_2, -\omega_1 - \omega_2, \omega, -\omega) g^{+-z}(-\omega_2, -\omega_1, \omega_1 + \omega_2) \\
& - \frac{1}{N^2 \beta^2} \sum_{\mathbf{q}_1 \mathbf{q}_2} J_{\mathbf{q}_1} J_{\mathbf{q}_2} J_{\mathbf{q}_1 + \mathbf{q}_2 + \mathbf{k}} \sum_{\omega_1 \omega_2} \left[g^{+-zz}(\omega_1, \omega_2, \omega, -\omega_1 - \omega_2 - \omega) g^{+-zz}(-\omega_2, -\omega_1, -\omega, \omega_1 + \omega_2 + \omega) \right. \\
& \quad \left. + \frac{1}{6} g^{zzzz}(\omega_1, \omega_2, \omega, -\omega_1 - \omega_2 - \omega) g^{zzzz}(-\omega_2, -\omega_1, -\omega, \omega_1 + \omega_2 + \omega) \right] \\
& - \frac{1}{2\beta N} \sum_{\mathbf{q}} J_{\mathbf{q}}^3 \sum_{\omega'} g^2(\omega') \left[2g^{+-zz}(\omega', -\omega', \omega, -\omega) + g^{zzzz}(\omega', -\omega', \omega, -\omega) \right] \\
& - \frac{2}{\beta N} \sum_{\mathbf{q}} J_{\mathbf{q}}^2 J_{\mathbf{q} + \mathbf{k}} \sum_{\omega'} g(\omega') g^{+-z}(\omega', -\omega - \omega', \omega) g^{+-z}(\omega + \omega', -\omega', -\omega). \tag{B7}
\end{aligned}$$

It is convenient to perform the frequency sums for finite field and then take the limit $\beta H \rightarrow 0$ using the identities

$$\begin{aligned} & \lim_{H \rightarrow 0} \frac{1}{\beta^2} \sum_{\omega_1 \omega_2} \left[g^{+-zz}(\omega_1, \omega_2, \omega, -\omega_1 - \omega_2 - \omega) g^{+--z}(-\omega_2, -\omega_1, -\omega, \omega_1 + \omega_2 + \omega) \right. \\ & \quad \left. + \frac{1}{6} g^{zzzz}(\omega_1, \omega_2, \omega, -\omega_1 - \omega_2 - \omega) g^{zzzz}(-\omega_2, -\omega_1, -\omega, \omega_1 + \omega_2 + \omega) \right] \\ & = \delta_{\omega,0} \frac{\beta^4}{72} [b_1^2 + 20b_3^2] + (1 - \delta_{\omega,0}) \frac{\beta^2 b_1^2}{2\omega^2}, \end{aligned} \quad (\text{B8})$$

and

$$\begin{aligned} & \lim_{H \rightarrow 0} \frac{1}{\beta^2} \sum_{\omega_1 \omega_2} g^{+--z}(\omega_1, \omega_2, -\omega_1 - \omega_2, \omega, -\omega) g^{+--z}(-\omega_2, -\omega_1, \omega_1 + \omega_2) \\ & = \delta_{\omega,0} \frac{\beta^4}{72} [b_1^2 - 10b_1 b_3] - (1 - \delta_{\omega,0}) \frac{\beta^2 b_1^2}{2\omega^2}. \end{aligned} \quad (\text{B9})$$

Finally, let us give some technical details for the derivation of our second-order result for chiral non-linear susceptibility $G^{+-z(2)}(K_1, K_2, K_3)$ in Eq. (3.40). All diagrams contributing up to second order in the exchange couplings are shown in Fig. 7. Assuming $H = 0$ and that none of the frequencies ω_1, ω_2 and ω_3 vanishes, only four second-order diagrams shown in Fig. 7 contribute: the diagram where the five-spin block is connected with the two-spin block in the third line, and the three diagrams where the four-spin block is connected with the three-spin block in the fourth line of Fig. 7. The contribution from the diagram containing the five-spin block is

$$G^{+-z(2a)}(K_1, K_2, K_2) = \frac{1}{2\beta N} \sum_{\mathbf{q}} J_{\mathbf{q}}^2 \sum_{\omega} g(\omega) [2g^{+---z}(\omega_1, \omega, \omega_2, -\omega, \omega_3) + g^{+--zz}(\omega_1, \omega_2, \omega_2, \omega, -\omega)]. \quad (\text{B10})$$

For $H \rightarrow 0$ we may replace $g(\omega) \rightarrow \beta b_1 \delta_{\omega,0}$ so that the frequency sum can be carried out trivially. Using our results for the five-spin blocks given in Eqs. (A9) and (A10) we obtain for $H \rightarrow 0$,

$$G^{+-z(2a)}(K_1, K_2, K_2) = i\beta b_1^2 \left(\frac{1}{N} \sum_{\mathbf{q}} J_{\mathbf{q}}^2 \right) \left[\frac{2}{\omega_1 \omega_2} \left(\frac{1}{\omega_1} - \frac{1}{\omega_2} \right) - \frac{1}{\omega_1 \omega_2 \omega_3} - \frac{2}{\omega_2 \omega_3^2} \right]. \quad (\text{B11})$$

Finally, using the constraint $\omega_1 + \omega_2 + \omega_3 = 0$ we may write the frequency-dependent term in a more symmetric form,

$$\begin{aligned} & \frac{2}{\omega_1 \omega_2} \left(\frac{1}{\omega_1} - \frac{1}{\omega_2} \right) - \frac{1}{\omega_1 \omega_2 \omega_3} - \frac{2}{\omega_2 \omega_3^2} = \frac{(\omega_1 - \omega_2)(\omega_2 - \omega_3)(\omega_3 - \omega_1)}{\omega_1^2 \omega_2^2 \omega_3^2} \\ & = \frac{1}{\omega_1^2} \left(\frac{1}{\omega_2} - \frac{1}{\omega_3} \right) + \frac{1}{\omega_2^2} \left(\frac{1}{\omega_3} - \frac{1}{\omega_1} \right) + \frac{1}{\omega_3^2} \left(\frac{1}{\omega_1} - \frac{1}{\omega_2} \right), \end{aligned} \quad (\text{B12})$$

which gives the momentum-independent contribution to $G^{xyz(2)}(K_1, K_2, K_3)$ in Eq. (3.40). The momentum-dependent part involving the form factors $\lambda_{\mathbf{k}}$ defined in Eq. (3.41) arises from the three diagrams in the fourth row of Fig. 7 where the four-spin block is connected to the three-spin block. The explicit evaluation of these

diagrams is straightforward. To write the resulting frequency dependence in the form given in Eq. (3.40) we use again some identities implied by frequency conservation such as

$$\frac{2}{\omega_1 \omega_3^2} + \frac{1}{\omega_1 \omega_2 \omega_3} = \frac{1}{\omega_3^2} \left(\frac{1}{\omega_1} - \frac{1}{\omega_2} \right). \quad (\text{B13})$$

-
- [1] C. Domb and M. S. Green, *Phase Transitions and Critical Phenomena, Volume 3: Series Expansions for Lattice Models*, (Academic Press, London, 1974).
 [2] J. Oitmaa, C. Hamer, and W. Zhang, *Series Expansion Methods for Strongly Interacting Lattice Models*, (Cam-

- bridge University Press, Cambridge, 2006).
 [3] A. Lohmann, H.-J. Schmidt, and J. Richter, *Tenth-order high-temperature expansion for the susceptibility and the specific heat of spin-s Heisenberg models with arbitrary exchange patterns: Applications to pyrochlore and*

- kagome magnets*, Phys. Rev. B **89**, 014415 (2014).
- [4] P. Müller, A. Lohmann, J. Richter, O. Menchyshyn, and O. Derzhko, *Thermodynamics of the pyrochlore Heisenberg ferromagnet with arbitrary spin S* , Phys. Rev. B **96**, 174419 (2017).
- [5] A. Hehn, N. van Well, and M. Troyer, *High-temperature series expansion for spin 1/2 Heisenberg models*, Computer Physics Communications **212**, 180 (2017).
- [6] M. G. Gonzalez, M. Bernu, L. Pierre, and L. Messio, *Finite-temperature phase-transitions in $s01/2$ three-dimensional Heisenberg magnets from high-temperature series expansion*, Phys. Rev. B **107**, 235151 (2023).
- [7] L. Pierre, B. Bernu, and L. Messio, *High temperature series expansions of $S = 1/2$ Heisenberg spin models: algorithm to include the magnetic field with optimized complexity*, arXiv:2404.02271v1 [cond-mat.str-el] 2 Apr 2024.
- [8] M. Campostrini, A. Pelissetto, M. Rossi, and E. Vicari, *25-th order high-temperature expansion results for three-dimensional Ising-like systems on the simple cubic lattice*, Phys. Rev. E **65**, 066127 (2002).
- [9] J. Krieg and P. Kopietz, *Exact renormalization group for quantum spin systems*, Phys. Rev. B **99**, 060403(R) (2019).
- [10] P. Kappl, F. Krien, C. Watzenböck, and K. Held, *Non-linear responses and three-particle correlators in correlated electron systems exemplified by the Anderson impurity model*, Phys. Rev. B **107**, 205108 (2023).
- [11] R. Goll, D. Tarasevych, J. Krieg, and P. Kopietz, *Spin functional renormalization group for quantum Heisenberg ferromagnets: Magnetization and magnon damping in two dimensions*, Phys. Rev. B **100**, 174424 (2019).
- [12] R. Goll, A. Rückriegel, and P. Kopietz, *Zero-magnon sound in quantum Heisenberg ferromagnets*, Phys. Rev. B **102**, 224437 (2020).
- [13] D. Tarasevych and P. Kopietz, *Dissipative spin dynamics in hot quantum paramagnets*, Phys. Rev. B **104**, 024423 (2021).
- [14] D. Tarasevych, A. Rückriegel, S. Keupert, V. Mitsioannou, and P. Kopietz, *Spin-functional renormalization group for the $J_1J_2J_3$ Heisenberg model*, Phys. Rev. B **106**, 174412 (2022); *ibid.* **107**, 019904(E) (2023).
- [15] A. Rückriegel, D. Tarasevych, and P. Kopietz, *Phase diagram of the J_1 - J_2 quantum Heisenberg model for arbitrary spin*, Phys. Rev. B **109**, 184410 (2024).
- [16] P. Kopietz, L. Bartosch, and F. Schütz, *Introduction to the Functional Renormalization Group*, (Springer, Berlin, 2010).
- [17] Yu. A. Izyumov and Yu. N. Skryabin, *Statistical Mechanics of Magnetically Ordered Systems*, (Springer, Berlin, 1988).
- [18] V. G. Vaks, A. I. Larkin, and S. A. Pikin, *Thermodynamics of an ideal ferromagnetic substance*, Zh. Eksp. Teor. Fiz. **53**, 281 (1967) [Sov. Phys. JETP **26**, 188 (1968)].
- [19] V. G. Vaks, A. I. Larkin, and S. A. Pikin, *Spin waves and correlation functions in a ferromagnetic*, Zh. Eksp. Teor. Fiz. **53**, 1089 (1967) [Sov. Phys. JETP **26**, 647 (1968)].
- [20] J. Halbinger, B. Schneider, and B. Sbierski, *Spectral representation of Matsubara n -point functions: Exact kernel functions and applications*, SciPost Phys. **15.5**, 193 (2023).
- [21] J. G. Rau, E. K.-H. Lee, and H.-Y. Kee, *Generic Spin Model for the Honeycomb Iridates beyond the Kitaev Limit*, Phys. Rev. Lett. **112**, 077204 (2014).
- [22] L. Smejkal, A. Marmodoro, K.-H. Ahn, R. Gonzalez-Hernandez, I. Turek, S. Mankovsky, H. Ebert, S. W. D'Souza, O. Sipr, J. Sinova, and T. Jungwirth, *Chiral Magnons in altermagnetic RuO_2* , Phys. Rev. Lett. **131**, 256703 (2023).
- [23] F. Garcia-Gaitan, A. Kefayati, J. Q. Xiao, and B. K. Nikolic, *Magnon spectrum of altermagnets: Time-dependent matrix product states vs. linearized Holstein-Primakoff calculations unravelling spontaneous magnon decay*, arXiv: 2402.19433.
- [24] R. A. Tahir-Kheli and D. G. McFadden, *Short-Range-Order Effects in Neutron Scattering from Heisenberg Paramagnets: Application to $RbMnF_3$* , Phys. Rev. B **1**, 3178 (1970).
- [25] M. F. Collins, *Series Expansions for High-Temperature Dynamics of Heisenberg Paramagnets*, Phys. Rev. B **4**, 1588 (1971).
- [26] J. Krieg, *Functional renormalization group approach to classical and quantum spin systems*, (PhD-Thesis, Goethe-Universität Frankfurt, 2019).
- [27] A. W. Sandvik, *Critical Temperature and the Transition from Quantum to Classical Order Parameter Fluctuations in the Three-Dimensional Heisenberg Antiferromagnet*, Phys. Rev. Lett. **80**, 5196 (1998).
- [28] M. Troyer, F. Alet, and S. Wessel, *Histogram Methods for Quantum Systems: from reweighting to Wang-Landau Sampling*, Braz. J. Phys. **34**, 377 (2004).
- [29] J. Oitmaa and W. Zheng, *Curie and Néel temperatures of quantum magnets*, J. Phys. Condens. Matter **16**, 8653 (2004).
- [30] S. McKenzie, C. Domb and D. L. Hunter *The high-temperature susceptibility of the classical Heisenberg model in four dimensions*, J. Phys. A: Math. Gen. **15**, 3909 (1982).
- [31] A. Cuccoli, R. Maciocco, and R. Vaia, *Transition temperature of three-dimensional quantum Heisenberg ferro- and antiferromagnets*, J. Magn. Magn. Mater. **226-230**, 566 (2001).
- [32] D. Tarasevych, J. Krieg, and P. Kopietz, *A rich man's derivation of scalings laws for the Kondo model*, Phys. Rev. B **98**, 235133 (2018).