Jordan degree type for codimension three Gorenstein algebras of small Sperner number

Nancy Abdallah¹, Nasrin Altafi², Anthony I
arrobino³, and Joachim Yaméogo⁴

¹University of Borås, Borås, Sweden

²Department of Mathematics and Statistics, Queen's University, Kingston, Ontario, Canada, and Department of Mathematics, KTH Royal Institute of Technology, Stockholm, Sweden
³Department of Mathematics, Northeastern University, Boston, MA 02115, USA ⁴Université Côte d'Azur, CNRS, LJAD, France.

June 10, 2024

Abstract

The Jordan type $P_{A,\ell}$ of a linear form ℓ acting on a graded Artinian algebra A over a field k is the partition describing the Jordan block decomposition of the multiplication map m_{ℓ} , which is nilpotent. The Jordan degree type $S_{A,\ell}$ is a finer invariant, describing also the initial degrees of the simple submodules of A in a decomposition of A as $k[\ell]$ modules. The set of Jordan types of A or Jordan degree types (JDT) of A as ℓ varies, is an invariant of the algebra. This invariant has been studied for codimension two graded algebras. We here extend the previous results to certain codimension three graded Artinian Gorenstein

Keywords: Artinian Gorenstein algebra, codimension three, Hilbert function, Jordan type, Jordan degree type, rank matrix, Sperner number.

²⁰²⁰ Mathematics Subject Classification: Primary: 13E10; Secondary: 13H10, 14C05.

Email addresses: nancy.abdallah@hb.se, nasrinar@kth.se, a.iarrobino@neu.edu, joachim.yameogo@univ-cotedazur.fr

(AG) algebras - those of small Sperner number. Given a Gorenstein sequence T - one possible for the Hilbert function of a codimension three AG algebra - the irreducible variety $\operatorname{Gor}(T)$ parametrizes all Gorenstein algebras of Hilbert function T. We here completely determine the JDT possible for all pairs $(A, \ell), A \in \operatorname{Gor}(T)$, for Gorenstein sequences T of the form $T = (1, 3, s^k, 3, 1)$ for Sperner number s = 3, 4, 5 and arbitrary multiplicity k. For s = 6 we delimit the prospective JDT, without verifying that each occurs.

Contents

1	Intr	coduction.	3
	1.1	Lefschetz properties and Jordan type	3
	1.2	Overview of results	5
ก	Der	al matrices and lander demos tures	G
4	Rai	ik matrices and Jordan degree type.	0
	2.1	Jordan basis and Jordan degree type	6
	2.2	Rank matrices.	10
		2.2.1 Map $Gor(T)$ to the punctual Hilbert scheme	10
		2.2.2 Macaulay duality.	13
		2.2.3 Jordan Degree Type matrix	14
0	D 4		1 4
3	Pot	ential Jordan degree types.	15
	3.1	Codimension two	15
	3.2	Codimension three.	16
		3.2.1 Case $s = 3$	17
		3.2.2 Case $s = 4$	19
		3.2.3 Case $s = 5$	20
		3.2.4 Case $s = 6$	24
	•		•
4	Act	Lual JDT for algebras $A \in Gor(T)$.	28
	4.1	Three kinds of growth of JDT parts	28
	4.2	Classification of JDT for $s \leq 5$	32
	4.3	Examples and Methods.	33
	4.4	Questions and problems	37

A Tables of JDT in codimension 3 for almost constant T, when $s \leq 5$. 39

1 Introduction.

Let R = k[x, y, z], the polynomial ring over an infinite field k. We primarily consider here codimension three graded Artinian Gorenstein (AG) algebras A = R/I, I an ideal of R, that have almost constant Hilbert functions H(A) = T with Sperner number at most 6, that is T satisfies

$$T = (1, 3, s^k, 3, 1)$$
 where $3 \le s \le 6, 1 \le k$ (1)

and j = k+3 is the socle degree. For a graded Artinian algebra A, the Jordan type $P_{A,\ell}$ where ℓ is a linear form in R is the partition giving the Jordan block decomposition of the multiplication m_{ℓ} by ℓ on A, which is nilpotent. The algebra A has a decomposition as $k[\ell]$ module: a direct sum of simple modules isomorphic to $k[\ell]/(\ell^{p_i})$, where $P_{A,\ell} = (p_1, p_2, \ldots, p_t), p_1 \ge p_2 \ge \cdots \ge p_t$. The Jordan degree type (JDT) also specifies the degrees of the generators of the simple modules, which is an invariant of A (Definition 2.1).

1.1 Lefschetz properties and Jordan type.

In determining the JDT possible for codimension three Artinian Gorenstein algebras A, a main tool is the Pfaffian structure theorem of D. Buchsbaum and D. Eisenbud, that generators for the ideal I defining A are Pfaffians of an antisymmetric matrix occurring in the minimal resolution [BuEi]. A consequence, realized after work by many, is that for k an infinite field, the graded codimension three AG algebras of given Hilbert function T are parametrized by a smooth irreducible variety Gor(T) of known dimension (see [Di],[IK, Theorem 5.25]). A Hilbert function sequence $T = (1, t_1, \ldots, t_j)$ possible for a graded AG algebra is termed a *Gorenstein sequence*. Those occurring for a codimension three graded Artinian Gorenstein algebra satisfy

$$T = (1, 3, T_1, s^k, T_1^r, 3, 1),$$
(2)

where T_1^r is the reverse of T_1 and $(1, 3, T_1, s)$ is an O-sequence, meaning that its first difference occurs as the Hilbert function of a codimension two graded Artinian algebra [BuEi, St, Di]. The maximum value s of T is termed the Sperner number of T.

Nevertheless, the possible Jordan types and Jordan degree types of these AG algebras, given their Hilbert function, are not well understood. 1

 $^{^1\}mathrm{In}$ higher codimension than three, the set of Gorenstein sequences themselves are not fully known.

Several authors have studied the Jordan type for codimension two graded complete intersections [AIIK], codimension two Artinian algebras [AIKY], and certain graded Artinian Gorenstein (AG) algebras of low socle degree [CGo]. These studies determined the possible Jordan types for the pair (A, ℓ) , for an arbitrary Artinian algebra A in the class studied, and an arbitrary linear form ℓ . We here extend this study to height three graded AG algebras having Sperner number - maximum value of the Hilbert function - 3, 4, 5 or 6, and whose Hilbert function is almost constant; i.e. is of the form $T = (1, 3, s^k, 3, 1)$. For s = 3, 4, 5, we determine all possible JDT for pairs $(A, \ell), A \in \text{Gor}(T)$. These JDT correspond 1-1 to the prospective JDT matrices arising from the rank matrices studied by the second author in [A11] (see Section 2.2). For s = 6 we delimit the prospective JDT, without verifying that each occurs. A key step is a pattern in the rank matrices for such T when $k \geq 3$: the pattern arises from a projection morphism taking Gor(T)to the punctual Hilbert scheme Hilb^s(\mathbb{P}^2) (Lemma 2.8).

We say that the pair (A, ℓ) where A is a graded Artinian algebra and $\ell \in A$ is a linear form is a *weak Lefschetz pair* (WL) if the multiplication map by ℓ from A_i to A_{i+1} has maximal rank for any non-negative integer *i*. If for any pair of non-negative integers (u, i) the multiplication map by ℓ^u from A_i to A_{i+u} has maximal rank, we say that the pair (A, ℓ) is strong Lefschetz (SL). For an Artinian algebra A if there exists a linear form ℓ such that (A, ℓ) is WL (or SL) pair we sometimes say A satisfies the WL (or SL) property and ℓ is a WL (or SL) element. The weak and strong Lefschetz pairs (A, ℓ) can be determined from the Jordan type $P_{A,\ell}$. A pair (A,ℓ) is WL if and only if the number of parts of $P_{A,\ell}$ is the Sperner number (maximum value) of the Hilbert function T. It is SL if and only if $P_{A,\ell}$ is the conjugate T^{\vee} of T, regarded as a partition. Artinian algebras of codimension two satisfy the SL property when the characteristic of k is zero or greater than the socle degree [Br]. In particular, Artinian Gorenstein algebras of codimension two (these are complete intersections) over a field of characteristic zero have the SL property. We know that all codimension three graded *complete intersections* are weak Lefschetz [HMNW] when char k = 0, a result that already uses the Grauert Mülich theorem for vector bundles on \mathbb{P}^2 . The question of whether all codimension three AG algebras A have the weak Lefschetz property equivalently, whether for a generic linear form ℓ the partition $P_{A,\ell}$ has the Sperner number of parts - has been reduced to the case of A compressed

of odd socle degree [BMMNZ].² In a recent paper, the authors with A. Seceleanu showed that when char k = 0, all codimension three AG algebras of Sperner number at most six are strong Lefschetz [AAISY]. Here, in contrast, we consider all JDT consistent with each Hilbert function we study.

1.2 Overview of results.

In Section 2 we define the rank matrices, and give preliminary results, in particular the properties of a new invariant Δ we associate to the rank matrices, that arises from a map from Gor(T) to the punctual Hilbert scheme. Then, after a review of results we will use from codimension two (Section 3.1), we in Section 3.2 determine the potential rank matrices for all AG codimension three algebras whose Hilbert function is almost constant, of Sperner number 3,4,5 and 6.

Theorem 1. Let $T = (1, 3, s^k, 3, 1)$ be almost constant ((1)), and let the field k be infinite. Satisfying the rank conditions of Corollary 2.10 there are for s = 3 and $k \ge 3$ eight rank matrices, for s = 4 and $k \ge 4$ twenty-six rank matrices, for s = 5 and $k \ge 3$ forty-seven rank matrices, and for s = 6 at most sixty-five rank matrices.

This is shown in Theorem 3.3 and Table 3 for s = 3; Theorem 3.5 and Table 4 for s = 4; Theorems 3.6, 3.8 and Table 5 for s = 5; in Theorems 3.9, 3.10 and Table 6 for s = 6.

We then in Section 4.1 first describe three kinds of growth of JDT sequences for a given Sperner number s, as the socle degree j (or, equivalently, the multiplicity k of s in T) increases: in particular *lengthening parts*, and *repeating parts* (Definition 4.1) and in Proposition 4.3 show the connection between the Δ invariant of the rank matrix and the number of these parts. In Section 4.2 and the tables of Section A we exhibit pairs (A, ℓ) having the actually occurring JDT for almost constant T and s = 3, 4, 5 (this is the new aspect of Theorem 2 over Theorem 1).

Theorem 2. Assume the characteristic of k is zero. There are 8 actually occurring JDT for $s = 3, k \ge 3$; 26 for $s = 4, k \ge 4$; and 47 for $s = 5, k \ge 3$. We also specify the numbers of actually occurring JDT for smaller k. We

 $^{^2\}mathbf{A}$ compressed AG algebra quotient A of $\mathsf{k}[x,y,z]$ is one having the maximum possible Hilbert function given j.

exhibit these JDT, along with added data about an example for each - usually a Macaulay dual generator and ideal - in the Tables 7-12 of Appendix A.

Theorem 2 is shown in Theorem 4.6 for s = 3, Theorem 4.7 for s = 4, and in Theorem 4.8 for s = 5. A notable feature of these results is that we give the upper bound on the number of Jordan degree types (Theorem 1, and we specify pairs (A, ℓ) having each occurring JDT (Theorem 2 and Tables 7-12) for arbitrary $k \ge 1$, in particular for arbitrarily large k or socle degree j.

For the case of finite characteristic see Remark 4.9. In Section 4.3 we illustrate how we verified the tables, giving examples of the several methods used. In Section 4.4 we pose further questions.

2 Rank matrices and Jordan degree type.

We introduce Jordan basis and the Jordan degree type (JDT) in Section 2.1 and then in Section 2.2 establish the connection with rank matrices and the JDT matrix studied by the second author in [Al1, Al3]. A map from Gor(T) to the punctual Hilbert scheme Hilb^s(\mathbb{P}^2) (Lemma 2.8) imposes conditions on a new Δ sequence associated to rank matrices for the more general Hilbert function of Equation (2) (Corollary 2.9), and in particular for the almost constant Hilbert functions of Equation (1) (Corollary 2.10). After introducing the Macaulay duality (Section 2.2.2) we specify the construction of the JDT matrix (Section 2.2.3) we use throughout. Remark 2.17 recalls how to write down the Jordan degree type from the JDT matrix [Al3].

2.1 Jordan basis and Jordan degree type.

For the following definition: see [IMM, Definition 2.28(i)], [AIMM, §2.3], or [Al3]. That the JDT can be defined for any graded Artinian algebra is a consequence of results of T. Harima and J. Watanabe [HaW1] concerning the invariants of central simple modules - see [IMM, Lemma 2.2iv, Def. 2.28], also [CGo, Definition 4.1]. A further discussion of this connection is in [IMM, Definition 2.33 and Lemma 2.34].

Definition 2.1 (Jordan basis, JDT). A. Let A = R/I be an Artinian graded algebra, let M be a finite graded A-module, and let $\ell \in A_1$ be a homogeneous linear element of A. Then M may be written as a direct sum of simple $k[\ell]$

modules M(i) that we call ℓ -strings,

$$M = \oplus M(i) \text{ where } M(i) = \langle v_i, \ell v_i, \dots, \ell^{p_i - 1} v_i \rangle, 1 \le i \le t, \text{ satisfying } \ell^{p_i} v_i = 0.$$
(3)

We call this set of vector spaces $\mathsf{k}[\ell]v_i$ a *Jordan basis* of M, with respect to ℓ . The set of lengths of these strings is uniquely determined by (M, ℓ) and form the partition $P_{M,\ell}$ with t parts, the *Jordan type* of (M, ℓ) . These are simply the dimensions of the Jordan blocks of the multiplication m_{ℓ} on M.

B. The Jordan degree type $S_{M,\ell}$ supplements the Jordan type by specifying the degrees ν_i of the generators v_i of each $k[\ell]$ -module M(i). Given the decomposition of M in Equation (3) we set

$$\mathcal{S}_{M,\ell} = \{(p_i, \nu_i), \nu_i = \text{ degree of } v_i.\}$$

We may also write the Jordan degree type in list manner:

$$\mathcal{S}_{M,\ell} = \sum_{p} \sum_{\nu} (p,\nu)^{\eta(p,\nu)},\tag{4}$$

where $\eta(p,\nu)$ is the multiplicity of (p,ν) , and the sum is over distinct pairs.

For convenience we may write the pair (p, ν) as p_{ν} . Also we will write $p \uparrow_{\nu}^{\nu+\alpha}$ in place of $\sum_{i=0}^{\alpha} (p, \nu+i)$.

Although the decomposition of the $k[\ell]$ -module M into simples is not unique, for M graded the JDT is uniquely determined up to order by the pair (M, ℓ) {[IMM, Lemma 2.2(iv)].

There is a notable symmetry for Jordan degree types that occur for standard graded Gorenstein algebras. This was first shown by T. Harima and J. Watanabe [HaW1]; it is described also in [CGo, Lemma 4.6], and, in the language of JDT, in [IMM, Proposition 2.28]. We take our statement from [AIMM, Proposition 2.21].

Lemma 2.2. (Symmetry of JDT) Let A be a graded AG algebra and $\ell \in A_1$, and write the Jordan degree type in list manner $S_{A,\ell} = \sum_p \sum_{\nu} (p,\nu)^{\eta(p,\nu)}$ as above. Then we have for $\nu \leq j/2$

$$\eta(p,\nu) = \eta(p, j+1-\nu-p).$$
(5)

B. Costa and R. Gondim give a nice visualization of this symmetry in their string diagrams, see [CGo, p. 11-12 and §5].

The second author has specified a smallest socle-degree pair of graded AG algebras in codimension three having the same Jordan type, but different JDT: two AG algebras $\{A, B\}$ of Hilbert function T = (1, 3, 6, 9, 9, 9, 6, 3, 1) of socle degree 8 [Al3, Example 4.2].

Example 2.3 (Which symmetric S occur in codimension three?). Consider $S_1 = (5_0, 3_1, 3_1, 1_2)$ and $S_2 = (5_0, 3_1, 2_1, 2_2)$ that are two potential JDT having the needed symmetry for an AG algebra A of Hilbert function T = (1, 3, 4, 3, 1), and $S_3 = (4_0, (2_1)^2)$ is consistent with T = (1, 3, 3, 1). These three actually occur (Theorem 4.7 and table 7, respectively). However, the sequence $S_4 = (4 \uparrow_0^2, \mathbf{1}_{1,4})$ which is symmetric in the sense of Equation (5) and consistent with the Hilbert function $T = (1, 3^4, 1)$, does not actually occur as the JDT of a graded AG algebra, as we shall see (Theorem 4.6 and Table 7). See Figure 1.

Given a potential JDT sequence S, we denote by T(In(S)) the Hilbert function of the initial elements of the strings of S: that is $T(In(S))_i$ is the number of initial elements of the strings of S having degree i. For S_1 we have $T(In S_1) = T(In S_2) = (1, 2, 1, 0)$ but $T(In S_4) = (1, 2, 1, 0, 1)$.

Lemma 2.4. A JDT sequence S can occur for a codimension three AG algebra only if T(In S) is an O-sequence in codimension two. That is, T(In S) must occur as the Hilbert function of a graded quotient of k[y, z].

Proof. Let $I \subset R = \mathsf{k}[x, y, z]$ be an ideal such that the codimension three AG algebra A = R/I has JDT $\mathcal{S}_{A,x}$. One can compute a Gröbner basis (standard basis) of I using homogeneous degree rvlex order with z > y > x. As stated in Definition 2.1, the algebra A = R/I can be written as a direct sum: $A = \bigoplus A_m, A_m = \langle v_m, xv_m, \ldots, x^{p_{\nu_m}-1}v_m \rangle, 1 \leq m \leq t$, satisfying $x^{p_{\nu_m}}v_m = 0$ and ν_m is the degree of the form v_m . Using homogeneous degree rvlex (reverse lex) order with z > y > x, let w_m be the leading monomial of v_m . It is clear that $w_m \in \mathsf{k}[y, z]$. Since by definition, the set $\{x^{\alpha_m}v_m\}_{1\leq m\leq t, 0\leq \alpha_m\leq p_{\nu_m}-1}$ is a cobasis of the ideal I, we have that $\{w_m\}_{1\leq m\leq t}$ is a cobasis of the ideal (I + (x)). Now, let $\gamma_i = \#\{w_m, degree(w_m) = i\}$. Then the sequence of the γ_i 's is the Hilbert function of the ideal (I + (x)).

The condition of Lemma 2.4 is not sufficient for a potential JDT sequence: that is, the rank conditions of Lemma 2.7 below impose further restrictions.



Figure 1: Sequences S_1, S_2, S_3 occur as JDT (Table 8 #1,3 and Table 7 #1) but S_4 does not.

Example 2.5. Consider $S = (5 \uparrow_0^2, 2_1, 1_3, 2_4)$. The sequence S is symmetric and compatible with the Hilbert function $(1, 3, 4^3, 3, 1)$. Let γ_i be the number of strings of S starting at degree i. We have $\gamma_0 = 1, \gamma_1 = 2, \gamma_2 = 1, \gamma_3 = 1, \gamma_4 = 1$. That gives an O-sequence in codimension two: (1, 2, 1, 1, 1). One can check in Table 8 that $S = (5 \uparrow_0^2, 2_1, 1_3, 2_4)$ does not occur as JDT of an AG algebra.

Question 2.6. Let $S = \sum_{p} \sum_{\nu} (p, \nu)^{\eta(p,\nu)}$ be a symmetric sequence potential for a JDT (notation of Lemma 2.2). Let $p_1 = \text{Max} \{p\}$. Then S can occur as a JDT if and only if for all $i \in [0, p_1]$, $S = \sum_{p-i} \sum_{\nu} (p-i, \nu)^{\eta(p-i,\nu)}$ satisfies the property described in Lemma 2.4: that is, if we shorten each string by i, with the convention that we omit $(p - i, \nu)^{\eta(p-i,\nu)}$ whenever $p - i \leq 0$ then the initial terms of the strings determine an O-sequence.

Note that the question is based on the fact that if an ideal I defines an algebra whose JDT is $S = \sum_{p} \sum_{\nu} (p, \nu)^{\eta(p,\nu)}$, then the ideal quotient (I : x) defines an algebra whose JDT should be $S = \sum_{p=1} \sum_{\nu} (p-1, \nu)^{\eta(p-1,\nu)}$ (the length of each string is shortened by 1).

2.2 Rank matrices.

A result of the second author (Lemma 2.7, below, see [A11]) is a key result in determining the Jordan degree type of Artinian Gorenstein algebras. Assume that $T = (1, 3, T_1, s^k, T_1^r, 3, 1)$ where $(1, 3, T_1, s)$ has first difference an Osequence, and T_1^r is the reverse of T_1 , as in Equation (2). Any height three Gorenstein sequence can be written in this way. For any AG algebra Ain Gor(T) recall from [A11] the rank matrix $M_{A,\ell}$ for the pair (A, ℓ) where $\ell \in A_1$. Here $M_{A,\ell}$ is a symmetric $(j + 1) \times (j + 1)$ triangular matrix whose entry $r_{a,b}$ for $0 \le a \le b \le j$ is

$$r_{a,b}(\ell) = \operatorname{rk}(m_{\ell^{j-b-a}} : A_a \to A_{j-b}).$$
(6)

The following Lemma applies to all Gorenstein sequences - Hilbert functions of AG algebras - but we state it for codimension three.

Lemma 2.7 (Rank Matrix conditions). [Al1, Corollary 3.9] Let T be a codimension three Gorenstein sequence as in Equation (2). The rank matrix $M_{A,\ell}$ satisfies the following:

- (i.) The diagonal parallel to the main diagonal through (0, i) is the Hilbert function of the AG algebra $R/(I : \ell^i)$ so is a Gorenstein sequence.
- (ii.) The differences between adjacent diagonals are O-sequences [Al1, Lemma 3.6].
- *(iii.)* The rows are non-increasing, the columns non-decreasing.
- (iv.) Furthermore, each adjacent square of entries in $M_{A,\ell}$ satisfies [Al1, Lemma 3.7].

$$\begin{pmatrix} \alpha & \beta \\ \gamma & \zeta \end{pmatrix} \subset M_{A,\ell} \Rightarrow \gamma + \beta \ge \alpha + \zeta.$$
(7)

2.2.1 Map Gor(T) to the punctual Hilbert scheme.

When the Sperner number occurs at least 3 times; i.e. $k \geq 3$ there is a relation between the family Gor(T) and the punctual Hilbert scheme $\text{Hilb}^s(\mathbb{P}^2)$, that parametrizes length-s subschemes of the projective plane \mathbb{P}^2 . We assume for all parametrization results that k is an infinite field.

Lemma 2.8 (Map Gor(T) to Hilb^s(\mathbb{P}^2)). Fix a codimension three Gorenstein sequence T as in Equation (2) with Sperner number (maximum value) s that

occurs at least 3 times, $t_a = t_{a+1} = \cdots = t_b = s$. Fix $A \in \text{Gor}(T)$ and fix any linear form $\ell \in A_1$. Then the rank of $m_{\ell^k} : A_i \to A_{i+k}$ is independent of i if $a \leq i \leq b - k$. Equivalently, in the rank matrix each diagonal of $M_{A,\ell}$ parallel to the main diagonal is constant within the square corresponding to those degrees i for which $t_i = s$ (see Equation (10) for the special case of almost constant T).

Proof. Let A = R/I. By [IK, Theorem 5.31] there is a scheme $\mathfrak{Z} \subset \mathbb{P}^2$ such that $(I_3)_i = I_i$ for $a \leq i \leq b$. Fix a non-zero-divisor $\alpha \in R/I_3$. That $\ell^k \alpha = \alpha \ell^k$ acting on R/I_3 for $i \geq a$ shows the desired equality of ranks. \square

Assuming $k \geq 3$, the Lemma 2.8 implies that the entries of $M_{A,\ell}$ satisfy identities

$$\begin{aligned} r_1 &:= r_{a,j-a-1} = r_{a+1,j-a-2} = \cdots = r_{\lfloor j/2 \rfloor, j-\lfloor j/2 \rfloor - 1} \\ r_2 &:= r_{a,j-a-2} = r_{a+1,j-a-3} = \cdots = r_{\lfloor j/2 \rfloor, j-\lfloor j/2 \rfloor - 2} \\ \vdots \\ r_{k-1} &:= r_{a,k-1}. \end{aligned}$$

Note that $r_1 \ge 1$ since otherwise $\ell \in I$. By Lemma 2.7 (iv), we have

$$2r_i \ge r_{i-1} + r_{i+1}$$
 for all $i = 1, \dots, k-2$. (8)

We define the following sequences of non-negative integers (taking $r_0 = s$)

$$\vec{r} := (r_1, \dots, r_{k-1}); \vec{\Delta} := \{ \Delta r_i, i = 1, \dots, k-1 \}, \quad \Delta r_i = r_{i-1} - r_i.$$
(9)

We let p(u, t) denote the number of partitions of u whose largest part is less or equal t, and p(u) is the number of partitions of u. We will sometimes denote $\overrightarrow{\Delta}$ by Δr or Δ .

Corollary 2.9 (The Δ sequence). Assume that $T = (1, 3, T_1, s^k, T_1^r, 3, 1)$ with $k \geq 3$ is a codimension three Gorenstein sequence as in Equation (2). We have the following:

- (a) the sequence $\overrightarrow{\Delta}$ is non-increasing;
- (b) $\Delta r_i \leq s 1$ for all *i*.

- (c) $\Sigma_{i=1}^{k-1} \Delta r_i = s r_{k-1} \leq s;$
- (d) $\Delta r_i = 0$ for all $i \ge r_0 + 1 = s + 1$, so $r_s = r_{s+1} = \cdots = r_{k-1}$ when $k \ge s + 1$.

Proof. Item (a) is equivalent to Equation (8). Parts (b),(c) follow from Lemma 2.7: Part (b) $\Delta r_i \leq s - 1$ because $r_1 \geq 1$ and $\overrightarrow{\Delta}$ is non-increasing; and Part (c) because we have $\sum_i \Delta r_i + r_{k-1} = r_0$. From (a),(b) and (c), we conclude that the sequence $\overrightarrow{\Delta}$ is a partition of some integer $r \leq s$, which proves part (d).

In this paper henceforth we study the almost constant Hilbert functions of Equation (1), that is

$$T = (1, 3, s^k, 3, 1)$$
 with $3 \le s \le 6$ and $1 \le k$.

Lemma 2.8 for such T when $k \ge 3$ implies the following

Corollary 2.10 (The Δ sequence for an almost constant Hilbert function). Assume T satisfies (1). Then the rank matrix $M_{A,\ell}$ of any pair (A, ℓ) where $A \in \text{Gor}(T)$ and $\ell \in \mathfrak{m}$ has the following form

$$M_{A,\ell} = \begin{pmatrix} 1 & r_{0,j-1} & r_{0,j-2} & r_{0,j-3} & r_{0,j-4} & r_{0,j-5} & \dots & r_{0,2} & r_{0,1} & r_{0,0} \\ 3 & r_{1,j-2} & r_{1,j-3} & r_{1,j-4} & r_{1,j-5} & \dots & r_{1,2} & r_{1,1} & r_{0,1} \\ & s & r_1 & r_2 & r_3 & \dots & r_{k-1} & r_{1,2} & r_{0,2} \\ & s & r_1 & r_2 & \dots & r_{k-2} & r_{1,3} & r_{0,3} \\ & & s & r_1 & \dots & r_{k-3} & r_{1,4} & r_{0,4} \\ & & & \ddots & \ddots & \vdots & \vdots & \vdots \\ & & & & \ddots & r_1 & r_{1,j-3} & r_{0,j-3} \\ & & & s & r_{1,j-2} & r_{0,j-2} \\ & & & & 3 & r_{0,j-1} \\ & & & & 1 \end{pmatrix},$$

$$(10)$$

where the entries below the main diagonal are all zero. Because of the connection of such an AG algebra with a punctual Hilbert scheme (Lemma 2.8), the $k \times k$ upper triangular matrix in the center (colored black) is a Toeplitz matrix whose main diagonal is s and whose constant diagonals parallel to the main diagonal are r_1, \ldots, r_{k-1} . **Remark 2.11.** Note that if T satisfies Equation 1 for k at least 3, then $\Delta r_1 \leq 3$ since otherwise the first difference of the diagonals in the rank matrix would not be an O-sequence. Take for example s = 6, $\Delta r_1 = 4$ that the first two diagonals of $M_{A,\ell}$ are

	1	2	4	4	
_	0	1	2	2	· · · ,
+	1	3	6	6	

whose difference (1, 2, 4, 4, ...) is not an O-sequence.

2.2.2 Macaulay duality.

Definition 2.12 (Macaulay dual). We let the ring R = k[x, y, z] act on S = k[X, Y, Z] as contraction, that is

$$x^{i} \circ X^{j} = \begin{cases} X^{j-i} \text{ if } j \ge i;\\ 0 \text{ otherwise,} \end{cases}$$
(11)

similarly for $y^i \circ Y^j$, $z^i \circ Z^j$, to monomials of R acting on monomials of S, and we extend linearly to an action of R on S. Given $F \in S$ we denote by $I_F = \operatorname{Ann}(F) = \{h \in R \mid h \circ F = 0\}$. Given a vector subspace $V \subset R_j$ we denote by $V^{\perp} \subset S_j$ the space $V^{\perp} = \{w \in S_j \mid v \circ w = 0 \text{ for all } v \in V\}$.

For example $x^2yz^2 \circ (X^3YZ^3 - X^2Y^3Z^2) = XZ - Y^2$, and for $F = X^3 + Y^3 + Z^3$, $A_F = R/I_F$ where $I_F = \text{Ann}(F) = (xy, xz, yz, x^3 - y^3, x^3 - z^3)$.

Lemma 2.13. There is a 1-1 correspondence between

- *i.* Elements $F \in S_j$, up to non-zero constant multiple, and
- ii. AG quotients A = R/I of socle-degree j, given by

$$F \to A_F = R/I_F, I_F = \operatorname{Ann}(F), \text{ and}$$
$$A = R/I \text{ of socle degree } j \to F = I_j^{\perp} \in S_j.$$
(12)

For $\ell \in R_1$ we have $\operatorname{Ann}(\ell \circ F) = I_F : \ell$.

Corollary 2.14. The rank matrix $M_{A_F,\ell}$ for an AG algebra has main diagonal the Hilbert function $H(A_F)$, $A_F = R/I_F$ and parallel diagonals the Hilbert functions of $A_{\ell^a \circ F} a = 1, 2, ...$

Proof. This follows from Lemma 2.7 (a), and from $\operatorname{Ann}(\ell \circ F) = I_F : \ell$ from Lemma 2.13.

Example 2.15. For $F = X^3 + Y^3 + Z^3$ and $\ell = x + y + z$ the diagonals of $M_{A,\ell}$ are (1,3,3,1), (1,3,1), (1,1) and 1, respectively arising from $F, \ell \circ F = X^2 + Y^2 + Z^2, \ell^2 \circ F = X + Y + Z$ and $\ell^3 \circ F = (3)$ - unless the characteristic of k is three, in which case the last diagonal is 0.

2.2.3 Jordan Degree Type matrix.

Definition 2.16. Suppose that $M_{A,\ell}$ is the rank matrix of an AG algebra A and the linear form ℓ . Then the *Jordan degree type matrix (JDT matrix)*, $J_{A,\ell}$, of the pair (A, ℓ) is defined to be the upper triangular matrix with the following entries

$$(J_{A,\ell})_{u,v} := (M_{A,\ell})_{u,v} + (M_{A,\ell})_{u-1,v+1} - (M_{A,\ell})_{u-1,v} - (M_{A,\ell})_{u,v+1},$$
(13)

where we set $(M_{A,\ell})_{u,v} = 0$ if either u < 0 or v < 0. We may write this as $J_{A,\ell} = J_M$, as it depends only on the matrix M.

Lemma 2.7(iv.) implies that the entries of $J_{A,\ell}$ are all non-negative. For every $0 \le u \le v \le j$ if $J_{u,v} \ne 0$ then in the Jordan degree type $\mathcal{S}_{A,\ell}$ the multiplicity of the pair (p,ν) where p = v - u + 1 and $\nu = u$ is $\eta(p,\nu) = J_{u,v}$. See [Al1, Al2] for more detail.

Remark 2.17 (Relation between JDT matrix $(J_{A,\ell})$ and JDT $(\mathcal{S}_{A,\ell})$). The row of the JDT matrix gives the index, which is the initial degree of the corresponding entry of $\mathcal{S}_{A,\ell}$. The diagonal (with main diagonal counted as 1) gives the value, so the *w*-th diagonal parallel to the main diagonal gives an entry *w* of \mathcal{S} with subscript the row (first row counted as zero).

Example 2.18. Let $R = k[x_1, x_2, x_3], A = R/I, I = Ann(X_1^2 X_2 X_3) = (x_1^3, x_2^2, x_3^2)$ and $\ell = x_1$. The Hilbert function of A is T = (1, 3, 4, 3, 1). The following are the rank matrix and JDT matrix for (A, ℓ)

The JDT for $\ell = x_1$ and A is $S_{A,\ell} = (3_0, 3_1^2, 3_2)$, which can be seen from $A \cong k[x_1]/(x_1^3) \otimes k[x_2, x_3]/(x_2^2, x_3^2)$, so $S_{A,\ell} = (3_0) \otimes (1_0, 1_1^2, 1_2^2)$. Or we can recover $S_{A,\ell}$ directly from $M_{A,\ell}$ working consecutively, from larger to smaller a, since $T(a) := T(R/(I : x_1^a))$ is the a-th diagonal above and parallel to the main diagonal T(A) of M_{A,x_1} [Al1, Proposition 3.4]. So we have from Equation (14) that $T(3) = T(R/(I : x_1^3)) = 0$ implying $x_1^3 \in I$. The diagonal $(1, 2, 1) = T(R/(I : x_1^2))$; since $x_1^3 \in I$ this implies $\mathcal{S}_{R/(I:x_1^2),x_1} = (1_0, 1_1^2, 1_2)$. Then $T(R/(I : x_1)) = (1, 3, 3, 1)$ implies $\mathcal{S}_{R/(I:x_1),x_1} = (2_0, 2_1^2, 2_2)$, and we conclude from T(A) = (1, 3, 4, 3, 1) that $S_{A,x_1} = (3_0, 3_1^2, 3_2)$. An alternative approach is to determine $\mathcal{S}_{A,\ell}$ from $J_{A,\ell}$ using Definition 2.16 and the discussion after it. By Equation (13) we get that the only non-zero entries of $J_{A,\ell}$ are $(J_{A,\ell})_{0,2} = 1, (J_{A,\ell})_{1,3} = 2$, and $(J_{A,\ell})_{2,4} = 1$. Therefore, $\mathcal{S}_{A,\ell} = (3_0, 3_1^2, 3_2)$.

An example of constructing an algebra from a particular rank matrix for T = (1, 3, 4, 3, 1) is found in [Al3, Example 4.1]. See also Section 4.3 below.

As in the Example 2.18 the rank matrix of a graded AG algebra determines the Jordan degree type, and vice-versa [Al1, Proposition 3.12]. That the rank matrix determines the JDT is essentially the result that the ranks of mixed Hessians determine JDT (see [GonZa, CGo]): this rests on the work of T. Harima and J. Watanabe on central simple modules [HaW1].

3 Potential Jordan degree types.

In this section we assume that k is an infinite field of arbitrary characteristic. We denote by $\mathcal{J}(T)$ the set of Jordan degree types (JDT) possible for pairs $(A, \ell), A \in \text{Gor}(T), \ell \in \mathfrak{m}$. We recall the Jordan degree types for complete intersections of codimension two, then in codimension three determine the potential Jordan types for almost constant Hilbert functions of each Sperner number 3, 4, 5, 6 (Section 3.2).

3.1 Codimension two.

The number of possible rank matrices, equivalently JDT, for codimension two complete intersection (CI) Hilbert functions was obtained explicitly in [AlIK]. We first state a portion of that result - giving the count. We then in an Example list the rank matrices in codimension two corresponding to Sperner numbers s = 2, 3.

#	$\overrightarrow{\Delta}$	\overrightarrow{r}	# rk matrices	JDT
1_a	(0, 0)	$(2,2,\ldots,2)$	2	$((j+1)_0, (j-1)_1)$
1_b				(j_0, j_1)
2_a	(1, 0)	$(2,1,\ldots,1)$	1	$((j+1)_0, 1\uparrow_1^{j-1})$
3_b	(1, 1)	$(2, 1, 0, \ldots, 0)$	1	$(2\uparrow_0^{j-1})$

Table 1: Case $s = 2, k \ge 2$; codimension two: the action of ι takes 1_b to 3_b .

Recall that the Hilbert function of a codimension two complete intersection of Sperner number d and socle degree j = k + 2d - 2 satisfies

$$T = (1, 2, \dots, d - 1, d^k, d - 1, \dots, 2, 1) \text{ where } k \ge 1.$$
(15)

For such T, the automorphism ι on $\mathcal{J}(T)$ flips (switches rows and columns) in the smallest rectangle in the Jordan type, and makes the corresponding change for the JDT (see [AlIK, Theorem 3.23]). For example, the Jordan degree type $\mathcal{S} = ((j+1)_0, (j-1)_1, (j-1)_2)$ satisfies $\iota(\mathcal{S}) = ((j+1)_0, 2\uparrow_1^{j-1})$.

The following is adapted from [AlIK, Theorems 2.21, 2.35], which have further very detailed combinatorial data.

Theorem 3.1. [AlIK] Let T satisfy Equation (15). When k = 1 there are 2^{d-1} Jordan types possible for $A \in CI(T)$, all weak-Lefschetz - with d parts. When $k \ge 2$ there are 2^d Jordan types possible for $A \in CI(T)$: half are weak Lefschetz - with d parts, and half, with d + k - 1 parts are of the form $\iota(S)$. When k = 1, the smallest part α of S occurs α times, so $\iota(S) = S$.

Example 3.2. In Tables 1 and 2, respectively, we list the possible rank matrices for codimension two, almost constant CI sequences of Sperner numbers 2 and 3, respectively.

Note that in Table 1, the first row $\Delta = 0$ corresponds to the two JDT $((j+1)_0, (j-1)_1)$ and (j_0, j_1) . The second row corresponds to $\iota(((j+1)_0, (j-1)_1) = (j+1)_0, 1\uparrow_1^{j-1})$ and the third row to $\iota((j_0, j_1)) = (2\uparrow_0^{j-1})$.

Another aspect of this case, the role of Δ in determining repeated parts of $S_{A,\ell}$, is shown in Proposition 4.3 and Example 4.4(c) below.

3.2 Codimension three.

In this section we count the number of potential rank matrices and Jordan degree types for codimension three AG algebras with almost constant Hilbert

#	$\overrightarrow{\Delta}$	\overrightarrow{r}	# rk matrices	JDT
1_a	(0, 0, 0)	$(3,3,3,3\ldots,3)$	4	$((j+1)_0, (j-1)_1, (j-3)_2)$
1_b				$(j_0, j_1, (j-3)_2)$
1_c				$((j+1)_0, (j-2)_{1,2})$
1_d				$((j-2)\uparrow_0^2)$
2_a	(1, 0, 0)	$(3, 2, 2, 2, \ldots, 2)$	2	$((j+1)_0, (j-1)_1, 1\uparrow_2^{j-2})$
2_b				$(j_0, j_1, 1\uparrow_2^{j-2})$
3_c	(1, 1, 0)	$(3, 2, 1, 1, \dots, 1)$	1	$((j+1)_0, 2\uparrow_1^{j-2})$
4_d	(1, 1, 1)	$(3, 2, 1, 0, \dots, 0)$	1	$3\uparrow_0^{j-2}$

Table 2: Case $s = 3, k \ge 2$, codimension 2. The action of ι takes 1_a to 2_a . See [AlIK, Figure 10].

functions. More precisely, for each Sperner number $3 \le s \le 6$ we count all possible rank matrices for each possible sequence $\overrightarrow{\Delta}$ of Equation (9) satisfying the conditions in Lemma 2.7 and Corollary 2.9.

3.2.1 Case s = 3.

Let $T = (1, 3^k, 1), k \ge 3$. Here we assume that T is the Hilbert function of codimension three AG algebras.

In this case the rank matrix has the following form.

$$M = \begin{pmatrix} 1 & r_{0,j-1} & r_{0,j-2} & r_{0,j-3} & r_{0,j-4} & \dots & r_{0,1} & r_{0,0} \\ 3 & r_1 & r_2 & r_3 & \dots & r_{k-1} & r_{0,1} \\ & 3 & r_1 & r_2 & \dots & r_{k-2} & r_{0,2} \\ & & 3 & r_1 & \dots & r_{k-3} & r_{0,3} \\ & & & \ddots & \ddots & \vdots & \vdots \\ & & & & \ddots & r_1 & r_{0,j-2} \\ & & & & & & 1 \end{pmatrix}.$$
(16)

Theorem 3.3. Let $T = (1, 3^k, 1)$ for $k \ge 3$. There are 8 potential Jordan degree types for pairs (A, ℓ) , where $A \in \text{Gor}(T)$ and $\ell \in A_1$.

Proof. First, we count possible sequences $\overrightarrow{\Delta}$ that is equal to p(0) + p(1) + p(2) + p(3,2) = 1 + 1 + 2 + 2 = 6. We list them in Table 3. Every rank matrix

is determined by the sequence \overrightarrow{r} (equivalently $\overrightarrow{\Delta}$ and entries $r_{0,0}, \ldots, r_{0,j-1}$ in the matrix given in Equation (16). But since each diagonal of the rank matrix is a Gorenstein sequence the entries $r_{0,2}, r_{0,3}, \ldots, r_{0,j-1}$ are determined by the sequence \overrightarrow{r} . Therefore, for a given sequence \overrightarrow{r} and values $r_{0,1}$ and $r_{0,0}$ the rank matrix is determined. Observe that $r_{0,1}$ and $r_{0,0}$ are the entries of the top right 2×2 submatrix of M as in Equation (16). Thus to find the possible rank matrices in this case we find the possible sequences \overrightarrow{r} and for each of these sequences we find the possible submatrices $\binom{r_{0,1} \quad r_{0,0}}{r_{k-1} \quad r_{0,1}}$.

For $\overrightarrow{\Delta} = (0, 0, 0)$, we have $\overrightarrow{r} = (3, 3, \dots, 3)$. So in particular $r_{k-1} = 3$ and there are two possible choices for the top right corner of the rank matrix:

$$\begin{pmatrix} 1 & 1 \\ 3 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 3 & 1 \end{pmatrix}.$$

For $\overrightarrow{\Delta} = (1,0,0), (1,1,0)$, and (1,1,1) the second diagonal of the rank matrix is $(1,2^{k-1},1)$ that is a codimension two Gorenstein Hilbert function. Therefore, the number of possible rank matrices can be obtained from the corresponding row of Table 1.

For $\overrightarrow{\Delta} = (2, 0, 0)$ we have $r_{k-1} = 1$ and this forces $r_{0,1} = r_{0,0} = 1$. Also, clearly for $\overrightarrow{\Delta} = (2, 1, 0)$ we have $r_{0,1} = r_{0,0} = 0$ forced by $r_{k-1} = 0$.

#	$\overrightarrow{\Delta}$	$\overrightarrow{r} = (r_0, \dots, r_{k-1})$	#rk matrices	JDT	Table 7
1a	(0,0,0)	$(3,3,3,3\ldots,3)$	2	$((j+1)_0, ((j-1)_1)^2)$	#1
1b				$(j_0, j_1, (j-1)_1)$	#5
2a	(1, 0, 0)	$(3,2,2,2,\ldots,2)$	2	$((j+1)_0, (j-1)_1, 1\uparrow_1^{j-1})$	#2
2b				$(j_0, j_1, 1\uparrow_1^{j-1})$	#6
3	(2, 0, 0)	$(3, 1, 1, 1, \dots, 1)$	1	$((j+1)_0, (1\uparrow_1^{j-1})^2)$	#4
4	(1, 1, 0)	$(3, 2, 1, 1, \ldots, 1)$	1	$((j+1)_0, 2\uparrow_1^{j-2}, 1_{1,j-1})$	#3
5	(2, 1, 0)	$(3,1,0,0,\ldots,0)$	1	$(2\uparrow_0^{j-1},(1\uparrow_1^{j-1})$	#8
6	(1, 1, 1)	$(3,2,1,0,\ldots,0)$	1	$(3\uparrow_0^{j-2}, 1_{1,j-1})$	#7

Table 3: Case s = 3, $T = (1, 3^k, 1)$, $k \ge 3$, codimension three; list of JDTs and reference to corresponding entry of Table 7.

Example 3.4. (i) When s = 3, j = 4 and $\overrightarrow{\Delta} = (0, 0, 0)$ then $\overrightarrow{r} = (3, 3, 3)$ and the upper right corner of the rank matrix M is (they are respectively #1a, #1b in Table 3, for j = 4.)

$$\begin{pmatrix} 1 & 1 \\ 3 & 1 \end{pmatrix}$$
 $\mathcal{S} = (5_0, (3_1)^2);$ or $\begin{pmatrix} 1 & 0 \\ 3 & 1 \end{pmatrix}$ $\mathcal{S} = (4_0, 4_1, 3_1).$

(ii) When s = 3, j = 4 and $\overrightarrow{\Delta} = (1, 0, 0)$ then $\overrightarrow{r} = (3, 2, 2, ...)$ and the upper right corner of M is (we give the JDT for j = 4 here, they are, respectively, #2a, #2b in Table 3)

$$\begin{pmatrix} 1 & 1 \\ 2 & 1 \end{pmatrix} \quad \mathcal{S} = ((5)_0, (3)_1, 1\uparrow_1^3); \quad \text{or} \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix} \quad \mathcal{S} = (4_0, 4_1, 1\uparrow_1^3).$$

(iii) When s = 3, j = 4 and $\overrightarrow{\Delta} = (2, 0, 0)$, then $\overrightarrow{r} = (3, 1, 1)$ we have

$$\mathbf{M} = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 \\ 3 & 1 & 1 & 1 \\ & 3 & 1 & 1 \\ & & 3 & 1 \\ & & & 1 \end{pmatrix}, \quad J = \begin{pmatrix} 0 & 0 & 0 & 0 & 1 \\ 2 & 0 & 0 & 0 \\ & 2 & 0 & 0 \\ & & & 2 & 0 \\ & & & & 0 \end{pmatrix}, \quad \mathcal{S} = (5_0, (1 \uparrow_1^3)^2),$$

as in the row 3 of Table 3 for j = 4. Here the Jordan degree type matrix J is that of Definition 2.16, from which the JDT can be read off simply.

3.2.2 Case s = 4.

Let $T = (1, 3, 4^k, 3, 1), k \ge 3$. The number of possible sequences $\overrightarrow{\Delta}$ (so \overrightarrow{r}) is equal to

$$p(0) + p(1) + p(2) + p(3) + p(4, 3) = 1 + 1 + 2 + 3 + (5 - 1) = 11.$$

In the following theorem, we find the number of rank matrices, which are of the form (10), for each of those sequences and list them in Table 4 below.

Theorem 3.5. Let $T = (1, 3, 4^k, 3, 1)$ for $k \ge 3$. There are 26 potential Jordan degree types for (A, ℓ) where $A \in Gor(T)$ and $\ell \in A_1$.

Proof. To get the possible rank matrices for $\overrightarrow{\Delta} = (0, 0, 0, 0)$ we note that since the diagonals are Hilbert functions for AG algebras, we have $r_{1,j-2} =$

 $\cdots = r_{1,3} = 3$, and $r_{0,4} = \cdots = r_{0,j-1} = 1$. It is left to count the possibilities of the matrix

 $\begin{pmatrix} 1 & r_{0,3} & r_{0,2} & r_{0,1} & r_{0,0} \\ 3 & 3 & r_{1,2} & r_{1,1} & r_{0,1} \\ 4 & 4 & 4 & r_{1,2} & r_{0,2} \\ 4 & 4 & 4 & 3 & r_{0,3} \\ 4 & 4 & 4 & 3 & 1 \end{pmatrix}.$

By Lemma 2.8 (iv), $r_{1,2} = 3$ and $r_{0,3} = r_{0,2} = 1$, and we have the following 5 choices for the upper right corner:

$$\begin{pmatrix} 1 & 0 & 0 \\ 3 & 2 & 0 \\ 4 & 3 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 0 \\ 3 & 2 & 1 \\ 4 & 3 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 1 \\ 3 & 2 & 1 \\ 4 & 3 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 0 \\ 3 & 3 & 1 \\ 4 & 3 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 1 \\ 3 & 3 & 1 \\ 4 & 3 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 1 \\ 3 & 3 & 1 \\ 4 & 3 & 1 \end{pmatrix}$$

Now suppose the first entry of $\overrightarrow{\Delta}$ is one, or equivalently $r_1 = 3$, so we have either $r_{1,j-2} = 2$ or 3. If $r_{1,j-2}$ is 3, the number of possible rank matrices follows from Table 3. If $r_{1,j-2} = 2$, the number of possible rank matrices follows from the case $T(1) = (1, 2, 3^{k-1}, 2, 1)$, (see Table 2). For instance, in #2 in Table 4, the number of rank matrices is 6 such that 2 of them correspond to having $T(1) = (1, 3^{k+1}, 1)$ for which we look at #1 in Table 3 that counts 2 possible rank matrices. Similarly, if $T(1) = (1, 2, 3^{k-1}, 2, 1)$ the 4 rank matrices are obtained from #1 in Table 2. These two sequences contribute to cases #2, 4, 7 and 11 (when $r_1 = 3$).

When $r_1 = 2$, $r_{1,j-2}$ is also equal to 2, and the number of possible rank matrices follows from Table 1.

When $r_{k-1} = 1$ (respectively $r_{k-1} = 0$), the top right 3×3 -submatrix of $M_{A,\ell}$ is forced to be $\begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}$, respectively $\begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$.

3.2.3 Case s = 5.

Let $T = (1, 3, 5^k, 3, 1), k \ge 3$. The number of possible sequences $\overrightarrow{\Delta}$ (equivalently \overrightarrow{r}) is equal to p(0) + p(1) + p(2) + p(3) + p(4, 3) + p(5, 3) = 16. For each of the sequences $\overrightarrow{\Delta}$ and equivalently \overrightarrow{r} such that r_1 (second entry of \overrightarrow{r}) is less than 5 we count the number of potential rank matrices recursively using previous results for s = 3, 4 and for sequences $(r_1, r_2, \ldots, r_{k-1})$; i.e. removing

#	$\overrightarrow{\Delta} = (\Delta r_1, \dots, \Delta r_4)$	$\overrightarrow{r} = (r_0, r_1, \dots, r_{k-1})$	#rk m	atrices
1	(0, 0, 0, 0)	$(4, 4, 4, 4, 4, \ldots, 4)$	5	0
2	(1, 0, 0, 0)	$(4, 3, 3, 3, 3, \ldots, 3)$	2	4
3	(2, 0, 0, 0)	$(4, 2, 2, 2, 2, \ldots, 2)$	2	0
4	(1, 1, 0, 0)	$(4, 3, 2, 2, 2, \ldots, 2)$	2	2
5	(3,0,0,0)	$(4, 1, 1, 1, 1, \dots, 1)$	1	0
6	(2, 1, 0, 0)	$(4, 2, 1, 1, 1, \dots, 1)$	1	0
7	(1, 1, 1, 0)	$(4,3,2,1,1,\ldots,1)$	1	1
8	(3, 1, 0, 0)	$(4, 1, 0, 0, 0, \dots, 0)$	1	0
9	(2,2,0,0)	$(4, 2, 0, 0, 0, \ldots, 0)$	1	0
10	(2, 1, 1, 0)	$(4, 2, 1, 0, 0, \dots, 0)$	1	0
11	(1, 1, 1, 1)	$(4, 3, 2, 1, 0, \dots, 0)$	1	1

Table 4: Case s = 4: the sequences Δ and number of rank matrices.

 r_0 from \overrightarrow{r} , and the appropriate T for which the Sperner number occurs k-1 times. Thus we have the following for s = 5 and $k \ge 4$.

Theorem 3.6. Let $T = (1, 3, 5^k, 3, 1)$ for $k \ge 4$. There are 47 potential Jordan degree types for (A, ℓ) where $A \in Gor(T)$ and $\ell \in A_1$.

Proof. The potential sequences for $\overrightarrow{\Delta}$ and \overrightarrow{r} are listed in Table 5. Since by assumption $r_0 = 5$ we must have $0 \leq \Delta r_0 = r_0 - r_1 < 4$. If $\Delta r_0 = r_0$ we have $r_1 = 0$ which is equivalent to having the AG algebra $A/(0:\ell) = 0$ which is impossible for $\ell \neq 0$. Also if we have $\Delta r_0 = 4$ when $r_0 = 5$ we have $r_1 = 1$ which means that all the entries on the second diagonal of the rank matrix $M_{A,\ell}$, which gives the Hilbert function of $A/(0:\ell)$, are equal to one. By Lemma 2.7 part (ii), the difference between any two consecutive diagonals of the rank matrix must be an O-sequence but $(1,3,5^k,3,1) - (1^{k+3}) = (1,2,4,4,\ldots)$ clearly is not an O-sequence. It is left to study the possible values of $r_{0,0}, \ldots r_{0,j-1}$ and $r_{1,1}, \ldots, r_{1,j-2}$. Note that if r_u is zero, then $r_{0,j-u} = r_{1,j-u-1} = 0$ (the whole diagonal containing r_u is zero.)

For case #1, since the diagonals are Hilbert functions for AG algebras, then $r_{1,j-2} = \cdots = r_{1,3} = 3$, and $r_{0,4} = \cdots = r_{0,j-1} = 1$. It is left to count the possibilities for the matrix

#	$\overrightarrow{\Delta} = (\Delta r_1, \dots, \Delta r_5)$	$\overrightarrow{r} = (r_0, r_1, \dots, r_{k-1})$	#rk m	atrices
1	$\left(0,0,0,0,0 ight)$	$(5, 5, 5, 5, 5, 5, 5, \dots, 5)$	7	0
2	(1, 0, 0, 0, 0)	$(5, 4, 4, 4, 4, 4, \ldots, 4)$	5	0
3	$\left(2,0,0,0,0 ight)$	$(5,3,3,3,3,3,3,\ldots,3)$	2	4
4	(1, 1, 0, 0, 0)	$(5, 4, 3, 3, 3, 3, \ldots, 3)$	6	0
5	(3, 0, 0, 0, 0)	$(5, 2, 2, 2, 2, 2, \ldots, 2)$	2	0
6	(2, 1, 0, 0, 0)	$(5, 3, 2, 2, 2, 2, \ldots, 2)$	2	2
7	(1, 1, 1, 0, 0)	$(5, 4, 3, 2, 2, 2, \ldots, 2)$	4	0
8	(3, 1, 0, 0, 0)	$(5, 2, 1, 1, 1, 1, \dots, 1)$	1	0
9	$\left(2,2,0,0,0 ight)$	$(5,3,1,1,1,1,\dots,1)$	1	0
10	(2, 1, 1, 0, 0)	$(5, 3, 2, 1, 1, 1, \dots, 1)$	1	1
11	(1, 1, 1, 1, 0)	$(5, 4, 3, 2, 1, 1, \dots, 1)$	2	0
12	$\left(3,2,0,0,0 ight)$	$(5, 2, 0, 0, 0, 0, \dots, 0)$	1	0
13	(3, 1, 1, 0, 0)	$(5, 2, 1, 0, 0, 0, \dots, 0)$	1	0
14	(2, 2, 1, 0, 0)	$(5,3,1,0,0,0,\ldots,0)$	1	0
15	(2, 1, 1, 1, 0)	$(5, 3, 2, 1, 0, 0, \dots, 0)$	1	1
16	(1, 1, 1, 1, 1)	$(5, 4, 3, 2, 1, 0, \dots, 0)$	2	0

Table 5: Case s = 5: the sequences Δ and number of rank matrices.

(1)	$r_{0,3}$	$r_{0,2}$	$r_{0,1}$	$r_{0,0}$
3	3	$r_{1,2}$	$r_{1,1}$	$r_{0,1}$
5	5	5	$r_{1,2}$	$r_{0,2}$
5	5	5	3	$r_{0,3}$
$\sqrt{5}$	5	5	3	1

By Lemma 2.8 (iv), $r_{1,2} = 3$ and $r_{0,3} = r_{0,2} = 1$. We have the following choices:

$$\begin{pmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 5 & 3 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 1 \\ 3 & 1 & 1 \\ 5 & 3 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 3 & 2 & 0 \\ 5 & 3 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 0 \\ 3 & 2 & 1 \\ 5 & 3 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 1 \\ 3 & 2 & 1 \\ 5 & 3 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 0 \\ 3 & 3 & 1 \\ 5 & 3 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 1 \\ 3 & 3 & 1 \\ 5 & 3 & 1 \end{pmatrix}$$

For cases #2,4,7,11, and 16, (i.e. $r_1 = 4$, and therefore $r_{1,j-2} = 3$) the second diagonal is $(1,3,4^{k-1},3,1)$. Hence, for $k \ge 4$, the possible rank matrices follow from the case $(1,3,4^k,3,1)$, (see Table 4).

For cases #3,6,10, and 15 $(r_1 = 3 \text{ and } r_2 \ge 2)$, $r_{1,j-2}$ is either 2 or 3. If it is equal to 3 (respectively 2), then the possible rank matrices follow from the case $(1, 3^{k+1}, 1), k \ge 4$ (Table 3), (respectively $(1, 2, 3^{k-3}, 2, 1), k \ge 4$, (Table 2)).

Where we have $r_1 = 3$ and $r_2 = 1$, cases #9 and #14, we must have $r_{1,j-2} = 3$. Since otherwise if $r_{1,j-2} = 2$ the difference between the second and third diagonal of the rank matrix is given by (1, 2, 3, 3, ...) - (1, 1, 1, ...) = (1, 1, 2, 2, ...); that is not an O-sequence contradicting the condition (ii) in Lemma 2.7. Thus the number of rank matrices in these two cases are given by the number of corresponding rank matrices for $(1, 3^{k+1}, 1)$; i.e. cases #3 and #5 in Table 3.

For the remaining cases, #5,8,12, and #13 where $r_1 = 2$, we have $r_{1,j-2} = 2$, the possible rank matrices follow from the case $(1, 2^{k-1}, 1)$ (Table 1).

Remark 3.7. The number of rank matrices for each row in Tables 4 and 5 is divided into two columns where the left (respectively, right) column indicates the number of rank matrices for which the diagonal after the main diagonal is the Hilbert function of some codimension three (respectively, codimension two) AG algebra.

Now we count the number of rank matrices when s = 5 and k = 3.

Theorem 3.8. Let $T = (1, 3, 5^k, 3, 1)$ and k = 3 then there are 43 potential Jordan degree types for pairs (A, ℓ) .

Proof. When k = 3 the vector \overrightarrow{r} has 3 non-zero entries. The number of rank matrices for $\overrightarrow{r} = (5, 2, 1)$ is equal to the sum of the number of rank matrices in rows #8 and #13 in Table 5. More precisely, the rank matrices for $\overrightarrow{r} = (5, 2, 1)$ such that $(r_{0,3}, r_{1,2}, r_{1,2}, r_{0,3})$ is the one or zero vector corresponding to #8 and #13 respectively.

Similarly, the number of rank matrices for $\overrightarrow{r} = (5, 3, 1)$ is equal to the number of rank matrices in the rows #9 and #14 where we have $(r_{0,3}, r_{1,2}, r_{1,2}, r_{0,3})$ equal to one and zero vector respectively.

The number of rank matrices for $\overrightarrow{r} = (5,3,2)$ is given by the sum of rank matrices counted in rows #6, #10, and #15. The number of rank matrices in #6 corresponds to the case where $(r_{0,3}, r_{1,2}, r_{1,2}, r_{0,3}) = (1,2,2,1)$ and $(r_{0,2}, r_{1,1}, r_{0,2}) = (1,2,1)$. Row #10 counts the number of rank matrices where $(r_{0,3}, r_{1,2}, r_{1,2}, r_{0,3}) = (1,1,1,1)$ and $(r_{0,2}, r_{1,1}, r_{0,2}) = (1,1,1)$ and row #15 counts the rank matrices where $(r_{0,3}, r_{1,2}, r_{1,2}, r_{0,3}) = (1,1,1,1)$ and $(r_{0,2}, r_{1,1}, r_{0,2}) = (1,1,1,1)$ and $(r_{0,2}, r_{1,1}, r_{0,2}) = (0,0,0)$.

Finally, rows #4,#7,#11, and #16 correspond to $\overrightarrow{r} = (5,4,3)$. We notice that there are only 2 rank matrices where $(r_{0,3}, r_{1,2}, r_{1,2}, r_{0,3}) = (1,3,3,1)$ and $(r_{0,2}, r_{1,1}, r_{0,2}) = (1,3,1)$ which is either $r_{0,0} = 1$ or $r_{0,0} = 0$. Therefore, the number of rank matrices where $\overrightarrow{r} = (5,4,3)$ is 4 less than the sum of the rank matrices equal to the sum of the number of rank matrices in #4,#7,#11, and #16 that is equal to 10. We conclude that when k = 3there are 43 potential rank matrices.

3.2.4 Case s = 6.

Let $T = (1, 3, 6^k, 3, 1)$. The partition $\Delta = (3, 3, 0, 0, ...)$ that corresponds to r = (6, 3, 0, 0, ...) cannot occur in this case since otherwise, if $T(1) = (1, 2, 3^{k-1}, 2, 1)$ the difference between the first two diagonals would not be an *O*-sequence, and if $T(1) = (1, 3^{k+1}, 1)$, then the submatrix

$$\begin{pmatrix} r_{0,j-1} & r_{0,j-2} \\ s & r_1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 3 & 3 \end{pmatrix}$$

would contradict Lemma 2.7. The number of possible sequences for $\overrightarrow{\Delta}$ is therefore

$$p(0) + p(1) + p(2) + p(3) + p(4,3) + p(5,3) + (p(6,3)-1) = 1 + 1 + 2 + 3 + 4 + 5 + (7-1) = 22$$

Similar to previous cases we count potential JDT or rank matrices for s = 6 and $k \ge 5$ recursively using previous results that count the rank matrices for s = 3, 4, 5 and $k \ge 4$.

Theorem 3.9. Let $T = (1, 3, 6^k, 3, 1)$, $k \ge 5$. There are 65 potential Jordan degree types for (A, ℓ) where $A \in Gor(T)$ and $\ell \in A_1$.

Proof. The possible sequences for $\overrightarrow{\Delta}$ are given in Table 6. We first note that we count the rank matrices for AG algebras A and linear forms $\ell \neq 0$ so we have $r_1 \neq 0$ and $\Delta r_0 < r_0 = 6$. Therefore, $r_{0,j-1} = 1$. We claim that the second diagonal, $(1, r_{1,j-2}, r_1, \ldots, r_1, r_{1,j-2}, 1)$ is the Hilbert function of some codimension three AG algebra. Since otherwise the difference between the main and second diagonal is $(1, 2, h, \ldots)$ where $h = r_0 - r_{1,j-2} = 4, 5$ meaning that this difference is not an O-sequence contradicting Lemma 2.7 (ii).

It is left to study the possible values of $r_{0,0}, \ldots, r_{0,j-1}$ and $r_{1,1}, \ldots, r_{1,j-2}$. Note that if r_u is zero, then $r_{0,j-u} = r_{1,j-u-1} = 0$ (the whole diagonal containing r_u is zero).

For case #1, since the diagonals are Hilbert functions for AG algebras, then $r_{1,j-2} = \cdots = r_{1,3} = 3$, and $r_{0,4} = \cdots = r_{0,j-1} = 1$. By Lemma 2.8 (iv), $r_{1,2} = 3$ and $r_{0,3} = r_{0,2} = 1$. It is left to count the possibilities of the matrix

 $\begin{pmatrix} 1 & 1 & r_{0,2} & r_{0,1} & r_{0,0} \\ 3 & 3 & 3 & r_{1,1} & r_{0,1} \\ 6 & 6 & 6 & 3 & r_{0,2} \\ 6 & 6 & 6 & 3 & 1 \\ 6 & 6 & 6 & 3 & 1 \end{pmatrix}$

and we have the following choices for the upper right corner:

$$\begin{pmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 6 & 3 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 1 \\ 3 & 1 & 1 \\ 6 & 3 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 3 & 2 & 0 \\ 6 & 3 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 1 \\ 3 & 2 & 1 \\ 6 & 3 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 1 \\ 3 & 2 & 1 \\ 6 & 3 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 0 \\ 3 & 3 & 1 \\ 6 & 3 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 1 \\ 3 & 3 & 1 \\ 6 & 3 & 1 \end{pmatrix}$$

For cases #2,4,7,12,19 and #29, (i.e. $r_1 = 5$, $r_2 \ge 4$ the second diagonal is $(1,3,5^{k-1},3,1)$. Hence, for $k \ge 5$, the possible rank matrices follow from the case $(1,3,5^k,3,1)$.

If $r_1 = 4$, the second diagonal is $(1, 3, 4^{k-1}, 3, 1)$. Therefore, the possible rank

#	$\overrightarrow{\Delta} = (\Delta r_1, \dots, \Delta r_6)$	$\overrightarrow{r} = (r_0, r_1, \dots, r_{k-1})$	#rk matrices
1	(0,0,0,0,0,0)	$(6, 6, 6, 6, 6, 6, 6, \dots, 6)$	7
2	$\left(1,0,0,0,0,0\right)$	$(6, 5, 5, 5, 5, 5, 5, \dots, 5)$	7
3	$\left(2,0,0,0,0,0\right)$	$(6, 4, 4, 4, 4, 4, 4, \ldots, 4)$	5
4	$\left(1,1,0,0,0,0\right)$	$(6, 5, 4, 4, 4, 4, 4, \ldots, 4)$	5
5	$\left(3,0,0,0,0,0 ight)$	$(6, 3, 3, 3, 3, 3, 3, 3, \ldots, 3)$	2
6	(2, 1, 0, 0, 0, 0)	$(6, 4, 3, 3, 3, 3, 3, 3, \ldots, 3)$	6
7	(1, 1, 1, 0, 0, 0)	$(6, 5, 4, 3, 3, 3, 3, \ldots, 3)$	6
8	$\left(3,1,0,0,0,0\right)$	$(6, 3, 2, 2, 2, 2, 2, \dots, 2)$	2
9	$\left(2,2,0,0,0,0\right)$	$(6, 4, 2, 2, 2, 2, 2, \dots, 2)$	2
10	(2, 1, 1, 0, 0, 0)	$(6, 4, 3, 2, 2, 2, 2, \ldots, 2)$	4
11	(1, 1, 1, 1, 0, 0)	$(6, 5, 4, 3, 2, 2, 2, \ldots, 2)$	4
12	$\left(3,2,0,0,0,0 ight)$	$(6, 3, 1, 1, 1, 1, 1, \dots, 1)$	1
13	$\left(3,1,1,0,0,0 ight)$	$(6, 3, 2, 1, 1, 1, 1, \dots, 1)$	1
14	$\left(2,2,1,0,0,0\right)$	$(6, 4, 2, 1, 1, 1, 1, \dots, 1)$	1
15	(2, 1, 1, 1, 0, 0)	$(6, 4, 3, 2, 1, 1, 1, \dots, 1)$	2
16	(1, 1, 1, 1, 1, 0)	$(6, 5, 4, 3, 2, 1, 1, \dots, 1)$	2
17	$\left(3,2,1,0,0,0 ight)$	$(6, 3, 1, 0, 0, 0, 0, \dots, 0)$	1
18	$\left(3,1,1,1,0,0 ight)$	$(6, 3, 2, 1, 0, 0, 0, \dots, 0)$	1
19	$\left(2,2,2,0,0,0\right)$	$(6, 4, 2, 0, 0, 0, 0, \dots, 0)$	1
20	(2, 2, 1, 1, 0, 0)	$(6, 4, 2, 1, 0, 0, 0, \dots, 0)$	1
21	(2, 1, 1, 1, 1, 0)	$(6, 4, 3, 2, 1, 0, 0, \dots, 0)$	2
22	$\left(1,1,1,1,1,1\right)$	$(6, 5, 4, 3, 2, 1, 0, \dots, 0)$	2

Table 6: Case s = 6: the sequences Δ and number of rank matrices.

matrices follow from the table of $(1, 3, 4^k, 3, 1)$. These are cases #3,6,10,11, 17,18, 26,27 and 28 (see Table 4)

For cases #5,9,15,16,24 and 25, $r_1 = 3$, then the possible rank matrices follow from the case $(1, 3^{k+1}, 1)$ (Table 3).

Theorem 3.10. Let $T = (1, 3, 6^k, 3, 1)$. For k = 3 there are 52 and for k = 4 there are 61 potential Jordan degree types for pairs (A, ℓ) .

Proof. Let k = 3 and note that \overrightarrow{r} has three non-zero entries. Similar to

the proof of Theorem 3.8 we consider the rows of Table 6 having the same (r_0, r_1, r_2) .

Suppose $\overrightarrow{r} = (6,3,1)$ that is given by first three entries of \overrightarrow{r} in #12 and #17 in Table 6. So the number of rank matrices in this case is equal to the sum of the rank matrices given in #12 and #17 that is equal to two.

The first three entries of #8, 13 and 18 are equal to $\overrightarrow{r} = (6, 3, 2)$. So the number of rank matrices in this case is equal to the sum of rank matrices of the corresponding rows that is equal to 4.

Similarly, for $\overrightarrow{r} = (6, 4, 2)$ there are 5 rank matrices that is the sum of the rank matrices in #9, 14, 19, and 20.

The case $\overrightarrow{r} = (6, 4, 3)$ corresponds to #6, 10, 15, and 21. Notice that the rank matrices for k = 3 and row #6 correspond to the cases where we have $(r_{0,3}, r_{1,2}, r_{1,2}, r_{0,3}) = (1, 3, 3, 1)$ and $(r_{0,2}, r_{1,1}, r_{0,2}) = (1, 3, 1)$ and there are only two possible rank matrices in this case. Thus the total number of rank matrices for $\overrightarrow{r} = (6, 4, 3)$ is equal to 10.

Finally, $\overrightarrow{r} = (6, 5, 4)$ is given in rows #4, 7, 11, 16, and 22. Rank matrices corresponding to #4 cannot occur for k = 3, since they correspond to having $(r_{0,3}, r_{1,2}, r_{1,2}, r_{0,3}) = (1, 4, 4, 1)$ that is not possible. And similar to the previous case there are only 2 rank matrices corresponding to #6 for k = 3. Thus the numbe of possible rank matrices for $\overrightarrow{r} = (6, 4, 3)$ is equal to 10.

We conclude that for k = 3 there are 52 potential rank matrices or equivalently Jordan degree types for pairs (A, ℓ) .

Now we show there are 61 potential rank matrices when k = 4 using previous theorems for s = 3, 4, 5 and k = 3. We consider the first four entries of \overrightarrow{r} in Table 6. Thus rows #14 and 20 correspond to $(r_0, r_1, r_2, r_3) =$ (6, 4, 2, 1) and using the count for s = 4 and k = 3 we get that the number of rank matrices in this case is equal to the sum of rank matrices in in rows #6 and 10 in Table 4 that is equal to 1 + 1 = 2. Rows #10, 2 and 21 correspond to (6, 4, 3, 2) and the count it equal to the sum of rank matrices of rows #4, 7, and 11 in Table 4 which correspond to (4, 3, 2) and is equal to 4+2+2=8. Finally, the first four entries of \overrightarrow{r} of rows #7, 11, 16, and 22 is (6, 5, 4, 3). The number of rank matrices in this case is equal to the rank matrices for s = 5, k = 3 and having the $\overrightarrow{r} = (5, 4, 3)$. Thus using the last part of the proof of Theorem 3.8 we get that in this case there are 10 possible rank matrices (that is equal to the sum of number of rank matrices in #7, 11, 16, and 22 minus 4). We conclude that the number of rank matrices for s = 6 and k = 4 is 4 less than the number of rank matrices for s = 6 and $k \geq 5$ that is equal to 61.

4 Actual JDT for algebras $A \in Gor(T)$.

We first classify three kinds of parts for JDT occurring for algebras in Gor(T), $T = (1, 3, s^k, 3, 1)$ as k increases (Section 4.1). We then state the second part of our main results, that the JDT occurring for almost constant T with $s \leq 5$ are completely determined, and are exhibited in the appended Tables (Section 4.2). We then illustrate the methods we used to verify the tables (Section 4.3), and set out some questions and problems (Section 4.4).

4.1 Three kinds of growth of JDT parts.

Recall that the JDT for any $A \in \text{Gor}(T)$ for any T is symmetric (Lemma 2.2). As the multiplicity k of the Sperner number increases for the almost constant T of Equation (1), there are three kinds of growth of the parts that may occur for the potential JDT for $A \in \text{Gor}(T)$.

Definition 4.1. Let j = socle degree of T. Here j = k + 3 for $s \ge 4$, or j = k + 1 for s = 3.

- (a) lengthening parts of the form $(j-a)_i$ where i and a are fixed as j increases;
- (b) repeated parts, of the form $(w \uparrow_i^{j+1-i-w})$, where w and i are fixed as j increases, and $w \leq s$;
- (c) symmetric fixed parts of the form $(w_{i,j+1-i})$ where w and i are fixed as j increases.

Example 4.2. When $T = (1, 3^k, 1)$ the Table 3 entry #4 is $\overrightarrow{\Delta} = (1, 1, 0)$ and $\mathcal{S} = ((j+1)_0, 2\uparrow_1^{j-2}, 1_{1,j-1})$ which has the lengthening part $(j+1)_0$, the repeated parts $(2\uparrow_1^{j-2})$ and the symmetric fixed parts $(1_{1,j-1})$.

Given T from Equation (1) and a rank matrix $M_{A,\ell}$ for $A \in \text{Gor}(T)$, recall from Equation 9 the sequence $\overrightarrow{\Delta} = (\Delta r_1, \ldots, \Delta r_{k-1})$, the first differences $\Delta r_i = r_{i-1} - r_i$ of the sequence $\overrightarrow{r} = (r_0, r_1, \ldots, r_{k-1})$ where $r_0 = s$. We consider a set of related JDT $\mathcal{S}(j) = \mathcal{S}_{F(j),x}$ where $F(j) \in S_j$ and $F(j-i) = x^i \circ F(j)$, of Hilbert function $H(F(u)) = (1, 3, s^{u-3}, 3, 1)$. We denote by $W_{\mathcal{S}(j)}$ the multi-set (with multiplicities)

$$W_{\mathcal{S}(j)} = \{ w \mid \exists (w \uparrow_i^{j+1-i-w}) \text{ in the JDT } \mathcal{S}(j) \}.$$
(17)

Recall that the conjugate partition P^{\vee} to P is obtained by switching rows and columns in the Ferrers diagram of P. We denote the sum of entries of $\overrightarrow{\Delta}$ by $|\Delta|$.

Proposition 4.3. Let $s \leq 6$ be fixed, let T satisfy Equation (1), and fix a sequence S(j) of related JDT for F(j) as for Equation (17) above. Consider $\overrightarrow{\Delta} = \overrightarrow{\Delta}(j)$ formed by Equation (9) from an AG algebra $A(j) = R/\operatorname{Ann}(F(j))$ for some $j \geq 6$. Assume that $k \geq s + 1$.

- (a) We have $W_{\mathcal{S}(j)} = (\overrightarrow{\Delta})^{\vee}$, the conjugate partition of $\overrightarrow{\Delta}$, independent of j.
- (b) The total number of lengthening parts in the set of JDT S(j) is $(s |\Delta|)$. Also, a lengthening part is at least j - 3.

Proof. By Corollary 2.9 $\Delta_i = 0$ for $i \geq s+1$, so we assume $i \geq s+1$. We prove (a) by induction on $(s-r_1)$, the largest part of $\overrightarrow{\Delta}$. When this is 1, so $\overrightarrow{\Delta} = (1, 1, \ldots, 1_p, 0, \ldots)$, then the $k \times k$ submatrix at the center of $M_{A,\ell}$ in Equation (10) satisfies

$$\begin{pmatrix}
s & s-1 & s-2 & s-3 & \dots & s-p+1 & s-p & s-p & \dots \\
s & s-1 & s-2 & \dots & s-p+1 & s-p & s-p\dots \\
s & s-1 & \dots & s-p+1 & s-p\dots \\
\ddots & \ddots & \ddots & & \ddots \\
s & s & \dots & s-p+1 & \\
\end{pmatrix}$$
(18)

The JDT matrix $JDT_{A,\ell}$ of Definition 2.16 will have in this $k \times k$ region

$$1 = (s - p + 1) + (s - p) - 2(s - p)$$

on the locations shown red in the above matrix of Equation (18) (after the first row). These are on the p-1 labelled diagonal of $M_{A,\ell}$ so each corresponds to a part p_i , that is, p in degree i where i is the row index in $M_{A,\ell}$: thus, $\mathcal{S}_{A,\ell}$ has corresponding parts $(p \uparrow_3^{j-p-3})$.

Now fix $t \geq 2$ and assume (a) is true for first entries of $\overrightarrow{\Delta}$ less than t, and consider a particular $\overrightarrow{\Delta}(0)$ with first entry t, that is nonzero in degrees $0, \ldots, p-1$, then zero. We can see that the last entry of each row of the $k \times k$ submatrix of $JDT_{A,\ell}$ is one: that is, on its p-1 labelled diagonal (analogous to that shown red in the $k \times k$ matrix $M_{A,\ell}$ of Equation (18)), so this corresponds to a repeating part $(p \uparrow_i^{j+1-p-i})$ in \mathcal{S} . Removing that layer, using the induction hypothesis, we see that the repeating parts of $S_{A,\ell}$ correspond to the conjugate partition Δ^{\vee} to Δ .

We now show (b). The remaining entries of the JDT matrix $JDT_{A,\ell}$ outside of the repeating parts, are symmetrically located. Since the maximum height of T due to repeated parts is just the length $|\Delta|$, by the construction of the matrix M and the related JDT matrix, the only way to attain the height s is for there to be $(s - |\Delta|)$ lengthening parts.

The last statement of (b), that a lengthening part is at least j-3, follows from the next result, Lemma 4.5.

Example 4.4. (a). Entry #34 in Table 11 is $\mathcal{S}_{A,x} = ((j_{0,1}, 2 \uparrow_2^{j-3}, 1 \uparrow_1^{j-1}, 1_{2,j-2})$ for $A = R/I, I = \operatorname{Ann}(X^{j-1}Y + XY^{j-1} + Y^{j-2}Z^2)$. Here s = 5, the repeated parts are 2 and 1, and the two lengthening parts are $j_{0,1}$, while $\Delta = (2, 1, 0)$ whose dual is (also) (2, 1).

b. Entry #43 in Table 12 has $S_{A,x} = ((4 \uparrow_0^{j-3}, 2_{1,j-2}, 1 \uparrow_2^{j-2}))$ for $A = R/\operatorname{Ann}(X^3Y^{j-3} + XY^{j-2}Z + Z^j)$. Here the repeated parts are 4 and 1, while $\Delta(A) = (2, 1, 1, 1)$, and there are no lenghtening parts.

(c). In codimension two, the repeating parts of the associated JDT are also obtained as the conjugate partition to $\overrightarrow{\Delta} = (\Delta r_1, \Delta r_2, \Delta r_3)$. For example $\overrightarrow{\Delta} = (1, 1, 0)$ has repeating part 2, and $\overrightarrow{\Delta} = (2, 0, 0)$ has repeating parts (1, 1). See Example 3.2, and Tables 1 and 2 above.

Lemma 4.5. Let T be an almost constant Gorenstein sequence, with Sperner number s = 3, 4, 5, or 6. Then the possible lengthening parts for $S(A, \ell), A \in$ Gor(T) satisfy the following restrictions,

- in degree 0: $(j a)_0$ with a = -1, 0, 1,
- in degree 1: $(j a)_1$ with a = 0, 1, 2,
- in degree 2: $(j a)_2$ with a = 1, 2, 3.

No other lengthening parts can occur.

Proof. Since the sequence Δ is non-increasing, non-negative and its elements add up to at most s, then $\Delta_i = 0$ for $i \geq s$, and therefore $r_s = r_{s+1} = \cdots = r_{k-1}$. Furthermore, if $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$ is an adjacent submatrix of the rank matrix such that $\gamma = \delta$, then $\alpha = \beta$. The rank matrix can then be written as follows:

$$\begin{pmatrix} 1 & r_{0,j-1} & \dots & r_{0,j-s} & r_{0,j-s-1} & r_{0,j-s-2} & r_{0,j-s-2} & \dots & r_{0,j-s-2} & r_{0,1} & r_{0,0} \\ 3 & r_{1,j-2} & \dots & r_{1,j-s-1} & r_{1,j-s-2} & r_{1,j-s-2} & r_{1,1} & r_{0,1} \\ s & r_1 & \dots & r_s & r_s & \dots & r_s & r_{1,j-s-2} & r_{0,j-s-2} \\ s & r_1 & \dots & r_s & \ddots & \vdots & \vdots & \vdots \\ & \ddots & \ddots & & \ddots & & r_s & r_{1,j-s-2} & r_{0,j-s-2} \\ & & \ddots & \ddots & & \ddots & & r_s & r_{1,j-s-2} & r_{0,j-s-2} \\ & & & \ddots & \ddots & & r_s & r_{1,j-s-2} & r_{0,j-s-2} \\ & & & \ddots & \ddots & & r_s & r_{1,j-s-2} & r_{0,j-s-2} \\ & & & & \ddots & \ddots & & r_s & r_{1,j-s-2} & r_{0,j-s-2} \\ & & & & \ddots & \ddots & & r_s & r_{1,j-s-1} & r_{0,j-s-1} \\ & & & & s & r_1 & \vdots & r_{0,j-s-1} \\ & & & & s & r_{1,j-2} & \vdots \\ & & & & & s & r_{1,j-2} & \vdots \\ & & & & & & s & r_{1,j-2} & \vdots \\ & & & & & & & 1 \end{pmatrix}$$

Furthermore, since $r_{0,a} = 0$ if one of the elements on the same diagonal is 0 and 1 otherwise, then $r_{0,j-s} = r_{0,j-s-1} = r_{0,j-s-2}$ Define $J_i = r_i + r_{i+2} - 2r_{i+1}$. The Jordan degree type matrix is therefore

Note that the lengthening parts come from the top right 3×3 -submatrix of the Jordan degree type matrix. This completes the proof of Lemma 4.5. \Box

4.2 Classification of JDT for $s \leq 5$.

We here report on the classification of all occurring JDT for graded Artinian algebras of almost constant Hilbert function T of codimension three with Sperner number 3, 4, 5 - that are given in the Tables Section A. The results of this section assume char $\mathbf{k} = 0$. For char \mathbf{k} finite see Remark 4.9.

We first determine the occurring JDT for T of Sperner number 3.

Theorem 4.6. A. Let $T = (1, 3^k, 1)$. Suppose first that $k \ge 3$. Then #1-8 in Table 7 is a complete table of the 8 occurring JDT for algebras $A \in Gor(T)$.

B. When k = 2, # 2 is the same as # 3 and #6 is the same as #7 but there is an extra # 9. When k = 1 only # 1 and # 8 are distinct.

We next consider Sperner number 4.

Theorem 4.7. ³ Let $T = (1, 3, 4^k, 3, 1)$. When $k \ge 3$ there are exactly 26 JDT that actually occur, as presented in Table 8. When k = 2, there are exactly 22 JDT, as then #6=#7, #10=#11, #16=#18, and #23=#24 in Table 8. When k = 1 there are exactly 12 JDT arising from Table 8 #1, #3=#5, 7, 8, 12, 13, 15, 19, 20, 24, 25, and #26.

We next consider Sperner number 5.

Theorem 4.8. Let $T = (1,3,5^k,3,1)$. When $k \ge 4$ (so $j \ge 7$) there are exactly 47 JDT, all distinct, that occur for T, listed inTables 11,12, where we give the JDT, a suitable dual generator (in all but one case) and the ideal, in terms of j. When k = 3 there are identities #9=#11, #10=#12, $\#29=\#30, \text{ and } \#39=\#42, \text{ and there are exactly 43 distinct JDT. When$ <math>k = 1 there are 12 distinct, occurring JDT - (Table 9). When k = 2 there are 35 distinct JDT: see Table 10, where we give in the last column the corresponding Table 11/12 entry).

Proof of Theorem 4.6, Theorem 4.7 and Theorem 4.8. The restriction on the number of potential JDT are shown in each case of almost constant T for $3 \le s \le 5$ in Section 3, Theorem 3.3 for (s = 3), Theorem 3.5 for (s = 4), Theorems 3.6 and 3.8 for s = 5 and Theorem 3.9 for s = 6. The tables - and

³In Table 8 we indicate the Jordan degree types that occur for Complete intersection algebras by CI and the ones that occur for connected sum algebras by cs, for more details on connected sum algebras see [IMM].

especially the dual generators we found were constructed using several techniques, and verified case by case. We give examples of the techniques we used to verify table entries in the next Section. We of course used MACAULAY2 in verifying some dual generators for specific j, as a check, but our techniques apply - as they must - to arbitrary socle degree j.

For Sperner number 6, $T = (1, 3, 6^k, 3, 1)$ we did not construct tables of the actually occurring JDT so have not verified that each potential JDT from Theorem 3.9 actually occurs.

Remark 4.9 (When the characteristic char k is finite). Many results in this area require a restriction to fields of characteristic zero, or of characteristic p higher than the socle degree j of A. For example, in characteristic 2 the algebra $A = R/I, I = (x^2, y^2, z^2) = Ann(F), F = XYZ, H(A) = (1, 3, 3, 1)$ is not strong Lefschetz: any linear form ℓ satisfies $\ell^2 = 0$, and for a generic ℓ , we have $\mathcal{S}_{A,\ell} = (2\uparrow_0^2, 2_1)$, rather than the strong Lefschetz JDT $H^{\vee} = (4_0, (2_1)^2)$. Example 2.15 above with $F = X^3 + Y^3 + Z^3$ shows that $\ell = x + y + z$, although a strong Lefschetz element when the characteristic of k is not 3, is not a strong Lefschetz element over characteristic 3. Thus, ostensibly, the set $\mathcal{J}(T)$ of JDT for a given Gorenstein sequence T might depend upon the characteristic of k: that is, for certain fields of small characteristic, some of the entries in our Tables for s = 3, 4, 5 might not appear. However we do not have any such example. Our Macaulay dual generators are notable in that, with the exception of one case, their coefficients on monomials are one; and the linear form $\ell = x$ in all but the two cases s = 3, Table 7, entries # 1, 2, where $F = X^{j} + Y^{j} + Z^{j}$ and $\ell = x + y + z$ or $\ell = x + y$. Both these JDT occur for suitable Artinian algebras $A(j) = R/\operatorname{Ann}(F(j))$ and the linear form x.

4.3 Examples and Methods.

We here give examples illustrating the methods we used to determine the JDT in the tables. The following example illustrates how we obtain the JDT $\mathcal{J}(T)$ for a given T and sequence $\overrightarrow{\Delta}$.

Example 4.10. Consider the Hilbert function $H = (1, 3, 4^k, 3, 1)$ and the sequence $(r_0, r_1, \ldots, r_k) = (4, 3, 3, \ldots, 3)$, so $\overrightarrow{\Delta} = (1, 0, \ldots, 0)$. The second diagonal of the rank matrix can be either the Hilbert function of a codimension 2 algebra $(1, 2, 3^{k-1}, 2, 1)$, or that of a codimension 3 algebra $(1, 3^{k+1}, 1)$. These two cases are illustrated by the below examples. The matrices $M_{A,\ell}$

and $J_{A,\ell}$ below are an example of a rank matrix and its corresponding JDT matrix for the first case. The JDT is $((j + 1)_0, (j - 1)_1, (j - 3)_2, 1 \uparrow_1^{j-1})$. The matrices $M'_{A,\ell}$ and $J'_{A,\ell}$ are an example for the second case. The Jordan degree type in this case is $((j + 1)_0, ((j - 1)_1)^2, 1 \uparrow_2^{j-2})$.

$$M_{A,\ell} = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & \dots & 1 & 1 & 1 \\ 3 & 2 & 2 & 2 & \dots & 2 & 2 & 1 \\ 4 & 3 & 3 & \dots & 3 & 2 & 1 \\ 4 & 3 & \dots & 3 & 2 & 1 \\ & & \ddots & \ddots & \vdots & \vdots & \vdots \\ & & & \ddots & \ddots & \vdots & \vdots & \vdots \\ & & & & 3 & 2 & 1 \\ 4 & 2 & 1 \\ & & & & & 1 \end{pmatrix}, J_{A,\ell} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & \dots & 0 & 0 & 1 \\ 1 & 0 & 0 & \dots & 1 & 0 & 0 \\ & & & 1 & 0 & \dots & 0 & 0 & 0 \\ & & & & \ddots & \ddots & \vdots & \vdots & \vdots \\ & & & & \ddots & 0 & 0 & 0 \\ & & & & & & 1 & 0 & 0 \\ & & & & & & & 1 & 0 \\ & & & & & & & 1 & 0 \\ & & & & & & & 0 \end{pmatrix},$$

In both examples, the lengthening parts are given by the nonzero elements of the 3×3 top right corner submatrix of the Jordan degree type matrix. For instance, the entries 1 on the first row and 2 on the second row of the $\begin{pmatrix} 0 & 0 & 1 \end{pmatrix}$

submatrix $\begin{pmatrix} 0 & 0 & 1 \\ 0 & 2 & 0 \\ 0 & 0 & 0 \end{pmatrix}$ of $J'_{A,\ell}$ give $(j+1)_0$ and twice $(j-1)_1$. The repeated

entries 1 on the diagonal of each JDT give the repeated portion $1 \uparrow_2^{j-2}$ for the first example, and $1 \uparrow_1^{j-1}$ for the second.

For all of the table entries for s = 3 and s = 4, and all but one for s = 5we are able to provide a dual generator for an algebra having the required JDT. An exception is # 3 for s = 5, where the dual generator can be found for small cases (we calculated up to j = 20 using MACAULAY2, but were unable to find a pattern). In this case we instead simply show (next) that for arbitrary socle degree j, the algebra A = R/I(j) is Gorenstein of the expected Hilbert function.

Example 4.11. For each of the ideals in the third column of the table 11 we can compute a Gröbner basis (standard basis) using homogeneous degree rvlex order with z > y > x.

We consider the case #3 for $T = (1, 3, 5^k, 3, 1)$ of socle degree j = k + 3, where the ideal I(j) (we will call I for simplicity) is

$$I = (xy - yz, x^{3} - z^{3}, y^{3} - x^{2}z - y^{2}z - xz^{2} + yz^{2} - z^{3}, x^{j-7}yz^{5}, x^{j-5}z^{4}).$$
(19)

It is straightforward to verify that the Hilbert function H(R/I) = T. Since $yz \equiv xy$ and $z^3 \equiv x^3 \mod I$, one can check that

$$I = \left(z^3 - x^3, zy - yx, y^3 - z^2x - y^2x - zx^2 + yx^2 - x^3, zx^{j-2}, yx^{j-2}\right)$$

and a standard basis for I is given by

$$(z^3 - x^3, zy - yx, y^3 - z^2x - y^2x - zx^2 + yx^2 - x^3, zx^{j-2}, yx^{j-2}, x^{j+1}).$$

The strings describing the JDT of the algebra R/I are the following:

- a length (j+1) string starting at degree 0: $(1, x, \ldots, x^i, \ldots, x^j)$
- two length (j-2) strings starting both at degree one:

$$(z, zx, ..., zx^{i}, ..., zx^{j-3})$$
 and $(y, yx, ..., yx^{i}, ..., yx^{j-3})$

• two length (j-2) strings, both starting at degree two:

$$(z^2, z^2x, \dots, z^2x^i, \dots, z^2x^{j-3})$$
 and $(y^2, y^2x, \dots, y^2x^i, \dots, y^2x^{j-3})$.

These strings give the JDT $((j+1)_0, ((j-2)_{1,2})^2)$ for the algebra A = R/I.

It remains to show that A is Gorenstein. We check that the socle of the graded algebra A has dimension one.

From the five strings that describe the JDT of A, one can see that for $2 \leq i \leq j-2$, a basis of A_i is given by $(z^2x^{i-2}, y^2x^{i-2}, zx^{i-1}, yx^{i-1}, x^i)$, $A_{j-1} = \langle z^2x^{j-3}, y^2x^{j-3}, x^{j-1} \rangle$, $A_j = \langle x^j \rangle$.

It is obvious that for $0 \le i \le j-3$, the multiplication by $x, m_x : A_i \to A_{i+1}$ is injective. This means that any non zero homogeneous element f of the socle of A has degree at least j-2.

We have $A_{j-2} = \langle z^2 x^{j-4}, y^2 x^{j-4}, zx^{j-3}, yx^{j-3}, x^{j-2} \rangle$. Let $(a_1, \dots, a_5) \in k^5$, $f = a_1 z^2 x^{j-4} + a_2 y^2 x^{j-4} + a_3 z x^{j-3} + a_4 y x^{j-3} + a_5 x^{j-2} \in A_{j-2}$ and suppose xf = yf = zf = 0.

- i. Since $zx^{j-2} \in I$ and $yx^{j-2} \in I$, xf = 0 gives $a_1z^2x^{j-3} + a_2y^2x^{j-3} + a_5x^{j-2} = 0$. Since $(z^2x^{j-3}, y^2x^{j-3}, x^{j-1})$ is a basis of A_{j-1} , we get $a_1 = a_2 = a_5 = 0$, so $f = a_3zx^{j-3} + a_4yx^{j-3}$.
- ii. Let $f = a_3 z x^{j-3} + a_4 y x^{j-3}$ and suppose $yf = a_3 z y x^{j-3} + a_4 y^2 x^{j-3} = 0$. We know that $zy - yx \in I$, so $yf = a_3 y x^{j-2} + a_4 y^2 x^{j-3} = a_4 y^2 x^{j-3}$. This means that yf = 0 gives $a_4 y^2 x^{j-3} = 0$, so $a_4 = 0$ and $f = a_3 z x^{j-3}$.

iii. For $f = a_3 z x^{j-3}$ one can see that $z f = a_3 z^2 x^{j-3} = 0$ if and only if $a_3 = 0$.

Now, looking at degree j-1, let $(a_1, a_2, a_3) \in k^3$, $f = a_1 z^2 x^{j-3} + a_2 y^2 x^{j-3} + a_3 x^{j-1}$ such that xf = yf = zf = 0. One can easily check that $xf = a_3 x^j$, $yf = a_2 x^j$ and $zf = a_1 x^j$ so xf = yf = zf = 0 if and only if $a_1 = a_2 = a_3 = 0$.

In conclusion, the socle of A is one dimensional, generated by x^{j} .

Example 4.12. Consider the Hilbert function $T = (1,3,5^k,3,1)$ and #5 in Table 11. We want to show that $F = X^j + X^{j-1}Y + X^{j-3}Z^3 + Y^j$ and $A = R/\operatorname{Ann}(F)$ has associated JDT $((j+1)_0, (j-1)_1, (j-2)_{1,2}, 1\uparrow_2^{j-2});$ Consider $F_x = x \circ F = X^{j-1} + X^{j-2}Y + X^{j-4}Z^3$, which corresponds to #3 in Table 8 for $T_1 = (1,3,4^{k-1},3,1)$ (or $F_1 = X^j + X^{j-1}Y + X^{j-3}Z^3$ for $T = (1,3,4^k,3,1)$). If we have a proof that the dual generator F_1 has JDT $((j+1)_0, (j-1)_1, (j-2)_{1,2})$ and we know that F gives the Hilbert function $H(R/\operatorname{Ann}(F)) = T$, then we are done, since the Jordan degree type matrix for $B = R/\operatorname{Ann}(F_1)$ is

These imply that the corresponding matrices for A are (first find $M_{A,x}$ then $J_{A,x}$) are, adding respectively a first diagonal $(1, 3, 5^k, 3, 1)$ to $M_{A,x}$ and a first diagonal $(0, 0, 1, 1, \ldots, 1, 0, 0)$ to $J_{A,x}$

The JDT matrix $J_{A,x}$ imples that the JDT for A is $((j+1)_0, (j-1)_1, (j-2)_{1,2}, 1\uparrow_2^{j-2})$.

4.4 Questions and problems.

First, can we extend these results to more complex Gorenstein sequences of codimension three, where the Sperner number s is repeated k times?

Question 4.13. Consider $T(T_1, s, k) = (1, 3, T_1, s^k, T_1^r, 3, 1)$ where T_1 is an SI sequence, and T_1^r is its reversal, As k increases, with T_1, s held constant, we expect

i. a stable constant limit number of potential Jordan degree types for $T(T_1, s, k)$ as T_1 and s are fixed but k increases;

ii. there to be a list of potential Jordan degree types for these $T(T_1, s, k)$ as T_1 , s are fixed and k increases, with structure analogous to those we found for almost constant T, that is, with the three kinds of parts of Definition 4.1 satisfying Proposition 4.3.

iii. that the potential JDT found in (ii) all actually occur.

Remark. We can do similar operations on the possible rank matrices as in the T almost constant case: thus, we can bound the number of potential Jordan types, so answering (i).

We expect this question parts (ii),(iii) could be approached inductively, from simpler T_1 to more complex ones. For example, to understand even $T = (1, 3, 5, 6^k, 5, 3, 1)$ we might need to have first determined the answers for simpler Hillbert functions as $T = (1, 3, 4, 5^k, 4, 3, 1)$.

Question 4.14 (Limits of Gorenstein algebras). Let T be a codimension three Gorenstein sequence. There is an inclusion $Gor(T) \subset G_T$ of the family of graded Gorenstein algebras of Hilbert function T in the family G_T of all graded algebra quotients of R = k[x, y, z] having Hilbert function T.

a. For which T is the Zariski closure $\overline{\text{Gor}(T)} = G_T$? It is known that for $T = (1, 3^k, 1)$ the closure $\overline{\text{Gor}(T)} = G_T$, but for T = (1, 3, 5, 3, 1), (1, 3, 6, 3, 1) or (1, 3, 6, 6, 3, 1) the closure of Gor(T) is proper in G_T [AEIY].

b. What are limits of JDT strata of Gor(T) in G_T ? Note the JDT for G_T need not be symmetric, even for algebras A in the closure $\overline{Gor(T)}$ [AEIY, Example 4.16].

Question 4.15. Do the actual tables - the set $\mathcal{J}(T)$ of JDT for T - depend on the characteristic of k? See Remark 4.9.

Acknowledgment. The work was begun at the CIRM Conference "Lefschetz Properties in Algebra, Geometry, and Combinatorics, II" in 2019 at Luminy, France. We appreciate contributions to our discussions by Alexandra Seceleanu, and Leila Khatami, who were members of our codimension three working group at Lumini, and participated further. We thank Pedro Macias Marques for comments and questions. The second author was supported by the grant VR2021-00472. Some part of the work was done when the second author was an NSF postdoc at the program in Commutative Algebra, during Spring 2024 semester at Simons Laufer Mathematical Sciences Institute (formerly MSRI).

A Tables of JDT in codimension 3 for almost constant T, when $s \le 5$.

#	S	F	$I = \operatorname{Ann}(F)$	ℓ
1	$((j+1)_0, ((j-1)_1)^2)$	$X^j + Y^j + Z^j$	$xy, xz, yz, x^j - y^j, x^j - z^j$	x + y + z
2	$((j+1)_0, (j-1)_1, 1\uparrow_1^{j-1})$	$X^j + Y^j + Z^j$	11	x + y
3	$((j+1)_0, 2\uparrow_1^{j-2}, 1_1, 1_{j-1})$	$X^j + XY^{j-1} + ZY^{j-1}$	$xz, xy - yz, z^2, x^j - xy^{j-1}, x^j - zy^{j-1}$	x
4	$((j+1)_0, (1\uparrow_1^{j-1})^2)$	$X^j + Y^j + Z^j$	$xy, xz, yz, x^j - y^j, x^j - z^j$	x
5	$(j_0, j_1, (j-1)_1)$	$X^{j-1}Z + X^{j-2}Y^2$	$xz-y^2, yz, z^2, x^{j-1}y, x^j$	x
6	$(j_0, j_1, 1\uparrow_1^{j-1})$	$X^{j-1}Y + Z^j$	$xz, yz, y^2, x^j, x^{j-1}y - z^j$	x
7	$(3\uparrow_0^{j-2},1_1,1_{j-1})$	$X^2 Y^{j-2} + Y^{j-1} Z$	$x^2 - yz, xz, z^2, xy^{j-1}, y^j$	x
8	$(2\uparrow_0^{j-1},1\uparrow_1^{j-1})$	$XZ^{j-1} + Y^2Z^{j-2}$	$x^2, xy, y^2 - xz, yz^{j-1}, z^j$	x
*9	$(2\uparrow_0^2, 2_1)$ if $k=2$	$X(Z^2 + Y^2)$	$x^2, yz, y^2 - z^2, x(y^2 - z^2)$	x
	:	#9 only occurs for k = 2		

Table 7: Complete 8 JDT for $T = (1, 3^k, 1), k \ge 3, j = k + 1$ (Theorem 4.6).

#	S	F	$I = \operatorname{Ann}(F)$
1	$((j+1)_0, ((j-1)_1)^2, (j-3)_2)$	$X^{j} + X^{j-2}(Y^{2} + Z^{2}) CI$	$yz, y^2 - z^2, x^{j-1} - x^{j-3}y^2$
2	$((j+1)_0, ((j-1)_1)^2, 1\uparrow_2^{j-2})$	$X^{j} + X^{j-1}Y + X^{j-2}Z^{2} + Z^{j}$	$y^2, yz, x^2y - xz^2, x^{j-2}z - z^{j-1}, x^j - z^j$
3	$((j+1)_0, (j-1)_1, (j-2)\uparrow_1^2)$	$X^{j} + X^{j-1}Y + X^{j-3}Z^{3}$	$yz, y^2, x^2y - z^3, x^{j-2}z, x^j - x^{j-3}z^3$
4	$((j+1)_0, (j-1)_1, (j-3)_2, 1\uparrow_1^{j-1})$	$X^j + X^{j-2}Y^2 + Z^j cs$	$yz, xz, y^3, x^{j-1} - x^{j-3}y^2, x^{j-2}y^2 - z^j$
5	$((j+1)_0, (j-1)_1, 2\uparrow_1^{j-2})$	$X^j + X^{j-1}Y + XZ^{j-1}cs$	$yz, y^2, x^2z, x^{j-2}y - z^{j-1}, x^j - xz^{j-1}$
6	$((j+1)_0, (j-1)_1, 2\uparrow_2^{j-3}, 1_{1,2,j-2,j-1})$	$X^{j} + X^{j-1}Y + XY^{j-1} + Y^{j-1}Z$	$xz, z^2, xy^2 - y^2z, x^{j-1} - x^{j-2}y - y^{j-2}z, y^j$
7	$((j+1)_0, (j-1)_1, 1_1, (1\uparrow_2^{j-2})^2, 1_{j-1})$	$X^j + X^{j-1}Y + Y^j + Z^j cs$	$yz, xz, xy^2, x^{j-1} - x^{j-2}y - y^{j-1}, y^j - z^j$
8	$((j+1)_0,(j-2)\uparrow_1^2,1\uparrow_1^{j-1})$	$\sum_{i=0}^{\lfloor j/3 \rfloor} X^{j-3i} Y^{3i} + Z^j cs$	$yz, xz, x^3 - y^3, x^{j-2}y, x^j - z^j$
9	$((j+1)_0, 3\uparrow_1^{j-3}, 2_{1,j-2})$	$X^{j} + X^{2}Y^{j-2} + XY^{j-2}Z + Y^{j-2}Z^{2}$	$xz - z^2, xy - yz, z^3, y^{j-1}, x^j - y^{j-2}z^2$
10	$((j+1)_0, 3\uparrow_1^{j-3}, 1_{1,2,j-2,j-1})$	$X^{j} + X^{2}Y^{j-2} + XY^{j-1} + Y^{j-1}Z$	$xz, z^2, x^2y - y^2z, x^{j-1} - xy^{j-2} + y^{j-2}z, y^j$
11	$((j+1)_0, 2\uparrow_1^{j-2}, 1\uparrow_1^{j-1})$	$X^j + XY^{j-1} + Z^j cs$	$yz, xz, x^2y, x^{j-1} - y^{j-1}, xy^{j-1} - z^j$
12	$((j+1)_0, (1\uparrow_1^{j-1})^2, 1\uparrow_2^{j-2})$	$X^j + Z^j + ZY^{j-1} cs$	$xz, yz, yz^2, y^{j-1} - z^{j-1}, x^j - z^j$
13	$(j\uparrow_0^1, (j-1)_1, (j-3)_2)$	$X^{j-1}Y + X^{j-2}(Y^2 + Z^2) CI$	$yz, y^2 - z^2, x^{j-1} - x^{j-2}y + x^{j-3}z^2$
14	$(j\uparrow^1_0,(j-1)_1,1\uparrow^{j-2}_2)$	$X^{j-1}Y + X^{j-2}Z^2 + Z^j$	$yz, y^2, x^2y - xz^2, x^{j-2}z - z^{j-1}, x^j$
15	$(j\uparrow_0^1, (j-2)_{1,2})$	$X^{j-1}Y + X^{j-3}Z^3$	$yz, y^2, x^2y - z^3, z^4, x^{j-2}z, x^j$
16	$(j\uparrow_0^1, (j-3)_2, 1\uparrow_1^{j-1})$	$\sum_{i=0}^{\lfloor (j-1)/3 \rfloor} X^{j-1-3i} Y^{3i+1} + Z^j cs$	$yz, xz, x^3 - y^3, x^{j-3}y^2, x^{j-1}y - z^j$
17	$(j\uparrow_0^1,2\uparrow_1^{j-2})$	$X^{j-1}Y + XZ^{j-1}$	$yz, y^2, x^2z, x^{j-2}y - z^{j-2}, x^j$
18	$(j\uparrow_0^1, 2\uparrow_2^{j-3}, 1_{1,2,j-2,j-1})$	$X^{j-1}Y + XY^{j-1} + Y^{j-1}Z$	$xz, z^2, xy^2 - y^2z, x^{j-1} - xy^{j-2}, y^j$
19	$(j\uparrow_0^1, 1_1, (1\uparrow_2^{j-2})^2, 1_{j-1})$	$X^{j-1}Y + Y^j + Z^j cs$	$yz, xz, xy^2, x^{j-1} - y^{j-1}, y^j - z^j$
20	$((j-1)_0, ((j-1)_1)^2, (j-1)_2)$	$X^{j-2}(Y^2+Z^2) CI$	$yz, y^2 - z^2, x^{j-1}$
21	$((j-1)\uparrow_0^2,1\uparrow_1^{j-1})$	$X^{j-2}Y^2 + Z^j cs$	$yz, xz, y^3, x^{j-1}, x^{j-2}y^2 - z^j$
22	$(4\uparrow_0^{j-3}, 2_1, 2_{j-2})$	$X^3 Y^{j-3} + X Y^{j-2} Z C I$	$x^2 - yz, z^2, y^{j-1}$
23	$(4\uparrow_0^{j-3},1_{1,2,j-2,j-1})$	$X^{3}Y^{j-3} + Y^{j} + Y^{j-1}Z$	$xz, z^2, x^3 - y^2z, xy^{j-2}, y^j - y^{j-1}z$
24	$(3\uparrow_0^{j-2},1\uparrow_1^{j-1})$	$\overline{X^2Y^{j-2} + Z^j cs}$	$\overline{yz, xz, x^3, y^{j-1}, x^2y^{j-2} - z^j}$
25	$(2\uparrow_0^{j-1},2\uparrow_1^{j-2})$	$X(Y^{j-1} + Z^{j-1}) CI$	$yz, x^2, y^{j-1} - z^{j-1}$
26	$(2\uparrow_0^{j-1}, 1_1, (1\uparrow_2^{j-2})^2, 1_{j-1})$	$XY^{j-1} + Y^{j-3}Z^3$	$xz, x^2, xy^2 - z^3, y^{j-2}z, y^j$

Table 8: Complete 26 JDT for $T = (1, 3, 4^k, 3, 1), k \ge 3$ with the corresponding Macaulay dual generator (Theorem 4.7).

	S	F	$I = \operatorname{Ann}(F)$
1	$(5_0, (3_1)^2, (1_2)^2)$	$X^4 + X^3Z + 2X^2Y^2 + XZ^3 + Y^4$	$yz, x^3 - xz^2 - z^3, x^2y - 2y^3, x^2z - z^3, xy^2 - 2z^3$
2	$(5_0, 3_1, 2\uparrow_1^2, 1_2)$	$X^4 + X^2 Z^2 + X Y^3$	$yz, x^2y, x^3 - xz^2, y^3 - xz^2, z^3$
3	$(5_0, 3_1, 1\uparrow_1^3, (1_2)^2)$	$X^4 + X^2 Z^2 + Y Z^3 + Y^4$	$xy, x^2z - yz^2, x^3 - xz^2, y^3 - z^3$
4	$(5_0, (2\uparrow_1^2)^2)$	$X^4 + XYZ^2$	$y^2, x^3 - yz^2, x^2y, x^2z, z^3$
5	$(5_0, 2\uparrow_1^2, 1\uparrow_1^3, 1_2)$	$X^4 + XY^3 + Y^2Z^2 + Z^4$	$xz, x^2y, x^3 - y^3, xy^2 - yz^2, y^2z - z^3$
6	$(4\uparrow_0^1, 3_1, (1_2)^2)$	$X^3Y + X^2Z^2 + Y^3Z$	$xy - z^2, x^3 - y^2 z, y^3 - x^2 z, y z^2, z^3$
7	$(4\uparrow_0^1, 2_{1,2}, 1_2)$	$X^3Y + XY^2Z + Z^4$	$x^2 - yz, xy^2 - z^3, xz^2, yz^2, y^3$
8	$(4\uparrow_0^1,1\uparrow_1^3,(1_2)^2)$	$X^3Y + Y^2Z^2$	$xz, x^3 - yz^2, xy^2, y^3, z^3$
9	$(3\uparrow_0^2,3_1,1_2)$	$X^{2}(Y^{2} + Z^{2}) + Y^{4}$	$yz, x^3, z^3, xy^2 - xz^2, x^2y - y^3$
10	$(3\uparrow_0^2,2\uparrow_1^2)$	$X^2Y^2 + XZ^3$	$yz, x^2z, x^3, y^3, xy^2 - z^3$
11	$(3\uparrow_0^2, 1\uparrow_1^3, 1_2)$	$X^2Y^2 + YZ^3$	$xz, y^2z, x^3, y^3, x^2y - z^3$
12	$(2\uparrow_0^3, 2\uparrow_1^2, 1_2)$	$X(Y^3 + Z^3) + Y^2 Z^2$	$x^2, y^2z - xz^2, xyz, y^3 - z^3, xy^2 - yz^2$

Table 9: The 12 JDT for T = (1, 3, 5, 3, 1) with the Macaulay dual generator, and ideal (Theorem 4.8).

	S	F	Table 11
1	$(6_0, (4_1)^2, (2_2)^2)$	$X^5 + X^3 Y Z + X Z^4$	1
2	$(6_0, (4_1)^2, 2_2, 1_{2,3})$	$X^5 + X^3 Y Z + Z^5$	4
3	$(6_0, (4_1)^2, (1_{2,3})^2)$	$X^5 + X^4Y + X^3Z^2 + Y^5 + Z^5$	3
4	$(6_0, 4_1, 3_{1,2}, 2_2)$	$X^5 + X^3 Y^2 + X^2 Z^3$	2
5	$(6_0, 4_1, 3_{1,2}, 1_{2,3})$	$X^5 + X^4Y + X^2Z^3 + Y^5$	5,7
6	$(6_0, 4_1, 2_2, 1 \uparrow_1^4, 1_{2,3})$	$X^5 + X^3Y^2 + Y^2Z^3$	17
7	$(6_0, 4_1, 2\uparrow_1^3, 1_{2,3})$	$X^5 + X^4Y + XZ^4 + Y^2Z^3$	14
8	$(6_0, 4_1, 2\uparrow_1^3, 2_2)$	$X^5 + X^3Y^2 + XZ^4$	9
9	$(6_0, 4_1, 1_1, (1\uparrow_2^3)^3, 1_4)$	$X^5 + X^4Y + Y^2Z^3$	21
10	$(6_0, 3_{1,2}, 2_{1,3}, 1_{2,3})$	$X^{5} + X^{2}Y^{3} + XY^{3}Z$	15
11	$(6_0, 3_{1,2}, 2\uparrow_1^3)$	$X^{5} + X^{2}Y^{3} + XZ^{4}$	10
12	$(6_0, (3_{1,2})^2)$	$X^{5} + X^{2}YZ^{2}$	3
13	$(6_0, 3\uparrow_1^2, 1\uparrow_1^4, 1\uparrow_2^3)$	$X^{5} + X^{2}Y^{3} + YZ^{4}$	18
14	$(6_0, (2\uparrow_1^3)^2)$	$\frac{X^3 + XY^4 + XZ^4}{X^5 + XY^4 + XZ^4}$	16
15	$(6_0, 2\uparrow_1^3, 1\uparrow_1^4, 1_{2,3})$	$X^3 + XY^4 + YZ^4$	22
16	$(5_{0,1}, 4_1, (2_2)^2)$	$X^4Y + X^3Z^2 + XYZ^3$	23
17	$(5_{0,1}, 4_1, 2_2, 1_{2,3})$	$X^4Y + X^3Z^2 + XZ^4 + Y^5$	25
18	$(5_{0,1}, 4_1, (1_{2,3})^2)$	$X^4Y + X^3Z^2 + YZ^4$	31
19	$(5_{0,1}, 3_{1,2}, 2_2)$	$\frac{X^4Y + X^2Z^3 + XY^4}{4}$	24
20	$(5_{0,1},3\uparrow_1^2,1\uparrow_2^3)$	$X^4Y + X^2Z^3 + YZ^4$	28
21	$(5_{0,1}, 2\uparrow_1^3, 2_2)$	$\frac{X^4Y + XYZ^3}{2747}$	29
22	$(5_{0,1}, 2_{1,2,3}, 1_{2,3})$	$X^{4}Y + XZ^{4} + Y^{2}Z^{3}$	32
23	$(5_{0,1}, 2_2, 1 \uparrow_1^4, 1_{2,3})$	$\frac{X^4Y + XY^4 + Y^2Z^3}{X^4Y}$	33,34
24	$(5_{0,1}, 1_1, (1\uparrow_2^3)^3, 1_4)$	$X^4Y + Y^2Z^3$	35
25	$(4\uparrow_0^2,3\uparrow_1^2)$	$X^3Y^2 + X^2Z^3$	37
26	$(4\uparrow_0^2, 4_1, 2_2)$	$X^{3}YZ + XZ^{4}$	36
27	$(4\uparrow_0^2,4_1,1\uparrow_2^3)$	$X^{3}Y^{2} + X^{3}Z^{2} + Z^{5}$	38
28	$(4\uparrow_0^2, 2\uparrow_1^3)$	$X^{3}Y^{2} + XZ^{4}$	39
29	$(4\uparrow_0^2, 2_{1,3}, 1_{2,3})$	$X^{3}Y^{2} + XY^{3}Z + Z^{5}$	43
30	$(4\uparrow_0^2,1\uparrow_1^4,1\uparrow_2^3)$	$X^3Y^2 + YZ^4$	40,44
31	$(3\uparrow_0^3,3_{1,2})$	$X^2Y^3 + X^2Z^3$	none
32	$(3\uparrow_0^3, 2\uparrow_1^3)$	$X^2Y^3 + XZ^4$	45
33	$(3\uparrow_0^3,1\uparrow_1^4,1_{2,3})$	$X^2Y^3 + YZ^4$	46
34	$(2\uparrow_0^4, 2\uparrow_1^3, 2_2)$	XY^2Z^2	none
35	$(2\uparrow_0^4, 2\uparrow_1^3, 1_{2,3})$	$\overline{XY^4 + XZ^4 + Y^2Z^3}$	47

Table 10: The 35 JDT for T = (1, 3, 5, 5, 3, 1), with Macaulay dual generator and reference to corresponding entry of Table 11 (Theorem 4.8). 42

$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	i-5 4
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	i - 5 4
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	i-5 4
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	$x^{j} z$
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	1
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	-1
$\begin{array}{ c c c c c c c c }\hline 9 & ((j+1)_0, (j-1)_1, (j-3)_2, 2\uparrow_1^{j-1}) & X^j + X^{j-2}Y^2 + XZ^{j-1}Z^2 & yz, x^2z, y^3, x^{j-3}y^2 - z^{j-1}, x^{j-1} - z^{j-1} \\\hline 10 & ((j+1)_0, (j-2)_{1,2}, 2\uparrow_1^{j-2}) & X^j + X^{j-1}Y + X^{j-2}Y^2 + XZ^{j-1} & yz, x^2z, y^3, x^{j-3}y^2 - z^{j-1}, x^{j-1} - x^{j-2}y \\\hline 11 & ((j+1)_0, (j-1)_{1,3}\uparrow_2^{j-4}, 2_{1,2}, z_{1,2}, z_{1,3}) & X^j + X^{j-1}Y + X^2Y^{j-2} + XY^{j-2}Z + Y^{j-2}Z^2 & xz - z^2, z^3, x^{j-2} - y^2z, y^{j-1}, x^{j-1} - x^{j-2}y - y^{j-3}z \\\hline \end{array}$	$^{1} - y^{j-3}z^{2}$
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	
$\begin{bmatrix} 11 \\ ((j+1)_0, (j-1)_{1,3} \uparrow_2^{j-4}, 2_{1,2},, 2_{j} \end{bmatrix} = \begin{bmatrix} X^j + X^{j-1}Y + X^2Y^{j-2} + XY^{j-2}Z + Y^{j-2}Z^2 \\ X^j + X^{j-1}Y + X^2Y^{j-2} + XY^{j-2}Z + Y^{j-2}Z^2 \end{bmatrix} = \begin{bmatrix} x_1 - x_1^2 + y_1 - x_2^2 + y_1 + y_2 + y_2 \\ x_2 - x_2^2 + y_1^2 + y_2 + y_2 + y_2 \end{bmatrix}$	
$ \begin{array}{c} \\ (J + -)(J $	1
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	
$ \begin{array}{ c c c c c c c c } \hline 13 & (j+1)_0, ((j-1)_1)^2, (1\uparrow_2^{j-2})^2) & X^j + X^{j-1}Y + X^{j-2}Z^2 + Y^j + Z^j & yz, yx^2 - z^2x, xy^2, x^{j-2}z - z^{j-1}, x^{j-1} - y^{j-1} - x^{j-1}Z^{j-1} & y^{j-1}Z^{j-1}Z^{j-1} & y^{j-1}Z^{j-1}Z^{j-1} & y^{j-1}Z^{j-1}Z^{j-1}Z^{j-1} & y^{j-1}Z^{j-1}Z^{j-1}Z^{j-1}Z^{j-1} & y^{j-1}Z^{j-1}Z^{j-1}Z^{j-1}Z^{j-1}Z^{j-1}Z^{j-1} & y^{j-1}Z^{j$	z^2
$ \begin{array}{ c c c c c c c c } 14 & ((j+1)_0, (j-1)_1, 2\uparrow_1^{j-2}, 1\uparrow_2^{j-2}) & X^j + X^{j-1}Y + XZ^{j-1} + Y^j & yz, x^2z, xy^2, x^{j-2}y - z^{j-1}, x^{j-1} - y^{j-1} - z^{j-1} \\ \end{array} $	
$15 \qquad (((j+1)_0, 3\uparrow_1^{j-3}, 2_{1,j-2}, 1\uparrow_2^{j-2}) \qquad \qquad X^j + X^2 Y^{j-2} + XZY^{j-2} \qquad \qquad z^2, x^2 z, x^2 y - xyz, y^{j-1}, x^{j-1} - zy^{j-2}$	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	<i>j</i> -1
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	²
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	
$ \begin{array}{c} 21 \\ ((j+1)_0, (j-1)_1, 1_{1,j-1}, (1+j^{2-2})^3) \\ X^j + X^{j-1}Y + Y^{j-2}Z^2 \\ \end{array} \\ \begin{array}{c} X^j + X^{j-1} - x^{j-2}y - y^{j-3}z^2 \\ xz, z^3, xy^2, y^{j-1}, x^{j-1} - x^{j-2}y - y^{j-3}z^2 \\ \end{array} \\ \end{array} $	
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	$x^{j-3}yz$
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	2
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	<i>z</i> ²
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	i-3 2
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	c ^{7 - 5} z ²
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	

Table 11: The JDT #1-40 for $T = (1, 3, 5^k, 3, 1), j = k + 3, |T| = 5j - 7, k \ge 4$ (Theorem 4.8).

#	S	F	$I = \operatorname{Ann}(F)$
41	$(5\uparrow_0^{j-4}, 3_{1,j-3}, 1_{2,j-2})$	$Y^{j-4}(X^4 + X^2YZ + Y^2Z^2)$	$x^2 - yz, z^3, xz^2, y^{j-1}, xy^{j-2}$
42	$(5\uparrow_0^{j-4}, 2_{1,2,j-3,j-2})$	$X^4Y^{j-4} + XY^{j-2}Z$	$z^2, x^2z, x^3 - y^2z, y^{j-1}, x^2y^{j-3}$
43	$(4\uparrow_0^{j-3}, 2_{1,j-2}, 1\uparrow_2^{j-2})$	$X^3Y^{j-3} + XY^{j-2}Z + Z^j$	$x^2 - yz, xz^2, yz^2, y^{j-1}, xy^{j-2} - z^{j-1}$
44	$(4\uparrow_0^{j-3},1\uparrow_1^{j-1},1_{2,j-2})$	$X^{3}Y^{j-3} + (Y+Z)^{[j]} - Y^{j} - ZY^{j-1}$	$xz, yz^2 - z^3, x^3 + y^2z - z^3, y^{j-1}, xy^{j-2}$
45	$(3\uparrow_0^{j-2},2\uparrow_1^{j-2})$	$X^2Y^{j-2} + XZ^{j-2}$	$yz, x^2z, x^3, y^{j-1}, xy^{j-2} - z^{j-1}$
46	$(3\uparrow_0^{j-2},1\uparrow_1^{j-1},1\uparrow_2^{j-2})$	$X^2 Y^{j-2} + Y Z^{j-1}$	$xz, y^2z, x^3, y^{j-1}, x^2y^{j-3} - z^{j-1}$
47	$(2\uparrow_0^{j-1}, 2\uparrow_1^{j-2}, 1\uparrow_2^{j-2})$	$XYZ^{j-2} + Y^j$	$x^2, y^2z, xy^2, z^{j-1}, y^{j-1} - xz^{j-2}$

Table 12: The JDT #41-47 for $T = (1, 3, 5^k, 3, 1), j = k + 3, |T| = 5j - 7, k \ge 4.$

List of Tables

- 1 Case $s = 2, k \ge 2$; codimension two: the action of ι takes 1_b to 3_b . 16
- 2 Case $s = 3, k \ge 2$, codimension 2. The action of ι takes 1_a to 2_a . 17
- 3 Case $s = 3, T = (1, 3^k, 1), k \ge 3$, codimension three; list of JDTs and reference to corresponding
- 4 Case s = 4: the sequences Δ and number of rank matrices. 21
- 5 Case s = 5: the sequences Δ and number of rank matrices. 22
- 6 Case s = 6: the sequences Δ and number of rank matrices. 26
- 7 Complete 8 JDT for $T = (1, 3^k, 1), k \ge 3, j = k + 1$ (Theorem 4.6). 39
- 8 Complete 26 JDT for $T = (1, 3, 4^k, 3, 1), k \ge 3$ with the corresponding Macaulay dual generator
- 9 The 12 JDT for T = (1, 3, 5, 3, 1) with the Macaulay dual generator, and ideal (Theorem 4.8). 4
- 10 The 35 JDT for T = (1, 3, 5, 5, 3, 1), with Macaulay dual generator and reference to corresponding
- 11 The JDT #1-40 for $T = (1, 3, 5^k, 3, 1), j = k + 3, |T| = 5j 7, k \ge 4$ (Theorem 4.8). 43
- 12 The JDT #41-47 for $T = (1, 3, 5^k, 3, 1), j = k + 3, |T| = 5j 7, k \ge 4.$ 44

References

- [AAISY] N. Abdallah, N. Altafi, A. Iarrobino, A. Secealanu, and J. Yaméogo: Lefschetz properties of some codimension three Artinian Gorenstein algebras, J. Algebra 625 (2023) 28-45. [AEIY] N. Abdallah, J. Emsalem, A. Iarrobino and J. Yaméogo: Limits of graded Gorenstein algebras of Hilbert function $(1, 3^k, 1)$, Eur. J. Math. 10 (2024), no. 1, Paper No. 9, 42 pp. [Al1] N. Altafi: Jordan types with small parts for Artinian Gorenstein algebras of codimension three, Linear Algebra Appl. 646 (2022), 54-83. [Al2]N. Altafi: Hilbert functions of Artinian Gorenstein algebras with the strong Lefschetz property, Proc. Amer. Math. Soc. 150 (2022), no. 2, 499-513. [Al3] N. Altafi: A note on Jordan types and Jordan degree types, to appear in "Deformations of Artinian algebras and Jordan type," A. Iarrobino, P. Macias Margues, M. E. Rossi, J. Vallès, eds., A.M.S. Contemporary Mathematics #805 (2024). arXiv:math.AC/2303.13673. [AlIK] N. Altafi, A. Iarrobino, and L. Khatami: Complete intersection
- [AIIK] N. Altafi, A. Iarrobino, and L. Khatami: Complete intersection Jordan types in height two, J. Algebra 557 (2020), 224–277.

- [AIMM] N. Altafi, A. Iarrobino, and P. Macias Marques: Jordan type of an Artinian algebra, a survey, to be published in "Lefschetz Properties: current and new directions", Springer INdAM series. arXiv:math.AC/2307.00957 (2023).
- [AIKY] N. Altafi, A. Iarrobino, L. Khatami, and J. Yaméogo: Number of generators of ideals in Jordan cells of the family of graded Artinian algebras of height two, J. Pure Appl. Algebra 227(2023), no.12, Paper No. 107419, 43 pp.
- [BMMNZ] M. Boij, J. Migliore, R.M. Miró-Roig, U. Nagel, F. Zanello: On the weak Lefschetz property for Artinian Gorenstein algebras of codimension three. J. Algebra 403 (2014), 48–68.
- [BMMN] M. Boij, J. Migliore, R. Miró-Roig, and U. Nagel: The non-Lefschetz locus, J. Algebra 505 (2018), 288–320.
- [BrKa] H. Brenner, and A. Kaid: A note on the weak Lefschetz property of monomial complete intersections in positive characteristic, Collect. Math. 62 (2011), no. 1, 85–93.
- [Br] J. Briançon: Description de Hilbⁿ $\mathbb{C}\{x, y\}$, Invent. Math. 41 (1977), 45–89.
- [BruH] W. Brums and J. Herzog: *Cohen-Macaulay rings*, Cambridge Studies in Advanced Mathematics #39, Cambridge University Press, Cambridge, UK, (1953); revised paperback edition 1998.
- [BuEi] D. Buchsbaum and D. Eisenbud: Algebra structures for finite free resolutions, and some structure theorems for codimension three, Amer. J. Math. **99** (1977), 447–485.
- [CoV] A. Conca, and G. Valla: Betti numbers and lifting of Gorenstein codimension three ideals, Comm. Algebra 28 (2000), no. 3, 1371-1386.
- [CGo] B. Costa and R. Gondim: *The Jordan type of graded Artinian Gorenstein algebras*, Adv. in Appl. Math. 111 (2019), 101941, 27 pp.
- [Di] S. J. Diesel: Some irreducibility and dimension theorems for families of height 3 Gorenstein algebras, Pacific J. Math. 172 (1996), 365–397.

[Gon] R. Gondim: On higher Hessians and the Lefschetz properties, J. Algebra 489 (2017), 241–263. [GonZa] R. Gondim and G. Zappalà: On mixed Hessians and the Lefschetz properties, J. Pure Appl. Algebra 223 (2019), no. 10, 4268–4282. [GorNoe] P. Gordan and M. Noether: Uber die algebraischen Formen deren Hesse'sche Determinante identisch verschwindet, Math. Ann. 10 (1876), no. 4, 547-568. [HScSe] B. Harbourne, H. Schenck, A. Seceleanu: Inverse systems, Gelfand-Tsetlin patterns and the weak Lefschetz property, J. Lond. Math. Soc. (2) 84 (2011), no. 3, 712–730. [Ha1] T. Harima: Characterization of Hilbert functions of Gorenstein Artin algebras with the weak Stanley property, Proc. Amer. Math. Soc. 123 (1995), no. 12, 3631–3638. [Ha2] T. Harima: A note on Artinian Gorenstein algebras of codimension three, J. Pure Appl. Algebra 135 (1999), no. 1, 45–56. [HaW1] T. Harima and J. Watanabe: The central simple modules of Artinian Gorenstein algebras, J. Pure Appl. Algebra 210 (2007), no. 2, 447-463. [HaW2] T. Harima and J. Watanabe: The strong Lefschetz property for Artinian algebras with non-standard grading, J. Algebra 311 (2007), 511 - 537.[HMNW] T. Harima, J. Migliore, U. Nagel, and J. Watanabe: The weak and strong Lefschetz properties for Artinian K-algebras, J. Algebra 262 (2003), no. 1, 99–126. [H-W] T. Harima, T. Maeno, H. Morita, Y. Numata, A. Wachi, and J. Watanabe: The Lefschetz properties. Lecture Notes in Mathematics, 2080. Springer, Heidelberg, 2013. xx+250 pp. ISBN: 978-3-642-38205-5. [IK] A. Iarrobino and V. Kanev: Power Sums, Gorenstein Algebras, and Determinantal Varieties, Appendix C by Iarrobino and Steven L. Kleiman The Gotzmann Theorems and the Hilbert scheme, Lecture Notes in Mathematics, 1721. Springer-Verlag, Berlin, 1999. xxxii+345 pp. ISBN: 3-540-66766-0 (1999), 345+xxvii p.

[IMM]	A. Iarrobino, P. Macias Marques, and C. McDaniel: Artinian algebras and Jordan type, J. Commut. Algebra 14 (3), 365-414, (2022).
[Mac]	F.H.S. Macaulay: <i>The algebraic theory of modular systems</i> , Cambridge Mathematical Library. Cambridge University Press, Cambridge, 1916. Reissued with an introduction by P. Roberts, 1994. xxxii+112 pp. ISBN: 0-521-45562-6.
[MW]	T. Maeno and J. Watanabe: Lefschetz elements of Artinian Goren- stein algebras and Hessians of homogeneous polynomials, Illinois J. Math. 53 (2009) 593–603.
[St]	R. Stanley: <i>Hilbert functions of graded algebras</i> , Advances in Math. 28 (1978), 58-73.