

The 2024 release of the ExoMol database: molecular line lists for exoplanet and other hot atmospheres

Jonathan Tennyson^{a,*}, Sergei N. Yurchenko^a, Jingxin Zhang^a, Charles A. Bowesman^a, Ryan P. Brady^a, Jeanna Buldyreva^g, Katy L. Chubb^b, Robert R. Gamache^h, Maire N. Gorman^{b,j}, Elizabeth R. Guest^a, Christian Hill^e, Kyriaki Kefala^a, A. E. Lynas-Gray^{a,c,d}, Thomas M. Mellor^a, Laura K. McKemmish^f, Georgi B. Mitev^a, Irina I. Mizus^k, Alec Owens^a, Zhijian Peng^a, Armando N. Perri^a, Marco Pezzella^{a,1}, Oleg L. Polyansky^{a,k}, Qianwei Qu^a, Mikhail Semenov^a, Oleksiy Smola^a, Andrei Solokov^a, Wilfrid Somogyi^a, Apoorva Upadhyay^a, Samuel O.M. Wright^a, Nikolai F. Zobov^k

^aDepartment of Physics and Astronomy, University College London, Gower Street, London WC1E 6BT, UK

^bSchool of Physics, HH Wills Physics Laboratory, Tyndall Avenue, Bristol, BS8 1TL, UK

^cDepartment of Physics, University of Oxford, Keble Road, Oxford OX1 3RH, UK

^dDepartment of Physics and Astronomy, University of the Western Cape, Private Bag X17, Bellville 7535, South Africa

^eNuclear Data Section, International Atomic Energy Agency, Vienna A-1400, Austria

^fSchool of Chemistry, University of New South Wales, 2052 Sydney, Australia

^gInstitut UTINAM UMR CNRS 6213, Université de Franche-Comté, 16 Route de Gray, 25030 Besançon cedex, France

^hDepartment of Environmental, Earth, and Atmospheric Sciences University of Massachusetts Lowell, Lowell, MA 01854 USA

ⁱDipartimento di Fisica e Geologia, Università di Perugia, Via Pascoli SNC, 06123 Perugia, Italy

^jSchool of Mathematical and Physical Sciences, University of Sussex, Brighton, BN1 9QH, UK

^kInstitute of Applied Physics, Russian Academy of Sciences, 46 Ulyanov Street, Nizhny Novgorod, 603950, Russia

Abstract

The ExoMol database (www.exomol.com) provides molecular data for spectroscopic studies of hot atmospheres. These data are widely used to model atmospheres of exoplanets, cool stars and other astronomical objects, as well as a variety of terrestrial applications. The 2024 data release reports the current status of the database which contains recommended line lists for 91 molecules and 224 isotopologues giving a total of almost 10^{12} individual transitions. New features of the database include extensive "MARVELization" of line lists to allow them to be used for high resolution studies, extension of several line lists to ultraviolet wavelengths, provision of photodissociation cross sections and extended provision of broadening parameters. Some of the in-house data specifications have been rewritten in JSON and moved to conformity with other international standards. Data products, including specific heats, a database of lifetimes for plasma studies, and the ExoMolHR web app which allows exclusively high resolution data to be extracted, are discussed.

Keywords: infrared, visible, Einstein A coefficients, transition frequencies, partition functions, cooling functions, lifetimes, cross sections, k coefficients, pressure broadening, photodissociation, ultraviolet

*Corresponding author.

Email address: j.tennyson@ucl.ac.uk (Jonathan Tennyson)

1. Introduction

The ExoMol project was founded in 2011 to provide molecular line lists for exoplanet and other atmospheres [1] with a particular emphasis on providing results for elevated temperatures which are not adequately covered by databases such as HITRAN [2] or GEISA [3]. ExoMol has followed the practice of providing data release papers every four years with the first two in 2016 [4] and 2020 [5]; the present paper is the third such release. A short review of the first decade of activity by the ExoMol project has been given by the project founders Tennyson and Yurchenko [6].

Since the launch of the project, ExoMol has created line lists for 67 molecules (approaching 150 isoptopologues) details of which are given below. Also discussed below are line lists created by other teams, which are included in the database to ensure coverage of key molecular species as far as possible. A major goal of the original ExoMol project was to provide line lists which were as complete as possible as these have been shown to be essential for correctly reproducing broad band spectra [7] and also opacities. Since that time, high resolution Doppler-shift cross-correlation spectroscopy has been pioneered as a very successful tool for detecting atoms and molecules in exoplanet atmospheres [8–10]. Key line lists constructed with an emphasis on completeness lacked the necessary accuracy for these high resolution studies [11, 12], see also Yurchenko *et al.* [13], thus addressing the needs of high resolution observation led to a significant extension in the scope of the ExoMol project.

The tension between completeness and accuracy in providing hot line lists for astronomical modelling is well-known [14]. ExoMol has developed techniques based on the use of the MARVEL (measured active vibration-rotation energy levels) algorithm [15] to provide empirical energy levels from high resolution laboratory spectra. High accuracy energy levels plus extrapolation from them [16] can be used to provide accurate transitions while retaining the completeness of the computed line list. These procedures are discussed in Sect. 2.1.

The ExoMol project has continued to provide line lists for new species and has updated and/or extended existing line lists for a number of other molecules. The database has also developed in other directions: notably there has been increasing provision for photoabsorption at shorter wavelengths. Extending the database to ultraviolet wavelengths has led to the need to consider not only line absorption at these wavelengths but also continuum absorption, line broadening due to predissociation and photodissociation rates. As a result we have had to both develop new methods for treating photodissociation [17], predissociation [18–20] and continuum absorption [19], and to extend the ExoMol data model to allow for these processes [21].

Since the ExoMol 2020 release, ExoMol line lists have been used extensively in the modelling of gas giant atmospheres, such as sub-Neptunes [22–37], Neptune-like exoplanets [38–42], so-called Hycean planets [43–45], Saturn-like exoplanets [46–59], young Jupiters [60], cool Jupiters [61], warm Jupiters [27, 62–66], hot Jupiters [29, 67–132] and ultra-hot Jupiters [12, 133–168]. They have also been employed in characterising the atmospheres of terrestrial planets [169–179], particularly hot rocky exoplanets [180–188], those containing volatiles [189] and those undergoing photoevaporation [190, 191]. Within our Solar System, they have been used to study the Venusian atmosphere [192, 193], volcanic plumes on Io [194] and comets [195]. In all kinds of planetary atmospheres they have been utilised in the modelling of clouds and hazes [22, 26, 54, 71, 102, 104, 115, 196–203] and photochemistry [204]. ExoMol line lists have been used to test the detection capabilities of current and future space based telescopes [27, 178, 205–213] and have frequently been incorporated in a variety of radiative transfer, forward-modelling, spectral synthesis and opacity codes [95, 214–224]. They have also been employed to study the detectability of biosignatures [225] and prebiosignatures [175]. In stars, ExoMol line lists have also aided in the study of molecular species in the atmospheres of main sequence dwarfs [226–232], giants [233–235], brown dwarfs [60, 200, 202, 236–254], eruptive Young Stellar Objects [255], in stellar winds from Oxygen-rich [256–258] and Carbon-rich [259, 260] Asymptotic Giant Branch stars, in cool Carbon stars [261, 262], in supergiants [256], in stellar merger remnants [263, 264] and modelling supernova ejecta [265]. They have also been used to look at molecules in the interstellar medium (ISM) [266, 267] and to study the isotope exchange rates of molecules important in those environments [268]. Outside of astronomy, ExoMol line lists have been employed in laser absorption spectroscopy [269–272], laser induced breakdown spectroscopy [273, 274], laser fragmentation [275], laser induced fluorescence [276], plasma stoichiometry [277], fusion plasmas [278], the design of gas sensors [279], combustion [280] and explosions [281]. ExoMol line lists are often used as a benchmark in *ab initio* calculations [282–287] and to compare line strengths [288–295].

The ExoMol database is complete enough to be used to generate other data products. Examples include the creation of the Lifetimes DataBase (LiDB) of vibronic state lifetimes, primarily for use in plasma modelling [296], see Sect. 8.2; a database of NASA polynomial fits to specific heats generated from ExoMol data [297] and a

database of high resolution transitions suitable for spectral assignment [298], see Sect. 8.1. Importantly ExoMol line lists have been used to give molecular opacities in the ExoMolOP database [299] discussed below; opacities based on ExoMol data have also been generated by several other groups [137, 248, 300–310]. The generation of opacities, see Sect. 3.5, requires treatment of line broadening, progress on this topic for the ExoMol project is discussed in Sect. 4. ExoMol is providing input to JWST (James Webb Space Telescope) through the MAESTRO (Molecules and Atoms in Exoplanet Science: Tools and Resources for Opacities) database, see <https://science.data.nasa.gov/opacities/> and the exoplanet characterisation Ariel Space Mission [311], which is due to launch in 2029.

Of course ExoMol is not the only source of spectroscopic data for studies of atmospheres. HITRAN [312] is a well established database [313] which is designed for use at temperatures in the region of 296 K and contains molecules of importance in the Earth’s atmosphere. HITEMP [314] extends HITRAN to higher temperatures, albeit currently for only eight molecules. Where appropriate the extension of HITRAN or HITEMP is based on ExoMol data [315], and ExoMol uses HITRAN line lists for the diatomic molecules HF, HCl, HBr and CO for which the HITRAN compilation is appropriate for use at higher temperatures. The TheoRets database [316] contains very extensive computed line lists for nine polyatomic molecules with five or more atoms; not all of these line lists are suitable for high temperature studies. The NASA Ames group provides very extensive line lists for 6 molecules; 5 triatomics plus ammonia. Their CO₂ line lists, which include isotopologues, are designed for use at temperatures up to 3000 K [317]. The SO₂ ExoMol line list [318] was produced in collaboration with the NASA Ames group. The MoLLIST [319] line lists due to Bernath and co-workers provides empirical line lists designed for use at higher temperatures for 26 diatomics, ammonia and methane as well as cross-sections for eight polyatomic molecules with six or more atoms. A number of the MoLLIST diatomic line lists are also available from the ExoMol database, see Sect. 2.6. The VALD [320] and Kurucz [321] databases, which are largely aimed at stellar atmospheres, contain data on some diatomic species.

As discussed in Sect. 6, with this release ExoMol has started providing temperature-dependent photodissociation cross sections. The main current provider of photodissociation cross sections for astronomical studies is the Leiden VUV cross section database [322] whose data are largely designed for use in studies of the (cold) interstellar medium. Thus far Leiden have not considered the temperature of the molecule in their photodissociation cross sections as the data are largely aimed for use at temperature of 300 K or below.

2. Line lists

Table 1 summarizes the recommended ExoMol line lists present in the database. Note that a number of ExoMol line lists have been replaced by either extended and/or improved upon line lists for the same species; these older line lists are no longer recommended and are not listed in the table although they are still available via the ExoMol website. The 2020 data release introduced the use of uncertainties in the energies given in the .states file to allow high resolution transitions to be identified. This feature has now been introduced for all ExoMol-generated recommended line lists, although uncertainties are generally not provided for old line lists or those provided from other sources. At the same time, the previously optional lifetime column in the .states file has been made mandatory and all .states files for recommended line lists include lifetimes; this feature is important for modelling line-broadening effects due to predissociation [21].

Below, specific notes for some individual line lists are given. Those line lists which were included in the 2020 release and have not been updated are not explicitly discussed, please refer to the 2020 manuscript or the cited original line list paper for further details. An overview of the temperature-dependent cross sections for earlier ExoMol line lists is already available [323] and below we give similar plots for many of the newly computed line lists. Before discussing individual line lists we discuss our strategy for improving the accuracy of calculated line positions.

Table 1: Datasets created by the ExoMol project and included in the ExoMol database: recommended line lists only. Line lists denoted with a ✓ are suitable for high resolution studies.

Paper	Molecule	N_{iso}	T_{max}	N_{elec}	N_{lines}^a	DSName	Reference
III	HCN/HNC	1 ^a	4000	1	34 418 408	Harris	✓ [324]
V	NaCl	2	3000	1	702 271	Barton	[325]
V	KCl	4	3000	1	1 326 765	Barton	[325]
VII	PH ₃	1	1500	1	16 803 703 395	SAITY	[326]
VIII	H ₂ CO	1	1500	1	12 688 112 669	AYTY	✓ [327]
IX	AlO	4	8000	3	4 945 580	ATP	✓ [328]
X	NaH	2	7000	2	79 898	Rivlin	[329]
XI	HNO ₃	1	500	1	6 722 136 109	AlJS	[330]
XII	CS	8	3000	1	548 312	JnK	✓ [331]
XIII	CaO	1	5000	5	21 279 299	VBATHY	✓ [332]
XIV	SO ₂	1	2000	1	1 300 000 000	ExoAmes	✓ [318]
XV	H ₂ O ₂	1	1250	1	20 000 000 000	APTY	[333]
XVI	H ₂ S	1	2000	1	115 530 373	AYT2	✓ [334]
XVII	SO ₃	1	800	1	21 413 927 818	UYT2	✓ [335]
XIX	H ₂ ^{17,18} O	2	3000	1	519 461 789	HotWat78	✓ [336]
XX	H ₃ ⁺	1	3000	1	127 542 657	MiZATeP	✓ [337]
XXII	SiH ₄	1	1200	1	62 690 449 078	OYT2	[338]
XXIII	PO	1	5000	1	2 096 289	POPS	[339]
XXIII	PS	1	5000	3	30 394 544	POPS	[339]
XXIV	SiH	4	5000	3	1 724 841	SIGHTLY	[340]
XXV	SiS	12	5000	1	91 715	UCTY	[341]
XXVI	NS	6	5000	1	3 479 067	SNaSH	[342]
XXVII	C ₂ H ₄	1	700	1	49 841 085 051	MaYTY	[343]
XXIX	CH ₃ Cl	2	1200	1	166 279 593 333	OYT	[344]
XXX	H ₂ ¹⁶ O	1	5000	1	5 745 071 340	POKAZATEL	✓ [345]
XXXI	C ₂	3	5000	8	6 080 920	8states	✓ [346]
XXXII	MgO	5	5000	5	72 833 173	LiTY	✓ [347]
XXXIII	TiO	5	5000	13	58 983 952	Toto	✓ [348]
XXXIV	PH	1	4000	2	65 055	LaTY	✓ [349]
XXXV	NH ₃	1	1500	1	16 941 637 250	CoYuTe	✓ [350]
XXXVI	SH	2	3000	2	572 145	GYT	✓ [351]
XXXVII	HCCH	1	2000	1	4 347 381 911	aCeTY	✓ [352]
XXXVIII	SiO ₂	1	3000	1	32 951 275 437	OYT3	[353]
XXXIX	CO ₂	1	3000	1	7 996 570 390	UCL-4000	✓ [354]
XL	H ₃ O ⁺	1	1500	1	2 089 331 073	eXeL	✓ [355]
XLI	KOH	1	3500	1	38 362 078 911	OYT4	[356]
XLI	NaOH	1	3500	1	49 663 923 092	OYT5	[356]
XLII	NO	1	3500	4	4 596 666	XABC	✓ [357]
XLIII	NaO	1	2500	2	4 726 283	NaOUCMe	✓ [358]
XLIV	SiO	1	3500	10	91 395 763	SiOUVenIR	✓ [359]
XLV	CaH	1	5000	3	293 151	XAB	✓ [360]
XLV	MgH	3	5000	3	88 575	XAB	✓ [360]
XLVI	SiN	4	3000	6	43 646 806	SiNful	✓ [360]
XLVII	CaOH	1	3000	3	24 215 753 701	OYT6	✓ [360]
XLVIII	H ₂ CS	1	2000	1	43 56 116 660	MOTY	✓ [361]
XLIX	AlCl	2	5000	4	4 722 048	YNAT	✓ [362]
L	H ₃ ⁺	4	3000	1	-	MiZo	✓ [363]
LI	LiOH	2	3000	1	331 274 717	OYT7	[364]
LII	CH ⁺	2	5000	2	34 194	PYT	[365]
LIII	YO	3	5000	6	60 678 140	BRYTS	[366]
LIV	AlIH	4	5000	2	36 152	AloHa	✓ [19]
LV	VO	1	5400	15	58 904 173 243	HyVO	✓ [367]
LVI	SO	1	5000	8	7 008 190	SOLIS	✓ [368]
LVII	CH ₄	1	2000	1	50 395 644 806	MM	✓ [369]
LVIII	OCS	1	2000	1	2 482 380 391	OYT8	✓ [370]
LIX	N ₂ O	5	2000	1	1 360 351 722	TYM	✓ [371]
LX	¹⁵ NH ₃	1	1000	1	929 795 249	CoYuTe-15	✓ [372]
LXI	OH	1	5000	8	1 685 102	MYTHOS	✓ [373]
LXII	C ₃	3	3000	1	6 797 725 272	AtLast	✓ [374]
LXIII	HDO	1	3000	1	3 000 000	TBD	✓ [375]
LXIV	PN	1	5000	1	634 243	PaiN	✓ [376]

Paper: Number in series published in Mon. Not. R. Astron. Soc.; N_{iso} : the number of isotopologues considered; T_{max} : maximum temperature for which the line list is complete; N_{elec} : number of electronic states considered; N_{lines} : number of lines, the value is for the main (parent) isotopologue.

^a The Larner line list [377] for H¹³CN/HN¹³C is recommended.

2.1. MARVELization

In terms of the resolving power, defined via wavelength λ and its uncertainty $\Delta\lambda$ as $R = \frac{\lambda}{\Delta\lambda}$, calculated ExoMol line lists are generally of sufficient accuracy to be useful for $R < 10\,000$. This is sufficient for current spaceborne observations of exoplanets, for example JWST has a maximum $R = 3000$. However, ground based observations, and in particular high resolution Doppler-shift cross-correlation spectroscopy, work at much higher resolutions, typically $R = 100\,000$ or even higher. It is clear that the standard theoretical method, even using empirical data to improve the spectroscopy model [378], struggles to approach the level of accuracy required for this resolution. A different approach has therefore been adopted based on the explicit use of empirically derived energy levels.

To provide empirical energy levels we used the MARVEL algorithm [15]. Originally developed to improve representations of water vibration-rotation spectra [379, 380], the MARVEL methodology is actually agnostic about the type of spectra being studied and we have applied it widely to rovibronic problems as well as, for VO, to hyperfine-resolved spectra [381]. In essence MARVEL takes a list of assigned high resolution transitions with uncertainties and inverts them to give a list of empirical energies with associated uncertainties. MARVEL does this by constructing a spectroscopic network [382] composed of (all available) assigned, measured transitions with uncertainties; this network is inverted using the so-called X-matrix method [383] to give empirical energy levels each with an associated uncertainty.

The ExoMol data structure gives a single set of energy levels in the `.states` file and a set of Einstein A coefficients in the transitions (`.trans`) file, which are processed together to compute transition wavenumbers and intensities. This structure lends itself to straightforward improvement by replacing calculated energy levels with empirical ones, increasing the accuracy of predicted transition wavenumbers. This approach has the advantage over simply replacing computed transition wavenumbers with measured ones because it produces accurate predictions for the transition wavenumbers of many yet to be observed transitions. For example, Al-Derzi *et al.* [384] performed a MARVEL project for formaldehyde (H_2CO) which used a network of 16 403 unique transitions to determine 5029 energy levels. Substituting these 5029 energies into the AYTY ExoMol H_2CO line list [327] resulted in 367 779 transitions whose wavenumbers were determined using these empirical energies. We refer to this process as MARVELization. In practice there are a number of ways of improving on the energy levels provided by the variational calculations as discussed by McKemmish *et al.* [16]. Even so, it is usually only possible to MARVELize a small proportion of the energy levels (for example the AYTY line list contains over 12 billion transitions) although in general the MARVELized levels include the ones involved in the strongest transitions. We note that the A in MARVEL stands for active which means that MARVEL datasets can be actively updated when new high resolution measurements become available. Thus, for example, a new optical frequency comb Fourier transform spectrum of formaldehyde by Germann *et al.* [385] was recently used to update the MARVEL results of Al-Derzi *et al.* [384] and the AYTY line list; use of this very high resolution spectrum added a further 82 new energy levels to the MARVEL compilation and, perhaps more importantly, substantially improved the accuracy of many of the energy levels. Line lists that have been MARVELized are identified by ✓ marks in Table 1.

To use the uncertainties in the energy levels it is assumed that the uncertainty in the transition wavenumber, $\Delta\omega$, is given by

$$\Delta\omega = \sqrt{(\Delta E')^2 + (\Delta E'')^2}, \quad (1)$$

where $\Delta E'$ and $\Delta E''$ are the uncertainties in the upper and lower state energy levels, respectively. Note that on occasion this uncertainty will be an overestimate as there are situations where a precisely determined transition has been measured between two states whose energies are both less well determined. However, we suggest that the current implementation should be adequate for most practical purposes. As part of this data release we have ensured that all ExoMol-generated recommended datasets contain uncertainties in their energy levels and the source of these energies are marked using the codes given below in Sect. 3.1. For cases where these levels have not been MARVELized, these uncertainties can be quite large. Conversely, transitions between two states with MARVELized energies should be accurate enough to use in high resolution studies. Our post-processing codes, ExoCross [386] and PyExoCross [387], give the option of generating spectra or cross sections only using transitions with low uncertainties and our new web portal ExoMolHR, see Sect. 8, provides a database of high resolution transitions which can be interrogated interactively.

An underlying assumption of the MARVELization process is that the relatively small shifts in the line positions do not significantly influence the line intensities. In most cases this is probably a reasonable assumption but there is one set of circumstances where things are more complicated. Resonances caused by the accidental interaction of levels in different vibrational or vibronic states can lead to severe perturbations of the intensities




Fig. 1: Cross sections generated using the ExoMol line list XAB for NO at $T = 300$ K and 2000 K.

caused by so-called intensity stealing between transitions. This effect is known to be very sensitive to the details of the calculation; Lodi and Tennyson [388] designed a method for identifying those transitions which are sensitive to these interactions by performing repeat calculations with different potential energy and dipole moment surfaces. This method has proved important for high accuracy calculations [389]. However, the Lodi-Tennyson method requires the generation of, at least, four line lists for given species and thus far has not been employed for the large and usually expensive ExoMol line lists.

2.2. Diatomics

2.2.1. AlO, paper XVIII

The ATP line list for aluminium monoxide, AlO, has been updated using MARVEL [390]. AlO has been detected in a number of objects such as hot Jupiter exoplanets WASP-43b [391] and HAT-P-41b [77], as well as in eruptive young stellar objects (YSOs) [255] using the ATP line list. The ATP line list has also been used in plasma studies [274, 392, 393].

2.2.2. NO, papers XXI and XLII

The full rovibronic $^{14}\text{N}^{16}\text{O}$ XABC line list [357] replaces the ground state NOName [394] line list. A MARVEL study was undertaken as part of the XABC line list construction. However, NOName should still be used for the minor isotopologues of NO.

The XABC line list gives thorough coverage of the so-called γ , β and δ band systems, which correspond, respectively, to the $\text{A}^2\Sigma^+ - \text{X}^2\Pi$, $\text{B}^2\Pi - \text{X}^2\Pi$ and $\text{C}^2\Pi - \text{X}^2\Pi$ electronic bands. We note that the γ band system has recently been proposed as a potential biomarker in exoplanetary atmospheres [395]. Cross sections of NO computed using XAB are shown in Fig. 1.

2.2.3. AlH, papers XXI and XLIV

Pavlenko et al. [396] used the AlHambra line list [397] to analyse the spectrum of AlH in Proxima Cen. While the line list reproduced the majority of the observed spectrum well, it did not reproduce the predissociative lines accurately either in their position or their widths. As a result it was decided to implement procedures [18] and data structures [21] to represent line broadening due to predissociation within the project. A new line list, AloHa [19], superseded the AlHambra line list. AloHa has been MARVELized for both AlH and AlD, and also provides an accurate representation of the predissociation lifetime. The line list also provides continuum absorption cross sections describing photoabsorption, i.e. absorption to continuum (unbound) states. The structure of the continuum absorption cross sections is introduced in Sect. 3.3.

2.2.4. MgO, paper XXXII

The LiTY line list [347] for magnesium monoxide, MgO, has been updated for the four lowest electronic states using MARVEL [16]. Additionally, the Predicted Shift (PS) methodology was used to improve the calculated energies and estimate energy uncertainties [16]. The updated line list was used to update the line lists for the $^{24}\text{Mg}^{17}\text{O}$, $^{24}\text{Mg}^{18}\text{O}$, $^{25}\text{Mg}^{16}\text{O}$, and $^{26}\text{Mg}^{16}\text{O}$ isotopologues with the isotopologue extrapolation procedure (IE) [16]. MgO has been suggested as one of the constituents of the so-called lava planets [181, 398].




Fig. 2: Cross sections generated using the ExoMol line list SiOUVenIR for SiO at $T = 300$ K and 2000 K.

2.2.5. *TiO, paper XXXIII*

The Toto line list for TiO [348] has been widely used. Notably ToTo was successfully used by Pavlenko et al. [399] to extract Ti isotopic abundance using band heads of ^{50}TiO , ^{49}TiO , ^{48}TiO , and ^{47}TiO in the spectra of M dwarfs. This work provides endorsement of the isotopologue extrapolation procedure (IE) [16] used to improve the line lists for minor isotopologues by using data from the main isotopologue. The ToTo line list has recently been updated using new MARVEL energy levels derived incorporating recent laboratory measurements [16]. The .states file was updated using the Predicted Shift (PS) methodology [16]. ToTo has been used for the detection of TiO in a number of exoplanets, such as the ultra-hot Jupiter WASP-189b [145], WASP-33b [141], WASP-76b [400] and WASP-77Ab [126].

2.2.6. *SH, paper XXXVI*

The GYT linelist [401] remains as the recommended ExoMol linelist for the mercapto radical and is recommended for all applications [5]. As in hydroxyl (sec. 2.2.13), SH exhibits strong and complex predissociation due to spin orbit interactions between the $\text{A}^2\Sigma^+$ state and the dissociative, $1^2\Sigma^-$, $1^4\Sigma^-$, and $1^4\Pi$ states. SH/SD predissociation lifetimes will be released soon [18] and lifetimes in the GYT linelist will be updated as per Sect. 5.

2.2.7. *NaO, paper XLIII*

A new line list for NaO called NaOUCMe [358] has been provided; there is very limited high resolution spectroscopic data on NaO so the line list is not suitable for high resolution studies.

2.2.8. *SiO, paper XLIV*

A new line list for silicon monoxide, SiO, called SiOUVenIR, has been provided [359] and replaces the well-used EJBT rovibrational line list [402]. The new line list has been MARVELized and also provides much broader wavelength coverage as it also covers rovibronic transitions. Like MgO, SiO has been suggested as one of the important sources of opacity in lava planets as well as sub-Neptunes [181, 186, 189, 190, 398, 403, 404]. Cross sections of SiO computed using SiOUVenIR are shown in Fig. 2.

2.2.9. *CaH and MgH, paper XLV*

New rovibronic line lists for CaH and MgH called XAB have been computed [360]. The line lists have been MARVELized and consider the important isotopologues of both species. The line lists replace the more restricted rovibrational Yadin line lists [405]. As MARVELized line lists for BeH, as well as BeD and BeT, are also available [406], the Yadin linelists [405] for these alkaline earth monohydrides are no longer recommended. Cross sections of MgH computed using the XAB line list are shown in Fig. 3.

2.2.10. *VO, paper XLV*

The new HyVO line list for Vanadium monoxide [367] replaces the previous ExoMol VOMYT line list [407]. VOMYT was built on a relatively crude theoretical model and known to be inadequate for high resolution studies [12]. The spectrum of VO shows significant splittings due to the hyperfine structure induced by the high spin




Fig. 3: Cross sections of MgH generated using the ExoMol line list XAB at $T = 300\text{ K}$ and 2000 K .

($I = \frac{7}{2}$) and large magnetic moment of the ^{51}V nucleus. To create a VO line list suitable for high resolution studies, it was necessary to perform a MARVEL study which explicitly considered hyperfine effects where possible [381], extend our diatomic variational nuclear motion code Duo to include hyperfine effects [408] and then perform extensive model building studies [409, 410]. HyVO is a fully hyperfine-resolved line list which contains nearly 59 billion transitions, a huge number for a diatomic molecule. The new HyVO line list has been used to detect VO in the atmosphere of WASP-76b via high-resolution cross-correlation spectroscopy [168], strengthening the findings of the previous study of this planet which utilised the old VOMYT line list [159].

2.2.11. SO, paper LVI

The new SOLIS line list for the main sulfur monoxide isotopologue $^{32}\text{S}^{16}\text{O}$ has been computed and MARVELized [368]. The electronic structure of SO exhibits non-adiabatic behavior where an earlier study [411] builds a quasi-diabatic spectroscopic *ab initio* model before refinement to MARVEL data.

SO is expected to be abundant within exoplanets whose atmospheres undergo warm chemistry or have regions with prolific photochemical reactions. It was recently detected in a JWST MIR transmission spectrum of WASP-39b, a Saturn-mass exoplanet, using the SOLIS line list [130].

2.2.12. SiN, paper XLVI

A new line list for silicon mononitride, SiN, called Sinful, has been computed [360] for the main isotopologues, $^{28}\text{Si}^{14}\text{N}$, $^{29}\text{Si}^{14}\text{N}$, $^{30}\text{Si}^{14}\text{N}$, $^{28}\text{Si}^{15}\text{N}$. The line list includes 6 lowest-lying electronic states. The line list has been MARVELized and is in good agreement with previously reported spectra. SiN is considered to be a good indicator for planet-metallicity [412] and has been detected in different media in space [413–419].

2.2.13. OH, paper LXI

The new MYTHOS OH line list [373] replaces the one provided by MoLLIST [420] as the recommended ExoMol line list. The rovibronic line list considers transitions within the ground X $^2\Pi$ electronic state to the A $^2\Sigma^+$, B $^2\Sigma^+$ and C $^2\Sigma^+$ states which all support bound states. The model also includes four dissociative states, 1 $^2\Sigma^-$, 1 $^4\Sigma^-$, 1 $^4\Pi$, and 1 $^2\Delta$, the first three of which causes significant predissociation broadening of levels within the A $^2\Sigma^+$ state [20]; the lifetime broadening effects are explicitly allowed for in the line list. Transitions from the ground state to the dissociative $^2\Sigma^-$ and $^2\Delta$ states are also calculated here and are included in the temperature dependent photoabsorption cross sections. The line list is of a high accuracy having used the empirical energies from a previous MARVEL study [421]. Two datasets are made available: a state bound to bound line list and temperature dependent photoabsorption cross sections which include the continuum transitions. Temperature-dependent photodissociation cross sections for OH are currently being computed.

2.2.14. PN, paper LXIV

A rovibrational line list for phosphorus nitride, named YYLT, was computed early in the ExoMol project [422]. A new, MARVELized rovibronic line list PaiN [376] covering the A-X band system and providing improvements for the X-X transitions has been constructed starting from the *ab initio* spectroscopic model of Semenov et al. [423].

PaiN has replaced YYLT as the recommended line list. We note the PN is likely to be hard to detect in exoplanet atmospheres because it has a strong vibrational fundamental in the $10\text{ }\mu\text{m}$ region, which is too long wavelength for current observational capabilities, and the A–X band system lies in the UV around 250 nm which is also hard to observe and unlikely to be important around cool stars.

2.2.15. Other new diatomic line lists

New line lists are provided for AlCl (YNAT [362]), CH⁺ (PYT [365]) and YO (AloHa [366]), all of which include rovibronic transitions as well as rotation-vibration transitions.

2.3. Triatomics

2.3.1. H₃⁺, Papers XX and L

Bowesman et al. [363] updated the MiZATEP H₃⁺ line list [337] and the ST line list for H₂D⁺ [424] using MARVEL. New MiZo line lists for D₂H⁺ and D₃⁺ were generated [363]; the MiZo D₂H⁺ line list also uses MARVEL energy levels while the energy levels of the more limited D₃⁺ line list were improved using effective Hamiltonian data. All the line lists are suitable for high resolution studies at long wavelengths. A review of the role of H₃⁺ in astrophysics has been provided by Miller et al. [425].

2.3.2. Water, Papers XIX, XXX and LXIII

POKAZATEL [345] remains the recommended line list for H₂¹⁶O, it has been widely and successfully used. Water is of course widely observed in exoplanet spectra and several other line lists including BT2 [426], HITRAN 2010 [314] which is based on BT2, and the pioneering but now dated Partridge and Schwenke [427]. We strongly recommend the adoption of POKAZATEL over these older, less complete and less accurate line lists for this key molecule. Indeed a near infrared laboratory high temperature ($T = 1723\text{ K}$) comparison suggested that POKAZATEL reproduced the observed spectrum roughly six times better than BT2 [428].

Isotopologues H₂¹⁷O and H₂¹⁸O are represented by the HotWat78 line lists [336]. While POKAZATEL is very complete and should give a good representation of water absorption up to at least 5000 K, the HotWat78 line lists are only complete up to 3000 K. We plan to extend the temperature range covered for H₂¹⁷O and H₂¹⁸O.

All H₂O line lists discussed above have been MARVELized [429, 430] and are therefore suitable for high accuracy studies.

A new line list TBD has been computed for deuterated water, HDO [375]. This replaces the old VTT line list [431]. TBD is more complete than VTT. In particular, the potential energy surface (PES) used in the line list calculations was obtained by fitting to the data [432] up to 35 000 cm⁻¹. Energies up to 41 000 cm⁻¹ were used in the linelist calculations. Unlike VTT, the TBD energy levels have been MARVELized using energies from an IUPAC study [433]; we note that a MARVEL project to update these energy levels is in progress which will lead to the release of an updated version of TBD. The spectrum of HDO is significantly different from that of H₂O and the TBD line list should be suitable for use in high resolution studies.

In addition, room temperature line lists for the isotopologues H₂¹⁴O [434], H₂¹⁵O [435] and H₂¹⁹O [436] have recently been added to the databases. These isotopologues, in which the radioactive oxygen atom has a half-life in the region of a minute, may form in thunderstorms [437], in water cooled nuclear and fusion reactors, and as a result of using isotopes in medical procedures.

2.3.3. CO₂, Paper XXXIX

The UCL-4000 line list [354] for carbon dioxide has been produced an accurate PES AMES-1 [438] and a high-level *ab initio* dipole moment surface [439] of CO₂ using the TROVE variational program [440]. UCL-4000 has been widely used but suffers from two issues. It is only for the parent, ¹²C¹⁶O₂, isotopologue and at visible wavelengths it was found to overestimate transition intensities at higher wavenumbers, a problem found with other hot CO₂ line lists [441]. We are therefore in the process of constructing an improved hot line list for this important molecule which will consider all important isotopologues and be MARVELized [442, 443]. Cross sections of CO₂ generated using the UCL-4000 line list are illustrated in Fig. 4.

The same methodology used for UCL-4000 was subsequently used to produce an electric quadrupole (E2) line list for CO₂ [444]. The theoretical E2 transitions were validated experimentally using in the $3.3\text{ }\mu\text{m}$ region [445] and used to detect E2 lines in the Martian atmospheric spectra from the ExoMars ACS observations [446].




Fig. 4: Cross sections generated using the ExoMol line list UCL-4000 for CO_2 at $T = 300 \text{ K}$, 2000 K and 3000 K .




Fig. 5: Cross sections of LiOH generated using the ExoMol line list OYT7 at $T = 300 \text{ K}$ and 2000 K .

2.3.4. LiOH , NaOH , KOH , Papers XLI and LI

New line lists for the alkali metal hydroxides KOH , NaOH and LiOH named OYT4 [356], OYT5 [356] and OYT7 [364], respectively, have been calculated. There is a lack of high resolution spectroscopic data for these systems so these are not MARVELized and therefore not suitable for high resolution studies. Cross sections of LiOH generated using the OYT7 line list are illustrated in Fig. 5.

2.3.5. CaOH , Paper XLVII

A new line list called OYT6 is provided for calcium monohydroxide, CaOH [360]. This is the first rovibronic line list for a triatomic molecule which includes consideration of the important $\tilde{\Lambda}^2\Pi - \tilde{X}^2\Sigma^+$ electronic band system as well as ground state rotation-vibration transitions. This line list was built on a MARVEL study [447], then constructed a spectroscopic model [448] using the rovibronic nuclear motion code for triatomics EVEREST [449]. CaOH is known to be an important component of the atmosphere of cool stars [450]; the OYT6 line list has been used as a thermometer in combustion and explosion applications [281]. Cross sections of CaOH computed using the OYT6 line list are shown in Fig. 6.

2.3.6. OCS , Paper LVIII

A new line list called OYT8 is provided for carbonyl sulphide, OCS [370]; this line list uses a previous MARVEL study [451] and therefore should be suitable for high resolution studies. Given that CO [452], CO_2 [51], SO_2 [53, 130], and now SO have been detected by JWST in exoplanetary atmospheres, the presence of OCS would also appear likely. Major studies are already searching for its spectroscopic signature [56, 453].




Fig. 6: Cross sections of CaOH generated using the ExoMol line list OYT6 at $T = 300\text{ K}$ and 2000 K .

2.3.7. N_2O , Paper LIX

A new line list for nitrous oxide, N_2O , has been computed [371]. N_2O is prominent in the Earth's atmosphere and considered to be an observable potential biomarker on exoplanets [454–456]. Five isotopologues are considered, $^{14}\text{N}_2^{16}\text{O}$, $^{15}\text{N}^{14}\text{N}^{16}\text{O}$, $^{14}\text{N}^{15}\text{N}^{16}\text{O}$, $^{14}\text{N}_2^{17}\text{O}$ and $^{14}\text{N}_2^{18}\text{O}$. The parent isotopologue $^{14}\text{N}_2^{16}\text{O}$ was MARVELized using the MARVEL energies of Tennyson *et al.* [457]. MARVEL studies on the four isotopologues arising from substituting a single atom are in progress.

2.3.8. C_3 , Paper LXII

A new line list for C_3 has been calculated, which considers ro-vibrational transitions within the ground state. Three isotopologues are considered: $^{12}\text{C}_3$, $^{12}\text{C}^{13}\text{C}^{12}\text{C}$ and the asymmetric $^{12}\text{C}^{12}\text{C}^{13}\text{C}$. The $^{12}\text{C}_3$ line list was MARVELized using results from a previous MARVEL study [458].

2.4. Tetratomics

2.4.1. H_2CO , Paper VIII

The AYT line list for formaldehyde [327], $\text{H}_2^{12}\text{C}^{16}\text{O}$, has been updated using MARVEL [384, 385] by replacing the calculated AYT energy levels with MARVEL ones. Based on this update, 373 160 transition frequencies with experimental accuracy were determined. These are illustrated in Fig. 7, where they are compared to 40 670 transitions of $\text{H}_2^{12}\text{C}^{16}\text{O}$ in HITRAN 2020. A line list for H_2^{13}CO is under construction.

2.4.2. NH_3 , Papers XXXV and LX

The CoYuTe line list for $^{14}\text{NH}_3$ ammonia [350] has been updated using new MARVEL data [459]. The CoYuTe line list has been and is continuing to be used to assign ammonia spectra in the near-infrared and to analyze plasma emissions spectra [460]. Laboratory spectra of ammonia need to be analyzed in this region and into the visible not least because pronounced ammonia features are observed in Jupiter in the near-infrared and visible. In particular, the feature at 6474 Å is modelled by CoYuTe [461, 462] better than other available compilations but for which CoYuTe still shows a noticeable shift in wavelength compared to the observations.

A new $^{15}\text{NH}_3$ line list called CoYuTe-15 was generated using the same procedure as for the $^{14}\text{NH}_3$ CoYuTe line list [350], including MARVELization; it replaces the previous room-temperature $^{15}\text{NH}_3$ line list [463] which is also less accurate and not MARVELized. The recent detection of $^{15}\text{NH}_3$ in the atmosphere of a cool brown dwarf [247] suggests that this isotopologue could also be detected in the atmospheres of exoplanets.

2.4.3. HCCH , Paper XXXVII

The aCeTY line list for acetylene, C_2H_2 , has been updated using MARVEL [464].

2.4.4. H_2CS , Paper XLVIII

A new line list for thioformaldehyde, H_2CS , called MOTY has been computed by Mellor *et al.* [361]; the energy levels were on the basis of a MARVEL study [465] to make it suitable for high resolution studies. Figure 8 illustrates the cross sections of H_2CS computed using MOTY.




Fig. 7: Room temperature ($T = 296$ K) spectra of H_2CO from three different sources: lower panel, HITRAN; upper panel, MARVELized transitions [384, 385] overlaid with the AYTY line list [327].




Fig. 8: Cross sections generated using the ExoMol line list MOTY for H_2CS at $T = 300$ K and 2000 K.




Fig. 9: Cross sections of CH_4 generated using the ExoMol line list MM at $T = 300 \text{ K}$ and 2000 K .

2.5. Pentatomics

2.5.1. CH_4 , Paper LVII

A completely new line list for methane, CH_4 , called MM [466] replaces the previous ExoMol methane line lists 10to10 [467] and 30to10 [468]. MM is based on the use of improved theory, an improved potential energy surface plus extensive MARVELization based on a comprehensive, parallel MARVEL study [369]. Line lists for hot methane are also available from Theorets [469, 470] and HITEMP [471]. MM is broadly comparable with the Theorets line list but at high temperatures ($> 1000 \text{ K}$) is significantly more complete than the HITEMP line list. Cross sections of CH_4 generated using the MM line list are illustrated in Fig. 9.

There are many unassigned or partially assigned published high resolution spectra of methane available [369]. We are actively using MM to assign these spectra [472]. Any new assignments we make, or indeed new high resolution spectra, will be used in an extended MARVEL study, the results of which can be used to re-MARVELize and further improve the accuracy of the MM line list.

2.6. Other line lists

Table 2: Datasets recommended as part of the ExoMol project but imported from other sources.

Molecule	N_{iso}	T_{max}	N_{elec}	N_{lines}	DSName	Reference	Methodology
CH	1	5000	4	52201	MoLLIST	[?]	Empirical
AlF	1	5000	1	40 490	MoLLIST	[?]	Empirical
CaF	1	5000	6	14 817	MoLLIST	[?]	Empirical
MgF	1	5000	3	8136	MoLLIST	[?]	Empirical
KF	1	5000	2	10 572	MoLLIST	[?]	Empirical
NaF	1	5000	1	7884	MoLLIST	[?]	Empirical
LiCl	1	5000	4	26 260	MoLLIST	[?]	Empirical
LiF	1	5000	2	10 621	MoLLIST	[?]	Empirical
TiH	1	5000	3	181 080	MoLLIST	[?]	Empirical
CrH	1	5000	2	13 824	MoLLIST	[?]	Empirical
FeH	1	5000	2	93 040	MoLLIST	[?]	Empirical
LaO	1	8000	2	2 066 535	BDL	[475]	Empirical
HF	2	5000	1	7956	Coxon-Hajig	[? ?]	Empirical
HCl	4	5000	1	2588	HITRAN	[? ?]	Empirical
HBr	1	5000	1	3039	HITRAN-HBr	[? ?]	Empirical
CO	9	9000	1	125 496	Li2015	[?]	Empirical
CP	1	5000	2	28752	MoLLIST	[473]	Empirical
ScH	1	5000	6	1 152 827	LYT	[?]	Ab initio
LiH	1	12000	1	18 982	CLT	[484]	Ab initio
LiH ⁺	1	12000	1	332	CLT	[484]	Ab initio
HeH ⁺	4	9000	1	1430	ADSJAAM	[483]	Ab initio
H ₂	1	10000	1	4712	RACPPK	[481]	Ab initio
H ₂	1	10000	3	32 915	ARLR	[482]	Ab initio
HD	1	9000	1	10285	ADSJAAM	[483]	Ab initio
HD ⁺	1	9000	1	10285	ADSJAAM	[483]	Ab initio
CH ₃ F	1	300	1	139 188 215	OYKYT	[?]	Ab initio
AsH ₃	1	300	1	3 600 000	CYT18	[?]	Ab initio
P ₂ H ₂ ^a	2	300	1	10 667 208 951	OY-Trans	[?]	Ab initio
P ₂ H ₂ ^a	2	300	1	11 020 092 365	OY-Cis	[?]	Ab initio
PF ₃	1	300	1	68 000 000 000	MCYTY	[?]	Ab initio
CH ₃	1	1500	1	2 058 655 166	AYYJ	[?]	Ab initio
BeH	3	5000	2	592 308	Darby-Lewis	[406]	ExoMol
SiH ₂	1	2000	1	254 061 207	CATS	[?]	ExoMol
CN	4	10000	3	2 285 103	Trihybrid	✓ [486]	ExoMol
ZrO	6	10000	10	47 662 773	ZorrO	✓ [490]	ExoMol
NH	4	10000	5	327 014	kNigHt	✓ [493]	ExoMol
HBO	2	3000	1	142 038 890	LQL	[498]	ExoMol
N ₂	1	10000	4	7 182 000	WCCRMT	[499]	Empirical
H ₂ O ^b	1	300	1	109 263	CKYKKY	[485]	Empirical

N_{iso} Number of isotopologues considered;

T_{max} Maximum temperature for which the line list is complete;

N_{elec} Number of electronic states considered;

N_{lines} Number of lines: value is for the main isotope.

✓ indicates line list that provide energy levels with individual uncertainties and contain lines suitable for use in high resolution studies.

^a There are separate line lists for cis and trans P₂H₂.

^b An electric quadrupole (E2) line list for H₂¹⁶O.

Table 2 gives an overview of recommended line lists available through the ExoMol website but not generated by the ExoMol project. As for Table 1, an indication is given as to whether the data as presented are appropriate for high resolution studies. For Table 2 this information may be less useful because while most of the sources do indeed give accurate predictions for the transition wavenumbers, the line lists do not contain uncertainties for their individual energy levels which means that within the ExoMol data structure wavenumber uncertainties cannot be

estimated. These datasets have not been marked as suitable for high resolution studies and it is necessary to check the original (cited) reference to establish the accuracy of the transitions.

A number of the line lists presented in Table 2 are taken from Bernath's MoLLIST project [319] including the one for the CP radical [473]. We note that an alternative line list for CP has been computed by Qin *et al.* [474]. A new MoLLIST line list covering the $B\ ^2\Sigma^+ - X\ ^2\Sigma^+$ band system of LaO by Bernath *et al.* [475] has been added. Compared to the 2020 release a number of MoLLIST line lists, namely OH, AlCl, CaH and NH, have been replaced by ones generated by ExoMol.

Diatomeric line lists for four molecules have also been taken from HITRAN [2], namely those for CO, HCl, HF and HBr. These line lists provide extensive compilations of rotation-vibration transitions within the electronic ground states and are appropriate for elevated temperatures. In the case of CO, a number of improvements to the treatment of the already very accurate ground state line list [476, 477] and extensions to consider rovibronic transitions [478] have been undertaken. These will be used to provide updated and greatly extended line lists.

HITRAN data have also been used to create opacities for the oxygen molecule; these data do not extend to high temperature. An extensive, ExoMol high-temperature line list for O₂ should be available soon. To this end, new methodology to treat magnetic dipole and electric quadrupole transitions in diatomic molecules has been developed and implemented in Duo [479, 480].

Molecular hydrogen, H₂, is one of the relatively few molecules for which the database provides more than one recommended line list. The weak, dipole-forbidden infrared spectrum of H₂ is given by the RACPPK line list [481] which includes electric quadrupole transitions, for which $\Delta J = 0, \pm 2$, and magnetic dipole transitions with $\Delta J = 0$ only. The dipole-allowed Lyman ($B\ ^1\Sigma_u^+ - X\ ^1\Sigma_g^+$) and Werner ($C\ ^1\Pi_u - X\ ^1\Sigma_g^+$) H₂ electronic band systems are provided by the ARLR [482] line list. An infrared line list of HD is provided from high accuracy *ab initio* calculations by Amaral *et al.* (ADSJAAM) [483]; in this case it is also necessary to consider weak dipole-allowed transitions which arise from failure of Born-Oppenheimer approximation. *Ab initio* infrared line lists for HD⁺ and HeH⁺ from Amaral *et al.* [483] are also included. *Ab initio* line lists for LiH⁺ and LiH, designed primarily for studies of the early Universe, due to Coppola *et al.* [484] are also included.

An electric quadrupole (E2) line list CKYKKY for H₂¹⁶O has been produced [485] using the POKAZATEL PES and a new *ab initio* electric quadrupole moment surface of H₂O. The line list was used to identify E2 transitions of H₂¹⁶O experimentally for the first time [485].

The UNSW (University of New South Wales) group of McKemmish have generated a number of ExoMol style diatomic line lists which are available and recommended in the database. All these line lists have been the subject of an associated MARVEL study and therefore provide high accuracy data and are suitable for high resolution studies. They are:

- CN: Syme and McKemmish [486] ExoMol style line list from UNSW calculated using Duo [487] hybridised with MARVEL [488] and MoLLIST [489] data.
- ZrO: Perri *et al.* [490] ExoMol style line list from UNSW calculated using Duo [487] hybridised with MARVEL [491] and MoLLIST [492] data.
- NH: Perri and McKemmish [493] ExoMol style line list from UNSW calculated using Duo [487] hybridised with MARVEL [494] and MoLLIST [495–497] data.

In addition Li *et al.* [498] have recently used ExoMol methodology to compute line lists for two isotopologues of HBO. This line list has been added to the database.

Finally the website provides partial data for two other diatomic molecules, N₂ and NiH.

Though the nitrogen molecule has a singlet ground state, at present data are only provided for transitions between excited triplet states; these transitions are prominent in nitrogen and air plasmas. The empirical dataset WCCRMT line list due to Western *et al.* [499] covering the 4 500 – 15 700 cm⁻¹ region is available. Recently, Jans [500] extended this work to consider the N₂ ($C\ ^3\Pi_u - B\ ^3\Pi_g$) second positive system, providing the data in ExoMol format. It is planned to merge these two datasets to form a single N₂ triplet dataset.

The website currently provides a link to a list of observed nickel monohydride (NiH) transitions recorded by Vallon *et al.* [501] and Harker *et al.* [502]. These datasets which consider the isotopologues ⁵⁸NiH, ⁶⁰NiH and ⁶²NiH as well as the effects of magnetic fields, are not presented in ExoMol format. However, Havalyova [503] has constructed a high accuracy, empirical spectroscopic model for NiH which is currently being used as the starting point for constructing an ExoMol line list for this system.

2.7. Atoms

Thus far the ExoMol database has concentrated on molecular spectra. However, atomic spectra are also important for exoplanet studies. So far the website contains data on two neutral atoms:

- Potassium: Kurucz-Allard dataset for ^{39}K [321, 504]
- Sodium: Kurucz-Allard dataset for ^{23}Na [321, 505]

These data are given only in the form of opacities [299]. We note that Allard *et al.* [506] have recently updated the opacities for broadening of K by He.

2.8. Partition functions

Partition functions are provided for all molecular line lists in two column format of temperature in K and partition function. The data are provided in 1 K steps up to the temperature maximum specified by the line list and given in the .def file. ExoMol follows the HITRAN convention [507] and includes the full nuclear spin degeneracy contribution in the partition function unlike many astronomy oriented compilations of partition functions [508–510] which use a reduced nuclear spin factor. Our convention makes the treatment of hyperfine resolved spectra, as done in the HyVO VO line list [367], straightforward.

It is possible to compute partition functions by direct summation of the levels provided in the .states file; both ExoCross [386] and PyExoCross [387] offer this service. However, we recommend using the partition functions provided in the .pf file as part of the line list, as for some molecules these represent more reliable values than those will be obtained by direct summation [511].

3. Data provided

Table 3 provides a summary of the various datasets available for each isotopologue on the ExoMol website. These data can be either downloaded manually or accessed through the API (applications program interface) described in Sec. 3.2 below.

The following subsections describe some of these datasets, again concentrating mainly on changes since the 2020 release.

Table 3: Summary of data provided by the ExoMol Database.

Data type
Line lists
Absorption cross sections
Pressure broadening coefficients
Temperature dependent super-lines (histograms)
Partition functions
Cooling functions
Specific heat - heat capacity
Temperature and pressure dependent opacities
Photo-absorption continuum cross section*
Photo-dissociation cross sections including VUV absorption*
Spectroscopic Models

* Denotes a new or updated data type, see text for details.

3.1. ExoMol format

Table 4 gives an overview of the ExoMol file structure. Note that ExoCross/pyExocross generates other file types such as .os for oscillator strengths but these files are not distributed via the ExoMol website/database.

The core of the ExoMol data structure is the provision of a .states and .trans file [512] which give, respectively, the energy levels with associated quantum numbers and Einstein A coefficients identified by upper and lower state indices (counting numbers) which point to the .states file. The format of the .trans file is unchanged

Table 4: Specification of the ExoMol file types. (Contents in brackets are optional.)

File extension	N_{files}	File DSname	Contents
.all	1	Master	Single file defining contents of the ExoMol database.
.def	N_{tot}	Definition	Defines contents of other files for each isotopologue.
.states	N_{tot}	States	Energy levels, quantum numbers, Uncertainties, lifetimes, (Landé g -factors).
.trans	a	Transitions	Einstein A coefficients, (wavenumber).
.broad	N_{mol}	Broadening	Parameters for pressure-dependent line profiles.
.cross	b	Cross sections	Temperature or temperature and pressure-dependent cross sections.
.kcoef	c	k -coefficients	Temperature and pressure-dependent k -coefficients.
.pf	N_{tot}	Partition function	Temperature-dependent partition function.
.cf	N_{tot}	Cooling function	Temperature-dependent cooling function.
.super	d	Super-lines	Temperature dependent super-lines (histograms) on a wavenumber grid.
.nm	e	VUV cross sections	Temperature and pressure dependent VUV cross-sections (wavelength, nm).
.fits, .h5, .kta	f	Opacities	Temperature and pressure dependent opacities for radiative-transfer applications.
.overview	N_{mol}	Overview	Overview of datasets available.
.readme	N_{iso}	Readme	Specifies data formats.
.model	N_{iso}	Model	Model specification.

N_{files} total number of possible files;

N_{mol} Number of molecules in the database;

N_{tot} is the sum of N_{iso} for the N_{mol} molecules in the database;

N_{iso} Number of isotopologues considered for the given molecule.

^a There are N_{tot} sets of .trans files but for molecules with large numbers of transitions the .trans files are subdivided into wavenumber regions.

^b There are N_{cross} sets of .cross files for each isotopologue.

^c There are N_{kcoef} sets of .kcoef files for each isotopologue.

^d There are N_T sets of T -dependent super-lines.

^e There are N_{VUV} sets of VUV cross sections.

^f Set of opacity files in the format native to specific radiative-transfer programs.

Table 5: Transitions file specification.

Field	Fortran Format	C Format	Description
i	I12	%12d	Upper state ID
f	I12	%12d	Lower state ID
A	ES10.4	%10.4e	Einstein A coefficient in s^{-1}
$\tilde{\nu}_{fi}$	E15.6	%15.6e	Transition wavenumber in cm^{-1} (optional).

Fortran format: (I12,1x,I12,1x,ES10.4,1x,ES15.6).

from previous releases while there have been some minor changes to data structure of the .states. Tables 5 and 6 present the structure of these two files. The states and, particularly, the .trans files can be very large so both files are supplied compressed using .bz2 format. For larger line lists the transitions are provided as a series of files which store data for a wavenumber region specified in the file name. Wavenumbers are only provided in the .trans file of smaller line list and even in these cases it is recommended to recompute them using the energies in the .states file as they may not reflect improvements to the states file through various corrections.

A sample extract from a .states file is given in Table 7. The .states file gives the state ID, which should correspond to the line number, and is used by the .trans file to identify initial and final states. The state energy, degeneracy and total angular momentum are then specified. For most species the total angular momentum corresponds to the quantum number J but for hyperfine resolved line lists this is F where $F = |J+I|$ and I is the nuclear spin. The uncertainty, introduced in the 2020 release, is now a compulsory element of recommended ExoMol line lists, although it is still not available for older line lists and most non-ExoMol ones. The uncertainty provides important information for high resolution studies and can also help to inform future experimental priorities. The state lifetime, τ , is also now compulsory for recommended ExoMol line lists; its definition is now generalised so that τ gives the lifetime obtained by considering both radiative decay [513] and predissociation effects, see Tennyson *et al.* [21].

The optional fields in the .states file are defined in the .def discussed below. These fields generally comprise a set of quantum numbers which characterize the state. These always include the total symmetry of the state, either

Table 6: Specification of the .states file including extra data options; the formats at the end of the table are for the compulsory section only.

Field	Fortran Format	C Format	Description
i	I12	%12d	State ID
\tilde{E}	F12.6	%12.6f	Recommended state energy in cm^{-1}
g_{tot}	I6	%6d	State degeneracy
J	I7/F7.1	%7d/%7.1f	Total angular momentum quantum number, J or F (integer/half-integer)
Unc	F12.6	%12.6f	Uncertainty in the state energy in cm^{-1}
τ	ES12.4	%12.4e	State lifetime (aggregated radiative and predissociative lifetimes) in s
(g)	F10.6	%10.6f	Landé g -factor (optional)
(QN)	See .def file		State quantum numbers, may be several columns (optional)
(Abbr)	A2	%2s	Abbreviation giving source of state energy, see Table 8.
(\tilde{E}_0)	F12.6	%12.6f	Calculated state energy in cm^{-1} (optional)

Fortran format, J integer: (I12,1x,F12.6,1x,I6,I7,1x,F12.6,1x,ES12.4,1x,F10.6)

or J half-integer: (I12,1x,F12.6,1x,I6,F7.1,1x,F12.6,1x,ES12.4,1x,F10.6).

in the form of a total parity (denoted $+/-$), a rotationless parity (denoted e/f) or an irreducible representation of the molecular symmetry group the molecule is associated with. The latter is more typical for polyatomic molecules with complex symmetry groups. Other quantum numbers are always approximate; for many polyatomic molecules, two or even three sets of quantum numbers are available. For those line lists denoted as being high resolution in Table 1 two extra columns have been added to the .states file. High resolution line lists have energy levels that have been updated either using MARVEL energies (denoted Ma) or energies from some other source with very high accuracy/ low uncertainties. The penultimate column of the line list gives the original calculated energy level; of course if the state energy remains the calculated one (denoted Ca) then these two levels will be the same. The final auxiliary column gives the source of this updated energy level, see Table 8; the various means of improving on the original energy levels are discussed by McKemmish *et al.* [16].

3.2. API, versioning, master and def files

The ExoMol database has an application programming interface (API) which allows users to extract data computer to computer without the need to manually download datasets from the website. Details of how to use the API are given in the ExoMol2020 release [5].

It is ExoMol practice to give each line list a name, denoted <DATASET> below. Alongside the name, each line list has a version number given by the version date in YYYYMMDD format. MARVELization or other energy level updates to the .states file just lead to a new version of the line list. Conversely, a newly calculated line list, and hence a new .trans file, is always given a new name. Both old versions and old (retired) line lists are retained in the database to allow users to compare with previous results. The most recent version is stored without a version number (date); all previous versions are also retained and have the version in YYYYMMDD format added to their name.

To facilitate the use of the API and to provide complete metadata, the database contains a .master file, which is located at www.exomol.com/exomol.all. The .master file points to a .def for each recommended line list. Note that there are .def and _def.json files for each isotopologue for which ExoMol provides a line list. The .master file itself has a version number in YYYYMMDD format; this date corresponds the latest update of a .def. The version dates of the individual line lists are given in the .master file.

JSON (JavaScript Object Notation) [516] is a lightweight human-readable, data-interchange format. ExoMol database newly provides JSON format .master (exomol.json) and definition (<ISOTOPOLYQUE>_<DATASET>_def.json) files. In the appendix, Listing 1 gives an extract from the current, JSON format .master file (exomol.json) with keywords defined in Table 9 and Listing 2 shows the JSON format definition file (_def.json) of the POKAZATEL H_2^{16}O line list. Table 10 gives the JSON keyword structure for the .def file.

Table 7: Extract from the state file for XAB line list for $^{24}\text{Ca}^1\text{H}$ [360].

<i>i</i>	\tilde{E}	<i>g</i>	<i>J</i>	unc	τ	<i>g</i>	+/-	<i>e/f</i>	State	<i>v</i>	Λ	Σ	Ω	Abbr	\tilde{E}_0
33	24428.044063	4	0.5	0.007010	1.7190E-07	2.002233	+	e	B'(2SIGMA+)	3	0	0.5	0.5	Ma	24428.027169
34	24986.629171	4	0.5	2.507500	5.4398E-08	-0.000637	+	e	A(2PI)	4	1	-0.5	0.5	Ca	24986.629171
35	25163.713092	4	0.5	0.022010	2.0005E-07	2.002233	+	e	B'(2SIGMA+)	4	0	0.5	0.5	Ma	25163.694359
36	25876.111435	4	0.5	0.101011	3.1000E-07	2.002217	+	e	B'(2SIGMA+)	5	0	0.5	0.5	Ma	25876.060619
37	26224.114885	4	0.5	3.007500	5.8534E-08	-0.000612	+	e	A(2PI)	5	1	-0.5	0.5	Ca	26224.114885

i: State counting number.

\tilde{E} : State energy in cm^{-1} .

g: State degeneracy.

J: Total angular momentum.

unc: Energy uncertainty in cm^{-1} .

τ : Lifetime in s.

g: Landé *g*-factor.

+/-: Total parity.

e/f: rotationless-parity.

State: state term value using PyValem format [514, 515].

v: State vibrational quantum number.

Λ : Projection of the electronic angular momentum.

Σ : Projection of the electronic spin.

Ω : $\Omega = \Lambda + \Sigma$, projection of the total angular momentum.

Abbr: Abbreviation giving source of state energy, see Table 8.

\tilde{E}_0 : Original (calculated) state energy in cm^{-1} .

Table 8: Source type abbreviations used to describe energy levels in hybrid (MARVELized) line lists; see McKemmish *et al.* [16] for further details.

Abbr	Meaning
Ca	Calculated
Ma	MARVEL
EH	Effective Hamiltonian
Mo	MOLLIST
HI	HITRAN
PS	Predicted Shift
IE	Isotopologue Extrapolation

Table 9: Keywords used in the JSON format .master (exomol.json) file.

Keywords	Meaning
ID	Always the string "EXOMOL.master"
version	Version number with format YYYYMMDD
num_molecules	Number of molecules in the database
num_isotopologues	Number of isotopologues in the database
num_datasets	Number of datasets in the database
num_species	Number of species in the database
molecules	List of molecules
formula*	Molecule chemical formula
num_molecule_names	Number of molecule names listed
molecule_names	List of the molecule names
num_isotopologues	Number of isotopologues considered
linelist	Line list
inchikey	InChIKey of isotopologue
iso_slug	Iso-slug
iso_formula	Formula of isotopologue
dataset_name	Isotopologue dataset name
version	Version number with format YYYYMMDD

*: The keywords with * are variable.

Table 10: Keywords use in the JSON format definition (<ISOTOPOLOGUE>_<DATASET>.json) file.

Keywords	Meaning
isotopologue	Isotopologue section
iso_formula	Formula of isotopologue
iso_slug	Iso-slug
inchikey	InChIKey of isotopologue
inchi	InChI of isotopologue
mass_in_Da	Mass of isotopologue (sum of atom masses) in Da
point_group	Symmetry group
atoms	Atoms section
number_of_atoms	Number of atoms
element	Element symbol
atom*	In format: Element symbol : Isotope number
irreducible_representations	Irreducible representation section
label*	In format: Irreducible representation label : Nuclear spin degeneracy
dataset	Dataset line list files section
name	Isotopologue dataset name
version	Version number with format YYYYMMDD
doi	DOI of dataset paper
max_temperature	Maximum temperature of linelist
num_pressure_broadeners	Number of pressure broadeners available
nxsec_files	Number of cross section files available
nkcoeff_files	Number of <i>k</i> -coefficient files available
dipole_available	Dipole availability (true=yes, false=no, null=undefined)
cooling_function_available	Cooling function availability (true=yes, false=no, null=undefined)
specific_heat_available	Specific heat availability (true=yes, false=no, null=undefined)
continuum	Photo-absorption continuum cross sections availability (true=yes, false=no, null=undefined)
predis	Photo-dissociation cross sections availability (true=yes, false=no, null=undefined)
states	States file part
number_of_states	Number of states in .states file
max_energy	Maximum energy (in cm ⁻¹) in the states file
uncertainty_description	Uncertainty description
uncertainties_available	Uncertainty availability (true=yes, false=no, null=undefined)
lifetime_available	Lifetime availability (true=yes, false=no, null=undefined)
lande_g_available	Lande g-factor availability (true=yes, false=no, null=undefined)
num_quanta	Number of quanta defined
states_file_fields	Columns list in the states file
name	Column name in .states file. There are three formats: 1. Main column name; 2. Quantum case label : Quantum label; 3. Auxiliary : Auxiliary title
desc	Description of column in the states file
ffmt	Fortran format
cfmt	C format
transitions	Transitions file(s) part
number_of_transitions	Total number of transitions in .trans file(s)
number_of_transition_files	Number of .trans file(s)
max_wavenumber	Maximum wavenumber (in cm ⁻¹)
partition_function	Partition function file section
max_partition_function_temperature	Maximum temperature of partition function
partition_function_step_size	Step size of temperature
broad	Broadening files section
default_Lorentzian_half-width	Default value of Lorentzian half-width for all lines (in cm ⁻¹ /bar)
default_temperature_exponent	Default value of temperature exponent for all lines
label*	A label for a particular broadener
filename	Filename of particular broadened
max_J	Maximum <i>J</i> for which pressure broadening parameters provided
Lorentzian_half_width	Value of Lorentzian half-width for <i>J</i> "> <i>J</i> _{max}
temperature_exponent	Value of temperature exponent for lines with <i>J</i> "> <i>J</i> _{max}
num_quantum_number_sets	Number of defined quantum number sets
quantum_number_sets	Defined quantum number sets
code	A code that defines this set of quantum numbers
num_lines	Number of lines in the broad that contain this code
num_quantum_numbers	Number of quantum numbers defined
quantum_numbers	Defined quantum number(s)

*: The keywords with * are variable.

3.3. Cross sections and continuum absorption

Table 11 gives the general ExoMol format for cross sections (.cross) files. Cross sections for line lists are supplied [323] to allow users to get an overall picture of the line lists contained in the database. These are provided for different temperatures but at zero pressure (*i.e.* there is no allowance for pressure broadening) via the ExoMol cross section app allowing to select the temperature, wavenumber range and the resolution (in cm^{-1}).

The extension of line lists into the UV means that molecules often have a continuum contribution to their spectrum alongside their characteristic, sharp line absorption. This temperature-dependent continuum component of the absorption is provided as a set of .cont files given at 100 K intervals from 100 K to the stated T_{\max} for the given line list in same format as the .cross file. The continuum functions were generated neglecting effects due to pressure broadening which are unlikely to be significant for continuum absorptions. The methodology has been introduced in Tennyson *et al.* [21]. .cont files are currently available for AlH and AlD [19], and OH [373].

Table 11: Specification of the .cross cross section and .cont file format

Field	Fortran Format	C Format	Description
$\tilde{\nu}_i$	F12.6	%12.6f	Central bin wavenumber, cm^{-1}
σ_i	ES14.8	%14.8e	Absorption cross section, $\text{cm}^2 \text{ molec}^{-1}$

Fortran format: (F12.6,1x,ES14.8)

3.4. Specific heats and cooling functions

The specific heat capacity at constant pressure (isobaric) C_p is supplied on a 1 K grid from $T = 0$ K to T_{\max} in units of $\text{J mol}^{-1} \text{ K}^{-1}$, where available. Otherwise, it can be computed from the available partition function by [517]

$$C_p(T) = R \left[\frac{Q''}{Q} - \left(\frac{Q'}{Q} \right)^2 \right] + \frac{3R}{2}, \quad (2)$$

where the second term is the translational contribution, and

$$Q'(T) = T \frac{dQ}{dT}, \quad (3)$$

$$Q''(T) = T^2 \frac{d^2Q}{dT^2} + 2Q', \quad (4)$$

and R is the gas constant. A complete set of specific heats was generated for the main (parent) isotopologue for each recommended ExoMol line list as part of the project to provide these data in standard (NASA) polynomial form [297].

Cooling functions can be important for astronomical studies [404, 484, 518, 519] and can be computed from ExoMol line lists. The temperature-dependent cooling function, $W(T)$ (units $\text{erg} (\text{s sr molecule})^{-1} = 10^{-7}$ Watts ($\text{molecule sr})^{-1}$) is the total energy per unit solid angle emitted by a molecule:

$$W(T) = \frac{1}{4\pi Q(T)} \sum_{f,i} A_{fi} h c \tilde{\nu}_{fi} g'_f e^{-c_2 \tilde{E}'_f / T}. \quad (5)$$

where Q is the partition function, A_{fi} the relevant Einstein-A coefficient, $\tilde{\nu}_{fi}$ is the transition wavenumber, g'_f is the degeneracy of the upper state and \tilde{E}'_f is the energy of the upper state. Finally, c_2 is the second radiation coefficient which allows for an energy level given in terms of wavenumbers. This can be done using either ExoCross [386] or PyExoCross [387]. At present the ExoMol website provides a small number of precomputed cooling functions.

3.5. ExoMolOP: Opacities

Chubb *et al.* [299] computed opacity cross sections for all molecules available from the ExoMol database plus the oxygen molecule, and atoms Na and K (using line position data from Kurucz [321] and data for the broadening of the strong doublets from Allard *et al.* [504, 505]) for a grid of temperatures and pressures and broadening regime appropriate for hot Jupiters, *i.e.* broadening by H₂ and He. The data are provided in formats appropriate for direct use in exoplanetary retrieval and modelling codes ARCiS [222], which uses the well established .fits format [520], petitRADTRANS [521, 522], NEMESIS [523] and Tau-REx [216]. The data format and gridding is

code dependent with opacities for ARCiS, petitRADTRANS, and NEMESIS given as k -tables at a resolving power of $R = \frac{\lambda}{\Delta\lambda} = 1000$ while cross-sections are provided for Tau-REx with $R = 15\,000$. The work of Chubb et al. has been updated for the newly created ExoMol line lists listed in Table 1. At present these opacities are typically only for the main (parent) isotopologue for each molecule (with some exceptions, for example CO is available with isotopologues mixed at terrestrial natural abundances).

The plan for the future will be to provide a broader range of opacities which (a) allow for mixtures of isotopologues and (b) using our newly derived broadening parameters (see below) to provide opacities for a variety of atmospheric compositions.

4. Collisional line broadening

The generation of opacities requires knowledge of line shape parameters, with collisional line broadening being a key process [524] in determining the frequency distribution of a line's intensity. In the course of previous works in the group, an ExoMol diet was introduced [525] which provided parameters characterising the collisional line broadening required for the Voigt profile. The reference collisional line width γ_0 (in $\text{cm}^{-1}\text{atm}^{-1}$) and its temperature exponent n are assumed to obey the following power law:

$$\gamma(T, P) = \gamma_0(T_{\text{ref}}, P_{\text{ref}}) \left(\frac{T_{\text{ref}}}{T} \right)^n \frac{P}{P_{\text{ref}}}, \quad (6)$$

where T_{ref} , P_{ref} are some reference temperature and pressure, which in ExoMol are chosen as 1 atm and 296 K, respectively. The ExoMol diet tabulates Lorentzian half-width, γ_0 , and its temperature dependence, n , for each perturbing species, and the final half-width is calculated as a sum of contributions from all the perturbers. The ExoMol diet is organised in the form of 'recipes', tabulating the dependence of γ_0 and n on the rotational and vibrational quantum numbers.

The pressure broadening .broad file structures used by the ExoMol diet are illustrated in Table 12. For each radiator-perturber pair, the diet files contain one or more recipes indicated by a unique name in the first column (a0, a1, etc.). This name is followed by collisional broadening parameters, for instance $\gamma_0(T_{\text{ref}}, P_{\text{ref}})$ and n from Eq. (6), and by a set of quantum numbers used in the recipe's parametrization scheme. For example, J'' is used in a0, while a1 additionally uses K'' . Note that a file might have a varying number of columns if different recipes require it. The naming convention for all pressure broadening files is to give broadening data for the main isotopologue, however pressure broadening data are not isotope specific in the current iteration of the database. If a user wishes to include pressure broadening for minor isotopologues, then they should use the data provided for the major isotopologue.

Table 12: Examples of the .broad ExoMol diet files for the 'a0', 'a1' and 'm0' recipes.

Label	γ	n	J''/m	K''
24Mg-16O_O2.broad				
a0	0.043	0.500	0	
...				
1H2-16O_H2.broad				
a1	0.0916	0.790	0	1
a1	0.0852	0.608	1	2
a1	0.0764	0.541	2	3
a1	0.0699	0.502	3	4
a1	0.0635	0.458	4	5
...				
27Al-1H_H2.broad				
m0	0.1554	0.6406	1	
m0	0.1489	0.6235	2	
m0	0.1450	0.6128	3	
m0	0.1410	0.6002	4	
...				

Table 13 list the current menu of ExoMol diets. As a new diet option, the spectroscopic index $|m|$ has been introduced to better describe the rotational dependence of collisional parameters γ_0 and n , where m is defined as

$m = -J''$ for P-branch transitions, $m = J'' + 1$ for R-branch, and $m = J''$ for Q-branch (i.e. the same as done in HITRAN). The “m0” option (see Table 13) has been used to define γ_L for AlH as part of the recent AloHa line list [19] (labeled in the diet .broad file as m0); the “m1” recipe was introduced to describe the dependence on the lower state K'' quantum number for symmetric top molecules in addition to the dependence on $|m|$ [372].

Table 13: Parameters defined in the ExoMol broadening diets.

diet code	quantum numbers	systems
a0	J''	general use (e.g. [525])
a1	J'', K''	general use (e.g. [525])
m0	$ m $	general use (e.g. [19])
m1	$ m , K''$	general use (e.g. [372])
a5	$J', K'_a, K'_c, J'', K''_a, K''_c$	asymmetric tops (e.g. SO ₂ [525])

The ExoMol database contains empirical broadening data for seven species: H₂O, NH₃, SO₂, CH₄, PH₃, HCN, and H₂CO broadened by H₂ and He. In addition, for most of the other ExoMol molecules, especially exotic ones, the ExoMol diet now provides theoretical line broadening parameters computed using at least one of three methods, simple semi-classical calculations, explicit semi-classical Modified Complex Robert-Bonamy (MCRB) approach and Machine Learning described below.

The MCRB semi-classical approach [526] is capable of producing broadening parameters accurate to within experimental uncertainty if an accurate interaction potential between the radiating and perturbing molecule is available. If only rough model potential data are available, fitting of the interaction potential to experimental line broadening coefficients can be used to improve the description for a range of transitions and temperatures. Calculations of vibrational contributions to broadening coefficients additionally require the knowledge of ro-vibrational state-dependent molecular quantities for the radiating molecule (dipole moment, polarizability). Broadening calculations have been done for AlH broadened by H₂, He, N₂, and AlH [19], and work on C₂H₂ broadened by H₂, He, N₂, C₂H₂, and CO₂ is currently under way [527]. In these cases, the vibrational contributions have been found to be negligible, so the respective .broad files only contain the dependence on the rotational quantum number J . Future work includes producing $J(m)$ -dependent broadening parameters for a number of diatomic molecules from the ExoMol database broadened by various perturbers using model interaction potentials, where the calculations are not too demanding.

A novel Machine Learning (ML) method has been developed [528] to predict the line broadening parameters γ_0 . This ML approach is suitable for the cheap mass production of pressure broadening parameters for any active neutral molecule; the ML code and associated data available on the ExoMol Zenodo area [529]. The algorithm considers the rotational dependence of the broadening but ignores any dependence on vibrational and/or electronic states.

Using this ML code, J -dependent air-broadening parameters γ_{air} have been produced for all molecules in the ExoMol database, suitable for room temperature and pressure environments, where air is the primary source of broadening. The empirical γ_0^{air} values for training were taken from the HITRAN database [2]. These data are available on the ExoMol website, where the temperature exponent value of $n = 0.5$ is assumed. Further work using machine learning to predict γ_0 for other perturbing species is ongoing. The molecules for which J -dependent air-broadening data are available is summarised in Table 14. High quality data have simply been copied from the HITRAN database [2] while other values are machine learned predictions [528].

The need for at least approximate values of pressure-broadened rovibrational line widths for “exotic” molecular pairs relevant to exoplanetary atmospheres initiated a theoretical study [530] giving access to simple semi-classical (rotationally-independent) estimates from the knowledge of molecular masses and kinetic diameters. Being valid for neutral active molecules but also for spectroscopically active molecular ions, this method allowed population of the ExoMol database with estimates of pressure-broadening coefficients for perturbation by H₂, O₂, N₂, He, Ar, CO, CS, NO, CO₂, H₂O, CH₄, NH₃ and self-perturbation for 45 active molecules (main species) [531] in ExoMol, see Table 15; work is in progress on producing similar data for their most abundant isotopologues. Available on the ExoMol website are the reference-temperature (296 K) predictions issued from data considered as the most reliable/recent kinetic diameter values. Results for CH₂, FeO, PO₂, SH₃, SiC, SiH₃ and TiF are not currently available on the ExoMol website, as there are no corresponding line lists available for these molecules. Results for temperatures above 296 K are easily obtained by a simple scaling factor $\sqrt{296/T}$ but the temperature considered should

Table 14: Availability of new air broadening data, γ_{air} . The molecules with ML data were generated with [528]. Other air broadening data are taken from the HITRAN database [2].

ML data					HITRAN data	
MgH	NaH	NiH	AlH	CrH	HCl	CH ₃ Cl
CaH	BeH	TiH	FeH	LiH	HF	C ₂ H ₄
ScH	NH	CH	OH	SiH	HBr	C ₂ H ₂
SH	PH	VO	ALO	YO	CO	PH ₃
MgO	TiO	SiO	CaO	NaO	NO	H ₂ O
LaO	ZrO	SO	PO	PN	O ₂	CO ₂
KCl	NaCl	LiCl	CN	C ₂	CS	SO ₂
H ₂	CP	PS	NS	SiS	N ₂	HCN
NaF	AlCl	AlF	KF	LiF	CH ₄	N ₂ O
CaF	MgF	SiN	H ₂ O ₂	SO ₃	NH ₃	H ₂ S
SiH ₄	AsH ₃	PF ₃	CH ₃	P ₂ H ₂	HNO ₃	OCS
H ₂ CS	CaOH	KOH	NaOH	SiH ₂	H ₂ CO	
SiO ₂	LiOH				CH ₃ F	

Table 15: New (rotationally-independent) line broadening data γ_0 in the ExoMol database generated using the semi-classical method by Buldyreva *et al.* [530].

Active molecules				Perturbers
AlCl	CrH	NaCl	SH	self
AlH	CS	NaF	SiH	Ar
AlO	KCl	NaH	SiH ₂	CH ₄
AsH ₃	KF	NaOH	SiO	CO
BeH	KOH	NH	SiO ₂	CO ₂
C ₃	LiCl	NS	SiS	H ₂
CaF	LiF	PF ₃	TiH	H ₂ O
CaH	LiH	PH	TiO	He
CaO	LiH ⁺	PN	VO	N ₂
CaOH	MgF	PO		NH ₃
CH ₃	MgH	PS		NO
CP	MgO	ScH		O ₂

not be too low as classical-path approximation has to remain valid (i.e. the kinetic energy of the relative molecular motion should be higher than the isotropic intermolecular potential depth), see the discussion by Buldyreva *et al.* [531].

Pressure-induced line-shape parameters have also been addressed for the case of vibronic transitions connecting low-lying vibrational levels in the ground and an excited (bound) electronic states [532]. In this case the collisional (pressure) broadening is still not dominated by predissociative effects. Pre-computed potential-energy surfaces for interactions of some representative molecular pairs enabled both line-width and line-shift calculations comparing favourably with available measurements from room temperature to 2800 K [532]. Further attempts at improvements are in progress, in order to establish a reliable computational scheme for providing line widths and shifts for rovibronic transitions of active molecular species listed in ExoMol.

In instances where a new set of broadening data is introduced for a system with already available data, a version number is added to the filename of the older data set in order to distinguish multiple versions of the same system. The version number is a timestamp indicating the date the dataset was created (uploaded to the data base) in the YYYYMMDD format as described in Sect. 3.2. The most recent or recommended version does not contain a version number. Although, at present, there is only one molecule, AlH, with broadening parameters computed by more than one method, we expect this number to grow. Apart from the experimental sources as e.g. provided by HITRAN, line broadening parameters have being provided by different computational methods such as ML [528], MCRB [526], simple semi-classical [530, 531] as well phase shift [532] approaches.

5. Predissociative line Broadening

As introduced in Sect. 3, the definition of the lifetimes in ExoMol has been extended to include the predissociative lifetimes as part of the `.states` file provision. The practical purpose of this extension is to be able to model the predissociative broadening together with the collisional and Doppler broadening. The lifetime broadening is described by Lorentzian line shape like pressure broadening with γ_τ , the half-width in cm^{-1} , given by

$$\gamma_\tau = \frac{\hbar}{2\tau hc}, \quad (7)$$

where τ is the half-harmonic mean of the radiative τ_r and predissociative τ_p lifetimes via

$$\tau = \frac{\tau_r \tau_p}{\tau_r + \tau_p}. \quad (8)$$

The half-width γ_τ can be added to the pressure-broadening half-width, γ in Eq. (6), to give the total Lorentzian component of the line profile.

Predissociation due to rotational barrier effects has been studied for AlH, see sec. 2.2.3. Predissociative line broadening due to spin-orbit interactions with the continuum has also been studied for OH [20], see 2.2.13, using a modified stabilization method [18, 533–535]. Predissociation lifetimes will also soon be released for SH and SD using the spectroscopic model from current linelist [342, 401] with the addition of the dissociative states, $1^2\Sigma^-$, $1^4\Sigma^-$, and $1^4\Pi$.

6. Photodissociation and continuum absorption

The extension of the database to consider photodissociation means we define a new file type `.photo`, which provides temperature-dependent photodissociation cross sections. The format of the photodissociation cross section has been introduced in Tennyson *et al.* [21] and specified in Table 16. In contrast to other data in the database, `.photo` files give results as a function of wavelength in nm. The project is only just starting to generate photodissociation cross sections, and so far, calculated cross sections are available for HF and HCl [536]. A major expansion of this part of the database is planned.

The previous section in the database/on the website called photo-absorption continuum cross sections, which contained measured UV photoabsorption cross sections, had been moved into the photodissociation area. These data are presented in the same format as the computed photodissociation cross sections, but the temperature, pressure and broadening parameters are determined by the experimental conditions used and generally do not form a regular grid. Measured cross sections are available for H_2O , CO_2 , SO_2 , NH_3 , H_2CO , C_2H_4 and CO. These cross sections probably represent a good approximation to the photodissociation cross section for the given species [537]. Table 4 of the ExoMol2020 release [5] gives more details on these cross sections provided which are due to Venot *et al.* [537] and Fateev and co-workers (private communication, 2020).

Table 16: Specification of the wavelength-dependent `.photo` cross section file format

Field	Fortran Format	C Format	Description
λ_i	F6.2	%6.2f	Central bin length, nm
σ_i	ES14.8	%14.8e	Photodissociation cross section, $\text{cm}^2 \text{ molec}^{-1}$

Fortran format: (F6.2,1x,ES14.8)

7. Post-processing

ExoMol provides post-processing capabilities through the Fortran program ExoCross (GitHub: <https://github.com/ExoMol/ExoCross>) [386] and the Python program PyExoCross [387] (GitHub: <https://github.com/ExoMol/PyExoCross>). There are example jobs for both ExoCross and PyExoCross on their GitHub pages.

ExoCross has many functions such as generating pressure and temperature dependent cross sections, partition functions, specific heat, state-resolved radiative lifetimes, non-local thermodynamic equilibrium (non-LTE) spectra, electric dipole, electric quadrupole and magnetic dipole spectra. ExoCross can read data in both ExoMol and HITRAN [538] formats and output them in these formats as well as SPECTRA (<http://spectra.iao.ru/>) and

Phoenix formats [539]. Should data be required in HITRAN format, it is strongly recommended that the data are downloaded to a local computer in the much more compact ExoMol format and then processed using ExoCross.

As a Python adaptation of the post-processing program ExoCross, PyExoCross supports importing and exporting line lists in the ExoMol and HITRAN format [315] and can convert data format between ExoMol and HITRAN formats. PyExoCross provides functions for calculating partition functions, specific heats, radiative lifetimes, cooling functions, oscillator strengths, LTE and non-LTE absorption and emission stick spectra and cross sections. Users can extract low uncertainties, high intensities and specified quantum number labels and values by using uncertainty, threshold and quantum number filters when calculating stick spectra and cross sections. PyExoCross can also help users automatically download line lists files from the ExoMol database in batches. The user instruction manual is available on <https://pyexocross.readthedocs.io/>.

8. New web services

8.1. ExoMolHR

ExoMolHR (ExoMol High-Resolution) [298] is a new high-resolution molecular spectroscopic database based on the ExoMol line list database; it is available at www.exomol.com/exomolhr. The ExoMolHR database is focused on providing high accuracy line positions for high-resolution studies such as line identification or simulations of high-resolution spectra. Initially reduced line lists are constructed by scraping the ExoMol database for energy levels (and hence transitions) which are determined to high accuracy (uncertainty $\leq 0.01 \text{ cm}^{-1}$). After this all transitions which can be predicted at high resolution ($R > 100\,000$) are stored; these transitions provide the core of the database. For each transition, ExoMolHR provides the frequency, uncertainty, Einstein A -coefficient, intensity (at a user specified temperature), lower energy, total level degeneracy, angular momentum and quantum numbers for upper and lower states. Users can download datasets using the application programming interface (API) directly with the intensities computed at a reference temperature $T_{\text{ref}} = 296 \text{ K}$. Alternatively, they can use the interactive web interface where one can set parameters to filter the wavenumber range, extract low uncertainties and lines with strong intensities, and then store the results generated for downloading. The ExoMolHR database currently contains data for 55 isotopologues from 32 molecules, see Table 17; this number is growing quite rapidly as more MARVEL studies are completed.

8.2. LiDB

LiDB (Lifetimes Database) is a newly developed database of molecular vibrational and vibronic state radiative lifetimes [296], created to enable radiative effects to be properly captured in low-temperature plasma models. The main data output of LiDB is radiative lifetimes at vibrational and electronic state resolution. Partial lifetimes, which give information on the dominant decay channels in a molecule, are also provided. Datasets for 36 molecules are available, produced from the respective molecular line list in the ExoMol database. LiDB is freely available and hosted at www.exomol.com/lidb. Users can dynamically view molecular datasets or use a specially-designed API to perform data requests. LiDB is linked directly to the Quantemol DataBase (QDB) [540] which aims to provide comprehensive datasets for plasma modelling. LiDB will expand in the future with the addition of more molecules, important isotopologues, and neutral and singly-charged atomic species.

9. Future development

The ExoMol project will continue not only to provide high quality data on more molecules, but the type of data available will be expanded to maximise the scientific richness that can be extracted from observations possible with new and upcoming telescopes. We are also open to hosting high quality, relevant datasets (both computed and measured) obtained by other groups.

The start in computing photodissociation cross sections and rates, as well as the treatment of rovibronic spectra signals our intention to extend the database to cover processes which occur at ultraviolet (UV) wavelengths. This will mean developing methods of performing calculations involving many electronic states for molecules larger than diatomics.

The calculations on the Renner-Teller system of CaOH [541] were performed using the variational rovibronic nuclear motion code EVEREST [449]. We are currently using EVEREST to explore the calculation of photodissociation spectra of triatomic systems using HCN as a test case [542]. The plan will be to start to provide extended

Table 17: Contents of the ExoMolHR database [298]; lines are extracted from the ExoMol database on the basis that they have low uncertainties.

ID	Molecule	Isotopologue	Dataset	N_{states}	N_{files}	N_{trans}	N_{HRstates}	N_{HRlines}
1	AlCl	$^{27}\text{Al}^{35}\text{Cl}$	YNAT	65869	1	4722048	41	101
2	AlCl	$^{27}\text{Al}^{37}\text{Cl}$	YNAT	67507	1	5748704	41	121
3	AlH	$^{27}\text{Al}^1\text{H}$	AloHa	1364	1	29725	135	692
4	AlO	$^{26}\text{Al}^{16}\text{O}$	ATP	93350	1	4866540	4783	143197
5	AlO	$^{27}\text{Al}^{16}\text{O}$	ATP	94862	1	4945580	4980	149577
6	AlO	$^{27}\text{Al}^{17}\text{O}$	ATP	96350	1	5148996	4787	142905
7	AlO	$^{27}\text{Al}^{18}\text{O}$	ATP	98269	1	5365592	4799	142976
8	C ₂	$^{12}\text{C}_2$	8states	44189	1	6080920	8376	445682
9	C ₂ H ₂	$^{12}\text{C}_2^{12}\text{H}_2$	aCeTY	5160803	100	4347381911	8898	473850
10	CH ₄	$^{12}\text{C}^4\text{H}_4$	MM	9155208	121	50395644806	21021	7649736
11	CN	$^{12}\text{C}^{14}\text{N}$	Trihybrid	28004	1	2285103	4833	244808
12	CO ₂	$^{12}\text{C}^{16}\text{O}_2$	UCL-4000	3480477	20	2557549946	18881	2600218
13	CaH	$^{40}\text{Ca}^1\text{H}$	XAB	6825	1	293151	1165	12341
14	CaOH	$^{40}\text{Ca}^{16}\text{O}^1\text{H}$	OYT6	3187522	18	23384729495	1424	12984
15	H ₂ CO	$^{1\text{H}_2}^{12}\text{C}^{16}\text{O}$	AYTY	10297025	100	12688112669	4813	317729
16	H ₂ CS	$^{1\text{H}_2}^{12}\text{C}^{32}\text{S}$	MOTY	52292454	8	43561116660	3625	72218
17	H ₂ O	$^{1\text{H}_2}^{16}\text{O}$	POKAZATEL	810269	412	5745071340	14395	3520554
18	H ₂ S	$^{1\text{H}_2}^{32}\text{S}$	AYT2	220631	35	115032941	2061	63719
19	H ₃ O ⁺	$^{1\text{H}_3}^{16}\text{O}^+$	eXeL	1173114	100	2089331073	232	1785
20	H ₃ ⁺	$^{1\text{H}_2}\text{H}^+$	ST	33330	1	22164810	109	646
21	H ₃ ⁺	$^{1\text{H}_3}\text{H}^+$	MiZATeP	158721	1	127542657	994	13606
22	H ₃ ⁺	$^{2\text{H}_1}\text{H}^+$	MiZo	369500	32	2290235000	115	683
23	H ₃ ⁺	$^{2\text{H}_2}\text{H}^+$	MiZo	37410	21	36078183	115	225
24	LiOH	$^{6}\text{Li}^{16}\text{O}^1\text{H}$	OYT7	192412	5	294573305	255	840
25	LiOH	$^{7}\text{Li}^{16}\text{O}^1\text{H}$	OYT7	203762	5	331274717	240	749
26	MgH	$^{24}\text{Mg}^1\text{H}$	XAB	3148	1	88575	237	2462
27	MgH	$^{25}\text{Mg}^1\text{H}$	XAB	3156	1	88776	548	5850
28	MgH	$^{26}\text{Mg}^1\text{H}$	XAB	3160	1	88891	537	5339
29	N ₂ O	$^{14}\text{N}_2^{16}\text{O}$	TYM	1759068	21	1360351722	17018	3459640
30	NH	$^{14}\text{N}^1\text{H}$	kNigHt	4076	1	327014	1030	26131
31	NH	$^{14}\text{N}^2\text{H}$	kNigHt	7406	1	778105	118	943
32	NH	$^{15}\text{N}^1\text{H}$	kNigHt	4089	1	327877	118	943
33	NH	$^{15}\text{N}^2\text{H}$	kNigHt	7465	1	785940	118	943
34	NH ₃	$^{14}\text{N}^1\text{H}_3$	CoYuTe	5095730	200	16941637250	4720	412149
35	NO	$^{14}\text{N}^{16}\text{O}$	XABC	30811	1	4596666	3044	106711
36	OCS	$^{16}\text{O}^{12}\text{C}^{32}\text{S}$	OYT8	2399110	10	2527364150	5198	279273
37	PN	$^{31}\text{P}^{14}\text{N}$	PaiN	30327	1	1333445	32	44
38	SO	$^{32}\text{S}^{16}\text{O}$	SOLIS	84114	1	7086100	536	2501
39	SO ₂	$^{32}\text{S}^{16}\text{O}_2$	ExoAmes	3270271	80	1300000000	14924	1504495
40	SiN	$^{28}\text{Si}^{14}\text{N}$	SiNfull	131936	1	43646806	99	670
41	SiN	$^{28}\text{Si}^{15}\text{N}$	SiNfull	133460	1	44816182	56	464
42	SiN	$^{29}\text{Si}^{14}\text{N}$	SiNfull	132335	1	43946969	56	464
43	SiN	$^{30}\text{Si}^{14}\text{N}$	SiNfull	132706	1	44223730	56	464
44	SiO	$^{28}\text{Si}^{16}\text{O}$	SiOUVenIR	174250	1	91395763	911	8729
45	TiO	$^{48}\text{Ti}^{16}\text{O}$	Toto	301245	1	58983952	8725	499775
46	VO	$^{51}\text{V}^{16}\text{O}$	HyVO	3410598	90	58904173243	7043	635722
47	YO	$^{89}\text{Y}^{16}\text{O}$	BRYTS	173621	1	60678140	28	25
48	YO	$^{89}\text{Y}^{17}\text{O}$	BRYTS	182598	1	62448157	28	25
49	YO	$^{89}\text{Y}^{18}\text{O}$	BRYTS	182547	1	64164605	28	25
50	ZrO	$^{90}\text{Zr}^{16}\text{O}$	Zorro	227118	1	47662773	5313	145317
51	ZrO	$^{91}\text{Zr}^{16}\text{O}$	Zorro	227118	1	47748501	1058	5164
52	ZrO	$^{92}\text{Zr}^{16}\text{O}$	Zorro	227124	1	47830250	1058	5164
53	ZrO	$^{93}\text{Zr}^{16}\text{O}$	Zorro	227126	1	47928979	1058	5164
54	ZrO	$^{94}\text{Zr}^{16}\text{O}$	Zorro	227128	1	47994352	1058	5164
55	ZrO	$^{96}\text{Zr}^{16}\text{O}$	Zorro	227134	1	48136388	1058	5164

N_{states} : Number of states in states file;

N_{files} : Number of transitions files;

N_{trans} : Number of transitions in transitions file(s);

N_{HRstates} : Number of states with uncertainties $\leq 0.01 \text{ cm}^{-1}$;

N_{HRlines} : Number of lines in ExoMolHR dataset with $R > 100\,000$.

coverage of temperature-dependent photoabsorption cross sections. For photodissociation we have already identified a number of studies on diatomics which provide temperature-dependent cross sections [282, 543–552] which we hope to add to the database.

The methodology for generating atomic as opposed to molecular data is actually very different and thus done by different research groups with information dispersed. However, astronomically, both species are often present together; for example, there is now a significant number of neutral and singly charged atoms that have been observed in exoplanetary atmospheres [553–555]. To support exoplanet studies, therefore, users have requested that atomic data also be provided in the same ExoMol format to facilitate more convenient analysis. We have started to address this need with the inclusion of the opacities for Na and K in a H₂/He atmosphere, see Sect. 2.7. Further expansions to include atomic data are being actively pursued with NIST [556, 557] and/or the Kurucz database [321] as possible sources.

As discussed in Sect. 4 there is a real need for an extended set of broadening parameters. Clearly this is a major focus of on-going work. At present ExoMol opacities and others are largely all focused on hot Jupiter planets. An extended set of broadening parameters would allow the generation of opacities for other atmospheres such as lava planets and rocky planets. What we do have as a result of our machine learning study is a full set of air-broadening parameters. This means we are already in a position to generate a full set of opacities for a planet whose atmosphere is Earth-like, ie close to 80% N₂ and 20% O₂. The use of different methodologies to generate broadening parameters, as discussed in Sect. 4, means that it is possible to have more than one file containing broadening parameters for a given perturber and molecule combination. Generalization of the ExoMol data structure to allow for this possibility will form part of an upcoming paper on the revised ExoMol broadening diet.

Currently, the ExoMol molecular opacities are generated assuming the LTE conditions, where the temperature dependent populations of the molecular states obey the Boltzmann distribution. The non-LTE processes assume a departure from LTE thus making the ExoMol opacities non-applicable for modelling, e.g., the solution of the radiative transfer equation at non-LTE conditions. Some work on using our data for non-LTE studies has already been performed [558]. However, a more general solution will require a new cross section format capable of preserving the information on the (rotation-)vibrational state that is therefore needed to allow for the solution of the statistical equilibrium equation. This will be addressed in the near future.

Finally but importantly, a major driver in our choice of species and processes to study is requests from interested users. We are always happy to receive such requests although we do not guarantee rapid delivery of results.

10. Conclusions

Since the ExoMol2020 data release [5], 27 new molecules have been added to the database. For many others the coverage for the line lists has been extended into the ultra violet and/or the energy levels, and hence transition wavenumbers, have been made significantly more accurate. Indeed, ExoMol-style line lists have proven extremely effective in providing very high quality data for astronomers, especially when supported by MARVEL studies of available experimental data.

The period since the last release of the ExoMol database in 2020 has seen the launch of JWST, the most important space observatory in decades, and its impact on astronomy research particularly in exoplanet atmospheric characterisation has been immediate. The ExoMol line list data have proven extremely powerful for modelling JWST observations when the molecule was predicted, with no identified shortcomings in the ExoMol databases' line list modelling of known molecular spectral bands at different temperatures. Yet spectral bands of unknown molecular origin are being identified regularly. Each of these unidentified spectral bands promises to shed light on physics, chemistry, geology (or maybe even biology) that was unsuspected in planetary models for exoplanetary systems. The set of molecules for which spectral data are required is clearly larger than already available in ExoMol, yet shortlisting the set of likely molecular candidates is not straightforward [117], especially for unusual non-terrestrial exoplanets. Computationally generated approximate spectral data for a very large number of potential atmospheric molecules [117, 559] is likely to be useful in helping to prioritise the molecules for which production of high-quality line lists suitable for the ExoMol database is warranted. Improved chemical models of a diverse set of atmospheres capable of predicting abundance of minor components with strong absorption would also be extremely valuable. With the planned launch of the Ariel space mission [560, 561] in 2029 producing a rapid increase in the number of exoplanet transit spectra available, we expect a corresponding increase in uniden-

tified spectral features and thus the need to produce high-quality molecular spectral data on an increasing number of molecules [311].

Yet, despite the launch of JWST, it is the ground-based observations that have generated the most substantial changes in the ExoMol database over the last four years as the high-resolution cross-correlation (HRCC) techniques has reached maturity. HRCC makes extremely high demands on the position accuracy of the strong spectral lines that are quite different from the high completeness required to model space observations. This difference has necessitated major changes in how line lists are created that had started in the 2020 ExoMol release with the introduction of MARVELized line lists but have become dominant in this 2024 release with the essential inclusion of energy level uncertainties within all line lists, major changes to line list construction methodology particularly the formalisation of the predicted shift and hybridisation approach, creation of the ExoMolHR web app, and even substantial updates to nuclear motion codes to enable modelling of hyperfine interactions.

ExoMol was originally designed to provide spectroscopic data at infrared and visible wavelengths. However, increasing demands to study the effects of ultraviolet radiation both in terms of photoabsorption and photodissociation has led to the scope of the database being extended to address these issues. Processes involving temperature-dependent photodissociation, line and continuum photoabsorption at UV wavelengths and predissociation are now being included in the database. These data are needed both to interpret observations and for chemical models of exoplanets. We are working on increasing our offering at UV wavelengths.

Data availability

The data discussed in this paper can be accessed at www.exomol.com. Programs associated with the ExoMol project including ExoCross and PyExoCross are available from <https://github.com/ExoMol/>.

Acknowledgments

This work was supported by the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme through Advance Grant numbers 267219 and 883830, and STFC Projects No. ST/I001050/1, ST/M001334/1 and ST/R000476/1, State Project IAP RAS No. FFUF-2024-0016

References

- [1] J. Tennyson, S. N. Yurchenko, ExoMol: molecular line lists for exoplanet and other atmospheres, *Mon. Not. Roy. Astron. Soc.* 425 (2012) 21–33. [doi:10.1111/j.1365-2966.2012.21440.x](https://doi.org/10.1111/j.1365-2966.2012.21440.x).
- [2] I. E. Gordon, L. S. Rothman, R. J. Hargreaves, R. Hashemi, E. V. Karlovets, F. M. Skinner, E. K. Conway, C. Hill, R. V. Kochanov, Y. Tan, P. Wcislo, A. A. Finenko, K. Nelson, P. F. Bernath, M. Birk, V. Boudon, A. Campargue, K. V. Chance, A. Coustenis, B. J. Drouin, J. Flaud, R. R. Gamache, J. T. Hodges, D. Jacquemart, E. J. Mlawer, A. V. Nikitin, V. I. Perevalov, M. Rotger, J. Tennyson, G. C. Toon, H. Tran, V. G. Tyuterev, E. M. Adkins, A. Baker, A. Barbe, E. Canè, A. G. Császár, A. Dudaryonok, O. Egorov, A. J. Fleisher, H. Fleurbaey, A. Foltynowicz, T. Furtenbacher, J. J. Harrison, J. Hartmann, V. Horneman, X. Huang, T. Karman, J. Karns, S. Kassi, I. Kleiner, V. Kofman, F. Kwabia-Tchana, N. N. Lavrentieva, T. J. Lee, D. A. Long, A. A. Lukashevskaya, O. M. Lyulin, V. Y. Makhnev, W. Matt, S. T. Massie, M. Melosso, S. N. Mikhailenko, D. Mondelain, H. S. P. Müller, O. V. Naumenko, A. Perrin, O. L. Polyansky, E. Raddaoui, P. L. Raston, Z. D. Reed, M. Rey, C. Richard, R. Tóbiás, I. Sadiek, D. W. Schwenke, E. Starikova, K. Sung, F. Tamassia, S. A. Tashkun, J. Vander Auwera, I. A. Vasilenko, A. A. Vigasin, G. L. Villanueva, B. Vispoel, G. Wagner, A. Yachmenev, S. N. Yurchenko, The HITRAN2020 molecular spectroscopic database, *J. Quant. Spectrosc. Radiat. Transf.* 277 (2022) 107949. [doi:10.1016/j.jqsrt.2021.107949](https://doi.org/10.1016/j.jqsrt.2021.107949).
- [3] J. Tennyson, L. Lodi, L. K. McKemmish, S. N. Yurchenko, The ab initio calculation of spectra of open shell diatomic molecules, *J. Phys. B-At. Mol. Opt. Phys.* 49 (2016) 102001. [doi:10.1088/0953-4075/49/10/102001](https://doi.org/10.1088/0953-4075/49/10/102001).
- [4] J. Tennyson, S. N. Yurchenko, A. F. Al-Refaie, E. J. Barton, K. L. Chubb, P. A. Coles, S. Diamantopoulou, M. N. Gorman, C. Hill, A. Z. Lam, L. Lodi, L. K. McKemmish, Y. Na, A. Owens, O. L. Polyansky, T. Rivlin, C. Sousa-Silva, D. S. Underwood, A. Yachmenev, E. Zak, The ExoMol database: molecular line lists for exoplanet and other hot atmospheres, *J. Mol. Spectrosc.* 327 (2016) 73–94. [doi:10.1016/j.jms.2016.05.002](https://doi.org/10.1016/j.jms.2016.05.002).
- [5] J. Tennyson, S. N. Yurchenko, A. F. Al-Refaie, V. H. J. Clark, K. L. Chubb, E. K. Conway, A. Dewan, M. N. Gorman, C. Hill, A. E. Lynas-Gray, T. Mellor, L. K. McKemmish, A. Owens, O. L. Polyansky, M. Semenov, W. Somogyi, G. Tinetti, A. Upadhyay, I. Waldmann, Y. Wang, S. Wright, O. P. Yurchenko, The 2020 release of the ExoMol database: Molecular line lists for exoplanet and other hot atmospheres, *J. Quant. Spectrosc. Radiat. Transf.* 255 (2020) 107228. [doi:10.1016/j.jqsrt.2020.107228](https://doi.org/10.1016/j.jqsrt.2020.107228).
- [6] J. Tennyson, S. N. Yurchenko, ExoMol at 10, *Astron. Geophys.* 62 (2021) 6.16–6.21. [doi:10.1093/astrogeo/atab102](https://doi.org/10.1093/astrogeo/atab102).
- [7] S. N. Yurchenko, J. Tennyson, J. Bailey, M. D. J. Hollis, G. Tinetti, Spectrum of hot methane in astronomical objects using a comprehensive computed line list, *Proc. Nat. Acad. Sci.* 111 (2014) 9379–9383. [doi:10.1073/pnas.1324219111](https://doi.org/10.1073/pnas.1324219111).
- [8] I. A. G. Snellen, A new method for probing the atmospheres of transiting exoplanets, *Mon. Not. Roy. Astron. Soc.* 353 (2004) L1–L6. [doi:10.1111/j.1365-2966.2004.08169.x](https://doi.org/10.1111/j.1365-2966.2004.08169.x).
- [9] I. Snellen, High-dispersion spectroscopy of extrasolar planets: from CO in hot Jupiters to O₂ in exo-Earths, *Phil. Trans. Royal Soc. London A* 372 (2014) 20130075. [doi:10.1098/rsta.2013.0075](https://doi.org/10.1098/rsta.2013.0075).

- [10] M. Brogi, J. Birkby, High-resolution Spectroscopy, in: ExoFrontiers, 2514-3433, IOP Publishing, 2021, pp. 8-1 to 8-10. [doi:10.1088/2514-3433/abfa8fch8](https://doi.org/10.1088/2514-3433/abfa8fch8).
- [11] H. J. Hoeijmakers, R. J. de Kok, I. A. G. Snellen, M. Brogi, J. L. Birkby, H. Schwarz, A search for TiO in the optical high-resolution transmission spectrum of HD 209458b: Hindrance due to inaccuracies in the line database, *Astron. Astrophys.* 575 (2015) A20. [doi:10.1051/0004-6361/201424794](https://doi.org/10.1051/0004-6361/201424794).
- [12] S. de Regt, A. Y. Kesseli, I. A. G. Snellen, S. R. Merritt, K. L. Chubb, A quantitative assessment of the VO line list: Inaccuracies hamper high-resolution VO detections in exoplanet atmospheres, *Astron. Astrophys.* 661 (2022) A109. [doi:10.1051/0004-6361/202142683](https://doi.org/10.1051/0004-6361/202142683).
- [13] S. N. Yurchenko, J. Tennyson, M. Brogi, High-resolution spectroscopy of exoplanets: data challenges and prospects, *Nature Rev. Phys.*(submitted).
- [14] J. Tennyson, The ExoMol project: Molecular opacity calculations at University College London, in: C. Mendoza, S. Turck-Chiéze, J. Colgan (Eds.), *Workshop on Astrophysical Opacities*, Vol. 515 of *Astron. Soc. Pac. Conf. Ser.*, Astronomical Society of the Pacific, 2018, pp. 137–144.
- [15] T. Furtenbacher, A. G. Császár, J. Tennyson, MARVEL: measured active rotational-vibrational energy levels, *J. Mol. Spectrosc.* 245 (2007) 115–125. [doi:10.1016/j.jms.2007.07.005](https://doi.org/10.1016/j.jms.2007.07.005).
- [16] L. K. McKemmish, C. A. Bowesman, K. Kefala, A. N. Perri, A. M. Syme, S. N. Yurchenko, J. Tennyson, A hybrid approach to generating diatomic line lists for high resolution studies of exoplanets and other hot astronomical objects: Updates to ExoMol MgO, VO and TiO line lists, *RAS Tech. Instr.*(Submitted).
- [17] M. Pezzella, S. N. Yurchenko, J. Tennyson, A method for calculating temperature-dependent photodissociation cross sections and rates, *Phys. Chem. Chem. Phys.* 23 (2021) 16390–16400. [doi:10.1039/D1CP02162A](https://doi.org/10.1039/D1CP02162A).
- [18] G. B. Mitev, R. P. Brady, O. Smola, J. Tennyson, S. N. Yurchenko, Calculation of spin-orbit induced predissociation lifetimes in rovibronic states of diatomic molecules, *Phys. Chem. Chem. Phys.* In preparation.
- [19] S. N. Yurchenko, W. Szajna, R. Hakalla, M. Semenov, A. Sokolov, J. Tennyson, Y. Pavlenko, M. R. Schmidt, ExoMol line lists – LIV. Empirical line lists for AIH and ALD and Emission spectroscopy of ALD in $A^1\Pi$ ($v = 0, 1, 2$), *Mon. Not. Roy. Astron. Soc.* 527 (2024) 9736–9756. [doi:10.1093/mnras/stad3802](https://doi.org/10.1093/mnras/stad3802).
- [20] G. B. Mitev, S. N. Yurchenko, J. Tennyson, Predissociation dynamics of the hydroxyl radical (OH) based on a five-state spectroscopic model, *J. Chem. Phys.* 160 (2024) 144110. [doi:10.1063/5.0198241](https://doi.org/10.1063/5.0198241).
- [21] J. Tennyson, M. Pezzella, J. Zhang, S. N. Yurchenko, Data structures for photoadsorption within the ExoMol project, *RAS Tech. Instr.* 2 (2023) 231–237. [doi:10.1093/rasti/rzad014](https://doi.org/10.1093/rasti/rzad014).
- [22] C. E. Hood, J. J. Fortney, M. R. Line, E. C. Martin, C. V. Morley, J. L. Birkby, Z. Rustamkulov, R. E. Lupu, R. S. Freedman, Prospects for Characterizing the Haziest Sub-Neptune Exoplanets with High-resolution Spectroscopy, *Astron. J.* 160 (2020) 198. [doi:10.3847/1538-3881/abb46b](https://doi.org/10.3847/1538-3881/abb46b).
- [23] A. A. A. Piette, N. Madhusudhan, On the Temperature Profiles and Emission Spectra of Mini-Neptune Atmospheres, *Astrophys. J.* 904 (2020) 154. [doi:10.3847/1538-4357/abbbfb1](https://doi.org/10.3847/1538-4357/abbbfb1).
- [24] S.-M. Tsai, H. Innes, T. Lichtenberg, J. Taylor, M. Malik, K. Chubb, R. Pierrehumbert, Inferring Shallow Surfaces on Sub-Neptune Exoplanets with JWST, *Astrophys. J. Lett.* 922 (2021) L27. [doi:10.3847/2041-8213/ac399a](https://doi.org/10.3847/2041-8213/ac399a).
- [25] D. Blain, B. Charnay, B. Bézard, 1D atmospheric study of the temperate sub-Neptune K2-18b, *Astron. Astrophys.* 646 (2021) A15. [doi:10.1051/0004-6361/202039072](https://doi.org/10.1051/0004-6361/202039072).
- [26] B. Charnay, D. Blain, B. Bézard, J. Leconte, M. Turbet, A. Falco, Formation and dynamics of water clouds on temperate sub-Neptunes: the example of K2-18b, *Astron. Astrophys.* 646. [doi:10.1051/0004-6361/202039525](https://doi.org/10.1051/0004-6361/202039525).
- [27] D. Gasman, M. Min, K. L. Chubb, Investigating the detectability of hydrocarbons in exoplanet atmospheres with JWST, *Astron. Astrophys.* 659 (2022) A114. [doi:10.1051/0004-6361/202141468](https://doi.org/10.1051/0004-6361/202141468).
- [28] N. E. Batalha, A. Wolfgang, J. Teske, M. K. Alam, L. Alderson, N. M. Batalha, M. López-Morales, H. R. Wakeford, Importance of Sample Selection in Exoplanet-atmosphere Population Studies, *Astron. J.* 165 (2022) 14. [doi:10.3847/1538-3881/ac9f45](https://doi.org/10.3847/1538-3881/ac9f45).
- [29] M. Zamyatina, E. Hébrard, B. Drummond, N. J. Mayne, J. Manners, D. A. Christie, P. Tremblin, D. K. Sing, K. Kohary, Observability of signatures of transport-induced chemistry in clear atmospheres of hot gas giant exoplanets, *Mon. Not. Roy. Astron. Soc.* 519 (2022) 3129–3153. [doi:10.1093/mnras/stac3432](https://doi.org/10.1093/mnras/stac3432).
- [30] S. Constantinou, N. Madhusudhan, Characterizing atmospheres of cloudy temperate mini-Neptunes with JWST, *Mon. Not. Roy. Astron. Soc.* 514 (2022) 2073–2091. [doi:10.1093/mnras/stac1277](https://doi.org/10.1093/mnras/stac1277).
- [31] A. Guzmán-Mesa, D. Kitzmann, C. Mordasini, K. Heng, Chemical diversity of the atmospheres and interiors of sub-Neptunes: a case study of GJ 436 b, *Mon. Not. Roy. Astron. Soc.* 513 (2022) 4015–4036. [doi:10.1093/mnras/stac1066](https://doi.org/10.1093/mnras/stac1066).
- [32] N. Lawson, G. Zhou, C. X. Huang, D. J. Wright, B. Edwards, E. Nabbie, A. Venner, S. N. Quinn, K. A. Collins, E. Gillen, M. Battley, A. Triaud, C. Hellier, S. Seager, J. N. Winn, J. M. Jenkins, B. Wohler, A. Shporer, R. P. Schwarz, F. Murgas, E. Pallé, D. R. Anderson, R. G. West, R. A. Wittenmyer, B. P. Bowler, J. Horner, S. R. Kane, J. Kielkopf, P. Plavchan, H. Zhang, T. Fairington, J. Okumura, M. W. Mengel, B. C. Addison, Two mini-Neptunes transiting the adolescent K-star HIP 113103 confirmed with TESS and CHEOPS, *Mon. Not. Roy. Astron. Soc.* 527 (2023) 1146–1162. [doi:10.1093/mnras/stac2756](https://doi.org/10.1093/mnras/stac2756).
- [33] P. Gao, A. A. A. Piette, M. E. Steinrueck, M. C. Nixon, M. Zhang, E. M.-R. Kempton, J. L. Bean, E. Rauscher, V. Parmentier, N. E. Batalha, A. B. Savel, K. E. Arnold, M. T. Roman, I. Malsky, J. Taylor, The Hazy and Metal-rich Atmosphere of GJ 1214 b Constrained by Near- and Mid-infrared Transmission Spectroscopy, *Astrophys. J.* 951 (2023) 96. [doi:10.3847/1538-4357/acd16f](https://doi.org/10.3847/1538-4357/acd16f).
- [34] E. M. R. Kempton, M. Zhang, J. L. Bean, M. E. Steinrueck, A. A. A. Piette, V. Parmentier, I. Malsky, M. T. Roman, E. Rauscher, P. Gao, T. J. Bell, Q. Xue, J. Taylor, A. B. Savel, K. E. Arnold, M. C. Nixon, K. B. Stevenson, M. Mansfield, S. Kendrew, S. Zieba, E. Ducrot, A. Dyrek, P.-O. Lagage, K. G. Stassun, G. W. Henry, T. Barman, R. Lupu, M. Malik, T. Kataria, J. Ih, G. Fu, L. Welbanks, P. McGill, A reflective, metal-rich atmosphere for GJ 1214b from its JWST phase curve, *Nature* 620 (2023) 67–71. [doi:10.1038/s41586-023-06159-5](https://doi.org/10.1038/s41586-023-06159-5).
- [35] T. Mikal-Evans, N. Madhusudhan, J. Dittmann, M. N. Günther, L. Welbanks, V. V. Eylen, I. J. M. Crossfield, T. Daylan, L. Kreidberg, Hubble Space Telescope Transmission Spectroscopy for the Temperate Sub-Neptune TOI-270 d: A Possible Hydrogen-rich Atmosphere Containing Water Vapor, *Astron. J.* 165 (2023) 84. [doi:10.3847/1538-3881/aca90b](https://doi.org/10.3847/1538-3881/aca90b).
- [36] Holmberg, Måns, Madhusudhan, Nikku, Possible Hycean conditions in the sub-Neptune TOI-270 d, *Astron. Astrophys.* 683 (2024) L2. [doi:10.1051/0004-6361/202348238](https://doi.org/10.1051/0004-6361/202348238).
- [37] S. Dash, M. Brogi, S. Gandhi, M. Lafarga, A. Meech, A. Bello-Arufe, P. J. Wheatley, Constraints on atmospheric water abundance and

- cloud deck pressure in the warm Neptune GJ 3470 b via CARMENES transmission spectroscopy, arXiv e-prints [arXiv:2404.06648](https://arxiv.org/abs/2404.06648).
- [38] A. Guzmán-Mesa, D. Kitzmann, C. Fisher, A. J. Burgasser, H. J. Hoeijmakers, P. Márquez-Neila, S. L. Grimm, A. M. Mandell, R. Snitman, K. Heng, Information Content of JWST NIRSpec Transmission Spectra of Warm Neptunes, *Astron. J.* 160 (2020) 15. [doi:10.3847/1538-3881/ab9176](https://doi.org/10.3847/1538-3881/ab9176).
- [39] G. Guilluy, A. Gressier, S. Wright, A. Santerne, A. Y. Jaziri, B. Edwards, Q. Changeat, D. Modirrousta-Galian, N. Skaf, A. Al-Refaie, R. Baeyens, M. F. Bieger, D. Blain, F. Kiefer, M. Morvan, L. V. Mugnai, W. Pluriel, M. Poveda, T. Zingales, N. Whiteford, K. H. Yip, B. Charnay, J. Leconte, P. Drossart, A. Sozzetti, E. Marcq, A. Tsiaras, O. Venot, I. Waldmann, J.-P. Beaulieu, ARES IV: Probing the Atmospheres of the Two Warm Small Planets HD 106315c and HD 3167c with the HST/WFC3 Camera*, *Astron. J.* 161 (2020) 19. [doi:10.3847/1538-3881/abc3c8](https://doi.org/10.3847/1538-3881/abc3c8).
- [40] N. A-thano, S. Awiphan, I.-G. Jiang, E. Kerins, A. Priyadarshi, I. McDonald, Y. C. Joshi, T. Chulikorn, J. J. C. Hayes, S. Charles, C.-K. Huang, R. Rattanamala, L.-C. Yeh, V. S. Dhillon, Revisiting the Transit Timing and Atmosphere Characterization of the Neptune-mass Planet HAT-P-26 b, *Astron. J.* 166 (2023) 223. [doi:10.3847/1538-3881/acfea](https://doi.org/10.3847/1538-3881/acfea).
- [41] B. Edwards, Q. Changeat, A. Tsias, A. Allan, P. Behr, S. R. Hagey, M. D. Himes, S. Ma, K. G. Stassun, L. Thomas, A. Thompson, A. Boley, L. Booth, J. Bouwman, K. France, N. Lowson, A. Meech, C. L. Phillips, A. A. Vidotto, K. H. Yip, M. Bieger, A. Gressier, E. Janin, I.-G. Jiang, P. Leonardi, S. Sarkar, N. Skaf, J. Taylor, M. Yang, D. Ward-Thompson, Characterizing a World Within the Hot-Neptune Desert: Transit Observations of LTT 9779 b with the Hubble Space Telescope/WFC3, *Astron. J.* 166 (2023) 158. [doi:10.3847/1538-3881/acea77](https://doi.org/10.3847/1538-3881/acea77).
- [42] M. Radica, L.-P. Coulombe, J. Taylor, L. Albert, R. Allart, B. Benneke, N. B. Cowan, L. Dang, D. Lafrenière, D. Thorngren, Étienne Artigau, R. Doyon, L. Flagg, D. Johnstone, S. Pelletier, P.-A. Roy, Muted Features in the JWST NIRISS Transmission Spectrum of Hot Neptune LTT 9779b, *Astrophys. J. Lett.* 962 (1) (2024) L20. [doi:10.3847/2041-8213/ad20e4](https://doi.org/10.3847/2041-8213/ad20e4).
- [43] N. Madhusudhan, A. A. A. Piette, S. Constantinou, Habitability and Biosignatures of Hycean Worlds, *Astrophys. J.* 918 (2021) 1. [doi:10.3847/1538-4357/abfd9c](https://doi.org/10.3847/1538-4357/abfd9c).
- [44] N. Madhusudhan, S. Sarkar, S. Constantinou, M. Holmberg, A. A. A. Piette, J. I. Moses, Carbon-bearing Molecules in a Possible Hycean Atmosphere, *Astrophys. J. Lett.* 956 (2023) L13. [doi:10.3847/2041-8213/acf577](https://doi.org/10.3847/2041-8213/acf577).
- [45] E. M.-R. Kempton, M. Lessard, M. Malik, L. A. Rogers, K. E. Futrowsky, J. Ih, N. Marounina, C. E. M. noz Romero, Where are the Water Worlds?: Self-consistent Models of Water-rich Exoplanet Atmospheres, *Astrophys. J.* 953 (1) (2023) 57. [doi:10.3847/1538-4357/ace10d](https://doi.org/10.3847/1538-4357/ace10d).
- [46] J. J. Spake, D. K. Sing, H. R. Wakeford, N. Nikolov, T. Mikal-Evans, D. Deming, J. K. Barstow, D. R. Anderson, A. L. Carter, M. Gillon, J. M. Goyal, G. Hebrard, C. Hellier, T. Kataria, K. W. F. Lam, A. H. M. J. Triaud, P. J. Wheatley, Abundance measurements of H₂O and carbon-bearing species in the atmosphere of WASP-127b confirm its supersolar metallicity, *Mon. Not. Roy. Astron. Soc.* 500 (2020) 4042–4064. [doi:10.1093/mnras/staa3116](https://doi.org/10.1093/mnras/staa3116).
- [47] L. O. Anisman, B. Edwards, Q. Changeat, O. Venot, A. F. Al-Refaie, A. Tsias, G. Tinetti, WASP-117 b: An Eccentric Hot Saturn as a Future Complex Chemistry Laboratory, *Astron. J.* 160 (2020) 233. [doi:10.3847/1538-3881/abb9b0](https://doi.org/10.3847/1538-3881/abb9b0).
- [48] Q. Changeat, B. Edwards, A. F. Al-Refaie, M. Morvan, A. Tsias, I. P. Waldmann, G. Tinetti, KELT-11 b: Abundances of Water and Constraints on Carbon-bearing Molecules from the Hubble Transmission Spectrum, *Astron. J.* 160 (2020) 260. [doi:10.3847/1538-3881/abbe12](https://doi.org/10.3847/1538-3881/abbe12).
- [49] K. D. Colón, L. Kreidberg, L. Welbanks, M. R. Line, N. Madhusudhan, T. Beatty, P. Tamburo, K. B. Stevenson, A. Mandell, J. E. Rodriguez, T. Barclay, E. D. Lopez, K. G. Stassun, D. Angerhausen, J. J. Fortney, D. J. James, J. Pepper, J. P. Ahlers, P. Plavchan, S. Awiphan, C. Kotnik, K. K. McLeod, G. Murawski, H. Chotani, D. LeBrun, W. Matzko, D. Rea, M. Vidaurri, S. Webster, J. K. Williams, L. S. Cox, N. Tan, E. A. Gilbert, An Unusual Transmission Spectrum for the Sub-Saturn KELT-11b Suggestive of a Subsolar Water Abundance, *Astron. J.* 160 (2020) 280. [doi:10.3847/1538-3881/abc1e9](https://doi.org/10.3847/1538-3881/abc1e9).
- [50] M. C. Maimone, M. Brogi, A. Chiavassa, M. E. van den Ancker, C. F. Manara, J. Leconte, S. Gandhi, W. Pluriel, Detecting H₂O with CRIRES+: WASP-20b, *Astron. Astrophys.* 667 (2022) A106. [doi:10.1051/0004-6361/202244383](https://doi.org/10.1051/0004-6361/202244383).
- [51] E.-M. Ahrer, L. Alderson, N. M. Batalha, N. E. Batalha, J. L. Bean, T. G. Beatty, T. J. Bell, B. Benneke, Z. K. Berta-Thompson, A. L. Carter, I. J. M. Crossfield, N. Espinoza, A. D. Feinstein, J. J. Fortney, N. P. Gibson, J. M. Goyal, E. M.-R. Kempton, J. Kirk, L. Kreidberg, M. Lopez-Morales, M. R. Line, J. D. Lothringer, S. E. Moran, S. Mukherjee, K. Ohno, V. Parmentier, C. Piaulet, Z. Rustamkulov, E. Schlawin, D. K. Sing, K. B. Stevenson, H. R. Wakeford, N. H. Allen, S. M. Birkmann, J. Brande, N. Crouzet, P. E. Cubillos, M. Damiano, J.-M. Desert, P. Gao, J. Harrington, R. Hu, S. Kendrew, H. A. Knutson, P.-O. Lagage, J. Leconte, M. Lendl, R. J. MacDonald, E. M. May, Y. Miguel, K. Molaverdikhani, J. Moses, I. C. A. Murray, M. Nehring, N. K. Nikolov, D. J. M. P. D. de la Roche, M. Radica, P.-A. Roy, K. G. Stassun, J. Taylor, W. C. Waalkes, P. Wachiraphan, L. Welbanks, P. J. Wheatley, K. Aggarwal, M. K. Alam, A. Banerjee, J. K. Barstow, J. Blecic, S. L. Casewell, Q. Changeat, K. L. Chubb, K. D. Colon, L.-P. Coulombe, T. Daylan, M. De Val-Borro, L. Decin, L. A. Dos Santos, L. Flagg, K. France, G. Fu, A. G. Munoz, J. E. Gizis, A. Glidden, D. Grant, K. Heng, T. Henning, Y.-C. Hong, J. Inglis, N. Iro, T. Kataria, T. D. Komacek, J. E. Krick, E. K. H. Lee, N. K. Lewis, J. Lillo-Box, J. Lustig-Yaeger, L. Mancini, A. M. Mandell, M. Mansfield, M. S. Marley, T. Mikal-Evans, G. Morello, M. C. Nixon, K. O. Ceballos, A. A. A. Piette, D. Powell, B. Rackham, V. L. Ramos-Rosado, E. Rauscher, S. Redfield, L. K. Rogers, M. T. Roman, G. M. Roudier, N. Scarsdale, E. L. Shkolnik, J. Southworth, J. J. Spake, M. E. Steinrueck, X. Tan, J. K. Teske, P. Tremblin, S.-M. Tsai, G. S. Tucker, J. D. Turner, J. A. Valenti, O. Venot, I. P. Waldmann, N. L. Wallack, X. Zhang, S. Zieba, Identification of carbon dioxide in an exoplanet atmosphere, *Nature* 614 (2023) 649–652. [doi:10.1038/s41586-022-05269-w](https://doi.org/10.1038/s41586-022-05269-w).
- [52] N. K. Nikolov, D. K. Sing, J. J. Spake, B. Smalley, J. M. Goyal, T. Mikal-Evans, H. R. Wakeford, Z. Rustamkulov, D. Deming, J. J. Fortney, A. Carter, N. P. Gibson, N. J. Mayne, Solar-to-supersolar sodium and oxygen absolute abundances for a ‘hot Saturn’ orbiting a metal-rich star, *Mon. Not. Roy. Astron. Soc.* 515 (2022) 3037–3058. [doi:10.1093/mnras/stac1530](https://doi.org/10.1093/mnras/stac1530).
- [53] S.-M. Tsai, E. K. H. Lee, D. Powell, P. Gao, X. Zhang, J. Moses, E. Hébrard, O. Venot, V. Parmentier, S. Jordan, R. Hu, M. K. Alam, L. Alderson, N. M. Batalha, J. L. Bean, B. Benneke, C. J. Bierson, R. P. Brady, L. Carone, A. L. Carter, K. L. Chubb, J. Inglis, J. Leconte, M. Line, M. López-Morales, Y. Miguel, K. Molaverdikhani, Z. Rustamkulov, D. K. Sing, K. B. Stevenson, H. R. Wakeford, J. Yang, K. Aggarwal, R. Baeyens, S. Barat, M. de Val-Borro, T. Daylan, J. J. Fortney, K. France, J. M. Goyal, D. Grant, J. Kirk, L. Kreidberg, A. Louca, S. E. Moran, S. Mukherjee, E. Nasedkin, K. Ohno, B. V. Rackham, S. Redfield, J. Taylor, P. Tremblin, C. Visscher, N. L. Wallack, L. Welbanks, A. Youngblood, E.-M. Ahrer, N. E. Batalha, P. Behr, Z. K. Berta-Thompson, J. Blecic, S. L. Casewell, I. J. M. Crossfield, N. Crouzet, P. E. Cubillos, L. Decin, J.-M. Désert, A. D. Feinstein, N. P. Gibson, J. Harrington, K. Heng, T. Henning, E. M.-R. Kempton, J. Krick, P.-O. Lagage, M. Lendl, J. D. Lothringer, M. Mansfield, N. J. Mayne, T. Mikal-Evans, E. Palle, E. Schlawin, O. Shorttle, P. J. Wheatley, S. N. Yurchenko, Photochemically produced SO₂ in the atmosphere of WASP-39b, *Nature* 617 (2023) 483–487. [doi:10.1038/s41586-023-05902-2](https://doi.org/10.1038/s41586-023-05902-2).
- [54] D. Samra, C. Helling, K. L. Chubb, M. Min, L. Carone, A. D. Schneider, Clouds form on the hot Saturn JWST ERO target WASP-96b,

- Astron. Astrophys. 669 (2023) 17. [doi:10.1051/0004-6361/202244939](https://doi.org/10.1051/0004-6361/202244939).
- [55] M. Radica, L. Welbanks, N. Espinoza, J. Taylor, L.-P. Coulombe, A. D. Feinstein, J. Goyal, N. Scarsdale, L. Albert, P. Baghel, J. L. Bean, J. Blecic, D. Lafrenière, R. J. MacDonald, M. Zamyatina, R. Allart, E. Artigau, N. E. Batalha, N. J. Cook, N. B. Cowan, L. Dang, R. Doyon, M. Fournier-Tondreau, D. Johnstone, M. R. Line, S. E. Moran, S. Mukherjee, S. Pelletier, P.-A. Roy, G. J. Talens, J. Filippazzo, K. Pontoppidan, K. Volk, Awesome SOSS: transmission spectroscopy of WASP-96b with NIRISS/SOSS, Mon. Not. Roy. Astron. Soc. 524 (2023) 835–856. [doi:10.1093/mnras/stad1762](https://doi.org/10.1093/mnras/stad1762).
- [56] L. Alderson, H. R. Wakeford, M. K. Alam, N. E. Batalha, J. D. Lothringer, J. A. Redai, S. Barat, J. Brande, M. Damiano, T. Daylan, N. Espinoza, L. Flagg, J. M. Goyal, D. Grant, R. Hu, J. Inglis, E. K. H. Lee, T. Mikal-Evans, L. Ramos-Rosado, P.-A. Roy, N. L. Wallack, N. M. Batalha, J. L. Bean, B. Benneke, Z. K. Berta-Thompson, A. L. Carter, Q. Changeat, K. D. Colon, I. J. M. Crossfield, J.-M. Desert, D. Foreman-Mackey, N. P. Gibson, L. Kreidberg, M. R. Line, M. Lopez-Morales, K. Molaverdikhani, S. E. Moran, G. Morello, J. I. Moses, S. Mukherjee, E. Schlawin, D. K. Sing, K. B. Stevenson, J. Taylor, K. Aggarwal, E.-M. Ahrer, N. H. Allen, J. K. Barstow, T. J. Bell, J. Blecic, S. L. Casewell, K. L. Chubb, N. Crouzet, P. E. Cubillos, L. Decin, A. D. Feinstein, J. J. Fortney, J. Harrington, K. Heng, N. Iro, E. M. R. Kempton, J. Kirk, H. A. Knutson, J. Krick, J. Leconte, M. Lendl, R. J. McDonald, L. Mancini, M. Mansfield, E. M. May, N. J. Mayne, Y. Miguel, N. K. Nikolov, K. Ohno, E. Palle, V. Parmentier, D. J. M. P. D. de la Roche, C. Piaulet, D. Powell, B. V. Rackham, S. Redfield, L. K. Rogers, Z. Rustamkulov, X. Tan, P. Tremblin, S.-M. Tsai, J. D. Turner, M. de Val-Borro, O. Venot, L. Welbanks, P. J. Wheatley, X. Zhang, Early Release Science of the exoplanet WASP-39b with JWST NIRSpec G395H, Nature 614 (2023) 664–669. [doi:10.1038/s41586-022-05591-3](https://doi.org/10.1038/s41586-022-05591-3).
- [57] M. Fournier-Tondreau, R. J. McDonald, M. Radica, D. Lafrenière, L. Welbanks, C. Piaulet, L.-P. Coulombe, R. Allart, K. Morel, E. Artigau, L. Albert, O. Lim, R. Doyon, B. Benneke, J. F. Rowe, A. Darveau-Bernier, N. B. Cowan, N. K. Lewis, N. J. Cook, L. Flagg, F. Genest, S. Pelletier, D. Johnstone, L. Dang, L. Kaltenegger, J. Taylor, J. D. Turner, Near-infrared transmission spectroscopy of HAT-P-18 b with NIRISS: Disentangling planetary and stellar features in the era of JWST, Mon. Not. Roy. Astron. Soc. 528 (2023) 3354–3377. [doi:10.1093/mnras/stad3813](https://doi.org/10.1093/mnras/stad3813).
- [58] J. Taylor, M. Radica, L. Welbanks, R. J. McDonald, J. Blecic, M. Zamyatina, A. Roth, J. L. Bean, V. Parmentier, L.-P. Coulombe, A. D. Feinstein, N. Espinoza, B. Benneke, D. Lafrenière, R. Doyon, E.-M. Ahrer, Awesome SOSS: atmospheric characterization of WASP-96b using the JWST early release observations, Mon. Not. Roy. Astron. Soc. 524 (2023) 817–834. [doi:10.1093/mnras/stad1547](https://doi.org/10.1093/mnras/stad1547).
- [59] A. Boucher, D. Lafrenière, S. Pelletier, A. Darveau-Bernier, M. Radica, R. Allart, e. Artigau, N. J. Cook, F. Debras, R. Doyon, E. Gaidos, B. Benneke, C. Cadieux, A. Carmona, R. Cloutier, P. Cortés-Zuleta, N. B. Cowan, X. Delfosse, J.-F. Donati, P. Fouqué, T. Forveille, K. Grankin, G. Hébrard, J. H. C. Martins, E. Martioli, A. Masson, S. Vinatier, CO or no CO? Narrowing the CO abundance constraint and recovering the H₂O detection in the atmosphere of WASP-127 b using SPIRou, Mon. Not. Roy. Astron. Soc. 522 (2023) 5062–5083. [doi:10.1093/mnras/stad1247](https://doi.org/10.1093/mnras/stad1247).
- [60] N. Whiteford, A. Glasse, K. L. Chubb, D. Kitzmann, S. Ray, M. W. Phillips, B. A. Biller, P. I. Palmer, K. Rice, I. P. Waldmann, Q. Changeat, N. Skaf, J. Wang, B. Edwards, A. Al-Refaie, Retrieval study of cool, directly imaged exoplanet 51 Eri b, Mon. Not. Roy. Astron. Soc. 525 (2023) 1375–1400. [doi:10.1093/mnras/stad670](https://doi.org/10.1093/mnras/stad670).
- [61] S. Xu, H. Diamond-Lowe, R. J. McDonald, A. Vanderburg, S. Blouin, P. Dufour, P. Gao, L. Kreidberg, S. K. Leggett, A. W. Mann, C. V. Morley, A. W. Stephens, C. E. O'Connor, P. C. Thao, N. K. Lewis, Gemini/GMOS Transmission Spectroscopy of the Grazing Planet Candidate WD 1856+534 b, Astron. J. 162 (2021) 296. [doi:10.3847/1538-3881/ac2d26](https://doi.org/10.3847/1538-3881/ac2d26).
- [62] G. Guilluy, P. Giacobbe, I. Carleo, P. E. Cubillos, A. Sozzetti, A. S. Bonomo, M. Brogi, S. Gandhi, L. Fossati, V. Nascimbeni, D. Turrini, E. Schisano, F. Borsa, A. F. Lanza, L. Mancini, A. Maggio, L. Malavolta, G. Micela, L. Pino, M. Rainer, A. Bignamini, R. Claudi, R. Cosentino, E. Covino, S. Desidera, A. Fiorenzano, A. Harutyunyan, V. Lorenzi, C. Knapic, E. Molinari, E. Pacetti, I. Pagano, M. Pedani, G. Piotto, E. Poretti, The GAPS Programme at TNG XXXVIII. Five molecules in the atmosphere of the warm giant planet WASP-69b detected at high spectral resolution, Astron. Astrophys. 665 (2022) A104. [doi:10.1051/0004-6361/202243854](https://doi.org/10.1051/0004-6361/202243854).
- [63] Z. Zhang, P. Mollière, K. Hawkins, C. Manea, J. J. Fortney, C. V. Morley, A. Skemer, M. S. Marley, B. P. Bowler, A. L. Carter, K. Franson, Z. G. Maas, C. Sneden, ELMental abundances of Planets and brown dwarfs Imaged around Stars (ELPIS). I. Potential Metal Enrichment of the Exoplanet AF Lep b and a Novel Retrieval Approach for Cloudy Self-luminous Atmospheres, Astron. J. 166 (2023) 198. [doi:10.3847/1538-3881/acf768](https://doi.org/10.3847/1538-3881/acf768).
- [64] I. Carleo, P. Giacobbe, G. Guilluy, P. E. Cubillos, A. S. Bonomo, A. Sozzetti, M. Brogi, S. Gandhi, L. Fossati, D. Turrini, K. Biazzo, F. Borsa, A. F. Lanza, L. Malavolta, A. Maggio, L. Mancini, G. Micela, L. Pino, E. Poretti, M. Rainer, G. Scandariato, E. Schisano, G. Andreuzzi, A. Bignamini, R. Cosentino, A. Fiorenzano, A. Harutyunyan, E. Molinari, M. Pedani, S. Redfield, H. Stoev, The GAPS Programme at TNG XXXIX. Multiple Molecular Species in the Atmosphere of the Warm Giant Planet WASP-80 b Unveiled at High Resolution with tIANO-B *, Astron. Astrophys. 164 (2022) 101. [doi:10.3847/1538-3881/ac80bf](https://doi.org/10.3847/1538-3881/ac80bf).
- [65] T. J. Bell, L. Welbanks, E. Schlawin, M. R. Line, J. J. Fortney, T. P. Greene, K. Ohno, V. Parmentier, E. Rauscher, T. G. Beatty, S. Mukherjee, L. S. Wiser, M. L. Boyer, M. J. Rieke, J. A. Stansberry, Methane throughout the atmosphere of the warm exoplanet WASP-80b, Nature 623 (2023) 709–712. [doi:10.1038/s41586-023-06687-0](https://doi.org/10.1038/s41586-023-06687-0).
- [66] S. Kanodia, S. Mahadevan, J. Libby-Roberts, G. Stefansson, C. I. Canas, A. A. A. Piette, A. Boss, J. Teske, J. Chambers, G. Zeimann, A. Monson, P. Robertson, J. P. Ninan, A. S. J. Lin, C. F. Bender, W. D. Cochran, S. A. Diddams, A. F. Gupta, S. Halverson, S. Hawley, H. A. Kobulnicky, A. J. Metcalf, B. A. Parker, L. Powers, L. W. Ramsey, A. Roy, C. Schwab, T. N. Swaby, R. C. Terrien, J. Wisniewski, TOI-5205b: A Short-period Jovian Planet Transiting a Mid-M Dwarf, Astron. J. 165 (2023) 120. [doi:10.3847/1538-3881/acabce](https://doi.org/10.3847/1538-3881/acabce).
- [67] M. Zhang, Y. Chachan, E. M.-R. Kempton, H. A. Knutson, W. H. Chang, PLATON II: New Capabilities and a Comprehensive Retrieval on HD 189733b Transit and Eclipse Data, Astrophys. J. 899 (2020) 27. [doi:10.3847/1538-4357/aba1e6](https://doi.org/10.3847/1538-4357/aba1e6).
- [68] A. A. A. Piette, N. Madhusudhan, L. K. McKemmish, S. Gandhi, T. Masseron, L. Welbanks, Assessing spectra and thermal inversions due to TiO in hot Jupiter atmospheres, Mon. Not. Roy. Astron. Soc. 496 (2020) 3870–3886. [doi:10.1093/mnras/staa1592](https://doi.org/10.1093/mnras/staa1592).
- [69] K. H. Yip, Q. Changeat, B. Edwards, M. Morvan, K. L. Chubb, A. Tsiaras, I. P. Waldmann, G. Tinetti, On the Compatibility of Ground-based and Space-based Data: WASP-96 b, an Example*, Astron. J. 161 (2020) 4. [doi:10.3847/1538-3881/abc179](https://doi.org/10.3847/1538-3881/abc179).
- [70] M. Lendl, S. Csizmadia, A. Deline, L. Fossati, D. Kitzmann, K. Heng, S. Hoyer, S. Salmon, W. Benz, C. Broeg, D. Ehrenreich, A. Fortier, D. Queloz, A. Bonfanti, A. Brandeker, A. Collier Cameron, L. Delrez, A. Garcia Muñoz, M. J. Hooton, P. F. L. Maxted, B. M. Morris, V. Van Grootel, T. G. Wilson, Y. Alibert, R. Alonso, J. Asquier, T. Bandy, T. Bárczy, D. Barrado, S. C. C. Barros, W. Baumjohann, M. Beck, T. Beck, A. Bekkelien, M. Bergomi, N. Billot, F. Biondi, X. Bonfils, V. Bourrier, M. D. Busch, J. Cabrera, V. Cessa, S. Charnoz, B. Chazelas, C. Corral Van Damme, M. B. Davies, M. Deleuil, O. D. S. Demangeon, B. O. Demory, A. Erikson, J. Farinato, M. Fridlund, D. Futyan, D. Gandolfi, M. Gillon, P. Guterman, J. Hasiba, E. Hernandez, K. G. Isaak, L. Kiss, T. Kuntzer, A. Lecavelier des Etangs, T. Lüftinger, J. Laskar, C. Lovis,

- D. Magrin, L. Malvasio, L. Marafatto, H. Michaelis, M. Munari, V. Nascimbeni, G. Olofsson, H. Ottacher, R. Ottensamer, I. Pagano, E. Pallé, G. Peter, D. Piazza, G. Piotto, D. Pollacco, F. Ratti, H. Rauer, R. Ragazzoni, N. Rando, I. Ribas, M. Rieder, R. Rohlfs, F. Safa, N. C. Santos, G. Scandariato, D. Ségransan, A. E. Simon, V. Singh, A. M. S. Smith, M. Sordet, S. G. Sousa, M. Steller, G. M. Szabó, N. Thomas, M. Tschentscher, S. Udry, V. Viotto, I. Walter, N. A. Walton, F. Wildi, D. Wolter, The hot dayside and asymmetric transit of WASP-189 b seen by CHEOPS, *Astron. Astrophys.* 643 (2020) A94. [doi:10.1051/0004-6361/202038677](https://doi.org/10.1051/0004-6361/202038677).
- [71] B. I. Lacy, A. Burrows, JWST Transit Spectra. I. Exploring Potential Biases and Opportunities in Retrievals of Tidally Locked Hot Jupiters with Clouds and Hazes, *Astrophys. J.* 905 (2020) 131. [doi:10.3847/1538-4357/abc01c](https://doi.org/10.3847/1538-4357/abc01c).
- [72] G. Chen, N. Casasayas-Barris, E. Palle, L. Welbanks, N. Madhusudhan, R. Luque, F. Murgas, Detection of Na in WASP-21b's lower and upper atmosphere, *Astron. Astrophys.* 642 (2020) A54. [doi:10.1051/0004-6361/202038661](https://doi.org/10.1051/0004-6361/202038661).
- [73] K. K. Wilcomb, Q. M. Konopacky, T. S. Barman, C. A. Theissen, J.-B. Ruffio, L. Brock, B. Macintosh, C. Marois, Moderate-resolution K-band Spectroscopy of Substellar Companion κ Andromedae b, *Astron. J.* 160 (2020) 207. [doi:10.3847/1538-3881/abb9b1](https://doi.org/10.3847/1538-3881/abb9b1).
- [74] P. Giacobbe, M. Brogi, S. Gandhi, P. E. Cubillos, A. S. Bonomo, A. Sozzetti, L. Fossati, G. Guilluy, I. Carleo, M. Rainer, A. Harutyunyan, F. Borsa, L. Pino, V. Nascimbeni, S. Benatti, K. Biazzo, A. Bignamini, K. L. Chubb, R. Claudi, R. Cosentino, E. Covino, M. Damasso, S. Desidera, A. F. M. Fiorenzano, A. Ghedina, A. F. Lanza, G. Leto, A. Maggio, L. Malavolta, J. Maldonado, G. Micela, E. Molinari, I. Pagano, M. Pedani, G. Piotto, E. Poretti, G. Scandariato, S. N. Yurchenko, D. Fantinel, A. Galli, M. Lodi, N. Sanna, A. Tozzi, Five carbon- and nitrogen-bearing species in a hot giant planet's atmosphere, *Nature* 592 (2021) 205–208. [doi:10.1038/s41586-021-03381-x](https://doi.org/10.1038/s41586-021-03381-x).
- [75] Casasayas-Barris, N., Palle, E., Stangret, M., Bourrier, V., Tabernero, H. M., Yan, F., Borsa, F., Allart, R., Zapatero Osorio, M. R., Lovis, C., Sousa, S. G., Chen, G., Oshagh, M., Santos, N. C., Pepe, F., Rebolo, R., Molaro, P., Cristiani, S., Adibekyan, V., Alibert, Y., Allende Prieto, C., Bouchy, F., Demangeon, O. D. S., Di Marcantonio, P., D'Odorico, V., Ehrenreich, D., Figueira, P., Génova Santos, R., González Hernández, J. I., Lavie, B., Lillo-Box, J., Lo Curto, G., Martins, C. J. A. P., Mehner, A., Micela, G., Nunes, N. J., Poretti, E., Sozzetti, A., Suárez Mascareño, A., Udry, S., The atmosphere of HD 209458b seen with ESPRESSO - No detectable planetary absorptions at high resolution, *Astron. Astrophys.* 647 (2021) A26. [doi:10.1051/0004-6361/202039539](https://doi.org/10.1051/0004-6361/202039539).
- [76] D. B. Serindag, I. A. G. Snellen, P. Mollière, Measuring titanium isotope ratios in exoplanet atmospheres, *Astron. Astrophys.* 655 (2021) A69. [doi:10.1051/0004-6361/202141941](https://doi.org/10.1051/0004-6361/202141941).
- [77] K. B. Sheppard, L. Welbanks, A. M. Mandell, N. Madhusudhan, N. Nikolov, D. Deming, G. W. Henry, M. H. Williamson, D. K. Sing, M. López-Morales, J. Ih, J. Sanz-Forcada, P. Lavvas, G. E. Ballester, T. M. Evans, A. G. Muñoz, L. A. dos Santos, The Hubble PanCET Program: A Metal-rich Atmosphere for the Inflated Hot Jupiter HAT-P-41b, *Astrophys. J.* 161 (2021) 51. [doi:10.3847/1538-3881/abc8f4](https://doi.org/10.3847/1538-3881/abc8f4).
- [78] M. Braam, F. F. S. van der Tak, K. L. Chubb, M. Min, Evidence for chromium hydride in the atmosphere of hot Jupiter WASP-31b, *Astron. Astrophys.* 646 (2021) A17. [doi:10.1051/0004-6361/202039509](https://doi.org/10.1051/0004-6361/202039509).
- [79] Q. Changeat, A. F. Al-Refaie, B. Edwards, I. P. Waldmann, G. Tinetti, An exploration of model degeneracies with a unified phase curve retrieval analysis: The light and dark sides of WASP-43 b, *Astrophys. J.* 913 (2021) 73. [doi:10.3847/1538-4357/abf2bb](https://doi.org/10.3847/1538-4357/abf2bb).
- [80] A. D. Rathcke, R. J. MacDonald, J. K. Barstow, J. M. Goyal, M. Lopez-Morales, J. ao M. Mendonça, J. Sanz-Forcada, G. W. Henry, D. K. Sing, M. K. Alam, N. K. Lewis, K. L. Chubb, J. Taylor, N. Nikolov, L. A. Buchhave, HST PanCET Program: A Complete Near-UV to Infrared Transmission Spectrum for the Hot Jupiter WASP-79b, *Astron. J.* 162 (2021) 138. [doi:10.3847/1538-3881/ac0e99](https://doi.org/10.3847/1538-3881/ac0e99).
- [81] M. Ishizuka, H. Kawahara, S. K. Nugroho, Y. Kawashima, T. Hirano, M. Tamura, Neutral Metals in the Atmosphere of HD 149026b, *Astron. J.* 161 (2021) 153. [doi:10.3847/1538-3881/abdb25](https://doi.org/10.3847/1538-3881/abdb25).
- [82] P. E. Cubillos, D. Keating, N. B. Cowan, J. M. Vos, B. Burningham, M. Ygouf, T. Karalidi, Y. Zhou, E. C. Gonzales, Longitudinally Resolved Spectral Retrieval (ReSpect) of WASP-43b, *Astrophys. J.* 915 (2021) 45. [doi:10.3847/1538-4357/abfe14](https://doi.org/10.3847/1538-4357/abfe14).
- [83] E. K. H. Lee, V. Parmentier, M. Hammond, S. L. Grimm, D. Kitzmann, X. Tan, S.-M. Tsai, R. T. Pierrehumbert, Simulating gas giant exoplanet atmospheres with Exo-FMS: comparing semigrey, picket fence, and correlated- k radiative-transfer schemes, *Mon. Not. Roy. Astron. Soc.* 506 (2021) 2695–2711. [doi:10.1093/mnras/stab1851](https://doi.org/10.1093/mnras/stab1851).
- [84] R. Baeyens, L. Decin, L. Carone, O. Venot, M. Agúndez, P. Mollière, Grid of pseudo-2D chemistry models for tidally locked exoplanets - I. The role of vertical and horizontal mixing, *Mon. Not. Roy. Astron. Soc.* 505 (2021) 5603–5653. [doi:10.1093/mnras/stab1310](https://doi.org/10.1093/mnras/stab1310).
- [85] J.-B. Ruffio, Q. M. Konopacky, T. Barman, B. Macintosh, K. K. W. Hoch, R. J. D. Rosa, J. J. Wang, I. Czekala, C. Marois, Deep Exploration of the Planets HR 8799 b, c, and d with Moderate-resolution Spectroscopy, *Astron. J.* 162 (2021) 290. [doi:10.3847/1538-3881/ac273a](https://doi.org/10.3847/1538-3881/ac273a).
- [86] E. Sedaghati, R. J. MacDonald, N. Casasayas-Barris, H. J. Hoeijmakers, H. M. J. Boffin, F. Rodler, R. Brahm, M. Jones, A. Sánchez-López, I. Carleo, P. Figueira, A. Mehner, M. López-Puertas, A spectral survey of WASP-19b with ESPRESSO, *Mon. Not. Roy. Astron. Soc.* 505 (2021) 435–458. [doi:10.1093/mnras/stab1164](https://doi.org/10.1093/mnras/stab1164).
- [87] M. R. Line, M. Brogi, J. L. Bean, S. Gandhi, J. Zalesky, V. Parmentier, P. Smith, G. N. Mace, M. Mansfield, E. M.-R. Kempton, J. J. Fortney, E. Shkolnik, J. Patience, E. Rauscher, J.-M. Desert, J. P. Wardenier, A solar C/O and sub-solar metallicity in a hot Jupiter atmosphere, *Nature* 598 (2021) 580+. [doi:10.1038/s41586-021-03912-6](https://doi.org/10.1038/s41586-021-03912-6).
- [88] J. J. Wang, J.-B. Ruffio, E. Morris, J.-R. Delorme, N. Jovanovic, J. Pezzato, D. Echeverri, L. Finnerty, C. Hood, J. J. Zanazzi, M. L. Bryan, C. Z. Bond, S. Cetre, E. C. Martin, D. Mawet, A. Skemer, A. Baker, J. W. Xuan, J. K. Wallace, J. Wang, R. Bartos, G. A. Blake, A. Boden, C. Buzard, B. Calvin, M. Chun, G. Doppmann, T. J. Dupuy, G. Duchêne, Y. K. Feng, M. P. Fitzgerald, J. Fortney, R. S. Freedman, H. Knutson, Q. Konopacky, S. Lilley, M. C. Liu, R. Lopez, R. Lupu, M. S. Marley, T. Meshkat, B. Miles, M. Millar-Blanchaer, S. Ragland, A. Roy, G. Ruane, B. Sappey, T. Schofield, L. Weiss, E. Wetherell, P. Wizinowich, M. Ygouf, Detection and Bulk Properties of the HR 8799 Planets with High-resolution Spectroscopy, *Astron. J.* 162 (2021) 148. [doi:10.3847/1538-3881/ac1349](https://doi.org/10.3847/1538-3881/ac1349).
- [89] L. Welbanks, N. Madhusudhan, On Atmospheric Retrievals of Exoplanets with Inhomogeneous Terminators, *Astrophys. J.* 933 (2022) 79. [doi:10.3847/1538-4357/ac6df1](https://doi.org/10.3847/1538-4357/ac6df1).
- [90] K. L. Chubb, M. Min, Exoplanet Atmosphere Retrievals in 3D Using Phase Curve Data with ARCIIS: Application to WASP-43b, *Astron. Astrophys.* 665 (2022) A2. [doi:10.1051/0004-6361/202142800](https://doi.org/10.1051/0004-6361/202142800).
- [91] A. Saba, A. Tsiaras, M. Morvan, A. Thompson, Q. Changeat, B. Edwards, A. Jolly, I. Waldmann, G. Tinetti, The Transmission Spectrum of WASP-17 b From the Optical to the Near-infrared Wavelengths: Combining STIS, WFC3, and IRAC Data Sets, *Astron. J.* 164 (2022) 2. [doi:10.3847/1538-3881/ac6c01](https://doi.org/10.3847/1538-3881/ac6c01).
- [92] Q. Changeat, B. Edwards, A. F. Al-Refaie, A. Tsiaras, J. W. Skinner, J. Y. K. Cho, K. H. Yip, L. Anisman, M. Ikoma, M. F. Bieger, O. Venot, S. Shibata, I. P. Waldmann, G. Tinetti, Five Key Exoplanet Questions Answered via the Analysis of 25 Hot-Jupiter Atmospheres in Eclipse, *Astrophys. J. Suppl. Ser.* 260 (2022) 3. [doi:10.3847/1538-4365/ac5cc2](https://doi.org/10.3847/1538-4365/ac5cc2).
- [93] A. D. Schneider, L. Carone, L. Decin, U. G. Jørgensen, P. Mollière, R. Baeyens, S. Kiefer, C. Helling, Exploring the deep atmospheres of

- HD 209458b and WASP-43b using a non-gray general circulation model, *Astron. Astrophys.* 664 (2022) A56. [doi:10.1051/0004-6361/202142728](https://doi.org/10.1051/0004-6361/202142728).
- [94] A. D. Schneider, L. Carone, L. Decin, U. G. Jørgensen, C. Helling, No evidence for radius inflation in hot Jupiters from vertical advection of heat, *Astron. Astrophys.* 666 (2022) L11. [doi:10.1051/0004-6361/202244797](https://doi.org/10.1051/0004-6361/202244797).
- [95] A. F. Al-Refaie, Q. Changeat, O. Venot, I. P. Waldmann, G. Tinetti, A Comparison of Chemical Models of Exoplanet Atmospheres Enabled by TauREx 3.1, *Astrophys. J.* 932 (2022) 123. [doi:10.3847/1538-4357/ac6dcf](https://doi.org/10.3847/1538-4357/ac6dcf).
- [96] D. Kasper, J. L. Bean, M. R. Line, A. Seifahrt, M. T. Brady, J. Lothringer, L. Pino, G. Fu, S. Pelletier, J. Stürmer, B. Benneke, M. Brogi, J.-M. Désert, Unifying High- and Low-resolution Observations to Constrain the Dayside Atmosphere of KELT-20b/MASCARA-2b, *Astron. J.* 165 (1) (2022) 7. [doi:10.3847/1538-3881/ac9f40](https://doi.org/10.3847/1538-3881/ac9f40).
- [97] R. K. Webb, S. Gandhi, M. Brogi, J. L. Birkby, E. de Mooij, I. Snellen, Y. Zhang, Water observed in the atmosphere of τ Boötis Ab with CARMENES/CAHA, *Mon. Not. Roy. Astron. Soc.* 514 (2022) 4160–4172. [doi:10.1093/mnras/stac1512](https://doi.org/10.1093/mnras/stac1512).
- [98] R. Deitrick, K. Heng, U. Schroffenegger, D. Kitzmann, S. L. Grimm, M. Malik, J. a. M. Mendonça, B. M. Morris, The THOR + HELIOS general circulation model: multiwavelength radiative transfer with accurate scattering by clouds/hazes, *Mon. Not. Roy. Astron. Soc.* 512 (2022) 3759–3787. [doi:10.1093/mnras/stac680](https://doi.org/10.1093/mnras/stac680).
- [99] M. Blažek, P. Kabáth, A. A. A. Piette, N. Madhusudhan, M. Skarka, J. Šubjak, D. R. Anderson, H. M. J. Boffin, C. C. Cáceres, N. P. Gibson, S. Hoyer, V. D. Ivanov, P. M. Rojo, Constraints on TESS albedos for five hot Jupiters, *Mon. Not. Roy. Astron. Soc.* 513 (2022) 3444–3457. [doi:10.1093/mnras/stac992](https://doi.org/10.1093/mnras/stac992).
- [100] W. Pluriel, J. Leconte, V. Parmentier, T. Zingales, A. Falco, F. Selsis, P. Borde, Toward a multidimensional analysis of transmission spectroscopy II. Day-night-induced biases in retrievals from hot to ultrahot Jupiters, *Astron. Astrophys.* 658 (2022) A42. [doi:10.1051/0004-6361/202141943](https://doi.org/10.1051/0004-6361/202141943).
- [101] M. C. Nixon, N. Madhusudhan, Aura-3D: A Three-dimensional Atmospheric Retrieval Framework for Exoplanet Transmission Spectra, *Astrophys. J.* 935 (2022) 73. [doi:10.3847/1538-4357/ac7c09](https://doi.org/10.3847/1538-4357/ac7c09).
- [102] P. Spyros, N. K. Nikolov, S. Constantinou, J. Southworth, N. Madhusudhan, E. Sedaghati, D. Ehrenreich, L. Mancini, A precise blue-optical transmission spectrum from the ground: evidence for haze in the atmosphere of WASP-74b, *Mon. Not. Roy. Astron. Soc.* 521 (2023) 2163–2180. [doi:10.1093/mnras/stad637](https://doi.org/10.1093/mnras/stad637).
- [103] L. Bazinet, S. Pelletier, B. Benneke, R. Salinas, G. N. Mace, A Subsolar Metallicity on the Ultra-short-period Planet HIP 65Ab, *Astron. J.* 167 (5) (2024) 206. [doi:10.3847/1538-3881/ad3071](https://doi.org/10.3847/1538-3881/ad3071).
- [104] A. D. Rathcke, L. A. Buchhave, J. a. M. Mendonça, D. K. Sing, M. López-Morales, M. K. Alam, G. W. Henry, N. K. Nikolov, A. García Muñoz, T. Mikal-Evans, H. R. Wakeford, L. A. Dos Santos, V. M. Rajpaul, HST PanCET programme: a flat optical transmission spectrum for the Hot Jupiter WASP-101b, *Mon. Not. Roy. Astron. Soc.* 522 (2023) 582–594. [doi:10.1093/mnras/stad1010](https://doi.org/10.1093/mnras/stad1010).
- [105] F. Wang, Q. Changeat, G. Tinetti, D. Turrini, S. O. M. Wright, Constraining the atmospheric elements in hot Jupiters with Ariel, *Mon. Not. Roy. Astron. Soc.* 523 (2023) 4365–4380. [doi:10.1093/mnras/stad1721](https://doi.org/10.1093/mnras/stad1721).
- [106] P. E. Cubillos, L. Fossati, T. Koskinen, C. Huang, A. G. Sreejith, K. France, P. Wilson Cauley, C. A. Haswell, The Hubble/STIS near-ultraviolet transmission spectrum of HD 189733 b, *Astron. Astrophys.* 671 (2023) A170. [doi:10.1051/0004-6361/202245064](https://doi.org/10.1051/0004-6361/202245064).
- [107] A. Bello-Arufe, H. A. Knutson, J. M. Mendonca, M. M. Zhang, S. H. C. Cabot, A. D. Rathcke, A. Ulla, S. Vissapragada, L. A. Buchhave, Transmission Spectroscopy of the Lowest-density Gas Giant: Metals and a Potential Extended Outflow in HAT-P-67b, *Astron. J.* 166 (2023) 69. [doi:10.3847/1538-3881/acd935](https://doi.org/10.3847/1538-3881/acd935).
- [108] K. L. Chubb, D. M. Stam, C. Helling, D. Samra, L. Carone, Modelling reflected polarized light from close-in giant exoplanet WASP-96b using PolHEEx (Polarization of hot exoplanets), *Mon. Not. Roy. Astron. Soc.* 527 (2023) 4955–4982. [doi:10.1093/mnras/stad3413](https://doi.org/10.1093/mnras/stad3413).
- [109] Z. Rustamkulov, D. K. Sing, S. Mukherjee, E. M. May, J. Kirk, E. Schlawin, M. R. Line, C. Piaulet, A. L. Carter, N. E. Batalha, J. M. Goyal, M. López-Morales, J. D. Lothringer, R. J. MacDonald, S. E. Moran, K. B. Stevenson, H. R. Wakeford, N. Espinoza, J. L. Bean, N. M. Batalha, B. Benneke, Z. K. Berta-Thompson, I. J. M. Crossfield, P. Gao, L. Kreidberg, D. K. Powell, P. E. Cubillos, N. P. Gibson, J. Leconte, K. Molaverdikhani, N. K. Nikolov, V. Parmentier, P. Roy, J. Taylor, J. D. Turner, P. J. Wheatley, K. Aggarwal, E. Ahrer, M. K. Alam, L. Alderson, N. H. Allen, A. Banerjee, S. Barat, D. Barrado, J. K. Barstow, T. J. Bell, J. Blecic, J. Brande, S. Casewell, Q. Changeat, K. L. Chubb, N. Crouzet, T. Daylan, L. Decin, J. Désert, T. Mikal-Evans, A. D. Feinstein, L. Flagg, J. J. Fortney, J. Harrington, K. Heng, Y. Hong, R. Hu, N. Iro, T. Kataria, E. M.-R. Kempton, J. Krick, M. Lendl, J. Lillo-Box, A. Louca, J. Lustig-Yaeger, L. Mancini, M. Mansfield, N. J. Mayne, Y. Miguel, G. Morello, K. Ohno, E. Palle, D. J. M. Petit dit de la Roche, B. V. Rackham, M. Radica, L. Ramos-Rosado, S. Redfield, L. K. Rogers, E. L. Shkolnik, J. Southworth, J. Teske, P. Tremblin, G. S. Tucker, O. Venot, W. C. Waalkes, L. Welbanks, X. Zhang, S. Zieba, Early Release Science of the exoplanet WASP-39b with JWST NIRSpec PRISM, *Nature* 614 (2023) 659–663. [doi:10.1038/s41586-022-05677-y](https://doi.org/10.1038/s41586-022-05677-y).
- [110] A. J. Louca, Y. Miguel, D. Kubyshkina, Metallicity and Spectral Evolution of WASP 39b: The Limited Role of Hydrodynamic Escape, *Astrophys. J. Lett.* 956 (2023) L19. [doi:10.3847/2041-8213/acfaec](https://doi.org/10.3847/2041-8213/acfaec).
- [111] A. Banerjee, J. K. Barstow, C. A. Haswell, S. R. Lewis, Effect of centrifugal force on transmission spectroscopy of exoplanet atmospheres, *Mon. Not. Roy. Astron. Soc.* 523 (2023) L64–L68. [doi:10.1093/mnrasl/slad058](https://doi.org/10.1093/mnrasl/slad058).
- [112] I. J. M. Crossfield, Volatile-to-sulfur Ratios Can Recover a Gas Giant's Accretion History, *Astrophys. J. Lett.* 952 (2023) L18. [doi:10.3847/2041-8213/ace35f](https://doi.org/10.3847/2041-8213/ace35f).
- [113] S. Constantinou, N. Madhusudhan, S. Gandhi, Early Insights for Atmospheric Retrievals of Exoplanets Using JWST Transit Spectroscopy, *Astrophys. J. Lett.* 943 (2023) L10. [doi:10.3847/2041-8213/acaead](https://doi.org/10.3847/2041-8213/acaead).
- [114] A. D. Feinstein, M. Radica, L. Welbanks, C. A. Murray, K. Ohno, L.-P. Coulombe, N. Espinoza, J. L. Bean, J. K. Teske, B. Benneke, M. R. Line, Z. Rustamkulov, A. Saba, A. Tsiaras, J. K. Barstow, J. J. Fortney, P. Gao, H. A. Knutson, R. J. MacDonald, T. Mikal-Evans, B. V. Rackham, J. Taylor, V. Parmentier, N. M. Batalha, Z. K. Berta-Thompson, A. L. Carter, Q. Changeat, L. A. dos Santos, N. P. Gibson, J. M. Goyal, L. Kreidberg, M. López-Morales, J. D. Lothringer, Y. Miguel, K. Molaverdikhani, S. E. Moran, G. Morello, S. Mukherjee, D. K. Sing, K. B. Stevenson, H. R. Wakeford, E.-M. Ahrer, M. K. Alam, L. Alderson, N. H. Allen, N. E. Batalha, T. J. Bell, J. Blecic, J. Brande, C. Caceres, S. L. Casewell, K. L. Chubb, I. J. M. Crossfield, N. Crouzet, P. E. Cubillos, L. Decin, J.-M. Désert, J. Harrington, K. Heng, T. Henning, N. Iro, E. M.-R. Kempton, S. Kendrew, J. Kirk, J. Krick, P.-O. Lagage, M. Lendl, L. Mancini, M. Mansfield, E. M. May, N. J. Mayne, N. K. Nikolov, E. Palle, D. J. M. Petit dit de la Roche, C. Piaulet, D. Powell, S. Redfield, L. K. Rogers, M. T. Roman, P.-A. Roy, M. C. Nixon, E. Schlawin, X. Tan, P. Tremblin, J. D. Turner, O. Venot, W. C. Waalkes, P. J. Wheatley, X. Zhang, Early Release Science of the exoplanet WASP-39b with JWST NIRISS, *Nature* 614 (2023) 670–675. [doi:10.1038/s41586-022-05674-1](https://doi.org/10.1038/s41586-022-05674-1).

- [115] D. Grant, N. K. Lewis, H. R. Wakeford, N. E. Batalha, A. Glidden, J. Goyal, E. Mullens, R. J. MacDonald, E. M. May, S. Seager, K. B. Stevenson, J. A. Valenti, C. Visscher, L. Alderson, N. H. Allen, C. I. C. nas, K. Colón, M. Clampin, N. Espinoza, A. Gressier, J. Huang, Z. Lin, D. Long, D. R. Louie, M. P. na Guerrero, S. Ranjan, K. S. Sotzen, D. Valentine, J. Anderson, W. O. Balmer, A. Bellini, K. K. W. Hoch, J. Kammerer, M. Libralato, C. M. Mountain, M. D. Perrin, L. Pueyo, E. Rickman, I. Rebollido, S. T. Sohn, R. P. van der Marel, L. L. Watkins, JWST-TST DREAMS: Quartz Clouds in the Atmosphere of WASP-17b, *Astrophys. J. Lett.* 956 (2023) L32. [doi:10.3847/2041-8213/acfc3b](https://doi.org/10.3847/2041-8213/acfc3b).
- [116] S. Gandhi, S. de Regt, I. Snellen, Y. Zhang, B. Rutgers, N. van Leur, Q. Bosschaart, JWST Measurements of ^{13}C , ^{18}O , and ^{17}O in the Atmosphere of Super-Jupiter VHS 1256 b, *Astrophys. J. Lett.* 957 (2023) L36. [doi:10.3847/2041-8213/ad07e2](https://doi.org/10.3847/2041-8213/ad07e2).
- [117] J. C. Zapata Trujillo, M. M. Pettyjohn, L. K. McKemmish, High-throughput quantum chemistry: empowering the search for molecular candidates behind unknown spectral signatures in exoplanetary atmospheres, *Mon. Not. Roy. Astron. Soc.* 524 (2023) 361–376. [doi:10.1093/mnras/stad1717](https://doi.org/10.1093/mnras/stad1717).
- [118] L. Welbanks, P. McGill, M. Line, N. Madhusudhan, On the Application of Bayesian Leave-one-out Cross-validation to Exoplanet Atmospheric Analysis, *Astron. J.* 165 (2023) 112. [doi:10.3847/1538-3881/acab67](https://doi.org/10.3847/1538-3881/acab67).
- [119] J. Taylor, V. Parmentier, Another look at the dayside spectra of WASP-43b and HD 209458b: Are there scattering clouds?, *Mon. Not. Roy. Astron. Soc.* 526 (2023) 2133–2140. [doi:10.1093/mnras/stad2287](https://doi.org/10.1093/mnras/stad2287).
- [120] Y.-X. Chen, A. Burrows, A. Sur, R. T. Arevalo, Jupiter atmospheric models and outer boundary conditions for giant planet evolutionary calculations, *Astrophys. J.* 957 (2023) 36. [doi:10.3847/1538-4357/acf456](https://doi.org/10.3847/1538-4357/acf456).
- [121] J. Hayoz, G. Cugno, S. P. Quanz, P. Patapis, E. Alei, M. J. Bonse, F. A. Dannert, E. O. Garvin, T. D. Gebhard, B. S. Konrad, L. F. Sartori, Incorporating medium-resolution spectroscopy of close-in directly imaged exoplanets into atmospheric retrievals via cross-correlation, *Astron. Astrophys.* 678 (2023) 22. [doi:10.1051/0004-6361/202245752](https://doi.org/10.1051/0004-6361/202245752).
- [122] J. Yang, P. G. J. Irwin, J. K. Barstow, Testing 2D temperature models in Bayesian retrievals of atmospheric properties from hot Jupiter phase curves, *Mon. Not. Roy. Astron. Soc.* 525 (2023) 5146–5167. [doi:10.1093/mnras/stad2555](https://doi.org/10.1093/mnras/stad2555).
- [123] C. J. Cheverall, N. Madhusudhan, M. Holmberg, Robustness measures for molecular detections using high-resolution transmission spectroscopy of exoplanets, *Mon. Not. Roy. Astron. Soc.* 522 (2023) 661–677. [doi:10.1093/mnras/stad648](https://doi.org/10.1093/mnras/stad648).
- [124] E.-M. Ahrer, K. B. Stevenson, M. Mansfield, S. E. Moran, J. Brande, G. Morello, C. A. Murray, N. K. Nikolov, D. J. M. P. D. de la Roche, E. Schlawin, P. J. Wheatley, S. Zieba, N. E. Batalha, M. Damiano, J. M. Goyal, M. Lendl, J. D. Lothringer, S. Mukherjee, K. Ohno, N. M. Batalha, M. P. Battley, J. L. Bean, T. G. Beatty, B. Benneke, Z. K. Berta-Thompson, A. L. Carter, P. E. Cubillos, T. Daylan, N. Espinoza, P. Gao, N. P. Gibson, S. Gill, J. Harrington, R. Hu, L. Kreidberg, N. K. Lewis, M. R. Line, M. Lopez-Morales, V. Parmentier, D. K. Powell, D. K. Sing, S.-M. Tsai, H. R. Wakeford, L. Welbanks, M. K. Alam, L. Alderson, N. H. Allen, D. R. Anderson, J. K. Barstow, D. Bayliss, T. J. Bell, J. Blecic, E. M. Bryant, M. R. Burleigh, L. Carone, S. L. Casewell, Q. Changeat, K. L. Chubb, I. J. M. Crossfield, N. Crouzet, L. Decin, J.-M. Desert, A. D. Feinstein, L. Flagg, J. J. Fortney, J. E. Gizis, K. Heng, N. Iro, E. M. R. Kempton, S. Kendrew, J. Kirk, H. A. Knutson, T. D. Komacek, P.-O. Lagage, J. Leconte, J. Lustig-Yaeger, R. J. MacDonald, L. Mancini, E. M. May, N. J. Mayne, Y. Miguel, T. Mikal-Evans, K. Molaverdikhani, E. Palle, C. Piaulet, B. V. Rackham, S. Redfield, L. K. Rogers, P.-A. Roy, Z. Rustamkulov, E. L. Shkolnik, K. S. Sotzen, J. Taylor, P. Tremblin, G. S. Tucker, J. D. Turner, M. de Val-Borro, O. Venot, X. Zhang, Early Release Science of the exoplanet WASP-39b with JWST NIRCam, *Nature* 614 (2023) 653–658. [doi:10.1038/s41586-022-05590-4](https://doi.org/10.1038/s41586-022-05590-4).
- [125] F. Lesjak, L. Nortmann, F. Yan, D. Cont, A. Reiners, N. Piskunov, A. Hatzes, L. Boldt-Christmas, S. Czesla, U. Heiter, O. Kochukhov, A. Lavail, E. Nagel, A. D. Rains, M. Rengel, F. Rodler, U. Seemann, D. Shulyak, Retrieval of the dayside atmosphere of WASP-43b with CRIRES⁺, *Astron. Astrophys.* 678 (2023) A23. [doi:10.1051/0004-6361/202347151](https://doi.org/10.1051/0004-6361/202347151).
- [126] B. Edwards, Q. Changeat, Measuring Tracers of Planet Formation in the Atmosphere of WASP-77A b: Substellar O/H and C/H Ratios, with a Stellar C/O Ratio and a Potentially Superstellar Ti/H Ratio, *Astrophys. J. Lett.* 962. [doi:10.3847/2041-8213/ad2000](https://doi.org/10.3847/2041-8213/ad2000).
- [127] P. C. B. Smith, M. R. Line, J. L. Bean, M. Brogi, P. August, L. Welbanks, J.-M. Desert, J. Lunine, J. Sanchez, M. Mansfield, L. Pino, E. Rauscher, E. Kempton, J. Zalesky, M. Fowler, A Combined Ground-based and JWST Atmospheric Retrieval Analysis: Both IGRINS and NIRSpec Agree that the Atmosphere of WASP-77A b Is Metal-poor, *Astron. J.* 167 (2024) 110. [doi:10.3847/1538-3881/ad17bf](https://doi.org/10.3847/1538-3881/ad17bf).
- [128] A. D. Schneider, P. Mollière, G. Louppe, L. Carone, U. G. Jørgensen, L. Decin, C. Helling, Harnessing machine learning for accurate treatment of overlapping opacity species in general circulation models, *Astron. Astrophys.* 682 (2024) A79. [doi:10.1051/0004-6361/202348221](https://doi.org/10.1051/0004-6361/202348221).
- [129] T. J. Bell, N. Crouzet, P. E. Cubillos, L. Kreidberg, A. A. A. Piette, M. T. Roman, J. K. Barstow, J. Blecic, L. Carone, L.-P. Coulombe, E. Ducrot, M. Hammond, J. ao M. Mendonça, J. I. Moses, V. Parmentier, K. B. Stevenson, L. Teinturier, M. Zhang, N. M. Batalha, J. L. Bean, B. Benneke, B. Charnay, K. L. Chubb, B.-O. Demory, P. Gao, E. K. H. Lee, M. López-Morales, G. Morello, E. Rauscher, D. K. Sing, X. Tan, O. Venot, H. R. Wakeford, K. Aggarwal, E.-M. Ahrer, M. K. Alam, R. Baeyens, D. Barrado, C. Caceres, A. L. Carter, S. L. Casewell, R. C. Challener, I. J. M. Crossfield, L. Decin, J.-M. Desert, I. Dobbs-Dixon, A. Dyrek, N. Espinoza, A. D. Feinstein, N. P. Gibson, J. Harrington, C. Helling, R. Hu, N. Iro, E. M. R. Kempton, S. Kendrew, T. D. Komacek, J. Krick, P.-O. Lagage, J. Leconte, M. Lendl, N. T. Lewis, J. D. Lothringer, I. Malsky, L. Mancini, M. Mansfield, N. J. Mayne, T. Mikal-Evans, K. Molaverdikhani, N. K. Nikolov, M. C. Nixon, E. Palle, D. J. M. P. dit de la Roche, C. Piaulet, D. Powell, B. V. Rackham, A. D. Schneider, M. E. Steinrueck, J. Taylor, L. Welbanks, S. N. Yurchenko, X. Zhang, S. Zieba, Nightside clouds and disequilibrium chemistry on the hot Jupiter WASP-43b, *arXiv e-prints* [arXiv:2401.13027](https://arxiv.org/abs/2401.13027).
- [130] D. Powell, A. D. Feinstein, E. K. H. Lee, M. Zhang, S.-M. Tsai, J. Taylor, J. Kirk, T. Bell, J. K. Barstow, P. Gao, J. L. Bean, J. Blecic, K. L. Chubb, I. J. M. Crossfield, S. Jordan, D. Kitzmann, S. E. Moran, G. Morello, J. I. Moses, L. Welbanks, J. Yang, X. Zhang, E.-M. Ahrer, A. Bello-Arufe, J. Brande, S. L. Casewell, N. Crouzet, P. E. Cubillos, B.-O. Demory, A. Dyrek, L. Flagg, R. Hu, J. Inglis, K. D. Jones, L. Kreidberg, M. López-Morales, P.-O. Lagage, E. A. Meier Valdés, Y. Miguel, V. Parmentier, A. A. A. Piette, B. V. Rackham, M. Radica, S. Redfield, K. B. Stevenson, H. R. Wakeford, K. Aggarwal, M. K. Alam, N. M. Batalha, N. E. Batalha, B. Benneke, Z. K. Berta-Thompson, R. P. Brady, C. Caceres, A. L. Carter, J.-M. Desert, J. Harrington, N. Iro, M. R. Line, J. D. Lothringer, R. J. MacDonald, L. Mancini, K. Molaverdikhani, S. Mukherjee, M. C. Nixon, A. V. Oza, E. Palle, Z. Rustamkulov, D. K. Sing, M. E. Steinrueck, O. Venot, P. J. Wheatley, S. N. Yurchenko, Sulphur dioxide in the mid-infrared transmission spectrum of WASP-39b, *Nature* 626 (2024) 979–983. [doi:10.1038/s41586-024-07040-9](https://doi.org/10.1038/s41586-024-07040-9).
- [131] K. H. Yip, Q. Changeat, A. Al-Refaie, I. P. Waldmann, To Sample or Not to Sample: Retrieving Exoplanetary Spectra with Variational Inference and Normalizing Flows, *Astrophys. J.* 961 (2024) 30. [doi:10.3847/1538-4357/ad063f](https://doi.org/10.3847/1538-4357/ad063f).
- [132] R. Landman, T. Stolker, I. A. G. Snellen, J. Costes, S. de Regt, Y. Zhang, S. Gandhi, P. Molliere, A. Kesseli, A. Vigan, A. Sanchez-Lopez, β pictoris b through the eyes of the upgraded cries plus atmospheric composition, spin rotation, and radial velocity, *Astron. Astrophys.* 682 (2024) A48. [doi:10.1051/0004-6361/202347846](https://doi.org/10.1051/0004-6361/202347846).
- [133] J. Wilson, N. P. Gibson, N. Nikolov, S. Constantinou, N. Madhusudhan, J. Goyal, J. K. Barstow, A. L. Carter, E. J. W. de Mooij, B. Drum-

- mond, T. Mikal-Evans, C. Helling, N. J. Mayne, D. K. Sing, Ground-based transmission spectroscopy with FORS2: A featureless optical transmission spectrum and detection of H₂O for the ultra-hot Jupiter WASP-103b, *Mon. Not. Roy. Astron. Soc.* 497 (2020) 5155–5170. [doi:10.1093/mnras/staa2307](https://doi.org/10.1093/mnras/staa2307).
- [134] S. Gandhi, A. S. Jermyn, Coupled day-night models of exoplanetary atmospheres, *Mon. Not. Roy. Astron. Soc.* 499 (2020) 4984–5003. [doi:10.1093/mnras/staa3143](https://doi.org/10.1093/mnras/staa3143).
- [135] H. M. Tabernero, M. R. Zapatero Osorio, R. Allart, F. Borsa, N. Casasayas-Barris, O. Demangeon, D. Ehrenreich, J. Lillo-Box, C. Lovis, E. Palle, S. G. Sousa, R. Rebolo, N. C. Santos, F. Pepe, S. Cristiani, V. Adibekyan, C. Allende Prieto, Y. Alibert, S. C. C. Barros, F. Bouchy, V. Bourrier, V. D'Odorico, X. Dumusque, J. P. Faria, P. Figueira, R. Genova Santos, J. Gonzalez Hernandez, I. S. Hojjatpanah, G. Lo Curto, B. Lavie, C. J. A. P. Martins, J. H. C. Martins, A. Mehner, G. Micela, P. Molaro, N. J. Nunes, E. Poretti, J. Seidel, V. A. Sozzetti, A. Suarez Mascareno, S. Udry, M. Aliverti, M. Affolter, D. Alves, M. Amate, G. Avila, T. Bandy, W. Benz, A. Bianco, C. Broeg, A. Cabral, P. Conconi, J. Coelho, C. Cumani, S. Deiries, H. Dekker, B. Delabre, A. Fragoso, M. Genoni, L. Genolet, I. Hughes, J. Knudstrup, F. Kerber, M. Landoni, J. L. Lizon, C. Maire, A. Manescu, P. Di Marcantonio, D. Megevand, M. Monteiro, M. Moschetti, E. Mueller, A. Modigliani, L. Oggioni, A. Oliveira, G. Pariani, L. Pasquini, J. L. Rasilla, E. Redaelli, M. Riva, S. Santana-Tschudi, P. Santin, P. Santos, A. Segovia, D. Sosnowska, P. Spano, F. Tenegi, O. Iwert, A. Zanutta, F. Zerbi, ESPRESSO high-resolution transmission spectroscopy of WASP-76 b, *Astron. Astrophys.* 646. [doi:10.1051/0004-6361/202039511](https://doi.org/10.1051/0004-6361/202039511).
- [136] E. M. May, T. D. Komacek, K. B. Stevenson, E. M.-R. Kempton, J. L. Bean, M. Malik, J. Ih, M. Mansfield, A. B. Savel, D. Deming, J.-M. Desert, Y. K. Feng, J. J. Fortney, T. Kataria, N. Lewis, C. Morley, E. Rauscher, A. Showman, Spitzer Phase-curve Observations and Circulation Models of the Inflated Ultrahot Jupiter WASP-76b, *Astron. J.* 162 (2021) 158. [doi:10.3847/1538-3881/ac0e30](https://doi.org/10.3847/1538-3881/ac0e30).
- [137] E. Gharib-Nezhad, A. R. Iyer, M. R. Line, R. S. Freedman, M. S. Marley, N. E. Batalha, EXOPLANES: Molecular Absorption Cross-section Database for Brown Dwarf and Giant Exoplanet Atmospheres, *Astrophys. J. Suppl. Ser.* 254 (2021) 34. [doi:10.3847/1538-4365/abf504](https://doi.org/10.3847/1538-4365/abf504).
- [138] M. Stangret, E. Pallé, N. Casasayas-Barris, M. Oshagh, A. Bello-Arufe, R. Luque, V. Nascimbeni, F. Yan, J. Orell-Miquel, D. Sicilia, L. Malavolta, B. C. Addison, L. A. Buchhave, A. S. Bonomo, F. Borsa, S. H. C. Cabot, M. Cecconi, D. A. Fischer, A. Harutyunyan, J. M. Mendonça, G. Nowak, H. Parviainen, A. Sozzetti, R. Tronsgaard, The obliquity and atmosphere of the ultra-hot Jupiter TOI-1431b (MASCARA-5b): A misaligned orbit and no signs of atomic or molecular absorptions, *Astron. Astrophys.* 654. [doi:10.1051/0004-6361/202040100](https://doi.org/10.1051/0004-6361/202040100).
- [139] D. Kasper, J. L. Bean, M. R. Line, A. Seifahrt, J. Stürmer, L. Pino, J.-M. Désert, M. Brogi, Confirmation of Iron Emission Lines and Nondetection of TiO on the Dayside of KELT-9b with MAROON-X, *Astrophys. J. Lett.* 921 (1) (2021) L18. [doi:10.3847/2041-8213/ac30e1](https://doi.org/10.3847/2041-8213/ac30e1).
- [140] A. Roth, B. Drummond, E. Hebrard, P. Tremblin, J. Goyal, N. Mayne, Pseudo-2D modelling of heat redistribution through H₂ thermal dissociation/recombination: consequences for ultra-hot Jupiters, *Mon. Not. Roy. Astron. Soc.* 505 (2021) 4515–4530. [doi:10.1093/mnras/stab1256](https://doi.org/10.1093/mnras/stab1256).
- [141] D. B. Serindag, S. K. Nugroho, P. Mollière, E. J. W. de Mooij, N. P. Gibson, I. A. G. Snellen, Is TiO emission present in the ultra-hot Jupiter WASP-33b? A reassessment using the improved ExoMol TOTO line list, *Astron. Astrophys.* 645 (2021) A90. [doi:10.1051/0004-6361/202039135](https://doi.org/10.1051/0004-6361/202039135).
- [142] J. Kirk, B. V. Rackham, R. J. MacDonald, M. López-Morales, N. Espinoza, M. Lendl, J. Wilson, D. J. Osip, P. J. Wheatley, I. Skillen, D. Apai, A. Bixel, N. P. Gibson, A. Jordán, N. K. Lewis, T. Louden, C. D. McGruder, N. Nikolov, F. Rodler, I. C. Weaver, ACCESS and LRG-BEASTS: A Precise New Optical Transmission Spectrum of the Ultrahot Jupiter WASP-103b, *Astron. J.* 162 (2021) 34. [doi:10.3847/1538-3881/abfcdd](https://doi.org/10.3847/1538-3881/abfcdd).
- [143] Q. Changeat, B. Edwards, The hubble WFC3 emission spectrum of the extremely hot jupiter KELT-9b, *Astrophys. J. Lett.* 907 (2021) L22. [doi:10.3847/2041-8213/abd84f](https://doi.org/10.3847/2041-8213/abd84f).
- [144] E. K. H. Lee, B. Prinith, D. Kitzmann, S.-M. Tsai, J. Hoeijmakers, N. W. Borsato, K. Heng, The Mantis Network II: examining the 3D high-resolution observable properties of the UHJs WASP-121b and WASP-189b through GCM modelling, *Mon. Not. Roy. Astron. Soc.* 517 (2022) 240–256. [doi:10.1093/mnras/stac2246](https://doi.org/10.1093/mnras/stac2246).
- [145] B. Prinith, H. J. Hoeijmakers, D. Kitzmann, E. Sandvik, J. Seidel, V. M. Lendl, N. W. Borsato, B. Thorsbro, D. R. Anderson, D. Barrado, K. Kravchenko, R. Allart, V. Bourrier, H. M. Cegla, D. Ehrenreich, C. Fisher, C. Lovis, A. Guzman-Mesa, S. Grimm, M. Hooton, B. M. Morris, M. Oreshenko, L. Pino, K. Heng, Titanium oxide and chemical inhomogeneity in the atmosphere of the exoplanet WASP-189 b, *Nature Astr.* 6 (2022) 449–457. [doi:10.1038/s41550-021-01581-z](https://doi.org/10.1038/s41550-021-01581-z).
- [146] T. Mikal-Evans, D. K. Sing, J. K. Barstow, T. Kataria, J. Goyal, N. Lewis, J. Taylor, N. J. Mayne, T. Daylan, H. R. Wakeford, M. S. Marley, J. J. Spake, Diurnal variations in the stratosphere of the ultrahot giant exoplanet WASP-121b, *Nat. Astron.* 6 (2022) 471 – 479. [doi:10.1038/s41550-021-01592-w](https://doi.org/10.1038/s41550-021-01592-w).
- [147] A. B. Savel, E. M.-R. Kempton, M. Malik, T. D. Komacek, J. L. Bean, E. M. May, K. B. Stevenson, M. Mansfield, E. Rauscher, No Umbrella Needed: Confronting the Hypothesis of Iron Rain on WASP-76b with Post-processed General Circulation Models, *Astron. J.* 926 (2022) 85. [doi:10.3847/1538-4357/ac423f](https://doi.org/10.3847/1538-4357/ac423f).
- [148] A. Bello-Arufe, S. H. C. Cabot, J. M. Mendonca, L. A. Buchhave, A. D. Rathcke, Mining the Ultrahot Skies of HAT-P-70b: Detection of a Profusion of Neutral and Ionized Species, *Astron. J.* 163 (2022) 96. [doi:10.3847/1538-3881/ac402e](https://doi.org/10.3847/1538-3881/ac402e).
- [149] Q. Changeat, On Spectroscopic Phase-curve Retrievals: H₂ Dissociation and Thermal Inversion in the Atmosphere of the Ultrahot Jupiter WASP-103 b, *Astron. J.* 163 (2022) 106. [doi:10.3847/1538-3881/ac4475](https://doi.org/10.3847/1538-3881/ac4475).
- [150] M. Holmberg, N. Madhusudhan, A First Look at CRIRES+: Performance Assessment and Exoplanet Spectroscopy, *Astron. J.* 164 (2022) 79. [doi:10.3847/1538-3881/ac77eb](https://doi.org/10.3847/1538-3881/ac77eb).
- [151] L. Finnerty, T. Schofield, B. Sappey, J. W. Xuan, J.-B. Ruffio, J. J. Wang, J.-R. Delorme, G. A. Blake, C. Buzard, M. P. Fitzgerald, A. Baker, R. Bartos, C. Z. Bond, B. Calvin, S. Cetre, G. Doppmann, D. Echeverri, N. Jovanovic, J. Liberman, R. A. López, E. C. Martin, D. Mawet, E. Morris, J. Pezzato, C. L. Phillips, S. Ragland, A. Skemer, T. Venenciano, J. K. Wallace, N. L. Wallack, J. Wang, P. Wizinowich, Keck Planet Imager and Characterizer Emission Spectroscopy of WASP-33b, *Astron. J.* 166 (2023) 31. [doi:10.3847/1538-3881/acda91](https://doi.org/10.3847/1538-3881/acda91).
- [152] Y. Yang, G. Chen, S. Wang, F. Yan, High-resolution Transmission Spectroscopy of Ultrahot Jupiter WASP-33b with NEID, *Astron. J.* 167 (2023) 36. [doi:10.3847/1538-3881/ad10a3](https://doi.org/10.3847/1538-3881/ad10a3).
- [153] N. Lowson, G. Zhou, D. J. J. Wright, C. X. X. Huang, J. M. Mendonca, S. H. C. Cabot, C. Pudmenzky, R. A. A. Wittenmyer, D. W. W. Latham, A. Bieryla, G. A. A. Esquerdo, P. Berlind, M. L. L. Calkins, Multiphase Detections of the Extended Atmosphere and Transmission Spectra of KELT-9b with a 1.5 m Telescope, *Astron. J.* 165. [doi:10.3847/1538-3881/acacf3](https://doi.org/10.3847/1538-3881/acacf3).
- [154] J. P. Wardenier, V. Parmentier, M. R. Line, E. K. H. Lee, Modelling the effect of 3D temperature and chemistry on the cross-correlation

- signal of transiting ultra-hot Jupiters: a study of five chemical species on WASP-76b, *Mon. Not. Roy. Astron. Soc.* 525 (2023) 4942–4961. [doi:10.1093/mnras/stad2586](https://doi.org/10.1093/mnras/stad2586).
- [155] L.-P. Coulombe, B. Benneke, R. Challener, A. A. A. Piette, L. S. Wiser, M. Mansfield, R. J. MacDonald, H. Beltz, A. D. Feinstein, M. Radica, A. B. Savel, L. A. Dos Santos, J. L. Bean, V. Parmentier, I. Wong, E. Rauscher, T. D. Komacek, E. M. R. Kempton, X. Tan, M. Hammond, N. T. Lewis, M. R. Line, E. K. H. Lee, H. Shrivikumar, I. J. M. Crossfield, M. C. Nixon, B. V. Rackham, H. R. Wakeford, L. Welbanks, X. Zhang, N. M. Batalha, Z. K. Berta-Thompson, Q. Changeat, J.-M. Desert, N. Espinoza, J. M. Goyal, J. Harrington, H. A. Knutson, L. Kreidberg, M. Lopez-Morales, A. Shporer, D. K. Sing, K. B. Stevenson, K. Aggarwal, E.-M. Ahrer, M. K. Alam, T. J. Bell, J. Blecic, C. Caceres, A. L. Carter, S. L. Casewell, N. Crouzet, P. E. Cubillos, L. Decin, J. J. Fortney, N. P. Gibson, K. Heng, T. Henning, N. Iro, S. Kendrew, P.-O. Lagage, J. Leconte, M. Lendl, J. D. Lothringer, L. Mancini, T. Mikal-Evans, K. Molaverdikhani, N. K. Nikolov, K. Ohno, E. Palle, C. Piaulet, S. Redfield, P.-A. Roy, S.-M. Tsai, O. Venot, P. J. Wheatley, A broadband thermal emission spectrum of the ultra-hot Jupiter WASP-18b, *Nature* 620 (2023) 292–298. [doi:10.1038/s41586-023-06230-1](https://doi.org/10.1038/s41586-023-06230-1).
- [156] M. C. Johnson, J. Wang, A. P. Asnodkar, A. S. Bonomo, B. S. Gaudi, T. Henning, I. Ilyin, E. Keles, L. Malavolta, M. Mallonn, K. Molaverdikhani, V. Nascimbeni, J. Patience, K. Poppenhaeger, G. Scandariato, E. Schlawin, E. Shkolnik, D. Sicilia, A. Sozzetti, K. G. Strassmeier, C. Veillet, F. Yan, The PEPSI Exoplanet Transit Survey (PETS). II. A Deep Search for Thermal Inversion Agents in KELT-20 b/MASCARA-2 b with Emission and Transmission Spectroscopy, *Astron. J.* 165. [doi:10.3847/1538-3881/acb7e2](https://doi.org/10.3847/1538-3881/acb7e2).
- [157] S. Petz, M. C. Johnson, A. P. Asnodkar, J. Wang, B. S. Gaudi, T. Henning, E. Keles, K. Molaverdikhani, K. Poppenhaeger, G. Scandariato, E. K. Shkolnik, D. Sicilia, K. G. Strassmeier, F. Yan, The PEPSI Exoplanet Transit Survey (PETS) - IV. Assessing the atmospheric chemistry of KELT-20b, *Mon. Not. Roy. Astron. Soc.* 527 (2023) 7079–7092. [doi:10.1093/mnras/stad3481](https://doi.org/10.1093/mnras/stad3481).
- [158] D. Kitzmann, H. J. Hoeijmakers, S. L. Grimm, N. W. Borsato, A. Lueber, B. Prinoth, The Mantis network I. A standard grid of templates and masks for cross-correlation analyses of ultra-hot Jupiter transmission spectra, *Astron. Astrophys.* 669 (2023) A113. [doi:10.1051/0004-6361/202142969](https://doi.org/10.1051/0004-6361/202142969).
- [159] S. Pelletier, B. Benneke, M. Ali-Dib, B. Prinoth, D. Kasper, A. Seifahrt, J. L. Bean, F. Debras, B. Klein, L. Bazinet, H. J. Hoeijmakers, A. Y. Kesseli, O. Lim, A. Carmona, L. Pino, N. Casasayas-Barris, T. Hood, J. Stürmer, Vanadium oxide and a sharp onset of cold-trapping on a giant exoplanet, *Nature* 619 (2023) 491–494. [doi:10.1038/s41586-023-06134-0](https://doi.org/10.1038/s41586-023-06134-0).
- [160] Y. Shi, W. Wang, G. Zhao, M. Zhai, G. Chen, Z. Jiang, Q. Ouyang, T. Henning, J. Zhao, N. Crouzet, R. van Boekel, L. Esteves, Thermal emission from the hot Jupiter WASP-103 b in J and Ks bands, *Mon. Not. Roy. Astron. Soc.* 522 (2023) 1491–1503. [doi:10.1093/mnras/stad891](https://doi.org/10.1093/mnras/stad891).
- [161] A. Ridden-Harper, E. de Mooij, R. Jayawardhana, N. Gibson, R. Karjalainen, M. Karjalainen, High-resolution Emission Spectroscopy of the Ultrahot Jupiter KELT-9b: Little Variation in Day- and Nightside Emission Line Contrasts, *Astron. J.* 165 (5) (2023) 211. [doi:10.3847/1538-3881/acc654](https://doi.org/10.3847/1538-3881/acc654).
- [162] S. Gandhi, A. Kesseli, Y. Zhang, A. Louca, I. Snellen, M. Brogi, Y. Miguel, N. Casasayas-Barris, S. Pelletier, R. Landman, C. Maguire, N. P. Gibson, Retrieval Survey of Metals in Six Ultrahot Jupiters: Trends in Chemistry, Rain-out, Ionization, and Atmospheric Dynamics, *Astron. J.* 165 (6) (2023) 242. [doi:10.3847/1538-3881/accd65](https://doi.org/10.3847/1538-3881/accd65).
- [163] M. Brogi, V. Emeka-Okafor, M. R. Line, S. Gandhi, L. Pino, E. M.-R. Kempton, E. Rauscher, V. Parmentier, J. L. Bean, G. N. Mace, N. B. Cowan, E. Shkolnik, J. P. Wardenier, M. Mansfield, L. Welbanks, P. Smith, J. J. Fortney, J. L. Birkby, J. A. Zalesky, L. Dang, J. Patience, J.-M. Désert, The Roasting Marshmallows Program with IGRINS on Gemini South I: Composition and Climate of the Ultrahot Jupiter WASP-18 b, *Astron. J.* 165 (2023) 91. [doi:10.3847/1538-3881/acaf5c](https://doi.org/10.3847/1538-3881/acaf5c).
- [164] B. Prinoth, H. J. Hoeijmakers, S. Pelletier, D. Kitzmann, B. M. Morris, A. Seifahrt, D. Kasper, H. H. Korhonen, M. Burheim, J. L. Bean, B. Benneke, N. W. Borsato, M. Brady, S. L. Grimm, R. Luque, J. Stuermer, B. Thorsbro, Time-resolved transmission spectroscopy of the ultra-hot Jupiter WASP-189 b, *Astron. Astrophys.* 678. [doi:10.1051/0004-6361/202347262](https://doi.org/10.1051/0004-6361/202347262).
- [165] L. van Sluijs, J. L. Birkby, J. Lothringer, E. K. H. Lee, I. J. M. Crossfield, V. Parmentier, M. Brogi, C. Kulesa, D. McCarthy, D. Charbonneau, Carbon monoxide emission lines reveal an inverted atmosphere in the ultra hot Jupiter WASP-33 b consistent with an eastward hot spot, *Mon. Not. Roy. Astron. Soc.* 522 (2023) 2145–2170. [doi:10.1093/mnras/stad1103](https://doi.org/10.1093/mnras/stad1103).
- [166] Q. Changeat, J. W. Skinner, J. y.-k. Cho, J. Nättilä, I. P. Waldmann, A. F. Al-Refaie, A. Dyrek, B. Edwards, T. Mikal-Evans, M. Joshua, G. Morello, N. Skaf, A. Tsiaras, O. Venot, K. H. Yip, Is the Atmosphere of the Ultra-hot Jupiter WASP-121 b Variable?, *Astrophys. J. Suppl. Ser.* 270. [doi:10.3847/1538-4365/ad1191](https://doi.org/10.3847/1538-4365/ad1191).
- [167] L. Finnerty, J. W. Xuan, Y. Xin, J. Liberman, T. Schofield, M. P. Fitzgerald, S. Agrawal, A. Baker, R. Bartos, G. A. Blake, B. Calvin, S. Cetre, J.-R. Delorme, G. Doppmann, D. Echeverri, C.-C. Hsu, N. Jovanovic, R. A. López, E. C. Martin, D. Mawet, E. Morris, J. Pezzato, J.-B. Ruffio, B. Sappey, A. Skemer, T. Venenciano, J. K. Wallace, N. L. Wallack, J. J. Wang, J. Wang, Atmospheric Metallicity and C/O of HD 189733 b from High-resolution Spectroscopy, *Astron. J.* 167 (2024) 43. [doi:10.3847/1538-3881/ad1180](https://doi.org/10.3847/1538-3881/ad1180).
- [168] C. Maguire, N. P. Gibson, S. K. Nugroho, M. Fortune, S. Ramkumar, S. Gandhi, E. de Mooij, High-resolution atmospheric retrievals of wasp-76b transmission spectroscopy with espresso: Monitoring limb asymmetries across multiple transits, *arXiv e-prints*[arXiv:2404.10463](https://arxiv.org/abs/2404.10463).
- [169] B. Edwards, Q. Changeat, M. Mori, L. O. Anisman, M. Morvan, K. H. Yip, A. Tsias, A. Al-Refaie, I. Waldmann, G. Tinetti, Hubble WFC3 Spectroscopy of the Habitable-zone Super-Earth LHS 1140 b, *Astron. J.* 161 (2020) 44. [doi:10.3847/1538-3881/abc6a5](https://doi.org/10.3847/1538-3881/abc6a5).
- [170] A. A. A. Piette, N. Madhusudhan, A. M. Mandell, HyDRO: atmospheric retrieval of rocky exoplanets in thermal emission, *Mon. Not. Roy. Astron. Soc.* 511 (2021) 2565–2584. [doi:10.1093/mnras/stab3612](https://doi.org/10.1093/mnras/stab3612).
- [171] L. V. Mugnai, D. Modirrousta-Galian, B. Edwards, Q. Changeat, J. Bouwman, G. Morello, A. Al-Refaie, R. Baeyens, M. F. Bieger, D. Blain, A. Gressier, G. Guilluy, Y. Jaziri, F. Kiefer, M. Morvan, W. Pluriel, M. Poveda, N. Skaf, N. Whiteford, S. Wright, K. H. Yip, T. Zingales, B. Charnay, P. Drossart, J. Leconte, O. Venot, I. Waldmann, J.-P. Beaulieu, ARES. V. No Evidence For Molecular Absorption in the HST WFC3 Spectrum of GJ 1132 b, *Astron. J.* 161 (2021) 284. [doi:10.3847/1538-3881/abf3c3](https://doi.org/10.3847/1538-3881/abf3c3).
- [172] A. Gressier, M. Mori, Q. Changeat, B. Edwards, J. P. Beaulieu, E. Marcq, B. Charnay, Near-infrared transmission spectrum of TRAPPIST-1 h using Hubble WFC3 G141 observations, *Astron. Astrophys.* 658. [doi:10.1051/0004-6361/202142140](https://doi.org/10.1051/0004-6361/202142140).
- [173] E. Alei, B. S. Konrad, D. Angerhausen, J. L. Grenfell, P. Molliere, S. P. Quanz, S. Rugheimer, F. Wunderlich, L. Collaboration, Large Interferometer For Exoplanets (LIFE) V. Diagnostic potential of a mid-infrared space interferometer for studying Earth analogs, *Astron. Astrophys.* 665. [doi:10.1051/0004-6361/202243760](https://doi.org/10.1051/0004-6361/202243760).
- [174] D. J. Bower, K. Hakim, P. A. Sossi, P. Sanan, Retention of Water in Terrestrial Magma Oceans and Carbon-rich Early Atmospheres, *Planet. Sci. J.* 3 (2022) 93. [doi:10.3847/PSJ/ac5fb1](https://doi.org/10.3847/PSJ/ac5fb1).

- [175] A. B. Claringbold, P. B. Rimmer, S. Rugheimer, O. Shorttle, Prebiosignature Molecules Can Be Detected in Temperate Exoplanet Atmospheres with JWST, *Astron. J.* 166. [doi:10.3847/1538-3881/acdacc](https://doi.org/10.3847/1538-3881/acdacc).
- [176] H. Diamond-Lowe, J. ao M. Mendonça, D. Charbonneau, L. A. Buchhave, Ground-based Optical Transmission Spectroscopy of the Nearby Terrestrial Exoplanet LTT 1445Ab, *Astron. J.* 165 (2023) 169. [doi:10.3847/1538-3881/acbf39](https://doi.org/10.3847/1538-3881/acbf39).
- [177] L. Zhou, B. Ma, Y.-H. Wang, Y.-N. Zhu, Hubble WFC3 Spectroscopy of the Terrestrial Planets L 98-59 c and d: No Evidence for a Clear Hydrogen Dominated Primary Atmosphere, *Res. Astron. Astrophys.* 23 (2023) 025011. [doi:10.1088/1674-4527/acaceb](https://doi.org/10.1088/1674-4527/acaceb).
- [178] C. L. Phillips, J. Wang, B. Edwards, R. R. Martínez, A. P. Asnodkar, B. S. Gaudi, Exploring the potential of Twinkle to unveil the nature of LTT 1445 Ab, *Mon. Not. Roy. Astron. Soc.* 526 (2023) 2251–2264. [doi:10.1093/mnras/stad2822](https://doi.org/10.1093/mnras/stad2822).
- [179] S. E. Moran, K. B. Stevenson, D. K. Sing, R. J. MacDonald, J. Kirk, J. Lustig-Yaeger, S. Peacock, L. C. Mayorga, K. A. Bennett, M. López-Morales, E. M. May, Z. Rustamkulov, J. A. Valenti, J. I. A. Redai, M. K. Alam, N. E. Batalha, G. Fu, J. Gonzalez-Quiles, A. N. Highland, E. Kruse, J. D. Lothringer, K. N. O. Ceballos, K. S. Sotzen, H. R. Wakeford, High Tide or Riptide on the Cosmic Shoreline? A Water-rich Atmosphere or Stellar Contamination for the Warm Super-Earth GJ 486b from JWST Observations, *Astrophys. J. Lett.* 948 (2023) L11. [doi:10.3847/2041-8213/accb9c](https://doi.org/10.3847/2041-8213/accb9c).
- [180] L. Zhou, B. Ma, Y. Wang, Y. Zhu, Hubble WFC3 Spectroscopy of the Rocky Planet L 98-59 b: No Evidence for a Cloud-free Primordial Atmosphere, *Astron. J.* 164. [doi:10.3847/1538-3881/ac8fe9](https://doi.org/10.3847/1538-3881/ac8fe9).
- [181] M. Zilinskas, C. P. A. van Buchem, Y. Miguel, A. Louca, R. Lupu, S. Zieba, W. van Westrenen, Observability of evaporating lava worlds star, *Astron. Astrophys.* 661 (2022) A126. [doi:10.1051/0004-6361/202142984](https://doi.org/10.1051/0004-6361/202142984).
- [182] E. A. Whittaker, M. Malik, J. Ih, E. M.-R. Kempton, M. Mansfield, J. L. Bean, E. S. Kite, D. D. B. Koll, T. W. Cronin, R. Hu, The Detectability of Rocky Planet Surface and Atmosphere Composition with the JWST: The Case of LHS 3844b, *Astron. J.* 164 (2022) 258. [doi:10.3847/1538-3881/ac9ab3](https://doi.org/10.3847/1538-3881/ac9ab3).
- [183] I. J. M. Crossfield, M. Malik, M. L. Hill, S. R. Kane, B. Foley, A. S. Polanski, D. Coria, J. Brande, Y. Zhang, K. Wienke, L. Kreidberg, N. B. Cowan, D. Dragomir, V. Gorjian, T. Mikal-Evans, B. Benneke, J. L. Christiansen, D. Deming, F. Y. Morales, GJ 1252b: A Hot Terrestrial Super-Earth with No Atmosphere, *Astrophys. J. Lett.* 937. [doi:10.3847/2041-8213/ac886b](https://doi.org/10.3847/2041-8213/ac886b).
- [184] S. Zieba, M. Zilinskas, L. Kreidberg, T. G. Nguyen, Y. Miguel, N. B. Cowan, R. Pierrehumbert, L. Carone, L. Dang, M. Hammond, T. Louden, R. Lupu, L. Malavolta, K. B. Stevenson, K2 and Spitzer phase curves of the rocky ultra-short-period planet k2-141 b hint at a tenuous rock vapor atmosphere, *Astron. Astrophys.* 664. [doi:10.1051/0004-6361/202142912](https://doi.org/10.1051/0004-6361/202142912).
- [185] L. J. Janssen, P. Woitke, O. Herbort, M. Min, K. L. Chubb, C. Hellings, L. Carone, The sulfur species in hot rocky exoplanet atmospheres, *ASTRONOMISCHE NACHRICHTEN* 344. [doi:10.1002/asna.20230075](https://doi.org/10.1002/asna.20230075).
- [186] A. A. A. Piette, P. Gao, K. Brugman, A. Shahar, T. Lichtenberg, F. Miozzi, P. Driscoll, Rocky Planet or Water World? Observability of Low-density Lava World Atmospheres, *Astrophys. J.* 954. [doi:10.3847/1538-4357/acdef2](https://doi.org/10.3847/1538-4357/acdef2).
- [187] K. Heng, The Transient Outgassed Atmosphere of 55 Cancri e, *Astrophys. J. Lett.* 956 (2023) L20. [doi:10.3847/2041-8213/acfe05](https://doi.org/10.3847/2041-8213/acfe05).
- [188] A. Ridder-Harper, S. K. Nugroho, L. Flagg, R. Jayawardhana, J. D. Turner, E. de Mooij, R. MacDonald, E. Deibert, M. Tamura, T. Kotani, T. Hirano, M. Kuzuhara, M. Omiya, N. Kusakabe, High-resolution Transmission Spectroscopy of the Terrestrial Exoplanet GJ 486b, *Astron. J.* 165 (2023) 170. [doi:10.3847/1538-3881/acbd39](https://doi.org/10.3847/1538-3881/acbd39).
- [189] M. Zilinskas, Y. Miguel, C. P. A. van Buchem, I. A. G. Snellen, Observability of silicates in volatile atmospheres of super-Earths and sub-Neptunes Exploring the edge of the evaporation desert, *Astron. Astrophys.* 671. [doi:10.1051/0004-6361/202245521](https://doi.org/10.1051/0004-6361/202245521).
- [190] R. A. Booth, J. E. Owen, M. Schulik, Dust formation in the outflows of catastrophically evaporating planets, *Mon. Not. Roy. Astron. Soc.* 518 (2023) 1761–1775. [doi:10.1093/mnras/stac3121](https://doi.org/10.1093/mnras/stac3121).
- [191] D. Z. Seligman, A. D. Feinstein, D. Lai, L. Welbanks, A. G. Taylor, J. Becker, F. C. Adams, M. Morgan, J. B. Bergner, Potential Melting of Extrasolar Planets by Tidal Dissipation, *Astrophys. J.* 961 (2024) 22. [doi:10.3847/1538-4357/ad0b82](https://doi.org/10.3847/1538-4357/ad0b82).
- [192] J. S. Greaves, P. B. Rimmer, A. M. S. Richards, J. J. Petkowski, W. Bains, S. Ranjan, S. Seager, D. L. Clements, C. S. Silva, H. J. Fraser, Low levels of sulphur dioxide contamination of Venusian phosphine spectra, *Mon. Not. Roy. Astron. Soc.* 514 (2) (2022) 2994–3001. [doi:10.1093/mnras/stac1438](https://doi.org/10.1093/mnras/stac1438).
- [193] J. A. Dias, P. Machado, J. Ribeiro, From Atmospheric Evolution to the Search of Species of Astrobiological Interest in the Solar System-Case Studies Using the Planetary Spectrum Generator, *Atmosphere* 13 (3). [doi:10.3390/atmos13030461](https://doi.org/10.3390/atmos13030461).
- [194] R. Koga, T. Suzuki, F. Tsuchiya, T. Sakanoi, Y. Hirahara, ALMA Observation of SO₂ Gas Originating from Io's Volcanic Plume and Lava Areas, *Astrophys. J.* 907 (2021) L6. [doi:10.3847/2041-8213/abd39f](https://doi.org/10.3847/2041-8213/abd39f).
- [195] P. Cambianica, G. Cremonese, G. Munaretto, M. T. Capria, M. Fulle, W. Boschin, L. Di Fabrizio, A. Harutyunyan, A high-spectral-resolution catalog of emission lines in the visible spectrum of comet C/2020 F3 (NEOWISE), *Astron. Astrophys.* 656 (2021) A160. [doi:10.1051/0004-6361/202140309](https://doi.org/10.1051/0004-6361/202140309).
- [196] B. I. Lacy, A. Burrows, JWST Transit Spectra. II. Constraining Aerosol Species, Particle-size Distributions, Temperature, and Metallicity for Cloudy Exoplanets, *Astrophys. J.* 904 (2020) 25. [doi:10.3847/1538-4357/abbc6c](https://doi.org/10.3847/1538-4357/abbc6c).
- [197] L. Welbanks, N. Madhusudhan, Aurora: A Generalized Retrieval Framework for Exoplanetary Transmission Spectra, *Astrophys. J.* (2021) 114. [doi:10.3847/1538-4357/abee94](https://doi.org/10.3847/1538-4357/abee94).
- [198] P. Lavvas, A. Arfaux, Impact of photochemical hazes and gases on exoplanet atmospheric thermal structure, *Mon. Not. Roy. Astron. Soc.* 502 (2021) 5643–5657. [doi:10.1093/mnras/stab456](https://doi.org/10.1093/mnras/stab456).
- [199] S. Mukherjee, N. E. Batalha, M. S. Marley, Cloud Parameterizations and their Effect on Retrievals of Exoplanet Reflection Spectroscopy, *Astrophys. J.* 910 (2021) 158. [doi:10.3847/1538-4357/abe53b](https://doi.org/10.3847/1538-4357/abe53b).
- [200] M. Lefèvre, X. Tan, E. K. H. Lee, R. T. Pierrehumbert, Cloud-convection Feedback in Brown Dwarf Atmospheres, *Astrophys. J.* 929 (2022) 153. [doi:10.3847/1538-4357/ac5e2d](https://doi.org/10.3847/1538-4357/ac5e2d).
- [201] C. Jiang, G. Chen, E. Palle, F. Murgas, H. Parviainen, Y. Ma, Featureless transmission spectra of 12 giant exoplanets observed by GTC/OSIRIS, *Astron. Astrophys.* 675. [doi:10.1051/0004-6361/202346091](https://doi.org/10.1051/0004-6361/202346091).
- [202] B. Lacy, A. Burrows, Self-consistent Models of Y Dwarf Atmospheres with Water Clouds and Disequilibrium Chemistry, *Astrophys. J.* 950 (2023) 8. [doi:10.3847/1538-4357/acc8cb](https://doi.org/10.3847/1538-4357/acc8cb).
- [203] S. Ma, Y. Ito, A. F. Al-Refaie, Q. Changeat, B. Edwards, G. Tinetti, YunMa: Enabling Spectral Retrievals of Exoplanetary Clouds, *Astrophys. J.* 957 (2) (2023) 104. [doi:10.3847/1538-4357/acf8ca](https://doi.org/10.3847/1538-4357/acf8ca).
- [204] R. Baeyens, T. Konings, O. Venot, L. Carone, L. Decin, Grid of pseudo-2D chemistry models for tidally locked exoplanets - II. The role of photochemistry, *Mon. Not. Roy. Astron. Soc.* 512 (2022) 4877–4892. [doi:10.1093/mnras/stac809](https://doi.org/10.1093/mnras/stac809).

- [205] B. Edwards, I. Stotesbury, Terminus: A Versatile Simulator for Space-based Telescopes, *Astron. J.* 161 (2021) 266. [doi:10.3847/1538-3881/abdf4d](https://doi.org/10.3847/1538-3881/abdf4d).
- [206] L. V. Mugnai, A. Al-Refaie, A. Bocchieri, Q. Changeat, E. Pascale, G. Tinetti, Alfnoor: Assessing the Information Content of Ariel's Low-resolution Spectra with Planetary Population Studies, *Astron. J.* 162 (2021) 288. [doi:10.3847/1538-3881/ac2e92](https://doi.org/10.3847/1538-3881/ac2e92).
- [207] B. Charnay, J. M. Mendonca, L. Kreidberg, N. B. Cowan, J. Taylor, T. J. Bell, O. Demangeon, B. Edwards, C. A. Haswell, G. Morello, L. V. Mugnai, E. Pascale, G. Tinetti, P. Tremblin, R. T. Zellem, A survey of exoplanet phase curves with Ariel, *Exp. Astron.* 53 (2022) 417–446. [doi:10.1007/s10686-021-09715-x](https://doi.org/10.1007/s10686-021-09715-x).
- [208] G. Guilluy, A. Sozzetti, P. Giacobbe, A. S. Bonomo, G. Micela, On the synergy between Ariel and ground-based high-resolution spectroscopy, *Exp. Astron.* 53 (2022) 655–677. [doi:10.1007/s10686-021-09824-7](https://doi.org/10.1007/s10686-021-09824-7).
- [209] B. Edwards, G. Tinetti, The Ariel Target List: The Impact of TESS and the Potential for Characterizing Multiple Planets within a System, *Astron. J.* 164 (2022) 15. [doi:10.3847/1538-3881/ac6bf9](https://doi.org/10.3847/1538-3881/ac6bf9).
- [210] J. K. Barstow, Q. Changeat, K. L. Chubb, P. E. Cubillos, B. Edwards, R. J. MacDonald, M. Min, I. P. Waldmann, A retrieval challenge exercise for the Ariel mission, *Exp. Astron.* 53 (2022) 447–471. [doi:10.1007/s10686-021-09821-w](https://doi.org/10.1007/s10686-021-09821-w).
- [211] B. Edwards, Q. Changeat, A. Tsiaras, K. H. Yip, A. F. Al-Refaie, L. Anisman, M. F. Bieger, A. Gressier, S. Shibata, N. Skaf, J. Bouwman, J. Y.-K. Cho, M. Ikoma, O. Venot, I. Waldmann, P.-O. Lagage, G. Tinetti, Exploring the Ability of Hubble Space Telescope WFC3 G141 to Uncover Trends in Populations of Exoplanet Atmospheres through a Homogeneous Transmission Survey of 70 Gaseous Planets, *Astrophys. J. Suppl.* 269. [doi:10.3847/1538-4365/ac9f1a](https://doi.org/10.3847/1538-4365/ac9f1a).
- [212] A. Bocchieri, L. V. Mugnai, E. Pascale, Q. Changeat, G. Tinetti, Detecting molecules in Ariel low resolution transmission spectra, *Exp. Astron.* 56 (2023) 605–644. [doi:10.1007/s10686-023-09911-x](https://doi.org/10.1007/s10686-023-09911-x).
- [213] Q. Changeat, Y. Ito, A. F. Al-Refaie, K. H. Yip, T. Lueftinger, Toward Atmospheric Retrievals of Panchromatic Light Curves: ExPLOR-ing Generalized Inversion Techniques for Transiting Exoplanets with JWST and Ariel, *Astron. J.* 167 (5) (2024) 195. [doi:10.3847/1538-3881/ad3032](https://doi.org/10.3847/1538-3881/ad3032).
- [214] MacDonald, Ryan J. and Lewis, Nikole K., Trident: A rapid 3d radiative-transfer model for exoplanet transmission spectra, *Astrophys. J.* 929 (2022) 20. [doi:10.3847/1538-4357/ac47fe](https://doi.org/10.3847/1538-4357/ac47fe).
- [215] I. P. Waldmann, M. Rocchetto, G. Tinetti, E. J. Barton, S. N. Yurchenko, J. Tennyson, τ -REx II: Retrieval of Emission Spectra, *Astrophys. J.* 813 (2015) 13.
- [216] A. F. Al-Refaie, Q. Changeat, I. P. Waldmann, G. Tinetti, TauREx 3: A Fast, Dynamic, and Extendable Framework for Retrievals, *Astrophys. J.* 917 (2021) 37. [doi:10.3847/1538-4357/ac0252](https://doi.org/10.3847/1538-4357/ac0252).
- [217] J. Harrington, M. D. Himes, P. E. Cubillos, J. Blecic, P. M. Rojo, R. C. Challener, N. B. Lust, M. O. Bowman, S. D. Blumenthal, I. Dobbs-Dixon, A. S. D. Foster, A. J. Foster, M. R. Green, T. J. Loredo, K. J. McIntyre, M. M. Stemm, D. C. Wright, An Open-source Bayesian Atmospheric Radiative Transfer (BART) Code. I. Design, Tests, and Application to Exoplanet HD 189733b, *Planet. Sci. J.* 3 (2022) 80. [doi:10.3847/PSJ/ac3513](https://doi.org/10.3847/PSJ/ac3513).
- [218] P. E. Cubillos, J. Harrington, J. Blecic, M. D. Himes, P. M. Rojo, T. J. Loredo, N. B. Lust, R. C. Challener, A. J. Foster, M. M. Stemm, A. S. D. Foster, S. D. Blumenthal, An Open-source Bayesian Atmospheric Radiative Transfer (BART) Code. II. The Transit Radiative Transfer Module and Retrieval of HAT-P-11b, *Planet. Sci. J.* 3 (2022) 81. [doi:10.3847/PSJ/ac348b](https://doi.org/10.3847/PSJ/ac348b).
- [219] E. K. H. Lee, J. P. Wardenier, B. Prinoth, V. Parmentier, S. L. Grimm, R. Baeyens, L. Carone, D. Christie, R. Deitrick, D. Kitzmann, N. Mayne, M. Roman, B. Thorsbro, 3D Radiative Transfer for Exoplanet Atmospheres. gCMCRT: A GPU-accelerated MCRT Code, *Astrophys. J.* 929 (2022) 180. [doi:10.3847/1538-4357/ac61d6](https://doi.org/10.3847/1538-4357/ac61d6).
- [220] P. E. Cubillos, J. Blecic, The pyrat bay framework for exoplanet atmospheric modelling: a population study of Hubble/WFC3 transmission spectra, *Mon. Not. Roy. Astron. Soc.* 505 (2021) 2675–2702. [doi:10.1093/mnras/stab1405](https://doi.org/10.1093/mnras/stab1405).
- [221] G. L. Villanueva, T. J. Fauchez, V. Kofman, E. Alei, E. K. H. Lee, E. Janin, M. D. Himes, J. Leconte, M. Leung, S. Faggi, M. T. Mak, D. E. Sergeev, T. Kozakis, J. Manners, N. Mayne, E. W. Schwerterman, A. R. Howe, N. Batalha, Modeling Atmospheric Lines by the Exoplanet Community (MALBEC) Version 1.0: A CUISINES Radiative Transfer Intercomparison Project, *Planet Space Sci.* 5 (2024) 64. [doi:10.3847/PSJ/ad2681](https://doi.org/10.3847/PSJ/ad2681).
- [222] M. Min, C. W. Ormel, K. Chubb, C. Helling, Y. Kawashima, The ARCiS framework for exoplanet atmospheres - Modelling philosophy and retrieval, *Astron. Astrophys.* 642 (2020) A28. [doi:10.1051/0004-6361/201937377](https://doi.org/10.1051/0004-6361/201937377).
- [223] L. V. Mugnai, D. Modirrousta-Galian, RAPOC: The Rosseland and Planck opacity converter A user-friendly and fast opacity program for Python, *Exp. Astron.* 55 (2023) 521–539. [doi:10.1007/s10686-022-09869-2](https://doi.org/10.1007/s10686-022-09869-2).
- [224] A. J. Wheeler, A. R. Casey, M. W. Abruzzo, Korg: Fitting, Model Atmosphere Interpolation, and Brackett Lines, *Astron. J.* 167 (2024) 83. [doi:10.3847/1538-3881/ad19cc](https://doi.org/10.3847/1538-3881/ad19cc).
- [225] Z. Zhan, S. Seager, J. J. Petkowski, C. Sousa-Silva, S. Ranjan, J. Huang, W. Bains, Assessment of Isoprene as a Possible Biosignature Gas in Exoplanets with Anoxic Atmospheres, *Astrobiology* 21 (2021) 765–792. [doi:10.1089/ast.2019.2146](https://doi.org/10.1089/ast.2019.2146).
- [226] H. T. Ishikawa, W. Aoki, T. Kotani, M. Kuzuhara, M. Omiya, A. Reiners, M. Zechmeister, Elemental abundances of M dwarfs based on high-resolution near-infrared spectra: Verification by binary systems, *Publ. Astron. Soc. Japan* 72 (2020) 102. [doi:10.1093/pasj/pssaa101](https://doi.org/10.1093/pasj/pssaa101).
- [227] G. Contursi, P. de Laverny, A. Recio-Blanco, P. A. Palicio, GSP-spec line list for the parametrisation of Gaia-RVS stellar spectra, *Astron. Astrophys.* 654. [doi:10.1051/0004-6361/202140912](https://doi.org/10.1051/0004-6361/202140912).
- [228] E. Marfil, H. M. Tabernero, D. Montes, J. A. Caballero, F. J. Lazaro, J. Gonzalez Hernandez, I. E. Nagel, V. M. Passegger, A. Schweitzer, I. Ribas, A. Reiners, A. Quirrenbach, P. J. Amado, C. Cifuentes, M. Cortes-Contreras, S. Dreizler, C. Duque-Arribas, D. Galadi-Enriquez, T. Henning, S. Jeffers, V. A. Kaminski, M. Kuerster, M. Lafarga, A. Lopez-Gallifa, J. C. Morales, Y. Shan, M. Zechmeister, The CARMENES search for exoplanets around M dwarfs Stellar atmospheric parameters of target stars with STEPARYN, *Astron. Astrophys.* 656. [doi:10.1051/0004-6361/202141980](https://doi.org/10.1051/0004-6361/202141980).
- [229] Y. P. Lyubchyk, Y. Pavlenko, V. O. K. Lyubchyk, H. R. A. Jones, Bands of NaH lines in Spectra of Late Type Stars, *Kinemat. Phys. Celest.* 38 (2022) 159–165. [doi:10.3103/S0884591322030059](https://doi.org/10.3103/S0884591322030059).
- [230] H. R. A. Jones, Y. Pavlenko, Y. Lyubchik, M. Bessell, N. Allard, D. J. Pinfield, A blue depression in the optical spectra of M dwarfs, *Mon. Not. Roy. Astron. Soc.* 523 (2023) 1297–1309. [doi:10.1093/mnras/stad1391](https://doi.org/10.1093/mnras/stad1391).
- [231] P. I. Cristofari, J.-F. Donati, C. P. Folsom, T. Masseron, P. Fouqué, C. Moutou, E. Artigau, A. Carmona, P. Petit, X. Delfosse, E. Martioli, the SLS consortium, Constraining atmospheric parameters and surface magnetic fields with ZeeTurbo: an application to SPIRou spectra, *Mon. Not. Roy. Astron. Soc.* 522 (2023) 1342–1357. [doi:10.1093/mnras/stad865](https://doi.org/10.1093/mnras/stad865).

- [232] J. W. Xuan, J. Wang, L. Finnerty, K. Horstman, S. Grimm, A. E. Peck, E. Nielsen, H. A. Knutson, D. Mawet, H. Isaacson, A. W. Howard, M. C. Liu, S. Walker, M. W. Phillips, G. A. Blake, J.-B. Ruffio, Y. Zhang, J. Inglis, N. L. Wallack, A. Sanghi, E. J. Gonzales, F. Dai, A. Baker, R. Bartos, C. Z. Bond, M. L. Bryan, B. Calvin, S. Cetre, J.-R. Delorme, G. Doppmann, D. Echeverri, M. P. Fitzgerald, N. Jovanovic, J. Liberman, R. A. Lopez, E. C. Martin, E. Morris, J. Pezzato, G. Ruane, B. Sappey, T. Schofield, A. Skemer, T. Venenciano, J. K. Wallace, J. Wang, P. Wizinowich, Y. Xin, S. Agrawal, C. R. Do, C.-C. Hsu, C. L. Phillips, Validation of Elemental and Isotopic Abundances in Late-M Spectral Types with the Benchmark HIP 55507 AB System, *Astrophys. J.* 962. doi:[10.3847/1538-4357/ad1243](https://doi.org/10.3847/1538-4357/ad1243).
- [233] Y. V. Pavlenko, A. Evans, D. P. K. Banerjee, T. R. Geballe, U. Munari, R. D. Gehrz, C. E. Woodward, S. Starrfield, Isotopic ratios in the red giant component of the recurrent nova T Coronae Borealis, *Mon. Not. Roy. Astron. Soc.* 498 (2020) 4853–4863. doi:[10.1093/mnras/staa2658](https://doi.org/10.1093/mnras/staa2658).
- [234] D. Gasman, I. Argyriou, G. C. Sloan, B. Aringer, J. Alvarez-Marquez, O. Fox, A. Glasse, A. Glauser, O. C. Jones, K. Justtanont, P. J. Kavanagh, P. Klaassen, A. Labiano, K. Larson, D. R. Law, M. Mueller, O. Nayak, A. Noriega-Crespo, P. Patapis, P. Royer, B. Vandenbussche, JWST MIRI/MRS in-flight absolute flux calibration and tailored fringe correction for unresolved sources, *Astron. Astrophys.* 673. doi:[10.1051/0004-6361/202245633](https://doi.org/10.1051/0004-6361/202245633).
- [235] C. R. Hayes, K. A. Venn, F. Waller, J. Jensen, A. W. Mcconnachie, J. Pazder, F. Sestito, A. Anthony, G. Baker, J. Bassett, J. Bento, T. Berg, G. Burley, J. Brzeski, S. Case, E. Chapin, T. Chin, E. Chisholm, V. Churilov, A. Densmore, R. Diaz, J. Dunn, M. Edgar, T. Farrell, V. Firpo, J. Fitzsimmons, J. Font-Serra, J. Fuentes, C. Ganton, M. Gomez-Jimenez, T. Hardy, D. Henderson, A. Hill, B. Hoff, M. Ireland, V. Kalari, N. Kelly, U. Klauser, Y. Kondrat, K. Labrie, S. Lambert, L. Luval, J. Lawrence, J. Lothrop, G. S. Macdonald, S. Mali, S. Margheim, R. Mcdermid, H. Mcgregor, B. Miller, F. Miranda, R. Muller, J. Nielsen, R. Norbury, O. Oberdorf, N. Pai, G. Perez, P. Prado, I. Price, C. Quiroz, V. Reshetov, G. Robertson, R. Ruiz-Carmona, R. Salinas, K. M. Sebo, A. Sheinis, M. Shetrone, K. Shortridge, K. Silversides, K. Silva, C. Simpson, G. Smith, K. Szeto, J. Tims, E. Toro, C. Urrutia, S. Venkatesan, L. Waller, I. Wevers, R. Wierzbicki, M. White, P. Young, R. Zhelem, GHOST Commissioning Science Results: Identifying a New Chemically Peculiar Star in Reticulum II, *Astrophys. J.* 955. doi:[10.3847/1538-4357/acebc0](https://doi.org/10.3847/1538-4357/acebc0).
- [236] T. D. Brandt, T. J. Dupuy, B. P. Bowler, D. C. B. Gagliuffi, J. Faherty, G. M. Brandt, D. Michalik, A Dynamical Mass of 70 ± 5 $M_{J_{\text{up}}}$ for Gliese 229B, the First T Dwarf, *Astron. J.* 160 (2020) 196. doi:[10.3847/1538-3881/abb45e](https://doi.org/10.3847/1538-3881/abb45e).
- [237] A. A. A. Piette, N. Madhusudhan, Considerations for atmospheric retrieval of high-precision brown dwarf spectra, *Mon. Not. Roy. Astron. Soc.* 497 (2020) 5136–5154. doi:[10.1093/mnras/staa2289](https://doi.org/10.1093/mnras/staa2289).
- [238] A. M. Meisner, A. C. Schneider, A. J. Burgasser, F. Marocco, M. R. Line, J. K. Faherty, J. D. Kirkpatrick, D. Caselden, M. J. Kuchner, C. R. Gelino, J. Gagne, C. Theissen, R. Gerasimov, C. Aganze, C.-c. Hsu, J. P. Wisniewski, S. L. Casewell, D. C. Bardalez Gagliuffi, S. E. Logsdon, P. R. M. Eisenhardt, K. Allers, J. H. Debes, M. B. Allen, N. Stevnbak Andersen, S. Goodman, L. Grammaize, D. W. Martin, A. Sainio, M. C. Cushing, The Backyard Worlds Planet 9 Collaboration, New Candidate Extreme T Subdwarfs from the Backyard Worlds: Planet 9 Citizen Science Project, *Astrophys. J.* 915. doi:[10.3847/1538-4357/ac013c](https://doi.org/10.3847/1538-4357/ac013c).
- [239] C. von Essen, M. Mallonn, A. Piette, N. B. Cowan, N. Madhusudhan, E. Agol, V. Antoci, K. Poppenhaeger, K. G. Stassun, S. Khalafinejad, G. Tautvaisiene, TESS unveils the optical phase curve of KELT-1b - Thermal emission and ellipsoidal variation from the brown dwarf companion along with the stellar activity, *Astron. Astrophys.* 648 (2021) A71. doi:[10.1051/0004-6361/202038524](https://doi.org/10.1051/0004-6361/202038524).
- [240] S. Mukherjee, J. J. Fortney, R. Jensen-Clem, X. Tan, M. S. Marley, N. E. Batalha, Modeling Polarization Signals from Cloudy Brown Dwarfs Luhman 16 A and B in Three Dimensions, *Astron. J.* 923 (2021) 113. doi:[10.3847/1538-4357/ac2d92](https://doi.org/10.3847/1538-4357/ac2d92).
- [241] M. S. Marley, D. Saumon, C. Visscher, R. Lupu, R. Freedman, C. Morley, J. J. Fortney, C. Seay, A. J. R. W. Smith, D. J. Teal, R. Wang, The Sonora Brown Dwarf Atmosphere and Evolution Models. I. Model Description and Application to Cloudless Atmospheres in Rainout Chemical Equilibrium, *Astrophys. J.* 920 (2021) 85. doi:[10.3847/1538-4357/ac141d](https://doi.org/10.3847/1538-4357/ac141d).
- [242] M. C. Cushing, A. C. Schneider, J. D. Kirkpatrick, C. V. Morley, M. S. Marley, C. R. Gelino, G. N. Mace, E. L. Wright, P. R. Eisenhardt, M. F. Skrutskie, K. A. Marsh, An Improved Near-infrared Spectrum of the Archetype Y Dwarf WISEP J182831.08+265037.8, *Astrophys. J.* 920 (2021) 20. doi:[10.3847/1538-4357/ac12cb](https://doi.org/10.3847/1538-4357/ac12cb).
- [243] A. Lueber, D. Kitzmann, B. P. Bowler, A. J. Burgasser, K. Heng, Retrieval Study of Brown Dwarfs across the L-T Sequence, *Astrophys. J.* 930 (2022) 136. doi:[10.3847/1538-4357/ac63b9](https://doi.org/10.3847/1538-4357/ac63b9).
- [244] M. E. Tannock, S. Metchev, C. E. Hood, G. N. Mace, J. J. Fortney, C. V. Morley, D. T. Jaffe, R. Lupu, A 1.46–2.48 μm spectroscopic atlas of a T6 dwarf (1060 K) atmosphere with IGRINS: First detections of H_2S and H_2 , and verification of H_2O , CH_4 , and NH_3 line lists, *Mon. Not. Roy. Astron. Soc.* 514. doi:[10.1093/mnras/stac1412](https://doi.org/10.1093/mnras/stac1412).
- [245] J. W. Xuan, J. Wang, J.-B. Ruffio, H. Knutson, D. Mawet, P. Molli'ere, J. Kolecki, A. Vigan, S. Mukherjee, N. Wallack, J. Wang, A. Baker, R. Bartos, G. A. Blake, C. Z. Bond, M. Bryan, B. Calvin, S. Cetre, M. Chun, J.-R. Delorme, G. Doppmann, D. Echeverri, L. Finnerty, M. P. Fitzgerald, K. Horstman, J. Inglis, N. Jovanovic, R. López, E. C. Martin, E. Morris, J. Pezzato, S. Ragland, B. Ren, G. Ruane, B. Sappey, T. Schofield, A. Skemer, T. Venenciano, J. K. Wallace, P. Wizinowich, A Clear View of a Cloudy Brown Dwarf Companion from High-resolution Spectroscopy, *Astrophys. J.* 937 (2022) 54. doi:[10.3847/1538-4357/ac8673](https://doi.org/10.3847/1538-4357/ac8673).
- [246] M. J. Rowland, C. V. Morley, M. R. Line, Toward Robust Atmospheric Retrieval on Cloudy L Dwarfs: the Impact of Thermal and Abundance Profile Assumptions, *Astrophys. J.* 947 (2023) 6. doi:[10.3847/1538-4357/acbb07](https://doi.org/10.3847/1538-4357/acbb07).
- [247] D. Barrado, P. Mollière, P. Patapis, M. Min, P. Tremblin, F. Ardevol Martinez, N. Whiteford, M. Vasist, I. Argyriou, M. Samland, P.-O. Lagage, L. Decin, R. Waters, T. Henning, M. Morales-Calderón, M. Guedel, B. Vandenbussche, O. Absil, P. Baudoz, A. Boccaletti, J. Bouwman, C. Cossou, A. Coulais, N. Crouzet, R. Gastaud, A. Glasse, A. M. Glauser, I. Kamp, S. Kendrew, O. Krause, F. Lahuis, M. Mueller, G. Olofsson, J. Pye, D. Rouan, P. Royer, S. Scheithauer, I. Waldmann, L. Colina, E. F. van Dishoeck, T. Ray, G. Östlin, G. Wright, $^{15}\text{NH}_3$ in the atmosphere of a cool brown dwarf, *Nature* 624 (2023) 263–266. doi:[10.1038/s41586-023-06813-y](https://doi.org/10.1038/s41586-023-06813-y).
- [248] S. Mukherjee, N. E. Batalha, J. J. Fortney, M. S. Marley, PICASO 3.0: A One-dimensional Climate Model for Giant Planets and Brown Dwarfs, *Astrophys. J.* 942 (2023) 71. doi:[10.3847/1538-4357/ac9f48](https://doi.org/10.3847/1538-4357/ac9f48).
- [249] C. E. Hood, J. J. Fortney, M. R. Line, J. K. Faherty, Brown Dwarf Retrievals on FIRE!: Atmospheric Constraints and Lessons Learned from High Signal-to-noise Medium-resolution Spectroscopy of a T9 Dwarf, *Astrophys. J.* 953 (2023) 170. doi:[10.3847/1538-4357/ace32e](https://doi.org/10.3847/1538-4357/ace32e).
- [250] K. Franson, B. P. Bowler, M. Bonavita, T. D. Brandt, M. Chen, M. Samland, Z. Zhang, A. Lueber, K. Heng, D. Kitzmann, T. Wolf, B. A. Jones, Q. H. Tran, D. C. B. Gagliuffi, B. Biller, J. Chilcote, J. R. Crepp, T. J. Dupuy, J. Faherty, C. Fontanive, T. D. Groff, R. Gratton, O. Guyon, R. Jensen-Clem, N. Jovanovic, N. J. Kasdin, J. Lozi, E. A. Magnier, K. Mužić, A. Sanghi, C. A. Theissen, Astrometric Accelerations as Dynamical Beacons: Discovery and Characterization of HIP 21152 B, the First T-dwarf Companion in the Hyades*, *Astron. J.* 165 (2023) 39. doi:[10.3847/1538-3881/aca408](https://doi.org/10.3847/1538-3881/aca408).

- [251] E. K. H. Lee, X. Tan, S.-M. Tsai, Dynamically coupled kinetic chemistry in brown dwarf atmospheres - I. Performing global scale kinetic modelling, Mon. Not. Roy. Astron. Soc. 523 (2023) 4477–4491. [doi:10.1093/mnras/stad1715](https://doi.org/10.1093/mnras/stad1715).
- [252] S. Mukherjee, J. J. Fortney, C. V. Morley, N. E. Batalha, M. S. Marley, T. Karalidi, C. Visscher, R. Lupu, R. Freedman, E. Gharib-Nezhad, The Sonora Substellar Atmosphere Models. IV. Elf Owl: Atmospheric Mixing and Chemical Disequilibrium with Varying Metallicity and C/O Ratios, Astrophys. J. 963 (2024) 73. [doi:10.3847/1538-4357/ad18c2](https://doi.org/10.3847/1538-4357/ad18c2).
- [253] A. J. Burgasser, R. Bezanson, I. Labbe, G. Brammer, S. E. Cutler, L. J. Furtak, J. E. Greene, R. Gerasimov, J. Leja, R. Pan, S. H. Price, B. Wang, J. R. Weaver, K. E. Whitaker, S. Fujimoto, V. Kokorev, P. Dayal, T. Nanayakkara, C. C. Williams, D. Marchesini, A. Zitrin, P. van Dokkum, UNCOVER: JWST Spectroscopy of Three Cold Brown Dwarfs at Kiloparsec-scale Distances, Astrophys. J. 962 (2024) 177. [doi:10.3847/1538-4357/ad206f](https://doi.org/10.3847/1538-4357/ad206f).
- [254] B. W. P. Lew, T. Roellig, N. E. Batalha, M. Line, T. Greene, S. Murkherjee, R. Freedman, M. Meyer, C. Beichman, C. A. de Oliveira, M. De Furio, D. Johnstone, A. Z. Greenbaum, M. Marley, J. J. Fortney, E. T. Young, J. Leisenring, M. Boyer, K. Hodapp, K. Misselt, J. Stansberry, M. Rieke, High-precision Atmospheric Characterization of a Y Dwarf with JWST NIRSpec G395H Spectroscopy: Isotopologue, C/O Ratio, Metallicity, and the Abundances of Six Molecular Species, Astron. J. 167 (2024) 237. [doi:10.3847/1538-3881/ad3425](https://doi.org/10.3847/1538-3881/ad3425).
- [255] Z. Guo, P. W. Lucas, R. Kurtev, J. Borissova, C. Contreras Peña, S. N. Yurchenko, L. C. Smith, D. Minniti, R. K. Saito, A. Bayo, M. Catelan, J. Alonso-García, A. Caratti o Garatti, C. Morris, D. Froebrich, J. Tennyson, K. Maucó, A. Aguayo, N. Miller, H. D. S. Muthu, Spectroscopic confirmation of high-amplitude eruptive YSOs and dipping giants from the VVV survey, Mon. Not. Roy. Astron. Soc. 528 (2024) 1769–1788. [doi:10.1093/mnras/stad3700](https://doi.org/10.1093/mnras/stad3700).
- [256] T. Danilovich, C. A. Gottlieb, L. Decin, A. M. S. Richards, K. L. K. Lee, T. Kaminski, N. A. Patel, K. H. Young, K. M. Menten, Rotational Spectra of vibrationally excited AlO and TiO in oxygen-rich stars, Astron. J. 904 (2020) 110. [doi:10.3847/1538-4357/abc079](https://doi.org/10.3847/1538-4357/abc079).
- [257] A. Coenegrachts, T. Danilovich, F. De Ceuster, L. Decin, The unusual 3D distribution of NaCl around the asymptotic giant branch star IK Tau, Astron. Astrophys. 678. [doi:10.1051/0004-6361/202346116](https://doi.org/10.1051/0004-6361/202346116).
- [258] C. Sandin, L. Mattsson, K. L. Chubb, M. Ergon, P. M. Weilbacher, Three-component modelling of O-rich AGB star winds I. Effects of drift using forsterite, Astron. Astrophys. 677. [doi:10.1051/0004-6361/202345841](https://doi.org/10.1051/0004-6361/202345841).
- [259] G. Pastorelli, P. Marigo, L. Girardi, B. Aringer, Y. Chen, S. Rubele, M. Trabucchi, S. Bladh, M. L. Boyer, A. Bressan, J. J. Dalcanton, M. A. T. Groenewegen, T. Lebzelter, N. Mowlavi, K. L. Chubb, M.-R. L. Cioni, R. de Grijs, V. D. Ivanov, A. Nanni, J. T. van Loon, S. Zaggia, Constraining the thermally pulsing asymptotic giant branch phase with resolved stellar populations in the Large Magellanic Cloud, Mon. Not. Roy. Astron. Soc. 498 (2020) 3283–3301. [doi:10.1093/mnras/staa2565](https://doi.org/10.1093/mnras/staa2565).
- [260] K. Eriksson, S. Hofner, B. Aringer, Synthetic photometry for carbon-rich giants V. Effects of grain-size-dependent dust opacities, Astron. Astrophys. 673. [doi:10.1051/0004-6361/202245206](https://doi.org/10.1051/0004-6361/202245206).
- [261] V. Karambelkar, M. M. Kasliwal, P. Tisserand, G. C. Clayton, C. L. Crawford, S. G. Anand, T. R. Geballe, E. Montiel, R Coronae Borealis and dustless hydrogen-deficient carbon stars likely have different oxygen isotope ratios, Astron. Astrophys. 667. [doi:10.1051/0004-6361/202142918](https://doi.org/10.1051/0004-6361/202142918).
- [262] D. A. García-Hernández, N. Kameswara Rao, D. L. Lambert, K. Eriksson, A. B. S. Reddy, T. Masseron, The Carbon Star DY Persei May Be a Cool R Coronae Borealis Variable, Astrophys. J. 948 (2023) 15. [doi:10.3847/1538-4357/acc574](https://doi.org/10.3847/1538-4357/acc574).
- [263] M. Z. Mobeen, T. Kaminski, A. Matter, M. Wittkowski, C. Paladini, The mid-infrared environment of the stellar merger remnant V838 Monocerotis, Astron. Astrophys. 655 (2021) A100. [doi:10.1051/0004-6361/202142297](https://doi.org/10.1051/0004-6361/202142297).
- [264] T. Steinmetz, T. Kaminski, M. Schmidt, A. Kiljan, A bipolar structure and shocks surrounding the stellar-merger remnant V1309 Scorpii, Astron. Astrophys. 682. [doi:10.1051/0004-6361/202347818](https://doi.org/10.1051/0004-6361/202347818).
- [265] S. Liljegren, A. Jerkstrand, P. S. Barklem, G. Nyman, R. Brady, S. N. Yurchenko, The molecular chemistry of Type Ibc supernovae and diagnostic potential with the James Webb Space Telescope, Astron. Astrophys. 674 (2023) A184. [doi:10.1051/0004-6361/202243491](https://doi.org/10.1051/0004-6361/202243491).
- [266] S. Nickerson, N. Rangwala, S. W. J. Colgan, C. DeWitt, X. Huang, K. Acharyya, M. Drozdovskaya, R. C. Fortenberry, E. Herbst, T. J. Lee, The First Mid-infrared Detection of HNC in the Interstellar Medium: Probing the Extreme Environment toward the Orion Hot Core, Astrophys. J. 907 (1) (2021) 51. [doi:10.3847/1538-4357/abca36](https://doi.org/10.3847/1538-4357/abca36).
- [267] S. Nickerson, N. Rangwala, S. W. J. Colgan, C. DeWitt, J. S. S. Monzon, X. Huang, K. Acharyya, M. N. N. Drozdovskaya, R. C. C. Fortenberry, E. Herbst, T. J. J. Lee, The Mid-infrared Molecular Inventory toward Orion IRc2, Astrophys. J. 945 (2023) 26. [doi:10.3847/1538-4357/aca6e8](https://doi.org/10.3847/1538-4357/aca6e8).
- [268] K. P. Bowen, P.-M. Hillenbrand, J. Lievin, D. W. Savin, X. Urbain, Dynamics of the isotope exchange reaction of d with h_3^+ , h_2d^+ , and d_2h^+ , J. Chem. Phys. 154. [doi:10.1063/5.0038434](https://doi.org/10.1063/5.0038434).
- [269] A. C. LoCurto, M. A. Welch, T. R. Sippel, J. B. Michael, High-speed visible supercontinuum laser absorption spectroscopy of metal oxides, Opt. Lett. 46 (2021) 3288–3291. [doi:10.1364/OL.428456](https://doi.org/10.1364/OL.428456).
- [270] K. A. Daniel, C. M. Murzyn, D. J. Allen, K. P. Lynch, C. R. Downing, J. L. Wagner, Coaxial laser absorption and optical emission spectroscopy of high-pressure aluminum monoxide, Opt. Lett. 47 (9) (2022) 2350–2353. [doi:10.1364/OL.456342](https://doi.org/10.1364/OL.456342).
- [271] C. M. Murzyn, D. J. Allen, A. N. Baca, M. L. Ching, R. T. Marinis, Tunable infrared laser absorption spectroscopy of aluminum monoxide $\text{A}^2\Pi_i - \text{X}^2\Sigma^+$, J. Quant. Spectrosc. Radiat. Transf. 279 (2022) 108029. [doi:10.1016/j.jqsrt.2021.108029](https://doi.org/10.1016/j.jqsrt.2021.108029).
- [272] J. J. Gilvey, M. D. Ruesch, K. A. Daniel, C. R. Downing, K. P. Lynch, J. L. Wagner, C. S. Goldenstein, Quantum-cascade-laser-absorption-spectroscopy diagnostic for temperature, pressure, and NO $\text{X}^2\Pi_{1/2}$ at 500 kHz in shock-heated air at elevated pressures, Appl. Optics 62 (2023) A12–A24. [doi:10.1364/AO.464623](https://doi.org/10.1364/AO.464623).
- [273] A. Abraham, P. Lynch, N. Glumac, On using ab initio calibration to fit temperature from AlO B-X emission, Proceedings of the Combustion Institute 39 (2023) 1249–1257. [doi:10.1016/j.proci.2022.09.050](https://doi.org/10.1016/j.proci.2022.09.050).
- [274] J. Riedel, J. Hufgard, Y. You, LIBS at high duty-cycles: effect of repetition rate and temporal width on the excitation laser pulses, Frontiers in Physics 11 (2023) 1241533. [doi:10.3389/fphy.2023.1241533](https://doi.org/10.3389/fphy.2023.1241533).
- [275] S. M. Bobrovnikov, E. V. Gorlov, V. I. Zharkov, Efficiency of Laser Excitation of PO Photofragments of Organophosphates, Atmos. Ocean. Opt. 35 (2022) 329–340. [doi:10.1134/S1024856022040017](https://doi.org/10.1134/S1024856022040017).
- [276] S. M. Bobrovnikov, E. V. Gorlov, V. I. Zharkov, Laser-induced fluorescence of PO-photofragments of dimethyl methylphosphonate, Appl. Optics 61 (2022) 6322–6329. [doi:10.1364/AO.456005](https://doi.org/10.1364/AO.456005).
- [277] J. Merten, E. Nicholas, S. Ethridge, H. Bariola, S. Chestnut, A. Anders, J. Brees, M. Foster, Following laser-induced plasma stoichiometry with atomic absorption spectroscopy, Spectroc. Acta Pt. A-Molec. Biomolec. Spectr. 200 (2023) 106600. [doi:10.1016/j.sab.2022.106600](https://doi.org/10.1016/j.sab.2022.106600).

- [278] E. Pawelec, D. Borodin, S. Brezinsek, T. Dittmar, D. Douai, D. Mazur, A. Meigs, A. Shaw, B. Thomas, J. Contributors, E. T. Exploitation, Internal energy distributions of beh, bed, and bet molecules created during chemically assisted physical sputtering in jet tokamak plasma, *Phys. Plasmas* 31 (2024) 042516. [doi:10.1063/5.0199084](https://doi.org/10.1063/5.0199084).
- [279] M. Yekta, M. A. Zanjanchi, H. Roohi, Adsorption of HCN, HNC and CH₃CN toxic gases on the M-doped (M = Cr, Fe, Ni and Zn) GaNNs: A DFT-D study, *Colloids And Surfaces A-Physicochemical And Engineering Aspects* 684 (2024) 133120. [doi:10.1016/j.colsurfa.2023.133120](https://doi.org/10.1016/j.colsurfa.2023.133120).
- [280] M. D. Ruesch, A. J. McDonald, G. C. Mathews, S. F. Son, C. S. Goldenstein, Characterization of the influence of aluminum particle size on the temperature of composite-propellant flames using CO absorption and AlO emission spectroscopy, *Proc. Combust. Inst.* 38 (2021) 4365-4372. [doi:10.1016/j.proci.2020.06.163](https://doi.org/10.1016/j.proci.2020.06.163).
- [281] A. Fateev, From optimising waste incineration to internal combustion engines and defence applications, *Johnson Matthey Technol. Rev.* 67 (2023) 25–35. [doi:10.1595/205651323x16643556587827](https://doi.org/10.1595/205651323x16643556587827).
- [282] Z. Qin, T. Bai, L. Liu, Rovibrationally Resolved Photodissociation of AlH via Excited Electronic States, *Astrophys. J.* 917 (2021) 87. [doi:{10.3847/1538-4357/ac06d1}](https://doi.org/10.3847/1538-4357/ac06d1).
- [283] R. S. da Silva, M. Y. Ballester, L. R. Ventura, C. E. Fellows, Theoretical study of the spin-orbit coupling in the X²Π state of NO, *Chem. Phys. Lett.* 780 (2021) 138896. [doi:https://doi.org/10.1016/j.cplett.2021.138896](https://doi.org/10.1016/j.cplett.2021.138896).
- [284] M. C. Zammit, J. A. Leiding, J. Colgan, W. Even, C. J. Fontes, E. Timmermans, A comprehensive study of the radiative properties of NO-a first step toward a complete air opacity, *J. Phys. B-At. Mol. Opt. Phys.* 55. [doi:10.1088/1361-6455/ac8213](https://doi.org/10.1088/1361-6455/ac8213).
- [285] A. P. Charmet, P. Stoppa, A. De Lorenzi, M. Melosso, A. Achilli, L. Dore, C. Puzzarini, E. Cane, F. Tamassia, Computational, rotational and ro-vibrational experimental investigation of monodeuterated chloromethane, *J. Quant. Spectrosc. Radiat. Transf.* 305. [doi:10.1016/j.jqsrt.2023.108624](https://doi.org/10.1016/j.jqsrt.2023.108624).
- [286] T. Bai, Z. Qin, L. Liu, Ultraviolet spectroscopy of AlO from first principle, *J. Quant. Spectrosc. Radiat. Transf.* 302 (2023) 108587. [doi:10.1016/j.jqsrt.2023.108587](https://doi.org/10.1016/j.jqsrt.2023.108587).
- [287] A. M. Velasco, C. Lavín, Absorption Line Oscillator Strengths for the C²Π(0)-A²Σ+(0) Band of Nitric Oxide, *Astrophys. J.* 954 (2023) 144. [doi:10.3847/1538-4357/acebcb](https://doi.org/10.3847/1538-4357/acebcb).
- [288] O. N. Ulenikov, O. V. Gromova, E. S. Bektereva, N. I. Raspopova, E. A. Sklyarova, C. Sydow, K. Berezkin, C. Maul, S. Bauerecker, Line strengths, widths and shifts analysis of the 2ν₂, ν₂ + ν₄ and 2ν₄ bands in ²⁸SiH₄, ²⁹SiH₄ and ³⁰SiH₄, *J. Quant. Spectrosc. Radiat. Transf.* 270. [doi:10.1016/j.jqsrt.2021.107683](https://doi.org/10.1016/j.jqsrt.2021.107683).
- [289] D. Shapko, P. Dohnal, S. Roucka, L. Uvarova, M. Kassayova, R. Plasil, J. Glosik, Cavity ring-down spectroscopy study of neon assisted recombination of H₃⁺ ions with electrons, *J. Mol. Spectrosc.* 378 (2021) 111450. [doi:10.1016/j.jms.2021.111450](https://doi.org/10.1016/j.jms.2021.111450).
- [290] A. Znotins, F. Grussie, A. Wolf, X. Urbain, H. Kreckel, An approach for multi-color action spectroscopy of highly excited states of H₃⁺, *J. Mol. Spectrosc.* 378 (2021) 111476. [doi:10.1016/j.jms.2021.111476](https://doi.org/10.1016/j.jms.2021.111476).
- [291] A. Foltynowicz, L. Rutkowski, I. Silander, A. C. Johansson, V. S. de Oliveira, O. Axner, G. Sobon, T. Martynkien, P. Mergo, K. K. Lehmann, Measurement and assignment of double-resonance transitions to the 8900-9100-cm⁻¹ levels of methane, *Phys. Rev. A* 103. [doi:10.1103/PhysRevA.103.022810](https://doi.org/10.1103/PhysRevA.103.022810).
- [292] C. Lavín, A. M. Velasco, Transition Energies and Line Oscillator Strengths of the C²Π(0)-X²Π(0-6) Absorption Bands of Nitric Oxide. A Theoretical Study, *Astrophys. J.* 941 (2022) 29. [doi:10.3847/1538-4357/ac9e56](https://doi.org/10.3847/1538-4357/ac9e56).
- [293] A. Perrin, L. Manceron, R. Armante, F. Kwabia-Tchana, P. Roy, G. Toon, First investigation of the ν₁, ν₁-ν₉, ν₁+ν₉, and ν₁+ν₇ absorption bands of nitric acid (H¹⁴N¹⁶O₃) at 3551.766 cm⁻¹, 3092.708 cm⁻¹, 4006.974 cm⁻¹, and 4127.782 cm⁻¹, respectively, *J. Mol. Spectrosc.* 392 (2023) 111741. [doi:10.1016/j.jms.2023.111741](https://doi.org/10.1016/j.jms.2023.111741).
- [294] A. Perrin, L. Manceron, R. Armante, F. Kwabia-Tchana, P. Roy, G. Toon, Part 2: Validation for line lists generated for nitric acid (H¹⁴N¹⁶O₃) for the ν₁ band and its first two associated hot bands (ν₁+ν₉-ν₉, ν₁+ν₇-ν₇) in the 2.8 μm region, the ν₁-ν₉, ν₁+ν₉ and ν₁+ν₇ bands at 3.2 μm, 2.5 μm and 2.4 μm, respectively, *J. Mol. Spectrosc.* 392 (2023) 111740. [doi:10.1016/j.jms.2023.111740](https://doi.org/10.1016/j.jms.2023.111740).
- [295] I. A. Vasilenko, O. V. Naumenko, V.-M. Horneman, Expert List of Absorption Lines of the ³²S¹⁶O₂ Molecule in the 0-4200 cm⁻¹ Spectral Region, *Atmos. Ocean. Opt.* 36 (2023) 199–206. [doi:10.1134/S102485602303020X](https://doi.org/10.1134/S102485602303020X).
- [296] A. Owens, T. He, M. Hanicinec, C. Hill, S. Mohr, J. Tennyson, LiDB: Database of molecular vibronic state radiative lifetimes for plasma processes, *Plasma Sources Sci. Technol.* 32 (2023) 085015. [doi:10.1088/1361-6595/aceeb0](https://doi.org/10.1088/1361-6595/aceeb0).
- [297] R. Wang, U. Balciunaite, J. Chen, C. Yuan, A. Owens, J. Tennyson, NASA polynomial representation of molecular specific heats, *J. Quant. Spectrosc. Radiat. Transf.* 306 (2023) 108617. [doi:10.1016/j.jqsrt.2023.108617](https://doi.org/10.1016/j.jqsrt.2023.108617).
- [298] J. Zhang, C. Hill, J. Tennyson, S. N. Yurchenko, ExoMolHR: A Relational Database of Empirical High-Resolution Molecular Spectra, *Astrophys. J. Suppl. Ser.*
- [299] K. L. Chubb, M. Rocchetto, S. N. Yurchenko, M. Min, I. Waldmann, J. K. Barstow, P. Molliere, A. F. Al-Refaie, M. Phillips, J. Tennyson, The ExoMolOP Database: Cross-sections and K-tables for Molecules of Interest in High-Temperature Exoplanet Atmospheres, *Astron. Astrophys.* 646 (2021) A21. [doi:10.1051/0004-6361/202038350](https://doi.org/10.1051/0004-6361/202038350).
- [300] R. S. Freedman, J. Lustig-Yaeger, J. J. Fortney, R. E. Lupu, M. S. Marley, K. Lodders, Gaseous mean opacities for giant planet and ultracool dwarf atmospheres over a range of metallicities and temperatures, *Astrophys. J. Suppl. Ser.* 214 (2014) 25. [doi:10.1088/0067-0049/214/2/25](https://doi.org/10.1088/0067-0049/214/2/25).
- [301] D. S. Amundsen, N. J. Mayne, I. Baraffe, J. Manners, P. Tremblin, B. Drummond, C. Smith, D. M. Acreman, D. Homeier, The uk met office global circulation model with a sophisticated radiation scheme applied to the hot jupiter hd 209458b, *Astron. Astrophys.* 595 (2016) A36. [doi:10.1051/0004-6361/201629183](https://doi.org/10.1051/0004-6361/201629183).
- [302] M. Min, Random sampling technique for ultra-fast computations of molecular opacities for exoplanet atmospheres, *Astron. Astrophys.* 607 (2017) A9. [doi:10.1051/0004-6361/201731612](https://doi.org/10.1051/0004-6361/201731612).
- [303] J. M. Goyal, N. Mayne, D. K. Sing, B. Drummond, P. Tremblin, D. S. Amundsen, T. Evans, A. L. Carter, J. Spake, I. Baraffe, N. Nikolov, J. Manners, G. Chabrier, E. Hebrard, A library of atmo forward model transmission spectra for hot jupiter exoplanets, *Mon. Not. Roy. Astron. Soc.* 474 (2018) 5158–5185. [doi:10.1093/mnras/stx3015](https://doi.org/10.1093/mnras/stx3015).
- [304] G. L. Villanueva, M. D. Smith, S. Protopapa, S. Faggi, A. M. Mandell, Planetary Spectrum Generator: An accurate online radiative transfer suite for atmospheres, comets, small bodies and exoplanets, *J. Quant. Spectrosc. Radiat. Transf.* 217 (2018) 86–104. [doi:10.1016/j.jqsrt.2018.05.023](https://doi.org/10.1016/j.jqsrt.2018.05.023).
- [305] S. Gandhi, M. Brogi, S. N. Yurchenko, J. Tennyson, P. A. Coles, R. K. Webb, J. L. Birkby, G. Guilluy, N. Madhusudhan, Molecular Cross

- Sections for High Resolution Spectroscopy of super Earths, warm Neptunes and Hot Jupiters, *Mon. Not. Roy. Astron. Soc.* 495 (2020) 224–237. [doi:10.1093/mnras/staa981](https://doi.org/10.1093/mnras/staa981).
- [306] S. L. Grimm, M. Malik, D. Kitzmann, A. Guzmán-Mesa, H. J. Hoeijmakers, C. Fisher, J. ao M. Mendonça, S. N. Yurchenko, J. Tennyson, R. L. Kurucz, K. Heng, HELIOS-K 2.0 and an Open-source Opacity Database for Exoplanetary Atmospheres, *Astrophys. J. Suppl. Ser.* 253 (2021) 30. [doi:10.3847/1538-4365/abd773](https://doi.org/10.3847/1538-4365/abd773).
- [307] P. E. E. Cubillos, J. Harrington, J. Bleicic, M. D. D. Himes, P. M. M. Rojo, T. J. J. Loredo, N. B. B. Lust, R. C. C. Challener, A. J. J. Foster, M. M. M. Stemm, A. S. D. Foster, S. D. D. Blumenthal, An Open-source Bayesian Atmospheric Radiative Transfer (BART) Code. II. The Transit Radiative Transfer Module and Retrieval of HAT-P-11b, *Planet. Sci. J.* 3 (2022) 81. [doi:10.3847/PSJ/ac348b](https://doi.org/10.3847/PSJ/ac348b).
- [308] S. Hirose, P. Hauschildt, T. Minoshima, K. Tomida, T. Sano, OPTAB: Public code for generating gas opacity tables for radiation hydrodynamics simulations, *Astron. Astrophys.* 659 (2022) A87. [doi:10.1051/0004-6361/202141076](https://doi.org/10.1051/0004-6361/202141076).
- [309] P. Marigo, B. Aringer, L. Girardi, A. Bressan, Updated Low-temperature Gas Opacities with AESOPUS 2.0, *Astrophys. J.* 940 (2022) 129. [doi:10.3847/1538-4357/ac9b40](https://doi.org/10.3847/1538-4357/ac9b40).
- [310] A. R. Iyer, M. R. Line, P. S. Muirhead, J. J. Fortney, E. Gharib-Nezhad, The SPHINX M-dwarf Spectral Grid. I. Benchmarking New Model Atmospheres to Derive Fundamental M-dwarf Properties, *Astrophys. J.* 944 (2023) 41. [doi:10.3847/1538-4357/acabc2](https://doi.org/10.3847/1538-4357/acabc2).
- [311] K. L. Chubb, S. Robert, C. Sousa-Silva, S. N. Yurchenko, N. F. Allard, V. Boudon, J. Buldyreva, B. Bultel, A. Coustenis, A. Foltynowicz, I. E. Gordon, R. J. Hargreaves, C. Hellings, C. Hill, H. R. Hrodmarsson, T. Karman, H. Lecoq-Molinos, A. Migliorini, M. Rey, C. Richard, I. Sadiek, F. Schmidt, A. Sokolov, S. Stefani, J. Tennyson, O. Venot, S. O. M. Wright, R. Arenales-Lope, J. K. Barstow, A. Bocchieri, N. Carrasco, D. Dubey, O. Egorov, A. G. Muñoz, Ehsan, Gharib-Nezhad, L. Gkouvelis, F. Grübel, P. G. J. Irwin, A. Knížek, D. A. Lewis, M. G. Lodge, S. Ma, Z. Martins, K. Molaverdikhani, G. Morello, A. Nikitin, E. Panek, M. Rengel, G. Rinaldi, J. W. Skinner, G. Tinetti, T. A. van Kempen, J. Yang, T. Zingales, Data availability and requirements relevant for the Ariel space mission and other exoplanet atmosphere applications, *RAS Tech. Instr.arXiv:2404.02188*.
- [312] E. V. Karlavets, I. E. Gordon, L. S. Rothman, R. Hashemi, R. J. Hargreaves, G. Toon, A. Campargue, V. I. Perevalov, P. Čermak, M. Birk, G. Wagner, J. T. Hodges, J. Tennyson, S. N. Yurchenko, The update of the line positions and intensities in the line list of carbon dioxide for the HITRAN2020 spectroscopic database, *J. Quant. Spectrosc. Radiat. Transf.* 276 (2021) 107896. [doi:10.1016/j.jqsrt.2021.107896](https://doi.org/10.1016/j.jqsrt.2021.107896).
- [313] L. S. Rothman, History of the HITRAN Database, *Nature Revs. Phys.* 3 (2021) 302–304. [doi:10.1038/s42254-021-00309-2](https://doi.org/10.1038/s42254-021-00309-2).
- [314] L. S. Rothman, I. E. Gordon, R. J. Barber, H. Dothe, R. R. Gamache, A. Goldman, V. I. Perevalov, S. A. Tashkun, J. Tennyson, HITEMP, the high-temperature molecular spectroscopic database, *J. Quant. Spectrosc. Radiat. Transf.* 111 (2010) 2139–2150. [doi:10.1016/j.jqsrt.2010.05.001](https://doi.org/10.1016/j.jqsrt.2010.05.001).
- [315] R. J. Hargreaves, I. E. Gordon, L. S. Rothman, S. A. Tashkun, V. I. Perevalov, A. A. Lukashevskaya, S. N. Yurchenko, J. Tennyson, H. S. P. Müller, Spectroscopic line parameters of NO, NO₂, and N₂O for the HITEMP database, *J. Quant. Spectrosc. Radiat. Transf.* 232 (2019) 35 – 53. [doi:10.1016/j.jqsrt.2019.04.040](https://doi.org/10.1016/j.jqsrt.2019.04.040).
- [316] M. Rey, A. V. Nikitin, Y. L. Babikov, V. G. Tyuterev, TheoReTS – an information system for theoretical spectra based on variational predictions from molecular potential energy and dipole moment surfaces, *J. Mol. Spectrosc.* 327 (2016) 138–158. [doi:10.1016/j.jms.2016.04.006](https://doi.org/10.1016/j.jms.2016.04.006).
- [317] X. Huang, R. S. Freedman, S. Tashkun, D. W. Schwenke, T. J. Lee, AI-3000K infrared line list for hot CO₂, *J. Mol. Spectrosc.* 392 (2023) 111748. [doi:10.1016/j.jms.2023.111748](https://doi.org/10.1016/j.jms.2023.111748).
- [318] D. S. Underwood, J. Tennyson, S. N. Yurchenko, X. Huang, D. W. Schwenke, T. J. Lee, S. Clausen, A. Fateev, ExoMol line lists XIV: A line list for hot SO₂, *Mon. Not. Roy. Astron. Soc.* 459 (2016) 3890–3899. [doi:10.1093/mnras/stw849](https://doi.org/10.1093/mnras/stw849).
- [319] P. F. Bernath, MOLLIST: Molecular Line Lists, Intensities and Spectra, *J. Quant. Spectrosc. Radiat. Transf.* 240 (2020) 106687. [doi:10.1016/j.jqsrt.2019.106687](https://doi.org/10.1016/j.jqsrt.2019.106687).
- [320] T. Ryabchikova, N. Piskunov, R. L. Kurucz, H. C. Stempels, U. Heiter, Y. Pakhomov, P. S. Barklem, A major upgrade of the VALD database, *Phys. Scr.* 90 (2015) 054005. [doi:\[10.1088/0031-8949/90/5/054005\]](https://doi.org/10.1088/0031-8949/90/5/054005).
- [321] R. L. Kurucz, Including all the lines, *Can. J. Phys.* 89 (2011) 417–428. [doi:10.1139/p10-104](https://doi.org/10.1139/p10-104).
- [322] H. R. Hrodmarsson, E. F. van Dishoeck, Photodissociation and photoionization of molecules of astronomical interest updates to the leiden photodissociation and photoionization cross section database, *Astron. Astrophys.* 675 (2023) A25. [doi:10.1051/0004-6361/202346645](https://doi.org/10.1051/0004-6361/202346645).
- [323] J. Tennyson, S. N. Yurchenko, The ExoMol atlas of molecular opacities, *Atoms* 6 (2018) 26. [doi:10.3390/atoms6020026](https://doi.org/10.3390/atoms6020026).
- [324] R. J. Barber, J. K. Strange, C. Hill, O. L. Polyansky, G. C. Mellau, S. N. Yurchenko, J. Tennyson, ExoMol line lists – III. An improved hot rotation-vibration line list for HCN and HNC, *Mon. Not. Roy. Astron. Soc.* 437 (2014) 1828–1835. [doi:10.1093/mnras/stt2011](https://doi.org/10.1093/mnras/stt2011).
- [325] E. J. Barton, C. Chiu, S. Golpayegani, S. N. Yurchenko, J. Tennyson, D. J. Froehman, P. F. Bernath, ExoMol Molecular linelists – V. The ro-vibrational spectra of NaCl and KCl, *Mon. Not. Roy. Astron. Soc.* 442 (2014) 1821–1829. [doi:10.1093/mnras/stu944](https://doi.org/10.1093/mnras/stu944).
- [326] C. Sousa-Silva, A. F. Al-Refaie, J. Tennyson, S. N. Yurchenko, ExoMol line lists - VII. the rotation-vibration spectrum of phosphine up to 1500 K, *Mon. Not. Roy. Astron. Soc.* 446 (2015) 2337–2347. [doi:10.1093/mnras/stu2246](https://doi.org/10.1093/mnras/stu2246).
- [327] A. F. Al-Refaie, S. N. Yurchenko, A. Yachmenev, J. Tennyson, ExoMol line lists - VIII: A variationally computed line list for hot formaldehyde, *Mon. Not. Roy. Astron. Soc.* 448 (2015) 1704–1714. [doi:10.1093/mnras/stv091](https://doi.org/10.1093/mnras/stv091).
- [328] A. T. Patrascu, J. Tennyson, S. N. Yurchenko, ExoMol molecular linelists - IX: The spectrum of AlO, *Mon. Not. Roy. Astron. Soc.* 449 (2015) 3613–3619. [doi:10.1093/mnras/stv507](https://doi.org/10.1093/mnras/stv507).
- [329] T. Rivlin, L. Lodi, S. N. Yurchenko, J. Tennyson, R. J. Le Roy, ExoMol line lists X: The spectrum of sodium hydride, *Mon. Not. Roy. Astron. Soc.* 451 (2015) 5153–5157. [doi:10.1093/mnras/stv979](https://doi.org/10.1093/mnras/stv979).
- [330] A. I. Pavlyuchko, S. N. Yurchenko, J. Tennyson, ExoMol line lists XI: A Hot Line List for nitric acid, *Mon. Not. Roy. Astron. Soc.* 452 (2015) 1702–1706. [doi:10.1093/mnras/stv1376](https://doi.org/10.1093/mnras/stv1376).
- [331] G. Paulose, E. J. Barton, S. N. Yurchenko, J. Tennyson, ExoMol Molecular linelists – XII. Line lists for eight isotopologues of CS, *Mon. Not. Roy. Astron. Soc.* 454 (2015) 1931–1939. [doi:10.1093/mnras/stv1543](https://doi.org/10.1093/mnras/stv1543).
- [332] S. N. Yurchenko, A. Blissett, U. Asari, M. Vasilios, C. Hill, J. Tennyson, ExoMol Molecular linelists – XIII. The spectrum of CaO, *Mon. Not. Roy. Astron. Soc.* 456 (2016) 4524–4532. [doi:10.1093/mnras/stv2858](https://doi.org/10.1093/mnras/stv2858).
- [333] A. F. Al-Refaie, O. L. Polyansky, R. I. Ovsyannikov, J. Tennyson, S. N. Yurchenko, ExoMol line lists XV: A hot line-list for hydrogen peroxide, *Mon. Not. Roy. Astron. Soc.* 461 (2016) 1012–1022. [doi:10.1093/mnras/stv1295](https://doi.org/10.1093/mnras/stv1295).
- [334] A. A. Azzam, S. N. Yurchenko, J. Tennyson, O. V. Naumenko, ExoMol line lists XVI: A Hot Line List for H₂S, *Mon. Not. Roy. Astron. Soc.* 460 (2016) 4063–4074. [doi:10.1093/mnras/stw1133](https://doi.org/10.1093/mnras/stw1133).

- [335] D. S. Underwood, J. Tennyson, S. N. Yurchenko, S. Clausen, A. Fateev, ExoMol line lists XVII: A line list for hot SO₃, Mon. Not. Roy. Astron. Soc. 462 (2016) 4300–4313. [doi:10.1093/mnras/stw1828](https://doi.org/10.1093/mnras/stw1828).
- [336] O. L. Polyansky, A. A. Kyuberis, L. Lodi, J. Tennyson, R. I. Ovsyannikov, N. Zobov, ExoMol molecular line lists XIX: high accuracy computed line lists for H₂¹⁷O and H₂¹⁸O, Mon. Not. Roy. Astron. Soc. 466 (2017) 1363–1371. [doi:10.1093/mnras/stw3125](https://doi.org/10.1093/mnras/stw3125).
- [337] I. I. Mizus, A. Aljah, N. F. Zobov, A. A. Kyuberis, S. N. Yurchenko, J. Tennyson, O. L. Polyansky, ExoMol molecular line lists XX: a comprehensive line list for H₃⁺, Mon. Not. Roy. Astron. Soc. 468 (2017) 1717–1725. [doi:10.1093/mnras/stx502](https://doi.org/10.1093/mnras/stx502).
- [338] A. Owens, S. N. Yurchenko, A. Yachmenev, W. Thiel, J. Tennyson, ExoMol molecular line lists XXII. The rotation-vibration spectrum of silane up to 1200 K, Mon. Not. Roy. Astron. Soc. 471 (2017) 5025–5032. [doi:10.1093/mnras/stx1952](https://doi.org/10.1093/mnras/stx1952).
- [339] L. Prajapat, P. Jagoda, L. Lodi, M. N. Gorman, S. N. Yurchenko, J. Tennyson, ExoMol molecular line lists XXIII. Spectra of PO and PS, Mon. Not. Roy. Astron. Soc. 472 (2017) 3648–3658. [doi:10.1093/mnras/stx2229](https://doi.org/10.1093/mnras/stx2229).
- [340] S. N. Yurchenko, F. Sinden, L. Lodi, C. Hill, M. N. Gorman, J. Tennyson, ExoMol Molecular linelists – XXIV: A new hot line list for silicon monohydride, SiH, Mon. Not. Roy. Astron. Soc. 473 (2018) 5324–5333. [doi:10.1093/mnras/stx2738](https://doi.org/10.1093/mnras/stx2738).
- [341] A. Upadhyay, E. K. Conway, J. Tennyson, S. N. Yurchenko, ExoMol Molecular linelists – XXV: A hot line list for silicon sulphide, SiS, Mon. Not. Roy. Astron. Soc. 477 (2018) 1520–1527. [doi:10.1093/mnras/sty998](https://doi.org/10.1093/mnras/sty998).
- [342] S. N. Yurchenko, W. Bond, M. N. Gorman, L. Lodi, L. K. McKemmish, W. Nunn, R. Shah, J. Tennyson, ExoMol Molecular linelists – XXVI: spectra of SH and NS, Mon. Not. Roy. Astron. Soc. 478 (2018) 270–282. [doi:10.1093/mnras/sty939](https://doi.org/10.1093/mnras/sty939).
- [343] B. P. Mant, A. Yachmenev, J. Tennyson, S. N. Yurchenko, ExoMol molecular line lists - XXVII: spectra of C₂H₄, Mon. Not. Roy. Astron. Soc. 478 (2018) 3220 – 3232. [doi:10.1093/mnras/sty1239](https://doi.org/10.1093/mnras/sty1239).
- [344] A. Owens, A. Yachmenev, J. Tennyson, W. Thiel, S. N. Yurchenko, ExoMol Molecular line lists XXIX: The rotation-vibration spectrum of methyl chloride up to 1200 K, Mon. Not. Roy. Astron. Soc. 479 (2018) 3002–3010. [doi:10.1093/mnras/sty1542](https://doi.org/10.1093/mnras/sty1542).
- [345] O. L. Polyansky, A. A. Kyuberis, N. F. Zobov, J. Tennyson, S. N. Yurchenko, L. Lodi, ExoMol molecular line lists XXX: a complete high-accuracy line list for water, Mon. Not. Roy. Astron. Soc. 480 (2018) 2597–2608. [doi:10.1093/mnras/sty1877](https://doi.org/10.1093/mnras/sty1877).
- [346] S. N. Yurchenko, I. Szabo, E. Pyatenko, J. Tennyson, ExoMol Molecular line lists XXXI: The spectrum of C₂, Mon. Not. Roy. Astron. Soc. 480 (2018) 3397–3411. [doi:10.1093/mnras/sty2050](https://doi.org/10.1093/mnras/sty2050).
- [347] H.-Y. Li, J. Tennyson, S. N. Yurchenko, ExoMol molecular line lists XXXII: the rovibronic spectrum of MgO, Mon. Not. Roy. Astron. Soc. 486 (2019) 2351–2365. [doi:10.1093/mnras/stz912](https://doi.org/10.1093/mnras/stz912).
- [348] L. K. McKemmish, T. Masseron, J. Hoeijmakers, V. V. Pérez-Mesa, S. L. Grimm, S. N. Yurchenko, J. Tennyson, ExoMol Molecular line lists – XXXIII. The spectrum of Titanium Oxide, Mon. Not. Roy. Astron. Soc. 488 (2019) 2836–2854. [doi:10.1093/mnras/stz1818](https://doi.org/10.1093/mnras/stz1818).
- [349] J. Langleben, , S. N. Yurchenko, J. Tennyson, ExoMol line list XXXIV: A Rovibrational Line List for Phosphinidene (PH) in its X ³Σ⁻ and a ¹Δ Electronic States, Mon. Not. Roy. Astron. Soc. 488 (2019) 2332. [doi:10.1093/mnras/stz1856-2342](https://doi.org/10.1093/mnras/stz1856-2342).
- [350] P. A. Coles, , S. N. Yurchenko, J. Tennyson, ExoMol molecular line lists XXXV: a rotation-vibration line list for hot ammonia, Mon. Not. Roy. Astron. Soc. 490 (2019) 4638–4647. [doi:10.1093/mnras/stz2778](https://doi.org/10.1093/mnras/stz2778).
- [351] M. Gorman, S. N. Yurchenko, J. Tennyson, ExoMol Molecular linelists – XXXVI. X ²Π – X ²Π and A ²Σ⁺ – X ²Π transitions of SH, Mon. Not. Roy. Astron. Soc. 490 (2019) 1652–1665. [doi:10.1093/mnras/stz2517/5565070](https://doi.org/10.1093/mnras/stz2517/5565070).
- [352] K. L. Chubb, J. Tennyson, S. N. Yurchenko, ExoMol Molecular linelists – XXXVII: spectra of acetylene, Mon. Not. Roy. Astron. Soc. 493 (2020) 1531–1545. [doi:10.1093/mnras/staa229](https://doi.org/10.1093/mnras/staa229).
- [353] A. Owens, E. K. Conway, J. Tennyson, S. N. Yurchenko, ExoMol Molecular linelists – XXXVIII: High-temperature molecular line list of silicon dioxide (SiO₂), Mon. Not. Roy. Astron. Soc. 495 (2020) 1927–1933. [doi:10.1093/mnras/staa1287](https://doi.org/10.1093/mnras/staa1287).
- [354] S. N. Yurchenko, T. M. Mellor, R. S. Freedman, J. Tennyson, ExoMol molecular line lists XXXIX: Ro-vibrational molecular line list for CO₂, Mon. Not. Roy. Astron. Soc. 496 (2020) 5282–5291. [doi:10.1093/mnras/staa1874](https://doi.org/10.1093/mnras/staa1874).
- [355] S. N. Yurchenko, J. Tennyson, S. Miller, V. V. Melnikov, J. O'Donoghue, L. Moore, ExoMol molecular line lists XL: Ro-vibrational molecular line list for the hydronium ion (H₃O⁺), Mon. Not. Roy. Astron. Soc. 497 (2020) 2340–2351. [doi:10.1093/mnras/staa2034](https://doi.org/10.1093/mnras/staa2034).
- [356] A. Owens, J. Tennyson, S. N. Yurchenko, ExoMol line lists – XLI. High-temperature molecular line lists for the alkali metal hydroxides KOH and NaOH, Mon. Not. Roy. Astron. Soc. 502 (2021) 1128–1135. [doi:10.1093/mnras/staa4041](https://doi.org/10.1093/mnras/staa4041).
- [357] Q. Qu, S. N. Yurchenko, J. Tennyson, ExoMol molecular line lists – XLII: Rovibronic molecular line list for the low-lying states of NO, Mon. Not. Roy. Astron. Soc. 504 (2021) 5768–5777. [doi:10.1093/mnras/stab1154](https://doi.org/10.1093/mnras/stab1154).
- [358] G. B. Mitev, S. Taylor, J. Tennyson, S. N. Yurchenko, A. A. Buchachenko, A. V. Stolyarov, ExoMol molecular line lists – XLIII: Rovibronic transitions corresponding to the close-lying X ²Π and A ²Σ⁺ states of NaO, Mon. Not. Roy. Astron. Soc. 511 (2022) 2349–2355. [doi:10.1093/mnras/stab3357](https://doi.org/10.1093/mnras/stab3357).
- [359] S. N. Yurchenko, J. Tennyson, A.-M. Syme, A. Y. Adam, V. H. J. Clark, B. Cooper, C. P. Dobney, S. T. E. Donnelly, M. N. Gorman, A. E. Lynas-Gray, T. Meltzer, A. Owens, Q. Qu, M. Semenov, W. Somogyi, A. Upadhyay, S. Wright, J. C. Zapata Trujillo, ExoMol line lists – XLIV. IR and UV line list for silicon monoxide (²⁸Si¹⁶O), Mon. Not. Roy. Astron. Soc. 510 (2022) 903–919. [doi:10.1093/mnras/stab3267](https://doi.org/10.1093/mnras/stab3267).
- [360] A. Owens, S. Dooley, L. McLaughlin, B. Tan, G. Zhang, S. N. Yurchenko, J. Tennyson, ExoMol line lists – XLV. Rovibronic molecular line lists of calcium monohydride (CaH) and magnesium monohydride (MgH), Mon. Not. Roy. Astron. Soc. 511 (2022) 5448–5461. [doi:10.1093/mnras/stac371](https://doi.org/10.1093/mnras/stac371).
- [361] T. Mellor, A. Owens, S. N. Yurchenko, J. Tennyson, ExoMol line lists – LI. Rvibronic molecular line list for thioformaldehyde (H₂CS), Mon. Not. Roy. Astron. Soc. 520 (2022) 1997–2008. [doi:10.1093/mnras/stad111](https://doi.org/10.1093/mnras/stad111).
- [362] S. N. Yurchenko, E. Nogué, A. A. A. Azzam, J. Tennyson, ExoMol line lists – IL. Rovibronic spectrum of aluminium monochloride (AlCl), Mon. Not. Roy. Astron. Soc. 520 (2023) 5183–5191. [doi:10.1093/mnras/stac3757](https://doi.org/10.1093/mnras/stac3757).
- [363] C. A. Bowesman, I. I. Mizus, N. F. Zobov, O. L. Polyansky, J. Sarka, B. Poirier, M. Pezzella, S. N. Yurchenko, J. Tennyson, ExoMol line lists - L: High-resolution line lists of H₃⁺, H₂D⁺, D₂H⁺ and D₃⁺, Mon. Not. Roy. Astron. Soc. 519 (2023) 6333–6348. [doi:10.1093/mnras/stad050](https://doi.org/10.1093/mnras/stad050).
- [364] A. Owens, S. O. M. Wright, Y. Pavlenko, A. Mitrushchenkov, J. Koput, S. N. Yurchenko, J. Tennyson, ExoMol line lists - LI. Molecular line list for lithium hydroxide (LiOH), Mon. Not. Roy. Astron. Soc. 527 (2024) 731–738. [doi:10.1093/mnras/stad3226](https://doi.org/10.1093/mnras/stad3226).
- [365] O. Pearce, S. N. Yurchenko, J. Tennyson, ExoMol line lists - LII. Line Lists for the Methyldyne Cation (CH⁺), Mon. Not. Roy. Astron. Soc. 527 (2024) 10726–10736. [doi:10.1093/mnras/stad3909](https://doi.org/10.1093/mnras/stad3909).
- [366] S. N. Yurchenko, R. P. Brady, J. Tennyson, A. N. Smirnov, O. A. Vasilyev, V. G. Solomonik, ExoMol line lists – LIII. Empirical Rovibronic spectra Yttrium Oxide (YO), Mon. Not. Roy. Astron. Soc. 527 (2024) 4899–4912. [doi:10.1093/mnras/stad3225](https://doi.org/10.1093/mnras/stad3225).
- [367] C. A. Bowesman, Q. Qu, L. K. McKemmish, S. N. Yurchenko, J. Tennyson, ExoMol line lists - LV: Hyperfine-resolved molecular line list for

- vanadium monoxide ($^{51}\text{V}^{16}\text{O}$), Mon. Not. Roy. Astron. Soc. 529 (2024) 1321–1332. [doi:10.1093/mnras/stae542](https://doi.org/10.1093/mnras/stae542).
- [368] R. P. Brady, S. N. Yurchenko, J. Tennyson, G.-S. Kim, ExoMol line lists – LVI. The SO line list, MARVEL analysis of experimental transition data and refinement of the spectroscopic model, Mon. Not. Roy. Astron. Soc. 527 (2024) 6675–6690. [doi:10.1093/mnras/stad3508](https://doi.org/10.1093/mnras/stad3508).
- [369] K. Kefala, V. Boudon, S. N. Yurchenko, J. Tennyson, Empirical rovibrational energy levels for methane, J. Quant. Spectrosc. Radiat. Transf. 316 (2024) 108897. [doi:10.1016/j.jqsrt.2024.108897](https://doi.org/10.1016/j.jqsrt.2024.108897).
- [370] A. Owens, , S. N. Yurchenko, J. Tennyson, ExoMol line lists - LVIII. High-temperature molecular line list of carbonyl sulphide (OCS), Mon. Not. Roy. Astron. Soc. 530 (2024) 4004–4015. [doi:10.1093/mnras/stae110](https://doi.org/10.1093/mnras/stae110).
- [371] S. N. Yurchenko, T. Mellor, J. Tennyson, ExoMol line lists – LIX. High-temperature line list for N_2O , Mon. Not. Roy. Astron. Soc.
- [372] S. N. Yurchenko, C. A. Bowesman, R. P. Brady, E. R. Guest, K. Kefala, G. B. Mitev, A. Owens, A. N. Perri, M. Pezzella, O. Smola, A. Solokov, J. Zhang, J. Tennyson, ExoMol Line Lists – LX: Molecular line list for the ammonia isotopologue $^{15}\text{NH}_3$, Mon. Not. Roy. Astron. Soc.
- [373] G. B. Mitev, C. A. Bowesman, S. N. Yurchenko, J. Tennyson, ExoMol Line Lists – LXI: A trihybrid linelist for rovibronic transitions of the hydroxyl radical (OH), Mon. Not. Roy. Astron. Soc.
- [374] A. E. Lynas-Gray, O. L. Polyansky, J. Tennyson, S. N. Yurchenko, N. F. Zobov, ExoMol Line Lists – LXII: Ro-Vibrational Energy Levels and Line-Strengths for the Propadienediylidene (C_3) electronic Ground-State, Mon. Not. Roy. Astron. Soc.
- [375] I. I. Mizus, N. F. Zobov, M. Pezzella, O. V. Boyarkin, M. A. Koshelev, D. S. Makarov, S. N. Yurchenko, J. Tennyson, O. L. Polyansky, ExoMol Line Lists – LXIII: Ro-Vibrational Energy Levels and Line-Strengths HDO, Mon. Not. Roy. Astron. Soc.
- [376] M. Semenov, N. El-Kork, S. N. Yurchenko, J. Tennyson, ExoMol Line Lists – LXIV: Empirical rovibronic spectra of phosphorous mononitride (PN) covering the IR and UV regions, Mon. Not. Roy. Astron. Soc.
- [377] G. J. Harris, F. C. Larmer, J. Tennyson, B. M. Kaminsky, Y. V. Pavlenko, H. R. A. Jones, A $\text{H}^{13}\text{CN}/\text{HN}^{13}\text{C}$ linelist, model atmospheres and synthetic spectra for carbon stars, Mon. Not. Roy. Astron. Soc. 390 (2008) 143–148.
- [378] J. Tennyson, Accurate variational calculations for line lists to model the vibration rotation spectra of hot astrophysical atmospheres, WIREs Comput. Mol. Sci. 2 (2012) 698–715. [doi:10.1002/wcms.94](https://doi.org/10.1002/wcms.94).
- [379] J. Tennyson, P. F. Bernath, L. R. Brown, A. Campargue, M. R. Carleer, A. G. Császár, R. R. Gamache, J. T. Hodges, A. Jenouvrier, O. V. Naumenko, O. L. Polyansky, L. S. Rothman, R. A. Toth, A. C. Vandaele, N. F. Zobov, L. Daumont, A. Z. Fazliev, T. Furtenbacher, I. E. Gordon, S. N. Mikhailenko, S. V. Shirin, IUPAC critical evaluation of the rotational-vibrational spectra of water vapor. Part I. Energy levels and transition wavenumbers for H_2^{17}O and H_2^{18}O , J. Quant. Spectrosc. Radiat. Transf. 110 (2009) 573–596. [doi:10.1016/j.jqsrt.2009.02.014](https://doi.org/10.1016/j.jqsrt.2009.02.014).
- [380] J. Tennyson, P. F. Bernath, L. R. Brown, A. Campargue, A. G. Császár, L. Daumont, R. R. Gamache, J. T. Hodges, O. V. Naumenko, O. L. Polyansky, L. S. Rothman, A. C. Vandaele, N. F. Zobov, A Database of Water Transitions from Experiment and Theory (IUPAC Technical Report), Pure Appl. Chem. 86 (2014) 71–83. [doi:10.1515/pac-2014-5012](https://doi.org/10.1515/pac-2014-5012).
- [381] C. A. Bowesman, H. Akbari, S. Hopkins, S. N. Yurchenko, J. Tennyson, Fine and hyperfine resolved empirical energy levels for VO, J. Quant. Spectrosc. Radiat. Transf. 289 (2022) 108295. [doi:10.1016/j.jqsrt.2022.108295](https://doi.org/10.1016/j.jqsrt.2022.108295).
- [382] A. G. Császár, T. Furtenbacher, Spectroscopic networks, J. Mol. Spectrosc. 266 (2011) 99 – 103. [doi:10.1016/j.jms.2011.03.031](https://doi.org/10.1016/j.jms.2011.03.031).
- [383] J.-M. Flaud, C. Camy-Peyret, J.-P. Maillard, Higher Ro-Vibrational Levels Of H_2O Deduced From High-Resolution Oxygen-Hydrogen Flame Spectra Between 2800-6200 cm^{-1} , Mol. Phys. 32 (1976) 499–521. [doi:10.1080/00268977600103251](https://doi.org/10.1080/00268977600103251).
- [384] A. R. Al-Derzi, S. N. Yurchenko, J. Tennyson, M. Melosso, N. Jiang, C. Puzzarini, L. Dore, T. Furtenbacher, R. Tobias, A. G. Császár, An improved rovibrational linelist of formaldehyde, $\text{H}_2^{12}\text{C}^{16}\text{O}$, J. Quant. Spectrosc. Radiat. Transf. 266 (2021) 107563. [doi:10.1016/j.jqsrt.2021.107563](https://doi.org/10.1016/j.jqsrt.2021.107563).
- [385] M. Germann, A. Hjalten, J. Tennyson, S. N. Yurchenko, I. E. Gordon, C. Pett, I. Silander, K. Krzempek, A. Hudzikowski, A. Gluszek, G. Sobon, A. Foltynowicz, Optical frequency comb Fourier transform spectroscopy of formaldehyde in the 1250 to 1390 cm^{-1} range: experimental line list and MARVEL analysis, J. Quant. Spectrosc. Radiat. Transf. 312 (2024) 108782. [doi:10.1016/j.jqsrt.2023.108782](https://doi.org/10.1016/j.jqsrt.2023.108782).
- [386] S. N. Yurchenko, A. F. Al-Refaie, J. Tennyson, ExoCross: A general program for generating spectra from molecular line lists, Astron. Astrophys. 614 (2018) A131. [doi:10.1051/0004-6361/201732531](https://doi.org/10.1051/0004-6361/201732531).
- [387] J. Zhang, J. Tennyson, S. N. Yurchenko, PyExoCross: a Python program for generating spectra and cross-sections from molecular line lists, RAS Tech. 3 (2024) 257–287 Instr. [doi:10.1093/rasti/rzae016](https://doi.org/10.1093/rasti/rzae016).
- [388] L. Lodi, J. Tennyson, Line lists for H_2^{18}O and H_2^{17}O based on empirically-adjusted line positions and ab initio intensities, J. Quant. Spectrosc. Radiat. Transf. 113 (2012) 850–858. [doi:10.1016/j.jqsrt.2012.02.023](https://doi.org/10.1016/j.jqsrt.2012.02.023).
- [389] E. J. Zak, J. Tennyson, O. L. Polyansky, L. Lodi, S. A. Tashkun, V. I. Perevalov, A room temperature CO_2 line list with *ab initio* computed intensities, J. Quant. Spectrosc. Radiat. Transf. 177 (2016) 31–42. [doi:10.1016/j.jqsrt.2015.12.022](https://doi.org/10.1016/j.jqsrt.2015.12.022).
- [390] C. A. Bowesman, M. Shuai, S. N. Yurchenko, J. Tennyson, A high resolution line list for AlO, Mon. Not. Roy. Astron. Soc. 508 (2021) 3181–3193. [doi:10.1093/mnras/stab2525](https://doi.org/10.1093/mnras/stab2525).
- [391] K. L. Chubb, M. Min, Y. Kawashima, C. Helling, I. Waldmann, Aluminium oxide in the atmosphere of hot Jupiter WASP-43b, Astron. Astrophys. 639 (2020) A3. [doi:10.1051/0004-6361/201937267](https://doi.org/10.1051/0004-6361/201937267).
- [392] A. A. Bol'shakov, X. Mao, R. E. Russo, Spectral emission enhancement by an electric pulse for LIBS and LAMIS, J. Anal. At. Spectrom. 32 (2017) 657–670. [doi:10.1039/c6ja00436a](https://doi.org/10.1039/c6ja00436a).
- [393] H. Hou, X. Mao, V. Zorba, R. E. Russo, Laser Ablation Molecular Isotopic Spectrometry for Molecules Formation Chemistry in Femtosecond-Laser Ablated Plasmas, Anal. Chem. 89 (2017) 7750–7757. [doi:10.1021/acs.analchem.7b01750](https://doi.org/10.1021/acs.analchem.7b01750).
- [394] A. Wong, S. N. Yurchenko, P. Bernath, H. S. P. Mueller, S. McConkey, J. Tennyson, ExoMol Line List XXI: Nitric Oxide (NO), Mon. Not. Roy. Astron. Soc. 470 (2017) 882–897. [doi:10.1093/mnras/stx1211](https://doi.org/10.1093/mnras/stx1211).
- [395] G. N. Tsurikov, D. V. Bisikalo, NO Biomarker: Transmission and Emission Methods for Its Potential Detection in Exoplanet Atmospheres with Spektr-UF (WSO-UV), Astron. Rep. 67 (2023) 1123–1138. [doi:10.1134/S1063772923110100](https://doi.org/10.1134/S1063772923110100).
- [396] Y. Pavlenko, J. Tennyson, S. N. Yurchenko, M. R. Schmidt, H. R. A. Jones, Y. Lyubchik, A. Suárez Mascareño, AlH lines in the blue spectrum of Proxima Centauri, Mon. Not. Roy. Astron. Soc. 516 (2022) 5655–5673. [doi:10.1093/mnras/stac2588](https://doi.org/10.1093/mnras/stac2588).
- [397] S. N. Yurchenko, H. Williams, P. C. Leyland, L. Lodi, J. Tennyson, ExoMol line lists XXVIII: The rovibronic spectrum of AlH, Mon. Not. Roy. Astron. Soc. 479 (2018) 1401–1411. [doi:10.1093/mnras/sty1524](https://doi.org/10.1093/mnras/sty1524).
- [398] A. Falco, P. Tremblin, S. Charnoz, R. J. Ridgway, P.-O. Lagage, Hydrogenated atmospheres of lava planets: atmospheric structure and emission spectra, Astron. Astrophys. 683 (2024) A194. [doi:10.1051/0004-6361/202347650](https://doi.org/10.1051/0004-6361/202347650).

- [399] Y. V. Pavlenko, S. N. Yurchenko, L. K. McKemmish, J. Tennyson, Analysis of the TiO isotopologues in stellar optical spectra, *Astron. Astrophys.* 42 (2020) A77. [doi:10.1051/0004-6361/202037863](https://doi.org/10.1051/0004-6361/202037863).
- [400] B. Edwards, Q. Changeat, R. Baeyens, A. Tsiaras, A. Al-Refaie, J. Taylor, K. H. Yip, M. F. Bieger, D. Blain, A. Gressier, et al., ARES I: WASP-76 b, A Tale of Two HST Spectra, *Astrophys. J.* 160 (2020) 8. [doi:10.3847/1538-3881/ab9225](https://doi.org/10.3847/1538-3881/ab9225).
- [401] M. N. Gorman, S. N. Yurchenko, J. Tennyson, *ExoMol molecular line lists XXXVI: X²Π-X²Π and A²Σ-X²Π transitions of SH*, *Mon. Not. Roy. Astron. Soc.* 490 (2019) 1652–1665. [doi:10.1093/mnras/stz2517](https://doi.org/10.1093/mnras/stz2517). URL <https://doi.org/10.1093/mnras/stz2517>
- [402] E. J. Barton, S. N. Yurchenko, J. Tennyson, ExoMol Molecular linelists – II. The ro-vibrational spectrum of SiO, *Mon. Not. Roy. Astron. Soc.* 434 (2013) 1469–1475. [doi:10.1093/mnras/stt1105](https://doi.org/10.1093/mnras/stt1105).
- [403] Y. Ito, M. Ikoma, H. Kawahara, H. Nagahara, Y. Kawashima, T. Nakamoto, THEORETICAL EMISSION SPECTRA OF ATMOSPHERES OF HOT ROCKY SUPER-EARTHS, *Astrophys. J.* 801 (2015) 144. [doi:10.1088/0004-637X/801/2/144](https://doi.org/10.1088/0004-637X/801/2/144).
- [404] T. G. Nguyen, N. B. Cowan, R. T. Pierrehumbert, R. E. Lupu, J. E. Moores, The impact of ultraviolet heating and cooling on the dynamics and observability of lava planet atmospheres, *Mon. Not. Roy. Astron. Soc.* 513 (2022) 6125–6133. [doi:10.1093/mnras/stac1331](https://doi.org/10.1093/mnras/stac1331).
- [405] B. Yadin, T. Vaness, P. Conti, C. Hill, S. N. Yurchenko, J. Tennyson, ExoMol Molecular linelists: I The rovibrational spectrum of BeH, MgH and CaH the X²Σ⁺ state, *Mon. Not. Roy. Astron. Soc.* 425 (2012) 34–43.
- [406] D. Darby-Lewis, J. Tennyson, K. D. Lawson, S. N. Yurchenko, M. F. Stamp, A. Shaw, S. Brezinsek, JET Contributor, Synthetic spectra of BeH, BeD and BeT for emission modelling in JET plasmas, *J. Phys. B-At. Mol. Opt. Phys.* 51 (2018) 185701. [doi:10.1088/1361-6455/aad6d0](https://doi.org/10.1088/1361-6455/aad6d0).
- [407] L. K. McKemmish, S. N. Yurchenko, J. Tennyson, ExoMol line lists – XVIII. The high-temperature spectrum of VO, *Mon. Not. Roy. Astron. Soc.* 463 (2016) 771–793. [doi:10.1093/mnras/stw1969](https://doi.org/10.1093/mnras/stw1969).
- [408] Q. Qu, S. N. Yurchenko, J. Tennyson, A Method for the Variational Calculation of Hyperfine-Resolved Rovibronic Spectra of Diatomic Molecules, *J. Chem. Theory Comput.* 18 (2022) 1808–1820. [doi:10.1021/acs.jctc.1c01244](https://doi.org/10.1021/acs.jctc.1c01244).
- [409] Q. Qu, S. N. Yurchenko, J. Tennyson, A variational model for the hyperfine resolved spectrum of VO in its ground electronic state, *J. Chem. Phys.* 157 (2022) 124305. [doi:10.1063/5.0105965](https://doi.org/10.1063/5.0105965).
- [410] C. A. Bowesman, S. N. Yurchenko, J. Tennyson, A hyperfine-resolved spectroscopic model for vanadium monoxide (⁵¹V¹⁶O), *Mol. Phys.* 122 (2024) e2255299. [doi:10.1080/00268976.2023.2255299](https://doi.org/10.1080/00268976.2023.2255299).
- [411] R. P. Brady, S. N. Yurchenko, G.-S. Kim, W. Somogyi, J. Tennyson, An ab initio study of the rovibronic spectrum of sulphur monoxide (SO): diabatic vs. adiabatic representation, *Phys. Chem. Chem. Phys.* 24 (2022) 24076–24088. [doi:10.1039/D2CP03051A](https://doi.org/10.1039/D2CP03051A).
- [412] E. Brugamyer, S. E. Dodson-Robinson, W. D. Cochran, C. Sneden, SILICON AND OXYGEN ABUNDANCES IN PLANET-HOST STARS, *Astrophys. J.* 738 (2011) 97. [doi:10.1088/0004-637X/738/1/97](https://doi.org/10.1088/0004-637X/738/1/97).
- [413] C. Bordiu, J. R. Rizzo, F. Bufano, G. Quintana-Lacaci, C. Buemi, P. Leto, F. Cavallaro, L. Cerrigone, A. Ingallinera, S. Loru, S. Riggi, C. Trigilio, G. Umana, E. Sciacca, First Detection of Silicon-bearing Molecules in eta Car, *Astrophys. J. Lett.* 939 (2022) L30. [doi:10.3847/2041-8213/ac9b10](https://doi.org/10.3847/2041-8213/ac9b10).
- [414] P. Schilke, S. Leurini, K. Menten, J. Alcolea, Interstellar SiN, *Astron. Astrophys.* 412 (2003) L15–L18. [doi:{10.1051/0004-6361:20031649}](https://doi.org/10.1051/0004-6361:20031649).
- [415] J. L. Turner, A. Dalgarno, Chemistry Of Silicon In Interstellar Clouds, *Astrophys. J.* 213 (1977) 386–389. [doi:{10.1086/155167}](https://doi.org/10.1086/155167).
- [416] P. A. Feldman, H. E. Matthews, M. B. Bell, G. Herzberg, S. Saito, Y. Endo, E. Hirota, Limits To Interstellar SiN, *Journal of the Royal Astronomical Society of Canada* 77 (1983) 258.
- [417] D. N. Davis, The Spectrum Of Beta-Pegasi, *Astron. J.* 106 (1947) 28-&. [doi:{10.1086/144938}](https://doi.org/10.1086/144938).
- [418] L. Gratton, Studies Of The Spectra Of K-Giants .1. A Table Of Wave Lengths And Identifications Of Spectral Lines In The Region Lambda-Lambda-4000 5000, *Astron. J.* 115 (1952) 346–401. [doi:{10.1086/145554}](https://doi.org/10.1086/145554).
- [419] B. E. Turner, Detection of SiN in IRC + 10216, *Astrophys. J.* 388 (1992) L35. [doi:10.1086/186324](https://doi.org/10.1086/186324).
- [420] M. Yousefi, P. F. Bernath, J. Hodges, T. Mässerönen, A new line list for the A²Σ⁺ – X²Π electronic transition of OH, *J. Quant. Spectrosc. Radiat. Transf.* 217 (2018) 416–424. [doi:10.1016/j.jqsrt.2018.06.016](https://doi.org/10.1016/j.jqsrt.2018.06.016).
- [421] T. Furtenbacher, S. T. Hegedus, J. Tennyson, A. G. Császár, Analysis of the measured high-resolution doublet rovibronic spectra of ¹²CH and ¹⁶OH, *Phys. Chem. Chem. Phys.* 24 (2022) 19287. [doi:10.1039/D2CP02240K](https://doi.org/10.1039/D2CP02240K).
- [422] L. Yorke, S. N. Yurchenko, L. Lodi, J. Tennyson, ExoMol line lists VI: A high temperature line list for Phosphorus Nitride, *Mon. Not. Roy. Astron. Soc.* 445 (2014) 1383–1391. [doi:10.1093/mnras/stu1854](https://doi.org/10.1093/mnras/stu1854).
- [423] M. Semenov, N. El-Kork, S. N. Yurchenko, J. Tennyson, Rovibronic spectroscopy of PN from first principles, *Phys. Chem. Chem. Phys.* 23 (2021) 22057–220662. [doi:10.1039/D1CP02537F](https://doi.org/10.1039/D1CP02537F).
- [424] T. Sochi, J. Tennyson, A computed line list for the H₂D⁺ molecular ion, *Mon. Not. Roy. Astron. Soc.* 405 (2010) 2345–2350.
- [425] S. Miller, T. R. Geballe, T. Stallard, J. Tennyson, Thirty years of H₃⁺ astronomy, *Rev. Mod. Phys.* 92 (2020) 035003. [doi:10.1103/RevModPhys.92.035003](https://doi.org/10.1103/RevModPhys.92.035003).
- [426] R. J. Barber, J. Tennyson, G. J. Harris, R. N. Tolchenov, A high accuracy computed water line list, *Mon. Not. Roy. Astron. Soc.* 368 (2006) 1087–1094.
- [427] H. Partridge, D. W. Schwenke, The determination of an accurate isotope dependent potential energy surface for water from extensive ab initio calculations and experimental data, *J. Chem. Phys.* 106 (1997) 4618–4639. [doi:10.1063/1.473987](https://doi.org/10.1063/1.473987).
- [428] S. T. Melin, S. T. Sanders, E. F. Nasir, Comparison of ExoMol simulated spectra for H₂O to high-temperature low-pressure gas cell measurements at 1723K in the 7321–7598 cm(⁻¹) range, *J. Quant. Spectrosc. Radiat. Transf.* 253 (2020) 107079. [doi:{10.1016/j.jqsrt.2020.107079}](https://doi.org/10.1016/j.jqsrt.2020.107079).
- [429] T. Furtenbacher, R. Tóbiás, J. Tennyson, O. L. Polyansky, A. G. Császár, W2020: A Database of Validated Rovibrational Experimental Transitions and Empirical Energy Levels of H₂¹⁶O, *J. Phys. Chem. Ref. Data* 49 (2020) 033101. [doi:10.1063/5.0008253](https://doi.org/10.1063/5.0008253).
- [430] T. Furtenbacher, R. Tóbiás, J. Tennyson, O. L. Polyansky, A. A. Kyuberis, R. I. Ovsyannikov, N. F. Zobov, A. G. Császár, W2020: A Database of Validated Rovibrational Experimental Transitions and Empirical Energy Levels Part II. H₂¹⁷O and H₂¹⁸O with an Update to H₂¹⁶O, *J. Phys. Chem. Ref. Data* 49 (2020) 043103. [doi:10.1063/5.0030680](https://doi.org/10.1063/5.0030680).
- [431] B. A. Voronin, J. Tennyson, R. N. Tolchenov, A. A. Lugovskoy, S. N. Yurchenko, A high accuracy computed line list for the HDO molecule, *Mon. Not. Roy. Astron. Soc.* 402 (2010) 492–496.
- [432] N. F. Zobov, M. A. Koshelev, D. S. Makarov, O. V. Boyarkin, V. G. Tyuterev, J. Tennyson, O. L. Polyansky, A global line list for HDO

- between 0 and 35000 cm⁻¹ constructed using multiphoton spectra, *J. Quant. Spectrosc. Radiat. Transf.* 271 (2021) 107694. doi: [10.1016/j.jqsrt.2021.107694](https://doi.org/10.1016/j.jqsrt.2021.107694).
- [433] J. Tennyson, P. F. Bernath, L. R. Brown, A. Campargue, A. G. Császár, L. Daumont, R. R. Gamache, J. T. Hodges, O. V. Naumenko, O. L. Polyansky, L. S. Rothman, R. A. Toth, A. C. Vandaele, N. F. Zobov, S. Fally, A. Z. Fazliev, T. Furtenbacher, I. E. Gordon, S. N. Mikhailyenko, B. A, IUPAC critical evaluation of the rotational-vibrational spectra of water vapor. Part II. Energy levels and transition wavenumbers for HD¹⁶O, HD¹⁷O, and HD¹⁸O, *J. Quant. Spectrosc. Radiat. Transf.* 111 (2010) 2160–2184. doi: [10.1016/j.jqsrt.2010.06.012](https://doi.org/10.1016/j.jqsrt.2010.06.012).
- [434] B. A. Voronin, J. Tennyson, T. Y. Chesnokova, A. V. Chentsov, A. D. Bykov, The infrared absorption spectrum of the H₂¹⁴O radioactive isotopologue of water, *Mol. Phys.* doi: [10.1080/00268976.2024.233474](https://doi.org/10.1080/00268976.2024.233474).
- [435] B. A. Voronin, J. Tennyson, S. N. Yurchenko, T. Y. Chesnokova, A. V. Chentsov, A. D. Bykov, M. V. Makarova, S. S. Voronina, F. C. Cruz, The infrared absorption spectrum of radioactive water isotopologue H₂¹⁵O, *Spectroc. Acta Pt. A-Molec. Biomolec. Spectr.* 311 (2024) 124007. doi: [10.1016/j.saa.2024.124007](https://doi.org/10.1016/j.saa.2024.124007).
- [436] B. A. Voronin, J. Tennyson, S. N. Yurchenko, T. Y. Chesnokova, A. V. Chentsov, A. D. Bykov, F. C. Cruz, The spectrum of radioactive water vapor: the H₂¹⁹O radio-isotopologue, *J. Radioanal. Nucl. Chem.*
- [437] B. A. Voronin, M. V. Makarova, A. V. Poberovskii, A. D. Bykov, E. A. Dudnikova, J. Tennyson, The absorption spectrum of short-lived isotopic variant of water, H₂¹⁵O: Tentative detection at the Earth's atmosphere, *J. Quant. Spectrosc. Radiat. Transf.* 276 (2021) 107929. doi: [10.1016/j.jqsrt.2021.107929](https://doi.org/10.1016/j.jqsrt.2021.107929).
- [438] X. Huang, D. W. Schwenke, S. A. Tashkun, T. J. Lee, An isotopic-independent highly accurate potential energy surface for CO₂ isotopologues and an initial ¹²C¹⁶O₂ infrared line list, *J. Chem. Phys.* 136 (2012) 124311. doi: [10.1063/1.3697540](https://doi.org/10.1063/1.3697540).
- [439] O. L. Polyansky, K. Bielska, M. Ghysels, L. Lodi, N. F. Zobov, J. T. Hodges, J. Tennyson, High accuracy CO₂ line intensities determined from theory and experiment, *Phys. Rev. Lett.* 114 (2015) 243001. doi: [10.1103/PhysRevLett.114.243001](https://doi.org/10.1103/PhysRevLett.114.243001).
- [440] S. N. Yurchenko, W. Thiel, P. Jensen, Theoretical ROVibrational Energies (TROVE): A robust numerical approach to the calculation of rovibrational energies for polyatomic molecules, *J. Mol. Spectrosc.* 245 (2007) 126–140. doi: [10.1016/j.jms.2007.07.009](https://doi.org/10.1016/j.jms.2007.07.009).
- [441] A. A. Balashov, A. Raj, S. Wójtewicz, R. Ciuryłło, D. Lisak, K. Bielska, Theoretically predicted CO₂ lines near 700 nm not observed, *J. Quant. Spectrosc. Radiat. Transf.* (2024) 108978 doi: [10.1016/j.jqsrt.2024.108978](https://doi.org/10.1016/j.jqsrt.2024.108978).
- [442] M. T. I. Ibrahim, D. Alatoom, T. Furtenbacher, A. G. Csaszar, S. N. Yurchenko, A. A. A. Azzam, J. Tennyson, MARVEL analysis of high-resolution rovibrational spectra of ¹³C¹⁶O₂, *J. Comput. Chem.* 45 (2024) 969–984. doi: [10.1002/jcc.27266](https://doi.org/10.1002/jcc.27266).
- [443] D. Alatoom, M. T. I. Ibrahim, T. Furtenbacher, A. G. Csaszar, M. Alghizzawi, S. N. Yurchenko, A. A. A. Azzam, J. Tennyson, MARVEL analysis of high-resolution rovibrational spectra of ¹⁶O¹²C¹⁸O, *J. Comput. Chem.*
- [444] A. Yachmenev, A. Campargue, S. N. Yurchenko, J. Küpper, J. Tennyson, [Electric quadrupole transitions in carbon dioxide](#), *J. Chem. Phys.* 154 (2021) 211104. doi: [10.1063/5.0053279](https://doi.org/10.1063/5.0053279).
URL <https://doi.org/10.1063/5.0053279>
- [445] H. Fleurbaey, R. Grilli, D. Mondelain, S. Kassi, A. Yachmenev, S. N. Yurchenko, A. Campargue, Electric-quadrupole and magnetic-dipole contributions to the $\nu_2 + \nu_3$ band of carbon dioxide near 3.3 μm , *J. Quant. Spectrosc. Radiat. Transf.* 266 (2021) 107558. doi: [10.1016/j.jqsrt.2021.107558](https://doi.org/10.1016/j.jqsrt.2021.107558).
- [446] O. I. Korablev, D. A. Belyaev, Y. S. Dobrolenskiy, A. Y. Trokhimovskiy, Y. K. Kalinnikov, Acousto-optic tunable filter spectrometers in space missions (Invited), *Appl. Optics* 57 (10) (2018) C103–C119. doi: [10.1364/AO.57.00C103](https://doi.org/10.1364/AO.57.00C103).
- [447] Y. Wang, A. Owens, J. Tennyson, S. N. Yurchenko, MARVEL analysis of the measured high-resolution rovibronic spectra of the calcium monohydroxide radical (CaOH), *Astrophys. J. Suppl. Ser.* 248 (2020) 9. doi: [10.3847/1538-4365/ab85cb](https://doi.org/10.3847/1538-4365/ab85cb).
- [448] A. Owens, V. H. J. Clark, A. Mitrushchenkov, S. N. Yurchenko, J. Tennyson, Theoretical rovibronic spectroscopy of the calcium monohydroxide radical (CaOH), *J. Chem. Phys.* 154 (2021) 234302. doi: [10.1063/5.0052958](https://doi.org/10.1063/5.0052958).
- [449] A. O. Mitrushchenkov, A new general renner-teller (including $\epsilon \geq 1$) spectroscopic formalism for triatomic molecules, *J. Chem. Phys.* 136 (2012) 024108. doi: [10.1063/1.3672162](https://doi.org/10.1063/1.3672162).
- [450] I. N. Reid, J. D. Kirkpatrick, J. E. Gizis, C. C. Dahn, D. G. Monet, R. J. Williams, J. Liebert, A. J. Burgasser, Four nearby L dwarfs, *Astron. J.* 119 (2000) 369–377. doi: [10.1086/301177](https://doi.org/10.1086/301177).
- [451] E. Xu, J. Tennyson, Empirical rovibrational energy levels for carbonyl sulphide, *Mol. Phys.* (2023) e2279694 doi: [10.1080/00268976.2023.2279694](https://doi.org/10.1080/00268976.2023.2279694).
- [452] E. Esperza-Borges, M. López-Morales, J. I. Adams Redai, E. Pallé, J. Kirk, N. Casasayas-Barris, N. E. Batalha, B. V. Rackham, J. L. Bean, S. L. Casewell, L. Decin, L. A. Dos Santos, A. G. Muñoz, J. Harrington, K. Heng, R. Hu, L. Mancini, K. Molaverdikhani, G. Morello, N. K. Nikolov, M. C. Nixon, S. Redfield, K. B. Stevenson, H. R. Wakeford, M. K. Alam, B. Benneke, J. Blecic, N. Crouzet, T. Daylan, J. Inglis, L. Kreidberg, D. J. M. Petit dit de la Roche, J. D. Turner, [Detection of carbon monoxide in the atmosphere of WASP-39b applying standard cross-correlation techniques to jwst nirspec g395h data](#), *Astrophys. J. Lett.* 955 (1) (2023) L19. doi: [10.3847/2041-8213/acf27b](https://doi.org/10.3847/2041-8213/acf27b).
URL <http://dx.doi.org/10.3847/2041-8213/acf27b>
- [453] S.-M. Tsai, E. K. H. Lee, D. Powell, P. Gao, X. Zhang, J. Moses, E. Hébrard, O. Venot, V. Parmentier, S. Jordan, R. Hu, M. K. Alam, L. Alderson, N. M. Batalha, J. L. Bean, B. Benneke, C. J. Bierson, R. P. Brady, L. Carone, A. L. Carter, K. L. Chubb, J. Inglis, J. Leconte, M. Line, M. López-Morales, Y. Miguel, K. Molaverdikhani, Z. Rustamkulov, D. K. Sing, K. B. Stevenson, H. R. Wakeford, J. Yang, K. Aggarwal, R. Baeyens, S. Barat, M. de Val-Borro, T. Daylan, J. J. Fortney, K. France, J. M. Goyal, D. Grant, J. Kirk, L. Kreidberg, A. Louca, S. E. Moran, S. Mukherjee, E. Nasedkin, K. Ohno, B. V. Rackham, S. Redfield, J. Taylor, P. Tremblin, C. Visscher, N. L. Wallack, L. Welbanks, A. Youngblood, E.-M. Ahrer, N. E. Batalha, P. Behr, Z. K. Berta-Thompson, J. Blecic, S. L. Casewell, I. J. M. Crossfield, N. Crouzet, P. E. Cubillos, L. Decin, J.-M. Désert, A. D. Feinstein, N. P. Gibson, J. Harrington, K. Heng, T. Henning, E. M.-R. Kempton, J. Krick, P.-O. Lagage, M. Lendl, J. D. Lothringer, M. Mansfield, N. J. Mayne, T. Mikal-Evans, E. Palle, E. Schlawin, O. Shorttle, P. J. Wheatley, S. N. Yurchenko, [Photochemically produced SO₂ in the atmosphere of WASP-39b](#), *Nature* doi: [10.1038/s41586-023-05902-2](https://doi.org/10.1038/s41586-023-05902-2).
URL <https://doi.org/10.1038/s41586-023-05902-2>
- [454] J. L. Grenfell, A review of exoplanetary biosignatures, *Phys. Rep.* 713 (2017) 1–17. doi: [10.1016/j.physrep.2017.08.003](https://doi.org/10.1016/j.physrep.2017.08.003).
- [455] E. W. Schwieterman, S. L. Olson, D. Pidhorodetska, C. T. Reinhard, A. Ganti, T. J. Fauchez, S. T. Bastelberger, J. S. Crouse, A. Ridgwell, T. W. Lyons, Evaluating the Plausible Range of N₂O Biosignatures on Exo-Earths: An Integrated Biogeochemical, Photochemical, and Spectral Modeling Approach, *Astrophys. J.* 937 (2022) 109. doi: [10.3847/1538-4357/ac8cfb](https://doi.org/10.3847/1538-4357/ac8cfb).
- [456] D. Angerhausen, D. Pidhorodetska, M. Leung, J. Hansen, E. Alei, F. Dannert, J. Kammerer, S. P. Quanz, E. W. Schwieterman, L. Initiative, Large Interferometer For Exoplanets (LIFE). XII. The Detectability of Capstone Biosignatures in the Mid-infrared-Sniffing Exoplanetary

- Laughing Gas and Methylated Halogens, *Astron. J.* 167 (2024) 128. doi:[10.3847/1538-3881/ad1f4b](https://doi.org/10.3847/1538-3881/ad1f4b).
- [457] J. Tennyson, T. Furtenbacher, S. N. Yurchenko, A. G. Császár, Empirical rovibrational energy levels for nitrous oxide, *J. Quant. Spectrosc. Radiat. Transf.* 316 (2024) 108902. doi:[10.1016/j.jqsrt.2024.108902](https://doi.org/10.1016/j.jqsrt.2024.108902).
- [458] J. Tennyson, Empirical rovibronic energy levels of C₃, *Mol. Phys.* doi:[10.1080/00268976.2023.2276912](https://doi.org/10.1080/00268976.2023.2276912).
- [459] T. Furtenbacher, P. A. Coles, J. Tennyson, S. N. Yurchenko, S. Yu, B. Drouin, R. Tóbiás, A. G. Császár, Empirical rovibational energy of ammonia up to 7500 cm⁻¹, *J. Quant. Spectrosc. Radiat. Transf.* 251 (2020) 107027. doi:[10.1016/j.jqsrt.2020.107027](https://doi.org/10.1016/j.jqsrt.2020.107027).
- [460] I. Sadiek, A. J. Fleisher, J. Hayden, X. Huang, A. Hugi, R. Engeln, N. Lang, J.-P. H. van Helden, Dual-comb spectroscopy of ammonia formation in non-thermal plasmas, *Comms. Chem.* 7 (2024) 110. doi:[10.1038/s42004-024-01190-7](https://doi.org/10.1038/s42004-024-01190-7).
- [461] P. A. Coles, R. I. Ovsyannikov, O. L. Polyansky, S. N. Yurchenko, J. Tennyson, Improved potential energy surface and spectral assignments for ammonia in the near-infrared region, *J. Quant. Spectrosc. Radiat. Transf.* 219 (2018) 199–212. doi:[10.1016/j.jqsrt.2018.07.022](https://doi.org/10.1016/j.jqsrt.2018.07.022).
- [462] P. G. J. Irwin, N. Bowles, A. S. Braude, R. Garland, S. Calcutt, P. A. Coles, S. N. Yurchenko, J. Tennyson, Analysis of gaseous ammonia (NH₃) absorption in the visible spectrum of Jupiter - Update, *Icarus* 321 (2019) 572–582. doi:[10.1016/j.icarus.2018.12.008](https://doi.org/10.1016/j.icarus.2018.12.008).
- [463] S. N. Yurchenko, A theoretical room-temperature line list for ¹⁵NH₃, *J. Quant. Spectrosc. Radiat. Transf.* 152 (2015) 28–36. doi:[10.1016/j.jqsrt.2014.10.023](https://doi.org/10.1016/j.jqsrt.2014.10.023).
- [464] K. L. Chubb, M. Joseph, J. Franklin, N. Choudhury, T. Furtenbacher, A. G. Császár, G. Gaspard, P. Oguoko, A. Kelly, S. N. Yurchenko, J. Tennyson, C. Sousa-Silva, MARVEL analysis of the measured high-resolution spectra of C₂H₂, *J. Quant. Spectrosc. Radiat. Transf.* 204 (2018) 42–55. doi:[10.1016/j.jqsrt.2017.08.018](https://doi.org/10.1016/j.jqsrt.2017.08.018).
- [465] T. Mellor, A. Owens, S. N. Yurchenko, J. Tennyson, MARVEL analysis of high-resolution spectra of thioformaldehyde (H₂CS), *J. Mol. Spectrosc.* 391 (2023) 111732. doi:[10.1016/j.jms.2022.111732](https://doi.org/10.1016/j.jms.2022.111732).
- [466] S. N. Yurchenko, A. Owens, K. Kefala, J. Tennyson, ExoMol line lists – LVII. High accuracy ro-vibrational line list for methane (CH₄), *Mon. Not. Roy. Astron. Soc.* 528 (2024) 3719–3729. doi:[10.1093/mnras/stae148](https://doi.org/10.1093/mnras/stae148).
- [467] S. N. Yurchenko, J. Tennyson, ExoMol line lists IV: The rotation-vibration spectrum of methane up to 1500 K, *Mon. Not. Roy. Astron. Soc.* 440 (2014) 1649–1661. doi:[10.1093/mnras/stu326](https://doi.org/10.1093/mnras/stu326).
- [468] S. N. Yurchenko, D. S. Amundsen, J. Tennyson, I. P. Waldmann, A hybrid line list for CH₄ and hot methane continuum, *Astron. Astrophys.* 605 (2017) A95. doi:[10.1051/0004-6361/201731026](https://doi.org/10.1051/0004-6361/201731026).
- [469] M. Rey, A. V. Nikitin, V. G. Tyuterev, Theoretical hot methane line list up T=2000 K for astrophysical applications, *Astrophys. J.* 789 (2014) 2. doi:[10.1088/0004-637X/789/1/2](https://doi.org/10.1088/0004-637X/789/1/2).
- [470] M. Rey, A. V. Nikitin, V. G. Tyuterev, *Accurate Theoretical Methane Line Lists in the Infrared up to 3000 K and Quasi-continuum Absorption/Emission Modeling for Astrophysical Applications*, *Astrophys. J.* 847 (2017) 105. doi:[10.3847/1538-4357/aa8909](https://doi.org/10.3847/1538-4357/aa8909). URL [10.3847/1538-4357/aa8909](https://doi.org/10.3847/1538-4357/aa8909)
- [471] R. J. Hargreaves, I. E. Gordon, M. Rey, A. V. Nikitin, V. G. Tyuterev, R. V. Kochanov, L. S. Rothman, An accurate, extensive, and practical line list of methane for the HITEMP database, *Astrophys. J. Suppl. Ser.* 247 (2020) 55. doi:[10.3847/1538-4365/ab7a1a](https://doi.org/10.3847/1538-4365/ab7a1a).
- [472] K. Kefala, S. N. Yurchenko, J. Tennyson, Assigning spectra of methane, (work in progress) (2024).
- [473] R. S. Ram, J. S. A. Brooke, C. M. Western, P. F. Bernath, Einstein A-values and oscillator strengths of the A ²Π – X ²Σ⁺ system of CP, *J. Quant. Spectrosc. Radiat. Transf.* 138 (2014) 107–115. doi:[10.1016/j.jqsrt.2014.01.030](https://doi.org/10.1016/j.jqsrt.2014.01.030).
- [474] Z. Qin, T. Bai, L. Liu, Line lists for the X ²Σ⁺ – X ²Σ⁺, A ²Π – A ²Π and A2Π – X²Σ⁺ transitions of CP, *J. Quant. Spectrosc. Radiat. Transf.* 258 (2021) 107352. doi:[10.1016/j.jqsrt.2020.107352](https://doi.org/10.1016/j.jqsrt.2020.107352).
- [475] P. F. Bernath, R. Dodangodage, J. Liévin, S-type Stars: LaO Line List for the B ²Σ⁺ - X ²Σ⁺ Band System, *J. Mol. Spectrosc.* 933 (2022) 99. doi:[10.3847/1538-4357/ac731f](https://doi.org/10.3847/1538-4357/ac731f).
- [476] K. Bielska, A. A. Kyuberis, Z. D. Reed, G. Li, A. Cygan, R. Ciuryło, E. M. Adkins, L. Lodi, N. F. Zobov, V. Ebert, D. Lisak, J. T. Hodges, J. Tennyson, O. L. Polyansky, Subpromille Measurements and Calculations of CO (3–0) Overtone Line Intensities, *Phys. Rev. Lett.* 129 (2022) 043002. doi:[10.1103/PhysRevLett.129.043002](https://doi.org/10.1103/PhysRevLett.129.043002).
- [477] A. A. Balashov, G. L. K. Bielska, A. A. Kyuberis, S. Wójciewicz, J. Domyslawska, R. Ciuryło, J. T. N. F. Zobov and D. Lisak, O. L. Polyansky, Measurement and calculation of CO (7–0) overtone line intensities, *J. Chem. Phys.* 158 (2023) 234306. doi:[10.1063/5.0152996](https://doi.org/10.1063/5.0152996).
- [478] M. Khalil, S. Mahmoud, R. P. Brady, M. Almehairbi, M. Gacesa, S. N. Yurchenko, J. Tennyson, A. A. Ghaferi, N. El-Kork, Theoretical investigation of the A ¹Π – X ¹Σ⁺, B ¹Σ⁺ – X ¹Σ⁺, C ¹Σ⁺ – X ¹Σ⁺, D' ¹Σ⁺ – X ¹Σ⁺ and E' ¹Π – X ¹Σ⁺, *Phys. Chem. Chem. Phys.*
- [479] W. Somogyi, S. N. Yurchenko, A. Yachmenev, Calculation of electric quadrupole linestrengths for diatomic molecules: Application to the H₂, CO, HF, and O₂ molecules, *J. Chem. Phys.* 155 (21) (2021) 214303. doi:[10.1063/5.0063256](https://doi.org/10.1063/5.0063256).
- [480] W. Somogyi, S. N. Yurchenko, G.-S. Kim, An *ab initio* spectroscopic model of the molecular oxygen atmospheric and infrared bands, *Phys. Chem. Chem. Phys.*
- [481] E. Roueff, H. Abgrall, P. Czachorowski, K. Pachucki, M. Puchalski, J. Komasa, The full infrared spectrum of molecular hydrogen, *Astron. Astrophys.* 630 (2019) A58. doi:[10.1051/0004-6361/201936249](https://doi.org/10.1051/0004-6361/201936249).
- [482] H. Abgrall, E. Roueff, F. Launay, J.-Y. Roncin, The B ¹Σ_u⁺ → X ¹Σ_g⁺ and D ¹Π_u → X ¹Σ_g⁺ band systems of molecular hydrogen, *Can. J. Phys.* 72 (1994) 856–865. doi:[10.1139/p94-112](https://doi.org/10.1139/p94-112).
- [483] P. H. R. Amaral, L. G. Diniz, K. A. Jones, M. Stanke, A. Alijah, L. Adamowicz, J. R. Mohallem, Benchmark rovibrational linelists and einstein a-coefficients for the primordial molecules and isotopologues, *Astrophys. J.* 878 (2019) 95. doi:[10.3847/1538-4357/ab1f65](https://doi.org/10.3847/1538-4357/ab1f65).
- [484] C. M. Coppola, L. Lodi, J. Tennyson, Radiative cooling functions for primordial molecules, *Mon. Not. Roy. Astron. Soc.* 415 (2011) 487–493.
- [485] A. Campargue, A. M. Solodov, A. A. Solodov, A. Yachmenev, S. N. Yurchenko, Detection of electric-quadrupole transitions in water vapour near 5.4 and 2.5 μm, *Phys. Chem. Chem. Phys.* 22 (2020) 12476–12481. doi:[10.1039/DQCP01667E](https://doi.org/10.1039/DQCP01667E).
- [486] A.-M. Syme, L. K. McKemmish, Full spectroscopic model and trihybrid experimental-perturbative-variational line list for CN, *Mon. Not. Roy. Astron. Soc.* 505 (2021) 4383–4395. doi:[10.1093/mnras/stab1551](https://doi.org/10.1093/mnras/stab1551).
- [487] S. N. Yurchenko, L. Lodi, J. Tennyson, A. V. Stolyarov, Duo: A general program for calculating spectra of diatomic molecules, *Comput. Phys. Commun.* 202 (2016) 262 – 275. doi:[10.1016/j.cpc.2015.12.021](https://doi.org/10.1016/j.cpc.2015.12.021).
- [488] A.-M. Syme, L. K. McKemmish, Experimental energy levels of ¹²C¹⁴N through MARVEL analysis, *Mon. Not. Roy. Astron. Soc.* 499 (2020) 25–39. doi:[10.1093/mnras/staa2791](https://doi.org/10.1093/mnras/staa2791).
- [489] J. S. A. Brooke, R. S. Ram, C. M. Western, G. Li, D. W. Schwenke, P. F. Bernath, Einstein A coefficients and oscillator strengths for the A ²Π – X ²Σ⁺ (red) and B ²Σ⁺ – X ²Σ⁺ violet systems and rovibrational transitions in the X ²Σ⁺ state of CN, *Astrophys. J. Suppl. Ser.* 210

- (2014) 23. doi:[10.1088/0067-0049/210/2/23](https://doi.org/10.1088/0067-0049/210/2/23).
- [490] A. N. Perri, F. Taher, L. K. McKemmish, Full spectroscopic model and trihybrid experimental-perturbative-variational line list for ZrO, Mon. Not. Roy. Astron. Soc. 524 (2023) 4631–4641. doi:[10.1093/mnras/stad2103](https://doi.org/10.1093/mnras/stad2103).
- [491] L. K. McKemmish, J. Borsovszky, K. L. Goodhew, S. Sheppard, A. F. V. Bennett, A. D. J. Martin, A. Singh, C. A. J. Sturgeon, T. Furtenbacher, A. G. Császár, J. Tennyson, Marvel Analysis of the Measured High-resolution Rovibronic Spectra of $^{90}\text{Zr}^{16}\text{O}$, Astrophys. J. 867 (2018) 33. doi:[10.3847/1538-4357/aadd19](https://doi.org/10.3847/1538-4357/aadd19).
- [492] J. J. Sorensen, P. F. Bernath, Near-infrared and Visible Opacities of S-type Stars: The $\text{B } ^1\Pi - \text{X } ^1\Sigma^+$ Band System of ZrO, Astrophys. J. 923 (2021) 234. doi:[10.3847/1538-4357/ac300d](https://doi.org/10.3847/1538-4357/ac300d).
- [493] A. N. Perri, L. K. McKemmish, Full spectroscopic model and trihybrid experimental-perturbative-variational line list for NH, Mon. Not. Roy. Astron. Soc. 531 (3) (2024) 3023–3033. doi:[10.1093/mnras/stae1340](https://doi.org/10.1093/mnras/stae1340).
- [494] D. Darby-Lewis, H. Shah, D. Joshi, F. Khan, M. Kauwo, N. Sethi, P. F. Bernath, T. Furtenbacher, R. Tóbiás, A. G. Császár, J. Tennyson, MARVEL analysis of the measured high-resolution spectra of ^{14}NH , J. Mol. Spectrosc. 362 (2019) 69–76. doi:[10.1016/j.jms.2019.06.002](https://doi.org/10.1016/j.jms.2019.06.002).
- [495] J. S. A. Brooke, P. F. Bernath, C. M. Western, M. C. van Hemert, G. C. Groenenboom, Line strengths of rovibrational and rotational transitions within the $\text{X } ^3\Sigma^-$ ground state of NH, J. Chem. Phys. 141 (2014) 054310. doi:[10.1063/1.4891468](https://doi.org/10.1063/1.4891468).
- [496] J. S. A. Brooke, P. F. Bernath, C. M. Western, Note: Improved line strengths of rovibrational and rotational transitions within the $\text{X } ^3\Sigma^-$ ground state of NH, J. Chem. Phys. 143 (2015) 026101. doi:[10.1063/1.4923422](https://doi.org/10.1063/1.4923422).
- [497] A. M. Fernando, P. F. Bernath, J. N. Hodges, T. Masseron, A new linelist for the $\text{A } ^3\Pi - \text{X } ^3\Sigma^-$ transition of the NH free radical, J. Quant. Spectrosc. Radiat. Transf. 217 (2018) 29–34. doi:[10.1016/j.jqsrt.2018.05.021](https://doi.org/10.1016/j.jqsrt.2018.05.021).
- [498] X. Li, Z. Qin, L. Liu, High-temperature molecular line list of hydroboron monoxide (hbo), Phys. Chem. Chem. Phys. 26 (2024) 12838–12843. doi:[10.1039/d3cp05997a](https://doi.org/10.1039/d3cp05997a).
- [499] C. M. Western, L. Carter-Blatchford, P. Crozet, A. J. Ross, J. Morville, D. W. Tokaryk, The spectrum of N₂ from 4,500 to 15,700 cm⁻¹ revisited with PGOPHER, J. Quant. Spectrosc. Radiat. Transf. 219 (2018) 127–141. doi:[10.1016/j.jqsrt.2018.07.017](https://doi.org/10.1016/j.jqsrt.2018.07.017).
- [500] E. R. Jans, Rovibronic molecular line list for the N₂(C $^3\Pi_u \rightarrow \text{B } ^3\Pi_g$) second positive system, J. Quant. Spectrosc. Radiat. Transf. 312 (2024) 108809. doi:[10.1016/j.jqsrt.2023.108809](https://doi.org/10.1016/j.jqsrt.2023.108809).
- [501] R. Vallon, C. Richard, P. Crozet, G. Wannous, A. Ross, Laboratory measurements of NiH by Fourier transform dispersed fluorescence, Astrophys. J. 696 (2009) 172–175. doi:[10.1088/0004-637X/696/1/172](https://doi.org/10.1088/0004-637X/696/1/172).
- [502] H. Harker, C. Richard, G. Tourasse, P. Crozet, A. J. Ross, Zeeman spectroscopy of NiH: Lande factors of three $\Omega = \frac{3}{2}$ excited electronic states, J. Mol. Spectrosc. 292 (2013) 28–34. doi:[10.1016/j.jms.2013.09.005](https://doi.org/10.1016/j.jms.2013.09.005).
- [503] I. Havalyova, I. Bozhinova, A. Pashov, A. J. Ross, P. Crozet, A coupled-channels model describing the low-lying $^2\Delta$, $^2\Sigma^+$ and $^2\Pi$ electronic states of nickel monohydride with experimental accuracy, J. Quant. Spectrosc. Radiat. Transf. 272 (2021) 107800. doi:[10.1016/j.jqsrt.2021.107800](https://doi.org/10.1016/j.jqsrt.2021.107800).
- [504] N. F. Allard, F. Spiegelman, J. F. Kielkopf, K-H₂ line shapes for the spectra of cool brown dwarfs, Astron. Astrophys. 589 (2016) A21. doi:[10.1051/0004-6361/201628270](https://doi.org/10.1051/0004-6361/201628270).
- [505] N. F. Allard, F. Spiegelman, T. Leininger, P. Molliere, New study of the line profiles of sodium perturbed by H₂, Astron. Astrophys. 628 (2019) A120. doi:[10.1051/0004-6361/201935593](https://doi.org/10.1051/0004-6361/201935593).
- [506] N. F. Allard, J. F. Kielkopf, K. Myneni, J. N. Blakely, New theoretical study of potassium perturbed by He and a comparison to laboratory spectra, Astron. Astrophys. 683 (2024) A188. doi:[10.1051/0004-6361/202348711](https://doi.org/10.1051/0004-6361/202348711).
- [507] R. R. Gamache, C. Roller, E. Lopes, I. E. Gordon, L. S. Rothman, O. L. Polyansky, N. F. Zobov, A. A. Kyuberis, J. Tennyson, S. N. Yurchenko, A. G. Császár, T. Furtenbacher, X. Huang, D. W. Schwenke, T. J. Lee, B. J. Drouin, S. A. Tashkun, V. I. Perevalov, R. V. Kochanov, Total Internal Partition Sums for 167 isotopologues of 53 molecules important in planetary atmospheres: application to HITRAN2016 and beyond, J. Quant. Spectrosc. Radiat. Transf. 203 (2017) 70–87. doi:[10.1016/j.jqsrt.2017.03.045](https://doi.org/10.1016/j.jqsrt.2017.03.045).
- [508] A. W. Irwin, Polynomial partition-function approximations of 344 atomic species and molecular-species, Astrophys. J. Suppl. Ser. 45 (4) (1981) 621–633.
- [509] A. J. Sauval, J. B. Tatum, A set of partition functions and equilibrium constants for 300 diatomic molecules of astrophysical interest, Astrophys. J. Suppl. Ser. 56 (1984) 193–209. doi:[10.1086/190980](https://doi.org/10.1086/190980).
- [510] P. S. Barklem, R. Collet, Partition functions and equilibrium constants for diatomic molecules and atoms of astrophysical interest, Astron. Astrophys. 588 (2016) A96. doi:[10.1051/0004-6361/201526961](https://doi.org/10.1051/0004-6361/201526961).
- [511] C. Sousa-Silva, N. Hesketh, S. N. Yurchenko, C. Hill, J. Tennyson, High Temperature partition functions and thermodynamic data for ammonia and phosphine, J. Quant. Spectrosc. Radiat. Transf. 142 (2014) 66–74. doi:[10.1016/j.jqsrt2014.03.012](https://doi.org/10.1016/j.jqsrt2014.03.012).
- [512] J. Tennyson, C. Hill, S. N. Yurchenko, Data structures for ExoMol: Molecular line lists for exoplanet and other atmospheres, in: 6th international conference on atomic and molecular data and their applications ICAMDATA-2012, Vol. 1545 of AIP Conference Proceedings, AIP, New York, 2013, pp. 186–195. doi:[10.1063/1.4815853](https://doi.org/10.1063/1.4815853).
- [513] J. Tennyson, K. Hulme, O. K. Naim, S. N. Yurchenko, Radiative lifetimes and cooling functions for astrophysically important molecules, J. Phys. B-At. Mol. Opt. Phys. 49 (2016) 044002. doi:[10.1088/0953-4075/49/4/044002](https://doi.org/10.1088/0953-4075/49/4/044002).
- [514] C. Hill, M. Hanicinec, A Python package for managing simple chemical species and states, <https://github.com/xnx/pyvalem>, version 2.5.12 (2022).
- [515] C. Hill, , Dipti, M. Hanicinec, et al., PyValem: a machine-readable notation and Python library for atomic and molecular structures and states Manuscript in preparation.
- [516] F. Pezoa, J. L. Reutter, F. Suarez, M. Ugarte, D. Vrgoč, Foundations of json schema, in: Proceedings of the 25th International Conference on World Wide Web, International World Wide Web Conferences Steering Committee, 2016, pp. 263–273.
- [517] T. Furtenbacher, T. Szidarovszky, J. Hruba, A. A. Kyuberis, N. F. Zobov, O. L. Polyansky, J. Tennyson, A. G. Császár, Definitive high-temperature ideal-gas thermochemical functions of the H₂¹⁶O molecule, J. Phys. Chem. Ref. Data 45 (2016) 043104. doi:[10.1063/1.4967723](https://doi.org/10.1063/1.4967723).
- [518] S. Miller, T. Stallard, H. Melin, J. Tennyson, H₃⁺ cooling in planetary atmospheres, Faraday Discuss. 147 (2010) 283–291. doi:[10.1039/C004152C](https://doi.org/10.1039/C004152C).
- [519] S. Miller, T. Stallard, J. Tennyson, H. Melin, Cooling by H₃⁺ emission, J. Phys. Chem. A 117 (2013) 9770–9777.
- [520] D. C. Wells, E. W. Greisen, R. H. Harten, FITS - a Flexible Image Transport System, Astron. Astrophys. Suppl. 44 (1981) 363.

- [521] P. Molière, J. P. Wardenier, R. van Boekel, T. Henning, K. Molaverdikhani, I. A. G. Snellen, petitRADTRANS - A Python radiative transfer package for exoplanet characterization and retrieval, *Astron. Astrophys.* 627 (2019) A67. doi:[10.1051/0004-6361/201935470](https://doi.org/10.1051/0004-6361/201935470).
- [522] P. Molière, E. Nasedkin, D. Blain, *petitRADTRANS* (2023). doi:[10.5281/zenodo.8337871](https://doi.org/10.5281/zenodo.8337871).
URL <https://doi.org/10.5281/zenodo.8337871>
- [523] P. G. J. Irwin, N. A. Teanby, R. de Kok, L. N. Fletcher, C. J. A. Howett, C. C. C. Tsang, C. F. Wilson, S. B. Calcutt, C. A. Nixon, P. D. Parrish, The NEMESIS planetary atmosphere radiative transfer and retrieval tool, *J. Quant. Spectrosc. Radiat. Transf.* 109 (2008) 1136 – 1150. doi:[10.1016/j.jqsrt.2007.11.006](https://doi.org/10.1016/j.jqsrt.2007.11.006).
- [524] J. J. Fortney, T. D. Robinson, S. Domagal-Goldman, A. D. Del Genio, I. E. Gordon, E. ib Nezhad, N. Lewis, C. Sousa-Silva, V. Airapetian, B. Drouin, R. J. Hargreaves, X. Huang, T. Karman, R. M. Ramirez, G. B. Rieker, J. Tennyson, R. Wordsworth, S. N. Yurchenko, A. V. Johnson, T. J. Lee, C. Dong, S. Kane, M. Lopez-Morales, T. Fauchez, T. Lee, M. S. Marley, K. Sung, N. Haghhighipour, T. Robinson, S. Horst, P. Gao, D. you Kao, C. Dressing, R. Lupu, D. W. Savin, B. Fleury, O. Venot, D. Ascenzi, S. Milam, H. Linnartz, M. Gudipati, G. Gronoff, F. Salama, L. Gavilan, J. Bouwman, M. Turbet, Y. Benilan, B. Henderson, N. Batalha, R. Jensen-Clem, T. Lyons, R. Freedman, E. Schwertner, J. Goyal, L. Mancini, P. Irwin, J.-M. Desert, K. Molaverdikhani, J. Gizis, J. Taylor, J. Lothringer, R. Pierrehumbert, R. Zellem, N. Batalha, S. Rugheimer, J. Lustig-Yaeger, R. Hu, E. Kempton, G. Arney, M. Line, M. Alam, J. Moses, N. Iro, L. Kreidberg, J. Bleicic, T. Louden, P. Moliere, K. Stevenson, M. Swain, K. Bott, N. Madhusudhan, J. Krissansen-Totton, D. Deming, I. Kitashvili, E. Shkolnik, Z. Rustamkulov, L. Rogers, L. Close, The need for laboratory measurements and ab initio studies to aid understanding of exoplanetary atmospheres, arXiv:1905.07064 (2019).
- [525] E. J. Barton, C. Hill, M. Czurylo, H.-Y. Li, A. Hyslop, S. N. Yurchenko, J. Tennyson, The ExoMol diet: H₂ and He line-broadening parameters, *J. Quant. Spectrosc. Radiat. Transf.* 203 (2017) 490–495. doi:[10.1016/j.jqsrt.2017.01.028](https://doi.org/10.1016/j.jqsrt.2017.01.028).
- [526] Q. Ma, R. Tipping, C. Boulet, **Modification of the Robert-Bonamy formalism in calculating Lorentzian half-widths and shifts**, *Journal of Quantitative Spectroscopy and Radiative Transfer* 103 (3) (2007) 588–596. doi:[10.1016/j.jqsrt.2006.08.001](https://doi.org/10.1016/j.jqsrt.2006.08.001).
URL <https://linkinghub.elsevier.com/retrieve/pii/S0022407306001968>
- [527] A. Sokolov, S. N. Yurchenko, J. Tennyson, R. R. Gamache, B. Vispoel, Calculation of collisional line-broadening and shifting of acetylene using the Modified Complex Robert-Bonamy approach , *J. Quant. Spectrosc. Radiat. Transf.*(in preparation).
- [528] E. R. Guest, J. Tennyson, S. N. Yurchenko, Modelling the Rotational Dependence of Line Broadening using Machine Learning, *J. Mol. Spectrosc.* 401 (2024) 111901. doi:[10.1016/j.jms.2024.111901](https://doi.org/10.1016/j.jms.2024.111901).
- [529] E. Guest, **Predicting the Rotational Dependence of Line Broadening using Machine Learning** (2024). doi:[10.5281/zenodo.10631728](https://doi.org/10.5281/zenodo.10631728).
URL <https://doi.org/10.5281/zenodo.10631728>
- [530] J. Buldyreva, S. N. Yurchenko, J. Tennyson, Simple semi-classical model of pressure-broadened infrared/microwave linewidths in the temperature range 200–3000 K, *RAS Tech. Instr. 1* (2022) 43–47. doi:[10.1093/rasti/rzac004](https://doi.org/10.1093/rasti/rzac004).
- [531] J. Buldyreva, K. Stehlin, S. N. Yurchenko, E. R. Guest, J. Tennyson, Semi-classical estimates of pressure-induced linewidths for infrared absorption by hot (exo)planetary atmospheres, *Icarus*.
- [532] J. Buldyreva, R. P. Brady, S. N. Yurchenko, J. Tennyson, Collisional broadening of molecular rovibronic lines, *J. Quant. Spectrosc. Radiat. Transf.* 313 (2024) 108843. doi:[10.1016/j.jqsrt.2023.108843](https://doi.org/10.1016/j.jqsrt.2023.108843).
- [533] A. U. Hazi, H. S. Taylor, Stabilization method of calculating resonance energies: Model problem, *Phys. Rev. A* 1 (1970) 1109–1120. doi:[10.1103/PhysRevA.1.1109](https://doi.org/10.1103/PhysRevA.1.1109).
- [534] Z. Bacic, J. Simons, Resonance energies and lifetimes from stabilization-based methods, *J. Phys. Chem.* 86 (7) (1982) 1192–1200. doi:[10.1021/j100396a027](https://doi.org/10.1021/j100396a027).
- [535] V. A. Mandelshtam, T. R. Ravuri, H. S. Taylor, Calculation of the density of resonance states using the stabilization method, *Phys. Rev. Lett.* 70 (13) (1993) 1932–1935. doi:[10.1103/PhysRevLett.70.1932](https://doi.org/10.1103/PhysRevLett.70.1932).
- [536] M. Pezzella, S. N. Yurchenko, J. Tennyson, ExoMol photodissociation cross sections I – HCl and HF, *Mon. Not. Roy. Astron. Soc.* 514 (2022) 4413–4425. doi:[10.1093/mnras/stac1634](https://doi.org/10.1093/mnras/stac1634).
- [537] O. Venot, Y. Bénilan, N. Fray, M.-C. Gazeau, F. Lefèvre, E. Es-sebbar, E. Hébrard, M. Schwell, C. Bahrini, F. Montmessin, M. Lefèvre, I. P. Waldmann, VUV-absorption cross section of carbon dioxide from 150 to 800 K and applications to warm exoplanetary atmospheres, *Astron. Astrophys.* 609 (2018) A34. doi:[10.1051/0004-6361/201731295](https://doi.org/10.1051/0004-6361/201731295).
- [538] L. S. Rothman, I. E. Gordon, Y. Babikov, A. Barbe, D. C. Benner, P. F. Bernath, M. Birk, L. Bizzocchi, V. Boudon, L. R. Brown, A. Campargue, K. Chance, E. A. Cohen, L. H. Coudert, V. M. Devi, B. J. Drouin, A. Fayt, J.-M. Flaud, R. R. Gamache, J. J. Harrison, J.-M. Hartmann, C. Hill, J. T. Hodges, D. Jacquemart, A. Jolly, J. Lamouroux, R. J. Le Roy, G. Li, D. A. Long, O. M. Lyulin, C. J. Mackie, S. T. Massie, S. Mikhailenko, H. S. P. Müller, O. V. Naumenko, A. V. Nikitin, J. Orphal, V. Perevalov, A. Perrin, E. R. Polovtseva, C. Richard, M. A. H. Smith, E. Starikova, K. Sung, S. Tashkun, J. Tennyson, G. C. Toon, V. G. Tyuterev, G. Wagner, The HITRAN 2012 molecular spectroscopic database, *J. Quant. Spectrosc. Radiat. Transf.* 130 (2013) 4 – 50. doi:[10.1016/j.jqsrt.2013.07.002](https://doi.org/10.1016/j.jqsrt.2013.07.002).
- [539] D. Jack, P. H. Hauschildt, E. Baron, Time-dependent radiative transfer with PHOENIX, *Astron. Astrophys.* 502 (2009) 1043–1049.
- [540] J. Tennyson, S. Mohr, M. Hanicinec, A. Dzarasova, C. Smith, S. Waddington, B. Liu, L. L. Alves, K. Bartschat, A. Bogaerts, S. U. Engelmann, T. Gans, A. R. Gibson, S. Hamaguchi, K. R. Hamilton, C. Hill, D. O’Connell, S. Rauf, K. van ’t Veer, O. Zatsarinsky, The 2021 release of the Quantemol database (QDB) of plasma chemistries and reactions, *Plasma Sources Sci. Technol.* 31 (2022) 095020. doi:[10.1088/1361-6595/ac907e](https://doi.org/10.1088/1361-6595/ac907e).
- [541] A. Owens, A. Mitrushchenkov, S. N. Yurchenko, J. Tennyson, ExoMol line lists – XLVII. Rovibronic molecular line list of the calcium monohydroxide radical (CaOH), *Mon. Not. Roy. Astron. Soc.* 516 (2022) 3995–4002. doi:[10.1093/mnras/stac2462](https://doi.org/10.1093/mnras/stac2462).
- [542] M. Pezzella, G. B. Mitev, S. N. Yurchenko, J. Tennyson, A. O. Mitrushchenkov, A variational method for studying the photodissociation of triatomic molecules, *Phys. Chem. Chem. Phys.*
- [543] P. F. Weck, A. Schweitzer, P. C. Stancil, P. H. Hauschildt, K. Kirby, The molecular continuum opacity of ²⁴MgH in cool stellar atmospheres, *ApJ* 584 (2003) 459–464. doi:[10.1086/345524](https://doi.org/10.1086/345524).
- [544] S. Miyake, C. D. Gay, P. C. Stancil, Rovibrationally Resolved Photodissociation Of HeH⁺, *Astrophys. J.* 735 (2011) 21. doi:[10.1088/0004-637X/735/1/21](https://doi.org/10.1088/0004-637X/735/1/21).
- [545] W. H. el Qadi, P. C. Stancil, Photodissociation Of CN, *Astrophys. J.* 779 (2013) 97. doi:[10.1088/0004-637X/779/2/97](https://doi.org/10.1088/0004-637X/779/2/97).
- [546] E. C. McMillan, G. Shen, J. F. McCann, B. M. McLaughlin, P. C. Stancil, Rovibrationally resolved photodissociation of SH⁺, *J. Phys. B-At. Mol. Opt. Phys.* 49 (2016) 084001. doi:[10.1088/0953-4075/49/8/084001](https://doi.org/10.1088/0953-4075/49/8/084001).
- [547] R. J. Pattillo, R. Cieszewski, P. C. Stancil, R. C. Forrey, J. F. Babb, J. F. McCann, B. M. McLaughlin, Photodissociation of CS from Excited

- Rovibrational Levels, *Astrophys. J.* 858 (2018) 10. doi:[10.3847/1538-4357/aab5b9](https://doi.org/10.3847/1538-4357/aab5b9).
- [548] T. Bai, Z. Qin, L. Liu, Rovibrationally resolved direct photodissociation of MgO, *Mon. Not. Roy. Astron. Soc.* 505 (2021) 2177–2185. doi:[10.1093/mnras/stab1426](https://doi.org/10.1093/mnras/stab1426).
- [549] Z. Qin, T. Bai, L. Liu, Temperature-dependent direct photodissociation cross sections and rates of AlCl, *MNRAS* 508 (2021) 2848–2854. doi:[10.1093/mnras/stab2655](https://doi.org/10.1093/mnras/stab2655).
- [550] Z. Qin, T. Bai, L. Liu, Destruction of AlF: a quantum study of its ground -state photodissociation, *Mon. Not. Roy. Astron. Soc.* 510 (2022) 3011–3018. doi:[10.1093/mnras/stab3598](https://doi.org/10.1093/mnras/stab3598).
- [551] T. Bai, X. Yang, Z. Qin, L. Liu, Photodissociation cross sections and rates of NaO, *Mon. Not. Roy. Astron. Soc.* 527 (2023) 3847–3857. doi:[10.1093/mnras/stad3447](https://doi.org/10.1093/mnras/stad3447).
- [552] Z. Qin, P. Hu, T. Bai, L. Liu, A Theoretical Study of Temperature-dependent Photodissociation Cross Sections and Rates for O₂, *Astrophys. J. Suppl. Ser.* 269 (2023) 48. doi:[10.3847/1538-4365/ad03ed](https://doi.org/10.3847/1538-4365/ad03ed).
- [553] J. J. Spake, D. K. Sing, T. M. Evans, A. Oklopčić, V. Bourrier, L. Kreidberg, B. V. Rackham, J. Irwin, D. Ehrenreich, A. Wyttenbach, H. R. Wakeford, Y. Zhou, K. L. Chubb, N. Nikolov, J. M. Goyal, G. W. Henry, M. H. Williamson, S. Blumenthal, D. R. Anderson, C. Hellier, D. Charbonneau, S. Udry, N. Madhusudhan, Helium in the eroding atmosphere of an exoplanet, *Nature* 557 (2018) 68–70. doi:[10.1038/s41586-018-0067-5](https://doi.org/10.1038/s41586-018-0067-5).
- [554] Hoeijmakers, H. J., Seidel, J. V., Pino, L., Kitzmann, D., Sindel, J. P., Ehrenreich, D., Oza, A. V., Bourrier, V., Allart, R., Gebek, A., Lovis, C., Yurchenko, S. N., Astudillo-Defru, N., Bayliss, D., Cegla, H., Lavie, B., Lendl, M., Melo, C., Murgas, F., Nascimbeni, V., Pepe, F., Ségransan, D., Udry, S., Wyttenbach, A., Heng, K., Hot Exoplanet Atmospheres Resolved with Transit Spectroscopy (HEARTS) - IV. A spectral inventory of atoms and molecules in the high-resolution transmission spectrum of WASP-121 b, *Astron. Astrophys.* 641 (2020) A123. doi:[10.1051/0004-6361/202038365](https://doi.org/10.1051/0004-6361/202038365).
- [555] D. Cont, F. Yan, A. Reiners, N. Casasayas-Barris, P. Mollière, E. Pallé, T. Henning, L. Nortmann, M. Stangret, S. Czesla, M. López-Puertas, A. Sánchez-López, F. Rodler, I. Ribas, A. Quirrenbach, J. A. Caballero, P. J. Amado, L. Carone, J. Khaimova, L. Kreidberg, K. Molaverdikhani, D. Montes, G. Morello, E. Nagel, M. Oshagh, M. Zechmeister, Detection of Fe and evidence for TiO in the dayside emission spectrum of WASP-33b, *Astron. Astrophys.* 651 (2021) A33. doi:[10.1051/0004-6361/202140732](https://doi.org/10.1051/0004-6361/202140732).
- [556] A. Kramida, Yu. Ralchenko, J. Reader, and NIST ASD Team, NIST Atomic Spectra Database (ver. 5.6.1), [Online]. Available: <https://physics.nist.gov/asd> [2019, April 13]. National Institute of Standards and Technology, Gaithersburg, MD. (2019).
- [557] A. Kramida, Y. Ralchenko, J. Reader, and NIST ASD Team, Development of nist atomic databases and online tools, *Atoms* 8 (2020) 56. doi:[10.3390/atoms8030056](https://doi.org/10.3390/atoms8030056).
- [558] S. O. M. Wright, S. K. Nugroho, N. P. Gibson, E. J. W. de Mooij, I. Waldmann, J. Tennyson, H. Kawahara, M. Kuzuhara, T. Hirano, T. Kotani, Y. Kawashima, C. A. Watson, M. Tamura, K. Zwintz, H. Harakawa, K. Hodapp, S. Jacobson, M. Konishi, T. Kurokawa, J. Nishikawa, M. Omiya, T. Serizawa, T. Serizawa, A. Ueda, S. Vievard, S. N. Yurchenko, A spectroscopic thermometer: individual vibrational band spectroscopy with the example of OH in the atmosphere of WASP-33b, *Astron. J.* 166 (2023) 41. doi:[10.3847/1538-3881/acdb75](https://doi.org/10.3847/1538-3881/acdb75).
- [559] C. Sousa-Silva, J. J. Petkowski, S. Seager, Molecular simulations for the spectroscopic detection of atmospheric gases, *Phys. Chem. Chem. Phys.* 21 (2019) 18970–18987. doi:[10.1039/C8CP07057A](https://doi.org/10.1039/C8CP07057A).
- [560] G. Tinetti, P. Drossart, P. Eccleston, P. Hartogh, A. Heske, J. Leconte, G. Micela, M. Ollivier, G. Pilbratt, L. Puig, D. Turrini, B. Vandenbussche, P. Wolkenberg, J.-P. Beaulieu, L. A. Buchave, M. Ferus, M. Griffin, M. Guedel, K. Justtanont, P.-O. Lagage, P. Machado, G. Malaguti, M. Min, H. U. Nørgaard-Nielsen, M. Rataj, T. Ray, I. Ribas, M. Swain, R. Szabo, S. Werner, J. Barstow, M. Burleigh, J. Cho, V. C. du Foresto, A. Coustenis, L. Decin, T. Encrenaz, M. Galand, M. Gillon, R. Helled, J. C. Morales, A. G. Muñoz, A. Moneti, I. Pagano, E. Pascale, G. Piccioni, D. Pinfield, S. Sarkar, F. Selsis, J. Tennyson, A. Triaud, O. Venot, I. Waldmann, D. Walther, G. Wright, J. Amiaux, J.-L. Auguères, M. Berthé, N. Bezáwada, G. Bishop, N. Bowles, D. Coffey, J. Colomé, M. Crook, P.-E. Crouzet, V. Da Peppo, I. E. Sanz, M. Focardi, M. Frericks, T. Hunt, R. Kohley, K. Middleton, G. Morgante, R. Ottensamer, E. Pace, C. Pearson, R. Stamper, K. Symonds, M. Rengel, E. Renotte, P. Ade, L. Affer, C. Alard, N. Allard, F. Altieri, Y. André, C. Arena, I. Argyriou, A. Aylward, C. Baccani, G. Bakos, M. Banaszkiewicz, M. Barlow, V. Batista, G. Bellucci, S. Benatti, P. Bernardi, B. Bézard, M. Blecka, E. Bolmont, B. Bonfond, R. Bonito, A. S. Bonomo, J. R. Brucato, A. S. Brun, I. Bryson, W. Bujwan, S. Casewell, B. Charnay, C. C. Pestellini, G. Chen, A. Ciaravella, R. Claudi, R. Clédassou, M. Damasso, M. Damiano, C. Danielski, P. Deroo, A. M. Di Giorgio, C. Dominik, V. Doublier, S. Doyle, R. Doyon, B. Drummond, B. Duong, S. Eales, B. Edwards, M. Farina, E. Flaccomio, L. Fletcher, F. Forget, S. Fossey, M. Fränz, Y. Fujii, Á. García-Piquer, W. Gear, H. Geoffray, J. C. Gérard, L. Gesa, H. Gomez, R. Graczyk, C. Griffith, D. Grodent, M. G. Guarcello, J. Gustin, K. Hamano, P. Hargrave, Y. Hello, K. Heng, E. Herreiro, A. Hornstrup, B. Hubert, S. Ida, M. Ikoma, N. Iro, P. Irwin, C. Jarchow, J. Jaubert, H. Jones, Q. Julien, S. Kameda, F. Kerschbaum, P. Kervella, T. Koskinen, M. Krijger, N. Krupp, M. Lafarga, F. Landini, E. Lellouch, G. Leto, A. Luntzer, T. Rank-Lüftinger, A. Maggio, J. Maldonado, J.-P. Maillard, U. Mall, J.-B. Marquette, S. Mathis, P. Maxted, T. Matsuo, A. Medvedev, Y. Miguel, V. Minier, G. Morello, A. Mura, N. Narita, V. Nascimbeni, N. Nguyen Tong, V. Noce, F. Oliva, E. Palle, P. Palmer, M. Pancrazzi, A. Papageorgiou, V. Parmentier, M. Perger, A. Petralia, S. Pezzuto, R. Pierrehumbert, I. Pillitteri, G. Piotto, G. Pisano, L. Prisinzano, A. Radioti, J.-M. Réess, L. Rezac, M. Rocchetto, A. Rosich, N. Sanna, A. Santerne, G. Savini, G. Scandariato, B. Sicardy, C. Sierra, G. Sindoni, K. Skup, I. Snellen, M. Sobiecki, L. Soret, A. Sozzetti, A. Stiepen, A. Strugarek, J. Taylor, W. Taylor, L. Terenzi, M. Tessenyi, A. Tsiaras, C. Tucker, D. Valencia, G. Vasisht, A. Vazan, F. Vilardell, S. Vinatier, S. Viti, R. Waters, P. Wawer, A. Wawrzaszek, A. Whitworth, Y. L. Yung, S. N. Yurchenko, M. R. Z. Osorio, R. Zelle, T. Zingales, F. Zwart, A chemical survey of exoplanets with ARIEL, *Exp. Astron.* 46 (2018) 135–209. doi:[10.1007/s10686-018-9598-x](https://doi.org/10.1007/s10686-018-9598-x).
- [561] E. Pascale, N. Bezáwada, J. Barstow, J.-P. Beaulieu, N. Bowles, V. C. du Foresto, A. Coustenis, L. Decin, P. Drossart, P. Eccleston, T. Encrenaz, F. Forget, M. Griffin, M. Güdel, P. Hartogh, A. Heske, P.-O. Lagage, J. Leconte, P. Malaguti, G. Micela, K. Middleton, M. Min, A. Moneti, J. C. Morales, L. Mugnai, M. Ollivier, E. Pace, A. Papageorgiou, G. Pilbratt, L. Puig, M. Rataj, T. Ray, I. Ribas, M. Rocchetto, S. Sarkar, F. Selsis, W. Taylor, J. Tennyson, G. Tinetti, D. Turrini, B. Vandenbussche, O. Venot, I. Waldmann, P. Wolkenberg, G. Wright, M.-R. Z. Osorio, T. Zingales, The ARIEL space mission, in: M. Lystrup, H. A. MacEwen, G. G. Fazio, N. Batalha, N. Siegler, E. C. Tong (Eds.), *Space Telescopes and Instrumentation 2018: Optical, Infrared, and Millimeter Wave*, Vol. 10698, International Society for Optics and Photonics, SPIE, 2018, p. 106980H. doi:[10.1117/12.2311838](https://doi.org/10.1117/12.2311838).

Appendix A. Sample JSON files

Listing 1: JSON format master file exomol.json.

```
{  
    "ID": "EXOMOL.master",  
    "version": 20240603,  
    "num_molecules": 91,  
    "num_isotopologues": 224,  
    "num_datasets": 106,  
    "num_species": 269,  
    "molecules": {  
        "H2O": {  
            "num_molecule_names": 11,  
            "molecule_names": [  
                "Water",  
                "Oxidane",  
                "Hydrogen oxide",  
                "Dihydrogen monoxide",  
                "Hydrogen monoxide",  
                "Dihydrogen oxide",  
                "Hydrogen hydroxide",  
                "Hydric acid",  
                "Hydrohydroxic acid",  
                "Hydroxic acid",  
                "Hydrol"  
            ],  
            "num_isotopologues": 7,  
            "linelist": [  
                {  
                    "inchikey": "XLYOFNOQVPJJNP-UHFFFAOYSA-N",  
                    "iso_slug": "1H2-160",  
                    "iso_formula": "(1H)2(160)",  
                    "dataset_name": "POKAZATEL",  
                    "version": 20180501  
                },  
                {  
                    "inchikey": "XLYOFNOQVPJJNP-OUBTZVSYSA-N",  
                    "iso_slug": "1H2-170",  
                    "iso_formula": "(1H)2(170)",  
                    "dataset_name": "HotWat78",  
                    "version": 20161222  
                },  
                {  
                    "inchikey": "XLYOFNOQVPJJNP-NJFSPNSNSA-N",  
                    "iso_slug": "1H2-180",  
                    "iso_formula": "(1H)2(180)",  
                    "dataset_name": "HotWat78",  
                    "version": 20161222  
                },  
                {  
                    "inchikey": "XLYOFNOQVPJJNP-DYCDLGHISA-N",  
                    "iso_slug": "1H-2H-160",  
                    "iso_formula": "(1H)(2H)(160)",  
                    "dataset_name": "VTT",  
                    "version": 20160726  
                },  
                {  
                    "inchikey": "XLYOFNOQVPJJNP-DYCDLGHISA-N",  
                    "iso_slug": "1H-2H-160",  
                    "iso_formula": "(1H)(2H)(160)",  
                    "dataset_name": "Hewitt",  
                    "version": 20161222  
                },  
                {  
                    "inchikey": "XLYOFNOQVPJJNP-DYCDLGHISA-N",  
                    "iso_slug": "1H-2H-160",  
                    "iso_formula": "(1H)(2H)(160)",  
                    "dataset_name": "TDB",  
                    "version": 20240603  
                },  
                {  
                    "inchikey": "XLYOFNOQVPJJNP-ZSJDYOACSA-N",  
                    "iso_slug": "2H2-160",  
                    "iso_formula": "(2H)2(160)",  
                    "dataset_name": "Hewitt",  
                    "version": 20161222  
                }  
            ]  
        }  
    }  
}
```

Listing 2: JSON format definition (.json) file of $^1\text{H}_2^{16}\text{O}$ POKAZATEL dataset.

```
{
  "isotopologue": {
    "iso_formula": "(1H)2(160)",
    "iso_slug": "1H2-160",
    "inchikey": "XLYOFNQQVPJJNP-UHFFFAOYSA-N",
    "inchi": "1S/H20/h1H2",
    "mass_in_Da": 18.010565,
    "point_group": "C2v"
  },
  "atoms": {
    "number_of_atoms": 3,
    "element": {
      "H": 1,
      "O": 16
    }
  },
  "nuclear_spin_degeneracy": {
    "A1": 1,
    "A2": 1,
    "B1": 3,
    "B2": 3
  },
  "dataset": {
    "name": "POKAZATEL",
    "version": 20230621,
    "doi": "10.1093/mnras/sty1877",
    "max_temperature": 6000.0,
    "num_pressure_broadeners": 4,
    "nxsec_files": 0,
    "nkcoeff_files": 0,
    "dipole_available": false,
    "cooling.function_available": false,
    "specific_heat_available": true,
    "continuum": null,
    "predis": null,
    "states": {
      "number_of_states": 810269,
      "max_energy": 41200.00,
      "uncertainties_available": true,
      "lifetime_available": false,
      "lande_g_available": false,
      "num_quanta": 6,
      "states_file_fields": [
        {
          "name": "ID",
          "desc": "Unique integer identifier for the energy level",
          "ffmt": "I12",
          "cfmt": "%12d"
        },
        {
          "name": "E",
          "desc": "State energy in cm-1",
          "ffmt": "F12.6",
          "cfmt": "%12.6f"
        },
        {
          "name": "gtot",
          "desc": "Total energy level degeneracy",
          "ffmt": "I6",
          "cfmt": "%6d"
        },
        {
          "name": "J",
          "desc": "Total rotational quantum number, excluding nuclear spin",
          "ffmt": "I7",
          "cfmt": "%7d"
        },
        {
          "name": "unc",
          "desc": "Energy uncertainty in cm-1",
          "ffmt": "F12.6",
          "cfmt": "%12.6f"
        },
        {
          "name": "nltcs:Ka",
          "ffmt": "I2",
          "cfmt": "%2d",
          "desc": "Ka rotational quantum number"
        }
      ]
    }
  }
}
```

```

    {
      "name": "nltcs:Kc",
      "ffmt": "I2",
      "cfmt": "%2d",
      "desc": "Kc rotational quantum number"
    },
    {
      "name": "nltcs:v1",
      "ffmt": "I2",
      "cfmt": "%2d",
      "desc": "v1 symmetric stretch quantum number(3)"
    },
    {
      "name": "nltcs:v2",
      "ffmt": "I2",
      "cfmt": "%2d",
      "desc": "v2 bend quantum number"
    },
    {
      "name": "nltcs:v3",
      "ffmt": "I2",
      "cfmt": "%2d",
      "desc": "v3 asymmetrics stretch quantum number"
    },
    {
      "name": "nltcs:Grve",
      "ffmt": "A3",
      "cfmt": "%3s",
      "desc": "Rovibrational symmetry label"
    },
    {
      "name": "Auxiliary:Ecal",
      "desc": "Energy in cm-1 from variational spectroscopic model",
      "ffmt": "F12.6",
      "cfmt": "%12.6f"
    },
    {
      "name": "Auxiliary:SourceType",
      "ffmt": "A2",
      "cfmt": "%2s",
      "desc": "Indicates if the value is from MARVEL (Ma) or calculated (Ca)"
    }
  ]
},
"transitions": {
  "number_of_transitions": 5745071340,
  "number_of_transition_files": 412,
  "max_wavenumber": 41200.00
},
"partition_function": {
  "max_partition_function_temperature": 10000.0,
  "partition_function_step_size": 1.0
},
"broad": {
  "default_Lorentzian_half-width": 0.07,
  "default_temperature_exponent": 0.5,
  "H2": {
    "filename": "1H2-160__H2.broad",
    "max_J": 50,
    "Lorentzian_half_width": 0.0209,
    "temperature_exponent": 0.027,
    "quantum_number_sets": [
      {
        "code": "a3",
        "num_lines": 23731,
        "quantum_numbers": [
          "J''",
          "Ka''",
          "Ka'"
        ]
      },
      {
        "code": "a1",
        "num_lines": 100,
        "quantum_numbers": [
          "J''"
        ]
      }
    ]
  }
}

```

```

        "code": "a0",
        "num_lines": 51,
        "quantum_numbers": []
    },
    ],
},
"He": {
    "filename": "1H2-160__He.broad",
    "max_J": 50,
    "Lorentzian_half_width": 0.0042,
    "temperature_exponent": 0.02,
    "quantum_number_sets": [
        {
            "code": "b1",
            "num_lines": 253,
            "quantum_numbers": [
                "J'",
                "Ka\"",
                "Ka'",
                "Kc\"",
                "Kc'",
                "v1\"",
                "v2\"",
                "v3\"",
                "v1'",
                "v2'",
                "v3'"
            ]
        },
        {
            "code": "a3",
            "num_lines": 23731,
            "quantum_numbers": [
                "J'",
                "Ka\"",
                "Ka'"
            ]
        },
        {
            "code": "a1",
            "num_lines": 100,
            "quantum_numbers": [
                "J'"
            ]
        },
        {
            "code": "a0",
            "num_lines": 51,
            "quantum_numbers": []
        }
    ]
}
}

```