

A Field-Theoretic Example for Hodge Theory in 3D

A. K. Rao^(a), R. P. Malik^(a,b)

^(a) *Department of Physics, Institute of Science,
Banaras Hindu University (BHU), Varanasi - 221 005, (U.P.), India*

^(b) *DST Centre for Interdisciplinary Mathematical Sciences,
Institute of Science, Banaras Hindu University (BHU), Varanasi - 221 005, India*
e-mails: amit.akrao@gmail.com; rpmalik1995@gmail.com

Abstract: We focus on the continuous symmetry transformations for the three $(2 + 1)$ -dimensional (3D) system of a combination of the *free* Abelian 1-form and 2-form gauge theories within the framework of Becchi-Rouet-Stora-Tyutin (BRST) formalism. We establish that this *combined* system is a tractable field-theoretic model of Hodge theory. The symmetry operators of our present system provide the physical realizations of the de Rham cohomological operators of differential geometry at the algebraic level. Our present investigation is important in the sense that, for the first time, we are able to establish an *odd* dimensional (i.e. $D = 3$) field-theoretic system to be an example for Hodge theory (besides earlier works on a few interesting $(0 + 1)$ -dimensional (1D) toy models as well as a set of well-known $\mathcal{N} = 2$ SUSY quantum mechanical systems of physical interest). For the sake of brevity, we have *not* taken into account the 3D Chern-Simon term for the Abelian 1-form gauge field in our theory which allows the mass and gauge-invariance to co-exist together.

PACS numbers: 11.15.-q, 03.70.+k

Keywords: 3D Abelian 1-form and 2-form gauge theories; off-shell nilpotent (anti-)BRST symmetries; off-shell nilpotent (anti-)co-BRST symmetries; a unique bosonic symmetry; ghost-scale symmetry; cohomological operators; algebraic structures

1 Introduction

The (super)string theories (see, e.g. [1-3] and references therein) are the forefront areas of research activities in the modern day theoretical high energy physics (THEP) which have brought together researchers in the domains of THEP and mathematics on a single intellectual platform. The modern developments in these theories have led to many other areas of research activities in THEP and pure mathematics. One of them is the study of the higher p -form ($p = 2, 3, \dots$) gauge theories because the higher p -form basic fields appear in the quantum excitations of (super)strings. Hence, very naturally, the reach and range of the (super)string theories go beyond the realm of the standard model of particle physics which is based on the interacting non-Abelian 1-form (i.e. $p = 1$) gauge theory. In this connection, it is pertinent to point out that, during the last few years, we have devoted time on the research activities that are connected with the Becchi-Rouet-Stora-Tyutin (BRST) approach to the higher p -form ($p = 2, 3, \dots$) gauge theories and established that the massless and Stückelberg-modified massive Abelian p -form ($p = 1, 2, 3, \dots$) gauge theories in $D = 2p$ dimensions of spacetime are the field-theoretic examples for Hodge theory where there is confluence of ideas from the physics part of the BRST formalism [4-7] and mathematical aspects of differential geometry at the algebraic level. Hence, we observe that, in our studies on the BRST approach to the models of Hodge theory (see, e.g. [8-10] and reference therein), the ideas from theoretical physics and pure mathematics are beautifully intertwined together in a meaningful fashion.

It is clear, from the above discussions, that the *above* field-theoretic models of Hodge theory are defined only in the even (i.e. $D = 2p$) dimensions of spacetime. The purpose of this Letter is to establish that the continuous symmetry transformation operators of the BRST-quantized *combined* system of the free Abelian 1-form and 2-form gauge theories provides a field-theoretic example for Hodge theory (within the framework of BRST formalism) in the three ($2 + 1$)-dimensional (3D) spacetime. This is for the first time, we are able to show an odd-dimensional ($D = 3$) field-theoretic system to be a model for Hodge theory (besides earlier works [11-13] on the one ($0 + 1$)-dimensional (1D) systems of a rigid rotor [11], the celebrated FLPR model [12] and the dynamics of a particle on a torus [13]). These 1D systems have also been shown to be the examples for Hodge theory. In addition, we have also devoted time on the proof that the $\mathcal{N} = 2$ SUSY quantum mechanical systems *also* belong to this class of Hodge theory (see, e.g. [14,15] and references therein). In our present investigation, we show that our present 3D field-theoretical model provides the physical realizations of the de Rham cohomological operators* of differential geometry (see, e.g. [16-18]) in terms of the continuous symmetry transformations (where there are total *six* of them). If the beauty of a theory is defined in terms of the numbers and varieties of symmetries it respects, the models of Hodge theory belong to this distinguished class.

Some of the key observations of our present endeavor are as follows. We note that the kinetic terms (owing their origin to the exterior derivative of differential geometry) for

*On a compact manifold without a boundary, a set of three operators (d, δ, Δ) belong to the de Rham cohomological operators which satisfy the algebra: $d^2 = \delta^2 = 0$, $\Delta = (d + \delta)^2 = \{d, \delta\}$, $[\Delta, d] = [\Delta, \delta] = 0$ where $(\delta)d$ are christened as the (co-)exterior derivatives and Δ is known as the Laplacian operator of differential geometry [16-18]. The (co-)exterior derivatives are connected by the relationship: $\delta = \pm * d *$ where $*$ is the Hodge duality operator (that is defined on the above compact manifold without a boundary).

the Abelian 1-form and 2-form theories remain invariant under the (anti-)BRST symmetry transformations. On the other hand, the gauge-fixing terms (owing their origin to the co-exterior derivative of differential geometry) for the Abelian 1-form and 2-form gauge-fields remain invariant under the (anti-)co-BRST symmetry transformations. The (anti-)ghost fields *either* do not transform at all *or* transform up to a U(1) gauge-type transformation under the *unique* bosonic symmetry transformation (which is nothing but an appropriate anticommutator of the nilpotent (anti-)BRST and (anti-)co-BRST symmetry transformations). The characteristic feature of the ghost-scale symmetry transformation is the observation that *only* the fields of the ghost-sector transform under it (and fields from the non-ghost sector do *not* transform at all). The algebraic structures, obeyed by these continuous symmetry operators, are reminiscent of the Hodge algebra that is respected by the de Rham cohomological operators of differential geometry. Hence, our present 3D system (of the BRST-invariant free Abelian 1-form and 2-form gauge theories without the Chern-Simon term) provides a field-theoretic example for Hodge theory.

The theoretical contents of our present endeavor are organized in the following order. First of all, in section two, we recapitulate the essentials of the (anti-)BRST symmetry transformations $[s_{(a)b}]$ from our earlier work [19]. Our section three is devoted to the discussion on the (anti-)co-BRST symmetry transformations $[s_{(a)d}]$. The subject matter of our section four is connected with the derivation of a unique bosonic symmetry transformation (s_w) from the appropriate anticommutators between the (anti-)BRST and (anti-)co-BRST transformations. In section five, we devote time on the derivation of the ghost-scale symmetry transformations (and their infinitesimal versions). Our section six deals with the algebraic structures of the symmetry operators of our theory and their connections with the Hodge algebra of cohomological operators. Finally, in section seven, we summarize our key results and point out the future prospective of our present investigation.

2 (Anti-)BRST Symmetries: Key Aspects

We begin with the BRST-invariant Lagrangian density[†] which is the sum (i.e. $\mathcal{L}_B = \mathcal{L}_B^{(1)} + \mathcal{L}_B^{(2)}$) of the BRST-invariant Lagrangian densities for the free Abelian 1-form and 2-form gauge theories as follows

$$\begin{aligned} \mathcal{L}_B &= -\frac{1}{4}F^{\mu\nu}F_{\mu\nu} - B(\partial \cdot A) + \frac{B^2}{2} - \partial_\mu \bar{C} \partial^\mu C + \mathcal{B} \left(\frac{1}{2} \varepsilon^{\mu\nu\sigma} \partial_\mu B_{\nu\sigma} \right) - \frac{\mathcal{B}^2}{2} \\ &+ B^\mu (\partial^\nu B_{\nu\mu} - \partial_\mu \phi) - \frac{B^\mu B_\mu}{2} + \partial_\mu \bar{\beta} \partial^\mu \beta + (\partial_\mu \bar{C}_\nu - \partial_\nu \bar{C}_\mu) (\partial^\mu C^\nu) \\ &+ (\partial \cdot \bar{C} + \rho) \lambda + (\partial \cdot C - \lambda) \rho, \end{aligned} \tag{1}$$

[†]For the flat 3D Minkowskian background spacetime manifold, we choose the metric tensor $\eta_{\mu\nu} = \text{diag}(+1, -1, -1)$ so that the dot product between two non-null vectors P_μ and Q_μ is $P \cdot Q = \eta_{\mu\nu} P^\mu Q^\nu \equiv P_0 Q_0 - P_i Q_i$ where the Greek indices $\mu, \nu, \sigma, \dots = 0, 1, 2$ stand for the time and space directions of our 3D spacetime manifold and the Latin indices $i, j, k = 1, 2$ correspond to space directions only. We have chosen 3D Levi-Civita tensor $\varepsilon_{\mu\nu\sigma}$ such that $\varepsilon_{012} = +1 = \varepsilon^{012}$ and it satisfies the standard relationships: $\varepsilon^{\mu\nu\sigma} \varepsilon_{\mu\nu\sigma} = 3!$, $\varepsilon^{\mu\nu\sigma} \varepsilon_{\mu\nu\eta} = 2! \delta_\eta^\sigma$, $\varepsilon^{\mu\nu\sigma} \varepsilon_{\mu\kappa\eta} = 1! (\delta_\kappa^\nu \delta_\eta^\sigma - \delta_\eta^\nu \delta_\kappa^\sigma)$ on the 3D flat spacetime manifold.

where $\mathcal{L}_B^{(1)} = -\frac{1}{4}F^{\mu\nu}F_{\mu\nu} - B(\partial \cdot A) + \frac{1}{2}B^2 - \partial_\mu \bar{C} \partial^\mu C$ is the (anti-)BRST invariant Lagrangian density for the free Abelian 1-form gauge theory and rest of the terms of the above Lagrangian density (1) are for the BRST-invariant Lagrangian density in the case of a 3D free Abelian 2-form theory. As a side remark, we would like to mention that the above *combined* system of the BRST-invariant Lagrangian density for the Abelian 1-form and 2-form gauge theories is the limiting case of the BRST invariant Lagrangian density that has been considered in our recent work [19] on the BRST approach to the Stückelberg-modified 3D massive Abelian 2-form theory with the conditions: $\tilde{\varphi} = 0$, $m = 0$ and $\phi_\mu = A_\mu$ so that $\Sigma_{\mu\nu} = F_{\mu\nu}$. Here the field $\tilde{\varphi}$ corresponds to the pseudo-scalar field, m stands for the rest mass of the Abelian 2-form field and the Lorentz vector field ϕ_μ denotes the Stückelberg field (with its field-strength tensor as: $\Sigma_{\mu\nu} = \partial_\mu \phi_\nu - \partial_\nu \phi_\mu$). The special feature of the 3D Abelian 2-form theory is that the kinetic term for the antisymmetric ($B_{\mu\nu} = -B_{\nu\mu}$) tensor field $B_{\mu\nu}$ turns out to be the following

$$\frac{1}{12} H^{\mu\nu\lambda} H_{\mu\nu\lambda} = \frac{1}{2} H^{012} H_{012} = \frac{1}{2} (H_{012})^2, \quad (2)$$

where the covariant form of $H_{012} = \frac{1}{2} \varepsilon^{\mu\nu\sigma} \partial_\mu B_{\nu\sigma}$ has been taken into account in (1). A set of three Nakanishi-Lautrup type auxiliary fields (\mathcal{B} , B , B_μ) have been invoked to linearize the kinetic term and the gauge-fixing terms for the gauge fields A_μ and $B_{\mu\nu}$, respectively. In the above, the 2-form: $B^{(2)} = \frac{1}{2} B_{\mu\nu} (dx^\mu \wedge dx^\nu)$ defines the antisymmetric ($B_{\mu\nu} = -B_{\nu\mu}$) tensor gauge field $B_{\mu\nu}$ and the 3-form: $H^{(3)} = dB^{(2)} = \frac{1}{3!} H_{\mu\nu\lambda} (dx^\mu \wedge dx^\nu \wedge dx^\lambda)$ defines the field-strength tensor $H_{\mu\nu\lambda} = \partial_\mu B_{\nu\lambda} + \partial_\nu B_{\lambda\mu} + \partial_\lambda B_{\mu\nu}$ for the antisymmetric tensor gauge field. Here $d = \partial_\mu dx^\mu$ [with $d^2 = \frac{1}{2!} (\partial_\mu \partial_\nu - \partial_\nu \partial_\mu) (dx^\mu \wedge dx^\nu) = 0$] is the exterior derivative of differential geometry. The scalar field ϕ is present in the theory due to the stage-one reducibility for our 2-form field $B_{\mu\nu}$. Similarly, for the Abelian 1-form theory, we have the field strength tensor $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ which is derived from the 2-form $F^{(2)} = dA^{(1)} = \frac{1}{2!} F_{\mu\nu} (dx^\mu \wedge dx^\nu)$ where d is the exterior derivative and the 1-form $A^{(1)} = A_\mu dx^\mu$ defines the vector potential A_μ of our Abelian 1-form gauge theory..

In the BRST invariant Lagrangian density (1), we have the Lorentz vector fermionic (i.e. $C_\mu^2 = \bar{C}_\mu^2 = 0$, $C_\mu C_\nu + C_\nu C_\mu = 0$, $C_\mu \bar{C}_\nu + \bar{C}_\nu C_\mu = 0$, etc.) (anti-)ghost fields $(\bar{C}_\mu)C_\mu$ with ghost numbers $(-1) + 1$, respectively. On the other hand, the (anti-)ghost fields $(\bar{\beta})\beta$ are the ghost-for-ghost fields which are bosonic (i.e. $\beta^2 \neq 0$, $\bar{\beta}^2 \neq 0$) in nature and they carry the ghost numbers $(-2) + 2$, respectively. The fermionic (i.e. $C^2 = 0$, $\bar{C}^2 = 0$, $C\bar{C} + \bar{C}C = 0$) (anti-)ghost fields $(\bar{C})C$ are endowed with ghost numbers $(-1) + 1$, respectively. These *latter* (anti-)ghost fields correspond to the Abelian 1-form gauge field (within the framework of BRST formalism). The auxiliary (anti-)ghost fields $(\rho)\lambda$ of our system also carry the ghost numbers $(-1)+1$, respectively, because we note that $\rho = -(1/2)(\partial \cdot \bar{C})$ and $\lambda = (1/2)(\partial \cdot C)$. These (anti-)ghost fields are required to maintain the property of unitarity in our BRST-invariant theory which is valid at any arbitrary order of perturbative computation for the physical processes that are allowed by our BRST-quantized theory. These (anti-)ghost fields appear in the nilpotent versions of the (anti-)BRST symmetry transformations which are nothing but the generalizations of the *classical* local gauge symmetry transformations of the original theory to the *quantum* level.

The above Lagrangian density (1) respects the following, continuous and off-shell nilpo-

tent ($s_b^2 = 0$) BRST symmetry transformations

$$\begin{aligned}
s_b B_{\mu\nu} &= -(\partial_\mu C_\nu - \partial_\nu C_\mu), & s_b C_\mu &= -\partial_\mu \beta, & s_b \bar{C}_\mu &= -B_\mu, \\
s_b A_\mu &= \partial_\mu C, & s_b \bar{C} &= B, & s_b \bar{\beta} &= -\rho, & s_b \phi &= +\lambda, & s_b F_{\mu\nu} &= 0, \\
s_b [\rho, \lambda, C, \beta, B_\mu, B, \mathcal{B}, H_{\mu\nu\lambda}] &= 0,
\end{aligned} \tag{3}$$

because we observe that:

$$s_b \mathcal{L}_B = -\partial_\mu \{(\partial^\mu C^\nu - \partial^\nu C^\mu) B_\nu + \lambda B^\mu + \rho \partial^\mu \beta + B \partial^\mu C\}. \tag{4}$$

As a consequence, the action integral $S = \int d^3x \mathcal{L}_B$ remains invariant (i.e. $s_b S = 0$) because all the physical fields vanish off as $x \rightarrow \pm\infty$. The analogous to the Lagrangian density \mathcal{L}_B , we have an anti-BRST invariant Lagrangian density ($\mathcal{L}_{\bar{B}}$) as

$$\begin{aligned}
\mathcal{L}_{\bar{B}} &= \mathcal{B} \left(\frac{1}{2} \varepsilon^{\mu\nu\sigma} \partial_\mu B_{\nu\sigma} \right) - \frac{\mathcal{B}^2}{2} + \bar{B}^\mu (\partial^\nu B_{\nu\mu} + \partial_\mu \phi) - \frac{\bar{B}^\mu \bar{B}_\mu}{2} - \frac{1}{4} F^{\mu\nu} F_{\mu\nu} - B(\partial \cdot A) + \frac{B^2}{2} \\
&- \partial_\mu \bar{C} \partial^\mu C + \partial_\mu \bar{\beta} \partial^\mu \beta + (\partial_\mu \bar{C}_\nu - \partial_\nu \bar{C}_\mu) (\partial^\mu C^\nu) + (\partial \cdot \bar{C} + \rho) \lambda + (\partial \cdot C - \lambda) \rho,
\end{aligned} \tag{5}$$

where \bar{B}_μ is a new Nakanishi-Lautrup type auxiliary field that has been invoked to linearize the gauge-fixing term for the Abelian antisymmetric tensor field $B_{\mu\nu}$ where we have taken into account $\phi \rightarrow -\phi$ for the sake of generality in the gauge-fixing term. The above Lagrangian density ($\mathcal{L}_{\bar{B}}$) respects the following infinitesimal, continuous and off-shell nilpotent ($s_{ab}^2 = 0$) anti-BRST symmetry transformations

$$\begin{aligned}
s_{ab} B_{\mu\nu} &= -(\partial_\mu \bar{C}_\nu - \partial_\nu \bar{C}_\mu), & s_{ab} \bar{C}_\mu &= -\partial_\mu \bar{\beta}, & s_{ab} C_\mu &= +\bar{B}_\mu, \\
s_{ab} A_\mu &= \partial_\mu \bar{C}, & s_{ab} C &= -B, & s_{ab} \beta &= -\lambda, & s_{ab} \phi &= +\rho, & s_{ab} F_{\mu\nu} &= 0, \\
s_{ab} [\rho, \lambda, \bar{C}, \bar{\beta}, \bar{B}_\mu, B, \mathcal{B}, H_{\mu\nu\lambda}] &= 0,
\end{aligned} \tag{6}$$

because we observe that ($\mathcal{L}_{\bar{B}}$) transforms to a total spacetime derivative as follows:

$$s_{ab} \mathcal{L}_{\bar{B}} = -\partial_\mu \{(\partial^\mu \bar{C}^\nu - \partial^\nu \bar{C}^\mu) \bar{B}_\nu - \rho \bar{B}^\mu + \lambda \partial^\mu \bar{\beta} + B \partial^\mu \bar{C}\}. \tag{7}$$

As a consequence, the action integral $S = \int d^3x \mathcal{L}_{\bar{B}}$ remains invariant (i.e. $s_{ab} S = 0$) due to the Gauss divergence theorem because all the physical fields of our theory vanish off as $x \rightarrow \pm\infty$. Thus, we note that $\mathcal{L}_{\bar{B}}$ respects the nilpotent anti-BRST transformations s_{ab} .

We end this section with the following remarks. First of all, we note that the field strength tensors $H_{\mu\nu\lambda}$ and $F_{\mu\nu}$ (owing their origin to the exterior derivative d of differential geometry) remain invariant under the (anti-)BRST symmetry transformations. Second, the nilpotent versions of the (anti-)BRST symmetry transformations are valid in any arbitrary dimension of spacetime. Third, it is well-known that the Lagrangian density $\mathcal{L}_B^{(1)} = -\frac{1}{4} F^{\mu\nu} F_{\mu\nu} + \frac{1}{2} B^2 - B(\partial \cdot A) - \partial_\mu \bar{C} \partial^\mu C$ for the Abelian 1-form gauge theory is invariant under the (anti-)BRST symmetry transformations which are off-shell nilpotent and absolutely anticommuting in nature[‡] without invoking any kind of CF-type restriction.

[‡]The nilpotent (anti-)BRST symmetry transformations: $s_{ab} A_\mu = \partial_\mu \bar{C}$, $s_{ab} \bar{C} = 0$, $s_{ab} C = -B$, $s_{ab} B = 0$ and $s_b A_\mu = \partial_\mu C$, $s_b C = 0$, $s_b \bar{C} = +B$, $s_b B = 0$ satisfy the absolute anticommutativity property (i.e. $\{s_b, s_{ab}\} = 0$) without any recourse to the CF-type restriction because the *latter* is trivial in this case.

On the other hand, the (anti-)BRST symmetry transformations for the Abelian 2-form theory are absolutely anticommuting only when one invokes the validity[§] of the CF-type restriction: $B_\mu - \bar{B}_\mu + 2\partial_\mu\phi = 0$ which emerges from the EL-EoMs w.r.t. the auxiliary fields B_μ and \bar{B}_μ from the Lagrangian densities \mathcal{L}_B and $\mathcal{L}_{\bar{B}}$, respectively (see, e.g. [19]). Finally, we would like to mention that there are (anti-)co-BRST symmetry transformations for the free Abelian 1-form theory in the two $(1 + 1)$ -dimensions of spacetime (see, e.g. [20]). We have also shown that for the Abelian 2-form theory, there is existence of the nilpotent (anti-)co-BRST transformations in the four $(3 + 1)$ -dimensions of the spacetime (see, e.g. [10]). We shall see, in our next section, that there is existence of the (anti-)co-BRST symmetry transformations for our present system (which is a combination of the Abelian 1-form and 2-form theories) in the *three* $(2 + 1)$ -dimensions of flat spacetime.

3 (Anti-)co-BRST Symmetries: Salient Features

We point out that the gauge-fixing terms $(\partial \cdot A)$ and $(\partial^\nu B_{\nu\mu})$ for the Abelian 1-form and 2-form gauge fields owe their origin to the co-exterior derivative $\delta = \mp * d *$ (with $\delta^2 = 0$) where $*$ is the Hodge duality operator on a given spacetime manifold. It is straightforward to note that $\delta A^{(1)} = + * d * A^{(1)} = (\partial \cdot A)$ and $\delta B^{(2)} = - * d * B^{(2)} = (\partial^\nu B_{\nu\mu}) dx^\mu$ because the operation of the co-exterior derivative of differential geometry on a given form reduces the degree of the form by *one*. We demand that, under the (anti-)co-BRST [or (anti-)dual-BRST] symmetry transformations $[s_{(a)d}]$, the total gauge-fixing term $(\partial \cdot A)$ and $(\partial^\nu B_{\nu\mu} \mp \partial_\mu\phi)$ should remain invariant. Against the backdrop of the above discussion, it can be checked that the under the following infinitesimal, continuous and off-shell nilpotent (anti-)co-BRST symmetry transformations $[s_{(a)d}]$, namely;

$$\begin{aligned} s_{ad} B_{\mu\nu} &= -\varepsilon_{\mu\nu\sigma} \partial^\sigma C, & s_{ad} C_\mu &= \partial_\mu \beta, & s_{ad} \bar{C} &= \mathcal{B}, & s_{ad} \bar{\beta} &= \rho, \\ s_{ad} [\rho, \lambda, C, \beta, \bar{C}_\mu, (\partial^\nu B_{\nu\mu}), \phi, A_\mu, F_{\mu\nu}, B, B_\mu, \bar{B}_\mu] &= 0, \end{aligned} \quad (8)$$

$$\begin{aligned} s_d B_{\mu\nu} &= -\varepsilon_{\mu\nu\sigma} \partial^\sigma \bar{C}, & s_d \bar{C}_\mu &= -\partial_\mu \bar{\beta}, & s_d C &= -\mathcal{B}, & s_d \beta &= -\lambda, \\ s_d [\rho, \lambda, B, B_\mu, \bar{B}_\mu, \bar{C}, \bar{\beta}, C_\mu, A_\mu, (\partial^\nu B_{\nu\mu}), F_{\mu\nu}, \phi] &= 0, \end{aligned} \quad (9)$$

both the Lagrangian densities \mathcal{L}_B and $\mathcal{L}_{\bar{B}}$ transform in a similar fashion[¶]. To be precise, we note that \mathcal{L}_B and $\mathcal{L}_{\bar{B}}$ transform under the above (anti-)co-BRST symmetry transformations $[s_{(a)d}]$ to the total spacetime derivatives as follows:

$$\begin{aligned} s_d \mathcal{L}_B &= -\partial_\mu [\mathcal{B} \partial^\mu \bar{C} + \lambda \partial^\mu \bar{\beta}] \equiv s_d \mathcal{L}_{\bar{B}}, \\ s_{ad} \mathcal{L}_B &= -\partial_\mu [\mathcal{B} \partial^\mu C - \rho \partial^\mu \beta] \equiv s_{ad} \mathcal{L}_{\bar{B}}. \end{aligned} \quad (10)$$

[§]It can be readily checked that $\{s_b, s_{ab}\} B_{\mu\nu} = \partial_\mu(B_\nu - \bar{B}_\nu) - \partial_\nu(B_\mu - \bar{B}_\mu)$ by taking into account the (anti-)BRST symmetry transformations in (6) and (3), respectively. The absolute anticommutativity property (i.e. $\{s_b, s_{ab}\} = 0$) is satisfied if and only if we utilize the expression for the CF-type restriction.

[¶]It is pertinent to point out that there is a non-trivial choice of the (anti-)co-BRST symmetry transformations for the vector field A_μ under which the gauge-fixing term $(\partial \cdot A)$ remains invariant. These are: $s_{ad} A_\mu = \pm \varepsilon_{\mu\nu\sigma} \partial^\nu C^\sigma$, $s_d A_\mu = \pm \varepsilon_{\mu\nu\sigma} \partial^\nu \bar{C}^\sigma$. However, this kind of choice does *not* lead to the overall symmetry of the Lagrangian densities \mathcal{L}_B and $\mathcal{L}_{\bar{B}}$. Moreover, we find that the kinetic term (i.e. $-\frac{1}{4} F_{\mu\nu} F^{\mu\nu}$) for the vector field A_μ transforms to an expression that contains *three* derivatives (which is pathological).

The above observation establish that the action integrals $S_1 = \int d^3x \mathcal{L}_{\bar{B}}$ and $S_2 = \int d^3x \mathcal{L}_{\bar{B}}$ both remain invariant under the (anti-)dual-BRST symmetry transformations (i.e. $s_{(a)d} S_1 = s_{(a)d} S_2 = 0$). This happens due to the Gauss divergence theorem because all the physically well-defined fields vanish off as $x \rightarrow \pm\infty$.

There are a few interesting observations when we focus on the infinitesimal, continuous and off-shell nilpotent (anti-)BRST symmetry transformations [cf. Eqs. (6), (3)] and the (anti-)co-BRST symmetry transformations [cf. Eqs. (9), (8)] in their operator forms. For instance, we note that the following anticommutators are true, namely;

$$\{s_d, s_{ad}\} = 0, \quad \{s_d, s_{ab}\} = 0, \quad \{s_b, s_{ad}\} = 0, \quad \{s_d, s_{ab}\} = 0. \quad (11)$$

All the above anticommutators are automatically satisfied *but* for the validity of $\{s_b, s_{ab}\} = 0$. In fact, the *latter* anticommutator (i.e. $\{s_b, s_{ab}\} = 0$) is *also* automatically satisfied for *all* the fields of the theory except the antisymmetric tensor gauge field $B_{\mu\nu}$. In other words, the anticommutator: $\{s_b, s_{ab}\} B_{\mu\nu} = 0$ is satisfied if and only if the CF-type restriction: $B_\mu - \bar{B}_\mu + 2\partial_\mu\phi = 0$ is invoked for its validity. Ultimately, we note that out of the *four* nilpotent (anti-)BRST and (anti-)co-BRST symmetry transformations of our theory, only the following *two* non-trivial anticommutators, namely;

$$\{s_b, s_d\} \neq 0, \quad \{s_{ad}, s_{ab}\} \neq 0, \quad (12)$$

are the ones that lead to the definitions of the bosonic symmetry transformations.

We wrap-up this section with the following concluding remarks. First of all, we note that the gauge-fixing terms [owing their origin to the co-exterior (or dual-exterior) derivative of differential geometry] remain invariant under the (anti-)co-BRST symmetry transformations. Second, the absolute anticommutativity property (i.e. $\{s_d, s_{ad}\} = 0$) is automatically satisfied without invoking any kind of CF-type restriction. Third, both the coupled Lagrangian densities respect both (i.e. co-BRST and anti-co-BRST) symmetry transformations. Fourth, it is interesting to point out that the variation of the kinetic term for the Abelian 2-form field under the (anti-)co-BRST symmetry transformations is compensated by the variation of the ghost term of the Abelian 1-form theory. Finally, only *two* of the anticommutators between the (anti-)BRST and (anti-)co-BRST symmetry transformations are non-trivial [cf. Eq. (12)] which lead to the definition of a *unique* bosonic symmetry transformation which we discuss in our next section.

4 Bosonic Symmetry Transformations: Uniqueness

We have noted that there are two non-trivial anticommutators [cf. Eq. (12)] between the off-shell nilpotent (anti-)BRST and (anti-)co-BRST symmetry transformations which define the following bosonic symmetry transformations:

$$s_w = \{s_b, s_d\}, \quad s_{\bar{w}} = \{s_{ad}, s_{ab}\}. \quad (13)$$

We shall see that only one of the above bosonic symmetry transformations is the independent bosonic symmetry of our theory which turns out to be *unique*. To corroborate

this statement, we note that the following infinitesimal and continuous bosonic symmetry transformations (i.e. $s_w = \{s_b, s_d\}$) are true, namely;

$$\begin{aligned} s_w B_{\mu\nu} &= -\varepsilon_{\mu\nu\sigma} \partial^\sigma B, & s_w \bar{C}_\mu &= \partial_\mu \rho, & s_w C_\mu &= \partial_\mu \lambda, \\ s_w A_\mu &= -\partial_\mu \mathcal{B}, & s_w [\rho, \lambda, B, B_\mu, \bar{B}_\mu, \mathcal{B}, C, \bar{C}, \beta, \bar{\beta}, \phi] &= 0, \end{aligned} \quad (14)$$

where we have used the symmetry transformations that have been listed in (3) and (9) to determine the transformation (14) which corresponds to $s_w = \{s_b, s_d\}$. On the other hand, we obtain the following symmetry transformations for the fields of our theory, corresponding to the bosonic symmetry transformations: $s_{\bar{w}} = \{s_{ab}, s_{ad}\}$, namely;

$$\begin{aligned} s_{\bar{w}} B_{\mu\nu} &= \varepsilon_{\mu\nu\sigma} \partial^\sigma B, & s_{\bar{w}} \bar{C}_\mu &= -\partial_\mu \rho, & s_{\bar{w}} C_\mu &= -\partial_\mu \lambda, \\ s_{\bar{w}} A_\mu &= \partial_\mu \mathcal{B}, & s_{\bar{w}} [\rho, \lambda, B, B_\mu, \bar{B}_\mu, \mathcal{B}, C, \bar{C}, \beta, \bar{\beta}, \phi] &= 0. \end{aligned} \quad (15)$$

A close and careful look at (14) and (15) shows that we have the validity of $s_w + s_{\bar{w}} = 0$. In other words, we have a *unique* bosonic symmetry in the theory. This claim is corroborated, once again, in the following observations:

$$\begin{aligned} s_w \mathcal{L}_B &= \partial_\mu \left[B \partial^\mu \mathcal{B} - \mathcal{B} \partial^\mu B - \rho \partial^\mu \lambda + (\partial^\mu \rho) \lambda \right], \\ s_{\bar{w}} \mathcal{L}_B &= -\partial_\mu \left[\mathcal{B} \partial^\mu B - B \partial^\mu \mathcal{B} - \rho \partial^\mu \lambda + (\partial^\mu \rho) \lambda \right], \end{aligned} \quad (16)$$

which establish that $(s_w + s_{\bar{w}}) \mathcal{L}_B = 0$. In other words the operator equation: $s_w + s_{\bar{w}} = 0$ implies that only *one* bosonic symmetry transformation is independent. This proves the *uniqueness* of the infinitesimal and continuous bosonic symmetry transformations.

We end this section with the following concluding remarks. First of all, we note that, under the bosonic symmetry transformations (14) and/or (15), all the ghost fields either do *not* transform at all *or* transform upto a U(1) gauge symmetry type of transformations. The *latter* observation is true *only* for the Lorentz vector (anti-)ghost fields $(\bar{C}_\mu) C_\mu$. Second, out of *four* off-shell nilpotent symmetries, only *two* anticommutators (out of *six* possible anticommutators [cf. Eqs. (11), (12)]) are non-trivial [cf. Eq. (12)] which define the bosonic symmetry in our theory. Out of these two bosonic symmetry transformations, only *one* of them is independent (e.g. s_w) which proves its uniqueness. Finally, a close look at (11), (12) and (13) establishes the fact that the following commutators are true, namely;

$$[s_w, s_b] = 0, \quad [s_w, s_{ab}] = 0, \quad [s_w, s_d] = 0, \quad [s_w, s_{ad}] = 0, \quad (17)$$

where we have taken into account the off-shell nilpotency ($s_{(a)b}^2 = 0$, $s_{(a)d}^2 = 0$) property of the (anti-)BRST and (anti-)co-BRST symmetry transformations.

5 Ghost-Scale Transformations: Key Observations

It is straightforward to check that under the following ghost-scale transformations

$$\begin{aligned} \beta &\longrightarrow e^{+2\Sigma} \beta, & \bar{\beta} &\longrightarrow e^{-2\Sigma} \bar{\beta}, & C_\mu &\longrightarrow e^{+\Sigma} C_\mu, & \bar{C}_\mu &\longrightarrow e^{-\Sigma} \bar{C}_\mu, \\ C &\longrightarrow e^{+\Sigma} C, & \bar{C} &\longrightarrow e^{-\Sigma} \bar{C}, & \lambda &\longrightarrow e^{+\Sigma} \lambda, & \rho &\longrightarrow e^{-\Sigma} \rho, \\ \Phi &\longrightarrow e^0 \Phi & (\Phi = A_\mu, B_{\mu\nu}, \phi, \bar{B}_\mu, B_\mu, \mathcal{B}, B), \end{aligned} \quad (18)$$

where Σ is a global (i.e. spacetime independent) scale transformation parameter, the Lagrangian densities \mathcal{L}_B and $\mathcal{L}_{\bar{B}}$ remain invariant. For the sake of brevity, if we choose the scale parameter $\Sigma = 1$, the infinitesimal version (s_g) of the above ghost-scale symmetry transformations reduce to the following simple forms, namely;

$$\begin{aligned} s_g \beta &= +2\beta, & s_g \bar{\beta} &= -2\bar{\beta}, & s_g C_\mu &= +C_\mu, & s_g \bar{C}_\mu &= -\bar{C}_\mu, \\ s_g C &= +C, & s_g \bar{C} &= -\bar{C}, & s_g \lambda &= +\lambda, & s_g \rho &= -\rho, & s_g \Phi &= 0. \end{aligned} \quad (19)$$

It is straightforward to check that, under the above infinitesimal ghost scale symmetry transformations, we obtain: $s_g \mathcal{L}_B = 0$, $s_g \mathcal{L}_{\bar{B}} = 0$. Hence, the action integrals corresponding to the Lagrangian densities \mathcal{L}_B and $\mathcal{L}_{\bar{B}}$, also remain invariant under s_g .

We end this section with a couple of clinching remarks. First of all, we note that only the (anti-)ghost basic and auxiliary [i.e. $\lambda = \frac{1}{2}(\partial \cdot C)$, $\rho = -\frac{1}{2}(\partial \cdot \bar{C})$] fields transform under the ghost-scale symmetry transformations and rest of the fields (e.g. $\Phi = B_{\mu\nu}$, A_μ , ϕ , B , \mathcal{B} , B_μ , \bar{B}_μ) of the theory do *not* transform at all. Second, we note that the operator form of the symmetries obey the following very useful relationships:

$$\begin{aligned} [s_g, s_w] &= 0, & [s_g, s_b] &= +s_b, & [s_g, s_{ab}] &= -s_{ab}, \\ [s_g, s_d] &= -s_d, & [s_g, s_{ad}] &= +s_{ad}. \end{aligned} \quad (20)$$

The above commutators establish the fact that the pairs (s_b, s_{ad}) and (s_{ab}, s_d) obey exactly similar kinds of algebra w.r.t. s_g and s_w commutes with s_g . Thus, we note that the bosonic symmetry transformation s_w commutes with *all* the symmetry operators of our theory.

6 Algebraic Structures: Cohomological Operators

In this section, we provide the physical realizations of the de Rham cohomological operators of the differential geometry at the algebraic level and establish that there is two-to-one mapping between the continuous symmetry operators of our theory and the cohomological operators. The *latter* are defined on a compact spacetime manifold without a boundary and they form a set of three operators (d, δ, Δ) where $d = \partial_\mu dx^\mu$ [with $d^2 = \frac{1}{2!}(\partial_\mu \partial_\nu - \partial_\nu \partial_\mu)(dx^\mu \wedge dx^\nu) = 0$] is the exterior derivative, $\delta = \pm * d *$ (with $\delta^2 = 0$) is the co-exterior derivative and $\Delta = (d + \delta)^2 = \{d, \delta\}$ is the Laplacian operator. They obey an algebra which is properly known as the Hodge algebra of differential geometry. This algebra is:

$$\begin{aligned} d^2 &= 0, & \delta^2 &= 0, & \Delta &= (d + \delta)^2 = \{d, \delta\}, \\ [\Delta, d] &= 0, & [\Delta, \delta] &= 0, & \{d, \delta\} &\neq 0. \end{aligned} \quad (21)$$

Thus, we note that the Laplacian operator commute with both the nilpotent ($d^2 = \delta^2 = 0$) exterior and co-exterior derivatives of the differential geometry.

At this juncture, we collect all the algebraic structures which we have discussed in our previous sections. In their full blaze of glory, the algebraic structures are as follows:

$$\begin{aligned} s_b^2 &= 0, & s_d^2 &= 0, & s_{ab}^2 &= 0, & s_{ad}^2 &= 0, & s_w &= \{s_b, s_d\} = -\{s_{ad}, s_{ab}\}, \\ [s_w, s_r] &= 0, & r &= b, ab, d, ad, g, w, & \{s_b, s_{ad}\} &= 0, & \{s_b, s_{ab}\} &= 0, \\ \{s_{ab}, s_d\} &= 0, & \{s_d, s_{ad}\} &= 0, & [s_g, s_b] &= +s_b, & [s_g, s_{ab}] &= -s_{ab}, \\ [s_g, s_{ad}] &= +s_{ad}, & [s_g, s_d] &= -s_d, & [s_g, s_w] &= 0. \end{aligned} \quad (22)$$

A comparison between the equations (21) and (22) establishes the two-to-one mapping between the symmetry operators of (22) and the cohomological operators (21) as

$$(s_b, s_{ad}) \longrightarrow d, \quad (s_d, s_{ab}) \longrightarrow \delta, \quad (s_w, s_{\bar{w}}) \longrightarrow \Delta, \quad (23)$$

where, as has been established in section four, the bosonic symmetry operator $s_{\bar{w}} = -s_w$.

We conclude this section with the remarks that the algebraic relationships between the infinitesimal and continuous ghost-scale symmetry operator (s_g) and other infinitesimal continuous symmetry operator (e.g. $s_b, s_{ab}, s_d, s_{ad}, s_w$) in equation (22) are important because they encompass in their folds the fact that the symmetry operators (s_b, s_{ad}) operating on a specific field raise the ghost number by one. On the other hand, the symmetry operators (s_d, s_{ab}) lower the ghost number by one when they operate on a specific field. In contrast to the above observations, the unique bosonic symmetry operator (s_w) does not change the ghost number of a specific field when it operates on that field. In terms of the Noether conserved charges, corresponding to the above infinitesimal continuous symmetries, the above observations are translated into the quantum Hilbert space of states.

7 Conclusions

In our present investigation, we have concentrated on the continuous symmetry transformations for the coupled (but equivalent) Lagrangian densities of the *combined* system of the free 3D Abelian 1-form and 2-form gauge theories within the framework of BRST formalism and established that the algebra, obeyed by the symmetry transformation operators, is reminiscent of the Hodge algebra that is respected by the de Rham cohomological operators of the differential geometry. We have also demonstrated that the continuous transformations of our theory correspond to the cohomological operators (d, δ, Δ) at the algebraic level and there is two-to-one mapping between them (cf. Eq. (26)]. Despite our sincere efforts, we have *not* yet been able to show the existence of the discrete symmetries in our theory to establish the physical realizations of the Hodge duality $*$ operation which connects the (anti-)co-BRST and (anti-)BRST symmetry transformations (see, e.g. [8-10],20]). In other words, we have not been able to provide the physical realizations of the connection between the (co-)exterior derivatives: $\delta = \pm * d *$ in the language of symmetry operators.

We have reasons to believe that our present *odd* dimensional (i.e. $D = 3$) example for Hodge theory might *not* be endowed with the discrete duality symmetry transformations because there is *no* existence of a pseudo-scalar field in our theory which is dual to the scalar field ϕ that is present in the gauge-fixing term for the Abelian 2-form field (cf. Eqs. (1),(5)]. This is *not* the case with the 2D Stückelberg-modified Proca theory (see, e.g. [20] for details) as well as the 4D Stückelberg-modified massive as well as massless Abelian 2-form theories (see, e.g. [8-10] for details) which are a set of tractable field-theoretic models of Hodge theory. To be precise, we have the existence of (i) the scalar and pseudo-scalar fields in the context of the 2D modified Proca theory [20], and (ii) the pair scalar and pseudo-scalar as well as the pair vector and axial-vector in the context of the 4D Abelian 2-form *modified* massive gauge theory (see, e.g. [10]). This is why there is existence of a set of duality symmetry transformations in these *even* dimensional models of Hodge theory.

We would like to lay emphasis on the fact that in all the *even* dimensional examples for Hodge theory [9-10,20], there is existence of some bosonic fields with negative kinetic terms which obey the usual Klein-Gordon equation of motion. The *latter* observation demonstrates that they are endowed with the well-defined rest masses. Such kinds of “exotic” fields have become quite popular in the realm of THEP as well as in the context of cosmological models of the Universe. For instance, the fields with negative kinetic terms with or without mass are (i) a possible set of candidates for dark matter and dark energy (see. e.g. [21,22] for details), and (ii) a set of fields that has been christened as the “phantom” or “ghost” fields in the contexts of cyclic, bouncing and self-accelerated cosmological models of the Universe (see, e.g. [23-25]) which are responsible for the modern observation of the accelerated expansion of the Universe. It is interesting to point out that, for our present system of an odd-dimensional field-theoretic example for Hodge theory, there is *no* room for the existence of such kinds of fields with negative kinetic terms.

We have, in our present endeavor, *not* computed the conserved Noether charges and *not* studied the algebra followed by them. In addition, we have *not* yet been able to provide the physical realization of the Hodge duality * operation of differential geometry. Thus, at our current level of understanding, our present 3D system should be treated as the field-theoretic example for *quasi*-Hodge theory. In our future investigation [26], we plan to concentrate on these issues and establish, from all aspects of the theoretical angles, that our present *combined* system of the free 3D Abelian 1-form and 2-form theories is indeed a perfect field-theoretic example for Hodge theory in the odd (i.e. $D = 3$) dimension of spacetime. If this happens, it will be a completely *new* result as far as our studies on the BRST approach to the existence of field-theoretic models of Hodge theory are concerned.

Acknowledgments

Both the authors, very humbly and respectfully, dedicate their present work to the memory of Prof. C. P. Singh who was responsible for the revival of the Nuclear and High Energy Physics Group at the Physics Department of BHU and who passed away in the recent past.

Conflicts of Interest

The authors declare that there are no conflicts of interest.

Data Availability

No data were used to support this study.

References

- [1] M. B. Green, J. H. Schwarz, E. Witten, *Superstring Theory*, Vols. 1 and 2, Cambridge University Press, Cambridge (1987)

- [2] J. Polchinski, *String Theory*, Vols. 1 and 2, Cambridge University Press, Cambridge (1998)
- [3] D. Lust, S. Theisen, *Lectures in String Theory*, Springer-Verlag, New York (1989)
- [4] C. Becchi, A. Rouet, R. Stora, *Phys. Lett. B* **52**, 344 (1974)
- [5] C. Becchi, A. Rouet, R. Stora, *Comm. Math. Phys.* **42**, 127 (1975)
- [6] C. Becchi, A. Rouet, R. Stora, *Ann. Phys. (N. Y.)* **98**, 287 (1976)
- [7] I. V. Tyutin, Lebedev Institute Preprint, Report Number: FIAN-39 (1975) (unpublished), arXiv:0812.0580 [hep-th]
- [8] R. Kumar, S. Krishna, A. Shukla, R. P. Malik, *Int. J. Mod. Phys. A* **29**, 1450135 (2014) (*A Brief Review*)
- [9] S. Gupta, R. P. Malik, *Eur. Phys. J. C* **58**, 517 (2008)
- [10] S. Krishna, R. Kumar, R. P. Malik, *Ann. Phys.* **414**, 168087 (2020)
- [11] S. Gupta, R. P. Malik, *Eur. Phys. J. C* **68**, 325 (2010)
- [12] S. Krishna, R. P. Malik, *Ann. Phys.* **464**, 169657 (2024)
- [13] V. K. Pandey, B. P. Mandal, *Adv. High Energy Phys.* **2017**, 6124189 (2017)
- [14] R. Kumar, R. P. Malik, *Eur. Phys. J. C* **73**, 2535 (2013)
- [15] S. Krishna, A. Shukla, R. P. Malik, *Ann. Phys.* **351**, 558 (2014)
- [16] T. Eguchi, P. B. Gilkey, A. Hanson, *Physics Reports* **66**, 213 (1980)
- [17] S. Mukhi, N. Mukunda, *Introduction to Topology, Differential Geometry and group theory for physicists*, Wiley Eastern Private Limited, New Delhi (1990)
- [18] M. Göckeler, T. Schücker, *Differential Geometry, Gauge Theories and Gravity*, Cambridge University Press, Cambridge (1987)
- [19] S. K. Panja, E. Harikumar, R. P. Malik, **arXiv: 2405.15588 [hep-th]**
- [20] B. Chauhan, S. Kumar, A. Tripathi, R. P. Malik, *Adv. High Energy Phys.* **2020**, 3495168 (2020)
- [21] V. M. Zhuravlev, D. A. Kornilov, E. P. Savelova, *Gen. Relat. Gravity* **36**, 1736 (2004)
- [22] Y. Aharonov, S. Popescu, D. Rohrlich, L. Vaidman, *Phys. Rev. A* **48**, 4084 (1993)
- [23] P. J. Steinhardt, N. Turok, *Science* **296**, 1436 (2002)
- [24] Y. F. Cai, A. Marcian, D.-G. Wang, E. Wilson-Ewing, *Universe* **3**, 1 (2017)
- [25] K. Koyama, *Class. Quantum Gravity* **24**, R231 (2007)
- [26] R. P. Malik, *et al.*, in preparation