

Differentially Private Best-Arm Identification

Achraf Azize

ACHRAF.AZIZE@INRIA.FR

Univ. Lille, Inria, CNRS, Centrale Lille, UMR 9189 - CRISTAL, F-59000 Lille, France

Marc Jourdan

MARC.JOURDAN@INRIA.FR

Univ. Lille, Inria, CNRS, Centrale Lille, UMR 9189 - CRISTAL, F-59000 Lille, France

Aymen Al Marjani*

ALMARJAN@AMAZON.LU

Amazon

Debabrota Basu

DEBABROTA.BASU@INRIA.FR

Univ. Lille, Inria, CNRS, Centrale Lille, UMR 9189 - CRISTAL, F-59000 Lille, France

Abstract

Best Arm Identification (BAI) problems are progressively used for data-sensitive applications, such as designing adaptive clinical trials, tuning hyper-parameters, and conducting user studies. Motivated by the data privacy concerns invoked by these applications, we study the problem of *BAI with fixed confidence* in both the local and central models, i.e. ϵ -local and ϵ -global Differential Privacy (DP). First, to quantify the cost of privacy, we derive *lower bounds on the sample complexity* of any δ -correct BAI algorithm satisfying ϵ -global DP or ϵ -local DP. Our lower bounds suggest the existence of two privacy regimes. In the high-privacy regime, the hardness depends on a coupled effect of privacy and novel information-theoretic quantities involving the *Total Variation*. In the low-privacy regime, the lower bounds reduce to the non-private lower bounds. We propose ϵ -local DP and ϵ -global DP variants of a Top Two algorithm, namely CTB-TT and AdaP-TT*, respectively. For ϵ -local DP, CTB-TT is asymptotically optimal by plugging in a private estimator of the means based on Randomised Response. For ϵ -global DP, our private estimator of the mean runs in *arm-dependent adaptive episodes* and adds *Laplace noise* to ensure a good privacy-utility trade-off. By adapting the transportation costs, the expected sample complexity of AdaP-TT* reaches the asymptotic lower bound up to multiplicative constants.

Keywords: differential privacy, multi-armed bandits, best arm identification, fixed confidence, top two algorithm

1 Introduction

We study the stochastic multi-armed *bandit* problem (Lattimore and Szepesvári, 2020), which allows us to reflect on fundamental information-utility trade-offs involved in interactive sequential learning. Specifically, in a bandit problem, a learning *agent* is exposed to interact with K unknown probability distributions $\{\nu_1, \dots, \nu_K\}$ with bounded expectations, referred to as the *reward distributions* (or *arms*). $\boldsymbol{\nu} \triangleq \{\nu_1, \dots, \nu_K\}$ is called a *bandit instance*. At every step $n > 0$, the agent chooses to interact with one of the reward

*. This work was done when Aymen Al Marjani was a PhD student at ENS Lyon.

distributions ν_{a_n} for an arm $a_n \in [K]$, and obtains a sample (or *reward*) r_n from it. The goal of the agent can be of two types: (a) maximise the reward accumulated over time, or equivalently to minimise the regret, and (b) to find the reward distribution (or arm) with the highest expected reward. The first problem is called the regret-minimisation problem (Auer et al., 2002), while the second one is called the *Best Arm Identification (BAI)* problem (Kaufmann et al., 2016). In this paper, we focus on the BAI problem, i.e. to compute $a^*(\boldsymbol{\nu}) \triangleq \arg \max_{a \in [K]} \mathbb{E}_{r \sim \nu_a} [r] \triangleq \arg \max_{a \in [K]} \mu_a$.

With its advent in 1950s (Bechhofer, 1954, 1958) and recent resurgence (Mannor and Tsitsiklis, 2004; Gabillon et al., 2012; Jamieson et al., 2014; Kaufmann et al., 2016), BAI has been extensively studied with different structural assumptions: Fixed-confidence (Jamieson and Nowak, 2014); Fixed-budget (Carpentier and Locatelli, 2016); Non-stochastic (Jamieson and Talwalkar, 2016); Best-of-both-worlds (Abbasi-Yadkori et al., 2018); Linear (Soare et al., 2014). In this paper, we specifically investigate the *Fixed Confidence BAI problem*, in brief FC-BAI, that yields a δ -correct recommendation $\hat{a} \in [K]$, i.e. the probability that the algorithm stops and returns $\hat{a} \neq a^*(\boldsymbol{\nu})$ is upper bounded by δ . FC-BAI is increasingly deployed for different applications, such as clinical trials (Aziz et al., 2021), hyper-parameter tuning (Li et al., 2017), communication networks (Lindståhl et al., 2022), online advertisement (Chen et al., 2014), crowd-sourcing (Zhou et al., 2014), user studies (Losada et al., 2022), and pandemic mitigation (Libin et al., 2019) to name a few. All of these applications often involve the sensitive and personal data of users, which raises serious data privacy concerns (Tucker et al., 2016), as illustrated in Example 1.

Example 1 (Adaptive dose finding trial) *In a dose-finding trial, one physician decides K possible dose levels of a medicine based on preliminary studies— $K \in \{3, \dots, 10\}$ in practice (Aziz et al., 2021). At each step n , a patient is chosen from a local pool of volunteers and a dose level $a_n \in [K]$ is applied to the patient. Following that, the effectiveness of the dose on the patient, i.e. $r_n \in \mathbb{R}$ is observed. The goal of the physician is to recommend after the trial, which dose level is most effective on average, i.e. the dose level a^* that maximises the expected reward. Here, every application of a dose level and the patient’s reaction to it exposes information regarding the medical conditions of the patient. Additionally, at each step n of an adaptive sequential trial, the physician can use an FC-BAI algorithm that observes the previous history of dose levels $\{a_t\}_{t < n}$ and their effectiveness $\{r_t\}_{t < n}$ to decide on the next dose level a_n to test. When releasing the experimental findings of the trial to health authorities, the physician should thoroughly detail the experimental protocol. This includes the dose allocated to each patient $\{a_t\}_{t \leq n}$ and the final recommended dose level a^* . Thus, even if the sequence of reactions to doses $\{r_t\}_{t \leq n}$ is kept secret, publishing the sequence of chosen dose levels $\{a_t\}_{t \leq n}$ and the final recommended dose level a^* computed using the history can leak information regarding patients involved in the trial.*

This example demonstrates the need for privacy in best-arm identification. In this paper, we investigate *privacy-utility trade-offs for a privacy-preserving algorithm in FC-BAI*. Specifically, we use the celebrated Differential Privacy (DP) (Dwork and Roth, 2014) as the framework to preserve data privacy. DP ensures that an algorithm’s output is unaffected by changes in input by a single data point. By limiting the amount of sensitive information that an adversary can deduce from the output, DP renders an individual corresponding

to a data point ‘*indistinguishable*’. Popular ways to achieve DP include Randomised Response (Warner, 1965) or injecting a calibrated amount of noise, from a Laplace (Dwork and Roth, 2014) or Gaussian distribution (Dong et al., 2022), into the algorithm. The scale of the noise is set to be proportional to the algorithm’s sensitivity and inversely proportional to the privacy budget ϵ . Specifically, we study ϵ -*local DP*, where users do not trust the data curator, and ϵ -*global DP*, where users trust the centralised decision-maker with access to the raw sensitive rewards. For example, in an adaptive dose-finding trial, the patients could trust the physician conducting the trial. In that case, at any time n , she has access to all the true history $\{a_t, r_t\}_{t < n}$, and it is her duty to design an algorithm such that *publishing* $\{a_t\}_{t \leq n}$ and the recommended optimal dose a^* obeys ϵ -*global DP* given the sensitive input, i.e. the effectiveness of the dose levels on the patients $\{r_t\}_{t \leq n}$. Without this trust from the user, she has only access to a perturbed history $\{a_t, \tilde{r}_t\}_{t < n}$, where \tilde{r}_t is a perturbed observation of the true observation r_t which ensures ϵ -*local DP*. We define the notions of ϵ -*local DP* and ϵ -*global DP* for BAI rigorously in Section 2.

For different settings of bandits, the costs of ϵ -*local DP* or ϵ -*global DP* and optimal algorithm design techniques are widely studied in the regret-minimisation problem (Mishra and Thakurta, 2015; Tossou and Dimitrakakis, 2016; Sajed and Sheffet, 2019; Shariff and Sheffet, 2018; Neel and Roth, 2018; Basu et al., 2019; Azize and Basu, 2022, 2024). Recently, a problem-dependent lower bound on regret of stochastic multi-armed bandits with ϵ -*global DP* and an algorithm matching the regret lower bound is proposed by Azize and Basu (2022). In contrast, DP is meagerly studied in the FC-BAI problem of bandits (Sajed and Sheffet, 2019; Kalogerias et al., 2021). Though *efficient* algorithm design in FC-BAI literature is traditionally propelled by deriving tight lower bounds, we do not have any explicit sample complexity lower bound for FC-BAI satisfying ϵ -*local DP* or ϵ -*global DP*. By “efficient” algorithm, we refer to the FC-BAI algorithms that aim to minimise the expected number of samples required (i.e. *sample complexity*) to find a δ -correct recommendation. Presently, we know neither the minimal cost in terms of sample complexity for ensuring DP in FC-BAI, nor the feasibility of efficient algorithm design to achieve the minimal cost.

1.1 Contributions

Motivated by this gap in the literature, this paper answers the following two questions:

- A. *How many additional samples a BAI strategy must need to ensure ϵ -local DP?*
- B. *How many additional samples a BAI strategy must need to ensure ϵ -global DP?*

1.1.1 LOWER BOUNDS

First, we derive a lower bound on the expected sample complexity of any δ -correct FC-BAI algorithm to ensure either ϵ -*local DP* (Theorem 9 and Corollary 10) or ϵ -*global DP* (Theorem 13 and Corollary 14). Due to the δ -correctness and the DP constraints, each of the lower bounds corresponds to the minimum of two characteristic times. The first one is the KL characteristic time $T_{\text{KL}}^*(\nu)$ of the non-private FC-BAI (Kaufmann et al., 2016) (Lemma 4). The second one depends on the privacy ϵ and novel information-theoretic quantities depending on the Total Variation (TV) distance: the TV^2 characteristic time $T_{\text{TV}^2}^*(\nu)$ for ϵ -*local DP* and the TV characteristic time $T_{\text{TV}}^*(\nu)$ for ϵ -*global DP*. As for ϵ -*global DP* regret minimisation (Azize and Basu, 2022), the lower bound indicates that

Algorithm	Trust model	Mean estimator	Transportation cost	Stopping threshold
TTUCB	None	MLE (Alg. 2)	W^G as in (3)	c^G as in (4)
CTB-TT	ϵ -local DP	CTB (Alg. 4)	W^G as in (3)	c^G as in (4)
AdaP-TT	ϵ -global DP	DAF (Alg. 5)	W^G as in (3)	$c^{G,\epsilon}$ as in (11)
AdaP-TT*	ϵ -global DP	DAF (Alg. 5)	$W^{G,\epsilon}$ as in (12)	$\tilde{c}^{G,\epsilon}$ as in (14)

Table 1: Instances of the TTUCB meta-algorithm.

there are two regimes of hardness depending on ϵ and the aforementioned characteristic times. For lower levels of privacy (i.e. higher ϵ), the expected sample complexity matches the non-private FC-BAI. However, for higher levels of privacy (i.e. lower ϵ), the expected sample complexity depends both on the privacy budget ϵ and the TV^2 or TV characteristic time. To derive the lower bound for ϵ -global DP, we provide an ϵ -global DP version of the “change-of-measure” lemma (Kaufmann and Kalyanakrishnan, 2013) (Lemma 12), which we prove using a sequential coupling argument.

1.1.2 ALGORITHM DESIGN

We propose algorithms which are δ -correct, and either ϵ -local DP or ϵ -global DP. While most existing asymptotically optimal FC-BAI algorithms can be modified to tackle DP, we consider the class of Top Two algorithms (Russo, 2016) due to their good empirical performances, low computational cost, and easy implementation. As a case study, we consider the TTUCB meta-algorithm based on the work of (Jourdan and Degenne, 2024). Table 1 summarises the different instances that we propose. We highlight that our private wrappers could be used for other FC-BAI algorithms.

A. For ϵ -local DP, we propose the CTB-TT algorithm. CTB-TT plugs in the CTB(ϵ) estimator of the means, which is ϵ -local DP (Ren et al., 2020, Lemma 11), into the TTUCB algorithm for σ -sub-Gaussian distributions, which is δ -correct.

B. For ϵ -global DP, we propose the DAF(ϵ) estimator of the means, which is ϵ -global DP (Lemma 15). It relies on three ingredients: *adaptive episodes with doubling per arm*, *forgetting*, and *adding calibrated Laplacian noise*. Using the DAF(ϵ) estimator in the TTUCB meta-algorithm, we propose the AdaP-TT and the AdaP-TT* algorithms. As a plug-in approach, AdaP-TT uses the non-private transportation costs both in the TC challenger and in the GLR stopping rule, which is shown to be δ -correct by adding a privacy term to the stopping threshold (Lemma 16). As a lower bound based approach, AdaP-TT* adapts the transportation costs to account for ϵ -global privacy both in the TC challenger and in the GLR stopping rule, which is shown to be δ -correct by modifying the privacy term in the stopping threshold (Lemma 18).

1.1.3 UPPER BOUNDS

We show that the proposed algorithms exhibit upper bounds that match the lower bounds up to multiplicative constants. We highlight that our generic asymptotic analysis can be applied to any Top Two algorithms, since it builds on the one of Jourdan et al. (2022).

A. As CTB-TT is equivalent to running TTUCB on a modified Bernoulli instance ν_ϵ , it recovers the asymptotic and non-asymptotic upper bounds on the expected sample complexity derived in Jourdan and Degenne (2024). The asymptotic upper bound matches the asymptotic lower bound up to a constant multiplicative term, 2 when $\epsilon \rightarrow +\infty$ and 4 when $\epsilon \rightarrow 0$. Our experiments confirm the good performance of CTB-TT, and the existence of two hardness regimes for ϵ -local DP (Section 5.1).

B. Using the DAF(ϵ) estimator yields a batched algorithm with adaptive and data-dependent changes of episodes. While our analysis is inspired by the one of Jourdan et al. (2022), studying AdaP-TT and AdaP-TT* requires carefully quantifying the effects of doubling, forgetting, and adding noise. We derive an asymptotic upper bound on the expected sample complexity of AdaP-TT (Theorem 17) and AdaP-TT* (Theorem 19). In the non-private regime of $\epsilon \rightarrow +\infty$, both algorithms recover the asymptotic lower bound up to multiplicative constants (16 and 8 respectively), with solely $\mathcal{O}(K \log_2(T_{\text{KL}}^*(\nu) \log(1/\delta)))$ rounds of adaptivity. When $\epsilon \rightarrow 0$, AdaP-TT* achieves the asymptotic lower bound up to a multiplicative constant 48, while AdaP-TT only recovers it for instances where the mean gaps have the same order of magnitude. Our experiments show the good performance of our algorithms compared to DP-SE (Sajed and Sheffet, 2019), which can be adapted for FC-BAI (see Section 4.4 for a detailed comparison). They confirm the existence of two hardness regimes for ϵ -global DP, as well as the empirical superiority of AdaP-TT* over AdaP-TT when $\epsilon \rightarrow 0$ (Section 5.2).

1.2 Outline

After presenting Differential Privacy and Best-Arm Identification in the Fixed-Confidence Setting in Section 2, we formulate the problem of private best-arm identification. We present lower bounds and matching upper bounds for ϵ -local DP FC-BAI (Section 3) and ϵ -global DP FC-BAI (Section 4). Our algorithms are studied empirically in Section 5.

2 Differential Privacy and Best-Arm Identification

In this section, we provide relevant background information on Differential Privacy (DP) in Section 2.1, and Best-Arm Identification in the Fixed-Confidence Setting (FC-BAI) in Section 2.2. Then, we formulate the problem of private best-arm identification (FC-BAI with DP) in Section 2.3, under both the local and global trust models.

2.1 Background: Differential Privacy

Differential Privacy (DP) ensures the protection of an individual’s sensitive information when her data is used for analysis. A randomised algorithm satisfies DP if the output of the algorithm stays almost the same, regardless of whether any single individual’s data is included in or excluded from the input. One way of achieving DP is by adding controlled noise to the algorithm’s output.

Definition 1 ((ϵ, δ)-DP (Dwork and Roth, 2014)) *A randomised algorithm \mathcal{A} satisfies (ϵ, δ)-Differential Privacy (DP) if for any two neighbouring datasets \mathcal{D} and \mathcal{D}' that*

differ only in one entry, i.e. $d_{Ham}(\mathcal{D}, \mathcal{D}') = 1$, and for all sets of output $\mathcal{O} \subseteq \text{Range}(\mathcal{A})$,

$$\Pr[\mathcal{A}(\mathcal{D}) \in \mathcal{O}] \leq e^\epsilon \Pr[\mathcal{A}(\mathcal{D}') \in \mathcal{O}] + \delta,$$

where the probability space is over the coin flips of the mechanism \mathcal{A} , and $(\epsilon, \delta) \in \mathbb{R}^{\geq 0} \times \mathbb{R}^{\geq 0}$. If $\delta = 0$, we say that \mathcal{A} satisfies ϵ -DP. A lower privacy budget ϵ implies higher privacy.

The Laplace mechanism (Dwork et al., 2010a; Dwork and Roth, 2014) ensures ϵ -DP by injecting controlled random noise into the output of the algorithm, which is sampled from a calibrated Laplace distribution (as specified in Theorem 2). We use $Lap(b)$ to denote the Laplace distribution with mean 0 and variance $2b^2$.

Theorem 2 (Laplace mechanism, Theorem 3.6 (Dwork and Roth, 2014)) *Let $f : \mathcal{X} \rightarrow \mathbb{R}^d$ be an algorithm with sensitivity $s(f) \triangleq \max_{\mathcal{D}, \mathcal{D}' \text{ s.t. } |\mathcal{D} - \mathcal{D}'|_{\text{Hamming}} = 1} \|f(\mathcal{D}) - f(\mathcal{D}')\|_1$, where $\|\cdot\|_1$ is the L_1 norm. If samples $\{N_i\}_{i=1}^d$ are generated independently from $Lap\left(\frac{s(f)}{\epsilon}\right)$, then the output injected with the noise, i.e. $f(\mathcal{D}) + [N_1, \dots, N_d]$, satisfies ϵ -DP.*

We also study the setting of *local differential privacy*, where users do not trust the data curator, i.e. the entity collecting the data. Local DP is one of the oldest formulations of privacy, dating back to Warner (1965), who advocated it as a solution to what he called “evasive answer bias” in survey sampling.

Definition 3 (ϵ -local DP (Duchi et al., 2013)) *A randomised algorithm \mathcal{M} satisfies ϵ -local DP if for any pair of input values $x, x' \in \mathcal{D}$, and for all sets of output $\mathcal{O} \subseteq \text{Range}(\mathcal{M})$,*

$$\Pr[\mathcal{M}(x) \in \mathcal{O}] \leq e^\epsilon \Pr[\mathcal{M}(x') \in \mathcal{O}],$$

where the probability space is over the coin flips of the mechanism \mathcal{M} , and for some $\epsilon \in \mathbb{R}^{\geq 0}$. The perturbation mechanism \mathcal{M} is applied to each user record independently.

For binary attributes, the Randomised Response (RR) mechanism (Warner, 1965) is a popular way to achieve ϵ -local DP. The idea is to output the true value of a user’s response with probability $e^\epsilon/(e^\epsilon + 1)$ and output the opposite value with probability $1/(e^\epsilon + 1)$. To make it suitable for larger discrete domains, a Generalised Randomised Response (GRR) is proposed in Kairouz et al. (2016). For continuous numerical data statistics, adding Laplace noise to each data record achieves local DP as well.

2.2 Background: Best Arm Identification in the Fixed-Confidence Setting

In this section, we first present the Best-arm identification (BAI) problem, a BAI strategy and δ -correctness. Then, we present a lower bound on the sample complexity of any δ -correct BAI strategy. Finally, we discuss algorithms in the BAI literature which match the sample complexity lower bound. We focus on the Top Two family of algorithms since they enjoy both theoretical optimality and good empirical performance.

2.2.1 THE BEST ARM IDENTIFICATION PROBLEM

Best-arm identification (BAI) is a pure exploration problem that aims to identify the optimal arm. It has been studied in two major theoretical frameworks (Audibert et al., 2010; Gabillon et al., 2012; Jamieson and Nowak, 2014; Garivier and Kaufmann, 2016): the fixed-confidence and fixed-budget setting. In the fixed-budget setting, the objective is to minimise the probability of misidentifying a correct answer with a fixed number of samples T . We consider the fixed-confidence setting (FC-BAI), in which the learner aims at minimising the number of samples used to identify a correct answer with confidence $1 - \delta \in (0, 1)$.¹ To achieve this, the learner defines an FC-BAI strategy to interact with the bandit instance $\boldsymbol{\nu} = \{\nu_a\}_{a \in [K]} \in \mathcal{F}^K$, consisting of K arms with finite means $\{\mu_a\}_{a \in [K]} \in (0, 1)^K$. We assume that there is a unique best arm $a^*(\boldsymbol{\nu})$ defined as $a^*(\boldsymbol{\nu}) = \arg \max_{a \in [K]} \mu_a$. The set of distributions \mathcal{F} will depend on the considered result, e.g. Bernoulli distributions, bounded distributions on $[0, 1]$ or σ -sub-Gaussian distributions. A distribution κ is σ -sub-Gaussian if it satisfies $\mathbb{E}_{X \sim \kappa}[e^{\lambda(X - \mathbb{E}_{X \sim \kappa}[X])}] \leq e^{\sigma^2 \lambda^2 / 2}$ for all $\lambda \in \mathbb{R}$.

We denote the action played at step n by a_n , and the corresponding observed reward by $r_n \sim \nu_{a_n}$. The σ -algebra $\mathcal{H}_n = \sigma(a_1, r_1, \dots, a_n, r_n)$ is the history of actions played and rewards collected at the end of time n . We augment the action set by a *stopping action* \top , and write $a_n = \top$ to denote that the algorithm has stopped before step n . A FC-BAI strategy π is composed of

i. A pair of sampling and stopping rules $(S_n : \mathcal{H}_{n-1} \rightarrow \mathcal{P}([1, K] \cup \{\top\}))_{n \geq 1}$. For an action $a \in [K]$, $S_n(a \mid \mathcal{H}_{n-1})$ denotes the probability of playing action a given history \mathcal{H}_{n-1} . On the other hand, $S_n(\top \mid \mathcal{H}_{n-1})$ is the probability of the algorithm halting given \mathcal{H}_{n-1} . For any history \mathcal{H}_{n-1} , a consistent sampling and stopping rule S_n satisfies $S_n(\top \mid \mathcal{H}_{n-1}) = 1$ if \top has been played before n .

ii. A recommendation rule $(\text{Rec}_n : \mathcal{H}_{n-1} \rightarrow \mathcal{P}([1, K]))_{n > 1}$. A recommendation rule dictates $\text{Rec}_n(a \mid \mathcal{H}_{n-1})$, i.e. the probability of returning action a as a guess for the best action given \mathcal{H}_{n-1} .

We denote by τ_δ the **stopping time** (or **sample complexity**) of the algorithm, i.e. the first step n demonstrating $a_n = \top$. A FC-BAI strategy π is called **δ -correct** for a class of bandit instances $\mathcal{M} \subseteq \mathcal{F}^K$, if for every instance $\boldsymbol{\nu} \in \mathcal{M}$, π recommends \hat{a} as the optimal action $a^*(\boldsymbol{\nu})$ with probability at least $1 - \delta$, i.e. $\mathbb{P}_\nu(\tau_\delta < +\infty, \hat{a} = a^*(\boldsymbol{\nu})) \geq 1 - \delta$.

2.2.2 LOWER BOUND ON THE EXPECTED SAMPLE COMPLEXITY

Being δ -correct imposes a lower bound on the expected sample complexity on any instance.

Lemma 4 (Garivier and Kaufmann (2016)) *Let $\delta \in (0, 1)$. For all δ -correct FC-BAI strategy and all instances $\boldsymbol{\nu} \in \mathcal{M}$, we have that $\mathbb{E}_\nu[\tau_\delta] \geq T_{\text{KL}}^*(\boldsymbol{\nu}) \log(1/(2.4\delta))$ with*

$$T_d^*(\boldsymbol{\nu})^{-1} \triangleq \sup_{\omega \in \Sigma_K} \inf_{\boldsymbol{\lambda} \in \text{Alt}(\boldsymbol{\nu})} \sum_{a \in [K]} \omega_a \mathbf{d}(\nu_a, \lambda_a), \quad (1)$$

where the probability simplex is denoted by $\Sigma_K \triangleq \{\omega \in [0, 1]^K \mid \sum_{a=1}^K \omega_a = 1\}$ and the set of alternative instances is $\text{Alt}(\boldsymbol{\nu}) \triangleq \{\boldsymbol{\lambda} \in \mathcal{M} \mid a^*(\boldsymbol{\lambda}) \neq a^*(\boldsymbol{\nu})\}$, i.e. the bandit instances with

1. We remind not to confuse risk level δ with the δ of (ϵ, δ) -DP. Hereafter, we consider ϵ -global DP as the privacy definition, and δ always represents the risk (or probability of mistake) of the BAI strategy.

Algorithm 1 TTUCB Meta-algorithm

Input: $(\beta, \delta) \in (0, 1)^2$, confidence bonuses $(b_a)_{a \in [K]}$, transportation costs $(W_{a,b})_{(a,b) \in [K]^2}$, estimator mechanisms $(\text{ESTIMATOR}_a)_{a \in [K]}$ and stopping conditions $(\text{STOP}_{a,b})_{(a,b) \in [K]^2}$.
 Observe $r_a \sim \nu_a$ for all $a \in [K]$, initialise $\tilde{\mu}_{n,a}$ and $N_{n,a} = \tilde{N}_{n,a} = 1$ where $n = K + 1$;
for $n > K$ **do**
 Set $\hat{a}_n = \arg \max_{a \in [K]} \tilde{\mu}_{n,a}$;
 If $\text{STOP}_{\hat{a}_n, a}(\tilde{\mu}_n, \tilde{N}_n, \delta) \geq 0$ for all $a \neq \hat{a}_n$ **then** set $a_n = \top$ and **return** \hat{a}_n , **otherwise**;
 Set $B_n = \arg \max_{a \in [K]} \left\{ \tilde{\mu}_{n,a} + b_a(\tilde{N}_n) \right\}$ and $C_n = \arg \min_{a \neq B_n} W_{B_n, a}(\tilde{\mu}_n, N_n)$;
 Store $r_n \sim \nu_{a_n}$ by pulling arm $a_n = B_n$ if $N_{n, B_n}^{B_n} \leq \beta L_{n+1, B_n}$, else $a_n = C_n$;
 Set $N_{n+1, a_n} \leftarrow N_{n, a_n} + 1$, $N_{n+1, a_n}^{B_n} \leftarrow N_{n, a_n}^{B_n} + 1$, $L_{n+1, B_n} \leftarrow L_{n, B_n} + 1$ and $n \leftarrow n + 1$;
 Get $(\tilde{\mu}_{n+1, a_n}, \tilde{N}_{n+1, a_n}) = \text{ESTIMATOR}_{a_n}(\mathcal{H}_n)$;
end for

a different optimal arm than ν . For two probability distributions \mathbb{P}, \mathbb{Q} on (Ω, \mathcal{F}) , the KL divergence is $\text{KL}(\mathbb{P} \parallel \mathbb{Q}) \triangleq \int \log \left(\frac{d\mathbb{P}}{d\mathbb{Q}}(\omega) \right) d\mathbb{P}(\omega)$, when $\mathbb{P} \ll \mathbb{Q}$, and $+\infty$ otherwise.

Early FC-BAI algorithms failed to reach the lower bound of Theorem 4, e.g. Successive Elimination (SE) based algorithms (Even-Dar et al., 2006) or confidence bounds based algorithms, e.g. LUCB (Kalyanakrishnan et al., 2012) or lil'UCB (Jamieson et al., 2014). Inspired by this lower bound, many algorithms have been designed to tackle FC-BAI. The Track-and-Stop algorithm (Garivier and Kaufmann, 2016) is the first algorithm to reach asymptotic optimality, by sequentially solving the optimisation problem $T_{\text{KL}}^*(\nu_n)$ and tracking the associated optimal weights. To reduce the computational cost of Track-and-Stop, several asymptotically optimal algorithms have been proposed recently: online optimisation-based approach, e.g. game-based algorithm (Degenne et al., 2019) or FWS (Wang et al., 2021), and Top Two algorithms (Russo, 2016). While most algorithms can be modified to tackle ϵ -DP, we consider the Top Two algorithms due to their great empirical performance and easy implementation. At every step, a Top Two sampling rule selects the next arm to sample from among two candidate arms, a leader and a challenger. In recent years, numerous variants of Top Two algorithms have been analysed and shown to be asymptotically optimal (Russo, 2016; Qin et al., 2017; Shang et al., 2020; Jourdan et al., 2022; You et al., 2023b; Jourdan et al., 2024). In particular, we consider one particular case study, i.e. the TTUCB algorithm (Jourdan and Degenne, 2024), but our approach can be directly adapted to any other Top Two algorithms.

Remark 5 *The private estimators of the means and the private stopping rules presented in Sections 3 and 4 could be used with most existing existing FC-BAI algorithm. The resulting algorithms will be private, δ -correct and have near-optimal asymptotic sample complexity. In this work, we consider and rigorously analyse the Top Two algorithms since they are simple algorithms enjoying both strong theoretical guarantees and empirical performance.*

Algorithm 2 Maximum Likelihood Estimator (MLE)

Input: History \mathcal{H}_n , arm $a \in [K]$.

Return $(\hat{\mu}_{n,a}, N_{n,a})$ where $\hat{\mu}_{n,a} = N_{n,a}^{-1} \sum_{t \in [n-1]} r_t \mathbb{1}\{a_t = a\}$;

2.2.3 THE TTUCB META-ALGORITHM

Since we will propose several private algorithm building on top of TTUCB, we propose a TTUCB meta-algorithm (Algorithm 1). To instantiate it, one should specify: a parameter $\beta \in (0, 1)$ (e.g. $\beta = 1/2$), confidence bonuses $(b_a)_{a \in [K]}$ where $b_a : \mathbb{N}^K \rightarrow \mathbb{R}^+$, transportation costs $(W_{a,b})_{(a,b) \in [K]^2}$ where $W_{a,b} : \mathbb{N}^K \times \mathbb{N}^K \rightarrow \mathbb{R}^+$, estimator mechanisms $(\text{ESTIMATOR}_a)_{a \in [K]}$ computing data-dependent estimators $(\mu_{n,a})_{(n,a) \in \mathbb{N} \times [K]}$ based on local counts $(\tilde{N}_{n,a})_{(n,a) \in \mathbb{N} \times [K]}$, and GLR stopping conditions

$$\forall (a, b) \in [K]^2, \quad \text{STOP}_{a,b}(\tilde{\mu}, \omega, \delta) = W_{a,b}(\tilde{\mu}, \omega) - c_{a,b}(\omega, \delta), \quad (2)$$

where $(c_{a,b})_{(a,b) \in [K]^2}$ where $c_{a,b} : \mathbb{N}^K \times (0, 1) \rightarrow \mathbb{R}^+$ are stopping thresholds.

For σ -sub-Gaussian distributions, TTUCB in Jourdan and Degenne (2024) is an instance of Algorithm 1 using the MLE (Algorithm 2) and

$$W_{a,b}^G(\tilde{\mu}, \omega) = \frac{(\tilde{\mu}_a - \tilde{\mu}_b)_+^2}{2\sigma^2(1/\omega_a + 1/\omega_b)} \quad \text{and} \quad b_a^G(\omega) = \sqrt{\frac{2\sigma^2\alpha(1+s)\log\|\omega\|_1}{\omega_a}} \quad \text{with } s, \alpha > 1. \quad (3)$$

In practice, they take $s = \alpha = 1.2$. The generalised likelihood ratio (GLR) stopping rule has to ensure δ -correctness. This is done by choosing the stopping threshold as

$$c_{a,b}^G(\omega, \delta) = 2\mathcal{C}_G(\log((K-1)/\delta)/2) + 2\log(4 + \log\omega_a) + 2\log(4 + \log\omega_b), \quad (4)$$

where the function \mathcal{C}_G is defined in (20). It satisfies $\mathcal{C}_G(x) \approx x + \log(x)$. For bounded distributions on $[0, 1]$ such as Bernoulli, we take $\sigma = 1/2$.

At each step, a Top Two algorithm selects two arms called leader and challenger, and samples one arm among them. TTUCB uses a UCB-based leader and a Transportation Cost (TC) challenger. The theoretical motivation behind the TC challenger comes from the theoretical lower bound in FC-BAI (Lemma 4), which involves the KL-characteristic time $T_{\text{KL}}^*(\boldsymbol{\mu}) = \min_{\beta \in (0,1)} T_{\text{KL},\beta}^*(\boldsymbol{\mu})$. For Gaussian distributions $\nu_a = \mathcal{N}(\mu_a, \sigma^2)$, it writes as

$$T_{\text{KL},\beta}^*(\boldsymbol{\nu})^{-1} = \max_{\omega \in \Sigma_K, \omega_{a^*} = \beta} \frac{(\mu_{a^*} - \mu_a)^2}{2\sigma^2(1/\beta + 1/\omega_a)} \quad \text{and} \quad T_{\text{KL},1/2}^*(\boldsymbol{\nu}) \leq 2T_{\text{KL}}^*(\boldsymbol{\nu}). \quad (5)$$

Note that $H(\boldsymbol{\nu}) \leq T_{\text{KL}}^*(\boldsymbol{\nu}) \leq 2H(\boldsymbol{\nu})$ where $H(\boldsymbol{\nu}) = 2\sigma^2 \sum_{a \in [K]} \Delta_a^{-2}$ with $\Delta_a = \mu_{a^*} - \mu_a$ for all $a \neq a^*$ and $\Delta_{a^*} = \Delta_{\min} = \min_{a \neq a^*} (\mu_{a^*} - \mu_a)$ for all $a \neq a^*$. The maximiser of (5) is denoted by $\omega_{\text{KL},\beta}^*(\boldsymbol{\mu})$, and is further referred to as the β -optimal allocation as it is unique. Let $N_{n,b}^a$ denote the number of times arm b was pulled when a was the leader, and $L_{n,a}$ denotes the number of times arm a was the leader. In order to select the next arm to sample I_n , TTUCB relies on K tracking procedures, i.e. set $I_n = B_n$ if $N_{n,B_n}^{B_n} \leq \beta L_{n+1,B_n}$, else $I_n = C_n$. This ensures that $\max_{a \in [K], n > K} |N_{n,a}^a - \beta L_{n,a}| \leq 1$ (Degenne et al., 2020).

2.3 Problem Statement: FC-BAI with DP

Now, we formally extend DP to BAI. We consider two trust models: (1) ϵ -local DP BAI, where each user sends her reward to the BAI strategy, using an ϵ -local DP perturbation mechanism, and (2) ϵ -global DP BAI, where the BAI strategy, a.k.a. the centralised decision maker, is trusted with all the intermediate rewards. We summarise the BAI strategy-Users interaction in Algorithm 3, under global DP and local DP.

2.3.1 LOCAL DP FC-BAI

We represent each user u_t by the vector $\mathbf{x}_t \triangleq (x_{t,1}, \dots, x_{t,K}) \in \mathbb{R}^K$, where $x_{t,a}$ represents the **potential** reward observed, if action a was recommended to user u_t . Due to the bandit feedback, only $r_t = x_{t,a_t} \sim \nu_{a_t}$ is observed at step t . The user observes the real reward $r_t = \mathbf{x}_{t,a_t}$ but only sends a noisy version z_t to the BAI strategy, by sampling z_t from the perturbation mechanism, i.e. $z_t \sim \mathcal{M}(r_t)$. The BAI strategy only has access to the noisy rewards (z_t) to make its decisions.

Definition 6 (ϵ -local DP for BAI) *A pair (\mathcal{M}, π) of perturbation mechanism and BAI strategy satisfies ϵ -local DP, if they satisfy*

- (a) *The perturbation mechanism \mathcal{M} is ϵ -local DP with respect to each reward record, i.e. for all T , all rewards r_t, r'_t and all noisy outputs z_t , $\Pr[\mathcal{M}(r_t) = z_t] \leq e^\epsilon \Pr[\mathcal{M}(r'_t) = z_t]$.*
- (b) *The BAI strategy only has access to the noisy rewards $z_t \sim \mathcal{M}(r_t)$ to make its decisions.*

For a pair (\mathcal{M}, π) to be δ -correct with respect to an environment ν , under a local DP interaction protocol, the pair should verify: (a) the perturbation mechanism \mathcal{M} should not change the identity of the optimal arm, i.e. $a^*(\nu) = a^*(\nu^{\mathcal{M}})$ and (b) the BAI strategy π should be δ -correct for the noisy environment $\nu^{\mathcal{M}}$. The goal in ϵ -local DP FC-BAI is to design a δ -correct ϵ -local DP pair (\mathcal{M}, π) of perturbation mechanism and BAI strategy, with $\mathbb{E}[\tau_\delta]$ as small as possible.

2.3.2 GLOBAL DP BAI

Again, we represent each user u_t by the vector $\mathbf{x}_t \triangleq (x_{t,1}, \dots, x_{t,K}) \in \mathbb{R}^K$, where $x_{t,a}$ represents the **potential** reward observed, if action a was recommended to user u_t . Due to the bandit feedback, only $r_t = x_{t,a_t} \sim \nu_{a_t}$ is observed at step t . We use an underline to denote any sequence. Thus, we denote the sequence of sampled actions until T as $\underline{a}^T = (a_1, \dots, a_T)$. We further represent a set of users $\{u_t\}_{t=1}^T$ until T by **the table of potential rewards** $\underline{\mathbf{d}}^T \triangleq \{\mathbf{x}_1, \dots, \mathbf{x}_T\} \in (\mathbb{R}^K)^T$. First, we observe that $\underline{\mathbf{d}}^T$ is the sensitive input dataset to be made private, and $(\underline{a}^T, \hat{a}, T)$ is the output of the BAI strategy. Hence, we define the probability that the BAI strategy π samples the action sequence \underline{a}^T , recommends the action \hat{a} , and halts at time T , as

$$\pi(\underline{a}^T, \hat{a}, T \mid \underline{\mathbf{d}}^T) \triangleq \text{Rec}_{T+1}(\hat{a} \mid \mathcal{H}_T) S_{T+1}(\top \mid \mathcal{H}_T) \prod_{t \in [T]} S_t(a_t \mid \mathcal{H}_{t-1}), \quad (6)$$

where T users under interaction are represented by the table of potential rewards $\underline{\mathbf{d}}^T$. A BAI strategy satisfies ϵ -global DP if the probability in Eq. (6) is similar when the BAI strategy interacts with two neighbouring tables of rewards differing by one user (i.e. a row

Algorithm 3 Sequential Interaction Between a BAI Strategy and Users

Input: A BAI strategy π , Users $\{u_t\}_{n \geq 1}$ represented by the table $\underline{\mathbf{d}}$ and a perturbation mechanism \mathcal{M}

Output: A stopping time τ , a sequence of samples actions $\underline{a}^\tau = (a_1, \dots, a_\tau)$ and a recommendation \hat{a} satisfying ϵ -DP

for $t = 1, \dots$ **do**

π recommends action $a_t \sim S_t(\cdot \mid a_1, z_1, \dots, a_{t-1}, z_{t-1})$

if $a_t = \top$ **then**

Halt. Return $\tau = t$ and $\hat{a} \sim \text{Rec}_t(\cdot \mid a_1, z_1, \dots, a_{t-1}, z_{t-1})$

else

if Global DP **then**

u_t observes the **sensitive** reward $r_t \triangleq \underline{\mathbf{d}}_{t,a_t}$

u_t sends the **sensitive** reward $z_t \triangleq r_t$ to π

else Local DP

u_t observes the **sensitive** reward $r_t \triangleq \underline{\mathbf{d}}_{t,a_t}$

u_t sends the **noisy** reward $z_t \sim \mathcal{M}(r_t)$ to π

end if

end if

end for

in $\underline{\mathbf{d}}^T$). Definition 7 can be seen as a BAI counterpart of the ϵ -global DP definition proposed in Azize and Basu (2022) for regret minimisation.

Definition 7 (ϵ -global DP for BAI) *A BAI strategy satisfies ϵ -global DP, if for all $T \geq 1$, all neighbouring table of rewards $\underline{\mathbf{d}}^T$ and $\underline{\mathbf{d}}'^T$, i.e. $d_{\text{Ham}}(\underline{\mathbf{d}}^T, \underline{\mathbf{d}}'^T) = 1$, all sequences of sampled actions $\underline{a}^T \in [K]^T$ and recommended actions $\hat{a} \in [K]$ we have that*

$$\pi(\underline{a}^T, \hat{a}, T \mid \underline{\mathbf{d}}^T) \leq e^\epsilon \pi(\underline{a}^T, \hat{a}, T \mid \underline{\mathbf{d}}'^T).$$

The goal in ϵ -global DP FC-BAI is to design a δ -correct ϵ -global DP BAI strategy π , with $\mathbb{E}[\tau_\delta]$ as small as possible.

Remark 8 *It is possible to consider that the output of a BAI strategy is only the final recommended action \hat{a} , i.e. not publishing the intermediate actions \underline{a}^T . This gives a weaker definition of privacy compared to Definition 7, since the latter defends against adversaries that may look inside the execution of the BAI strategy, i.e. pan-privacy (Dwork et al., 2010b). Also, Definition 7 is needed in practice. For example, in the case of dose-finding (Example 1), the experimental protocol, i.e. the intermediate actions, needs to be published too.*

3 Local Differentially Private Best-Arm Identification

In this section, we answer the following question: *How many additional samples a BAI strategy must select to ensure ϵ -local DP?* We provide a lower bound on the expected sample complexity of any δ -correct ϵ -local DP pair of perturbation mechanism and BAI strategy. We complement the sample complexity lower bound with a matching upper bound.

3.1 Lower Bound on the Expected Sample Complexity

We derive a lower bound on the expected sample complexity in ϵ -local DP FC-BAI, which features problem-dependent characteristic times as in the FC-BAI setting.

Theorem 9 *Let $\delta \in (0, 1)$ and $\epsilon > 0$. For any δ -correct ϵ -local DP pair (\mathcal{M}, π) of perturbation mechanism and BAI strategy, we have that $\mathbb{E}_{\nu}[\tau_{\delta}] \geq T_{\ell}^*(\nu; \epsilon) \log(1/(2.4\delta))$ with*

$$T_{\ell}^*(\nu; \epsilon)^{-1} \triangleq \sup_{\omega \in \Sigma_K} \inf_{\lambda \in \text{Alt}(\nu)} \sum_{a \in [K]} \omega_a \min \left\{ \text{KL}(\nu_a \parallel \lambda_a), c(\epsilon) (\text{TV}(\nu_a \parallel \lambda_a))^2 \right\},$$

where $c(\epsilon) = \min\{4, e^{2\epsilon}\} (e^{\epsilon} - 1)^2$ is a privacy term. For two probability distributions \mathbb{P}, \mathbb{Q} on the measurable space (Ω, \mathcal{F}) , the TV distance is $\text{TV}(\mathbb{P} \parallel \mathbb{Q}) \triangleq \sup_{A \in \mathcal{F}} \{\mathbb{P}(A) - \mathbb{Q}(A)\}$.

Proof sketch. To prove this theorem, we first use the KL-decomposition of Garivier and Kaufmann (2016) applied on the "noisy" bandit environment. Then, Theorem 1 of Duchi et al. (2013) is applied to relate the KL of rewards in the "noisy" bandit environment to the original environment. In Appendix B, we formally define the bandit canonical model under local DP, and provide a formal proof of the theorem. \blacksquare

Similar to the lower bound for the non-private BAI (Garivier and Kaufmann, 2016), the lower bound of Theorem 9 is the value of a two-player zero-sum game between a MIN player and MAX player. MIN plays an alternative instance λ close to ν in order to confuse MAX. The latter plays an allocation $\omega \in \Sigma_K$ to explore the different arms, to maximise the divergence between ν and the confusing instance λ that MIN played.

Corollary 10 (Relaxing the local DP lower bound) *Let $T_{\ell}^*(\nu; \epsilon)$ as in Theorem 9 and $T_{\text{d}}^*(\nu)$ as in Eq. (1). Then, we have*

$$T_{\ell}^*(\nu; \epsilon) \geq \max \left\{ T_{\text{KL}}^*(\nu), c(\epsilon)^{-1} T_{\text{TV}^2}^*(\nu) \right\} \quad \text{with} \quad c(\epsilon) = \min\{4, e^{2\epsilon}\} (e^{\epsilon} - 1)^2. \quad (7)$$

Let ν_{G} be the Gaussian instance with unit variances and the same means as the Bernoulli instance ν . Then, we have $T_{\text{TV}^2}^*(\nu) \geq 2T_{\text{KL}}^*(\nu)$ and $T_{\text{TV}^2}^*(\nu) = T_{\text{KL}}^*(\nu_{\text{G}})/2$.

Proof The first part is true since $T_{\ell}^*(\nu; \epsilon) \geq T_{\text{KL}}^*(\nu)$ and $T_{\ell}^*(\nu; \epsilon) \geq \frac{T_{\text{TV}^2}^*(\nu)}{\min\{4, e^{2\epsilon}\} (e^{\epsilon} - 1)^2}$. The second part uses that $\text{KL}(\mathcal{N}(p, 1) \parallel \mathcal{N}(q, 1)) = \frac{1}{2}|p - q|^2 = \frac{1}{2}\text{TV}(\text{Ber}(p) \parallel \text{Ber}(q))^2$. \blacksquare

Corollary 10 relates the TV^2 characteristic time for Bernoulli to the KL characteristic times for Bernoulli and Gaussian. The sample complexity of FC-BAI with local DP on Bernoulli instances is reduced to the characteristic time of the non-private FC-BAI on Gaussian instances, up to a multiplicative factor which only depends on ϵ .

Two privacy regimes. The sample complexity lower bound in Eq. (7) suggests the existence of two hardness regimes depending on ϵ , $T_{\text{KL}}^*(\nu)$ and $T_{\text{TV}^2}^*(\nu)$. In the high privacy regime, as $\epsilon \rightarrow 0$, the lower bound reduces to $\epsilon^{-2} T_{\text{TV}^2}^*(\nu)$. In the low privacy regime, as $\epsilon \rightarrow \infty$, the lower bound reduces to the non-private complexity $T_{\text{KL}}^*(\nu)$. The switch between the low and the high privacy regimes happens at the ϵ verifying $\min\{4, e^{2\epsilon}\} (e^{\epsilon} - 1)^2 = \frac{T_{\text{TV}^2}^*(\nu)}{T_{\text{KL}}^*(\nu)}$. For example, for environments where the Pinsker inequality is tight, i.e. $T_{\text{TV}^2}^*(\nu) \approx 2T_{\text{KL}}^*(\nu)$, then the switch happens at $\epsilon \approx 0.582$.

Algorithm 4 Convert-To-Bernoulli(ϵ) Estimator (CTB) (Ren et al., 2020)

Input: History \mathcal{H}_n with past perturbations $(\tilde{r}_t)_{t \in [n-2]}$, arm $a \in [K]$.

Return $(\tilde{\mu}_{n,a}, N_{n,a})$ with $\tilde{\mu}_{n,a} = \frac{1}{N_{n,a}} \sum_{t=1}^{n-1} \tilde{r}_t \mathbb{1}\{a_t = a\}$ and $\tilde{r}_{n-1} \sim \text{Ber}\left(\frac{r_{n-1}(e^\epsilon-1)+1}{e^\epsilon+1}\right)$;

3.2 A Plug-In Approach: the CTB-TT Algorithm

Ren et al. (2020) proposed the Convert-To-Bernoulli (CTB, Algorithm 4) estimator of the means, which relies on the Randomised Response mechanism to ensure ϵ -local DP on $[0, 1]$.

Lemma 11 (Ren et al. 2020) *CTB(ϵ) ensures ϵ -local DP on $[0, 1]$. For all n , we have $\tilde{r}_n \mid a_n \sim \mathcal{B}(\mu_{\epsilon,a_n})$ where $\mu_{\epsilon,a} = (2\mu_a - 1) \frac{e^\epsilon - 1}{2(e^\epsilon + 1)} + 1/2$.*

CTB-TT algorithm. To solve ϵ -local DP FC-BAI, we propose the CTB-TT algorithm. CTB-TT is an instance of Algorithm 1 using the CTB(ϵ) estimator (Algorithm 4), $(W_{a,b}^G, b_a^G)$ as in (3) with $\sigma = 1/2$ and $c_{a,b}^G$ as in (4).

Using Lemma 11, the CTB-TT algorithm is ϵ -local DP and is equivalent to running the non-private TTUCB algorithm on a modified bandit instance ν_ϵ , where $\nu_{\epsilon,a} = \text{Ber}(\mu_{\epsilon,a})$ with $\mu_{\epsilon,a} = (2\mu_a - 1) \frac{e^\epsilon - 1}{2(e^\epsilon + 1)} + 1/2$ for all $a \in [K]$. While the analysis in Jourdan and Degenne (2024) is written for Gaussian distributions with unit variance, their Section 3.2 shows that the same results can be obtained for σ -sub-Gaussian distributions. As such, the theoretical guarantees obtained in Jourdan and Degenne (2024) apply to our algorithm. In particular, CTB-TT is δ -correct and satisfies that, for all $\nu \in \mathcal{M}$ such that $\min_{a \neq b} |\mu_a - \mu_b| > 0$,

$$\limsup_{\delta \rightarrow 0} \frac{\mathbb{E}_\nu[\tau_\delta]}{\log(1/\delta)} \leq T_{\text{KL},\beta}^*(\nu_\epsilon) = \left(1 + \frac{2}{e^\epsilon - 1}\right)^2 T_{\text{KL},\beta}^*(\nu),$$

where $T_{\text{KL},\beta}^*$ as in (5) for $\sigma = 1/2$. For $\beta = 1/2$, combining Lemma 10 and (5) yields $\limsup_{\delta \rightarrow 0} \mathbb{E}_\nu[\tau_\delta]/\log(1/\delta) \leq (1 + 2/(e^\epsilon - 1))^2 T_{\text{TV}^2}^*(\nu)$. On top of its asymptotic guarantees, CTB-TT enjoys guarantees on its expected sample complexity at any confidence level (non-asymptotic regime). For all $\delta \in (0, 1)$ and all $\nu \in \mathcal{M}$ such that $|a^*(\nu)| = 1$,

$$\mathbb{E}_\nu[\tau_\delta] = \mathcal{O}((H(\nu_\epsilon) \log H(\nu_\epsilon))^\alpha) \quad \text{with} \quad H(\nu_\epsilon) = (1 + 2/(e^\epsilon - 1))^2 H(\nu) \quad \text{and} \quad \alpha > 1.$$

The notation \mathcal{O} gives the dominating term when $H(\nu) \rightarrow +\infty$.

In the non-private regime where $\epsilon \rightarrow +\infty$, our upper bound recovers the result of Jourdan and Degenne (2024). It matches the non-private lower bound for Gaussian distributions $T_{\text{KL}}^*(\nu)$ up to a multiplicative factor 2. Our upper bound matches the lower bound of Theorem 9 up to a multiplicative factor of $(e^\epsilon + 1)^2 \min\{4, e^{2\epsilon}\}$, whose limit is 4 when $\epsilon \rightarrow 0$. Instead of a fixed design β , we could use the optimal design IDS (You et al., 2023a) which sets β_n adaptively, i.e. $\beta_n = \frac{N_{n,C_n}}{N_{n,C_n} + N_{n,B_n}}$ for Gaussian distributions. Since this modification yields $T_{\text{KL}}^*(\nu_\epsilon)$ as an asymptotic upper bound, it shaves a multiplicative factor 2. In the limit of $\epsilon \rightarrow 0$, it leaves a multiplicative gap of 2 between the lower and the upper bound. Closing this gap is an interesting direction for future research.

4 Global Differentially Private Best-Arm Identification

The central question that we address in this section is: *How many additional samples a BAI strategy must select for ensuring ϵ -global DP?* In response, we prove a lower bound on the expected sample complexity of any δ -correct ϵ -global DP BAI strategy (Section 4.1). In order to obtain a matching upper bound, we propose a private mean estimator (Section 4.2) which can be used by our TTUCB-based algorithms (Sections 4.3 and 4.4).

4.1 Lower Bound on the Expected Sample Complexity

First, we derive an ϵ -global DP version of the “change-of-measure” lemma.

Lemma 12 *Let $\delta \in (0, 1)$ and $\epsilon > 0$. Let ν be a bandit instance and $\lambda \in \text{Alt}(\nu)$. For any δ -correct ϵ -global DP BAI strategy, we have that*

$$6\epsilon \sum_{a \in [K]} \mathbb{E}_{\nu, \pi} [N_{\tau_\delta, a}] \text{TV}(\nu_a \parallel \lambda_a) \geq \text{kl}(1 - \delta, \delta),$$

where $\text{kl}(1 - \delta, \delta) \triangleq x \log \frac{x}{y} + (1 - x) \log \frac{1-x}{1-y}$ for $x, y \in (0, 1)$.

Proof We use Sequential Karwa-Vadhan Lemma (Azize and Basu, 2022, Lemma 2) with a data-processing inequality in the BAI canonical model. Extra care is needed *to deal with the stopping times in the coupling, compared to a fixed horizon T in regret minimisation.* ■

Then, we derive a lower bound on the expected sample complexity in ϵ -global DP FC-BAI, which features problem-dependent characteristic times. The proof employs a similar sequential coupling argument as in the regret lower bound for bandits with ϵ -global DP (Shariff and Sheffet, 2018; Azize and Basu, 2022), see Appendix B for more details.

Theorem 13 *Let $\delta \in (0, 1)$ and $\epsilon > 0$. For any δ -correct and ϵ -global DP FC-BAI algorithm, we have that $\mathbb{E}_\nu[\tau_\delta] \geq T_g^*(\nu; \epsilon) \log(1/(2.4\delta))$ with*

$$T_g^*(\nu; \epsilon)^{-1} \triangleq \sup_{\omega \in \Sigma_K} \inf_{\lambda \in \text{Alt}(\nu)} \min \left\{ \sum_{a \in [K]} \omega_a \text{KL}(\nu_a \parallel \lambda_a), 6\epsilon \sum_{a \in [K]} \omega_a \text{TV}(\nu_a \parallel \lambda_a) \right\}.$$

As for the non-private BAI (Garivier and Kaufmann, 2016), Theorem 13 is the value of a two-player zero-sum game between a MIN player and MAX player. On top of the KL divergence present in the non-private lower bound, our bound features the TV distance that appears naturally when incorporating the ϵ -global DP constraint. The TV characteristic time $T_{\text{TV}}^*(\nu)$ serves as the BAI counterpart to the TV-distinguishability gap (t_{inf}) in the problem-dependent regret lower bound for bandits with ϵ -global DP as in Azize and Basu (2022, Theorem 3).

Corollary 14 (Relaxing the global DP lower bound) *Let $T_g^*(\nu; \epsilon)$ as in Theorem 13 and $T_d^*(\nu)$ as in Eq. (1). Then, we have*

$$T_g^*(\nu; \epsilon) \geq \max\{T_{\text{KL}}^*(\nu), T_{\text{TV}}^*(\nu)/(6\epsilon)\}. \tag{8}$$

Let $\boldsymbol{\nu}$ be a Bernoulli instance with mean gaps $\Delta_a = \mu_{a^*} - \mu_a$ for all $a \neq a^*$ and $\Delta_{a^*} = \Delta_{\min} = \min_{a \neq a^*} (\mu_{a^*} - \mu_a)$ for all $a \neq a^*$. Then, we have

$$T_{\text{TV}}^*(\boldsymbol{\nu}) = \sum_{a \in [K]} \Delta_a^{-1} \quad \text{and} \quad T_g^*(\boldsymbol{\nu}; \epsilon) \leq T_\epsilon^*(\boldsymbol{\nu}) \leq H_\epsilon(\boldsymbol{\nu}), \quad \text{with}$$

$$T_\epsilon^*(\boldsymbol{\nu})^{-1} \triangleq \max_{\omega \in \Sigma_K} \min_{a \neq a^*} \frac{2\Delta_a \min\{3\epsilon, \Delta_a\}}{1/\omega_{a^*} + 1/\omega_a} \quad \text{and} \quad H_\epsilon(\boldsymbol{\nu}) = \sum_{a \in [K]} (\Delta_a \min\{\Delta_a, 3\epsilon\})^{-1}. \quad (9)$$

Proof The first part is a direct consequence of the definition of $T^*(\boldsymbol{\nu}; \epsilon)$. The second part uses $\text{TV}(\text{Ber}(p) \parallel \text{Ber}(q)) = |p - q|$ to solve the optimisation problem and is detailed in Appendix B.3.3. The last part is obtained by using Pinsker's inequality. \blacksquare

Two privacy regimes. The sample complexity lower bound in (8) suggests the existence of *two hardness regimes depending on ϵ , $T_{\text{KL}}^*(\boldsymbol{\nu})$ and $T_{\text{TV}}^*(\boldsymbol{\nu})$* . (1) *Low-privacy regime:* When $\epsilon > T_{\text{TV}}^*(\boldsymbol{\nu})/(6T_{\text{KL}}^*(\boldsymbol{\nu}))$, the lower bound retrieves the non-private lower bound, i.e. $T_{\text{KL}}^*(\boldsymbol{\nu})$, and thus, **privacy can be achieved for free**. (2) *High-privacy regime:* When $\epsilon < T_{\text{TV}}^*(\boldsymbol{\nu})/(6T_{\text{KL}}^*(\boldsymbol{\nu}))$, the lower bound becomes $T_{\text{TV}}^*(\boldsymbol{\nu})/6\epsilon$ and ϵ -global DP δ -BAI requires more samples than non-private ones. Using Pinsker's inequality, one can connect the TV and KL characteristic times by $T_{\text{TV}}^*(\boldsymbol{\nu}) \geq \sqrt{2T_{\text{KL}}^*(\boldsymbol{\nu})}$.

The global trade-off between low and high privacy regimes at the instance level given by (8) does not give any information at the level of a specific arm. For each sub-optimal arm, the transition from low to high privacy is better understood by considering (9), even though it only upper bounds $T^*(\boldsymbol{\nu}; \epsilon)$. For any arm $a \neq a^*(\boldsymbol{\nu})$, the high-privacy regime corresponds to a mean gap such that $\epsilon < \Delta_a/3$, and the low-privacy regime to $\epsilon > \Delta_a/3$.

4.2 Private Mean Estimator

To define a sequence of mean estimators, we propose the $\text{DAF}(\epsilon)$ update (Algorithm 5) which relies on three ingredients: *adaptive episodes with doubling, forgetting, and adding calibrated Laplacian noise*. (1) DAF maintains K episodes, i.e. one per arm. The private empirical estimate of the mean of an arm is only updated at the end of an episode, that means when the number of times that a particular arm was played doubles. (2) For each arm a , DAF forgets rewards from previous phases of arm a , i.e. the private empirical estimate of arm a is only computed using the rewards collected in the last phase of arm a . This assures that the means of each arm are estimated using a non-overlapping sequence of rewards. (3) Thanks to this *doubling* and *forgetting*, DAF is ϵ -global DP as soon as each empirical mean is made ϵ -DP, and thus, avoiding any use of privacy composition. This is achieved by adding Laplace noise. We formalise this intuition in Lemma 22 of Appendix C.

Lemma 15 *Any algorithm relying solely on the $\text{DAF}(\epsilon)$ update is ϵ -global DP on $[0, 1]$.*

Proof A change in one user *only affects* the empirical mean calculated at one episode of an arm, which is made private using the Laplace Mechanism and Lemma 22. Since the sampled actions, recommended action, and stopping time are computed only using the private empirical means, the algorithm satisfies ϵ -global DP thanks to the post-processing lemma. \blacksquare

Algorithm 5 Doubling-And-Forgetting(ϵ) Estimator (DAF)

Input: History \mathcal{H}_n , arm $a \in [K]$.
Initialisation: $T_1(a) = K + 1$ and $k_{K+1,a} = 1$;
if $N_{n,a} \geq 2N_{T_{k_{n,a}}(a),a}$ **then**
 Change phase $k_{n,a} \leftarrow k_{n,a} + 1$ for arm a ;
 Set $T_{k_{n,a}}(a) = n$ and $\tilde{N}_{k_{n,a},a} = N_{T_{k_{n,a}}(a),a} - N_{T_{k_{n,a}-1}(a),a}$;
 Set $\hat{\mu}_{k_{n,a},a} = \tilde{N}_{k_{n,a},a}^{-1} \sum_{t=T_{k_{n,a}-1}(a)}^{T_{k_{n,a}}(a)-1} r_t \mathbb{1}\{a_t = a\}$;
 Set $\tilde{\mu}_{k_{n,a},a} = \hat{\mu}_{k_{n,a},a} + Y_{k_{n,a},a}$ where $Y_{k_{n,a},a} \sim \text{Lap}((\epsilon \tilde{N}_{k_{n,a},a})^{-1})$
end if
Return $(\tilde{\mu}_{n,a}, \tilde{N}_{n,a})$;

4.3 A Plug-In Approach: the AdaP-TT Algorithm

A natural approach is to simply plug in the private mean estimator in the non-private algorithm. The Plug-In approach is successful for ϵ -local DP FC-BAI (Section 3) and for ϵ -global DP regret minimisation (Azize and Basu, 2022).

AdaP-TT algorithm. To solve ϵ -global DP FC-BAI, we propose the AdaP-TT algorithm. AdaP-TT is an instance of Algorithm 1 using the DAF(ϵ) estimator (Algorithm 5),

$$W_{a,b}^G \text{ as in (3) for } \sigma = 1/2, b_a^{G,\epsilon}(\omega) = \sqrt{\frac{k(\omega_a)}{\omega_a} + \frac{k(\omega_a)}{\epsilon\omega_a}} \text{ with } k(x) = \log_2 x + 2, \quad (10)$$

and $c_{a,b}^{G,\epsilon}$ as in (11) which yields δ -correctness for any sampling rule (Lemma 16).

Lemma 16 *Let $\delta \in (0, 1)$, $\epsilon > 0$. Let $s > 1$ and ζ be the Riemann ζ function. Let $c_{a,b}^G$ as in (4) and $k(x) = \log_2 x + 2$. Given any sampling rule, combining the DAF(ϵ) estimator with the GLR stopping rule as in Eq. (2) with $W_{a,b}^G$ as in (3) and the stopping threshold*

$$c_{a,b}^{G,\epsilon}(\omega, \delta) = 2c_{a,b}^G(\omega, \delta(2\zeta(s)^2 k(\omega_a)^s k(\omega_b)^s)^{-1}) + \frac{1}{\epsilon^2 \sigma^2} \sum_{c \in \{a,b\}} \frac{1}{\omega_c} \left(\log \frac{2K\zeta(s)k(\omega_c)^s}{\delta} \right)^2, \quad (11)$$

yields a δ -correct algorithm for σ -sub-Gaussian distributions.

Asymptotically, our threshold is $c_{a,b}^{G,\epsilon}(\omega, \delta) \approx_{\delta \rightarrow 0} 2 \log(1/\delta) + (1/\omega_a + 1/\omega_b) \log(1/\delta)^2 / (\epsilon^2 \sigma^2)$.

Proof Proving δ -correctness of a GLR stopping rule is done by leveraging concentration results. Specifically, we start by decomposing the failure probability $\mathbb{P}_\mu(\tau_\delta < +\infty, \hat{a} \neq a^*)$ into a non-private and a private part using the basic property of $\mathbb{P}(X + Y \geq a + b) \leq \mathbb{P}(X \geq a) + \mathbb{P}(Y \geq b)$. The two-factor in front of $c_{a,b}^G$ originates from the looseness of this decomposition, and we improve on it in Section 4.4. We conclude using concentration results from σ -sub-Gaussian and Laplace random variables. The proof is detailed in Appendix D.2. ■

Theorem 17 *Let $(\delta, \beta) \in (0, 1)^2$ and $\epsilon > 0$. The AdaP-TT algorithm is ϵ -global DP, δ -correct and satisfies that, for all $\mu \in \mathbb{R}^K$ such that $\min_{a \neq b} |\mu_a - \mu_b| > 0$,*

$$\limsup_{\delta \rightarrow 0} \frac{\mathbb{E}_\nu[\tau_\delta]}{\log(1/\delta)} \leq 4T_{\text{KL},\beta}^*(\nu) \left(1 + \sqrt{1 + \frac{\Delta_{\max}^2}{2\sigma^4 \epsilon^2}} \right) \text{ with } \sigma = 1/2.$$

We adapt the asymptotic proof of the TTUCB algorithm, which is based on the unified analysis of Top Two algorithms from Jourdan et al. (2022). We sketch high-level ideas of the proof and specify the effect of the DAF(ϵ) estimator on the expected sample complexity.

Proof (1) The *non-private TTUCB algorithm* (Jourdan and Degenne, 2024) achieves a sample complexity of $T_{\text{KL},\beta}^*(\boldsymbol{\mu})$ for sub-Gaussian random variables. The proof relies on showing that the empirical pulling counts are converging towards the β -optimal allocation $\omega_{\text{KL},\beta}^*(\boldsymbol{\mu})$. (2) The *effect of doubling and forgetting* is a multiplicative four-factor, i.e. $4T_{\text{KL},\beta}^*(\boldsymbol{\mu})$. The first two-factor is due to forgetting since we throw away half of the samples. The second two-factor is due to doubling since we have to wait for the end of an episode to evaluate the stopping condition. (3) The *Laplace noise* only affects the empirical estimate of the mean. Since the Laplace noise has no bias and a sub-exponential tail, the private means will still converge towards their true values. Therefore, the empirical counts will also converge to $\omega_{\text{KL},\beta}^*(\boldsymbol{\mu})$ asymptotically. (4) While the *Laplace noise has little effect on the sampling rule* itself, it *changes the dependency in $\log(1/\delta)$ of the threshold* used in the GLR stopping rule. The private threshold $c_{a,b}^{G,\epsilon}$ has an extra factor $\mathcal{O}(\log^2(1/\delta))$ compared to the non-private one $c_{a,b}^G$. Using the convergence towards $\omega_{\text{KL},\beta}^*(\boldsymbol{\mu})$, the stopping condition is met as soon as $\frac{n}{T_{\text{KL},\beta}^*(\boldsymbol{\mu})} \lesssim 2 \log(1/\delta) + \frac{\Delta_{\max}^2}{2\sigma^4\epsilon^2} \frac{T_{\text{KL},\beta}^*(\boldsymbol{\mu})}{n} \log^2(1/\delta)$. Solving the inequality for n concludes the proof while adding a multiplicative four-factor. \blacksquare

Discussion. In the non-private regime where $\epsilon \rightarrow +\infty$, our upper bound recovers the non-private lower bound for Gaussian distributions $T_{\text{KL}}^*(\boldsymbol{\nu})$ up to a multiplicative factor 16. For Bernoulli distributions (or bounded distributions in $[0, 1]$), there is still a mismatch between the upper and lower bounds due to the mismatch between the KL divergence of Bernoulli distributions and that of Gaussian (e.g. large ratio when the means are close to 0 or 1). This is in essence, similar to the mismatch between UCB and KL-UCB in the regret-minimisation literature (e.g. Chapter 10 in Lattimore and Szepesvári 2020). To overcome this mismatch, it is necessary to adapt the transportation costs to the family of distributions considered. While the Top Two algorithms for Bernoulli distributions (or bounded distributions in $[0, 1]$) have been studied in Jourdan et al. (2022), the analysis is more involved. Therefore, it would obfuscate where and how privacy is impacting the expected sample complexity.

In the asymptotic highly privacy regime where $\epsilon \rightarrow 0$, our upper bound becomes $\mathcal{O}(T_{\text{KL}}^*(\boldsymbol{\nu})\Delta_{\max}/\epsilon)$ while the lower bound is $\Omega(T_{\text{TV}}^*(\boldsymbol{\nu})/\epsilon)$. Therefore, our upper bound is only asymptotically tight for instances such that $T_{\text{KL}}^*(\boldsymbol{\nu}) = \mathcal{O}(T_{\text{TV}}^*(\boldsymbol{\nu})/\Delta_{\max})$, e.g. instances where the mean gaps have the same order of magnitude. In all the other cases, the plug-in approach is sub-optimal due to a problem-dependent gap.

4.4 A Lower Bound Based Approach: the AdaP-TT* Algorithm

To overcome the limitation of AdaP-TT, one should adapt the transportation costs to reflect the lower bound (Theorem 13) instead of “ignoring” the privacy constraint by using the transportation costs $W_{a,b}^G$ as in (3) which are tailored for non-private FC-BAI (Lemma 4).

AdaP-TT* algorithm. Therefore, we propose the AdaP-TT* algorithm. AdaP-TT* is an instance of Algorithm 1 using the DAF(ϵ) estimator (Algorithm 5), $b_{a,b}^{G,\epsilon}$ as in (10),

$$W_{a,b}^{G,\epsilon}(\tilde{\mu}, \omega) = \frac{(\tilde{\mu}_a - \tilde{\mu}_b)_+ \min\{3\epsilon, (\tilde{\mu}_a - \tilde{\mu}_b)_+\}}{2\sigma^2(1/\omega_a + 1/\omega_b)} \quad \text{with } \sigma = 1/2, \quad (12)$$

and $\tilde{c}_{a,b}^{G,\epsilon}(\tilde{\mu}, \omega, \delta)$ as in (14), which yields δ -correctness for any sampling rule (Lemma 18). Compared to previous stopping thresholds, $\tilde{c}_{a,b}^{G,\epsilon}$ depends on the mean estimator $\tilde{\mu}$. The transportation cost $W_{a,b}^{G,\epsilon}$ is inspired by the relaxed β -characteristic time (Corollary 14)

$$T_{\epsilon,\beta}^*(\boldsymbol{\nu})^{-1} \triangleq \max_{\omega \in \Sigma_K, \omega_{a^*} = \beta} \min_{a \neq a^*} \frac{(\mu_{a^*} - \mu_a) \min\{3\epsilon, \mu_{a^*} - \mu_a\}}{2\sigma^2(1/\beta + 1/\omega_a)} \quad \text{with } \sigma = 1/2. \quad (13)$$

Lemma 18 *Let $\delta \in (0, 1)$, $\epsilon > 0$. Let $s > 1$ and ζ be the Riemann ζ function. Let $\overline{W}_{-1}(x) = -W_{-1}(-e^{-x})$ for all $x \geq 1$, where W_{-1} is the negative branch of the Lambert W function. It satisfies $\overline{W}_{-1}(x) \approx x + \log x$. Let $c_{a,b}^{G,\epsilon}$ as in (11), $k(x) = \log_2 x + 2$ and*

$$h(x, \delta) = \overline{W}_{-1}(2 \log(3K\zeta(s)k(x)^s/\delta) + 4 \log(4 + \log x) + 1/2) / 2.$$

Given any sampling rule, combining the DAF(ϵ) estimator with the GLR stopping rule as in Eq. (2) with $W_{a,b}^{G,\epsilon}$ as in (12) and the stopping threshold $\tilde{c}_{a,b}^{G,\epsilon}(\tilde{\mu}, \omega, \delta)$ which is equal to

$$\begin{cases} \frac{1}{2}c_{a,b}^{G,\epsilon}(\omega, 2\delta/3) + \frac{\sqrt{2}}{\epsilon\sigma} \sum_{c \in \{a,b\}} \sqrt{\frac{h(\omega_c, \delta)}{\omega_c}} \log\left(\frac{3K\zeta(s)k(\omega_c)^s}{\delta}\right) & \text{if } (\tilde{\mu}_a - \tilde{\mu}_b)_+ < 3\epsilon \\ \frac{3}{\sigma^2} \log(3K\zeta(s) \max_{c \in \{a,b\}} k(w_c)/\delta) + \frac{3\epsilon}{\sqrt{2}\sigma} \sum_{c \in \{a,b\}} \sqrt{\omega_c h(\omega_c, \delta)} & \end{cases}, \quad (14)$$

yields a δ -correct algorithm for σ -sub-Gaussian distributions.

Our threshold is $\frac{3}{\sigma^2} \log(1/\delta) + \frac{3\epsilon}{\sqrt{2}\sigma} (\sqrt{\omega_b} + \sqrt{\omega_a}) \sqrt{\log(1/\delta)}$ when $\tilde{\mu}_a - \tilde{\mu}_b \geq 3\epsilon$, and

$$\log(1/\delta) + \frac{1}{2\epsilon^2\sigma^2} (1/\omega_a + 1/\omega_b) \log(1/\delta)^2 + \frac{\sqrt{2}}{\epsilon\sigma} (\sqrt{1/\omega_a} + \sqrt{1/\omega_b}) \log(1/\delta)^{3/2} \quad \text{otherwise.}$$

Proof The proof is similar to the one of Lemma 16 with tighter manipulations allowing to divide $c_{a,b}^{G,\epsilon}$ by 2. It is detailed in Appendix D.3. \blacksquare

Theorem 19 *Let $(\delta, \beta) \in (0, 1)^2$ and $\epsilon > 0$. The AdaP-TT* algorithm is ϵ -global DP, δ -correct and satisfies that, for all $\mu \in \mathbb{R}^K$ such that $\min_{a \neq b} |\mu_a - \mu_b| > 0$,*

$$\limsup_{\delta \rightarrow 0} \frac{\mathbb{E}_{\boldsymbol{\nu}}[\tau_\delta]}{\log(1/\delta)} \leq \begin{cases} 4T_{\text{KL},\beta}^*(\boldsymbol{\nu})g_1(\Delta_{\max}/(\sigma^2\epsilon)) & \text{if } \Delta_{\max} < 3\epsilon \\ 12T_{\epsilon,\beta}^*(\boldsymbol{\nu})g_2(3\epsilon^2T_{\epsilon,\beta}^*(\boldsymbol{\nu}) \max\{\beta, 1-\beta\}/2)/\sigma^2 & \text{otherwise} \end{cases},$$

where $T_{\epsilon,\beta}^(\boldsymbol{\nu})$ as in (13) with $\sigma = 1/2$. The function $g_1(y) = \sup\{x \mid x^2 < x + y\sqrt{2x} + \frac{y^2}{4}\}$ is increasing on $[0, 12]$ and satisfies that $g_1(0) = 1$ and $g_1(12) \leq 10$. The function $g_2(y) = 1 + 2(\sqrt{1 + 1/y} - 1)^{-1}$ is increasing on \mathbb{R}_+^* and satisfies that $\lim_{y \rightarrow 0} g_2(y) = 1$.*

Proof The proof is similar to the one of Theorem 17 with tighter manipulations. \blacksquare

Discussion. When $\Delta_{\max} < 3\epsilon$, our upper bound recovers the non-private lower bound for Gaussian distributions $T_{\text{KL}}^*(\boldsymbol{\nu})$ up to a multiplicative factor $8g_1(4\Delta_{\max}/\epsilon) \in [8, 80]$, whose limit is 8 in non-private regime where $\epsilon \rightarrow +\infty$. When $\Delta_{\min} \geq 3\epsilon$, we have $12T_{\epsilon, \beta}^*(\boldsymbol{\nu}) \leq 8T_{\text{TV}}^*(\boldsymbol{\nu})/\epsilon$. In the asymptotic highly privacy regime where $\epsilon \rightarrow 0$, our upper bound matches the lower bound up to a multiplicative factor 48. Therefore, we close the gap left open by the algorithm in Section 4.3. While the regime $\Delta_{\max} \geq 3\epsilon > \Delta_{\min}$ is relevant for practical application, it is harder to understand how the different quantities interact in the upper/lower bounds in transitional phases. Thus, it is harder to claim optimality in those phases. Having matching upper and lower bounds *only* for high privacy regimes is an interesting phenomenon that appears in different settings of differential privacy literature, such as regret minimisation (Azize and Basu, 2022), parameter estimation (Cai et al., 2021) and hidden probabilistic graphical models (Nikolakakis et al., 2019).

Comparison to DP-SE. DP-SE (Sajed and Sheffet, 2019) is an ϵ -global DP version of the Successive Elimination algorithm introduced for the regret minimisation setting. The algorithm samples active arms uniformly during phases of geometrically increasing length. Based on the private confidence bounds, DP-SE eliminates provably sub-optimal arms at the end of each phase. Due to its phased-elimination structure, DP-SE can be easily converted into an ϵ -global DP FC-BAI algorithm, where we stop once there is only one active arm left. In particular, the proof of Theorem 4.3 of Sajed and Sheffet (2019) shows that with high probability any sub-optimal arm $a \neq a^*$ is sampled no more than $\mathcal{O}(\Delta_a^2 + (\epsilon\Delta_a)^{-1})$. From this result, it is straightforward to extract a sample complexity upper bound for DP-SE, i.e. $\mathcal{O}(\sum_{a \neq a^*} \Delta_a^{-2} + \sum_{a \neq a^*} (\epsilon\Delta_a)^{-1})$. This shows that DP-SE too achieves (ignoring constants) the high-privacy lower bound $T_{\text{TV}}^*(\boldsymbol{\nu})/\epsilon$ for Bernoulli instances. However, due to its uniform sampling within the phases, DP-SE is less adaptive than TTUCB. Inside a phase, DP-SE continues to sample arms that might already be known to be bad, while TTUCB adapts its sampling rule based on the transportation costs that reflect the amount of evidence collected in favour of the hypothesis that the leader is the best arm. Finally, TTUCB has the advantage of being anytime, i.e. its sampling strategy does not depend on the risk δ .

Another adaptation of DP-SE, namely DP-SEQ, is proposed in Kalogerias et al. (2021) for the problem of privately finding the arm with the highest quantile at a fixed level, hence it is different from BAI. For multiple agents, Rio et al. (2023) studies privacy for BAI under fixed confidence. They propose and analyse the sample complexity of DP-MASE, a multi-agent version of DP-SE. They show that multi-agent collaboration leads to better sample complexity than independent agents, even under privacy constraints. While the multi-agent setting with federated learning allows tackling large-scale clinical trials taking place at several locations simultaneously, we study the single-agent setting, which is relevant for many small-scale clinical trials (see Example 1).

Remark 20 (On the number of rounds of adaptivity) *Used on any existing FC-BAI algorithm, the DAF update yields a batched algorithm, which satisfies ϵ -global DP. At the end of the episode of arm a (after updating its mean), it is possible to compute the sequence of all the arms to be pulled before the end of the next episode (for another arm), without*




Figure 1: Empirical stopping time τ_δ (mean \pm std. over 1000 runs, $\delta = 10^{-2}$) with respect to the privacy budget ϵ for ϵ -local DP on Bernoulli instance μ_1 (left) and μ_2 (right). The shaded vertical line separates the two privacy regimes.

taking the collected observations into account. In contrast to the classical batched setting where the batch size is fixed, the size of the resulting batches is adaptive and data-dependent. In the non-private setting ($\epsilon = +\infty$), we recover Batched Best-Arm Identification (BBAI) in the fixed-confidence setting. AdaP-TT and AdaP-TT* are asymptotically optimal up to a multiplicative factor 4 with solely $\mathcal{O}(K \log_2(T_{\text{KL}}^*(\nu) \log(1/\delta)))$ rounds of adaptivity. We refer the reader to Appendix F for more details, including comparison to existing works.

5 Experimental Analysis

We perform experiments for both ϵ -local DP and ϵ -global DP. The code is available here.

5.1 Local DP

We run CTB-TT in different Bernoulli instances as in Sajed and Sheffet (2019). As a benchmark, we also compare to the non-private TTUCB. As for ϵ -global DP, we set the risk $\delta = 10^{-2}$, implement all the algorithms in Python (version 3.8) and run each algorithm 1000 times. We plot the corresponding average and standard deviations of the empirical stopping times in Figure 1. We also test the algorithms on other Bernoulli instances and report the results in Appendix G.

Figure 1 shows that CTB-TT performance has two regimes. In the low privacy regime ($\epsilon > 4$ for μ_1 and $\epsilon > 2$ for μ_2), the CTB(ϵ) estimator reduces to the MLE, and CTB-TT matches exactly the performance of the non-private TTUCB. In the high privacy regime ($\epsilon < 4$ for μ_1 and $\epsilon < 2$ for μ_2), the price of privacy on the stopping time is a multiplicative ϵ^{-2} . Therefore, the sample complexity is prohibitively large to be computed numerically for $\epsilon < 0.1$. The switching value of ϵ between the low and high privacy regimes is an order of magnitude higher for ϵ -local DP compared to the one for ϵ -global DP. This is predictable since local DP provides a “stronger” privacy guarantee at the cost of worse performance.




Figure 2: Empirical stopping time τ_δ (mean \pm std. over 1000 runs) with respect to the privacy budget ϵ for ϵ -global DP on Bernoulli instance μ_1 (left) and μ_2 (right). The shaded vertical line separates the two privacy regimes.

5.2 Global DP

We compare the performances of AdaP-TT, AdaP-TT* and DP-SE for FC-BAI in different Bernoulli instances as in Sajed and Sheffet (2019). The first instance has means $\mu_1 = (0.95, 0.9, 0.9, 0.9, 0.5)$ and the second instance has means $\mu_2 = (0.75, 0.7, 0.7, 0.7, 0.7)$. As a benchmark, we also compare to the non-private TTUCB. We set the risk $\delta = 10^{-2}$ and implement all the algorithms in Python (version 3.8). We run each algorithm 1000 times, and plot corresponding average and standard deviations of the empirical stopping times in Figure 2. We also test the algorithms on other Bernoulli instances and report the results in Appendix G.

Figure 2 shows that: (a) AdaP-TT and AdaP-TT* require fewer samples than DP-SE to provide a δ -correct answer, for different values of ϵ and in all the instances tested. AdaP-TT and AdaP-TT* have the same performance in the low privacy regimes, while AdaP-TT* improves the sample complexity in the high privacy regime, as predicted theoretically. (b) The experimental performance of AdaP-TT and AdaP-TT* demonstrate two regimes. A high-privacy regime (for $\epsilon < 0.1$ for μ_1 and $\epsilon < 0.4$ for μ_2), where the stopping time depends on the privacy budget ϵ , and a low privacy regime (for $\epsilon > 0.1$ for μ_1 and $\epsilon > 0.4$ for μ_2), where the performance of AdaP-TT and AdaP-TT* does not depend on ϵ , and is four times the samples required by TTUCB in the worst case, as shown theoretically.

6 Perspectives

We study FC-BAI with ϵ -local DP and ϵ -global DP. In both settings, we derive a lower bound on the expected sample complexity which quantifies the additional samples needed by a δ -correct BAI strategy to ensure DP. The lower bounds further suggest the existence of two privacy regimes. In the *low-privacy regime*, no additional samples are needed, and *privacy can be achieved for free*. For the *high-privacy regime*, the lower bound reduces to $\Omega(\epsilon^{-2}T_{TV^2}^*(\nu))$ for ϵ -local DP, and to $\Omega(\epsilon^{-1}T_{TV}^*(\nu))$ for ϵ -global DP. To match those lower

bounds up to multiplicative constants, we propose ϵ -local DP and ϵ -global DP variants of a Top Two algorithm. For ϵ -local DP, the CTB-TT algorithm reaches asymptotic optimality by plugging in a private estimator of the means based on Randomised Response. For ϵ -global DP, our private estimator of the mean runs in *arm-dependent adaptive episodes* and adds *Laplace noise* to ensure a good privacy-utility trade-off. By solely plugging in this estimator, the AdaP-TT algorithm fails to recover the asymptotic lower bound for instances with highly different mean gaps. The AdaP-TT* algorithm overcomes this limitation by adapting the transportation costs.

The upper bound matches the lower bound by a multiplicative constant in the high privacy regime, and is also loose in some instances in the low privacy regime, due to the mismatch between the KL divergence of Bernoulli distributions and that of Gaussian. One possible direction to solve this issue is to use transportation costs tailored to Bernoulli for both the Top Two Sampling and the stopping. Since our bounds only give a clear picture in the high and low privacy regimes, it would be interesting to provide better insights for the regime in-between where both the δ -correctness and the DP constraints are of the same order. An interesting direction would be to extend the proposed technique to other variants of pure DP, namely (ϵ, δ) -DP and Rényi-DP (Mironov, 2017), or other trust models, e.g. shuffle DP (Cheu, 2021; Girgis et al., 2021).

Acknowledgments and Disclosure of Funding

This work has been partially supported by the THIA ANR program “AI.PhD@Lille”. A. Al-Marjani acknowledges the support of the Chaire SeqALO (ANR-20-CHIA-0020). D. Basu acknowledges the Inria-Kyoto University Associate Team “RELIANT” for supporting the project, the ANR JCJC for the REPUBLIC project (ANR-22-CE23-0003-01), and the PEPR project FOUNDRY (ANR23-PEIA-0003). We thank Emilie Kaufmann and Aurélien Garivier for the interesting conversations. We also thank Philippe Preux for his support.

Appendix A. Outline

The appendices are organised as follows:

- In Appendix B, we detail the proofs of our lower bounds for ϵ -local DP (Theorem 9 and Corollary 10) and ϵ -global DP (Theorem 13 and Corollary 14).
- In Appendix C, we show that AdaP-TT and AdaP-TT* are ϵ -global DP since they use the DAF(ϵ) estimator of the means.
- In Appendix D, we prove that the ϵ -global DP GLR stopping rules yields δ -correctness regardless of the sampling rule, both when using non-private transportation costs (Lemma 16) and adapted transportation costs (Lemma 18).
- In Appendix E, we detail the proofs of the asymptotic upper bound on the expected sample complexity of AdaP-TT (Theorem 17) and AdaP-TT* (Theorem 19).
- In Appendix F, we discuss in more details the number of rounds of adaptivity.
- Extended experiments are presented in Appendix G.

Appendix B. Lower Bounds on the Expected Sample Complexity

In this section, we provide the proofs for the sample complexity lower bounds. First, we present the canonical model for BAI to introduce the relevant quantities. Then, we prove the ϵ -local DP sample complexity lower bound. Finally, for global-DP, we first prove an ϵ -global version of the transportation lemma, i.e. Lemma 12. Using this lemma, we prove the ϵ -global DP sample complexity lower bound of Theorem 13. We also prove the formula expressing the TV characteristic time for Bernoulli instances.

B.1 Canonical Model for BAI

Let $\nu \triangleq \{\nu_a : a \in [K]\}$ be a bandit instance, consisting of K arms with finite means $\{\mu_a\}_{a \in [K]}$. Now, we recall the interaction between a BAI strategy π and the bandit instance ν in the Protocol 3. The BAI strategy π halts at τ , samples a sequence of actions \underline{A}^τ , and recommends the action \hat{A} . Let $\mathbb{P}_{\nu, \pi}$ be the probability distribution over the triplets $(\tau, \underline{A}^\tau, \hat{A})$, when the BAI strategy π interacts with the bandit instance ν .

For a fixed $T > 1$, a sequence of actions $\underline{a}^T = (a_1, \dots, a_T) \in [K]^T$ and a recommendation $\hat{a} \in [K]$, we define the event $E = \{\tau = T, \underline{A}^\tau = \underline{a}^T, \hat{A} = \hat{a}\}$. We have that

$$\mathbb{P}_{\nu, \pi}(E) = \int_{\underline{r}^T = (r_1, \dots, r_T) \in \mathbb{R}^T} \pi(\underline{a}^T, \hat{a}, T \mid \underline{r}^T) \prod_{t=1}^T d\nu_{a_t}(r_t) dr_t$$

where

$$\pi(\underline{a}^T, \hat{a}, T \mid \underline{r}^T) \triangleq \text{Rec}_{T+1}(\hat{a} \mid \mathcal{H}_T) S_{T+1}(\top \mid \mathcal{H}_T) \prod_{t=1}^T S_t(a_t \mid \mathcal{H}_{t-1})$$

and $\mathcal{H}_t = (a_1, r_1, \dots, a_t, r_t)$.

Remark on the bandit feedback for global DP. Let π be an ϵ -DP BAI strategy. Let $T \geq 1$, $\underline{a}^T \in [K]^T$ a sequence sampled actions and $\hat{a} \in [K]$ a recommended actions. This time, let $\underline{r}^T = \{r_1, \dots, r_T\} \in \mathbb{R}^T$ and $\underline{r}'^T \in \mathbb{R}^T$ two neighbouring sequence of rewards, i.e. $d_{\text{Ham}}(\underline{r}^T, \underline{r}'^T) \triangleq \sum_{t=1}^T \mathbb{1}\{r_t \neq r'_t\} = 1$. Consider the table of rewards \underline{d}^T consisting of concatenating \underline{r}^T colon-wise K times, i.e. $\underline{d}_{t,i}^T = r_t^T$ for all $i \in [K]$ and all $t \in [T]$. Define \underline{d}'^T similarly with respect to \underline{r}'^T .

In this case, by definition of π , \underline{d}^T and \underline{d}'^T , it is direct that

$$\pi(\underline{a}^T, \hat{a}, T \mid \underline{r}^T) = \pi(\underline{a}^T, \hat{a}, T \mid \underline{d}^T)$$

and $d_{\text{Ham}}(\underline{d}^T, \underline{d}'^T) = 1$.

Which means that

$$\pi(\underline{a}^T, \hat{a}, T \mid \underline{r}^T) \leq e^\epsilon \pi(\underline{a}^T, \hat{a}, T \mid \underline{r}'^T).$$

In other words, if π is ϵ -pure DP for neighbouring table of rewards \underline{d}^T , then π is also ϵ -pure DP for neighbouring sequence of observed rewards \underline{r}^T .

Remark on the local DP canonical model. Let (\mathcal{M}, π) be a pair of perturbation mechanism and BAI satisfying ϵ -local DP. Let $\nu \triangleq \{\nu_a : a \in [K]\}$ be a bandit instance. In the local DP interaction protocol, the BAI strategy π only accesses the noisy rewards from the perturbation mechanism, i.e. $z_t \sim \mathcal{M}(r_t)$, where $r_t \sim \nu_{a_t}$. Thus, we can define an environment $\nu^{\mathcal{M}} \triangleq \{\nu_a^{\mathcal{M}} : a \in [K]\}$ induced by the perturbation mechanism, where

$$\nu_a^{\mathcal{M}}(Z) = \int_{r \in \mathbb{R}} \mathcal{M}(Z \mid r) d\nu_a(r) dr$$

is the marginal over the noisy rewards of arm a .

Thus, the local DP canonical model of the interaction between (\mathcal{M}, π) and an environment ν is equivalent to the “classical” canonical model between π and the induced environment $\nu^{\mathcal{M}}$.

B.2 Expected Sample Complexity Lower Bound under ϵ -local DP

Theorem 9 1 (Sample complexity lower bound for ϵ -local DP FC-BAI) *Let $\delta \in (0, 1)$ and $\epsilon > 0$. For any δ -correct ϵ -local DP pair (\mathcal{M}, π) of perturbation mechanism and BAI strategy, we have that $\mathbb{E}_\nu[\tau_\delta] \geq T_\ell^*(\nu; \epsilon) \log(1/(2.4\delta))$ with*

$$T_\ell^*(\nu; \epsilon)^{-1} \triangleq \sup_{\omega \in \Sigma_K} \inf_{\lambda \in \text{Alt}(\nu)} \sum_{a \in [K]} \omega_a \min \left\{ \text{KL}(\nu_a \parallel \lambda_a), c(\epsilon) (\text{TV}(\nu_a \parallel \lambda_a))^2 \right\},$$

where $c(\epsilon) = \min\{4, e^{2\epsilon}\} (e^\epsilon - 1)^2$ is a privacy term. For two probability distributions \mathbb{P}, \mathbb{Q} on the measurable space (Ω, \mathcal{F}) , the TV divergence is $\text{TV}(\mathbb{P} \parallel \mathbb{Q}) \triangleq \sup_{A \in \mathcal{F}} \{\mathbb{P}(A) - \mathbb{Q}(A)\}$.

Proof Let (\mathcal{M}, π) a perturbation mechanism and BAI strategy pair that it ϵ -local DP.

We suppose that π is δ -correct.

Using the remark in the local DP canonical model, π is δ -correct with respect to the environment $\nu^{\mathcal{M}} \triangleq \{\nu_a^{\mathcal{M}} : a \in [K]\}$ induced by the perturbation mechanism \mathcal{M} , where

$$\nu_a^{\mathcal{M}}(Z) = \int_{r \in \mathbb{R}} \mathcal{M}(Z \mid r) d\nu_a(r) dr$$

is the marginal over the noisy rewards of arm a .

Thus using Lemma in Kaufmann et al. (2016), we get that

$$\sum_{a=1}^K \mathbb{E} [N_a(\tau)] \text{KL} (\nu_a^{\mathcal{M}} \parallel \lambda_a^{\mathcal{M}}) \geq \text{kl}(1 - \delta, \delta)$$

for any alternative environment $\boldsymbol{\lambda} \in \text{Alt}(\boldsymbol{\nu})$.

Using Theorem 1 in Duchi et al. (2013), we have that

$$\begin{aligned} \text{KL} (\nu_a^{\mathcal{M}} \parallel \lambda_a^{\mathcal{M}}) &\leq \text{KL} (\nu_a^{\mathcal{M}} \parallel \lambda_a^{\mathcal{M}}) + \text{KL} (\lambda_a^{\mathcal{M}} \parallel \nu_a^{\mathcal{M}}) \\ &\leq \min\{4, e^{2\epsilon}\}(e^\epsilon - 1)^2 (\text{TV} (\nu_a \parallel \lambda_a))^2 \\ &= c(\epsilon)(\text{TV} (\nu_a \parallel \lambda_a))^2 \end{aligned}$$

where $c(\epsilon) \triangleq \min\{4, e^{2\epsilon}\}(e^\epsilon - 1)^2$.

On the other hand, using the data-processing inequality, we also have that

$$\text{KL} (\nu_a^{\mathcal{M}} \parallel \lambda_a^{\mathcal{M}}) \leq \text{KL} (\nu_a \parallel \lambda_a)$$

Thus, combining the two inequalities gives that

$$\begin{aligned} \text{kl}(1 - \delta, \delta) &\leq \inf_{\boldsymbol{\lambda} \in \text{Alt}(\boldsymbol{\nu})} \sum_{a=1}^K \mathbb{E} [N_a(\tau)] \min \{ \text{KL} (\nu_a \parallel \lambda_a), c(\epsilon)(\text{TV} (\nu_a \parallel \lambda_a))^2 \} \\ &= \mathbb{E}[\tau] \inf_{\boldsymbol{\lambda} \in \text{Alt}(\boldsymbol{\nu})} \sum_{a=1}^K \frac{\mathbb{E} [N_a(\tau)]}{\mathbb{E}[\tau]} \min \{ \text{KL} (\nu_a \parallel \lambda_a), c(\epsilon)(\text{TV} (\nu_a \parallel \lambda_a))^2 \} \\ &\leq \mathbb{E}[\tau] \left(\sup_{\omega \in \Sigma_K} \inf_{\boldsymbol{\lambda} \in \text{Alt}(\boldsymbol{\nu})} \sum_{a=1}^K \omega_a \min \{ \text{KL} (\nu_a \parallel \lambda_a), c(\epsilon)(\text{TV} (\nu_a \parallel \lambda_a))^2 \} \right) \\ &= \mathbb{E}[\tau] (T_\ell^*(\boldsymbol{\nu}, \epsilon))^{-1} \end{aligned}$$

The theorem follows by noting that for $\delta \in (0, 1)$, $\text{kl}(1 - \delta, \delta) \geq \log(1/2.4\delta)$. ■

B.3 Expected Sample Complexity Lower Bound under ϵ -global DP

B.3.1 TRANSPORTATION LEMMA UNDER ϵ -GLOBAL DP: PROOF OF LEMMA 12

Lemma 12 2 (Transportation lemma under ϵ -global DP) *Let $\delta \in (0, 1)$ and $\epsilon > 0$. Let $\boldsymbol{\nu}$ be a bandit instance and $\boldsymbol{\lambda} \in \text{Alt}(\boldsymbol{\nu})$. For any δ -correct ϵ -global DP BAI strategy, we have that*

$$6\epsilon \sum_{a=1}^K \mathbb{E}_{\boldsymbol{\nu}, \pi} [N_a(\tau)] \text{TV} (\nu_a \parallel \lambda_a) \geq \text{kl}(1 - \delta, \delta),$$

where $\text{kl}(x, y) \triangleq x \log \frac{x}{y} + (1 - x) \log \frac{1-x}{1-y}$ for $x, y \in (0, 1)$.

Proof Step 1: Distinguishability due to δ -correctness. Let π be a δ -correct ϵ -global DP BAI strategy. Let ν be a bandit instance and $\lambda \in \text{Alt}(\nu)$.

Let $\mathbb{P}_{\nu, \pi}$ denote the probability distribution of $(\underline{A}, \hat{A}, \tau)$ when the BAI strategy π interacts with ν . For any alternative instance $\lambda \in \text{Alt}(\nu)$, the data-processing inequality gives that

$$\begin{aligned} \text{KL}(\mathbb{P}_{\nu, \pi} \parallel \mathbb{P}_{\lambda, \pi}) &\geq \text{kl}\left(\mathbb{P}_{\nu, \pi}\left(\hat{A} = a^*(\nu)\right), \mathbb{P}_{\lambda, \pi}\left(\hat{A} = a^*(\nu)\right)\right) \\ &\geq \text{kl}(1 - \delta, \delta). \end{aligned} \quad (15)$$

where the second inequality is because π is δ -correct i.e. $\mathbb{P}_{\nu, \pi}\left(\hat{A} = a^*(\nu)\right) \geq 1 - \delta$ and $\mathbb{P}_{\lambda, \pi}\left(\hat{A} = a^*(\nu)\right) \leq \delta$, and the monotonicity of the kl.

Step 2: Connecting KL and TV under ϵ -global DP. On the other hand, by the definition of the KL, we have that

$$\text{KL}(\mathbb{P}_{\nu, \pi} \parallel \mathbb{P}_{\lambda, \pi}) = \mathbb{E}_{\tau, \underline{A}^T, \hat{A} \sim \mathbb{P}_{\nu, \pi}} \left[\log \left(\frac{\mathbb{P}_{\nu, \pi}(\tau, \underline{A}^T, \hat{A})}{\mathbb{P}_{\lambda, \pi}(\tau, \underline{A}^T, \hat{A})} \right) \right]$$

where

$$\mathbb{P}_{\nu, \pi}(\tau = T, \underline{A}^T = \underline{a}^T, \hat{A} = \hat{a}) = \int_{\underline{r} \in \mathbb{R}^T} \pi(\underline{a}^T, \hat{a}, T \mid \underline{r}) \prod_{t=1}^T d\nu_{a_t}(r_t).$$

Since π is ϵ -global DP, using the sequential Karwa-Vadhan lemma (Azize and Basu, 2022, Lemma 2), we get that

$$\begin{aligned} \log \left(\frac{\mathbb{P}_{\nu, \pi}(\tau = T, \underline{A}^T = \underline{a}^T, \hat{A} = \hat{a})}{\mathbb{P}_{\lambda, \pi}(\tau = T, \underline{A}^T = \underline{a}^T, \hat{A} = \hat{a})} \right) &\leq 6\epsilon \sum_{t=1}^T \text{TV}(\nu_{a_t} \parallel \lambda_{a_t}) \\ &= 6\epsilon \sum_{a=1}^K N_a(T) \text{TV}(\nu_a \parallel \lambda_a) \end{aligned}$$

Which gives that

$$\text{KL}(\mathbb{P}_{\nu, \pi} \parallel \mathbb{P}_{\lambda, \pi}) \leq 6\epsilon \mathbb{E}_{\nu, \pi} \left[\sum_{a=1}^K N_a(\tau) \text{TV}(\nu_a \parallel \lambda_a) \right]. \quad (16)$$

Combining Inequalities 15 and 16 concludes the proof. \blacksquare

B.3.2 PROOF OF THEOREM 13

Theorem 13 3 *Let $\delta \in (0, 1)$ and $\epsilon > 0$. For any δ -correct and ϵ -global DP FC-BAI algorithm, we have that $\mathbb{E}_{\nu}[\tau_{\delta}] \geq T_g^*(\nu; \epsilon) \log(1/(2.4\delta))$ with*

$$T_g^*(\nu; \epsilon)^{-1} \triangleq \sup_{\omega \in \Sigma_K} \inf_{\lambda \in \text{Alt}(\nu)} \min \left\{ \sum_{a \in [K]} \omega_a \text{KL}(\nu_a \parallel \lambda_a), 6\epsilon \sum_{a \in [K]} \omega_a \text{TV}(\nu_a \parallel \lambda_a) \right\}.$$

Proof Let π be a δ -correct ϵ -global DP BAI strategy. Let ν be a bandit instance and $\lambda \in \text{Alt}(\nu)$.

Let \mathbb{E} denote the expectation under $\mathbb{P}_{\nu, \pi}$, ie $\mathbb{E} \triangleq \mathbb{E}_{\nu, \pi}$.

By Lemma 12, we have that $6\epsilon \sum_{a=1}^K \mathbb{E}[N_a(\tau)] \text{TV}(\nu_a \parallel \lambda_a) \geq \text{kl}(1 - \delta, \delta)$.

Lemma 1 from Kaufmann et al. (2016) gives that $\sum_{a=1}^K \mathbb{E}[N_a(\tau)] \text{KL}(\nu_a \parallel \lambda_a) \geq \text{kl}(1 - \delta, \delta)$.

Since these two inequalities hold for all $\lambda \in \text{Alt}(\nu)$, we get

$$\begin{aligned} \text{kl}(1 - \delta, \delta) &\leq \inf_{\lambda \in \text{Alt}(\nu)} \min \left(6\epsilon \sum_{a=1}^K \mathbb{E}[N_a(\tau)] \text{TV}(\nu_a \parallel \lambda_a), \sum_{a=1}^K \mathbb{E}[N_a(\tau)] \text{KL}(\nu_a \parallel \lambda_a) \right) \\ &\stackrel{(a)}{=} \mathbb{E}[\tau] \inf_{\lambda \in \text{Alt}(\nu)} \min \left(6\epsilon \sum_{a=1}^K \frac{\mathbb{E}[N_a(\tau)]}{\mathbb{E}[\tau]} \text{TV}(\nu_a \parallel \lambda_a), \sum_{a=1}^K \frac{\mathbb{E}[N_a(\tau)]}{\mathbb{E}[\tau]} \text{KL}(\nu_a \parallel \lambda_a) \right) \\ &\stackrel{(b)}{\leq} \mathbb{E}[\tau] \left(\sup_{\omega \in \Sigma_K} \inf_{\lambda \in \text{Alt}(\nu)} \min \left(6\epsilon \sum_{a=1}^K \omega_a \text{TV}(\nu_a \parallel \lambda_a), \sum_{a=1}^K \omega_a \text{KL}(\nu_a \parallel \lambda_a) \right) \right). \end{aligned}$$

(a) is due to the fact that $\mathbb{E}[\tau]$ does not depend on λ . (b) is obtained by noting that the vector $(\omega_a)_{a \in [K]} \triangleq \left(\frac{\mathbb{E}_{\nu, \pi}[N_a(\tau)]}{\mathbb{E}_{\nu, \pi}[\tau]} \right)_{a \in [K]}$ belongs to the simplex Σ_K .

The theorem follows by noting that for $\delta \in (0, 1)$, $\text{kl}(1 - \delta, \delta) \geq \log(1/3\delta)$. \blacksquare

B.3.3 TV CHARACTERISTIC TIME FOR BERNOULLI INSTANCES: PROOF OF COROLLARY 14

Proposition 21 (TV characteristic time for Bernoulli instances) *Let ν be a bandit instance, i.e. such that $\nu_a = \text{Bernoulli}(\mu_a)$ and $\mu_1 > \mu_2 \geq \dots \geq \mu_K$. Let $\Delta_a \triangleq \mu_1 - \mu_a$ and $\Delta_{\min} \triangleq \min_{a \neq 1} \Delta_a$. We have that*

$$T_{\text{TV}}^*(\nu) = \frac{1}{\Delta_{\min}} + \sum_{a=2}^K \frac{1}{\Delta_a}, \quad \text{and} \quad \frac{1}{\Delta_{\min}} \leq T_{\text{TV}}^*(\nu) \leq \frac{K}{\Delta_{\min}}.$$

Proof Step 1: Let ν be a bandit instance, i.e. such that $\nu_a \triangleq \text{Bernoulli}(\mu_a)$ and $\mu_1 > \mu_2 \geq \dots \geq \mu_K$.

For the alternative bandit instance λ , we refer to the mean of arm a as ρ_a , i.e. $\lambda_a \triangleq \text{Bernoulli}(\rho_a)$.

By the definition of T_{TV}^* , we have that

$$\begin{aligned} (T_{\text{TV}}^*(\nu))^{-1} &= \sup_{\omega \in \Sigma_K} \inf_{\lambda \in \text{Alt}(\nu)} \sum_{a=1}^K \omega_a \text{TV}(\nu_a \parallel \lambda_a) \\ &\stackrel{(a)}{=} \sup_{\omega \in \Sigma_K} \min_{a \neq 1} \inf_{\lambda: \rho_a > \rho_1} \omega_1 |\mu_1 - \rho_1| + \omega_a |\mu_a - \rho_a| \\ &\stackrel{(b)}{=} \sup_{\omega \in \Sigma_K} \min_{a \neq 1} \min(\omega_1, \omega_a) \Delta_a \end{aligned}$$

$$\begin{aligned} & \stackrel{(c)}{=} \sup_{\omega \in \Sigma_K} \omega_1 \min_{a \neq 1} \min(1, \frac{\omega_a}{\omega_1}) \Delta_a \\ & \stackrel{(d)}{=} \sup_{(x_2, \dots, x_K) \in (\mathbb{R}^+)^{K-1}} \frac{\min_{a \neq 1} g_a(x_a)}{1 + x_2 + \dots + x_K}, \end{aligned}$$

where $g_a(x_a) \triangleq \min(1, x_a) \Delta_a$.

Equality (a) is obtained due to the fact that $\text{Alt}(\boldsymbol{\nu}) = \bigcup_{a \neq 1} \{\boldsymbol{\lambda} : \rho_a > \rho_1\}$, and for Bernoullis, $\text{TV}(\nu_a \parallel \lambda_a) = |\mu_a - \rho_a|$.

Equality (b) is true, since $\inf_{\boldsymbol{\lambda}: \rho_a > \rho_1} \omega_1 |\mu_1 - \rho_1| + \omega_a |\mu_a - \rho_a| = \min(\omega_1, \omega_a) \Delta_a$.

Equality (c) holds true, since $\omega_1 \neq 1$ (if $\omega_1 = 0$, the value of the objective is 0).

Equality (d) is obtained by the change of variable $x_a \triangleq \frac{\omega_a}{\omega_1}$

Step 2: Let $(x_2, \dots, x_K) \in (\mathbb{R}^+)^{K-1}$. By the definition of g_a , we have that

$$g_a(x_a) \leq x_a \Delta_a \quad \text{and} \quad g_a(x_a) \leq \Delta_a.$$

This leads to the inequalities

$$\min_{a \neq 1} g_a(x_a) \leq g_a(x_a) \leq x_a \Delta_a \quad \text{and} \quad \min_{a \neq 1} g_a(x_a) \leq \Delta_{\min}.$$

Thus,

$$\begin{aligned} \left(\min_{a \neq 1} g_a(x_a) \right) \left(\frac{1}{\Delta_{\min}} + \sum_{a=2}^K \frac{1}{\Delta_a} \right) &= \frac{\min_{a \neq 1} g_a(x_a)}{\Delta_{\min}} + \sum_{a=2}^K \frac{\min_{a \neq 1} g_a(x_a)}{\Delta_a} \\ &\leq 1 + \sum_{a=2}^K x_a. \end{aligned}$$

This means that for every $(x_2, \dots, x_K) \in (\mathbb{R}^+)^{K-1}$,

$$\frac{\min_{a \neq 1} g_a(x_a)}{1 + x_2 + \dots + x_K} \leq \frac{1}{\frac{1}{\Delta_{\min}} + \sum_{a=2}^K \frac{1}{\Delta_a}}.$$

Here, the upper bound is achievable for $x_a^* = \frac{\Delta_{\min}}{\Delta_a}$, since $g_a(x_a^*) = \Delta_{\min}$ for all $a \neq 1$.

This concludes that

$$T_{\text{TV}}^*(\boldsymbol{\nu})^{-1} = \frac{1}{\frac{1}{\Delta_{\min}} + \sum_{a=2}^K \frac{1}{\Delta_a}} \quad \implies \quad T_{\text{TV}}^*(\boldsymbol{\nu}) = \Delta_{\min} + \sum_{a=2}^K \frac{1}{\Delta_a}.$$

Step 3: The lower and upper bounds on $T_{\text{TV}}^*(\boldsymbol{\nu})$ follow from the fact that $\frac{1}{\Delta_a} \geq 0$ for all a , and $\frac{1}{\Delta_a} \leq \frac{1}{\Delta_{\min}}$ for all $a \neq 1$.

Hence, we conclude the proof. ■

B.3.4 ON THE TOTAL VARIATION DISTANCE AND THE HARDNESS OF PRIVACY

Our lower bound suggests that the hardness of the DP-FC-BAI problem is characterized by $T_{TV}^*(\boldsymbol{\nu})$, which is a total variation counterpart of the classic KL-based characteristic time $T_{KL}^*(\boldsymbol{\nu})$ in FC-BAI Garivier and Kaufmann (2016). The total variation distance appears to be the natural measure to quantify the hardness of privacy in other settings such as regret minimization Azize and Basu (2022), Karwa-Vadhan lemma Karwa and Vadhan (2018) and Differentially Private Assouad, Fano, and Le Cam Acharya et al. (2021). The high-level intuition is that: Pure DP can be seen as a multiplicative stability constraint of e^ϵ when one data point changes. With group privacy, if two datasets differ in d_{ham} points, then one incurs a factor $e^{d_{ham} \epsilon}$. Now, by sampling n i.i.d points from a distribution P and n i.i.d points from a distribution Q , the Karwa-Vadhan lemma states that the incurred factor is $e^{(nTV(P,Q)) \epsilon}$. This is proved by building a maximal coupling, which is the coupling that minimizes the Hamming distance in expectation. In brief, *the total variation naturally appears in lower bounds since it is the quantity that characterises the hardness of the optimal transport problem minimizing the hamming distance*, i.e $TV(P, Q) = \inf_{(X,Y) \sim (P,Q)} E(1_{X \neq Y})$. However, it is possible that the problem can be characterized by other f-divergences. Finally, one can always go from TV to KL using Pinsker's inequality, though that would always be less tight than the TV-based lower bound.

On the relation between $T_{TV}^(\boldsymbol{\nu})$ and $T_{KL}^*(\boldsymbol{\nu})$.* A direct application of Pinsker's inequality gives that $T_{TV}^*(\boldsymbol{\nu}) \geq \sqrt{2T_{KL}^*(\boldsymbol{\nu})}$. For completeness, we present here the exact calculations:

For every alternative mean parameter λ and every arm a , using Pinsker's inequality, we have that $d_{TV}(\mu_a, \lambda_a) \leq \sqrt{\frac{1}{2}d_{KL}(\mu_a, \lambda_a)}$. Therefore, for every allocation over arms ω , we have

$$\sum_a \omega_a d_{TV}(\mu_a, \lambda_a) \leq \sum_a \omega_a \sqrt{\frac{1}{2}d_{KL}(\mu_a, \lambda_a)} \leq \sqrt{\frac{1}{2} \sum_a \omega_a d_{KL}(\mu_a, \lambda_a)}.$$

Taking the supremum over the simplex and the infimum over the set of alternative mean parameters yields $T_{TV}^*(\boldsymbol{\nu})^{-1} \leq \sqrt{\frac{1}{2}T_{KL}^*(\boldsymbol{\nu})^{-1}}$. This concludes the proof.

Appendix C. Privacy analysis

We prove that AdaP-TT and AdaP-TT* satisfy ϵ -global DP. We first provide the privacy lemma that justifies using doubling and forgetting. Using the privacy lemma and the post-processing property of DP, we conclude the privacy analysis of AdaP-TT and AdaP-TT*.

C.1 Privacy Lemma for Non-overlapping Sequences

Lemma 22 (Privacy of non-overlapping sequence of empirical means) *Let \mathcal{M} be a mechanism that takes a set as input and outputs the private empirical mean, i.e.*

$$\mathcal{M}(\{r_i, \dots, r_j\}) \triangleq \frac{1}{j-i} \sum_{t=i}^j r_t + \text{Lap}\left(\frac{1}{(j-i)\epsilon}\right). \quad (17)$$

Let $\ell < T$ and $t_1, \dots, t_\ell, t_{\ell+1}$ be in $[1, T]$ such that $1 = t_1 < \dots < t_\ell < t_{\ell+1} - 1 = T$. Let's define the following mechanism

$$\mathcal{G} : \{r_1, \dots, r_T\} \rightarrow \bigotimes_{i=1}^{\ell} \mathcal{M}_{\{r_{t_i}, \dots, r_{t_{i+1}-1}\}} \quad (18)$$

In other words, \mathcal{G} is the mechanism we get by applying \mathcal{M} to the non-overlapping partition of the sequence $\{r_1, \dots, r_T\}$ according to $t_1 < \dots < t_\ell < t_{\ell+1}$, i.e.

$$\begin{pmatrix} r_1 \\ r_2 \\ \vdots \\ r_T \end{pmatrix} \xrightarrow{\mathcal{G}} \begin{pmatrix} \mu_1 \\ \vdots \\ \mu_\ell \end{pmatrix}$$

where $\mu_i \sim \mathcal{M}_{\{r_{t_i}, \dots, r_{t_{i+1}-1}\}}$.

For $r_t \in [0, 1]$, the mechanism \mathcal{G} is ϵ -DP.

Proof Let $r^T \triangleq (r_1, \dots, r_T)$ and $r'^T \triangleq (r'_1, \dots, r'_T)$ be two neighbouring reward sequences in $[0, 1]$. This implies that $\exists j \in [1, T]$ such that $r_j \neq r'_j$ and $\forall t \neq j, r_t = r'_t$.

Let ℓ' be such that $t_{\ell'} \leq j \leq t_{\ell'+1} - 1$, and follows the convention that $t_0 = 1$ and $t_{\ell'+1} = T + 1$.

Let $\mu \triangleq (\mu_1, \dots, \mu_\ell)$ a fixed sequence of outcomes. Then,

$$\frac{\mathbb{P}(\mathcal{G}(r^T) = \mu)}{\mathbb{P}(\mathcal{G}(r'^T) = \mu)} = \frac{\mathbb{P}(\mathcal{M}(\{r_{t_{\ell'}}, \dots, r_{t_{\ell'+1}-1}\}) = \mu_{\ell'})}{\mathbb{P}(\mathcal{M}(\{r'_{t_{\ell'}}, \dots, r'_{t_{\ell'+1}-1}\}) = \mu_{\ell'})} \leq e^\epsilon,$$

where the last inequality holds true because \mathcal{M} satisfies ϵ -DP following Theorem 2. \blacksquare

C.2 Privacy Analysis of AdaP-TT and AdaP-TT*

Theorem 23 (Privacy analysis) For rewards in $[0, 1]$, AdaP-TT and AdaP-TT* satisfy ϵ -global DP.

Remark 24 The following proof is valid for any BAI strategy that only uses the DAF(ϵ) to estimate the means.

Proof Let $T \geq 1$. Let $\mathbf{d}^T = \{\mathbf{x}_1, \dots, \mathbf{x}_T\}$ and $\mathbf{d}'^T = \{\mathbf{d}'_1, \dots, \mathbf{d}'_T\}$ two neighbouring reward tables in $(\mathbb{R}^K)^T$. Let $j \in [1, T]$ such that, for all $t \neq j, d_t = d'_t$.

We also fix a sequence of sampled actions $\underline{a}^T = \{a_1, \dots, a_T\} \in [K]^T$ and a recommended action $\hat{a} \in K$.

Let π be a BAI strategy that only uses DAF(ϵ) to estimate the means, i.e. either AdaP-TT or AdaP-TT*.

We want to show that: $\pi(\underline{a}^T, \hat{a}, T \mid \mathbf{d}^T) \leq e^\epsilon \pi(\underline{a}^T, \hat{a}, T \mid \mathbf{d}'^T)$.

The main idea is that the change of reward in the j -th reward only affects the empirical mean computed in one episode, which is made private using the Laplace Mechanism and Lemma 22.

Step 1. Sequential decomposition of the output probability

We observe that due to the sequential nature of the interaction, the output probability can be decomposed to a part that depends on $\underline{\mathbf{d}}^{j-1} \triangleq \{\mathbf{x}_1, \dots, \mathbf{x}_{j-1}\}$, which is identical for both $\underline{\mathbf{d}}^T$ and $\underline{\mathbf{d}}'^T$ and a second conditional part on the history.

Specifically, we have that

$$\begin{aligned} \pi(\underline{a}^T, \hat{a}, T \mid \underline{\mathbf{d}}^T) &\triangleq \text{Rec}_{T+1}(\hat{a} \mid \mathcal{H}_T) S_{T+1}(\top \mid \mathcal{H}_T) \prod_{t=1}^T S_t(a_t \mid \mathcal{H}_{t-1}) \\ &\triangleq \mathcal{P}_{\underline{\mathbf{d}}^{j-1}}^\pi(\underline{a}^j) \mathcal{P}_{\underline{\mathbf{d}}}^\pi(a_{>j}, \hat{a}, T \mid \underline{a}^j) \end{aligned}$$

where

- $a_{>j} \triangleq (a_{j+1}, \dots, a_T)$
- $\mathcal{P}_{\underline{\mathbf{d}}^{j-1}}^\pi(\underline{a}^j) \triangleq \prod_{t=1}^j S_t(a_t \mid \mathcal{H}_{t-1})$
- $\mathcal{P}_{\underline{\mathbf{d}}}^\pi(a_{>j}, \hat{a}, T \mid \underline{a}^j) \triangleq \text{Rec}_{T+1}(\hat{a} \mid \mathcal{H}_T) S_{T+1}(\top \mid \mathcal{H}_T) \prod_{t=j+1}^T S_t(a_t \mid \mathcal{H}_{t-1})$

Similarly

$$\pi(\underline{a}^T, \hat{a}, T \mid \underline{\mathbf{d}}'^T) \triangleq \mathcal{P}_{\underline{\mathbf{d}}^{j-1}}^\pi(\underline{a}^j) \mathcal{P}_{\underline{\mathbf{d}}'}^\pi(a_{>j}, \hat{a}, T \mid \underline{a}^j)$$

since $\underline{\mathbf{d}}'^{j-1} = \underline{\mathbf{d}}^{j-1}$.

Which means that

$$\frac{\pi(\underline{a}^T, \hat{a}, T \mid \underline{\mathbf{d}}^T)}{\pi(\underline{a}^T, \hat{a}, T \mid \underline{\mathbf{d}}'^T)} = \frac{\mathcal{P}_{\underline{\mathbf{d}}}^\pi(a_{>j}, \hat{a}, T \mid \underline{a}^j)}{\mathcal{P}_{\underline{\mathbf{d}}'}^\pi(a_{>j}, \hat{a}, T \mid \underline{a}^j)} \quad (19)$$

Step 2. The adaptive episodes are the same, before step j

Let ℓ such that $t_\ell \leq j < t_{\ell+1}$ when π interacts with $\underline{\mathbf{d}}^T$. Let us call it $\psi_{\underline{\mathbf{d}}^T}^\pi(j) \triangleq \ell$.

Similarly, let ℓ' such that $t_{\ell'} \leq j < t_{\ell'+1}$ when π interacts with $\underline{\mathbf{d}}'^T$. Let us call it $\psi_{\underline{\mathbf{d}}'^T}^\pi(j) \triangleq \ell'$.

Since $\psi_{\underline{\mathbf{d}}^T}^\pi(j)$ only depends on $\underline{\mathbf{d}}^{j-1}$, which is identical for $\underline{\mathbf{d}}^T$ and $\underline{\mathbf{d}}'^T$, we have that $\psi_{\underline{\mathbf{d}}^T}^\pi(j) = \psi_{\underline{\mathbf{d}}'^T}^\pi(j)$ with probability 1.

We call ξ_j the last **time-step** of the episode $\psi_{\underline{\mathbf{d}}^T}^\pi(j)$, i.e $\xi_j \triangleq t_{\psi_{\underline{\mathbf{d}}^T}^\pi(j)+1} - 1$.

Step 3. Private sufficient statistics

Let $r_t \triangleq \underline{\mathbf{d}}_{t, a_t}^T$, be the reward corresponding to the action a_t in the table $\underline{\mathbf{d}}^T$. Similarly, $r'_t \triangleq \underline{\mathbf{d}}'_{t, a_t}$ for $\underline{\mathbf{d}}'^T$.

Let us define $L_j \triangleq \mathcal{G}_{\{r_1, \dots, r_{\xi_j}\}}$ and $L'_j \triangleq \mathcal{G}_{\{r'_1, \dots, r'_{\xi_j}\}}$, where \mathcal{G} is defined as in Eq. 18, using the same episodes for d and d' . In other words, L_j is the list of private empirical means computed on a non-overlapping sequence of rewards before step ξ_j .

Using the forgetting structure of π , there exists a randomised mapping $f_{\underline{\mathbf{d}}_{>\xi_j}}$ such that $\mathcal{P}_{\underline{\mathbf{d}}}^\pi(\cdot | \underline{\mathbf{a}}^j) = f_{\underline{\mathbf{d}}_{>\xi_j}} \circ L_j$ and $\mathcal{P}_{\underline{\mathbf{d}'}}^\pi(\cdot | \underline{\mathbf{a}}^j) = f_{\underline{\mathbf{d}}_{>\xi_j}} \circ L'_j$.

In other words, the interaction of π with $\underline{\mathbf{d}}$ and $\underline{\mathbf{d}'}$ from step $\xi_j + 1$ until T only depends on the sufficient statistics L_j , which summarises what happened before ξ_j , and the new inputs $\underline{\mathbf{d}}_{>\xi_j}$, which are the same for $\underline{\mathbf{d}}$ and $\underline{\mathbf{d}'}$.

Step 4. Concluding with Lemma 22 and the post-processing lemma

Since rewards are in $[0, 1]$, using Lemma 22, we have that \mathcal{G} is ϵ -DP.

Since $\mathcal{P}_{\underline{\mathbf{d}}}^\pi(\cdot | \underline{\mathbf{a}}^j)$ is just a post-processing of the output of \mathcal{G} , we have that

$$\frac{\mathcal{P}_{\underline{\mathbf{d}}}^\pi(a_{>j}, \hat{a}, T | \underline{\mathbf{a}}^j)}{\mathcal{P}_{\underline{\mathbf{d}'}}^\pi(a_{>j}, \hat{a}, T | \underline{\mathbf{a}}^j)} \leq e^\epsilon,$$

and Eq. (19) concludes the proof. \blacksquare

Appendix D. Globally Differentially Private GLR Stopping Rules

After studying the non-private GLR stopping rule with phases (Appendix D.1), we study the private GLR stopping rule with non-private transportation costs $W_{a,b}^G$ in Appendix D.2 (Lemma 16) and with adapted transportation costs $W_{a,b}^{G,\epsilon}$ in Appendix D.3 (Lemma 18).

D.1 Non-private GLR Stopping Rule with Per-arm Phases

Before accounting for the privacy (i.e. Laplace noise), we first highlight the price of $\text{DAF}(\epsilon)$ for $\epsilon = +\infty$. This stopping condition is only evaluated at the beginning of each phase for each arm since it involves quantities that are fixed until we switch phase again, and it recommends $\hat{a}_n = \arg \max_{a \in [K]} \hat{\mu}_{k_{n,a},a}$ which is the best arm for the non-private empirical means. Lemma 25 yields a threshold function ensuring δ -correctness.

Lemma 25 *Let $\delta \in (0, 1)$. Let $s > 1$, ζ be the Riemann ζ function, $c_{a,b}^G$ as in (4) and $k(x) = \log_2 x + 2$. Combining the $\text{DAF}(\epsilon)$ estimator for $\epsilon = +\infty$ with the GLR stopping rule with $W_{a,b}^G$ as in (3) and the stopping threshold $c_{a,b}^G(\omega, \delta(\zeta(s)^2 k(\omega_a)^s k(\omega_b)^s)^{-1})$ yields a δ -correct algorithm for σ -sub-Gaussian distributions regardless of the sampling rule.*

Proof The non-private GLR stopping rule matches the one used for Gaussian bandits. Proving δ -correctness of a GLR stopping rule is done by leveraging concentration results.

Lemma 26 (Theorem 9 in Kaufmann and Koolen (2021)) *Let ν be a sub-Gaussian bandit with means $\boldsymbol{\mu} \in \mathbb{R}^K$ and variance proxy σ . Let $S \subseteq [K]$ and $x > 0$.*

$$\mathbb{P}_\nu \left(\exists n \in \mathbb{N}, \sum_{a \in S} \frac{N_{n,a}}{2\sigma^2} (\mu_{n,a} - \mu_a)^2 > \sum_{a \in S} 2 \log(4 + \log(N_{n,a})) + |S| \mathcal{C}_G \left(\frac{x}{|S|} \right) \right) \leq e^{-x},$$

where \mathcal{C}_G is defined in Kaufmann and Koolen (2021) as

$$\mathcal{C}_G(x) \triangleq \min_{\lambda \in [1/2, 1]} \frac{g_G(\lambda) + x}{\lambda} \text{ and } g_G(\lambda) \triangleq 2\lambda - 2\lambda \log(4\lambda) + \log \zeta(2\lambda) - \frac{1}{2} \log(1 - \lambda). \quad (20)$$

Here, ζ is the Riemann ζ function and $\mathcal{C}_G(x) \approx x + \log(x)$.

We consider the concentration event $\mathcal{E}_\delta^{(1)} = \bigcap_{a \neq a^*} \bigcap_{n \in \mathbb{N}} \mathcal{E}_\delta^{(1)}(a, n)$ with $\mathcal{E}_\delta^{(1)}(a, n) =$

$$\left\{ \frac{\tilde{N}_{k_n, a}}{2\sigma^2} (\hat{\mu}_{k_n, a} - \mu_a)^2 + \frac{\tilde{N}_{k_n, a^*, a^*}}{2\sigma^2} (\hat{\mu}_{k_n, a^*, a^*} - \mu_{a^*})^2 < c_{a, a^*}^G(\tilde{N}_{k_n}, \frac{\delta}{\zeta(s)^2 k_{n, a}^s k_{n, a^*}^s}) \right\}. \quad (21)$$

For all $a \neq a^*$ and all $(k_a, k_{a^*}) \in \mathbb{N}^2$, the estimators $(\hat{\mu}_{k_c, c})_{c \in \{a, a^*\}}$ are based solely on the observations collected for arm a (resp. arm a^*) between times $n \in \{T_{k_a-1}(a), \dots, T_{k_a}(a)-1\}$ (resp. $n \in \{T_{k_{a^*}-1}(a^*), \dots, T_{k_{a^*}}(a^*)-1\}$) with local counts $(\tilde{N}_{k_c, c})_{c \in \{a, a^*\}}$, i.e. dropping past observations. Using a direct union bound, we obtain that $\mathbb{P}_\nu((\mathcal{E}_\delta^{(1)})^c)$ is smaller than

$$\begin{aligned} & \sum_{a \neq a^*} \sum_{k_a, k_{a^*} \in \mathbb{N}} \mathbb{P}_\nu \left(\frac{\tilde{N}_{k_a, a}}{2\sigma^2} (\hat{\mu}_{k_a, a} - \mu_a)^2 + \frac{\tilde{N}_{k_{a^*}, a^*}}{2\sigma^2} (\hat{\mu}_{k_{a^*}, a^*} - \mu_{a^*})^2 \geq c_{a, a^*}^G(\tilde{N}_{k_n}, \frac{\delta}{\zeta(s)^2 k_a^s k_{a^*}^s}) \right) \\ & \leq \frac{\delta}{K-1} \frac{1}{\zeta(s)^2} \sum_{a \neq a^*} \sum_{(k_a, k_{a^*}) \in \mathbb{N}^2} \frac{1}{(k_a k_{a^*})^s} = \delta. \end{aligned}$$

where the last inequality uses Lemma 26 for all $a \neq a^*$ and all $(k_a, k_{a^*}) \in \mathbb{N}^2$. Therefore,

$$\mathbb{P}_\nu(\tau_\delta < +\infty, \hat{a}_{\tau_\delta} \neq a^*) \leq \delta + \mathbb{P}_\nu(\mathcal{E}_\delta^{(1)} \cap \{\tau_\delta < +\infty, \hat{a}_{\tau_\delta} \neq a^*\}).$$

Under $\mathcal{E}_\delta^{(1)} \cap \{\tau_\delta < +\infty, \hat{a}_{\tau_\delta} \neq a^*\}$, we have $\hat{a}_{\tau_\delta} = \arg \max_{b \in [K]} \hat{\mu}_{k_{\tau_\delta}, a, a} \neq a^*$ and

$$\begin{aligned} c_{\hat{a}_{\tau_\delta}, a^*}^G(\tilde{N}_{k_{\tau_\delta}}, \frac{\delta}{\zeta(s)^2 k_{\tau_\delta, a}^s k_{\tau_\delta, a^*}^s}) & \leq \frac{(\hat{\mu}_{k_{\tau_\delta}, \hat{a}_{\tau_\delta}, \hat{a}_{\tau_\delta}} - \hat{\mu}_{k_{\tau_\delta}, a^*, a^*})^2}{2\sigma^2(1/\tilde{N}_{k_{\tau_\delta}, \hat{a}_{\tau_\delta}, \hat{a}_{\tau_\delta}} + 1/\tilde{N}_{k_{\tau_\delta}, a^*, a^*})} \\ & = \inf_{y \geq x} \left\{ \frac{\tilde{N}_{k_{\tau_\delta}, \hat{a}_{\tau_\delta}, \hat{a}_{\tau_\delta}}}{2\sigma^2} (\hat{\mu}_{k_{\tau_\delta}, \hat{a}_{\tau_\delta}, \hat{a}_{\tau_\delta}} - x)^2 + \frac{\tilde{N}_{k_{\tau_\delta}, a^*, a^*}}{2\sigma^2} (\hat{\mu}_{k_{\tau_\delta}, a^*, a^*} - y)^2 \right\} \\ & \leq \frac{\tilde{N}_{k_{\tau_\delta}, \hat{a}_{\tau_\delta}, \hat{a}_{\tau_\delta}}}{2\sigma^2} (\hat{\mu}_{k_{\tau_\delta}, \hat{a}_{\tau_\delta}, \hat{a}_{\tau_\delta}} - \mu_{\hat{a}_{\tau_\delta}})^2 + \frac{\tilde{N}_{k_{\tau_\delta}, a^*, a^*}}{2\sigma^2} (\hat{\mu}_{k_{\tau_\delta}, a^*, a^*} - \mu_{a^*})^2 \\ & < c_{\hat{a}_{\tau_\delta}, a^*}^G(\tilde{N}_{k_{\tau_\delta}}, \frac{\delta}{\zeta(s)^2 k_{\tau_\delta, a}^s k_{\tau_\delta, a^*}^s}). \end{aligned}$$

This is a contradiction, hence $\mathcal{E}_\delta^{(1)} \cap \{\tau_\delta < +\infty, \hat{a}_{\tau_\delta} \neq a^*\} = \emptyset$. This concludes the proof. \blacksquare

D.2 Private GLR with Non-private Transportation Cost: Proof of Lemma 16

The proof of Lemma 16 is similar as the one detailed in Appendix D.1, with the added difficulty of controlling the Laplace noise. We consider the concentration event $\mathcal{E}_\delta = \mathcal{E}_{\delta/2}^{(1)} \cap \mathcal{E}_{\delta/2}^{(2)}$ where $\mathcal{E}_\delta^{(1)}$ as in (21) and $\mathcal{E}_\delta^{(2)} = \bigcap_{a \in [K]} \bigcap_{n \in \mathbb{N}} \mathcal{E}_\delta^{(2)}(a, n)$ with

$$\mathcal{E}_\delta^{(2)}(a, n) = \left\{ \epsilon \tilde{N}_{k_n, a, a} |Y_{k_n, a, a}| < \log \left(\frac{K \zeta(s) k_{n, a}^s}{\delta} \right) \right\}. \quad (22)$$

Since $Y_{k_{n,a},a} \sim \text{Lap}\left((\epsilon \tilde{N}_{k_{n,a},a})^{-1}\right)$, we have that $\tilde{N}_{k_{n,a},a} | Y_{k_{n,a},a} \sim \mathcal{E}(\epsilon)$ for all $a \in [K]$ and all $n \in \mathbb{N}$, where $\mathcal{E}(\cdot)$ denotes the exponential distribution. Using concentration results for exponential distribution, a direct union bound yields that $\mathbb{P}_\nu((\mathcal{E}_\delta^{(2)})^c) \leq \delta$, hence

$$\mathbb{P}_\nu(\tau_\delta < +\infty, \hat{a}_{\tau_\delta} \neq a^*) \leq \delta + \mathbb{P}_\nu(\mathcal{E}_\delta \cap \{\tau_\delta < +\infty, \hat{a}_{\tau_\delta} \neq a^*\}).$$

Under $\mathcal{E}_\delta \cap \{\tau_\delta < +\infty, \hat{a}_{\tau_\delta} \neq a^*\}$, we have $\hat{a}_{\tau_\delta} = \arg \max_{b \in [K]} \tilde{\mu}_{k_{\tau_\delta},a} \neq a^*$ and

$$\begin{aligned} c_{\hat{a}_{\tau_\delta}, a^*}^{G, \epsilon}(\tilde{N}_{k_{\tau_\delta}}, \delta) &\leq \frac{\tilde{N}_{k_{\tau_\delta}, \hat{a}_{\tau_\delta}, \hat{a}_{\tau_\delta}}}{2\sigma^2} (\tilde{\mu}_{k_{\tau_\delta}, \hat{a}_{\tau_\delta}, \hat{a}_{\tau_\delta}} - \mu_{\hat{a}_{\tau_\delta}})^2 + \frac{\tilde{N}_{k_{\tau_\delta}, a^*, a^*}}{2\sigma^2} (\tilde{\mu}_{k_{\tau_\delta}, a^*, a^*} - \mu_{a^*})^2 \\ &\leq \frac{\tilde{N}_{k_{\tau_\delta}, \hat{a}_{\tau_\delta}, \hat{a}_{\tau_\delta}}}{\sigma^2} (\hat{\mu}_{k_{\tau_\delta}, \hat{a}_{\tau_\delta}, \hat{a}_{\tau_\delta}} - \mu_{\hat{a}_{\tau_\delta}})^2 + \frac{\tilde{N}_{k_{\tau_\delta}, a^*, a^*}}{\sigma^2} (\hat{\mu}_{k_{\tau_\delta}, a^*, a^*} - \mu_{a^*})^2 \\ &\quad + \frac{\tilde{N}_{k_{\tau_\delta}, \hat{a}_{\tau_\delta}, \hat{a}_{\tau_\delta}}}{\sigma^2} Y_{k_{\tau_\delta}, \hat{a}_{\tau_\delta}, \hat{a}_{\tau_\delta}}^2 + \frac{\tilde{N}_{k_{\tau_\delta}, a^*, a^*}}{\sigma^2} Y_{k_{\tau_\delta}, a^*, a^*}^2 \\ &< 2c_{\hat{a}_{\tau_\delta}, a^*}^G(\tilde{N}_{k_{\tau_\delta}}, \frac{\delta}{2\zeta(s)^2 k_{\tau_\delta, a}^s k_{\tau_\delta, a^*}^s}) + \frac{1}{\epsilon^2 \sigma^2} \sum_{c \in \{a, a^*\}} \frac{1}{\tilde{N}_{k_{\tau_\delta}, c, c}} \left(\log \frac{2K\zeta(s)k_{\tau_\delta, c}^s}{\delta} \right)^2. \end{aligned}$$

where we used that $\tilde{\mu}_{k_{\tau_\delta}, a} = \hat{\mu}_{k_{\tau_\delta}, a} + Y_{k_{\tau_\delta}, a}$ and $(x - y)^2 \leq 2x^2 + 2y^2$. This is a contradiction, hence $\mathcal{E}_\delta \cap \{\tau_\delta < +\infty, \hat{a}_{\tau_\delta} \neq a^*\} = \emptyset$. This concludes the proof.

D.3 Private GLR with Adapted Transportation Cost: Proof of Lemma 18

The proof of Lemma 18 is similar as the one detailed in Appendix D.2. The main difference lies in the considered transportation costs, namely $W_{a,b}^{G, \epsilon}$ instead of $W_{a,b}^G$.

Lemma 27 (Lemma 28 in Jourdan et al. (2023)) *Let $\delta \in (0, 1)$. For all $x \geq 1$, let $\bar{W}_{-1}(x) = -W_{-1}(-e^{-x})$ (see Lemma 31), where W_{-1} is the negative branch of the Lambert W function. Let $c(x, \delta) = \frac{1}{2} \bar{W}_{-1}(2 \log(K/\delta) + 4 \log(4 + \log x) + 1/2)$. Consider σ -sub-Gaussian bandits with means $\mu \in \mathbb{R}^K$. Then,*

$$\mathbb{P}\left(\exists n \in \mathbb{N}, \exists a \in [K], \frac{N_{n,a}}{2\sigma^2} (\mu_{n,a} - \mu_a)^2 > c(N_{n,a}, \delta)\right) \leq \delta.$$

Recall that $h(\tilde{N}_{k_{n,a},a}, \delta) = c(\tilde{N}_{k_{n,a},a}, \frac{\delta}{3\zeta(s)k_{n,a}^s})$. We use the concentration event $\mathcal{E}_\delta = \mathcal{E}_{\delta/3}^{(1)} \cap \mathcal{E}_{\delta/3}^{(2)} \cap \mathcal{E}_{\delta/3}^{(3)}$ where $\mathcal{E}_\delta^{(1)}$ as in (21), $\mathcal{E}_\delta^{(2)}$ as in (22) and $\mathcal{E}_\delta^{(3)} = \bigcap_{a \in [K]} \bigcap_{n \in \mathbb{N}} \mathcal{E}_\delta^{(3)}(a, n)$ with

$$\mathcal{E}_\delta^{(3)}(a, n) = \left\{ \frac{\tilde{N}_{k_{n,a},a}}{2\sigma^2} (\hat{\mu}_{k_{n,a},a} - \mu_a)^2 < h(\tilde{N}_{k_{n,a},a}, 3\delta) \right\}. \quad (23)$$

Using Lemma 27, a direct union bound yields that $\mathbb{P}_\nu((\mathcal{E}_\delta^{(3)})^c) \leq \delta$, hence

$$\mathbb{P}_\nu(\tau_\delta < +\infty, \hat{a}_{\tau_\delta} \neq a^*) \leq \delta + \mathbb{P}_\nu(\mathcal{E}_\delta \cap \{\tau_\delta < +\infty, \hat{a}_{\tau_\delta} \neq a^*\}).$$

Under $\mathcal{E}_\delta \cap \{\tau_\delta < +\infty, \hat{a}_{\tau_\delta} \neq a^*\}$, we have $\hat{a}_{\tau_\delta} = \arg \max_{b \in [K]} \tilde{\mu}_{k_{\tau_\delta},a} \neq a^*$.

Case 1. Under $\mathcal{E}_\delta \cap \{\tau_\delta < +\infty, \hat{a}_{\tau_\delta} \neq a^*, (\hat{\mu}_{k_{\tau_\delta, \hat{a}_{\tau_\delta}}, \hat{a}_{\tau_\delta}} - \hat{\mu}_{k_{\tau_\delta, a^*}, a^*})_+ < 3\epsilon\}$, we have

$$\begin{aligned}
 & \frac{1}{2} c_{\hat{a}_{\tau_\delta, a^*}}^{G, \epsilon}(\tilde{N}_{k_{\tau_\delta}}, \delta) + \frac{\sqrt{2}}{\epsilon \sigma} \sum_{c \in \{\hat{a}_{\tau_\delta, a^*}\}} \sqrt{\frac{h(\tilde{N}_{k_{\tau_\delta, c}}, \delta)}{\tilde{N}_{k_{\tau_\delta, c}}} \log\left(\frac{3K\zeta(s)k_{\tau_\delta, c}^s}{\delta}\right)} \\
 & \leq \frac{\tilde{N}_{k_{\tau_\delta, \hat{a}_{\tau_\delta}}, \hat{a}_{\tau_\delta}}}{2\sigma^2} (\tilde{\mu}_{k_{\tau_\delta, \hat{a}_{\tau_\delta}}, \hat{a}_{\tau_\delta}} - \mu_{\hat{a}_{\tau_\delta}})^2 + \frac{\tilde{N}_{k_{\tau_\delta, a^*}, a^*}}{2\sigma^2} (\tilde{\mu}_{k_{\tau_\delta, a^*}, a^*} - \mu_{a^*})^2 \\
 & = \frac{\tilde{N}_{k_{\tau_\delta, \hat{a}_{\tau_\delta}}, \hat{a}_{\tau_\delta}}}{2\sigma^2} (\hat{\mu}_{k_{\tau_\delta, \hat{a}_{\tau_\delta}}, \hat{a}_{\tau_\delta}} - \mu_{\hat{a}_{\tau_\delta}})^2 + \frac{\tilde{N}_{k_{\tau_\delta, a^*}, a^*}}{2\sigma^2} (\hat{\mu}_{k_{\tau_\delta, a^*}, a^*} - \mu_{a^*})^2 \\
 & \quad + \frac{\tilde{N}_{k_{\tau_\delta, \hat{a}_{\tau_\delta}}, \hat{a}_{\tau_\delta}}}{2\sigma^2} Y_{k_{\tau_\delta, \hat{a}_{\tau_\delta}}, \hat{a}_{\tau_\delta}}^2 + \frac{\tilde{N}_{k_{\tau_\delta, a^*}, a^*}}{2\sigma^2} Y_{k_{\tau_\delta, a^*}, a^*}^2 \\
 & \quad + \frac{\tilde{N}_{k_{\tau_\delta, \hat{a}_{\tau_\delta}}, \hat{a}_{\tau_\delta}}}{\sigma^2} Y_{k_{\tau_\delta, \hat{a}_{\tau_\delta}}, \hat{a}_{\tau_\delta}} (\hat{\mu}_{k_{\tau_\delta, \hat{a}_{\tau_\delta}}, \hat{a}_{\tau_\delta}} - \mu_{\hat{a}_{\tau_\delta}}) + \frac{\tilde{N}_{k_{\tau_\delta, a^*}, a^*}}{\sigma^2} Y_{k_{\tau_\delta, a^*}, a^*} (\hat{\mu}_{k_{\tau_\delta, a^*}, a^*} - \mu_{a^*}) \\
 & < \frac{1}{2} c_{\hat{a}_{\tau_\delta, a^*}}^{G, \epsilon}(\tilde{N}_{k_{\tau_\delta}}, \delta) + \frac{\sqrt{2}}{\epsilon \sigma} \sum_{c \in \{\hat{a}_{\tau_\delta, a^*}\}} \sqrt{\frac{h(\tilde{N}_{k_{\tau_\delta, c}}, \delta)}{\tilde{N}_{k_{\tau_\delta, c}}} \log\left(\frac{3K\zeta(s)k_{\tau_\delta, c}^s}{\delta}\right)}.
 \end{aligned}$$

This is a contradiction, hence $\mathcal{E}_\delta \cap \{\tau_\delta < +\infty, \hat{a}_{\tau_\delta} \neq a^*, (\hat{\mu}_{k_{\tau_\delta, \hat{a}_{\tau_\delta}}, \hat{a}_{\tau_\delta}} - \hat{\mu}_{k_{\tau_\delta, a^*}, a^*})_+ < 3\epsilon\} = \emptyset$.

Case 2. Under $\mathcal{E}_\delta \cap \{\tau_\delta < +\infty, \hat{a}_{\tau_\delta} \neq a^*, (\hat{\mu}_{k_{\tau_\delta, \hat{a}_{\tau_\delta}}, \hat{a}_{\tau_\delta}} - \hat{\mu}_{k_{\tau_\delta, a^*}, a^*})_+ \geq 3\epsilon\}$, we have

$$\begin{aligned}
 & \frac{3}{\sigma^2} \log\left(\frac{3K\zeta(s) \max_{c \in \{\hat{a}_{\tau_\delta, a^*}\}} k_{\tau_\delta, c}}{\delta}\right) + \frac{3\epsilon}{\sqrt{2}\sigma} \sum_{c \in \{\hat{a}_{\tau_\delta, a^*}\}} \sqrt{\tilde{N}_{k_{\tau_\delta, c}} h(\tilde{N}_{k_{\tau_\delta, c}}, \delta)} \\
 & \leq \frac{3\epsilon(\tilde{\mu}_{k_{\tau_\delta, \hat{a}_{\tau_\delta}}, \hat{a}_{\tau_\delta}} - \tilde{\mu}_{k_{\tau_\delta, a^*}, a^*})}{2\sigma^2(1/\tilde{N}_{k_{\tau_\delta, \hat{a}_{\tau_\delta}}, \hat{a}_{\tau_\delta}} + 1/\tilde{N}_{k_{\tau_\delta, a^*}, a^*})} \\
 & \leq \frac{3\epsilon}{2\sigma^2} \min\{\tilde{N}_{k_{\tau_\delta, \hat{a}_{\tau_\delta}}, \hat{a}_{\tau_\delta}}, \tilde{N}_{k_{\tau_\delta, a^*}, a^*}\} (\tilde{\mu}_{k_{\tau_\delta, \hat{a}_{\tau_\delta}}, \hat{a}_{\tau_\delta}} - \tilde{\mu}_{k_{\tau_\delta, a^*}, a^*}) \\
 & = \frac{3\epsilon}{2\sigma^2} \inf_{y \geq x} \left\{ \tilde{N}_{k_{\tau_\delta, \hat{a}_{\tau_\delta}}, \hat{a}_{\tau_\delta}} |\tilde{\mu}_{k_{\tau_\delta, \hat{a}_{\tau_\delta}}, \hat{a}_{\tau_\delta}} - x| + \tilde{N}_{k_{\tau_\delta, a^*}, a^*} |\tilde{\mu}_{k_{\tau_\delta, a^*}, a^*} - y| \right\} \\
 & \leq \frac{3\epsilon}{2\sigma^2} \tilde{N}_{k_{\tau_\delta, \hat{a}_{\tau_\delta}}, \hat{a}_{\tau_\delta}} |\tilde{\mu}_{k_{\tau_\delta, \hat{a}_{\tau_\delta}}, \hat{a}_{\tau_\delta}} - \mu_{\hat{a}_{\tau_\delta}}| + \frac{3\epsilon}{2\sigma^2} \tilde{N}_{k_{\tau_\delta, a^*}, a^*} |\tilde{\mu}_{k_{\tau_\delta, a^*}, a^*} - \mu_{a^*}| \\
 & \leq \frac{3\epsilon}{2\sigma^2} \tilde{N}_{k_{\tau_\delta, \hat{a}_{\tau_\delta}}, \hat{a}_{\tau_\delta}} |\hat{\mu}_{k_{\tau_\delta, \hat{a}_{\tau_\delta}}, \hat{a}_{\tau_\delta}} - \mu_{\hat{a}_{\tau_\delta}}| + \frac{3\epsilon}{2\sigma^2} \tilde{N}_{k_{\tau_\delta, a^*}, a^*} |\hat{\mu}_{k_{\tau_\delta, a^*}, a^*} - \mu_{a^*}| \\
 & \quad + \frac{3\epsilon}{2\sigma^2} \tilde{N}_{k_{\tau_\delta, \hat{a}_{\tau_\delta}}, \hat{a}_{\tau_\delta}} |Y_{k_{\tau_\delta, \hat{a}_{\tau_\delta}}, \hat{a}_{\tau_\delta}}| + \frac{3\epsilon}{2\sigma^2} \tilde{N}_{k_{\tau_\delta, a^*}, a^*} |Y_{k_{\tau_\delta, a^*}, a^*}| \\
 & < \frac{3}{\sigma^2} \log\left(\frac{3K\zeta(s) \max_{c \in \{\hat{a}_{\tau_\delta, a^*}\}} k_{\tau_\delta, c}}{\delta}\right) + \frac{3\epsilon}{\sqrt{2}\sigma} \sum_{c \in \{\hat{a}_{\tau_\delta, a^*}\}} \sqrt{\tilde{N}_{k_{\tau_\delta, c}} h(\tilde{N}_{k_{\tau_\delta, c}}, \delta)}.
 \end{aligned}$$

This is a contradiction, hence $\mathcal{E}_\delta \cap \{\tau_\delta < +\infty, \hat{a}_{\tau_\delta} \neq a^*, (\hat{\mu}_{k_{\tau_\delta, \hat{a}_{\tau_\delta}}, \hat{a}_{\tau_\delta}} - \hat{\mu}_{k_{\tau_\delta, a^*}, a^*})_+ \geq 3\epsilon\} = \emptyset$.

Summary. Putting both cases together yields the result.

Appendix E. Expected Sample Complexity of AdaP-TT and AdaP-TT*

Let $\beta \in (0, 1)$, $\epsilon \in \mathbb{R}_+^*$, and ν be a bandit instance consisting of σ -sub-Gaussian distributions with distinct means $\mu \in \mathbb{R}^K$, i.e. $\min_{a \neq b} |\mu_a - \mu_b| > 0$. For conciseness, we denote $\Delta_a \triangleq \mu_{a^*} - \mu_a$, $\Delta_{\min} \triangleq \min_{a \neq a^*} \Delta_a$, and $\Delta_{\max} \triangleq \max_{a \neq a^*} \Delta_a$. For Gaussian, we define the unique β -optimal allocations $\omega_{\text{KL},\beta}^*(\nu) = \{(\omega_{\beta,a}^*)_{a \in [K]}\}$ and $\omega_{\epsilon,\beta}^*(\nu) = \{(\omega_{\epsilon,\beta,a}^*)_{a \in [K]}\}$ as

$$\omega_{\text{KL},\beta}^*(\nu) \triangleq \arg \max_{\omega \in \Sigma_K, \omega_{a^*} = \beta} \min_{a \neq a^*} \frac{\Delta_a^2}{1/\beta + 1/\omega_a} \quad , \quad \omega_{\epsilon,\beta}^*(\nu) \triangleq \arg \max_{\omega \in \Sigma_K, \omega_{a^*} = \beta} \min_{a \neq a^*} \frac{\Delta_a \min\{3\epsilon, \Delta_a\}}{1/\beta + 1/\omega_a} . \quad (24)$$

At equilibrium, we have equality of the transportation costs (see Jourdan and Degenne (2024) for example), namely

$$\forall a \neq a^* , \quad \frac{\Delta_a^2}{1/\beta + 1/\omega_{\beta,a}^*} = 2\sigma^2 T_{\text{KL},\beta}^*(\nu)^{-1} \quad , \quad \frac{\Delta_a \min\{3\epsilon, \Delta_a\}}{1/\beta + 1/\omega_{\epsilon,\beta,a}^*} = 2\sigma^2 T_{\epsilon,\beta}^*(\nu)^{-1} . \quad (25)$$

Our proof follows the unified sample complexity analysis of Top Two algorithms from Jourdan et al. (2022).

Let $\gamma > 0$. Let $\omega \in \Sigma_K$ be any allocation over arms such that $\min_a \omega_a > 0$. We denote by $T_{\mu,\gamma}(\omega)$ the *convergence time* towards ω , which is a random variable quantifying the number of samples required for the global empirical allocations $N_n/(n-1)$ to be γ -close to ω for any subsequent time, namely

$$T_{\mu,\gamma}(\omega) \triangleq \inf \left\{ T \geq 1 \mid \forall n \geq T, \left\| \frac{N_n}{n-1} - \omega \right\|_{\infty} \leq \gamma \right\} . \quad (26)$$

As the AdaP-TT and AdaP-TT* algorithms share the same leader, all results solely on the leader applies to both of them. As the AdaP-TT and AdaP-TT* algorithms consider a TC challenger with different transportation costs, all results involving the challenger should be slightly modified. Except if specified otherwise, all the results presented in the following hold for both algorithms.

The rest of Appendix E is organised as follows. After recalling some technical results (Appendix E.1), we prove sufficient exploration (Appendix E.2) Second, we prove that convergence towards the β -optimal allocation (Appendix E.3) in finite time. Third, we explicit the cost of doubling and forgetting (Appendix E.4). Finally, we conclude the proof of Theorems 17 and 19 (Appendix E.5).

E.1 Technical Results

Before delving into the proofs, we first recall some useful technical results.

Doubling trick. Due to the doubling, the growth of the counts is exponential (Lemma 28).

Lemma 28 *For all $(a, k) \in [K] \times \mathbb{N}$ s.t. $\mathbb{E}_{\nu}[T_k(a)] < +\infty$, $N_{T_k(a),a} = 2^{k-1}$ and $\tilde{N}_{k,a} = 2^{k-2}$.*

Proof Let $a \in [K]$. After initialisation, we have $k = 1$, $T_1(a) = K + 1$ and $N_{T_1(a),a} = 1$. Using the definition of the phase switch, it is direct to see that $N_{T_2(a),a} = 2$ and $\tilde{N}_{2,a} = 1$ when $\mathbb{E}_{\nu}[T_2(a)] < +\infty$.

Now, we proceed by recurrence. Suppose that $N_{T_k(a),a} = 2^{k-1}$ and $\tilde{N}_{k,a} = 2^{k-2}$ when $\mathbb{E}_\nu[T_k(a)] < +\infty$. If $\mathbb{E}_\nu[T_{k+1}(a)] < +\infty$, then it means that the phase k ends for arm a almost surely. Since we sample only one arm at each round, at the beginning of phase $k+1$ for arm a , we have $N_{T_{k+1}(a),a} = 2N_{T_k(a),a} = 2^k$ by using the definition of the phase switch. Then, we have directly that $\tilde{N}_{k+1,a} = N_{T_{k+1}(a),a} - N_{T_k(a),a} = 2^k - 2^{k-1} = 2^{k-1}$. \blacksquare

Tracking. We denote by $N_{n,b}^a \triangleq \sum_{t \in [n-1]} \mathbb{1}(B_t = a, a_t = C_t = b)$ the number of times the arm b was pulled while the arm a was the leader, and by $L_{n,a} \triangleq \sum_{t \in [n-1]} \mathbb{1}(B_t = a)$ the number of times arm a was the leader.

Lemma 29 (Lemma 2.2 in Jourdan and Degenne (2024)) *For all $n > K$ and all $a \in [K]$, we have $-1/2 \leq N_{n,a}^a - \beta L_{n,a} \leq 1$.*

Concentration results. In order to control the randomness of $(\tilde{\mu}_{k_a,a})_{a \in [K]}$, we use a standard concentration result on the empirical mean of sub-Gaussian random variables and on sub-exponential observations (Lemma 30). Since Bernoulli distributions are $1/2$ -sub-Gaussian and the absolute value of a Laplace is an exponential distribution, Lemma 30 applies to our setting.

Lemma 30 *There exists a sub-Gaussian random variable W_μ such that, almost surely,*

$$\forall a \in [K], \forall k_a \in \mathbb{N}, \quad |\hat{\mu}_{k_a,a} - \mu_a| \leq W_\mu \sqrt{\frac{\log(e + \tilde{N}_{k_a,a})}{\tilde{N}_{k_a,a}}}.$$

There exists a sub-exponential random variable W_ϵ such that, almost surely,

$$\forall a \in [K], \forall k_a \in \mathbb{N}, \quad |Y_{k_a,a}| \leq W_\epsilon \frac{\log(e + k_a)}{\tilde{N}_{k_a,a}}.$$

In particular, any random variable which is polynomial in (W_ϵ, W_μ) has a finite expectation.

Proof The first part is a known result, e.g. Appendix E.2 in Jourdan et al. (2022). Let

$$W_\epsilon \triangleq \sup_{a \in [K]} \sup_{k_a \in \mathbb{N}} \frac{\tilde{N}_{k_a,a} |Y_{k_a,a}|}{\log(e + k_a)}.$$

By definition, we have that, almost surely,

$$\forall a \in [K], \forall k_a \in \mathbb{N}, \quad |Y_{k_a,a}| \leq W_\epsilon \frac{\log(e + k_a)}{\tilde{N}_{k_a,a}}.$$

Since $\tilde{N}_{k,i} |Y_{k,i}| \sim \mathcal{E}(\epsilon)$, Lemma 72 in Jourdan et al. (2022) yields that W_ϵ is a sub-exponential random variable. Since W_μ is sub-Gaussian and W_ϵ is a sub-exponential, any random variable which is polynomial in (W_ϵ, W_μ) has a finite expectation. \blacksquare

Inversion results. Lemma 31 gathers properties on the function \overline{W}_{-1} , which is used in the literature to obtain concentration results.

Lemma 31 (Jourdan et al. (2023)) *Let $\bar{W}_{-1}(x) \triangleq -W_{-1}(-e^{-x})$ for all $x \geq 1$, where W_{-1} is the negative branch of the Lambert W function. The function \bar{W}_{-1} is increasing on $(1, +\infty)$ and strictly concave on $(1, +\infty)$. In particular, $\bar{W}'_{-1}(x) = \left(1 - \frac{1}{\bar{W}_{-1}(x)}\right)^{-1}$ for all $x > 1$. Then, for all $y \geq 1$ and $x \geq 1$,*

$$\bar{W}_{-1}(y) \leq x \iff y \leq x - \log(x).$$

Moreover, for all $x > 1$,

$$x + \log(x) \leq \bar{W}_{-1}(x) \leq x + \log(x) + \min\left\{\frac{1}{2}, \frac{1}{\sqrt{x}}\right\}.$$

Lemma 32 is an inversion result to upper bound a time, which is implicitly defined. It is a direct consequence of Lemma 31.

Lemma 32 *Let \bar{W}_{-1} defined in Lemma 31. Let $A > 0$, $B > 0$ such that $B/A + \log A > 1$ and*

$$C(A, B) = \sup\{x \mid x < A \log x + B\}.$$

Then, $C(A, B) < h_1(A, B)$ with $h_1(z, y) = z\bar{W}_{-1}(y/z + \log z)$.

Proof Since $B/A + \log A > 1$, we have $C(A, B) \geq A$, hence

$$C(A, B) = \sup\{x \mid x < A \log(x) + B\} = \sup\{x \geq A \mid x < A \log(x) + B\}.$$

Using Lemma 31 yields that

$$x \geq A \log x + B \iff \frac{x}{A} - \log\left(\frac{x}{A}\right) \geq \frac{B}{A} + \log A \iff x \geq A\bar{W}_{-1}\left(\frac{B}{A} + \log A\right).$$

■

E.2 Sufficient Exploration

The first step of in the generic analysis of Top Two algorithms Jourdan et al. (2022) consists in showing sufficient exploration. The main idea is that, if there are still undersampled arms, either the leader or the challenger will be among them. Therefore, after a long enough time, no arm can still be undersampled. We emphasise that there are multiple ways to select the leader/challenger pair in order to ensure sufficient exploration. Therefore, other choices of leader/challenger pair would yield similar results.

Given an arbitrary phase $p \in \mathbb{N}$, we define the sampled enough set, i.e. the arms having reached phase p , and the arm with highest mean in this set (when not empty) as

$$S_n^p = \{a \in [K] \mid N_{n,a} \geq 2^{p-1}\} \quad \text{and} \quad a_n^* = \arg \max_{a \in S_n^p} \mu_a. \quad (27)$$

Since $\min_{a \neq b} |\mu_a - \mu_b| > 0$, a_n^* is unique. Let $p \in \mathbb{N}$ such that $(p-1)/4 \in \mathbb{N}$. We define the highly and the mildly under-sampled sets as

$$U_n^p \triangleq \{a \in [K] \mid N_{n,a} < 2^{(p-1)/2}\} \quad \text{and} \quad V_n^p \triangleq \{a \in [K] \mid N_{n,a} < 2^{3(p-1)/4}\}. \quad (28)$$

Those arms have not reached phase $(p-1)/2$ and phase $3(p-1)/4$, respectively.

Lemma 33 shows that, when the leader is sampled enough, it is the arm with highest true mean among the sampled enough arms.

Lemma 33 *Let S_n^p and a_n^* as in (27). There exists p_0 with $\mathbb{E}_\nu[\exp(\alpha p_0)] < +\infty$ for all $\alpha > 0$ such that if $p \geq p_0$, for all n such that $S_n^p \neq \emptyset$, $B_n \in S_n^p$ implies that $B_n = a_n^* = \arg \max_{a \in S_n^p} \tilde{\mu}_{k_{n,a},a}$.*

Proof Let p_0 to be specified later. Let $p \geq p_0$. Let $n \in \mathbb{N}$ such that $S_n^p \neq \emptyset$, where S_n^p and a_n^* as in Equation (27). Let $(k_{n,a})_{a \in [K]}$ be the phases indices for all arms. Since $N_{n,a} \geq 2^{p-1}$ for all $a \in S_n^p$, we have $k_{n,a} \geq p$ and $\tilde{N}_{k_{n,a},a} \geq 2^{p-2}$ by using Lemma 28. Using Lemma 30, we obtain that

$$\begin{aligned} \tilde{\mu}_{k_{n,a_n^*},a_n^*} &\geq \mu_{a_n^*} - W_\mu \sqrt{\frac{\log(e + 2^{p-2})}{2^{p-2}}} - W_\epsilon \frac{\log(e + p)}{2^{p-2}}, \\ \tilde{\mu}_{k_{n,a},a} &\leq \mu_a + W_\mu \sqrt{\frac{\log(e + 2^{p-2})}{2^{p-2}}} + W_\epsilon \frac{\log(e + p)}{2^{p-2}}, \quad \forall a \in S_n^p \setminus \{a_n^*\}. \end{aligned}$$

Here, we use that $x \rightarrow \log(e+x)/x$ is decreasing.

Let $\bar{\Delta}_{\min} = \min_{a \neq b} |\mu_a - \mu_b|$. By assumption on the considered instances, we know that $\bar{\Delta}_{\min} > 0$. Let $p_1 = \lceil \log_2(X_1 - e) \rceil + 2$ and $p_2 = \lceil \log_2((X_2 - e - 2) \log 2 + 1) \rceil + 2$ with

$$\begin{aligned} X_1 &= \sup \left\{ x > 1 \mid x \leq 64 \bar{\Delta}_{\min}^{-2} W_\mu^2 \log x + e \right\} \leq h_1(64 \bar{\Delta}_{\min}^{-2} W_\mu^2, e), \\ X_2 &= \sup \left\{ x > 1 \mid x \leq \frac{8}{\log 2} \bar{\Delta}_{\min}^{-1} W_\epsilon \log x + e + 2 - 1/\log 2 \right\} \leq h_1(8 \bar{\Delta}_{\min}^{-1} W_\epsilon / \log 2, 4), \end{aligned}$$

where we used Lemma 32, and h_1 defined therein. Then, for all $p \in \mathbb{N}$ such that $p \geq \max\{p_1, p_2\} + 1$ and all $n \in \mathbb{N}$ such that $S_n^p \neq \emptyset$, we have $\tilde{\mu}_{k_{n,a_n^*},a_n^*} \geq \mu_{a_n^*} - \bar{\Delta}_{\min}/4$ and $\tilde{\mu}_{k_{n,a},a} \leq \mu_a + \bar{\Delta}_{\min}/4$ for all $a \in S_n^p \setminus \{a_n^*\}$, hence $a_n^* = \arg \max_{a \in [K]} \tilde{\mu}_{k_{n,a},a}$.

We have, for all $\alpha \in \mathbb{R}_+$,

$$\exp(\alpha p_1) \leq e^{3\alpha} (X_1 - e)^{\alpha / \log 2} \quad \text{hence} \quad \mathbb{E}_\nu[\exp(\alpha p_1)] < +\infty,$$

where we used Lemma 30 and $h_1(x, e) \sim_{x \rightarrow +\infty} x \log x$ to obtain that $\exp(\alpha p_1)$ is at most polynomial in W_μ . Likewise, we obtain that $\mathbb{E}_\nu[\exp(\alpha p_2)] < +\infty$ for all $\alpha \in \mathbb{R}_+$.

Let us define the UCB indices by $I_{k_{n,a},a} = \tilde{\mu}_{k_{n,a},a} + \sqrt{k_{n,a}/\tilde{N}_{k_{n,a},a}} + k_{n,a}/(\epsilon \tilde{N}_{k_{n,a},a})$. Using the above, we have

$$\begin{aligned} I_{k_{n,a_n^*},a_n^*} &\geq \mu_{a_n^*} - W_\mu \sqrt{\frac{\log(e + 2^{p-2})}{2^{p-2}}} - W_\epsilon \frac{\log(e + p)}{2^{p-2}}, \\ \forall a \in S_n^p \setminus \{a_n^*\}, \quad I_{k_{n,a},a} &\leq \mu_a + W_\mu \sqrt{\frac{\log(e + 2^{p-2})}{2^{p-2}}} + W_\epsilon \frac{\log(e + p)}{2^{p-2}} + \sqrt{\frac{p}{2^{p-2}}} + \frac{p}{\epsilon 2^{p-2}}, \end{aligned}$$

where we used Lemma 28 and the fact that $x \rightarrow \log(e+x)/x$ and $x \rightarrow x2^{2-x}$ are decreasing function for $x \geq 2$. Let $p_3 = \lceil \log_2 X_3 \rceil + 2$ and $p_4 = \lceil \log_2 X_4 \rceil + 2$ with

$$\begin{aligned} X_3 &= \sup \left\{ x > 1 \mid x \leq 64\bar{\Delta}_{\min}^{-2}(\log_2 x + 2) \right\} \leq h_1(64\bar{\Delta}_{\min}^{-2}/\log 2, 128\bar{\Delta}_{\min}^{-2}), \\ X_4 &= \sup \left\{ x > 1 \mid x \leq 8\epsilon^{-1}\bar{\Delta}_{\min}^{-1}(\log_2 x + 2) \right\} \leq h_1(8\epsilon^{-1}\bar{\Delta}_{\min}^{-1}/\log 2, 16\bar{\Delta}_{\min}^{-1}\epsilon^{-1}), \end{aligned}$$

where we used Lemma 32, and h_1 defined therein. We highlight that (p_3, p_4) are deterministic values, hence their expectation is finite. Then, for all $p \in \mathbb{N}$ such that $p \geq p_0 = \max\{p_1, p_2, p_3, p_4\} + 1$ and all $n \in \mathbb{N}$ such that $S_n^p \neq \emptyset$, we have $I_{k_n, a_n^*, a_n^*} \geq \mu_{a_n^*} - \bar{\Delta}_{\min}/4$ and $I_{k_n, a, a} \leq \mu_a + \bar{\Delta}_{\min}/2$ for all $a \in S_n^p \setminus \{a_n^*\}$, hence $a_n^* = B_n$ since we have $B_n = \arg \max_{a \in [K]} I_{k_n, a, a}$.

Since we have $\mathbb{E}_\nu[\exp(\alpha p_0)] < +\infty$ for all $\alpha \in \mathbb{R}_+$, this concludes the proof. \blacksquare

Lemma 34 shows that the transportation costs between the sampled enough arms with largest true means and the other sampled enough arms are increasing fast enough.

Lemma 34 *Let S_n^p and a_n^* as in (27). There exists p_1 with $\mathbb{E}_\nu[\exp(\alpha p_1)] < +\infty$ for all $\alpha > 0$ such that if $p \geq p_1$, for all n such that $S_n^p \neq \emptyset$, for all $b \in S_n^p \setminus \{a_n^*\}$, we have*

$$\begin{aligned} [\text{AdaP-TT}] \quad & \frac{\tilde{\mu}_{k_n, a_n^*, a_n^*} - \tilde{\mu}_{k_n, b, b}}{\sqrt{1/\tilde{N}_{k_n, a_n^*, a_n^*} + 1/\tilde{N}_{k_n, b, b}}} \geq 2^{p/2} C_\mu, \\ [\text{AdaP-TT}^*] \quad & \frac{(\tilde{\mu}_{k_n, a_n^*, a_n^*} - \tilde{\mu}_{k_n, b, b}) \min\{3\epsilon, \tilde{\mu}_{k_n, a_n^*, a_n^*} - \tilde{\mu}_{k_n, b, b}\}}{1/\tilde{N}_{k_n, a_n^*, a_n^*} + 1/\tilde{N}_{k_n, b, b}} \geq 2^p C_\mu, \end{aligned}$$

where $C_\mu > 0$ is a problem dependent constant.

Proof Let p_1 to be specified later. Let $p \geq p_1$. Let $n \in \mathbb{N}$ such that $S_n^p \neq \emptyset$, where S_n^p and a_n^* as in Equation (27). Let $(k_{n,a})_{a \in [K]}$ be the phases indices for all arms. Since $N_{n,a} \geq 2^{p-1}$ for all $a \in S_n^p$, we have $k_{n,a} \geq p$ and $\tilde{N}_{k_{n,a}, a} \geq 2^{p-2}$ by using Lemma 28. Let $\bar{\Delta}_{\min} = \min_{a \neq b} |\mu_a - \mu_b|$, which satisfies $\bar{\Delta}_{\min} > 0$ by assumption on the instance considered.

Using Lemma 30, for all $b \in S_n^p \setminus \{a_n^*\}$, we obtain

$$\tilde{\mu}_{k_n, a_n^*, a_n^*} - \tilde{\mu}_{k_n, b, b} \geq \bar{\Delta}_{\min} - W_\mu \sqrt{\frac{\log(e + 2^{p-2})}{2^{p-4}}} - W_\epsilon \frac{\log(e + p)}{2^{p-3}}.$$

Let $p_3 = \lceil \log_2((X_3 - e)/4) \rceil + 4$ and $p_2 = \lceil \log_2((X_2 - e - 3) \log 2 + 1) \rceil + 3$ with

$$\begin{aligned} X_3 &= \sup \left\{ x > 1 \mid x \leq 64\bar{\Delta}_{\min}^{-2} W_\mu^2 \log x + e \right\} \leq h_1(64\bar{\Delta}_{\min}^{-2} W_\mu^2, e), \\ X_2 &= \sup \left\{ x > 1 \mid x \leq 4\bar{\Delta}_{\min}^{-1} W_\epsilon \log x + e + 3 - 1/\log 2 \right\} \leq h_1(4\bar{\Delta}_{\min}^{-1} W_\epsilon, 5), \end{aligned}$$

where we used Lemma 32, and h_1 defined therein. Then, for all $p \in \mathbb{N}$ such that $p \geq p_1 = \max\{p_3, p_2\} + 1$ and all $n \in \mathbb{N}$ such that $S_n^p \neq \emptyset$, we have, for all $b \in S_n^p \setminus \{a_n^*\}$,

$$\tilde{\mu}_{k_n, a_n^*, a_n^*} - \tilde{\mu}_{k_n, b, b} \geq \bar{\Delta}_{\min}/2.$$

As in the proof of Lemma 33, we obtain that $\mathbb{E}_\nu[\exp(\alpha p_1)] < +\infty$ for all $\alpha \in \mathbb{R}_+$.

Then, for all $b \in S_n^p \setminus \{a_n^*\}$, we have

$$\frac{\tilde{\mu}_{k_n, a_n^*, a_n^*} - \tilde{\mu}_{k_n, b, b}}{\sqrt{1/\tilde{N}_{k_n, a_n^*, a_n^*} + 1/\tilde{N}_{k_n, b, b}}} \geq 2^{p/2} \frac{\bar{\Delta}_{\min}}{2^{5/2}},$$

where we used that $\min\{\tilde{N}_{k_n, a_n^*, a_n^*}, \tilde{N}_{k_n, b, b}\} \geq 2^{p-2}$. Setting $C_\mu = \bar{\Delta}_{\min}/2^{5/2}$ yields the first result.

The second result is obtained similarly by taking $C_\mu = \frac{\bar{\Delta}_{\min}}{16} \min\{3\epsilon, \frac{\bar{\Delta}_{\min}}{2}\}$ \blacksquare

Lemma 35 shows that the transportation costs between sampled enough arms and under-sampled arms are not increasing too fast.

Lemma 35 *Let S_n^p be as in (27). For all $p \geq 1$ and all n such that $S_n^p \neq \emptyset$, for all $a \in S_n^p$ and $b \notin S_n^p$,*

$$\begin{aligned} [\text{AdaP-TT}] \quad & \frac{\tilde{\mu}_{k_n, a, a} - \tilde{\mu}_{k_n, b, b}}{\sqrt{1/\tilde{N}_{k_n, a, a} + 1/\tilde{N}_{k_n, b, b}}} \leq 2^{p/2} D_\mu + 2W_\mu \sqrt{\log(e + 2^{p-2})} + 2W_\epsilon \log(e + p), \\ [\text{AdaP-TT}^*] \quad & \frac{(\tilde{\mu}_{k_n, a, a} - \tilde{\mu}_{k_n, b, b}) \min\{3\epsilon, \tilde{\mu}_{k_n, a, a} - \tilde{\mu}_{k_n, b, b}\}}{1/\tilde{N}_{k_n, a, a} + 1/\tilde{N}_{k_n, b, b}} \\ & \leq 2^p D_\mu + 8W_\mu^2 \log(e + 2^{p-2}) + 8W_\epsilon^2 \log(e + p)^2, \end{aligned}$$

where $D_\mu > 0$ is a problem dependent constant and (W_μ, W_ϵ) are the random variables defined in Lemma 30.

Proof Let $p \geq 1$. Let $n \in \mathbb{N}$ such that $S_n^p \neq \emptyset$, where S_n^p as in Equation (27). Let $(k_{n,a})_{a \in [K]}$ be the phases indices for all arms. Since $N_{n,a} \geq 2^{p-1}$ for all $a \in S_n^p$, we have $k_{n,a} \geq p$ and $\tilde{N}_{k_n, a, a} \geq 2^{p-2}$ by using Lemma 28. Likewise, $N_{n,a} < 2^{p-1}$ for all $a \notin S_n^p$, we have $k_{n,a} < p$ and $\tilde{N}_{k_n, a, a} < 2^{p-2}$. Let $\bar{\Delta}_{\max} = \min_{a \neq b} |\mu_a - \mu_b|$, which satisfies $\bar{\Delta}_{\max} > 0$ by assumption on the instance considered. Using Lemma 30, for all $a \in S_n^p$ and $b \notin S_n^p$, we obtain

$$\begin{aligned} \frac{\tilde{\mu}_{k_n, a, a} - \tilde{\mu}_{k_n, b, b}}{\sqrt{1/\tilde{N}_{k_n, a, a} + 1/\tilde{N}_{k_n, b, b}}} & \leq \sqrt{\tilde{N}_{k_n, b, b}} (\tilde{\mu}_{k_n, a, a} - \tilde{\mu}_{k_n, b, b}) \\ & \leq \sqrt{\tilde{N}_{k_n, b, b}} (\mu_a - \mu_b) + 2W_\mu \sqrt{\log(e + \tilde{N}_{k_n, b, b})} + 2W_\epsilon \frac{\log(e + k_{n,b})}{\sqrt{\tilde{N}_{k_n, b, b}}} \\ & \leq 2^{(p-2)/2} \bar{\Delta}_{\max} + 2W_\mu \sqrt{\log(e + 2^{p-2})} + 2W_\epsilon \log(e + p) \end{aligned}$$

where we used that $\tilde{N}_{k_n, b, b} \geq 1$, $k_{n,b} < p$, $\tilde{N}_{k_n, b, b} < 2^{p-2} \leq \tilde{N}_{k_n, a, a}$ and $x \rightarrow \log(e + x)/x$ is decreasing. Taking $D_\mu = \bar{\Delta}_{\max}/2$ yields the first result.

The proof of the second result follows along the same line by noting that this transportation cost is lower than the other:

$$\frac{(\tilde{\mu}_{k_n, a, a} - \tilde{\mu}_{k_n, b, b}) \min\{3\epsilon, \tilde{\mu}_{k_n, a, a} - \tilde{\mu}_{k_n, b, b}\}}{1/\tilde{N}_{k_n, a, a} + 1/\tilde{N}_{k_n, b, b}}$$

$$\begin{aligned}
 &\leq 2\tilde{N}_{k_n,b}(\hat{\mu}_{k_n,a} - \hat{\mu}_{k_n,b})^2 + 2\tilde{N}_{k_n,b}(Y_{k_n,a} - Y_{k_n,b})^2 \\
 &\leq 2^p \bar{\Delta}_{\max}^2 + 8W_\mu^2 \log(e + 2^{p-2}) + 8W_\epsilon^2 \log(e + p)^2.
 \end{aligned}$$

Taking $D_\mu = \bar{\Delta}_{\max}^2$ yields the result. \blacksquare

Lemma 36 shows that the challenger is mildly undersampled if the leader is not mildly undersampled.

Lemma 36 *Let V_n^p be as in Equation (28). There exists p_2 with $\mathbb{E}_\nu[\exp(\alpha p_2)] < +\infty$ for all $\alpha > 0$ such that if $p \geq p_2$, for all n such that $U_n^p \neq \emptyset$, $B_n \notin V_n^p$ implies $C_n \in V_n^p$.*

Proof Let p_2 to be specified later. Let $p \geq p_2$. Let $n \in \mathbb{N}$ such that $U_n^p \neq \emptyset$ and $V_n^p \neq [K]$, where $U_n^p \subseteq V_n^p$ are defined in Equation (28). In the following, we suppose that $B_n \notin V_n^p$.

Let $(k_{n,a})_{a \in [K]}$ be the phases indices for all arms. Let p_0 as in Lemma 33. Let $b_n^* = \arg \max_{b \notin V_n^p} \mu_b$. Then, for all $p \geq 4p_0/3 - 1/3$ and all n such that $B_n \notin V_n^p$, Lemma 33 yields that $B_n = b_n^* = \arg \max_{a \notin V_n^p} \tilde{\mu}_{k_{n,a},a}$.

Let p_1 and C_μ as in Lemma 34, and D_μ as in Lemma 35. Then, for all $p \geq \frac{4}{3} \max\{p_0, p_1\} - 1/3$ and all n such that $B_n \notin V_n^p$, we have $B_n = b_n^*$ and

$$\begin{aligned}
 \forall b \notin V_n^p, \quad &\frac{\tilde{\mu}_{k_n,b_n^*,b_n^*} - \tilde{\mu}_{k_n,b,b}}{\sqrt{1/\tilde{N}_{k_n,b_n^*,b_n^*} + 1/\tilde{N}_{k_n,b,b}}} \geq 2^{(3p+1)/8} C_\mu, \\
 \forall b \in U_n^p, \quad &\frac{\tilde{\mu}_{k_n,b_n^*,b_n^*} - \tilde{\mu}_{k_n,b,b}}{\sqrt{1/\tilde{N}_{k_n,b_n^*,b_n^*} + 1/\tilde{N}_{k_n,b,b}}} \leq 2^{(p+1)/4} D_\mu + 2W_\mu \sqrt{\log(e + 2^{(p+1)/2-2})} \\
 &\quad + 2W_\epsilon \log(e + (p+1)/2),
 \end{aligned}$$

where we used the first results of Lemmas 34 and 35. Let $p_3 = 16 \lceil \log_2(2D_\mu/C_\mu) \rceil + 1$, then we have $2^{(p-1)/16} > \frac{D_\mu}{C_\mu}$ for all $p \geq p_3$. Let $p_4 = \frac{16}{9} \lceil \log_2 X_4 \rceil + 25$ and $p_5 = \frac{32}{9} \lceil \log_2 X_5 \rceil + 7$ where

$$\begin{aligned}
 X_4 &= \sup \left\{ x > 1 \mid x \leq \frac{W_\mu^2}{C_\mu^2} \log(e + x^{8/9} 2^{25/18-3/4}) \right\}, \\
 X_5 &= \sup \left\{ x > 1 \mid x \leq \frac{2W_\epsilon}{C_\mu} \log(e + 4 + 32 \log_2(x)/18) \right\}.
 \end{aligned}$$

As in the proof of Lemma 33, using Lemma 30 yields that $\mathbb{E}_\nu[\exp(\alpha p_4)] < +\infty$ and $\mathbb{E}_\nu[\exp(\alpha p_5)] < +\infty$ for all $\alpha \in \mathbb{R}_+$. Let $p_2 = \max\{p_3, p_4, p_5, 4 \max\{p_0, p_1\}/3 - 1/3\} + 1$. Then, we have shown that for all $p \geq p_2$, for all n such that $B_n \notin V_n^p$, we have $B_n = b_n^*$ and

$$\min_{b \notin V_n^p} \frac{\tilde{\mu}_{k_n,b_n^*,b_n^*} - \tilde{\mu}_{k_n,b,b}}{\sqrt{1/\tilde{N}_{k_n,b_n^*,b_n^*} + 1/\tilde{N}_{k_n,b,b}}} > \max_{b \in U_n^p} \frac{\tilde{\mu}_{k_n,b_n^*,b_n^*} - \tilde{\mu}_{k_n,b,b}}{\sqrt{1/\tilde{N}_{k_n,b_n^*,b_n^*} + 1/\tilde{N}_{k_n,b,b}}},$$

Therefore, by definition of the TC challenger $C_n = \arg \min_{b \neq b_n^*} \frac{\tilde{\mu}_{k_n,b_n^*,b_n^*} - \tilde{\mu}_{k_n,b,b}}{\sqrt{1/\tilde{N}_{k_n,b_n^*,b_n^*} + 1/\tilde{N}_{k_n,b,b}}}$, we obtain that $C_n \in V_n^p$. Otherwise, there would be a contradiction given that we assumed that

$U_n^p \neq \emptyset$. Given all the condition exhibited above, it is direct to see that $\mathbb{E}_\nu[\exp(\alpha p_2)] < +\infty$ for all $\alpha > 0$. This concludes the proof for the AdaP-TT algorithm.

For the AdaP-TT* algorithm, the proof is done similarly based on the second results of Lemmas 34 and 35. As above, we can construct \tilde{p}_3 , with $\mathbb{E}_\nu[\exp(\alpha \tilde{p}_3)] < +\infty$ for all $\alpha \in \mathbb{R}_+$, such that for all $p \geq \tilde{p}_3$, we have

$$2^{(3p+1)/4} C_\mu > 2^{(p+1)/2} D_\mu + 8W_\mu^2 \log(e + 2^{(p+1)/2-2}) + 8W_\epsilon^2 \log(e + (p+1)/2)^2.$$

Let $\tilde{p}_2 = \max\{\tilde{p}_3, 4 \max\{p_0, p_1\}/3 - 1/3\} + 1$. Then, we have shown that for all $p \geq \tilde{p}_2$, for all n such that $B_n \notin V_n^p$, we have $B_n = b_n^*$ and

$$\begin{aligned} & \min_{b \notin V_n^p} \frac{(\tilde{\mu}_{k_n, b_n^*, b_n^*} - \tilde{\mu}_{k_n, b, b}) \min\{3\epsilon, \tilde{\mu}_{k_n, b_n^*, b_n^*} - \tilde{\mu}_{k_n, b, b}\}}{1/\tilde{N}_{k_n, b_n^*, b_n^*} + 1/\tilde{N}_{k_n, b, b}} \\ & > \max_{b \in U_n^p} \frac{(\tilde{\mu}_{k_n, b_n^*, b_n^*} - \tilde{\mu}_{k_n, b, b}) \min\{3\epsilon, \tilde{\mu}_{k_n, b_n^*, b_n^*} - \tilde{\mu}_{k_n, b, b}\}}{1/\tilde{N}_{k_n, b_n^*, b_n^*} + 1/\tilde{N}_{k_n, b, b}}. \end{aligned}$$

Then, we conclude similarly by using the definition of the TC challenger. \blacksquare

Lemma 37 shows that all the arms are sufficient explored for large enough n .

Lemma 37 *There exists N_0 with $\mathbb{E}_\nu[N_0] < +\infty$ such that for all $n \geq N_0$ and all $a \in [K]$,*

$$N_{n,a} \geq \sqrt{n/K} \quad \text{and} \quad k_{n,a} \geq \frac{\log(n/K)}{2 \log 2} + 1.$$

Proof Let p_0 and p_2 as in Lemmas 33 and 36. Combining Lemmas 33 and 36 yields that, for all $p \geq p_3 = \max\{p_2, 4p_0/3 - 1/3\}$ and all n such that $U_n^p \neq \emptyset$, we have $B_n \in V_n^p$ or $C_n \in V_n^p$. We have $\mathbb{E}_\nu[2^{p_2}] < +\infty$. We have $2^{p-1} \geq K 2^{3(p-1)/4}$ for all $p \geq p_4 = 4\lceil \log_2 K \rceil + 1$. Let $p \geq \max\{p_3, p_4\}$.

Suppose towards contradiction that $U_{K 2^{p-1}}^p$ is not empty. Then, for any $1 \leq t \leq K 2^{p-1}$, U_t^p and V_t^p are non empty as well. Using the pigeonhole principle, there exists some $a \in [K]$ such that $N_{2^{p-1}, a} \geq 2^{3(p-1)/4}$. Thus, we have $|V_{2^{p-1}}^p| \leq K - 1$. Our goal is to show that $|V_{2^p}^p| \leq K - 2$. A sufficient condition is that one arm in $V_{2^{p-1}}^p$ is pulled at least $2^{3(p-1)/4}$ times between 2^{p-1} and $2^p - 1$.

Case 1. Suppose there exists $a \in V_{2^{p-1}}^p$ such that $L_{2^p, a} - L_{2^{p-1}, a} \geq \frac{2^{3(p-1)/4}}{\beta} + 3/(2\beta)$. Using Lemma 29, we obtain

$$N_{2^p, a}^a - N_{2^{p-1}, a}^a \geq \beta(L_{2^p, a} - L_{2^{p-1}, a}) - 3/2 \geq 2^{3(p-1)/4},$$

hence a is sampled $2^{3(p-1)/4}$ times between 2^{p-1} and $2^p - 1$.

Case 2. Suppose that for all $a \in V_{2^{p-1}}^p$, we have $L_{2^p, a} - L_{2^{p-1}, a} < 2^{3(p-1)/4}/\beta + 3/(2\beta)$. Then,

$$\sum_{a \notin V_{2^{p-1}}^p} (L_{2^p, a} - L_{2^{p-1}, a}) \geq 2^{p-1} - K \left(2^{3(p-1)/4}/\beta + 3/(2\beta) \right)$$

Using Lemma 29, we obtain

$$\left| \sum_{a \notin V_{2^{p-1}}^p} (N_{2^p,a}^a - N_{2^{p-1},a}^a) - \beta \sum_{a \notin V_{2^{p-1}}^p} (L_{2^p,a} - L_{2^{p-1},a}) \right| \leq 3(K-1)/2.$$

Combining all the above, we obtain

$$\begin{aligned} & \sum_{a \notin V_{2^{p-1}}^p} (L_{2^p,a} - L_{2^{p-1},a}) - \sum_{a \notin V_{2^{p-1}}^p} (N_{2^p,a}^a - N_{2^{p-1},a}^a) \\ & \geq (1-\beta) \sum_{a \notin V_{2^{p-1}}^p} (L_{2^p,a} - L_{2^{p-1},a}) - 3(K-1)/2 \\ & \geq (1-\beta) \left(2^{p-1} - K \left(2^{3(p-1)/4}/\beta + 3/(2\beta) \right) \right) - 3(K-1)/2 \geq K2^{3(p-1)/4}, \end{aligned}$$

where the last inequality is obtained for $p \geq p_5$ with

$$p_5 = \sup \left\{ p \in \mathbb{N} \mid (1-\beta) \left(2^{p-1} - K \left(2^{3(p-1)/4}/\beta + \frac{3}{2\beta} \right) \right) - \frac{3}{2}(K-1) < K2^{3(p-1)/4} \right\}.$$

The LHS summation is exactly the number of times where an arm $a \notin V_{2^{p-1}}^p$ was leader but wasn't sampled, hence

$$\sum_{t=2^{p-1}}^{2^p-1} \mathbb{1}(B_t \notin V_{2^{p-1}}^p, a_t = C_t) \geq K2^{3(p-1)/4}$$

For any $2^{p-1} \leq t \leq 2^p - 1$, U_t^p is non-empty, hence we have $B_t \notin V_{2^{p-1}}^p$ (hence $B_t \notin V_t^p$) implies $C_t \in V_t^p \subseteq V_{2^{p-1}}^p$. Therefore, we have shown that

$$\sum_{t=2^{p-1}}^{2^p-1} \mathbb{1}(a_t \in V_{2^{p-1}}^p) \geq \sum_{t=2^{p-1}}^{2^p-1} \mathbb{1}(B_t \notin V_{2^{p-1}}^p, a_t = C_t) \geq K2^{3(p-1)/4}.$$

Therefore, there is at least one arm in $V_{2^{p-1}}^p$ that is sampled $2^{3(p-1)/4}$ times between 2^{p-1} and $2^p - 1$.

In summary, we have shown $|V_{2^p}^p| \leq K-2$ for all $p \geq p_6 = \max\{p_3, p_4, p_5\}$. By induction, for any $1 \leq k \leq K$, we have $|V_{k2^{p-1}}^p| \leq K-k$, and finally $U_{K2^{p-1}}^p = \emptyset$ for all $p \geq p_6$. Defining $N_0 = K2^{p_6-1}$, we have $\mathbb{E}_\nu[N_0] < +\infty$ by using Lemmas 33 and 36 for $p_3 = \max\{p_2, 4p_0/3 - 1/3\}$ and p_4 and p_5 are deterministic. For all $n \geq N_0$, we let $2^{p-1} = \frac{n}{K}$. Then, by applying the above, we have $U_{K2^{p-1}}^p = U_n^{\log_2(n/K)+1}$ is empty, which shows that $N_{n,a} \geq \sqrt{n/K}$ for all $a \in [K]$. Using Lemma 28, we obtain that $k_{n,a} \geq \frac{\log(n/K)}{2 \log 2} + 1$ for all $a \in [K]$. This concludes the proof. \blacksquare

E.3 Convergence Towards β -optimal Allocation

The second step of in the generic analysis of Top Two algorithms Jourdan et al. (2022) is to show the convergence of the empirical proportions towards the β -optimal allocation. First, we show that the leader coincides with the best arm. Hence, the tracking procedure will ensure that the empirical proportion of time we sample it is exactly β . Second, we show that a sub-optimal arm whose empirical proportion overshoots its β -optimal allocation will not be sampled next as challenger. Therefore, this ‘‘overshoots implies not sampled’’ mechanism will ensure the convergence towards the β -optimal allocation. We emphasise that there are multiple ways to select the leader/challenger pair in order to ensure convergence towards the β -optimal allocation. Therefore, other choices of leader/challenger pair would yield similar results. Note that our results heavily rely on having obtained sufficient exploration first.

Convergence for the best arm. Lemma 38 exhibits a random phase which ensures that the leader and the candidate answer are equal to the best arm for large enough n .

Lemma 38 *Let N_0 be as in Lemma 37. There exists $N_1 \geq N_0$ with $\mathbb{E}_\nu[N_1] < +\infty$ such that, for all $n \geq N_1$, we have $\hat{a}_n = B_n = a^*$.*

Proof Let $k \geq 1$. Suppose that $\mathbb{E}_\nu[\max_{a \in [K]} T_k(a)] < +\infty$. Then, Lemma 28 yields that $N_{T_k(a),a} = 2^{k-1}$ and $\tilde{N}_{k,a} = 2^{k-2}$. Using Lemma 30, we obtain that

$$\begin{aligned} \tilde{\mu}_{k,a^*} &\geq \mu_{a^*} - W_\mu \sqrt{\frac{\log(e + 2^{k-2})}{2^{k-2}}} - W_\epsilon \frac{\log(e + k)}{2^{k-2}}, \\ \forall a \neq a^*, \quad \tilde{\mu}_{k,a} &\leq \mu_a + W_\mu \sqrt{\frac{\log(e + 2^{k-2})}{2^{k-2}}} + W_\epsilon \frac{\log(e + k)}{2^{k-2}}. \end{aligned}$$

Let $p_1 = \lceil \log_2(X_1 - e) \rceil + 2$ and $p_2 = \lceil \log_2(X_2 - e - 1) \rceil + 2$ with

$$\begin{aligned} X_1 &= \sup \{x > 1 \mid x \leq 64\Delta_{\min}^{-2} W_\mu^2 \log x + e\} \leq h_1(64\Delta_{\min}^{-2} W_\mu^2, e), \\ X_2 &= \sup \{x > 1 \mid x \leq 8\Delta_{\min}^{-1} W_\epsilon \log x + e + 1\} \leq h_1(8\Delta_{\min}^{-1} W_\epsilon, e + 1), \\ X_2 &\geq \sup \{x > 1 \mid x \leq 8\Delta_{\min}^{-1} W_\epsilon \log(e + 2 + \log x)\}, \end{aligned}$$

where we used Lemma 32, and h_1 defined therein. Then, for all $k \in \mathbb{N}^K$ such that $\min_{a \in [K]} k_a > p_0 = \max\{p_1, p_2\}$ such that $\mathbb{E}_\nu[\max_{a \in [K]} T_{k_a}(a)] < +\infty$, we have $\tilde{\mu}_{k,a^*} \geq \mu_{a^*} - \Delta_{\min}/4$ and $\tilde{\mu}_{k,a} \leq \mu_a + \Delta_{\min}/4$ for all $a \neq a^*$, hence $a^* = \arg \max_{a \in [K]} \tilde{\mu}_{k,a}$. We have, for all $\alpha \in \mathbb{R}_+$,

$$\exp(\alpha p_1) \leq e^{3\alpha}(X_1 - e)^{\alpha/\log 2} \quad \text{hence} \quad \mathbb{E}_\nu[\exp(\alpha p_1)] < +\infty,$$

where we used Lemma 30 and $h_1(x, e) \sim_{x \rightarrow +\infty} x \log x$ to obtain that $\exp(\alpha p_1)$ is at most polynomial in W_μ . Likewise, we obtain that $\mathbb{E}_\nu[\exp(\alpha p_2)] < +\infty$ for all $\alpha \in \mathbb{R}_+$. Therefore, we have $\mathbb{E}_\nu[\exp(\alpha p_0)] < +\infty$ for all $\alpha \in \mathbb{R}_+$.

Let us define the UCB indices by $I_{k,a} = \tilde{\mu}_{k,a} + \sqrt{k/\tilde{N}_{k,a}} + k/(\epsilon \tilde{N}_{k,a})$. Using the above, we have

$$I_{k,a^*} \geq \mu_{a^*} - W_\mu \sqrt{\frac{\log(e + 2^{k-2})}{2^{k-2}}} - W_\epsilon \frac{\log(e + k)}{2^{k-2}} + \frac{k}{\epsilon 2^{k-2}},$$

$$\forall a \neq a^*, \quad I_{k,a} \leq \mu_a + W_\mu \sqrt{\frac{\log(e + 2^{k-2})}{2^{k-2}}} + W_\epsilon \frac{\log(e + k)}{2^{k-2}} + \frac{k}{\epsilon 2^{k-2}}.$$

Therefore, we have $a^* = \arg \max_{a \in [K]} I_{k,a}$ for all $k \in \mathbb{N}^K$ such that $\min_a k_a > \max\{p_1, p_2\}$ such that $\mathbb{E}_\nu[\max_{a \in [K]} T_{k_a}(a)] < +\infty$.

Let N_0 as in Lemma 37. Using Lemma 37, we obtain that, for all $n \geq N_0$ and all $a \in [K]$, $k_{n,a} \geq \log_2(n/K)/2 + 1$. Therefore, we obtain $\min_{a \in [K]} k_{n,a} > \max\{p_1, p_2\}$ is implied by $n \geq N_1 = \max\{K4^{\max\{p_1, p_2\}}, N_0\}$. Using the above, we conclude that $\mathbb{E}_\nu[N_1] < +\infty$ and $\hat{a}_n = B_n = a^*$ for all $n \geq N_1$. \blacksquare

Lemma 39 shows that that the pulling proportion of the best arm converges towards β , provided the phase defined in Lemma 38 is reached in finite time for all arms.

Lemma 39 *Let $\gamma > 0$, and N_1 be as in Lemma 38. There exists a deterministic constant $C_0 \geq 1$ such that, for all $n \geq C_0 N_1$,*

$$\left| \frac{N_{n,a^*}}{n-1} - \beta \right| \leq \gamma.$$

Proof Let $\gamma > 0$. Let N_1 as in Lemma 38. Let $M \geq N_1$. Using Lemma 38, we obtain $B_n = a^*$ for all $n \geq M$. Therefore, we obtain $L_{n,a^*} \geq n - M$ and $\sum_{a \neq a^*} N_{n,a^*}^a \leq M$ for all $n \geq M$. Using Lemma 29 yields that

$$\begin{aligned} \left| \frac{N_{n,a^*}}{n-1} - \beta \right| &\leq \frac{|N_{n,a^*}^{a^*} - \beta L_{n,a^*}|}{n-1} + \beta \left| \frac{L_{n,a^*}}{n-1} - 1 \right| + \frac{1}{n-1} \sum_{a \neq a^*} N_{n,a^*}^a \\ &\leq \frac{1}{2(n-1)} + \beta \frac{2(M-1)}{n-1} \leq \gamma, \end{aligned}$$

where the last inequality is obtained by taking $n \geq \max\{M, (1/2 + 2\beta(M-1))/\gamma + 1\}$. \blacksquare

Convergence for the sub-optimal arms. Lemma 40 exhibits a random phase which ensures that if a sub-optimal arm overshoots its β -optimal allocation then it cannot be selected as challenger for large enough n .

Lemma 40 *Let $\gamma > 0$. Let N_1 and C_0 be as in Lemma 38 and 39. There exists $N_2 \geq C_0 N_1$ with $\mathbb{E}_\nu[N_2] < +\infty$ such that, for all $n \geq N_2$,*

$$\exists a \neq a^*, \quad \frac{N_{n,a}}{n-1} \geq \gamma + \begin{cases} \omega_{\beta,a}^* & [\text{AdaP-TT}] \\ \omega_{\epsilon,\beta,a}^* & [\text{AdaP-TT}^*] \end{cases} \implies C_n \neq a,$$

Proof Let $\gamma > 0$ and $\tilde{\gamma} > 0$. Let N_1 as in Lemma 38 and C_0 as in Lemma 39 for $\tilde{\gamma}$. Let $n \geq C_0 N_1$.

Let $a \neq a^*$ such that $\frac{N_{n,a}}{n-1} \geq \omega_{\beta,a}^* + \gamma$. Suppose towards contradiction that $\frac{N_{n,b}}{n-1} > \omega_{\beta,a}^*$ for all $b \notin \{a^*, a\}$. Then, for all $n \geq C_0 N_1$, we have

$$1 - \beta + \tilde{\gamma} \geq 1 - \frac{N_{n,a^*}}{n-1} = \sum_{b \neq a^*} \frac{N_{n,b}}{n-1} > \gamma + \sum_{b \neq a^*} \omega_{\beta,b}^* = 1 - \beta + \gamma,$$

which yields a contradiction for $\tilde{\gamma} \leq \gamma$. Therefore, for all $n \geq C_0 N_1$, we have

$$\exists a \neq a^*, \quad \frac{N_{n,a}}{n-1} \geq \omega_{\beta,a}^* + \gamma \quad \implies \quad \exists b \notin \{a^*, a\}, \quad \frac{N_{n,b}}{n-1} \leq \omega_{\beta,b}^*.$$

Then, we have

$$\sqrt{\frac{1 + N_{n,a^*}/N_{n,b}}{1 + N_{n,a^*}/N_{n,a}}} \geq \sqrt{\frac{1 + (\beta - \tilde{\gamma})/\omega_{\beta,b}^*}{1 + (\beta + \tilde{\gamma})/(\omega_{\beta,a}^* + \gamma)}}.$$

In the following, we use Lemma 30 and similar manipulations as in the proof of Lemma 38. Therefore, we obtain that, for all $c \neq a^*$,

$$\begin{aligned} \left| \tilde{\mu}_{k_{n,a^*},a^*} - \tilde{\mu}_{k_{n,c},c} - \Delta_c \right| &\leq W_\mu \left(\sqrt{\frac{\log(e + 2^{k_{n,a^*}-2})}{2^{k_{n,a^*}-2}}} + \sqrt{\frac{\log(e + 2^{k_{n,c}-2})}{2^{k_{n,c}-2}}} \right) \\ &\quad + W_\epsilon \left(\frac{\log(e + k_{n,a^*})}{2^{k_{n,a^*}-2}} + \frac{\log(e + k_{n,c})}{2^{k_{n,c}-2}} \right). \end{aligned}$$

Let $p_3 = \lceil \log_2(X_1 - e) \rceil + 2$ and $p_2 = \lceil \log_2(X_2 - e - 1) \rceil + 2$ with

$$\begin{aligned} X_3 &= \sup \{x > 1 \mid x \leq 16\eta^{-2}W_\mu^2 \log x + e\} \leq h_1(16\eta^{-2}W_\mu^2, e), \\ X_2 &= \sup \{x > 1 \mid x \leq 4\eta^{-1}W_\epsilon \log x + e + 1\} \leq h_1(4\eta^{-1}W_\epsilon, e + 1), \\ X_2 &\geq \sup \{x > 1 \mid x \leq 4\eta^{-1}W_\epsilon \log(e + 2 + \log x)\}, \end{aligned}$$

where we used Lemma 32, and h_1 defined therein. We have, for all $\alpha \in \mathbb{R}_+$,

$$\exp(\alpha p_3) \leq e^{3\alpha}(X_3 - e)^{\alpha/\log 2} \quad \text{hence} \quad \mathbb{E}_\nu[\exp(\alpha p_3)] < +\infty,$$

where we used Lemma 30 and $h_1(x, e) \sim_{x \rightarrow +\infty} x \log x$ to obtain that $\exp(\alpha p_3)$ is at most polynomial in W_μ . Likewise, we obtain that $\mathbb{E}_\nu[\exp(\alpha p_2)] < +\infty$ for all $\alpha \in \mathbb{R}_+$.

Using Lemma 37 (with $C_0 N_1 \geq N_1 \geq N_0$), we obtain that, for all $n \geq C_0 N_1$ and all $a \in [K]$, $k_{n,a} \geq \log_2(n/K)/2 + 1$. Therefore, we obtain $\min_{a \in [K]} k_{n,a} > \max\{p_2, p_3\}$ is implied by $n \geq N_2 = \max\{K4^{\max\{p_3, p_2\}}, C_0 N_1\}$. Using the above, we conclude that $\mathbb{E}_\nu[N_2] < +\infty$ and $\max_{c \neq a^*} |\tilde{\mu}_{k_{n,a^*},a^*} - \tilde{\mu}_{k_{n,c},c} - \Delta_c| \leq \eta$ for all $n \geq N_2$.

Then, for all $n \geq N_2$, we have $B_n = a^*$ and

$$\frac{\tilde{\mu}_{k_{n,a^*},a^*} - \tilde{\mu}_{k_{n,a},a}}{\tilde{\mu}_{k_{n,a^*},a^*} - \tilde{\mu}_{k_{n,b},b}} \sqrt{\frac{1 + N_{n,a^*}/N_{n,b}}{1 + N_{n,a^*}/N_{n,a}}} \geq \frac{\Delta_a - \eta}{\Delta_b + \eta} \sqrt{\frac{1 + (\beta - \tilde{\gamma})/\omega_{\beta,b}^*}{1 + (\beta + \tilde{\gamma})/(\omega_{\beta,a}^* + \gamma)}} > 1,$$

where the last inequality is obtained by taking η and $\tilde{\gamma}$ sufficiently small and by using (25)

$$\frac{\Delta_a}{\Delta_b} \sqrt{\frac{1 + \beta/\omega_{\beta,b}^*}{1 + \beta/\omega_{\beta,a}^*}} = 1.$$

Therefore, we have shown that $B_n = a^*$ and

$$\frac{\tilde{\mu}_{k_{n,a^*},a^*} - \tilde{\mu}_{k_{n,a},a}}{\sqrt{1/N_{n,a^*} + 1/N_{n,a}}} > \frac{\tilde{\mu}_{k_{n,a^*},a^*} - \tilde{\mu}_{k_{n,b},b}}{\sqrt{1/N_{n,a^*} + 1/N_{n,b}}} \quad \text{hence} \quad C_n \neq a.$$

This concludes the proof of the first result.

For the AdaP-TT^{*} algorithm, the proof is done similarly. As above, we can construct \tilde{N}_2 with $\mathbb{E}_\nu[\tilde{N}_2] < +\infty$ such that, for all $n \geq \tilde{N}_2$, we have $B_n = a^*$ and

$$\begin{aligned} & \frac{(\tilde{\mu}_{k_n, a^*, a^*} - \tilde{\mu}_{k_n, a, a}) \min\{3\epsilon, \tilde{\mu}_{k_n, a^*, a^*} - \tilde{\mu}_{k_n, a, a}\} 1 + N_{n, a^*}/N_{n, b}}{(\tilde{\mu}_{k_n, a^*, a^*} - \tilde{\mu}_{k_n, b, b}) \min\{3\epsilon, \tilde{\mu}_{k_n, a^*, a^*} - \tilde{\mu}_{k_n, b, b}\} 1 + N_{n, a^*}/N_{n, a}} \\ & \geq \frac{(\Delta_a - \eta) \min\{3\epsilon, \Delta_a - \eta\}}{(\Delta_b + \eta) \min\{3\epsilon, \Delta_b + \eta\}} \frac{1 + (\beta - \tilde{\gamma})/\omega_{\epsilon, \beta, b}^*}{1 + (\beta + \tilde{\gamma})/(\omega_{\epsilon, \beta, a}^* + \gamma)} > 1, \end{aligned}$$

where the last inequality is obtained by taking η and $\tilde{\gamma}$ sufficiently small and by using (25)

$$\frac{\Delta_a \min\{3\epsilon, \Delta_a\}}{\Delta_b \min\{3\epsilon, \Delta_b\}} \sqrt{\frac{1 + \beta/\omega_{\epsilon, \beta, b}^*}{1 + \beta/\omega_{\epsilon, \beta, a}^*}} = 1.$$

Then, we conclude similarly by using the definition of the TC challenger. \blacksquare

Lemma 41 shows that the pulling proportion of the best arm converges towards β for large enough n .

Lemma 41 *Let $\gamma > 0$ and $T_{\mu, \gamma}(w)$ as in (26). Then, we have $\mathbb{E}_\nu[T_{\mu, \gamma}(\omega_\beta^*)] < +\infty$ (AdaP-TT) and $\mathbb{E}_\nu[T_{\mu, \gamma}(\omega_{\epsilon, \beta}^*)] < +\infty$ (AdaP-TT^{*}).*

Proof Let $\gamma > 0$ and $\tilde{\gamma} > 0$. Let N_2 as in Lemma 40 for $\tilde{\gamma}$. Let $M \geq N_2$. Using Lemmas 38, 39 and 40 for all $n \geq M$, we obtain that $B_n = a^*$, $\left| \frac{N_{n, a^*}}{n-1} - \beta \right| \leq \tilde{\gamma}$ and

$$\exists a \neq a^*, \quad \frac{N_{n, a}}{n-1} \geq \omega_{\beta, a}^* + \tilde{\gamma} \quad \implies \quad C_n \neq a^*.$$

For all $a \neq a^*$, let us define $t_{n, a}(\tilde{\gamma}) = \max\{t \mid M \leq t \leq n, N_{t, a}/(n-1) < \omega_{\beta, a}^* + \tilde{\gamma}\}$. Since $N_{t, a}/(n-1) \leq N_{t, a}/(t-1)$ for $t \leq n$, we have

$$\begin{aligned} \frac{N_{n, a}}{n-1} & \leq \frac{M-1}{n-1} + \frac{1}{n-1} \sum_{t=M}^n \mathbb{1}(a_t = C_t = a) \\ & \leq \frac{M-1}{n-1} + \frac{1}{n-1} \sum_{t=M}^n \mathbb{1}\left(\frac{N_{t, a}}{n-1} < \omega_{\beta, a}^* + \tilde{\gamma}, a_t = C_t = a\right) \\ & \leq \frac{M-1}{n-1} + \frac{N_{t_{n, a}(\tilde{\gamma}), a}}{n-1} < \frac{M-1}{n-1} + \omega_{\beta, a}^* + \tilde{\gamma}. \end{aligned}$$

The second inequality uses Lemma 40, and the two last inequalities use the definition of $t_{n, a}(\tilde{\gamma})$. Using that $\sum_{a \in [K]} \frac{N_{n, a}}{n-1} = \sum_{a \in [K]} \omega_{\beta, a}^* = 1$, we obtain

$$\frac{N_{n, a}}{n-1} = 1 - \sum_{b \neq a} \frac{N_{n, b}}{n-1} \geq 1 - \sum_{b \neq a} \left(\omega_{\beta, b}^* + \tilde{\gamma} + \frac{M-1}{n-1} \right) = \omega_{\beta, a}^* - (K-1) \left(\tilde{\gamma} + \frac{M-1}{n-1} \right).$$

Taking $\tilde{\gamma} \leq \gamma/(2(K-1))$ and $n \geq \max\{M, 2(K-1)(M-1)/\gamma + 1\}$ yields that

$$\left\| \frac{N_n}{n-1} - \omega_\beta^* \right\|_\infty \leq \gamma.$$

Let $T_{\mu,\gamma}(w)$ as in (26). Then, we showed that $T_{\mu,\gamma}(\omega_\beta^*) \leq \max\{M, 2(K-1)(M-1)/\gamma + 1\}$. Therefore, we have

$$\mathbb{E}_\nu[T_{\mu,\gamma}(\omega_\beta^*)] \leq \mathbb{E}_\nu[\max\{M, 2(K-1)(M-1)/\gamma + 1\}] < +\infty,$$

which concludes the proof of the first result.

For the AdaP-TT* algorithm, the proof is exactly the same by replacing ω_β^* by $\omega_{\epsilon,\beta}^*$. ■

E.4 Cost of Doubling and Forgetting

Compared to the generic analysis of Top Two algorithms Jourdan et al. (2022), we need to control the sample complexity cost of the DAF(ϵ) update (Algorithm 5). Due to this reason, we have to pay a multiplicative four-factor: one two-factor due to doubling, and another two-factor due to forgetting. It is possible to show that this cost exists when adapting any BAI algorithm in which the empirical proportions are converging towards an allocation ω such that $\min_a \omega_a > 0$, i.e. there exists ω such that $\mathbb{E}_\nu[T_{\mu,\gamma}(\omega)] < +\infty$. As shown in Lemma 41, this is the case for the AdaP-TT and AdaP-TT* algorithms.

Lemma 42 shows that the phase switches of the arms happen in a round-robin fashion, which means that an arm switches phase for a second time after all other arms first switch their own phases.

Lemma 42 *Let $\omega \in \Sigma_K$ such that $\min_a \omega_a > 0$. Assume that there exists $\gamma_\mu > 0$ such that for $\mathbb{E}_\nu[T_{\mu,\gamma}(\omega)] < +\infty$ for all $\gamma \in (0, \gamma_\mu)$, where $T_{\mu,\gamma}(\omega)$ is defined in Equation (26). Let $\eta > 0$. There exists $\tilde{\gamma}_\mu \in (0, \gamma_\mu)$ such that, for all $\gamma \in (0, \tilde{\gamma}_\mu)$, there exists $N_3 \geq T_{\mu,\gamma}(\omega)$ with $\mathbb{E}_\nu[N_3] < +\infty$ which satisfies*

$$\forall n \geq N_3, \quad \frac{\max_{a \in [K]} T_{k_{n,a}}(a) - 1}{\min_{a \in [K]} T_{k_{n,a}}(a) - 1} \leq 2 + \eta.$$

Proof Let $\eta > 0$. Let $\tilde{\gamma}_\mu \in (0, \gamma_\mu)$ such that $2 \max_{a \in [K]} (\omega_a + \gamma) / (\omega_a - \gamma) \leq 2 + \eta$, which is possible since $\min_a \omega_a > 0$. Let $\gamma \in (0, \tilde{\gamma}_\mu)$. By assumption, we have $\mathbb{E}_\nu[T_{\mu,\gamma}(\omega)] < +\infty$. Then, for all $n \geq T_{\mu,\gamma}(\omega)$,

$$\left\| \frac{N_n}{n-1} - \omega \right\|_\infty \leq \gamma.$$

Let $M \geq T_{\mu,\gamma}(\omega)$. Let us denote by $k_M = (k_{M,a})_{a \in [K]}$ the current phases for all arms $a \in [K]$ at time M . Then, for all $n \geq M$ and all $a \in [K]$, we have $N_{n,a} \geq (n-1)(\omega_a - \gamma)$. Therefore, taking $n \geq \max_{a \in [K]} 2^{k_{M,a}} (\omega_a - \gamma)^{-1} + 1$, we obtain that $N_{n,a} \geq 2^{k_{M,a}}$ for all $a \in [K]$, hence we have $\max_{a \in [K]} T_{k_{M,a}+1}(a) \leq n$. Since $\min_{a \in [K]} T_{k_{M,a}+1}(a) \geq M$, we have

$$\max_{a \in [K]} \left| \frac{N_{T_{k_{M,a}+1}(a),a}}{n-1} - \omega_a \right| \leq \gamma.$$

Likewise, taking $n \geq \max_{a \in [K]} 2^{k_{M,a}+1}(\omega_a - \gamma)^{-1} + 1$, we obtain that $N_{n,a} \geq 2^{k_{M,a}+1}$ for all $a \in [K]$, hence we have $\max_{a \in [K]} T_{k_{M,a}+2}(a) \leq n$. Let $a_1 = \arg \min_{a \in [K]} T_{k_{M,a}+2}(a)$. By definition and using Lemma 28, we have

$$\begin{aligned} 2^{k_{M,a_1}+1} &= N_{T_{k_{M,a_1}+2}(a_1), a_1} \leq (T_{k_{M,a_1}+2}(a_1) - 1)(\omega_{a_1} + \gamma), \\ \forall a \neq a_1, \quad 2^{k_{M,a}} &\leq N_{T_{k_{M,a_1}+2}(a_1), a} \leq (T_{k_{M,a_1}+2}(a_1) - 1)(\omega_a + \gamma). \end{aligned}$$

Let $a_2 = \arg \max_{a \in [K]} T_{k_{M,a}+2}(a)$. By definition and using Lemma 28, we have

$$2^{k_{M,a_2}+1} = N_{T_{k_{M,a_2}+2}(a_2), a_2} \geq (T_{k_{M,a_2}+2}(a_2) - 1)(\omega_{a_2} - \gamma),$$

Therefore, combining the above yields

$$(T_{k_{M,a_2}+2}(a_2) - 1) \leq (T_{k_{M,a_1}+2}(a_1) - 1) 2 \frac{\omega_{a_2} + \gamma}{\omega_{a_2} - \gamma} \leq (T_{k_{M,a_2}+2}(a_2) - 1)(2 + \eta),$$

where the last inequality uses that $\gamma \in (0, \tilde{\gamma}_\mu)$ and $\tilde{\gamma}_\mu \in (0, \gamma_\mu)$ is such that $2 \max_{a \in [K]} (\omega_a + \gamma)/(\omega_a - \gamma) \leq 2 + \eta$. We take $n \geq N_3 = \max_{a \in [K]} T_{k_{M,a}+2}(a)$, hence we have $k_{n,a} \geq k_{M,a} + 2$ for all $a \in [K]$. Since $\mathbb{E}_\nu[T_{\mu,\gamma}(\omega)] < +\infty$ (i.e. arms are sampled linearly), it is direct to see that $\mathbb{E}_\nu[\max_{a \in [K]} T_{k_{M,a}+2}(a)] < +\infty$. This concludes the proof. \blacksquare

E.5 Asymptotic Upper Bound on the Expected Sample Complexity

The final step of the generic analysis of Top Two algorithms (Jourdan et al., 2022) is to invert the private GLR stopping rule by leveraging the convergence of the empirical proportions towards the β -optimal allocation. Provided this convergence is shown, the asymptotic upper bound on the expected sample complexity only depends on the dependence in $\log(1/\delta)$ of the threshold that ensures δ -correctness. Compared to the non-private GLR stopping rule, the private GLR stopping rules pay an extra cost to ensure privacy. In Section 4.3, the stopping threshold is adapted with an additive term in $\mathcal{O}(\log(1/\delta)^2)$. In Section 4.4, both the stopping threshold and the transportation costs are modified.

Lemma 43 *Let $(\delta, \beta) \in (0, 1)^2$. Assume that there exists $\gamma_\mu > 0$ such that $\mathbb{E}_\nu[T_{\mu,\gamma}(\omega_\beta^*)] < +\infty$ for all $\gamma \in (0, \gamma_\mu)$, where $T_{\mu,\gamma}(w)$ is defined in (26). Combining such a sampling rule, using the $DAF(\epsilon)$ update, with the GLR stopping rule with $W_{a,b}^G$ as in (3) and the stopping threshold $c_{a,b}^{G,\epsilon}$ as in (11) yields a δ -correct algorithm which satisfies that, for all ν with mean μ such that $|a^*(\mu)| = 1$,*

$$\limsup_{\delta \rightarrow 0} \frac{\mathbb{E}_\nu[\tau_\delta]}{\log(1/\delta)} \leq 4T_{\text{KL},\beta}^*(\nu) \left(1 + \sqrt{1 + \frac{\Delta_{\max}^2}{2\epsilon^2\sigma^4}} \right).$$

where $T_{\text{KL},\beta}^*(\nu)$ as in (5) with $\sigma = 1/2$.

Assume that there exists $\gamma_\mu > 0$ such that $\mathbb{E}_\nu[T_{\mu,\gamma}(\omega_{\epsilon,\beta}^*)] < +\infty$. Combining such a sampling rule, using the $DAF(\epsilon)$ update, with the GLR stopping rule with $W_{a,b}^{G,\epsilon}$ as in (12)

and the stopping threshold $\tilde{c}_{a,b}^{G,\epsilon}$ as in (14) yields a δ -correct algorithm which satisfies that, for all ν with mean μ such that $|a^*(\mu)| = 1$,

$$\limsup_{\delta \rightarrow 0} \frac{\mathbb{E}_\nu[\tau_\delta]}{\log(1/\delta)} \leq \begin{cases} 4T_{\text{KL},\beta}^*(\nu)g_1(\Delta_{\max}/(\sigma^2\epsilon)) & \text{if } \Delta_{\max} < 3\epsilon \\ 12T_{\epsilon,\beta}^*(\nu)g_2(3\epsilon^2T_{\epsilon,\beta}^*(\nu)\max\{\beta, 1-\beta\}/2)/\sigma^2 & \text{otherwise} \end{cases},$$

where $T_{\epsilon,\beta}^*(\nu)$ as in (13) with $\sigma = \frac{1}{2}$. The function $g_1(y) = \sup\{x \mid x^2 < x + y\sqrt{2x} + \frac{y^2}{4}\}$ is increasing on $[0, 12]$ and satisfies that $g_1(0) = 1$ and $g_1(12) \leq 10$. The function $g_2(y) = 1 + 2(\sqrt{1+1/y} - 1)^{-1}$ is increasing on \mathbb{R}_+^* and satisfies that $\lim_{y \rightarrow 0} g_2(y) = 1$.

Proof Lemma 16 and Lemma 18 yields the δ -correctness of both algorithms.

AdaP-TT algorithm. Let $\zeta > 0$, a^* be the unique best arm. Using (25) and the continuity of

$$(\mu, w) \mapsto \min_{a \neq a^*(\mu)} \frac{(\mu_{a^*}(\mu) - \mu_a)^2}{2\sigma^2(1/w_{a^*}(\mu) + 1/w_a)}$$

yields that there exists $\gamma_\zeta > 0$ such that $\left\| \frac{N_n}{n-1} - \omega_\beta^* \right\|_\infty \leq \gamma_\zeta$ and $\max_{a \in [K]} |\tilde{\mu}_{k_{n,a}+1,a} - \mu_a| \leq \gamma_\zeta$ implies that

$$\begin{aligned} \forall a \neq a^*, \quad & \frac{(\tilde{\mu}_{k_{n,a^*}+1,a^*} - \tilde{\mu}_{k_{n,a}+1,a})^2}{(n-1)/N_{n,a^*} + (n-1)/N_{n,a}} \geq \frac{2\sigma^2(1-\zeta)}{T_{\text{KL},\beta}^*(\nu)}, \\ & \frac{n-1}{N_{n,a^*}} + \frac{n-1}{N_{n,a}} \leq \frac{\Delta_a^2}{2\sigma^2}(1+\zeta)T_{\text{KL},\beta}^*(\nu). \end{aligned}$$

We choose such a γ_ζ . Let $\gamma_\mu > 0$ be such that for $\mathbb{E}_\nu[T_{\mu,\gamma}(\omega_\beta^*)] < +\infty$ for all $\gamma \in (0, \gamma_\mu)$, where $T_{\mu,\gamma}(\omega)$ is defined in (26). Let $\eta > 0$. Let $\tilde{\gamma}_\mu \in (0, \gamma_\mu)$ as in Lemma 42 for this η . In the following, let us consider $\gamma \in (0, \min\{\tilde{\gamma}_\mu, \gamma_\zeta, \beta/4, \Delta_{\min}/4\})$.

Let $N_3 \geq T_{\mu,\gamma}(\omega_\beta^*)$ with $\mathbb{E}_\nu[N_3] < +\infty$ as Lemma 42 for those (γ, η) . Then, we have $\mathbb{E}_\nu[T_{\mu,\gamma}(\omega_\beta^*)] < +\infty$ and

$$\forall n \geq N_3, \quad \frac{\max_{a \in [K]} T_{k_{n,a}}(a) - 1}{\min_{a \in [K]} T_{k_{n,a}}(a) - 1} \leq 2 + \eta.$$

Since arms are sampled linearly, it is direct to construct $N_4 \geq N_3$ with $\mathbb{E}_\nu[N_4] < +\infty$ such that, for all $n \geq N_4$, we have $\max_{a \in [K]} \max_{k \in \{k_{n,a}, k_{n,a+1}\}} |\tilde{\mu}_{k,a} - \mu_a| \leq \gamma$, Therefore, we have $\hat{a}_n = a^*$.

Let $\kappa \in (0, 1)$. Let $n \geq N_4/\kappa$ and $(k_{n,a})_{a \in [K]}$ be the current phases at time n . Combining the above, we have $\hat{a}_n = a^*$ and

$$\max_{a \in [K]} |\tilde{\mu}_{k_{n,a}+1,a} - \mu_a| \leq \gamma, \quad \left\| \frac{N_n}{n-1} - \omega_\beta^* \right\|_\infty \leq \gamma \quad \text{and} \quad \frac{\max_{a \in [K]} T_{k_{n,a}}(a) - 1}{\min_{a \in [K]} T_{k_{n,a}}(a) - 1} \leq 2 + \eta.$$

Let $a_1 = \arg \min_{a \in [K]} T_{k_{n,a}}(a)$ and $a_2 = \arg \max_{a \in [K]} T_{k_{n,a}}(a)$. Therefore, we obtain

$$\forall a \neq a^*, \quad \frac{(\tilde{\mu}_{k_{n,\hat{a}_n}+1,\hat{a}_n} - \tilde{\mu}_{k_{n,a}+1,a})^2}{1/\tilde{N}_{k_{n,\hat{a}_n}+1,\hat{a}_n} + 1/\tilde{N}_{k_{n,a}+1,a}} = \frac{(\tilde{\mu}_{k_{n,a^*}+1,a^*} - \tilde{\mu}_{k_{n,a}+1,a})^2}{1/N_{T_{k_{n,a^*}}(a^*),a^*} + 1/N_{T_{k_{n,a}}(a),a}}$$

$$\begin{aligned}
 &\geq \frac{(\tilde{\mu}_{k_n, a^*+1, a^*} - \tilde{\mu}_{k_n, a+1, a})^2}{1/N_{T_{k_n, a_1}(a_1), a^*} + 1/N_{T_{k_n, a_1}(a_1), a}} \\
 &\geq \left(\min_{a \in [K]} T_{k_n, a}(a) - 1 \right) \frac{2\sigma^2(1-\zeta)}{T_{\text{KL}, \beta}^*(\boldsymbol{\nu})}.
 \end{aligned}$$

Similarly, we can show that, for all $a \neq a^*$,

$$\begin{aligned}
 \frac{1}{\tilde{N}_{k_n, a^*+1, a^*}} + \frac{1}{\tilde{N}_{k_n, a+1, a}} &= \frac{1}{N_{T_{k_n, a^*}(a^*), a^*}} + \frac{1}{N_{T_{k_n, a}(a), a}} \\
 &\leq \frac{1}{N_{T_{k_n, a_1}(a_1), a^*}} + \frac{1}{N_{T_{k_n, a_1}(a_1), a}} \\
 &\leq \frac{1}{\min_{a \in [K]} T_{k_n, a}(a) - 1} \frac{\Delta_a^2}{2\sigma^2} (1 + \zeta) T_{\text{KL}, \beta}^*(\boldsymbol{\nu}) \\
 &\leq \frac{1}{\min_{a \in [K]} T_{k_n, a}(a) - 1} \frac{\Delta_{\max}^2}{2\sigma^2} (1 + \zeta) T_{\text{KL}, \beta}^*(\boldsymbol{\nu}).
 \end{aligned}$$

Let $c_{a,b}^G$ as in (4). Using Lemma 28, we obtain, for all $a \neq a^*$,

$$\begin{aligned}
 c_{a^*, a}^G (\tilde{N}_{k_n+1}, \delta (2\zeta(s)^2 (k_n, a^* + 1)^s (k_n, a + 1)^s)^{-1}) &\leq 4 \log(4 + (\max_{b \in [K]} k_n, b - 1) \log 2) \\
 + 2\mathcal{C}_G \left(\log(1/\delta)/2 + s \log(\max_{b \in [K]} k_n, b - 1) + \log(2(K-1)\zeta(s)^2)/2 \right)
 \end{aligned}$$

Likewise, we obtain, for all $a \in [K]$,

$$\begin{aligned}
 &\frac{1}{\epsilon^2 \sigma^2} \sum_{c \in \{a^*, a\}} \frac{1}{\tilde{N}_{k_n, c, c}} \left(\log \frac{2K\zeta(s)(k_n, c + 1)^s}{\delta} \right)^2 \\
 &\leq \frac{\Delta_{\max}^2}{2\epsilon^2 \sigma^4} \frac{(1 + \zeta) T_{\text{KL}, \beta}^*(\boldsymbol{\nu})}{\min_{b \in [K]} T_{k_n, b}(b) - 1} \left(\log(1/\delta) + s \log(\max_{b \in [K]} k_n, b + 1) + \log(2K\zeta(s)) \right)^2
 \end{aligned}$$

Let us denote by $T_{k_n+1}^+ = \max_{b \in [K]} T_{k_n, b+1}(b)$, $T_{k_n+2}^+ = \max_{b \in [K]} T_{k_n, b+2}(b)$, $T_{k_n+1}^- = \min_{b \in [K]} T_{k_n, b+1}(b)$, $T_{k_n}^- = \min_{b \in [K]} T_{k_n, b}(b)$. Let T be a time such that $T \geq T_{k_n+1}^+ \geq \kappa T$. Using Lemmas 28 and 42, we have

$$(k_n, b - 1) \log 2 = \log N_{T_{k_n, b}(b), b} \leq \log T_{k_n, b}(b) \leq \log T_{k_n}^+ \leq \log T_{k_n}^- + \log(2 + \eta).$$

Using the DAF(ϵ) update with the GLR stopping rule with $W_{a,b}^G$ as in (3) and the stopping threshold $c_{a,b}^{G, \epsilon}$ as in (11), we have

$$\begin{aligned}
 \min\{\tau_\delta, T\} - \kappa T &\leq \sum_{T \geq T_{k_n}^+ \geq \kappa T} (T_{k_n+2}^+ - T_{k_n+1}^+) \mathbb{1}(\tau_\delta > T_{k_n+1}^+) \\
 &\leq \sum_{T_{k_n}^+ = \kappa T}^T (T_{k_n+2}^+ - T_{k_n+1}^+) \mathbb{1} \left(\exists a \neq a^*, \frac{(\tilde{\mu}_{k_n, a^*+1, a^*} - \tilde{\mu}_{k_n, a+1, a})^2}{2\sigma^2 \left(\frac{1}{\tilde{N}_{k_n, a^*+1, a^*}} + \frac{1}{\tilde{N}_{k_n, a+1, a}} \right)} < c_{a^*, a}^{G, \epsilon} (\tilde{N}_{k_n+1}, \delta) \right)
 \end{aligned}$$

$$\begin{aligned}
 &\leq \sum_{T \geq T_{k_n}^+ \geq \kappa T} (T_{k_n+2}^+ - T_{k_n+1}^+) \mathbf{1} \left((T_{k_n}^- - 1) \frac{1 - \zeta}{T_{\text{KL},\beta}^*(\boldsymbol{\nu})} < 8 \log(4 + \log T_{k_n}^- + \log(2 + \eta)) \right) \\
 &+ 4\mathcal{C}_G (\log(1/\delta)/2 + s \log(2 + \log_2 T_{k_n}^- + \log_2(2 + \eta)) + \log(2(K - 1)\zeta(s)^2)/2) \\
 &+ \frac{\Delta_{\max}^2}{2\epsilon^2\sigma^4} \frac{(1 + \zeta)T_{\text{KL},\beta}^*(\boldsymbol{\nu})}{T_{k_n}^- - 1} (\log(1/\delta) + s \log(2 + \log_2 T_{k_n}^- + \log_2(2 + \eta)) + \log(2K\zeta(s)))^2 \Big),
 \end{aligned}$$

Let $T_\zeta(\delta)$ defined as the largest deterministic time such that the above condition is satisfied when replacing $T_{k_n}^-$ by $(1 - \kappa)T$. Let k_δ be the largest random vector of phases such that that $T_{k_\delta+1}^+ \leq T_\zeta(\delta)$ almost surely, hence $T_{k_\delta+2}^+ > T_\zeta(\delta)$ almost surely. Then, using the above yields that $\tau_\delta \leq T_{k_\delta+2}^+$ almost surely, hence

$$\limsup_{\delta \rightarrow 0} \frac{\mathbb{E}_\nu[\tau_\delta]}{\log(1/\delta)} \leq \limsup_{\delta \rightarrow 0} \frac{\mathbb{E}_\nu[T_{k_\delta+2}^+]}{\log(1/\delta)} \leq (2+\eta)^2 \limsup_{\delta \rightarrow 0} \frac{\mathbb{E}_\nu[T_{k_\delta+1}^+]}{\log(1/\delta)} \leq (2+\eta)^2 \limsup_{\delta \rightarrow 0} \frac{T_\zeta(\delta)}{\log(1/\delta)}$$

where the second inequality uses Lemma 42 twice, i.e. $T_{k_\delta+2}^+ \leq (2+\eta)T_{k_\delta+2}^- \leq (2+\eta)^2 T_{k_\delta+1}^+$, and the last one used the definition of k_δ and that $T_\zeta(\delta)$ is deterministic.

Since we are only interested in upper bounding $\limsup_{\delta \rightarrow 0} \frac{T_\zeta(\delta)}{\log(1/\delta)}$, we can safely drop the second orders terms in T and $\log(1/\delta)$. This allows us to remove the terms in $\mathcal{O}(\log \log T)$ and in $\mathcal{O}(\log \log(1/\delta))$. Using that $\mathcal{C}_G(x) = x + \mathcal{O}(\log x)$, tedious manipulations yields that

$$\limsup_{\delta \rightarrow 0} \frac{T_\zeta(\delta)}{\log(1/\delta)} \leq \frac{T_{\text{KL},\beta}^*(\boldsymbol{\nu})}{1 - \kappa} D_\zeta(\mu, \epsilon),$$

where

$$D_\zeta(\mu, \epsilon) = \sup \left\{ x \mid x^2 < \frac{2}{1 - \zeta}x + \frac{1 + \zeta}{1 - \zeta} \frac{\Delta_{\max}^2}{2\epsilon^2\sigma^4} \right\} \leq \frac{1}{1 - \zeta} \left(1 + \sqrt{1 + (1 - \zeta^2) \frac{\Delta_{\max}^2}{2\epsilon^2\sigma^4}} \right).$$

The last inequality uses that $x^2 - 2bx - c < 0$ for all $x \in [0, b(1 + \sqrt{1 + c/b^2})]$. Therefore, we have shown that

$$\limsup_{\delta \rightarrow 0} \frac{\mathbb{E}_\nu[\tau_\delta]}{\log(1/\delta)} \leq (2 + \eta)^2 \frac{T_{\text{KL},\beta}^*(\boldsymbol{\nu})}{(1 - \kappa)(1 - \zeta)} \left(1 + \sqrt{1 + (1 - \zeta^2) \frac{\Delta_{\max}^2}{2\epsilon^2\sigma^4}} \right).$$

Letting κ, η and ζ goes to zero concludes the proof of the first result.

AdaP-TT* algorithm. For the AdaP-TT* algorithm, the proof is done with similar arguments. Using (25) and the continuity of $(\boldsymbol{\mu}, w) \rightarrow \min_{a \neq a^*} W_{a^*,a}^{G,\epsilon}(\boldsymbol{\mu}, w)$, defined in (12), we obtain another $\gamma_\zeta > 0$ such that $\left\| \frac{N_n}{n-1} - \omega_{\epsilon,\beta}^* \right\|_\infty \leq \gamma_\zeta$ and $\max_{a \in [K]} |\tilde{\mu}_{k_n,a+1,a} - \mu_a| \leq \gamma_\zeta$ implies that

$$\begin{aligned}
 \forall a \neq a^*, \quad &\frac{(\tilde{\mu}_{k_n,a^*+1,a^*} - \tilde{\mu}_{k_n,a+1,a}) \min\{3\epsilon, \tilde{\mu}_{k_n,a^*+1,a^*} - \tilde{\mu}_{k_n,a+1,a}\}}{(n-1)/N_{n,a^*} + (n-1)/N_{n,a}} \geq \frac{2\sigma^2(1 - \zeta)}{T_{\epsilon,\beta}^*(\boldsymbol{\nu})}, \\
 &\frac{n-1}{N_{n,a^*}} + \frac{n-1}{N_{n,a}} \leq \frac{\Delta_a \min\{3\epsilon, \Delta_a\}}{2\sigma^2} (1 + \zeta) T_{\epsilon,\beta}^*(\boldsymbol{\nu}).
 \end{aligned}$$

We choose such a γ_ζ . Let $\gamma_\mu > 0$ be such that for $\mathbb{E}_\nu[T_{\mu,\gamma}(\omega_{\epsilon,\beta}^*)] < +\infty$ for all $\gamma \in (0, \gamma_\mu)$. Let $\eta > 0$. Let $\tilde{\gamma}_\mu \in (0, \gamma_\mu)$ as in Lemma 42 for this η . In the following, let us consider $\gamma \in (0, \min\{\tilde{\gamma}_\mu, \gamma_\zeta, \beta/4, \Delta_{\min}/4, (3\epsilon - \max_{a \in [K]} \Delta_a)/2\})$.

Let $\kappa \in (0, 1)$. As above, we can construct N_3 with Lemma 42 and $N_4 \geq N_3$ such that $\mathbb{E}_\nu[N_4] < +\infty$. Let $n \geq N_4/\kappa$ and $(k_{n,a})_{a \in [K]}$ the current phases. Then, we have $\hat{a}_n = a^*$,

$$\max_{a \in [K]} |\tilde{\mu}_{k_{n,a}+1,a} - \mu_a| \leq \gamma, \quad \left\| \frac{N_n}{n-1} - \omega_{\epsilon,\beta}^* \right\|_\infty \leq \gamma \text{ and } \frac{\max_{a \in [K]} T_{k_{n,a}}(a) - 1}{\min_{a \in [K]} T_{k_{n,a}}(a) - 1} \leq 2 + \eta.$$

Depending on the value of the private empirical gap, the stopping condition that is checked is different. For all $a \neq a^*$ such that $\Delta_a < 3\epsilon$, we have $\tilde{\mu}_{k_{n,a^*}+1,a^*} - \tilde{\mu}_{k_{n,a}+1,a} \leq \Delta_a + 2\gamma < 3\epsilon$. For all $a \neq a^*$ such that $\Delta_a \geq 3\epsilon$, we have either $\tilde{\mu}_{k_{n,a^*}+1,a^*} - \tilde{\mu}_{k_{n,a}+1,a} \geq 3\epsilon$ or $\tilde{\mu}_{k_{n,a^*}+1,a^*} - \tilde{\mu}_{k_{n,a}+1,a} \leq 3\epsilon$ and $\Delta_a \leq 3\epsilon + 2\gamma$. Let $a_1 = \arg \min_{a \in [K]} T_{k_{n,a}}(a)$ and $a_2 = \arg \max_{a \in [K]} T_{k_{n,a}}(a)$. Therefore, we obtain similarly that, for all $a \neq a^*$,

$$\frac{(\tilde{\mu}_{k_{n,a^*}+1,a^*} - \tilde{\mu}_{k_{n,a}+1,a}) \min\{3\epsilon, \tilde{\mu}_{k_{n,a^*}+1,a^*} - \tilde{\mu}_{k_{n,a}+1,a}\}}{1/\tilde{N}_{k_{n,a^*}+1,a^*} + 1/\tilde{N}_{k_{n,a}+1,a}} \geq (\min_{b \in [K]} T_{k_{n,b}}(b) - 1) \frac{2\sigma^2(1-\zeta)}{T_{\epsilon,\beta}^*(\nu)}.$$

Similarly, for all $a \neq a^*$ such that $\tilde{\mu}_{k_{n,a^*}+1,a^*} - \tilde{\mu}_{k_{n,a}+1,a} \leq 3\epsilon$, hence $\Delta_a \leq 3\epsilon + 2\gamma$, we have

$$\begin{aligned} \frac{1}{\tilde{N}_{k_{n,a^*}+1,a^*}} + \frac{1}{\tilde{N}_{k_{n,a}+1,a}} &\leq \frac{1}{\min_{b \in [K]} T_{k_{n,b}}(b) - 1} \frac{\Delta_a \min\{3\epsilon, \Delta_a\}}{2\sigma^2} (1 + \zeta) T_{\epsilon,\beta}^*(\nu), \\ \frac{1}{\sqrt{\tilde{N}_{k_{n,a^*}+1,a^*}}} + \frac{1}{\sqrt{\tilde{N}_{k_{n,a}+1,a}}} &\leq \sqrt{\frac{1}{(\min_{b \in [K]} T_{k_{n,b}}(b) - 1)} \frac{\Delta_a \min\{3\epsilon, \Delta_a\}}{\sigma^2} (1 + \zeta) T_{\epsilon,\beta}^*(\nu)}, \\ \frac{1}{2\epsilon^2\sigma^2} \sum_{c \in \{a^*, a\}} \frac{1}{\tilde{N}_{k_{n,c}+1,c}} &\left(\log \frac{3K(k_{n,c} + 1)^s \zeta(s)}{\delta} \right)^2 \\ &\leq \frac{\Delta_a \min\{3\epsilon, \Delta_a\} (1 + \zeta) T_{\epsilon,\beta}^*(\nu)}{4\epsilon^2\sigma^4 (\min_{b \in [K]} T_{k_{n,b}}(b) - 1)} \left(\log(1/\delta) + s \log(\max_{b \in [K]} k_{n,b} + 1) + \log(3K\zeta(s)) \right)^2 \\ \frac{\sqrt{2}}{\epsilon\sigma} \sum_{c \in \{a^*, a\}} \sqrt{\frac{h(\tilde{N}_{k_{n,c}+1,c}, \delta)}{\tilde{N}_{k_{n,c}+1,c}}} \log \left(\frac{3K\zeta(s)(k_{n,c} + 1)^s}{\delta} \right) &\leq \sqrt{\frac{2\Delta_a \min\{3\epsilon, \Delta_a\} (1 + \zeta) T_{\epsilon,\beta}^*(\nu)}{\epsilon^2\sigma^4 (\min_{b \in [K]} T_{k_{n,b}}(b) - 1)}} \\ &\sqrt{h(2^{\max_{b \in [K]} k_{n,b} - 1}, \delta)} \left(\log(1/\delta) + s \log(\max_{b \in [K]} k_{n,b} + 1) + \log(3K\zeta(s)) \right). \end{aligned}$$

Moreover, for all $a \neq a^*$ such that $\tilde{\mu}_{k_{n,a^*}+1,a^*} - \tilde{\mu}_{k_{n,a}+1,a} \geq 3\epsilon$, hence $\Delta_a \geq 3\epsilon$, we have

$$\begin{aligned} \sqrt{\tilde{N}_{k_{n,a}+1,a}} + \sqrt{\tilde{N}_{k_{n,a^*}+1,a^*}} &\leq \sqrt{T_{k_{n,a_2}}(a_2) - 1} \left(\sqrt{\beta + \gamma} + \sqrt{\omega_{\epsilon,\beta,a}^* + \gamma} \right) \\ &\leq \sqrt{\min_{b \in [K]} T_{k_{n,b}}(b) - 1} \sqrt{2(2 + \eta)(\max\{\beta, 1 - \beta\} + \gamma)}, \\ \frac{3\epsilon}{\sqrt{2}\sigma^2} \sum_{c \in \{a^*, a\}} \sqrt{\tilde{N}_{k_{n,c}+1,c} h(\tilde{N}_{k_{n,c}+1,c}, \delta)} & \end{aligned}$$

$$\leq \sqrt{h(2^{\max_{b \in [K]} k_{n,b} - 1}, \delta)} \frac{3\epsilon}{\sigma} \sqrt{\min_{b \in [K]} T_{k_{n,b}}(b) - 1} \sqrt{(2 + \eta)(\max\{\beta, 1 - \beta\} + \gamma)}.$$

Let $T_{k_{n+1}}^+ = \max_b T_{k_{n,b}+1}(b)$, $T_{k_{n+2}}^+ = \max_b T_{k_{n,b}+2}(b)$, $T_{k_{n+1}}^- = \min_b T_{k_{n,b}+1}(b)$, $T_{k_n}^- = \min_b T_{k_{n,b}}(b)$. Let T be a time such that $T \geq T_{k_{n+1}}^+ \geq \kappa T$. Then, $(\max_b k_{n,b} - 1) \log 2 \leq \log T_{k_n}^- + \log(2 + \eta)$. As above, using the DAF(ϵ) update with the GLR stopping rule with $W_{a,b}^{G,\epsilon}$ as in (12) and the stopping threshold $\tilde{c}_{a,b}^{G,\epsilon}$ as in (14), we have

$$\begin{aligned} \min\{\tau_\delta, T\} - \kappa T &\leq \sum_{T \geq T_{k_n}^+ \geq \kappa T} (T_{k_{n+2}}^+ - T_{k_{n+1}}^+) \mathbb{1}(\tau_\delta > T_{k_{n+1}}^+) \\ &\leq \sum_{T \geq T_{k_n}^+ \geq \kappa T} (T_{k_{n+2}}^+ - T_{k_{n+1}}^+) \mathbb{1}(\exists a \neq a^*), \\ &\left(\tilde{\mu}_{k_{n,a^*+1}, a^*} - \tilde{\mu}_{k_{n,a+1}, a} < 3\epsilon, \frac{(\tilde{\mu}_{k_{n,a^*+1}, a^*} - \tilde{\mu}_{k_{n,a+1}, a})^2}{2\sigma^2(1/\tilde{N}_{k_{n,a^*+1}, a^*} + 1/\tilde{N}_{k_{n,a+1}, a})} < \tilde{c}_{a^*, a}^{G,\epsilon}(\tilde{N}_{k_{n+1}}, \delta) \right) \vee \\ &\left(\tilde{\mu}_{k_{n,a^*+1}, a^*} - \tilde{\mu}_{k_{n,a+1}, a} \geq 3\epsilon, \frac{3\epsilon(\tilde{\mu}_{k_{n,a^*+1}, a^*} - \tilde{\mu}_{k_{n,a+1}, a})}{2\sigma^2(1/\tilde{N}_{k_{n,a^*+1}, a^*} + 1/\tilde{N}_{k_{n,a+1}, a})} < \tilde{c}_{a^*, a}^{G,\epsilon}(\tilde{N}_{k_{n+1}}, \delta) \right). \end{aligned}$$

Leveraging the inequalities explicited above, we can upper bound it by a condition which only involves $T_{k_n}^-$ and problem dependent quantities (in a highly convoluted fashion). As above, we define $T_\zeta(\delta)$ as the largest deterministic time such that the above condition is satisfied when replacing $T_{k_n}^-$ by $(1 - \kappa)T$. Then, we obtain similarly that

$$\limsup_{\delta \rightarrow 0} \frac{\mathbb{E}_\nu[\tau_\delta]}{\log(1/\delta)} \leq (2 + \eta)^2 \limsup_{\delta \rightarrow 0} \frac{T_\zeta(\delta)}{\log(1/\delta)}.$$

Dropping the second orders terms in T and $\log(1/\delta)$ and using that $\mathcal{C}_G(x) = x + \mathcal{O}(\log x)$ and $\overline{W}_{-1}(x) = x + \mathcal{O}(\log x)$, tedious manipulations yields that

$$\limsup_{\delta \rightarrow 0} \frac{T_\zeta(\delta)}{\log(1/\delta)} \leq \frac{T_{\epsilon, \beta}^*(\boldsymbol{\nu})}{1 - \kappa} \max\{D_{\gamma, \zeta}^{(1)}(\mu, \epsilon), \mathbb{1}(\Delta_{\max} \geq 3\epsilon) D_{\gamma, \zeta, \eta}^{(2)}(\mu, \epsilon)\},$$

where $g(x, y) = \sqrt{2xy} + y/4$ and

$$\begin{aligned} D_{\gamma, \zeta}^{(1)}(\mu, \epsilon) &= \sup \left\{ x \mid x^2(1 - \zeta) < x + g\left(x, \frac{1 + \zeta}{\epsilon^2 \sigma^4} \max_{\Delta_a \leq 3\epsilon + 2\gamma} \Delta_a \min\{\Delta_a, 3\epsilon\}\right) \right\}, \\ D_{\gamma, \zeta, \eta}^{(2)}(\mu, \epsilon) &= \sup \left\{ x \mid x\sigma(1 - \zeta)/3 < \epsilon\sqrt{x} \sqrt{(2 + \eta)T_{\epsilon, \beta}^*(\boldsymbol{\nu})(\max\{\beta, 1 - \beta\} + \gamma) + \sigma^{-1}} \right\}. \end{aligned}$$

Letting κ, γ, η and ζ goes to zero yields that

$$\limsup_{\delta \rightarrow 0} \frac{\mathbb{E}_\nu[\tau_\delta]}{\log(1/\delta)} \leq 4T_{\epsilon, \beta}^*(\boldsymbol{\nu}) \max\{D_{0,0}^{(1)}(\mu, \epsilon), \mathbb{1}(\Delta_{\max} \geq 3\epsilon) D_{0,0,0}^{(2)}(\mu, \epsilon)\}.$$

Using that $x^2 - 2bx - c < 0$ for all $x \in [0, b(1 + \sqrt{1 + c/b^2})]$, we obtain that

$$D_{0,0,0}^{(2)}(\mu, \epsilon) = \left(\sup \left\{ x \mid x^2 < \sqrt{18\epsilon^2 \sigma^{-2} T_{\epsilon, \beta}^*(\boldsymbol{\nu}) \max\{\beta, 1 - \beta\}} x + 3\sigma^{-2} \right\} \right)^2$$

$$\leq \frac{9\epsilon^2}{2\sigma^2} T_{\epsilon,\beta}^*(\boldsymbol{\nu}) \max\{\beta, 1 - \beta\} \left(1 + \sqrt{1 + \frac{2}{3\epsilon^2 T_{\epsilon,\beta}^*(\boldsymbol{\nu}) \max\{\beta, 1 - \beta\}}} \right)^2,$$

$$D_{0,0}^{(1)}(\mu, \epsilon) = g_1 \left(\max_{\Delta_a \leq 3\epsilon} \frac{\Delta_a}{\sigma^2 \epsilon} \right) \quad \text{with} \quad g_1(y) = \sup \left\{ x \mid x^2 < x + y\sqrt{2x} + y^2/4 \right\}.$$

In more details, we have

$$g_1(0) = \sup \{x \mid x^2 < x\} = 1 \quad \text{and} \quad g_1(12) = \sup \left\{ x \mid x^2 < x + 12\sqrt{2x} + 36 \right\} \leq 10,$$

where the last inequality is obtained by numerical analysis. The function g_2 is obtained by noting that $y(1 + \sqrt{1 + 1/y})^2 = 1 + 2(\sqrt{1 + 1/y} - 1)^{-1}$. When $\Delta_{\max} < 3\epsilon$, we have $T_{\epsilon,\beta}^*(\mu) = T_{\text{KL},\beta}^*(\boldsymbol{\nu})$ where $T_{\text{KL},\beta}^*(\boldsymbol{\nu})$ as in (5) with $\sigma = 1/2$. This concludes the proof of the second result. \blacksquare

Concluding the proof of Theorems 17 and 19. Combining Lemmas 37, 41, 42 and 43 concludes the proof of Theorems 17 and 19. We restrict the result to instances such that $\min_{a \neq b} |\mu_a - \mu_b| > 0$ in order for Lemma 37 to hold. Note that this is an artifact of the asymptotic proof which could be alleviated with more careful considerations. \blacksquare

Appendix F. On the Number of Rounds of Adaptivity

Due to its generality, using the DAF update yields a batched version of any existing FC-BAI algorithm, which satisfies ϵ -global DP. At the end of the episode of arm a (after updating its mean), it is possible to compute the sequence of all the arms to be pulled before the end of the next episode (for another arm), without taking the collected observations into account. In contrast to the classical batched setting where the batch size is fixed, the size of the resulting batches is adaptive and data-dependent.

Let $C(\tau_\delta) = \sum_{a \in [K]} k_{\tau_\delta, a}$ be the number of rounds of adaptivity, where $k_{\tau_\delta, a}$ denotes the number of episodes of arm $a \in [K]$ at stopping time. Using Jensen's inequality, the number of rounds of adaptivity is upper bounded by $\mathbb{E}_\nu [C(\tau_\delta)] \leq K \log_2 \mathbb{E}_\nu [\tau_\delta]$. Therefore, any upper bound on the expected sample complexity directly implies an upper bound on the number of rounds of adaptivity.

One global episode. The multiplicative factor K is incurred because DAF maintains one episode per arm. Alternatively, one can consider one global episode k_n . Formally, we switch phase as soon as all the arms have doubled their empirical counts, i.e. $N_{n,a} \geq 2N_{T_{k_n}, a}$ for all $a \in [K]$. This modification allows to shave the K factor since $\mathbb{E}_\nu [C(\tau_\delta)] \leq \log_2 \mathbb{E}_\nu [\tau_\delta]$. When using one global episode, one can show the same asymptotic upper bound as when we used one episode per arm.

Empirically, the performance is worsen by considering one global episode, hence we recommend to use one episode per arm. A sub-optimal arm a might be sampled more than a^* in early stage due to unlucky first draws. When there is only one global episode, the learner will always have to double the counts of this sub-optimal arm before updating its estimators of the other arms. After realizing that this arm is sub-optimal, it won't be sampled frequently, hence many samples should be collected before ending the episode.

Algorithm 6 Doubling-Per-Arm (DPA)

Input: History \mathcal{H}_n , arm $a \in [K]$.
Initialization: For all $a \in [K]$, $T_1(a) = K + 1$ and $k_{K+1,a} = 1$;
if $N_{n,a} \geq 2N_{T_{k_{n,a}}(a),a}$ **then**
 Change phase $k_{n,a} \leftarrow k_{n,a} + 1$ for this arm a ;
 Set $T_{k_{n,a}}(a) = n$ and $\hat{\mu}_{k_{n,a},a} = N_{T_{k_{n,a}}(a),a}^{-1} \sum_{t \in [T_{k_{n,a}}(a)-1]} r_t \mathbb{1}\{a_t = a\}$;
end if
Return $(\hat{\mu}_{n,a}, N_{n,a})$;

Batched best arm identification In the non-private setting ($\epsilon = +\infty$), we recover Batched Best-Arm Identification (BBAI) in the fixed-confidence setting. One of the question arising in this setting is the following: *Can we solve the BBAI problem with asymptotically optimal sample complexity (up to a constant factor) and a small number of batches?* A slight modification of the above result provides a positive answer.

Without the privacy constraint, there is no need to forget about past observations or to add Laplacian noise. Therefore, the DPA update is better suited for BBAI than the DAF one. Using the DPA update yields an adaptive batched version of any existing FC-BAI algorithm. It is direct to see that the same analysis can be used to study TTUCB with DPA update. Namely, it yields a δ -correct algorithm such that, for all μ with distinct means,

$$\limsup_{\delta \rightarrow 0} \frac{\mathbb{E}_{\nu} [\tau_{\delta}]}{\log(1/\delta)} \leq 2T_{\text{KL},\beta}^*(\nu), \quad \limsup_{\delta \rightarrow 0} (\mathbb{E}_{\nu} [C(\tau_{\delta})] - K \log_2 \log(1/\delta)) \leq K \log_2(2T_{\text{KL},\beta}^*(\nu)).$$

For $\beta = 1/2$, the algorithm is asymptotically optimal (up to a multiplicative factor 4) with solely $\mathcal{O}(K \log_2(T_{\text{KL}}^*(\nu) \log(1/\delta)))$ rounds of adaptivity.

There are already several works studying BBAI (Karnin et al., 2013; Jin et al., 2019, 2023), see Table 1 in Jin et al. (2023) for a detailed comparison. Building on the Exponential-Gap Elimination algorithm (Karnin et al., 2013), Jin et al. (2019) proposed an algorithm achieving an expected sample complexity of the order of $\mathcal{O}(\sum_{a \neq a^*} \Delta_a^{-2} \log(\log(\Delta_a^{-1})/\delta))$ with $\mathcal{O}(\log_{1/\delta}^*(K) \log(\Delta_{\min}^{-1}))$ batches, where $\log_{1/\delta}^*$ is the iterated logarithm function with base $1/\delta$. To the best of our knowledge, existing lower bound on the number of rounds are worst-case bounds. For constant $\delta \in (0, 1)$, Tao et al. (2019) proved that for certain bandit instances, any algorithm that achieves the sample complexity bound obtained in Jin et al. (2019) requires at least $\Omega(\log(\Delta_{\min}^{-1})/\log \log \Delta_{\min}^{-1})$ batches. Jin et al. (2023) proposed the Tri-BBAI algorithm which achieves asymptotic optimality with two rounds of adaptivity (i.e. three phases). An important remark here is that the analysis of Tri-BBAI is purely asymptotic, and it is only δ -correct for sufficiently small δ . As an improvement with similar asymptotic guarantees as well as non-asymptotic ones, they propose Opt-BBAI which uses the same first two phases as Tri-BBAI, then uses successive elimination and checks for best arm elimination.

Appendix G. Extended Experimental Analysis

For both local DP and global DP, we perform additional experiments on six bandit environments with Bernoulli distributions, as defined by (Sajed and Sheffet, 2019), namely

$$\begin{aligned} \mu_1 &= (0.95, 0.9, 0.9, 0.9, 0.5), & \mu_2 &= (0.75, 0.7, 0.7, 0.7, 0.7), \\ \mu_3 &= (0, 0.25, 0.5, 0.75, 1), & \mu_4 &= (0.75, 0.625, 0.5, 0.375, 0.25)\}, \\ \mu_5 &= (0.75, 0.53125, 0.375, 0.28125, 0.25), & \mu_6 &= (0.75, 0.71875, 0.625, 0.46875, 0.25)\}. \end{aligned}$$

For each Bernoulli instance, we implement the algorithms with

$$\epsilon \in \{0.001, 0.005, 0.01, 0.05, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 10, 100, 1000\},$$

for global DP, and

$$\epsilon \in \{0.01, 0.05, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 10, 100\},$$

for local DP.

The risk level is set at $\delta = 0.01$. We verify empirically that the algorithms are δ -correct by running each algorithm 1000 times.

The additional results for local DP are presented in Figure 3. For global DP, the additional results are provided in Figure 4. To show the difference between AdaP-TT and AdaP-TT*, we plot the stopping time not in a logarithmic scale in Figure 5. The additional experiments validate the same conclusions as the ones reached in Section 5.

Remark 44 *To implement the thresholds of AdaP-TT and AdaP-TT*, we use empirical thresholds that we get by approximating the theoretical thresholds. The expressions of the empirical thresholds used can be found in the code here.*




Figure 3: Evolution of the stopping time τ (mean \pm std. over 1000 runs) of CTB-TT and TTUCB with respect to the privacy budget ϵ for $\delta = 10^{-2}$ on different Bernoulli instances. The shaded vertical line separates the two privacy regimes.




Figure 4: Evolution of the stopping time τ (mean \pm std. over 1000 runs) of Imp-AdaP-TT, AdaP-TT, DP-SE, and TTUCB with respect to the privacy budget ϵ for $\delta = 10^{-2}$ on different Bernoulli instances. The shaded vertical line separates the two privacy regimes. Both the x -axis and y -axis are in logarithmic scale.




Figure 5: Evolution of the stopping time τ (mean \pm std. over 1000 runs) of AdaP-TT*, AdaP-TT, DP-SE, and TTUCB with respect to the privacy budget ϵ for $\delta = 10^{-2}$ on different Bernoulli instances. The shaded vertical line separates the two privacy regimes. Only the x -axis is in logarithmic scale.

References

- Y. Abbasi-Yadkori, P. Bartlett, V. Gabillon, A. Malek, and M. Valko. Best of both worlds: Stochastic & adversarial best-arm identification. In *Conference on Learning Theory*, pages 918–949. PMLR, 2018.
- J. Acharya, Z. Sun, and H. Zhang. Differentially private assouad, fano, and le cam. In *Algorithmic Learning Theory*, pages 48–78. PMLR, 2021.
- J.-Y. Audibert, S. Bubeck, and R. Munos. Best Arm Identification in Multi-armed Bandits. In *Conference on Learning Theory*, 2010.
- P. Auer, N. Cesa-Bianchi, and P. Fischer. Finite-time analysis of the multiarmed bandit problem. *Machine learning*, 47(2-3):235–256, 2002.
- M. Aziz, E. Kaufmann, and M.-K. Riviere. On multi-armed bandit designs for dose-finding clinical trials. *The Journal of Machine Learning Research*, 22(1):686–723, 2021.
- A. Azize and D. Basu. When privacy meets partial information: A refined analysis of differentially private bandits. *Advances in Neural Information Processing Systems*, 35:32199–32210, 2022.
- A. Azize and D. Basu. Concentrated differential privacy for bandits. In *2nd IEEE Conference on Secure and Trustworthy Machine Learning*, 2024.
- D. Basu, C. Dimitrakakis, and A. Tossou. Differential privacy for multi-armed bandits: What is it and what is its cost? *arXiv preprint arXiv:1905.12298*, 2019.
- R. E. Bechhofer. A single-sample multiple decision procedure for ranking means of normal populations with known variances. *The Annals of Mathematical Statistics*, pages 16–39, 1954.
- R. E. Bechhofer. A sequential multiple-decision procedure for selecting the best one of several normal populations with a common unknown variance, and its use with various experimental designs. *Biometrics*, 14(3):408–429, 1958.
- T. T. Cai, Y. Wang, and L. Zhang. The cost of privacy: Optimal rates of convergence for parameter estimation with differential privacy. *The Annals of Statistics*, 49(5):2825–2850, 2021.
- A. Carpentier and A. Locatelli. Tight (lower) bounds for the fixed budget best arm identification bandit problem. In *Conference on Learning Theory*, pages 590–604. PMLR, 2016.
- S. Chen, T. Lin, I. King, M. R. Lyu, and W. Chen. Combinatorial pure exploration of multi-armed bandits. *Advances in neural information processing systems*, 27, 2014.
- A. Cheu. Differential privacy in the shuffle model: A survey of separations. *arXiv preprint arXiv:2107.11839*, 2021.

- R. Degenne, W. M. Koolen, and P. Ménard. Non-asymptotic pure exploration by solving games. *Advances in Neural Information Processing Systems*, 32, 2019.
- R. Degenne, H. Shao, and W. Koolen. Structure adaptive algorithms for stochastic bandits. In *International Conference on Machine Learning*, pages 2443–2452. PMLR, 2020.
- J. Dong, A. Roth, and W. J. Su. Gaussian differential privacy. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 84(1):3–37, 2022.
- J. C. Duchi, M. I. Jordan, and M. J. Wainwright. Local privacy and statistical minimax rates. In *2013 IEEE 54th Annual Symposium on Foundations of Computer Science*, pages 429–438. IEEE, 2013.
- C. Dwork and A. Roth. The algorithmic foundations of differential privacy. *Foundations and Trends® in Theoretical Computer Science*, 9(3–4):211–407, 2014.
- C. Dwork, M. Naor, T. Pitassi, and G. N. Rothblum. Differential privacy under continual observation. In *ACM symposium on Theory of computing*, pages 715–724. ACM, 2010a.
- C. Dwork, M. Naor, T. Pitassi, G. N. Rothblum, and S. Yekhanin. Pan-private streaming algorithms. In *Innovations in Computer Science*, pages 66–80, 2010b.
- E. Even-Dar, S. Mannor, Y. Mansour, and S. Mahadevan. Action elimination and stopping conditions for the multi-armed bandit and reinforcement learning problems. *Journal of machine learning research*, 7(6), 2006.
- V. Gabillon, M. Ghavamzadeh, and A. Lazaric. Best arm identification: A unified approach to fixed budget and fixed confidence. *Advances in Neural Information Processing Systems*, 25, 2012.
- A. Garivier and E. Kaufmann. Optimal best arm identification with fixed confidence. In *Conference on Learning Theory*, pages 998–1027. PMLR, 2016.
- A. M. Girgis, D. Data, S. Diggavi, A. T. Suresh, and P. Kairouz. On the renyi differential privacy of the shuffle model. In *ACM SIGSAC Conference on Computer and Communications Security*, pages 2321–2341, 2021.
- K. Jamieson and R. Nowak. Best-arm identification algorithms for multi-armed bandits in the fixed confidence setting. In *Conference on Information Sciences and Systems (CISS)*, pages 1–6. IEEE, 2014.
- K. Jamieson and A. Talwalkar. Non-stochastic best arm identification and hyperparameter optimization. In *Artificial intelligence and statistics*, pages 240–248. PMLR, 2016.
- K. Jamieson, M. Malloy, R. Nowak, and S. Bubeck. lil’ucb: An optimal exploration algorithm for multi-armed bandits. In *Conference on Learning Theory*, pages 423–439. PMLR, 2014.
- T. Jin, J. Shi, X. Xiao, and E. Chen. Efficient pure exploration in adaptive round model. *Advances in Neural Information Processing Systems*, 32, 2019.

- T. Jin, Y. Yang, J. Tang, X. Xiao, and P. Xu. Optimal batched best arm identification. *arXiv preprint arXiv:2310.14129*, 2023.
- M. Jourdan and R. Degenne. Non-asymptotic analysis of a ucb-based top two algorithm. *Advances in Neural Information Processing Systems*, 36, 2024.
- M. Jourdan, R. Degenne, D. Baudry, R. de Heide, and E. Kaufmann. Top two algorithms revisited. *Advances in Neural Information Processing Systems*, 35:26791–26803, 2022.
- M. Jourdan, R. Degenne, and E. Kaufmann. Dealing with unknown variances in best-arm identification. *International Conference on Algorithmic Learning Theory*, 2023.
- M. Jourdan, R. Degenne, and E. Kaufmann. An ε -best-arm identification algorithm for fixed-confidence and beyond. *Advances in Neural Information Processing Systems*, 36, 2024.
- P. Kairouz, K. Bonawitz, and D. Ramage. Discrete distribution estimation under local privacy. In *International Conference on Machine Learning*, pages 2436–2444. PMLR, 2016.
- D. S. Kalogerias, K. E. Nikolakakis, A. D. Sarwate, and O. Sheffet. Quantile multi-armed bandits: Optimal best-arm identification and a differentially private scheme. *IEEE Journal on Selected Areas in Information Theory*, 2(2):534–548, 2021.
- S. Kalyanakrishnan, A. Tewari, P. Auer, and P. Stone. Pac subset selection in stochastic multi-armed bandits. In *International Conference on Machine Learning*, volume 12, pages 655–662, 2012.
- Z. Karnin, T. Koren, and O. Somekh. Almost optimal exploration in multi-armed bandits. In *International Conference on Machine Learning*. PMLR, 2013.
- V. Karwa and S. Vadhan. Finite Sample Differentially Private Confidence Intervals. In *9th Innovations in Theoretical Computer Science Conference (ITCS 2018)*, volume 94. Schloss Dagstuhl – Leibniz-Zentrum für Informatik, 2018.
- E. Kaufmann and S. Kalyanakrishnan. Information complexity in bandit subset selection. In *Conference on Learning Theory*, pages 228–251. PMLR, 2013.
- E. Kaufmann and W. M. Koolen. Mixture martingales revisited with applications to sequential tests and confidence intervals. *Journal of Machine Learning Research*, 22(246):1–44, 2021.
- E. Kaufmann, O. Cappé, and A. Garivier. On the complexity of best arm identification in multi-armed bandit models. *Journal of Machine Learning Research*, 17:1–42, 2016.
- T. Lattimore and C. Szepesvári. *Bandit algorithms*. Cambridge University Press, 2020.
- L. Li, K. Jamieson, G. DeSalvo, A. Rostamizadeh, and A. Talwalkar. Hyperband: A novel bandit-based approach to hyperparameter optimization. *The Journal of Machine Learning Research*, 18(1):6765–6816, 2017.

- P. J. Libin, T. Verstraeten, D. M. Roijers, J. Grujic, K. Theys, P. Lemey, and A. Nowé. Bayesian best-arm identification for selecting influenza mitigation strategies. In *Machine Learning and Knowledge Discovery in Databases: European Conference, ECML PKDD 2018*, 2019.
- S. Lindståhl, A. Proutiere, and A. Johnsson. Measurement-based admission control in sliced networks: A best arm identification approach. In *GLOBECOM 2022-2022 IEEE Global Communications Conference*, pages 1484–1490. IEEE, 2022.
- D. E. Losada, D. Elswiler, M. Harvey, and C. Trattner. A day at the races: using best arm identification algorithms to reduce the cost of information retrieval user studies. *Applied Intelligence*, 52(5):5617–5632, 2022.
- S. Mannor and J. N. Tsitsiklis. The sample complexity of exploration in the multi-armed bandit problem. *Journal of Machine Learning Research*, 5(Jun):623–648, 2004.
- I. Mironov. Rényi differential privacy. In *2017 IEEE 30th computer security foundations symposium (CSF)*, pages 263–275. IEEE, 2017.
- N. Mishra and A. Thakurta. (Nearly) optimal differentially private stochastic multi-arm bandits. In *Conference on Uncertainty in Artificial Intelligence*, 2015.
- S. Neel and A. Roth. Mitigating bias in adaptive data gathering via differential privacy. In *International Conference on Machine Learning*, pages 3720–3729. PMLR, 2018.
- K. E. Nikolakakis, D. S. Kalogerias, and A. D. Sarwate. Optimal rates for learning hidden tree structures. *arXiv preprint arXiv:1909.09596*, 2019.
- C. Qin, D. Klabjan, and D. Russo. Improving the expected improvement algorithm. *Advances in Neural Information Processing Systems*, 30, 2017.
- W. Ren, X. Zhou, J. Liu, and N. B. Shroff. Multi-armed bandits with local differential privacy. *arXiv preprint arXiv:2007.03121*, 2020.
- A. Rio, M. Barlier, I. Colin, and M. Soare. Multi-agent best arm identification with private communications. In *International Conference on Machine Learning*, 2023.
- D. Russo. Simple bayesian algorithms for best arm identification. In *Conference on Learning Theory*, pages 1417–1418. PMLR, 2016.
- T. Sajed and O. Sheffet. An optimal private stochastic-mab algorithm based on optimal private stopping rule. In *International Conference on Machine Learning*, pages 5579–5588. PMLR, 2019.
- X. Shang, R. Heide, P. Menard, E. Kaufmann, and M. Valko. Fixed-confidence guarantees for bayesian best-arm identification. In *International Conference on Artificial Intelligence and Statistics*, pages 1823–1832. PMLR, 2020.
- R. Shariff and O. Sheffet. Differentially private contextual linear bandits. In *Advances in Neural Information Processing Systems*, pages 4296–4306, 2018.

- M. Soare, A. Lazaric, and R. Munos. Best-arm identification in linear bandits. *Advances in Neural Information Processing Systems*, 27, 2014.
- C. Tao, Q. Zhang, and Y. Zhou. Collaborative learning with limited interaction: Tight bounds for distributed exploration in multi-armed bandits. In *2019 IEEE 60th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 126–146, 2019.
- A. C. Tossou and C. Dimitrakakis. Algorithms for differentially private multi-armed bandits. In *Thirtieth AAAI Conference on Artificial Intelligence*, 2016.
- K. Tucker, J. Branson, M. Dilleen, S. Hollis, P. Loughlin, M. J. Nixon, and Z. Williams. Protecting patient privacy when sharing patient-level data from clinical trials. *BMC medical research methodology*, 16(1):5–14, 2016.
- P.-A. Wang, R.-C. Tzeng, and A. Proutiere. Fast pure exploration via frank-wolfe. *Advances in Neural Information Processing Systems*, 34:5810–5821, 2021.
- S. L. Warner. Randomized response: A survey technique for eliminating evasive answer bias. *Journal of the American statistical association*, pages 63–69, 1965.
- W. You, C. Qin, Z. Wang, and S. Yang. Information-directed selection for top-two algorithms. In *Conference on Learning Theory*, pages 2850–2851. PMLR, 2023a.
- W. You, C. Qin, Z. Wang, and S. Yang. Information-directed selection for top-two algorithms. In *Conference on Learning Theory*, pages 2850–2851. PMLR, 2023b.
- Y. Zhou, X. Chen, and J. Li. Optimal pac multiple arm identification with applications to crowdsourcing. In *International Conference on Machine Learning*, pages 217–225. PMLR, 2014.