# A NORMAL VERSION OF BRAUER'S HEIGHT ZERO CONJECTURE

ALEXANDER MORETÓ AND A. A. SCHAEFFER FRY

ABSTRACT. The celebrated Itô–Michler theorem asserts that a prime p does not divide the degree of any irreducible character of a finite group G if and only if G has a normal and abelian Sylow p-subgroup. The principal block case of the recently-proven Brauer's height zero conjecture isolates the abelian part in the Itô– Michler theorem. In this paper, we show that the normal part can also be isolated in a similar way. This is a consequence of work on a strong form of the so-called Brauer's height zero conjecture for two primes of Malle and Navarro. Using our techniques, we also provide an alternate proof of this conjecture.

# 1. INTRODUCTION

The Itô-Michler theorem [43] was one of the first results obtained as a consequence of the announcement of the classification of finite simple groups. This fundamental result in representation theory of finite groups asserts that a prime p does not divide the degree of any complex irreducible character of a finite group G if and only if Ghas a normal abelian Sylow p-subgroup. This theorem was motivated by Brauer's height zero conjecture, which was proposed in 1955, long-before the classification was completed, and was recently proved in [40]. Brauer's height zero conjecture can be considered as a block version of the Itô-Michler theorem, asserting that if B is a Brauer p-block of G, then every irreducible character in B has height zero if and only if the defect group of the block is abelian. In particular, the principal block case proven in [39] says that a Sylow p-subgroup is abelian if and only if p does not divide the degree of any irreducible character in the principal p-block  $B_p(G)$  of G.

*Date*: June 18, 2024.

<sup>2010</sup> Mathematics Subject Classification. Primary 20C15, 20C20, 20C33.

Key words and phrases. Principal block, character degree, Sylow subgroup, Hall subgroup.

We thank Gunter Malle for his comments on an earlier draft of this manuscript, as well as for helpful conversations on this problem during the Isaac Newton Institute program "Groups, Representations, and Applications: New Perspectives", supported by EPSRC grant EP/R014604/1, where this project was started. The first author is supported by Ministerio de Ciencia e Innovación (Grants PID2019-103854GB-I00 and PID2022-137612NB-I00 funded by MCIN/AEI/10.13039/501100011033 and "ERDF A way of making Europe"). He also acknowledges support by Generalitat Valenciana CIAICO/2021/163. The second author gratefully acknowledges support from the National Science Foundation, Award No. DMS-2100912, and her former institution, Metropolitan State University of Denver, which holds the award and allows her to serve as PI.

Noting that Brauer's height zero conjecture isolates the "abelian" part of the Itô-Michler theorem, isolating the "normal" part is another interesting problem (see, e.g. [48]) that has also received attention. As pointed out by Navarro in [48, p. 1365] "when we restrict our attention to the characters in the principal block then the commutativity of P is characterized (by the principal block case of Brauer's height zero conjecture) while the normality of P is lost". Our first main result shows that, perhaps surprisingly, the normality of P is also characterized when we restrict our attention to the characters in the principal block for the primes different from p.

**Theorem A.** Let G be a finite group, let p be a prime and let  $P \in \text{Syl}_p(G)$ . Then  $P \trianglelefteq G$  if and only if for every prime  $q \neq p$  that divides |G| and every  $\chi \in \text{Irr}(B_q(G))$ , p does not divide  $\chi(1)$ .

This is therefore a dual version of Brauer's height zero conjecture for principal blocks. We remark that Malle and Navarro found a different subset of  $\operatorname{Irr}(G)$  that characterizes the normality of a Sylow *p*-subgroup. In [37] they proved that  $P \leq G$  if and only if *p* does not divide  $\chi(1)$  for every irreducible constituent  $\chi$  of the permutation character  $(1_P)^G$ . (See also [17] for a further refinement of the statement.) Note, however, that while the union of  $\operatorname{Irr}(B_q(G))$  for  $q \neq p$  prime divisor of |G| can be easily determined from the character table of *G*, it is still not known whether  $(1_P)^G$  is determined by the character table of *G*. (However, see [46] or [49, Thm. 1.2], where this is proved for *p*-solvable groups.)

Using Brauer's height zero conjecture for principal blocks [39] and Theorem A, we deduce the following refinement of the Itô–Michler theorem.

**Corollary B.** Let G be a finite group and let p be a prime. Then G has a normal and abelian Sylow p-subgroup if and only if p does not divide  $\chi(1)$  for every  $\chi$  that belongs to some principal block for some prime divisor of |G|.

This project was motivated by a conjecture of Malle and Navarro called Brauer's height zero conjecture for two primes: if G is a finite group and p and q are different prime divisors of |G|, then G has a nilpotent Hall  $\{p, q\}$ -subgroup if and only if p does not divide the degree of the characters in the principal q-block and q does not divide the degrees of the characters in the principal p-block. (Actually, they allowed the case p = q in their statement, which is exactly Brauer's height zero conjecture for principal blocks, proved in [39].) The "only if" direction of this conjecture was proved in [38], while the "if" direction has recently been proven in [30]. This is one of the recent results in modular representation theory that take two primes into account (see, for instance, [50, 38, 30]).

In [38], Malle and Navarro pointed out that it is an interesting question to characterize when the irreducible characters of the principal *p*-block  $B_p(G)$  have degree not divisible by *q*. Our original goal was to address this question. We obtain a block version of the Itô–Michler theorem. **Theorem C.** Let G be a finite group and let p and q be two different primes. Assume that S is not a composition factor of G if (S, p, q) is one of the 5 exceptions listed in Theorem 2.1 below. If q does not divide the degree of any irreducible character in  $B_p(G)$ , then a Sylow p-subgroup of G normalizes a Sylow q-subgroup of G.

We remark that the conclusion of Theorem C is equivalent to G having a p-nilpotent Hall  $\{p,q\}$ -subgroup. There does not seem to be much room for improvement in Theorem C. We will see in Theorem 2.1 below that the exceptions listed are necessary. It is not clear how to strengthen the conclusion either. As  $G = PSL_2(11)$  for p = 2and q = 3 shows, the hypothesis that q does not divide the degrees of the characters in the principal p-block does not imply that G has a nilpotent Hall  $\{p,q\}$ -subgroup. In fact, in this example G has a p-nilpotent Hall  $\{p,q\}$ -subgroup isomorphic to  $D_{12}$ but it also has Hall  $\{p,q\}$ -subgroups isomorphic to  $A_4$  which are not p-nilpotent. This shows that the hypothesis does not imply that Hall subgroups are conjugate.

Further, the converse of Theorem C does not hold. There are many simple groups with a Sylow *p*-subgroup that normalizes a Sylow *q*-subgroup but such that *q* divides the degree of some irreducible character in the principal *p*-block, for instance  $A_5$  for p = 3 and q = 2. The  $\{p, q\}$ -separable case of Theorem C for arbitrary blocks is the main result of [53]. As pointed out in [38], the group 6.A<sub>7</sub> shows that the conclusion of Theorem C does not hold for arbitrary non-principal *p*-blocks of full defect.

Theorem A, and hence Corollary B, are pleasant consequences of Theorem C. Using the techniques developed to prove Theorem C, we further provide a short proof of Brauer's height zero conjecture for two primes (see Theorem 4.1 below). Therefore, our work does not rely on [30].

In Section 2, we prove Theorem C for almost simple groups. In Section 3, we obtain some other results that will be used in the proof of Theorem C. We complete our proof of Brauer's height zero conjecture for two primes in Section 4 and the proof of Theorems A and C in Section 5. We also obtain the principal block case of a theorem that Navarro [47] proved in 2004 assuming the Alperin–McKay conjecture; see Theorem 5.3. We conclude in Section 6 with a version of Brauer's height zero conjecture for any number of primes and conjecturing a Galois version and a height-zero version of Theorem A, which we prove for solvable groups.

# 2. Almost simple groups

Throughout, we continue to let  $B_p(G)$  denote the principal *p*-block of a group G. If N is a normal subgroup of a group G and  $\theta$  is an irreducible character of N, we write  $Irr(G|\theta)$  to denote the set of irreducible constituents of the induced character  $\theta^G$ . Our notation is standard and follows [25, 27, 45]. Here we begin our discussion on almost simple groups.

First, we consider sporadic and alternating groups.

**Theorem 2.1.** Let G be an almost simple group with socle a sporadic simple group S or the Tits group  ${}^{2}F_{4}(2)'$  and let p and q be two different primes dividing |G|. Then the principal p-block of G has an irreducible character of degree divisible by q unless (S, p, q) is one of the following:  $(S, p, q) = (J_{1}, 2, 3), (J_{1}, 2, 5), (J_{1}, 3, 5), (J_{1}, 5, 3),$  $(J_{4}, 5, 7), (J_{4}, 7, 5), (J_{4}, 7, 11), (M_{11}, 5, 3), or (M_{22}, 7, 2)$ . Among these exceptional cases, G does not have a p-nilpotent Hall  $\{p, q\}$ -subgroup if and only if (S, p, q) is in the following list:

- $(J_1, 2, 3),$
- $(J_1, 2, 5),$
- $(J_4, 7, 11),$
- $(M_{11}, 5, 3)$ , or
- $(M_{22}, 7, 2).$

*Proof.* The first part can be checked with GAP [15]. The second part can be checked with GAP, [60], and [56]. Note that by [56, Thm. 8.2],  $M_{22}$  does not contain a Hall  $\{7, 2\}$ -subgroup, and hence neither does  $M_{22}$ .

**Theorem 2.2.** Let G be an almost simple group with socle an alternating group and let p and q be two different primes dividing |G|. Then the principal p-block contains an irreducible character of degree divisible by q.

*Proof.* Let  $S = A_n$ . If n = 6 this can be checked with GAP. If  $n \neq 6$  this was proved in [18, Thm. C] when q < p and in [19, Thm. A] when q > p.

We now turn our attention to groups S of Lie type. First, we consider the case that  $r_0 \in \{p, q\}$ , where  $r_0$  is the defining characteristic for S.

**Lemma 2.3.** Let p and q be two different primes. Let S be a simple group of Lie type such that p and q divide |S| and S is defined in characteristic p or q. Then there exists  $\chi \in \operatorname{Irr}(B_p(S))$  of degree divisible by q.

*Proof.* First, assume that S is defined in characteristic p. Then  $B_p(S) = \operatorname{Irr}(S) \setminus \{\operatorname{St}_S\}$ , where  $\operatorname{St}_S$  is the Steinberg character, by [6, Thm. 6.18]. Since S is simple, the Itô-Michler theorem [43, Thm. 2.3] tells us that either  $q \nmid |S|$ , or  $q \mid \chi(1)$  for some  $\chi \in \operatorname{Irr}(S)$  (and hence some  $\chi \in \operatorname{Irr}(B_p(S))$ ), since  $\operatorname{St}_S$  has degree a power of p).

Next, suppose that S is defined in characteristic q. Then by [33, Thm. 6.8], either every nontrivial unipotent character of S has degree divisible by q, or  $q \leq 3$ and S is  $B_n(2) \cong C_n(2)$  with  $n \geq 3$ ,  $B_2(2)'$ ,  $G_2(2)'$ ,  $F_4(2)$ , or  $G_2(3)$ . Since  $B_p(S)$ necessarily contains a nontrivial unipotent character (see, e.g. [40, Lem. 3.15]), we may assume that S is one of these exceptions. If  $S = B_2(2)'$ ,  $B_3(2)$ ,  $G_2(2)'$ ,  $F_4(2)$ , or  $G_2(3)$ , we may check in GAP that  $B_p(S)$  contains a character of degree divisible by 2, respectively 3, for every other prime p dividing |S|. So, assume  $S = B_n(2)$  with  $n \geq 4$ . Here [33, Thm. 6.8] yields that there are only five unipotent characters of odd degree, but  $B_p(S)$  contains  $k(2e, \lfloor \frac{n}{e} \rfloor) \geq 6$  unipotent characters (see, e.g., the

4

discussion before [35, Prop. 5.4]), where e is the order of 4 modulo p and k(2e, w) is the number of irreducible characters of the imprimitive complex reflection group G(2e, 1, w). Hence we see  $B_p(S)$  must contain a unipotent character of degree divisible by 2, as desired.

For the remainder of this section, the primary case is then that neither p nor q are the defining characteristic for S. We will rely on and build upon the work of [38] in this situation.

For an integer *m* relatively prime to a prime *p*, we will let  $d_p(m)$  denote the order of *m* modulo *p* if *p* is odd and modulo 4 if p = 2.

2.1. Exceptional Groups. We begin with the case of exceptional types, but excluding Suzuki and Ree groups for now. We remark that, throughout,  $E_6^{\epsilon}(r)$  will denote  $E_6(r)$  when  $\epsilon = 1$  and  ${}^2E_6(r)$  when  $\epsilon = -1$ .

**Lemma 2.4.** Let  $H = \mathbf{H}^F$  be a quasisimple group of Lie type such that  $\mathbf{H}$  is a simple, simply connected algebraic group of exceptional type (including the case  $H = {}^{3}\mathbf{D}_{4}(r)$ ) and  $F : \mathbf{H} \to \mathbf{H}$  is a Frobenius endomorphism defining H over  $\mathbb{F}_r$ . Let G = H/Z for some  $Z \leq \mathbf{Z}(H)$ . Let p and q be two primes dividing |G|. Assume that there is no  $\chi \in \operatorname{Irr}(B_p(G))$  with  $q \mid \chi(1)$ . Then neither p nor q divides r and:

- (i) q is odd;
- (ii)  $d_p(r) = d_q(r);$
- (iii) a Sylow q-subgroup Q of G is abelian; and
- (iv) there is a Sylow p-subgroup P of G such that P normalizes Q.
- (v) Further, either there is a Sylow p-subgroup P of G such that [P,Q] = 1 or  $(H, d_a(r) = d_p(r), p)$  is one of:
  - $(E_6(r), 2, 2)$
  - $(^{2}E_{6}(r), 1, 2)$
  - $({}^{3}D_{4}(r), 1, 3)$
  - $({}^{3}D_{4}(r), 2, 3)$

*Proof.* Throughout, the notation for unipotent characters is taken from [7, Sec. 13.9]. Further, we use  $\Phi_i := \Phi_i(r)$  to denote the *i*th cyclotomic polynomial in r.

Write  $S := H/\mathbb{Z}(H)$ , and note that  $G \in \{H, S\}$  in these cases. Further, letting  $H^* = (\mathbf{H}^*)^F$  with  $(\mathbf{H}^*, F)$  dual to  $(\mathbf{H}, F)$  (and hence  $\mathbf{H}^*$  is simple of adjoint type), we have  $S \cong [H^*, H^*]$  and  $\tilde{S} \cong H^*$ , where  $\tilde{S}$  is the group of inner-diagonal automorphisms of S. (See, e.g. [41, Prop. 24.21, Tab. 22.1] and the subsequent discussions, since S is of exceptional type.) Since we have  $\operatorname{Irr}(B_p(S)) \subseteq \operatorname{Irr}(B_p(G))$ , viewed by inflation, we have by Lemma 2.3 that  $r = r_0^f$  for some prime  $r_0 \notin \{p, q\}$ . Let  $d_p := d_p(r)$  and  $d_q := d_q(r)$ .

Now, we claim that q is odd. Indeed, if q = 2, then since the group  $[H^*, H^*]$  is simple, [1, Thm. 2.2] yields that there is a p-element  $s \in [H^*, H^*]$  such that its centralizer in  $[H^*, H^*]$  has even index. Since this group is index 2 or odd in  $H^*$ ,

the same is true for the centralizer in  $H^*$ . Let  $\chi_s$  be the semisimple character of H corresponding to the class of s, so that  $\chi_s(1) = [H^* : \mathbf{C}_{H^*}(s)]_{r'}$  is even. Further,  $\chi_s$  is trivial on  $\mathbf{Z}(H)$  by [51, Lem. 4.4(ii) and Rem. 4.6] and lies in  $B_p(H)$  using [23, Cor. 3.4] if  $\mathbf{Z}(\mathbf{H})$  is connected or [6, Thm. 21.13] if p is good for  $\mathbf{H}$  and  $\mathbf{C}_{\mathbf{H}^*}(s)$  is connected. Namely, the latter holds when  $p \nmid |\mathbf{Z}(H)|$  by [41, Exer. 20.16]. So, we are left to consider the case p = 3 and  $S = \mathbf{E}_6(r)$  or  ${}^2\mathbf{E}_6(r)$ . In these cases, we see by [7, Sec. 13.9] that there is a unipotent character whose degree is divisible by 2 but not by 3. (Namely, we may take the character  $\phi_{20,2}$  and  $\phi_{4,1}$ , respectively.) But since  $B_3(H)$  is the unique unipotent block containing unipotent characters of 3'-degree (this follows from [33, Thm. 6.6] and [11, Thm. A]) and unipotent characters are trivial on the center, we see these lie in  $B_3(G)$ .

So, we see that q is odd. Now, consider the case that  $q \mid |\mathbf{Z}(H)| = |S/S|$ . Then  $S = \mathrm{E}_{6}^{\epsilon}(r)$  with  $q = 3 \mid (r - \epsilon)$ . If  $S = \mathrm{E}_{6}(r)$  with  $q = 3 \mid (r - 1)$ , we see that the characters  $\phi_{81,10}, \phi_{6.1}$ , and  $\phi_{90,8}$  in the notation of [7, Sec. 13.9] have degree divisible by 3. However, using the explicit decompositions in [4, Tables 1-2] when  $d_{p}$  is non-regular, and the fact that  $B_{p}(H)$  is the unique unipotent block containing unipotent characters of p'-degree if  $d_{p}$  is regular (again this follows from [33, Thm. 6.6] and [11, Thm. A]), we see that for each p, at least one of these three characters lies in  $B_{p}(H)$ . When  $S = {}^{2}\mathrm{E}_{6}(r)$  with  $3 \mid (r + 1)$ , the same holds, with the characters  $\phi_{9,6}', \phi_{2,4}', \phi_{6,6}''$ .

Now, assume that  $q \nmid |\mathbf{Z}(H)| = [\widetilde{S} : S]$ , so that a Sylow q-subgroup Q of S can be identified with one of H and of  $H^* \cong \widetilde{S}$ . Suppose that there exists some p-element  $s \in H$  such that the corresponding semisimple character  $\chi_s \in \operatorname{Irr}(\widetilde{S})$  satisfies  $q \mid \chi_s(1)$ . By [23, Cor. 3.4], we have  $\chi_s \in \operatorname{Irr}(B_p(\widetilde{S}))$ . Then since  $q \nmid |\widetilde{S}/S|$ , we obtain that there is some  $\chi \in \operatorname{Irr}(B_p(S)) \subseteq \operatorname{Irr}(B_p(G))$  with  $q \mid \chi(1)$ . Since  $\chi_s(1) = [H : \mathbf{C}_H(s)]_{r'}$ , we may assume that every p-element  $s \in H$  centralizes a Sylow q-subgroup of H. It follows by [33, Thm. 5.9] that every such s also centralizes a Sylow  $d_q$ -torus of  $(\mathbf{H}, F)$ .

Now, by [1, Prop. 2.12] and also applying [34, Prop. 2.2], we have either Q is abelian,  $\mathbf{C}_H(Q) = \mathbf{Z}(H)\mathbf{Z}(Q)$ , or (H,q) is as in the list [1, Prop. 2.12](b)-(g) and  $d_p, d_q \in \{1, 2\}$ , with  $d_p = d_q$  unless possibly if p = 2. If  $\mathbf{C}_H(Q) = \mathbf{Z}(H)\mathbf{Z}(Q)$ , we obtain every *p*-element  $s \in H$  lies in  $\mathbf{Z}(H)$ , which is not true.

Next, suppose that (H,q) is as in the list [1, Prop. 2.12](b)-(g). In each case, we will exhibit a unipotent character whose degree is divisible by q and lies in  $B_p(H)$ , which will follow from arguments like above and [4, Thm. 3.10] and [28, Thm. A]. In case (b),(c), we have  $q = 5 | (r - \epsilon)$  and  $H = E_6^{\epsilon}(r)$  and the characters  $\phi_{20,2}$  or  $\phi_{4,1}$ , in the respective cases  $\epsilon = 1, \epsilon = -1$ , lie in  $B_p(H)$  and have degree divisible by 5. In case (d), we have  $(H,q) = (E_7(r),3)$ , and we see  $\phi_{27,2}$  has degree divisible by 3 and must lie in  $B_p(H)$ . In case (e), we have  $(H,q) = (E_7(r),5)$ , and  $\phi_{105,15}$  has degree divisible by 5 and lies in  $B_p(H)$ . In case (f) and (g), we have  $(H,q) = (E_7(r),7)$ , resp.  $(E_8(r),7)$ , and we see  $\phi_{7,1}$ , resp.  $\phi_{35,2}$ , has degree divisible by 7 and lies in  $B_p(H)$ .

Н	$d_q$	$d_p$	$\chi \in \operatorname{Irr}(B_p(H))$ with $q \mid \chi(1)$
${}^{3}\mathrm{D}_{4}(r)$	1	3	${}^{3}\mathrm{D}_{4}[1]$
${}^{3}\mathrm{D}_{4}(r)$	2	6	$\phi_{2,2}$
$E_6(r)$	2  or  4	1	$\phi_{64,4}$
$E_6(r)$	6	3	$\phi_{64,4}$
${}^{2}\mathrm{E}_{6}(r)$	1  or  4	2	${}^{2}A_{5}, 1$
${}^{2}\mathrm{E}_{6}(r)$	3	6	${}^{2}A_{5}, 1$
$E_7(r)$	3	1	$\phi_{27,2}$
$E_7(r)$	4	1  or  2	$\phi_{168,6}$
$E_7(r)$	6	2	$\phi_{27,2}$

TABLE 1. Possible  $(H, d_q, d_p)$  when  $d_p \neq d_q$  and  $\Phi_{d_p} \mid |\mathbf{C}_H(\mathbb{S}_{d_q})|$  for  $\mathbb{S}_{d_q}$  a Sylow  $d_q$ -torus of **H** that is not maximal

It follows that Q is abelian. Suppose first that P is also abelian. Note that both p and q divide a unique cyclotomic factor of the generic order of H, by [34, Lem. 2.1, Prop. 2.2]. If  $d_p = d_q$ , there is a Sylow  $d_q$ -torus containing a Sylow p-subgroup P and a Sylow q-subgroup Q, so that [P, Q] = 1.

So, assume  $d_p \neq d_q$ . Then by applying [34, Prop. 2.2], we have by [1, Prop. 2.13] that the Sylow  $d_q$ -tori of **H** are not maximal tori, and hence  $d_q$  is as in [1, Table 1]. Recall our assumption that any *p*-element centralizes a Sylow  $d_q$ -torus. Since  $d_p \neq d_q$  and *p* divides exactly one cyclotomic polynomial in the generic order of *H*, by the structure of centralizers of Sylow  $d_q$ -tori given in [16, Table 3.3], we then have  $(H, d_q, d_p)$  is as in Table 1. In the last column, we list a unipotent character in  $B_p(H)$  with degree divisible by *q*, which can be seen by arguing in the same ways as above.

So now assume that P is nonabelian. Then note that  $p \leq 7$ . Further, we have  $p \leq 3$  if  $H \in {G_2(r), {}^{3}D_4(r), F_4(r)}$ . If  $H = E_6^{\epsilon}(r)$ , we have either  $p \leq 3$  or p = 5 and  $d_p = 1$  or 2, depending on whether  $\epsilon = 1$  or -1. If  $H = E_7(r)$ , then  $p \leq 7$  and  $d_p \in {1,2}$ . If  $H = E_8(r)$  then  $p \leq 7$  and  $d_p \in {1,2}$  if p = 7.

First, suppose that  $H = {}^{3}\mathbb{D}_{4}(r)$ . Then there is a unique unipotent 2-block by [6, Thm. 21.14], and we see from the character degrees in [7, Sec. 13.9] that for each possible  $d_{q}$ , there is a unipotent character of degree divisible by q. So we assume p = 3. If  $d_{q} \in \{1, 2\}$ , we are in the situation that a Sylow  $d_{q}$ -torus is not maximal, and we see from Table 1 that  $d_{3} = d_{q}$ . (This gives one of the situations in part (v) of our conclusion.) If  $d_{q} = 3$  or 6, we have  $d_{q}$  is a regular number for H and a Sylow  $d_{q}$ -torus  $\mathbb{S}_{d_{q}}$  of  $\mathbf{H}$  is maximal, so that  $3 = p \mid |\mathbf{C}_{H}(\mathbb{S}_{d_{q}})| = |\mathbb{S}_{d_{q}}^{F}|$ , so  $3 = p \mid \Phi_{d_{q}}$ . In particular, this means that  $d_{q} = 3^{a}d_{3}$  for some  $a \geq 0$ . Then  $d_{3} = 1$ , respectively 2. In this case, we see that the element  $s_{5}$ , respectively  $s_{10}$  in [9], chosen to have 3-power order larger than 3, does not centralize a Sylow q-subgroup. If  $d_{q} = 12$ , we see again  $3 \mid \Phi_{12}$ , a contradiction. Next, suppose  $H = E_6(r)$ . If  $d_p = 1$ , we have the principal series unipotent characters  $\phi_{64,4}$ ,  $\phi_{60,5}$ , and  $\phi_{81,6}$  must lie in  $B_p(H)$ , and at least one of these has degree divisible by q if  $d_q \neq 1$ . If  $d_p = d_q = 1$ , we may consider a p-element s of maximal order, which does not divide the exponent of a Sylow 1-torus, and hence s does not centralize a Sylow  $d_q$ -torus.

Hence, we may assume that  $d_p = 2$  and  $p \in \{2,3\}$ . Suppose that  $d_q = 2, 4$ , or 6, so that a Sylow  $d_q$ -torus is not maximal. If  $d_q = 6$ , then we see in Table 1 that  $p \mid \Phi_3 \Phi_6$ and we have p = 3. Then applying the Ennola duality proved in [4, Thm. 3.3], we see it suffices to exhibit a principal series unipotent character for  $H^- := {}^2E_6(r)$  in the case  $d_p = 1$  whose degree is divisible by  $\Phi_3$ . The character  $\phi_{9,6'}$  of  $H^-$  satisfies these properties. If  $d_q = 4$ , then we see p = 2 and in this case it similarly suffices to note that the character  $\phi_{4,1}$  of  $H^-$  is a principal series unipotent character with degree divisible by  $\Phi_4$ . Now suppose  $d_q = 2$ . When p = 3, we again see that the maximal order of a 3-element does not divide the size of the centralizer of a Sylow 2-torus, whose order polynomial is divisible only by the cyclotomic polynomials  $\Phi_1$ and  $\Phi_2$ . This leaves only the case p = 2, as indicated in the statement.

Then we may assume that a Sylow  $d_q$ -torus is a maximal torus of **H**. Either  $d_q$  is regular, and hence  $d_q = p^a d_p$  for some  $a \ge 0$  since p divides the order of a self-centralizing Sylow  $d_q$ -torus, or  $d_q = 5$ . When  $d_q = 5$ , similar to before by the Ennola duality [4, Thm. 3.3], it suffices to note that the unipotent character  $\phi_{4,1}$  of  $H^- := {}^2E_6(r)$  is a principal series character in  $B_p(H^-)$  when  $d_p = 1$  whose degree is divisible by  $\Phi_{10}$ . So we finally suppose that  $d_q$  is regular, so  $d_q = p^a d_p$ . Since we are also assuming a Sylow  $d_q$ -torus is maximal, the only option is  $d_q = 8$ . Here we again appeal to the case of  $H^-$ , with  $d_q = 8$ . The principal series unipotent character  $\phi_{2,4'}$  of  $H^-$  has degree divisible by  $\Phi_8$ , yielding the result in this case.

The cases for  $H = {}^{2}E_{6}(r)$  not yet considered follow from the case of  $E_{6}(r)$  by the Ennola duality proved in [4, Thm. 3.3] and at times arguing as above with *p*-elements of maximal order.

Next, suppose  $H = E_7(r)$ . Recall that again  $d_p \in \{1, 2\}$  in this case. The case  $d_p = 1$  is completed very similar to the case of  $E_6(r)$  above. Namely, for each  $d_q \neq 1$  such that  $\Phi_{d_q}$  divides the order polynomial of H, there is a principal series unipotent character of degree divisible by  $\Phi_{d_q}$ , and when  $d_p = d_q = 1$ , we may again consider a *p*-element of maximal order. The case  $d_p = 2$  then follows by the Ennola duality proved in [4, Thm. 3.3]. The cases of  $H = F_4(r)$  and  $G_2(r)$  are completed similarly.

Finally, let  $H = E_8(r)$ . Here the Sylow  $d_q$ -torus is maximal. Then if  $d_q$  is regular, we have  $d_q = p^a d_p$  as before. If  $d_p \in \{1, 2\}$ , we may argue similarly to the preceeding cases. So, suppose p = 5 and  $d_p = 4$ . Here,  $d_p$  is a regular number, so as before,  $B_p(H)$  is the unique block containing p'-degree unipotent characters. If  $d_q$  is regular, we see that  $d_q \in \{4, 20\}$ . Again taking s to be a 5-element of maximal order, we see |s|does not divide the exponent of a Sylow  $\Phi_{d_q}$ -torus, and hence s does not centralize a Sylow q-subgroup. If  $d_q$  is not regular, then  $d_q \in \{7, 9, 14, 18\}$ , and we see the unipotent character  $\phi_{112,3}$  has degree divisible by  $\Phi_{d_q}$  if  $d_q \neq 9$ , but not divisible by p = 5. Similarly, the unipotent character  $\phi_{567,6}$  has degree divisible by  $\Phi_9$  but not by p = 5.

Now, suppose that we are in one of the situations in (v). Let  $d_p = d_q =: d$ . We may identify  $P \in \operatorname{Syl}_p(G)$  and  $Q \in \operatorname{Syl}_q(G)$  with Sylow subgroups of H. If p is odd, then by [20, Thm. 4.10.2], we have P is conjugate to a group of the form  $P_T \rtimes P_W$ , where  $P_T$  lies in a Sylow d-torus of H and  $P_W$  normalizes that torus. But similarly,  $Q = Q_T$  lies in a Sylow d-torus. Then these may be chosen such that  $[P_T, Q_T] = 1$ and  $P_W$  normalizes  $Q_T$ . If p = 2, then we still have  $P_T \triangleleft P$  and  $[P_T, Q_T] = 1$ , and  $P/P_T$  still normalizes the Sylow d-torus. This yields (iv), completing the proof.  $\Box$ 

We next address the case of Suzuki and Ree groups.

**Lemma 2.5.** Let G be a simple Suzuki or Ree group  ${}^{2}G_{2}(3^{2k+1})$ ,  ${}^{2}B_{2}(2^{2k+1})$ , or  ${}^{2}F_{4}(2^{2k+1})$ . Write  $r^{2} = 3^{2k+1}$ , respectively  $2^{2k+1}$ . Let p and q be primes dividing |G| such that  $Irr(B_{p}(G))$  contains no character of degree divisible by q. Then neither p nor q divides  $r^{2}$  and:

- q is odd;
- $d_p(r^2) = d_q(r^2);$
- a Sylow q-subgroup Q of G is abelian; and
- there is a Sylow p-subgroup P of G such that P normalizes Q.
- Further, either there is a Sylow p-subgroup P of G such that [P,Q] = 1 or  $G = {}^{2}G_{2}(r^{2})$  with p = 2 and  $q \mid (r^{2} + 1)$ .

*Proof.* Note that by Lemma 2.3, neither p nor q divides  $r^2$ , and as in the beginning of the proof of Lemma 2.4, we have  $q \neq 2$ .

First suppose that  $G = {}^{2}G_{2}(3^{2k+1})$ . If p = 2, we see by [57, p. 74] that there is a character of degree divisible by q in  $B_{2}(G)$  unless  $q \mid (r^{2} + 1)$ . Note that in the latter situation, we have Q is abelian and we also have  $4 \mid (r^{2} + 1)$  so that  $d_{2}(3^{2k+1}) = d_{q}(3^{2k+1})$ . Further, from [24, Sec. 2.1], we see that  $N_{G}(Q)$  contains a Sylow 2-subgroup in this case. Now suppose that both p and q are larger than 3. Then both P and Q are abelian and we see from [24, Sec. 4.1] that there is  $\chi \in Irr(B_{p}(G))$ with  $q \mid \chi(1)$  whenever  $d_{p} \neq d_{q}$ . So  $d_{p} = d_{q}$  and [P, Q] = 1 as both can be chosen to lie in the same cyclic torus.

Next, let  $G = {}^{2}B_{2}(2^{2k+1})$ . Then both p and q are at least 5 and the Sylow p and q-subgroups are cyclic. If  $d_{p}(r^{2}) \neq d_{q}(r^{2})$ , we see from [5] that  $B_{p}(G)$  contains a character of degree divisible by q. Hence we again have  $d_{p} = d_{q}$  and P and Q can be chosen to lie in the same cyclic torus, giving [P, Q] = 1.

Finally, let  $G = {}^{2}F_{4}(2^{2k+1})$ . Then we see using [32, Bem. 1] that  $B_{p}(G)$  contains a character of degree divisible by q except possibly when  $d_{p}(r^{2}) = d_{q}(r^{2})$  and neither p nor q is 3. In that situation, there again exist P and Q in the same torus and [P,Q] = 1. 2.2. Classical Groups. In the next several statements, we turn our attention to classical groups, aiming to obtain a statement similar to those for exceptional groups above. Throughout, we will continue to let  $p \neq q$  be two primes and let r be a power of some prime  $r_0$ . As usual,  $\operatorname{GL}_n(\epsilon r)$  will mean  $\operatorname{GL}_n(r)$  if  $\epsilon = 1$  and  $\operatorname{GU}_n(r)$  if  $\epsilon = -1$ , and similar for related groups like  $\operatorname{SL}_n(\epsilon r)$  and their projective versions.

As before, our strategy will often be to exhibit a unipotent character in the principal p-block whose degree is divisible by q. The next lemma covers an initial case.

**Lemma 2.6.** Let  $\widetilde{G} := \operatorname{GL}_n(\epsilon r)$  with  $n \geq 3$  and let  $p \neq q$  be primes not dividing r but dividing  $|\widetilde{G}|$ . Let  $e_p := d_p(\epsilon r)$  and  $e_q := d_q(\epsilon r)$ . Assume that  $e_p \in \{1, 2\}$  and that  $e_q > e_p$  but  $(n, e_q) \notin \{(3, 3), (4, 2)\}$ . Then  $B_p(\widetilde{G})$  contains a unipotent character of degree divisible by q.

*Proof.* The unipotent characters of  $\tilde{G}$  correspond to partitions of n. If p = 2, all unipotent characters lie in  $B_p(\tilde{G})$  by [6, Thm. 21.14]. Otherwise, by [13] the unipotent characters in  $B_p(\tilde{G})$  are those whose  $e_p$ -cores are the same as that of the trivial character, which corresponds to the partition (n). Namely,  $B_p(\tilde{G})$  contains all unipotent characters when  $e_p = 1$  and contains those with 2-core  $(n \mod 2)$  if  $e_p = 2$ .

First, assume  $e_q \ge 3$ . Then it therefore suffices to show that there is a unipotent character of degree divisible by q corresponding to a partition with 2-core  $(n \mod 2)$ .

Let  $d_q := d_q(r)$ . Now, by [33, Cor. 6.6], a unipotent character has degree divisible by q if it does not lie in a  $d_q$ -Harish-Chandra series  $(L, \lambda)$  with  $L = \mathbf{C}_{\tilde{G}}(\mathbb{S}_{d_q})$  the centralizer of a Sylow  $d_q$ -torus and  $\lambda$  a unipotent character of L. Further, letting  $n = we_q + m$  with  $0 \leq m < e_q$ , we have any unipotent character lying in such a  $d_q$ -Harish-Chandra series would correspond to a partition of n with  $e_q$ -core of size m, using [16, Cors. 4.6.5 and 4.6.7]. Hence, it suffices to exhibit a partition of n with  $e_q$ -core not of size m and 2-core  $(n \mod 2)$ .

If  $m \notin \{0, e_q - 1\}$ , then the partition  $\tau := (1^{m+1}, e_q - 1)$  is an  $e_q$ -core of size  $e_q + m > m$ . The partition  $\lambda_1 := (1^{e_q(w-1)+m+1}, e_q - 1)$  then has  $e_q$ -core  $\tau$  and has 2-core  $(n \mod 2)$  if  $e_q$  is even. If  $e_q$  and m are both odd, then the partition  $\lambda_2 := (1^{m+1}, e_q - 1 + e_q(w-1))$  instead has  $e_q$ -core  $\tau$  and has 2-core  $(n \mod 2)$ . Note that if  $e_q = m + 2$  or if m = 1, then one of  $\lambda_1$  or  $\lambda_2$  satisfies our statement.

Now suppose  $m \notin \{1, e_q - 2\}$  (and now including the cases m = 0 or  $e_q - 1$ ). Then the partition  $\mu := (1^m, 2, e_q - 2)$  is an  $e_q$ -core for  $e_q \ge 4$ , again of size  $e_q + m$ . If  $e_q \ge 4$  is even or m is even with  $e_q \ge 5$  odd, then  $\lambda_3 := (1^m, 2, e_q - 2 + e_q(w - 1))$ has  $e_q$ -core  $\mu$  and 2-core  $(n \mod 2)$ .

We are therefore left with the case  $e_q = 3$  and  $m \in \{0, 2\}$ . If m = 2 and  $e_q = 3$ , the partition (1, 1, n - 2) has 3-core (1, 1, 3) and 2-core  $(n \mod 2)$ . So, assume m = 0 and  $e_q = 3$ , so  $3 \mid n$ . Our assumption  $(n, e_q) \neq (3, 3)$  means that  $n \geq 6$ , so the partition (2, n - 2) has 3-core (2, 4) and 2-core  $(n \mod 2)$ . This completes the proof when  $e_q \geq 3$ .

Finally, assume  $e_p = 1$  and  $e_q = 2$ . From the discussions above, we have  $B_p(\tilde{G})$  contains every unipotent character and it suffices to know that there is a partition of n with 2-core different from  $(n \mod 2)$ . If n is odd, the partition (1, n - 1) has 2-core (1, 2), satisfying this condition. If  $n \ge 6$  is even, the partition (1, 2, n - 3) has 2-core (1, 2, 3), again satisfying our condition. Given our assumption  $(n, e_q) \ne (4, 2)$ , this completes the proof.

In the following, let r be a power of a prime  $r_0$  and let  $\widetilde{H}$  be one of  $\operatorname{GL}_n(\epsilon r)$  with  $n \geq 5$ ,  $\operatorname{Sp}_{2n}(r)$  with  $n \geq 2$ ,  $\operatorname{SO}_{2n+1}(r)$  with  $n \geq 3$ , or  $\operatorname{SO}_{2n}^{\pm}(r)$  with  $n \geq 4$ . Assume that both p and q divide  $|\widetilde{H}|$  and that  $r_0 \notin \{p,q\}$ . Let  $d_p := d_p(r)$  and  $d_q := d_q(r)$  and let  $e_p$  denote  $d_p(\epsilon r)$  if  $\widetilde{H} = \operatorname{GL}_n(\epsilon r)$  and  $e_p := d_p(r^2)$  otherwise, and similar for  $e_q$ . In the latter cases, let  $\epsilon_p$  and  $\epsilon_q$  be the numbers in  $\{\pm 1\}$  such that  $p \mid (r^{e_p} - \epsilon_p)$  and  $q \mid (r^{e_q} - \epsilon_q)$ . The next observation extends the last. (Note that the cases  $n \in \{3, 4\}$  for  $\operatorname{GL}_n(\epsilon r)$  are covered in the previous lemma.)

**Lemma 2.7.** Keep the hypotheses and notation in the preceding paragraph. Assume that  $e_p \mid e_q$  but that  $e_p \neq e_q$ . Further assume that  $\epsilon_p = \epsilon_q$  (when relevant) and that  $p \mid (r^{e_q} - \epsilon_q)$ . Then  $B_p(\widetilde{H})$  contains a unipotent character of degree divisible by q.

*Proof.* Note that by Lemma 2.6, we may assume that  $e_p \ge 3$  in the case that  $H = \operatorname{GL}_n(\epsilon r)$  with  $n \ge 5$ . Throughout, let  $n = we_q + m$  with  $0 \le m < e_q$ , and let  $m = w_1e_p + m_1$  with  $0 \le m_1 < e_p$ .

The unipotent characters of  $\widetilde{H}$  are labeled by partitions of n if  $\widetilde{H} = \operatorname{GL}_n(\epsilon r)$  and by certain symbols  $\binom{X}{Y}$  in the other cases (see e.g. [7, Section 13.8]). If p = 2, then every unipotent character lies in  $B_p(\widetilde{H})$ , by [6, Thm. 21.14]. If p is odd and  $\widetilde{H} = \operatorname{GL}_n(\epsilon r)$ , then two unipotent characters lie in the same p-block of  $\widetilde{H}$  if they correspond to partitions that have the same  $e_p$ -core. In the other cases, if p is odd then two unipotent characters lie in the same p-block of  $\widetilde{H}$  if they correspond to symbols with the same  $e_p$ -core, respectively  $e_p$ -cocore, if  $\epsilon_p = 1$ , respectively  $\epsilon_p = -1$  (see [13, Thm. (7A)] and [14, (12A) and (13B)]). Hence the unipotent characters in  $B_p(\operatorname{GL}_n(\epsilon r))$  are those whose corresponding partition has  $e_p$ -core  $(m_1)$ . For  $\widetilde{H} = \operatorname{Sp}_{2n}(r)$  or  $\operatorname{SO}_{2n+1}(r)$ , the unipotent characters in  $B_p(\widetilde{H})$  are those whose symbol has  $e_p$ -core (resp.  $e_p$ -cocore)  $\binom{m_1}{0}$ . In  $\operatorname{SO}_{2n}^+(r)$ , the unipotent characters in  $B_p(\widetilde{H})$  have  $e_p$ -core (resp.  $e_p$ -cocore) the same as that of  $\binom{0}{0}$  and in  $\operatorname{SO}_{2n}^-(r)$ , they have  $e_p$ -core (resp.  $e_p$ -cocore) the same as that of  $\binom{0,n}{0}$ . In all cases, we will simply say that a partition or symbol has trivial  $e_p$ -core (resp. trivial  $e_p$ -cocore) if this is the case.

On the other hand, we again have by [33, Cor. 6.6], that any unipotent character not lying in a  $d_q$ -Harish-Chandra series  $(L, \lambda)$  with L the centralizer in  $\tilde{H}$  of a Sylow  $d_q$ -torus of the underlying algebraic group must necessarily have degree divisible by q. Using [16, Cors. 4.6.5, 4.6.7, and 4.6.16], the characters in such a series would have  $e_q$ -core (resp.  $e_q$ -cocore) of rank m.

In Table 2, we illustrate in each case an  $e_q$ -core  $\mu$  of rank  $m + e_q$  that further has trivial  $e_p$ -core. (Recall here that  $e_p > 2$  in the  $\operatorname{GL}_n(\epsilon r)$  case.) If  $\mu = (\mu_1, \ldots, \mu_t)$ is a partition, one (or both) of either the partition  $(1^{e_q(w-1)}, \mu)$  or the partition  $(\mu_1, \ldots, \mu_{t-1}, \mu_t + e_q(w-1))$  then has trivial  $e_p$ -core and  $e_q$ -core  $\mu$ . If  $\mu = \binom{X}{Y}$  is a symbol with  $X = x_1, \ldots, x_t$ , then the symbol  $\binom{X'}{Y}$  with  $X' = x_1, \ldots, x_{t-1}, x_t + e_q(w-1)$ has trivial  $e_p$ -core and has  $e_q$ -core  $\mu$  (so in particular, of rank not m). Hence if  $\widetilde{H} = \operatorname{GL}_n(\epsilon r)$  or if  $\epsilon_p = 1 = \epsilon_q$ , we have shown that there is a unipotent character of degree divisible by q lying in  $B_p(\widetilde{H})$ .

Now suppose  $\epsilon_p = -1 = \epsilon_q$  with  $\widetilde{H}$  one of the symplectic or special orthogonal groups and  $p \mid (r^{e_q} + 1)$ . Note that in this situation,  $\frac{e_q}{e_p}$  must be odd. Let  $\mu$  be the symbol in Table 2. Then Olsson's theory of  $e_p$ -twists (see [54, p. 235]) yields a symbol  $\overline{\mu}$  (the " $e_p$ -twist of  $\mu$ ") with rank  $m + e_q$  and with trivial  $e_p$ -cocore. Further, in each case we can see that our  $\overline{\mu}$  is an  $e_q$ -cocore, using that  $e_q/e_p$  is odd and the definition of the  $e_p$ -twist. Now, write  $\overline{\mu} = \binom{x_1, \dots, x_t}{y_1, \dots, y_s}$  with (without loss of generality)  $x_t \geq y_s$ . Then we see that if w - 1 is even, then the symbol  $S := \binom{x_1, \dots, x_{t-1}, x_t + e_q(w-1)}{y_1, \dots, y_s}$  has rank  $n, e_q$ -cocore  $\overline{\mu}$ , and has trivial  $e_p$ -cocore since it can be converted to  $\overline{\mu}$  by removing  $e_p$ -cohooks. If w - 1 is instead odd, then since  $e_q/e_p$  is odd, the same properties hold for  $S' := \binom{x_1, \dots, x_{t-1}}{y_1, \dots, y_s, x_t + e_q(w-1)}$ . We remark that in the latter case, if  $\widetilde{H} = \operatorname{SO}_{2n}^{\epsilon}(r)$ , then  $\overline{\mu}$  corresponds to a unipotent character of  $\operatorname{SO}_{2(m+e_q)}^{-\epsilon}(r)$ , which is also the type of the  $d_q$ -split Levi subgroup corresponding to the symbol S'. Hence in all cases, we have exhibited a unipotent character in  $B_p(\widetilde{H})$  of degree divisible by q.

**Proposition 2.8.** Let  $H = \mathbf{H}^F$  be a quasisimple group of Lie type such that  $\mathbf{H}$  is a simple, simply connected algebraic group and  $F \colon \mathbf{H} \to \mathbf{H}$  is a Frobenius endomorphism defining H over  $\mathbb{F}_r$ . Let G be a group such that G = H/Z for some  $Z \leq \mathbf{Z}(H)$ . Let p and q be two primes dividing |G|. Assume that  $B_p(G)$  contains no irreducible character of degree divisible by q. Then neither p nor q divides r and:

- (i) q is odd;
- (ii)  $d_p(r) = d_q(r);$
- (iii) a Sylow q-subgroup Q of G is abelian; and
- (iv) there is a Sylow p-subgroup P of G such that P normalizes Q.
- (v) Further, either there is a Sylow p-subgroup P of G such that [P,Q] = 1, or  $G = \text{PSL}_p(\epsilon r)$  with  $p \mid (r \epsilon)$ , or  $(H, d_q(r), p)$  are as in (v) of Lemma 2.4.

*Proof.* First, note that if H is of exceptional type, the statement follows from Lemma 2.4. We again have by Lemma 2.3 that r is a power of  $r_0 \notin \{p, q\}$ .

Then we may assume H is of classical type and may write  $S := H/\mathbf{Z}(H)$  so that  $S = \mathrm{PSL}_n(\epsilon r)$  with  $n \ge 2$  and  $\epsilon \in \{\pm 1\}$ ;  $S = \mathrm{PSp}_{2n}(r)$  with  $n \ge 2$ ,  $S = \mathrm{P}\Omega_{2n+1}(r)$ 

	$\operatorname{GL}_n(\epsilon r)$	$SO_{2n+1}(r)$ or $Sp_{2n}(r)$
m = 0	$(e_p, e_q - e_p)$	$\begin{pmatrix} 0, e_q - e_p + 1 \\ e_p \end{pmatrix}$
	$e_p \neq 1$	r
$m_1 = 0, m > 0$	$(1^m, e_q)$	$\begin{pmatrix} 1,2,\dots,m,e_q+m\\ 0,1,\dots,m-1 \end{pmatrix}$
$m_1 > 0, w_1 = 0$	$(1^{e_q-e_p}, m+e_p)$	$\binom{1,2,\dots,e_q-e_p-1,e_q-e_p,e_q+m}{0,1,\dots,e_q-e_p-1}$
$m_1 > 0, w_1 = 1$	$(1^{e_p}, m_1 + 1, e_q - 1)$	$\binom{1,2,\dots,e_q-1,e_q+e_p}{0,1,\dots,e_q-1,m+e_q-e_p}$
$m_1 > 0, w_1 > 2$	$(1^{e_q-1}, e_p+1, m-e_p)$	$\begin{pmatrix} 1,2,\dots,e_q-1,e_q+e_p \\ 0,1,\dots,0 \end{pmatrix}$
1 / 1 -		$(0,1,,e_q-1,m+e_q-e_p)$
	$\frac{\mathrm{SO}_{2n}^{+}(r)}{\mathrm{SO}_{2n}^{+}(r)}$	$\frac{(0,1,\dots,e_q-1,m+e_q-e_p)}{SO_{2n}^{-}(r)}$
m = 0	$\frac{\operatorname{SO}_{2n}^{+}(r)}{\binom{e_{q}-e_{p}}{e_{p}}}$	$\frac{SO_{2n}^{-}(r)}{\begin{pmatrix} e_{p}, e_{q} - e_{p} \\ \emptyset \end{pmatrix}}$
m = 0 $m_1 = 0, m > 0$	$\frac{SO_{2n}^{+}(r)}{\begin{pmatrix} e_{q}-e_{p} \\ e_{p} \end{pmatrix}} \\ \frac{\begin{pmatrix} 1,2,,m,e_{q}+m \\ 0,1,,m \end{pmatrix}}{\begin{pmatrix} 1,2,,m \end{pmatrix}}$	$\begin{array}{c c} & & & \\ & & & \\ \hline & & & \\ & & \\ \hline & & \\ & & \\ & & \\ \hline & & \\ & &$
m = 0 $m_1 = 0, m > 0$ $m_1 > 0, w_1 = 0$	$\frac{SO_{2n}^{+}(r)}{\begin{pmatrix} e_{q}-e_{p} \\ e_{p} \end{pmatrix}} \\ \frac{\begin{pmatrix} 1,2,,m,e_{q}+m \\ 0,1,,m \end{pmatrix}}{\begin{pmatrix} 1,2,,e_{q}-e_{p},e_{q}+m \\ 0,1,,e_{q}-e_{p} \end{pmatrix}}$	$\begin{array}{c c} & & & \\ \hline & & & \\ \hline & & & \\ \hline \hline & & \\ \hline & & \\ \hline \hline \\ \hline \\$
m = 0 $m_1 = 0, m > 0$ $m_1 > 0, w_1 = 0$ $m_1 > 0, w_1 = 1$	$\frac{SO_{2n}^{+}(r)}{\begin{pmatrix} e_{q}-e_{p} \\ e_{p} \end{pmatrix}} \\ \hline \begin{pmatrix} (1,2,,m,e_{q}+m) \\ 0,1,,m \end{pmatrix}}{\begin{pmatrix} (1,2,,e_{q}-e_{p}-1,e_{q}-e_{p},e_{q}+m) \\ 0,1,,e_{q}-e_{p} \end{pmatrix}} \\ \hline \begin{pmatrix} (1,2,,e_{q}-e_{p}-1,e_{q}-e_{p},e_{q}+m) \\ 0,1,,e_{q}-1,e_{q} \end{pmatrix}} \\ \hline \end{pmatrix}$	$\begin{array}{c c} & (0,1,,e_q-1,m+e_q-e_p) \\ \hline & & SO_{2n}^-(r) \\ & (e_p,e_q-e_p) \\ & (0,1,2,,m,e_q+m) \\ & (1,,e_q-e_p-1,e_q-e_p,e_q+m) \\ & (0,1,,e_q-e_p-1,e_q+e_p) \\ & (1,2,,e_q-1,e_q+e_p) \\ & (0,1,,e_q-1,e_q,m+e_q-e_p) \end{array}$

TABLE 2. Some  $e_q$ -cores of rank  $e_q + m$  with trivial  $e_p$ -core when  $e_p \mid e_q, e_p \neq e_q$ 

with  $n \geq 3$ , or  $S = P\Omega_{2n}^{\pm}(r)$  with  $n \geq 4$ . Let  $(\mathbf{H}^*, F)$  be dual to  $(\mathbf{H}, F)$  and write  $H^* := (\mathbf{H}^*)^F$ . Note that  $[H^*, H^*]$  is also a simple group of Lie type by [41, Prop. 24.21].

Let  $s \in H^*$  with order a power of p. We claim that a semisimple character  $\chi_s \in \operatorname{Irr}(H)$  corresponding to the  $H^*$ -class of s lies in  $B_p(H)$ . Indeed, if  $p \nmid |\mathbf{Z}(H)|$ , then p > 2 in the cases of the orthogonal and symplectic types, so we have p is good for H,  $\mathbf{C}_{\mathbf{H}^*}(s)$  is connected by [41, Exer. 20.16], and  $\chi_s \in \operatorname{Irr}(B_p(H))$  by [6, Thm. 21.13]. If instead  $p \mid |\mathbf{Z}(H)|$ , then either p = 2 or  $H = \operatorname{SL}_n(\epsilon r)$  and  $p \mid (n, r - \epsilon)$ . In either of these cases, H has a unique unipotent block by [6, Thm. 21.14] and [13] and hence any semisimple character  $\chi_s \in \operatorname{Irr}(H)$  corresponding to the  $H^*$ -conjugacy class of s lies in  $B_p(H)$  by [6, Thm. 9.12].

Now, if s further lies in  $[H^*, H^*]$ , then  $\chi_s$  is trivial on  $\mathbf{Z}(H)$  by [51, Lem. 4.4], and we can view  $\chi_s \in \operatorname{Irr}(B_p(G))$  by deflation. (If  $p \nmid |\mathbf{Z}(H)| = |H^*/[H^*, H^*]|$ , then this condition automatically holds.) In such a case, since  $\chi_s$  has degree  $[H^* : \mathbf{C}_{H^*}(s)]_{r'}$ , we see s must centralize a Sylow q-subgroup of  $H^*$ .

By [1, Thm. 2.2], we see that for any odd p,  $[H^*, H^*]$  necessarily contains a pelement that does not centralize a Sylow 2-subgroup of  $[H^*, H^*]$ , and hence does not centralize a Sylow 2-subgroup of  $H^*$ . We therefore see that q is odd, giving (i).

Write  $d_p := d_p(r)$  and  $d_q := d_q(r)$ . We will next show (ii), that  $d_p = d_q$ . Now, let  $\widetilde{H} := \operatorname{GL}_n(\epsilon r)$ ,  $\operatorname{Sp}_{2n}(r)$ ,  $\operatorname{SO}_{2n+1}(r)$ , or  $\operatorname{SO}_{2n}^{\epsilon}(r)$ , respectively, in the cases  $S = \operatorname{PSL}_n(\epsilon r)$ ,  $\operatorname{PSp}_{2n}(r)$ ,  $\operatorname{PO}_{2n+1}(r)$  or  $\operatorname{PO}_{2n}^{\epsilon}(r)$ . Then S can naturally be viewed as a normal subgroup of  $\widetilde{H}/\mathbf{Z}(\widetilde{H})$ . First let  $S = \text{PSL}_n(\epsilon r)$  and write  $e_q := d_q(\epsilon r)$  and  $e_p := d_p(\epsilon r)$ . We claim that  $e_p \mid e_q$ . If  $e_p = 1$ , this is certainly true, so suppose  $e_p \neq 1$ , and hence  $p \nmid (r - \epsilon)$ . Then any semisimple *p*-element  $\tilde{s} \in [\tilde{H}^*, \tilde{H}^*]$  satisfies its image  $s \in H^*$  has connected centralizer  $\mathbf{C}_{\mathbf{H}^*}(s)$  by [41, Exer. 20.16], and therefore the corresponding semisimple character  $\chi_{\tilde{s}} \in \text{Irr}(B_p(\tilde{H}))$  restricts irreducibly to  $\text{Irr}(B_p(H))$ . Further, such a character is also trivial on  $\mathbf{Z}(\tilde{H})$  by [51, Lem. 4.4], so the restriction can be viewed in  $\text{Irr}(B_p(G))$ . Then such a  $\tilde{s}$  must centralize a Sylow *q*-subgroup of  $\tilde{H}^*$ , as otherwise  $\chi_{\tilde{s}}$  has degree divisible by *q*. Then  $\tilde{s}$  centralizes a Sylow  $d_q$ -torus  $\mathbb{S}_{d_q}$  of  $\tilde{H}^*$  by [33, Thm. 5.9].

Here  $\mathbf{C}_{\tilde{H}^*}(\mathbb{S}_{d_q}) \cong T \times \mathrm{GL}_m(\epsilon r)$  with  $n = we_q + m$ ,  $0 \leq m < e_q$ , and  $T \cong \mathrm{GL}_1((\epsilon r)^{e_q})^w$  a torus centralizing a Sylow  $d_q$ -torus of  $\mathrm{GL}_{we_q}(\epsilon r)$  (see [16, Ex. 3.5.14]). If  $p \mid |\mathrm{GL}_m(\epsilon r)|$  but  $p \nmid |T|$ , note that  $e_p \leq m < e_q$  and that any element of p-power order has at most m nontrivial eigenvalues. But writing  $m = w_1e_p + m_1$  with  $m_1 < e_p$ , we can find an element  $s_1$  of  $\mathrm{GL}_m(\epsilon r)$  with  $w_1e_p$  nontrivial eigenvalues of p-power order. Since 2m < n, we can embed the element  $\mathrm{diag}(s_1, s_1^{-1}, I)$  into  $\mathrm{GL}_n(\epsilon r)$ , and this element has  $2w_1e_p > m$  nontrivial eigenvalues, a contradiction. This forces  $p \mid |T|$  after all, and hence  $e_p \mid e_q$ .

If  $n \geq 5$ , then by Lemmas 2.6 and 2.7, we have  $e_p = e_q$ , since unipotent characters are trivial on  $\mathbf{Z}(\widetilde{H})$  and restrict irreducibly to S. But this forces also  $d_p = d_q$ .

Now suppose  $H = SL_2(r)$ . If  $d_p \neq d_q$ , then  $d_p = 1$  and  $d_q = 2$ . Any *p*-element  $s \in [H^*, H^*]$  yields a character  $\chi_s \in Irr(B_p(G))$  of degree (r+1)/2 or (r+1), and hence divisible by q, a contradiction.

Next let  $H = \text{SL}_3(\epsilon r)$ . If  $e_q \neq 3$ , we may again conclude using Lemma 2.6. So assume  $e_q = 3$  and  $e_p = 1$ . Here we see that  $B_p(H)$  is the unique unipotent *p*-block, and that for any nontrivial *p*-element  $s \in [H^*, H^*]$  with  $|s| \mid (r-\epsilon)_p$ , the corresponding semisimple character  $\chi_s \in \text{Irr}(H)$  has degree divisible by  $(r^2 + \epsilon r + 1)$ . Then any such character deflates to one in  $B_p(G)$  with degree divisible by q, a contradiction.

Now let  $H = \mathrm{SL}_4(\epsilon r)$ . Here we may assume by Lemma 2.6 that  $e_q = 2$  and  $e_p = 1$ . Again  $B_p(H)$  is the unique unipotent *p*-block, and the image in  $[H^*, H^*]$  of the *p*-element diag $(\zeta, \zeta^{-1}, 1, 1)$  of  $\tilde{H}^*$ , where  $|\zeta| = (r - \epsilon)_p$  gives a semisimple character  $\chi_s \in \mathrm{Irr}(B_p(H))$  trivial on the center and with degree divisible by  $r + \epsilon$ , and hence q, again giving a contradiction.

Now consider the remaining classical types. Since  $q \neq 2$ , again  $B_p(\tilde{H})$  does not contain a character trivial on the center with degree divisible by q, as this would again yield a character of  $B_p(G)$  with degree divisible by q. Here let  $e_p := d_p(r^2)$ and as before, let  $\epsilon_p \in \{\pm 1\}$  be such that  $p \mid (r^{e_p} - \epsilon_p)$ , and similar for  $e_q, \epsilon_q$ . Again write  $n = we_q + m$  with  $0 \leq m < e_q$ . The structure of  $\mathbf{C}_{\tilde{H}^*}(\mathbb{S}_{d_q})$  is described in [16, Ex. 3.5.15]. We have  $\mathbf{C}_{\tilde{H}^*}(\mathbb{S}_{d_q}) \cong T \times H_m$  where  $H_m$  is of classical type of rank mand  $|T| = (r^{e_q} - 1)^w$  if  $d_q = e_q$  is odd and  $(r^{e_q} + 1)^w$  if  $d_q = 2e_q$  is even. By similar arguments as above, we have  $p \mid (r^{e_q} - 1)$ , resp.  $p \mid (r^{e_q} + 1)$  and this forces  $e_p \mid e_q$ ,  $\epsilon_p = \epsilon_q$ , and  $p \mid (r^{e_q} + 1)$  when  $\epsilon_p = -1$ . Then we are in the situation of Lemma 2.7, and again  $e_p = e_q$ . Then  $d_p = d_q$  since  $d_p = e_p = e_q = d_q$  if  $d_q$  is odd, and  $d_p = 2e_p = 2e_q = d_q$  if  $d_q$  is even. Hence we have shown (ii).

We next turn to (iii), the claim that  $Q \in \text{Syl}_q(G)$  is abelian. Assume even that  $\hat{Q} \in \text{Syl}_q(H)$  is nonabelian. Then by [29, Lems. 2.8–2.10] and since  $d_p = d_q$ , we have  $B_p(H)$  contains a unipotent character of degree divisible by q, unless q = 3,  $H = \text{SL}_3(\epsilon r)$ , and  $e_p = e_3 = 1$ . Then assume we are in the latter situation. Here again  $B_p(H)$  is the unique unipotent p-block, and similar to the proof of (ii), we see that for any p-element  $s \in [H^*, H^*]$ , we have  $\chi_s(1)$  is divisible by 3. Hence any such character deflates to one in  $B_p(G)$  with degree divisible by 3, a contradiction, yielding that  $\hat{Q}$ , and thus Q, must be abelian as claimed.

The argument for (iv) is exactly as in Lemma 2.4. Let  $\hat{P} \in \text{Syl}_p(H)$  and  $\hat{Q} \in \text{Syl}_q(H)$ . If p is odd, then by [20, Thm. 4.10.2], we have  $\hat{P}$  is conjugate to a group of the form  $P_T \rtimes P_W$ , where  $P_T$  lies in a Sylow *d*-torus of H and  $P_W$  normalizes that torus. But similarly,  $\hat{Q} = Q_T$  lies in a Sylow *d*-torus. Then these may be chosen such that  $[P_T, Q_T] = 1$  and  $P_W$  normalizes  $Q_T$ . If p = 2, then we still have  $P_T \triangleleft \hat{P}$  and  $[P_T, Q_T] = 1$ , and  $\hat{P}/P_T$  still normalizes the Sylow *d*-torus. This yields (iv).

Finally, we prove (v). Assume that no  $P \in \text{Syl}_p(G)$  and  $Q \in \text{Syl}_q(G)$  exist satisfying [P,Q] = 1. From above, note that we may assume that  $\hat{P} \in \text{Syl}_p(H)$  is nonabelian, as otherwise  $\hat{P} = P_T$ .

Let  $d := d_p = d_q$  and  $e := e_p = e_q$  and write n = ew + m as before, so that  $p \leq w < q$  since  $\hat{Q}$  is abelian but  $\hat{P}$  is not. Then there exists a *p*-element  $s \in H^*$  such that |s| does not divide the exponent of  $|\mathbb{T}|$ , and hence cannot centralize a Sylow *q*-subgroup of  $H^*$ . Then the corresponding semisimple character  $\chi_s$  of *H* has degree divisible by *q* and lies in  $B_p(H)$ .

If such an s can be chosen in  $[H^*, H^*]$ , then  $\chi_s$  is trivial on  $\mathbf{Z}(H)$  and we obtain a contradiction. So, assume that this is not the case. Then  $p \mid |H^*/[H^*, H^*]| =$  $|\mathbf{Z}(H)|$ . Then either p = 2 or  $p \mid (n, r - \epsilon)$  with  $S = \text{PSL}_n(\epsilon r)$ . In the latter case when p is odd, note that e = 1, and assume that  $2p \leq n$ . Consider  $\zeta \in \mathbb{F}_{r^{2p}}^{\times}$  with order  $(r^p - \epsilon)_p > (r - \epsilon)_p$ . Then an element  $\tilde{s} \in \tilde{H}^*$  with nontrivial eigenvalues  $\{\zeta, \zeta^r, \zeta^{r^2}, \cdots, \zeta^{r^{p-1}}, \zeta^{-1}, \zeta^{-r}, \cdots, \zeta^{-r^{p-1}}\}$  lies in  $[\tilde{H}^*, \tilde{H}^*]$ , giving a character  $\chi_{\tilde{s}} \in$  $\text{Irr}(B_p(\tilde{H}))$  trivial on  $\mathbf{Z}(\tilde{H})$  with degree divisible by q, using the same considerations as above. Note that since q > n here, we have  $q \nmid [\tilde{S} : S]$ , where  $\tilde{S} = \tilde{H}/\mathbf{Z}(\tilde{H})$ . Then the constituents of  $\chi_{\tilde{s}}$  on H also have degree divisible by q, lie in  $B_p(H)$ , and are trivial on  $\mathbf{Z}(H)$ . This is a contradiction, so we must have n = p. This gives one of the listed groups in (v).

Now assume p = 2. Since H is a classical group, there is a unique unipotent 2block by [6, Thm. 21.14], so the characters indexed by any 2-element of  $H^*$  lie in  $B_2(H)$ . Let  $\zeta \in \mathbb{F}_{r^2}$  with  $|\zeta| = (r^2 - 1)_2$ . If  $S = \text{PSL}_n(\epsilon r)$ , we may argue exactly as in the previous paragraph, now using an element  $\tilde{s} \in \tilde{H}^*$  with nontrivial eigenvalues  $\{\zeta, \zeta^r, \zeta^{-1}, \zeta^{-r}\}$ , to see that n = 2. If  $S = \text{PSp}_4(r)$ ,  $\text{PSp}_6(r)$ , or  $\text{P}\Omega_7(r)$ , then since  $q \mid (r^2 - 1)$ , we see explicitly that there is a unipotent character (which therefore lies in  $B_2(S)$ ) of degree divisible by q.

Now suppose  $n \geq 4$  and  $S = P\Omega_{2n}^{\epsilon}(r)$ ,  $P\Omega_{2n+1}(r)$ , or  $PSp_{2n}(r)$ . Note that e = 1. We have  $GL_3(r)$  embeds naturally into  $\widetilde{H}^*$ , so that  $SL_3(r)$  embeds into  $[\widetilde{H}^*, \widetilde{H}^*]$ . An element of  $SL_3(r)$  with eigenvalues  $\{\zeta, \zeta^r, \zeta^{-r-1}\}$  and  $|\zeta| = (r^2 - 1)_2$  then yields a 2-element  $\widetilde{s} \in [\widetilde{H}^*, \widetilde{H}^*]$ , giving a character  $\chi_{\widetilde{s}} \in Irr(B_2(\widetilde{H}))$  trivial on  $\mathbf{Z}(\widetilde{H})$ . Since  $|\widetilde{s}| = (r^2 - 1)_2$  again does not divide the exponent of  $|\mathbb{T}|$ , it cannot centralize a Sylow q-subgroup. Hence, we have obtained (v), completing the proof.

2.3. Theorem C for Almost Simple Groups. Here, we finally prove Theorem C for certain almost simple groups. For S a simple group of Lie type defined in characteristic  $r_0$ , we have  $\operatorname{Aut}(S) = \widetilde{S} \rtimes D$  for an appropriate group D of graph-field automorphisms and  $\widetilde{S}$  the group of inner-diagonal automorphisms of S. (See, e.g. [20, Thms. 2.5.12 and 2.5.14].)

We start with an elementary lemma that will also be used in the proofs of Theorems A and C in the coming sections.

**Lemma 2.9.** Let  $p \neq q$  be two primes. Let G be a finite group and assume that q does not divide  $\chi(1)$  for every  $\chi \in \operatorname{Irr}(B_p(G))$ . If N is subnormal in G, then q does not divide  $\theta(1)$  for every  $\chi \in \operatorname{Irr}(B_p(N))$ .

Proof. Suppose first that N is normal in G. Let  $\theta \in \operatorname{Irr}(B_p(N))$ . Since  $B_p(N)$  is covered by  $B_p(G)$ , [45, Thm. 9.4] implies that there exists  $\chi \in \operatorname{Irr}(G|\theta)$  in the principal p-block. Therefore  $\chi$  has q'-degree and hence the same holds for  $\theta$ . The general case follows by induction on the subnormal length of N in G.

**Theorem 2.10.** Let p and q be two different primes. Let G be an almost simple group with socle S a simple group of Lie type. Suppose that G/S is a q-group and that  $B_p(G)$  contains no irreducible character of degree divisible by q. Then there is a Sylow p-subgroup  $\widetilde{P}$  of G and a Sylow q-subgroup  $\widetilde{Q}$  of G such that  $\widetilde{P}$  normalizes  $\widetilde{Q}$ .

Proof. Note that we may assume  $p \mid |S|$ , as otherwise a Sylow *p*-subgroup of *G* is trivial and the result holds. Assume that no irreducible characters in the principal *p*-block  $B_p(G)$  have degree divisible by *q*. Then every  $\chi \in \operatorname{Irr}(B_p(S))$  also satisfies  $q \nmid \chi(1)$  using Lemma 2.9. Hence by Lemmas 2.1 and 2.3, we may assume that  $S \neq {}^2F_4(2)'$  and *p* and *q* are nondefining primes for *S*, with the possibility that  $q \nmid |S|$ . Let  $r_0$  be the defining prime for *S*.

(I) First, assume that there is a Sylow *p*-subgroup P of S and Sylow *q*-subgroup Q of S such that [P, Q] = 1. (Note that this is trivially true if  $q \nmid |S|$ .) We claim that in this case, there exist Sylow *p*- and Sylow *q*- subgroups  $\widetilde{P}$  and  $\widetilde{Q}$  of G such that  $[\widetilde{P}, \widetilde{Q}] = 1$ . (Note that this follows from [30, Thm. 1.2], as by [38, Thm. 3.6],

this forces that  $p \nmid \chi(1)$  for every  $\chi \in \operatorname{Irr}(B_q(S))$  as well. Then since  $p \nmid |G/S|$ , we have further  $p \nmid \chi(1)$  for every  $\chi \in \operatorname{Irr}(B_q(G))$ . However, we aim to provide a proof independent of the main result of [30].)

We have  $S = H/\mathbb{Z}(H)$  for  $H = \mathbb{H}^F$  with  $\mathbb{H}$  a simple, simply connected algebraic group defined in characteristic  $r_0$  and  $F: \mathbb{H} \to \mathbb{H}$  a Steinberg endomorphism. Let  $r = r_0^f$  such that H is defined over  $\mathbb{F}_r$  in the case F is Frobenius, or let  $r^2 = r_0^{2n+1}$  in the case that H is a Suzuki or Ree group.

First suppose that  $q \mid |S|$ . By [38, Lem. 3.1 and Prop. 3.5], combined with [34, Lem. 2.1 and Prop. 2.2], we have  $d_p(r) = d_q(r) =: d$  (resp.  $d_p(r^2) = d_q(r^2) =: d$ ); p and q are odd, good for  $\mathbf{H}$ , larger than 3 if H is of type  ${}^{3}\mathbf{D}_{4}$ , and do not divide  $|\mathbf{Z}(\mathbf{H})^F : (\mathbf{Z}(\mathbf{H})^{\circ})^F| \cdot |\mathbf{Z}(\mathbf{H}^*)^F : (\mathbf{Z}(\mathbf{H}^*)^{\circ})^F|$ , where  $(\mathbf{H}^*, F)$  is dual to  $(\mathbf{H}, F)$ ; and there exist  $\hat{P} \in \text{Syl}_p(H)$  and  $\hat{Q} \in \text{Syl}_q(H)$  that are abelian with  $\hat{P}\hat{Q} \leq \mathbb{S}_d$  for an (F-stable) Sylow d-torus  $\mathbb{S}_d$  of  $\mathbf{H}$ .

With these conditions, we also see that G/S is induced by field automorphisms. Similarly, if  $q \nmid |S|$ , we see that the only option is that G/S is induced by field automorphisms. Say  $G = S\langle\varphi\rangle$  for some field automorphism  $\varphi$  of order a power of q. Namely, we may let  $\varphi \in \langle F_{r_0}^m \rangle$ , where the group of field automorphisms has order  $q^a \cdot m$  with  $q \nmid m$  and  $F_{r_0}$  denotes a generating field automorphism induced by a standard Frobenius.

Let  $s \in [H^*, H^*]$  be a *p*-element. As in the proofs of Lemma 2.4 and Proposition 2.8, we obtain that the corresponding semisimple character  $\chi_s$  lies in  $\operatorname{Irr}(B_p(H))$ , with its deflation lying in  $B_p(S)$ , and *s* must centralize a Sylow *q*-subgroup of  $H^*$ . Then *q* does not divide  $[H^* : \mathbf{C}_{H^*}(s)]$ . Further,  $\chi_s$  is stable under  $\varphi$ , as otherwise  $B_p(G)$  would contain some character lying above  $\chi_s$  with degree divisible by *q*.

This means that the  $H^*$ -conjugacy class of s is  $\varphi$ -stable. Since the number of  $H^*$ -conjugates of s is prime to q, we see there is even some  $H^*$ -conjugate that is invariant under  $\varphi$ . Without loss, we may assume that this is s itself. That is, we have  $s \in (\mathbf{H}^*)^{\varphi}$ , and therefore we see  $p \mid (r_0^{dm} - 1)$ . By comparing the order polynomials, we see this means  $\hat{P}$  can be chosen to be a Sylow p-subgroup of the corresponding group  $H(r_0^m)$  of the same type defined over  $\mathbb{F}_{r_0^m}$ . Then  $\hat{P}$  is centralized by  $\varphi$ . If  $q \nmid |S|$ , this proves the claim. Otherwise, by [33, Prop. 5.13], it follows that  $\mathbb{S}_d$  (and hence  $\hat{Q}$ ) is  $\varphi$ -stable. Then we can choose P and Q such that  $[P, Q\langle \varphi \rangle] = 1$ , again completing the claim.

(II) We now may assume that no such P and Q exist satisfying [P,Q] = 1. Then by Lemma 2.5 and Proposition 2.8, we still have q is odd,  $d_p(r) = d_q(r)$  (resp.  $d_p(r^2) = d_q(r^2)$ ), and there is a Sylow p-subgroup P of G (and hence of G) that normalizes an abelian Sylow q-subgroup Q of S. Further, either  $S = \text{PSL}_p(\epsilon r)$  with  $p \mid (r - \epsilon); p = 2$  and  $S = {}^2\text{G}_2(r^2); S = \text{E}_6^{\epsilon}(r), p = 2$ , and 4 and q divide  $(r + \epsilon);$ or p = 3 and  $S = {}^3\text{D}_4(r)$ . In each case, note that q > p and the only q-elements of Out(S) are field automorphisms. So, up to conjugation by inner automorphisms, we may again assume  $G \leq S\langle \varphi \rangle$  with  $\varphi$  a field automorphism of q-power order. Let  $r = r_1^{q^a}$  (resp.  $r^2 = r_1^{q^a}$  in the case of the Ree groups) for some  $a \geq 1$  and some power  $r_1$  of  $r_0$ . Note that in each case, we can again choose P to already be a Sylow p-subgroup of  $\text{PSL}_p(\epsilon r_1)$ , resp.  ${}^2\text{G}_2(r_1)$ ,  $\text{E}_6^{\epsilon}(r_1)$ ,  ${}^3\text{D}_4(r_1)$  for any such  $r_1$ . In particular, P is centralized by  $\varphi$ . Then since P normalizes Q, we see P normalizes the Sylow q-subgroup  $\widetilde{Q} = Q\langle \varphi \rangle$  of G as well.

#### 3. More on simple groups

We will also need the following results on simple groups.

**Theorem 3.1.** Let G be an almost simple group with socle S. Let p be a prime divisor of |S| and let  $q \neq p$  be another prime. Suppose that G/S is a p'-group and that  $B_p(G)$  contains no irreducible character of degree divisible by q. Then either G/S has a normal Sylow q-subgroup or  $S = \text{PSL}_n(\epsilon r)$  with Sylow p and q-subgroups of S such that [P,Q] = 1 and  $p, q \nmid (r - \epsilon)$ .

In the latter case, let  $T := \widetilde{S} \cap G$ , and otherwise let T := S. Then  $S \leq T \triangleleft G$  satisfies |T/S| is coprime to pq and G/T has a normal abelian Sylow q-subgroup.

*Proof.* First, note that if Out(S) is abelian, then G/S has a normal Sylow q-subgroup. So, we may assume that S is not an alternating, sporadic, or Tits group. The groups with exceptional Schur multipliers can further be checked in GAP [15].

Suppose for the rest of the proof that S is a simple group of Lie type defined in characteristic  $r_0$  with a nonexceptional Schur multiplier. Further, since  $\operatorname{Out}(S)$  is cyclic for the Suzuki, Ree, and triality groups, we assume that S is not one of these. As in the proof of Theorem 2.10, we may assume by Lemma 2.3 that either  $q \nmid |S|$ or that  $r_0 \notin \{p,q\}$ . We may find a quasi-simple group of Lie type H such that  $S = H/\mathbb{Z}(H)$  and  $H = \mathbb{H}^F$ , where  $\mathbb{H}$  is a simple algebraic group of simply connected type over  $\overline{\mathbb{F}}_{r_0}$  and  $F: \mathbb{H} \to \mathbb{H}$  is a Frobenius endomorphism endowing H with an  $\mathbb{F}_r$ -rational structure, where  $r = r_0^f$  for some positive integer f. Further, as there we obtain that  $d_p = d_q$  if  $q \mid |H|$  and either there exists  $P \in \operatorname{Syl}_p(S)$  and  $Q \in \operatorname{Syl}_q(S)$ such that [P,Q] = 1, or  $S = \operatorname{PSL}_p(\epsilon r)$  with  $p,q \mid (r-\epsilon)$ , or  $S = \operatorname{E}_6^\epsilon(r)$  with p = 2 and  $4, q \mid (r + \epsilon)$ . In the latter cases, recall that there is  $P \in \operatorname{Syl}_p(S)$  and  $Q \in \operatorname{Syl}_q(S)$ with P normalizing Q.

Let  $\hat{P}$  and  $\hat{Q}$  be Sylow p- and q-subgroups of H such that  $P = \hat{P}\mathbf{Z}(H)/\mathbf{Z}(H)$  and  $Q = \hat{Q}\mathbf{Z}(H)/\mathbf{Z}(H)$ . Now, recall  $\operatorname{Out}(S)$  can be realized as  $\tilde{S}/S \rtimes D$ , where  $\tilde{S}$  is the group of inner-diagonal automorphisms of S and D is an appropriate group of graph-field automorphisms. If [P,Q] = 1 and  $q \mid |H|$ , then using [38, Lem. 3.1 and Prop. 3.5], we have  $[\hat{P}, \hat{Q}] = 1$ ; p and q are odd;  $\hat{P}$  and  $\hat{Q}$  are abelian; and we have further from [34, Prop. 2.2 and Lem. 2.1], that p and q are good for  $\mathbf{H}$  and do not divide  $|\mathbf{Z}(H)|$ . The condition  $q \nmid |H|$  or the above conditions on q when  $q \mid |H|$  yield

that in either case,  $q \nmid |\tilde{S}/S|$ . In the cases when no such P and Q exist, we also see by our conditions on p and q that  $q \nmid |\tilde{S}/S|$ .

Further, in either situation, q does not divide the order of a graph automorphism in D unless possibly if  $\mathbf{H} = \mathbf{D}_4$  and q = 3, in which case  $\hat{Q} \neq 1$  is not abelian. Hence we see a Sylow q-subgroup X of  $\operatorname{Out}(S)$  is cyclic with order dividing f and can be chosen to be generated by a field automorphism. Since the graph automorphisms in D commute with the field automorphisms, we wish to show that X is also normalized by  $\tilde{S}/S$ .

Assume that **H** is not of type  $A_{n-1}$ . Then  $\tilde{S}/S$  is either cyclic of size at most 3 or is Klein four with **H** of type  $D_n$ ,  $n \ge 4$ . In the latter case, we again note that  $q \ne 3$  as above. Then in these cases, q does not divide the order of the automorphism group of  $\tilde{S}/S$ , and hence X acts trivially on  $\tilde{S}/S$ .

Finally, we consider the case that  $\mathbf{H} = A_{n-1}$ , so that  $H = \mathrm{SL}_n(\epsilon r)$  with  $n \ge 2$  and  $\epsilon \in \{\pm 1\}$ . First, assume that  $q \mid |H|$  and  $q \mid (r - \epsilon)$ . Then since Q is abelian, this forces q > n. If  $q \nmid |H|$ , we also have q > n. In particular,  $|\mathrm{Aut}(\widetilde{S}/S)| < n$  must be relatively prime to q. Then again we see X must act trivially on  $\widetilde{S}/S$ .

Hence, we may assume that  $q \mid |H|$  and  $q \nmid (r-\epsilon)$ . If n = p = 2, then again X acts trivially on  $\widetilde{S}/S$ . Then since  $d_p(r) = d_q(r)$ , we also have  $p \nmid (r-\epsilon)$ . Further, note that we are in the case [P,Q] = 1. Here let  $T := \widetilde{S} \cap G \triangleleft G$ . Then |T/S| is prime to both p and q and G/T is abelian. The statement then follows.

As a corollary to our work so far and that of [38], we obtain Brauer's Height Zero Conjecture for two primes ([38, Conj. A]) for almost simple groups. (Note that this of course would follow from [30, Thm. 1.2], but our goal is to provide an alternate proof in Theorem 4.1 below.)

**Corollary 3.2.** Let A be an almost simple group  $S \leq A \leq \operatorname{Aut}(S)$  with S a nonabelian simple group. Suppose that  $p \neq q$  are primes dividing |A| such that  $B_p(A)$ contains no character of degree divisible by q and  $B_q(A)$  contains no character of degree divisible by p. Then there are  $\widetilde{P} \in \operatorname{Syl}_p(A)$  and  $\widetilde{Q} \in \operatorname{Syl}_q(A)$  such that  $[\widetilde{P}, \widetilde{Q}] = 1$ .

Proof. Note that by Lemma 2.9,  $B_p(S)$  has no character of degree divisible by q and  $B_q(S)$  has no character of degree divisible by p. Then there is  $P \in \operatorname{Syl}_p(S)$  and  $Q \in \operatorname{Syl}_q(S)$  such that [P,Q] = 1 by [38, Thm. 5.1]. Note that we may therefore assume  $A \neq S$ . As in part (I) of the proof of Theorem 2.10, we may assume that S is a simple group of Lie type defined in characteristic  $r_0 \notin \{p,q\}$  and let  $S = H/\mathbb{Z}(H)$  as in there. We again have  $d_p(r) = d_q(r) =: d$  (resp.  $d_p(r^2) = d_q(r^2) =: d$ ) and the same constraints on p and q as there. In particular, we see that the Sylow p- and q-subgroups of A/S are induced by field automorphisms. Recall that  $\operatorname{Aut}(S) = \widetilde{S} \rtimes D$  with D abelian unless  $S = D_4(r)$ , in which case  $D \cong S_3 \times C_f$ . In each case, there is a normal subgroup  $A_{p'} \triangleleft A$  such that  $A/A_{p'}$  is a p-group and  $A_{p'}/S$  is a p'-group. By

Lemma 2.9, we have  $B_p(A_{p'})$  has no character of degree divisible by q and  $B_q(A_{p'})$  has no character of degree divisible by p, and therefore by Theorem 3.1, either  $A_{p'}/S$  has a normal Sylow q-subgroup or  $S = \text{PSL}_n(\epsilon r)$  with p and q not dividing  $r - \epsilon$ . In the latter case, we have  $T = \widetilde{S} \cap A_{p'} = \widetilde{S} \cap A$  satisfies T/S is a  $\{p, q\}'$ -group and  $A_{p'}/T$  has a normal Sylow q-subgroup. Similarly, there is a normal subgroup  $A_{q'} \triangleleft A$  such that  $A/A_{q'}$  is a q-group,  $A_{q'}/S'$  is a q'-group, and  $A_{q'}/T$  has a normal Sylow  $T = \widetilde{S} \cap A$  in the exceptional case of  $\text{PSL}_n(\epsilon r)$  from Theorem 3.1 or T = S in the other cases.

Let  $X_p$  and  $X_q$  be subgroups of A such that  $X_p/T$  and  $X_q/T$  are Sylow p- and q-subgroups of A/T with  $X_q \triangleleft A_{p'}$  and  $X_p \triangleleft A_{q'}$ . Applying Lemma 2.9 to each  $X \in \{X_p, X_q\}$ , we have  $B_p(X)$  has no character of degree divisible by q and  $B_q(X)$  has no character of degree divisible by p. Then by part (I) of the proof of Theorem 2.10 (note that the same proof works in the case  $T \neq S$  here), there is  $\tilde{Q} \in \text{Syl}_q(X_q)$  and  $\tilde{P} \in \text{Syl}_p(X_p)$  (which are therefore Sylow q- and p-subgroups of A) such that  $[\tilde{Q}, P] = 1 = [Q, \tilde{P}]$ . Since  $\tilde{Q} = Q\langle \varphi \rangle$  and  $\tilde{P} = P\langle \psi \rangle$  with  $\varphi, \psi$  (commuting) field automorphisms, we therefore have  $[\tilde{P}, \tilde{Q}] = 1$ .

**Theorem 3.3.** Let  $p \neq q$  be two primes. Let G be a quasisimple group with  $\mathbf{Z}(G) > 1$ a p-group. If q divides |G|, then there exists  $\chi \in \operatorname{Irr}(B_p(G))$  of degree a multiple of q.

Proof. Let H be the full Schur covering group of the simple group  $S := G/\mathbf{Z}(G)$ . Then G = H/Z for some  $Z \leq \mathbf{Z}(H)$  and our assumption that  $|\mathbf{Z}(G)|$  is a power of p implies that p divides  $|\mathbf{Z}(H)|$ . Note that  $\operatorname{Irr}(B_p(S)) \subseteq \operatorname{Irr}(B_p(G))$ , viewed by inflation. The cases that S is sporadic or that  $\mathbf{Z}(H)$  is an exceptional Schur multiplier can be checked in GAP [15]. By Theorem 2.2, we may assume that S is not an alternating group. Then we may assume that H is a group of Lie type and that  $\mathbf{Z}(H)$  is a nonexceptional Schur multiplier for S. Further, by Lemma 2.3, we may assume that H is defined in characteristic  $r_0 \notin \{p, q\}$ . Since we have assumed  $\mathbf{Z}(H)$  is nontrivial, H is also not a Suzuki or Ree group. Hence H is as in Proposition 2.8. Since  $|\mathbf{Z}(G)|$  is a power of p, G is not one of the exceptions listed in Proposition 2.8(v), and therefore there is  $P \in \operatorname{Syl}_p(G)$  and  $Q \in \operatorname{Syl}_q(G)$  with [P,Q] = 1. Then as in the proof of Theorem 3.1, we obtain that  $p \nmid |\mathbf{Z}(H)|$ , a contradiction.

#### 4. BRAUER'S HEIGHT ZERO CONJECTURE FOR TWO PRIMES

As we have mentioned, the so-called Brauer's height zero conjecture for two primes has been proven in [30]. The proof uses  $p^*$ -theory [59] and some specific results on simple groups. We present a proof with a more elementary reduction to (some of) the problems on simple groups that we have solved in this paper.

The following is the main theorem of [30]. Our arguments in this proof are part of the arguments that we will use in the proof of Theorem C. We refer the reader to

21

Section 9A of [27] for the elementary properties of the generalized Fitting subgroup  $\mathbf{F}^*(G)$  and the layer  $\mathbf{E}(G)$ .

**Theorem 4.1** (Liu-Wang-Willems-Zhang). Let G be a finite group and let  $p \neq q$  be two primes. If p does not divide  $\chi(1)$  for every  $\chi \in \operatorname{Irr}(B_q(G))$  and q does not divide  $\chi(1)$  for every  $\chi \in \operatorname{Irr}(B_p(G))$  then G has a nilpotent Hall  $\{p,q\}$ -subgroup.

Proof. First, we note that by Lemma 2.9 the hypotheses are inherited by normal subgroups and factor groups. Let G be a minimal counterexample. Let  $\pi = \{p, q\}$ . We may assume that  $\mathbf{O}^{\pi'}(G) = G$  and  $\mathbf{O}_{\pi'}(G) = 1$ . We claim that G has a unique minimal normal subgroup. If M and N are two different normal subgroups, then G is isomorphic to a subgroup of  $G/M \times G/N$ . By the minimality of G as a counterexample, G/M and G/N have a nilpotent Hall  $\{p, q\}$ -subgroup. By [55, Cor. 8],  $G = G/(M \cap N)$  has a nilpotent Hall  $\{p, q\}$ -subgroup. The claim follows.

Let  $N = S \times \cdots \times S$ , with S simple, be the minimal normal subgroup of G. Suppose first that S is abelian. Then we may assume that S is cyclic of order p. Suppose that  $E = \mathbf{E}(G) > 1$ , so that G has a quasisimple subnormal subgroup T. Since N is the unique minimal normal subgroup of G, this implies that  $Z = \mathbf{Z}(T) > 1$  is a p-group. Now, Theorem 3.3 yields that there exists  $\varphi \in \operatorname{Irr}(B_q(N))$  of degree divisible by p. This is a contradiction with Lemma 2.9. Therefore, E = 1 and  $\mathbf{F}^*(G) = \mathbf{O}_p(G)$ . Hence,  $\mathbf{C}_G(\mathbf{O}_p(G)) \leq \mathbf{O}_p(G)$  and G has a unique p-block by [12, Cor. V.3.11]. It follows that q does not divide the degree of any irreducible character of G and, by the Itô-Michler theorem, G has a normal abelian Sylow q-subgroup. By the uniqueness of N, this implies that q does not divide |G|, and the result is obvious.

Now, we may assume that S is simple nonabelian. Let n be the number of copies of S that appear in N. If n = 1, then G is almost simple and the result follows from Corollary 3.2. Therefore, we may assume that n > 1. Write  $N = S_1 \times \cdots \times S_n$ , with  $S_i \cong S$ . We have that G is isomorphic to a subgroup of  $\Gamma = \operatorname{Aut}(S) \wr S_n$ . Let  $B = \operatorname{Aut}(S)^n \cap G$  so that G/B is a transitive permutation group on  $\Omega = \{S_1, \ldots, S_n\}$ . Since  $\mathbf{O}^{\pi'}(G) = G$ , we may assume that p divides |G/B|.

Suppose first that q divides |S|. By [10, Lem. 3.2], there exist  $\Theta, \Delta \subseteq \Omega$  such that p divides  $|G : (G_{\Theta} \cap G_{\Delta})|$ . Since S is simple, the order of a Sylow 2-subgroup of S exceeds 2. By [3],  $|\operatorname{Irr}(B_q(S))| \geq 3$ , so we may take  $\gamma_1, \gamma_2, \gamma_3 \in \operatorname{Irr}(B_q(S))$ . If we choose  $\delta \in \operatorname{Irr}(N)$  to be an appropriate product of copies of  $\gamma_1, \gamma_2$  and  $\gamma_3$  (copies of  $\gamma_1$  in the copies of S corresponding to  $\Theta, \gamma_2$  in the copies of S corresponding to  $\Delta$  and of  $\gamma_3$  elsewhere), we have that p divides  $|G : G_{\delta}|$ . Therefore, p divides the degree of any irreducible character of G over  $\delta$ . Since some of them belongs to the principal q-block, we have a contradiction. We have thus seen that if p divides |G/B| then q does not divide |N|. Similarly, if q divides |G/B|, then p does not divide |N|. Since  $\mathbf{O}^{\pi'}(G) = G$  and  $\mathbf{O}_{\pi'}(G) = 1$  this means that one, and only one, of the primes in  $\pi$  divides |G/B||N|, and that prime divides both factors.

So suppose that p is the prime in  $\pi$  that divides |G/B| and |N|. By the inductive hypothesis, G/N and N have nilpotent Hall  $\pi$ -subgroups. By [21, Thm. D5], this implies that G satisfies  $D_{\pi}$ . Recall that this means that H has Hall  $\pi$ -subgroups, all of them are conjugate and any  $\pi$ -subgroup of G is contained in some Hall  $\pi$ -subgroup.

Let  $H_0 = P_0$  be a Sylow *p*-subgroup of *N*. Since *B* has nilpotent Hall  $\pi$ -subgroups, [58] implies that  $H_0 \leq H_1 = P_1 \times Q_1$  for some  $P_1 \in \operatorname{Syl}_p(B), Q_1 \in \operatorname{Syl}_q(B)$ . Since *G* is in  $D_{\pi}$ , there exists a Hall  $\pi$ -subgroup *H* of *G* such that *H* contains  $H_1$ . Since *q* does not divide  $|G/B|, Q_1$  is a Sylow *q*-subgroup of *G*. Thus  $H = PQ_1$  for some  $P \in \operatorname{Syl}_p(G)$ . Note that  $H_1 = H \cap B \leq H$ . By Frattini's argument,  $H = H_1 \mathbf{N}_H(Q_1)$ . Since  $\mathbf{N}_H(Q_1)$  contains  $H_1$ , it follows that  $Q_1$  is normal in *H*. Thus  $[Q_1, P] \leq Q_1$ . Furthermore,  $P_0 = H \cap N$  is normal in *H* and  $H/P_0$  is nilpotent. Therefore,  $[Q_1, P] \leq P_0$ . It follows that  $[Q_1, P] \leq Q_1 \cap P_0 = 1$ . Hence,  $H = Q_1 \times P$  is nilpotent, as wanted.

## 5. Proof of the main results

In this section, we prove the main results. We start with the following elementary group theoretical lemma.

**Lemma 5.1.** Let  $\pi$  be a set of primes and let  $p \in \pi$ . Suppose that N is a subnormal subgroup of a finite group G and G has a p-nilpotent Hall  $\pi$ -subgroup H. Then  $H \cap N$  is a p-nilpotent Hall  $\pi$ -subgroup of N.

*Proof.* First, we note that any subgroup of a *p*-nilpotent group is *p*-nilpotent so it suffices to prove that  $H \cap N$  is a Hall  $\pi$ -subgroup of N. If  $N \trianglelefteq G$  this follows from the facts that  $N \cap H$  is a  $\pi$ -group and  $|N : (N \cap H)| = |NH : H|$  is a  $\pi$ '-number and the general case follows by induction on the subnormal length.  $\Box$ 

We will use several times the following.

**Lemma 5.2.** Let p and q be two different primes. Suppose that G = AN is the semidirect product of a q-group A acting on a q'-group N. If q does not divide  $\chi(1)$  for every  $\chi \in Irr(B_p(G))$  then every character in  $Irr(B_p(N))$  is G-invariant.

Proof. By way of contradiction, suppose that there exists  $\theta \in \operatorname{Irr}(B_p(N))$  that is not G-invariant. If follows from Clifford's correspondence [25, Thm. 6.11] that q divides the degree of any character in  $\operatorname{Irr}(G|\theta)$ . Since  $B_p(G)$  covers  $B_p(N)$ , [45, Thm. 9.4] implies that there exists  $\chi \in \operatorname{Irr}(G|\theta)$  in the principal p-block of G. This contradicts the hypothesis.

The following was proved for arbitrary blocks in [47] assuming the Alperin–McKay conjecture. Here we will prove the case of principal blocks using Theorem 4.1.

**Theorem 5.3.** Suppose that a finite group A acts coprimely on G. Let p be a prime and let P be an A-invariant Sylow p-subgroup of G. If every  $\chi \in \text{Irr}(B_p(G))$  is A-invariant, then [A, P] = 1.

22

*Proof.* We argue by induction on |AG|. It suffices to see that for  $a \in A$ , P commutes with a. Thus, we may assume that  $A = \langle a \rangle$  is cyclic of q-power order for some prime q. Since AG is q-solvable, it follows from [45, Thm. 10.20] that  $Irr(B_q(AG)) = Irr(A)$  is a set of characters of p'-degree.

On the other hand, let  $\chi \in \operatorname{Irr}(B_p(AG))$ . Let  $\theta \in \operatorname{Irr}(G)$  lying under  $\chi$ . Since  $\theta$  belongs to  $B_p(G)$ , it follows that  $\theta$  is A-invariant, so  $\chi = \lambda \theta$  for some  $\lambda \in \operatorname{Irr}(AG/G)$ . In particular,  $\chi$  has q'-degree. Now, Theorem 4.1 implies that A centralizes P, as wanted.

The following is Theorem C, which we restate.

**Theorem 5.4.** Let p, q be two different primes. Let G be a finite group and assume that S is not a composition factor of G if (S, p, q) is one of the 5 exceptions listed in Theorem 2.1. Suppose that q does not divide  $\chi(1)$  for every  $\chi \in Irr(B_p(G))$ . Then Ghas a p-nilpotent Hall  $\{p, q\}$ -subgroup.

*Proof.* We argue by induction on |G|. We split the proof in a series of steps. Step 1: We may assume that  $\mathbf{O}_{p'}(G) = 1$  and  $\mathbf{O}^{q'}(G) = G$ .

Suppose that  $K = \mathbf{O}_{p'}(G) > 1$ . By the inductive hypothesis, there exists  $P \in \operatorname{Syl}_p(G)$  such that PK/K normalizes QK/K for some  $Q \in \operatorname{Syl}_q(G)$ , i.e.,

$$PK/K \leq \mathbf{N}_{G/K}(QK/K) = \mathbf{N}_G(Q)K/K.$$

Hence,  $P \leq \mathbf{N}_G(Q)K$ . Since K is a p'-group, we conclude that  $\mathbf{N}_G(Q)$  contains a Sylow p-subgroup of G, as wanted.

Now, suppose that N is a proper normal subgroup of G of q'-index. Let P be a Sylow p-subgroup of G, so that  $|PN|_{\{p,q\}} = |G|_{\{p,q\}}$ . By Lemma 2.9 and the inductive hypothesis, N has a p-nilpotent Hall  $\{p,q\}$ -subgroup. It follows from Lemma 2.1 of [30] that G has a Sylow p-subgroup that normalizes a Sylow q-subgroup of G, as desired.

Now, let  $E = \mathbf{E}(G)$  be the layer of G.

Step 2: We may assume that E > 1.

Suppose that E = 1. Then  $\mathbf{F}^*(G) = \mathbf{O}_p(G)$  and by [27, Thm. 9.8]  $\mathbf{C}_G(\mathbf{O}_p(G)) \subseteq \mathbf{O}_P(G)$ . Now, [12, Cor. V.3.11] implies that G has a unique p-block. Therefore, q does not divide  $\chi(1)$  for every  $\chi \in \operatorname{Irr}(G)$  and by the Itô–Michler theorem G has a normal abelian Sylow q-subgroup. The result follows.

Let  $Z = \mathbf{Z}(E)$ , so that Z is a p-group (by Step 1) and E/Z is a direct product of nonabelian minimal normal subgroups of G/Z of order divisible by p. Write  $E/Z = N_1/Z \times \cdots \times N_t/Z$ , where  $N_i/Z$  are the non-abelian minimal normal subgroups of G/Z. Let  $N_i/Z$  be the product of  $j_i$  copies  $S_{j_1}, \ldots, S_{j_i}$  of a non-abelian simple group  $S_i$ . Let  $C_i/Z = \mathbf{C}_{G/Z}(N_i/Z)$ , so that  $G/C_i$  is isomorphic to a subgroup of  $\operatorname{Aut}(N_i/Z) = \operatorname{Aut}(S) \wr S_{j_i}$  that contains  $M_i/C_i \cong N_i/Z$ . Step 3: We claim that  $j_i = 1$  for every  $i = 1, \ldots, t$ .

Let  $L_i/C_i = \operatorname{Aut}(S)^{j_i} \cap G/C_i$ . Suppose that q divides  $|G/L_i|$ . Note that  $\tilde{G} = G/L_i$ acts on  $N_i/Z$  as a permutation group on  $\Omega = \{S_{j_1}, \ldots, S_{j_i}\}$ . By [10, Lem. 3.2], there exist  $\Gamma, \Delta \subseteq \Omega$  such that q divides  $|\tilde{G} : (\tilde{G}_{\Gamma} \cap \tilde{G}_{\Delta})|$ . It follows from [45, Thm. 3.18] and [3] that the principal p-block of  $S_i$  has at least 3 irreducible characters (because a Sylow 2-subgroup of S has order at least 4). Let  $\gamma_1, \gamma_2, \gamma_3 \in \operatorname{Irr}(B_p(S_i))$ . Choosing  $\psi \in \operatorname{Irr}(B_p(M_i/C_i))$  as a suitable product of copies of  $\gamma_1, \gamma_2$  and  $\gamma_3$ , we can obtain that q divides  $|G : G_{\psi}|$ . It follows from Clifford's correspondence [25, Thm. 6.11] that for every  $\chi \in \operatorname{Irr}(G|\psi)$ , q divides  $\chi(1)$ . Using again [45, Thm. 9.4], we see that there exists  $\chi \in \operatorname{Irr}(G|\psi)$  in  $B_p(G)$ . This contradicts the hypothesis.

Therefore, we may assume that  $G/L_i$  is a q'-group. Since by Step 1  $G = \mathbf{O}^{q'}(G)$ , we deduce that  $G = L_i$ . This implies that  $G/C_i$  is isomorphic to a subgroup of  $\operatorname{Aut}(S)^{j_i}$ . Since  $M_i/C_i$  is a minimal normal subgroup of  $G/C_i$ , this forces  $j_i = 1$ .

In other words, we have shown that the non-abelian minimal normal subgroups of G/Z are simple groups. Write  $E/Z = X_1 \times \cdots \times X_s$  where  $X_1, \ldots, X_s$  are the non-abelian (simple) minimal normal subgroups of G/Z. Let  $C/Z = \mathbf{C}_{G/Z}(E/Z)$ , so that G/C is isomorphic to a subgroup of  $\operatorname{Aut}(X_i) \times \cdots \times \operatorname{Aut}(X_s)$  that contains  $EC/C \cong E/Z$ . Put M = EC.

Step 4: G/M contains a normal subgroup T/M such that G/T is an abelian q-group and |T/M| is prime to p and q.

Note that G/M is isomorphic to a subgroup of  $\operatorname{Out}(X_1) \times \cdots \times \operatorname{Out}(X_s)$  and by Schreier's conjecture G/M is solvable. Write  $\overline{G} = G/M$ . By [45, Thm. 10.20], we have that  $\operatorname{Irr}(\overline{G}/\mathbf{O}_{p'}(\overline{G})) = \operatorname{Irr}(B_p(\overline{G})) \subseteq \operatorname{Irr}(B_p(G))$  is a set of characters of q'-degree. By Itô's theorem [25, Thm. 12.33]  $\overline{G}/\mathbf{O}_{p'}(\overline{G})$  has a normal Sylow q-subgroup. This implies that q does not divide  $\overline{G}/\mathbf{O}_{p'}(\overline{G})$ , so by Step 1  $\overline{G} = \mathbf{O}_{p'}(\overline{G})$  is a p'-group.

Recall that G/C is isomorphic to a subgroup of  $\operatorname{Aut}(X_1) \times \cdots \times \operatorname{Aut}(X_s)$  that contains  $X_1 \times \cdots \times X_s$ . Let  $\pi_i : G/C \longrightarrow \operatorname{Aut}(X_i)$  be the projection homomorphism, so that  $\pi_i(G/C)$  is isomorphic to a factor group of G. Thus,  $\operatorname{Irr}(B_p(\pi_i(G/C))) \subseteq$  $\operatorname{Irr}(B_p(G))$  is a set of characters of q'-degree. Note also that G/C is isomorphic to a subgroup of  $\pi_1(G/C) \times \cdots \times \pi_s(G/C)$ .

On the other hand,  $\pi_i(G/C)$  is an almost simple group with socle  $X_i$ . It follows from Theorem 3.1 that  $\pi_i(G/C)$  has a normal subgroup  $T_i$  such that  $X_i \leq T_i, T_i/X_i$ is a  $\{p,q\}'$ -group and  $\pi_i(G/C)/T_i$  has a normal abelian Sylow q-subgroup. Therefore,  $\frac{\pi_1(G/C) \times \cdots \times \pi_s(G/C)}{X_1 \times \cdots \times X_s}$  is an extension of a  $\{p,q\}'$ -group by a group with a normal abelian Sylow q-subgroup. Since G/M is a subgroup of this group, we conclude that the same holds for G/M. Let  $T/M \leq G/M$  be a  $\{p,q\}'$ -group such that G/T has a normal abelian Sylow q-subgroup. By Step 1  $\mathbf{O}^{q'}(G/T) = G/T$ . We deduce that G/T is an abelian q-group, as wanted. Step 5: q does not divide |C| and C has a unique p-block.

Since E contains all the components of G, and  $E \cap C = Z$ , it follows that  $\mathbf{E}(C) = 1$ so  $\mathbf{F}^*(C) = \mathbf{F}(C) = \mathbf{O}_p(C)$ . By [27, Thm. 9.8] we conclude that  $\mathbf{C}_C(\mathbf{O}_p(C)) \subseteq \mathbf{O}_p(C)$ so C contains a unique p-block again by [12, Cor. V.3.11]. Hence q does not divide the degree of any irreducible character in C by Lemma 2.9. Now, the Itô–Michler theorem implies that C has a normal Sylow q-subgroup. By Step 1,  $\mathbf{O}_q(C) = 1$  so qdoes not divide |C|.

Step 6: Final Step.

By Step 1, Z is a p-group. Suppose first that Z > 1. Then there exists a quasisimple subnormal subgroup X of G with  $\mathbf{Z}(X)$  a nontrivial p-group. By Step 3, X is normal in G. Let  $L \leq \mathbf{Z}(X)$  be a minimal normal subgroup of G. Let  $D = \mathbf{C}_G(L)$ . We claim that G = D. By way of contradiction, suppose that D < G. By Step 1, every nontrivial factor group of G/D has order divisible by q.

Suppose first that D is cyclic. Therefore, G/D is an abelian p'-group that acts faithfully on L. By [42, Prop. 12.1] it also acts faithfully on Irr(L). Thus, there exists  $\lambda \in Irr(L)$  in a G/D-orbit of size divisible by q. It follows from Clifford's correspondence that q divides the degree of any character in  $Irr(G|\lambda)$ . But since  $\lambda$ belongs to the principal p-block of the p-group L, some of these characters belong to  $B_p(G)$ . This contradicts the hypothesis.

Hence, we may assume that L is not cyclic. It follows from [8] that  $L = C_p \times C_p$  for p = 2 or 3.

Suppose that p = 2. It follows that q = 3 and  $G/D \cong C_3$ . We reach a contradiction as in the case L cyclic.

Finally, suppose that p = 3, so that G/D is isomorphic to a subgroup of  $GL_2(3)$ . Hence q = 2. Arguing as before, we may assume that every character in Irr(L) is invariant by a Sylow 3-subgroup of G/D. It follows from [42, Thms. 9.3 and 10.4] that  $G/D \cong GL_2(3)$ . Note that  $B_3(G/D)$  has an irreducible character of degree 2. Since  $Irr(B_0(G/D)) \subseteq Irr(B_0(G))$ , this contradicts the hypothesis. This completes the proof of the claim.

We have thus seen that L is central in G. By the inductive hypothesis, G/L has a p-nilpotent Hall  $\{p,q\}$ -subgroup. Let  $P \in \operatorname{Syl}_p(G)$  and  $Q \in \operatorname{Syl}_q(G)$  be such that PQ/L is a p-nilpotent Hall subgroup of G/L (so PQ is a Hall  $\{p,q\}$ -subgroup of G). Therefore,  $QL \leq PQ$ . Since L is central in G, Q is characteristic in QL and we conclude that Q is normal in PQ, as desired.

Suppose now that Z = 1, so that  $M = C \times E$ . Therefore,  $M/E \cong C$ . Since T/Eis a q'-group by Steps 4 and 5, we deduce using the Schur-Zassenhaus theorem again that G/E is the semidirect product of R/E acting on T/E, where R/E is a Sylow q-subgroup of G/E. Therefore, G/M is the semidirect product of the q-group RM/Macting on the  $\{p,q\}'$ -group T/M. Since p does not divide |T/M|,  $Irr(B_p(T/M)) =$ Irr(T/M). Further,  $B_p(G/M) \subseteq B_p(G)$ , so q does not divide the degree of any character in  $\operatorname{Irr}(B_p(G/M))$ . By Lemma 5.2, we conclude that every character in  $\operatorname{Irr}(T/M)$  is *G*-invariant. By coprime action (see [26, Lem 3.2], for instance), this implies that the action of RM/M on T/M is trivial. In particular,  $RM \leq G$ . By Lemma 2.9, q does not divide the degree of any character in  $\operatorname{Irr}(B_p(RM))$ .

Now, note that RM/E is the semidirect product of the q-group R/E acting on the q'-group M/E. On the other hand,  $M/E \cong C$  has a unique p-block, by Step 5, so using Lemma 5.2 again, we obtain that all irreducible characters of M/E are R/E-invariant. As before, this implies that R/E acts trivially on M/E. We deduce that  $R/E \cong RM/M$  acts trivially and coprimely on both T/M and M/E. It follows from [25, Lem. 4.29] that R/E acts trivially on T/E. In particular,  $R \leq G$ . Since  $G/R \cong T/E$  is a q'-group and  $G = \mathbf{O}^{q'}(G)$ , this implies that G = R.

It follows that G/E is a q-group. By Step 4, this implies that C = 1. Hence G is isomorphic to a subgroup of  $\operatorname{Aut}(X_1) \times \cdots \times \operatorname{Aut}(X_s)$  that contains  $E = X_1 \times \cdots \times X_s$ . Therefore, if  $Q_i/X_i \in \operatorname{Syl}_q(\operatorname{Aut}(X_i)/X_i)$  for every *i*, then G is isomorphic to a subgroup of  $Q_1 \times \cdots \times Q_s$ .

Let  $\pi_i : G \longrightarrow Q_i$  be the restriction to G of the projection from  $Q_1 \times \cdots \times Q_s$  onto  $Q_i$ , so that  $G/\operatorname{Ker} \pi_i \cong \pi_i(G)$  and  $X_i \le \pi_i(G) \le Q_i$ . Notice that G is isomorphic to a subgroup of  $\Gamma = \pi_1(G) \times \cdots \times \pi_s(G)$ .

Since  $B_p(G/\operatorname{Ker} \pi_i) \subseteq B_p(G)$ , q does not divide  $\chi(1)$  for every  $\chi \in \operatorname{Irr}(B_p(\pi_i(G)))$ .

By Theorem 2.2, this implies that the socle of  $\pi_i(G)$  is not an alternating group. In the remaining cases, when the socle of  $\pi_i(G)$  is a sporadic group or a group of Lie type, Theorems 2.1 and 2.10 imply that  $\pi_i(G)$  has a *p*-nilpotent Hall  $\{p, q\}$ -subgroup. Therefore,  $\Gamma$  has a *p*-nilpotent Hall  $\{p, q\}$ -subgroup. Since  $\Gamma/E$  is a *q*-group and  $G/E \leq \Gamma/E$ , we conclude that G is subnormal in  $\Gamma$ . By Lemma 5.1, it follows that G has a *p*-nilpotent Hall  $\{p, q\}$ -subgroup.

Next, we conclude the proof of Theorem A, which we restate, with p and q interchanged. As mentioned in the Introduction, it is an easy consequence of Theorem C.

**Theorem 5.5.** Let G be a finite group, let q be a prime and let  $Q \in \text{Syl}_q(G)$ . Then  $Q \leq G$  if and only if for every prime  $p \neq q$  that divides |G| and every  $\chi \in \text{Irr}(B_p(G))$ , q does not divide  $\chi(1)$ .

Proof. Suppose first that Q is a normal Sylow q-subgroup of G. Then for every prime  $p \neq q$ ,  $Q \subseteq \mathbf{O}_{p'}(G)$  so  $\operatorname{Irr}(B_p(G)) \subseteq \operatorname{Irr}(G/Q)$  is a set of characters of q'degree. Conversely, suppose that for every  $p \neq q$ , q does not divide  $\chi(1)$  for every  $\chi \in \operatorname{Irr}(B_p(G))$ . Let  $Q \in \operatorname{Syl}_q(G)$ . Assume first that G does not have composition factors isomorphic to any of the 4 sporadic groups listed in Theorem 2.1. Then Theorem C implies that for any  $p \neq q$  a Sylow p-subgroup of G normalizes Q. Thus pdoes not divide  $|G: \mathbf{N}_G(Q)|$  for any prime  $p \neq q$  and we conclude that  $G = \mathbf{N}_G(Q)$ , as wanted. Finally, assume that G has some composition factor S isomorphic to  $J_1, J_4, M_{11}$  or  $M_{22}$ . Since the hypothesis is inherited by factor groups and by normal subgroups, it is also inherited by composition factors. Therefore, for any  $p \neq q$ , q does not divide  $\chi(1)$  for every  $\chi \in \operatorname{Irr}(B_p(S))$ . This contradicts Theorem 2.1. (For instance, if  $S = J_1$  and q = 3, we can take p = 7.)

Finally, we prove Corollary B.

**Corollary 5.6.** Let G be a finite group and let p be a prime. Then G has a normal and abelian Sylow p-subgroup if and only if p does not divide  $\chi(1)$  for every  $\chi$  that belongs to some principal block for some prime divisor of |G|.

*Proof.* This follows from Theorem A and Brauer's height zero conjecture for principal blocks [39].  $\Box$ 

We remark that the union of the irreducible characters in the principal blocks of a finite group G is usually a proper subset of Irr(G) (see [2, Thm. 3.7] and [22, Thm. 1]). It was proved in [2] however that if G is simple then  $A_{11}$  and  $A_{13}$  are the unique simple groups with this property.

## 6. Further directions

It makes sense to ask what happens if we just consider characters of height zero (as was done for instance in [50], refining the conjecture in [31]). For instance, if  $G = PSL_3(5)$  or  $S_5$ , p = 2 and q = 3, then q does not divide the degree of any irreducible character of p-height zero in  $B_p(G)$  but G does not have a p-nilpotent Hall  $\{p, q\}$ -subgroup. It may be worth remarking that there do not seem to be many counterexamples among simple groups, and  $PSL_3(5)$  is the smallest one.

We do not know any counterexamples to the following.

**Conjecture 6.1.** Let G be a finite group and let q be a prime. Then G has a normal Sylow q-subgroup if and only if for any  $p \neq q$ , q does not divide the degree of any p-height zero irreducible character in  $B_p(G)$ .

We can prove this conjecture, and the p-height zero version of Theorem C, for solvable groups.

**Proposition 6.2.** Let p and q be two different primes. Let G be a  $\{p,q\}$ -solvable group. If q does not divide the degree of any p-height zero irreducible character in  $B_p(G)$ , then a Sylow p-subgroup of G normalizes a Sylow q-subgroup of G. In particular, if G is solvable then G has a normal Sylow q-subgroup if and only if for every prime  $p \neq q$  that divides |G|, q does not divide  $\chi(1)$  for every  $\chi \in Irr(B_p(G))$  of p-height zero.

*Proof.* Since G is p-solvable,  $\operatorname{Irr}(B_p(G)) = \operatorname{Irr}(G/\mathbf{O}_{p'}(G))$ . Therefore, the hypothesis implies that q does not divide the degree of any irreducible character of p'-degree of  $G/\mathbf{O}_{p'}(G)$ . Now, the first part follows from Theorem A of [52]. For the second part we can argue as in the proof of Theorem A.

Recently, along with Malle and Rizo, we have proposed a strengthening of Brauer's height zero conjecture with Galois automorphisms [36]. We think that it should be possible to strengthen Theorem A in a similar way. Given a prime p, let  $\mathcal{J}_p$  be the subgroup of  $\operatorname{Gal}(\mathbb{Q}^{\mathrm{ab}}/\mathbb{Q})$  consisting of the automorphisms of order p that fix all p-power order roots of unity. Given a group G, let  $\operatorname{Irr}_{\mathcal{J}_p}(B_p(G))$  be the set of  $\mathcal{J}_p$ -invariant irreducible characters of the principal p-block of G.

**Conjecture 6.3.** Let G be a finite group and let q be a prime. Then G has a normal Sylow q-subgroup if and only if for any  $p \neq q$  that divides |G|, q does not divide  $\chi(1)$  for every  $\chi \in \operatorname{Irr}_{\mathcal{J}_p}(B_p(G))$ .

Again, we can prove this conjecture for solvable groups.

**Proposition 6.4.** Let G be a p-solvable group. If q does not divide the degree of any character in  $\operatorname{Irr}_{\mathcal{J}_p}(B_p(G))$  then a Sylow p-subgroup of G normalizes a Sylow qsubgroup of G. In particular, if G is solvable, then G has a normal Sylow q-subgroup if and only if for every prime  $p \neq q$  that divides |G|, q does not divide  $\chi(1)$  for every  $\chi \in \operatorname{Irr}_{\mathcal{J}_p}(B_p(G))$ .

Proof. Since G is p-solvable,  $\operatorname{Irr}(B_p(G)) = \operatorname{Irr}(G/\mathbf{O}_{p'}(G))$ . Hence, q does not divide the degree of any character in  $\operatorname{Irr}_{\mathcal{J}_p}(G/\mathbf{O}_{p'}(G))$ . By [36, Thm 3], this implies that  $G/\mathbf{O}_{p'}(G)$  has a normal Sylow q-subgroup. Thus q does not divide  $G/\mathbf{O}_{p'}(G)$ . Let P be a Sylow p-subgroup of G. Since P acts coprimely on  $\mathbf{O}_{p'}(G)$ , using [27, Thm. 3.23] we conclude that there exists a P-invariant Sylow q-subgroup of G. The first part of the result follows. For the second part we can argue as in the proof of Theorem A.

It makes sense to ask if there is a version of Brauer's height zero conjecture for an arbitrary number of primes. This has an easier solution, as a consequence of Brauer's height zero conjecture for two primes and [44].

**Theorem 6.5.** Let G be a finite group and let  $\pi = \{p_1, \ldots, p_n\}$  be a set of primes. Then G has a nilpotent Hall  $\pi$ -subgroup if and only if for every i and every  $j \neq i$ ,  $p_j$  does not divide the degree of any irreducible character in  $B_{p_i}(G)$ .

Proof. Let  $\tau = \{p_i, p_j\}$  be a subset of cardinality 2 of  $\pi$ . The "only if" part follows immediately from the fact that if G has a nilpotent Hall  $\pi$ -subgroup, then G has a nilpotent Hall  $\tau$ -subgroup and the "only if" part of Brauer's height zero conjecture for two primes [38]. Conversely, if for every i and every  $j \neq i$ ,  $p_j$  does not divide the degree of any irreducible character in  $B_{p_i}(G)$ , then Theorem 4.1 implies that G has nilpotent Hall  $\tau$ -subgroups for every  $\tau \subseteq \pi$  with  $|\tau| = 2$ . Now, we deduce that G has a nilpotent Hall  $\pi$ -subgroup using [44, Lem. 3.4] (which relies on the classification) and the comment that follows it.

28

#### References

- A. Beltrán, M. J. Felipe, G. Malle, A. Moretó, G. Navarro, L. Sanus, R. Solomon, and P.H. Tiep, Nilpotent and abelian Hall subgroups in finite groups, *Trans. Amer. Math. Soc.* 368 (2016), 2497–2513. 5, 6, 7, 13
- [2] C. Bessenrodt and J. Zhang, Character separation and principal covering, J. Algebra 327 (2011), 170–185. 27
- [3] J. Brandt, A lower bound for the number of irreducible characters in a block, J. Algebra 74 (1982), 509–515. 21, 24
- [4] M. Broué, G. Malle, and J. Michel, Generic blocks of finite reductive groups, Astérisque 212 (1993), 7–92. 6, 8
- [5] R. Burkhardt, Über die Zerlegungszahlen der Suzukigruppen Sz(q), J. Algebra 59 (1979), 421–433. 9
- [6] M. Cabanes and M. Enguehard, Representation theory of finite reductive groups. New Mathematical Monographs 1, Cambridge University Press, Cambridge, 2004. 4, 6, 7, 10, 11, 13, 15
- [7] R. W. Carter, Finite groups of Lie type. Conjugacy classes and complex characters, John Wiley and Sons, Inc., New York, 1985, 544 pp. 5, 6, 7, 11
- [8] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, and R. A. Wilson, An ATLAS of Finite Groups, Clarendon Press, Oxford, 1985. 25
- [9] D. I. Deriziotis and G. O. Michler, Character table and blocks of finite simple triality groups <sup>3</sup>D<sub>4</sub>(q), Trans. Amer. Math. Soc. **303** (1987), 39–70. 7
- [10] S. Dolfi, A. Moretó, and G. Navarro, The groups with exactly one class of size a multiple of p, J. Group Theory 12 (2009), 219–234. 21, 24
- [11] M. Enguehard, Sur les *l*-blocs unipotents des groupes rèductifs finis quand *l* est mauvais, *J. Algebra* 230 (2000), 334–377. 6
- [12] W. Feit, The representation theory of finite groups. North-Holland Mathematical Library, 25. North-Holland Publishing Co., Amsterdam-New York, 1982. 21, 23, 25
- P. Fong and B. Srinivasan, The blocks of finite general linear and unitary groups, *Invent. Math.* 69 (1982), 109–153. 10, 11, 13
- [14] P. Fong and B. Srinivasan, The blocks of finite classical groups, J. reine angew. Math. 396 (1989), 122–191. 11
- [15] The GAP group, GAP Groups, Algorithms, and Programming. Version 4.12.2, 2022, http://www.gap-system.org. 4, 18, 20
- [16] M. GECK, G. MALLE, The character theory of finite groups of Lie type: A guided tour. Cambridge University Press, Cambridge, 2020. 7, 10, 12, 14
- [17] E. Giannelli, S. Law, J. Long, and C. Vallejo, Sylow branching coefficients and a conjecture of Malle and Navarro. Bull. London Math. Soc. 54 (2022), 552-567. 2
- [18] E. Giannelli, G. Malle, and C. Vallejo, Even degree characters in principal blocks, J. Pure Appl. Algebra 223 (2019), 900–907. 4
- [19] E. Giannelli and E. Meini, Degrees and rationality of characters in the principal block of  $A_n$ , Arch. Math. 116 (2021), 161–170. 4
- [20] D. Gorenstein, R. Lyons, and R. Solomon, *The classification of the finite simple groups*, Number 3, Part I, Chapter A, Almost simple K-groups, Mathematical Surveys and Monographs, 40.3, American Mathematical Society, Providence, RI, 1998. 9, 15, 16
- [21] P. Hall, Theorems like Sylow's, Proc. London Math. Soc. 6 (1956), 286–304. 22
- [22] M. E. Harris, On the p-deficiency class of a finite group, J. Algebra 94 (1985), 411–424. 27

- [23] G. Hiss, Regular and semisimple blocks of finite reductive groups, J. London Math. Soc. (2) 41 (1990), 63–68. 6
- [24] G. Hiss, The Brauer trees of the Ree groups, Comm. Alg. 19(3) (1991), 871-888. 9
- [25] I. M. Isaacs, Character theory of finite groups. Pure and Applied Mathematics, No. 69. Academic Press [Harcourt Brace Jovanovich, Publishers], New York-London, 1976. 3, 22, 24, 26
- [26] I. M. Isaacs, Coprime group actions fixing all nonlinear irreducible characters, Canad. J. Math 41 (1989), 68–82. 26
- [27] I. M. Isaacs, *Finite group theory*. Graduate Studies in Mathematics, 92. American Mathematical Society, Providence, RI, 2008. 3, 21, 23, 25, 28
- [28] R. Kessar and G. Malle, Lusztig induction and *l*-blocks of finite reductive groups. *Pacific J. Math.* 279 (2015), 269–298. 6
- [29] R. Kessar and G. Malle, Brauer's height zero conjecture for quasi-simple groups. J. Algebra 475 (2017), 43–60. 15
- [30] Y. Liu, L. Wang, W. Willems, and J. Zhang, Brauer's height zero conjecture for two primes holds true, *Math. Ann.* 388 (2024), 1677–1690. 2, 3, 16, 17, 19, 20, 23
- [31] Y. Liu, W. Willems, H. Xiong, and J. Zhang, Trivial intersection of blocks and nilpotent subgroups, J. Algebra 559 (2020), 510–528. 27
- [32] G. Malle, Die unipotenten Charaktere von  ${}^{2}F_{4}(q^{2})$ , Comm. Algebra 18 (1990), 2361–2381. 9
- [33] G. Malle, Height 0 characters of finite groups of Lie type. *Represent. Theory* 11 (2007), 192–220.
  4, 6, 10, 11, 14, 17
- [34] G. Malle, On the inductive Alperin–McKay and Alperin weight conjecture for groups with abelian Sylow subgroups. J. Algebra 397 (2014), 190–208. 6, 7, 17, 18
- [35] G. Malle, On a minimal counterexample to Brauer's k(B)-conjecture. Israel J. Math. 228 (2018), 527–556. 5
- [36] G. Malle, A. Moretó, N. Rizo, and A. A. Schaeffer Fry, A Brauer-Galois height zero conjecture, arXiv:2402.08361. 28
- [37] G. Malle and G. Navarro, Characterizing normal Sylow p-subgroups by character degrees, J. Algebra 370 (2012), 402–406. 2
- [38] G. Malle and G. Navarro, Brauer's height zero conjecture for two primes, Math. Z. 295 (2020), 1723–1732. 2, 3, 5, 16, 17, 18, 19, 28
- [39] G. Malle and G. Navarro, Brauer's height zero conjecture for principal blocks, J. Reine Angew. Math. 778 (2021), 119–125. 1, 2, 27
- [40] G. Malle, G. Navarro, A. A. Schaeffer Fry, and P. H. Tiep, Brauer's Height Zero Conjecture, Annals of Math. (2) 200 No. 2 (to appear Sept. 2024), arXiv:2209.04736. 1, 4
- [41] G. Malle and D. Testerman, Linear algebraic groups and finite groups of Lie type. Cambridge Studies in Advanced Mathematics, 133, Cambridge University Press, Cambridge, 2011. 5, 6, 13, 14
- [42] O. Manz and T. R. Wolf, Representations of solvable groups. Cambridge University Press, Cambridge, 1993. 25
- [43] G. O. Michler, Brauer's conjectures and the classification of finite simple groups. *Representation theory*, II (Ottawa, Ont., 1984), 129–142. Lecture Notes in Math., 1178, Springer-Verlag, Berlin, 1986. 1, 4
- [44] A. Moretó, Sylow numbers and nilpotent Hall subgroups, J. Algebra 379 (2013), 80-84. 28
- [45] G. Navarro, Characters and blocks of finite groups. Cambridge University Press, Cambridge, 1998. 3, 16, 22, 23, 24
- [46] G. Navarro, Fusion in the character table, *Proc. Amer. Math. Soc.* **126** (1998), 65–66. 2
- [47] G. Navarro, Actions and characters in blocks, J. Algebra 275 (2004), 471–480. 3, 22

- [48] G. Navarro, Variations on the Itô–Michler theorem on character degrees, Rocky Mountain J. Math. 46 (2016), 1363–1376. 2
- [49] G. Navarro and N. Rizo, A Brauer-Wielandt formula (with an application to character tables), Proc. Amer. Math. Soc. 144 (2016), 4199–4204. 2
- [50] G. Navarro, N. Rizo, and A. A. Schaeffer Fry, Principal blocks for different primes, I. J. Algebra 610 (2022), 632–654. 2, 27
- [51] G. Navarro and P. H. Tiep, Characters of relative p'-degree over normal subgroups, Ann. Math. 178:3 (2013) 1135–1171. 6, 13, 14
- [52] G. Navarro and T. R. Wolf, Character degrees and local subgroups of π-separable groups, Proc. Amer. Math. Soc. 126, (1998), 2599–2605. 27
- [53] G. Navarro and T. R. Wolf, Character degrees and blocks of finite groups, J. reine angew. Math. 531, (2001), 141–146. 3
- [54] J. Olsson, Remarks on symbols, hooks and degrees of unipotent characters. J. Comb. Theory Series A 42 (1986), 223–238. 12
- [55] E.P. Vdovin and D. O. Revin, An existence criterion for Hall subgroups of finite groups, J. Group Theory 14 (2011), 93–101. 21
- [56] E.P. Vdovin and D. O. Revin, Theorems of Sylow type, Russian Math. Surveys 66:5 (2011), 829–870. 4
- [57] H. N. Ward, On Ree's series of simple groups, Trans. Amer. Math. Soc. 121 (1966), 62-89. 9
- [58] H. Wielandt, Zum Satz von Sylow, Math. Z. 60 (1954), 407–408. 22
- [59] W. Willems, p\*-theory and modular representation theory, J. Algebra 104 (1986), 135–140. 20
- [60] R. A. Wilson, The McKay conjecture is true for the sporadic simple groups, J. Algebra 207 (1998), 294–305. 4

(A. Moretó) Departament de Matemàtiques, Universitat de València, 46100 Burjassot, València, Spain

Email address: alexander.moreto@uv.es

(A. A. Schaeffer Fry) DEPT. MATHEMATICS - UNIVERSITY OF DENVER, DENVER, CO 80210, USA

Email address: mandi.schaefferfry@du.edu