A Note on Vectorial Boolean Functions as Embeddings

Augustine Musukwa¹ and Massimiliano Sala²

¹Mzuzu University, P/Bag 201, Mzuzu 2, Malawi ^{1,2}University of Trento, Via Sommarive, 14, 38123 Povo, Trento, Italy ¹augustinemusukwa@gmail.com ²maxsalacodes@gmail.com

Abstract

Let F be a vectorial Boolean function from \mathbb{F}^n to \mathbb{F}^m , where $m \ge n$. We define F as an embedding if F is injective. In this paper, we examine the component functions of F, focusing on constant and balanced components. Our findings reveal that at most $2^m - 2^{m-n}$ components of F can be balanced, and this maximum is achieved precisely when F is an embedding, with the remaining 2^{m-n} components being constants. Additionally, for quadratic embeddings, we demonstrate that there are always at least $2^n - 1$ balanced components when n is even, and $2^{m-1} + 2^{n-1} - 1$ balanced components when n is odd.

Keywords: vectorial Boolean functions; component functions; balanced function; injective function; embeddings

MSC 2010: 06E30, 94A60, 14G50

1 Introduction

Vectorial Boolean functions are extensively studied because they play a pivotal role in many fields such as cryptography and coding theory [5].

In this note, we study vectorial Boolean functions mapping from a lower-dimensional vector space to a higher-dimensional vector space. Although few papers have been published on these functions, some studies have suggested their possible application in ciphers. Recently, some researchers have considered these functions, with a focus on studying APN functions that satisfy the so-called D-property (see, for example, [1, 3]). These functions

were also partially studied in [2], where the authors examined certain cryptographic properties of Feistel networks using (n, m)-functions as S-boxes. Specifically, they provided an example of a cipher using an injective APN function from 4 bits to 5 bits as an S-box. This suggests further investigation of injective maps, that is, "embeddings." In our study of these embeddings, we are also motivated by geometric reasons. Indeed, while a linear embedding just provides a vector subspace, a non-linear embeddings will provide a sort of a nonlinear copy of a smaller space. This is especially true for low-degree embeddings, which deserve a deeper understanding.

The paper is structured as follows. Section 2 provides an overview of known results and notations essential for our study. In Section 3, we present some preliminary results, with a primary focus on the weight of first-order derivatives of any Boolean function. Section 4 presents our main results, among which is Theorem 13, where we demonstrate that any injective quadratic embedding must possess many balanced components.

2 Preliminaries and Notations

In this section, we provide definitions, known results and notations essential to our work. For further details, we recommend referring to [5, 6, 7, 8].

Let n and m be positive integers. The field of two elements, 0 and 1, is denoted by \mathbb{F} . The vector space of dimension n over \mathbb{F} is represented as \mathbb{F}^n . The size of any finite set A is denoted by |A|.

A function F from \mathbb{F}^n to \mathbb{F}^m is referred to as a vectorial Boolean function (vBf). We simply say a Boolean function when m = 1 and it is represented with lowercase letters, such as f. This paper exclusively focusses on studying vBf's from \mathbb{F}^n to \mathbb{F}^m , where $m \geq n$.

A Boolean function in algebraic normal form (ANF) is given by: $f(x_1, ..., x_n) = \sum_{I \subseteq \mathcal{P}} a_I(\prod_{i \in I} x_i)$, where $\mathcal{P} = \{1, ..., n\}$ and $a_I \in \mathbb{F}$ for all $I \subseteq P$. The algebraic degree (or simply degree) of f, denoted by deg(f), is $\max_{I \subseteq \mathcal{P}} \{|I| \mid a_I \neq 0\}$.

A Boolean function f is called *constant* if deg(f) = 0, affine if deg $(f) \le 1$, and quadratic if deg(f) = 2. We define the weight of a Boolean function f as wt $(f) = |\{x \in \mathbb{F}^n \mid f(x) = 1\}|$, and we say that f is balanced if wt $(f) = 2^{n-1}$.

A vBf F from \mathbb{F}^n to \mathbb{F}^m can be represented by $F = (f_1, ..., f_m)$, where $f_1, ..., f_m$ are Boolean functions called *coordinate functions*. The Boolean functions given by $\lambda \cdot F$, with $\lambda \in \mathbb{F}^m$ and " \cdot " denoting the dot product, are called *component functions* (or *components*) of F and they are denoted by F_{λ} . A vBf F is *balanced* if all its components are balanced. The degree of F is given by $\deg(F) = \max_{\lambda \in \mathbb{F}^m} \deg(F_{\lambda})$. Components F_{λ} with $\lambda \neq 0_m$ are called *nontrivial components*. The image of a vBf F is defined as $\text{Im}(F) = \{F(x) : x \in \mathbb{F}^n\}$. The evaluation of a Boolean function f is defined by $\text{ev}(f) = (f(x))_{x \in \mathbb{F}^n}$. We say that a vBf F is *injective* if $|\text{Im}(F)| = 2^n$. We call an injective vBf from \mathbb{F}^n into \mathbb{F}^m an *embedding*.

The Walsh transform of a Boolean function f is defined as the function $\mathcal{W}_f : \mathbb{F}^n \to \mathbb{Z}$ given by: $\mathcal{W}_f(a) = \sum_{x \in \mathbb{F}^n} (-1)^{f(x) + a \cdot x}$, for any $a \in \mathbb{F}^n$.

Remark 1. We denote the Walsh transform of f in zero by $\mathcal{F}(f)$. Observe that $\mathcal{F}(f) = \mathcal{W}_f(0_n) = \sum_{x \in \mathbb{F}^n} (-1)^{f(x)} = 2^n - 2 \operatorname{wt}(f)$.

Note that f is balanced if and only if $\mathcal{F}(f) = 0$. We mention that $\mathcal{W}_f(a)$ and $\mathcal{F}(f)$ are powerful tools in solving many problems in Boolean functions.

The nonlinearity of a Boolean function f is $\mathcal{N}(f) = 2^{n-1} - \frac{1}{2} \max_{a \in \mathbb{F}^n} |\mathcal{W}_f(a)|$. It is well-known that $\mathcal{N}(f) \leq 2^{n-1} - 2^{\frac{n}{2}-1}$. A Boolean function f is called *bent* if $\mathcal{N}(f) = 2^{n-1} - 2^{\frac{n}{2}-1}$ and this happens only when n is even. A Boolean function f is called *semi-bent* if $\mathcal{N}(f) = 2^{n-1} - 2^{\frac{n-1}{2}}$ and this happens only when n is odd.

We define the first-order derivative of f in the direction of $a \in \mathbb{F}^n$ by $D_a f(x) = f(x+a) + f(x)$. Similarly, the first-order derivative of F in the direction of $a \in \mathbb{F}^n$ is defined by $D_a F(x) = F(x+a) + F(x)$. An element $a \in \mathbb{F}^n$ is called a *linear structure* of f if $D_a f$ is constant. The set of all linear structures of f, denoted by V(f), is well-known to be a subspace of \mathbb{F}^n .

A Boolean function f is bent if and only if $D_a f$ is balanced for all nonzero $a \in \mathbb{F}^n$. It is well-known that for any quadratic Boolean function f with n odd, dim $V(f) \ge 1$, with equality occurring when f is a semi-bent. For n even, dim $V(f) \ge 0$, with equality occurring when f is bent. A Boolean function f is *partially-bent* if there exists a linear subspace E of \mathbb{F}^n such that the restriction of f to E is affine and the restriction of f to any complementary subspace E' of E (where $E \oplus E' = \mathbb{F}^n$) is bent. In [4] it is shown that E = V(f).

The quantity $\mathcal{F}(f)$ of any unbalanced quadratic Boolean function f is linked to the dimension V(f) (as mentioned in Remark 2.7 in [4]). We present this result in the following.

Remark 2. For any unbalanced quadratic Boolean function f, we have $\mathcal{F}(f) = \pm 2^{\frac{n+k}{2}}$, where $k = \dim V(f)$.

3 Preliminary results on Boolean functions

In this section, we focus on Boolean functions and analyze the sum of the weights of their first-order derivatives. We begin with the following result, which can also be found on page 1496 of [6].

Lemma 3. For any Boolean function f, we have

$$\mathcal{F}^2(f) = \sum_{a \in \mathbb{F}^n} \mathcal{F}(D_a f).$$

Proof. We have the following:

$$\mathcal{F}^{2}(f) = \left(\sum_{x \in \mathbb{F}^{n}} (-1)^{f(x)}\right)^{2} = \sum_{x,y \in \mathbb{F}^{n}} (-1)^{f(x)+f(y)}$$
$$= \sum_{x,a \in \mathbb{F}^{n}} (-1)^{f(x)+f(x+a)} = \sum_{x,a \in \mathbb{F}^{n}} (-1)^{D_{a}f(x)}$$
$$= \sum_{a \in \mathbb{F}^{n}} \sum_{x \in \mathbb{F}^{n}} (-1)^{D_{a}f(x)}$$
$$= \sum_{a \in \mathbb{F}^{n}} \mathcal{F}(D_{a}f).$$

From Lemma 3, we can easily infer the following.

Lemma 4. Let f be a Boolean function. Then we have

$$\sum_{a \in \mathbb{F}^n} \operatorname{wt}(D_a f) = 2^{2n-1} - \frac{1}{2} \mathcal{F}^2(f).$$

Proof. By Lemma 3 and Remark 1, we can obtain the sum of weight of all the first-order derivatives of f in terms of $\mathcal{F}^2(f)$ as follows:

$$\mathcal{F}^{2}(f) = \sum_{a \in \mathbb{F}^{n}} \mathcal{F}(D_{a}f) = \sum_{a \in \mathbb{F}^{n}} [2^{n} - 2\mathrm{wt}(D_{a}f)]$$
$$= 2^{2n} - 2\sum_{a \in \mathbb{F}^{n}} \mathrm{wt}(D_{a}f)$$

from which we deduce the following

$$\sum_{a\in\mathbb{F}^n} \operatorname{wt}(D_a f) = 2^{2n-1} - \frac{1}{2}\mathcal{F}^2(f).$$

We can use Lemma 4 to derive a bound on the sum of the weights of the first-order derivatives of f as follows.

Lemma 5. Let f be a Boolean function on n variables. Then

$$\sum_{a\in \mathbb{F}^n\backslash \{0_n\}} \operatorname{wt}(D_a f) = \sum_{a\in \mathbb{F}^n} \operatorname{wt}(D_a f) \leq 2^{2n-1}.$$

Furthermore, equality holds if and only if f is balanced.

Proof. Since wt $(D_{0_n}f) = 0$, $\sum_{a \in \mathbb{F}^n \setminus \{0_n\}} \operatorname{wt}(D_a f) = \sum_{a \in \mathbb{F}^n} \operatorname{wt}(D_a f)$. Recall that $\mathcal{F}(f) = 0$ if and only if f is balanced. By Lemma 4 and the fact that $\mathcal{F}^2(f) \ge 0$, the result is obtained.

We now specialize our results in Lemma 4 and Lemma 5 to quadratic Boolean functions. **Proposition 6.** Let f be any quadratic Boolean function. Then

$$\sum_{a \in \mathbb{F}^n \setminus \{0_n\}} \operatorname{wt}(D_a f) = \begin{cases} 2^{2n-1} & \text{if } f \text{ is balanced} \\ 2^{2n-1} - 2^{n+k-1} & \text{otherwise,} \end{cases}$$

where $k = \dim V(f)$.

Proof. If f is balanced, we just apply Lemma 5. Otherwise, by Lemma 4 and Remark 2, we obtain our result as follows:

$$\sum_{a \in \mathbb{F}^n} \operatorname{wt}(D_a f) = 2^{2n-1} - \frac{1}{2} \mathcal{F}^2(f) = 2^{2n-1} - \frac{1}{2} \left(2^{\frac{n+k}{2}}\right)^2 = 2^{2n-1} - 2^{n+k-1}.$$

4 Main results for vectorial Boolean functions

This section explores vBf's from \mathbb{F}^n to \mathbb{F}^m , mostly with $m \ge n$. We investigate the balancedness of their component functions, with a particular emphasis on injective functions (embeddings). Our analysis begins with the following lemma, where m and n are arbitrary.

Lemma 7. Let $F : \mathbb{F}^n \longrightarrow \mathbb{F}^m$ be any vBf. Then, for any $a, x \in \mathbb{F}^n$,

$$\sum_{\lambda \in \mathbb{F}^m} (-1)^{\lambda \cdot [F(x) + F(x+a)]} = \begin{cases} 2^m & \text{if } F(x) = F(x+a) \\ 0 & \text{otherwise.} \end{cases}$$
(1)

Proof. Observe that F(x) = F(x+a) implies that $F(x) + F(x+a) = 0_m$ and in this case we have

$$\sum_{\lambda \in \mathbb{F}^m} (-1)^{\lambda \cdot [F(x) + F(x+a)]} = \sum_{\lambda \in \mathbb{F}^m} (-1)^{\lambda \cdot 0_m} = 2^m.$$

On the other hand, $F(x) \neq F(x+a)$ implies that $F(x) + F(x+a) = \varepsilon \neq 0$ and so we have

$$\sum_{\lambda \in \mathbb{F}^m} (-1)^{\lambda \cdot [F(x) + F(x+a)]} = \sum_{\lambda \in \mathbb{F}^m} (-1)^{\lambda \cdot \varepsilon} = 0_m.$$

Our next step is to establish a relation involving the Walsh transforms of components evaluated at zero. For the remainder of this section, $m \ge n$.

Theorem 8. Let F be a vBf from \mathbb{F}^n to \mathbb{F}^m , with $m \ge n$. Then

$$\sum_{\lambda \in \mathbb{F}^m} \mathcal{F}^2(F_\lambda) \ge 2^{n+m}.$$

Moreover, equality holds if and only if F is an embedding.

Proof. Observe that F is an embedding if and only if, for any nonzero $a \in \mathbb{F}^n$, we have $F(x) \neq F(x+a)$ for all $x \in \mathbb{F}^n$. In other words, F is an embedding if and only if, for any nonzero $a \in \mathbb{F}^n$, we have $F(x) + F(x+a) \neq 0$ for all $x \in \mathbb{F}^n$.

Therefore, when F is an embedding, we can rewrite Lemma 7 more so that for any $a, x \in \mathbb{F}^n$, we have

$$\sum_{\lambda \in \mathbb{F}^m} (-1)^{\lambda \cdot [F(x) + F(x+a)]} = \begin{cases} 2^m & \text{if } a = 0\\ 0 & \text{otherwise.} \end{cases}$$
(2)

We can apply Equation (2) to deduce the following:

$$\sum_{a \in \mathbb{F}^n \setminus \{0_n\}} \sum_{x \in \mathbb{F}^n} \sum_{\lambda \in \mathbb{F}^m} (-1)^{\lambda \cdot [F(x) + F(x+a)]} \ge 0.$$
(3)

Next, we apply Equation (2) and Lemma 3 in the following:

$$\begin{split} \sum_{\lambda \in \mathbb{F}^m} \mathcal{F}^2(F_\lambda) &= \sum_{\lambda \in \mathbb{F}^m} \left(\sum_{a \in \mathbb{F}^n} \mathcal{F}(D_a F_\lambda) \right) = \sum_{\lambda \in \mathbb{F}^m} \sum_{a \in \mathbb{F}^n} \sum_{x \in \mathbb{F}^n} (-1)^{D_a F_\lambda(x)} \\ &= \sum_{a \in \mathbb{F}^n} \sum_{x \in \mathbb{F}^n} \sum_{\lambda \in \mathbb{F}^m} (-1)^{\lambda \cdot [F(x) + F(x+a)]} \\ &= \sum_{x \in \mathbb{F}^n} \sum_{\lambda \in \mathbb{F}^m} (-1)^{\lambda \cdot [F(x) + F(x+0_n)]} + \sum_{a \in \mathbb{F}^n \setminus \{0_n\}} \sum_{x \in \mathbb{F}^n} \sum_{\lambda \in \mathbb{F}^m} (-1)^{\lambda \cdot [F(x) + F(x+a)]} \\ &= 2^{n+m} + \sum_{a \in \mathbb{F}^n \setminus \{0_n\}} \sum_{x \in \mathbb{F}^n} \sum_{\lambda \in \mathbb{F}^m} (-1)^{\lambda \cdot [F(x) + F(x+a)]} \\ &\geq 2^{n+m}. \end{split}$$

It is clear that this equality holds if and only if

$$\sum_{a \in \mathbb{F}^n \setminus \{0_n\}} \sum_{x \in \mathbb{F}^n} \sum_{\lambda \in \mathbb{F}^m} (-1)^{\lambda \cdot [F(x) + F(x+a)]} = 0.$$
(4)

Thanks to Lemma 7, the sum in Equation (4) is zero if and only if all the addends

$$\sum_{\lambda \in \mathbb{F}^m} (-1)^{\lambda \cdot [F(x) + F(x+a)]}$$

are zero, which in turn is true if and only if, for any $a \neq 0$ in \mathbb{F}^n , $F(x) \neq F(x+a)$ for all $x \in \mathbb{F}^n$ (see the relation (2)). However, the latter statement is equivalent to F being an embedding.

Lemma 3 and Theorem 8 can be used to deduce the following.

Corollary 9. Let $F : \mathbb{F}^n \to \mathbb{F}^m$, with $m \ge n$, be any vBf. Then we have

$$\sum_{\lambda \in \mathbb{F}^m} \sum_{a \in \mathbb{F}^n} \mathcal{F}(D_a F_\lambda) \ge 2^{n+m}$$

Moreover, equality holds if and only if F is an embedding.

Corollary 9 carries significant implications for the weights of the first-order derivatives of components. This can be seen in the following.

Theorem 10. Let F be a vBf from \mathbb{F}^n into \mathbb{F}^m , with $m \ge n$. Then

$$\sum_{\substack{\in \mathbb{F}^m \setminus \{0_m\}}} \sum_{a \in \mathbb{F}^n} \operatorname{wt}(D_a F_{\lambda}) \le 2^{2n-1} (2^m - 2^{m-n}).$$

Moreover, equality holds if and only if F is an embedding.

Proof. We begin by first observing that

λ

$$\sum_{\lambda \in \mathbb{F}^m} \sum_{a \in \mathbb{F}^n} \mathcal{F}(D_a F_\lambda) = \sum_{a \in \mathbb{F}^n} \mathcal{F}(D_a F_{0_m}) + \sum_{\lambda \in \mathbb{F}^m \setminus \{0_m\}} \sum_{a \in \mathbb{F}^n} \mathcal{F}(D_a F_\lambda)$$
$$= 2^{2n} + \sum_{\lambda \in \mathbb{F}^m \setminus \{0_m\}} \sum_{a \in \mathbb{F}^n} \mathcal{F}(D_a F_\lambda).$$
(5)

So, by Corollary 9 and Equation (5), we obtain the following:

$$\sum_{\lambda \in \mathbb{F}^m \setminus \{0_m\}} \sum_{a \in \mathbb{F}^n} \mathcal{F}(D_a F_\lambda) \ge 2^{n+m} - 2^{2n} = 2^{2n} (2^{m-n} - 1).$$
(6)

and equality holds if and only if F is an embedding. We infer the following:

$$2^{2n}(2^{m-n}-1) \leq \sum_{\lambda \in \mathbb{F}^m \setminus \{0_m\}} \sum_{a \in \mathbb{F}^n} \mathcal{F}(D_a F_\lambda)$$
$$= \sum_{\lambda \in \mathbb{F}^m \setminus \{0_m\}} \sum_{a \in \mathbb{F}^n} [2^n - 2 \operatorname{wt}(D_a F_\lambda)]$$
$$= \sum_{\lambda \in \mathbb{F}^m \setminus \{0_m\}} \left[2^{2n} - 2 \sum_{a \in \mathbb{F}^n} \operatorname{wt}(D_a F_\lambda) \right]$$
$$= 2^{2n}(2^m - 1) - 2 \sum_{\lambda \in \mathbb{F}^m \setminus \{0_m\}} \sum_{a \in \mathbb{F}^n} \operatorname{wt}(D_a F_\lambda)$$

from which we conclude that

$$\sum_{\lambda \in \mathbb{F}^m \setminus \{0_m\}} \sum_{a \in \mathbb{F}^n} \operatorname{wt}(D_a F_\lambda) \le 2^{2n-1} (2^m - 2^{m-n})$$

and equality holds if and only if F is an embedding.

Next, we apply Theorem 10 to determine the maximum number of balanced components.

Corollary 11. Let F be a vBf from \mathbb{F}^n to \mathbb{F}^m , with $m \ge n$. Then, F can have at most $2^m - 2^{m-n}$ balanced components. Moreover, equality is achieved if and only if 2^{m-n} are constant components and F is an embedding.

Proof. By Lemma 5, a component F_{λ} is balanced if $\sum_{a \in \mathbb{F}^n} \operatorname{wt}(D_a F_{\lambda}) = 2^{2n-1}$. Thus, by Theorem 10, we conclude that F can have at most $2^m - 2^{m-n}$ balanced components. Equality is achieved if and only if F is an embedding and exactly 2^{m-n} vectors $\mu \in \mathbb{F}^m$ are such that $\sum_{a \in \mathbb{F}^n} \operatorname{wt}(D_a F_{\mu}) = 0$, which occurs only when the corresponding components F_{μ} are constants.

Remark 12. From Corollary 11, we observe that a vBf from \mathbb{F}^n to \mathbb{F}^m , where m > n, cannot have all its nontrivial components balanced.

Let B(F) represent the set of all balanced components of a vBf F. We are now ready to determine the minimum number of balanced components for any quadratic embedding. With "quadratic embedding" we mean that $\max_{\lambda \in \mathbb{F}^m} \deg(F_\lambda) \leq 2$. In the following statement, m is strictly larger than n.

Theorem 13. Let F be a quadratic embedding from \mathbb{F}^n to \mathbb{F}^m , with $m \ge n+1$. Then

- (i) $|B(F)| \ge 2^n 1$, for n even and equality holds if and only if all the other nontrivial components are bent,
- (ii) $|B(F)| \ge 2^{m-1} + 2^{n-1} 1$, for n odd and equality holds if and only if all the other nontrivial components are unbalanced semi-bent.

Proof. Let $k_{\lambda} = \dim V(F_{\lambda})$. Observe that $0 \le k_{\lambda} \le n$. By Proposition 6, we can derive the following equality:

$$\sum_{\lambda \in \mathbb{F}^m \setminus \{0_m\}} \sum_{a \in \mathbb{F}^n} \operatorname{wt}(D_a F_{\lambda}) = 2^{2n-1} |B(F)| + \sum_{\lambda \in \mathbb{F}^m \setminus (B(F) \cup \{0_m\})} \sum_{a \in \mathbb{F}^n} \operatorname{wt}(D_a F_{\lambda})$$

$$= 2^{2n-1} |B(F)| + \sum_{\lambda \in \mathbb{F}^m \setminus (B(F) \cup \{0_m\})} (2^{2n-1} - 2^{n+k_{\lambda}-1})$$

$$= 2^{2n-1} |B(F)| + 2^{2n-1} (2^m - |B(F)| - 1) - \sum_{\lambda \in \mathbb{F}^m \setminus (B(F) \cup \{0_m\})} 2^{n+k_{\lambda}-1}$$

$$= 2^{2n-1} (2^m - 1) - 2^{n-1} \sum_{\lambda \in \mathbb{F}^m \setminus (B(F) \cup \{0_m\})} 2^{k_{\lambda}}$$
(7)

Since F is an embedding, by relation (7) and Theorem 10, we have

$$2^{2n-1}(2^m-1) - 2^{n-1} \sum_{\lambda \in \mathbb{F}^m \setminus (B(F) \cup \{0_m\})} 2^{k_\lambda} = 2^{2n-1}(2^m - 2^{m-n})$$

from which we deduce the following:

$$2^{n-1} \sum_{\lambda \in \mathbb{F}^m \setminus (B(F) \cup \{0_m\})} 2^{k_\lambda} = 2^{2n-1}(2^m - 1) - 2^{2n-1}(2^m - 2^{m-n}) = 2^{2n-1}(2^{m-n} - 1)$$

$$\implies \sum_{\lambda \in \mathbb{F}^m \setminus (B(F) \cup \{0_m\})} 2^{k_\lambda} = 2^n(2^{m-n} - 1)$$

$$\implies \sum_{\lambda \in \mathbb{F}^m \setminus (B(F) \cup \{0_m\})} 2^{k_\lambda} = 2^m - 2^n.$$
(8)

Observe that

$$2^{m} - 2^{n} = \sum_{\lambda \in \mathbb{F}^{m} \setminus (B(F) \cup \{0_{m}\})} 2^{k_{\lambda}} \ge \sum_{\lambda \in \mathbb{F}^{m} \setminus (B(F) \cup \{0_{m}\})} 2^{0} = 2^{m} - |B(F)| - 1$$

$$\implies |B(F)| \ge 2^{n} - 1.$$

 $|B(F)| = 2^n - 1$ if and only if $k_{\lambda} = 0$ for all $\lambda \notin B(F) \cup \{0_m\}$ if and only if F_{λ} is bent for all $\lambda \notin B(F) \cup \{0_m\}$. So *n* must be even in this case. For *n* odd, we know $k_{\lambda} \ge 1$. So in this case, we have

$$2^{m} - 2^{n} = \sum_{\lambda \in \mathbb{F}^{m} \setminus (B(F) \cup \{0_{m}\})} 2^{k_{\lambda}} \ge \sum_{\lambda \in \mathbb{F}^{m} \setminus (B(F) \cup \{0_{m}\})} 2^{1} = 2(2^{m} - |B(F)| - 1)$$
$$\implies |B(F)| \ge 2^{m-1} + 2^{n-1} - 1.$$

 $|B(F)| = 2^{m-1} + 2^{n-1} - 1$ if and only if $k_{\lambda} = 1$ for all $\lambda \notin B(F) \cup \{0_m\}$ if and only if F_{λ} is unbalanced semi-bent for all $\lambda \notin B(F) \cup \{0_m\}$.

We now present two examples of quadratic embeddings. The first example is an embedding from \mathbb{F}^3 to \mathbb{F}^4 , and the second example is an embedding from \mathbb{F}^4 to \mathbb{F}^5 .

Example 14. The following are the coordinate functions of a quadratic embedding from \mathbb{F}^3 to \mathbb{F}^4 that has 11 balanced components and 4 unbalanced semi-bent components: $f_1 = x_1x_2 + x_1 + x_2 + x_3$, $f_2 = x_1x_3 + x_1 + x_2 + x_3$, $f_3 = x_2x_3 + x_1 + x_2$, $f_4 = x_1 + x_3$. Note that 14 components are quadratic, while only one component is linear.

Example 15. The following are the coordinate functions of a quadratic embedding from \mathbb{F}^4 to \mathbb{F}^5 that has 15 balanced components and 16 bent components: $f_1 = x_1x_2 + x_4, f_2 = x_1x_3 + x_3 + x_4, f_3 = x_1x_4 + x_3x_4 + x_2, f_4 = x_2x_3 + x_3x_4 + x_1 + x_4, f_5 = x_1x_3 + x_2x_4$. Note that all nontrivial components are quadratic.

A careful examination of Theorem 13 reveals that all arguments used can be generalized from quadratic functions to partially-bent functions, in particular:

- (i) the properties of V(f), in particular $k \ge 0$ in general and $k \ge 1$ for n odd,
- (ii) Proposition 6 holds also for partially-bent functions.

Therefore we state without proof an obvious generalization of Theorem 13, with "partiallybent embedding" we mean an embedding where all components are partially-bent.

Theorem 16. Let F be a partially-bent embedding from \mathbb{F}^n to \mathbb{F}^m , with $m \ge n+1$. Then

- (i) $|B(F)| \ge 2^n 1$, for n even and equality holds if and only if all the other nontrivial components are bent,
- (ii) $|B(F)| \ge 2^{m-1} + 2^{n-1} 1$, for n odd and equality holds if and only if all the other nontrivial components are unbalanced semi-bent.

5 Special Case for embeddings

In this section, we consider two special cases, where we see that the bound of Corollary 11 can be actually tight.

In the first case we restrict to m = n + 1. The second case considers a situation where we see a behaviour in the embedding similar to that of an affine map. We now examine the first case.

Proposition 17. Let $F : \mathbb{F}^n \to \mathbb{F}^{n+1}$ be an embedding. Then

(i) either there is one and only one constant nontrivial component and the other nontrivial components are balanced, or

(ii) there is no constant nontrivial component.

Proof. The statement is obvious when n = 1, so we can assume that $n \ge 2$. Let T be the number of constant nontrivial components of F. Let us call t the dimension of the vector subspace V_t of \mathbb{F}^{n+1} spanned by the v's such that F_v is constant. If we have at least two nonzero distinct vectors, then the dimension of V_t is at least 2, therefore if $T \ge 2$ then $t \ge 2$. Since $t \ge 2$, we can apply an affine change of output to obtain $F = (f_1, \ldots, f_{n-t+1}, \underbrace{0, \ldots, 0})$.

So in this case we have $|Im(F)| \le 2^{n-t+1} \le 2^{n-1} < 2^n$.

If t = 1, then by an affine change of output we can write $F = (f_1, \ldots, f_n, 0)$. Let ε be $\varepsilon = (\underbrace{0, \ldots, 0}_{n-\text{times}}, 1)$ in \mathbb{F}^{n+1} . Since $|\text{Im}(F)| = 2^n$, F_v is balanced for all $v \in \mathbb{F}^m \setminus \{0_n, \varepsilon\}$ and F_{ε}

is constant. Indeed, if $\overline{\lambda} \in \mathbb{F}^n$, all components of F are

$$\{(\lambda,0)\cdot(f_1,\ldots,f_n,0)\}_{\bar{\lambda}\in\mathbb{F}^n}\cup\{(\lambda,1)\cdot(f_1,\ldots,f_n,0)\}_{\bar{\lambda}\in\mathbb{F}^n},$$

the two sets are the same. So we are in case (i).

If t = 0, then we are in case (ii).

Finally, we examine a special case where the image of an embedding F from \mathbb{F}^n to \mathbb{F}^m forms a subspace of \mathbb{F}^n . We present the result in the following.

Theorem 18. Let $F : \mathbb{F}^n \longrightarrow \mathbb{F}^m$ be an embedding, with $m \ge n$, and $\operatorname{Im}(F)$ be an affine subspace of \mathbb{F}^m . Then, for all $v \in \mathbb{F}^m$, there are only two cases: either $v \cdot F$ is constant or $v \cdot F$ is balanced. Moreover, there are precisely $2^m - 2^{m-n}$ balanced components and 2^{m-n} constant components of F.

Proof. Let U = Im(F). We can assume w.l.o.g. that F(0) = 0 and so U is a vector subspace. It is clear that dim U = n. Observe that

$$\operatorname{ev}(v \cdot F) = (v \cdot F(x))_{x \in \mathbb{F}^n} = (v \cdot u)_{u \in U}.$$

We can view U as a binary linear code. Denote any parity-check matrix of U by H(U). Observe that if $v \in U^{\perp}$ then $v \cdot F$ is constant. We now suppose that $v \notin U^{\perp}$. Let $W = \{u \in U \mid v \cdot u = 0\}$ and so $W \subsetneq U$. So the parity-check matrix of W is

$$H(W) = \left(\begin{array}{c} H(U) \\ v \end{array}\right)$$

So we have dim $W = \dim U - 1 = n - 1$. That is, $|W| = 2^{n-1}$. Hence, we have

$$|\{x \in \mathbb{F} \mid v \cdot F(x) = 0\}| = |W| = 2^{n-1} = |\{x \in \mathbb{F} \mid v \cdot F(x) = 1\}| = |W| = \operatorname{wt}(F_v).$$

This implies that F_v is balanced.

Since dim $U^{\perp} = m - n$, then the number of constant components is 2^{m-n} and the balanced components are $2^m - 2^{m-n}$.

References

- Abbondati, M., Calderini, M., Villa, I.: On Dillon's property of (n, m)-functions. https://doi.org/10.48550/arXiv.2302.13922.
- [2] Aragona, R., Calderini, M., Civino, R., Sala, M., Zappatore, I.: Wave shaped round functions and primitive groups. Adv. Math. Commun. 13(1), (2019), 67-88.
- [3] Taniguchi, H.: D-property for APN functions from \mathbb{F}_2^n to \mathbb{F}_2^{n+1} . Cryptography and Communications, 15 (2023), 627–647.
- [4] Calderini, M., Sala, M and Villa, I.: A note on APN permutations in even dimension. Finite Fields Their Appl. 46, (2015) 1-6.
- [5] Carlet, C.: Boolean Functions for Cryptography and Coding Theory. Cambridge University Press, Cambridge, 2021.
- [6] Canteaut, A., Carlet, C., Charpin, P., Fontaine, C.: On cryptographic properties of the cosets of R(1, m). IEEE Transactions on Information Theory, 47(4),(2001), 1494-1513.
- [7] MacWilliams, F.-J., Sloane, N.-J.-A.: The Theory of Error-Correcting Codes. Elsevier, New York (1977).
- [8] Musukwa, A., Sala, M.: On the linear structures of balanced functions and quadratic APN functions. Cryptogr. Commun. 12, 859–880 (2020).