

# Multiple SLEs for $\kappa \in (0, 8)$ : Coulomb gas integrals and pure partition functions

Yu Feng<sup>\*1</sup>, Mingchang Liu<sup>†1,2</sup>, Eveliina Peltola<sup>‡3</sup>, and Hao Wu<sup>§1</sup>

<sup>1</sup>Tsinghua University, China

<sup>2</sup>KTH Royal Institute of Technology, Sweden

<sup>3</sup>Aalto University, Finland, and University of Bonn, Germany

## Abstract

In this article, we give an explicit relationship of SLE partition functions with Coulomb gas formalism of conformal field theory. We first construct a family of  $SLE_\kappa$  partition functions as Coulomb gas integrals and derive their various properties. In accordance with an interpretation as probabilistic correlations in loop  $O(n)$  models, they are always positive when  $\kappa \in (8/3, 8)$ , while they may have zeroes for  $\kappa \leq 8/3$ . They also admit a Fröbenius series expansion that matches with the algebraic content from CFT. Moreover, we check that at the first level of fusion, they have logarithmic asymptotic behavior when  $\kappa = 8/3$  and  $\kappa = 8$ , in accordance with logarithmic minimal models  $M(2, 1)$  and  $M(2, 3)$ , respectively.

Second, we construct  $SLE_\kappa$  pure partition functions and show that they are continuous in  $\kappa \in (0, 8)$  and they decay to zero as a polynomial of  $(8 - \kappa)$  when  $\kappa \rightarrow 8$ . We explicitly relate the Coulomb gas integrals and pure partition functions together in terms of the meander matrix. As a by-product, our results yield a construction of global non-simple multiple chordal  $SLE_\kappa$  measures ( $\kappa \in (4, 8)$ ) uniquely determined by their re-sampling property.

**Keywords:** Coulomb gas integral, partition function, Schramm-Loewner evolution (SLE)

**MSC:** 60J67, 81T40, 35C15, 33C70

## Contents

<b>1</b>	<b>Introduction</b>	<b>3</b>
1.1	Critical loop $O(n)$ models on planar domains . . . . .	3
1.2	Coulomb gas integrals . . . . .	7
1.3	Pure partition functions . . . . .	12
1.4	Relation between pure partition functions and Coulomb gas integrals . . . . .	13
1.5	Asymptotics and Fröbenius series . . . . .	15
1.6	Global multiple SLEs . . . . .	19

---

<sup>\*</sup>yufeng\_proba@163.com

<sup>†</sup>liumc\_prob@163.com

<sup>‡</sup>eveliina.peltola@hcm.uni-bonn.de

<sup>§</sup>hao.wu\_proba@gmail.com.

<b>2</b>	<b>Coulomb gas integrals</b>	<b>21</b>
2.1	General Möbius covariance . . . . .	22
2.2	Integral removals . . . . .	25
2.3	Cyclic permutation symmetry . . . . .	27
2.4	Contour deformations . . . . .	28
2.5	Asymptotics properties . . . . .	30
2.6	Coulomb gas integrals for exceptional values of $\kappa$ . . . . .	34
<b>3</b>	<b>Constructing variants of multiple <math>SLE_\kappa</math> measures for <math>\kappa \in (4, 8)</math></b>	<b>35</b>
3.1	Conformally invariant measures on curves . . . . .	35
3.2	Construction of a probability measure $P_N$ . . . . .	37
3.3	Marginal law: proof of Proposition 3.3 . . . . .	41
<b>4</b>	<b>Multiple <math>SLE_\kappa</math> and pure partition functions for <math>\kappa \in (4, 8)</math></b>	<b>47</b>
4.1	First properties of pure partition functions . . . . .	48
4.2	Cascade property . . . . .	49
4.3	General Möbius covariance . . . . .	52
4.4	Smoothness and hypoelliptic BPZ PDEs . . . . .	53
4.5	Relations of pure partition functions to Coulomb gas integrals . . . . .	56
4.6	Asymptotics properties . . . . .	58
4.7	Reversibility . . . . .	59
4.8	Global multiple $SLE_\kappa$ for $\kappa \in (4, 8)$ : proof of Theorem 1.21 . . . . .	60
<b>5</b>	<b>Tying up loose ends</b>	<b>63</b>
5.1	Pure partition functions: linear independence . . . . .	63
5.2	Pure partition functions: continuity in $\kappa$ . . . . .	63
5.3	Finishing the proof of Theorem 1.5 and Proposition 1.11 . . . . .	66
5.4	Renormalized partition functions . . . . .	67
5.5	Finishing the proof of Theorem 1.10 . . . . .	69
<b>6</b>	<b>Fröbenius series</b>	<b>70</b>
6.1	Generic case: proof of Proposition 1.16 . . . . .	70
6.2	$\kappa = 8/3$ : proof of Proposition 1.17 . . . . .	75
6.3	$\kappa = 8$ : proof of Proposition 1.18 . . . . .	77
<b>A</b>	<b>Hypoelliptic BPZ PDEs</b>	<b>78</b>
A.1	Distributions and hypoellipticity . . . . .	79
A.2	Dirichlet problem . . . . .	79
A.3	Application to SLE partition functions . . . . .	80
<b>B</b>	<b>Some integral identities</b>	<b>82</b>
<b>C</b>	<b>Additional formulas for Coulomb gas integrals</b>	<b>84</b>
<b>D</b>	<b>Examples with <math>N = 1</math> and <math>N = 2</math></b>	<b>88</b>
D.1	Asymptotics for $N = 2$ . . . . .	88
D.2	Positivity for $N = 2$ . . . . .	91
D.3	Fröbenius series for $N = 2$ . . . . .	92
<b>E</b>	<b>Fröbenius series for <math>\kappa = 4</math></b>	<b>95</b>

# 1 Introduction

In the present work, we are interested in geometric and algebraic properties of natural probability measures on conformally invariant planar curves: variants of the celebrated Schramm-Loewner evolution (SLE). They have become a success story in the description of critical planar interfaces in statistical physics [Sch00, SW01, LSW04, Smi06, SS09, CDCH<sup>+</sup>14], sharing an intimate connection to conformal field theory (CFT) [BB03, FK04, BBK05, Dub15b, Pel19], and playing an important role in various constructions in planar random geometry: properties of random fractals [LSW01a, LSW01b], constructions for models of gravity [She16, DMS21, MSW21, AHSY23, SY23] and so on.

Concretely, our goal is to highlight the explicit relationship of the SLE partition functions with the Coulomb gas formalism of CFT: certain natural SLE partition functions have Coulomb gas integral form, while others do not seem to admit very simple integral expressions and can alternatively be constructed probabilistically. We show that both approaches result in the same objects, which unifies various results presented in prior literature, e.g., [BBK05, Dub06, KL07, Law09, FK15c, KP16, PW19, BPW21, Wu20, SY23, AHSY23, Zha23]. We interpret them in terms of crossing formulas for critical loop models.

The introduction to this article presents the big picture, gathers the key results of our work, and summarizes some known results and consequences. It is organized as follows. We first discuss critical loop  $O(n)$  models and formulate a conjecture concerning general connection probabilities in these models (Conjecture 1.2). This formula is known to hold in a few cases, and from the theory of  $SLE_\kappa$  it is plausible that it should hold for all  $\kappa \in (8/3, 8)$ . The same formula should also describe hook-up probabilities for loops in a suitable conformal loop ensemble ( $CLE_\kappa$ ), in the spirit of [MW18].

After setting up the conjecture, we turn to discussing the key players in it. First of all, we define the *Coulomb gas integrals* in Section 1.2 (the denominator in Conjecture 1.2) and briefly relate our definition to CFT constructions. We also collect the main properties of these functions: they are, in fact, examples of multiple  $SLE_\kappa$  partition functions and have intricate behavior in terms of certain special values of  $\kappa$ , that correspond to certain CFT (possibly logarithmic) minimal models. They also admit a Fröbenius series expansion that matches with the algebraic content from CFT (see Theorem 1.5 and Propositions 1.7 & 1.8, as well as Proposition 1.16 in Section 1.5). We also illustrate the logarithmic behavior of the Coulomb gas integrals in the special cases of  $\kappa = 8/3$  and  $\kappa = 8$  (see Propositions 1.17 & 1.18, respectively), which correspond to the two endpoints of the interval  $(8/3, 8) \ni \kappa$  relevant to Conjecture 1.2.

Second of all, we recall the definition of multiple  $SLE_\kappa$  *pure partition functions* in Section 1.3 (the numerator in Conjecture 1.2). They have been intensively studied in the recent years, but to date have lacked a complete construction, which we finish here. Moreover, we show that the multiple  $SLE_\kappa$  pure partition functions are continuous in  $\kappa \in (0, 8)$  and that they decay to zero as a polynomial of  $(8 - \kappa)$  when  $\kappa \rightarrow 8-$ . We summarize the properties of the pure partition functions in Theorem 1.10. As a by-product, we also obtain a construction of global non-simple multiple chordal  $SLE_\kappa$  measures ( $\kappa \in (4, 8)$ ) uniquely determined by a re-sampling property. We show in Section 4.8 that our construction agrees with the recent works [AHSY23, Zha23] — see Theorem 1.21.

Third, we relate the Coulomb gas integrals and multiple  $SLE_\kappa$  pure partition functions together (in Propositions 1.11 and 1.15) in terms of the *meander matrix* (the third key player, giving combinatorial coefficients, in Conjecture 1.2). This relation extends our earlier results in [FPW24a].

## 1.1 Critical loop $O(n)$ models on planar domains

Our first main objective is to give explicit conjectures on connection probabilities for loop  $O(n)$  models at their critical regime. It is believed that the  $O(n)$  loops converge to  $SLE_\kappa$  and  $CLE_\kappa$  processes, where  $\kappa$  is related to  $n$  as in (1.1). As part of the novel content in the present work, we derive key properties of the quantities appearing in the predicted formula, phrased in terms of Coulomb gas on the one hand, and  $SLE_\kappa$  partition function theory on the other hand. Some of these properties are also needed in the




Figure 1.1: The  $O(n)$  model has an interesting phase diagram. This figure plots the critical edge-weight  $n \mapsto p_c(n) = (\sqrt{2 + \sqrt{2 - n}})^{-1}$  (in blue) and the critical edge-weight  $n \mapsto \hat{p}_c(n) = (\sqrt{2 - \sqrt{2 - n}})^{-1}$  (in orange). Although the scaling limit when  $p > p_c(n)$  is conjecturally the same, at the line  $\hat{p}_c(n)$  one should see higher order corrections to the critical behavior at the lattice level [Nie82]. From the scaling limit, or SLE perspective, one can however only observe the critical phenomena at the critical line  $p_c(n)$  (dilute phase) and above it (dense phase).

second part of our work, where we construct the pure partition functions and global multiple chordal  $SLE_\kappa$  measures with the parameter range  $\kappa \in (4, 8)$ .

### Definition of loop $O(n)$ models and boundary conditions

Denote by  $\diamond$  the hexagonal lattice. A loop is a finite subgraph of  $\diamond$  which is isomorphic to a simple cycle. A loop configuration is a subgraph of  $\diamond$  in which every vertex has even degree. For a loop configuration  $\omega$ , we denote by  $\ell(\omega)$  the number of loops in  $\omega$  and by  $o(\omega)$  the number of edges of  $\omega$ . Suppose  $\Omega$  is a non-empty simply connected induced subgraph of  $\diamond$ . Let  $n$  and  $p$  be positive real numbers. The loop  $O(n)$  measure on  $\Omega$  with edge weight  $p$  is the probability measure on loop configurations defined by

$$\mathbb{P}[\omega] \propto p^{o(\omega)} n^{\ell(\omega)}.$$

For recent progress on loop  $O(n)$  models, see the survey [PS19] and references therein.

It has been predicted by Nienhuis and Smirnov [Nie82, KN04, Smi06] that the loop  $O(n)$  model exhibits critical behavior when  $n \in [0, 2]$  with a critical edge-weight parameter  $p_c(n)$  given by

$$p_c(n) = \frac{1}{\sqrt{2 + \sqrt{2 - n}}}.$$

Furthermore, there should be two critical regimes:  $p = p_c$  and  $p > p_c$ . In both cases, the loops should scale in a suitable limit to  $SLE_\kappa$  processes with parameter  $\kappa$  satisfying

$$n = -2 \cos(4\pi/\kappa). \tag{1.1}$$

See Figure 1.1 for the phase diagram. The case  $p = p_c(n)$  is called *dilute phase* and the scaling limits of loops are simple, while the case  $p > p_c(n)$  is called *dense phase* and the scaling limits of loops are

non-simple. Using these predictions, we will give a precise conjecture concerning connection probabilities of critical loop  $O(n)$  interfaces, see Conjecture 1.2, which also has an  $\text{CLE}_\kappa$  counterpart, see Remark 1.3. We begin with the setup.

For  $N \geq 1$ , by a (*topological*) *polygon* we refer to a simply connected domain  $\Omega \subsetneq \mathbb{C}$  with  $2N$  distinct points  $x_1, \dots, x_{2N} \in \partial\Omega$  lying counterclockwise along the boundary, and we normally<sup>1</sup> assume that  $\partial\Omega$  is locally connected. Throughout, we shall denote a generic polygon as  $(\Omega; x_1, \dots, x_{2N})$ . Let  $(\Omega^\delta; x_1^\delta, \dots, x_{2N}^\delta)$  be a sequence of discrete polygons on  $\delta\mathbb{C}$  approximating a given polygon  $(\Omega; x_1, \dots, x_{2N})$  as  $\delta \rightarrow 0$ . Consider the loop  $O(n)$  model in the discrete polygon  $(\Omega^\delta; x_1^\delta, \dots, x_{2N}^\delta)$  with the following boundary conditions, illustrated in Figure 1.2(a): the  $2N$  marked boundary points  $x_1^\delta, \dots, x_{2N}^\delta$  are connected pairwise outside of  $\Omega^\delta$  according to a planar link pattern (non-crossing pair partition)

$$\begin{aligned} \beta &= \{\{a_1, b_1\}, \{a_2, b_2\}, \dots, \{a_N, b_N\}\} \in \text{LP}_N \\ &\text{with link endpoints ordered as } a_1 < a_2 < \dots < a_N \text{ and } a_r < b_r, \text{ for all } 1 \leq r \leq N, \\ &\text{and such that there are no indices } 1 \leq r, s \leq N \text{ with } a_r < a_s < b_r < b_s, \end{aligned} \quad (1.2)$$

where  $\{a_1, b_1, \dots, a_N, b_N\} = \{1, 2, \dots, 2N\}$  and for convenience, we have chosen a particular ordering of the endpoints of the links  $\{a_r, b_r\}$ . We also denote

$$\text{LP} := \bigsqcup_{N \geq 0} \text{LP}_N.$$

Let  $\omega$  be a loop  $O(n)$  configuration on  $\Omega^\delta$  with boundary condition  $\beta$ , and denote its law by

$$\mathbb{P}_\beta^\delta = \mathbb{P}_{\beta; (\Omega; x_1, \dots, x_{2N})}^{\delta; (n, p)}.$$

In such a loop  $O(n)$  configuration, there are  $N$  curves (interfaces) on  $\Omega^\delta$  connecting the marked points  $x_1^\delta, \dots, x_{2N}^\delta$  pairwise inside  $\Omega^\delta$ , as illustrated in Figure 1.2(b). The connectivity of these  $N$  interfaces induces a random link pattern  $\zeta^\delta$  in  $\text{LP}_N$ . The connection probabilities  $\mathbb{P}_\beta^\delta[\zeta^\delta = \alpha]$ , for  $\alpha \in \text{LP}_N$ , are expected to have an explicit conformally invariant formula in the scaling limit  $\delta \rightarrow 0$  expressed in terms of combinatorial numbers encoded in the meander matrix (defined below) and ratios of specific  $\text{SLE}_\kappa$  partition functions (or certain CFT correlation functions [Pel19]), which we will introduce and analyze thoroughly in this article. A similar but more restricted conjecture appeared in [FSKZ17, FPW24a].

### Conjecture for critical loop $O(n)$ models

**Definition 1.1.** A *meander* formed from two link patterns  $\alpha, \beta \in \text{LP}_N$  is the planar diagram obtained by placing  $\alpha$  and the horizontal reflection  $\beta$  on top of each other. We denote by  $\ell(\alpha, \beta)$  the number of loops in the meander formed from  $\alpha$  and  $\beta$ . We define the *meander matrix*  $\{\mathcal{M}_\nu(\alpha, \beta) : \alpha, \beta \in \text{LP}_N\}$  via


$$\mathcal{M}_\nu(\alpha, \beta) := \nu^{\ell(\alpha, \beta)}, \quad \nu \in \mathbb{C}. \quad (1.3)$$

**Conjecture 1.2.** Consider the loop  $O(n)$  model with  $n \in (0, 2]$  and with boundary condition  $\beta \in \text{LP}_N$ . Denote by  $\zeta^\delta$  the random connectivity in  $\text{LP}_N$  induced by the  $N$  discrete interfaces in  $\Omega^\delta$ .


1. For the critical dilute phase with  $p = p_c$ , let  $\kappa$  be the solution to (1.1) with  $\kappa \in (8/3, 4]$ .
2. For the critical dense phase with  $p > p_c$ , let  $\kappa$  be the solution to (1.1) with  $\kappa \in [4, 8)$ .

---

<sup>1</sup>At times, we might require more regularity, while on the other hand, one could also consider the boundary points in terms of prime ends, which however would not bring anything more to the present discussion.



(a) Boundary condition  $\beta = \{\{1, 2\}, \{3, 4\}, \{5, 6\}\}$ .



(b)

(c)

(d)

(e)

(f)

Figure 1.2: In (a), we illustrate a boundary condition  $\beta = \{\{1, 2\}, \{3, 4\}, \{5, 6\}\}$  for the  $O(n)$  loop model. In (b)-(f), we have five possible internal link patterns of the interfaces (in red):  $\{\{1, 2\}, \{3, 4\}, \{5, 6\}\}$ ,  $\{\{1, 6\}, \{2, 3\}, \{4, 5\}\}$ ,  $\{\{1, 2\}, \{3, 6\}, \{4, 5\}\}$ ,  $\{\{1, 6\}, \{2, 5\}, \{3, 4\}\}$ , and  $\{\{1, 4\}, \{2, 3\}, \{5, 6\}\}$ , respectively. The numbers of loops in the meander formed from the boundary condition and the internal link patterns are 3,1,2,2, and 2, respectively.

Then, the following conformally invariant connection probability formula holds:

$$\lim_{\delta \rightarrow 0} \mathbb{P}_\beta^\delta[\zeta^\delta = \alpha] = \mathcal{M}_n(\alpha, \beta) \frac{\mathcal{Z}_\alpha^{(\kappa)}(\hat{x}_1, \dots, \hat{x}_{2N})}{\mathcal{F}_\beta^{(\kappa)}(\hat{x}_1, \dots, \hat{x}_{2N})}, \quad \text{for any } \alpha \in \text{LP}_N, \quad (1.4)$$

where  $\mathcal{M}_n(\alpha, \beta)$  is the meander matrix in Definition 1.1 with  $\nu = n$ , and  $\mathcal{Z}_\alpha^{(\kappa)}(\hat{x}_1, \dots, \hat{x}_{2N})$  is the pure partition function in Definition 1.9, and  $\mathcal{F}_\beta^{(\kappa)}(\hat{x}_1, \dots, \hat{x}_{2N})$  is the Coulomb gas integral in Definition 1.4, and  $\hat{x}_j = \varphi(x_j)$  for  $1 \leq j \leq 2N$  are images of the marked points under any conformal map  $\varphi: \Omega \rightarrow \mathbb{H} = \{z \in \mathbb{C} : \text{Im}(z) > 0\}$ .

The connection probabilities in Conjecture 1.2 are also predicted in [FSKZ17]. The aim of this article is to present them in a framework that is directly amenable to investigations using SLE $_\kappa$  and CLE $_\kappa$  processes. Some cases of Conjecture 1.2 have been already proved rigorously in the literature:

- Critical loop  $O(n)$  model with  $n = 1$  and  $p = 1$  corresponds to critical *Bernoulli percolation*, for which the conformal invariance is known [Smi01, CN06, CN07]. In this case (1.4) reads

$$\lim_{\delta \rightarrow 0} \mathbb{P}_\beta^\delta[\zeta^\delta = \alpha] = \mathcal{Z}_\alpha^{(6)}(\hat{x}_1, \dots, \hat{x}_{2N}), \quad (1.5)$$

because  $\mathcal{M}_n(\alpha, \beta) \equiv 1$  and  $\mathcal{F}_\beta^{(6)} \equiv 1$  as in (1.26). The case with  $N = 2$  is known as Cardy-Smirnov formula [Car92, Smi01]. The convergence in (1.5) with  $N \geq 3$  was proven in [FPW24b].

- Critical loop  $O(n)$  model with  $n = \sqrt{2}$  and  $p = 1$  corresponds to the critical *random-cluster model* with cluster-weight 2, for which the conformal invariance is known on  $\mathbb{Z}^2$  [CS12, CDCH<sup>+</sup>14]. Chelkak & Smirnov

proved (1.4) in the case  $N = 2$  [CS12], and the case  $N \geq 3$  was proven in [FPW24a, Theorem 1.8]. In this case, the denominator  $\mathcal{F}_\beta^{(16/3)}$  has an explicit expression (1.28).

- Critical loop  $O(n)$  model with  $n \rightarrow 0$  and  $p = 1$  can be understood as a *uniform spanning tree* (UST), for which the conformal invariance is also known on  $\mathbb{Z}^2$  [LSW04]. An analogue of the convergence in (1.4) is proved in [LPW24, Theorem 1.3], where one replaces  $\mathcal{M}_n(\alpha, \beta)$  by the renormalized meander matrix  $\widehat{\mathcal{M}}(\alpha, \beta)$  in (1.29), and one replaces  $\mathcal{Z}_\alpha^{(\kappa)}$  and  $\mathcal{F}_\beta^{(\kappa)}$  by renormalized partition functions  $\widehat{\mathcal{Z}}_\alpha^{(8)}$  and  $\widehat{\mathcal{F}}_\beta^{(8)}$  in (1.23) and (1.31), respectively. Related probabilities for UST models were also considered by Dubédat [Dub06] and Kenyon & Wilson [KW11a].
- Critical loop  $O(n)$  model with  $n = 1$  and  $p = p_c(1) = 1/\sqrt{3}$  corresponds to the critical (spin) *Ising model*, for which the conformal invariance on  $\mathbb{Z}^2$  is known as well [CS12, CDCH<sup>+</sup>14]. The convergence in (1.4) with  $N = 2$  was proven by Chelkak & Smirnov in [CS12], and the case  $N \geq 3$  was treated in [PW23]. In this case, we have  $\mathcal{M}_n(\alpha, \beta) = 1$  and  $\mathcal{F}_\beta^{(3)}$  has an explicit expression (1.27).

**Remark 1.3.** Conformal loop ensemble,  $\text{CLE}_\kappa$ , is argued [Nie82, KN04, SW12] to be the only possible conformally invariant scaling limit for the joint law of all of the macroscopic loops for critical loop  $O(n)$  model with “uniform” free boundary condition, where  $\kappa \in (8/3, 8)$  and  $n \in (0, 2]$  are related through (1.1). In [MW18], Miller & Werner introduced  $\text{CLE}_\kappa$  in a rectangle with alternating boundary conditions, and derived connection probabilities for  $\text{CLE}_\kappa$  loops which coincide with the right-hand side of (1.4) with  $N = 2$ . In general, we can introduce the  $\text{CLE}_\kappa$  in arbitrary polygons with alternating boundary conditions, by utilizing a similar exploration process as described in [MW18]. We can then ask what the general connection probabilities are. In light of the conjecture that critical  $O(n)$  loops should converge to  $\text{CLE}_\kappa$  loops, it is natural to expect that the corresponding probabilities are given by the right-hand side of (1.4) for all  $N \geq 2$ .

Finally, let us discuss the loop  $O(n)$  model as  $n \rightarrow 0$  and  $p = p_c(0) = (\sqrt{2 + \sqrt{2}})^{-1}$ , which is closely related to self-avoiding walk. In the discrete polygon  $(\Omega^\delta; x_1^\delta, \dots, x_{2N}^\delta)$ , we consider configurations  $\omega$  in  $\Omega^\delta$  such that there are  $N$  disjoint self-avoiding paths connecting pairwise among the marked points  $(x_1^\delta, \dots, x_{2N}^\delta)$  and there are no loops. Denote by  $\zeta^\delta(\omega)$  the connectivity in  $\text{LP}_N$  induced by these  $N$  discrete paths. Denote by  $\mathcal{W}_\alpha(\Omega^\delta; x_1^\delta, \dots, x_{2N}^\delta)$  the set of all configurations  $\omega$  such that  $\zeta^\delta(\omega) = \alpha$ . As a generalization of [DCS12] for  $N = 1$ , the partition function of these walks (the number of configurations realizing the event  $\{\zeta^\delta = \alpha\}$ ) should behave in the following manner:

$$\delta^{-\frac{5}{4}N} \sum_{\omega \in \mathcal{W}_\alpha(\Omega^\delta; x_1^\delta, \dots, x_{2N}^\delta)} \left( \frac{1}{\sqrt{2 + \sqrt{2}}} \right)^{o(\omega)} \stackrel{\delta \rightarrow 0}{\approx} \mathcal{Z}_\alpha^{(8/3)}(\hat{x}_1, \dots, \hat{x}_{2N}), \quad \alpha \in \text{LP}_N, \quad (1.6)$$

whose right-hand side is conformally covariant by Theorem 1.10. One possible way to understand the notation “ $\approx$ ” is as

$$\lim_{\delta \rightarrow 0} \frac{1}{\log \delta} \left( \log(\text{LHS of (1.6)}) - \log \mathcal{Z}_\alpha^{(8/3)}(\hat{x}_1, \dots, \hat{x}_{2N}) \right) = \frac{5}{4}N.$$

## 1.2 Coulomb gas integrals

Representation theory of the Virasoro algebra provides a way to find solutions to PDEs appearing in conformal field theory and its applications to random geometry. In CFT, this approach is commonly known as “Coulomb gas formalism” [DFMS97], and it formally builds upon the free boson (Gaussian free field, GFF). While we will not use the GFF in the present work, we use the Coulomb gas formalism as a motivating framework to write down explicit formulas for  $\text{SLE}_\kappa$  partition functions, in the spirit of [Dub06, KP16, FK15c]. In this context, it is customary<sup>2</sup> to introduce parameters depending on  $\kappa > 0$ :

<sup>2</sup>Here, we opt for the notation that we find most commonly in CFT literature. Note that the charges  $\alpha_0$  and  $\alpha_i$  have nothing to do with link patterns, and we trust that the notation is clear from context.




Figure 1.3: The left panel depicts a plot of the function  $\kappa \mapsto h(\kappa)$ . Note that  $h(6) = 0$ . The right panel depicts a plot of the central charge  $\kappa \mapsto c(\kappa)$ . Note that  $c(8/3) = c(6) = 0$ .

- *central charge*  $c = c(\kappa)$  and *background charge*  $\alpha_0 = \alpha_0(\kappa)$ :

$$c = \frac{(3\kappa - 8)(6 - \kappa)}{2\kappa} = 1 - 24\alpha_0^2 \quad \text{and} \quad \alpha_0 = \frac{\kappa - 4}{4\sqrt{\kappa}}; \quad (1.7)$$

- *Coulomb charges*  $\alpha_j = \alpha_j(\kappa) \in \mathbb{R}$  at the marked points  $x_j \in \{x_1, x_2, \dots, x_{2N}\}$ ;
- *screening charges*  $\alpha_- = \alpha_-(\kappa)$  and  $\alpha_+ = \alpha_+(\kappa)$ :

$$\alpha_- = -\frac{2}{\sqrt{\kappa}} \quad \text{and} \quad \alpha_+ = \frac{\sqrt{\kappa}}{2} \quad \text{such that} \quad \alpha_- \alpha_+ = -1;$$

- *conformal weights*  $h_j = h_j(\kappa)$  at the marked points  $x_j \in \{x_1, x_2, \dots, x_{2N}\}$  are given by  $\alpha_j(\kappa)$  as

$$h_j = \alpha_j^2 - 2\alpha_j\alpha_0.$$

Choosing  $\alpha_j = 1/\sqrt{\kappa}$  for all  $j \in \{1, 2, \dots, 2N\}$  (this is the charge associated to a degenerate CFT field at level two, known to be closely related to  $\text{SLE}_\kappa$  curves), the Coulomb gas integral

$$\int \cdots \int \prod_{1 \leq i < j \leq 2N} (x_j - x_i)^{2\alpha_i \alpha_j} \prod_{1 \leq r < s \leq N} (u_s - u_r)^{2\alpha^2} \prod_{\substack{1 \leq i \leq 2N \\ 1 \leq r \leq N}} (u_r - x_i)^{2\alpha - \alpha_i} du_1 \cdots du_N,$$

plays an important role in SLE theory. The conformal weight for a degenerate field at level two is

$$h_j = h(\kappa) := h_{1,2}(\kappa) := \frac{6 - \kappa}{2\kappa}. \quad (1.8)$$

**Definition 1.4.** For each  $\beta \in \text{LP}_N$  ordered as in (1.2), we define the *generic Coulomb gas integral* functions on the configuration space

$$\mathfrak{X}_{2N} := \{\mathbf{x} := (x_1, \dots, x_{2N}) \in \mathbb{R}^{2N} : x_1 < \cdots < x_{2N}\} \quad (1.9)$$

as  $\mathcal{F}_\beta^{(\kappa)} : \mathfrak{X}_{2N} \rightarrow \mathbb{C}$ ,

$$\mathcal{F}_\beta^{(\kappa)}(\mathbf{x}) := C(\kappa)^N \oint_{\vartheta_1^\beta} du_1 \oint_{\vartheta_2^\beta} du_2 \cdots \oint_{\vartheta_N^\beta} du_N f_\beta^{(\kappa)}(\mathbf{x}; \mathbf{u}), \quad \kappa \in (0, \infty) \setminus \left\{ \frac{8}{m} : m \in \mathbb{Z}_{>0} \right\}, \quad (1.10)$$






Figure 1.4: The two pairs of Pochhammer integration contours  $\{\vartheta_1^{\overleftarrow{\curvearrowright}}, \vartheta_2^{\overleftarrow{\curvearrowright}}\} = \{\vartheta(x_1, x_2), \vartheta(x_3, x_4)\}$  and  $\{\vartheta_1^{\overrightarrow{\curvearrowright}}, \vartheta_2^{\overrightarrow{\curvearrowright}}\} = \{\vartheta(x_1, x_4), \vartheta(x_2, x_3)\}$  used to define the functions  $\mathcal{F}_\beta^{(\kappa)}$  and  $\mathcal{F}_\beta^{(\kappa)}$  associated to the link patterns  $\{\{1, 2\}, \{3, 4\}\}$  and  $\{\{1, 4\}, \{2, 3\}\}$  in Definition 1.4. The branch choice (1.13) is illustrated by a red cross.

where the normalization factor is given by (see also Figure 1.5)

$$C(\kappa) := \frac{\nu(\kappa)}{4 \sin^2(4\pi/\kappa)} \frac{\Gamma(2 - 8/\kappa)}{\Gamma(1 - 4/\kappa)^2}, \quad \text{with fugacity} \quad \nu(\kappa) := -2 \cos(4\pi/\kappa), \quad (1.11)$$

the (multivalued) integrand is

$$f_\beta^{(\kappa)}(\mathbf{x}; \mathbf{u}) := \prod_{1 \leq i < j \leq 2N} (x_j - x_i)^{2/\kappa} \prod_{1 \leq r < s \leq N} (u_s - u_r)^{8/\kappa} \prod_{\substack{1 \leq i \leq 2N \\ 1 \leq r \leq N}} (u_r - x_i)^{-4/\kappa}, \quad (1.12)$$

with  $\mathbf{u} = (u_1, \dots, u_N)$  disjoint and with branch chosen to be real and positive when

$$x_{a_r} < \text{Re}(u_r) < x_{a_r+1}, \quad \text{for all } 1 \leq r \leq N, \quad (1.13)$$

and where each  $\vartheta_r^\beta$  in (1.10) is a Pochhammer contour which surrounds each of the points  $x_{a_r}, x_{b_r}$  once in the positive direction and once in the negative direction (using the notation (1.2)):

$$\vartheta_r^\beta = \vartheta(x_{a_r}, x_{b_r}) = \text{Diagram} \quad (1.14)$$

and which does not surround any other marked point among  $\{x_1, \dots, x_{2N}\}$ , and lastly, all of the contours  $\vartheta_1^\beta, \vartheta_2^\beta, \dots, \vartheta_N^\beta$  are assumed to be disjoint — see Figure 1.4.

In other words, the integration contours for  $\mathcal{F}_\beta^{(\kappa)}$  in (1.10) are determined by pairwise non-intersecting Pochhammer contours surrounding the marked points pairwise according to the connectivity  $\beta$ . Note that since the integration contours  $\vartheta_r^\beta$  avoid the marked points  $x_1, \dots, x_{2N}$ , the integral in (1.10) is convergent for all  $\kappa > 0$ . We can also extend  $\mathcal{F}_\beta^{(\kappa)}$  to a multivalued function on the larger set

$$\mathfrak{V}_{2N} := \{\mathbf{x} = (x_1, \dots, x_{2N}) \in \mathbb{C}^{2N} : x_i \neq x_j \text{ for all } i \neq j\}. \quad (1.15)$$

In a few special cases, the Coulomb gas integrals  $\mathcal{F}_\beta^{(\kappa)}$  reduce to algebraic expressions — see Remark 1.13.

We have excluded particular values of  $\kappa$  in (1.10) because the constant  $C(\kappa)$  is not well-defined at those exceptional points. Nevertheless, it turns out that the function  $\mathcal{F}_\beta^{(\kappa)}$  is well-defined for all  $\kappa \in (0, 8)$ . It follows from [FPW24a, Propositions 2.2 & 2.3] that each  $\mathcal{F}_\beta^{(\kappa)}$  satisfies the following properties.

(PDE) **BPZ equations**, important in both CFT and SLE theory:

$$\left[ \frac{\kappa}{2} \frac{\partial^2}{\partial x_j^2} + \sum_{i \neq j} \left( \frac{2}{x_i - x_j} \frac{\partial}{\partial x_i} - \frac{2h(\kappa)}{(x_i - x_j)^2} \right) \right] F(x_1, \dots, x_{2N}) = 0, \quad \text{for all } j \in \{1, \dots, 2N\}. \quad (\text{PDE})$$

These PDEs appear in SLE theory as a manifestation of martingales yielding SLE variants obtained from the chordal  $\text{SLE}_\kappa$  (as in Section 1.6) [BBK05, Dub06, Dub07a, Gra07, KL07, Law09], as well as in CFT literature [BPZ84a, BPZ84b] emerging from the algebraic structure resulting from the infinitesimal conformal symmetry of planar critical models. Notably, these PDEs are hypoelliptic (see Appendix A) in terms of the Hörmander bracket condition. We include in Appendix A a proof of the fact that typical local martingales for the  $\text{SLE}_\kappa$  indeed give rise to (smooth) solutions to (PDE).

(COV) **Möbius covariance:**

$$F(x_1, \dots, x_{2N}) = \prod_{i=1}^{2N} \varphi'(x_i)^{h(\kappa)} \times F(\varphi(x_1), \dots, \varphi(x_{2N})), \quad (\text{COV})$$

for all Möbius maps  $\varphi$  of the upper half-plane  $\mathbb{H}$  such that  $\varphi(x_1) < \dots < \varphi(x_{2N})$ .

Such a covariance property gives a version of global conformal symmetry, manifest in the conformal invariance of the  $\text{SLE}_\kappa$  process on the one hand, and for CFT primary fields on the other hand.

(PLB) **Power-law bound:** There exist constants  $C > 0$  and  $p > 0$  such that for all  $N \geq 1$  and  $\mathbf{x} \in \mathfrak{X}_{2N}$ ,

$$|F(\mathbf{x})| \leq C \prod_{1 \leq i < j \leq 2N} (x_j - x_i)^{\mu_{ij}(p)}, \quad \text{where} \quad \mu_{ij}(p) := \begin{cases} p, & \text{if } |x_j - x_i| > 1, \\ -p, & \text{if } |x_j - x_i| < 1. \end{cases} \quad (\text{PLB})$$

The bound (PLB) appears in the works of Flores & Kleban [FK15b, FK15c], where uniqueness results for solutions to the above PDEs were established using elliptic PDE techniques (cf. Lemma 2.11).

In particular, in order to argue that each  $\mathcal{F}_\beta^{(\kappa)}$  is a multiple  $\text{SLE}_\kappa$  partition function, it only remains to check whether  $\mathcal{F}_\beta^{(\kappa)}(\mathbf{x}) > 0$  for all  $\mathbf{x} \in \mathfrak{X}_{2N}$ . The positivity of the partition function is necessary for its usage in Loewner theory (indeed, it should arise from an exponential martingale in a Girsanov transform). This positivity property has been, however, a substantial difficulty in the literature thus far. Indeed, from the iterated integral formulas (1.10) of the complex, multivalued integrand (1.12) it is very hard to see in general nothing more than that the functions  $\mathcal{F}_\beta^{(\kappa)}$  are real-valued (by a simple reflection argument). We settle this problem in Theorem 1.5. Interestingly, the Coulomb gas integral functions are only positive when  $\kappa$  avoids certain special intervals, and we also analyze the behavior of the zeros in a special case. The positivity does however hold in the range  $\kappa \in (8/3, 8)$  relevant to critical loop  $O(n)$  models.

**Theorem 1.5.** *The Coulomb gas integral functions  $\{\mathcal{F}_\beta^{(\kappa)} : \beta \in \text{LP}\}$  satisfy the following properties.*

1. *They can be continuously extended to  $(0, 8) \ni \kappa$ : for each  $\kappa \in (0, 8)$ , we have*

$$\lim_{\kappa' \rightarrow \kappa} \mathcal{F}_\beta^{(\kappa')} = \mathcal{F}_\beta^{(\kappa)}, \quad \beta \in \text{LP}_N. \quad (1.16)$$

2. *They satisfy (PDE), (COV), and (PLB). Moreover,*

(POS) *when  $\kappa \in \bigcup_{m \in \mathbb{Z}_{>0}} \left(\frac{8}{4m-1}, \frac{8}{4m-3}\right)$ , we have  $\mathcal{F}_\beta^{(\kappa)}(\mathbf{x}) > 0$  for all  $\mathbf{x} \in \mathfrak{X}_{2N}$  and  $\beta \in \text{LP}_N$ ;*

(NUL) *when  $\kappa \in \left\{\frac{8}{2m+1} : m \in \mathbb{Z}_{>0}\right\}$ , we have  $\mathcal{F}_\beta^{(\kappa)}(\mathbf{x}) = 0$  for all  $\mathbf{x} \in \mathfrak{X}_{2N}$  and  $\beta \in \text{LP}_N$ ;*

(SGN) *when  $N \geq 2$  and  $\kappa \in \bigcup_{m \in \mathbb{Z}_{>0}} \left(\frac{8}{4m+1}, \frac{8}{4m-1}\right)$ , there exist points*

$$\mathbf{x}^+ = \mathbf{x}^+(\kappa, \beta) \in \mathfrak{X}_{2N}, \quad \mathbf{x}^0 = \mathbf{x}^0(\kappa, \beta) \in \mathfrak{X}_{2N}, \quad \mathbf{x}^- = \mathbf{x}^-(\kappa, \beta) \in \mathfrak{X}_{2N},$$

*such that*

$$\mathcal{F}_\beta^{(\kappa)}(\mathbf{x}^+) > 0, \quad \mathcal{F}_\beta^{(\kappa)}(\mathbf{x}^0) = 0, \quad \mathcal{F}_\beta^{(\kappa)}(\mathbf{x}^-) < 0. \quad (1.17)$$




Figure 1.5: Plot of the fugacity  $\kappa \mapsto \nu(\kappa)$  (orange) and the constant  $\kappa \mapsto C(\kappa)$  (blue). Note that  $\nu(\kappa) = 0$  at points of the form  $\kappa = \frac{8}{m}$  with  $m \in 2\mathbb{Z}_{>0} + 1$  odd, and  $\nu(\kappa) \in \{\pm 2\}$  at points of the form  $\kappa = \frac{8}{m}$  with  $m \in 2\mathbb{Z}_{>0}$  even. The constant  $C(\kappa)$  diverges at points of the form  $\kappa = \frac{8}{m}$  with  $m \in 2\mathbb{Z}_{>0}$  even (where  $\nu(\kappa) \in \{\pm 2\}$ ).

We prove Theorem 1.5 in Section 5 by combining analysis of the Coulomb gas integrals and multiple SLE theory. In particular, we shall use the relation between the Coulomb gas integrals and the pure partition functions of multiple  $SLE_\kappa$ , discussed in Section 1.4. The novel parts of Theorem 1.5 are Item 1 and the analysis of the zeroes in (POS), (NUL), (SGN) of Item 2, which we find quite curious.

**Remark 1.6.** In the case where  $N = 2$ , we give a detailed analysis of the zeroes of the functions  $\{\mathcal{F}_{\beta}^{(\kappa)}, \mathcal{F}_{\beta}^{(\kappa)}\}$  in Proposition D.4 in terms of the cross-ratio  $\text{CR}(x_1, x_2, x_3, x_4)$ . Interestingly, the cross-ratio of the zeroes oscillates as  $\kappa \rightarrow 0$ , see Eq. (D.15).

Let us elaborate on the exceptional values of  $\kappa$  where the constant  $C(\kappa)$  is not well-defined. The Coulomb gas integrals have different behavior for

$$\kappa \in \left\{ \frac{4}{m} : m \in \mathbb{Z}_{>0} \right\} \quad \text{or} \quad \kappa \in \left\{ \frac{8}{2m+1} : m \in \mathbb{Z}_{>0} \right\} \quad \text{or} \quad \kappa = 8. \quad (1.18)$$

- For  $\kappa \in \left\{ \frac{4}{m} : m \in \mathbb{Z}_{>0} \right\}$ , the constant  $C(\kappa)$  diverges (see Figure 1.5) and the integral of  $f_{\beta}^{(\kappa)}(\mathbf{x}; \mathbf{u})$  is zero (see Lemma 2.14). Thus, it is not clear what the value of the product of the two is. We verify in Theorem 1.5 that  $\mathcal{F}_{\beta}^{(\kappa)}$  is finite as a continuous extension in this case.
- For  $\kappa \in \left\{ \frac{8}{2m+1} : m \in \mathbb{Z}_{>0} \right\}$ , the constant  $C(\kappa)$  is finite (as a continuous extension, see Figure 1.5) and the integral of  $f_{\beta}^{(\kappa)}(\mathbf{x}; \mathbf{u})$  is zero (see Lemma 2.14), so one expects that the product of the two is zero. We verify in Theorem 1.5 that  $\mathcal{F}_{\beta}^{(\kappa)}$  is indeed zero (as a continuous extension) in this case. Furthermore, we will show that  $\mathcal{F}_{\beta}^{(\kappa')}$  tends to zero at rate  $|\kappa' - \kappa|$  as  $\kappa' \rightarrow \kappa$ , see Proposition 1.7.
- For  $\kappa = 8$ , the constant  $C(8)$  is zero and the integral of  $f_{\beta}^{(8)}(\mathbf{x}; \mathbf{u})$  is finite. Thus,  $\mathcal{F}_{\beta}^{(8)}$  is zero. Furthermore, we will show that  $\mathcal{F}_{\beta}^{(\kappa')}$  tends to zero at rate  $(8 - \kappa')^N$  as  $\kappa' \rightarrow 8-$ , see Proposition 1.8.

**Proposition 1.7.** Fix  $\kappa \in \left\{ \frac{8}{2m+1} : m \in \mathbb{Z}_{>0} \right\}$ . Each Coulomb gas integral  $\mathcal{F}_{\beta}^{(\kappa')}$  tends to zero at rate  $(\kappa' - \kappa)$  as  $\kappa' \rightarrow \kappa$ :

$$\widehat{\mathcal{F}}_{\beta}^{(\kappa)}(\mathbf{x}) := \lim_{\kappa' \rightarrow \kappa} \frac{-\kappa^2}{8\pi \sin(4\pi/\kappa)} \frac{\mathcal{F}_{\beta}^{(\kappa')}(\mathbf{x})}{(\kappa' - \kappa)} \in (0, \infty), \quad \mathbf{x} \in \mathfrak{X}_{2N}, \beta \in \text{LP}_N. \quad (1.19)$$

In particular,  $\{\widehat{\mathcal{F}}_{\beta}^{(\kappa)} : \beta \in \text{LP}_N\}$  defined in (1.19) are multiple  $SLE_\kappa$  partition functions.

**Proposition 1.8.** Fix  $\kappa = 8$ . Each Coulomb gas integral  $\mathcal{F}_\beta^{(\kappa')}$  tends to zero at rate  $(8 - \kappa')^N$  as  $\kappa' \rightarrow 8-$ :

$$\widehat{\mathcal{F}}_\beta^{(8)}(\mathbf{x}) := \lim_{\kappa' \rightarrow 8-} \frac{8}{\pi} \frac{\mathcal{F}_\beta^{(\kappa')}(\mathbf{x})}{(8 - \kappa')^N} \in (0, \infty), \quad \mathbf{x} \in \mathfrak{X}_{2N}, \beta \in \text{LP}_N. \quad (1.20)$$

In particular,  $\{\widehat{\mathcal{F}}_\beta^{(8)} : \beta \in \text{LP}_N\}$  defined in (1.20) are multiple  $\text{SLE}_8$  partition functions.

### 1.3 Pure partition functions

We now turn to SLE theory. Following Oded Schramm’s seminal idea to describe critical interfaces in planar models dynamically as Loewner evolutions, whose behavior is governed by uniformizing conformal maps and a one-dimensional driving process (a variant of Brownian motion), one is led to analyzing the set of various natural driving processes. For multiple chordal curves, these processes are classified in terms of multiple  $\text{SLE}_\kappa$  *partition functions*: smooth positive functions  $\mathcal{Z} : \mathfrak{X}_{2N} \rightarrow (0, \infty)$  satisfying the hypoelliptic partial differential equations (PDE) (which essentially follow because the partition function describes a tilting of the Brownian motion by a local martingale), and the full Möbius covariance (COV), a consequence of the conformal invariance of the curves.

In the present work, we will give a complete classification and construction of multiple chordal SLE curves and their partition functions for the full parameter range  $\kappa \in (0, 8)$  in general simply connected domains. The extremal points in the convex set of multiple chordal SLE measures are provided by the “pure” partition functions, which also appear in the numerator in Conjecture 1.2.

**Definition 1.9.** Fix  $\kappa \in (0, 8)$ . The *pure partition functions* of multiple  $\text{SLE}_\kappa$  are the recursive collection  $\{\mathcal{Z}_\alpha^{(\kappa)} : \alpha \in \text{LP}\}$  of positive functions  $\mathcal{Z}_\alpha^{(\kappa)} : \mathfrak{X}_{2N} \rightarrow (0, \infty)$ , indexed by planar link patterns  $\alpha = \{\{a_1, b_1\}, \dots, \{a_N, b_N\}\}$  as in (1.2), uniquely determined by the PDE system (PDE), Möbius covariance (COV), the power-law bound (PLB), as well as the following recursive asymptotics property (ASY):

(ASY) **Asymptotics:** With  $\mathcal{Z}_\emptyset^{(\kappa)} \equiv 1$  for the empty link pattern  $\emptyset \in \text{LP}_0$ , the collection  $\{\mathcal{Z}_\alpha^{(\kappa)} : \alpha \in \text{LP}_N\}$  satisfies the following recursive asymptotics property. Fix  $N \geq 1$  and  $j \in \{1, 2, \dots, 2N - 1\}$ . Then,

$$\lim_{x_j, x_{j+1} \rightarrow \xi} \frac{\mathcal{Z}_\alpha^{(\kappa)}(\mathbf{x})}{(x_{j+1} - x_j)^{-2h(\kappa)}} = \begin{cases} \mathcal{Z}_{\alpha/\{j, j+1\}}^{(\kappa)}(\check{\mathbf{x}}_j), & \text{if } \{j, j+1\} \in \alpha, \\ 0, & \text{if } \{j, j+1\} \notin \alpha, \end{cases} \quad (\text{ASY})$$

where  $\alpha/\{j, j+1\} \in \text{LP}_{N-1}$  denotes the link pattern obtained from  $\alpha$  by removing the link  $\{j, j+1\}$  and relabeling the remaining indices by  $1, 2, \dots, 2N - 2$ , and  $\xi \in (x_{j-1}, x_{j+2})$  and where

$$\mathbf{x} = (x_1, \dots, x_{2N}) \in \mathfrak{X}_{2N} \quad \text{and} \quad \check{\mathbf{x}}_j = (x_1, \dots, x_{j-1}, x_{j+2}, \dots, x_{2N}) \in \mathfrak{X}_{2N-2}. \quad (1.21)$$

Relaxing the positivity requirement, the collection  $\{\mathcal{Z}_\alpha^{(\kappa)} : \alpha \in \text{LP}_N\}$  was first constructed in [FK15c] indirectly by using Coulomb gas integrals for all  $\kappa \in (0, 8)$ , and explicitly for all  $\kappa \in (0, 8) \setminus \mathbb{Q}$  in [KP16], following the conjectures from [BBK05]. For  $\kappa \in (0, 6]$ , an explicit probabilistic construction was given in [Wu20, Theorem 1.7]. Recently, Zhan constructed global multiple chordal  $\text{SLE}_\kappa$  measures for  $\kappa \in (0, 6]$ , thus also giving another construction of the pure partition functions in this range [Zha23].

Also recently, Ang, Holden, Sun & Yu gave a construction of some SLE partition functions using LQG theory [AHSY23] for  $\kappa \in (6, 8)$ . In this construction, they however do not derive the asymptotic properties of their partition functions. Nevertheless, we will show that using global multiple SLE measures as a bridge, we indirectly know that these partition functions agree with the pure partition functions (up to a multiplicative constant) — see Remark 4.23 in Section 4.8.

Now, the second main purpose of the present article is to give a direct SLE theory construction for the pure partition functions with the parameter range  $\kappa \in (4, 8)$  (Theorem 1.10) and use it to construct global multiple SLEs (Theorem 1.21). Let us emphasize a couple of crucial aspects of our results:

- The pure partition functions are well-defined for all  $\kappa \in (0, 8)$  and continuous in  $\kappa$ . Moreover, the limit  $\kappa \rightarrow 8-$  can be normalized to obtain a pure partition function associated to  $\text{SLE}_8$ .
- From the probabilistic construction it readily follows that the pure partition functions are *positive*. This seems not obvious at all from the Coulomb gas perspective.
- The recursive asymptotics property (ASY) of the pure partition functions is their most important, characterizing property, providing the required boundary values for the unique determination of the hypoelliptic PDE boundary value problem (PDE)–(COV). It is important to note that the proof of the asymptotics properties (ASY) for  $\kappa \in (6, 8)$  is quite hard (see Section 4.6), because  $h(\kappa) < 0$  and there are no bounds available to exchange the limit and expected value (cf. [Pel19, Appendix B]).
- As a consequence of their asymptotics (ASY), the pure partition functions are linearly independent. Hence, they can be regarded as the “extremal” partition functions for multiple  $\text{SLE}_\kappa$  processes.

**Theorem 1.10.** *The collection  $\{\mathcal{Z}_\alpha^{(\kappa)} : \alpha \in \text{LP}\}$  of pure partition functions exists and is unique for all  $\kappa \in (0, 8)$ . They are linearly independent and have the following further properties.*

1. *They are continuous in  $(0, 8) \ni \kappa$ : for each  $\kappa \in (0, 8)$ , we have*

$$\lim_{\kappa' \rightarrow \kappa} \mathcal{Z}_\alpha^{(\kappa')}(\mathbf{x}) = \mathcal{Z}_\alpha^{(\kappa)}(\mathbf{x}), \quad \mathbf{x} \in \mathfrak{X}_{2N}, \alpha \in \text{LP}_N. \quad (1.22)$$

2. *They tend to zero at rate  $(8 - \kappa')^{N-1}$  as  $\kappa' \rightarrow 8-$ , i.e., the following limit exists:*

$$\widehat{\mathcal{Z}}_\alpha^{(8)}(\mathbf{x}) := \lim_{\kappa' \rightarrow 8-} \frac{\mathcal{Z}_\alpha^{(\kappa')}(\mathbf{x})}{(8 - \kappa')^{N-1}} \in (0, \infty), \quad \mathbf{x} \in \mathfrak{X}_{2N}, \alpha \in \text{LP}_N. \quad (1.23)$$

We prove Theorem 1.10 in Section 5.5 using the relation between the pure partition functions and Coulomb gas integrals (see Section 1.4). Note that (1.23) tells that the pure partition functions decay as a polynomial in  $\kappa$  as  $\kappa \rightarrow 8-$  when  $N \geq 2$ . In contrast to this polynomial decay, the pure partition functions explode exponentially fast as  $\kappa \rightarrow 0+$ : the quantity  $\kappa \log \mathcal{Z}_\alpha^{(\kappa)}$  has a finite limit as  $\kappa \rightarrow 0$ , see [PW24, Eq. (1.10) & Theorem 1.5]. Interestingly, the limit functions also satisfy the semiclassical BPZ equations, by [PW24, Proposition 1.8]. See also [ABKM20].

## 1.4 Relation between pure partition functions and Coulomb gas integrals

In fact, both collections  $\{\mathcal{F}_\alpha^{(\kappa)} : \alpha \in \text{LP}_N\}$  and  $\{\mathcal{Z}_\alpha^{(\kappa)} : \alpha \in \text{LP}_N\}$  are solutions to the same system (PDE), and they are related by a linear transformation encoded in the meander matrix (Definition 1.1) — which is however not always invertible (see [DFGG97, Eq. (5.6)] and Lemma 5.5).

**Proposition 1.11.** *Fix  $\kappa \in (0, 8)$  and  $\nu = \nu(\kappa)$  as in (1.11). The Coulomb gas integrals in Definition 1.4 are linear combinations of pure partition functions:*

$$\mathcal{F}_\beta^{(\kappa)}(\mathbf{x}) = \sum_{\alpha \in \text{LP}_N} \mathcal{M}_\nu(\alpha, \beta) \mathcal{Z}_\alpha^{(\kappa)}(\mathbf{x}), \quad \mathbf{x} \in \mathfrak{X}_{2N}, \beta \in \text{LP}_N. \quad (1.24)$$

*In particular, the functions  $\{\mathcal{F}_\beta^{(\kappa)} : \beta \in \text{LP}_N\}$  of Definition 1.10 are linearly independent for*

$$\kappa \in (0, 8) \setminus \left\{ \frac{4p}{p'} : p, p' \in \mathbb{Z}_{>0} \text{ coprime and } 2 \leq p \leq N + 1 \right\}. \quad (1.25)$$

Proposition 1.11 can be proven by similar arguments as [FPW24a, Proposition 1.10]. Indeed, the asserted equation (1.24) follows from knowing that both sides of it satisfy the same PDE boundary value problem (PDE, COV,  $\mathcal{F}$ -ASY), where the asymptotic boundary conditions ( $\mathcal{F}$ -ASY) are specified in Corollary 2.13 in Section 2.5. Invoking [FK15b, Lemma 1] shows that such a function is unique as long as the power-law bound (PLB) satisfied. The exceptional values of  $\kappa$  require slightly more care.

**Example 1.12.** For instance, the meander matrix with  $N = 3$  reads

$$\left( \begin{array}{c|ccccc} & \text{---} & \text{---} & \text{---} & \text{---} & \text{---} \\ \hline \text{---} & \nu^3 & \nu^2 & \nu^2 & \nu & \nu^2 \\ \text{---} & \nu^2 & \nu^3 & \nu & \nu^2 & \nu \\ \text{---} & \nu^2 & \nu & \nu^3 & \nu^2 & \nu \\ \text{---} & \nu & \nu^2 & \nu^2 & \nu^3 & \nu^2 \\ \text{---} & \nu^2 & \nu & \nu & \nu^2 & \nu^3 \end{array} \right)$$

Its determinant is  $\nu^5(\nu^2 - 2)(\nu^2 - 1)^4$ , which is zero when  $\nu = 0$  or  $\nu = \pm\sqrt{2}$  or  $\nu = \pm 1$ .

Let us discuss three exceptional values in (1.25):  $\kappa \in \{3, 16/3, 6\}$ . In particular,  $\nu(6) = \nu(3) = 1$  and  $\nu(16/3) = \sqrt{2}$ , which are roots of the determinant of meander matrix with  $N = 3$ . In these three cases, the Coulomb gas integrals are linearly dependent in general, but they have explicit expressions.

**Remark 1.13.**

- When  $\kappa = 6$ , we have  $\nu(6) = 1$ . The meander matrix  $\mathcal{M}_\nu$  is not invertible as long as  $N \geq 2$ . All Coulomb gas integrals are equal, given by

$$\mathcal{F}_\beta^{(6)}(\mathbf{x}) = 1, \quad \mathbf{x} \in \mathfrak{X}_{2N}, \beta \in \text{LP}_N. \quad (1.26)$$

This is a feature of the target-independence of  $\text{SLE}_6$ , or the fact that Bernoulli percolation is insensitive to boundary conditions (cf. Conjecture 1.2).

- When  $\kappa = 3$ , we have  $\nu(3) = 1$ . The meander matrix  $\mathcal{M}_\nu$  is not invertible as long as  $N \geq 2$ . All Coulomb gas integrals are equal, given by a well-known explicit Pfaffian formula (related to fermion, or boundary energy density correlations in the Ising model [HS13, Izy15], see also [PW23]),

$$\mathcal{F}_\beta^{(3)}(\mathbf{x}) = \text{pf} \left( \frac{1}{x_j - x_i} \right)_{i,j=1}^{2N}, \quad \mathbf{x} \in \mathfrak{X}_{2N}, \beta \in \text{LP}_N. \quad (1.27)$$

- When  $\kappa = 16/3$ , we have  $\nu(16/3) = \sqrt{2}$ . The meander matrix  $\mathcal{M}_\nu$  is not invertible as long as  $N \geq 3$ . In this case, the Coulomb gas integrals are linearly dependent [FPW24a, Theorem 1.5]

$$\mathcal{F}_\beta^{(16/3)}(\mathbf{x}) = \prod_{s=1}^N |x_{b_s} - x_{a_s}|^{-1/8} \left( \sum_{\sigma \in \{\pm 1\}^N} \prod_{1 \leq s < t \leq N} \left( \frac{|x_{a_t} - x_{a_s}| |x_{b_s} - x_{b_t}|}{|x_{b_t} - x_{a_s}| |x_{b_s} - x_{a_t}|} \right)^{\sigma_s \sigma_t / 4} \right)^{1/2}, \quad (1.28)$$

$$\mathbf{x} \in \mathfrak{X}_{2N}, \beta \in \text{LP}_N.$$

where  $\sigma = (\sigma_1, \dots, \sigma_N) \in \{\pm 1\}^N$ , reminiscent to spin correlation functions in the Ising model [Izy22].

**Remark 1.14.** The (Duplantier) duality of  $\text{SLE}_\kappa$  is a symmetry in the parameter  $\kappa$  preserving the central charge  $c(\kappa) = (3\kappa - 8)(6 - \kappa)/2\kappa$ : if  $\kappa \leq 4$  and  $\kappa' \geq 4$ , then Eq. (1.7) shows that

$$\kappa\kappa' = 16 \quad \iff \quad c(\kappa) = c(\kappa').$$

That this relation has a deeper meaning for  $\text{SLE}_\kappa$  curves was predicted by Bertrand Duplantier using ideas from quantum gravity (see, e.g., [Dup06]). This duality is now well-understood in terms of Hausdorff dimensions of SLE curves: the frontier of  $\text{SLE}_{\kappa'}$  curve is locally the same as  $\text{SLE}_\kappa$  curve, and in particular, the frontier of  $\text{SLE}_{\kappa'}$  has the same Hausdorff dimension as  $\text{SLE}_\kappa$  (see, e.g., [Dub07b, Zha08]).

In contrast, there is another symmetry that is much more mysterious. Namely, for the fugacity  $\nu(\kappa) = -2 \cos(4\pi/\kappa)$ , dilute and dense models with the same fugacity share similar algebraic structure. Indeed, Eq. (1.11) shows that if  $\kappa \in (8/3, 4]$  and  $\hat{\kappa} \in [4, 8)$ , then

$$\frac{1}{\kappa} + \frac{1}{\hat{\kappa}} = \frac{1}{2} \quad \iff \quad \nu(\kappa) = \nu(\hat{\kappa}).$$

The fugacity parameter  $\nu(\kappa)$  appears as the parameter of the Temperley-Lieb algebra (see, e.g., [FK15c, FP18, FP20, GSNJ<sup>+</sup>22]), and  $q(\kappa) = \exp(4\pi i/\kappa)$  as the parameter of the associated quantum group  $U_q(\mathfrak{sl}(2, \mathbb{C}))$  (see, e.g., [KP20]), where the symmetry  $\kappa \leftrightarrow \hat{\kappa}$  corresponds with  $q(\hat{\kappa}) = 1/q(\kappa) = \exp(-4\pi i/\kappa)$ . A special case of this symmetry is the connection between the Ising model and percolation:

$$\frac{1}{3} + \frac{1}{6} = \frac{1}{2}.$$

If one considers the critical-temperature Ising model as a special case of Conjecture 1.2 with  $n = 1$  and  $p = p_c(1)$ , then the high-temperature Ising model corresponds to the special case of Conjecture 1.2 with  $n = 1$  and  $p > p_c(1)$ . This is related to the conjecture that the universality class of high-temperature Ising model is the same as that of critical Bernoulli percolation, see, e.g. [KN04, Smi06, BCM10]. In (1.26) and (1.27), we see that Coulomb gas integrals are equal for both  $\kappa = 3$  and  $\kappa = 6$ , which consistent with the fact that  $\nu(3) = \nu(6) = 1$ .

In the cases where the Coulomb gas integrals identically vanish, we may renormalize them as in Propositions 1.7 & 1.8. We also define the *renormalized meander matrix*  $\{\widehat{\mathcal{M}}(\alpha, \beta) : \alpha, \beta \in \text{LP}_N\}$  via

$$\widehat{\mathcal{M}}(\alpha, \beta) := \lim_{\nu \rightarrow 0} \frac{1}{\nu} \mathcal{M}_\nu(\alpha, \beta) = \begin{cases} 1, & \text{if the meander formed from } \alpha \text{ and } \beta \text{ has one loop,} \\ 0, & \text{otherwise.} \end{cases} \quad (1.29)$$

Using the renormalized meander matrix, we can extend the relations in Proposition 1.11 to the exceptional values of  $\kappa$  (see Section 5.4). In this way, we obtain partition functions for  $\text{SLE}_\kappa$  processes with exceptional values of  $\kappa$ . It would be interesting to find probabilistic or CFT interpretations for these partition functions. We discuss some possible CFT relations in Section 1.5.

**Proposition 1.15.**

1. Fix  $\kappa \in \{\frac{8}{2m+1} : m \in \mathbb{Z}_{>0}\}$ . The renormalized Coulomb gas integral (1.19) is a linear combination of pure partition functions:

$$\widehat{\mathcal{F}}_\beta^{(\kappa)}(\mathbf{x}) = \sum_{\alpha \in \text{LP}_N} \widehat{\mathcal{M}}(\alpha, \beta) \mathcal{Z}_\alpha^{(\kappa)}(\mathbf{x}) \in (0, \infty), \quad \mathbf{x} \in \mathfrak{X}_{2N}, \beta \in \text{LP}_N. \quad (1.30)$$

In particular, the functions  $\{\widehat{\mathcal{F}}_\beta^{(\kappa)} : \beta \in \text{LP}_N\}$  defined in (1.19) are linearly independent.

2. Fix  $\kappa = 8$ . The renormalized Coulomb gas integral (1.20) is a linear combination of the renormalized pure partition functions (1.23):

$$\widehat{\mathcal{F}}_\beta^{(8)}(\mathbf{x}) = \sum_{\alpha \in \text{LP}_N} \widehat{\mathcal{M}}(\alpha, \beta) \widehat{\mathcal{Z}}_\alpha^{(8)}(\mathbf{x}) \in (0, \infty), \quad \mathbf{x} \in \mathfrak{X}_{2N}, \beta \in \text{LP}_N. \quad (1.31)$$

In particular, both  $\{\widehat{\mathcal{F}}_\beta^{(8)} : \beta \in \text{LP}_N\}$  and  $\{\widehat{\mathcal{Z}}_\alpha^{(8)} : \alpha \in \text{LP}_N\}$  are linearly independent.

### 1.5 Asymptotics and Fröbenius series

The pure partition functions (Section 1.3) are singled out as unique solutions to the PDE boundary value problem (PDE)–(COV) via their recursive asymptotics property (ASY), as in Theorem 1.10. In CFT parlance, the two possible asymptotics in (ASY) correspond to two possible fusion channels of the degenerate field  $\Phi_{1,2}$  (of conformal weight  $h_{1,2}$  (1.8)) into the identity field  $\Phi_{1,1}$  and the field  $\Phi_{1,3}$  which is degenerate at level three. Indeed, the corresponding exponents in (ASY) have the form

$$-2h(\kappa) = h_{1,1}(\kappa) - 2h_{1,2}(\kappa) \quad \text{and} \quad \frac{2}{\kappa} = h_{1,1}(\kappa) - 2h_{1,2}(\kappa), \quad (1.32)$$

where  $h_{1,1}(\kappa) = 0$  and  $h_{1,3}(\kappa) = (8 - \kappa)/\kappa$  (see [Pel19] and references therein for more background). The Coulomb gas integrals are generically (i.e., when  $\kappa \notin \{\frac{8}{m} : m \in \mathbb{Z}_{>0}\}$ ), also singled out as unique solutions to the PDE boundary value problem (PDE)–(COV) but with different asymptotic boundary conditions — see Corollary 2.13. In the present work, we will also investigate the subleading terms in the expansion of the Coulomb gas integrals when two of the variables collide. For simplicity of notation, and without loss of generality, we only consider the case where  $|x_2 - x_1| \rightarrow 0$ . Analogously to the solutions of the hypergeometric equation (see also Appendix D), solutions to (PDE) admit a Fröbenius series with the two indicial exponents (1.32). We investigate the explicit terms in this expansion in Proposition 1.16 (with some exceptional cases in Propositions 1.17 and 1.18). Notably, these terms agree with the findings in [FP18, FP20, KP20, FP24], where a quantum group structure on the solution space of (PDE)–(COV) was found and investigated, as well as with the fusion properties expected from (possibly logarithmic) minimal models in the physics literature [DFMS97, PRZ06, RS07]. Observe that it is evident from the Coulomb gas integrals in general (see [Pel20] for the generic case) that they describe correlation functions outside of CFT minimal models, because their arbitrary fusions are non-zero.

### 1.5.1 Loop $O(n)$ model — generic case

In the terms appearing in the Fröbenius series, we will use the Coulomb gas type integrands with general numbers of  $d \geq 2$  variables and  $\ell \geq 0$  integration variables (with any branch choice),

$$f^{(\kappa)}(\mathbf{x}; \mathbf{u}) := \prod_{1 \leq i < j \leq d} (x_j - x_i)^{2/\kappa} \prod_{1 \leq r < s \leq \ell} (u_s - u_r)^{8/\kappa} \prod_{\substack{1 \leq i \leq d \\ 1 \leq r \leq \ell}} (u_r - x_i)^{-4/\kappa} \quad (1.33)$$

having negative screening charges  $\alpha_- = -2/\sqrt{\kappa}$  at the integration variables  $\mathbf{u} = (u_1, u_2, \dots, u_\ell)$ , and symmetric charges  $\alpha_j = 1/\sqrt{\kappa}$  for all  $j \in \{1, 2, \dots, d\}$  at all of the variables  $\mathbf{x} = (x_1, x_2, \dots, x_d)$ .

**Proposition 1.16.** *Fix  $\kappa \in (8/3, 8)$  and  $N \geq 2$ . Then, each Coulomb gas integral  $\mathcal{F}_\beta^{(\kappa)}$ , with  $\beta \in \text{LP}_N$ , has the following expansion (using the notation (1.21)).*

- If  $\{1, 2\} \in \beta$ , then for all  $\xi \in (-\infty, x_3)$ ,

$$\mathcal{F}_\beta^{(\kappa)}(\mathbf{x}) = \nu(\kappa) \mathcal{F}_{\beta/\{1,2\}}^{(\kappa)}(\tilde{\mathbf{x}}_1) (x_2 - x_1)^{-2h(\kappa)} + o((x_2 - x_1)^{2/\kappa}), \quad \text{as } x_1, x_2 \rightarrow \xi, \quad (1.34)$$

where  $\beta/\{1, 2\} \in \text{LP}_{N-1}$  denotes the link pattern obtained from  $\beta$  by removing the link  $\{1, 2\}$  and relabeling the remaining indices by  $1, 2, \dots, 2N - 2$ .


- If  $\{1, 2\} \notin \beta$ , then for all  $\xi \in (-\infty, x_3)$ ,

$$\mathcal{F}_\beta^{(\kappa)}(\mathbf{x}) = \mathcal{F}_{\wp_1(\beta)/\{1,2\}}^{(\kappa)}(\tilde{\mathbf{x}}_1) (x_2 - x_1)^{-2h(\kappa)} + \mathcal{F}_{\beta/\vee_1}^{(\kappa)}(\xi, \tilde{\mathbf{x}}_1) (x_2 - x_1)^{2/\kappa} + o((x_2 - x_1)^{2/\kappa}), \quad \text{as } x_1, x_2 \rightarrow \xi, \quad (1.35)$$

where  $\wp_1$  is the “tying operation” defined by

$$\wp_1: \text{LP}_N \rightarrow \text{LP}_N, \quad \wp_1(\beta) = (\beta \setminus (\{1, k_1\}, \{2, k_2\})) \cup \{1, 2\} \cup \{k_1, k_2\},$$

where  $k_1$  (resp.  $k_2$ ) is the pair of 1 (resp. 2) in  $\beta$  (and  $\{1, k_1\}, \{2, k_2\}, \{k_1, k_2\}$  are unordered),



and where  $\mathcal{F}_{\beta/\vee_1}^{(\kappa)}(\xi, \tilde{\mathbf{x}}_1)$  is the “fused” Coulomb gas integral function defined as follows.



▷ If  $\kappa \in (8/3, 8) \setminus \{4\}$ , then

$$\mathcal{F}_{\beta/\nu_1}^{(\kappa)}(\xi, \mathbf{\ddot{x}}_1) := \frac{C(\kappa)^{N-2} (\nu(\kappa)^2 - 1)}{2F_1\left(\frac{4}{\kappa}, 1 - \frac{4}{\kappa}, \frac{8}{\kappa}; 1\right)} \oint_{\vartheta(x_{a_3}, x_{b_3})} du_3 \cdots \oint_{\vartheta(x_{a_N}, x_{b_N})} du_N \hat{f}_{\beta/\nu_1}^{(\kappa)}(\xi, \mathbf{\ddot{x}}_1; \mathbf{\ddot{u}}_1), \quad (1.36)$$

where  $\mathbf{\ddot{u}}_1 := (u_3, u_4, \dots, u_N)$  and

$$\hat{f}_{\beta/\nu_1}^{(\kappa)}(\xi, \mathbf{\ddot{x}}_1; \mathbf{\ddot{u}}_1) := \prod_{r=3}^N (u_r - \xi)^{16/\kappa-2} \prod_{j=3}^{2N} (x_j - \xi)^{1-8/\kappa} f^{(\kappa)}(\mathbf{\ddot{x}}_1; \mathbf{\ddot{u}}_1), \quad (1.37)$$

whose branch is chosen to be real and positive when  $x_{a_r} < \text{Re}(u_r) < x_{a_{r+1}}$  for  $r \in \{3, 4, \dots, N\}$ .

▷ If  $\kappa = 4$ , then  $\mathcal{F}_{\beta/\nu_1}^{(4)}(\xi, \mathbf{\ddot{x}}_1)$  is a linear combination of rational functions with explicit formula given by Eq. (E.3) in Proposition E.1.

We prove Proposition 1.16 in Section 6.1.

The Fröbenius series (1.34, 1.35) are consistent with the fusion rules in a CFT (minimal) model with two fields  $\Phi_{1,2}(x_1)$  and  $\Phi_{1,2}(x_2)$  merging to  $\Phi_{1,1}(\xi) = 1$  in the former case, or a combination of  $\Phi_{1,1}(\xi) = 1$  and  $\Phi_{1,3}(\xi)$  in the latter case. The factor  $\mathcal{F}_{\beta/\nu_1}^{(\kappa)}(\xi, \mathbf{\ddot{x}}_1)$  represents a correlation function of a product  $\Phi_{1,3}(\xi)\Phi_{1,2}(x_3) \cdots \Phi_{1,2}(x_{2N})$ , while the Coulomb gas integral functions  $\mathcal{F}_{\beta}^{(\kappa)}(\mathbf{x})$  represents a correlation function of a product  $\Phi_{1,2}(x_1)\Phi_{1,2}(x_2)\Phi_{1,2}(x_3) \cdots \Phi_{1,2}(x_{2N})$ . We also expect the same result to hold for all  $\kappa \in (0, 8) \setminus \{\frac{8}{m} : m \in \mathbb{Z}_{>0}\}$ . However, the detailed analysis of the Coulomb gas integral becomes increasingly complicated in the intervals  $\kappa \in (\frac{8}{4m-1}, \frac{8}{4m-3})$  with  $m = 2, 3, 4, \dots$

The CFT describing the geometry of critical  $O(n)$  loop models is expected to be logarithmic for all  $n \in (0, 2)$  [GZ20, GSNJ<sup>+</sup>22]. At the first level of fusion, however, we only see anomalous behavior in the borderline cases  $\kappa \in \{8/3, 8\}$ , that is,  $m \in \{1, 3\}$ , which correspond respectively to the logarithmic minimal models  $M(2, 1)$  and  $M(2, 3)$  with central charges  $c = -2$  (dense  $O(0)$  model, or UST, Proposition 1.18), and  $c = 0$  (dilute  $O(0)$  model, or self-avoiding walk, Proposition 1.17). More generally, the exceptional values of  $\kappa \in \{\frac{8}{m} : m \in \mathbb{Z}_{>0}\}$  correspond to central charges<sup>3</sup>

$$c(8/m) = 13 - 6(2/m + m/2) = 1 - 6 \frac{(p - p')^2}{pp'}, \quad m = \frac{2p'}{p},$$

with  $p, p' \geq 1$  coprime or  $(p, p') = (1, 1)$ . The case  $m = 2$  (loop  $O(2)$  model) corresponds to  $(p, p') = (1, 1)$  and  $c = 1$  (with  $\kappa = 4$ ). In the three cases (1.18), we have the following identifications<sup>4</sup>.

- For  $\kappa \in \{\frac{4}{m} : m \in \mathbb{Z}_{>0}\}$ , in which case  $\mathcal{F}_{\beta}^{(\kappa)}$  is finite, we see that  $p = 1$  and  $p' = m/2$ . The logarithmic minimal models  $M(1, p')$  were discussed, e.g., in [PRZ06] and references therein.
- For  $\kappa \in \{\frac{8}{2m+1} : m \in \mathbb{Z}_{>0}\}$ , in which case  $\mathcal{F}_{\beta}^{(\kappa)}$  is zero but the renormalized function  $\widehat{\mathcal{F}}_{\beta}^{(\kappa)}$  is well-defined via (1.19), we see that  $p = 2$  and  $p' = m$ . The logarithmic minimal models  $M(2, p')$  were discussed, e.g., in [PRZ06, RS07] and references therein. In Proposition 1.17 we consider the special case where  $m = 3$ , corresponding to the dilute  $O(0)$  model, or self-avoiding walk ( $c = 0$  and  $\kappa = 8/3$ ).
- For  $\kappa = 8$ , in which case  $\mathcal{F}_{\beta}^{(8)}$  is zero but the renormalized function  $\widehat{\mathcal{F}}_{\beta}^{(8)}$  is well-defined via (1.20), we see that  $p = 2$  and  $p' = 1$ . The logarithmic minimal model  $M(2, 1)$  was discussed, e.g., in [PRZ06, PR07, RS07] and references therein. We consider this case, corresponding to the dense  $O(0)$  model, or UST ( $c = -2$  and  $\kappa = 8$ ), in Proposition 1.18.

A minimal model  $M(p, p')$  comprises fields  $\{\Phi_{r,s} : r, s \in \mathbb{Z}_{>0}, 1 \leq r \leq p' - 1, 1 \leq s \leq p - 1, pr > p's\}$ . In all of the above cases, the field  $\Phi_{1,2}$  is absent from the corresponding minimal model. This could heuristically explain why  $\mathcal{F}_{\beta}^{(\kappa)}$  are identically zero in the cases where  $\kappa = 8/m$  with  $m$  odd. (The cases where  $\kappa = 8/m$  with  $m$  even are outside of the minimal models anyway, since  $p = 1$  in these cases.)

<sup>3</sup>Note also that with  $m = \frac{2p'}{p}$ , we have  $\kappa = \frac{4p}{p'}$ .

<sup>4</sup>Traditionally, the cases where  $p, p' \geq 2$  are coprime label the CFT *minimal models*  $M(p, p')$  — theories comprising a finite number of primary fields [DFMS97] — but Coulomb gas integrals describe correlation functions outside these models [Pel20].

### 1.5.2 Logarithmic minimal model $M(2,3)$ — dilute $O(0)$ loops with $\kappa = 8/3$ and $c(\kappa) = 0$

When  $\kappa = 8/3$ , the Coulomb gas integrals vanish (see Lemma 2.14). The renormalized integrals, however, make sense as in Proposition 1.7. We can derive the explicit Fröbenius series also in this case. From the physics perspective, models which are believed to be related to dilute  $O(0)$  loop models and logarithmic minimal models have been considered, e.g., in [RS07]. Gurarie [Gur13, Eq. (45)] computed the OPE of the anticipated field  $\Phi_{1,2}$  in such a model, finding the asymptotic expansion (see also [FSK15, Appendix B])

$$\begin{aligned} \langle \Phi_{1,2}(x_1)\Phi_{1,2}(x_2)\cdots \rangle &= c_1 (x_2 - x_1)^{-5/4} \langle \Phi_{1,1}(\xi)\cdots \rangle + c_2 (x_2 - x_1)^{3/4} \langle \Psi(\xi)\cdots \rangle \\ &\quad + (c_3 + c_2 \log(x_2 - x_1)) (x_2 - x_1)^{3/4} \langle T(\xi)\cdots \rangle \\ &\quad + o((x_2 - x_1)^{3/4}), \quad \text{as } x_1, x_2 \rightarrow \xi. \end{aligned}$$

where  $\Psi(\xi)$  denotes the so-called logarithmic partner of the stress-energy tensor  $T(\xi)$ , and  $c_1$ ,  $c_2$ , and  $c_3$  are some constants (which are explicit in Proposition 1.17, if the functions  $\widehat{\mathcal{F}}_{\beta}^{(8/3)}$ ,  $\widehat{\mathcal{F}}_{\beta/\sqrt{1}}^{(8/3)}$ , and  $\mathcal{G}_{\beta/\sqrt{1}}^{(8/3)}$  are interpreted suitably). The relationship of  $\text{SLE}_{8/3}$  with the stress-energy tensor was observed already by Friedrich & Werner [FW03], but we have not found a mathematically complete treatment.

**Proposition 1.17.** *Fix  $\kappa = 8/3$  and  $N \geq 2$ . Then, each renormalized Coulomb gas integral  $\widehat{\mathcal{F}}_{\beta}^{(8/3)}(\mathbf{x})$  in Proposition 1.7, with  $\beta \in \text{LP}_N$ , has the following expansion.*

- If  $\{1, 2\} \in \beta$ , then for all  $\xi \in (-\infty, x_3)$ ,

$$\widehat{\mathcal{F}}_{\beta}^{(8/3)}(\mathbf{x}) = \widehat{\mathcal{F}}_{\beta/\sqrt{1}}^{(8/3)}(\xi, \ddot{\mathbf{x}}_1) (x_2 - x_1)^{3/4} + o((x_2 - x_1)^{3/4}), \quad \text{as } x_1, x_2 \rightarrow \xi, \quad (1.38)$$

where

$$\widehat{\mathcal{F}}_{\beta/\sqrt{1}}^{(8/3)}(\xi, \ddot{\mathbf{x}}_1) := \frac{C(8/3)^{N-2}}{{}_2F_1\left(\frac{3}{2}, -\frac{1}{2}, 3; 1\right)} \oint_{\vartheta(x_{a_3}, x_{b_3})} du_3 \cdots \oint_{\vartheta(x_{a_N}, x_{b_N})} du_N \widehat{f}_{\beta/\sqrt{1}}^{(8/3)}(\xi, \ddot{\mathbf{x}}_1; \ddot{\mathbf{u}}_1), \quad (1.39)$$

and  $\widehat{f}_{\beta/\sqrt{1}}^{(8/3)}$  is defined in (1.37) with  $\kappa = 8/3$  and

$$C(8/3) = \lim_{\kappa \rightarrow 8/3} C(\kappa) = -\frac{1}{16} \quad \text{and} \quad {}_2F_1\left(\frac{3}{2}, -\frac{1}{2}, 3; 1\right) = \frac{32}{15\pi}.$$

- If  $\{1, 2\} \notin \beta$ , then for all  $\xi \in (-\infty, x_3)$ ,

$$\begin{aligned} \widehat{\mathcal{F}}_{\beta}^{(8/3)}(\mathbf{x}) &= \widehat{\mathcal{F}}_{\wp_1(\beta)\setminus\{1,2\}}^{(8/3)}(\ddot{\mathbf{x}}_1) (x_2 - x_1)^{-5/4} - \frac{1}{\pi} (-1)^{(b_2-1)/2} \widehat{\mathcal{F}}_{\wp_1(\beta)/\sqrt{1}}^{(8/3)}(\xi, \ddot{\mathbf{x}}_1) (x_2 - x_1)^{3/4} |\log(x_2 - x_1)| \\ &\quad + O((x_2 - x_1)^{3/4}), \quad \text{as } x_1, x_2 \rightarrow \xi. \end{aligned} \quad (1.40)$$

We prove Proposition 1.17 in Section 6.2.

### 1.5.3 Logarithmic minimal model $M(2,1)$ — dense $O(0)$ loops with $\kappa = 8$ and $c(\kappa) = -2$

Conjecturally, the geometry of dense  $O(0)$  loops with  $c = -2$  should be closely related to  $\text{SLE}_8$  processes and interfaces in the UST model. The scaling limit of the latter has been intensively studied (e.g., [LSW04, Dub06, KW11a, LPW24]). In the physics literature, possible conformal field theories relevant to this model are among the most investigated logarithmic CFTs — see, e.g., [GL02, RS07] and references therein. In particular, Gurarie [Gur93, Eq. (11,12)] computed the OPE of the anticipated field  $\Phi_{1,2}$ , finding the asymptotic expansion (see also [FSK15, Appendix B])

$$\begin{aligned} \langle \Phi_{1,2}(x_1)\Phi_{1,2}(x_2)\cdots \rangle &= (c_1 + c_2 \log(x_2 - x_1)) (x_2 - x_1)^{1/4} \langle \Phi_{1,1}(\xi)\cdots \rangle + c_2 (x_2 - x_1)^{1/4} \langle \Psi(\xi)\cdots \rangle \\ &\quad + o((x_2 - x_1)^{1/4}), \quad \text{as } x_1, x_2 \rightarrow \xi. \end{aligned}$$

where  $\Psi(\xi)$  denotes the so-called logarithmic partner of the identity field  $\Phi_{1,1}(\xi) = 1$ , and  $c_1$  and  $c_2$  are some constants (which are explicit in Proposition 1.18, if the functions  $\widehat{\mathcal{F}}_{\beta}^{(8)}$  are interpreted suitably).

**Proposition 1.18.** Fix  $\kappa = 8$  and  $N \geq 2$ . The renormalized Coulomb gas integral  $\widehat{\mathcal{F}}_\beta^{(8)}(\mathbf{x})$  in Proposition 1.8, with  $\beta \in \text{LP}_N$ , has the following expansion.

- If  $\{1, 2\} \in \beta$ , then for all  $\xi \in (-\infty, x_3)$ ,

$$\widehat{\mathcal{F}}_\beta^{(8)}(\mathbf{x}) = \frac{\pi}{8} \widehat{\mathcal{F}}_{\beta/\{1,2\}}^{(8)}(\ddot{\mathbf{x}}_1)(x_2 - x_1)^{1/4} + o((x_2 - x_1)^{1/4}), \quad \text{as } x_1, x_2 \rightarrow \xi. \quad (1.41)$$

- If  $\{1, 2\} \notin \beta$ , then for all  $\xi \in (-\infty, x_3)$ ,

$$\widehat{\mathcal{F}}_\beta^{(8)}(\mathbf{x}) = \frac{1}{8} \widehat{\mathcal{F}}_{\emptyset \cup \{1,2\}}^{(8)}(\ddot{\mathbf{x}}_1)(x_2 - x_1)^{1/4} |\log(x_2 - x_1)| + O((x_2 - x_1)^{1/4}), \quad \text{as } x_1, x_2 \rightarrow \xi. \quad (1.42)$$

The expansion (1.41) and a weaker version of (1.42) were proved in [LPW24, Proposition 2.9]. We prove Proposition 1.18 in Section 6.3.

**Remark 1.19.** Illustration for the terms appearing in Fröbenius series in Propositions 1.16, 1.17 and 1.18.

		$(x_2 - x_1)^{1-6/\kappa}$	$(x_2 - x_1)^{2/\kappa}  \log(x_2 - x_1) $	$(x_2 - x_1)^{2/\kappa}$
$\kappa \in (8/3, 8)$	$\{1, 2\} \in \beta$	✓	×	×
$\kappa \in (8/3, 8)$	$\{1, 2\} \notin \beta$	✓	×	✓
$\kappa = 8/3$	$\{1, 2\} \in \beta$	×	×	✓
$\kappa = 8/3$	$\{1, 2\} \notin \beta$	✓	✓	✓
$\kappa = 8$	$\{1, 2\} \in \beta$	✓	×	(✓)
$\kappa = 8$	$\{1, 2\} \notin \beta$	✓	✓	(✓)

## 1.6 Global multiple SLEs

As a by-product, our results also give a construction of a global  $N$ -SLE $_\kappa$  probability measure for all  $\kappa \in (4, 8)$ . Our proofs only rely on SLE techniques and some analysis and PDE theory (in particular, we do not use couplings with the Gaussian free field GFF, except that we do use the reversibility of non-simple chordal SLE $_\kappa$ , proven in [MS16c] via SLE/GFF couplings).

To define the multiple SLE measures, we denote by  $X_{\text{simple}}(\Omega; x_1, x_2)$  the set of continuous simple unparameterized curves in  $\Omega$  connecting  $x_1$  and  $x_2$  such that they only touch the boundary  $\partial\Omega$  in  $\{x_1, x_2\}$ . We denote by  $X_0(\Omega; x_1, x_2)$  the closure of  $X_{\text{simple}}(\Omega; x_1, x_2)$  in the metric topology of the set  $X$  of all planar oriented curves (i.e., continuous mappings from  $[0, 1]$  to  $\mathbb{C}$  modulo reparameterization) endowed with metric<sup>5</sup>

$$\text{dist}(\eta, \tilde{\eta}) := \inf_{\psi, \tilde{\psi}} \sup_{t \in [0, 1]} |\eta(\psi(t)) - \tilde{\eta}(\tilde{\psi}(t))|, \quad \eta, \tilde{\eta} \in X, \quad (1.43)$$

where the infimum is taken over all increasing homeomorphisms  $\psi, \tilde{\psi}: [0, 1] \rightarrow [0, 1]$ .

<sup>5</sup>The metric space  $(X, \text{dist})$  is complete and separable.

Note that curves in  $X_0(\Omega; x_1, x_2)$  may have multiple points. In particular, the chordal  $\text{SLE}_\kappa$  curves belong to this space almost surely when  $\kappa > 4$  [RS05]. (See Section 3.1 for a brief survey of the definition of SLE curves.) For  $N \geq 2$  and  $\alpha = \{\{a_1, b_1\}, \{a_2, b_2\}, \dots, \{a_N, b_N\}\} \in \text{LP}_N$ , we denote by<sup>6</sup>

$$X_0^\alpha(\Omega; x_1, \dots, x_{2N})$$

the collection of curves  $(\eta_1, \dots, \eta_N)$  such that, for each  $s \in \{1, \dots, N\}$ , we have  $\eta_s \in X_0(\Omega; x_{a_s}, x_{b_s})$  and  $\eta_s$  does not disconnect any two points  $x_a, x_b$  such that  $\{a, b\} \in \alpha$  from each other. We also set

$$X_0(\Omega; x_1, \dots, x_{2N}) = \bigsqcup_{\alpha \in \text{LP}_N} X_0^\alpha(\Omega; x_1, \dots, x_{2N}). \quad (1.44)$$

**Definition 1.20.** Fix  $\kappa \in (0, 8)$  and  $N \geq 2$  and  $\alpha = \{\{a_1, b_1\}, \dots, \{a_N, b_N\}\} \in \text{LP}_N$ . We call a probability measure on the families  $(\eta_1, \dots, \eta_N) \in X_0^\alpha(\Omega; x_1, \dots, x_{2N})$  a *global  $N$ -SLE $_\kappa$  associated to  $\alpha$*  if it satisfies the following resampling property: for each  $s \in \{1, \dots, N\}$ , the conditional law of the curve  $\eta_s$  given  $\{\eta_1, \dots, \eta_N\} \setminus \{\eta_s\}$  is the chordal  $\text{SLE}_\kappa$  connecting  $x_{a_s}$  and  $x_{b_s}$  in the connected component of the domain  $\Omega \setminus \bigcup_{r \neq s} \eta_r$  having the endpoints  $x_{a_s}$  and  $x_{b_s}$  of  $\eta_s$  on its boundary.

A Loewner chain *associated to the partition function  $\mathcal{Z}$*  starting from  $x_i$  (and with launching points  $(x_1, \dots, x_{2N})$ ) is the Loewner chain whose driving function  $W$  solves the following system of SDEs:

$$\begin{cases} dW_t = \sqrt{\kappa} dB_t + \kappa(\partial_i \log \mathcal{Z})(V_t^1, \dots, V_t^{i-1}, W_t, V_t^{i+1}, \dots, V_t^{2N}) dt, & W_0 = \varphi(x_i), \\ dV_t^j = \frac{2dt}{V_t^j - W_t}, & V_0^j = \varphi(x_j), \quad j \in \{1, \dots, i-1, i+1, \dots, 2N\}, \end{cases} \quad (1.45)$$

Note that the chordal  $\text{SLE}_\kappa$  in  $\mathbb{H}$  from  $x$  to  $y$  is the Loewner chain associated to the partition function  $\mathcal{Z}(x, y) = |x - y|^{-2h(\kappa)}$ . Note also that in (1.45) it suffices to consider the partition function  $\mathcal{Z}$  only up to a multiplicative constant — which however is canonically fixed by requiring the asymptotics (ASY).

**Theorem 1.21.** Fix  $\kappa \in (0, 8)$  and  $\alpha \in \text{LP}_N$ .

1. There exists a unique probability measure on global  $N$ -SLE $_\kappa$  associated to  $\alpha$ .
2. Suppose  $(\eta_1, \dots, \eta_N)$  has the law of global  $N$ -SLE $_\kappa$  associated to  $\alpha$ . Then, for each  $s \in \{1, 2, \dots, N\}$ , the marginal law of  $\eta_s$  is a Loewner chain associated to the pure partition function  $\mathcal{Z}_\alpha$ .

When  $\kappa \in (0, 4]$ , the existence of global  $N$ -SLE $_\kappa$  is given in [KL07] and [PW19, Theorem 1.3], and the uniqueness of such a measure was proven in [BPW21, Theorem 1.2] (see also [MS16b] and [MSW20, Appendix A] for the case  $N = 2$ ). The marginal law was also considered in [PW19, Proposition 3.5] for  $\kappa \in (0, 4]$ . The recent [AHSY23] gives a construction of global  $N$ -SLE $_\kappa$  and derives its marginal law for  $\kappa \in (4, 8)$ , and the recent [Zha23] concerns the uniqueness for  $\kappa \in (4, 8)$ . In this article, we obtain a different construction for  $\kappa \in (4, 8)$  and derive the marginal law in Section 4.8.

**Remark 1.22.** In the chordal case, the resampling property uniquely determines global multiple SLE $_\kappa$  with  $\kappa \in (0, 8)$ . Such a characterization fails in the radial setting: there exists a one-parameter continuous family of two-sided radial SLE with resampling property, which is called two-sided radial SLE with spiral — hence, the resampling property does not uniquely determine two-sided radial SLE due to the additional degree of freedom in the spiral, see [WW24, Section 1.4].

---

<sup>6</sup>The space  $X_0^\alpha(\Omega; x_1, \dots, x_{2N})$  is not complete.

**Outline of this article.** In Section 2, we derive various properties of Coulomb gas integrals, including a general Möbius covariance property, a cyclic permutation relation, and recursive asymptotics properties. In Section 3, we construct a conformally invariant measure on multiple  $\text{SLE}_\kappa$  curves which will be used to construct pure partition functions and global multiple SLEs for  $\kappa \in (4, 8)$  in Section 4. In particular, we complete the proof for Theorem 1.21 in Section 4. In Section 5, using the tools from preceding sections, we complete the proofs of Theorems 1.5 & 1.10, Propositions 1.7 & 1.8, and Propositions 1.11 & 1.15. In Section 6, prove Propositions 1.16–1.18 concerning Fröbenius series expansions.

The bulk of this article is supplemented by several appendices that contain technical material or illuminating examples. The most important ones are the first Appendix A, where we argue that general SLE local martingales give rise to smooth functions solving BPZ type differential equations, and the last Appendix E, where we derive explicit formulas for partition functions in the case  $\kappa = 4$ . The second Appendix B gathers some useful simple integral identities, and the third Appendix C gathers useful but technical Coulomb gas integral formulas for usage in Sections 3–4. The fourth Appendix D contains illuminating examples of properties of the partition functions.

## Acknowledgments.

- We thank Bernard Nienhuis for several interesting discussions related to  $O(n)$  models and Coulomb gas. We thank Nicolai Makarov and Jiaxin Zhang for inspiring discussions about Coulomb gas, and Xin Sun for stimulating discussions about multiple SLEs. We thank Masha Gordina and Dapeng Zhan for very useful correspondences for Appendix A.
- This material is part of a project that has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (101042460): ERC Starting grant “Interplay of structures in conformal and universal random geometry” (ISCoURaGe) and from the Academy of Finland grant number 340461 “Conformal invariance in planar random geometry.” E.P. is also supported by the Academy of Finland Centre of Excellence Programme grant number 346315 “Finnish centre of excellence in Randomness and Structures (FiRST)” and by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany’s Excellence Strategy EXC-2047/1-390685813, as well as the DFG collaborative research centre “The mathematics of emerging effects” CRC-1060/211504053.
- Part of this research was performed while Y.F., E.P., and H.W. were visiting the Institute for Pure and Applied Mathematics (IPAM), which is supported by the National Science Foundation (Grant No. DMS-1925919).
- M.L. is supported by Knut and Alice Wallenberg Foundation.
- Y.F. and H.W. are supported by Beijing Natural Science Foundation (JQ20001). H.W. is partly affiliated at Yanqi Lake Beijing Institute of Mathematical Sciences and Applications, Beijing, China.

## 2 Coulomb gas integrals

Throughout, we consider link patterns  $\beta \in \text{LP}_N$  with link endpoints ordered as in (1.2). Recall that

$$\mathcal{F}_\beta = \mathcal{F}_\beta^{(\kappa)} := C(\kappa)^N \mathcal{H}_\beta^{(\kappa)}, \quad \text{where} \quad C(\kappa) := \frac{\nu(\kappa)}{4 \sin^2(4\pi/\kappa)} \frac{\Gamma(2 - 8/\kappa)}{\Gamma(1 - 4/\kappa)^2}, \quad (2.1)$$

and where, using the branch choice (1.13) of the integrand  $f_\beta = f_\beta^{(\kappa)}$  defined in (1.12), we have

$$\mathcal{H}_\beta(\mathbf{x}) = \mathcal{H}_\beta^{(\kappa)}(\mathbf{x}) := \oint_{\vartheta_1^\beta} du_1 \oint_{\vartheta_2^\beta} du_2 \cdots \oint_{\vartheta_N^\beta} du_N f_\beta^{(\kappa)}(\mathbf{x}; \mathbf{u}), \quad \mathbf{x} \in \mathfrak{X}_{2N}. \quad (2.2)$$

Choosing  $\alpha_j = 1/\sqrt{\kappa}$  for all  $1 \leq j \leq 2N$  (this is the charge associated to a degenerate CFT field at level two), we see that the integral (2.2) can be written in the Coulomb gas form with screenings  $\alpha_- = -2/\sqrt{\kappa}$ ,

$$\mathcal{H}_\beta(\mathbf{x}) = \oint_{\vartheta_1^\beta \times \dots \times \vartheta_N^\beta} \prod_{1 \leq i < j \leq 2N} (x_j - x_i)^{2\alpha_i \alpha_j} \prod_{1 \leq r < s \leq N} (u_s - u_r)^{2\alpha_-^2} \prod_{\substack{1 \leq i \leq 2N \\ 1 \leq r \leq N}} (u_r - x_i)^{2\alpha_- \alpha_i} du_1 \cdots du_N,$$

where the branch choice is implicit. The conformal weight for a degenerate field at level two is given by  $h_j = h(\kappa) = \frac{6-\kappa}{2\kappa}$ . It gives the Möbius covariance factor as in Proposition 2.2.

Now, the main purpose of this section is to collect properties of  $\mathcal{H}_\beta$ .

- We prove a general Möbius covariance property in Proposition 2.2 in Section 2.1, involving also Möbius transformations that may move the point  $\infty$ .
- We prove a cyclic permutation relation in Proposition 2.6 in Section 2.3. This agrees with the deep relationship of the Coulomb gas integrals with representation theory of the quantum group  $U_q(\mathfrak{sl}(2, \mathbb{C}))$ , where  $q = \exp(4\pi i/\kappa)$ . Indeed, at least when  $\kappa \notin \mathbb{Q}$  so that  $q$  is not a root of unity, the monodromy of the Coulomb gas integrals is encoded into the (universal) R-matrix of  $U_q(\mathfrak{sl}(2, \mathbb{C}))$  [Kas95] acting on a tensor product of its type-one two-dimensional simple modules [FW91, KP20]. Our findings in Proposition 2.6 perfectly match with this structure. Note also that  $\nu(\kappa) = -q - \frac{1}{q}$ .

- We give an alternative formula for  $\mathcal{H}_\beta$  in Proposition 2.5 in Section 2.2, in terms of  $N - 1$  integrals satisfying charge neutrality [DFMS97, FK15c], including the case where the integrand (2.10) equals

$$\begin{aligned} \hat{f}(\mathbf{x}; u_1, \dots, u_{N-1}) := & \prod_{\substack{1 \leq i < j \leq 2N \\ i, j \neq 2N}} (x_j - x_i)^{2\alpha_i \alpha_j} \prod_{1 \leq i < 2N} (x_{2N} - x_i)^{2\alpha_i \alpha_{2N}} \\ & \times \prod_{1 \leq t < s \leq N-1} (u_s - u_t)^{2\alpha_-^2} \prod_{\substack{1 \leq i \leq 2N-1 \\ 1 \leq s \leq N-1}} (u_s - x_i)^{2\alpha_- \alpha_i} \prod_{1 \leq s \leq N-1} (u_s - x_{2N})^{2\alpha_- \alpha_{2N}}, \end{aligned}$$

where  $\alpha_j = 1/\sqrt{\kappa}$  for all  $1 \leq j \leq 2N - 1$  as before, and  $\alpha_{2N} = \frac{\kappa-6}{2\sqrt{\kappa}}$  is the so-called *conjugate charge*<sup>7</sup> of  $1/\sqrt{\kappa}$ , so that the sum of the charges equals the background charge:

$$\left( \sum_{j=1}^{2N-1} \alpha_j \right) + \left( \frac{\kappa-6}{2\sqrt{\kappa}} \right) + (N-1)\alpha_- = 2\alpha_0.$$

This shows in particular that the Coulomb gas integral (1.10) is consistent with the earlier works of Dubédat [Dub06], and Flores & Kleban [FK15c]. Hence, our formulas, which are symmetric in the marked points, indeed agree with those considered in the physics literature, and demonstrates that the charge neutrality assumption, commonly imposed in physics, is not actually needed here.

- We prove a recursive asymptotics property for  $\mathcal{H}_\beta$  in Proposition 2.12 in Section 2.5, from which we obtain the asymptotics for  $\mathcal{F}_\beta$  in the case where  $\kappa \in (0, 8) \setminus \{\frac{8}{m} : m \in \mathbb{Z}_{>0}\}$ , see Corollary 2.13.

## 2.1 General Möbius covariance

In this section, we prove a very general Möbius covariance property for general Coulomb gas integrals.

<sup>7</sup>The charges associated to degenerate CFT fields are commonly labeled as  $\alpha_{r,s} = \frac{1}{2}((1-r)\alpha_+ + (1-s)\alpha_-)$ . The conjugate charge is  $\alpha_{r,s}^* = \frac{1}{2}((1+s)\alpha_+ + (1+r)\alpha_-)$ . In this notation, we have  $\alpha_{1,2} = 1/\sqrt{\kappa}$  and  $\alpha_{1,2}^* = \frac{1}{2}((1+s)\alpha_+ + (1+r)\alpha_-)$ .

**Lemma 2.1.** Fix  $\beta \in \text{LP}_N$  and let  $-\beta \in \text{LP}_N$  be the link pattern obtained from  $\beta$  by reflecting it across the imaginary axis. Then, we have

$$\mathcal{H}_{-\beta}(-x_{2N}, \dots, -x_1) = \mathcal{H}_\beta(x_1, \dots, x_{2N}), \quad x_1 < \dots < x_{2N}. \quad (2.3)$$

*Proof.* Making in  $\mathcal{H}_{-\beta}(-x_1, \dots, -x_{2N})$  the change of variables  $v_r = -u_{N-r}$  for  $1 \leq r \leq N$ , whose Jacobian is 1, and collecting the phase factors relative to our branch choice (1.13) of the integrand  $f_\beta$  defined in (1.12) — to observe that they all cancel out — we readily obtain (2.3).  $\square$

**Proposition 2.2.** Fix  $\beta \in \text{LP}_N$ . The function  $\mathcal{H}_\beta$  satisfies the general Möbius covariance

$$\mathcal{H}_\beta(\mathbf{x}) = \prod_{i=1}^{2N} |\varphi'(x_i)|^{h(\kappa)} \times \mathcal{H}_{\varphi(\beta)}(\varphi(\mathbf{x})), \quad \mathbf{x} \in \mathfrak{X}_{2N}, \quad (2.4)$$

for all Möbius maps  $\varphi \in \text{SL}(2, \mathbb{R})$ , where  $\varphi(\mathbf{x}) := (\varphi(x_{j+1}), \dots, \varphi(x_{2N}), \varphi(x_1), \dots, \varphi(x_j))$ , so that

$$\varphi(x_{j+1}) < \varphi(x_{j+2}) < \dots < \varphi(x_{2N}) < \varphi(x_1) < \varphi(x_2) < \dots < \varphi(x_j)$$

with some  $j \in \{0, 1, 2, \dots, 2N-1\}$ , and where  $\varphi(\beta) \in \text{LP}_N$  is the link pattern obtained from  $\beta$  via permuting the indices according to the permutation of the boundary points induced by  $\varphi$  and then ordering the link endpoints appropriately<sup>8</sup>.

*Proof.* The proof is a generalization of the arguments appearing in [KP20, Proposition 4.15] (for  $\kappa \notin \mathbb{Q}$ ). Recall that the Möbius group  $\text{SL}(2, \mathbb{R})$  is generated by translations  $z \mapsto z + b$ , for  $b \in \mathbb{R}$ , dilations  $z \mapsto az$ , for  $a \in \mathbb{R} \setminus \{0\}$ , and special conformal transformations  $\varphi_c: z \mapsto \frac{z}{1+cz}$ , for  $c \in \mathbb{R}$ . The covariance (2.4) under translations and scalings  $z \mapsto az$ , for  $a > 0$ , is straightforward to check. The covariance (2.4) for dilations  $\delta_a: z \mapsto az$ , for  $a < 0$ , can be checked by considering the  $a$ -variation of the right-hand side of (2.4) with  $\varphi = \delta_a$  in the following manner. Fix  $a_0 \in \mathbb{R} \setminus \{0\}$  and a small  $\varepsilon > 0$ , and consider  $a \in (a_0 - \varepsilon, a_0 + \varepsilon)$ . Note that the integration contours  $\vartheta_1^{\delta_a(\beta)} \times \dots \times \vartheta_N^{\delta_a(\beta)}$  stay in the same homotopy class under a variation by  $\delta_a$  when  $\varepsilon$  is small enough. Hence, the  $a$ -variation of the right-hand side of (2.4) with  $\varphi = \delta_a$  equals

$$\begin{aligned} & \frac{d}{da} \left( \prod_{i=1}^{2N} |\delta'_a(x_i)|^{h(\kappa)} \times \oint_{\vartheta_1^{\delta_a(\beta)} \times \dots \times \vartheta_N^{\delta_a(\beta)}} f_{\delta_a(\beta)}(\delta_a(\mathbf{x}); \mathbf{u}) du_1 \dots du_N \right) \\ &= \frac{1}{a} \prod_{i=1}^{2N} |\delta'_a(x_i)|^{h(\kappa)} \times \oint_{\vartheta_1^{\delta_a(\beta)} \times \dots \times \vartheta_N^{\delta_a(\beta)}} \sum_{j=1}^{2N} (ax_j \partial_j + h(\kappa)) f_{\delta_a(\beta)}(\delta_a(\mathbf{x}); \mathbf{u}) du_1 \dots du_N. \end{aligned} \quad (2.5)$$

A short computation shows that

$$\sum_{j=1}^{2N} \left( y_j \frac{\partial}{\partial y_j} + h(\kappa) \right) f_{\delta_a(\beta)}(\mathbf{y}; \mathbf{u}) = - \sum_{r=1}^N \frac{\partial}{\partial u_r} (u_r f_{\delta_a(\beta)})(\mathbf{y}; \mathbf{u}).$$

Applying this to  $\mathbf{y} = \delta_a(\mathbf{x})$ , we see that (2.5) equals

$$(2.5) = - \frac{1}{a} \prod_{i=1}^{2N} |\delta'_a(x_i)|^{h(\kappa)} \times \oint_{\vartheta_1^{\delta_a(\beta)} \times \dots \times \vartheta_N^{\delta_a(\beta)}} \sum_{r=1}^N \frac{\partial}{\partial u_r} (u_r f_{\delta_a(\beta)})(\delta_a(\mathbf{x}); \mathbf{u}) du_1 \dots du_N,$$

<sup>8</sup>Note that since  $\varphi$  is a diffeomorphism on  $\mathbb{R}$ , for any  $\beta \in \text{LP}_N$ , also  $\varphi(\beta) \in \text{LP}_N$  is a planar link pattern.

which equals zero because each term in the sum vanishes by integration by parts, as the Pochhammer contours are homologically trivial. Therefore, the right-hand side of the asserted formula (2.4) with  $\varphi = \delta_a$  is constant in  $a \in (a_0 - \varepsilon, a_0 + \varepsilon)$  for any  $a_0 \in \mathbb{R} \setminus \{0\}$  and small enough  $\varepsilon > 0$ .

On the one hand, since at  $a_0 = 1$  we have  $\delta_1 = \text{id}$ , this constant equals  $\mathcal{H}_\beta(\mathbf{x})$  for all  $a > 0$  (as we already knew). On the other hand, since at  $a_0 = -1$  we have  $\delta_{-1} = -\text{id}$ , and  $\delta'_{-1}(z) = -1$ , to prove the asserted formula (2.4) for dilations it suffices to invoke Lemma 2.1.

Lastly, to prove the covariance property (2.4) for special conformal transformations  $\varphi_c: z \mapsto \frac{z}{1+cz}$ , for  $c \in \mathbb{R}$ , we use the same type of argument (already presented in [FPW24a, Proof of Proposition 2.2]). Note that  $\varphi'_c(z) > 0$ . We consider the  $c$ -variation of the right-hand side of (2.4) with  $\varphi = \varphi_c$ . Fix  $c_0 \in \mathbb{R}$  and a small  $\varepsilon > 0$ , and consider  $c \in (c_0 - \varepsilon, c_0 + \varepsilon)$ . Note that the integration contours  $\vartheta_1^{\varphi_c(\beta)} \times \cdots \times \vartheta_N^{\varphi_c(\beta)}$  stay in the same homotopy class under a variation by  $\varphi_c$  when  $\varepsilon$  is small enough. Hence, the  $c$ -variation of the right-hand side of (2.4) with  $\varphi = \varphi_c$  equals

$$\begin{aligned} & \frac{d}{dc} \left( \prod_{i=1}^{2N} \varphi'_c(x_i)^{h(\kappa)} \times \oint_{\vartheta_1^{\varphi_c(\beta)} \times \cdots \times \vartheta_N^{\varphi_c(\beta)}} f_{\varphi_c(\beta)}(\varphi_c(\mathbf{x}); \mathbf{u}) du_1 \cdots du_N \right) \\ &= - \prod_{i=1}^{2N} \varphi'_c(x_i)^{h(\kappa)} \times \oint_{\vartheta_1^{\varphi_c(\beta)} \times \cdots \times \vartheta_N^{\varphi_c(\beta)}} \sum_{j=1}^{2N} \left( \varphi_c(x_j)^2 \partial_j + 2h(\kappa) \varphi_c(x_j) \right) f_{\varphi_c(\beta)}(\varphi_c(\mathbf{x}); \mathbf{u}) du_1 \cdots du_N. \end{aligned} \quad (2.6)$$

This can be evaluated by observing (via a long calculation combined with Liouville theorem, as in [KP20, Lemma 4.14]) that the integrand function  $f_{\varphi_c(\beta)}$  defined in (1.12) satisfies the partial differential equation

$$\sum_{j=1}^{2N} \left( y_j^2 \frac{\partial}{\partial y_j} + 2h(\kappa) y_j \right) f(\mathbf{y}; \mathbf{u}) = \sum_{r=1}^N \frac{\partial}{\partial u_r} (g(u_r; \mathbf{y}; \dot{\mathbf{u}}_r) f_{\varphi_c(\beta)}(\mathbf{y}; \mathbf{u})),$$

where  $\dot{\mathbf{u}}_r = (u_1, \dots, u_{r-1}, u_{r+1}, \dots, u_N)$  and  $g$  is a rational function which is symmetric in its last  $N - 1$  variables, and whose only poles are where some of its arguments coincide. Applying this to  $\mathbf{y} = \varphi_c(\mathbf{x})$ , we obtain

$$(2.6) = - \prod_{i=1}^{2N} \varphi'_c(x_i)^{h(\kappa)} \times \oint_{\vartheta_1^{\varphi_c(\beta)} \times \cdots \times \vartheta_N^{\varphi_c(\beta)}} \sum_{r=1}^N \frac{\partial}{\partial u_r} (g(u_r; \varphi_c(\mathbf{x}); \dot{\mathbf{u}}_r) f_{\varphi_c(\beta)}(\varphi_c(\mathbf{x}); \mathbf{u})) du_1 \cdots du_N,$$

which equals zero because each term in the sum vanishes by integration by parts, as the Pochhammer contours are homologically trivial. Therefore, the right-hand side of the asserted formula (2.4) with  $\varphi = \varphi_c$  is constant in  $c \in (c_0 - \varepsilon, c_0 + \varepsilon)$  for any  $c_0 \in \mathbb{R}$  and small enough  $\varepsilon > 0$ . Since at  $c_0 = 0$  we have  $\varphi_0 = \text{id}$ , this constant equals  $\mathcal{H}_\beta(\mathbf{x})$ . This concludes the proof for all Möbius maps  $\varphi \in \text{SL}(2, \mathbb{R})$ .  $\square$

**Corollary 2.3.** *Fix  $\beta \in \text{LP}_N$ . The function  $\mathcal{F}_\beta$  satisfies the Möbius covariance*

$$\mathcal{F}_\beta(\mathbb{H}; \mathbf{x}) = \prod_{i=1}^{2N} \varphi'(x_i)^{h(\kappa)} \times \mathcal{F}_{\varphi(\beta)}(\mathbb{H}; \varphi(\mathbf{x})) = \prod_{i=1}^{2N} |\varphi'(x_i)|^{h(\kappa)} \times \mathcal{F}_{\varphi(\beta)}(\mathbb{H}; \varphi(\mathbf{x})), \quad \mathbf{x} \in \mathfrak{X}_{2N},$$

for all Möbius maps  $\varphi \in \text{PSL}(2, \mathbb{R})$  preserving the upper half-plane, where  $\varphi'(x_i) > 0$  for all  $1 \leq i \leq 2N$ .

*Proof.* For  $\kappa \in (0, 8) \setminus \{\frac{8}{m} : m \in \mathbb{Z}_{>0}\}$ , this is immediate from Proposition 2.2, since  $\mathcal{F}_\beta$  is a constant multiple of  $\mathcal{H}_\beta$ . For exceptional  $\kappa \in \{\frac{8}{m} : m \in \mathbb{Z}_{>0}\}$ , the claim then follows by continuity in  $\kappa \in \mathbb{C} \setminus \{0\}$ .  $\square$



**Remark 2.4.** We extend the definition of  $\mathcal{F}_\beta(\Omega; \mathbf{x})$  to more general (nice<sup>9</sup>) polygons  $(\Omega; \mathbf{x})$  by sending  $\Omega$  onto  $\mathbb{H}$  via any conformal map  $\varphi$  with  $\varphi(x_1) < \dots < \varphi(x_{2N})$ , using the conformal covariance

$$\mathcal{F}_\beta(\Omega; \mathbf{x}) := \prod_{j=1}^{2N} |\varphi'(x_j)|^{h(\kappa)} \times \mathcal{F}_\beta(\Omega; \varphi(\mathbf{x})) \quad (2.7)$$

Thanks to Corollary 2.3, this definition is independent of the choice of  $\varphi$ . Proposition 2.2 is also consistent:

$$\mathcal{F}_\beta(\mathbb{H}; \mathbf{x}) = \prod_{i=1}^{2N} |\varphi'(x_i)|^{h(\kappa)} \times \mathcal{F}_\beta(\mathbb{H}^*; \tilde{\varphi}(\mathbf{x})), \quad \mathbf{x} \in \mathfrak{X}_{2N},$$

for all Möbius maps  $\varphi: \mathbb{H} \rightarrow \mathbb{H}^* := \{z \in \mathbb{C} : \text{Im}(z) < 0\}$ , where  $\tilde{\varphi}(\mathbf{x}) := (\varphi(x_1), \dots, \varphi(x_{2N}))$ .

## 2.2 Integral removals

We next verify that our Coulomb gas integral formula (1.10) is consistent with the earlier works of Julien Dubédat [Dub06], and Steven Flores & Peter Kleban [FK15c], involving one less integration, in the spirit of the physics literature where the charge neutrality assumption is imperative. Our original formulas have the virtue that they do not (need to) satisfy any charge neutrality, and they are completely symmetric in the marked points. However, the alternative formulas in Proposition 2.5 are very useful in computations.

**Proposition 2.5.** Fix  $\beta \in \text{LP}_N$  and  $r \in \{1, 2, \dots, N\}$ . For  $\kappa \in (0, 8) \setminus \{\frac{8}{m} : m \in \mathbb{Z}_{>0}\}$ , we have

$$\mathcal{H}_\beta(\mathbf{x}) = \frac{\nu(\kappa)}{C(\kappa)} \oint_{\vartheta_1^\beta} du_1 \cdots \oint_{\vartheta_{r-1}^\beta} du_{r-1} \oint_{\vartheta_{r+1}^\beta} du_{r+1} \cdots \oint_{\vartheta_N^\beta} du_N \hat{f}_\beta^b(\mathbf{x}; \dot{\mathbf{u}}_r) \quad (2.8)$$

$$= \frac{\nu(\kappa)}{C(\kappa)} \oint_{\vartheta_1^\beta} du_1 \cdots \oint_{\vartheta_{r-1}^\beta} du_{r-1} \oint_{\vartheta_{r+1}^\beta} du_{r+1} \cdots \oint_{\vartheta_N^\beta} du_N \hat{f}_\beta^a(\mathbf{x}; \dot{\mathbf{u}}_r), \quad \mathbf{x} \in \mathfrak{X}_{2N}, \quad (2.9)$$

where  $\dot{\mathbf{u}}_r = (u_1, \dots, u_{r-1}, u_{r+1}, \dots, u_N)$  and the branches of the multivalued integrands

$$\begin{aligned} \hat{f}_\beta^b(\mathbf{x}; \dot{\mathbf{u}}_r) &:= \prod_{\substack{1 \leq i < j \leq 2N \\ i, j \neq b_r}} (x_j - x_i)^{2/\kappa} \prod_{1 \leq i < b_r} (x_{b_r} - x_i)^{-2h(\kappa)} \prod_{b_r < j \leq 2N} (x_j - x_{b_r})^{-2h(\kappa)} \\ &\times \prod_{\substack{1 \leq t < s \leq N \\ t, s \neq r}} (u_s - u_t)^{8/\kappa} \prod_{\substack{1 \leq i < 2N \\ i \neq b_r \\ 1 \leq s \leq N \\ s \neq r}} (u_s - x_i)^{-4/\kappa} \prod_{\substack{1 \leq s \leq N \\ s \neq r}} (u_s - x_{b_r})^{4h(\kappa)}, \end{aligned} \quad (2.10)$$

which carries the conjugate charge at  $x_{b_r}$ , and

$$\begin{aligned} \hat{f}_\beta^a(\mathbf{x}; \dot{\mathbf{u}}_r) &:= \prod_{\substack{1 \leq i < j \leq 2N \\ i, j \neq a_r}} (x_j - x_i)^{2/\kappa} \prod_{1 \leq i < a_r} (x_{a_r} - x_i)^{-2h(\kappa)} \prod_{a_r < j \leq 2N} (x_j - x_{a_r})^{-2h(\kappa)} \\ &\times \prod_{\substack{1 \leq t < s \leq N \\ t, s \neq r}} (u_s - u_t)^{8/\kappa} \prod_{\substack{1 \leq i < 2N \\ i \neq a_r \\ 1 \leq s \leq N \\ s \neq r}} (u_s - x_i)^{-4/\kappa} \prod_{\substack{1 \leq s \leq N \\ s \neq r}} (u_s - x_{a_r})^{4h(\kappa)}, \end{aligned} \quad (2.11)$$

which carries the conjugate charge at  $x_{a_r}$ , are chosen to be real and positive when  $x_{a_s} < \text{Re}(u_s) < x_{a_s+1}$  for all  $s \in \{1, 2, \dots, N\} \setminus \{r\}$ . (Here,  $h(\kappa) = \frac{6-\kappa}{2\kappa}$ .)

<sup>9</sup>By a *nice polygon*, we refer to a polygon such that the marked boundary points  $x_1, \dots, x_{2N}$  lie on sufficiently regular boundary segments for the derivatives of  $\varphi$  to be defined there, e.g.  $C^{1+\epsilon}$  for some  $\epsilon > 0$ .

Note that the omitted integral is associated to a variable carrying the conjugate charge. This is important in order to preserve the desired asymptotics properties (Proposition 2.12).

*Proof.* It suffices to show (2.8) when  $b_r = 2N$  and (2.9) when  $a_r = 1$ , because for the other cases the claim then follows after conjugating by a suitable Möbius map and using Proposition 2.2.

Fix  $\mathbf{x} = (x_1, \dots, x_{2N}) \in \mathfrak{X}_{2N}$ . Suppose  $b_r = 2N$  (in this case,  $a_r = 2r - 1$ ) and denote by  $\hat{\mathcal{H}}_\beta$  the right-hand side of (2.8). Note that if  $\mathcal{H}_\beta(\mathbf{x}) = \hat{\mathcal{H}}_\beta(\mathbf{x}) = 0$ , then (2.8) is clear, and otherwise we may assume without loss of generality that  $\mathcal{H}_\beta(\mathbf{x}) \neq 0$ , and consider the ratio  $\hat{\mathcal{H}}_\beta(\mathbf{x})/\mathcal{H}_\beta(\mathbf{x})$  of the right and left-hand sides. For large  $R > 0$ , let  $\varphi_R$  be the Möbius map of the upper half-plane  $\mathbb{H}$  such that

$$\begin{cases} -\infty < \varphi_R(x_1) < \varphi_R(x_2) < \dots < \varphi_R(x_{2N}) < \infty, \\ \varphi_R(x_1) = x_1, & \varphi_R(x_{2N-1}) = x_{2N-1}, & \varphi_R(x_{2N}) = R. \end{cases}$$

Denote  $\varphi_R(\mathbf{x}) := (\varphi_R(x_1), \dots, \varphi_R(x_{2N}))$  and  $\hat{x}_j := \lim_{R \rightarrow \infty} \varphi_R(x_j)$  for  $1 \leq j \leq 2N - 1$ . Note that

$$-\infty < x_1 = \hat{x}_1 < \hat{x}_2 < \dots < \hat{x}_{2N-1} = x_{2N-1} < R < \infty.$$

Because both  $\mathcal{H}_\beta$  and  $\hat{\mathcal{H}}_\beta$  are Möbius covariant (respectively by Proposition 2.2 and [Dub06, Lemma 2]), we may compute their ratio in the form

$$\begin{aligned} \frac{\hat{\mathcal{H}}_\beta(\mathbf{x})}{\mathcal{H}_\beta(\mathbf{x})} &= \lim_{R \rightarrow \infty} \frac{\hat{\mathcal{H}}_\beta(\varphi_R(\mathbf{x}))}{\mathcal{H}_\beta(\varphi_R(\mathbf{x}))} \\ &= \frac{\nu(\kappa)}{C(\kappa)} \lim_{R \rightarrow \infty} \frac{R^{2h(\kappa)} \oint_{\hat{\vartheta}_1^\beta} du_1 \cdots \oint_{\hat{\vartheta}_{r-1}^\beta} du_{r-1} \oint_{\hat{\vartheta}_{r+1}^\beta} du_{r+1} \cdots \oint_{\hat{\vartheta}_N^\beta} du_N \mathring{f}_\beta^b(\varphi_R(\mathbf{x}); \hat{\mathbf{u}}_r)}{R^{2h(\kappa)} \oint_{\hat{\vartheta}_1^\beta} du_1 \cdots \oint_{\hat{\vartheta}_{r-1}^\beta} du_{r-1} \oint_{\hat{\vartheta}_{r+1}^\beta} du_{r+1} \cdots \oint_{\hat{\vartheta}_N^\beta} du_N \mathring{f}_{\vartheta(\varphi_R(x_{a_r}), R)} du_r f_\beta(\varphi_R(\mathbf{x}); \mathbf{u})}. \end{aligned} \quad (2.12)$$

where the Pochhammer contours  $\hat{\vartheta}_1^\beta, \dots, \hat{\vartheta}_{r-1}^\beta, \hat{\vartheta}_{r+1}^\beta, \dots, \hat{\vartheta}_N^\beta$  are all disjoint, and each  $\hat{\vartheta}_s^\beta$  surrounds the points  $\hat{x}_{a_s}, \hat{x}_{b_s}$  and does not surround any other points among  $\{\hat{x}_1, \dots, \hat{x}_{r-1}, \hat{x}_{r+1}, \dots, \hat{x}_{2N-1}\}$ . Now, after collecting the powers  $0 = 2h(\kappa) - 2h(\kappa) + 4(N-1)h(\kappa) - 4(N-1)h(\kappa)$  of the blow-up parameter  $R$ , the limit of the numerator equals

$$\oint_{\hat{\vartheta}_1^\beta} du_1 \cdots \oint_{\hat{\vartheta}_{r-1}^\beta} du_{r-1} \oint_{\hat{\vartheta}_{r+1}^\beta} du_{r+1} \cdots \oint_{\hat{\vartheta}_N^\beta} du_N \mathring{f}_\beta(\hat{x}_1, \dots, \hat{x}_{2N-1}; \hat{\mathbf{u}}_r), \quad (2.13)$$

where  $\mathring{f}_\beta$  is the integrand function (1.33) with  $d = 2N - 1$  and  $\ell = N - 1$ , and branch chosen to be real and positive when  $\hat{x}_{a_s} < \text{Re}(u_s) < x_{a_s+1}$  for all  $s \neq r$ . For the denominator, using the change of variables  $v := \frac{u_r - \varphi_R(x_{a_r})}{R - \varphi_R(x_{a_r})}$  and Eq. (B.1) from Lemma B.1 to evaluate the  $r$ :th integral as

$$\oint_{\vartheta(0,1)} v^{-4/\kappa} (v-1)^{-4/\kappa} dv = \frac{\nu(\kappa)}{C(\kappa)}, \quad \text{for } \kappa \in (0, 8) \setminus \left\{ \frac{8}{m} : m \in \mathbb{Z}_{>0} \right\}, \quad (2.14)$$

we obtain

$$\begin{aligned} &\lim_{R \rightarrow \infty} R^{2h(\kappa)} \oint_{\hat{\vartheta}_1^\beta} du_1 \cdots \oint_{\hat{\vartheta}_{r-1}^\beta} du_{r-1} \oint_{\hat{\vartheta}_{r+1}^\beta} du_{r+1} \cdots \oint_{\hat{\vartheta}_N^\beta} du_N \oint_{\vartheta(\varphi_R(x_{a_r}), R)} du_r \mathring{f}_\beta(\varphi_R(\mathbf{x}); \mathbf{u}) \\ &= \left( \oint_{\vartheta(0,1)} v^{-4/\kappa} (v-1)^{-4/\kappa} dv \right) \oint_{\hat{\vartheta}_1^\beta} du_1 \cdots \oint_{\hat{\vartheta}_{r-1}^\beta} du_{r-1} \oint_{\hat{\vartheta}_{r+1}^\beta} du_{r+1} \cdots \oint_{\hat{\vartheta}_N^\beta} du_N \mathring{f}_\beta(\hat{x}_1, \dots, \hat{x}_{2N-1}; \hat{\mathbf{u}}_r) \\ &= \frac{\nu(\kappa)}{C(\kappa)} \oint_{\hat{\vartheta}_1^\beta} du_1 \cdots \oint_{\hat{\vartheta}_{r-1}^\beta} du_{r-1} \oint_{\hat{\vartheta}_{r+1}^\beta} du_{r+1} \cdots \oint_{\hat{\vartheta}_N^\beta} du_N \mathring{f}_\beta(\hat{x}_1, \dots, \hat{x}_{2N-1}; \hat{\mathbf{u}}_r). \end{aligned} \quad (2.15)$$

Plugging (2.13, 2.15) into (2.12), we obtain (2.8):

$$\frac{\hat{\mathcal{H}}_\beta(\mathbf{x})}{\mathcal{H}_\beta(\mathbf{x})} = \lim_{R \rightarrow \infty} \frac{\hat{\mathcal{H}}_\beta(\varphi_R(\mathbf{x}))}{\mathcal{H}_\beta(\varphi_R(\mathbf{x}))} = 1.$$

Similarly, the second asserted identity (2.9) when  $a_r = 1$  can be proven by sending  $x_1$  to  $-\infty$ .  $\square$

### 2.3 Cyclic permutation symmetry

We next record a simple property of  $\mathcal{H}_\beta$  when its variables are cyclically permuted within  $\mathfrak{Y}_{2N}$ . To state it, we denote by  $\sigma = \begin{pmatrix} 1 & 2 & 3 & \dots & 2N-1 & 2N \\ 2 & 3 & 4 & \dots & 2N & 1 \end{pmatrix}$  the cyclic counterclockwise permutation of the indices, and for each  $\beta \in \text{LP}_N$ , we denote by  $\sigma(\beta) \in \text{LP}_N$  the link pattern obtained from  $\beta$  via permuting the indices by  $\sigma$  and then ordering the link endpoints appropriately. For example, with  $N = 2$  we have

$$\begin{aligned} \beta = \text{---} \overbrace{\text{---}} \text{---} &= \{\{1, 2\}, \{3, 4\}\} & \mapsto & \sigma(\beta) = \text{---} \overbrace{\text{---}} \text{---} &= \{\{1, 4\}, \{2, 3\}\} \\ \beta = \text{---} \overbrace{\text{---}} \text{---} &= \{\{1, 4\}, \{2, 3\}\} & \mapsto & \sigma(\beta) = \text{---} \overbrace{\text{---}} \text{---} &= \{\{1, 2\}, \{3, 4\}\}. \end{aligned}$$

Note that this cannot be realized as an action by a Möbius transformation, unless there are only two (or three) points. Indeed, it is not possible to fix the images of more than three points under a Möbius transformation. Rather, the permutation should be thought of as an action of a *mapping class group* (of the upper half-plane with marked points on the boundary, say) or of a suitable subgroup<sup>10</sup> of the *braid group* (that is, the fundamental group of the punctured Riemann sphere). Also, this cannot be realized as an action by the group of cyclic permutations, unless  $q = \exp(4\pi i/\kappa)$  is a third root of unity, because according to Proposition 2.6, permuting  $2N$  times results in a multiplicative factor  $(-\exp(-6\pi i/\kappa))^{2N}$ .

**Proposition 2.6.** *Fix  $\beta \in \text{LP}_N$ . Let  $\rho: [0, 1] \rightarrow \mathfrak{Y}_{2N}$  be a path from  $\rho(0) = \mathbf{x} = (x_1, x_2, \dots, x_{2N-1}, x_{2N}) \in \mathfrak{X}_{2N}$  to  $\rho(1) := (x_2, x_3, \dots, x_{2N-1}, x_{2N}, x_1) \in \mathfrak{Y}_{2N}$  such that  $\rho = (\rho_1, \rho_2, \dots, \rho_{2N})$  satisfy*

$$\begin{cases} \text{Im}(\rho_j(t)) = 0, & \text{for all } j \in \{1, 2, \dots, 2N-1\}, \\ \text{Im}(\rho_j(t)) \geq 0, & \text{for } j = 2N, \end{cases} \quad \text{for all } t \in [0, 1]$$

see also Figure 2.1. Then, we have

$$\mathcal{H}_{\sigma(\beta)}(\rho(1)) = -e^{-\frac{6\pi i}{\kappa}} \mathcal{H}_\beta(\rho(0)) = -q^{3/2} \mathcal{H}_\beta(\rho(0)), \quad q = \exp(4\pi i/\kappa). \quad (2.16)$$




Figure 2.1: Illustration of the braid transformation  $\rho$  in Proposition 2.6.

*Proof.* Note that the path  $\rho$  transforms the integration contours associated to  $\beta$  in  $\mathcal{H}_\beta$  into the integration contours associated to  $\sigma(\beta)$  in  $\mathcal{H}_\beta$ . Hence, it is clear that (2.16) holds up to some multiplicative phase factor. To find it, we simplify  $\mathcal{H}_\beta$  using Proposition 2.5: choosing  $r$  in (2.8) such that  $b_r = 2N$  (in this case,  $a_r = 2r - 1$ ), we see that  $\rho$  gives rise to the following phase factors:

- the factor  $\prod_{1 \leq i \leq 2N-1} (x_{2N} - x_i)^{1-6/\kappa}$  in (2.10) gets a phase  $(-1)^{2N-1} \exp(-\frac{6\pi i}{\kappa}(2N-1))$ ;
- and the factor  $\prod_{\substack{1 \leq s \leq N \\ s \neq r}} (u_s - x_{2N})^{12/\kappa-2}$  in (2.10) gets a phase  $\exp(\frac{12\pi i}{\kappa}(N-1))$ .

<sup>10</sup>Cyclic permutations form a subgroup of the symmetric group and the latter is a quotient of the braid group.

The asserted equality (2.16) follows by collecting the overall phase factor.  $\square$

**Remark 2.7.** Let  $M_d$  denote the  $d$ -dimensional type-one simple module of the Hopf algebra  $U_q(\mathfrak{sl}(2, \mathbb{C}))$  (see, e.g., [Kas95] for definitions). A space of SLE $_{\kappa}$  partition functions and Coulomb gas integral solutions to the BPZ PDEs (PDE) was shown in [FW91, KP16, KP20] in the generic case to coincide with the set of highest-weight vectors in the tensor product module  $M_2^{\otimes 2N}$ . The solutions satisfying the full Möbius covariance belong to the trivial submodule  $(\frac{1}{N+1} \binom{2N}{N}) M_1$  of this tensor product module, whose multiplicity is the  $N$ :th Catalan number. The (inverse) R-matrix of  $U_q(\mathfrak{sl}(2, \mathbb{C}))$  represents monodromy transformations that braid the variables  $\mathbf{x} = (x_1, \dots, x_{2N})$  in the Riemann sphere. It acts on the tensor product  $M_2 \otimes M_2 \cong M_1 \oplus M_3$  in the block diagonal form

$$= \begin{pmatrix} -q^{3/2} & 0 & 0 & 0 \\ 0 & q^{-1/2} & 0 & 0 \\ 0 & 0 & q^{-1/2} & 0 \\ 0 & 0 & 0 & q^{-1/2} \end{pmatrix},$$

and its eigenvalue on  $M_1$  equals  $-q^{3/2}$ . Note that (2.16) in Proposition 2.6 is consistent with this.

## 2.4 Contour deformations

We now consider Coulomb gas integrals with general numbers of  $d \geq 2$  variables and  $\ell \geq 0$  screening variables (with any branch choice),

$$\int_{\Gamma} \dots \int \underbrace{\prod_{1 \leq i < j \leq d} (x_j - x_i)^{2\alpha_i \alpha_j} \prod_{1 \leq r < s \leq \ell} (u_s - u_r)^{2\alpha^2} \prod_{\substack{1 \leq i \leq d \\ 1 \leq r \leq \ell}} (u_r - x_i)^{2\alpha - \alpha_i}}_{=: f(\mathbf{x}; \mathbf{u})} du_1 \dots du_{\ell}, \quad (2.17)$$

with Coulomb charges  $\alpha_j = \alpha_j(\kappa) \in \mathbb{R}$  at the variables  $\mathbf{x} = (x_1, x_2, \dots, x_d)$ , and negative screening charges  $\alpha_- = -2/\sqrt{\kappa}$  at the integration variables  $\mathbf{u} = (u_1, u_2, \dots, u_{\ell})$ . We assume that the integration contours in (2.17) have the form  $\Gamma = \Gamma_1 \times \Gamma_2 \times \dots \times \Gamma_{\ell}$  such that  $\Gamma_r \cap \Gamma_s = \emptyset$  for all  $r \neq s$ . For example, they could be Pochhammer contours<sup>11</sup>  $\vartheta(x_a, x_b)$  as in (1.14), line integrals, or simple loops surrounding some of the marked points. Specifically, we let  $\varrho(x_a, x_b)$  denote a clockwise oriented simple loop on  $\mathbb{C} \setminus \{x_1, \dots, x_d\}$  surrounding two points  $x_a$  and  $x_b$  and none of the other points among  $\{x_1, \dots, x_d\}$ . (A limiting case of such a loop as  $x_b \rightarrow +\infty$  is illustrated in Eq. (B.7) in Lemma B.2.) We also denote by  $\partial B(x, \epsilon)$  the clockwise oriented circle of radius  $\epsilon$  around a point  $x$ .

The next simple but useful lemma shows how to transform an integration around a Pochhammer contour into an integration involving line integrals between the marked points together with integrations around two small loops surrounding these points. To avoid any ambiguity, by the integration symbol  $\int_{x+\epsilon}^{y-\epsilon} du$ , we indicate that the integration is performed from  $(x + \epsilon)$  to  $(y - \epsilon)$  in the upper half-plane.

**Lemma 2.8.** *Fix  $r \in \{1, 2, \dots, \ell\}$  such that  $\Gamma_r = \vartheta(x_a, x_b)$  for some  $a, b \in \{1, 2, \dots, d\}$ ,  $a \neq b$ . Then,*

$$\begin{aligned} & \int_{\Gamma_1} du_1 \dots \int_{\Gamma_{\ell}} du_{\ell} f(\mathbf{x}; \mathbf{u}) \\ &= \int_{\Gamma_1} du_1 \dots \int_{\Gamma_{r-1}} du_{r-1} \left( (1 - e^{-4\pi i \alpha - \alpha_b})(1 - e^{4\pi i \alpha - \alpha_a}) \int_{x_a + \epsilon}^{x_b - \epsilon} du_r \right. \\ & \quad \left. + (1 - e^{-4\pi i \alpha - \alpha_b}) \oint_{\partial B(x_a, \epsilon)} du_r + (1 - e^{4\pi i \alpha - \alpha_a}) \oint_{\partial B(x_b, \epsilon)} du_r \right) \int_{\Gamma_{r+1}} du_{r+1} \dots \int_{\Gamma_{\ell}} du_{\ell} f(\mathbf{x}; \mathbf{u}), \end{aligned}$$

for all  $\epsilon > 0$  small enough.

<sup>11</sup>Throughout, we always assume that  $\vartheta(x_a, x_b)$  does not surround any other marked point among  $\{x_1, \dots, x_d\}$ .

*Proof.* The claim follows in a straightforward manner by contour deformation, collecting the phase factors of the integrand in (2.17) around the points  $x_a$  and  $x_b$ .  $\square$




Figure 2.2: Illustration of Corollary 2.9: when  $\kappa = \frac{8}{2m-1}$  for  $m \in \mathbb{Z}_{>0}$ , the integral of  $f(\mathbf{x}; \mathbf{u})$  along the Pochhammer contour  $\vartheta(x_{a_r}, x_{b_r})$  is the same as 2 times its integral along the simple clockwise loop  $\varrho(x_{a_r}, x_{b_r})$ .

**Corollary 2.9.** Fix  $\kappa \in \{\frac{8}{2m-1} : m \in \mathbb{Z}_{>0}\}$ ,  $\beta \in \text{LP}_N$ , and  $\mathbf{x} \in \mathfrak{X}_{2N}$ . Suppose that the charges in (2.17) are  $\alpha_j = 1/\sqrt{\kappa}$  for all  $j \in \{1, 2, \dots, d\}$ . Fix  $r \in \{1, 2, \dots, \ell\}$  such that  $\Gamma_r = \vartheta(x_{a_r}, x_{b_r})$ . Then, we have

$$\begin{aligned} & \int_{\Gamma_1} du_1 \cdots \int_{\Gamma_{r-1}} du_{r-1} \oint_{\vartheta(x_{a_r}, x_{b_r})} du_r \int_{\Gamma_{r+1}} du_{r+1} \cdots \int_{\Gamma_\ell} du_\ell f(\mathbf{x}; \mathbf{u}) \\ &= 2 \int_{\Gamma_1} du_1 \cdots \int_{\Gamma_{r-1}} du_{r-1} \oint_{\varrho(x_{a_r}, x_{b_r})} du_r \int_{\Gamma_{r+1}} du_{r+1} \cdots \int_{\Gamma_\ell} du_\ell f(\mathbf{x}; \mathbf{u}). \end{aligned}$$

*Proof.* Similarly as when proving Lemma 2.8, a straightforward contour deformation shows that

$$\begin{aligned} \oint_{\varrho(x_{a_r}, x_{b_r})} du_r f(\mathbf{x}; \mathbf{u}) &= (1 - e^{-4\pi i \alpha_- \alpha_{b_r}}) \int_{x_{a_r} + \epsilon}^{x_{b_r} - \epsilon} du_r f(\mathbf{x}; \mathbf{u}) \\ &\quad + \oint_{\partial B(x_{a_r}, \epsilon)} du_r f(\mathbf{x}; \mathbf{u}) + \oint_{\partial B(x_{b_r}, \epsilon)} du_r f(\mathbf{x}; \mathbf{u}). \end{aligned}$$

The claim now follows from Lemma 2.8, after noting that where  $1 - e^{-4\pi i \alpha_- \alpha_{b_r}} = 2 = 1 - e^{4\pi i \alpha_- \alpha_{a_r}}$  due to the choices  $\alpha_{a_r} = \alpha_{b_r} = 1/\sqrt{\kappa}$  and  $\alpha_- = -2/\sqrt{\kappa}$ , since  $\kappa = 8/m$  with  $m \geq 1$  odd.  $\square$

We next record a very useful general property of Coulomb gas integrals of type (2.17) with  $\ell \in \{2, 3, \dots, d\}$  and  $\kappa \in \{\frac{8}{2m-1} : m \in \mathbb{Z}_{>0}\}$  and integration contours  $\Gamma = \varrho_1 \times \varrho_2 \times \cdots \times \varrho_\ell$ , where  $\varrho_1, \varrho_2, \dots, \varrho_\ell$  are clockwise<sup>12</sup> oriented simple mutually non-intersecting loops on  $\mathbb{C} \setminus \mathcal{C}$ , where  $\mathcal{C} = \gamma_1 \times \gamma_2 \times \cdots \times \gamma_{d'}$  are a choice of branch cuts for the multivalued integrand in (2.17) (with  $d' = d/2$  if  $d$  is even, and  $d' = (d+1)/2$  if  $d$  is odd). Namely, for any fixed  $s \in \{1, 2, \dots, \ell\}$ , we can replace the integration along  $\varrho_s$  by an integration along another simple loop  $\hat{\varrho}_s$  obtained from  $\varrho_s$  by pulling it over some of the other loops in  $\Gamma$  (as specified in Lemma 2.10 below). This property hold regardless of the branch choice for the integrand  $f$  defined in (2.17), as long as  $\kappa \in \{\frac{8}{2m-1} : m \in \mathbb{Z}_{>0}\}$ . The setup is illustrated in Figure 2.3. This is a generalization of [LPW24, Lemma 2.12] which deals with the case  $\kappa = 8$ .

**Lemma 2.10.** Fix  $\kappa \in \{\frac{8}{2m-1} : m \in \mathbb{Z}_{>0}\}$ . Fix  $\ell \in \{2, 3, \dots, d\}$  and  $s \in \{1, 2, \dots, \ell\}$ . Let  $\hat{\varrho}_s$  be a clockwise oriented simple loop on  $\mathbb{C} \setminus \gamma_1 \times \gamma_2 \times \cdots \times \gamma_{d'}$  obtained from  $\varrho_s$  by pulling  $\varrho_s$  over some of the other loops in  $\{\varrho_1, \varrho_2, \dots, \varrho_\ell\} \setminus \{\varrho_s\}$ :

$$\hat{\varrho}_s = \varrho_s + \sum_{r \in I_s} \varrho_r, \tag{2.18}$$

<sup>12</sup>One could also orient them counterclockwise — what is important is that all loops have the same orientation.




Figure 2.3: Illustration of Lemma 2.10: when  $\kappa = \frac{8}{2m-1}$  for  $m \in \mathbb{Z}_{>0}$ , the integral of  $f(\mathbf{x}; \mathbf{u})$  along the simple clockwise loop  $\varrho_r$  is the same as its integral along a loop  $\hat{\varrho}_r$  possibly surrounding more marked points.

where  $I_s \subset \{1, 2, \dots, \ell\} \setminus \{s\}$ . Then, writing  $\mathbf{u} = (u_1, \dots, u_\ell)$ , we have

$$\begin{aligned} & \oint_{\varrho_1} du_1 \oint_{\varrho_2} du_2 \cdots \oint_{\varrho_\ell} du_\ell f(\mathbf{x}; \mathbf{u}) \\ &= \oint_{\varrho_1} du_1 \cdots \oint_{\varrho_{s-1}} du_{s-1} \oint_{\hat{\varrho}_s} du_s \oint_{\varrho_{s+1}} du_{s+1} \cdots \oint_{\varrho_\ell} du_\ell f(\mathbf{x}; \mathbf{u}), \end{aligned} \quad (2.19)$$

where  $f$  is the integrand defined in (2.17).

Note that the replacement in Lemma 2.10 is only valid when we integrate  $f(\mathbf{x}; u_1, \dots, u_\ell)$  along all  $\ell$  loops  $\varrho_1, \varrho_2, \dots, \varrho_\ell$ , in which case we can use antisymmetry to get cancellations.

*Proof.* Note that since  $\kappa \in \{\frac{8}{2m-1} : m \in \mathbb{Z}_{>0}\}$ , the integrand (2.17) reads

$$f(\mathbf{x}; \mathbf{u}) = \prod_{1 \leq i < j \leq d} (x_j - x_i)^{m/4} \prod_{1 \leq r < s \leq \ell} (u_s - u_r)^m \prod_{\substack{1 \leq i \leq d \\ 1 \leq r \leq \ell}} (u_r - x_i)^{-m/2}. \quad (2.20)$$

After deforming and decomposing the loop  $\hat{\varrho}_s$  into the linear combination (2.18), we see by antisymmetry of the integrand (2.20) (since  $m$  is odd) and Fubini's theorem that only  $\varrho_s$  can give a non-zero contribution on the right-hand side of (2.19): indeed, for any  $r \neq s$ , the double-integral of (2.20) along  $\varrho_r$  vanishes (here, we use the assumption that  $\ell \geq 2$ ),

$$\oint_{\varrho_r} du_r \oint_{\varrho_r} du_s f(\mathbf{x}; u_1, \dots, u_\ell) = 0,$$

by antisymmetry of the integrand (2.20) with respect to the exchange  $u_r \leftrightarrow u_s$ .  $\square$

## 2.5 Asymptotics properties

It was proved in the series [FK15a, FK15b, FK15c, FK15d] of articles that when  $\kappa \in (0, 8)$ , the space

$$\mathcal{S}_N := \{F: \mathfrak{X}_{2N} \rightarrow \mathbb{C} : F \text{ satisfies (PDE), (COV), and (PLB)}\} \quad (2.21)$$

has dimension  $\dim \mathcal{S}_N = \frac{1}{N+1} \binom{2N}{N}$ , the  $N$ :th Catalan number. One can uniquely specify functions in this space in terms of their asymptotic behavior as two of the variables in  $\mathfrak{X}_{2N}$  tend together.

**Lemma 2.11.** [FK15b, Lemma 1] *Fix  $\kappa \in (0, 8)$ . Let  $F: \mathfrak{X}_{2N} \rightarrow \mathbb{C}$  be a function satisfying (PDE), (COV), and (PLB). If  $F$  also has the asymptotics property*

$$\lim_{x_j, x_{j+1} \rightarrow \xi} \frac{F(x_1, \dots, x_{2N})}{(x_{j+1} - x_j)^{-2h(\kappa)}} = 0, \quad \text{for all } j \in \{1, 2, 3, \dots, 2N-1\} \text{ and } \xi \in (x_{j-1}, x_{j+2})$$

(with the convention that  $x_0 = -\infty$  and  $x_{2N+1} = \infty$ ), then  $F \equiv 0$ .

This is a version of an operator product expansion (OPE) for the corresponding CFT correlation functions, important for identifying solutions of interest to problems in random geometry, such as the SLE pure partition functions and interface crossing probabilities [BBK05, KP16, Pel19, PW23, FPW24a]. In this section, we consider asymptotics of the Coulomb gas integral solutions in  $\mathcal{S}_N$ .

**Proposition 2.12.** *Fix  $\kappa \in (0, 8) \setminus \{\frac{8}{m} : m \in \mathbb{Z}_{>0}\}$ . With  $\mathcal{H}_\emptyset \equiv 1$  for the empty link pattern  $\emptyset \in \text{LP}_0$ , the collection  $\{\mathcal{H}_\beta : \beta \in \text{LP}_N\}$  satisfies the following recursive asymptotics property. Fix  $N \geq 1$  and  $j \in \{1, 2, \dots, 2N - 1\}$ . Then, for all  $\xi \in (x_{j-1}, x_{j+2})$ , using the notation (1.21),*

$$\lim_{x_j, x_{j+1} \rightarrow \xi} \frac{\mathcal{H}_\beta(\mathbf{x})}{(x_{j+1} - x_j)^{-2h(\kappa)}} = \frac{1}{C(\kappa)} \begin{cases} \nu(\kappa) \mathcal{H}_{\beta/\{j, j+1\}}(\ddot{\mathbf{x}}_j), & \text{if } \{j, j+1\} \in \beta, \\ \mathcal{H}_{\wp_j(\beta)/\{j, j+1\}}(\ddot{\mathbf{x}}_j), & \text{if } \{j, j+1\} \notin \beta, \end{cases} \quad (2.22)$$

where  $\beta/\{j, j+1\} \in \text{LP}_{N-1}$  denotes the link pattern obtained from  $\beta$  by removing the link  $\{j, j+1\}$  and relabeling the remaining indices by  $1, 2, \dots, 2N - 2$ , and  $\wp_j$  is the ‘‘tying operation’’ defined by

$$\wp_j : \text{LP}_N \rightarrow \text{LP}_N, \quad \wp_j(\beta) = (\beta \setminus (\{j, k_1\}, \{j+1, k_2\})) \cup \{j, j+1\} \cup \{k_1, k_2\}, \quad (2.23)$$

where  $k_1$  (resp.  $k_2$ ) is the pair of  $j$  (resp.  $j+1$ ) in  $\beta$  (and  $\{j, k_1\}, \{j+1, k_2\}, \{k_1, k_2\}$  are unordered).



Moreover, we have (using the cyclic indexing convention  $2N + 1 \mapsto 1$  to define the tying operation  $\wp_{2N}$ )

$$\lim_{\substack{x_1 \rightarrow -\infty \\ x_{2N} \rightarrow +\infty}} \frac{\mathcal{H}_\beta(\mathbf{x})}{|x_{2N} - x_1|^{-2h(\kappa)}} = \frac{1}{C(\kappa)} \begin{cases} \nu(\kappa) \mathcal{H}_{\beta/\{1, 2N\}}(x_2, \dots, x_{2N-1}), & \text{if } \{1, 2N\} \in \beta, \\ \mathcal{H}_{\wp_{2N}(\beta)/\{1, 2N\}}(x_2, \dots, x_{2N-1}), & \text{if } \{1, 2N\} \notin \beta. \end{cases} \quad (2.24)$$

[FPW24a, Appendix C] provides a proof for (2.22) for the case  $\{j, j+1\} \notin \beta$  with  $\kappa \in (4, 8)$ , which is rather long and technical. The proof presented here uses Proposition 2.5 and works for all  $\kappa \in (0, 8) \setminus \{\frac{8}{m} : m \in \mathbb{Z}_{>0}\}$  and is much shorter. Readers may also have a look at proof of Proposition D.2, where we prove the asymptotics for  $N = 2$  and the notations there are simpler.

*Proof.* The normalization property  $\mathcal{H}_\emptyset \equiv 1$  is understood as an empty product of integrals in the definition (2.2). The asymptotics in (2.24) is a consequence of (2.22) and Proposition 2.2.

In the case  $\{j, j+1\} \in \beta$ , let  $\vartheta_r^\beta \ni u_r$  be the Pochhammer contour (1.14) which surrounds the points  $x_j$  and  $x_{j+1}$ . Because the integration contours remain bounded away from each other, we see that the integral relevant for evaluating the limit is

$$\lim_{x_j, x_{j+1} \rightarrow \xi} \oint_{\vartheta_r^\beta} du_r \frac{f_\beta(\mathbf{x}; \mathbf{u})}{(x_{j+1} - x_j)^{-2h(\kappa)}}.$$

After making the change of variables  $v = \frac{u_r - x_j}{x_{j+1} - x_j}$  in this integral and collecting all the factors, carefully noting that no branch cuts are crossed, and after taking into account cancellations, we obtain (2.22):

$$\begin{aligned} & \lim_{x_j, x_{j+1} \rightarrow \xi} \frac{\mathcal{H}_\beta(\mathbf{x})}{(x_{j+1} - x_j)^{-2h(\kappa)}} \\ &= \left( \oint_{\vartheta(0,1)} du (u-1)^{-4/\kappa} u^{-4/\kappa} \right) \oint_{\vartheta_1^\beta} du_1 \cdots \oint_{\vartheta_{r-1}^\beta} du_{r-1} \oint_{\vartheta_{r+1}^\beta} du_{r+1} \cdots \oint_{\vartheta_N^\beta} du_N f_{\beta/\{j, j+1\}}(\ddot{\mathbf{x}}_j; \dot{\mathbf{u}}_r) \\ &= \frac{\nu(\kappa)}{C(\kappa)} \mathcal{H}_{\beta/\{1, 2\}}(\ddot{\mathbf{x}}_j) \end{aligned}$$

where  $\dot{\mathbf{u}}_r := (u_1, u_2, \dots, u_{r-1}, u_{r+1}, \dots, u_N)$ , and the prefactor evaluates to (2.14).

In the case  $\{j, j+1\} \notin \beta$ , thanks to Proposition 2.2, by conjugating by a suitable Möbius map it suffices without loss of generality to check (2.22) in the case  $j = 1$ . Write  $\hat{\beta} := \beta \setminus (\{\{1, b_1\}, \{2, b_2\}\})$  (we do not relabel the indices here), and denote by  $\Gamma_{\hat{\beta}}$  the integration contours in  $\mathcal{H}_\beta$  other than the two Pochhammer contours surrounding the pairs  $\{x_1, x_{b_1}\}$  and  $\{x_2, x_{b_2}\}$ , respectively. Proposition 2.5, gives

$$\begin{aligned} \mathcal{H}_\beta(\mathbf{x}) &= \frac{\nu(\kappa)}{C(\kappa)} \oint_{\vartheta_2^\beta} du_2 \cdots \oint_{\vartheta_N^\beta} du_N \hat{f}_\beta^b(\mathbf{x}; \mathbf{u}_1) \\ &= \frac{\nu(\kappa)}{C(\kappa)} (x_2 - x_1)^{2/\kappa} (x_{b_1} - x_1)^{-2h(\kappa)} (x_{b_1} - x_2)^{-2h(\kappa)} (x_{b_2} - x_1)^{2/\kappa} (x_{b_2} - x_2)^{2/\kappa} \\ &\quad \times \prod_{\substack{3 \leq i \leq 2N \\ i \notin \{b_1, b_2\}}} ((x_i - x_1)^{2/\kappa} (x_i - x_2)^{2/\kappa}) \\ &\quad \times \int_{\Gamma_{\hat{\beta}}} d\mathbf{u}_1 \oint_{\vartheta(x_2, x_{b_2})} du_2 \tilde{f}(\mathbf{x}_1; \mathbf{u}_1) (u_2 - x_{b_1})^{4h(\kappa)} (u_2 - x_{b_2})^{-4/\kappa} (u_2 - x_1)^{-4/\kappa} (u_2 - x_2)^{-4/\kappa} \\ &\quad \times \prod_{\substack{3 \leq i \leq 2N \\ i \notin \{b_1, b_2\}}} (u_2 - x_i)^{-4/\kappa} \prod_{3 \leq s \leq N} (u_s - u_2)^{8/\kappa} \prod_{3 \leq s \leq N} ((u_s - x_1)^{-4/\kappa} (u_s - x_2)^{-4/\kappa}) \end{aligned}$$

where  $\mathbf{u}_1 = (u_3, u_4, \dots, u_N)$  and  $\hat{f}_\beta^b$  is defined in (2.10) with  $r = 1$  and the conjugate charge at  $x_{b_1}$ , and

$$\begin{aligned} \tilde{f}(\mathbf{x}_1; \mathbf{u}_1) &:= \prod_{\substack{3 \leq i < l \leq 2N \\ i, l \neq b_1}} (x_l - x_i)^{2/\kappa} \prod_{\substack{3 \leq i \leq 2N \\ i \neq b_1}} |x_{b_1} - x_i|^{-2h(\kappa)} \\ &\quad \times \prod_{3 \leq t < s \leq N} (u_s - u_t)^{8/\kappa} \prod_{\substack{3 \leq i < 2N \\ i \neq b_1 \\ 3 \leq s \leq N}} (u_s - x_i)^{-4/\kappa} \prod_{3 \leq s \leq N} (u_s - x_{b_1})^{4h(\kappa)}, \end{aligned}$$

and the branch of  $\tilde{f}(\mathbf{x}_1; \cdot)$  and the rest of the products in the integrand is chosen to be real and positive when  $x_{a_s} < \text{Re}(u_s) < x_{a_{s+1}}$  for all  $s \in \{3, 4, \dots, N\}$ . Next, we use the change of variables

$$v = \frac{\text{CR}(x_{b_2}, x_{b_1}, x_2, u_2)}{\text{CR}(x_1, x_2, x_{b_2}, x_{b_1})},$$

where CR is the cross-ratio

$$\text{CR}(y_1, y_2, y_3, y_4) := \frac{(y_2 - y_1)(y_4 - y_3)}{(y_3 - y_1)(y_4 - y_2)}, \quad (2.25)$$

to obtain

$$\begin{aligned} \mathcal{H}_\beta(\mathbf{x}) &= \frac{\nu(\kappa)}{C(\kappa)} (x_2 - x_1)^{2/\kappa} (x_{b_1} - x_{b_2})^{8/\kappa - 1} (x_{b_1} - x_1)^{-2h(\kappa)} (x_{b_2} - x_2)^{-2h(\kappa)} (x_{b_1} - x_2)^{-2/\kappa} (x_{b_2} - x_1)^{-2/\kappa} \\ &\quad \times \chi^{1-8/\kappa} \int_{\Gamma_{\hat{\beta}}} d\mathbf{u}_1 \oint_{\vartheta(0, \frac{1}{\chi})} dv \tilde{f}(\mathbf{x}_1; \mathbf{u}_1) \tilde{g}(x_1, x_2, v; \mathbf{x}_1; \mathbf{u}_1) k(\chi, v), \end{aligned} \quad (2.26)$$

denoting  $\chi := \text{CR}(x_1, x_2, x_{b_2}, x_{b_1})$  and

$$\begin{aligned} \tilde{g}(x_1, x_2, v; \mathbf{x}_1; \mathbf{u}_1) &:= \prod_{\substack{3 \leq i \leq 2N \\ i \notin \{b_1, b_2\}}} \left| x_{b_1} - (x_{b_1} - x_2) \left( 1 + v \chi \frac{x_{b_2} - x_2}{x_{b_1} - x_{b_2}} \right)^{-1} - x_i \right|^{-4/\kappa} \\ &\quad \times \prod_{3 \leq s \leq N} \left( u_s - x_{b_1} + (x_{b_1} - x_2) \left( 1 + v \chi \frac{x_{b_2} - x_2}{x_{b_1} - x_{b_2}} \right)^{-1} \right)^{8/\kappa} \end{aligned}$$



$$\times \prod_{3 \leq s \leq N} ((u_s - x_1)^{-4/\kappa} (u_s - x_2)^{-4/\kappa}) \prod_{\substack{3 \leq i \leq 2N \\ i \notin \{b_1, b_2\}}} ((x_i - x_1)^{2/\kappa} (x_i - x_2)^{2/\kappa}),$$

whose branch is chosen to be real and positive when

$$0 < \operatorname{Re}(v) < \frac{\operatorname{CR}(x_{b_2}, x_{b_1}, x_2, x_3)}{\operatorname{CR}(x_1, x_2, x_{b_2}, x_{b_1})}, \quad \text{and} \quad x_{a_s} < \operatorname{Re}(u_s) < x_{a_s+1}, \quad \text{for all } s \in \{3, 4, \dots, N\},$$

and where the branch of

$$k(\chi, v) := (1 + (1 - \chi)v)^{-4/\kappa} v^{-4/\kappa} (1 - \chi v)^{-4/\kappa},$$

is chosen to be real and positive when  $0 < \operatorname{Re}(v) < 1/\chi$ . Note that  $k(\chi, v) = f(0, \frac{1}{\chi-1}, \frac{1}{\chi}; v)$  has the Coulomb gas integrand form (2.17) with symmetric charges  $\alpha = 1/\sqrt{\kappa}$  at the variables  $(0, \frac{1}{\chi-1}, \frac{1}{\chi})$  and the negative screening charge  $\alpha_- = -2/\sqrt{\kappa}$  at  $v$ . Hence, using Lemma 2.8, we obtain

$$\begin{aligned} \oint_{\partial(0, \frac{1}{\chi})} k(\chi, v) dv &= (1 - e^{8\pi i/\kappa})(1 - e^{-8\pi i/\kappa}) \int_{\epsilon}^{\frac{1}{\chi} - \epsilon} k(\chi, v) dv \\ &\quad + (1 - e^{8\pi i/\kappa}) \oint_{\partial B(0, \epsilon)} k(\chi, v) dv + (1 - e^{-8\pi i/\kappa}) \oint_{\partial B(\frac{1}{\chi}, \epsilon)} k(\chi, v) dv, \end{aligned}$$

for all  $\epsilon > 0$  small enough. Now, we take  $\chi \rightarrow 0+$  (while keeping  $\epsilon$  fixed). On the one hand, using the fact that  $8/\kappa > 1$  and the dominated convergence theorem, we see that the first two terms have a limit,

$$\begin{aligned} &(1 - e^{8\pi i/\kappa})(1 - e^{-8\pi i/\kappa}) \lim_{\chi \rightarrow 0+} \int_{\epsilon}^{\frac{1}{\chi} - \epsilon} k(\chi, v) dv + (1 - e^{8\pi i/\kappa}) \lim_{\chi \rightarrow 0+} \oint_{\partial B(0, \epsilon)} k(\chi, v) dv \\ &= (1 - e^{8\pi i/\kappa}) \oint_{\varrho(0, \infty)} (1 + v)^{-4/\kappa} v^{-4/\kappa} dv, \end{aligned}$$

where  $\varrho(0, \infty)$  is a contour which starts from  $\infty$ , then winds around 0 once in clockwise direction and finally goes back to  $\infty$ , illustrated in Eq. (B.7) in Lemma B.2. On the other hand, the third term is

$$\lim_{\chi \rightarrow 0+} \oint_{\partial B(\frac{1}{\chi}, \epsilon)} k(\chi, v) dv = 0,$$

since  $|k(\chi, v)| \sim v^{-8/\kappa}$  as  $v \rightarrow \infty$ . Consequently, we have

$$\begin{aligned} \lim_{x_1, x_2 \rightarrow \xi} \frac{\mathcal{H}_\beta(\mathbf{x})}{(x_2 - x_1)^{-2h(\kappa)}} &= \frac{\nu(\kappa)}{C(\kappa)} \int_{\Gamma_\beta} \tilde{f}(\tilde{\mathbf{x}}_1; \tilde{\mathbf{u}}_1) d\tilde{\mathbf{u}}_1 \left( \lim_{\chi \rightarrow 0+} \oint_{\partial(0, \frac{1}{\chi})} k(\chi, v) dv \right) \\ &= \frac{\nu(\kappa)}{C(\kappa)} \int_{\Gamma_\beta} \tilde{f}(\tilde{\mathbf{x}}_1; \tilde{\mathbf{u}}_1) d\tilde{\mathbf{u}}_1 \left( (1 - e^{8\pi i/\kappa}) \int_{\varrho(0, \infty)} (1 + v)^{-4/\kappa} v^{-4/\kappa} dv \right). \end{aligned}$$

Using Eq. (B.4) from Lemma B.2 to evaluate the integral

$$\int_{\varrho(0, \infty)} (1 + v)^{-4/\kappa} v^{-4/\kappa} dv = \frac{(1 - e^{-8\pi i/\kappa})}{\nu(\kappa)} \frac{\Gamma(1 - 4/\kappa)^2}{\Gamma(2 - 8/\kappa)}$$

and Proposition 2.5, we conclude that

$$\begin{aligned} \lim_{x_1, x_2 \rightarrow \xi} \frac{\mathcal{H}_\beta(\mathbf{x})}{(x_2 - x_1)^{-2h(\kappa)}} &= \frac{\nu(\kappa)}{C(\kappa)} (1 - e^{8\pi i/\kappa}) \frac{(1 - e^{-8\pi i/\kappa})}{\nu(\kappa)} \frac{\Gamma(1 - 4/\kappa)^2}{\Gamma(2 - 8/\kappa)} \int_{\Gamma_\beta} \tilde{f}(\tilde{\mathbf{x}}_1; \tilde{\mathbf{u}}_1) d\tilde{\mathbf{u}}_1 \\ &= \frac{1}{C(\kappa)} \left( \frac{\nu(\kappa)}{C(\kappa)} \int_{\Gamma_\beta} \tilde{f}(\tilde{\mathbf{x}}_1; \tilde{\mathbf{u}}_1) d\tilde{\mathbf{u}}_1 \right) = \frac{1}{C(\kappa)} \mathcal{H}_{\varphi_1(\beta)/\{1,2\}}(\tilde{\mathbf{x}}_1), \end{aligned}$$

which shows (2.22) for  $j = 1$  and  $\{1, 2\} \notin \beta$  and finishes the proof.  $\square$

**Corollary 2.13.** Fix  $\kappa \in (0, 8) \setminus \{\frac{8}{m} : m \in \mathbb{Z}_{>0}\}$ . The collection  $\{\mathcal{F}_\beta : \beta \in \text{LP}_N\}$  satisfies the properties (PDE), (COV), and (PLB). Furthermore, with  $\mathcal{F}_\emptyset \equiv 1$  for the empty link pattern  $\emptyset \in \text{LP}_0$ , the following recursive asymptotics property holds. Fix  $N \geq 1$  and  $j \in \{1, 2, \dots, 2N - 1\}$ . Then, for all  $\xi \in (x_{j-1}, x_{j+2})$ , using the notation (1.21),

$$\lim_{x_j, x_{j+1} \rightarrow \xi} \frac{\mathcal{F}_\beta(\mathbf{x})}{(x_{j+1} - x_j)^{-2h(\kappa)}} = \begin{cases} \nu(\kappa) \mathcal{F}_{\beta/\{j, j+1\}}(\tilde{\mathbf{x}}_j), & \text{if } \{j, j+1\} \in \beta, \\ \mathcal{F}_{\emptyset_{2N}(\beta)/\{j, j+1\}}(\tilde{\mathbf{x}}_j), & \text{if } \{j, j+1\} \notin \beta. \end{cases} \quad (\mathcal{F}\text{-ASY})$$

Moreover, we have

$$\lim_{\substack{x_1 \rightarrow -\infty \\ x_{2N} \rightarrow +\infty}} \frac{\mathcal{F}_\beta(\mathbf{x})}{|x_{2N} - x_1|^{-2h(\kappa)}} = \begin{cases} \nu(\kappa) \mathcal{F}_{\beta/\{1, 2N\}}(x_2, \dots, x_{2N-1}), & \text{if } \{1, 2N\} \in \beta, \\ \mathcal{F}_{\emptyset_{2N}(\beta)/\{1, 2N\}}(x_2, \dots, x_{2N-1}), & \text{if } \{1, 2N\} \notin \beta. \end{cases} \quad (\mathcal{F}\text{-ASY}_\infty)$$

*Proof.* (PDE) & (COV) are well-known, see, e.g., [FPW24a, Propositions 2.2 & 2.3]. (See also Proposition 2.2 for a stronger covariance under  $\varphi \in \text{SL}(2, \mathbb{R})$ .) The bound (PLB) is clear from the definition. The asymptotics properties ( $\mathcal{F}\text{-ASY}$ ,  $\mathcal{F}\text{-ASY}_\infty$ ) follow from Proposition 2.12 and the relation (2.1).  $\square$

## 2.6 Coulomb gas integrals for exceptional values of $\kappa$

Let us record here the observation that the Coulomb gas type integrals vanish for the special values of the parameter of the form  $\kappa = 8/m$  for integers  $m \geq 2$ , including  $\kappa = 4$ ,  $\kappa = 8/3$ , and  $\kappa = 2$ .

**Lemma 2.14.** For  $\kappa \in \{\frac{8}{m} : m \in \mathbb{Z}_{>0} + 1\}$ , we have

$$\mathcal{H}_\beta(\mathbf{x}) \equiv 0, \quad \text{for all } \beta \in \text{LP}_N \text{ and } \mathbf{x} \in \mathfrak{X}_{2N}.$$

*Proof.* We first check the case  $\kappa = \frac{4}{m}$  for  $m \in \mathbb{Z}_{>0}$ . Note that the integrand (1.12) reads

$$f_\beta(\mathbf{x}; \mathbf{u}) = \prod_{1 \leq i < j \leq 2N} (x_j - x_i)^{m/2} \prod_{1 \leq r < s \leq N} (u_s - u_r)^{2m} \prod_{\substack{1 \leq i \leq 2N \\ 1 \leq r \leq N}} (u_r - x_i)^{-m}, \quad \kappa = \frac{4}{m}.$$

For any fixed  $s \in \{1, 2, \dots, N\}$ , and  $\mathbf{x} \in \mathfrak{X}_{2N}$ , and  $\mathbf{u}_s \in \vartheta_1^\beta \times \vartheta_2^\beta \times \dots \times \vartheta_{s-1}^\beta \times \vartheta_{s+1}^\beta \times \dots \times \vartheta_N^\beta$ , the function  $u_s \mapsto f_\beta(\mathbf{x}; \mathbf{u})$  is holomorphic on  $\mathbb{C} \setminus \{x_1, \dots, x_{2N}, u_1, \dots, u_{s-1}, u_{s+1}, \dots, u_N\}$ . Because the Pochhammer contour  $\vartheta_s^\beta$  surrounds both of its poles  $\{x_{a_s}, x_{b_s}\}$  once in the positive direction and once in the negative direction, we see by the residue theorem that the integral of  $f_\beta$  around  $\vartheta_s^\beta$  vanishes:

$$\oint_{\vartheta_s^\beta} du_s f_\beta(\mathbf{x}; \mathbf{u}) = 2\pi i \left( \text{Res}_{u_s=x_{a_s}} f_\beta(\mathbf{x}; \mathbf{u}) \right) (1-1) + 2\pi i \left( \text{Res}_{u_s=x_{b_s}} f_\beta(\mathbf{x}; \mathbf{u}) \right) (1-1) = 0.$$

Consequently, using Fubini's theorem, we conclude that  $\mathcal{H}_\beta(\mathbf{x}) \equiv 0$  for  $\kappa \in \{\frac{8}{2m} : m \in \mathbb{Z}_{>0}\}$ .

Next, we check the case  $\kappa = \frac{8}{2m+1}$  for  $m \in \mathbb{Z}_{>0}$ . Note that the integrand (1.12) reads

$$f_\beta(\mathbf{x}; \mathbf{u}) = \prod_{1 \leq i < j \leq 2N} (x_j - x_i)^{(2m+1)/4} \prod_{1 \leq r < s \leq N} (u_s - u_r)^{2m+1} \prod_{\substack{1 \leq i \leq 2N \\ 1 \leq r \leq N}} (u_r - x_i)^{-(2m+1)/2}, \quad \kappa = \frac{8}{2m+1}.$$

Using Corollary 2.9, we can replace the Pochhammer contours  $\vartheta_r^\beta = \vartheta(x_{a_r}, x_{b_r})$  (1.14) in the integral  $\mathcal{H}_\beta$  (2.2) by simple clockwise loops  $\varrho_r^\beta$ , each surrounding the two points  $\{x_{a_r}, x_{b_r}\}$  and no other points (see Figure 2.2). Moreover, a repeated application of Lemma 2.10 shows that we can replace these loops  $\varrho_r^\beta$  by nested loops  $\tilde{\varrho}_r^\beta$  that are symmetric with respect to the real axis — see Figure 2.3. Moreover, we can then deform the outermost such loop into a loop surrounding the point  $\infty$  at infinity.

Now, thanks to Proposition 2.2, we may assume that  $\{1, 2N\} \in \beta$  without loss of generality. Then, for any fixed  $\mathbf{x} \in \mathfrak{X}_{2N}$  and  $\mathbf{u}_1 \in \vartheta_2^\beta \times \vartheta_3^\beta \times \cdots \times \vartheta_N^\beta$ , the function  $u_1 \mapsto f_\beta(\mathbf{x}; \mathbf{u})$  is holomorphic on  $\mathbb{C} \setminus [x_1, x_{2N}]$ . In this case, the aforementioned outermost loop is  $\varrho_1^\beta$ , and using the residue theorem, we see that

$$\begin{aligned}
\mathcal{H}_\beta(\mathbf{x}) &= \oint_{\vartheta_1^\beta} du_1 \oint_{\vartheta_2^\beta} du_2 \cdots \oint_{\vartheta_N^\beta} du_N f_\beta(\mathbf{x}; \mathbf{u}) && \text{[by (2.2)]} \\
&= 2^N \oint_{\varrho_2^\beta} du_2 \cdots \oint_{\varrho_N^\beta} du_N \oint_{\varrho_1^\beta} du_1 f_\beta(\mathbf{x}; \mathbf{u}) && \text{[by Corollary 2.9]} \\
&= 2^N \oint_{\varrho_2^\beta} du_2 \cdots \oint_{\varrho_N^\beta} du_N \oint_{\varrho_1^\beta} du_1 f_\beta(\mathbf{x}; \mathbf{u}) && \text{[by Lemma 2.10]} \\
&= -2^{N+1} \pi i \oint_{\varrho_2^\beta} du_2 \cdots \oint_{\varrho_N^\beta} du_N \left( \operatorname{Res}_{u_1=\infty} f_\beta(\mathbf{x}; \mathbf{u}) \right) && \text{[by residue theorem]} \\
&= 0,
\end{aligned}$$

since  $\operatorname{Res}_{u_1=\infty} f_\beta(\mathbf{x}; \mathbf{u}) = -\lim_{|u_1| \rightarrow \infty} u_1 f_\beta(\mathbf{x}; \mathbf{u}) = 0$ . This concludes the proof.  $\square$

**Remark 2.15.** Note that Lemma 2.14 does not apply to the case where  $\kappa = 8$ , because the residue of the integrand  $u_1 f_\beta(\mathbf{x}; \mathbf{u})$  is not zero in that case.

### 3 Constructing variants of multiple $\operatorname{SLE}_\kappa$ measures for $\kappa \in (4, 8)$

In this section, we fix  $\kappa \in (4, 8)$  and omit the dependence on  $\kappa$  from the notation, while we shall highlight  $N \geq 1$ . In this and the next Section 4, we make use of two particular Coulomb gas integrals:  $\mathcal{F}_{\overbrace{\cup \cup \cup}_N}$  and  $\mathcal{F}_{\overbrace{\cup \cup \cup}_N}$ , where

$$\begin{aligned}
\overbrace{\cup \cup \cup}_N &= \{\{1, 2\}, \{3, 4\}, \dots, \{2N-1, 2N\}\} \in \operatorname{LP}_N, \\
\overbrace{\cup \cup \cup}_N &= \{\{1, 2N\}, \{2, 3\}, \{4, 5\}, \dots, \{2N-2, 2N-1\}\} \in \operatorname{LP}_N.
\end{aligned}$$

#### 3.1 Conformally invariant measures on curves

The goal of this section is to construct a certain probability measure on the curve space  $X_0(\Omega; x_1, \dots, x_{2N})$  defined in (1.44). It serves as a tool to construct the global  $N$ - $\operatorname{SLE}_\kappa$  measures of Theorem 1.21. To this end, we will first have to introduce some notation and terminology<sup>13</sup>.

**Schramm-Loewner evolution.** To begin, recall that the chordal  $\operatorname{SLE}_\kappa$  is a random curve  $\eta$ , parameterized by speed  $\kappa > 0$  and a simply connected domain  $\Omega$  together with two distinct endpoints  $x_1, x_2 \in \partial\Omega$  of the curve  $\eta$  on the boundary (so  $\operatorname{SLE}_\kappa$  is a random chord in  $(\Omega; x_1, x_2)$ ). It is a conformally invariant process (as in Definition 3.1), and taking as the reference domain the upper half-plane  $\mathbb{H} = \{z \in \mathbb{C} : \operatorname{Im}(z) > 0\}$ , i.e.,  $(\Omega; x_1, x_2) = (\mathbb{H}; 0, \infty)$ , the chordal  $\operatorname{SLE}_\kappa$  curve  $\eta$  can be constructed as a random Loewner chain via solving the ODE

$$\partial_t g_t(z) = \frac{2}{g_t(z) - W_t}, \quad \text{with initial condition} \quad g_0(z) = z, \quad z \in \overline{\mathbb{H}}, \quad (3.1)$$

where the driving function  $W_t = \sqrt{\kappa} B_t$  is given by the standard one-dimensional Brownian motion. For each  $z \in \overline{\mathbb{H}}$ , the evolution  $(g_t(z), t \geq 0)$  is well-defined only up to the blow-up time of the equation (3.1),  $\tau(z) := \sup\{t \geq 0 : \min_{s \in [0, t]} |g_s(z) - W_s| > 0\}$ , when the denominator becomes zero, and the domain of

<sup>13</sup>As usual, we write  $\mathbf{x} = (x_1, \dots, x_{2N})$  and  $\varphi(\mathbf{x}) = (\varphi(x_1), \dots, \varphi(x_{2N}))$ .

the conformal map  $g_t$  is thus given by the complement of the SLE $_{\kappa}$  hull  $K_t := \{z \in \mathbb{H} : \tau(z) \leq t\}$ . The conformal maps  $g_t: \mathbb{H} \setminus K_t \rightarrow \mathbb{H}$  are uniquely determined and normalized as  $\lim_{z \rightarrow \infty} |g_t(z) - z| = 0$ . When  $\kappa \in (0, 4]$ , we have  $K_t = \eta[0, t]$  while when  $\kappa > 4$ , the curve  $\eta$  has self-touchings and  $K_t$  is the filling of  $\eta[0, t]$  in the sense that the domain  $\mathbb{H} \setminus K_t$  is the unbounded connected component of  $\mathbb{H} \setminus \eta[0, t]$ . Notably, when  $\kappa \geq 8$ , the chordal SLE $_{\kappa}$  curve is space-filling — thus, in the present article we shall be concerned with the range  $\kappa \in (0, 8)$ . Detailed introductions can be found in textbooks, e.g. [Wer03, Law05, Kem17].

**Conformal invariance.** The global multiple SLE measures of Theorem 1.21 are conformally invariant.

**Definition 3.1.** A collection  $\{\mathbb{P}(\mathbb{H}; \mathbf{x}) : \mathbf{x} \in \mathfrak{X}_{2N}\}$  of probability measures on  $X_0(\mathbb{H}; \mathbf{x})$  is *Möbius invariant* if the following property holds. Suppose  $(\eta_1, \dots, \eta_N) \sim \mathbb{P}(\mathbb{H}; \mathbf{x})$  and  $\varphi: \mathbb{H} \rightarrow \mathbb{H}$  is a Möbius map such that  $\varphi(x_1) < \dots < \varphi(x_{2N})$ . Then, the law of the collection  $(\varphi(\eta_1), \dots, \varphi(\eta_N))$  is  $\mathbb{P}(\mathbb{H}; \varphi(\mathbf{x}))$ .

Whenever the collection  $\{\mathbb{P}(\mathbb{H}; \mathbf{x}) : \mathbf{x} \in \mathfrak{X}_{2N}\}$  is Möbius invariant, we may extend its definition for general polygons  $(\Omega; \mathbf{y}) = (\Omega; y_1, \dots, y_{2N})$  via conformal invariance:  $\mathbb{P}(\Omega; \mathbf{y})$  is defined to be the pushforward law of  $(\varphi(\eta_1), \dots, \varphi(\eta_N))$ , where  $\varphi: \mathbb{H} \rightarrow \Omega$  is any conformal bijection such that  $\varphi^{-1}(y_1) < \dots < \varphi^{-1}(y_{2N})$  and  $(\eta_1, \dots, \eta_N) \sim \mathbb{P}(\mathbb{H}; \varphi^{-1}(\mathbf{y}))$ . We then say that the collection  $\{\mathbb{P}(\Omega; \mathbf{y}) : (\Omega; \mathbf{y}) \text{ is a polygon}\}$  of probability measures is *conformally invariant*.

**SLE with force points.** The SLE $_{\kappa}(\underline{\rho})$  *process* is a variant of the chordal SLE $_{\kappa}$  where one keeps track of additional points on the boundary. We will encounter the following version of it. Fix  $y_1 < \dots < y_{2N}$  and  $j \in \{1, \dots, 2N\}$ . Let  $\underline{y}^L = (y_1, \dots, y_{j-1})$  and  $\underline{y}^R = (y_{j+1}, \dots, y_{2N})$  and  $\underline{\rho}^L = (\rho_1, \dots, \rho_{j-1}) \in \mathbb{R}^{j-1}$  and  $\underline{\rho}^R = (\rho_{j+1}, \dots, \rho_{2N}) \in \mathbb{R}^{2N-j}$ . An SLE $_{\kappa}(\underline{\rho}^L; \underline{\rho}^R)$  process in  $(\mathbb{H}; y_j, \infty)$  started at  $y_j$  with force points  $(\underline{y}^L; \underline{y}^R)$  is the Loewner evolution driven by  $W_t$  that solves the following system of integrated SDEs:

$$\begin{aligned} W_t &= y_j + \sqrt{\kappa} B_t + \sum_{i=1}^{j-1} \int_0^t \frac{\rho_i ds}{W_s - g_s(y_i)} + \sum_{i=j+1}^{2N} \int_0^t \frac{\rho_i ds}{W_s - g_s(y_i)}, \\ g_t(y_i) &= y_i + \int_0^t \frac{2 ds}{g_s(y_i) - W_s}, \quad i \in \{1, 2, \dots, 2N\} \setminus \{j\}, \end{aligned} \tag{3.2}$$

where  $B_t$  is the one-dimensional Brownian motion started at  $B_0 = 0$ . Note that the process  $g_t(y_i)$  is the time-evolution of the point  $y_i$  for each  $i \neq j$ . We define the *continuation threshold* of the SLE $_{\kappa}(\underline{\rho}^L; \underline{\rho}^R)$  to be the infimum of the time  $t$  for which

$$\text{either } \sum_{\substack{i \in \{1, \dots, j-1\} \\ g_t(y_i) = W_t}} \rho_i \leq -2, \quad \text{or} \quad \sum_{\substack{i \in \{j+1, \dots, 2N\} \\ g_t(y_i) = W_t}} \rho_i \leq -2.$$

By [MS16a], the SLE $_{\kappa}(\underline{\rho}^L; \underline{\rho}^R)$  process is well-defined up to the continuation threshold, and it is almost surely generated by a continuous curve up to and including the continuation threshold. Concretely, it is the Loewner chain associated to the partition function

$$\mathcal{Z}(y_1, \dots, y_{2N}) = \prod_{\substack{1 \leq i < 2N \\ i \neq j}} (y_i - y_j)^{\rho_i / \kappa} \prod_{\substack{1 \leq i < l < 2N \\ i, l \neq j}} (y_l - y_i)^{\rho_l \rho_i / 2\kappa}.$$

**Poisson kernel.** (Boundary) Poisson kernel  $H(\Omega; x, y)$  is defined for nice Dobrushin domains  $(\Omega; x, y)$ . When  $\Omega = \mathbb{H}$ , we have

$$H(\mathbb{H}; x, y) = |y - x|^{-2}, \quad x, y \in \mathbb{R}.$$

For a nice Dobrushin domain  $(\Omega; x, y)$ , we define the Poisson kernel via conformal covariance:

$$H(\Omega; x, y) = |\varphi'(x)\varphi'(y)|H(\mathbb{H}; \varphi(x), \varphi(y)),$$

where  $\varphi$  is any conformal map from  $\Omega \rightarrow \mathbb{H}$ . Note that the Poisson kernel satisfies the following monotonicity. Let  $(\Omega; x, y)$  be a nice Dobrushin domain and let  $U \subset \Omega$  be a subdomain that agrees with  $\Omega$  in a neighborhood of the boundary arc  $(yx)$ . Then, we have

$$H(U; x, y) \leq H(\Omega; x, y). \quad (3.3)$$

### 3.2 Construction of a probability measure $P_N$

In this section, we will construct probability measures  $P_N = P_N(\Omega; \mathbf{x})$  on  $N$ -tuples of chordal curves  $\boldsymbol{\eta} := (\eta_1, \dots, \eta_N) \in X_0(\Omega; \mathbf{x})$  for every  $N \geq 1$  by induction. When  $N = 1$ , the measure  $P_1 = P_1(\Omega; x_1, x_2)$  on the Dobrushin domain equals the law of  $\text{SLE}_\kappa$  on  $\Omega$  from  $x_1$  to  $x_2$ . Note that by the reversibility of  $\text{SLE}_\kappa$  [MS16c], we have  $P_1(\Omega; x_1, x_2) = P_1(\Omega; x_2, x_1)$ . We use this fact in the proof of Proposition 3.3. When  $N = 2$ , a similar construction was used in [FLW23] to derive a decomposition of global 2-SLE. As  $N$  becomes larger, there are more possibilities for the curves to hit different segments of the boundary. To address all of them, we introduce the following construction which involves heavy notation. Unfortunately, we find it unavoidable. We shall highlight the orientation of the curves by denoting

$$(\vec{\eta}_1, \dots, \vec{\eta}_N) \sim \vec{P}_N = \vec{P}_N(\Omega; \mathbf{x}),$$

to emphasize that  $\vec{\eta}_s$  starts from  $x_{2s-1}$  for each  $1 \leq s \leq N$ . The construction is given in Definition 3.6.

Given the construction of  $\vec{P}_N$ , we will also define another probability measure  $\tilde{P}_N = \tilde{P}_N(\Omega; \mathbf{x})$  as follows. Choose an anti-conformal map  $\zeta$  from  $\Omega$  onto  $\Omega$  and sample  $(\vec{\eta}_1, \dots, \vec{\eta}_N) \sim \vec{P}_N(\Omega; \zeta(x_{2N}), \dots, \zeta(x_1))$ . Then, we define  $\tilde{P}_N$  to be the law of  $(\vec{\eta}_1, \dots, \vec{\eta}_N) := (\zeta^{-1}(\vec{\eta}_N), \dots, \zeta^{-1}(\vec{\eta}_1))$ . Note that under  $\tilde{P}_N$ , each curve  $\vec{\eta}_s$  starts from  $x_{2s}$  for  $1 \leq s \leq N$ . (Corollary 4.22 implies that these measures agree.)

Now, fix  $N \geq 2$  and suppose we have constructed  $\vec{P}_n$  for all  $n \leq N - 1$ . We will then construct the measure  $\vec{P}_N$  in this section — see Definition 3.6. Throughout, let  $\phi$  be the Möbius transformation sending<sup>14</sup>  $x_1 \mapsto -\infty$ , and  $x_2 \mapsto 0$ , and  $x_3 \mapsto 1$ ,

$$\phi(z) := \left( \frac{z - x_2}{z - x_1} \right) \left( \frac{x_3 - x_1}{x_3 - x_2} \right), \quad z \in \mathbb{H}. \quad (3.4)$$

We first construct  $N$  conformally invariant probability measures  $P_N^{x_1 \rightarrow x_{2b}}$  on  $X_0(\mathbb{H}; \mathbf{x})$  with  $\mathbf{x} \in \mathfrak{X}_{2N}$ , indexed by  $b \in \{1, 2, \dots, N\}$ , such that the random curve  $\vec{\eta}_1$  under  $(\vec{\eta}_1, \dots, \vec{\eta}_N) \sim P_N^{x_1 \rightarrow x_{2b}}$  ends at  $x_{2b}$  almost surely (Definition 3.2). The probability measure  $\vec{P}_N$  will then be a convex combination of  $\{P_N^{x_1 \rightarrow x_{2b}} : b \in \{1, \dots, N\}\}$  weighted by the probabilities for the curve  $\vec{\eta}_1$  to end at  $x_2, x_4, \dots, x_{2N}$ .

**Definition 3.2.** Fix  $b \in \{1, 2, \dots, N\}$ . For each polygon  $(\Omega; \mathbf{y})$ , the probability measure  $P_N^{y_1 \rightarrow y_{2b}}(\Omega; \mathbf{y})$  on  $X_0(\Omega; \mathbf{y})$  is constructed uniquely in the following manner. Under  $(\vec{\eta}_1, \dots, \vec{\eta}_N) \sim P_N^{y_1 \rightarrow y_{2b}}(\Omega; \mathbf{y})$ , the curve  $\vec{\eta}_1$  traverses from  $y_1$  to  $y_{2b}$ , and each  $\vec{\eta}_s$  starts from  $y_{2s-1}$ , for  $2 \leq s \leq N$ . The measure  $P_N^{y_1 \rightarrow y_{2b}}(\Omega; \mathbf{y})$  is the pushforward of  $P_N^{x_1 \rightarrow x_{2b}} := P_N^{x_1 \rightarrow x_{2b}}(\mathbb{H}; \mathbf{x})$ , which is constructed uniquely in the following manner.

1. Sample a random point  $\dot{U}_b = (U_1, \dots, U_{b-1}, U_{b+1}, \dots, U_N)$  such that<sup>15</sup>

$$U_s \in \begin{cases} (\phi(x_{2s-1}), \phi(x_{2s})), & \text{for } 1 \leq s \leq b-1, \\ (\phi(x_{2s}), \phi(x_{2s+1})), & \text{for } b+1 \leq s \leq N-1, \\ (\phi(x_{2N}), +\infty), & \text{for } s = N, \end{cases} \quad (3.5)$$

whose joint law has the following density with respect to the Lebesgue measure (see Figure 3.1):

$$\mathcal{U}(\dot{U}_b) := \frac{|f(\phi(\dot{\mathbf{x}}_1); \dot{U}_b)|}{\int_{-\infty}^{\phi(x_2)} du_1 \cdots \int_{\phi(x_{2b-3})}^{\phi(x_{2b-2})} du_{b-1} \int_{\phi(x_{2b+2})}^{\phi(x_{2b+3})} du_{b+1} \cdots \int_{\phi(x_{2N})}^{+\infty} du_N |f(\phi(\dot{\mathbf{x}}_1); \dot{\mathbf{u}}_b)|} \quad (3.6)$$

<sup>14</sup>Here, it is useful to require that  $\phi(x_1) = -\infty$  and  $\phi(x_2) = 0$ , while the normalization  $\phi(x_3) = 1$  could be arbitrary.

<sup>15</sup>Throughout, we use the convention that if  $b = 1$ , we do not sample any point on the left side of  $\phi(x_2)$ ; and if  $b = N$ , we do not sample any point on the right side of  $\phi(x_{2N})$ .

where  $\hat{\mathbf{x}}_1 = (x_2, x_3, \dots, x_{2N}) \in \mathfrak{X}_{2N-1}$  and  $\phi(\hat{\mathbf{x}}_1) = (\phi(x_2), \dots, \phi(x_{2N})) \in \mathfrak{X}_{2N-1}$ , and

$$f(\mathbf{y}; \mathbf{v}) := \prod_{1 \leq i < j \leq 2N-1} (y_j - y_i)^{2/\kappa} \prod_{1 \leq r < s \leq N-1} (v_s - v_r)^{8/\kappa} \prod_{\substack{1 \leq i \leq 2N-1 \\ 1 \leq r \leq N-1}} (v_r - y_i)^{-4/\kappa}. \quad (3.7)$$

2. Given  $\hat{U}_b$ , sample a curve  $\gamma_b$  as an  $\text{SLE}_\kappa(\underline{\rho}^L; \underline{\rho}^R)$  process in  $(\mathbb{H}; \phi(x_{2b}), \infty)$  with force points

$$\begin{aligned} \underline{x}^L &= (U_1, \phi(x_2), \phi(x_3), \dots, U_{b-1}, \phi(x_{2b-2}), \phi(x_{2b-1})), \\ \underline{x}^R &= (\phi(x_{2b+1}), \phi(x_{2b+2}), U_{b+1}, \dots, \phi(x_{2N-1}), \phi(x_{2N}), U_N), \end{aligned}$$

and weights  $\underline{\rho}^L = (-4, 2, 2, \dots, -4, 2, 2)$  on its left and  $\underline{\rho}^R = (2, 2, -4, \dots, 2, 2, -4)$  on its right (so the weights are  $-4$  for each  $U_s$  and  $2$  for each  $\phi(x_i)$ ). It follows from [Dub07b, Lemma 15] that

- the curve  $\gamma_b$  does not hit the intervals  $\bigcup_{s=1}^{b-1} [U_s, \phi(x_{2s+1})]$  nor the intervals  $\bigcup_{s=b+1}^N [\phi(x_{2s-1}), U_s]$ ;
- the curve  $\gamma_b$  hits both intervals  $(-\infty, U_1)$  and  $(U_N, +\infty)$ , almost surely.

3. Given  $\gamma_b$ , we denote by  $\Omega_1^R, \dots, \Omega_r^R$  the (random) connected components of  $\mathbb{H} \setminus \gamma_b$  whose boundaries contain some of the points  $\phi(x_i)$  for  $2 \leq i \leq 2b-1$ . For each  $1 \leq m \leq r$ , let  $\phi(x_{2s_m^R}), \dots, \phi(x_{2s_{m+1}^R-1})$  be the boundary points contained in  $\partial\Omega_m^R$ . Using the induction hypothesis that we have already constructed  $\bar{P}_n$  for all  $n \leq N-1$ , sample independently for each  $1 \leq m \leq l$  the curves

$$\gamma_m^R := (\gamma_{s_m^R}, \dots, \gamma_{s_{m+1}^R-1}) \sim \bar{P}_{s_{m+1}^R - s_m^R}(\Omega_m^R; \phi(x_{2s_m^R}), \dots, \phi(x_{2s_{m+1}^R-1})),$$

where each  $\gamma_s$  is a curve starting from  $\phi(x_{2s-1})$  for  $2 \leq s \leq b$ . See also Figure 3.1.

4. Given  $\gamma_b$ , we denote by  $\Omega_1^L, \dots, \Omega_l^L$  the (random) connected components of  $\mathbb{H} \setminus \gamma_b$  whose boundaries contain some of the points  $\phi(x_i)$  for  $2b+1 \leq i \leq 2N$ . For each  $1 \leq m \leq l$ , let  $\phi(x_{2s_m^L+1}), \dots, \phi(x_{2s_{m+1}^L})$  be the boundary points contained in  $\partial\Omega_m^L$ . Using the induction hypothesis that we have already constructed  $\bar{P}_n$  for all  $n \leq N-1$ , sample independently for each  $1 \leq m \leq l$  the curves

$$\gamma_m^L := (\gamma_{s_m^L+1}, \dots, \gamma_{s_{m+1}^L}) \sim \bar{P}_{s_{m+1}^L - s_m^L}(\Omega_m^L; \phi(x_{2s_m^L+1}), \dots, \phi(x_{2s_{m+1}^L})),$$

where each  $\gamma_s$  is a curve starting from  $\phi(x_{2s-1})$  for  $b+1 \leq s \leq N$ . See also Figure 3.1.

5. Denote by  $\tilde{\gamma}_b$  the time-reversal of  $\gamma_b$ . Set  $\bar{P}_N^{x_1 \rightarrow x_{2b}}$  to be the law of  $(\vec{\eta}_1, \dots, \vec{\eta}_N)$ , where<sup>16</sup>


$$\vec{\eta}_1 := \phi^{-1}(\tilde{\gamma}_b), \quad \vec{\eta}_{s+1} := \phi^{-1}(\gamma_s), \quad \text{for } 1 \leq s \leq b-1, \quad \text{and} \quad \vec{\eta}_s := \phi^{-1}(\gamma_s), \quad \text{for } b+1 \leq s \leq N,$$

Note that under the measure  $P_N^{x_1 \rightarrow x_{2b}}$ , the curve  $\vec{\eta}_1$  traverses from  $x_1$  to  $x_{2b}$ , and each  $\vec{\eta}_s$  starts from  $x_{2s-1}$ , for  $2 \leq s \leq N$ .


6. Lastly, by conformal invariance of  $\text{SLE}_\kappa$ , the collection  $\{P_N^{x_1 \rightarrow x_{2b}}(\mathbb{H}; \mathbf{x}) : \mathbf{x} \in \mathfrak{X}_{2n}\}$  is Möbius invariant. Hence, it uniquely determines the probability measures  $P_N^{y_1 \rightarrow y_{2b}}(\Omega; \mathbf{y})$  for all polygons  $(\Omega; \mathbf{y})$ .

We will next gather properties of the measure  $P_N^{y_1 \rightarrow y_{2b}}(\Omega; \mathbf{y})$ , before we continue the construction of  $\bar{P}_N$ . At this point, we know that the construction of the probability measures  $P_n^{y_1 \rightarrow y_{2b}}(\Omega; \mathbf{y})$  is complete for all  $n \leq N$ , and the construction of the probability measures  $\bar{P}_n(\Omega; \mathbf{y})$  is complete for all  $n \leq N-1$ .

<sup>16</sup>Importantly, only  $\gamma_b$  is time-reversed in this construction.



(a) We first sample points  $(U_1, U_3, U_4)$  according to (3.6). Second, given  $(U_1, U_3, U_4)$ , we sample  $\gamma_2$  from  $\phi(x_4)$  to  $\infty$  as  $\text{SLE}_\kappa(-4, 2, 2; 2, 2, -4, 2, 2, -4)$  with force points  $(U_1, \phi(x_2), \phi(x_3); \phi(x_5), \phi(x_6), U_3, \phi(x_7), \phi(x_8), U_4)$ . Note that the curve  $\gamma_2$  does not hit the intervals marked in grey, because the accumulated force point weight on these intervals exceeds  $2 \geq \kappa/2 - 2$  (which guarantees that  $\gamma_2$  does not hit these intervals). Third, we sample the curves  $\gamma_1, \gamma_3, \gamma_4$  in connected components of  $\mathbb{H} \setminus \gamma_2$  using the induction hypothesis.



(b) Denote by  $\tilde{\gamma}_2$  the time-reversal of  $\gamma_2$ . We set  $\vec{\eta}_1 := \phi^{-1}(\tilde{\gamma}_2)$ , which traverses from  $x_1$  to  $x_4$ . We set  $\vec{\eta}_2 := \phi^{-1}(\gamma_1)$ , which starts from  $x_3$ , and  $\vec{\eta}_3 := \phi^{-1}(\gamma_3)$ , which starts from  $x_5$ , and  $\vec{\eta}_4 := \phi^{-1}(\gamma_4)$ , which starts from  $x_7$ .

Figure 3.1: Example with  $N = 4$  and  $b = 2$  in Definition 3.2.

**Proposition 3.3.** Fix  $b \in \{1, 2, \dots, N\}$ . The marginal law of  $\vec{\eta}_1$  under  $\mathbb{P}_N^{x_1 \rightarrow x_{2b}}$  can be described as follows. Let  $\eta$  be the chordal  $\text{SLE}_\kappa$  on  $(\mathbb{H}; x_1, x_{2b})$ , and define the event

$$A_b(\eta) := \{\eta \cap (x_{2s}, x_{2s+1}) = \emptyset \text{ for all } 1 \leq s \leq b-1\} \cap \{\eta \cap (x_{2s-1}, x_{2s}) = \emptyset \text{ for all } b+1 \leq s \leq N\}.$$

On the event  $A_b(\eta)$ , denote by  $H_1, \dots, H_p$  the connected components of  $\mathbb{H} \setminus \eta$  whose boundaries contain points in  $\{x_1, \dots, x_{2N}\} \setminus \{x_1, x_{2b}\}$ . For each  $1 \leq m \leq p$ , denote by  $x_{j_m}, \dots, x_{j_{m+1}-1}$  the points contained in  $\partial H_m$ . Then, using the notation (1.21), the marginal law of  $\vec{\eta}_1$  under  $\mathbb{P}_N^{x_1 \rightarrow x_{2b}}$  is the same as the law of  $\eta$  weighted by

$$\frac{(x_{2b} - x_1)^{-2h(\kappa)}}{\hat{\mathcal{R}}_{\omega_b}(\mathbf{x})} \mathbb{1}\{A_b(\eta)\} \prod_{m=1}^p \frac{1}{\nu(\kappa)} \mathcal{F}_{\underbrace{\quad}_{\frac{j_{m+1}-j_m}{2}}} (H_m; x_{j_m}, \dots, x_{j_{m+1}-1}) > 0, \quad (3.8)$$

where  $\hat{\mathcal{R}}_{\omega_b}$  are the functions appearing in Lemma C.4.

We will prove Proposition 3.3 in Section 3.3. Note that  $\mathcal{F}_{\underbrace{\quad}_m} > 0$  for all  $m \in \mathbb{Z}_{>0}$  by Lemma C.2 and  $\hat{\mathcal{R}}_{\omega_b} > 0$  by Lemma C.4.

**Lemma 3.4.** Fix  $b \in \{1, 2, \dots, N\}$ . Denote by  $\varsigma(\vec{\eta})$  the random link pattern in  $\text{LP}_N$  formed by the curves  $\vec{\eta} := (\vec{\eta}_1, \dots, \vec{\eta}_N) \sim \mathbb{P}_N^{x_1 \rightarrow x_{2b}}$ . For any  $\alpha \in \text{LP}_N$  such that  $\{1, 2b\} \in \alpha$ , we have

$$\mathbb{P}_N^{x_1 \rightarrow x_{2b}}[\varsigma(\vec{\eta}) = \alpha] > 0. \quad (3.9)$$

*Proof.* We prove the claim (3.9) by induction on the number of curves. The base case is obvious<sup>17</sup>. We shall perform the induction step at the same time as we prove Lemma 3.7, using the following assumptions.

<sup>17</sup>In this case,  $\mathbb{P}_1^{x_1 \rightarrow x_2}$  equals the law of  $\text{SLE}_\kappa$  from  $x_1$  to  $x_2$ .

- The construction of the measures  $\mathbb{P}_n^{y_1 \rightarrow y_{2b}}(\Omega; \mathbf{y})$  is complete and Proposition 3.3 holds for all  $n \leq N$ .
- (Induction hypothesis): For all  $n \leq N - 1$ , we have  $\mathbb{P}_n^{x_1 \rightarrow x_{2b}}[\zeta(\vec{\eta}) = \alpha] > 0$  for all  $b \in \{1, \dots, n\}$  and for all  $\alpha \in \text{LP}_n$  such that  $\{1, 2b\} \in \alpha$ .
- The construction of the measures  $\vec{\mathbb{P}}_n(\Omega; \mathbf{y})$  is complete and Lemma 3.7 holds for all  $n \leq N - 1$ .

We will complete the proof in the course of the proof of Lemma 3.7.  $\square$

Before finishing the proofs of Proposition 3.3 and Lemma 3.4, we continue the construction of  $\vec{\mathbb{P}}_N$ . For  $\alpha \in \text{LP}_N$  such that  $\{1, 2b\} \in \alpha$ , we define  $\alpha^R \in \text{LP}_{b-1}$  to be the link pattern induced by the links of  $\alpha$  on  $\{2, 3, \dots, 2b-1\}$  nested by the link  $\{1, 2b\}$ . Using Lemma 3.4, we set

$$p_\alpha(\mathbf{x}) := \frac{\hat{\mathcal{R}}_{\omega_b}(\mathbf{x}) w_b(\alpha) \mathbb{P}_N^{x_1 \rightarrow x_{2b}}[\zeta(\vec{\eta}) = \alpha]}{\sum_{1 \leq s \leq N} \hat{\mathcal{R}}_{\omega_s}(\mathbf{x}) \sum_{\substack{\beta \in \text{LP}_N \\ \{1, 2s\} \in \beta}} w_s(\beta) \mathbb{P}_N^{x_1 \rightarrow x_{2s}}[\zeta(\vec{\eta}) = \beta]}, \quad \mathbf{x} \in \mathfrak{X}_{2N}, \quad (3.10)$$

where  $\hat{\mathcal{R}}_{\omega_s}$  are the functions appearing in Lemma C.4, and

$$w_s(\alpha) := \begin{cases} 1, & s = 1, \\ \nu(\kappa) \frac{\mathcal{M}_\nu(\underbrace{\text{---}\text{---}\text{---}}_{s-1}, \alpha^R)}{\mathcal{M}_\nu(\underbrace{\text{---}\text{---}\text{---}}_{s-1}, \alpha^R)}, & 2 \leq s \leq N. \end{cases} \quad (3.11)$$

The reason for the choice of  $w_s(\alpha)$  will become clear in Lemma 4.8. (Note that the proof of Lemma 3.4 only relies on the induction hypothesis which implies that  $p_\alpha$  is well-defined for  $\alpha \in \text{LP}_n$  with  $n \leq N - 1$ . The induction then shows that  $p_\alpha$  is well-defined for  $\alpha \in \text{LP}_N$ .)

**Lemma 3.5.** *The function  $p_\alpha$  defined in (3.10) is Möbius invariant, that is,  $p_\alpha \circ \varphi = p_\alpha$  for all Möbius maps  $\varphi$  of the upper half-plane  $\mathbb{H}$  such that  $\varphi(x_1) < \dots < \varphi(x_{2N})$ .*

*Proof.* The functions  $\mathcal{R}_{\omega_b}$  are Möbius covariant by Lemma C.3 and their covariance factors cancel out in the ratio (3.10). The other terms in (3.10) are Möbius invariant.  $\square$

**Definition 3.6.** Denote by  $\zeta(\vec{\eta})$  the random link pattern in  $\text{LP}_N$  formed by the curves  $\vec{\eta}$ . Set

$$\vec{\mathbb{P}}_N[\cdot] = \vec{\mathbb{P}}_N(\mathbb{H}; \mathbf{x})[\cdot] := \sum_{b=1}^N \sum_{\substack{\alpha \in \text{LP}_N \\ \{1, 2b\} \in \alpha}} p_\alpha(\mathbf{x}) \mathbb{P}_N^{x_1 \rightarrow x_{2b}}[\cdot \mid \zeta(\vec{\eta}) = \alpha], \quad (3.12)$$

where the weight  $p_\alpha(\mathbf{x})$  is defined in (3.10), and  $\mathbb{P}_N^{x_1 \rightarrow x_{2b}}$  is the probability measure from Definition 3.2. Since  $p_\alpha$  is Möbius invariant by Lemma 3.4, the collection  $\{\vec{\mathbb{P}}_N(\mathbb{H}; \mathbf{x}) : \mathbf{x} \in \mathfrak{X}_{2N}\}$  is Möbius invariant. Hence, it determines a conformally invariant collection  $\{\vec{\mathbb{P}}_N : (\Omega; \mathbf{y}) \text{ is a polygon}\}$  of probability measures.

**Lemma 3.7.** *Denote by  $\zeta(\vec{\eta})$  the random link pattern in  $\text{LP}_N$  formed by the curves  $\vec{\eta} \sim \vec{\mathbb{P}}_N$ . Then,*

$$\vec{\mathbb{P}}_N[\zeta(\vec{\eta}) = \alpha] > 0 \quad \text{for any } \alpha \in \text{LP}_N. \quad (3.13)$$

During the proof of Lemma 3.7 we also complete the proof of Lemma 3.4.

*Proof.* We prove the claim (3.13) by induction on the number of curves. The base case is obvious<sup>18</sup>. For the induction step, we use the following assumptions.

- The construction of the measures  $\mathbb{P}_n^{y_1 \rightarrow y_{2b}}(\Omega; \mathbf{y})$  is complete and Proposition 3.3 holds for all  $n \leq N$ .

<sup>18</sup>In this case,  $\vec{\mathbb{P}}_1^{x_1 \rightarrow x_2}$  equals the law of  $\text{SLE}_\kappa$  from  $x_1$  to  $x_2$ .



- The construction of the measures  $\vec{P}_n(\Omega; \mathbf{y})$  is complete for all  $n \leq N - 1$ .
- (Induction hypothesis): For all  $n \leq N - 1$ , the following hold:

Lemma 3.4:  $P_n^{x_1 \rightarrow x_{2b}}[\zeta(\vec{\eta}) = \alpha] > 0$  for all  $b \in \{1, \dots, n\}$  and for all  $\alpha \in \text{LP}_n$  such that  $\{1, 2b\} \in \alpha$ ;

Lemma 3.7:  $\vec{P}_n[\zeta(\vec{\eta}) = \alpha] > 0$  for all  $\alpha \in \text{LP}_n$ .

Consider the event  $E := \{\vec{\eta}_1 \cap (x_2, x_{2b-1}) = \emptyset\} \cap \{\vec{\eta}_1 \cap (x_{2b+1}, x_{2n}) = \emptyset\}$ . Note that  $P_n^{x_1 \rightarrow x_{2b}}[E] > 0$ , for otherwise, Proposition 3.3 with  $N = 3$  and the marked boundary points  $(x_1, x_2, x_{2b-1}, x_{2b}, x_{2b+1}, x_{2n})$  would imply that the expectation of (3.8) is zero. For  $\alpha \in \text{LP}_n$  such that  $\{1, 2b\} \in \alpha$ , we define  $\alpha^L \in \text{LP}_{n-b}$  to be the link pattern induced by the links of  $\alpha$  on  $\{2b+1, \dots, 2n\}$  outside of the link  $\{1, 2b\}$ . Write  $\vec{\eta}^R := (\vec{\eta}_2, \dots, \vec{\eta}_{b-1})$  and  $\vec{\eta}^L := (\vec{\eta}_{b+1}, \dots, \vec{\eta}_n)$ . By the construction in Definitions 3.2 & 3.6, conditional on  $\vec{\eta}_1$  (and on the event  $E$ ), under  $P_n^{x_1 \rightarrow x_{2b}}$  the law of  $\vec{\eta}^R$  equals  $\vec{P}_n(\Omega^R; x_2, \dots, x_{2b-1})$ , and the law of  $\vec{\eta}^L$  equals  $\vec{P}_n(\Omega^L; x_{2b+1}, \dots, x_{2n})$ , where  $\Omega^R$  (resp.  $\Omega^L$ ) is the connected component of  $\mathbb{H} \setminus \vec{\eta}_1$  containing the relevant curves  $\vec{\eta}^R$  (resp.  $\vec{\eta}^L$ ). Hence, we have

$$\begin{aligned}
& P_n^{x_1 \rightarrow x_{2b}}[\zeta(\vec{\eta}) = \alpha] \\
& \geq P_n^{x_1 \rightarrow x_{2b}} \left[ P_n^{x_1 \rightarrow x_{2b}}[\zeta(\vec{\eta}^R) = \alpha^R \text{ and } \zeta(\vec{\eta}^L) = \alpha^L \mid \vec{\eta}_1] \mathbb{1}\{E\} \right] \\
& = P_n^{x_1 \rightarrow x_{2b}} \left[ P_{b-1}^{x_1 \rightarrow x_{2b}}[\zeta(\vec{\eta}^R) = \alpha^R \mid \vec{\eta}_1] P_{n-b}^{x_1 \rightarrow x_{2b}}[\zeta(\vec{\eta}^L) = \alpha^L \mid \vec{\eta}_1] \mathbb{1}\{E\} \right] \\
& = P_n^{x_1 \rightarrow x_{2b}} \left[ \vec{P}_{b-1}(\Omega^R; x_2, \dots, x_{2b-1})[\zeta(\gamma^R) = \alpha^R] \vec{P}_{n-b}(\Omega^L; x_{2b+1}, \dots, x_{2n})[\zeta(\gamma^L) = \alpha^L] \mathbb{1}\{E\} \right] \\
& > 0,
\end{aligned}$$

where  $\gamma^R := (\gamma_2, \dots, \gamma_{b-1})$  and  $\gamma^L := (\gamma_{b+1}, \dots, \gamma_n)$ . This proves Lemma 3.4. To complete the induction step for Lemma 3.7, it just remains to note that  $\vec{P}_n[\zeta(\vec{\eta}) = \alpha] = p_\alpha(\mathbf{x}) > 0$ , by Lemma 3.4 for  $N = n$ .  $\square$

### 3.3 Marginal law: proof of Proposition 3.3

In this section, we derive the marginal law of  $\vec{\eta}_1$ . For simplicity of notation, let us assume that  $b = 1$ ; the general case can be proved similarly. Recall the definition of  $\phi$  in (3.4), with  $\phi(x_1) = \infty$ , and  $\phi(x_2) = 0$ , and  $\phi(x_3) = 1$ . Recall that in Item 5 in the construction of  $P_N^{x_1 \rightarrow x_2}$ , the curve  $\vec{\eta}_1 := \phi^{-1}(\tilde{\gamma}_1)$  is defined via the time-reversal of the  $\text{SLE}_\kappa(\rho^R)$  process  $\gamma_1$  in  $(\mathbb{H}; 0, \infty)$ . Hence, it suffices to derive the marginal law of  $\tilde{\gamma}_1$ . To this end, we will compare the law of  $\gamma_1$  to the law of the chordal  $\text{SLE}_\kappa$  curve  $\eta$  in  $(\mathbb{H}; 0, \infty)$ .

**Lemma 3.8.** *Let  $\eta$  be the chordal  $\text{SLE}_\kappa$  in  $(\mathbb{H}; 0, \infty)$ , with associated conformal maps  $g_t: \mathbb{H} \setminus K_t \rightarrow \mathbb{H}$  and driving function  $W_t = \sqrt{\kappa}B_t$  given by the standard one-dimensional Brownian motion. Fix points  $\dot{\mathbf{u}}_1 = (u_2, \dots, u_N)$  satisfying (3.5). Let  $\tau$  be the hitting time of  $\eta$  at*

$$I^{\dot{\mathbf{u}}_1} := \bigcup_{s=2}^N (\phi(x_{2s-1}), u_s),$$

and let  $T$  be the hitting time of  $\eta$  at  $(\phi(x_3), +\infty) \setminus I^{\dot{\mathbf{u}}_1}$ . Next, define

$$\begin{aligned}
M_t^{\dot{\mathbf{u}}_1} = M_t & := \mathbb{1}\{t < T \wedge \tau\} \prod_{j=3}^{2N} g_t'(\phi(x_j))^h \prod_{s=2}^N g_t'(u_s) \\
& \times |f(W_t, g_t(\phi(x_3)), g_t(\phi(x_4)), \dots, g_t(\phi(x_{2N}))); g_t(u_2), \dots, g_t(u_N)|,
\end{aligned} \tag{3.14}$$

where  $f$  is the function (3.7). Then, the following two properties hold.

- The process  $(M_t)_{t \geq 0}$  is a uniformly integrable martingale for  $\eta$ . Moreover:

1. On the event  $\{T < \tau\}$ , denote by  $\Omega_\eta$  the bounded connected component of  $\mathbb{H} \setminus \eta[0, T]$  whose boundary contains  $\phi(x_3)$ . Suppose  $\eta(T) \in (u_p, \phi(x_{2p+1}))$  for some (random)  $p \in \{2, 3, \dots, N\}$ . Choose any conformal map  $\varphi_\eta$  from  $\Omega_\eta$  onto  $\mathbb{H}$  such that  $\varphi_\eta(\eta(T)) = \infty$ . Define

$$\begin{aligned} S_T &:= \prod_{j=3}^{2p} |\varphi'_\eta(\phi(x_j))|^h \prod_{s=2}^p |\varphi'_\eta(u_s)| \\ &\quad \times \left| f(\varphi_\eta(\phi(x_3)), \varphi_\eta(\phi(x_4)), \dots, \varphi_\eta(\phi(x_{2p})); \varphi_\eta(u_2), \dots, \varphi_\eta(u_p)) \right|, \\ V_T &:= \prod_{j=2p+1}^{2N} g'_T(\phi(x_j))^h \prod_{s=p+1}^N g'_T(u_s) \\ &\quad \times \left| f(W_T, g_T(\phi(x_{2p+1})), \dots, g_T(\phi(x_{2N})); g_T(u_{p+1}), \dots, g_T(u_N)) \right|. \end{aligned}$$

Then, we have

$$M_T := \lim_{t \rightarrow T} M_t = \mathbb{1}\{T < \tau\} S_T V_T. \quad (3.15)$$

2. On the event  $\{\tau < T\}$ , we have

$$\lim_{t \rightarrow \tau} M_t = 0. \quad (3.16)$$

- Let  $\gamma^{\dot{u}^1}$  be an  $\text{SLE}_\kappa(\underline{\rho}^R)$  process in  $(\mathbb{H}; 0, \infty)$  with all force points on its right side,

$$\underline{x}^R = (\phi(x_3), \phi(x_4), u_2, \dots, \phi(x_{2N-1}), \phi(x_{2N}), u_N)$$

and with weights  $\underline{\rho}^R = (2, 2, -4, \dots, 2, 2, -4)$  (so the weights are  $-4$  for each  $u_s$  and  $2$  for each  $\phi(x_i)$ ). Then, the law of  $\eta$  weighted by  $(M_{t \wedge T \wedge \tau})_{t \geq 0}$  is the same as the law of  $\gamma^{\dot{u}^1}$ .

*Proof.* We know from [SW05, Theorem 6],  $M_t$  in (3.14) is a local martingale for  $\eta$  and the law of  $\eta$  weighted by  $(M_t)_{t \geq 0}$  is the same as the law of  $\gamma^{\dot{u}^1}$  up to time  $T \wedge \tau$ . It remains to show that  $(M_{t \wedge T \wedge \tau})_{t \geq 0}$  is uniformly integrable and to derive the terminal values (3.15) and (3.16).

**Terminal value (3.15).** Let us first assume that  $T < \tau$  and it hits the set  $(\phi(x_3), +\infty) \setminus I^{\dot{u}^1}$  at the point  $\eta(T) \in (u_p, \phi(x_{2p+1}))$  for  $p \in \{2, 3, \dots, N\}$ . Note that on the event  $\{T < \tau\}$ , the curve  $\eta$  has a positive distance from  $I^{\dot{u}^1}$  almost surely. Write  $M_t = M_{1,t} M_{2,t} M_{3,t} M_{4,t}$ , where

$$\begin{aligned} M_{1,t} &:= \prod_{j=3}^{2p} g'_t(\phi(x_j))^h \prod_{s=2}^p g'_t(u_s) \times \left| f(g_t(\phi(x_3)), g_t(\phi(x_4)), \dots, g_t(\phi(x_{2p})); g_t(u_2), \dots, g_t(u_p)) \right|, \\ M_{2,t} &:= \prod_{j=2p+1}^{2N} g'_t(\phi(x_j))^h \prod_{s=p+1}^N g'_t(u_s) \times \left| f(W_t, g_t(\phi(x_{2p+1})), \dots, g_t(\phi(x_{2N})); g_t(u_{p+1}), \dots, g_t(u_N)) \right|, \\ M_{3,t} &:= \prod_{s=2}^p \prod_{j=2p+1}^{2N} \left( \frac{(g_t(\phi(x_j)) - g_t(u_s))^2}{(g_t(\phi(x_j)) - g_t(\phi(x_{2s-1}))) (g_t(\phi(x_j)) - g_t(\phi(x_{2s})))} \right)^{-2/\kappa} \\ &\quad \times \prod_{s=2}^p \prod_{q=p+1}^N \left( \frac{(g_t(u_q) - g_t(u_s))^2}{(g_t(u_q) - g_t(\phi(x_{2s-1}))) (g_t(u_q) - g_t(\phi(x_{2s})))} \right)^{4/\kappa}, \\ M_{4,t} &:= \prod_{s=2}^p \left( \frac{(g_t(\phi(x_{2s-1})) - W_t) (g_t(\phi(x_{2s})) - W_t)}{(g_t(u_s) - W_t)^2} \right)^{2/\kappa}. \end{aligned}$$

We derive the terminal value (3.15) by considering the limits of  $M_{1,t}, \dots, M_{4,t}$  one by one.

1. We write  $M_{1,t}$  in the following form:

$$M_{1,t} = \prod_{3 \leq i < j \leq 2p} \left( \frac{g'_t(\phi(x_j)) g'_t(\phi(x_i))}{(g_t(\phi(x_j)) - g_t(\phi(x_i)))^2} \right)^{-1/\kappa} \times \prod_{\substack{3 \leq i \leq 2p \\ 2 \leq s \leq p}} \left( \frac{g'_t(\phi(x_i)) g'_t(u_s)}{(g_t(u_s) - g_t(\phi(x_i)))^2} \right)^{2/\kappa} \quad (3.17)$$

$$\times \prod_{2 \leq s < q \leq p} \left( \frac{g'_t(\phi(u_q)) g'_t(\phi(u_s))}{(g_t(\phi(u_q)) - g_t(\phi(u_s)))^2} \right)^{-4/\kappa} \times \prod_{2 \leq s \leq p} \left( \frac{g'_t(u_s)^2}{g'_t(\phi(x_{2s-1})) g'_t(\phi(x_{2s}))} \right)^{(\kappa-4)/2\kappa}.$$

Note that the quantity

$$\frac{g'_t(\phi(x_j)) g'_t(\phi(x_i))}{(g_t(\phi(x_j)) - g_t(\phi(x_i)))^2}$$

equals the Poisson kernel for the connected component of  $\mathbb{H} \setminus \eta[0, t]$  whose boundary contains  $\phi(x_j)$  and  $\phi(x_i)$ . As  $t \rightarrow T$ , it converges to the Poisson kernel  $H(\Omega_\eta; \phi(x_j), \phi(x_i))$ , so

$$\prod_{3 \leq i < j \leq 2p} \left( \frac{g'_t(\phi(x_j)) g'_t(\phi(x_i))}{(g_t(\phi(x_j)) - g_t(\phi(x_i)))^2} \right)^{-1/\kappa} \xrightarrow{t \rightarrow T} \prod_{3 \leq i < j \leq 2p} \left| \frac{\varphi'_\eta(\phi(x_j)) \varphi'_\eta(\phi(x_i))}{(\varphi_\eta(\phi(x_j)) - \varphi_\eta(\phi(x_i)))^2} \right|^{-1/\kappa},$$

and similarly,

$$\prod_{\substack{3 \leq i \leq 2p \\ 2 \leq s \leq p}} \left( \frac{g'_t(\phi(x_i)) g'_t(u_s)}{(g_t(u_s) - g_t(\phi(x_i)))^2} \right)^{2/\kappa} \xrightarrow{t \rightarrow T} \prod_{\substack{3 \leq i \leq 2p \\ 2 \leq s \leq p}} \left| \frac{\varphi'_\eta(\phi(x_i)) \varphi'_\eta(u_s)}{(\varphi_\eta(u_s) - \varphi_\eta(\phi(x_i)))^2} \right|^{2/\kappa},$$

$$\prod_{2 \leq s < q \leq p} \left( \frac{g'_t(\phi(u_q)) g'_t(\phi(u_s))}{(g_t(\phi(u_q)) - g_t(\phi(u_s)))^2} \right)^{-4/\kappa} \xrightarrow{t \rightarrow T} \prod_{2 \leq s < q \leq p} \left| \frac{\varphi'_\eta(\phi(u_q)) \varphi'_\eta(\phi(u_s))}{(\varphi_\eta(\phi(u_q)) - \varphi_\eta(\phi(u_s)))^2} \right|^{-4/\kappa}$$

Next, recall that  $g_t$  is the conformal map from the unbounded component of  $\mathbb{H} \setminus \eta[0, t]$  onto  $\mathbb{H}$  normalized at  $\infty$ . Let  $\xi_t$  to be the conformal map from the unbounded component of  $\mathbb{H} \setminus \eta[0, t]$  onto  $\mathbb{H}$  such that  $\xi_t(\phi(x_3)) = 0$ , and  $\xi_t(\phi(x_4)) = 1$ , and  $\xi_t(\eta(T)) = \infty$ ,

$$\xi_t(z) := \left( \frac{g_t(z) - g_t(\phi(x_3))}{g_t(\eta(T)) - g_t(z)} \right) \left( \frac{g_t(\eta(T)) - g_t(\phi(x_4))}{g_t(\phi(x_4)) - g_t(\phi(x_3))} \right).$$

Then, we have

$$\frac{g'_t(u_s)^2}{g'_t(\phi(x_{2s-1})) g'_t(\phi(x_{2s}))} = \frac{\xi'_t(u_s)^2}{\xi'_t(\phi(x_{2s-1})) \xi'_t(\phi(x_{2s}))} \frac{(g_t(\eta(T)) - g_t(u_s))^2}{(g_t(\eta(T)) - g_t(\phi(x_{2s-1}))) (g_t(\eta(T)) - g_t(\phi(x_{2s})))}.$$

A standard argument (for instance, see [LW21, Eq. (4.6)]) then shows that

$$\prod_{2 \leq s \leq p} \left( \frac{g'_t(u_s)^2}{g'_t(\phi(x_{2s-1})) g'_t(\phi(x_{2s}))} \right)^{(\kappa-4)/2\kappa} \xrightarrow{t \rightarrow T} \prod_{2 \leq s \leq p} \left| \frac{\varphi'_\eta(u_s)^2}{\varphi'_\eta(\phi(x_{2s-1})) \varphi'_\eta(\phi(x_{2s}))} \right|^{(\kappa-4)/2\kappa}. \quad (3.18)$$

Note that the right side of (3.18) is independent of the choice of  $\varphi_\eta$  as long as  $\varphi_\eta(\eta(T)) = \infty$ . Combining these observations, we find that, almost surely on the event  $\{T < \tau\}$ ,

$$\lim_{t \rightarrow T} M_{1,t} = S_T.$$

2. For  $M_{2,t}$ , by continuity, we have  $\lim_{t \rightarrow T} M_{2,t} = V_T$ .

3. For  $M_{3,t}$ , by continuity, we have  $\lim_{t \rightarrow T} M_{3,t} = 1$ .

4. By a standard argument (see for instance [PW19, Eq. (B.2)]), we find that  $\lim_{t \rightarrow T} M_{4,t} = 1$ .

Combining Items 1–4, we obtain (3.15) on the event  $\{T < \tau\}$ .

**Uniform integrability.** We argue next that  $\{M_{t \wedge T \wedge \tau}\}_{t \geq 0}$  is uniformly integrable. For  $\epsilon > 0$ , denote by  $\tau_\epsilon$  the hitting time of  $\eta$  at the union of the  $\epsilon$ -neighborhood of  $I^{\mathbf{u}_1}$  and  $\partial B(0, 1/\epsilon) := \{z \in \mathbb{C} : |z| = 1/\epsilon\}$ .

- For  $M_{1,t}$ , we consider the decomposition in (3.17). Because the Poisson kernel is monotone by (3.3), we see that there exists a constant  $C = C(\epsilon)$  such that for all  $t \leq \tau_\epsilon$ , we have

$$\frac{1}{C} \leq \left\{ \prod_{3 \leq i < j \leq 2p} \frac{g'_t(\phi(x_j)) g'_t(\phi(x_i))}{(g_t(\phi(x_j)) - g_t(\phi(x_i)))^2}, \prod_{\substack{3 \leq i \leq 2p \\ 2 \leq s \leq p}} \frac{g'_t(\phi(x_i)) g'_t(u_s)}{(g_t(u_s) - g_t(\phi(x_i)))^2}, \prod_{2 \leq s < q \leq p} \frac{g'_t(\phi(u_q)) g'_t(\phi(u_s))}{(g_t(\phi(u_q)) - g_t(\phi(u_s)))^2} \right\} \leq C.$$

It thus remains to bound the fourth product in (3.17). To this end, note that

$$\partial_t \log \left( \frac{g'_t(u_s)^2}{g'_t(\phi(x_{2s-1})) g'_t(\phi(x_{2s}))} \right) = \frac{-4}{(g_t(u_s) - W_t)^2} + \frac{2}{(g_t(\phi(x_{2s-1})) - W_t)^2} + \frac{2}{(g_t(\phi(x_{2s})) - W_t)^2} \geq 0.$$

Therefore, there exists a constant  $C' = C'(\epsilon)$  such that for all  $t \leq \tau_\epsilon$ , we have

$$\frac{g'_t(u_s)^2}{g'_t(\phi(x_{2s-1})) g'_t(\phi(x_{2s}))} \leq \frac{|\varphi'_\eta(u_s)|^2}{|\varphi'_\eta(\phi(x_{2s-1})) \varphi'_\eta(\phi(x_{2s}))|} \leq C'.$$

We thus see that  $|M_{1,t}|$  is bounded on the event  $\{t \leq \tau_\epsilon\}$  by a constant depending on  $\epsilon$ .

- Similarly, we find that  $|M_{2,t}|$  is bounded on the event  $\{t \leq \tau_\epsilon\}$  by a constant depending on  $\epsilon$ .
- Let us then consider  $M_{3,t}$ . For all  $2 \leq s \leq p$  and  $p+1 \leq q \leq N$ , we have

$$\left| \frac{(g_t(u_q) - g_t(u_s))^2}{(g_t(u_q) - g_t(\phi(x_{2s-1}))) (g_t(u_q) - g_t(\phi(x_{2s})))} \right| \leq 1.$$

For all  $2 \leq s \leq p$  and  $2p+1 \leq j \leq 2N$ , note that the process

$$Y_{s,j}(t) := \frac{(g_t(\phi(x_j)) - g_t(u_s))^2}{(g_t(\phi(x_j)) - g_t(\phi(x_{2s-1}))) (g_t(\phi(x_j)) - g_t(\phi(x_{2s})))}$$

satisfies

$$\partial_t \log Y_{s,j}(t) = \frac{2}{W_t - g_t(\phi(x_j))} \left( \frac{-2}{W_t - g_t(u_s)} + \frac{1}{W_t - g_t(\phi(x_{2s-1}))} + \frac{1}{W_t - g_t(\phi(x_{2s}))} \right) \geq 0,$$

which implies that

$$Y_{s,j}(t) \geq \frac{(\phi(x_j) - u_s)^2}{(\phi(x_j) - \phi(x_{2s-1})) (\phi(x_j) - \phi(x_{2s}))},$$

and we may conclude that  $|M_{3,t}|$  is bounded on the event  $\{t \leq \tau_\epsilon\}$  by a constant depending on  $\epsilon$ .

- Lastly, note that  $|M_{4,t}| \leq 1$  is bounded.

Combining these observations, we see that  $|M_t|$  is bounded on the event  $\{t \leq \tau_\epsilon\}$  by a constant depending on  $\epsilon$ . We will next take  $\epsilon \rightarrow 0$ . Denote by  $P_\epsilon$  the law of  $\eta$  weighted by  $M_{\tau_\epsilon}/M_0$ . Then, by construction,  $P_\epsilon$  is the same as the law of  $\gamma^{\dot{\mathbf{u}}_1}$  up to the hitting time at the union of  $\epsilon$ -neighborhoods of  $I^{\dot{\mathbf{u}}_1}$  and  $\partial B(0, 1/\epsilon)$ . On the one hand, as the collection  $\{P_\epsilon\}_{\epsilon>0}$  is consistent in  $\epsilon$ , by Kolmogorov's extension theorem there exists a probability measure  $P_*$  such that, for each  $\epsilon$ , the curve  $\eta$  has the law of  $\gamma^{\dot{\mathbf{u}}_1}$  up to the hitting time at the union of  $\epsilon$ -neighborhoods of  $I^{\dot{\mathbf{u}}_1}$  and  $\partial B(0, 1/\epsilon)$ . On the other hand, we know that  $\gamma^{\dot{\mathbf{u}}_1}$  has a positive distance from the intervals in  $I^{\dot{\mathbf{u}}_1}$  almost surely. Hence, the measure  $P_*$  is exactly the law of  $\gamma^{\dot{\mathbf{u}}_1}$  stopped at the hitting time at  $(\phi(x_3), +\infty) \setminus I^{\dot{\mathbf{u}}_1}$ . This implies that  $(M_{t \wedge T \wedge \tau})_{t \geq 0}$  is uniformly integrable and the law of  $\eta$  weighted by  $(M_{t \wedge T \wedge \tau})_{t \geq 0}$  is the same as the law of  $\gamma^{\dot{\mathbf{u}}_1}$  stopped at the hitting time at  $(\phi(x_3), +\infty) \setminus I^{\dot{\mathbf{u}}_1}$ .

**Terminal value (3.16).** To finish the proof, we derive the limit of  $M_t$  on the event  $\{\tau < T\}$ : since  $\gamma^{\dot{\mathbf{u}}_1}$  has a positive distance from the intervals in  $I^{\dot{\mathbf{u}}_1}$  almost surely, we have

$$\mathbb{E} \left[ \mathbb{1}_{\{\tau < T\}} \frac{M_\tau}{M_0} \right] = \mathbb{P}^{\dot{\mathbf{u}}_1}[\gamma^{\dot{\mathbf{u}}_1} \text{ hits } I^{\dot{\mathbf{u}}_1}] = 0,$$

where  $\mathbb{E}$  is the expected value under the chordal  $\text{SLE}_\kappa$  in  $(\mathbb{H}; 0, \infty)$ , and  $\mathbb{P}^{\dot{\mathbf{u}}_1}$  is the  $\text{SLE}_\kappa(\underline{\rho}^R)$  measure in  $(\mathbb{H}; 0, \infty)$ . This proves (3.16) and completes the proof.  $\square$

*Proof of Proposition 3.3.* Assume that  $\flat = 1$ . Let  $\mathbb{E}_N^{x_1 \rightarrow x_2}$  be the expected value under the measure  $\mathbb{P}_N^{x_1 \rightarrow x_2}$  from Definition 3.2. Recall that  $\vec{\eta}_1 = \phi^{-1}(\vec{\gamma}_1)$ , where the curve  $\vec{\gamma}_1$  is the time-reversal of an  $\text{SLE}_\kappa(\underline{\rho}^R)$  process  $\gamma_1$  in  $(\mathbb{H}; 0, \infty)$  with force points

$$\underline{x}^R = (\phi(x_3), \phi(x_4), U_2, \dots, \phi(x_{2N-1}), \phi(x_{2N}), U_N),$$

and weights  $\underline{\rho}^R = (2, 2, -4, \dots, 2, 2, -4)$  on its right; the points  $(U_2, U_2, \dots, U_N)$  being random, with density (3.6). Hence, for any bounded continuous function  $F$ , we have

$$\mathbb{E}_N^{x_1 \rightarrow x_2}[F(\vec{\eta}_1)] = \frac{\int_{\phi(x_4)}^{\phi(x_5)} du_2 \cdots \int_{\phi(x_{2N})}^{+\infty} du_N \frac{|f(\phi(\dot{\mathbf{x}}_1); \dot{\mathbf{u}}_1)| \mathbb{E}^{\dot{\mathbf{u}}_1}[F(\phi^{-1}(\vec{\gamma}^{\dot{\mathbf{u}}_1}))]}{\int_{\phi(x_4)}^{\phi(x_5)} dv_2 \cdots \int_{\phi(x_{2N})}^{+\infty} dv_N |f(\phi(\dot{\mathbf{x}}_1); \dot{\mathbf{v}}_1)|} \quad (3.19)$$

where  $\mathbb{E}^{\dot{\mathbf{u}}_1}$  is the expected value under the  $\text{SLE}_\kappa(\underline{\rho}^R)$  measure in  $(\mathbb{H}; 0, \infty)$ . We will compare this to the law of the chordal  $\text{SLE}_\kappa$  curve  $\eta$  in  $(\mathbb{H}; 0, \infty)$ . We denote by  $\Omega_1, \dots, \Omega_p$  the connected components of  $\mathbb{H} \setminus \eta$  whose boundaries contain points in  $\{\phi(x_3), \phi(x_4), \dots, \phi(x_{2N})\}$ , and we denote by  $\phi(x_{j_m}), \dots, \phi(x_{j_{m+1}-1})$  the points contained in  $\partial\Omega_m$ , for  $1 \leq m \leq p$ . We also define  $T_m$  to be the hitting time of  $\eta$  at  $(\phi(x_{j_{m+1}-1}), +\infty)$  and choose conformal maps  $\varphi_m$  from  $\Omega_m$  onto  $\mathbb{H}$  such that  $\varphi_m(\eta(T_m)) = \infty$ . Then, by applying Lemma 3.8 finitely many times iteratively, we obtain

$$\mathbb{E}^{\dot{\mathbf{u}}_1}[F(\phi^{-1}(\vec{\gamma}^{\dot{\mathbf{u}}_1}))] = \frac{1}{M_0^{\dot{\mathbf{u}}_1}} \mathbb{E} \left[ F(\phi^{-1}(\vec{\eta})) \mathbb{1}_{\{A^{\dot{\mathbf{u}}_1}\}} \prod_{m=1}^p R_m^{\dot{\mathbf{u}}_1} \right], \quad (3.20)$$

where  $\mathbb{E}$  is the expected value for the chordal  $\text{SLE}_\kappa$  curve  $\eta$  in  $(\mathbb{H}; 0, \infty)$ , and  $\vec{\eta}$  is its time-reversal, and  $A^{\dot{\mathbf{u}}_1}(\eta) := \{\eta \cap I^{\dot{\mathbf{u}}_1} = \emptyset\}$ , and

$$\begin{aligned} R_m^{\dot{\mathbf{u}}_1} &= \prod_{j=j_m}^{j_{m+1}-1} |\varphi'_m(\phi(x_j))|^h \prod_{s=(j_m+1)/2}^{(j_{m+1}-1)/2} |\varphi'_m(u_s)| \\ &\quad \times \left| f(\varphi_m(\phi(x_{j_m})), \dots, \varphi_m(\phi(x_{j_{m+1}-1}))); \varphi_m(u_{(j_m+1)/2}), \dots, \varphi_m(u_{(j_{m+1}-1)/2}) \right|. \end{aligned}$$

Note that  $M_0^{\dot{\mathbf{u}}_1} = |f(\phi(\dot{\mathbf{x}}_1); \dot{\mathbf{u}}_1)|$  by (3.14). Combining (3.19, 3.20), we obtain

$$\begin{aligned} \mathbb{E}_N^{x_1 \rightarrow x_2} [F(\tilde{\eta}_1)] &= \frac{\int_{\phi(x_4)}^{\phi(x_5)} dv_2 \cdots \int_{\phi(x_{2N})}^{+\infty} dv_N \mathbb{E} \left[ F(\phi^{-1}(\tilde{\eta})) \mathbb{1}\{A^{\dot{\mathbf{u}}_1}\} \prod_{m=1}^p R_m^{\dot{\mathbf{u}}_1} \right]}{\int_{\phi(x_4)}^{\phi(x_5)} dv_2 \cdots \int_{\phi(x_{2N})}^{+\infty} dv_N |f(\phi(\dot{\mathbf{x}}_1); \dot{\mathbf{v}}_1)|} \\ &= \frac{\mathbb{E} \left[ \mathbb{1}\{A_1(\eta)\} F(\phi^{-1}(\tilde{\eta})) \prod_{m=1}^p \int_{\phi(x_{j_m+1})}^{\phi(x_{j_m+2})} du_{(j_m+1)/2} \cdots \int_{\phi(x_{j_{m+1}-1})}^{\eta(T_m)} du_{(j_{m+1}-1)/2} R_m^{\dot{\mathbf{u}}_1} \right]}{\int_{\phi(x_4)}^{\phi(x_5)} dv_2 \cdots \int_{\phi(x_{2N})}^{+\infty} dv_N |f(\phi(\dot{\mathbf{x}}_1); \dot{\mathbf{v}}_1)|}, \end{aligned} \quad (3.21)$$

where

$$A_1(\eta) := \{\eta \cap (\phi(x_{2s-1}), \phi(x_{2s})) = \emptyset \text{ for all } 2 \leq s \leq N\}.$$

We will simplify the numerator and denominator of (3.21) separately.

- For the numerator, using the change of variables  $v_s = \varphi_m(u_s)$  for all  $s$ , and invoking Lemma C.2 and the conformal covariance of  $\mathcal{F}_{\underbrace{\quad}_{j_{m+1}-j_m}}$ , we see that

$$\begin{aligned} & \nu(\kappa) \hat{C}(\kappa)^{(j_{m+1}-j_m)/2} \int_{\phi(x_{j_m+1})}^{\phi(x_{j_m+2})} du_{(j_m+1)/2} \cdots \int_{\phi(x_{j_{m+1}-1})}^{\eta(T_m)} du_{(j_{m+1}-1)/2} R_m^{\dot{\mathbf{u}}_1} \\ &= \prod_{j=j_m}^{j_{m+1}-1} |\varphi'_m(\phi(x_j))|^h \times \mathcal{F}_{\underbrace{\quad}_{j_{m+1}-j_m}}(\varphi_m(\phi(x_{j_m})), \dots, \varphi_m(\phi(x_{j_{m+1}-1}))) \quad [\text{by Lemma C.2}] \\ &= \mathcal{F}_{\underbrace{\quad}_{j_{m+1}-j_m}}(\Omega_m; \phi(x_{j_m}), \dots, \phi(x_{j_{m+1}-1})), \quad [\text{by (2.7)}] \\ &= \prod_{j=j_m}^{j_{m+1}-1} |\phi'(x_j)|^{-h} \times \mathcal{F}_{\underbrace{\quad}_{j_{m+1}-j_m}}(H_m; x_{j_m}, \dots, x_{j_{m+1}-1}), \end{aligned}$$

where  $H_m = \phi^{-1}(\Omega_m)$  and  $\hat{C}(\kappa) = 4 \sin^2(4\pi/\kappa) C(\kappa)$ .

- For the denominator, we make the change of variables  $u_1 = \phi^{-1}(v_N)$  and  $u_r = \phi^{-1}(v_{r-1})$  for all  $3 \leq r \leq N$ . Then, using the identity  $\frac{\phi(z)-\phi(w)}{z-w} = \sqrt{\phi'(z)}\sqrt{\phi'(w)}$  (see, e.g., [KP16, Lemma 4.7]) for the Möbius transformation (3.4) satisfying  $\phi^{-1}(\infty) = x_1$  and  $\phi'(z) = \frac{x_2-x_1}{(z-x_1)^2} \frac{x_3-x_1}{x_3-x_2}$ , we obtain

$$\begin{aligned} & \int_{\phi(x_4)}^{\phi(x_5)} dv_2 \cdots \int_{\phi(x_{2N})}^{+\infty} dv_N |f(\phi(\dot{\mathbf{x}}_1); \dot{\mathbf{v}}_1)| \\ &= \int_{x_{2N}}^{x_1} du_1 \int_{x_4}^{x_5} du_3 \cdots \int_{x_{2N-2}}^{x_{2N-1}} du_N |f(\dot{\mathbf{x}}_1; \dot{\mathbf{u}}_2)| \prod_{\substack{1 \leq r \leq N \\ r \neq 2}} |\phi'(u_r)|^{-2h(\kappa)} \\ &= \left( \frac{(x_2-x_1)(x_3-x_1)}{x_3-x_2} \right)^{-2(N-1)h(\kappa)} \prod_{j=2}^{2N} (x_j-x_1)^{2h(\kappa)} \int_{x_{2N}}^{x_1} du_1 \int_{x_4}^{x_5} du_3 \cdots \int_{x_{2N-2}}^{x_{2N-1}} du_N |\hat{f}^a(\mathbf{x}; \dot{\mathbf{u}}_2)| \\ &= (x_2-x_1)^{2h(\kappa)} \prod_{j=3}^{2N} |\phi'(x_j)|^{-h} \times \int_{x_{2N}}^{x_1} du_1 \int_{x_4}^{x_5} du_3 \cdots \int_{x_{2N-2}}^{x_{2N-1}} du_N |\hat{f}^a(\mathbf{x}; \dot{\mathbf{u}}_2)| \\ &= \frac{(x_2-x_1)^{2h(\kappa)}}{\hat{C}(\kappa)^{N-1}} \prod_{j=3}^{2N} |\phi'(x_j)|^{-h} \times \hat{\mathcal{R}}_{\omega_1}(\mathbf{x}), \end{aligned}$$

using Lemma C.4, where  $\hat{f}^a$  defined in (C.1) has the conjugate charge at  $x_1$ .

We conclude that (3.21) equals

$$\mathbb{E}_N^{x_1 \rightarrow x_2} [F(\vec{\eta}_1)] = \frac{\mathbb{E} \left[ \mathbb{1}\{A_1(\eta)\} F(\phi^{-1}(\vec{\eta})) \prod_{m=1}^p \frac{1}{\nu(\kappa)} \mathcal{F}_{\widehat{\mathcal{A}}_{j_{m+1} - j_m}}(H_m; x_{j_m}, \dots, x_{j_{m+1} - 1}) \right]}{(x_2 - x_1)^{2h(\kappa)} \widehat{\mathcal{R}}_{\omega_1}(\mathbf{x})}. \quad (3.22)$$

The asserted formula (3.8) follows by conformal invariance and reversibility [MS16c] of the chordal  $\text{SLE}_\kappa$  curve on  $(\mathbb{H}; 0, \infty)$ : since  $\phi: \mathbb{H} \rightarrow \mathbb{H}$  sends  $x_1 \mapsto \infty$  and  $x_2 \mapsto 0$ , the curve  $\phi^{-1}(\eta)$  has the law of chordal  $\text{SLE}_\kappa$  on  $(\mathbb{H}; x_2, x_1)$ , and the time-reversal  $\phi^{-1}(\vec{\eta})$  has the law of chordal  $\text{SLE}_\kappa$  on  $(\mathbb{H}; x_1, x_2)$ .  $\square$

## 4 Multiple $\text{SLE}_\kappa$ and pure partition functions for $\kappa \in (4, 8)$

We now use the measures constructed in Section 3.2 to construct global multiple  $\text{SLE}_\kappa$  processes and their pure partition functions for  $\kappa \in (4, 8)$ . Together with prior results [PW19, Wu20], this completes the construction of the pure partition functions for the full parameter range  $\kappa \in (0, 8)$ .

Recall that  $\mathbb{P}_N^{x_1 \rightarrow x_{2b}}$  (Definition 3.2) and  $\vec{\mathbb{P}}_N$  (Definition 3.6) are conformally invariant probability measures on (oriented) curves  $\vec{\eta} := (\vec{\eta}_1, \dots, \vec{\eta}_N) \in X_0(\Omega; \mathbf{x})$  on the curve space (1.44) on polygons  $(\Omega; \mathbf{x})$ . We will mainly work with them in the reference domain  $\Omega = \mathbb{H}$ , writing  $\mathbb{P}_N^{x_1 \rightarrow x_{2b}} = \mathbb{P}_N^{x_1 \rightarrow x_{2b}}(\mathbb{H}; \mathbf{x})$  and  $\mathbb{P}_N = \mathbb{P}_N(\mathbb{H}; \mathbf{x})$ . Using the normalization factor of (3.10), set

$$\mathcal{K}_{\widehat{\mathcal{A}}_N}(\mathbf{x}) := \nu(\kappa) \sum_{b=1}^N \widehat{\mathcal{R}}_{\omega_b}(\mathbf{x}) \sum_{\substack{\beta \in \text{LP}_N \\ \{1, 2b\} \in \beta}} w_b(\beta) \mathbb{P}_N^{x_1 \rightarrow x_{2b}}[\zeta(\vec{\eta}) = \beta], \quad \mathbf{x} \in \mathfrak{X}_{2N}, \quad (4.1)$$

where  $\zeta(\vec{\eta})$  denotes the random link pattern in  $\text{LP}_N$  formed by the curves  $\vec{\eta} \sim \mathbb{P}_N^{x_1 \rightarrow x_{2b}}$ , and where  $\widehat{\mathcal{R}}_{\omega_b}$  are the functions appearing in Lemma C.4 and the weights  $w_b(\beta)$  are defined in (3.11). In fact, we have  $\mathcal{K}_{\widehat{\mathcal{A}}_N} = \mathcal{F}_{\widehat{\mathcal{A}}_N}$ , see Lemma 4.4. To prove this, we need key properties of the Coulomb gas integral functions  $\mathcal{F}_{\widehat{\mathcal{A}}_N}$ : they are uniquely characterized as solutions to the PDE boundary value problem in the space  $\mathcal{S}_N$  (2.21) by their asymptotic properties ( $\mathcal{F}$ -ASY) in Corollary 2.13.

**Definition 4.1.** Fix  $\kappa \in (4, 8)$ . We define the the *pure partition function* of multiple  $\text{SLE}_\kappa$  by

$$\mathcal{Z}_\alpha(\mathbf{x}) = \vec{\mathcal{Z}}_\alpha(\mathbf{x}) := \mathcal{K}_{\widehat{\mathcal{A}}_N}(\mathbf{x}) \frac{\vec{\mathbb{P}}_N[\zeta(\vec{\eta}) = \alpha]}{\mathcal{M}_\nu(\widehat{\mathcal{A}}_N, \alpha)}, \quad \mathbf{x} \in \mathfrak{X}_{2N}, \alpha \in \text{LP}_N. \quad (4.2)$$

We define analogously

$$\vec{\mathcal{Z}}_\alpha(\mathbf{x}) := \mathcal{K}_{\widehat{\mathcal{A}}_N}(\mathbf{x}) \frac{\vec{\mathbb{P}}_N[\zeta(\vec{\eta}) = \alpha]}{\mathcal{M}_\nu(\widehat{\mathcal{A}}_N, \alpha)}, \quad \mathbf{x} \in \mathfrak{X}_{2N}, \alpha \in \text{LP}_N. \quad (4.3)$$

Recall that, under the measure  $\vec{\mathbb{P}}_N$ , each curve  $\vec{\eta}_s$  starts from  $x_{2s}$  for  $1 \leq s \leq N$  (while under the measure  $\vec{\mathbb{P}}_N$ , each curve  $\vec{\eta}_s$  starts from  $x_{2s-1}$  for  $1 \leq s \leq N$ ). We will prove that the two functions (4.2) and (4.3) are in fact the same (Lemma 4.3). This is a feature of reversibility for multiple  $\text{SLE}_\kappa$  curves.

**Remark 4.2.** When  $N = 1$ , we have  $\mathcal{Z}_\triangle(x_1, x_2) = (x_2 - x_1)^{-2h}$ . Further examples are discussed in Appendix D.

The function  $\mathcal{K}_{\widehat{\mathcal{A}}_N}(\mathbf{x})$  is defined in (4.1) in terms of the probability measure  $\mathbb{P}_N^{x_1 \rightarrow x_{2b}}$ . By definition of  $\vec{\mathbb{P}}_N$  in (3.12), and similarly for  $\vec{\mathcal{Z}}$ , we obtain the factorization formula

$$\vec{\mathcal{Z}}_\alpha(\mathbf{x}) = \nu(\kappa) \widehat{\mathcal{R}}_{\omega_b}(\mathbf{x}) \frac{w_b(\alpha) \mathbb{P}_N^{x_1 \rightarrow x_{2b}}[\zeta(\vec{\eta}) = \alpha]}{\mathcal{M}_\nu(\widehat{\mathcal{A}}_N, \alpha)}, \quad \{1, 2b\} \in \alpha. \quad (4.4)$$

The goal of this section is to prove that  $\{\vec{\mathcal{Z}}_\alpha: \alpha \in \text{LP}_N\}$  indeed are the *pure* partition functions of multiple  $\text{SLE}_\kappa$ , that is, they are positive (Lemma 4.10) and satisfy the PDE system (PDE) (see Proposition 4.12), Möbius covariance (COV) (see Lemma 4.5), the power-law bound in (PLB) (see Lemma 4.13), as well as the recursive asymptotics property (ASY) (see Proposition 4.16). These imply the existence part in Theorem 1.10. The uniqueness part in Theorem 1.10 is already well known by results of Flores and Kleban (see Lemma 2.11). We verify the additional properties in Theorem 1.10 in Section 5.

Before proceeding to the proofs, let us make a few remarks regarding the scope of these results.

- The BPZ PDE system (PDE) is a linear hypoelliptic system, for which smoothness of solutions can be derived from Hörmander’s theory (cf. [Bon69, Hör90, Dub15b, PW19]) — see Appendix A.
- The Möbius covariance (COV) can be verified immediately from the definitions, as the involved functions are Möbius invariant/covariant (Lemma 4.5). We also prove a stronger version in Lemma 4.9.
- The power-law bound (PLB) is immediate (Lemma 4.13). Positivity needs an argument (Lemma 4.10).
- The recursive asymptotics property (ASY) of the pure partition functions (addressed in Proposition 4.16) is not only their most important, characterizing property, but is also rather difficult to verify in the case where  $\kappa \in (6, 8)$ . Indeed, the proof in [Wu20, Section 6] fails for this range, as also discussed in [Pel19, Appendix B]. The main reason for this is that it is hard to establish sufficient control for exchanging the limit in the asymptotics in (ASY) with the expected value in the cascade relation in (4.7) in Lemma 4.7. We are not aware of any proof for the asymptotics property (ASY) of the pure partition functions for  $\kappa \in (6, 8) \cap \mathbb{Q}$  before the present work. In the case where  $\kappa \in (6, 8) \setminus \mathbb{Q}$ , the asymptotics property (ASY) can be verified using the Coulomb gas formalism [FK15d, KP16, KP20], which however fails to ensure positivity of the functions (crucial for their probabilistic meaning). Therefore, it seems to us that both the probabilistic arguments and the Coulomb gas are necessary.

#### 4.1 First properties of pure partition functions

To prove the key properties of the pure partition functions, we proceed by induction on the number  $N \geq 1$  of curves. The induction step will require a number of additional properties, gathered in the next lemmas.

**Lemma 4.3.** *We have  $\vec{\mathcal{Z}}_\alpha = \tilde{\mathcal{Z}}_\alpha$  for all  $\alpha \in \text{LP}_N$  and  $N \geq 1$ .*

*Proof.* We prove the claim by induction on the number of curves in the course of this section. In the base case  $N = 1$ , we have  $\vec{\mathcal{Z}}_\alpha(x_1, x_2) = \tilde{\mathcal{Z}}_\alpha(x_1, x_2) = (x_2 - x_1)^{-2h}$ . The proof is completed in Section 4.7.  $\square$

**Lemma 4.4.** *Fix  $N \geq 1$ . We have*

$$\mathcal{Z}_\alpha(\mathbf{x}) = \vec{\mathcal{Z}}_\alpha(\mathbf{x}) = \mathcal{F}_{\text{arc}_N}(\mathbf{x}) \frac{\vec{\mathbb{P}}_N[\varsigma(\vec{\eta}) = \alpha]}{\mathcal{M}_\nu(\text{arc}_N, \alpha)}, \quad \mathbf{x} \in \mathfrak{X}_{2N}, \alpha \in \text{LP}_N. \quad (4.5)$$

Note that the formula (4.5) is equivalent to the identity  $\mathcal{K}_{\text{arc}_N} = \mathcal{F}_{\text{arc}_N}$  in (4.1).

*Proof.* We prove this by induction on the number of curves in the course of this section. In the base case  $N = 1$ , we have  $\mathcal{F}_{\text{arc}_1}(x_1, x_2) = \nu(\kappa)(x_2 - x_1)^{-2h} = \mathcal{K}_{\text{arc}_1}(x_1, x_2)$ . The proof is completed in Section 4.5.  $\square$

**Induction hypothesis.** Fix  $N \geq 2$ . From now on, suppose that for all  $n \leq N - 1$ , the following hold:

Lemma 4.3:  $\vec{\mathcal{Z}}_\alpha = \tilde{\mathcal{Z}}_\alpha$  for all  $\alpha \in \text{LP}_n$  and  $n \leq N - 1$ ;

Lemma 4.4:  $\mathcal{K}_{\text{arc}_n} = \mathcal{F}_{\text{arc}_n}$  for all  $n \leq N - 1$ .

Given this induction hypothesis, we will prove Lemmas 4.3 and 4.4, as well as other salient properties of the functions  $\mathcal{Z}_\alpha$ . We begin with a fundamental “cascade property” in the next section.



## 4.2 Cascade property

The ‘‘cascade property’’ of the pure partition functions on polygons (Lemma 4.7) plays a crucial role not only in the verification of the defining properties of the pure partition functions, but also in applications of the pure partition functions in various problems in random geometry. We will prove it by induction on the number of curves, using also the induction hypothesis on Lemmas 4.3 and 4.4.

**Lemma 4.5.** *Fix  $\kappa \in (4, 8)$ . The collection  $\{\vec{Z}_\alpha : \alpha \in \text{LP}_N\}$  of Definition 4.1 satisfies (COV).*

*Proof.* The Möbius covariance property (COV) follows immediately from the conformal invariance of  $P_N^{x_1 \rightarrow x_{2b}}$  with the conformal covariance of  $\hat{\mathcal{R}}_\omega$ , from Lemma C.3, together with the factorization (4.4).  $\square$

**Remark 4.6.** We extend the definition of any partition function  $\vec{Z}_\alpha(\Omega; \mathbf{x})$  to general nice polygons  $(\Omega; \mathbf{x})$  by sending  $\Omega$  onto  $\mathbb{H}$  via any conformal map  $\varphi$  with  $\varphi(x_1) < \dots < \varphi(x_{2N})$ , using the conformal covariance

$$\vec{Z}_\alpha(\Omega; \mathbf{x}) := \prod_{j=1}^{2N} |\varphi'(x_j)|^{h(\kappa)} \times \vec{Z}_\alpha(\Omega; \varphi(\mathbf{x})). \quad (4.6)$$

Thanks to (COV) in Lemma 4.5, this definition is independent of the choice of  $\varphi$ .

**Lemma 4.7.** *Fix  $\kappa \in (4, 8)$ . Fix  $N \geq 1$ , a nice polygon  $(\Omega; \mathbf{x})$ , and a link pattern  $\alpha = \{\{a_1, b_1\}, \dots, \{a_N, b_N\}\}$  as in (1.2). Fix a link  $\{a, b\} \in \alpha$ . Let  $\eta$  be the chordal SLE $_\kappa$  on  $(\Omega; x_a, x_b)$ . Then, we have*

$$\vec{Z}_\alpha(\Omega; \mathbf{x}) = \vec{Z}_{\alpha \setminus \{a, b\}}(\Omega; x_a, x_b) \mathbb{E}(\Omega; x_a, x_b) [\mathbb{1}\{\mathcal{E}_\alpha(\eta)\} \vec{Z}_{\alpha/\{a, b\}}(\hat{\Omega}_\eta; \ddot{\mathbf{x}}_{a, b})], \quad (4.7)$$

where  $\mathbb{E}(\Omega; x_a, x_b)$  is the expected value under the chordal SLE $_\kappa$  in  $(\Omega; x_a, x_b)$ , and  $\mathcal{E}_\alpha(\eta)$  is the event that the curve  $\eta$  does not partition  $\Omega$  into components where some variables corresponding to a link in  $\alpha$  would belong to different components,  $\hat{\Omega}_\eta$  is the union of those connected components (c.c)  $D$  of  $\Omega \setminus \eta$  that contain some of the points  $\{x_1, \dots, x_{2N}\} \setminus \{x_a, x_b\}$  in  $\bar{D}$ ,

$$\hat{\Omega}_\eta := \bigsqcup_{\substack{D \text{ c.c of } \Omega \setminus \eta \\ \bar{D} \cap \{x_1, \dots, x_{2N}\} \setminus \{x_a, x_b\} \neq \emptyset}} D,$$

and  $\ddot{\mathbf{x}}_{a, b} = (x_1, \dots, x_{a-1}, x_{a+1}, \dots, x_{b-1}, x_{b+1}, \dots, x_{2N})$  are the remaining marked points and

$$\vec{Z}_{\alpha/\{a, b\}}(\hat{\Omega}_\eta; \ddot{\mathbf{x}}_{a, b}) := \prod_{\substack{D \text{ c.c of } \Omega \setminus \eta \\ \bar{D} \cap \{x_1, \dots, x_{2N}\} \setminus \{x_a, x_b\} \neq \emptyset}} \vec{Z}_{\alpha^D}(D; \dots), \quad (4.8)$$

with  $\alpha^D$  the sub-link pattern of  $\alpha/\{a, b\}$  associated to the component  $D \subset \hat{\Omega}_\eta$  and  $\vec{Z}_{\alpha^D}(D; \dots)$  the pure partition function in the polygon  $(D; \dots)$  with some of the points among  $\{x_1, \dots, x_{2N}\} \setminus \{x_a, x_b\}$  on  $\partial D$ .

Note that  $(\hat{\Omega}_\eta; \ddot{\mathbf{x}}_{a, b})$  is not necessarily a polygon, but it can be thought of as a union of polygons obtained from the different connected components of  $\Omega \setminus \eta$ . The function  $\vec{Z}_{\alpha/\{a, b\}}(\hat{\Omega}_\eta; \ddot{\mathbf{x}}_{a, b})$  is a partition function for a domain with multiple components, simply defined as the product of the partition functions on the various connected components. This is a feature of the domain Markov property of SLE $_\kappa$ , which also results in the fundamental cascade property (4.7) of the pure partition functions.

Before we prove Lemma 4.7, we first collect some combinatorial facts about the meander matrix and the weights  $w_s(\alpha)$  defined in (3.11).

**Lemma 4.8.** *The meander matrix  $\{\mathcal{M}_\nu(\alpha, \beta) : \alpha, \beta \in \text{LP}_N\}$  defined in (1.3) satisfies the following factorization properties. For any  $\alpha \in \text{LP}_N$  such that  $\{1, 2b\} \in \alpha$ , we have*

$$\mathcal{M}_\nu(\underbrace{\text{---}}_N, \alpha) = \begin{cases} \mathcal{M}_\nu(\underbrace{\text{---}}_{N-1}, \alpha^L), & b = 1, \\ \mathcal{M}_\nu(\underbrace{\text{---}}_{b-1}, \alpha^R) \mathcal{M}_\nu(\underbrace{\text{---}}_{N-b}, \alpha^L), & 2 \leq b \leq N-1, \\ \nu(\kappa) \mathcal{M}_\nu(\underbrace{\text{---}}_{N-1}, \alpha^R), & b = N, \end{cases} \quad (4.9)$$

where  $\alpha^L \in \text{LP}_{N-b}$  is the link pattern induced by the links of  $\alpha$  on  $\{2b+1, \dots, 2N\}$  outside of the link  $\{1, 2b\}$ , and  $\alpha^R \in \text{LP}_{b-1}$  the link pattern induced by the links of  $\alpha$  on  $\{2, 3, \dots, 2b-1\}$  nested by  $\{1, 2b\}$ . Combining (4.9) with the definition (3.11) of  $w_s(\alpha)$  shows that

- if  $b = 1$ , we have

$$w_b(\alpha) \frac{\nu(\kappa)}{\mathcal{M}_\nu(\underbrace{\text{---}}_N, \alpha)} = \frac{\nu(\kappa)}{\mathcal{M}_\nu(\underbrace{\text{---}}_{N-b}, \alpha^L)}; \quad (4.10)$$

- if  $2 \leq b \leq N-1$ , we have

$$w_b(\alpha) \frac{\nu(\kappa)}{\mathcal{M}_\nu(\underbrace{\text{---}}_N, \alpha)} = \frac{\nu(\kappa)}{\mathcal{M}_\nu(\underbrace{\text{---}}_{b-1}, \alpha^R)} \frac{\nu(\kappa)}{\mathcal{M}_\nu(\underbrace{\text{---}}_{N-b}, \alpha^L)}; \quad (4.11)$$

- if  $b = N$ , we have

$$w_b(\alpha) \frac{\nu(\kappa)}{\mathcal{M}_\nu(\underbrace{\text{---}}_N, \alpha)} = \frac{\nu(\kappa)}{\mathcal{M}_\nu(\underbrace{\text{---}}_{b-1}, \alpha^R)}. \quad (4.12)$$


*Proof.* The asserted properties are straightforward to check — see Figure 4.1.  $\square$

*Proof of Lemma 4.7.* Without loss of generality, we assume that  $\Omega = \mathbb{H}$  — the formula for general polygons follows by Möbius covariance (Lemma 4.5). Let  $\{1, 2b\} \in \alpha$ . Let us assume that  $2 \leq b \leq N-1$ ; the other two cases can be proved similarly. On the one hand, by the construction in Definitions 3.2 & 3.6, and by Proposition 3.3, using the notation from Items 3–4 in Definition 3.2, we have


$$\begin{aligned} \mathbb{P}_N^{x_1 \rightarrow x_{2b}}[\varsigma(\vec{\eta}) = \alpha] &= \mathbb{P}_N^{x_1 \rightarrow x_{2b}} \left[ \left( \prod_{m=1}^r \tilde{\mathbb{P}}_{s_{m+1}^R - s_m^R}[\varsigma(\gamma_m^R) = \alpha_m^R] \right) \left( \prod_{m=1}^l \tilde{\mathbb{P}}_{s_{m+1}^L - s_m^L}[\varsigma(\gamma_m^L) = \alpha_m^L] \right) \right] \\ &= \frac{(x_{2b} - x_1)^{-2h}}{\hat{\mathcal{R}}_{\omega_b}(\mathbf{x})} \mathbb{E}(\mathbb{H}; x_1, x_{2b}) \left[ \mathbb{1}\{\mathcal{E}_\alpha(\eta)\} \left( \prod_{m=1}^r \frac{\tilde{S}_m^R}{\nu(\kappa)} \right) \left( \prod_{m=1}^l \frac{\tilde{S}_m^L}{\nu(\kappa)} \right) \right] \\ &= \frac{\vec{\mathcal{Z}}_{\triangleleft}(\mathbb{H}; x_1, x_{2b})}{\hat{\mathcal{R}}_{\omega_b}(\mathbf{x})} \mathbb{E}(\mathbb{H}; x_1, x_{2b}) \left[ \mathbb{1}\{\mathcal{E}_\alpha(\eta)\} \left( \prod_{m=1}^r \frac{\tilde{S}_m^R}{\nu(\kappa)} \right) \left( \prod_{m=1}^l \frac{\tilde{S}_m^L}{\nu(\kappa)} \right) \right], \end{aligned} \quad (4.13)$$

where  $\{\alpha_1^R, \dots, \alpha_r^R, \alpha_1^L, \dots, \alpha_l^L\} = \{\alpha^D : D \text{ c.c. of } \mathbb{H} \setminus \eta\}$ , and where  $\mathbb{E}(\mathbb{H}; x_1, x_{2b})$  is the expectation under the chordal  $\text{SLE}_\kappa$  in  $(\mathbb{H}; x_1, x_{2b})$ , and we write


$$\begin{aligned} \tilde{\mathbb{P}}_{s_{m+1}^R - s_m^R}(H_m^R; x_{2s_m^R}, \dots, x_{2s_{m+1}^R-1})[\varsigma(\gamma_m^R) = \alpha_m^R] &= \frac{\tilde{S}_m^R}{\mathcal{F}_{\underbrace{\text{---}}_{s_{m+1}^R - s_m^R}}(H_m^R; x_{2s_m^R}, \dots, x_{2s_{m+1}^R-1})}, \\ \tilde{\mathbb{P}}_{s_{m+1}^L - s_m^L}(H_m^L; x_{2s_m^L+1}, \dots, x_{2s_{m+1}^L})[\varsigma(\gamma_m^L) = \alpha_m^L] &= \frac{\tilde{S}_m^L}{\mathcal{F}_{\underbrace{\text{---}}_{s_{m+1}^L - s_m^L}}(H_m^L; x_{2s_m^L+1}, \dots, x_{2s_{m+1}^L})}, \end{aligned}$$



(a) Case:  $b = 1$ .



(b) Case:  $2 \leq b \leq N - 1$ .



(c) Case:  $b = N$ .

Figure 4.1: Illustration of the property (4.9) in Lemma 4.8.

with  $H_m^R = \phi^{-1}(\Omega_m^R)$  and  $H_m^L = \phi^{-1}(\Omega_m^L)$ , where  $\phi$  is the Möbius transformation (3.4) (depending on  $\mathbf{x}$ ). On the other hand, using the factorization formula

$$\begin{aligned} \frac{\nu(\kappa)}{\mathcal{M}_\nu(\underbrace{\text{arc}_{b-1}}_{\text{arc}_{b-1}}, \alpha^R)} &= \prod_{1 \leq m \leq r} \frac{\nu(\kappa)}{\mathcal{M}_\nu(\underbrace{\text{arc}_{s_{m+1}^R - s_m^R}}_{\text{arc}_{s_{m+1}^R - s_m^R}}, \alpha_m^R)}, \\ \frac{\nu(\kappa)}{\mathcal{M}_\nu(\underbrace{\text{arc}_{N-b}}_{\text{arc}_{N-b}}, \alpha^L)} &= \prod_{1 \leq m \leq l} \frac{\nu(\kappa)}{\mathcal{M}_\nu(\underbrace{\text{arc}_{s_{m+1}^L - s_m^L}}_{\text{arc}_{s_{m+1}^L - s_m^L}}, \alpha_m^L)}. \end{aligned} \quad (4.14)$$

(which is easy to verify, see Figure 4.2 for an illustration), we obtain

$$\begin{aligned} \vec{Z}_\alpha(\mathbb{H}; \mathbf{x}) &= w_b(\alpha) \frac{\nu(\kappa)}{\mathcal{M}_\nu(\underbrace{\text{arc}_N}_{\text{arc}_N}, \alpha)} \hat{\mathcal{R}}_{\omega_b}(\mathbf{x}) \mathbb{P}_N^{x_1 \rightarrow x_{2b}}[\zeta(\vec{\eta}) = \alpha] && \text{[by (4.4)]} \\ &= \frac{\nu(\kappa)}{\mathcal{M}_\nu(\underbrace{\text{arc}_{b-1}}_{\text{arc}_{b-1}}, \alpha^R)} \frac{\nu(\kappa)}{\mathcal{M}_\nu(\underbrace{\text{arc}_{N-b}}_{\text{arc}_{N-b}}, \alpha^L)} \hat{\mathcal{R}}_{\omega_b}(\mathbf{x}) \mathbb{P}_N^{x_1 \rightarrow x_{2b}}[\zeta(\vec{\eta}) = \alpha] && \text{[by (4.11)]} \end{aligned}$$

$$= \vec{\mathcal{Z}}_{\triangleleft}(\mathbb{H}; x_1, x_{2b}) \mathbb{E}(\mathbb{H}; x_1, x_{2b}) \left[ \mathbb{1}\{\mathcal{E}_\alpha(\eta)\} \left( \prod_{m=1}^r \vec{\mathcal{Z}}_{\alpha_m^R} \right) \left( \prod_{m=1}^l \vec{\mathcal{Z}}_{\alpha_m^L} \right) \right],$$

where to derive the last equality, we combine (4.13) and (4.14) and using the induction hypothesis that Lemma 4.4 holds for all  $n \leq N - 1$ , writing

$$\begin{aligned} \vec{\mathcal{Z}}_{\alpha_m^R} &= \vec{\mathcal{Z}}_{\alpha_m^R}(H_m^R; x_{2s_m^R}, \dots, x_{2s_{m+1}^R-1}) = \frac{\vec{S}_m^R}{\mathcal{M}_\nu(\underbrace{\quad}_{s_{m+1}^R - s_m^R}, \alpha_m^R)} && \text{[by (4.5)]} \\ \vec{\mathcal{Z}}_{\alpha_m^L} &= \vec{\mathcal{Z}}_{\alpha_m^L}(H_m^L; x_{2s_m^L}, \dots, x_{2s_{m+1}^L-1}) = \frac{\vec{S}_m^L}{\mathcal{M}_\nu(\underbrace{\quad}_{s_{m+1}^L - s_m^L}, \alpha_m^L)}. && \text{[by (4.5)].} \end{aligned}$$

Lastly, invoking the induction hypothesis that the reversibility Lemma 4.3 holds for all  $n \leq N - 1$ , we have  $\vec{\mathcal{Z}}_{\alpha_m^R} = \vec{\mathcal{Z}}_{\alpha_m^L}$ , so we conclude that the asserted formula (4.7) with  $b = 2b$  holds. Induction on the number of curves and the same arguments as in the proof of [Wu20, Lemma 6.2] or [Pel19, Proposition B.1] then show that the cascade formula (4.7) also holds for the other links  $\{a, b\} \in \alpha$ .  $\square$




Figure 4.2: Illustration of the relation (4.14).

### 4.3 General Möbius covariance

We next strengthen the Möbius covariance (Lemma 4.9) into a covariance property under any Möbius map  $\varphi \in \text{PSL}(2, \mathbb{R})$  preserving the upper half-plane, which may rotate the marked boundary points.

**Lemma 4.9.** *Fix  $\kappa \in (4, 8)$  and  $\alpha \in \text{LP}_N$ . The function  $\vec{\mathcal{Z}}_\alpha$  of Definition 4.1 satisfies the Möbius covariance*

$$\vec{\mathcal{Z}}_\alpha(\mathbf{x}) = \prod_{i=1}^{2N} |\varphi'(x_i)|^h \times \vec{\mathcal{Z}}_{\varphi(\alpha)}(\varphi(\mathbf{x})), \quad \mathbf{x} \in \mathfrak{X}_{2N}, \quad (4.15)$$

for all Möbius maps  $\varphi \in \text{PSL}(2, \mathbb{R})$ , where  $\varphi(\mathbf{x}) := (\varphi(x_{j+1}), \dots, \varphi(x_{2N}), \varphi(x_1), \dots, \varphi(x_j))$ , so that

$$\varphi(x_{j+1}) < \varphi(x_{j+2}) < \dots < \varphi(x_{2N}) < \varphi(x_1) < \varphi(x_2) < \dots < \varphi(x_j)$$

with some  $j \in \{0, 1, 2, \dots, 2N - 1\}$ , and where  $\varphi(\alpha) \in \text{LP}_N$  is the link pattern obtained from  $\alpha$  via permuting the indices according to the permutation of the boundary points induced by  $\varphi$  and reordering.

*Proof.* We prove (4.15) by induction on  $N \geq 1$ . The base case  $N = 1$  is clear for  $\vec{\mathcal{Z}}_{\triangleleft}(x_1, x_2) = (x_2 - x_1)^{-2h}$ . Assuming that (4.15) holds for the collection  $\{\vec{\mathcal{Z}}_\alpha : \alpha \in \text{LP}_n, n \leq N - 1\}$  and under

the induction hypothesis that Lemmas 4.3 and 4.4 hold for all  $n \leq N - 1$ , we may apply the cascade property (4.7) from Lemma 4.7 to obtain

$$\begin{aligned} \vec{Z}_{\varphi(\alpha)}(\varphi(\mathbf{x})) &= (\varphi(x_{2b}) - \varphi(x_1))^{-2h} \mathbb{E}(\mathbb{H}; \varphi(x_1), \varphi(x_{2b})) [\mathbb{1}\{\mathcal{E}_{\varphi(\alpha)}(\eta)\} \vec{Z}_{\varphi(\alpha/\{1,2b\})}(\varphi(\hat{\mathbb{H}}_\eta); \varphi(\hat{\mathbf{x}}_{1,2b}))] \\ &= \prod_{i=1}^{2N} |\varphi'(x_i)|^{-h} \times (x_{2b} - x_1)^{-2h} \mathbb{E}(\mathbb{H}; x_1, x_{2b}) [\mathbb{1}\{\mathcal{E}_\alpha(\eta)\} \vec{Z}_{\alpha/\{1,2b\}}(\hat{\mathbb{H}}_\eta; \hat{\mathbf{x}}_{1,2b})] \\ &= \prod_{i=1}^{2N} |\varphi'(x_i)|^{-h} \times \vec{Z}_\alpha(\mathbf{x}), \quad \alpha \in \text{LP}_N, \{1, 2b\} \in \alpha, \end{aligned}$$

where  $\mathbb{E}(\mathbb{H}; x_1, x_{2b})$  is the expected value under the chordal  $\text{SLE}_\kappa$  in  $(\mathbb{H}; x_1, x_{2b})$ .  $\square$

#### 4.4 Smoothness and hypoelliptic BPZ PDEs

We first verify the continuity and positivity of the functions  $\vec{Z}_\alpha: \mathfrak{X}_{2N} \rightarrow (0, \infty)$  (Lemma 4.10). We will promote the continuity to smoothness by hypoellipticity arguments.

**Lemma 4.10.** *The functions  $\vec{Z}_\alpha: \mathfrak{X}_{2N} \rightarrow (0, \infty)$  are positive and continuous:*

$$\lim_{\substack{y_j \rightarrow x_j \\ \forall 1 \leq j \leq 2N}} \vec{Z}_\alpha(\mathbf{y}) = \vec{Z}_\alpha(\mathbf{x}) \quad \text{for any } \alpha \in \text{LP}_N \text{ and } \mathbf{x} \in \mathfrak{X}_{2N}. \quad (4.16)$$

*Proof.* By Definition 4.1, we have

$$\begin{aligned} \vec{Z}_\alpha(\mathbf{x}) &= \mathcal{K}_{\text{arc}_N}(\mathbf{x}) \frac{\bar{\mathbb{P}}_N[\varsigma(\vec{\eta}) = \alpha]}{\mathcal{M}_\nu(\text{arc}_N, \alpha)}, \quad \alpha \in \text{LP}_N, \\ \mathcal{K}_{\text{arc}_N}(\mathbf{x}) &= \nu(\kappa) \sum_{s=1}^N \hat{\mathcal{R}}_{\omega_s}(\mathbf{x}) \sum_{\substack{\beta \in \text{LP}_N \\ \{1, 2s\} \in \beta}} w_s(\beta) \mathbb{P}_N^{x_1 \rightarrow x_{2s}}[\varsigma(\vec{\eta}) = \beta], \quad \mathbf{x} \in \mathfrak{X}_{2N}. \end{aligned}$$

Now, the constants  $\nu(\kappa) > 0$  and  $w_s(\beta) > 0$  are positive in the current range  $\kappa \in (4, 8)$ , and the functions  $\hat{\mathcal{R}}_{\omega_s}(\mathbf{x}) > 0$  are positive on  $\mathfrak{X}_{2N} \ni \mathbf{x}$  by Lemma C.4. Moreover, we have  $\bar{\mathbb{P}}_N[\varsigma(\vec{\eta}) = \alpha] > 0$  and  $\mathbb{P}_N^{x_1 \rightarrow x_{2s}}[\varsigma(\vec{\eta}) = \beta] > 0$  for every  $\alpha \in \text{LP}_N$  and  $\mathbf{x} \in \mathfrak{X}_{2N}$  respectively by Lemmas 3.7 and 3.4. Hence, we have  $\mathcal{K}_{\text{arc}_N}(\mathbf{x}) > 0$ , which implies that  $\vec{Z}_\alpha(\mathbf{x}) > 0$ , for all  $\mathbf{x} \in \mathfrak{X}_{2N}$ .

We prove the continuity (4.16) by induction on  $N \geq 1$ . The base case  $\vec{Z}_\alpha(x_1, x_2) = (x_2 - x_1)^{-2h}$  is clear. Assuming that all functions in the collection  $\{\vec{Z}_\alpha: \alpha \in \text{LP}_n, n \leq N - 1\}$  are continuous on  $\mathfrak{X}_{2N}$ , we use the cascade property (Lemma 4.7) to perform the induction step. Fix  $\alpha \in \text{LP}_N$  and suppose  $\{1, 2\} \in \alpha$ , without loss of generality by Lemma 4.9. Let  $\varphi_{\mathbf{y}}$  be the Möbius transformation such that  $\varphi_{\mathbf{y}}(y_1) = x_1$ ,  $\varphi_{\mathbf{y}}(y_2) = x_2$ , and  $\varphi_{\mathbf{y}}(\infty) = \infty$ . Denote  $\tilde{\mathbf{x}} = \varphi_{\mathbf{y}}(\mathbf{y})$ . Note that if  $y_j \rightarrow x_j$  for all  $1 \leq j \leq 2N$ , then  $\varphi_{\mathbf{y}}(z) - z \rightarrow 0$  uniformly on  $\bar{\mathbb{H}}$ , and in particular  $\varphi'_{\mathbf{y}}(z) \rightarrow 1$  on  $\bar{\mathbb{H}}$ . Thus, by the Möbius covariance in Lemma 4.5, it suffices to prove that

$$\lim_{\substack{\tilde{x}_j \rightarrow x_j \\ \forall 3 \leq j \leq 2N}} \vec{Z}(\tilde{\mathbf{x}}) = \vec{Z}_\alpha(\mathbf{x}). \quad (4.17)$$

To this end, using the cascade property (4.7) from Lemma 4.7 and Fatou's lemma, we have

$$\liminf_{\substack{\tilde{x}_j \rightarrow x_j \\ \forall 3 \leq j \leq 2N}} \vec{Z}(\tilde{\mathbf{x}}) \geq (x_2 - x_1)^{-2h} \mathbb{E}(\mathbb{H}; x_1, x_2) \left[ \liminf_{\substack{\tilde{x}_j \rightarrow x_j \\ \forall 3 \leq j \leq 2N}} \mathbb{1}\{\mathcal{E}_\alpha(\eta)\} \vec{Z}_{\alpha/\{1,2\}}(\tilde{\Omega}_\eta; \tilde{x}_3, \dots, \tilde{x}_{2N}) \right], \quad (4.18)$$

where  $\tilde{\Omega}_\eta$  is the union of those connected components of  $\mathbb{H} \setminus \eta$  that contain some of the points  $\{\tilde{x}_3, \dots, \tilde{x}_{2N}\}$  on their boundary, and the event  $\mathcal{E}_\alpha(\eta)$  also implicitly depends on these marked points. By the induction hypothesis, we see that the right-hand side of (4.18) equals

$$(4.18) = (x_2 - x_1)^{-2h} \mathbb{E}(\mathbb{H}; x_1, x_2) \left[ \mathbb{1}\{\mathcal{E}_\alpha(\eta)\} \vec{\mathcal{Z}}_{\alpha/\{1,2\}}(\hat{\Omega}_\eta; x_3, \dots, x_{2N}) \right] = \vec{\mathcal{Z}}(\mathbf{x}) \quad (4.19)$$

for the marked points  $\{x_3, \dots, x_{2N}\}$ , again by Lemma 4.7. Hence, using the identity (4.4), because the explicit function  $\hat{\mathcal{R}}_{\omega_1}$  appearing in Lemma C.4 is clearly continuous, we obtain

$$\begin{aligned} \hat{\mathcal{R}}_{\omega_1}(\mathbf{x}) &= \lim_{\substack{\tilde{x}_j \rightarrow x_j \\ \forall 3 \leq j \leq 2N}} \hat{\mathcal{R}}_{\omega_1}(\tilde{\mathbf{x}}) = \lim_{\substack{\tilde{x}_j \rightarrow x_j \\ \forall 3 \leq j \leq 2N}} \hat{\mathcal{R}}_{\omega_1}(\tilde{\mathbf{x}}) \sum_{\substack{\alpha \in \text{LP}_N \\ \{1,2\} \in \alpha}} \mathbb{P}_N^{x_1 \rightarrow x_2}(\mathbb{H}; \tilde{\mathbf{x}}) [\varsigma(\vec{\eta}) = \alpha] \\ &= \frac{1}{\nu(\kappa)} \limsup_{\substack{\tilde{x}_j \rightarrow x_j \\ \forall 3 \leq j \leq 2N}} \sum_{\substack{\alpha \in \text{LP}_N \\ \{1,2\} \in \alpha}} \mathcal{M}_\nu(\underbrace{\quad}_N, \alpha) \vec{\mathcal{Z}}_\alpha(\tilde{\mathbf{x}}) \quad [\text{by (4.4) \& (3.11)}] \\ &\geq \frac{1}{\nu(\kappa)} \sum_{\substack{\alpha \in \text{LP}_N \\ \{1,2\} \in \alpha}} \mathcal{M}_\nu(\underbrace{\quad}_N, \alpha) \liminf_{\substack{\tilde{x}_j \rightarrow x_j \\ \forall 3 \leq j \leq 2N}} \vec{\mathcal{Z}}_\alpha(\tilde{\mathbf{x}}) \\ &\geq \frac{1}{\nu(\kappa)} \sum_{\substack{\alpha \in \text{LP}_N \\ \{1,2\} \in \alpha}} \mathcal{M}_\nu(\underbrace{\quad}_N, \alpha) \vec{\mathcal{Z}}(\mathbf{x}). \quad [\text{by (4.18) \& (4.19)}] \\ &= \hat{\mathcal{R}}_{\omega_1}(\mathbf{x}), \quad [\text{by (4.4) \& (3.11)}] \end{aligned}$$

which implies (4.17) since  $\mathcal{M}_\nu(\underbrace{\quad}_N, \alpha) > 0$  for all  $\alpha \in \text{LP}_N$  (for the current range  $\kappa \in (4, 8)$ ).  $\square$

Let us next record the intuitive fact that, if  $\{1, 2\} \in \alpha$ , then upon shrinking the points  $x_1$  and  $x_2$  together,  $\mathbb{P}_N^{x_1 \rightarrow x_2}[\varsigma(\vec{\eta}) = \alpha]$  tends to the corresponding probability for  $\vec{\mathbb{P}}_{N-1}$ . In the proof, we make use of the continuity of the pure partition functions, Lemma 4.10, and the induction hypothesis that Lemma 4.4 holds for all  $n \leq N - 1$ .

**Lemma 4.11.** *Suppose  $\{1, 2\} \in \alpha \in \text{LP}_N$  and write  $\hat{\vec{\eta}} = (\vec{\eta}_2, \dots, \vec{\eta}_N)$ . Then, for all  $\xi < x_3$ , using the notation (1.21), we have*

$$\lim_{x_1, x_2 \rightarrow \xi} \mathbb{P}_N^{x_1 \rightarrow x_2}(\mathbb{H}; \mathbf{x}) [\varsigma(\vec{\eta}) = \alpha] = \vec{\mathbb{P}}_{N-1}(\mathbb{H}; \mathbf{x}_1) [\varsigma(\hat{\vec{\eta}}) = \alpha/\{1, 2\}].$$

*Proof.* Write  $\hat{\alpha} = \alpha/\{1, 2\}$ . Using Proposition 3.3 (and Equation (3.22) from its proof), the construction of  $\mathbb{P}_N^{x_1 \rightarrow x_2}$  in Definition 3.6, and the asymptotics property of  $\hat{\mathcal{R}}_{\omega_1}$  from Lemma C.5, we obtain

$$\begin{aligned} &\lim_{x_1, x_2 \rightarrow \xi} \mathbb{P}_N^{x_1 \rightarrow x_2}(\mathbb{H}; \mathbf{x}) [\varsigma(\vec{\eta}) = \alpha] \\ &\geq \lim_{x_1, x_2 \rightarrow \xi} \mathbb{P}_N^{x_1 \rightarrow x_2}(\mathbb{H}; \mathbf{x}) [\vec{\eta}_1 \cap [x_3, +\infty) = \emptyset, \varsigma(\vec{\eta}) = \alpha] \\ &= \lim_{x_1, x_2 \rightarrow \xi} \frac{(x_2 - x_1)^{-2h}}{\nu(\kappa) \hat{\mathcal{R}}_{\omega_1}(\mathbf{x})} \mathbb{E} \left[ \mathbb{1}\{\phi^{-1}(\vec{\eta}) \cap [x_3, +\infty) = \emptyset\} \mathcal{F}_{\underbrace{\quad}_{N-1}}(H_\eta; \mathbf{x}_1) \vec{\mathbb{P}}_{N-1}(H_\eta; \mathbf{x}_1) [\varsigma(\hat{\vec{\eta}}) = \hat{\alpha}] \right] \\ &\geq \mathbb{E} \left[ \liminf_{x_1, x_2 \rightarrow \xi} \mathbb{1}\{\phi^{-1}(\vec{\eta}) \cap [x_3, +\infty) = \emptyset\} \frac{\mathcal{F}_{\underbrace{\quad}_{N-1}}(H_\eta; \mathbf{x}_1)}{\mathcal{F}_{\underbrace{\quad}_{N-1}}(\mathbf{x}_1)} \vec{\mathbb{P}}_{N-1}(H_\eta; \mathbf{x}_1) [\varsigma(\hat{\vec{\eta}}) = \hat{\alpha}] \right], \end{aligned}$$

where  $\mathbb{E}$  is the expected value for the chordal  $\text{SLE}_\kappa$  curve  $\eta$  in  $(\mathbb{H}; 0, \infty)$  and  $H_\eta = \phi^{-1}(\Omega_\eta)$ , where  $\phi$  is the Möbius transformation (3.4) (depending on  $\mathbf{x}$ ) and  $\Omega_\eta$  is the component of the complement  $\mathbb{H} \setminus \eta$  containing  $[0, \infty)$ . We analyze the limits of the three factors inside the expectation separately.

- As the points  $x_1$  and  $x_2$  shrink together, we have  $\mathbb{P}$ -almost surely  $\mathbb{1}\{\phi^{-1}(\vec{\eta}) \cap [x_3, +\infty) = \emptyset\} \rightarrow 1$  because the curve  $\phi^{-1}(\vec{\eta})$  shrinks to a point.

- Also, on this event the conformal covariance of  $\mathcal{F}_{\overbrace{\quad\quad\quad}^{N-1}}$  (Corollary 2.3 and Remark 2.4) yields

$$\lim_{x_1, x_2 \rightarrow \xi} \frac{\mathcal{F}_{\overbrace{\quad\quad\quad}^{N-1}}(H_\eta; \ddot{\mathbf{x}}_1)}{\mathcal{F}_{\overbrace{\quad\quad\quad}^{N-1}}(\ddot{\mathbf{x}}_1)} = \lim_{x_1, x_2 \rightarrow \xi} \prod_{j=3}^{2N} |\varphi'_\eta(x_j)|^h \times \frac{\mathcal{F}_{\overbrace{\quad\quad\quad}^{N-1}}(\varphi_\eta(\ddot{\mathbf{x}}_1))}{\mathcal{F}_{\overbrace{\quad\quad\quad}^{N-1}}(\ddot{\mathbf{x}}_1)} = 1,$$

$\mathbb{P}$ -almost surely, where  $\varphi_\eta$  is any conformal map from  $H_\eta$  onto  $\mathbb{H}$  (so  $\varphi_\eta$  converges to the identity map locally uniformly away from the boundary segment  $[x_1, x_2]$ ).

- Lastly, using the induction hypothesis that Lemma 4.4 holds for  $\mathcal{F}_{\overbrace{\quad\quad\quad}^{N-1}}(\ddot{\mathbf{x}}_1) = \mathcal{K}_{\overbrace{\quad\quad\quad}^{N-1}}(\ddot{\mathbf{x}}_1)$ , we have

$$\mathcal{M}_\nu(\overbrace{\quad\quad\quad}^{N-1}, \hat{\alpha}) \lim_{\substack{y_j \rightarrow x_j \\ \forall 1 \leq j \leq 2N-2}} \frac{\vec{\mathcal{Z}}_{\hat{\alpha}}(\mathbf{y})}{\mathcal{F}_{\overbrace{\quad\quad\quad}^{N-1}}(\mathbf{y})} = \lim_{\substack{y_j \rightarrow x_j \\ \forall 1 \leq j \leq 2N-2}} \vec{\mathbb{P}}_{N-1}(\mathbb{H}; \mathbf{y})[\varsigma(\hat{\boldsymbol{\eta}}) = \hat{\alpha}],$$

so it follows from Lemma 4.10 and the continuity of the explicit function  $\mathcal{F}_{\overbrace{\quad\quad\quad}^N}$  in (1.10) that

$$\lim_{\substack{y_j \rightarrow x_j \\ \forall 1 \leq j \leq 2N-2}} \vec{\mathbb{P}}_{N-1}(\mathbb{H}; \mathbf{y})[\varsigma(\hat{\boldsymbol{\eta}}) = \hat{\alpha}] = \vec{\mathbb{P}}_{N-1}(\mathbb{H}; \mathbf{x})[\varsigma(\hat{\boldsymbol{\eta}}) = \hat{\alpha}], \quad \mathbf{x} \in \mathfrak{X}_{2N}.$$

Combining this with conformal invariance shows that,  $\mathbb{P}$ -almost surely, we have

$$\lim_{x_1, x_2 \rightarrow \xi} \vec{\mathbb{P}}_{N-1}(H_\eta; \ddot{\mathbf{x}}_1)[\varsigma(\hat{\boldsymbol{\eta}}) = \hat{\alpha}] = \lim_{x_1, x_2 \rightarrow \xi} \vec{\mathbb{P}}_{N-1}(\mathbb{H}; \varphi_\eta(\ddot{\mathbf{x}}_1))[\varsigma(\hat{\boldsymbol{\eta}}) = \hat{\alpha}] = \vec{\mathbb{P}}_{N-1}(\mathbb{H}; \ddot{\mathbf{x}}_1)[\varsigma(\hat{\boldsymbol{\eta}}) = \hat{\alpha}].$$

We conclude that

$$\lim_{x_1, x_2 \rightarrow \xi} \mathbb{P}_N^{x_1 \rightarrow x_2}(\mathbb{H}; \mathbf{x})[\varsigma(\vec{\boldsymbol{\eta}}) = \alpha] \geq \vec{\mathbb{P}}_{N-1}(\mathbb{H}; \ddot{\mathbf{x}}_1)[\varsigma(\hat{\boldsymbol{\eta}}) = \alpha/\{1, 2\}]. \quad (4.20)$$

To finish, note that the sum of total probability yields

$$\sum_{\substack{\alpha \in \text{LP}_N \\ \{1, 2\} \in \alpha}} \mathbb{P}_N^{x_1 \rightarrow x_2}(\mathbb{H}; \mathbf{x})[\varsigma(\vec{\boldsymbol{\eta}}) = \alpha] = 1 = \sum_{\substack{\alpha \in \text{LP}_N \\ \{1, 2\} \in \alpha}} \vec{\mathbb{P}}_{N-1}(\mathbb{H}; \ddot{\mathbf{x}}_1)[\varsigma(\hat{\boldsymbol{\eta}}) = \alpha/\{1, 2\}],$$

which implies equality in (4.20).  $\square$

**Proposition 4.12.** *Fix  $\kappa \in (4, 8)$ . The collection  $\{\vec{\mathcal{Z}}_\alpha : \alpha \in \text{LP}_N\}$  of Definition 4.1 comprises smooth functions on  $\mathfrak{X}_{2N}$  which satisfy (PDE).*

*Proof.* Even though we already know that the functions  $\{\hat{\mathcal{R}}_{\omega_b}(\mathbf{x}) : 1 \leq b \leq N\}$  satisfy (PDE) (Lemma C.3), because the probability measures  $\mathbb{P}_N^{x_1 \rightarrow x_{2b}}(\mathbb{H}; \mathbf{x})$  and  $\vec{\mathbb{P}}_N(\mathbb{H}; \mathbf{x})$  are not known to immediately satisfy any differential equations, or even to be differentiable, the verification of (PDE) for the functions of Definition 4.1 requires a rather nontrivial argument. Under the induction hypothesis that Lemmas 4.3 & 4.4 hold for all  $n \leq N - 1$ , as detailed in the proof of [Pel19, Lemma B.4] or [Wu20, Lemmas 6.3 & 6.4], we may apply the Möbius covariance from Lemma 4.9 and the cascade property (4.7) from Lemma 4.7 to derive a local martingale for the evolution of the chordal  $\text{SLE}_\kappa$  curve. Together with the continuity from Lemma 4.10, this shows that  $\vec{\mathcal{Z}}_\alpha$  is a distributional solution to the PDE system (PDE) (see Proposition A.6 in Appendix A). Because the PDE system (PDE) comprises hypoelliptic PDEs, we may conclude that any distributional solution to it is smooth (Proposition A.2 in Appendix A).  $\square$

## 4.5 Relations of pure partition functions to Coulomb gas integrals

Our proof of the asymptotics property in Section 4.6 uses crucially the fact that the pure partition functions are intimately related to the Coulomb gas integrals discussed in Section 2 and used in Section 3. The purpose of this section is to make this relation precise, that is, to verify Lemma 4.4.

We begin by checking the power-law bound (PLB) for the pure partition functions (Lemma 4.13). This together with Lemma 4.5 and Proposition 4.12 show that  $\{\vec{Z}_\alpha: \alpha \in \text{LP}_N\} \subset \mathcal{S}_N$  belong to the PDE solution space (2.21), which enables us to invoke the uniqueness from Lemma 2.11 for elements in  $\mathcal{S}_N$ .

**Lemma 4.13.** *Fix  $\kappa \in (4, 8)$ . The collection  $\{\vec{Z}_\alpha: \alpha \in \text{LP}_N\}$  of Definition 4.1 satisfies the bound (PLB).*

*Proof.* By Definition 4.1, it suffices to bound  $\mathcal{K}_{\overbrace{\text{---}}_N}$  in terms of the Coulomb gas integral functions  $\hat{\mathcal{R}}_{\omega_s}$ , which satisfy the bound (PLB) by construction (Lemma C.3):

$$\mathcal{K}_{\overbrace{\text{---}}_N}(\mathbf{x}) \lesssim \max_{1 \leq s \leq N} \hat{\mathcal{R}}_{\omega_s}(\mathbf{x}),$$

where “ $\lesssim$ ” holds up to some finite constant depending on  $\kappa$  and  $N$ . □

Next, observe that by Equations (4.1) and (4.4), we have

$$\mathcal{K}_{\overbrace{\text{---}}_N}(\mathbf{x}) = \sum_{\alpha \in \text{LP}_N} \mathcal{M}_\nu(\overbrace{\text{---}}_N, \alpha) \vec{Z}_\alpha(\mathbf{x}), \quad \mathbf{x} \in \mathfrak{X}_{2N}. \quad (4.21)$$

We analogously define

$$\mathcal{K}_{\underbrace{\text{---}}_N}(\mathbf{x}) := \sum_{\alpha \in \text{LP}_N} \mathcal{M}_\nu(\underbrace{\text{---}}_N, \alpha) \vec{Z}_\alpha(\mathbf{x}), \quad \mathbf{x} \in \mathfrak{X}_{2N}. \quad (4.22)$$

These functions are naturally related by rotation.

**Lemma 4.14.** *Fix  $\kappa \in (4, 8)$ . The collections  $\{\mathcal{K}_{\overbrace{\text{---}}_N}: N \in \mathbb{Z}_{\geq 0}\}$  and  $\{\mathcal{K}_{\underbrace{\text{---}}_N}: N \in \mathbb{Z}_{\geq 0}\}$  in (4.21, 4.22) satisfy the Möbius covariance*

$$\mathcal{K}_{\overbrace{\text{---}}_N}(\mathbf{x}) = \prod_{i=1}^{2N} |\varphi'(x_i)|^h \times \begin{cases} \mathcal{K}_{\underbrace{\text{---}}_N}(\varphi(\mathbf{x})), & \text{if } j \in \{2, 4, \dots, 2N-2\}, \\ \mathcal{K}_{\overbrace{\text{---}}_N}(\varphi(\mathbf{x})), & \text{if } j \in \{1, 3, \dots, 2N-1\}, \end{cases} \quad \mathbf{x} \in \mathfrak{X}_{2N}, \quad (4.23)$$

for all Möbius maps  $\varphi \in \text{PSL}(2, \mathbb{R})$ , where  $\varphi(\mathbf{x}) := (\varphi(x_j) \dots, \varphi(x_{2N}), \varphi(x_1), \dots, \varphi(x_{j-1}))$ , so that

$$\varphi(x_j) < \varphi(x_{j+1}) < \dots < \varphi(x_{2N}) < \varphi(x_1) < \varphi(x_2) < \dots < \varphi(x_{j-1}).$$

*Proof.* For any Möbius maps  $\varphi \in \text{PSL}(2, \mathbb{R})$ , using rotation symmetry of the meander matrix (1.3) and the Möbius covariance (4.15) of  $\vec{Z}_\alpha$  from Lemma 4.9, using (4.21), we obtain

$$\begin{aligned} \mathcal{K}_{\overbrace{\text{---}}_N}(\mathbf{x}) &= \sum_{\alpha \in \text{LP}_N} \mathcal{M}_\nu(\overbrace{\text{---}}_N, \alpha) \vec{Z}_\alpha(\mathbf{x}) && \text{[by (4.21)]} \\ &= \prod_{i=1}^{2N} |\varphi'(x_i)|^h \times \sum_{\alpha \in \text{LP}_N} \mathcal{M}_\nu(\overbrace{\text{---}}_N, \alpha) \vec{Z}_{\varphi(\alpha)}(\varphi(\mathbf{x})) && \text{[by (4.15)]} \\ &= \prod_{i=1}^{2N} |\varphi'(x_i)|^h \times \sum_{\beta \in \text{LP}_N} \mathcal{M}_\nu(\overbrace{\text{---}}_N, \varphi^{-1}(\beta)) \vec{Z}_\beta(\varphi(\mathbf{x})). \end{aligned}$$

- If  $j$  is even, we have  $\mathcal{M}_\nu(\overbrace{\text{---}}_N, \varphi^{-1}(\beta)) = \mathcal{M}_\nu(\varphi(\overbrace{\text{---}}_N), \beta) = \mathcal{M}_\nu(\underbrace{\text{---}}_N, \beta)$ , and thus,

$$\begin{aligned} \prod_{i=1}^{2N} |\varphi'(x_i)|^{-h} \times \mathcal{K}_{\overbrace{\text{---}}_N}(\mathbf{x}) &= \sum_{\beta \in \text{LP}_N} \mathcal{M}_\nu(\underbrace{\text{---}}_N, \beta) \vec{Z}_\beta(\varphi(\mathbf{x})) \\ &= \mathcal{K}_{\underbrace{\text{---}}_N}(\varphi(\mathbf{x})). \end{aligned} \quad \text{[by (4.22)]}$$



- If  $j$  is odd, we have  $\mathcal{M}_\nu(\underbrace{\hspace{1cm}}_N, \varphi^{-1}(\beta)) = \mathcal{M}_\nu(\varphi(\underbrace{\hspace{1cm}}_N), \beta) = \mathcal{M}_\nu(\underbrace{\hspace{1cm}}_N, \beta)$ , and thus,

$$\begin{aligned} \prod_{i=1}^{2N} |\varphi'(x_i)|^{-h} \times \mathcal{K}_{\underbrace{\hspace{1cm}}_N}(\mathbf{x}) &= \sum_{\beta \in \text{LP}_N} \mathcal{M}_\nu(\underbrace{\hspace{1cm}}_N, \beta) \vec{\mathcal{Z}}_\beta(\varphi(\mathbf{x})) \\ &= \mathcal{K}_{\underbrace{\hspace{1cm}}_N}(\varphi(\mathbf{x})). \end{aligned} \quad [\text{by (4.21)}]$$

This proves the asserted covariance property (4.23).  $\square$

We are now ready to finish the induction step to prove Lemma 4.4.

**Lemma 4.15.** *Fix  $N \geq 2$ . Suppose that the following hold:*

Lemma 4.3:  $\vec{\mathcal{Z}}_\alpha = \tilde{\mathcal{Z}}_\alpha$  for all  $\alpha \in \text{LP}_n$  and  $n \leq N-1$ ;

Lemma 4.4:  $\mathcal{K}_{\underbrace{\hspace{1cm}}_n} = \mathcal{F}_{\underbrace{\hspace{1cm}}_n}$  for all  $n \leq N-1$ .

Then, we have

$$\mathcal{Z}_\alpha(\mathbf{x}) = \vec{\mathcal{Z}}_\alpha(\mathbf{x}) = \mathcal{F}_{\underbrace{\hspace{1cm}}_N}(\mathbf{x}) \frac{\tilde{\mathbb{P}}_N[\zeta(\vec{\eta}) = \alpha]}{\mathcal{M}_\nu(\underbrace{\hspace{1cm}}_N, \alpha)}, \quad \mathbf{x} \in \mathfrak{X}_{2N}, \alpha \in \text{LP}_N.$$

*Proof.* It suffices to prove that  $\mathcal{K}_{\underbrace{\hspace{1cm}}_N} = \mathcal{F}_{\underbrace{\hspace{1cm}}_N}$  in (4.1). To this end, we use the uniqueness from Lemma 2.11 of elements in the PDE solution space  $\mathcal{S}_N$  (2.21) singled out by their asymptotic properties. Because  $\mathcal{F}_{\underbrace{\hspace{1cm}}_N} \in \mathcal{S}_N$ , to prove that  $\mathcal{K}_{\underbrace{\hspace{1cm}}_N} = \mathcal{F}_{\underbrace{\hspace{1cm}}_N}$  it suffices to verify that  $\mathcal{K}_{\underbrace{\hspace{1cm}}_N} \in \mathcal{S}_N$  and they satisfy the same recursive asymptotics property ( $\mathcal{F}$ -ASY):

$$\lim_{x_j, x_{j+1} \rightarrow \xi} \frac{\mathcal{K}_{\underbrace{\hspace{1cm}}_N}(\mathbf{x})}{(x_{j+1} - x_j)^{-2h}} = \begin{cases} \nu(\kappa) \mathcal{K}_{\underbrace{\hspace{1cm}}_{N-1}}(\ddot{\mathbf{x}}_j), & \text{if } j \in \{2, 4, \dots, 2N-2\}, \\ \mathcal{K}_{\underbrace{\hspace{1cm}}_{N-1}}(\ddot{\mathbf{x}}_j), & \text{if } j \in \{1, 3, \dots, 2N-1\}, \end{cases} \quad (4.24)$$

for all  $\xi \in (x_{j-1}, x_{j+2})$  and for all  $N \geq 1$  and  $j \in \{1, 2, \dots, 2N-1\}$ . To this end, note first that  $\vec{\mathcal{Z}}_\alpha \in \mathcal{S}_N$  by Proposition 4.12 and Lemmas 4.5 & 4.13. Thus, we readily see that  $\mathcal{K}_{\underbrace{\hspace{1cm}}_N} \in \mathcal{S}_N$  as a linear combination (4.21) of them. To verify the asymptotics property (4.24), thanks to Lemma (4.23), by conjugating by a suitable Möbius map it suffices without loss of generality to check (4.24) in the case  $j = 1$ , and the following analogous property in the case  $j = 1$ :

$$\lim_{x_1, x_2 \rightarrow \xi} \frac{\mathcal{K}_{\underbrace{\hspace{1cm}}_N}(\mathbf{x})}{(x_2 - x_1)^{-2h}} = \begin{cases} \mathcal{K}_{\underbrace{\hspace{1cm}}_{N-1}}(\ddot{\mathbf{x}}_1), & \text{if } j \in \{2, 4, \dots, 2N-2\}, \\ \nu(\kappa) \mathcal{K}_{\underbrace{\hspace{1cm}}_{N-1}}(\ddot{\mathbf{x}}_1), & \text{if } j \in \{1, 3, \dots, 2N-1\}. \end{cases} \quad (4.25)$$

- For the first case ((4.24) with  $j = 1$ ), using the definition (4.1) and the asymptotics property of  $\hat{\mathcal{R}}_{\omega_s}$  from Lemma C.5, we have

$$\begin{aligned} & \lim_{x_1, x_2 \rightarrow \xi} \frac{\mathcal{K}_{\underbrace{\hspace{1cm}}_N}(\mathbf{x})}{(x_2 - x_1)^{-2h}} \\ &= \nu(\kappa) \lim_{x_1, x_2 \rightarrow \xi} \sum_{s=1}^N \frac{\hat{\mathcal{R}}_{\omega_s}(\mathbf{x})}{(x_2 - x_1)^{-2h}} \sum_{\substack{\beta \in \text{LP}_N \\ \{1, 2s\} \in \beta}} w_s(\beta) \mathbb{P}_N^{x_1 \rightarrow x_2s}[\zeta(\vec{\eta}) = \beta] \quad [\text{by (4.1)}] \\ &= \nu(\kappa) \lim_{x_1, x_2 \rightarrow \xi} \frac{\hat{\mathcal{R}}_{\omega_1}(\mathbf{x})}{(x_2 - x_1)^{-2h}} \sum_{\substack{\beta \in \text{LP}_N \\ \{1, 2\} \in \beta}} \mathbb{P}_N^{x_1 \rightarrow x_2}[\zeta(\vec{\eta}) = \beta] \quad [\text{by Lemma C.5 and (3.11)}] \\ &= \mathcal{F}_{\underbrace{\hspace{1cm}}_{N-1}}(\ddot{\mathbf{x}}_1) \quad [\text{by Lemma C.5}] \\ &= \mathcal{K}_{\underbrace{\hspace{1cm}}_{N-1}}(\ddot{\mathbf{x}}_1), \quad [\text{by Lemma 4.4}] \end{aligned}$$

due to the induction hypothesis that Lemma 4.4 holds for  $\mathcal{F}_{\underbrace{\hspace{1cm}}_{N-1}}(\ddot{\mathbf{x}}_1) = \mathcal{K}_{\underbrace{\hspace{1cm}}_{N-1}}(\ddot{\mathbf{x}}_1)$ .

- The second case ((4.25) with  $j = 1$ ) is slightly more elaborate. First, note that

$$\begin{aligned}
& \lim_{x_1, x_2 \rightarrow \xi} \frac{\mathcal{K}_{\underbrace{\quad\quad\quad}_N}(\mathbf{x})}{(x_2 - x_1)^{-2h}} \tag{4.26} \\
& := \lim_{x_1, x_2 \rightarrow \xi} \sum_{\alpha \in \text{LP}_N} \mathcal{M}_\nu(\underbrace{\quad\quad\quad}_N, \alpha) \frac{\vec{\mathcal{Z}}_\alpha(\mathbf{x})}{(x_2 - x_1)^{-2h}} \tag{by (4.22)} \\
& = \nu(\kappa) \lim_{x_1, x_2 \rightarrow \xi} \sum_{s=1}^N \frac{\hat{\mathcal{R}}_{\omega_s}(\mathbf{x})}{(x_2 - x_1)^{-2h}} \sum_{\substack{\alpha \in \text{LP}_N \\ \{1, 2s\} \in \alpha}} \frac{w_s(\alpha) \mathcal{M}_\nu(\underbrace{\quad\quad\quad}_N, \alpha)}{\mathcal{M}_\nu(\underbrace{\quad\quad\quad}_N, \alpha)} \mathbb{P}_N^{x_1 \rightarrow x_{2s}}[\zeta(\vec{\eta}) = \alpha]. \tag{by (4.4)}
\end{aligned}$$

By Lemma C.5, only the  $s = 1$  term in the sum survives in the limit, with  $w_1(\alpha) = 1$  by (3.11). From Lemmas C.5 and 4.11, upon shrinking the points  $x_1$  and  $x_2$  together, the limits of interest are

$$\begin{aligned}
\lim_{x_1, x_2 \rightarrow \xi} \mathbb{P}_N^{x_1 \rightarrow x_2}(\mathbb{H}; \mathbf{x})[\zeta(\vec{\eta}) = \alpha] &= \vec{\mathbb{P}}_{N-1}(\mathbb{H}; \dot{\mathbf{x}}_1)[\zeta(\hat{\vec{\eta}}) = \alpha/\{1, 2\}], \\
\lim_{x_1, x_2 \rightarrow \xi} \frac{\hat{\mathcal{R}}_{\omega_1}(\mathbf{x})}{(x_2 - x_1)^{-2h}} &= \frac{1}{\nu(\kappa)} \mathcal{F}_{\underbrace{\quad\quad\quad}_{N-1}}(\dot{\mathbf{x}}_1) = \frac{1}{\nu(\kappa)} \mathcal{K}_{\underbrace{\quad\quad\quad}_{N-1}}(\dot{\mathbf{x}}_1), \tag{4.27}
\end{aligned}$$

where we write  $\hat{\vec{\eta}} = (\vec{\eta}_2, \dots, \vec{\eta}_N)$ , and use again the induction hypothesis that Lemma 4.4 holds for  $\mathcal{F}_{\underbrace{\quad\quad\quad}_{N-1}}(\dot{\mathbf{x}}_1) = \mathcal{K}_{\underbrace{\quad\quad\quad}_{N-1}}(\dot{\mathbf{x}}_1)$ . Thus, we obtain

$$\begin{aligned}
(4.26) &= \mathcal{K}_{\underbrace{\quad\quad\quad}_{N-1}}(\dot{\mathbf{x}}_1) \sum_{\substack{\alpha \in \text{LP}_N \\ \{1, 2\} \in \alpha}} \frac{\mathcal{M}_\nu(\underbrace{\quad\quad\quad}_N, \alpha)}{\mathcal{M}_\nu(\underbrace{\quad\quad\quad}_N, \alpha)} \vec{\mathbb{P}}_{N-1}(\mathbb{H}; \dot{\mathbf{x}}_1)[\zeta(\hat{\vec{\eta}}) = \alpha/\{1, 2\}] \tag{by (4.27)} \\
&= \nu(\kappa) \mathcal{K}_{\underbrace{\quad\quad\quad}_{N-1}}(\dot{\mathbf{x}}_1) \sum_{\beta \in \text{LP}_{N-1}} \frac{\mathcal{M}_\nu(\underbrace{\quad\quad\quad}_{N-1}, \beta)}{\mathcal{M}_\nu(\underbrace{\quad\quad\quad}_{N-1}, \beta)} \vec{\mathbb{P}}_{N-1}(\mathbb{H}; \dot{\mathbf{x}}_1)[\zeta(\hat{\vec{\eta}}) = \beta] \\
&= \nu(\kappa) \sum_{\beta \in \text{LP}_{N-1}} \mathcal{M}_\nu(\underbrace{\quad\quad\quad}_{N-1}, \beta) \vec{\mathcal{Z}}_\beta(\dot{\mathbf{x}}_1) \tag{by (4.2)} \\
&= \nu(\kappa) \mathcal{K}_{\underbrace{\quad\quad\quad}_{N-1}}(\dot{\mathbf{x}}_1). \tag{by (4.22)}
\end{aligned}$$

This concludes the proof.  $\square$

## 4.6 Asymptotics properties

**Proposition 4.16.** *Fix  $\kappa \in (4, 8)$ . The collection  $\{\vec{\mathcal{Z}}_\alpha : \alpha \in \text{LP}_N\}$  of Definition 4.1 satisfies the recursive asymptotics (ASY).*

*Proof.* It suffices to show (ASY) when  $j = 1$ , that is, when  $x_1, x_2 \rightarrow \xi < x_3$ , because for the other cases the claim then follows after conjugating by a suitable Möbius map and using Lemma 4.9.

We first consider the case  $\{1, 2\} \notin \alpha$  in (ASY). By (4.4) and the asymptotics for  $\hat{\mathcal{R}}_{\omega_\nu}$  from Lemma C.5,

$$\lim_{x_1, x_2 \rightarrow \xi} \frac{\vec{\mathcal{Z}}_\alpha(\mathbf{x})}{(x_2 - x_1)^{-2h}} \leq \lim_{x_1, x_2 \rightarrow \xi} \sum_{\substack{\beta \in \text{LP}_N \\ \{1, 2\} \notin \beta}} \frac{\vec{\mathcal{Z}}_\beta(\mathbf{x})}{(x_2 - x_1)^{-2h}} \lesssim \lim_{x_1, x_2 \rightarrow \xi} \sum_{s=2}^N \frac{\hat{\mathcal{R}}_{\omega_s}(\mathbf{x})}{(x_2 - x_1)^{-2h}} = 0, \tag{4.28}$$

where “ $\lesssim$ ” holds up to some finite constant depending on  $\kappa$  and  $N$ .

We next consider the case  $\{1, 2\} \in \alpha$  in (ASY).

Under the induction hypothesis that Lemma 4.3 holds for all  $n \leq N - 1$ , and because we already verified Lemma 4.4 in the previous section, we may apply the cascade property (4.7) from Lemma 4.7,

induction on the number of curves, and the same arguments as in the proof of [Wu20, Lemma 6.6] or [Pel19, Lemma B.3 and Lemma C.1] to show that<sup>19</sup>

$$\liminf_{x_1, x_2 \rightarrow \xi} \frac{\vec{\mathcal{Z}}_\alpha(\mathbf{x})}{(x_2 - x_1)^{-2h}} \geq \vec{\mathcal{Z}}_{\alpha/\{1,2\}}(\ddot{\mathbf{x}}_1). \quad (4.29)$$

We will accomplish the existence of the limit (ASY) by using the relation of  $\vec{\mathcal{Z}}_\alpha$  with the Coulomb gas integral functions. Indeed, by Lemma 4.4, we have  $\mathcal{K}_{\text{arc}_{N-1}} = \mathcal{F}_{\text{arc}_{N-1}}$ , so we obtain

$$\begin{aligned} \mathcal{F}_{\text{arc}_{N-1}}(\ddot{\mathbf{x}}_1) &= \lim_{x_1, x_2 \rightarrow \xi} \frac{\mathcal{F}_{\text{arc}_N}(\mathbf{x})}{(x_2 - x_1)^{-2h}} && \text{[by } (\mathcal{F}\text{-ASY)]} \\ &= \lim_{x_1, x_2 \rightarrow \xi} \frac{\mathcal{K}_{\text{arc}_N}(\mathbf{x})}{(x_2 - x_1)^{-2h}} && \text{[by Lemma 4.4]} \\ &= \limsup_{x_1, x_2 \rightarrow \xi} \sum_{\substack{\alpha \in \text{LP}_N \\ \{1,2\} \in \alpha}} \mathcal{M}_\nu(\text{arc}_N, \alpha) \frac{\vec{\mathcal{Z}}_\alpha(\mathbf{x})}{(x_2 - x_1)^{-2h}} && \text{[by (4.2) \& (4.28)]} \\ &\geq \sum_{\substack{\alpha \in \text{LP}_N \\ \{1,2\} \in \alpha}} \mathcal{M}_\nu(\text{arc}_N, \alpha) \liminf_{x_1, x_2 \rightarrow \xi} \frac{\vec{\mathcal{Z}}_\alpha(\mathbf{x})}{(x_2 - x_1)^{-2h}} \\ &\geq \sum_{\substack{\alpha \in \text{LP}_N \\ \{1,2\} \in \alpha}} \mathcal{M}_\nu(\text{arc}_N, \alpha) \vec{\mathcal{Z}}_{\alpha/\{1,2\}}(\ddot{\mathbf{x}}_1) && \text{[by (4.28) \& (4.29)]} \\ &= \sum_{\beta \in \text{LP}_{N-1}} \mathcal{M}_\nu(\text{arc}_{N-1}, \beta) \vec{\mathcal{Z}}_\beta(\ddot{\mathbf{x}}_1) \\ &= \mathcal{K}_{\text{arc}_{N-1}}(\ddot{\mathbf{x}}_1) && \text{[by (4.21)]} \\ &= \mathcal{F}_{\text{arc}_{N-1}}(\ddot{\mathbf{x}}_1), && \text{[by Lemma 4.4]} \end{aligned}$$

Since  $\mathcal{M}_\nu(\text{arc}_N, \alpha) > 0$  for all  $\alpha \in \text{LP}_N$  (for the current range  $\kappa \in (4, 8)$ ), we conclude that

$$\limsup_{x_1, x_2 \rightarrow \xi} \frac{\vec{\mathcal{Z}}_\alpha(\mathbf{x})}{(x_2 - x_1)^{-2h}} = \liminf_{x_1, x_2 \rightarrow \xi} \frac{\vec{\mathcal{Z}}_\alpha(\mathbf{x})}{(x_2 - x_1)^{-2h}} = \vec{\mathcal{Z}}_{\alpha/\{1,2\}}(\ddot{\mathbf{x}}_1).$$

This shows (ASY) and finishes the proof.  $\square$

## 4.7 Reversibility

In this section, we finish the induction step to prove Lemma 4.3. We work under the remaining induction hypothesis that Lemma 4.3 holds for all  $n \leq N - 1$ .

**Lemma 4.17.** *Fix  $N \geq 2$ . Suppose that  $\vec{\mathcal{Z}}_\alpha = \vec{\mathcal{Z}}_{\hat{\alpha}}$  for all  $\hat{\alpha} \in \text{LP}_n$  and  $n \leq N - 1$ . Then, we have  $\vec{\mathcal{Z}}_\alpha = \vec{\mathcal{Z}}_\alpha$  for all  $\alpha \in \text{LP}_N$ .*

*Proof.* Using Lemmas 4.4 and 2.1 combined with the identity (2.2) for  $\mathcal{F}_{\text{arc}_N}$ , we have

$$\vec{\mathcal{Z}}_\alpha(\mathbf{x}) = \mathcal{F}_{\text{arc}_N}(\mathbf{x}) \frac{\tilde{\mathbb{P}}_N(\mathbb{H}; \mathbf{x})[\zeta(\vec{\eta}) = \alpha]}{\mathcal{M}_\nu(\text{arc}_N, \alpha)} \quad \text{[by Lemma 4.4]}$$

<sup>19</sup>It seems very hard to get equality of the limit in (4.29) without substantial additional work, since as far as we are aware, there are no known estimates available that would allow the exchange of the limit and expected value in the asymptotics (ASY) and the cascade relation (4.7) — see the discussion in [Pel19, Appendix B].

$$\begin{aligned}
&= \mathcal{F}_{-\overbrace{\quad\quad\quad}^N}(-\mathbf{x}) \frac{\tilde{\mathbb{P}}_N(\mathbb{H}; \mathbf{x})[\zeta(\vec{\eta}) = \alpha]}{\mathcal{M}_\nu(\overbrace{\quad\quad\quad}^N, \alpha)} && \text{[by (2.3) \& (2.2)]} \\
&= \mathcal{F}_{\overbrace{\quad\quad\quad}^N}(-\mathbf{x}) \frac{\tilde{\mathbb{P}}_N(\mathbb{H}; -\mathbf{x})[\zeta(\vec{\eta}) = -\alpha]}{\mathcal{M}_\nu(\overbrace{\quad\quad\quad}^N, -\alpha)} && \text{[by Definition of } \tilde{\mathbb{P}}_N\text{]} \\
&= \vec{\mathcal{Z}}_{-\alpha}(-\mathbf{x}) && \text{[by (4.5)]}
\end{aligned}$$

where  $-\mathbf{x} = (-x_{2N}, \dots, -x_1)$  and  $-\overbrace{\quad\quad\quad}^N = \overbrace{\quad\quad\quad}^N$  is reflection invariant (across the imaginary axis). Hence, in particular,  $\vec{\mathcal{Z}}_\alpha$  satisfies the properties (PDE), (COV), and (PLB) — so  $\vec{\mathcal{Z}}_\alpha \in \mathcal{S}_N$ . Therefore, we can prove that  $\vec{\mathcal{Z}}_\alpha = \vec{\mathcal{Z}}_\alpha$  by invoking the uniqueness of the PDE boundary value problem on  $\mathcal{S}_N$  from Lemma 2.11. Indeed, both collections  $\{\vec{\mathcal{Z}}_\alpha: \alpha \in \text{LP}_N\}$  and  $\{\vec{\mathcal{Z}}_\alpha: \alpha \in \text{LP}_N\}$  satisfy the same recursive asymptotics property (ASY): for the former collection this was the content of Proposition 4.16, and for the latter, the above identity  $\vec{\mathcal{Z}}_\alpha(\mathbf{x}) = \vec{\mathcal{Z}}_{-\alpha}(-\mathbf{x})$  implies that

$$\begin{aligned}
\lim_{x_j, x_{j+1} \rightarrow \xi} \frac{\vec{\mathcal{Z}}_\alpha(\mathbf{x})}{|x_{j+1} - x_j|^{-2h}} &= \lim_{x_j, x_{j+1} \rightarrow \xi} \frac{\vec{\mathcal{Z}}_{-\alpha}(-\mathbf{x})}{|x_{j+1} - x_j|^{-2h}} \\
&= \begin{cases} \vec{\mathcal{Z}}_{-\alpha/\{2N-j, 2N-j+1\}}(-\ddot{\mathbf{x}}_{2N-j}), & \text{if } \{j, j+1\} \in \alpha, \\ 0, & \text{if } \{j, j+1\} \notin \alpha, \end{cases} && \text{[by (ASY)]} \\
&= \begin{cases} \vec{\mathcal{Z}}_{\alpha/\{j, j+1\}}(\ddot{\mathbf{x}}_j), & \text{if } \{j, j+1\} \in \alpha, \\ 0, & \text{if } \{j, j+1\} \notin \alpha, \end{cases}
\end{aligned}$$

finishing the proof using the induction hypothesis that  $\vec{\mathcal{Z}}_\alpha = \vec{\mathcal{Z}}_\alpha$  holds for all  $\alpha \in \text{LP}_n$  and  $n \leq N-1$ .  $\square$

#### 4.8 Global multiple SLE $_\kappa$ for $\kappa \in (4, 8)$ : proof of Theorem 1.21

The goal of this section is to construct a probability measure on the curve space  $X_0(\Omega; \mathbf{x})$  defined in (1.44) which satisfies the defining property of a global multiple SLE $_\kappa$  measure (Definition 1.20), and to prove Theorem 1.21. Because the construction is already well-known for the range  $\kappa \in (0, 4]$  [KL07, PW19], and not very meaningful when  $\kappa \geq 8$  (cf. [LPW24]), we will focus on the case  $\kappa \in (4, 8)$ . Very recently a construction in the case  $\kappa \in (4, 8)$  was established using welding of LQG surfaces in [AHSY23], and the uniqueness of such measures was proven in [Zha23]. In this article, we obtain a different construction and derive the marginal law using the relation of Coulomb gas integrals and pure partition functions.

Fix  $\kappa \in (4, 8)$  and  $N \geq 1$ . Recall that in Section 3.2 we constructed a conformally invariant probability measure  $\tilde{\mathbb{P}}_N(\Omega; \mathbf{x})$  (Definition 3.6) on (oriented) curves  $\vec{\eta} := (\vec{\eta}_1, \dots, \vec{\eta}_N) \in X_0(\Omega; \mathbf{x})$  such that  $\vec{\eta}_s$  starts from  $x_{2s-1}$  for each  $1 \leq s \leq N$ . Similarly, the probability measure  $\tilde{\mathbb{P}}_N(\Omega; \mathbf{x})$  is obtained from  $\tilde{\mathbb{P}}_N(\Omega; \mathbf{x})$  by sending  $\Omega$  onto  $\Omega$  via an anti-conformal map  $\zeta$ : if one samples  $(\vec{\eta}_1, \dots, \vec{\eta}_N) \sim \tilde{\mathbb{P}}_N(\Omega; \zeta(x_{2N}), \dots, \zeta(x_1))$ , then  $\tilde{\mathbb{P}}_N(\Omega; \mathbf{x})$  is the law of the (oriented) curves  $(\vec{\eta}_1, \dots, \vec{\eta}_N) := (\zeta^{-1}(\vec{\eta}_N), \dots, \zeta^{-1}(\vec{\eta}_1))$ .

Now, Definition 1.20 states that a global  $N$ -SLE $_\kappa$  associated to a given link pattern

$$\alpha = \{\{a_1, b_1\}, \dots, \{a_N, b_N\}\} \in \text{LP}_N$$

must be supported on unoriented curves  $\eta := (\eta_1, \dots, \eta_N) \in X_0^\alpha(\Omega; \mathbf{x})$  and satisfy the resampling property. To construct it, we consider the regular conditional laws<sup>20</sup>

$$\begin{aligned}
\tilde{\mathbb{P}}_\alpha(\Omega; \mathbf{x}) &:= \tilde{\mathbb{P}}_N(\Omega; \mathbf{x})[\cdot \mid \zeta(\vec{\eta}) = \alpha], \\
\tilde{\mathbb{P}}_\alpha(\Omega; \mathbf{x}) &:= \tilde{\mathbb{P}}_N(\Omega; \mathbf{x})[\cdot \mid \zeta(\vec{\eta}) = \alpha], \quad \alpha \in \text{LP}_N.
\end{aligned} \tag{4.30}$$

We denote by  $\vec{\mathbb{E}}_\alpha(\Omega; \mathbf{x})$  and  $\vec{\mathbb{E}}_\alpha(\Omega; \mathbf{x})$  the corresponding expected values.

<sup>20</sup>Note that the probabilities  $\tilde{\mathbb{P}}_N(\Omega; \mathbf{x})[\zeta(\vec{\eta}) = \alpha]$  and  $\tilde{\mathbb{P}}_N(\Omega; \mathbf{x})[\zeta(\vec{\eta}) = \alpha]$  are positive by Lemma 3.7.

**Definition 4.18.** Fix  $\kappa \in (4, 8)$  and  $\alpha = \{\{a_1, b_1\}, \dots, \{a_N, b_N\}\} \in \text{LP}_N$ . Here, we index the link endpoints not as in (1.2), but we rather assume that each  $a_r$  is odd and each  $b_r$  is even. For each  $s \in \{1, 2, \dots, N\}$ , using the notation from Lemma 4.7, we construct a probability measure  $\text{P}_\alpha^{(s)}(\Omega; \mathbf{x})$  in the following manner.

1. Sample a random curve  $\eta_s$  as a chordal  $\text{SLE}_\kappa$  curve  $\eta$  on  $(\Omega; x_{a_s}, x_{b_s})$  weighted by (cf. (4.7))

$$\mathbb{1}\{\mathcal{E}_\alpha(\eta)\} \frac{\vec{\mathcal{Z}}_{\triangleleft}(\Omega; x_{a_s}, x_{b_s})}{\vec{\mathcal{Z}}_\alpha(\Omega; \mathbf{x})} \vec{\mathcal{Z}}_{\alpha/\{a_s, b_s\}}(\hat{\Omega}_\eta; \hat{\mathbf{x}}_{a_s, b_s}).$$

2. Given  $\eta_s$ , sample the other curves  $\{\eta_1, \dots, \eta_N\} \setminus \{\eta_s\}$  independently under the product measure of

$$\vec{\text{P}}_{\alpha^D}(D; \dots) \quad \text{or} \quad \check{\text{P}}_{\alpha^D}(D; \dots),$$

with  $\alpha^D$  the sub-link pattern of  $\alpha/\{a_s, b_s\}$  associated to a component  $D \subset \hat{\Omega}_\eta$  with some of the points among  $\{x_1, \dots, x_{2N}\} \setminus \{x_{a_s}, x_{b_s}\}$  on  $\partial D$ , where the orientation is chosen in such a way that curves in the components left of the curve  $\gamma_s$  are sampled from  $\vec{\text{P}}_{\alpha^D}(D; \dots)$  and curves in the components right of the curve  $\gamma_s$  are sampled from  $\check{\text{P}}_{\alpha^D}(D; \dots)$ .

We first note that  $\text{P}_\alpha^{(s)}(\Omega; \mathbf{x})$  is independent of  $s$ .

**Lemma 4.19.** Fix  $\kappa \in (4, 8)$ ,  $N \geq 1$ ,  $\alpha \in \text{LP}_N$ , and a polygon  $(\Omega; \mathbf{x})$ . For all  $1 \leq s \neq r \leq N$ , we have

$$\text{P}_\alpha^{(s)}(\Omega; \mathbf{x}) = \text{P}_\alpha^{(r)}(\Omega; \mathbf{x}).$$

*Proof.* This can be proven by induction on the number  $N \geq 1$  of curves similarly to [Wu20, Lemma 6.2], by applying the cascade property (4.7) from Lemma 4.7 and the arguments in its proof.  $\square$

Next, we show that  $\text{P}_\alpha^{(s)}(\Omega; \mathbf{x})$  coincides with  $\vec{\text{P}}_\alpha(\Omega; \mathbf{x})$  defined in (4.30).

**Lemma 4.20.** Fix  $\kappa \in (4, 8)$ ,  $N \geq 1$ ,  $\alpha \in \text{LP}_N$ , and a polygon  $(\Omega; \mathbf{x})$ . For all  $1 \leq s \leq N$ , we have

$$\text{P}_\alpha^{(s)}(\Omega; \mathbf{x}) = \vec{\text{P}}_\alpha(\Omega; \mathbf{x}).$$

*Proof.* Without loss of generality, we assume that  $\Omega = \mathbb{H}$  — the formula for general polygons follows by conformal invariance. Also, thanks to Lemma 4.19 it suffices to prove the asserted identity for  $s$  such that  $\{a_s, b_s\} = \{1, 2b\} \in \alpha$ . Now, it suffices to prove that the marginal law of  $\eta_s$  under  $\vec{\text{P}}_\alpha = \vec{\text{P}}_\alpha(\mathbb{H}; \mathbf{x})$  equals the marginal law of  $\eta_s$  under  $\text{P}_\alpha^{(s)} = \text{P}_\alpha^{(s)}(\mathbb{H}; \mathbf{x})$ . For any bounded continuous function  $F$ , we have

$$\begin{aligned} \vec{\text{E}}_\alpha[F(\eta_s)] &= \vec{\text{E}}_N[F(\eta_s) \mid \zeta(\vec{\eta}) = \alpha] = \frac{\vec{\text{E}}_N[F(\eta_s) \mathbb{1}\{\zeta(\vec{\eta}) = \alpha\}]}{\vec{\text{P}}_N[\zeta(\vec{\eta}) = \alpha]} && \text{[by (4.30)]} \\ &= \text{E}_N^{x_1 \rightarrow x_{2b}}[F(\eta_s) \mathbb{1}\{\zeta(\vec{\eta}) = \alpha\} \mid \zeta(\vec{\eta}) = \alpha] = \frac{\text{E}_N^{x_1 \rightarrow x_{2b}}[F(\eta_s) \mathbb{1}\{\zeta(\vec{\eta}) = \alpha\}]}{\text{P}_N^{x_1 \rightarrow x_{2b}}[\zeta(\vec{\eta}) = \alpha]} && \text{[by (3.12)]} \end{aligned}$$

using Definition 3.6. The computations made in the proof of Lemma 4.7 imply that

$$\begin{aligned} \frac{\text{E}_N^{x_1 \rightarrow x_{2b}}[F(\eta_s) \mathbb{1}\{\zeta(\vec{\eta}) = \alpha\}]}{\text{P}_N^{x_1 \rightarrow x_{2b}}[\zeta(\vec{\eta}) = \alpha]} &= \frac{\vec{\mathcal{Z}}_{\triangleleft}(\mathbb{H}; x_1, x_{2b})}{\vec{\mathcal{Z}}_\alpha(\mathbb{H}; \mathbf{x})} \mathbb{E}(\mathbb{H}; x_1, x_{2b}) \left[ F(\eta) \mathbb{1}\{\mathcal{E}_\alpha(\eta)\} \left( \prod_{m=1}^r \vec{\mathcal{Z}}_{\alpha_m^R} \right) \left( \prod_{m=1}^l \vec{\mathcal{Z}}_{\alpha_m^L} \right) \right] \\ &= \text{E}_\alpha^{(s)}[F(\eta_s)], \end{aligned}$$

where  $\text{E}_\alpha^{(s)}$  is the expected value under the measure  $\text{E}_\alpha^{(s)}$ . This shows that  $\vec{\text{E}}_\alpha[F(\eta_s)] = \text{E}_\alpha^{(s)}[F(\eta_s)]$ .  $\square$

**Corollary 4.21.** Fix  $\kappa \in (4, 8)$ . For any  $N \geq 1$  and  $\alpha \in \text{LP}_N$ , the measure  $\vec{\text{P}}_\alpha(\Omega; \mathbf{x})$  defined in (4.30) is a global  $N$ - $\text{SLE}_\kappa$  associated to  $\alpha$  (as in Definition 1.20).

*Proof.* When  $N = 1$ , the law  $\vec{P}_{\triangleleft}(\Omega; x_1, x_2) = \vec{P}_1(\Omega; x_1, x_2)$  is just the law of chordal  $\text{SLE}_\kappa$  in  $(\Omega; x_1, x_2)$ . The claim for  $N \geq 2$  then follows by induction: indeed, the resampling property follows directly from construction for the measures  $P_\alpha^{(s)}(\Omega; \mathbf{x})$ , which all coincide with  $\vec{P}_\alpha(\Omega; \mathbf{x})$  by Lemma 4.20.  $\square$

*Proof of Theorem 1.21.* Existence is given by Corollary 4.21 for  $\kappa \in (4, 8)$  and [KL07] and [PW19, Theorem 1.3] for  $\kappa \in (0, 4]$ ; and uniqueness follows from Zhan's recent work [Zha23, Theorem 4.2] for  $\kappa \in (4, 8)$  and [BPW21, Theorem 1.2] for  $\kappa \in (0, 4]$ . It remains to derive the marginal law of  $\eta_s$  for each  $1 \leq s \leq N$ , that is, to verify that it is a Loewner chain associated to the pure partition function  $\mathcal{Z}_\alpha$ .

By symmetry, it suffices to consider the case where  $\Omega = \mathbb{H}$  and  $\{a_s, b_s\} = \{1, 2b\} \in \alpha$ . Let  $\eta$  be a chordal  $\text{SLE}_\kappa$  curve  $\eta$  on  $(\mathbb{H}; x_1, x_{2b})$ , with Loewner driving function  $W$ . Define  $\tau_\epsilon$  to be the hitting time of  $\eta$  at the  $\epsilon$ -neighborhood of  $(x_2, +\infty)$ . We use the same notations as in Lemma 4.7. For any bounded continuous function  $F$ , using Lemma 4.20, Lemma 4.7, and conformal covariance, we obtain

$$\begin{aligned} & \vec{E}_\alpha(\mathbb{H}; \mathbf{x})[F(\eta_s[0, \tau_\epsilon])] \\ &= E_\alpha^{(s)}(\mathbb{H}; \mathbf{x})[F(\eta_s[0, \tau_\epsilon])] \\ &= \frac{\vec{Z}_{\triangleleft}(\mathbb{H}; x_1, x_{2b})}{\vec{Z}_\alpha(\mathbb{H}; \mathbf{x})} \mathbb{E}(\mathbb{H}; x_1, x_{2b}) \left[ F(\eta[0, \tau_\epsilon]) \mathbb{1}\{\mathcal{E}_\alpha(\eta)\} \left( \prod_{m=1}^r \vec{Z}_{\alpha_m^R} \right) \left( \prod_{m=1}^l \vec{Z}_{\alpha_m^L} \right) \right] \\ &= \frac{(x_{2b} - x_1)^{-2h}}{\vec{Z}_\alpha(\mathbb{H}; \mathbf{x})} \mathbb{E}(\mathbb{H}; x_1, x_{2b}) \left[ F(\eta[0, \tau_\epsilon]) \mathbb{1}\{\mathcal{E}_\alpha(\eta)\} \mathbb{E}(\mathbb{H}; x_1, x_{2b}) \left[ \left( \prod_{m=1}^r \vec{Z}_{\alpha_m^R} \right) \left( \prod_{m=1}^l \vec{Z}_{\alpha_m^L} \right) \mid \eta[0, \tau_\epsilon] \right] \right] \\ &= \mathbb{E}(\mathbb{H}; x_1, x_{2b}) \left[ F(\eta[0, \tau_\epsilon]) \mathbb{1}\{\mathcal{E}_\alpha(\eta)\} \frac{|x_{2b} - x_1|^{-2h}}{|g_{\tau_\epsilon}(x_{2b}) - W_{\tau_\epsilon}|^{-2h}} \frac{\mathcal{Z}_\alpha(\mathbb{H}; W_{\tau_\epsilon}, g_{\tau_\epsilon}(\dot{\mathbf{x}}_1))}{\vec{Z}_\alpha(\mathbb{H}; \mathbf{x})} \prod_{\substack{1 \leq j \leq 2N \\ j \neq 1, 2b}} g'_{\tau_\epsilon}(x_j)^h \right], \end{aligned}$$

where  $g_t: \mathbb{H} \setminus K_t \rightarrow \mathbb{H}$  are the conformal maps normalized such that  $g_t(x_{2b}) = x_{2b}$  for all  $t \leq \tau_\epsilon$ , with  $\mathbb{H} \setminus K_t$  being the unbounded component of  $\mathbb{H} \setminus \eta[0, t]$  (containing in particular the target point  $x_{2b}$ ).

Now, recall from, e.g., [SW05] that the law of  $\eta$  up to the stopping time  $\tau_\epsilon$  is the same as that of the chordal  $\text{SLE}_\kappa$  curve  $\tilde{\eta}$  on  $(\mathbb{H}; x_1, \infty)$  weighted by  $\{M_t/M_0\}_{0 \leq t \leq \tau_\epsilon}$  with the martingale

$$M_t := |\tilde{g}_t(x_{2b}) - \tilde{W}_t|^{-2h} \tilde{g}'_t(x_{2b})^h,$$

where  $(\tilde{g}_t: t \geq 0)$  are the conformal maps corresponding to  $\tilde{\eta}$  normalized at infinity<sup>21</sup> and  $\tilde{W}$  its driving function. Hence, we conclude that the law of  $\eta_s$  under  $\vec{P}_\alpha(\mathbb{H}; \mathbf{x})$  is the same as the law of  $\tilde{\eta}$  weighted by

$$\prod_{j=2}^{2N} \tilde{g}'_{\tau_\epsilon}(x_j)^h \times \frac{\mathcal{Z}_\alpha(\mathbb{H}; \tilde{W}_{\tau_\epsilon}, \tilde{g}_{\tau_\epsilon}(\dot{\mathbf{x}}_1))}{\mathcal{Z}_\alpha(\mathbb{H}; \mathbf{x})}.$$

In other words, the driving function of  $\eta_s$  under  $\vec{P}_\alpha(\mathbb{H}; \mathbf{x})$  solves the SDE system (1.45) with  $\mathcal{Z} = \mathcal{Z}_\alpha$  and  $i = 1$ , that is, the Loewner chain associated to  $\mathcal{Z}_\alpha$ .  $\square$

**Corollary 4.22.** *Fix  $\kappa \in (4, 8)$ . For any  $N \geq 1$ , the measures  $\vec{P}_N$  and  $\tilde{P}_N$  are the same.*

*Proof.* On the one hand, by Lemma 4.3 and Definition 4.1, we have

$$\vec{P}_N[\zeta(\vec{\eta}) = \alpha] = \tilde{P}_N[\zeta(\vec{\eta}) = \alpha], \quad \alpha \in \text{LP}_N.$$

On the other hand, we have

$$\vec{P}_\alpha(\Omega; \mathbf{x}) := \vec{P}_N(\Omega; \mathbf{x})[\cdot \mid \zeta(\vec{\eta}) = \alpha] = \tilde{P}_N(\Omega; \mathbf{x})[\cdot \mid \zeta(\vec{\eta}) = \alpha] =: \tilde{P}_\alpha(\Omega; \mathbf{x}), \quad \alpha \in \text{LP}_N,$$

since by Corollary 4.21 (and similar arguments for  $\tilde{P}_\alpha$ ), both measures  $\vec{P}_\alpha$  and  $\tilde{P}_\alpha$  defined in (4.30) are global  $N$ - $\text{SLE}_\kappa$  measures associated to  $\alpha$  — but such measures are unique by [Zha23, Theorem 4.2].  $\square$

<sup>21</sup>That is, solutions to (3.1) with  $\lim_{z \rightarrow \infty} |\tilde{g}_t(z) - z| = 0$ .

**Remark 4.23.** Let us argue that the SLE $_{\kappa}$  partition functions  $\tilde{\mathcal{Z}}_{\alpha}$  constructed in [AHSY23] are the same as pure partition functions  $\mathcal{Z}_{\alpha}$  in Definition 1.9, up to a multiplicative constant. In [AHSY23], the authors prove that any curve in a global multiple SLE $_{\kappa}$  has the same law as the Loewner chain associated to their partition functions  $\tilde{\mathcal{Z}}_{\alpha}$ . Combining with the uniqueness of global multiple SLE $_{\kappa}$  measures [Zha23] and Theorem 1.21, we see that  $\partial_i \log \tilde{\mathcal{Z}}_{\alpha}$  is the same as  $\partial_i \log \mathcal{Z}_{\alpha}$  in (1.45) for all  $i \in \{1, 2, \dots, 2N\}$ . This implies that  $\tilde{\mathcal{Z}}_{\alpha}$  is the same as  $\mathcal{Z}_{\alpha}$  up to a multiplicative constant.

## 5 Tying up loose ends

In this section, we investigate the solutions to the PDEs in the space  $\mathcal{S}_N$  defined in (2.21). Specifically, we finish the proofs of the main results: key properties of the pure partition functions with  $\kappa \in (0, 8)$  (main part of Theorem 1.10) in Sections 5.1–5.2; key properties of the Coulomb gas integral functions (Theorem 1.5) and their relation to the pure partition functions (Proposition 1.11) in Section 5.3; and the renormalization of these functions for special values of  $\kappa$  (Propositions 1.7, 1.8, and Item 2 of Theorem 1.10) in Section 5.4.

### 5.1 Pure partition functions: linear independence

In this section, we prove the linear independence of the pure partition functions (which is similar to [PW19, Proposition 4.5]). By [FK15b, Lemma 1], the asymptotics properties can be used to uniquely characterize elements in  $\mathcal{S}_N$ . To analyze them, we consider linear functionals  $\mathcal{L}_{\beta}: \mathcal{S}_N \rightarrow \mathbb{C}$  defined in terms of iterated limits (allowable sequences of limits in the terminology of [FK15a, KP16]). For a link pattern

$$\beta = \{\{a_1, b_1\}, \{a_2, b_2\}, \dots, \{a_N, b_N\}\}, \quad a_r < b_r, \text{ for all } 1 \leq r \leq N, \quad (5.1)$$

where the links  $\{a_r, b_r\}$  are not necessarily ordered as in (1.2), we say that the ordering of links is *allowable* if all links of  $\beta$  as in (5.1) can be removed in the order  $\{a_1, b_1\}, \{a_2, b_2\}, \dots, \{a_N, b_N\}$  in such a way that at each step, the link to be removed connects two consecutive indices (see [PW19, Figure 2.1] for an illustration and, e.g., [KP16, Section 3.5] for a more formal definition). Suppose the ordering (5.1) of the links in  $\beta$  is allowable. Fix points  $\xi_r \in (x_{a_r-1}, x_{b_r+1})$  for all  $r \in \{1, \dots, N\}$ , with the convention that  $x_0 = -\infty$  and  $x_{2N+1} = +\infty$ . It was proved in [FK15b, Lemma 10] that the following sequence of limits exists and is finite for any solution  $F \in \mathcal{S}_N$ :

$$\mathcal{L}_{\beta}(F) := \lim_{x_{a_N}, x_{b_N} \rightarrow \xi_N} \cdots \lim_{x_{a_1}, x_{b_1} \rightarrow \xi_1} (x_{b_N} - x_{a_N})^{2h(\kappa)} \cdots (x_{b_1} - x_{a_1})^{2h(\kappa)} F(x_1, \dots, x_{2N}). \quad (5.2)$$

Furthermore, by [FK15b, Lemma 12], any other allowable ordering of the links of  $\beta$  gives the same limit (5.2). Therefore, for each  $\beta \in \text{LP}_N$  with any choice of allowable ordering of links, (5.2) defines a linear functional  $\mathcal{L}_{\beta}: \mathcal{S}_N \rightarrow \mathbb{C}$ . Finally, it was proved in [FK15c, Theorem 8] that, for any  $\kappa \in (0, 8)$ , the collection  $\{\mathcal{L}_{\beta}: \beta \in \text{LP}_N\}$  is a basis for the dual space  $\mathcal{S}_N^*$  of the solution space  $\mathcal{S}_N$ .

**Corollary 5.1.** *For  $\kappa \in (0, 8)$ , the collection  $\{\mathcal{Z}_{\alpha}^{(\kappa)}: \alpha \in \text{LP}_N\}$  is linearly independent.*

*Proof.* By Proposition 4.16, we have  $\mathcal{L}_{\beta}(\mathcal{Z}_{\alpha}^{(\kappa)}) = \delta_{\alpha, \beta}$  for all  $\alpha, \beta \in \text{LP}_N$ . This shows that the set  $\{\mathcal{Z}_{\alpha}^{(\kappa)}: \alpha \in \text{LP}_N\}$  is linearly independent as a dual basis of the linear operators  $\{\mathcal{L}_{\beta}: \beta \in \text{LP}_N\}$ .  $\square$

### 5.2 Pure partition functions: continuity in $\kappa$

The main goal of this section is to show the continuity in  $\kappa$  of pure partition functions (Item 1 in Theorem 1.10). This is easier to derive for  $\kappa \in (0, 6)$  (Lemma 5.2), because in that case the pure partition functions admit a very strong upper bound (5.3), which is significantly better than (PLB). The range

$\kappa \in (4, 8)$  can be handled using Coulomb gas integrals and their relation to the pure partition functions from Proposition 1.11 (see Lemmas 5.3 & 5.4). We are not aware of a direct proof only using the probabilistic construction for the range  $\kappa \in (4, 8)$ .

**Lemma 5.2.** *The pure partition functions  $\{\mathcal{Z}_\alpha^{(\kappa)} : \alpha \in \text{LP}\}$  are continuous in  $(0, 6) \ni \kappa$ .*

*Proof.* We can use induction, the cascade property, and dominated convergence to verify the continuity of  $\mathcal{Z}_\alpha^{(\kappa)}$  in  $\kappa$ . Indeed, the cascade property (4.7) holds for all  $\kappa \in (0, 8)$  by Lemma 4.7 (for  $\kappa \in (4, 8)$ ), by [PW19, Proposition 3.5] (for  $\kappa \in (0, 4]$ ), and by [Wu20, Corollary 6.8] (for  $\kappa \in (0, 6]$ ). Moreover, when  $\kappa \in (0, 6]$ , the pure partition functions admit the strong upper bound [PW19, Lemma 4.1] (for  $\kappa \in (0, 4]$ ), [Wu20, Theorem 1.7] (for  $\kappa \in (0, 6]$ )

$$\mathcal{Z}_\alpha^{(\kappa)}(\Omega; x_1, \dots, x_{2N}) \leq \prod_{\{a,b\} \in \alpha} H(\Omega; x_a, x_b)^{h(\kappa)}, \quad (5.3)$$

where  $H(\Omega; x, y)$  is the (boundary) Poisson kernel. This upper bound plays an essential role in the following proof of continuity, which is obstructed for  $\kappa \geq 6$  because we do not have such a strong bound available, nor could we use the Poisson kernel monotonicity (3.3) for  $h(\kappa) \leq 0$ .

For the base case  $N = 1$ , the pure partition function is explicit,  $\mathcal{Z}_{\triangleleft}^{(\kappa)}(x_1, x_2) = (x_2 - x_1)^{-2h(\kappa)}$ , and obviously continuous. Fix  $N \geq 2$ , and suppose the members in the collection  $\{\mathcal{Z}_\alpha^{(\kappa)} : \alpha \in \text{LP}_n, n \leq N-1\}$  are continuous in  $(0, 6) \ni \kappa$ . Fix  $\kappa \in (0, 6)$  and  $\alpha \in \text{LP}_N$ , with  $\{1, 2b\} \in \alpha$ . Let  $\kappa_m \in (0, 6)$  be a converging sequence,  $\kappa_m \rightarrow \kappa$  as  $m \rightarrow \infty$ . Denote by  $\eta$  the  $\text{SLE}_\kappa$  curve from  $x_1$  to  $x_{2b}$  and by  $\eta_m$  the  $\text{SLE}_{\kappa_m}$  curve from  $x_1$  to  $x_{2b}$  for each  $m \geq 1$ . By [KS17, Theorem 1.10], we know that the law of  $\eta_m$  converges to the law of  $\eta$  weakly under the metric (1.43). By Skorokhod's representation theorem, there is a coupling of  $\eta$  and  $\{\eta_m\}_{m \geq 1}$ , which we denote by  $\mathbb{P}$ , such that under  $\mathbb{P}$ , random curve  $\eta_m$  converges to  $\eta$  almost surely as  $m \rightarrow \infty$ . Using the cascade property (4.7) or [Wu20, Corollary 6.8], we may thus conclude that

$$\mathcal{Z}_\alpha^{(\kappa_m)}(\mathbf{x}) = \mathcal{Z}_{\triangleleft}^{(\kappa_m)}(x_1, x_{2b}) \mathbb{E}[\mathbb{1}\{\mathcal{E}_\alpha(\eta_m)\} \mathcal{Z}_{\alpha/\{1,2b\}}^{(\kappa_m)}(\hat{\mathbb{H}}_{\eta_m}; \ddot{\mathbf{x}}_{1,2b})], \quad (5.4)$$

where  $\mathbb{E}$  is the expectation with respect to  $\mathbb{P}$ . Because the Poisson kernel is monotone by (3.3), and  $h(\kappa_m) > 0$  for all  $\kappa_m \in (0, 6)$ , we see from (5.3) that the functions inside the expected value are bounded:

$$\begin{aligned} \mathcal{Z}_{\alpha/\{1,2b\}}^{(\kappa_m)}(\hat{\mathbb{H}}_{\eta_m}; \ddot{\mathbf{x}}_{1,2b}) &\leq \prod_{\substack{D \text{ c.c. of } \mathbb{H} \setminus \eta_m \\ \bar{D} \cap \{x_1, \dots, x_{2N}\} \setminus \{x_1, x_{2b}\} \neq \emptyset}} \prod_{\{a,b\} \in \alpha^D} H(D; x_a, x_b)^{h(\kappa)} \\ &\leq \prod_{\substack{D \text{ c.c. of } \mathbb{H} \setminus \eta_m \\ \bar{D} \cap \{x_1, \dots, x_{2N}\} \setminus \{x_1, x_{2b}\} \neq \emptyset}} \prod_{\{a,b\} \in \alpha^D} H(\mathbb{H}; x_a, x_b)^{h(\kappa)}. \end{aligned}$$

Therefore, combined with the cascade property (4.7) or [Wu20, Corollary 6.8], the dominated convergence theorem and the induction hypothesis guarantee that

$$\lim_{m \rightarrow \infty} \mathcal{Z}_\alpha^{(\kappa_m)}(\mathbf{x}) = \mathcal{Z}_{\triangleleft}^{(\kappa)}(x_1, x_{2b}) \mathbb{E}[\mathbb{1}\{\mathcal{E}_\alpha(\eta)\} \mathcal{Z}_{\alpha/\{1,2b\}}^{(\kappa)}(\hat{\mathbb{H}}_{\eta_m}; \ddot{\mathbf{x}}_{1,2b})] = \mathcal{Z}_\alpha^{(\kappa)}(\mathbf{x}),$$

which proves that  $\mathcal{Z}_\alpha^{(\kappa)}(\mathbf{x})$  is continuous on  $(0, 6) \ni \kappa$ .  $\square$

**Lemma 5.3.** *The relation (1.24) holds for all  $\kappa \in (0, 8) \setminus \{\frac{8}{m} : m \in \mathbb{Z}_{>0}\}$ .*

*Proof.* We use the uniqueness from Lemma 2.11 of elements in the PDE solution space  $\mathcal{S}_N$  (2.21) singled out by their asymptotic properties. On the one hand, we already know from Corollary 2.13 that  $\{\mathcal{F}_\beta^{(\kappa)} : \beta \in \text{LP}_N\} \subset \mathcal{S}_N$ , and that they satisfy the recursive asymptotic property ( $\mathcal{F}$ -ASY). On the other hand, we



also know that  $\{\mathcal{Z}_\alpha^{(\kappa)} : \alpha \in \text{LP}_N\} \subset \mathcal{S}_N$ , thanks to Proposition 4.12, Lemma 4.5, and Lemma 4.13. It thus remains to check that the right-hand side of (1.24),

$$\mathcal{G}_\beta^{(\kappa)}(\mathbf{x}) := \sum_{\alpha \in \text{LP}_N} \mathcal{M}_\nu(\alpha, \beta) \mathcal{Z}_\alpha^{(\kappa)}(\mathbf{x}), \quad \mathbf{x} \in \mathfrak{X}_{2N}, \beta \in \text{LP}_N, \quad (5.5)$$

satisfy the same recursive asymptotics property property ( $\mathcal{F}$ -ASY) as the left-hand side  $\mathcal{F}_\beta^{(\kappa)}$ :

$$\begin{aligned} \lim_{x_j, x_{j+1} \rightarrow \xi} \frac{\mathcal{G}_\beta^{(\kappa)}(\mathbf{x})}{(x_{j+1} - x_j)^{-2h(\kappa)}} &= \sum_{\alpha \in \text{LP}_N} \mathcal{M}_\nu(\alpha, \beta) \lim_{x_j, x_{j+1} \rightarrow \xi} \frac{\mathcal{Z}_\alpha^{(\kappa)}(\mathbf{x})}{(x_{j+1} - x_j)^{-2h(\kappa)}} && \text{[by (1.24)]} \\ &= \sum_{\substack{\alpha \in \text{LP}_N \\ \{j, j+1\} \in \alpha}} \mathcal{M}_\nu(\alpha, \beta) \mathcal{Z}_{\alpha/\{j, j+1\}}^{(\kappa)}(\ddot{\mathbf{x}}_j) && \text{[by (ASY)]} \\ &= \begin{cases} \nu(\kappa) \sum_{\gamma \in \text{LP}_{N-1}} \mathcal{M}_\nu(\gamma, \beta/\{j, j+1\}) \mathcal{Z}_\gamma^{(\kappa)}(\ddot{\mathbf{x}}_j), & \text{if } \{j, j+1\} \in \beta, \\ \sum_{\gamma \in \text{LP}_{N-1}} \mathcal{M}_\nu(\gamma, \wp_j(\beta)/\{j, j+1\}) \mathcal{Z}_\gamma^{(\kappa)}(\ddot{\mathbf{x}}_j), & \text{if } \{j, j+1\} \notin \beta, \end{cases} \\ &= \begin{cases} \nu(\kappa) \mathcal{G}_{\beta/\{j, j+1\}}^{(\kappa)}(\ddot{\mathbf{x}}_j), & \text{if } \{j, j+1\} \in \beta, \\ \mathcal{G}_{\wp_j(\beta)/\{j, j+1\}}^{(\kappa)}(\ddot{\mathbf{x}}_j), & \text{if } \{j, j+1\} \notin \beta, \end{cases} && \text{[by (1.24)]} \end{aligned}$$

using the bijection  $\alpha \leftrightarrow \alpha/\{j, j+1\}$  between  $\{\alpha \in \text{LP}_N : \{j, j+1\} \in \alpha\}$  and  $\text{LP}_{N-1}$  and the facts that  $\mathcal{M}_\nu(\alpha, \beta) = \nu(\kappa) \mathcal{M}_\nu(\alpha/\{j, j+1\}, \beta/\{j, j+1\})$  and  $\mathcal{M}_\nu(\alpha, \beta) = \mathcal{M}_\nu(\alpha/\{j, j+1\}, \wp_j(\beta)/\{j, j+1\})$ .  $\square$

**Lemma 5.4.** *The pure partition functions  $\{\mathcal{Z}_\alpha^{(\kappa)} : \alpha \in \text{LP}\}$  are continuous in  $(4, 8) \ni \kappa$ .*

*Proof.* From Lemma 5.3, for  $\kappa \in (4, 8)$ , we have

$$\mathcal{F}_\beta^{(\kappa)}(\mathbf{x}) = \sum_{\alpha \in \text{LP}_N} \mathcal{M}_\nu(\alpha, \beta) \mathcal{Z}_\alpha^{(\kappa)}(\mathbf{x}), \quad \mathbf{x} \in \mathfrak{X}_{2N}, \beta \in \text{LP}_N. \quad (5.6)$$

This relation plays an essential role in the following proof of continuity by induction on  $N \geq 1$ .

For the base case  $N = 1$ , the pure partition function is explicit,  $\mathcal{Z}_\alpha^{(\kappa)}(x_1, x_2) = (x_2 - x_1)^{-2h(\kappa)}$ , and obviously continuous. Fix  $N \geq 2$ , and suppose the members in the collection  $\{\mathcal{Z}_\alpha^{(\kappa)} : \alpha \in \text{LP}_n, n \leq N-1\}$  are continuous in  $(4, 8) \ni \kappa$ . Fix  $\kappa \in (4, 8)$  and  $\alpha \in \text{LP}_N$ , with  $\{1, 2b\} \in \alpha$ . Let  $\kappa_m \in (4, 8)$  be a converging sequence,  $\kappa_m \rightarrow \kappa$  as  $m \rightarrow \infty$ . With (5.6) and the cascade property (5.4) (Lemma 4.7), we obtain:

- On the one hand, for the quantities on the right hand-side of (5.4), by the induction hypothesis, we have, on the event  $\bigcup_{i \geq 0} \bigcap_{m \geq i} \mathcal{E}_\alpha(\eta_m) \supset \mathcal{E}_\alpha(\eta)$ ,

$$\lim_{m \rightarrow \infty} \mathcal{Z}_{\alpha/\{1, 2b\}}^{(\kappa_m)}(\hat{\mathbb{H}}_{\eta_m}; \ddot{\mathbf{x}}_{1, 2b}) = \mathcal{Z}_{\alpha/\{1, 2b\}}^{(\kappa)}(\hat{\mathbb{H}}_\eta; \ddot{\mathbf{x}}_{1, 2b}),$$

so Fatou's lemma yields

$$\liminf_{m \rightarrow \infty} \mathcal{Z}_\alpha^{(\kappa_m)}(\mathbf{x}) \geq \mathcal{Z}_\alpha^{(\kappa)}(\mathbf{x}). \quad (5.7)$$

- On the other hand, as  $\mathcal{F}_\beta^{(\kappa)}$  is continuous in  $\kappa \in (4, 8)$ , we conclude from (5.6) that

$$\limsup_{m \rightarrow \infty} \sum_{\alpha \in \text{LP}_N} \mathcal{M}_\nu(\alpha, \beta) \mathcal{Z}_\alpha^{(\kappa_m)}(\mathbf{x}) = \sum_{\alpha \in \text{LP}_N} \mathcal{M}_\nu(\alpha, \beta) \mathcal{Z}_\alpha^{(\kappa)}(\mathbf{x}). \quad (5.8)$$

Combining (5.7) and (5.8) and the fact that  $\mathcal{M}_\nu(\alpha, \beta) > 0$  when  $\kappa \in (4, 8)$ , we obtain

$$\lim_{m \rightarrow \infty} \mathcal{Z}_\alpha^{(\kappa_m)}(\mathbf{x}) = \mathcal{Z}_\alpha^{(\kappa)}(\mathbf{x}),$$

which proves that  $\mathcal{Z}_\alpha^{(\kappa)}(\mathbf{x})$  is continuous on  $(4, 8) \ni \kappa$ .  $\square$

### 5.3 Finishing the proof of Theorem 1.5 and Proposition 1.11

The primary goal of this section is to derive the relation (1.24) between the pure partition functions and the Coulomb gas integrals (which is similar to [FPW24a, Proposition 1.10]) for all  $\kappa \in (0, 8)$ . We also prove the continuity properties of the Coulomb gas integrals, thereby completing the proof of Theorem 1.5.

**Lemma 5.5.** *The meander matrix (1.3) is invertible if and only if  $\kappa$  is not one of the exceptional values*

$$\kappa = \frac{4p}{p'}, \quad p, p' \in \mathbb{Z}_{>0} \text{ coprime and } 2 \leq p \leq N + 1. \quad (5.9)$$

*Proof.* By [DFGG97, Eq. (5.6)], the determinant of the meander matrix (1.3) is given by

$$\det \mathcal{M}_\nu = \prod_{n=1}^N U_n(\nu)^{a_{N,n}}, \quad \text{where } a_{N,n} = \binom{2N}{N-n} - 2 \binom{2N}{N-n-1} + \binom{2N}{N-n-2},$$

and  $U_n(\cdot)$  are Chebyshev polynomials of the second kind. From [DFGG97, Eq. (4.2)], we obtain

$$U_n(z + z^{-1}) = \frac{z^{n+1} - z^{-n-1}}{z - z^{-1}}.$$

Now, recall that  $\nu = \nu(\kappa) = -\exp(4\pi i/\kappa) - \exp(-4\pi i/\kappa)$ . Therefore, we have  $\det \mathcal{M}_\nu = 0$  if and only if

$$\exp(8(n+1)\pi i/\kappa) = 1 \quad \text{for some } n \in \{1, \dots, N\},$$

which is equivalent to (5.9). □

*Proof of Theorem 1.5 and Proposition 1.11.* As in (5.5), denote by  $\mathcal{G}_\beta^{(\kappa)}$  the right-hand side of (1.24). Since both  $\mathcal{M}_{\nu(\kappa)}(\alpha, \beta)$  and  $\mathcal{Z}_\alpha^{(\kappa)}$  are continuous in  $(0, 8) \ni \kappa$  (by Lemmas 5.2 & 5.4), we see that  $\mathcal{G}_\beta^{(\kappa)}$  is continuous in  $(0, 8) \ni \kappa$ . Therefore, the function  $\mathcal{F}_\beta^{(\kappa)} = \mathcal{G}_\beta^{(\kappa)}$  can be continuously extended to  $\kappa \in (0, 8)$  and the continuous extensions satisfy (PDE), (COV), (PLB), ( $\mathcal{F}$ -ASY), and (1.24) for all  $\kappa \in (0, 8)$ .

(POS): For all  $\kappa \in \bigcup_{m \in \mathbb{Z}_{>0}} \left( \frac{8}{4m-1}, \frac{8}{4m-3} \right)$ , since  $\mathcal{M}_\nu(\alpha, \beta) > 0$  and  $\mathcal{Z}_\alpha^{(\kappa)} > 0$ , we have

$$\mathcal{F}_\beta^{(\kappa)}(\mathbf{x}) = \sum_{\alpha \in \text{LP}_N} \mathcal{M}_\nu(\alpha, \beta) \mathcal{Z}_\alpha^{(\kappa)}(\mathbf{x}) > 0, \quad \mathbf{x} \in \mathfrak{X}_{2N}.$$

(NUL): For all  $\kappa \in \left\{ \frac{8}{2m+1} : m \in \mathbb{Z}_{>0} \right\}$ , since  $\mathcal{M}_\nu(\alpha, \beta) = 0$ , we have

$$\mathcal{F}_\beta^{(\kappa)}(\mathbf{x}) = \sum_{\alpha \in \text{LP}_N} \mathcal{M}_\nu(\alpha, \beta) \mathcal{Z}_\alpha^{(\kappa)}(\mathbf{x}) = 0, \quad \mathbf{x} \in \mathfrak{X}_{2N}.$$

(SGN): Note that when  $\kappa \in \bigcup_{m \in \mathbb{Z}_{>0}} \left( \frac{8}{4m+1}, \frac{8}{4m-1} \right)$ , the fugacity is  $\nu(\kappa) < 0$ . We claim that there are

$$\mathbf{x}^+ = \mathbf{x}^+(\kappa, \beta) \in \mathfrak{X}_{2N}, \quad \mathbf{x}^0 = \mathbf{x}^0(\kappa, \beta) \in \mathfrak{X}_{2N}, \quad \mathbf{x}^- = \mathbf{x}^-(\kappa, \beta) \in \mathfrak{X}_{2N},$$

such that

$$\mathcal{F}_\beta^{(\kappa)}(\mathbf{x}^+) > 0, \quad \mathcal{F}_\beta^{(\kappa)}(\mathbf{x}^0) = 0, \quad \mathcal{F}_\beta^{(\kappa)}(\mathbf{x}^-) < 0. \quad (5.10)$$

We prove (5.10) by induction on  $N \geq 2$ . The base case  $N = 2$  is governed by Proposition D.4 in Appendix D. Assuming that (1.17) holds for  $\{\mathcal{F}_\beta^{(\kappa)} : \beta \in \text{LP}_n, n \leq N-1\}$ , fix  $\beta \in \text{LP}_N$ . Without much loss of generality, suppose  $\{1, 2\} \in \beta$  (the case  $\{1, 2\} \notin \beta$  is similar). The induction hypothesis for  $\beta/\{1, 2\} \in \text{LP}_{N-1}$  guarantees that there exists a point  $\mathbf{y}^+ = \mathbf{y}^+(\kappa, \beta/\{1, 2\}) \in \mathfrak{X}_{2N-2}$  such that  $\mathcal{F}_{\beta/\{1, 2\}}^{(\kappa)}(\mathbf{y}^+) > 0$ . Now, the asymptotics property ( $\mathcal{F}$ -ASY) shows that

$$\lim_{x_1, x_2 \rightarrow \xi} \frac{\mathcal{F}_\beta^{(\kappa)}(\mathbf{x})}{(x_2 - x_1)^{-2h(\kappa)}} = \mathcal{F}_{\beta/\{1, 2\}}^{(\kappa)}(\mathbf{y}^+) > 0,$$

where  $\mathbf{x} = (x_1, x_2, y_1, \dots, y_{2N-2}) \in \mathfrak{X}_{2N}$  and  $\mathbf{y}^+ = (y_1, \dots, y_{2N-2}) \in \mathfrak{X}_{2N-2}$ . Therefore, there exists a point  $\mathbf{x}^+ = \mathbf{x}^+(\kappa, \beta) \in \mathfrak{X}_{2N}$  such that  $\mathcal{F}_\beta^{(\kappa)}(\mathbf{x}^+) > 0$ . In a similar manner, we see that there exists a point  $\mathbf{x}^- = \mathbf{x}^-(\kappa, \beta) \in \mathfrak{X}_{2N}$  such that  $\mathcal{F}_\beta^{(\kappa)}(\mathbf{x}^-) < 0$ . However, the function  $\mathbf{x} \mapsto \mathcal{F}_\beta^{(\kappa)}(\mathbf{x})$  is continuous on  $\mathfrak{X}_{2N}$ . Thus, there exists a point  $\mathbf{x}^0 \in \mathfrak{X}_{2N}$  such that  $\mathcal{F}_\beta^{(\kappa)}(\mathbf{x}^0) = 0$ . This completes the proof of (SGN) (5.10).

Finally,  $\{\mathcal{F}_\beta^{(\kappa)} : \beta \in \text{LP}_N\}$  is linearly independent for  $\kappa$  in (1.25) because  $\{\mathcal{Z}_\alpha^{(\kappa)} : \alpha \in \text{LP}_N\}$  is linearly independent (by Lemma 5.1) and the meander matrix  $\mathcal{M}_\nu$  is invertible for these  $\kappa$  (by Lemma 5.5).  $\square$

## 5.4 Renormalized partition functions

We will show in this section that null partition functions can be positive and finite after a suitable renormalization (1.19), (1.20), and (1.23). Moreover, the renormalized partition functions are linearly independent and have asymptotic properties analogous to those of the generic ones (see Lemma 5.7).

**Lemma 5.6.** *The renormalized meander matrix (1.29) is invertible.*

*Proof.* This is immediate from [DFGG97, Eq. (5.18)], since  $\det \widehat{\mathcal{M}} = (\partial_\nu \det \mathcal{M}_\nu) \big|_{\nu=0} \neq 0$ .  $\square$

*Proof of Proposition 1.7.* From (1.24), for all  $\kappa' \in (0, 8)$ , we have

$$\mathcal{F}_\beta^{(\kappa')}(\mathbf{x}) = \sum_{\alpha \in \text{LP}_N} \mathcal{M}_{\nu(\kappa')}(\alpha, \beta) \mathcal{Z}_\alpha^{(\kappa')}(\mathbf{x}), \quad \mathbf{x} \in \mathfrak{X}_{2N}, \beta \in \text{LP}_N.$$

Thus, for  $\kappa \in \{\frac{8}{m} : m \in 2\mathbb{Z}_{>0} + 1\}$ , we obtain

$$\begin{aligned} \lim_{\kappa' \rightarrow \kappa} \frac{1}{\nu(\kappa')} \mathcal{F}_\beta^{(\kappa')}(\mathbf{x}) &= \sum_{\alpha \in \text{LP}_N} \left( \lim_{\kappa' \rightarrow \kappa} \frac{1}{\nu(\kappa')} \mathcal{M}_{\nu(\kappa')}(\alpha, \beta) \right) \mathcal{Z}_\alpha^{(\kappa')}(\mathbf{x}) \\ &= \sum_{\alpha \in \text{LP}_N} \widehat{\mathcal{M}}(\alpha, \beta) \mathcal{Z}_\alpha^{(\kappa)}(\mathbf{x}), \end{aligned} \quad [\text{by (1.29) and Lemmas 5.2 \& 5.4}]$$

using the definition of  $\widehat{\mathcal{M}}$  and the continuity of pure partition functions in  $\kappa$ . Now, note that

$$\lim_{\kappa' \rightarrow \kappa} \frac{(\kappa' - \kappa)}{\nu(\kappa')} = \frac{-\kappa^2}{8\pi \sin(4\pi/\kappa)}.$$

This proves Proposition 1.7.  $\square$

Moreover, from the above proof, we obtain

$$\widehat{\mathcal{F}}_\beta^{(\kappa)}(\mathbf{x}) = \sum_{\alpha \in \text{LP}_N} \widehat{\mathcal{M}}(\alpha, \beta) \mathcal{Z}_\alpha^{(\kappa)}(\mathbf{x}) \in (0, \infty), \quad \mathbf{x} \in \mathfrak{X}_{2N}, \beta \in \text{LP}_N, \quad (5.11)$$

where the positivity follows from the facts that the renormalized meander matrix has non-negative entries (1.29) and is invertible (Lemma 5.6) and from the positivity of  $\mathcal{Z}_\alpha^{(\kappa)}$  (Lemma 4.10). We can also verify that the renormalized partition functions  $\widehat{\mathcal{F}}_\beta^{(\kappa)}$  satisfy recursive asymptotics analogous to ( $\mathcal{F}$ -ASY).

**Lemma 5.7.** Fix  $\kappa \in \{\frac{8}{m} : m \in 2\mathbb{Z}_{>0} + 1\}$ . For the renormalized Coulomb gas integral (1.19), we have

$$\widehat{\mathcal{F}}_{\emptyset}^{(\kappa)} \equiv 1 \quad \text{and} \quad \widehat{\mathcal{F}}_{\underline{\alpha}}^{(\kappa)}(x_1, x_2) = (x_2 - x_1)^{-2h(\kappa)},$$

and for any  $N \geq 2$  and  $j \in \{1, 2, \dots, 2N - 1\}$  and for all  $\xi \in (x_{j-1}, x_{j+2})$ ,

$$\lim_{x_j, x_{j+1} \rightarrow \xi} \frac{\widehat{\mathcal{F}}_{\beta}^{(\kappa)}(\mathbf{x})}{(x_{j+1} - x_j)^{-2h(\kappa)}} = \begin{cases} 0, & \text{if } \{j, j+1\} \in \beta, \\ \widehat{\mathcal{F}}_{\varphi_j(\beta)/\{j, j+1\}}^{(\kappa)}(\check{\mathbf{x}}_j), & \text{if } \{j, j+1\} \notin \beta. \end{cases} \quad (\widehat{\mathcal{F}}\text{-ASY})$$

*Proof.* Fix  $N \geq 2$ . The asymptotics ( $\widehat{\mathcal{F}}$ -ASY) follow from the asymptotics of the right-hand side of (5.11), which is obtained easily from the asymptotics properties (ASY) of  $\mathcal{Z}_{\alpha}^{(\kappa)}$  from [PW19, Lemma 4.3] (for  $\kappa \in (0, 4]$ ) and from Proposition 4.16 (for  $\kappa \in (4, 8)$ ):

- if  $\{j, j+1\} \in \beta$ , we have

$$\lim_{x_j, x_{j+1} \rightarrow \xi} \frac{\widehat{\mathcal{F}}_{\beta}^{(\kappa)}(\mathbf{x})}{(x_{j+1} - x_j)^{-2h(\kappa)}} = \sum_{\substack{\alpha \in \text{LP}_N \\ \{j, j+1\} \notin \alpha}} \widehat{\mathcal{M}}(\alpha, \beta) \lim_{x_j, x_{j+1} \rightarrow \xi} \frac{\mathcal{Z}_{\alpha}^{(\kappa)}(\mathbf{x})}{(x_{j+1} - x_j)^{-2h(\kappa)}} = 0; \quad [\text{by (ASY)}]$$

- if  $\{j, j+1\} \notin \beta$ , we have

$$\begin{aligned} \lim_{x_j, x_{j+1} \rightarrow \xi} \frac{\widehat{\mathcal{F}}_{\beta}^{(\kappa)}(\mathbf{x})}{(x_{j+1} - x_j)^{-2h(\kappa)}} &= \sum_{\substack{\alpha \in \text{LP}_N \\ \{j, j+1\} \in \alpha}} \widehat{\mathcal{M}}(\alpha, \beta) \lim_{x_j, x_{j+1} \rightarrow \xi} \frac{\mathcal{Z}_{\alpha}^{(\kappa)}(\mathbf{x})}{(x_{j+1} - x_j)^{-2h(\kappa)}} \\ &= \sum_{\substack{\alpha \in \text{LP}_N \\ \{j, j+1\} \in \alpha}} \widehat{\mathcal{M}}(\alpha, \beta) \mathcal{Z}_{\alpha/\{j, j+1\}}^{(\kappa)}(\check{\mathbf{x}}_j) && [\text{by (ASY)}] \\ &= \widehat{\mathcal{F}}_{\varphi_j(\beta)/\{j, j+1\}}^{(\kappa)}(\check{\mathbf{x}}_j), && [\text{by (5.11)}] \end{aligned}$$

using the bijection  $\alpha \leftrightarrow \alpha/\{j, j+1\}$  between  $\{\alpha \in \text{LP}_N : \{j, j+1\} \in \alpha\}$  and  $\text{LP}_{N-1}$  and the fact that  $\widehat{\mathcal{M}}(\alpha, \beta) = \widehat{\mathcal{M}}(\alpha/\{j, j+1\}, \varphi_j(\beta)/\{j, j+1\})$  to arrive at the last expression.  $\square$

*Proof of Proposition 1.8.* Recall from (1.11) and (2.2) that

$$\mathcal{H}_{\beta}^{(\kappa')}(\mathbf{x}) := \oint_{\vartheta_1^{\beta}} du_1 \oint_{\vartheta_2^{\beta}} du_2 \cdots \oint_{\vartheta_N^{\beta}} du_N f_{\beta}^{(\kappa')}(\mathbf{x}; \mathbf{u}),$$

and

$$\mathcal{F}_{\beta}^{(\kappa')}(\mathbf{x}) = C(\kappa')^N \mathcal{H}_{\beta}^{(\kappa')}(\mathbf{x}), \quad \text{where } C(\kappa') = \frac{\nu(\kappa')}{4 \sin^2(4\pi/\kappa')} \frac{\Gamma(2 - 8/\kappa')}{\Gamma(1 - 4/\kappa')^2}.$$

Note that

$$\lim_{\kappa' \rightarrow 8^-} \frac{(8 - \kappa')}{\nu(\kappa')} = \frac{8}{\pi} \quad \text{and} \quad \lim_{\kappa' \rightarrow 8^-} \frac{C(\kappa')}{\nu(\kappa')} = \frac{\Gamma(1)}{4\Gamma(1/2)^2} = \frac{1}{4\pi},$$

and that  $\mathcal{H}_{\beta}^{(\kappa')}$  is continuous in  $\kappa'$  and the limit  $\mathcal{H}_{\beta}^{(8)} = \lim_{\kappa' \rightarrow 8^-} \mathcal{H}_{\beta}^{(\kappa')}$  is positive and finite. Hence, we have

$$\widehat{\mathcal{F}}_{\beta}^{(8)}(\mathbf{x}) := \left(\frac{\pi}{8}\right)^{N-1} \lim_{\kappa' \rightarrow 8^-} \frac{\mathcal{F}_{\beta}^{(\kappa')}(\mathbf{x})}{\nu(\kappa')^N} = \frac{8}{32^N \pi} \mathcal{H}_{\beta}^{(8)}(\mathbf{x}) \in (0, \infty).$$

This proves Proposition 1.8.  $\square$

## 5.5 Finishing the proof of Theorem 1.10

*Proof of Theorem 1.10.* The existence and uniqueness of the collection  $\{\mathcal{Z}_\alpha^{(\kappa)} : \alpha \in \text{LP}\}$  follows from [PW19, Theorem 1.1] (for  $\kappa \in (0, 4]$ ) and from the results of Section 4 (for  $\kappa \in (4, 8)$ ):

- the uniqueness follows by results of Flores and Kleban (see Lemma 2.11); and
- the PDE system (PDE) is given by Proposition 4.12;
- the Möbius covariance (COV) is given by Lemma 4.5;
- the recursive asymptotics property (ASY) is given by Proposition 4.16;
- the power-law bound (PLB) is given by Lemma 4.13;
- and the positivity is given by Lemma 4.10.

The linear independence is the content of Lemma 5.1. The continuity (Item 1 of Theorem 1.10) is the content of Lemmas 5.2 & 5.4. The limit as  $\kappa \rightarrow 8-$  (Item 2 of Theorem 1.10) can be shown using the relation (1.24) (proven in Lemma 5.3): for all  $\kappa' \in (0, 8)$ , we have

$$\mathcal{F}_\beta^{(\kappa')}(\mathbf{x}) = \sum_{\alpha \in \text{LP}_N} \mathcal{M}_{\nu(\kappa')}(\alpha, \beta) \mathcal{Z}_\alpha^{(\kappa')}(\mathbf{x}), \quad \mathbf{x} \in \mathfrak{X}_{2N}, \beta \in \text{LP}_N. \quad (5.12)$$

As the renormalized meander matrix is invertible by Lemma 5.6, and there exists  $\kappa_0 = \kappa_0(N) \in (4, 8)$  such that the meander matrix  $\{\mathcal{M}_{\nu(\kappa')}(\alpha, \beta) : \alpha, \beta \in \text{LP}_N\}$  is invertible for all  $\kappa' \in (\kappa_0, 8)$  by Lemma 5.5, Cramer's rule shows that

$$\lim_{\kappa' \rightarrow 8-} \left( \frac{1}{\nu(\kappa')} \mathcal{M}_{\nu(\kappa')} \right)^{-1}(\alpha, \beta) = \widehat{\mathcal{M}}^{-1}(\alpha, \beta), \quad \text{for all } \alpha, \beta \in \text{LP}_N. \quad (5.13)$$

Plugging this into (5.12), we obtain (1.23):

$$\begin{aligned} \widehat{\mathcal{Z}}_\alpha^{(8)}(\mathbf{x}) &:= \lim_{\kappa' \rightarrow 8-} \frac{\mathcal{Z}_\alpha^{(\kappa')}(\mathbf{x})}{(8 - \kappa')^{N-1}} = \left( \frac{\pi}{8} \right)^{N-1} \lim_{\kappa' \rightarrow 8-} \frac{\mathcal{Z}_\alpha^{(\kappa')}(\mathbf{x})}{\nu(\kappa')^{N-1}} \\ &= \left( \frac{\pi}{8} \right)^{N-1} \lim_{\kappa' \rightarrow 8-} \sum_{\beta \in \text{LP}_N} \left( \frac{1}{\nu(\kappa')} \mathcal{M}_{\nu(\kappa')} \right)^{-1}(\alpha, \beta) \frac{\mathcal{F}_\beta^{(\kappa')}(\mathbf{x})}{\nu(\kappa')^N} \quad [\text{by (5.12)}] \\ &= \sum_{\beta \in \text{LP}_N} \widehat{\mathcal{M}}^{-1}(\alpha, \beta) \widehat{\mathcal{F}}_\beta^{(8)} \in (0, \infty), \quad [\text{by (1.20) and (5.13)}] \end{aligned}$$

where the positivity follows from [LPW24, Theorem 1.2].  $\square$

Moreover, from the above proof, we obtain

$$\widehat{\mathcal{Z}}_\alpha^{(8)}(\mathbf{x}) = \sum_{\beta \in \text{LP}_N} \widehat{\mathcal{M}}^{-1}(\alpha, \beta) \widehat{\mathcal{F}}_\beta^{(8)}. \quad (5.14)$$

*Proof of Proposition 1.15.* The first claim (1.30) follows from (5.11), and the linear independence follows from the fact that the renormalized meander matrix is invertible (Lemma 5.6) and the pure partition functions are linearly independent (Lemma 5.1). The second claim (1.31) follows from (5.14), and the linear independence is proved in [LPW24, Section 4.3].  $\square$

## 6 Fröbenius series

In this section, we refine the asymptotic properties considered in Section 2.5 to derive the Fröbenius series of the Coulomb gas integral functions. We thus finish the proofs of Propositions 1.16, 1.17 & 1.18.

It was proven in [Dub15a] and [FK15d] that solutions to the second order BPZ PDEs (PDE) have a Fröbenius series expansion. In this section, our aim is to make these expansions explicit for the Coulomb gas integral functions  $\mathcal{F}_\beta$ . Using Proposition 1.11, one can then derive the explicit Fröbenius series expansions also for the pure partition functions  $\mathcal{Z}_\alpha$ . From the algebraic points of view, it is known that the pure partition functions form a distinguished basis in a tensor product module of the quantum group (Hopf algebra)  $U_q(\mathfrak{sl}(2, \mathbb{C}))$  with  $q = \exp(4\pi i/\kappa)$  (cf. Remark 2.7 and [KP16]) — dual to the basis carrying the natural action of the Temperley-Lieb algebra, see, e.g., [FP18, FP20, GSNJ<sup>+</sup>22] (which for the functions is realized as iterated limits, or “fusions”, cf. Section 5.1). In light of the anticipated algebraic structure, the factors in the Fröbenius series in the generic cases addressed in Proposition 1.16 completely match our expectations (the cases where  $\kappa = 8/m$  for some integer  $m$  correspond to more complicated, non-semisimple representation theory, with logarithmic corrections to the Fröbenius series in Propositions 1.17 & 1.18).

### 6.1 Generic case: proof of Proposition 1.16

In this section, we use the notation from Proposition 1.16. The proof of Proposition 1.16 comprises the two Lemmas 6.1 & 6.2, together with the analysis of the exceptional case  $\kappa = 4$  in Appendix E:

*Proof of Proposition 1.16.* When  $\{1, 2\} \in \beta$ , the asserted expansion (1.34) is proven in Lemma 6.1. When  $\{1, 2\} \notin \beta$ , the asserted expansion (1.35) is proven in Lemma 6.2 for the generic case  $\kappa \in (8/3, 8) \setminus \{4\}$ , and in Proposition E.1 in Appendix E for the exceptional case  $\kappa = 4$ .  $\square$

**Lemma 6.1.** *Fix  $\kappa \in (0, 8) \setminus \{\frac{8}{2m+1} : m \in \mathbb{Z}_{>0}\}$ . Fix  $\beta \in \text{LP}_N$  and suppose  $\{1, 2\} \in \beta$ . Then, for all  $\xi \in (-\infty, x_3)$ , we have*

$$\mathcal{F}_\beta(\mathbf{x}) = \nu(\kappa) \mathcal{F}_{\beta/\{1,2\}}(\ddot{\mathbf{x}}_1) (x_2 - x_1)^{-2h(\kappa)} + O((x_2 - x_1)^{-2h(\kappa)+2}), \quad \text{as } x_1, x_2 \rightarrow \xi. \quad (6.1)$$

As a consequence of Lemma 6.1, we see that Eq. (1.34) in Proposition 1.16 holds for all  $\kappa \in (8/3, 8)$ . Indeed, when  $\kappa > 8/3$ , we have  $-2h(\kappa) + 2 = (3\kappa - 6)/\kappa > 2/\kappa$ , so (1.34) follows from (6.1).

*Proof.* Using the change of variables  $u = \frac{u_1 - x_1}{x_2 - x_1}$ , we obtain

$$\begin{aligned} \mathcal{H}_\beta(\mathbf{x}) &= (x_2 - x_1)^{-2h(\kappa)} \oint_{\vartheta_2^\beta} du_2 \cdots \oint_{\vartheta_N^\beta} du_N f_{\beta/\{1,2\}}(\ddot{\mathbf{x}}_1; \dot{\mathbf{u}}_1) \\ &\quad \times \oint_{\vartheta(0,1)} du (u - 1)^{-4/\kappa} u^{-4/\kappa} g(\ddot{\mathbf{x}}_1; \dot{\mathbf{u}}_1; x_1, x_2, u), \end{aligned} \quad (6.2)$$

where

$$\begin{aligned} g(x_1, x_2, u; \ddot{\mathbf{x}}_1; \dot{\mathbf{u}}_1) &:= \prod_{s=2}^N \left( (u_s - (x_2 - x_1)u - x_1)^{8/\kappa} (u_s - x_1)^{-4/\kappa} (u_s - x_2)^{-4/\kappa} \right) \\ &\quad \times \prod_{i=3}^{2N} (x_i - (x_2 - x_1)u - x_1)^{-4/\kappa} \prod_{i=3}^{2N} ((x_i - x_1)^{2/\kappa} (x_i - x_2)^{2/\kappa}), \end{aligned}$$

and its branch is chosen to be real and positive when  $0 < \text{Re}(u) < 1$  and  $x_{a_s} < \text{Re}(u_s) < x_{a_s+1}$  for all  $s \in \{3, 4, \dots, N\}$ . We expand the function  $g$  in  $(x_2 - x_1)$  exactly as

$$\begin{aligned} g(x_1, x_2, u; \ddot{\mathbf{x}}_1; \dot{\mathbf{u}}_1) &= 1 + (x_2 - x_1)(2u - 1) \left( \sum_{s=2}^N \frac{4/\kappa}{x_1 - u_s} - \sum_{i=3}^{2N} \frac{2/\kappa}{x_i - x_1} \right) \\ &\quad + (x_2 - x_1)^2 \hat{g}^{(\kappa)}(x_1, x_2, u; \ddot{\mathbf{x}}_1, \dot{\mathbf{u}}_1), \end{aligned} \quad (6.3)$$

where the function  $\hat{g}^{(\kappa)}(x_1, x_2, u; \mathbf{\ddot{x}}_1, \mathbf{\dot{u}}_1)$  is uniformly bounded in  $x_1, x_2 \in (\xi - (x_3 - \xi)/3, \xi + (x_3 - \xi)/3)$  and  $\kappa \in (\frac{8}{2m+1}, \frac{8}{2m-1})$  for each fixed  $m \in \mathbb{Z}_{>0}$ , and  $\mathbf{\ddot{x}}_1 \in \mathfrak{X}_{2N-2}$ , and  $\xi \in (-\infty, x_3)$ . Therefore, for each fixed  $m \in \mathbb{Z}_{>0}$ , plugging Eq. (6.3) and Eq. (B.1, B.2) from Lemma B.1 into (6.2) shows that

$$\begin{aligned} & \mathcal{F}_\beta(\mathbf{x}) - \nu(\kappa) \mathcal{F}_{\beta/\{1,2\}}(\mathbf{\ddot{x}}_1) (x_2 - x_1)^{-2h(\kappa)} \\ &= (x_2 - x_1)^{-2h(\kappa)+2} C(\kappa)^N \oint_{\vartheta_2^\beta} du_2 \cdots \oint_{\vartheta_N^\beta} du_N f_{\beta/\{1,2\}}(\mathbf{\ddot{x}}_1; \mathbf{\dot{u}}_1) \\ & \quad \times \oint_{\vartheta(0,1)} du (u-1)^{-4/\kappa} u^{-4/\kappa} \hat{g}^{(\kappa)}(x_1, x_2, u; \mathbf{\ddot{x}}_1; \mathbf{\dot{u}}_1) \\ &= O((x_2 - x_1)^{-2h(\kappa)+2}), \quad \text{as } x_1, x_2 \rightarrow \xi, \end{aligned} \tag{6.4}$$

for any  $\kappa \in (\frac{8}{2m+1}, \frac{8}{2m-1}) \setminus \{\frac{4}{m}\}$ . To extend (6.1) to all parameters  $\kappa \in (0, 8) \setminus \{\frac{8}{2m+1} : m \in \mathbb{Z}_{>0}\}$ , let us fix  $\mathbf{\ddot{x}}_1 \in \mathfrak{X}_{2N-2}$ , and  $\xi \in (-\infty, x_3)$ , and  $m \in \mathbb{Z}_{>0}$ , and consider the limit  $\kappa' \rightarrow \kappa_m := 4/m$ . Similarly as in the proof of Lemma 2.8, in the integral (6.4) over  $f_{\beta/\{1,2\}}^{(\kappa')}(\mathbf{\ddot{x}}_1; \mathbf{\dot{u}}_1) \hat{g}^{(\kappa')}(x_1, x_2, u; \mathbf{\ddot{x}}_1; \mathbf{\dot{u}}_1)$ , we can replace each Pochhammer contour  $\vartheta(x_{a_r}, x_{b_r})$  by the linear combination of the three contours  $(1 - e^{8\pi i/\kappa})(1 - e^{-8\pi i/\kappa}) \int_{x_a+\epsilon}^{x_b-\epsilon} du_r$ , and  $(1 - e^{8\pi i/\kappa}) \oint_{\partial B(x_a, \epsilon)} du_r$ , and  $(1 - e^{-8\pi i/\kappa}) \oint_{\partial B(x_b, \epsilon)} du_r$ ; and similarly, we can replace the Pochhammer contour  $\vartheta(0, 1)$  by the linear combination of the three contours  $(1 - e^{8\pi i/\kappa})(1 - e^{-8\pi i/\kappa}) \int_\epsilon^{1-\epsilon} du$ , and  $(1 - e^{8\pi i/\kappa}) \oint_{\partial B(0, \epsilon)} du$ , and  $(1 - e^{-8\pi i/\kappa}) \oint_{\partial B(1, \epsilon)} du$ . Therefore, since

$$\text{both } C(\kappa)(1 - e^{8\pi i/\kappa}) \text{ and } C(\kappa)(1 - e^{-8\pi i/\kappa}) \text{ are bounded in } \kappa \in (\frac{8}{2m+1}, \frac{8}{2m-1}) \setminus \{\frac{4}{m}\},$$

we conclude that

$$C(\kappa)^N \oint_{\vartheta_2^\beta} du_2 \cdots \oint_{\vartheta_N^\beta} du_N f_{\beta/\{1,2\}}(\mathbf{\ddot{x}}_1; \mathbf{\dot{u}}_1) \oint_{\vartheta(0,1)} du (u-1)^{-4/\kappa} u^{-4/\kappa} \hat{g}^{(\kappa)}(x_1, x_2, u; \mathbf{\ddot{x}}_1; \mathbf{\dot{u}}_1)$$

is uniformly bounded in  $\kappa \in (\frac{8}{2m+1}, \frac{8}{2m-1}) \setminus \{\frac{4}{m}\}$  and  $x_1, x_2 \in (\xi - (x_3 - \xi)/3, \xi + (x_3 - \xi)/3)$ . Finally, taking the limit  $\kappa \rightarrow \frac{4}{m}$  in (6.4) and using the continuity from Theorem 1.5, we get (6.1) for  $\kappa = \frac{4}{m}$ .  $\square$

**Lemma 6.2.** *Fix  $\kappa \in (8/3, 8) \setminus \{4\}$  and  $N \geq 2$ . Fix  $\beta \in \text{LP}_N$  and suppose  $\{1, 2\} \notin \beta$ . Then, for all  $\xi \in (-\infty, x_3)$ , we have*

$$\begin{aligned} \mathcal{F}_\beta(\mathbf{x}) &= \mathcal{F}_{\varphi_1(\beta)/\{1,2\}}(\mathbf{\ddot{x}}_1) (x_2 - x_1)^{-2h(\kappa)} + \mathcal{F}_{\beta/\vee_1}(\xi, \mathbf{\ddot{x}}_1) (x_2 - x_1)^{2/\kappa} \\ & \quad + o((x_2 - x_1)^{2/\kappa}), \quad \text{as } x_1, x_2 \rightarrow \xi, \end{aligned} \tag{6.5}$$

where (using the branch choice (1.37))

$$\begin{aligned} \mathcal{F}_{\beta/\vee_1}(\xi, \mathbf{\ddot{x}}_1) &= \frac{C(\kappa)^{N-2} (\nu(\kappa)^2 - 1)}{2F_1(\frac{4}{\kappa}, 1 - \frac{4}{\kappa}, \frac{8}{\kappa}; 1)} \oint_{\vartheta(x_{a_3}, x_{b_3})} du_3 \cdots \oint_{\vartheta(x_{a_N}, x_{b_N})} du_N \hat{f}_{\beta/\vee_1}(\xi, \mathbf{\ddot{x}}_1; \mathbf{\dot{u}}_1), \\ \hat{f}_{\beta/\vee_1}(\xi, \mathbf{\ddot{x}}_1; \mathbf{\dot{u}}_1) &= \prod_{r=3}^N (u_r - \xi)^{16/\kappa-2} \prod_{j=3}^{2N} (x_j - \xi)^{1-8/\kappa} f(\mathbf{\ddot{x}}_1; \mathbf{\dot{u}}_1). \end{aligned}$$

Note that Eq. (6.5) is exactly the second claim (1.35) in Proposition 1.16 for  $\kappa \in (8/3, 8) \setminus \{4\}$ .

*Proof.* Using Proposition 2.5, we obtain<sup>22</sup>

$$\mathcal{H}_\beta(\mathbf{x}) = \frac{\nu(\kappa)}{C(\kappa)} \oint_{\vartheta_2^\beta} du_2 \cdots \oint_{\vartheta_N^\beta} du_N \hat{f}_\beta^a(\mathbf{x}; \mathbf{\dot{u}}_1)$$

<sup>22</sup>The way we remove the integral surrounding  $x_1$  and  $x_{b_1}$  here is different from that in the proof of Proposition 2.12: in the latter, we use the integrand  $\hat{f}_\beta^b$  with conjugate charge at  $x_{b_1}$ , while here, we use  $\hat{f}_\beta^a$  with conjugate charge at  $x_1$ .

$$\begin{aligned}
&= \frac{\nu(\kappa)}{C(\kappa)} (x_2 - x_1)^{-2h(\kappa)} (x_{b_1} - x_1)^{-2h(\kappa)} (x_{b_1} - x_2)^{2/\kappa} (x_{b_2} - x_1)^{-2h(\kappa)} (x_{b_2} - x_2)^{2/\kappa} \\
&\quad \times \prod_{\substack{3 \leq i \leq 2N \\ i \notin \{b_1, b_2\}}} ((x_i - x_1)^{-2h(\kappa)} (x_i - x_2)^{2/\kappa}) \\
&\quad \times \int_{\Gamma_{\hat{\beta}}} d\mathbf{\ddot{u}}_1 f(\mathbf{\ddot{x}}_1; \mathbf{\ddot{u}}_1) \oint_{\vartheta(x_2, x_{b_2})} du_2 (u_2 - x_{b_1})^{-4/\kappa} (u_2 - x_{b_2})^{-4/\kappa} (u_2 - x_1)^{4h(\kappa)} (u_2 - x_2)^{-4/\kappa} \\
&\quad \times \prod_{\substack{3 \leq i \leq 2N \\ i \notin \{b_1, b_2\}}} (u_2 - x_i)^{-4/\kappa} \prod_{3 \leq s \leq N} (u_s - u_2)^{8/\kappa} \prod_{3 \leq s \leq N} ((u_s - x_1)^{4h(\kappa)} (u_s - x_2)^{-4/\kappa}),
\end{aligned}$$

where the integration  $\Gamma_{\hat{\beta}}$  comprises all of the other contours than the two Pochhammer contours surrounding the pairs  $\{x_1, x_{b_1}\}$  and  $\{x_2, x_{b_2}\}$ , and the integrand  $\hat{f}_{\hat{\beta}}^a$  is defined in (2.11) with  $r = 1$  and the conjugate charge at  $x_1$ , and  $f$  is the integrand function in (1.33), with the branch of  $f(\mathbf{\ddot{x}}_1; \cdot)$  and the rest of the products in the integrand chosen to be real and positive when  $x_{a_s} < \text{Re}(u_s) < x_{a_s} + 1$  for all  $s \in \{3, 4, \dots, N\}$ . Using the change of variables  $w = \text{CR}(x_{b_2}, x_{b_1}, x_2, u_2)$ , where  $\text{CR}$  is the cross-ratio (2.25), we obtain

$$\begin{aligned}
\mathcal{H}_{\beta}(\mathbf{x}) &= \frac{\nu(\kappa)}{C(\kappa)} (x_{b_1} - x_{b_2})^{1-8/\kappa} \left( \frac{(x_2 - x_1)(x_{b_1} - x_2)(x_{b_2} - x_1)}{(x_{b_1} - x_1)(x_{b_2} - x_2)} \right)^{-2h(\kappa)} \\
&\quad \times \int_{\Gamma_{\hat{\beta}}} d\mathbf{\ddot{u}}_1 f(\mathbf{\ddot{x}}_1; \mathbf{\ddot{u}}_1) \oint_{\vartheta(0,1)} dw \tilde{g}(x_1, x_2, w; \mathbf{\ddot{x}}_1; \mathbf{\ddot{u}}_1) \hat{k}(\chi, w),
\end{aligned} \tag{6.6}$$

writing  $\chi = \text{CR}(x_1, x_2, x_{b_2}, x_{b_1})$ , and  $\hat{k}(\chi, w) := (w + \frac{\chi}{1-\chi})^{4h(\kappa)} (1-w)^{-4/\kappa} w^{-4/\kappa}$ , and

$$\begin{aligned}
\tilde{g}(x_1, x_2, w; \mathbf{\ddot{x}}_1; \mathbf{\ddot{u}}_1) &:= \prod_{\substack{3 \leq i \leq 2N \\ i \notin \{b_1, b_2\}}} \left| x_{b_1} - (x_{b_1} - x_2) \left( 1 + w \frac{x_{b_2} - x_2}{x_{b_1} - x_{b_2}} \right)^{-1} - x_i \right|^{-4/\kappa} \\
&\quad \times \prod_{3 \leq s \leq N} \left( u_s - x_{b_1} + (x_{b_1} - x_2) \left( 1 + w \frac{x_{b_2} - x_2}{x_{b_1} - x_{b_2}} \right)^{-1} \right)^{8/\kappa} \\
&\quad \times \prod_{3 \leq s \leq N} ((u_s - x_1)^{4h(\kappa)} (u_s - x_2)^{-4/\kappa}) \prod_{\substack{3 \leq i \leq 2N \\ i \notin \{b_1, b_2\}}} ((x_i - x_1)^{-2h(\kappa)} (x_i - x_2)^{2/\kappa}),
\end{aligned} \tag{6.7}$$

whose branch is chosen to be real and positive when  $0 < \text{Re}(w) < 1$  and  $x_{a_s} < \text{Re}(u_s) < x_{a_s+1}$  for all  $s \in \{3, 4, \dots, N\}$ . Note that  $\tilde{g}$  stays bounded in the limit  $x_1, x_2 \rightarrow \xi$ , and  $\chi \rightarrow 0$  in this limit.

Let us prove (6.5) for  $\kappa \in (4, 8)$ . By (6.6), it suffices to consider the limit of the integral over  $\vartheta(0, 1)$  as  $x_1, x_2 \rightarrow \xi$ . Similarly as in Lemma 2.8, we see that

$$\begin{aligned}
&\oint_{\vartheta(0,1)} dw \tilde{g}(x_1, x_2, w; \mathbf{\ddot{x}}_1; \mathbf{\ddot{u}}_1) \hat{k}(\chi, w) \\
&= (1 - e^{8\pi i/\kappa})(1 - e^{-8\pi i/\kappa}) \underbrace{\int_{\epsilon \frac{\chi}{1-\chi}}^{1-\epsilon} dw \tilde{g}(x_1, x_2, w; \mathbf{\ddot{x}}_1; \mathbf{\ddot{u}}_1) \hat{k}(\chi, w)}_{:= I_1} \\
&\quad + (1 - e^{8\pi i/\kappa}) \underbrace{\oint_{\partial B(0, \epsilon \frac{\chi}{1-\chi})} dw \tilde{g}(x_1, x_2, w; \mathbf{\ddot{x}}_1; \mathbf{\ddot{u}}_1) \hat{k}(\chi, w)}_{:= I_2} \\
&\quad + (1 - e^{-8\pi i/\kappa}) \underbrace{\oint_{\partial B(1, \epsilon)} dw \tilde{g}(x_1, x_2, w; \mathbf{\ddot{x}}_1; \mathbf{\ddot{u}}_1) \hat{k}(\chi, w)}_{:= I_3},
\end{aligned}$$



when  $\epsilon > 0$  is small and fixed. We derive the limit as  $x_1, x_2 \rightarrow \xi$  by analyzing the terms  $I_1$ ,  $I_2$ , and  $I_3$  one by one. The second term vanishes in the limit: since  $|\hat{k}(\chi, w)| \sim \chi^{8/\kappa-2}$  for all  $w \in \partial B(0, \epsilon \frac{\chi}{1-\chi})$ , we have

$$I_2 = \oint_{\partial B(0, \epsilon \frac{\chi}{1-\chi})} dw \tilde{g}(x_1, x_2, w; \ddot{\mathbf{x}}_1; \ddot{\mathbf{u}}_1) \hat{k}(\chi, w) \xrightarrow{x_1, x_2 \rightarrow \xi} 0.$$

The third term has the limit

$$I_3 = \oint_{\partial B(1, \epsilon)} dw \tilde{g}(x_1, x_2, w; \ddot{\mathbf{x}}_1; \ddot{\mathbf{u}}_1) \hat{k}(\chi, w) \xrightarrow{x_1, x_2 \rightarrow \xi} \oint_{\partial B(1, \epsilon)} dw \tilde{g}(\xi, \xi, w; \ddot{\mathbf{x}}_1; \ddot{\mathbf{u}}_1) w^{8/\kappa-2} (1-w)^{-4/\kappa}.$$

The first term has the limit

$$I_1 = \int_{\epsilon \frac{\chi}{1-\chi}}^{1-\epsilon} dw \tilde{g}(x_1, x_2, w; \ddot{\mathbf{x}}_1; \ddot{\mathbf{u}}_1) \hat{k}(\chi, w) \xrightarrow{x_1, x_2 \rightarrow \xi} \int_0^{1-\epsilon} dw \tilde{g}(\xi, \xi, w; \ddot{\mathbf{x}}_1; \ddot{\mathbf{u}}_1) w^{8/\kappa-2} (1-w)^{-4/\kappa}.$$

Collecting these facts, we conclude that

$$\begin{aligned} \frac{1}{\nu(\kappa)} \mathcal{H}_{\varphi_1(\beta)/\{1,2\}}(\ddot{\mathbf{x}}_1) &= \frac{C(\kappa)}{\nu(\kappa)} \lim_{x_1, x_2 \rightarrow \xi} \frac{\mathcal{H}_\beta(\mathbf{x})}{(x_2 - x_1)^{-2h(\kappa)}} \\ &= (1 - e^{-8\pi i/\kappa}) (x_{b_1} - x_{b_2})^{1-8/\kappa} \int_{\Gamma_{\hat{\beta}}} d\ddot{\mathbf{u}}_1 f(\ddot{\mathbf{x}}_1; \ddot{\mathbf{u}}_1) \\ &\quad \times \left( (1 - e^{8\pi i/\kappa}) \int_0^{1-\epsilon} dw + \oint_{\partial B(1, \epsilon)} dw \right) \tilde{g}(\xi, \xi, w; \ddot{\mathbf{x}}_1; \ddot{\mathbf{u}}_1) w^{8/\kappa-2} (1-w)^{-4/\kappa}, \end{aligned}$$

using also Proposition 2.12 to identify the left-hand side. Note that  $-2(\kappa) + 1 = 2 - 6/\kappa > 2/\kappa$  when  $\kappa \in (4, 8)$ . Consequently,

$$\begin{aligned} &\frac{\mathcal{H}_\beta(\mathbf{x}) - (x_2 - x_1)^{-2h(\kappa)} \frac{1}{C(\kappa)} \mathcal{H}_{\varphi_1(\beta)/\{1,2\}}(\ddot{\mathbf{x}}_1)}{\frac{\nu(\kappa)}{C(\kappa)} (x_2 - x_1)^{-2h(\kappa)}} \\ &= \left( \frac{(x_{b_1} - \xi)(x_{b_2} - \xi)}{x_{b_2} - x_{b_1}} \right)^{8/\kappa-1} \int_{\Gamma_{\hat{\beta}}} d\ddot{\mathbf{u}}_1 \hat{f}_{\beta/\nu_1}(\xi, \ddot{\mathbf{x}}_1; \ddot{\mathbf{u}}_1) \underbrace{\left( (1 - e^{8\pi i/\kappa}) \oint_{\partial B(0, \epsilon \frac{\chi}{1-\chi})} dw \left( w + \frac{\chi}{1-\chi} \right)^{4h(\kappa)} w^{-4/\kappa} \right)}_{I_4} \\ &\quad + \underbrace{(1 - e^{-8\pi i/\kappa}) \int_{\epsilon \frac{\chi}{1-\chi}}^{10 \frac{\chi}{1-\chi}} dw \left( w + \frac{\chi}{1-\chi} \right)^{4h(\kappa)} w^{-4/\kappa} - (1 - e^{-8\pi i}) \int_0^{10 \frac{\chi}{1-\chi}} dw w^{8/\kappa-2}}_{I_5} \\ &\quad + \underbrace{4 \sin^2(4\pi/\kappa) \int_{10 \frac{\chi}{1-\chi}}^{1-\epsilon} dw \left( \left( w + \frac{\chi}{1-\chi} \right)^{4h(\kappa)} w^{-4/\kappa} - w^{8/\kappa-2} \right)}_{I_6} \\ &\quad + O(x_2 - x_1), \quad \text{as } x_1, x_2 \rightarrow \xi, \end{aligned} \tag{6.8}$$

where  $O(x_2 - x_1) = o((x_2 - x_1)^{8/\kappa-1})$  since  $1 > 8\kappa - 1$  when  $\kappa \in (4, 8)$ . For the terms  $I_4$  and  $I_5$ , using the change of variables  $u = w \frac{1-\chi}{\chi}$ , we see that

$$I_4 + I_5 = \chi^{8/\kappa-1} C_1(\kappa),$$

where

$$C_1(\kappa) := (1 - e^{8\pi i/\kappa}) \oint_{\partial B(0, \epsilon)} du (u+1)^{4h(\kappa)} u^{-4/\kappa} + (1 - e^{-8\pi i}) \left( \int_\epsilon^{10} (u+1)^{4h(\kappa)} u^{-4/\kappa} - \int_0^{10} du u^{8/\kappa-2} \right)$$

is a finite constant depending only on  $\kappa$  (and not on  $N$ ). For the term  $I_6$ , we Taylor expand

$$\left(w + \frac{\chi}{1-\chi}\right)^{12/\kappa-2} - w^{12/\kappa-2} = w^{12/\kappa-2} \sum_{n=1}^{\infty} \frac{(\frac{12}{\kappa}-2)(\frac{12}{\kappa}-3)\cdots(\frac{12}{\kappa}-n-1)}{n!} \left(\frac{\chi}{w(1-\chi)}\right)^n$$

for  $w \in (\frac{10\chi}{1-\chi}, \frac{9}{10})$ . Since  $1 > 8/\kappa - 1$  when  $\kappa \in (4, 8)$ , we see that

$$I_6 = \left(\frac{\chi}{1-\chi}\right)^{8/\kappa-1} C_2(\kappa) + o((x_2 - x_1)^{8/\kappa-1}),$$

where

$$C_2(\kappa) := -4 \sin^2(4\pi/\kappa) \sum_{n=1}^{\infty} \frac{(\frac{12}{\kappa}-2)(\frac{12}{\kappa}-3)\cdots(\frac{12}{\kappa}-n-1)}{n! (\frac{8}{\kappa}-n-1)} 10^{8/\kappa-1-n}$$

is a finite constant depending only on  $\kappa$  (and not on  $N$ ).

Collecting the above expansions, on the one hand, plugging them into (6.8) gives

$$\begin{aligned} \mathcal{H}_\beta(\mathbf{x}) &= (x_2 - x_1)^{-2h(\kappa)} \frac{1}{C(\kappa)} \mathcal{H}_{\varphi_1(\beta)/\{1,2\}}(\ddot{\mathbf{x}}_1) + o((x_2 - x_1)^{2/\kappa}) \\ &\quad + (x_2 - x_1)^{2/\kappa} (C_1(\kappa) + C_2(\kappa)) \frac{\nu(\kappa)}{C(\kappa)} \int_{\Gamma_{\hat{\beta}}} d\ddot{\mathbf{u}}_1 \hat{f}_{\beta/\vee_1}(\xi, \ddot{\mathbf{x}}_1; \ddot{\mathbf{u}}_1), \end{aligned} \quad (6.9)$$

On the other hand, when  $N = 2$ , using the relation (D.25) from Corollary D.7, we have, as  $x_1, x_2 \rightarrow \xi$ ,

$$\begin{aligned} \mathcal{H}_{\underbrace{\quad}}(x_1, x_2, x_3, x_4) &= (x_2 - x_1)^{-2h(\kappa)} \frac{1}{C(\kappa)} \mathcal{H}_{\underbrace{\quad}}(x_3, x_4) + o((x_2 - x_1)^{2/\kappa}) \\ &\quad + (x_2 - x_1)^{2/\kappa} \frac{1}{C(\kappa)^2} \frac{\nu(\kappa)^2 - 1}{{}_2F_1(\frac{4}{\kappa}, 1 - \frac{4}{\kappa}, \frac{8}{\kappa}; 1)} \left( \frac{(x_4 - x_3)^{2/\kappa}}{(x_4 - \xi)^{8/\kappa-1} (x_3 - \xi)^{8/\kappa-1}} \right). \end{aligned} \quad (6.10)$$

Comparing (6.9) with (6.10) then shows that

$$(C_1(\kappa) + C_2(\kappa)) \frac{\nu(\kappa)}{C(\kappa)} = \frac{1}{C(\kappa)^2} \frac{\nu(\kappa)^2 - 1}{{}_2F_1(\frac{4}{\kappa}, 1 - \frac{4}{\kappa}, \frac{8}{\kappa}; 1)},$$

and finally, plugging this into (6.9) and using the relation (2.1) gives (6.5) for  $\kappa \in (4, 8)$ .

Let us then discuss the case  $\kappa \in (8/3, 4)$ , where the analysis is essentially the same, with one important difference: in this case, the term of order  $(x_2 - x_1)^{-2h(\kappa)+1}$  in the expansion of  $\mathcal{H}_\beta(\mathbf{x})$  vanishes. Indeed, it follows from (2.26) and the analysis after it in the proof of Proposition 2.12 that

$$\begin{aligned} \mathcal{H}_\beta^{(\kappa)}(\mathbf{x}) &= \frac{1}{C(\kappa)} \mathcal{H}_{\varphi_1(\beta)/\{1,2\}}^{(\kappa)}(\ddot{\mathbf{x}}_1) (x_2 - x_1)^{-2h(\kappa)} + \nu(\kappa) o((x_2 - x_1)^{-2h(\kappa)+1}) \\ &\quad + (x_2 - x_1)^{-2h(\kappa)+1} \frac{\nu(\kappa)}{C(\kappa)} (1 - e^{8\pi i/\kappa}) \int_{\Gamma_{\hat{\beta}}} \tilde{f}(\ddot{\mathbf{x}}_1; \ddot{\mathbf{u}}_1) d\ddot{\mathbf{u}}_1 \\ &\quad \times \oint_{\varrho(0,\infty)} dv v^{-4/\kappa} (1+v)^{-4/\kappa} \left( \frac{6}{\kappa} - 1 + \frac{4}{\kappa} \left( \frac{v}{1+v} + v \right) + (2v+1) \left( \sum_{2 \leq s \leq N} \frac{4/\kappa}{\xi - u_s} + \sum_{\substack{3 \leq i \leq 2N \\ i \notin \{b_1, b_2\}}} \frac{2/\kappa}{\xi - x_i} \right) \right) \\ &= \frac{1}{C(\kappa)} \mathcal{H}_{\varphi_1(\beta)/\{1,2\}}^{(\kappa)}(\ddot{\mathbf{x}}_1) (x_2 - x_1)^{-2h(\kappa)} + \nu(\kappa) o((x_2 - x_1)^{-2h(\kappa)+1}), \quad \text{as } x_1, x_2 \rightarrow \xi, \end{aligned} \quad (6.11)$$

thanks to the identities (B.5) and (B.6) from Lemma B.2.  $\square$

## 6.2 $\kappa = 8/3$ : proof of Proposition 1.17

In this section, we fix  $\kappa = 8/3$  and use the notation from Proposition 1.17.

*Proof of Proposition 1.17.* When  $\{1, 2\} \in \beta$ , the asserted expansion (1.38) is proven in Lemma 6.3. When  $\{1, 2\} \notin \beta$ , the asserted expansion (1.40) is proven in Lemma 6.5.  $\square$

**Lemma 6.3.** *Fix  $\beta \in \text{LP}_N$  and suppose  $\{1, 2\} \in \beta$ . Then, for all  $\xi \in (-\infty, x_3)$ , we have*

$$\widehat{\mathcal{F}}_{\beta}^{(8/3)}(\mathbf{x}) = \widehat{\mathcal{F}}_{\beta/\vee_1}^{(8/3)}(\xi, \ddot{\mathbf{x}}_1) (x_2 - x_1)^{3/4} + o((x_2 - x_1)^{3/4}), \quad \text{as } x_1, x_2 \rightarrow \xi, \quad (6.12)$$

where (using the branch choice (1.37))

$$\begin{aligned} \widehat{\mathcal{F}}_{\beta/\vee_1}^{(8/3)}(\xi, \ddot{\mathbf{x}}_1) &= \frac{C(8/3)^{N-2}}{2F_1\left(\frac{3}{2}, -\frac{1}{2}, 3; 1\right)} \oint_{\vartheta(x_{a_3}, x_{b_3})} du_3 \cdots \oint_{\vartheta(x_{a_N}, x_{b_N})} du_N \hat{f}_{\beta/\vee_1}^{(8/3)}(\xi, \ddot{\mathbf{x}}_1; \ddot{\mathbf{u}}_1), \\ \hat{f}_{\beta/\vee_1}^{(8/3)}(\xi, \ddot{\mathbf{x}}_1; \ddot{\mathbf{u}}_1) &= \prod_{r=3}^N (u_r - \xi)^4 \prod_{j=3}^{2N} (x_j - \xi)^{-2} f(\ddot{\mathbf{x}}_1; \ddot{\mathbf{u}}_1). \end{aligned}$$

Note that Eq. (6.12) is exactly the first claim (1.38) in Proposition 1.17.

*Proof.* According to Proposition 2.5, we have

$$\begin{aligned} \widehat{\mathcal{H}}_{\beta}^{(8/3)}(\mathbf{x}) &:= \lim_{\kappa' \rightarrow 8/3} \frac{1}{\nu(\kappa')} \mathcal{H}_{\beta}^{(\kappa')}(\mathbf{x}) \\ &= \frac{1}{C(8/3)} \oint_{\vartheta_2^{\beta}} du_2 \cdots \oint_{\vartheta_N^{\beta}} du_N \hat{f}_{\beta}^a(\mathbf{x}; \mathbf{u}_1) \\ &= \frac{(x_2 - x_1)^{-5/4}}{C(8/3)} \int_{\Gamma_{\beta}} d\ddot{\mathbf{u}}_1 f^{(8/3)}(\ddot{\mathbf{x}}_1; \ddot{\mathbf{u}}_1) \oint_{\vartheta(x_3, x_{b_2})} du_2 g_1(x_1, x_2, u_2; \ddot{\mathbf{x}}_1, \ddot{\mathbf{u}}_1), \end{aligned} \quad (6.13)$$

where the integration  $\Gamma_{\beta}$  comprises all of the other contours than the two Pochhammer contours surrounding the pairs  $\{x_1, x_2\}$  and  $\{x_3, x_{b_2}\}$ , and the integrand  $\hat{f}_{\beta}^a$  is defined in (2.11) with  $r = 1$  and the conjugate charge at  $x_1$ , and  $f$  is the integrand function in (1.33) with  $\kappa = 8/3$ , and

$$\begin{aligned} g_1(x_1, x_2, u_2; \ddot{\mathbf{x}}_1, \ddot{\mathbf{u}}_1) &:= (u_2 - x_1)^{5/2} (u_2 - x_2)^{-3/2} \prod_{3 \leq s \leq N} (u_s - u_2)^3 \prod_{3 \leq i \leq 2N} (u_2 - x_i)^{-3/2} \\ &\quad \times \prod_{3 \leq s \leq N} ((u_s - x_1)^{5/2} (u_s - x_2)^{-3/2}) \prod_{3 \leq i \leq 2N} ((x_i - x_1)^{5/2} (x_i - x_2)^{-3/2}), \end{aligned}$$

with the branch of  $f(\ddot{\mathbf{x}}_1; \cdot)$  and the rest of the products in the integrand chosen to be real and positive when  $x_{a_s} < \text{Re}(u_s) < x_{a_s} + 1$  for all  $s \in \{2, \dots, N\}$ . Note that the function  $u_2 \mapsto g_1(x_1, x_2, u_2; \ddot{\mathbf{x}}_1, \ddot{\mathbf{u}}_1)$  is of order  $|u_2|^{-2}$  as  $|u_2| \rightarrow \infty$ , thus holomorphic at infinity. Using Corollary 2.9 (for  $\kappa = 8/3$ ) to transform all of the Pochhammer contours  $\vartheta(x_{a_r}, x_{b_r})$  in the integrals in (6.13) into simple clockwise loops  $\varrho(x_{a_r}, x_{b_r})$  (see also Figure 2.2), and then using Lemma 2.10 to pull the loop  $\varrho_2 = \varrho(x_3, x_{b_2})$  surrounding the points  $x_3$  and  $x_{b_2}$  (and originally, no other marked points) across the other loops so that it becomes a large loop  $\hat{\varrho}_2$  surrounding all of the points  $\{x_3, x_4, \dots, x_{2N}\}$  (see also Figure 2.3), and then pulling that loop across the point at infinity to become a loop  $\varrho(x_1, x_2)$  only surrounding the points  $x_1$  and  $x_2$ , and lastly, transforming all of the loops back to Pochhammer contours (so that in particular  $\varrho(x_1, x_2)$  becomes  $-\vartheta(x_1, x_2)$ ), we see that

$$\widehat{\mathcal{H}}_{\beta}^{(8/3)}(\mathbf{x}) = \frac{(x_2 - x_1)^{-5/4}}{C(8/3)} \int_{\Gamma_{\beta}} d\ddot{\mathbf{u}}_1 f^{(8/3)}(\ddot{\mathbf{x}}_1; \ddot{\mathbf{u}}_1) \oint_{\vartheta(x_1, x_2)} du_2 g_1(x_1, x_2, u_2; \ddot{\mathbf{x}}_1, \ddot{\mathbf{u}}_1), \quad (6.14)$$

where the branch of the product in the integrand is chosen to be real and positive when  $x_1 < \operatorname{Re} u_2 < x_2$  and  $x_{a_s} < u_s < x_{a_s+1}$  for all  $s \in \{3, 4, \dots, N\}$  (so that the change in the branch choice cancels the minus sign from  $-\vartheta(x_1, x_2)$ ). To finish, thanks to (6.14), we can use the change of variables  $u = \frac{u_2 - x_1}{x_2 - x_1}$  to obtain

$$\widehat{\mathcal{H}}_\beta^{(8/3)}(\mathbf{x}) = \frac{(x_2 - x_1)^{3/4}}{C(8/3)} \int_{\Gamma_\beta} d\ddot{\mathbf{u}}_1 \widehat{f}_{\beta/\vee_1}^{(8/3)}(\xi, \ddot{\mathbf{x}}_1; \ddot{\mathbf{u}}_1) \oint_{\vartheta(0,1)} du (u^{5/2}(u-1)^{-3/2} + o(1)), \quad \text{as } x_1, x_2 \rightarrow \xi.$$

Using (B.3) from Lemma B.1 to evaluate the integral

$$\oint_{\vartheta(0,1)} u^{5/2}(u-1)^{-3/2} du = 4 \frac{\Gamma(7/2)\Gamma(-1/2)}{\Gamma(3)} = -\frac{15}{2}\pi,$$

we conclude that

$$\begin{aligned} \lim_{x_1, x_2 \rightarrow \xi} \frac{\widehat{\mathcal{F}}_\beta^{(8/3)}(\mathbf{x})}{(x_2 - x_1)^{3/4}} &= C(8/3)^N \lim_{x_1, x_2 \rightarrow \xi} \frac{\widehat{\mathcal{H}}_\beta^{(8/3)}(\mathbf{x})}{(x_2 - x_1)^{3/4}} \\ &= C(8/3)^{N-2} \frac{15\pi}{32} \int_{\Gamma_\beta} d\ddot{\mathbf{u}}_1 \widehat{f}_{\beta/\vee_1}^{(8/3)}(\xi, \ddot{\mathbf{x}}_1; \ddot{\mathbf{u}}_1) = \widehat{\mathcal{F}}_{\beta/\vee_1}^{(8/3)}(\xi, \ddot{\mathbf{x}}_1), \end{aligned}$$

which proves the asserted expansion (6.12).  $\square$

**Remark 6.4.** Note that in the proof of Lemma 6.3, one can replace  $\{x_3, x_{b_2}\} = \{x_{a_2}, x_{b_2}\}$  by any  $\{x_{a_s}, x_{b_s}\}$  for  $s \in \{2, 3, \dots, N\}$ . The only difference is that one will get an extra  $(-1)^{(a_s-1)/2}$  phase factor on the right-hand side of (6.14) for a general  $s \in \{2, 3, \dots, N\}$ :

$$\begin{aligned} \widehat{\mathcal{F}}_{\beta/\vee_1}^{(8/3)}(\xi, \ddot{\mathbf{x}}_1) &= (-1)^{(a_s-1)/2} \frac{C(8/3)^{N-2}}{2F_1\left(\frac{3}{2}, -\frac{1}{2}, 3; 1\right)} \oint_{\vartheta(x_{a_2}, x_{b_2})} du_2 \cdots \oint_{\vartheta(x_{a_{s-1}}, x_{b_{s-1}})} du_{s-1} \\ &\quad \times \oint_{\vartheta(x_{a_{s+1}}, x_{b_{s+1}})} du_{s+1} \cdots \oint_{\vartheta(x_{a_N}, x_{b_N})} du_N \prod_{\substack{2 \leq r \leq N \\ r \neq s}} (u_r - \xi)^4 \prod_{j=3}^{2N} (x_j - \xi)^{-2} f(\ddot{\mathbf{x}}_1; \ddot{\mathbf{u}}_{1,s}), \end{aligned}$$

where  $\ddot{\mathbf{u}}_{1,s} := (u_2, \dots, u_{s-1}, u_{s+1}, \dots, u_N)$  and the branch of the integrand is chosen to be real and positive when  $x_{a_r} < \operatorname{Re} u_r < x_{a_r+1}$  for  $r \in \{2, 3, \dots, N\} \setminus \{s\}$ .

**Lemma 6.5.** Fix  $\beta \in \operatorname{LP}_N$  and suppose  $\{1, 2\} \notin \beta$ . Then, for all  $\xi \in (-\infty, x_3)$ , we have

$$\begin{aligned} \widehat{\mathcal{F}}_\beta^{(8/3)}(\mathbf{x}) &= \widehat{\mathcal{F}}_{\wp_1(\beta)/\{1,2\}}^{(8/3)}(\ddot{\mathbf{x}}_1) (x_2 - x_1)^{-5/4} - \frac{1}{\pi} (-1)^{(b_2-1)/2} \widehat{\mathcal{F}}_{\wp_1(\beta)/\vee_1}^{(8/3)}(\xi, \ddot{\mathbf{x}}_1) (x_2 - x_1)^{3/4} |\log(x_2 - x_1)| \\ &\quad + O((x_2 - x_1)^{3/4}), \quad \text{as } x_1, x_2 \rightarrow \xi. \end{aligned} \tag{6.15}$$

Note that Eq. (6.15) is exactly the second claim (1.40) in Proposition 1.17.

*Proof.* For  $\widehat{\mathcal{H}}_\beta^{(8/3)}(\mathbf{x})$  defined in (6.13), one can proceed as in the proof of (6.8) to show that

$$\begin{aligned} \widehat{\mathcal{H}}_\beta^{(8/3)}(\mathbf{x}) &= \frac{1}{C(8/3)} \widehat{\mathcal{H}}_{\wp_1(\beta)/\{1,2\}}^{(8/3)}(\ddot{\mathbf{x}}_1) + g_2(\xi, \ddot{\mathbf{x}}_1) (x_2 - x_1)^{-1/4} + O((x_2 - x_1)^{3/4}) \\ &\quad + \frac{4}{C(8/3)} (x_2 - x_1)^{-5/4} (x_{b_1} - x_{b_2})^{-2} (x_{b_1} - \xi)^2 (x_{b_2} - \xi)^2 \int_{\Gamma_\beta} d\ddot{\mathbf{u}}_1 \widehat{f}_{\beta/\vee_1}^{(8/3)}(\xi, \ddot{\mathbf{x}}_1; \ddot{\mathbf{u}}_1) \\ &\quad \times \int_{10\frac{\chi}{1-\chi}}^{\frac{9}{10}} dw \left( \left( w + \frac{\chi}{1-\chi} \right)^{5/2} - w^{5/2} \right) w^{-3/2} (1-w)^{-3/2}, \end{aligned} \tag{6.16}$$

as  $x_1, x_2 \rightarrow \xi$ , where the integration  $\Gamma_{\hat{\beta}}$  comprises all of the other contours than the two Pochhammer contours surrounding the pairs  $\{x_1, x_{b_1}\}$  and  $\{x_2, x_{b_2}\}$ , and  $\chi = \text{CR}(x_1, x_2, x_{b_2}, x_{b_1})$ . The Taylor expansion

$$\left(w + \frac{\chi}{1-\chi}\right)^{5/2} - w^{5/2} = w^{5/2} \left( \frac{5}{2} \frac{\chi}{w(1-\chi)} + \frac{15}{8} \left(\frac{\chi}{w(1-\chi)}\right)^2 + O\left(\left(\frac{\chi}{w(1-\chi)}\right)^3\right) \right), \quad \text{for } w \in \left(10 \frac{\chi}{1-\chi}, \frac{9}{10}\right),$$

shows that there exists a function  $\hat{g}_2$  such that

$$\begin{aligned} \widehat{\mathcal{H}}_{\beta}^{(8/3)}(\mathbf{x}) &= \frac{1}{C(8/3)} \widehat{\mathcal{H}}_{\wp_1(\beta)/\{1,2\}}^{(8/3)}(\ddot{\mathbf{x}}_1) + \hat{g}_2(\xi, \ddot{\mathbf{x}}_1) (x_2 - x_1)^{-1/4} + O((x_2 - x_1)^{3/4}) \\ &\quad + \frac{15}{2} \frac{1}{C(8/3)} (x_2 - x_1)^{3/4} \int_{\Gamma_{\hat{\beta}}} d\ddot{\mathbf{u}}_1 \hat{f}_{\beta/\vee_1}^{(8/3)}(\xi, \ddot{\mathbf{x}}_1; \ddot{\mathbf{u}}_1) \int_{10 \frac{\chi}{1-\chi}}^{\frac{9}{10}} \frac{dw}{w} \\ &= \frac{1}{C(8/3)} \widehat{\mathcal{H}}_{\wp_1(\beta)/\{1,2\}}^{(8/3)}(\ddot{\mathbf{x}}_1) + \hat{g}_2(\xi, \ddot{\mathbf{x}}_1) (x_2 - x_1)^{-1/4} + O((x_2 - x_1)^{3/4}) \\ &\quad + \frac{15}{2} \frac{1}{C(8/3)} (x_2 - x_1)^{3/4} |\log(x_2 - x_1)| \int_{\Gamma_{\hat{\beta}}} d\ddot{\mathbf{u}}_1 \hat{f}_{\beta/\vee_1}^{(8/3)}(\xi, \ddot{\mathbf{x}}_1; \ddot{\mathbf{u}}_1), \end{aligned} \quad (6.17)$$

as  $x_1, x_2 \rightarrow \xi$ . However, letting  $\kappa \rightarrow 8/3$  in Eq. (6.11) shows that  $\hat{g}_2(\xi, \ddot{\mathbf{x}}_1) \equiv 0$ . Combining this observation with (6.17) and Remark 6.4 gives the asserted expansion (6.15).  $\square$

### 6.3 $\kappa = 8$ : proof of Proposition 1.18

In this section, we fix  $\kappa = 8$  and use the notation from Proposition 1.18.

*Proof of Proposition 1.18.* When  $\{1, 2\} \in \beta$ , the asserted expansion (1.41) is proven in [LPW24, Proposition 2.9]. When  $\{1, 2\} \notin \beta$ , the asserted expansion (1.42) is proven in Lemma 6.6.  $\square$

**Lemma 6.6.** *Fix  $\beta \in \text{LP}_N$  and suppose  $\{1, 2\} \notin \beta$ . Then, for all  $\xi \in (-\infty, x_3)$ , we have*

$$\widehat{\mathcal{F}}_{\beta}^{(8)}(\mathbf{x}) = \frac{1}{8} \widehat{\mathcal{F}}_{\wp_1(\beta)/\{1,2\}}^{(8)}(\ddot{\mathbf{x}}_1) (x_2 - x_1)^{1/4} |\log(x_{j+1} - x_j)| + O((x_2 - x_1)^{1/4}), \quad \text{as } x_1, x_2 \rightarrow \xi. \quad (6.18)$$

Note that Eq. (6.18) is the second claim (1.42) in Proposition 1.18.

*Proof.* Using Proposition 1.8 and the identity (6.6) from the proof of Lemma 6.2, we obtain

$$\begin{aligned} \widehat{\mathcal{F}}_{\beta}^{(8)}(\mathbf{x}) &= \lim_{\kappa' \rightarrow 8-} \frac{8}{\pi} \frac{\mathcal{F}_{\beta}^{(\kappa')}(\mathbf{x})}{(8 - \kappa')^N} = \frac{8}{32^N \pi} \mathcal{H}_{\beta}^{(8)}(\mathbf{x}) = \frac{8}{32^N \pi} \lim_{\kappa' \rightarrow 8-} \mathcal{H}_{\beta}^{(\kappa')}(\mathbf{x}) \\ &= \frac{1}{32^{N-1}} \left( \frac{(x_2 - x_1)(x_{b_1} - x_2)(x_{b_2} - x_1)}{(x_{b_1} - x_1)(x_{b_2} - x_2)} \right)^{1/4} \\ &\quad \times \int_{\Gamma_{\hat{\beta}}} d\ddot{\mathbf{u}}_1 f(\ddot{\mathbf{x}}_1; \ddot{\mathbf{u}}_1) \oint_{\vartheta(0,1)} dw \tilde{g}(x_1, x_2, w; \ddot{\mathbf{x}}_1; \ddot{\mathbf{u}}_1) \left(w + \frac{\chi}{1-\chi}\right)^{-1/2} (1-w)^{-1/2} w^{-1/2}, \end{aligned}$$

where the integration  $\Gamma_{\hat{\beta}}$  comprises all of the other contours than the two Pochhammer contours surrounding the pairs  $\{x_1, x_{b_1}\}$  and  $\{x_2, x_{b_2}\}$ , and  $\chi = \text{CR}(x_1, x_2, x_{b_2}, x_{b_1})$ , and where  $\tilde{g}$  is defined in (6.7) with  $\kappa = 8$ . One can proceed as in the proof of (6.8) to show that

$$\begin{aligned} \widehat{\mathcal{F}}_{\beta}^{(8)}(\mathbf{x}) &= O((x_2 - x_1)^{1/4}) \\ &\quad + \frac{4}{32^{N-1}} (x_2 - x_1)^{1/4} \int_{\Gamma_{\hat{\beta}}} d\ddot{\mathbf{u}}_1 f(\ddot{\mathbf{x}}_1; \ddot{\mathbf{u}}_1) \int_{10 \frac{\chi}{1-\chi}}^{\frac{9}{10}} \left(w + \frac{\chi}{1-\chi}\right)^{-1/2} (1-w)^{-1/2} w^{-1/2} \end{aligned} \quad (6.19)$$

as  $x_1, x_2 \rightarrow \xi$ . On the one hand, plugging the Taylor expansion

$$(w + \frac{\chi}{1-\chi})^{-1/2} = w^{-1/2} (1 + O(\frac{\chi}{w(1-\chi)})), \quad \text{for } w \in (10\frac{\chi}{1-\chi}, \frac{9}{10}),$$

into (6.19) shows that

$$\widehat{\mathcal{F}}_\beta^{(8)}(\mathbf{x}) = O((x_2 - x_1)^{1/4}) + (x_2 - x_1)^{1/4} |\log(x_2 - x_1)| \frac{4}{32^{N-1}} \int_{\Gamma_\beta} d\mathbf{\ddot{u}}_1 f(\mathbf{\ddot{x}}_1; \mathbf{\ddot{u}}_1). \quad (6.20)$$

On the other hand, according to Proposition 2.5, we have

$$\mathcal{H}_{\varphi_1(\beta)/\{1,2\}}^{(\kappa')}(\mathbf{\ddot{x}}_1) = \frac{\nu(\kappa')}{C(\kappa')} \int_{\Gamma_\beta} d\mathbf{\ddot{u}}_1 \hat{f}_\beta^a(\mathbf{\ddot{x}}_1; \mathbf{\ddot{u}}_1), \quad \text{for all } \kappa' \in (0, 8) \setminus \{\frac{8}{m} : m \in \mathbb{Z}_{>0}\}, \quad (6.21)$$

where  $\hat{f}_\beta^a$  is defined in (2.11) (with  $\kappa$  there replaced by  $\kappa'$ ). Taking the limit  $\kappa' \rightarrow 8-$  in (6.21) then gives

$$\int_{\Gamma_\beta} d\mathbf{\ddot{u}}_1 f(\mathbf{\ddot{x}}_1; \mathbf{\ddot{u}}_1) = \frac{1}{4\pi} \mathcal{H}_{\varphi_1(\beta)/\{1,2\}}^{(8)} = 32^{N-2} \widehat{\mathcal{F}}_{\varphi_1(\beta)/\{1,2\}}^{(8)}(\mathbf{\ddot{x}}_1). \quad (6.22)$$

Combining (6.20) with (6.22) gives (6.18), as desired.  $\square$

## A Hypoelliptic BPZ PDEs

In this appendix, we prove that local martingales arising from SLE variants give smooth solutions to the Belavin-Polyakov-Zamolodchikov (BPZ) type partial differential equation [BPZ84a]  $\mathcal{D}^{(i)}F = 0$ , where

$$\mathcal{D}^{(i)} := \frac{\kappa}{2} \partial_i^2 + \sum_{\substack{1 \leq j \leq n \\ j \neq i}} \left( \frac{2}{x_j - x_i} \partial_j - \frac{2\Delta_j}{(x_j - x_i)^2} \right), \quad (A.1)$$

for a fixed  $i \in \{1, 2, \dots, n\}$ , where the system has some given distinct marked points  $(x_1, \dots, x_n)$ , with  $x_i$  being the starting point of the  $\text{SLE}_\kappa$  curve, and  $(\Delta_j : 1 \leq j \leq n)$  are scaling exponents  $\Delta_j \in \mathbb{R}$ , such that  $\Delta_i = h(\kappa)$  is the  $\text{SLE}_\kappa$  boundary exponent for the point  $x_i$ . A general argument for this was already given in [Dub15b], and a special case was discussed in [PW19, Sections 2.3.3 and 4.1]. The general idea can be separated into two steps:

1. Local martingales for SLE variants give distributional solutions to the PDE (A.1) (Proposition A.6).
2. Any distributional solution to the PDE (A.1) is smooth (see Proposition A.2).

Property 2 is well-known: it holds by the fact that the PDE (A.1) is hypoelliptic:  $\mathcal{D}^{(i)}$  satisfies the Hörmander bracket condition by [Dub15b, Lemma 5] and [PW19, Lemma 2.6 & Remark 2.7]. We refer readers, e.g., to the textbooks [Str08, Chapter 7] for details concerning hypoelliptic PDEs, and [Rud91, Chapters 6–7] or [Hör90] for general theory of distributions needed in the present setup.

Property 1 was established in a very general setup by Julien Dubédat [Dub15b]. In [PW19, Lemma 4.4], Property 1 was presented with a simpler argument for pure partition functions of multiple chordal SLEs. However, the argument in [PW19, Lemma 4.4] contains a gap. There, one wants to extend transition semigroup operators  $(P_t)_{t \geq 0}$  associated to an Itô diffusion to continuous mappings on the space of distributions in its Fréchet topology. The key idea is to argue via the fact that the space of smooth compactly supported test functions is dense in the space of distributions with respect to its *Fréchet topology*. However, it is only known that  $P_t$  is bounded (thus continuous) on the space of smooth compactly supported test functions with respect to the *sup-norm*. Hence, it is not clear whether one can actually extend  $P_t$  to a continuous mapping for the Fréchet topology. More precisely, we do not know whether it is possible to derive [PW19, Equation (4.5)] via such a simple argument. We therefore will provide another argument for Property 1 in the spirit of Dubédat's original one [Dub15b]. We would like to thank Dapeng Zhan for pointing out this issue and for very useful correspondences in the process of fixing it.

## A.1 Distributions and hypoellipticity

Let  $O \subset \mathfrak{X}_n := \{(x_1, \dots, x_n) \in \mathbb{R}^n : x_i \neq x_j \text{ for all } i \neq j\}$  be an open subset. We denote by  $C^\infty(O; \mathbb{C})$  the set of smooth functions from  $O$  to  $\mathbb{C}$ , and by  $C_c^\infty(O; \mathbb{C})$  the space of smooth compactly supported functions from  $O$  to  $\mathbb{C}$ , and by  $(C_c^\infty)^*(O; \mathbb{C})$  the space of distributions, that is, the dual space of  $C_c^\infty(O; \mathbb{C})$  consisting of continuous linear functionals  $C_c^\infty(O; \mathbb{C}) \rightarrow \mathbb{C}$ . Recall that any locally integrable (e.g., continuous) function  $F$  on  $O$  defines a distribution, also denoted by  $F \in (C_c^\infty)^*(O; \mathbb{C})$ , via the assignment

$$\langle F, \phi \rangle := \int_O F(\mathbf{x}) \phi(\mathbf{x}) d\mathbf{x}, \quad (\text{A.2})$$

for all test functions  $\phi \in C_c^\infty(O; \mathbb{C})$ . Furthermore, with this identification, the space  $C_c^\infty(O; \mathbb{C}) \ni F$  of test functions is a dense subset in the space  $(C_c^\infty)^*(O; \mathbb{C})$  of distributions (see, e.g., [Tao09, Lemma 1.13.5]).

The differential operator  $\mathcal{D}^{(i)}$  from (A.1) defines a linear operator on the space of distributions via its transpose (dual operator)  $(\mathcal{D}^{(i)})^*$ : for a distribution  $F \in (C_c^\infty)^*(O; \mathbb{C})$ , we have  $\mathcal{D}^{(i)} F \in (C_c^\infty)^*(O; \mathbb{C})$  and

$$\langle \mathcal{D}^{(i)} F, \phi \rangle := \int_O F(\mathbf{x}) (\mathcal{D}^{(i)})^* \phi(\mathbf{x}) d\mathbf{x}, \quad \text{for all } \phi \in C_c^\infty(O; \mathbb{C}). \quad (\text{A.3})$$

**Definition A.1.** We say that  $F \in (C_c^\infty)^*(O; \mathbb{C})$  is a solution to the Dirichlet problem for  $\mathcal{D}^{(i)}$  in distributional sense if (A.3) equals zero for all test functions  $\phi \in C_c^\infty(O; \mathbb{C})$ . Abusing notation, we write

$$\mathcal{D}^{(i)} F = 0. \quad (\text{A.4})$$

The hypoellipticity of the above PDE operators (A.1) is detailed, e.g., in [PW19, Section 2.3.3]:

**Proposition A.2.** Fix  $\kappa > 0$ . The partial differential operator  $\mathcal{D}^{(i)}$  is hypoelliptic. In particular, any distributional solution to (A.4) is smooth.

*Proof.* See [Dub15b, Lemma 5] or [PW19, Lemma 2.6 & Remark 2.7].  $\square$

## A.2 Dirichlet problem

We shall next summarize the results in the article [Bon69] of J.-M. Bony concerning existence and uniqueness of solutions to Dirichlet problems for hypoelliptic differential operators, of which the one in Definition A.1 is a special instance, relevant for SLE theory.

**Theorem A.3.** Fix  $\kappa > 0$ . Fix

$$\mathbf{x}_0 \in \mathfrak{X}_n := \{\mathbf{x} := (x_1, \dots, x_n) \in \mathbb{R}^n : x_1 < \dots < x_n\}. \quad (\text{A.5})$$

Then,  $\mathbf{x}_0$  has an open neighborhood  $O_0 \subset \mathfrak{X}_n$  such that  $\overline{O}_0 \subset \mathfrak{X}_n$  and the following holds. For any continuous function  $\varphi: \partial O_0 \rightarrow \mathbb{C}$ , the Dirichlet boundary value problem for  $\mathcal{D}^{(i)}$  in  $O_0$ ,

$$\begin{cases} \mathcal{D}^{(i)} F(\mathbf{x}) = 0, & \mathbf{x} \in O_0, \\ F(\mathbf{x}) = \varphi(\mathbf{x}), & \mathbf{x} \in \partial O_0, \end{cases} \quad (\text{A.6})$$

has a unique solution  $F$ . Moreover, the solution  $F$  is smooth on  $O_0$ .

*Proof.* As in [Bon69, Proposition 5.1], there exist  $M_0, \varepsilon_0 \in (0, \infty)$  such that for the operator

$$[\tilde{\mathcal{D}}^{(i)} F](\mathbf{x}) := H_{\mathbf{x}_0}(\mathbf{x}) \mathcal{D}^{(i)} (H_{\mathbf{x}_0} F)(\mathbf{x}),$$

where  $H_{\mathbf{x}_0}(\mathbf{x}) := 1 - M_0 |\mathbf{x} - \mathbf{x}_0|^2$ , the constant term is negative for all  $\mathbf{x} \in \mathfrak{X}_n$  with  $|\mathbf{x} - \mathbf{x}_0| < \varepsilon_0$ ,

$$[\tilde{\mathcal{D}}^{(i)} 1](\mathbf{x}) < 0, \quad \text{for all } \mathbf{x} \in B(\mathbf{x}_0, \varepsilon_0) \cap \mathfrak{X}_n,$$

and the second order term of  $\tilde{\mathcal{D}}^{(i)}$  equals  $\frac{\kappa}{2}(H_{\mathbf{x}_0}(\mathbf{x}))^2 > 0$  on  $B(\mathbf{x}_0, \varepsilon_0) \ni \mathbf{x}$ . Without loss of generality, we may also assume that  $B(\mathbf{x}_0, 2\varepsilon_0) \subset \mathfrak{X}_n$ . We can then apply [Bon69, Theorem 5.2] (see Theorem A.4 stated below in our setup) to the modified operator  $\tilde{\mathcal{D}}^{(i)}$ , after we find an open bounded subset  $O_0 \subset B(\mathbf{x}_0, \varepsilon_0)$  such that at any point in  $\partial O_0$  there exists an outer normal vector  $\boldsymbol{\nu} = (\nu_1, \dots, \nu_n)$  to  $\partial O_0$  satisfying  $\nu_i \neq 0$ . To this end, we can take the intersection of two balls  $B(\mathbf{x}_0 - \lambda \hat{e}_i, \varepsilon + \lambda)$  and  $B(\mathbf{x}_0 + \lambda \hat{e}_i, \varepsilon + \lambda)$ , where  $\hat{e}_i$  is the unit vector to the  $i$ :th direction (the index  $i$  is fixed in the statement to be the one for which the second order term in  $\tilde{\mathcal{D}}^{(i)}$ , and thus in  $\mathcal{D}^{(i)}$ , is non-zero), and  $\lambda, \varepsilon \in (0, \infty)$  are chosen so that

$$O_0 := B(\mathbf{x}_0 - \lambda \hat{e}_i, \varepsilon + \lambda) \cap B(\mathbf{x}_0 + \lambda \hat{e}_i, \varepsilon + \lambda) \subset B(\mathbf{x}_0, \varepsilon_0).$$

Then, clearly at any point on the smooth part of  $\partial O_0$  the sought outer normal vector exists, while any corner point of  $\partial O_0$  has several outer normal vectors (according to [Bon69, Definition 2.1]), and one of them satisfies  $\nu_i \neq 0$ . This concludes the proof, given [Bon69, Theorem 5.2] (Theorem A.4).  $\square$

In the version of [Bon69, Theorem 5.2] relevant to the present work, we assume for simplicity that the second order term in the differential operator of interest only comprises one derivative,  $\partial_i^2$ , like in SLE applications. This facilitates the application of the exterior ball condition (Assumption 2 in Theorem A.4).

**Theorem A.4.** (Special case of [Bon69, Theorem 5.2].) *Let  $\mathcal{D}$  be a hypoelliptic partial differential operator on a subset  $\mathfrak{U}_n$  such that its second order term equals a positive function times  $\partial_i^2$ . Consider an open bounded connected subset  $O \subset \mathfrak{X}_n$  such that  $\bar{O} \subset \mathfrak{X}_n$ . Suppose furthermore that the following hold.*

1. *The constant term of  $\mathcal{D}$  is strictly negative on  $O$ , i.e.,*

$$[\mathcal{D}1](\mathbf{x}) < 0, \quad \text{for all } \mathbf{x} \in O.$$

2. *At any point in  $\partial O$  there exists an outer normal vector  $\boldsymbol{\nu} = (\nu_1, \dots, \nu_n)$  to  $\partial O$  satisfying  $\nu_i \neq 0$ .*

*Then, for any continuous function  $\varphi: \partial O \rightarrow \mathbb{C}$ , the Dirichlet boundary value problem for  $\mathcal{D}$ ,*

$$\begin{cases} \mathcal{D}F(\mathbf{x}) = 0, & \mathbf{x} \in O, \\ F(\mathbf{x}) = \varphi(\mathbf{x}), & \mathbf{x} \in \partial O, \end{cases}$$

*has a unique solution  $F$  in distributional sense. Moreover, the solution  $F$  is smooth on  $O$ .*

### A.3 Application to SLE partition functions

Consider a chordal  $\text{SLE}_\kappa$  curve  $\eta$  in  $\mathbb{H}$  from  $x_i$  to  $\infty$ , and denote by  $(W_t, t \geq 0)$  its driving function and by  $(g_t, t \geq 0)$  the corresponding family of conformal maps normalized at infinity. Suppose that  $\kappa > 0$ . In applications, one often encounters local martingales for  $\eta$  of the form

$$\begin{aligned} M_t(\mathcal{Z}) &:= \prod_{j \neq i} g'_t(x_j)^{\Delta_j} \times \mathcal{Z}(g_t(x_1), \dots, g_t(x_{i-1}), W_t, g_t(x_{i+1}), \dots, g_t(x_n)) \\ &= \psi(X_t), \end{aligned} \tag{A.7}$$

where  $\mathcal{Z}: \mathfrak{X}_n \rightarrow \mathbb{C}$  is a continuous function (which will be an  $\text{SLE}_\kappa$  partition function),

$$X_t = (g_t(x_1), \dots, g_t(x_{i-1}), W_t, g_t(x_{i+1}), \dots, g_t(x_n), g'_t(x_1), \dots, g'_t(x_{i-1}), g'_t(x_{i+1}), \dots, g'_t(x_n)), \quad t \geq 0,$$

is an Itô process, whose infinitesimal generator, when acting on twice continuously differentiable functions, has the form

$$A = \frac{\kappa}{2} \partial_i^2 + \sum_{j \neq i} \left( \frac{2}{x_j - x_i} \partial_j - \frac{2y_j}{(x_j - x_i)^2} \partial_{n-1+j} \right), \tag{A.8}$$



and we use the shorthand notation

$$\psi(\mathbf{x}, \dot{\mathbf{y}}_i) = \prod_{j \neq i} y_j^{\Delta_j} \times \mathcal{Z}(\mathbf{x})$$

for the continuous function of  $(\mathbf{x}, \dot{\mathbf{y}}_i) := (x_1, \dots, x_n, y_1, \dots, y_{i-1}, y_{i+1}, \dots, y_n) \in \mathfrak{X}_n \times \mathbb{R}^{n-1}$  (independent of  $y_i$ ). A calculation shows that the two differential operators  $A$  and  $\mathcal{D}^{(i)}$  defined in (A.1) are related via

$$A \prod_{j \neq i} y_j^{\Delta_j} \times \phi(\mathbf{x}) = \prod_{j \neq i} y_j^{\Delta_j} \times \mathcal{D}^{(i)} \phi(\mathbf{x}), \quad (\text{A.9})$$

for any nice enough function  $\phi$  (to be determined later).

Let  $T$  be the first time when one of the marked points  $\{x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n\}$  is swallowed by  $\eta$ .

**Lemma A.5.** *There exists an open bounded set  $O_0 \subset \mathfrak{X}_n$  such that  $\overline{O_0} \subset \mathfrak{X}_n$  and the following holds. For any  $\mathbf{x} \in O_0$ , for the Itô process  $(X_t: t \geq 0)$  started at  $X_0 = (\mathbf{x}, \dot{\mathbf{i}})$ , the exit time*

$$\tau_0 := \inf\{s \geq 0: X_s \in \partial(O_0 \times \mathbb{R}^{n-1})\}$$

satisfies

1.  $\tau_0 < T$  almost surely;
2.  $(g_{\tau_0}(x_1), \dots, g_{\tau_0}(x_{i-1}), W_{\tau_0}, g_{\tau_0}(x_{i+1}), \dots, g_{\tau_0}(x_n)) \in \partial O_0$  almost surely;
3.  $\mathbb{E}_{(\mathbf{x}, \dot{\mathbf{i}})}[\tau_0] < \infty$ .

*Proof.* From the Loewner equation (3.1), we have

$$g'_t(z) = \exp\left(-\int_0^t \frac{2 ds}{(g_s(z) - W_s)^2}\right), \quad t \geq 0 \quad \text{and} \quad z \in H_t,$$

where  $H_t$  is the unbounded component of  $\mathbb{H} \setminus \eta[0, t]$ . In particular, because some of the first  $n$  coordinates in  $X_t$  must collide at the swallowing time  $T$  and are separated for all times before time  $T$ , by definition (A.5) of  $\mathfrak{X}_n$  we see that for any open bounded set  $O_0 \subset \overline{O_0} \subset \mathfrak{X}_n$ , the inequality  $\tau_0 < T$  holds almost surely (Items 1), and Item 2 holds, as the derivatives of  $g_t$  stay bounded up to and including the time  $\tau_0$ .

For Item 3, by Item 2 it suffices to note that for any open bounded  $O_0 \subset \overline{O_0} \subset \mathfrak{X}_n$ , the expected hitting time of  $(g_{\tau_0}(x_1), \dots, g_{\tau_0}(x_{i-1}), W_{\tau_0}, g_{\tau_0}(x_{i+1}), \dots, g_{\tau_0}(x_n))$  to  $\partial O_0$  is finite (cf. [KS98, Lemma 7.4]).  $\square$

The goal of this appendix is to show that in this situation,  $\mathcal{Z}: \mathfrak{X}_n \rightarrow \mathbb{C}$  is always a smooth solution to the PDE  $\mathcal{D}^{(i)} \mathcal{Z} = 0$ . To this end, by Proposition A.2 it suffices to prove the following result.

**Proposition A.6.** *Fix  $\mathbf{x}_0 \in \mathfrak{X}_n$ . Suppose  $\mathcal{Z}: \mathfrak{X}_n \rightarrow \mathbb{C}$  is a continuous function such that  $(M_t(\mathcal{Z}): t \geq 0)$  defined in (A.7) is a local martingale for the chordal SLE $_{\kappa}$  curve in  $\mathbb{H}$  from  $x_i$  to  $\infty$ . Then,  $\mathbf{x}_0$  has an open neighborhood  $O_0 \subset \mathfrak{X}_n$  such that*

$$\mathcal{D}^{(i)} \mathcal{Z}(\mathbf{x}) = 0 \quad \text{for all } \mathbf{x} \in O_0$$

(in distributional sense as in Definition A.1, and thus a fortiori also as a smooth function).

*Proof.* By Theorem A.3,  $\mathbf{x}_0$  has an open neighborhood  $O_0 \subset \mathfrak{X}_n$  such that  $\overline{O_0} \subset \mathfrak{X}_n$  and the Dirichlet boundary value problem (A.6) for  $\mathcal{D}^{(i)}$  in  $O_0$  with boundary condition

$$\varphi(\mathbf{x}) := \mathcal{Z}(\mathbf{x}) = \psi(\mathbf{x}, \dot{\mathbf{i}}), \quad \mathbf{x} \in \partial O_0$$

(here,  $\mathbf{i} = (1, 1, \dots, 1) \in \mathbb{R}^{n-1}$ ), has a unique (smooth) solution  $F \in C_c^\infty(O_0; \mathbb{C})$ . Write also

$$\tilde{Z}(\mathbf{x}, \mathbf{y}_i) := \prod_{j \neq i} y_j^{\Delta_j} \times F(\mathbf{x}),$$

which is a smooth function on  $O_0 \times \mathbb{R}^{n-1}$ . Then, taking  $\phi = F$  in (A.9), the right-hand side of (A.9) equals zero, which implies that  $A\tilde{Z}(\mathbf{x}, \mathbf{y}_i) = 0$  for all  $(\mathbf{x}, \mathbf{y}_i) \in O_0 \times \mathbb{R}^{n-1}$ .

Let  $\mathbb{E}_{(\mathbf{x}, \mathbf{y}_i)}$  denote the expected value with respect to the Itô process  $X_t$  with generator  $A$  given by (A.8) started at  $X_0 = (\mathbf{x}, \mathbf{i})$ . Dynkin's formula [Øks03, Theorem 7.4.1] gives

$$\mathbb{E}_{(\mathbf{x}, \mathbf{i})}[\tilde{Z}(X_\tau)] = \tilde{Z}(\mathbf{x}, \mathbf{i}) + \mathbb{E}_{(\mathbf{x}, \mathbf{i})} \left[ \int_0^\tau A\tilde{Z}(X_s) ds \right],$$

for any stopping time  $\tau$  such that  $\mathbb{E}_{(\mathbf{x}, \mathbf{i})}[\tau] < \infty$ . After possibly shrinking the neighborhood  $O_0$ , applying this to the exit time  $\tau_0 := \inf\{s \geq 0 : X_s \in \partial(O_0 \times \mathbb{R}^{n-1})\}$  from Lemma A.5 (given below), we obtain

$$\mathbb{E}_{(\mathbf{x}, \mathbf{i})}[\tilde{Z}(X_{\tau_0})] = \tilde{Z}(\mathbf{x}, \mathbf{i}) + \mathbb{E}_{(\mathbf{x}, \mathbf{i})} \left[ \int_0^{\tau_0} A\tilde{Z}(X_s) ds \right] = \tilde{Z}(\mathbf{x}, \mathbf{i}), \quad \mathbf{x} \in O_0,$$

since  $A\tilde{Z}(\mathbf{x}, \mathbf{y}_i) = 0$  for all  $(\mathbf{x}, \mathbf{y}_i) \in O_0 \times \mathbb{R}^{n-1}$ . This shows that

$$F(\mathbf{x}) = \mathbb{E}_{(\mathbf{x}, \mathbf{i})}[\tilde{Z}(X_{\tau_0})], \quad \mathbf{x} \in O_0, \quad (\text{A.10})$$

and in particular, that  $\mathbf{x} \mapsto \mathbb{E}_{(\mathbf{x}, \mathbf{i})}[\tilde{Z}(X_{\tau_0})]$  is smooth. Note also that

$$\begin{aligned} \tilde{Z}(X_{\tau_0}) &= \prod_{j \neq i} g'_{\tau_0}(x_j)^{\Delta_j} \times \mathcal{Z}(g_{\tau_0}(x_1), \dots, g_{\tau_0}(x_{i-1}), W_{\tau_0}, g_{\tau_0}(x_{i+1}), \dots, g_{\tau_0}(x_n)) \\ &= \psi(X_{\tau_0}) = M_{\tau_0}(\mathcal{Z}). \end{aligned}$$

Moreover, since  $\mathcal{Z}$  is continuous and  $\tau_0 < T$  almost surely by Lemma A.5, the martingale  $M_t(\mathcal{Z})$  is bounded for  $t \leq \tau_0$ . Optional stopping theorem thus gives

$$\mathbb{E}_{(\mathbf{x}, \mathbf{i})}[\tilde{Z}(X_{\tau_0})] = \mathbb{E}_{(\mathbf{x}, \mathbf{i})}[M_{\tau_0}(\mathcal{Z})] = M_0(\mathcal{Z}) = \mathcal{Z}(\mathbf{x}), \quad \mathbf{x} \in O_0. \quad (\text{A.11})$$

Hence, we conclude from (A.10, A.11) that the identity  $\mathcal{Z} = F$  holds not only on  $\partial O_0$  but also on  $O_0$ , so in particular,  $\mathcal{Z}$  is a smooth solution to the PDE  $\mathcal{D}^{(i)}\mathcal{Z} = 0$  in  $O_0$ , as claimed.  $\square$

**Remark A.7.** Note that uniqueness of solutions to hypoelliptic boundary value problems as in Theorem A.3 is not necessarily applicable on the whole set  $\mathfrak{X}_n$  in our situation, because the solutions that we consider cannot necessarily be continuously extended up to the boundary of  $\mathfrak{X}_n$ .

## B Some integral identities

The identities gathered in this appendix are used in Coulomb gas computations.

**Lemma B.1.** *We have*

$$\oint_{\partial(0,1)} u^{-4/\kappa} (u-1)^{-4/\kappa} du = \frac{\nu(\kappa)}{\mathcal{C}(\kappa)}, \quad \text{for } \kappa \in \mathbb{C} \setminus (\{\frac{8}{m} : m \in \mathbb{Z}_{>0}\} \cup \{0\}); \quad (\text{B.1})$$

$$\oint_{\partial(0,1)} u^{-4/\kappa} (u-1)^{-4/\kappa} (2u-1) du = 0, \quad \text{for } \kappa \in \mathbb{C} \setminus \{0\}; \quad (\text{B.2})$$

and

$$\begin{aligned} \oint_{\partial(0,1)} u^{12/\kappa-2} (1-u)^{-4/\kappa} du &= 4 \sin^2(4\pi/\kappa) (1-\nu(\kappa))^2 e^{16\pi i/\kappa} \frac{\Gamma(12/\kappa-1) \Gamma(1-4/\kappa)}{\Gamma(8/\kappa)}, \\ &\text{for } \kappa \in \mathbb{C} \setminus (\{\frac{8}{m} : m \in (2\mathbb{Z}_{>0}) \cup (\mathbb{Z}_{<0})\} \cup \{0\}); \end{aligned} \quad (\text{B.3})$$

*Proof.* We first verify (B.1): for  $\kappa \in (4, 8)$ , we have

$$\begin{aligned} \oint_{\vartheta(0,1)} u^{-4/\kappa} (u-1)^{-4/\kappa} du &= (1 - e^{8\pi i/\kappa} + 1 - e^{-8\pi i/\kappa}) \int_0^1 |u(u-1)|^{-4/\kappa} du \\ &= 4 \sin^2(4\pi/\kappa) \frac{\Gamma(1-4/\kappa)^2}{\Gamma(2-8/\kappa)} = \frac{\nu(\kappa)}{C(\kappa)}. \end{aligned}$$

The identity (B.1) for  $\kappa \in \mathbb{C} \setminus (\{\frac{8}{m} : m \in \mathbb{Z}_{>0}\} \cup \{0\})$  follows by analytic continuation.

Let us next verify (B.2): for  $\kappa \in (4, 8)$ , we have

$$\begin{aligned} \oint_{\vartheta(0,1)} u^{-4/\kappa} (u-1)^{-4/\kappa} (2u-1) du &= (1 - e^{8\pi i/\kappa} + 1 - e^{-8\pi i/\kappa}) \int_0^1 |u(u-1)|^{-4/\kappa} (2u-1) du \\ &= 4 \sin^2(4\pi/\kappa) \left( 2 \frac{\Gamma(2-4/\kappa)\Gamma(1-4/\kappa)}{\Gamma(3-8/\kappa)} - \frac{\Gamma(1-4/\kappa)^2}{\Gamma(2-8/\kappa)} \right) \\ &= 4 \sin^2(4\pi/\kappa) \frac{\Gamma(1-4/\kappa)^2}{\Gamma(2-8/\kappa)} (1-1) = 0. \end{aligned}$$

The identity (B.2) for  $\kappa \in \mathbb{C} \setminus \{0\}$  follows by analytic continuation.

Third, we prove (B.3): for  $\kappa \in (4, 8)$ , we have

$$\begin{aligned} \oint_{\vartheta(0,1)} u^{12/\kappa-1} (u-1)^{-4/\kappa} du &= (1 - e^{8\pi i/\kappa} + e^{32\pi i/\kappa} - e^{24\pi i/\kappa}) \int_0^1 u^{12/\kappa-1} (u-1)^{4/\kappa} du \\ &= 4 \sin^2(4\pi/\kappa) (1 - \nu(\kappa)^2) e^{16\pi i/\kappa} \frac{\Gamma(12/\kappa-1)\Gamma(1-4/\kappa)}{\Gamma(8/\kappa)}. \end{aligned}$$

The identity (B.3) for  $\kappa \in \mathbb{C} \setminus (\{\frac{8}{m} : m \in (2\mathbb{Z}_{>0}) \cup (\mathbb{Z}_{<0})\} \cup \{0\})$  follows by analytic continuation.  $\square$

**Lemma B.2.** *We have*

$$\begin{aligned} \int_{\varrho(0,\infty)} u^{-4/\kappa} (1+u)^{-4/\kappa} du &= \frac{(1 - e^{-8\pi i/\kappa})}{\nu(\kappa)} \frac{\Gamma(1-4/\kappa)^2}{\Gamma(2-8/\kappa)}, \\ &\text{for } \kappa \in \{\kappa' \in \mathbb{C} : 0 < |\kappa'| < 8\} \setminus \{\frac{8}{m} : m \in \mathbb{Z}_{>0}\}; \end{aligned} \quad (\text{B.4})$$

and

$$\int_{\varrho(0,\infty)} u^{-4/\kappa} (1+u)^{-4/\kappa} \left( \frac{6}{\kappa} - 1 + \frac{4}{\kappa} \left( \frac{u}{1+u} + u \right) \right) du = 0, \quad \text{for } 0 < |\kappa| < 8; \quad (\text{B.5})$$

$$\int_{\varrho(0,\infty)} u^{-4/\kappa} (1+u)^{-4/\kappa} (2u+1) du = 0, \quad \text{for } 0 < |\kappa| < 8; \quad (\text{B.6})$$

where  $\varrho(0, \infty)$  is a contour which starts from  $\infty$ , then winds around 0 once in clockwise direction and finally goes back to  $\infty$ ,

$$\varrho(0, \infty) := \text{---} \left( \text{---} \bigcirc \text{---} \right) \text{---} \quad (\text{B.7})$$

and the branch of  $u \mapsto u^{-4/\kappa} (1+u)^{-4/\kappa}$  is chosen to be real and positive when  $0 < \text{Re}(u) < \infty$ .

*Proof.* First, we prove (B.4): for  $\kappa \in (4, 8)$ , we have

$$\int_{\varrho(0,\infty)} u^{-4/\kappa} (1+u)^{-4/\kappa} du = (1 - e^{-8\pi i/\kappa}) \int_0^\infty u^{-4/\kappa} (1+u)^{-4/\kappa} du$$

$$\begin{aligned}
&= (1 - e^{-8\pi i/\kappa}) \frac{\Gamma(1 - 4/\kappa)\Gamma(8/\kappa - 1)}{\Gamma(4/\kappa)} \\
&= \frac{(1 - e^{-8\pi i/\kappa})}{\nu(\kappa)} \frac{\Gamma(1 - 4/\kappa)^2}{\Gamma(2 - 8/\kappa)}, \quad \text{since} \quad \Gamma(1 - z)\Gamma(z) = \frac{\pi}{\sin(\pi z)}.
\end{aligned}$$

The identity (B.4) for  $\kappa \in \{\kappa' \in \mathbb{C} : 0 < |\kappa'| < 8\} \setminus \{\frac{8}{m} : m \in \mathbb{Z}_{>0}\}$  follows by analytic continuation.

Next, we prove (B.5, B.6): for  $\kappa \in (4, 8)$ , we have

$$\begin{aligned}
&\frac{1}{4 \sin^2(4\pi/\kappa)} \int_{\varrho(0, \infty)} u^{-4/\kappa} (1+u)^{-4/\kappa} \left( \frac{6}{\kappa} - 1 + \frac{4}{\kappa} \left( \frac{u}{1+u} + u \right) \right) du \\
&= \left( \frac{6}{\kappa} - 1 \right) \frac{\Gamma(1 - 4/\kappa)\Gamma(8/\kappa - 1)}{\Gamma(4/\kappa)} + \frac{4}{\kappa} \left( \frac{\Gamma(2 - 4/\kappa)\Gamma(8/\kappa - 1)}{\Gamma(1 + 4/\kappa)} + \frac{\Gamma(2 - 4/\kappa)\Gamma(8/\kappa - 2)}{\Gamma(4/\kappa)} \right) \\
&= \frac{\Gamma(1 - 4/\kappa)\Gamma(8/\kappa - 1)}{\Gamma(4/\kappa)} \left( \frac{6}{\kappa} - 1 + 1 - \frac{4}{\kappa} - \frac{2}{\kappa} \right) = 0; \\
&\frac{1}{4 \sin^2(4\pi/\kappa)} \int_{\varrho(0, \infty)} u^{-4/\kappa} (1+u)^{-4/\kappa} (2u+1) du \\
&= \frac{2\Gamma(2 - 4/\kappa)\Gamma(8/\kappa - 2)}{\Gamma(4/\kappa)} + \frac{\Gamma(1 - 4/\kappa)\Gamma(8/\kappa - 1)}{\Gamma(4/\kappa)} \\
&= \frac{\Gamma(1 - 4/\kappa)\Gamma(8/\kappa - 1)}{\Gamma(4/\kappa)} (-1 + 1) = 0,
\end{aligned}$$

and for  $0 < |\kappa| < 8$  the identities (B.5, B.6) follow by analytic continuation.  $\square$

## C Additional formulas for Coulomb gas integrals

This section gathers some useful formulas for Sections 3–4. We will consider the Coulomb gas type integrands with  $d \geq 2$  variables and  $\ell \geq 0$  integration variables (with any branch choice),

$$f(\mathbf{x}; \mathbf{u}) := \prod_{1 \leq i < j \leq d} (x_j - x_i)^{2/\kappa} \prod_{1 \leq r < s \leq \ell} (u_s - u_r)^{8/\kappa} \prod_{\substack{1 \leq i \leq d \\ 1 \leq r \leq \ell}} (u_r - x_i)^{-4/\kappa}$$

having negative screening charges  $\alpha_- = -2/\sqrt{\kappa}$  at the integration variables  $\mathbf{u} = (u_1, u_2, \dots, u_\ell)$ , and symmetric charges  $\alpha_j = 1/\sqrt{\kappa}$  for all  $j \in \{1, 2, \dots, d\}$  at all of the variables  $\mathbf{x} = (x_1, x_2, \dots, x_d)$ , and

$$\begin{aligned}
\hat{f}(\mathbf{x}; \mathbf{v}) &:= \prod_{2 \leq i < j \leq d} (x_j - x_i)^{2/\kappa} \prod_{2 \leq j \leq d} (x_j - x_1)^{1-6/\kappa} \\
&\times \prod_{1 \leq t < s \leq \ell-1} (v_s - v_t)^{8/\kappa} \prod_{\substack{2 \leq i \leq d \\ 1 \leq s \leq \ell-1}} (v_s - x_i)^{-4/\kappa} \prod_{1 \leq s \leq \ell-1} (v_s - x_1)^{12/\kappa-2}, \tag{C.1}
\end{aligned}$$

having the conjugate charge  $\alpha_1 = \frac{\kappa-6}{2\sqrt{\kappa}}$  at  $x_1$  and charges  $\alpha_j = 1/\sqrt{\kappa}$  for all other  $j \in \{2, 3, \dots, d\}$ . When  $d = 2N$  and  $\ell = N$ , the latter integrand satisfies the charge neutrality [DFMS97, FK15c]:

$$\left( \sum_{j=2}^{2N} \alpha_j \right) + \left( \frac{\kappa-6}{2\sqrt{\kappa}} \right) + (N-1)\alpha_- = 2\alpha_0. \tag{C.2}$$

**Remark C.1.** Note that when  $\ell = m/2$ , the residue of  $\hat{f}$  at infinity is well-defined and equals zero:

$$\text{Res}_{v_r=\infty} \hat{f}(\mathbf{x}; \mathbf{v}) = - \lim_{|v_r| \rightarrow \infty} v_r \hat{f}(\mathbf{x}; \mathbf{v}) = - \lim_{|v_r| \rightarrow \infty} v_r^{-1} = 0, \quad 1 \leq r \leq \ell - 1.$$

In particular, the integrand function  $v_r \mapsto \hat{f}(\mathbf{x}; \mathbf{v})$  is holomorphic at infinity for all  $1 \leq r \leq \ell - 1$ .

**Lemma C.2.** Fix  $\kappa \in (4, 8)$ . We have

$$\begin{aligned} \mathcal{F}_{\overbrace{\underbrace{\quad}_N}(\mathbf{x})} &= e^{4\pi i/\kappa} \nu(\kappa) \hat{C}(\kappa)^N \int_{x_2}^{x_3} dv_1 \cdots \int_{x_{2N-2}}^{x_{2N-1}} dv_{N-1} \int_{x_{2N}}^{+\infty} dv_N f_{\overbrace{\underbrace{\quad}_N}(\mathbf{x}; \mathbf{v})} \\ &= \nu(\kappa) \hat{C}(\kappa)^N \int_{x_2}^{x_3} dv_1 \cdots \int_{x_{2N-2}}^{x_{2N-1}} dv_{N-1} \int_{x_{2N}}^{+\infty} dv_N |f(\mathbf{x}; \mathbf{v})|, \quad \mathbf{x} \in \mathfrak{X}_{2N}, \end{aligned} \quad (\text{C.3})$$

and

$$\begin{aligned} \mathcal{F}_{\underbrace{\overbrace{\quad}_N}(\mathbf{x})} &= e^{-4\pi i/\kappa} \nu(\kappa) \hat{C}(\kappa)^N \int_{-\infty}^{x_1} dv_1 \int_{x_3}^{x_4} dv_2 \cdots \int_{x_{2N-1}}^{x_{2N}} dv_N f_{\underbrace{\overbrace{\quad}_N}(\mathbf{x}; \mathbf{v})} \\ &= \nu(\kappa) \hat{C}(\kappa)^N \int_{-\infty}^{x_1} dv_1 \int_{x_3}^{x_4} dv_2 \cdots \int_{x_{2N-1}}^{x_{2N}} dv_N |f(\mathbf{x}; \mathbf{v})|, \quad \mathbf{x} \in \mathfrak{X}_{2N}, \end{aligned} \quad (\text{C.4})$$

where (see Figure C.1)

$$\hat{C}(\kappa) := 4 \sin^2(4\pi/\kappa) C(\kappa) = \nu(\kappa) \frac{\Gamma(2 - 8/\kappa)}{\Gamma(1 - 4/\kappa)^2}.$$

In particular, we have  $\mathcal{F}_{\overbrace{\underbrace{\quad}_N}(\mathbf{x})} > 0$  and  $\mathcal{F}_{\underbrace{\overbrace{\quad}_N}(\mathbf{x})} > 0$  for all  $\mathbf{x} \in \mathfrak{X}_{2N}$ .




Figure C.1: The left panel depicts a plot of the function  $\kappa \mapsto \hat{C}(\kappa)$ . Note that  $\hat{C}(4) = 0 = \hat{C}(8)$ . The right panel depicts a plot of the function  $\kappa \mapsto C(\kappa)$ . Note that  $C(8) = 0$  and  $C(\kappa)$  diverges as  $\kappa \rightarrow 4+$ .

*Proof.* The two lines on the right-hand side of (C.3) (resp. (C.4)) are obtained from each other by the branch choice (1.13) of the integrand  $f_{\overbrace{\underbrace{\quad}_N}(\mathbf{x}; \mathbf{v})}$  (resp.  $f_{\underbrace{\overbrace{\quad}_N}(\mathbf{x}; \mathbf{v})}$ ). To prove the identity (C.3), we note on the one hand that, by the limit property ( $\mathcal{F}$ -ASY $_{\infty}$ ) in Corollary 2.13,

$$\lim_{R \rightarrow \infty} R^{2h(\kappa)} \mathcal{F}_{\underbrace{\overbrace{\underbrace{\quad}_{N+1}}(-R, \mathbf{x}, R)} = \mathcal{F}_{\overbrace{\underbrace{\quad}_N}(\mathbf{x}).$$

On the other hand, using Proposition 2.5 for  $\beta = \underbrace{\overbrace{\quad}_{N+1}}$  and  $r = 1$  in (2.9) so that  $a_1 = 1$  and  $b_1 = 2$  and writing the Pochhammer integrations as line integrals<sup>23</sup> by taking  $\epsilon \rightarrow 0$  in Lemma 2.8,

$$\oint_{\vartheta(x_{a_s}, x_{b_s})} du_s \hat{f}(\mathbf{x}; \mathbf{u}) = 4 \sin^2(4\pi/\kappa) \int_{x_{a_s}}^{x_{b_s}} du_s \hat{f}(\mathbf{x}; \mathbf{u}), \quad (\text{C.5})$$

which holds for all  $s$  such that neither  $x_{a_s}$  nor  $x_{b_s}$  carries the conjugate charge (i.e.,  $s \in \{2, 3, \dots, \ell\}$  since  $\hat{f}$  in (C.1) has the conjugate charge at  $x_1$ ), and taking note of the branch choice of  $\hat{f}_{\underbrace{\overbrace{\quad}_{N+1}}^a}$  in (2.11), we obtain

$$\mathcal{F}_{\underbrace{\overbrace{\underbrace{\quad}_{N+1}}(-R, \mathbf{x}, R)} = \nu(\kappa) C(\kappa)^N \oint_{\vartheta(x_2, x_3)} du_2 \cdots \oint_{\vartheta(x_{2N}, R)} du_{N+1} \hat{f}_{\underbrace{\overbrace{\quad}_{N+1}}^a}(-R, \mathbf{x}, R; \mathbf{u}_1) \quad [\text{by (2.11)}]$$

<sup>23</sup>If necessary, we use the integration symbols  $\int_{x_{a_s}}^{x_{b_s}} du_s$  to indicate that the integration of the variable  $u_s$  is performed from  $x_{a_s}$  to  $x_{b_s}$  in the upper half-plane.

$$= \nu(\kappa) \hat{C}(\kappa)^N \int_{x_2}^{x_3} du_2 \cdots \int_{x_{2N}}^R du_{N+1} |\hat{f}(-R, \mathbf{x}, R; \dot{\mathbf{u}}_1)|. \quad [\text{by (C.5)}]$$

Multiplying this by  $(2R)^{2h(\kappa)}$  and taking  $R \rightarrow \infty$  gives

$$\lim_{R \rightarrow \infty} (2R)^{2h(\kappa)} \mathcal{F}_{\underbrace{\quad\quad\quad}_N}(-R, \mathbf{x}, R) = \nu(\kappa) \hat{C}(\kappa)^N \int_{x_2}^{x_3} dv_1 \cdots \int_{x_{2N}}^{+\infty} dv_N |f(\mathbf{x}; \mathbf{v})|,$$

using the definition (C.1). Hence, we obtain

$$\mathcal{F}_{\underbrace{\quad\quad\quad}_N}(\mathbf{x}) = \nu(\kappa) \hat{C}(\kappa)^N \int_{x_2}^{x_3} dv_1 \cdots \int_{x_{2N}}^{+\infty} dv_N |f(\mathbf{x}; \mathbf{v})|,$$

which proves (C.3). The other asserted identity (C.4) can be proven similarly.  $\square$

The following auxiliary Coulomb gas integral functions will be useful in Sections 3–4. They are integrals of the integrand function (C.1), which has the conjugate charge at  $x_1$ , and they differ from the functions in (2.9) in the sense that in (2.9), no integral contour surrounds the point  $x_1$  carrying the conjugate charge, while in the functions  $\hat{\mathcal{R}}_{\omega_s}$  below, an integral contour does surround the point  $x_1$ . We consider special link patterns  $\omega_s$  indexed by the pair  $2s - 1 \in \{1, 3, 5, \dots, 2N - 1\}$  of the last point  $2N$ ,

$$\begin{aligned} \omega_1 &:= \{\{1, 2N\}, \{2, 3\}, \{4, 5\}, \dots, \{2N - 2, 2N - 1\}\} = \underbrace{\quad\quad\quad}_N, \\ \omega_2 &:= \{\{1, 2\}, \{3, 2N\}, \{4, 5\}, \{6, 7\}, \dots, \{2N - 2, 2N - 1\}\}, \\ &\vdots \\ \omega_N &:= \{\{1, 2\}, \{3, 4\}, \dots, \{2N - 1, 2N\}\} = \underbrace{\quad\quad\quad}_N. \end{aligned}$$

Let the branch of  $\hat{f}_{\omega_s} = \hat{f}_{\omega_s}^a$  be chosen as in (2.11) with  $a = 1$ . For  $\kappa \in (0, 8) \setminus \{\frac{8}{m} : m \in \mathbb{Z}_{>0}\}$ , we define<sup>24</sup>

$$\begin{aligned} \hat{\mathcal{R}}_{\omega_1}(\mathbf{x}) &:= \frac{e^{-4\pi i/\kappa} C(\kappa)^{N-1}}{\nu(\kappa)^2 - 1} \oint_{\vartheta(x_1, x_{2N})} du_1 \oint_{\vartheta(x_4, x_5)} du_3 \cdots \oint_{\vartheta(x_{2N-2}, x_{2N-1})} du_N \hat{f}_{\omega_1}^a(\mathbf{x}; \dot{\mathbf{u}}_2), \\ \hat{\mathcal{R}}_{\omega_s}(\mathbf{x}) &:= -\frac{e^{-20\pi i/\kappa} C(\kappa)^{N-1}}{(\nu(\kappa)^2 - 1)^2} \oint_{\vartheta(x_1, x_2)} du_1 \cdots \oint_{\vartheta(x_{2s-3}, x_{2s-2})} du_{s-1} \oint_{\vartheta(x_1, x_{2N})} du_s \\ &\quad \times \oint_{\vartheta(x_{2s+2}, x_{2s+3})} du_{s+2} \cdots \oint_{\vartheta(x_{2N-2}, x_{2N-1})} du_N \hat{f}_{\omega_s}^a(\mathbf{x}; \dot{\mathbf{u}}_{s+1}), \quad 2 \leq s \leq N - 1, \\ \hat{\mathcal{R}}_{\omega_N}(\mathbf{x}) &:= -\frac{e^{-16\pi i/\kappa} C(\kappa)^{N-1}}{\nu(\kappa)^2 - 1} \oint_{\vartheta(x_1, x_2)} du_1 \oint_{\vartheta(x_3, x_4)} du_2 \cdots \oint_{\vartheta(x_{2N-3}, x_{2N-2})} du_{N-1} \hat{f}_{\omega_N}^a(\mathbf{x}; \dot{\mathbf{u}}_N), \end{aligned}$$

where the  $(s + 1)$ :st integral is omitted and for  $2 \leq s \leq N - 1$  the integration contours are given by  $\omega_s$  except for the  $s$ :th integral, which is a Pochhammer contour<sup>25</sup>  $\vartheta(x_1, x_{2N})$  around the points  $x_1, x_{2N}$ .

Now, because the integration contours for  $\hat{\mathcal{R}}_{\omega_s}$  are closed in a suitable homology and satisfy the charge neutrality (C.2) [DFMS97, Dub06, FK15c], we readily conclude that they provide solutions to the BPZ PDEs and are Möbius covariant, so they lie in the solution space  $\mathcal{S}_N$  defined in (2.21).

**Lemma C.3.** *The functions  $\{\hat{\mathcal{R}}_{\omega_s} : 1 \leq s \leq N\}$  satisfy the properties (PDE), (COV), and (PLB).*

*Proof.* (PDE) & (COV) are well-known [Dub06, Lemma 2]. The bound (PLB) is clear.  $\square$

We write  $\int_{x_{2N}}^{x_1} du = \int_{-\infty}^{x_1} du + \int_{x_{2N}}^{+\infty} du$  (cf. Remark C.1).

<sup>24</sup>When  $\kappa = 6$ , we have  $\nu(\kappa)^2 = 1$ , but also, the integrals give zero by (C.6, C.7), and  $\hat{\mathcal{R}}_{\omega_s}$  are well-defined by continuity.

<sup>25</sup>The integrals involving  $x_1$  are convergent, for the integrand  $\hat{f}$  in (C.1) has the conjugate charge at  $x_1$ , which implies that the divergence as both integration variables  $u_1, u_s$  are  $\varepsilon$ -close to  $x_1$  is of order  $\varepsilon^{1+2(12/\kappa-2)} \rightarrow 0$  as  $\varepsilon \rightarrow 0$  for all  $\kappa < 8$ .

**Lemma C.4.** Fix  $\kappa \in (4, 8)$ . We have

$$\begin{aligned}\hat{\mathcal{R}}_{\omega_1}(\mathbf{x}) &= \hat{C}(\kappa)^{N-1} \int_{x_{2N}}^{x_1} du_1 \int_{x_4}^{x_5} du_3 \cdots \int_{x_{2N-2}}^{x_{2N-1}} du_N |\hat{f}(\mathbf{x}; \dot{\mathbf{u}}_2)|, \\ \hat{\mathcal{R}}_{\omega_s}(\mathbf{x}) &= \hat{C}(\kappa)^{N-1} \int_{x_1}^{x_2} du_1 \cdots \int_{x_{2s-3}}^{x_{2s-2}} du_{s-1} \int_{x_{2N}}^{x_1} du_s \\ &\quad \times \int_{x_{2s+2}}^{x_{2s+3}} du_{s+2} \cdots \int_{x_{2N-2}}^{x_{2N-1}} du_N |\hat{f}(\mathbf{x}; \dot{\mathbf{u}}_{s+1})|, \\ \hat{\mathcal{R}}_{\omega_N}(\mathbf{x}) &= \hat{C}(\kappa)^{N-1} \int_{x_1}^{x_2} du_1 \cdots \int_{x_{2N-3}}^{x_{2N-2}} du_{N-1} |\hat{f}(\mathbf{x}; \dot{\mathbf{u}}_N)|, \quad \mathbf{x} \in \mathfrak{X}_{2N},\end{aligned}$$

In particular, we have  $\hat{\mathcal{R}}_{\omega_s}(\mathbf{x}) > 0$  for all  $s \in \{1, 2, \dots, N\}$  and  $\mathbf{x} \in \mathfrak{X}_{2N}$ .

*Proof.* This follows using (C.5) for the integrals not involving the point  $x_1$ , and the analogous formulas

$$\oint_{\vartheta(x_1, x_2)} du_1 \hat{f}(\mathbf{x}; \mathbf{u}) = -4 \sin^2(4\pi/\kappa) (\nu(\kappa)^2 - 1) e^{16\pi i/\kappa} \int_{x_1}^{x_2} du_1 \hat{f}(\mathbf{x}; \mathbf{u}), \quad (\text{C.6})$$

$$\oint_{\vartheta(x_1, x_{2N})} du_1 \hat{f}(\mathbf{x}; \mathbf{u}) = 4 \sin^2(4\pi/\kappa) (\nu(\kappa)^2 - 1) e^{-8\pi i/\kappa} \int_{x_{2N}}^{x_1} du_1 \hat{f}(\mathbf{x}; \mathbf{u}), \quad (\text{C.7})$$

for the integrals involving the point  $x_1$  obtained by taking  $\epsilon \rightarrow 0$  in Lemma 2.8.  $\square$

**Lemma C.5.** The functions  $\{\hat{\mathcal{R}}_{\omega_s} : 1 \leq s \leq N\}$  satisfy the asymptotics property

$$\lim_{x_1, x_2 \rightarrow \xi} \frac{\hat{\mathcal{R}}_{\omega_s}(\mathbf{x})}{(x_2 - x_1)^{-2h(\kappa)}} = \begin{cases} \frac{1}{\nu(\kappa)} \mathcal{F}_{\text{arc}_{N-1}}(\ddot{\mathbf{x}}_1), & \text{if } s = 1, \\ 0, & \text{if } s \neq 1, \end{cases} \quad (\text{C.8})$$

for any  $\xi < x_3 < x_4 < \cdots < x_{2N}$ .

*Proof.* First, we treat the case  $s = 1$ . Thanks to Lemma C.4, a direct computation shows that

$$\begin{aligned}\hat{\mathcal{R}}(\ddot{\mathbf{x}}_1; \xi) &:= \lim_{x_1, x_2 \rightarrow \xi} \frac{\hat{\mathcal{R}}_{\omega_1}(\mathbf{x})}{(x_2 - x_1)^{-2h(\kappa)}} \\ &= \hat{C}(\kappa)^{N-1} \prod_{j=3}^{2N} (x_j - \xi)^{1-4/\kappa} \int_{x_{2N}}^{\xi} du_1 \int_{x_4}^{x_5} du_3 \cdots \int_{x_{2N-2}}^{x_{2N-1}} du_N \prod_{\substack{1 < t \leq N \\ t \neq 2}} |u_t - \xi|^{8/\kappa-2} |f(\ddot{\mathbf{x}}_1; \dot{\mathbf{u}}_2)|.\end{aligned}$$

If  $\varphi$  is a Möbius map of the upper half-plane  $\mathbb{H}$  with  $\varphi(\xi) < \varphi(x_3) < \cdots < \varphi(x_{2N})$ , then the Möbius covariance of  $\hat{\mathcal{R}}_{\omega_1}$  in Lemma C.3 and the simple identity  $\frac{\varphi(x_2) - \varphi(x_1)}{x_2 - x_1} = \sqrt{\varphi'(x_1)} \sqrt{\varphi'(x_2)}$  imply that

$$\begin{aligned}\hat{\mathcal{R}}(\varphi(\ddot{\mathbf{x}}_1); \varphi(\xi)) &= \lim_{x_1, x_2 \rightarrow \xi} \frac{\hat{\mathcal{R}}_{\omega_1}(\varphi(x_1), \varphi(x_2), \varphi(\ddot{\mathbf{x}}_1))}{(\varphi(x_2) - \varphi(x_1))^{-2h(\kappa)}} = \lim_{x_1, x_2 \rightarrow \xi} \frac{\hat{\mathcal{R}}_{\omega_1}(\mathbf{x})}{(x_2 - x_1)^{-2h(\kappa)}} \times \prod_{j=3}^{2N} \varphi'(x_j)^{-h} \\ &= \hat{\mathcal{R}}_{\omega_1}(\ddot{\mathbf{x}}_1, \xi) \times \prod_{j=3}^{2N} \varphi'(x_j)^{-h},\end{aligned} \quad (\text{C.9})$$

where  $\varphi(\ddot{\mathbf{x}}_1) = (\varphi(x_3), \dots, \varphi(x_{2N}))$ . For large enough  $R > 0$ , take  $\varphi_R$  to be the Möbius transformation of  $\mathbb{H}$  such that

$$\varphi_R(\xi) = -R, \quad \varphi_R(x_3) = x_3, \quad \text{and} \quad \varphi_R(x_{2N}) = x_{2N}.$$

Denote  $\varphi_R(\tilde{\mathbf{x}}_1) := (\varphi_R(x_3), \dots, \varphi_R(x_{2N}))$ . Then, we have

$$\frac{\hat{\mathcal{R}}(\tilde{\mathbf{x}}_1; \xi)}{\mathcal{F}_{\overbrace{\text{---}}_{N-1}}(\tilde{\mathbf{x}}_1)} = \lim_{R \rightarrow \infty} \frac{\hat{\mathcal{R}}(\varphi_R(\tilde{\mathbf{x}}_1); \varphi_R(\xi))}{\mathcal{F}_{\overbrace{\text{---}}_{N-1}}(\varphi_R(\tilde{\mathbf{x}}_1))} = \frac{1}{\nu(\kappa)},$$

thanks to the Möbius covariance of  $\mathcal{F}_{\overbrace{\text{---}}_{N-1}}$  from Corollary 2.3, the Möbius covariance of  $\hat{\mathcal{R}}$  in (C.9), and the line-integral formula (C.3) for  $\mathcal{F}_{\overbrace{\text{---}}_{N-1}}$  from Lemma C.2. This shows (C.8) when  $s = 1$ .

Second, in the case where  $s \neq 1$ , using Lemma C.4 and the change of variables  $u = \frac{u_1 - x_1}{x_2 - x_1}$  gives

$$\begin{aligned} \frac{\hat{\mathcal{R}}_{\omega_s}(\mathbf{x})}{(x_2 - x_1)^{-2h(\kappa)}} &= (x_2 - x_1)^{8/\kappa - 1} \int_0^1 du \cdots \int_{x_{2s-3}}^{x_{2s-2}} du_{s-1} \int_{x_{2N}}^{x_1} du_s \\ &\quad \times \int_{x_{2s+2}}^{x_{2s+3}} du_{s+2} \cdots \int_{x_{2N-2}}^{x_{2N-1}} du_N |u|^{12/\kappa - 2} |u - 1|^{-4/\kappa} |f(\tilde{\mathbf{x}}_1; \tilde{\mathbf{u}})| \hat{g}_s(\mathbf{x}; \tilde{\mathbf{u}}, u), \\ &\xrightarrow{x_1, x_2 \rightarrow \xi} 0, \end{aligned}$$

where  $\tilde{\mathbf{u}} := (u_2, u_3, \dots, u_s, u_{s+2}, \dots, u_N)$  and the function  $\hat{g}_s(\mathbf{x}; \tilde{\mathbf{u}}, u)$  is bounded as  $x_2, x_1 \rightarrow \xi$ .  $\square$

## D Examples with $N = 1$ and $N = 2$

In this appendix, we investigate the Coulomb gas integrals for  $N = 1$  and  $N = 2$ . We denote

$$\overbrace{\text{---}} = \{\{1, 2\}\}, \quad \overbrace{\text{---}} \overbrace{\text{---}} = \{\{1, 2\}, \{3, 4\}\}, \quad \overbrace{\text{---}} \overbrace{\text{---}} \overbrace{\text{---}} = \{\{1, 4\}, \{2, 3\}\}.$$

It is straightforward to calculate  $\mathcal{F}_{\overbrace{\text{---}}^{(\kappa)}}$  for  $N = 1$  (Lemma D.1). We derive asymptotics properties of the Coulomb gas integrals with  $N = 2$  in Section D.1, and investigate their positivity in the case  $N = 2$  in Section D.2. Lastly, in Section D.3 we consider the Fröbenius series of the Coulomb gas integrals and the pure partition functions in the case  $N = 2$ .

**Lemma D.1.** *For  $\kappa \in (0, 8) \setminus \{\frac{8}{m} : m \in \mathbb{Z}_{>0}\}$ , we have*

$$\mathcal{H}_{\overbrace{\text{---}}^{(\kappa)}}(x_1, x_2) = \frac{\nu(\kappa)}{\mathbf{C}(\kappa)} (x_2 - x_1)^{-2h(\kappa)}, \quad (x_1, x_2) \in \mathfrak{X}_2. \quad (\text{D.1})$$

Consequently, for  $\kappa \in (0, 8)$ , we have

$$\mathcal{F}_{\overbrace{\text{---}}^{(\kappa)}}(x_1, x_2) = \nu(\kappa) (x_2 - x_1)^{-2h(\kappa)}, \quad (x_1, x_2) \in \mathfrak{X}_2.$$

*Proof.* This follows by a change of variables, Eq. (B.1) from Lemma B.1, and continuity in  $(0, 8) \ni \kappa$ .  $\square$

### D.1 Asymptotics for $N = 2$

This section contains an example computation for the asymptotics in Proposition 2.12 when  $N = 2$ .

**Proposition D.2.** *Fix  $\kappa \in (0, 8) \setminus \{\frac{8}{m} : m \in \mathbb{Z}_{>0}\}$ . The following asymptotics hold for  $(x_1, x_2, x_3, x_4) \in \mathfrak{X}_4$ .*

- For any  $\xi \in (-\infty, x_3)$ , we have

$$\lim_{x_1, x_2 \rightarrow \xi} \frac{\mathcal{H}_{\overbrace{\text{---}} \overbrace{\text{---}}^{(\kappa)}}(x_1, x_2, x_3, x_4)}{(x_2 - x_1)^{-2h(\kappa)}} = \frac{\nu(\kappa)}{\mathbf{C}(\kappa)} \mathcal{H}_{\overbrace{\text{---}}^{(\kappa)}}(x_3, x_4); \quad (\text{D.2})$$

$$\lim_{x_1, x_2 \rightarrow \xi} \frac{\mathcal{H}_{\overbrace{\text{---}} \overbrace{\text{---}} \overbrace{\text{---}}^{(\kappa)}}(x_1, x_2, x_3, x_4)}{(x_2 - x_1)^{-2h(\kappa)}} = \frac{1}{\mathbf{C}(\kappa)} \mathcal{H}_{\overbrace{\text{---}}^{(\kappa)}}(x_3, x_4). \quad (\text{D.3})$$



- For any  $\xi \in (x_1, x_4)$ , we have

$$\lim_{x_2, x_3 \rightarrow \xi} \frac{\mathcal{H}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa)}}(x_1, x_2, x_3, x_4)}{(x_3 - x_2)^{-2h(\kappa)}} = \frac{1}{C(\kappa)} \mathcal{H}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa)}}(x_1, x_4); \quad (\text{D.4})$$

$$\lim_{x_2, x_3 \rightarrow \xi} \frac{\mathcal{H}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa)}}(x_1, x_2, x_3, x_4)}{(x_3 - x_2)^{-2h(\kappa)}} = \frac{\nu(\kappa)}{C(\kappa)} \mathcal{H}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa)}}(x_1, x_4). \quad (\text{D.5})$$

- For any  $\xi \in (x_2, \infty)$ , we have

$$\lim_{x_3, x_4 \rightarrow \xi} \frac{\mathcal{H}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa)}}(x_1, x_2, x_3, x_4)}{(x_4 - x_3)^{-2h(\kappa)}} = \frac{\nu(\kappa)}{C(\kappa)} \mathcal{H}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa)}}(x_1, x_2); \quad (\text{D.6})$$

$$\lim_{x_3, x_4 \rightarrow \xi} \frac{\mathcal{H}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa)}}(x_1, x_2, x_3, x_4)}{(x_4 - x_3)^{-2h(\kappa)}} = \frac{1}{C(\kappa)} \mathcal{H}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa)}}(x_1, x_2). \quad (\text{D.7})$$

To prove Proposition D.2, we will use the following expressions for  $\mathcal{H}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa)}}$  in Lemma D.3.

**Lemma D.3.** Fix  $N = 2$ , and  $\underbrace{\quad}_{\underbrace{\quad}} = \{\{1, 2\}, \{3, 4\}\}$ , and  $\kappa \in (0, 8) \setminus \{\frac{8}{m} : m \in \mathbb{Z}_{>0}\}$ . We have

$$\begin{aligned} \mathcal{H}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa)}}(x_1, x_2, x_3, x_4) &= \frac{\nu(\kappa)}{C(\kappa)} (x_3 - x_1)^{-2/\kappa} (x_3 - x_2)^{2/\kappa} (x_4 - x_1)^{2/\kappa} (x_4 - x_3)^{1-6/\kappa} (x_4 - x_2)^{-2/\kappa} \\ &\quad \times (x_2 - x_1)^{1-6/\kappa} \oint_{\vartheta(0,1)} (w(1-w)(1-zw))^{-4/\kappa} dw \end{aligned} \quad (\text{D.8})$$

$$\begin{aligned} &= \frac{\nu(\kappa)}{C(\kappa)} (x_3 - x_1)^{-2/\kappa} (x_3 - x_2)^{2/\kappa} (x_4 - x_1)^{2/\kappa} (x_4 - x_3)^{1-6/\kappa} (x_4 - x_2)^{-2/\kappa} \\ &\quad \times (x_2 - x_1)^{1-6/\kappa} (1-z)^{1-8/\kappa} \int_{\vartheta(0, \frac{1}{1-z})} ((1+zv)v(1+(z-1)v))^{-4/\kappa} dv, \end{aligned} \quad (\text{D.9})$$

where  $z = \text{CR}(x_1, x_2, x_3, x_4)$  is the cross-ratio (2.25), the branch of  $w \mapsto (w(1-w)(1-zw))^{-4/\kappa}$  is chosen to be real and positive when  $0 < \text{Re}(w) < 1$ , and the branch of  $v \mapsto ((1+zv)v(1+(z-1)v))^{-4/\kappa}$  is chosen to be real and positive when  $0 < \text{Re}(v) < \frac{1}{1-z}$ .

*Proof.* First, we prove (D.8, D.9) for  $\kappa \in (4, 8)$ . According to Proposition 2.5, we have

$$\begin{aligned} \mathcal{H}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa)}}(x_1, x_2, x_3, x_4) &= \frac{\nu(\kappa)}{C(\kappa)} ((x_3 - x_1)(x_2 - x_1)(x_3 - x_2))^{2/\kappa} ((x_4 - x_1)(x_4 - x_3)(x_4 - x_2))^{1-6/\kappa} \\ &\quad \times \oint_{\vartheta(x_1, x_2)} (u_1 - x_1)^{-4/\kappa} (u_1 - x_2)^{-4/\kappa} (u_1 - x_3)^{-4/\kappa} (u_1 - x_4)^{12/\kappa-2} du_1, \end{aligned}$$

where the branch of  $(u_1 - x_1)^{-4/\kappa} (u_1 - x_2)^{-4/\kappa} (u_1 - x_3)^{-4/\kappa} (u_1 - x_4)^{12/\kappa-2}$  is chosen to be real and positive when  $x_1 < \text{Re}(u_1) < x_2$ . Thus, for  $\kappa \in (4, 8)$ , we have

$$\begin{aligned} \mathcal{H}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa)}}(x_1, x_2, x_3, x_4) &= \frac{\nu(\kappa)}{C(\kappa)} ((x_3 - x_1)(x_2 - x_1)(x_3 - x_2))^{2/\kappa} ((x_4 - x_1)(x_4 - x_3)(x_4 - x_2))^{1-6/\kappa} \\ &\quad \times 4 \sin^2(4\pi/\kappa) \int_{x_1}^{x_2} |(u_1 - x_1)(u_1 - x_2)(u_1 - x_3)|^{-4/\kappa} |u_1 - x_4|^{12/\kappa-2} du_1. \end{aligned}$$

We then use the change of variables  $u_1 = x_4 - (x_4 - x_1)(1 + w \frac{x_2 - x_1}{x_4 - x_2})^{-1}$  to obtain

$$\mathcal{H}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa)}}(x_1, x_2, x_3, x_4) = \frac{\nu(\kappa)}{C(\kappa)} (x_3 - x_1)^{-2/\kappa} (x_3 - x_2)^{2/\kappa} (x_4 - x_1)^{2/\kappa} (x_4 - x_3)^{1-6/\kappa} (x_4 - x_2)^{-2/\kappa}$$

$$\begin{aligned}
& \times (x_2 - x_1)^{1-6/\kappa} 4 \sin^2(4\pi/\kappa) \int_0^1 |w(1-w)(1-zw)|^{-4/\kappa} dw \quad (\text{D.10}) \\
& = \frac{\nu(\kappa)}{C(\kappa)} (x_3 - x_1)^{-2/\kappa} (x_3 - x_2)^{2/\kappa} (x_4 - x_1)^{2/\kappa} (x_4 - x_3)^{1-6/\kappa} (x_4 - x_2)^{-2/\kappa} \\
& \quad \times (x_2 - x_1)^{1-6/\kappa} \oint_{\vartheta(0,1)} (w(1-w)(1-zw))^{-4/\kappa} dw,
\end{aligned}$$

which gives (D.8) for  $\kappa \in (4, 8)$ . From (D.10), we use the change of variables  $v(1-z) = 1-w$  to obtain

$$\begin{aligned}
\mathcal{H}_{\underline{\circlearrowleft}}^{(\kappa)}(x_1, x_2, x_3, x_4) & = \frac{\nu(\kappa)}{C(\kappa)} (x_3 - x_1)^{-2/\kappa} (x_3 - x_2)^{2/\kappa} (x_4 - x_1)^{2/\kappa} (x_4 - x_3)^{1-6/\kappa} (x_4 - x_2)^{-2/\kappa} \\
& \quad \times (x_2 - x_1)^{1-6/\kappa} (1-z)^{1-8/\kappa} 4 \sin^2(4\pi/\kappa) \int_0^{\frac{1}{1-z}} |(1+zv)v(1+(z-1)v)|^{-4/\kappa} dv \\
& = \frac{\nu(\kappa)}{C(\kappa)} (x_3 - x_1)^{-2/\kappa} (x_3 - x_2)^{2/\kappa} (x_4 - x_1)^{2/\kappa} (x_4 - x_3)^{1-6/\kappa} (x_4 - x_2)^{-2/\kappa} \\
& \quad \times (x_2 - x_1)^{1-6/\kappa} (1-z)^{1-8/\kappa} \oint_{\vartheta(0, \frac{1}{1-z})} ((1+zv)v(1+(z-1)v))^{-4/\kappa} dv,
\end{aligned}$$

We just proved that (D.8, D.9) hold for  $\kappa \in (4, 8)$ , and (D.8, D.9) for  $\kappa \in (0, 8) \setminus \{\frac{8}{m} : m \in \mathbb{Z}_{>0}\}$  follow by analytic continuation.  $\square$

Now, we are ready to prove Proposition D.2.

*Proof of Proposition D.2.* First, we show (D.2). By (D.8) in Lemma D.3, for any  $\xi \in (-\infty, x_3)$ , we have

$$\begin{aligned}
\lim_{x_1, x_2 \rightarrow \xi} \frac{\mathcal{H}_{\underline{\circlearrowleft}}^{(\kappa)}(x_1, x_2, x_3, x_4)}{(x_2 - x_1)^{-2h(\kappa)}} & = \frac{\nu(\kappa)}{C(\kappa)} (x_4 - x_3)^{-2h(\kappa)} \lim_{z \rightarrow 0^+} \oint_{\vartheta(0,1)} (w(1-w)(1-zw))^{-4/\kappa} dw \\
& = \frac{\nu(\kappa)}{C(\kappa)} (x_4 - x_3)^{-2h(\kappa)} \oint_{\vartheta(0,1)} (w(1-w))^{-4/\kappa} dw \\
& = \left(\frac{\nu(\kappa)}{C(\kappa)}\right)^2 (x_4 - x_3)^{-2h(\kappa)} = \frac{\nu(\kappa)}{C(\kappa)} \mathcal{H}_{\underline{\circlearrowleft}}^{(\kappa)}(x_3, x_4), \quad [\text{by (B.1 \& D.1)}]
\end{aligned}$$

Second, we show (D.4). By (D.9) in Lemma D.3, for any  $\xi \in (x_1, x_4)$ , we have

$$\begin{aligned}
\lim_{x_2, x_3 \rightarrow \xi} \frac{\mathcal{H}_{\underline{\circlearrowleft}}^{(\kappa)}(x_1, x_2, x_3, x_4)}{(x_3 - x_2)^{-2h(\kappa)}} & = \frac{\nu(\kappa)}{C(\kappa)} (x_4 - x_3)^{-2h(\kappa)} \lim_{z \rightarrow 1^-} \oint_{\vartheta(0, \frac{1}{1-z})} ((1+zv)v(1+(z-1)v))^{-4/\kappa} dv \\
& = \frac{\nu(\kappa)}{C(\kappa)} (x_4 - x_3)^{-2h(\kappa)} (1 - e^{8\pi i/\kappa}) \int_{\varrho(0, \infty)} (1+v)^{-4/\kappa} v^{-4/\kappa} dv \\
& = \frac{\nu(\kappa)}{C(\kappa)} (x_4 - x_3)^{-2h(\kappa)} (1 - e^{8\pi i/\kappa}) (1 - e^{-8\pi i/\kappa}) \frac{\Gamma(1-4/\kappa)^2}{\Gamma(2-8/\kappa)} \frac{1}{\nu(\kappa)} \\
& = \frac{1}{C(\kappa)} \mathcal{H}_{\underline{\circlearrowleft}}^{(\kappa)}(x_1, x_4), \quad [\text{by (B.1)}]
\end{aligned}$$

where we use the identity (B.4) from Lemma B.2 to evaluate the integral over  $\varrho(0, \infty)$ . Third, the proof of (D.6) is essentially the same as that of (D.2). Finally, (D.3, D.5, D.7) follow using (D.2, D.4, D.6) and Proposition 2.2.  $\square$

## D.2 Positivity for $N = 2$

We shall next investigate the signs of the two Coulomb gas integral functions  $\mathcal{F}_{\underline{\circlearrowleft}\underline{\circlearrowright}}^{(\kappa)}$  and  $\mathcal{F}_{\underline{\circlearrowright}\underline{\circlearrowleft}}^{(\kappa)}$  when  $\mathbf{x} = (x_1, x_2, x_3, x_4) \in \mathfrak{X}_4$ , using their relation with the pure partition functions. Interestingly enough, we observe that when  $\kappa < 8/3$ , these functions are not always positive (thus, they are not true  $\text{SLE}_\kappa$  partition functions in that case) — unlike the pure partition functions  $\mathcal{Z}_{\underline{\circlearrowleft}\underline{\circlearrowright}}^{(\kappa)}$  and  $\mathcal{Z}_{\underline{\circlearrowright}\underline{\circlearrowleft}}^{(\kappa)}$ .

**Proposition D.4.** *Fix  $\kappa \in (0, 8)$ . The following identities hold.*

$$\mathcal{F}_{\underline{\circlearrowleft}\underline{\circlearrowright}}^{(\kappa)}(\mathbf{x}) = \nu(\kappa)^2 \mathcal{Z}_{\underline{\circlearrowleft}\underline{\circlearrowright}}^{(\kappa)}(\mathbf{x}) + \nu(\kappa) \mathcal{Z}_{\underline{\circlearrowright}\underline{\circlearrowleft}}^{(\kappa)}(\mathbf{x}), \quad \mathbf{x} \in \mathfrak{X}_4; \quad (\text{D.11})$$

$$\mathcal{F}_{\underline{\circlearrowright}\underline{\circlearrowleft}}^{(\kappa)}(\mathbf{x}) = \nu(\kappa) \mathcal{Z}_{\underline{\circlearrowleft}\underline{\circlearrowright}}^{(\kappa)}(\mathbf{x}) + \nu(\kappa)^2 \mathcal{Z}_{\underline{\circlearrowright}\underline{\circlearrowleft}}^{(\kappa)}(\mathbf{x}), \quad \mathbf{x} \in \mathfrak{X}_4; \quad (\text{D.12})$$

where

$$\begin{aligned} \mathcal{Z}_{\underline{\circlearrowleft}\underline{\circlearrowright}}^{(\kappa)}(x_1, x_2, x_3, x_4) &= (x_2 - x_1)^{-2h(\kappa)} (x_4 - x_3)^{-2h(\kappa)} (1 - \chi)^{2/\kappa} \frac{{}_2F_1\left(\frac{4}{\kappa}, 1 - \frac{4}{\kappa}, \frac{8}{\kappa}; 1 - \chi\right)}{{}_2F_1\left(\frac{4}{\kappa}, 1 - \frac{4}{\kappa}, \frac{8}{\kappa}; 1\right)}, \\ \mathcal{Z}_{\underline{\circlearrowright}\underline{\circlearrowleft}}^{(\kappa)}(x_1, x_2, x_3, x_4) &= (x_4 - x_1)^{-2h(\kappa)} (x_3 - x_2)^{-2h(\kappa)} \chi^{2/\kappa} \frac{{}_2F_1\left(\frac{4}{\kappa}, 1 - \frac{4}{\kappa}, \frac{8}{\kappa}; \chi\right)}{{}_2F_1\left(\frac{4}{\kappa}, 1 - \frac{4}{\kappa}, \frac{8}{\kappa}; 1\right)}, \end{aligned} \quad (\text{D.13})$$

and  $\chi = \text{CR}(\mathbf{x}) = \frac{(x_2 - x_1)(x_4 - x_3)}{(x_3 - x_1)(x_4 - x_2)}$  is the cross-ratio (2.25).

Moreover, the positivity of  $\mathcal{F}_{\underline{\circlearrowleft}\underline{\circlearrowright}}^{(\kappa)}(\mathbf{x})$  and  $\mathcal{F}_{\underline{\circlearrowright}\underline{\circlearrowleft}}^{(\kappa)}(\mathbf{x})$  depends on  $\kappa$  and  $\mathbf{x}$  in the following manner:

(POS) When  $\nu(\kappa) > 0$ , i.e.,  $\kappa \in \bigcup_{m \in \mathbb{Z}_{>0}} \left(\frac{8}{4m-1}, \frac{8}{4m-3}\right)$ , we have  $\mathcal{F}_{\underline{\circlearrowleft}\underline{\circlearrowright}}^{(\kappa)}(\mathbf{x}) > 0$  and  $\mathcal{F}_{\underline{\circlearrowright}\underline{\circlearrowleft}}^{(\kappa)}(\mathbf{x}) > 0$  on  $\mathfrak{X}_4$ .

(NUL) When  $\nu(\kappa) = 0$ , i.e.,  $\kappa \in \left\{\frac{8}{2m+1} : m \in \mathbb{Z}_{>0}\right\}$ , we have  $\mathcal{F}_{\underline{\circlearrowleft}\underline{\circlearrowright}}^{(\kappa)}(\mathbf{x}) = \mathcal{F}_{\underline{\circlearrowright}\underline{\circlearrowleft}}^{(\kappa)}(\mathbf{x}) = 0$  on  $\mathfrak{X}_4$ .

(SGN) When  $\nu(\kappa) < 0$ , i.e.,  $\kappa \in \bigcup_{m \in \mathbb{Z}_{>0}} \left(\frac{8}{4m+1}, \frac{8}{4m-1}\right)$ , there exists  $z(\kappa) \in (0, 1)$  such that

$$\begin{cases} \mathcal{F}_{\underline{\circlearrowleft}\underline{\circlearrowright}}^{(\kappa)}(\mathbf{x}) > 0, & \text{if } \text{CR}(\mathbf{x}) < z(\kappa); \\ \mathcal{F}_{\underline{\circlearrowleft}\underline{\circlearrowright}}^{(\kappa)}(\mathbf{x}) = 0, & \text{if } \text{CR}(\mathbf{x}) = z(\kappa); \\ \mathcal{F}_{\underline{\circlearrowleft}\underline{\circlearrowright}}^{(\kappa)}(\mathbf{x}) < 0, & \text{if } \text{CR}(\mathbf{x}) > z(\kappa); \end{cases} \quad \begin{cases} \mathcal{F}_{\underline{\circlearrowright}\underline{\circlearrowleft}}^{(\kappa)}(\mathbf{x}) > 0, & \text{if } \text{CR}(\mathbf{x}) > 1 - z(\kappa); \\ \mathcal{F}_{\underline{\circlearrowright}\underline{\circlearrowleft}}^{(\kappa)}(\mathbf{x}) = 0, & \text{if } \text{CR}(\mathbf{x}) = 1 - z(\kappa); \\ \mathcal{F}_{\underline{\circlearrowright}\underline{\circlearrowleft}}^{(\kappa)}(\mathbf{x}) < 0, & \text{if } \text{CR}(\mathbf{x}) < 1 - z(\kappa). \end{cases} \quad (\text{D.14})$$

Moreover, we have

$$\limsup_{\kappa \rightarrow 0} z(\kappa) = 1/2 \quad \text{and} \quad \liminf_{\kappa \rightarrow 0} z(\kappa) = 0. \quad (\text{D.15})$$

*Proof of Proposition D.4.* By continuity in  $\kappa \in (0, 8)$ , it suffices to check (D.11, D.12) when  $\kappa \neq 8/m$  for  $m \in \mathbb{Z}_{>0}$ . From Lemma D.1 and Proposition D.2, we see that both sides of (D.11, D.12) satisfy (PDE), (COV), (PLB), and ( $\mathcal{F}$ -ASY) respectively, and Lemma 2.11 guarantees that they are equal.

Next, let us consider the sign of  $\mathcal{F}_{\underline{\circlearrowleft}\underline{\circlearrowright}}^{(\kappa)}(\mathbf{x})$ . The cases  $\nu(\kappa) > 0$  and  $\nu(\kappa) = 0$  are clear from (D.11). It remains to investigate the case when  $\nu(\kappa) < 0$  and to prove (D.14) and (D.15). Suppose  $\nu(\kappa) < 0$ . Then, we have  $\mathcal{F}_{\underline{\circlearrowleft}\underline{\circlearrowright}}^{(\kappa)}(\mathbf{x}) > 0$  if and only if  $-\nu(\kappa) > \mathcal{Z}_{\underline{\circlearrowright}\underline{\circlearrowleft}}^{(\kappa)}(\mathbf{x})/\mathcal{Z}_{\underline{\circlearrowleft}\underline{\circlearrowright}}^{(\kappa)}(\mathbf{x})$ . Using (D.13), we see that

$$\mathcal{F}_{\underline{\circlearrowleft}\underline{\circlearrowright}}^{(\kappa)}(\mathbf{x}) > 0, \quad \text{if and only if} \quad 2 \cos(4\pi/\kappa) > \frac{\chi^{8/\kappa-1} {}_2F_1\left(\frac{4}{\kappa}, 1 - \frac{4}{\kappa}, \frac{8}{\kappa}; \chi\right)}{(1 - \chi)^{8/\kappa-1} {}_2F_1\left(\frac{4}{\kappa}, 1 - \frac{4}{\kappa}, \frac{8}{\kappa}; 1 - \chi\right)}, \quad (\text{D.16})$$

where  $\chi = \text{CR}(\mathbf{x})$  is the cross-ratio (2.25). Using the identity [AS64, Eq. 15.3.3],

$${}_2F_1\left(\frac{4}{\kappa}, 1 - \frac{4}{\kappa}, \frac{8}{\kappa}; z\right) = (1 - z)^{8/\kappa - 1} {}_2F_1\left(\frac{4}{\kappa}, \frac{12}{\kappa} - 1, \frac{8}{\kappa}; z\right),$$

we obtain

$$\mathcal{F}_{\underbrace{\quad}_{\quad}}^{(\kappa)}(\mathbf{x}) > 0, \quad \text{if and only if} \quad 2 \cos(4\pi/\kappa) > \frac{{}_2F_1\left(\frac{4}{\kappa}, \frac{12}{\kappa} - 1, \frac{8}{\kappa}; \chi\right)}{{}_2F_1\left(\frac{4}{\kappa}, \frac{12}{\kappa} - 1, \frac{8}{\kappa}; 1 - \chi\right)} =: G(\chi). \quad (\text{D.17})$$

Next note that  $z \mapsto {}_2F_1\left(\frac{4}{\kappa}, \frac{12}{\kappa} - 1, \frac{8}{\kappa}; z\right)$  is increasing for  $z \in (0, 1)$  and we have

$${}_2F_1\left(\frac{4}{\kappa}, \frac{12}{\kappa} - 1, \frac{8}{\kappa}; 0\right) = 1 \quad \text{and} \quad \lim_{z \rightarrow 1^-} {}_2F_1\left(\frac{4}{\kappa}, \frac{12}{\kappa} - 1, \frac{8}{\kappa}; z\right) = \infty.$$

Thus  $z \mapsto G(z)$  is also increasing, and  $\lim_{z \rightarrow 0^+} G(z) = 0$  and  $\lim_{z \rightarrow 1^-} G(z) = \infty$ . Therefore, there exists a unique point  $z(\kappa) \in (0, 1)$  such that

$$G(z(\kappa)) = 2 \cos(4\pi/\kappa). \quad (\text{D.18})$$

Combining this with (D.17), we obtain the conclusion (SGN) for  $\mathcal{F}_{\underbrace{\quad}_{\quad}}^{(\kappa)}$  in (D.14). The proof for  $\mathcal{F}_{\overbrace{\quad}^{\quad}}^{(\kappa)}$  in (D.14) is similar.

It remains to prove (D.15). Note that  $\kappa \mapsto z(\kappa)$  is a continuous function and we may extend its definition to  $\kappa \in (0, 8/3]$  by setting  $z(\kappa) = 0$  when  $\nu(\kappa) \geq 0$ , and such an extension  $\kappa \in (0, 8/3] \mapsto z(\kappa)$  is still continuous. Because  $G(0) = 0$  and  $G(1/2) = 1$ , we have

$$z\left(\frac{8}{2m+1}\right) = 0, \quad z\left(\frac{12}{6m \pm 1}\right) = 1/2, \quad \text{for } m \in \mathbb{Z}_{>0}. \quad (\text{D.19})$$

Let us consider the asymptotics of  $G(z)$  as  $\kappa \rightarrow 0$ . From direct analysis of the hypergeometric function  ${}_2F_1$  (see also [ABKM20, Section 7.3]), we obtain

$$U(z) := \lim_{\kappa \rightarrow 0} \kappa \log G(z) = 4 \log \left( \frac{(1 - z + z^2)^{3/2} + (1 + z)(z - 1/2)(2 - z)}{(1 - z + z^2)^{3/2} - (1 + z)(z - 1/2)(2 - z)} \right).$$

For any  $\epsilon > 0$ , we have  $U(\frac{1}{2} + \epsilon) > 0$ . Thus  $G(\frac{1}{2} + \epsilon) \geq 2$  when  $\kappa$  is small enough. Therefore,  $z(\kappa) \leq \frac{1}{2} + \epsilon$  when  $\kappa$  is small enough. In other words, we have  $\limsup_{\kappa \rightarrow 0} z(\kappa) \leq \frac{1}{2} + \epsilon$ . Sending  $\epsilon \rightarrow 0$ , we obtain

$$\limsup_{\kappa \rightarrow 0} z(\kappa) \leq \frac{1}{2}.$$

Combining this with (D.19), we obtain (D.15) and complete the proof.  $\square$

### D.3 Fröbenius series for $N = 2$

Using the explicit formulas (D.13) in terms of the hypergeometric function  ${}_2F_1$ , we derive in this section the Fröbenius series of the pure partition functions  $\mathcal{Z}_{\underbrace{\quad}_{\quad}}^{(\kappa)}$  (Lemma D.5) and  $\mathcal{Z}_{\overbrace{\quad}^{\quad}}^{(\kappa)}$  (Lemma D.6), and from this, also obtain the Fröbenius series of the Coulomb gas integral functions  $\mathcal{F}_{\underbrace{\quad}_{\quad}}^{(\kappa)}$  and  $\mathcal{F}_{\overbrace{\quad}^{\quad}}^{(\kappa)}$  (Corollary D.7) for  $\kappa \in (8/3, 8)$ . We also address the case where  $\kappa = 8/3$  in Lemma D.8 — it involves a logarithmic correction to the asymptotics, which is a special case of Proposition 1.17.

**Lemma D.5.** *For  $\kappa \in (0, 8)$  and  $\xi < x_3 < x_4$ , we have*

$$\mathcal{Z}_{\underbrace{\quad}_{\quad}}^{(\kappa)}(x_1, x_2, x_3, x_4) = (x_2 - x_1)^{2/\kappa} \mathcal{Z}_{\underbrace{\quad}_{\quad}}^{(\kappa)}(\xi, x_3, x_4) + o((x_2 - x_1)^{2/\kappa}), \quad \text{as } x_1, x_2 \rightarrow \xi. \quad (\text{D.20})$$

where

$$\mathcal{Z}_{\underbrace{\quad}_{\quad}}^{(\kappa)}(\xi, x_3, x_4) := \frac{1}{{}_2F_1\left(\frac{4}{\kappa}, 1 - \frac{4}{\kappa}, \frac{8}{\kappa}; 1\right)} \frac{(x_4 - x_3)^{2/\kappa}}{(x_3 - \xi)^{8/\kappa - 1} (x_4 - \xi)^{8/\kappa - 1}}. \quad (\text{D.21})$$

*Proof.* Denote by  $\chi = \text{CR}(x_1, x_2, x_3, x_4)$  the cross-ratio (2.25), the asserted (D.20) follows from (D.13):

$$\begin{aligned} \frac{\mathcal{Z}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa)}}(x_1, x_2, x_3, x_4)}{(x_3 - x_1)^{-2h(\kappa)}(x_4 - x_2)^{-2h(\kappa)}} &= \chi^{2/\kappa} (1 - \chi)^{-2h(\kappa)} \frac{{}_2F_1\left(\frac{4}{\kappa}, 1 - \frac{4}{\kappa}, \frac{8}{\kappa}; \chi\right)}{{}_2F_1\left(\frac{4}{\kappa}, 1 - \frac{4}{\kappa}, \frac{8}{\kappa}; 1\right)} \\ &= \chi^{2/\kappa} \frac{(1 + O(\chi))}{{}_2F_1\left(\frac{4}{\kappa}, 1 - \frac{4}{\kappa}, \frac{8}{\kappa}; 1\right)}, \quad \text{as } \chi \rightarrow 0. \end{aligned}$$

□

**Lemma D.6.** For  $\kappa \in (8/3, 8)$  and  $\xi < x_3 < x_4$ , we have

$$\begin{aligned} \mathcal{Z}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa)}}(x_1, x_2, x_3, x_4) &= (x_2 - x_1)^{-2h(\kappa)} \mathcal{Z}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa)}}(x_3, x_4) - \frac{1}{\nu(\kappa)} (x_2 - x_1)^{2/\kappa} \mathcal{Z}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa)}}(\xi, x_3, x_4) \\ &\quad + o((x_2 - x_1)^{2/\kappa}), \quad \text{as } x_1, x_2 \rightarrow \xi. \end{aligned} \tag{D.22}$$

*Proof.* In the case where  $\kappa = 4$ , the expansion (D.22) is immediate:

$$\mathcal{Z}_{\underbrace{\quad}_{\underbrace{\quad}}^{(4)}}(x_1, x_2, x_3, x_4) = (x_2 - x_1)^{-1/2} (x_4 - x_3)^{-1/2} (1 - \chi)^{1/2},$$

where  $\chi = \text{CR}(x_1, x_2, x_3, x_4)$ . Hence, in the rest of the proof, we assume  $\kappa \in (8/3, 4) \cup (4, 8)$ . From the explicit formula (D.13), we obtain

$$\frac{\mathcal{Z}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa)}}(x_1, x_2, x_3, x_4)}{(x_3 - x_1)^{-2h(\kappa)}(x_4 - x_2)^{-2h(\kappa)}} = \chi^{-2h(\kappa)} (1 - \chi)^{2/\kappa} \frac{{}_2F_1\left(\frac{4}{\kappa}, 1 - \frac{4}{\kappa}, \frac{8}{\kappa}; 1 - \chi\right)}{{}_2F_1\left(\frac{4}{\kappa}, 1 - \frac{4}{\kappa}, \frac{8}{\kappa}; 1\right)}, \tag{D.23}$$

and we can expand the hypergeometric function using [AS64, Eq. 15.3.6] as

$$\begin{aligned} \frac{{}_2F_1\left(\frac{4}{\kappa}, 1 - \frac{4}{\kappa}, \frac{8}{\kappa}; 1 - \chi\right)}{{}_2F_1\left(\frac{4}{\kappa}, 1 - \frac{4}{\kappa}, \frac{8}{\kappa}; 1\right)} &= {}_2F_1\left(\frac{4}{\kappa}, 1 - \frac{4}{\kappa}, 2 - \frac{8}{\kappa}; \chi\right) + \chi^{8/\kappa-1} \frac{\Gamma(12/\kappa - 1)\Gamma(1 - 8/\kappa)}{\Gamma(1 - 4/\kappa)\Gamma(8/\kappa - 1)} {}_2F_1\left(\frac{4}{\kappa}, \frac{12}{\kappa} - 1, \frac{8}{\kappa}; \chi\right) \\ &= 1 + \frac{2}{\kappa}\chi + O(\chi^2) + \chi^{8/\kappa-1} \frac{\Gamma(12/\kappa - 1)\Gamma(1 - 8/\kappa)}{\Gamma(1 - 4/\kappa)\Gamma(8/\kappa - 1)} \left(1 + \frac{12-\kappa}{2\kappa}\chi + O(\chi^2)\right), \quad \text{as } \chi \rightarrow 0. \end{aligned}$$

Plugging this into (D.23), since  $\kappa > 8/3$ , we obtain

$$\begin{aligned} \frac{\mathcal{Z}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa)}}(x_1, x_2, x_3, x_4)}{(x_3 - x_1)^{-2h(\kappa)}(x_4 - x_2)^{-2h(\kappa)}} &= \chi^{-2h(\kappa)} \left(1 + \chi^{8/\kappa-1} \frac{\Gamma(12/\kappa - 1)\Gamma(1 - 8/\kappa)}{\Gamma(1 - 4/\kappa)\Gamma(8/\kappa - 1)} + O(\chi^{8/\kappa}) + O(\chi^2)\right) \\ &= \chi^{-2h(\kappa)} + \chi^{2/\kappa} \frac{\Gamma(12/\kappa - 1)\Gamma(1 - 8/\kappa)}{\Gamma(1 - 4/\kappa)\Gamma(8/\kappa - 1)} + o(\chi^{2/\kappa}), \quad \text{as } \chi \rightarrow 0. \end{aligned}$$

This gives (D.22) for  $\kappa \in (8/3, 4) \cup (4, 8)$  because the constants can be written in the form

$$\begin{aligned} \frac{\Gamma(12/\kappa - 1)\Gamma(1 - 8/\kappa)}{\Gamma(1 - 4/\kappa)\Gamma(8/\kappa - 1)} &= \frac{\Gamma(12/\kappa - 1)\Gamma(4/\kappa)\Gamma(8/\kappa)\Gamma(1 - 8/\kappa)}{\Gamma(1 - 4/\kappa)\Gamma(4/\kappa)\Gamma(8/\kappa)\Gamma(8/\kappa - 1)} \\ &= \frac{\Gamma(12/\kappa - 1)\Gamma(4/\kappa) \sin(4\pi/\kappa)}{\Gamma(8/\kappa)\Gamma(8/\kappa - 1) \sin(8\pi/\kappa)} && \text{[by [AS64, Eq. 6.1.7]]} \\ &= \frac{\Gamma(12/\kappa - 1)\Gamma(4/\kappa)}{\Gamma(8/\kappa)\Gamma(8/\kappa - 1)} \frac{1}{2 \cos(4\pi/\kappa)} && \text{[by [AS64, Eq. 15.1.20]]} \\ &= \frac{1}{{}_2F_1\left(\frac{4}{\kappa}, 1 - \frac{4}{\kappa}, \frac{8}{\kappa}; 1\right)} \frac{1}{2 \cos(4\pi/\kappa)}. \end{aligned}$$

□

**Corollary D.7.** For  $\kappa \in (8/3, 8)$  and  $\xi < x_3 < x_4$ , we have

$$\mathcal{F}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa)}}(x_1, x_2, x_3, x_4) = (x_2 - x_1)^{-2h(\kappa)} \nu(\kappa)^2 \mathcal{Z}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa)}}(x_3, x_4) + o((x_2 - x_1)^{2/\kappa}), \quad (\text{D.24})$$

$$\begin{aligned} \mathcal{F}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa)}}(x_1, x_2, x_3, x_4) &= (x_2 - x_1)^{-2h(\kappa)} \nu(\kappa) \mathcal{Z}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa)}}(x_3, x_4) + (x_2 - x_1)^{2/\kappa} (\nu(\kappa)^2 - 1) \mathcal{Z}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa)}}(\xi, x_3, x_4) \\ &\quad + o((x_2 - x_1)^{2/\kappa}), \quad \text{as } x_1, x_2 \rightarrow \xi. \end{aligned} \quad (\text{D.25})$$

*Proof.* These are immediate from (D.11, D.12, D.22, D.20).  $\square$

**Lemma D.8.** Fix  $\kappa = 8/3$  and  $\xi < x_3 < x_4$ . For the renormalized Coulomb gas integrals in Proposition 1.15, we have

$$\widehat{\mathcal{F}}_{\underbrace{\quad}_{\underbrace{\quad}}^{(8/3)}}(x_1, x_2, x_3, x_4) = (x_2 - x_1)^{3/4} \mathcal{Z}_{\underbrace{\quad}_{\underbrace{\quad}}^{(8/3)}}(\xi, x_3, x_4) + o((x_2 - x_1)^{3/4}), \quad (\text{D.26})$$

$$\begin{aligned} \widehat{\mathcal{F}}_{\underbrace{\quad}_{\underbrace{\quad}}^{(8/3)}}(x_1, x_2, x_3, x_4) &= (x_2 - x_1)^{-5/4} \widehat{\mathcal{F}}_{\underbrace{\quad}_{\underbrace{\quad}}^{(8/3)}}(x_3, x_4) - \frac{1}{\pi} (x_2 - x_1)^{3/4} |\log(x_2 - x_1)| \mathcal{Z}_{\underbrace{\quad}_{\underbrace{\quad}}^{(8/3)}}(\xi, x_3, x_4) \\ &\quad + O((x_2 - x_1)^{3/4}), \quad \text{as } x_1, x_2 \rightarrow \xi. \end{aligned} \quad (\text{D.27})$$

*Proof.* First, we prove (D.26) by using a limiting argument. Using Lemma D.3, we obtain

$$\begin{aligned} \widehat{\mathcal{F}}_{\underbrace{\quad}_{\underbrace{\quad}}^{(8/3)}}(x_1, x_2, x_3, x_4) &= \lim_{\kappa' \rightarrow 8/3} \frac{1}{\nu(\kappa')} \mathcal{F}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa')}}(x_1, x_2, x_3, x_4) \\ &= C(8/3) (x_3 - x_1)^{-3/4} (x_3 - x_2)^{3/4} (x_4 - x_1)^{3/4} (x_4 - x_3)^{-5/4} (x_4 - x_2)^{-3/4} \\ &\quad \times (x_2 - x_1)^{-5/4} \oint_{\vartheta(0,1)} (w(1-w)(1-\chi w))^{-3/2} dw, \end{aligned} \quad (\text{D.28})$$

where  $\chi = \text{CR}(x_1, x_2, x_3, x_4)$ . On the one hand, we have the expansion

$$(w(1-w)(1-\chi w))^{-3/2} = (w(1-w))^{-3/2} \left(1 + \frac{3}{2}\chi w + \frac{15}{8}\chi^2 w^2 + o(\chi^2)\right), \quad \text{as } x_1, x_2 \rightarrow \xi. \quad (\text{D.29})$$

On the other hand, using Eq. (B.1) from Lemma B.1, we have

$$\oint_{\vartheta(0,1)} (w(1-w))^{-3/2} dw = \lim_{\kappa' \rightarrow 8/3} \oint_{\vartheta(0,1)} (w(1-w))^{-4/\kappa'} dw = \lim_{\kappa' \rightarrow 8/3} \frac{\nu(\kappa')}{C(\kappa')} = 0, \quad (\text{D.30})$$

Plugging (D.29, D.30) into (D.28) gives the asserted (D.26).

To prove (D.27), one can proceed as in the proof of (D.9) to show that

$$\begin{aligned} \mathcal{F}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa')}}(x_1, x_2, x_3, x_4) &= \nu(\kappa') C(\kappa') (x_4 - x_2)^{-2/\kappa'} (x_2 - x_1)^{2/\kappa'} (x_4 - x_3)^{2/\kappa'} (x_3 - x_2)^{1-6/\kappa'} (x_3 - x_1)^{-2/\kappa'} \\ &\quad \times (x_4 - x_1)^{1-6/\kappa'} \chi^{1-8/\kappa'} \oint_{\vartheta(0, \frac{1}{\chi})} ((1 + (1-\chi)v)v(1-\chi v))^{-4/\kappa} dv \end{aligned}$$

when  $\kappa' \notin \{\frac{8}{m} : m \in \mathbb{Z}_{>0}\}$ , which implies that

$$\begin{aligned} \frac{\widehat{\mathcal{F}}_{\underbrace{\quad}_{\underbrace{\quad}}^{(8/3)}}(x_1, x_2, x_3, x_4)}{(x_3 - x_1)^{-5/4} (x_4 - x_2)^{-5/4}} &= \lim_{\kappa' \rightarrow 8/3} \frac{\mathcal{F}_{\underbrace{\quad}_{\underbrace{\quad}}^{(\kappa')}}(x_1, x_2, x_3, x_4)}{\nu(\kappa') (x_3 - x_1)^{-2h(\kappa')} (x_4 - x_2)^{-2h(\kappa')}} \\ &= C(8/3) (1-\chi)^{-5/4} \chi^{-5/4} \oint_{\vartheta(0, \frac{1}{\chi})} ((1 + (1-\chi)v)v(1-\chi v))^{-3/2} dv. \end{aligned} \quad (\text{D.31})$$

Expanding

$$\begin{aligned} &(1-\chi)^{-5/4} ((1 + (1-\chi)v)v(1-\chi v))^{-3/2} \\ &= ((1+v)v)^{-3/2} \left(1 + \frac{5}{4}\chi + \frac{3}{2}\frac{\chi v}{1+v} + \frac{3}{2}\chi v + \frac{15}{8}\chi^2 v^2 + O(\chi^2) + O(\chi^3 v^3)\right), \end{aligned} \quad (\text{D.32})$$

as  $x_1, x_2 \rightarrow \xi$ , with the constants in  $O(\chi^2)$  and  $O(\chi^3 v^3)$  uniformly bounded in  $v$ , and using the change of variables  $w = \chi v$ , we obtain

$$\oint_{\vartheta(0, \frac{1}{\chi})} \chi^3 v^3 ((1+v)v)^{-3/2} dv = \chi^2 \oint_{\vartheta(0,1)} t^3 (t(\chi+t))^{-3/2} dt = O(\chi^2), \quad \text{as } x_1, x_2 \rightarrow \xi. \quad (\text{D.33})$$

Plugging (B.5, D.32, D.33) into (D.31), we obtain the asserted (D.27).  $\square$

## E Fröbenius series for $\kappa = 4$

In this appendix, we derive Fröbenius series for Coulomb gas integrals  $\mathcal{F}_\beta^{(4)}$  when  $\kappa = 4$ .

**Proposition E.1.** *Fix  $\kappa = 4$  and  $\beta \in \text{LP}_N$ . Let  $\mathbf{x} \in \mathfrak{X}_{2N}$  and  $\xi \in (-\infty, x_3)$ , and use the notation (1.21):*

$$\mathbf{x} = (x_1, \dots, x_{2N}) \in \mathfrak{X}_{2N} \quad \text{and} \quad \check{\mathbf{x}}_1 = (x_3, x_4, \dots, x_{2N}) \in \mathfrak{X}_{2N-2}.$$

- If  $\{1, 2\} \in \beta$ ,

$$\mathcal{F}_\beta^{(4)}(\mathbf{x}) = 2\mathcal{F}_{\beta/\{1,2\}}^{(4)}(\check{\mathbf{x}}_1)(x_2 - x_1)^{-1/2} + o\left((x_2 - x_1)^{1/2}\right), \quad \text{as } x_1, x_2 \rightarrow \xi. \quad (\text{E.1})$$

- If  $\{1, 2\} \notin \beta$ ,

$$\begin{aligned} \mathcal{F}_\beta^{(4)}(\mathbf{x}) &= \mathcal{F}_{\varphi_1(\beta)/\{1,2\}}^{(4)}(\check{\mathbf{x}}_1)(x_2 - x_1)^{-1/2} + \mathcal{F}_{\beta/\vee_1}^{(4)}(\xi, \check{\mathbf{x}}_1)(x_2 - x_1)^{1/2} \\ &\quad + o\left((x_2 - x_1)^{1/2}\right), \quad \text{as } x_1, x_2 \rightarrow \xi, \end{aligned} \quad (\text{E.2})$$

where

$$\mathcal{F}_{\beta/\vee_1}^{(4)}(\xi, \check{\mathbf{x}}_1) = \sum_{\alpha \in \text{LP}_N: \{1,2\} \in \alpha} \mathcal{M}_2(\alpha, \beta) \mathcal{Z}_{\alpha/\vee_1}^{(4)}(\xi, \check{\mathbf{x}}_1) + \sum_{\alpha \in \text{LP}_N: \{1,2\} \notin \alpha} \mathcal{M}_2(\alpha, \beta) \mathcal{Z}_{\alpha/\times_1}^{(4)}(\xi, \check{\mathbf{x}}_1), \quad (\text{E.3})$$

and  $\mathcal{Z}_{\alpha/\vee_1}^{(4)}$  and  $\mathcal{Z}_{\alpha/\times_1}^{(4)}$  are defined in (E.11) and (E.13) respectively.

When  $\kappa = 4$ , pure partition functions have explicit formulas and we will prove Proposition E.1 using these explicit formulas. We first introduce conformal block functions  $\{\mathcal{U}_\gamma : \gamma \in \text{LP}_N\}$  and incidence matrix  $\{\mathcal{K}(\alpha, \gamma) : \alpha, \gamma \in \text{LP}_N\}$ .

**Definition E.2.** For  $\gamma = \{\{a_1, b_1\}, \dots, \{a_N, b_N\}\} \in \text{LP}_N$  ordered as in (1.2), we define conformal block function  $\mathcal{U}_\gamma : \mathfrak{X}_{2N} \rightarrow \mathbb{R}_{>0}$  as follows:

$$\mathcal{U}_\gamma(x_1, \dots, x_{2N}) = \prod_{1 \leq i < j \leq 2N} (x_j - x_i)^{\frac{1}{2} \lambda_\gamma(i,j)}, \quad (\text{E.4})$$

where

$$\lambda_\gamma(i, j) := \begin{cases} +1, & \text{if } i, j \in \{a_1, \dots, a_N\} \text{ or } i, j \in \{b_1, \dots, b_N\}, \\ -1, & \text{otherwise.} \end{cases}$$

**Lemma E.3.** *Suppose  $\gamma \in \text{LP}_N$ . Let  $\mathbf{x} \in \mathfrak{X}_{2N}$  and  $\xi \in (-\infty, x_3)$ , and use the notation (1.21).*

- If  $\{1, 2\} \in \gamma$ ,

$$\begin{aligned} \mathcal{U}_\gamma(\mathbf{x}) = & \mathcal{U}_{\gamma/\{1,2\}}(\ddot{\mathbf{x}}_1)(x_2 - x_1)^{-1/2} + \mathcal{U}_{\gamma/\{1,2\}}(\ddot{\mathbf{x}}_1) \times \left( \sum_{i=3}^{2N} \frac{\frac{1}{2}\lambda_\gamma(i, 1)}{x_i - \xi} \right) (x_2 - x_1)^{1/2} \\ & + o\left((x_2 - x_1)^{1/2}\right), \quad \text{as } x_1, x_2 \rightarrow \xi. \end{aligned} \quad (\text{E.5})$$

- If  $\{1, 2\} \notin \gamma$ ,

$$\mathcal{U}_\gamma(\mathbf{x}) = \mathcal{U}_{\gamma/\times_1}(\xi, \ddot{\mathbf{x}}_1)(x_2 - x_1)^{1/2} + o\left((x_2 - x_1)^{1/2}\right), \quad \text{as } x_1, x_2 \rightarrow \xi, \quad (\text{E.6})$$

where

$$\mathcal{U}_{\gamma/\times_1}(\xi, \ddot{\mathbf{x}}_1) = \prod_{i=3}^{2N} (x_i - \xi)^{\lambda_\gamma(i,1)} \times \prod_{3 \leq i < j \leq 2N} (x_j - x_i)^{\frac{1}{2}\lambda_\gamma(i,j)}. \quad (\text{E.7})$$

*Proof.* The expansion (E.6) is proved in [LW21, Lemma 3.3]. It remains to show (E.5). Assume  $\{1, 2\} \in \gamma$ . From (E.4), we have

$$\mathcal{U}_\gamma(\mathbf{x}) = (x_2 - x_1)^{-1/2} \times \underbrace{\prod_{i=3}^{2N} \left( \frac{x_i - x_2}{x_i - x_1} \right)^{-\frac{1}{2}\lambda_\gamma(i,1)}}_P \times \prod_{3 \leq i < j \leq 2N} (x_j - x_i)^{\frac{1}{2}\lambda_\gamma(i,j)}. \quad (\text{E.8})$$

Note that

$$\begin{aligned} P &= \prod_{i=3}^{2N} \left( 1 - \frac{x_2 - x_1}{x_i - x_1} \right)^{-\frac{1}{2}\lambda_\gamma(i,1)} \\ &= \prod_{i=3}^{2N} \left( 1 + \frac{1}{2}\lambda_\gamma(i,1) \frac{x_2 - x_1}{x_i - x_1} \right) + o(x_2 - x_1) \\ &= 1 + \sum_{i=3}^{2N} \frac{1}{2}\lambda_\gamma(i,1) \frac{x_2 - x_1}{x_i - x_1} + o(x_2 - x_1). \end{aligned}$$

Plugging into (E.8), we obtain (E.5).  $\square$

**Definition E.4.** Let  $\alpha = \{\{a_1, b_1\}, \dots, \{a_N, b_N\}\} \in \text{LP}_N$  be ordered as in (1.2). For  $\gamma \in \text{LP}_N$ , we write  $\alpha \leftarrow \gamma$  if there exists a permutation  $\sigma$  of  $\{1, 2, \dots, N\}$  such that

$$\gamma = \{\{a_1, b_{\sigma(1)}\}, \dots, \{a_N, b_{\sigma(N)}\}\}.$$

We define the *incidence matrix*  $\{\mathcal{K}(\alpha, \gamma) : \alpha, \gamma \in \text{LP}_N\}$  via

$$\mathcal{K}(\alpha, \gamma) = \mathbb{1}\{\alpha \leftarrow \gamma\}.$$

The incidence matrix  $\mathcal{K}$  is invertible, see [KW11b, Theorem 1.5 and 1.6] and [KKP20, Theorem 2.9]. When  $\kappa = 4$ , pure partition functions and conformal block functions are related through incidence matrix.

**Lemma E.5.** [PW19, Theorem 1.5] Fix  $\kappa = 4$ .

$$\mathcal{Z}_\alpha^{(4)}(\mathbf{x}) = \sum_{\gamma \in \text{LP}_N} \mathcal{K}^{-1}(\alpha, \gamma) \mathcal{U}_\gamma(\mathbf{x}), \quad \text{for all } \mathbf{x} \in \mathfrak{X}_{2N}, \alpha \in \text{LP}_N. \quad (\text{E.9})$$



**Lemma E.6.** Fix  $\kappa = 4$  and  $\alpha \in \text{LP}_N$ . Let  $\mathbf{x} \in \mathfrak{X}_{2N}$  and  $\xi \in (-\infty, x_3)$ , and use the notation (1.21).

- If  $\{1, 2\} \in \alpha$ ,

$$\begin{aligned} \mathcal{Z}_\alpha^{(4)}(\mathbf{x}) &= \mathcal{Z}_{\alpha/\{1,2\}}^{(4)}(\ddot{\mathbf{x}}_1)(x_2 - x_1)^{-1/2} + \mathcal{Z}_{\alpha/\vee_1}^{(4)}(\xi, \ddot{\mathbf{x}}_1)(x_2 - x_1)^{1/2} \\ &\quad + o\left((x_2 - x_1)^{1/2}\right), \quad \text{as } x_1, x_2 \rightarrow \xi, \end{aligned} \quad (\text{E.10})$$

where

$$\begin{aligned} \mathcal{Z}_{\alpha/\vee_1}^{(4)}(\xi, \ddot{\mathbf{x}}_1) &= \sum_{\gamma \in \text{LP}_N: \{1,2\} \in \gamma} \mathcal{K}^{-1}(\alpha, \gamma) \mathcal{U}_{\gamma/\{1,2\}}(\ddot{\mathbf{x}}_1) \left( \sum_{i=3}^{2N} \frac{\frac{1}{2}\lambda_\gamma(i, 1)}{x_i - \xi} \right) \\ &\quad + \sum_{\gamma \in \text{LP}_N: \{1,2\} \notin \gamma} \mathcal{K}^{-1}(\alpha, \gamma) \mathcal{U}_{\gamma/\times_1}(\xi, \ddot{\mathbf{x}}_1). \end{aligned} \quad (\text{E.11})$$

- If  $\{1, 2\} \notin \alpha$ ,

$$\mathcal{Z}_\alpha^{(4)}(\mathbf{x}) = \mathcal{Z}_{\alpha/\times_1}^{(4)}(\xi, \ddot{\mathbf{x}}_1)(x_2 - x_1)^{1/2} + o\left((x_2 - x_1)^{1/2}\right), \quad \text{as } x_1, x_2 \rightarrow \xi, \quad (\text{E.12})$$

where

$$\mathcal{Z}_{\alpha/\times_1}^{(4)}(\xi, \ddot{\mathbf{x}}_1) = \sum_{\gamma \in \text{LP}_N: \{1,2\} \notin \gamma} \mathcal{K}^{-1}(\alpha, \gamma) \mathcal{U}_{\gamma/\times_1}(\xi, \ddot{\mathbf{x}}_1). \quad (\text{E.13})$$

*Proof.* The expansion (E.10) follows from Lemmas E.3 and E.5. The expansion (E.12) is proved in [LW21, Lemma 3.4].  $\square$

*Proof of Proposition E.1.* As  $x_1, x_2 \rightarrow \xi$ , from (E.12) and (E.10), we have

$$\begin{aligned} \mathcal{F}_\beta^{(4)}(\mathbf{x}) &= \sum_{\alpha \in \text{LP}_N} \mathcal{M}_2(\alpha, \beta) \mathcal{Z}_\alpha^{(4)}(\mathbf{x}) \\ &= \sum_{\alpha \in \text{LP}_N: \{1,2\} \in \alpha} \mathcal{M}_2(\alpha, \beta) \left( \mathcal{Z}_{\alpha/\{1,2\}}^{(4)}(\ddot{\mathbf{x}}_1)(x_2 - x_1)^{-1/2} + \mathcal{Z}_{\alpha/\vee_1}^{(4)}(\xi, \ddot{\mathbf{x}}_1)(x_2 - x_1)^{1/2} \right) \\ &\quad + \sum_{\alpha \in \text{LP}_N: \{1,2\} \notin \alpha} \mathcal{M}_2(\alpha, \beta) \mathcal{Z}_{\alpha/\times_1}^{(4)}(\xi, \ddot{\mathbf{x}}_1)(x_2 - x_1)^{1/2} + o\left((x_2 - x_1)^{1/2}\right) \\ &= \underbrace{\sum_{\alpha \in \text{LP}_N: \{1,2\} \in \alpha} \mathcal{M}_2(\alpha, \beta) \mathcal{Z}_{\alpha/\{1,2\}}^{(4)}(\ddot{\mathbf{x}}_1)(x_2 - x_1)^{-1/2}}_{S_1} \\ &\quad + \underbrace{\left( \sum_{\alpha \in \text{LP}_N: \{1,2\} \in \alpha} \mathcal{M}_2(\alpha, \beta) \mathcal{Z}_{\alpha/\vee_1}^{(4)}(\xi, \ddot{\mathbf{x}}_1) + \sum_{\alpha \in \text{LP}_N: \{1,2\} \notin \alpha} \mathcal{M}_2(\alpha, \beta) \mathcal{Z}_{\alpha/\times_1}^{(4)}(\xi, \ddot{\mathbf{x}}_1) \right)}_{\mathcal{F}_{\beta/\vee_1}^{(4)}(\xi, \ddot{\mathbf{x}}_1)} (x_2 - x_1)^{1/2} \\ &\quad + o\left((x_2 - x_1)^{1/2}\right). \end{aligned}$$

Note that

$$S_1 = \begin{cases} 2\mathcal{F}_{\beta/\{1,2\}}^{(4)}(\ddot{\mathbf{x}}_1), & \text{if } \{1, 2\} \in \beta; \\ \mathcal{F}_{\emptyset_1(\beta)/\{1,2\}}^{(4)}(\ddot{\mathbf{x}}_1), & \text{if } \{1, 2\} \notin \beta. \end{cases}$$

In particular, we obtain (E.2). It remains to show (E.1).

Suppose  $\{1, 2\} \in \beta$  and we will prove (E.1) using estimates in Lemma 6.1. Recall from (6.4) and the discussion after it, for  $\kappa \in (8/3, 8)$ ,

$$\mathcal{F}_\beta^{(\kappa)}(\mathbf{x}) - \nu(\kappa) \mathcal{F}_{\beta/\{1,2\}}^{(\kappa)}(\ddot{\mathbf{x}}_1) (x_2 - x_1)^{-2h(\kappa)} = (x_2 - x_1)^{-2h(\kappa)+2} G_\beta^{(\kappa)}(\mathbf{x}) \quad (\text{E.14})$$

where  $G_\beta^{(\kappa)}(\mathbf{x})$  is uniformly bounded in  $\kappa \in (8/3, 8)$  and  $x_1, x_2 \in (\xi - (x_3 - \xi)/3, \xi + (x_3 - \xi)/3)$ . Set  $\kappa = 4$  in (E.14), we obtain

$$\mathcal{F}_\beta^{(4)}(\mathbf{x}) - 2\mathcal{F}_{\beta/\{1,2\}}^{(4)}(\ddot{\mathbf{x}}_1) (x_2 - x_1)^{-1/2} = O\left((x_2 - x_1)^{3/2}\right), \quad \text{as } x_1, x_2 \rightarrow \xi,$$

and complete the proof of (E.1). □

## References

- [ABKM20] Tom Alberts, Sung-Soo Byun, Nam-Gyu Kang, and Nikolai Makarov. Pole dynamics and an integral of motion for multiple SLE(0). Preprint in [arXiv:2011.05714](#), 2020.
- [AHSY23] Morris Ang, Nina Holden, Xin Sun, and Pu Yu. Conformal welding of quantum disks and multiple SLE: the non-simple case. Preprint in [arXiv:2310.20583](#), 2023.
- [AS64] Milton Abramowitz and Irene A. Stegun. *Handbook of mathematical functions*. Dover Publications Inc., 1964.
- [BB03] Michel Bauer and Denis Bernard. Conformal field theories of stochastic Loewner evolutions. *Comm. Math. Phys.*, 239(3):493–521, 2003.
- [BBK05] Michel Bauer, Denis Bernard, and Kalle Kytölä. Multiple Schramm-Loewner evolutions and statistical mechanics martingales. *J. Stat. Phys.*, 120(5-6):1125–1163, 2005.
- [BCM10] András Bálint, Federico Camia, and Ronald Meester. The high temperature Ising model on the triangular lattice is a critical Bernoulli percolation model. *J. Stat. Phys.*, 139(1):122–138, 2010.
- [Bon69] Jean-Michel Bony. Principe du maximum, inégalité de Harnack et unicité du problème de Cauchy pour les opérateurs elliptiques dégénérés. *Ann. Inst. Fourier*, 19(1):277–304, 1969.
- [BPW21] Vincent Beffara, Eveliina Peltola, and Hao Wu. On the uniqueness of global multiple SLEs. *Ann. Probab.*, 49(1):400–434, 2021.
- [BPZ84a] Alexander A. Belavin, Alexander M. Polyakov, and Alexander B. Zamolodchikov. Infinite conformal symmetry in two-dimensional quantum field theory. *Nucl. Phys. B*, 241(2):333–380, 1984.
- [BPZ84b] Alexander A. Belavin, Alexander M. Polyakov, and Alexander B. Zamolodchikov. Infinite conformal symmetry of critical fluctuations in two dimensions. *J. Stat. Phys.*, 34(5-6):763–774, 1984.
- [Car92] John L. Cardy. Critical percolation in finite geometries. *J. Phys. A*, 25(4):L201–206, 1992.
- [CDCH<sup>+</sup>14] Dmitry Chelkak, Hugo Duminil-Copin, Clément Hongler, Antti Kemppainen, and Stanislav Smirnov. Convergence of Ising interfaces to Schramm’s SLE curves. *C. R. Acad. Sci. Paris Sér. I Math.*, 352(2):157–161, 2014.

- [CN06] F. Camia and C. M. Newman. Two-dimensional critical percolation: the full scaling limit. *Comm. Math. Phys.*, 268(1):1–38, 2006.
- [CN07] F. Camia and C. M. Newman. Critical percolation exploration path and SLE(6): a proof of convergence. *Probab. Theory Related Fields*, 139(3-4):473–519, 2007.
- [CS12] Dmitry Chelkak and Stanislav Smirnov. Universality in the 2D Ising model and conformal invariance of fermionic observables. *Invent. Math.*, 189(3):515–580, 2012.
- [DCS12] Hugo Duminil-Copin and Stanislav Smirnov. The connective constant of the honeycomb lattice equals  $\sqrt{2 + \sqrt{2}}$ . *Ann. of Math.*, 175(3):1653–1665, 2012.
- [DFGG97] P. Di Francesco, O. Golinelli, and E. Guitter. Meanders and the Temperley-Lieb algebra. *Comm. Math. Phys.*, 186(1):1–59, 1997.
- [DFMS97] Philippe Di Francesco, Pierre Mathieu, and David Sénéchal. *Conformal field theory*. Graduate Texts in Contemporary Physics. Springer-Verlag, New York, 1997.
- [DMS21] Bertrand Duplantier, Jason Miller, and Scott Sheffield. Liouville quantum gravity as a mating of trees. *Astérisque*, 427:258pp, 2021.
- [Dub06] Julien Dubédat. Euler integrals for commuting SLEs. *J. Stat. Phys.*, 123(6):1183–1218, 2006.
- [Dub07a] Julien Dubédat. Commutation relations for SLE. *Comm. Pure Appl. Math.*, 60(12):1792–1847, 2007.
- [Dub07b] Julien Dubédat. Duality of Schramm-Loewner Evolutions. *Ann. Sci. Éc. Norm. Supér.*, 42(5):697–724, 2007.
- [Dub15a] Julien Dubédat. SLE and Virasoro representations: fusion. *Comm. Math. Phys.*, 336(2):761–809, 2015.
- [Dub15b] Julien Dubédat. SLE and Virasoro representations: localization. *Comm. Math. Phys.*, 336(2):695–760, 2015.
- [Dup06] Bertrand Duplantier. Conformal random geometry. In *Mathematical Statistical Physics: Lecture Notes of the Les Houches Summer School (2005)*. Elsevier, 2006.
- [Eva10] Lawrence C. Evans. *Partial Differential Equations*. Graduate Studies in Mathematics. American Mathematical Society, 2 edition, 2010.
- [FH23] Juraj Földes and David P. Herzog. The method of stochastic characteristics for linear second-order hypoelliptic equations. *Probab. Surveys*, 20:113–169, 2023.
- [FK04] Roland Friedrich and Jussi Kalkkinen. On conformal field theory and stochastic Loewner evolution. *Nucl. Phys. B*, 687(3):279–302, 2004.
- [FK15a] Steven M. Flores and Peter Kleban. A solution space for a system of null-state partial differential equations 1. *Comm. Math. Phys.*, 333(1):389–434, 2015.
- [FK15b] Steven M. Flores and Peter Kleban. A solution space for a system of null-state partial differential equations 2. *Comm. Math. Phys.*, 333(1):435–481, 2015.
- [FK15c] Steven M. Flores and Peter Kleban. A solution space for a system of null-state partial differential equations 3. *Comm. Math. Phys.*, 333(2):597–667, 2015.

- [FK15d] Steven M. Flores and Peter Kleban. A solution space for a system of null-state partial differential equations 4. *Comm. Math. Phys.*, 333(2):669–715, 2015.
- [FLW23] Yu Feng, Mingchang Liu, and Hao Wu. Decomposition of global 2-SLE for  $\kappa \in (4, 8)$  and an application for critical FK-Ising model. Preprint in [arXiv:2310.10234](#), 2023.
- [FP18] Steven M. Flores and Eveliina Peltola. Standard modules, radicals, and the valenced Temperley-Lieb algebra. Preprint in [arXiv:1801.10003](#), 2018.
- [FP20] Steven M. Flores and Eveliina Peltola. Higher spin quantum and classical Schur-Weyl duality for  $\mathfrak{sl}(2)$ . Preprint in [arXiv:2008.06038](#), 2020.
- [FP24] Steven M. Flores and Eveliina Peltola. Solution spaces of the Benoit & Saint-Aubin partial differential equations. In preparation, 2024.
- [FPW24a] Yu Feng, Eveliina Peltola, and Hao Wu. Connection probabilities of multiple FK-Ising interfaces. *Probab. Theory Related Fields*, 189(1-2):281–367, 2024.
- [FPW24b] Yu Feng, Eveliina Peltola, and Hao Wu. Crossing probabilities of multiple percolation interfaces: generalizations of Cardy’s formula and Watt’s formula. In preparation, 2024.
- [FSK15] Steven M. Flores, Jacob J. H. Simmons, and Peter Kleban. Multiple-SLE $_{\kappa}$  connectivity weights for rectangles, hexagons, and octagons. Preprint in [arXiv:1505.07756](#), 2015.
- [FSKZ17] Steven M. Flores, Jacob J. H. Simmons, Peter Kleban, and Robert M. Ziff. A formula for crossing probabilities of critical systems inside polygons. *J. Phys. A*, 50(6):064005, 2017.
- [FW91] Giovanni Felder and Christian Wierczkowski. Topological representations of the quantum group  $U_q(\mathfrak{sl}_2)$ . *Comm. Math. Phys.*, 138(3):583–605, 1991.
- [FW03] Roland Friedrich and Wendelin Werner. Conformal restriction, highest weight representations and SLE. *Comm. Math. Phys.*, 243(1):105–122, 2003.
- [GL02] Victor Gurarie and Andreas W. W. Ludwig. Conformal algebras of two-dimensional disordered systems. *J. Phys. A: Math. Gen.*, 35:L377–L384, 2002.
- [Gra07] Kevin Graham. On multiple Schramm-Loewner evolutions. *J. Stat. Mech. Theory Exp.*, P03008, 2007.
- [GSNJ<sup>+</sup>22] Linnea Grans-Samuelsson, Rongvoram Nivesvivat, Jesper L. Jacobsen, Sylvain Ribault, and Hubert Saleur. Global symmetry and conformal bootstrap in the two-dimensional  $O(n)$  model. *SciPost Physics*, 12(5):1–45, 2022.
- [GT01] David Gilbarg and Neil S. Trudinger. *Elliptic partial differential equations of second order*. Classics in Mathematics. Springer-Verlag, 2001. Reprint of the 2nd ed. Berlin Heidelberg New York 1983.
- [Gur93] Victor Gurarie. Logarithmic operators in conformal field theory. *Nucl. Phys. B*, 410(3):535–549, 1993.
- [Gur13] Victor Gurarie. Logarithmic operators and logarithmic conformal field theories. *J. Phys. A: Math. Theor.*, 46:494003, 2013.
- [GZ20] Victor Gorbenko and Bernardo Zan. Two-dimensional  $O(n)$  models and logarithmic CFTs. *J. High Energy Phys.*, 10(99):1–54, 2020.

- [Hör90] Lars Hörmander. *The analysis of linear partial differential operators I: Distribution theory and Fourier analysis*, volume 256 of *Grundlehren der mathematischen Wissenschaften*. Springer-Verlag, Berlin Heidelberg, 2 edition, 1990.
- [HS13] Clément Hongler and Stanislav Smirnov. The energy density in the planar Ising model. *Acta Math.*, 211(2):191–225, 2013.
- [Izy15] Konstantin Izyurov. Smirnov’s observable for free boundary conditions, interfaces and crossing probabilities. *Comm. Math. Phys.*, 337(1):225–252, 2015.
- [Izy22] Konstantin Izyurov. On multiple SLE for the FK-Ising model. *Ann. Probab.*, 50(2):771–790, 2022.
- [Kas95] Christian Kassel. *Quantum groups*, volume 155 of *Graduate texts in Mathematics*. Springer-Verlag, 1995.
- [Kem17] Antti Kemppainen. *Schramm-Loewner evolution*, volume 24 of *SpringerBriefs in Mathematical Physics*. Springer International Publishing, 2017.
- [KKP20] Alex Karrila, Kalle Kytölä, and Eveliina Peltola. Boundary correlations in planar LERW and UST. *Comm. Math. Phys.*, 376(3):2065–2145, 2020.
- [KL07] Michael J. Kozdron and Gregory F. Lawler. The configurational measure on mutually avoiding SLE paths. *Fields Inst. Commun.*, 50:199–224, 2007.
- [KN04] Wouter Kager and Bernard Nienhuis. A guide to stochastic Löwner evolution and its applications. *J. Stat. Phys.*, 115(5):1149–1229, 2004.
- [KP16] Kalle Kytölä and Eveliina Peltola. Pure partition functions of multiple SLEs. *Comm. Math. Phys.*, 346(1):237–292, 2016.
- [KP20] Kalle Kytölä and Eveliina Peltola. Conformally covariant boundary correlation functions with a quantum group. *J. Eur. Math. Soc.*, 22(1):55–118, 2020.
- [KS98] Ioannis Karatzas and Steven E. Shreve. *Brownian Motion and Stochastic Calculus*. Springer, 2 edition, 1998.
- [KS17] Antti Kemppainen and Stanislav Smirnov. Random curves, scaling limits and Loewner evolutions. *Ann. Probab.*, 45(2):698–779, 2017.
- [KW11a] Richard W. Kenyon and David B. Wilson. Boundary partitions in trees and dimers. *Trans. Amer. Math. Soc.*, 363(3):1325–1364, 2011.
- [KW11b] Richard W. Kenyon and David B. Wilson. Double-dimer pairings and skew Young diagrams. *Electron. J. Combin.*, 18(1):130–142, 2011.
- [Law05] Gregory F. Lawler. *Conformally invariant processes in the plane*, volume 114 of *Mathematical Surveys and Monographs*. American Mathematical Society, 2005.
- [Law09] Gregory F. Lawler. Partition functions, loop measure, and versions of SLE. *J. Stat. Phys.*, 134(5-6):813–837, 2009.
- [LPW24] Mingchang Liu, Eveliina Peltola, and Hao Wu. Uniform spanning tree in topological polygons, partition functions for SLE(8), and correlations in  $c = -2$  logarithmic CFT. *Ann. Probab.*, to appear, 2024. Preprint in [arXiv:2108.04421](https://arxiv.org/abs/2108.04421).

- [LSW01a] Gregory F. Lawler, Oded Schramm, and Wendelin Werner. The dimension of the planar Brownian frontier is  $4/3$ . *Math. Res. Lett.*, 8:401–411, 2001.
- [LSW01b] Gregory F. Lawler, Oded Schramm, and Wendelin Werner. Values of Brownian intersection exponents I: Half-plane exponents. *Acta Math.*, 187(2):237–273, 2001.
- [LSW04] Gregory F. Lawler, Oded Schramm, and Wendelin Werner. Conformal invariance of planar loop-erased random walks and uniform spanning trees. *Ann. Probab.*, 32(1B):939–995, 2004.
- [LW21] Mingchang Liu and Hao Wu. Scaling limits of crossing probabilities in metric graph GFF. *Electron. J. Probab.*, 26(37):1–46, 2021.
- [MS16a] Jason Miller and Scott Sheffield. Imaginary geometry I: interacting SLEs. *Probab. Theory Related Fields*, 164(3-4):553–705, 2016.
- [MS16b] Jason Miller and Scott Sheffield. Imaginary geometry II: reversibility of  $SLE_\kappa(\rho_1, \rho_2)$  for  $\kappa \in (0, 4)$ . *Ann. Probab.*, 44(3):1647–1722, 2016.
- [MS16c] Jason Miller and Scott Sheffield. Imaginary geometry III: reversibility of  $SLE_\kappa$  for  $\kappa \in (4, 8)$ . *Ann. Math.*, 184(2):455–486, 2016.
- [MSW20] Jason Miller, Scott Shaffield, and Wendelin Werner. Non-simple SLE curves are not determined by their range. *J. Eur. Math. Soc.*, 22(3):669–716, 2020.
- [MSW21] Jason Miller, Scott Sheffield, and Wendelin Werner. Non-simple conformal loop ensembles on Liouville quantum gravity and the law of CLE percolation interfaces. *Probab. Theory Related Fields*, 181(1-3):669–710, 2021.
- [MW18] Jason Miller and Wendelin Werner. Connection probabilities for conformal loop ensembles. *Comm. Math. Phys.*, 362(2):415–453, 2018.
- [Nie82] Bernard Nienhuis. Exact critical point and exponents of the  $O(n)$  model in two dimensions. *Phys. Rev. Lett.*, 49:1062–1065, 1982.
- [Øks03] Bernt Øksendal. *Stochastic Differential Equations*. Springer, 2003.
- [Pel19] Eveliina Peltola. Towards a conformal field theory for Schramm-Loewner evolutions. *J. Math. Phys.*, 60(10):103305, 2019. Special issue (Proc. ICMP, Montreal, July 2018).
- [Pel20] Eveliina Peltola. Basis for solutions of the Benoit & Saint-Aubin PDEs with particular asymptotic properties. *Ann. Inst. Henri Poincaré D*, 7(1):1–73, 2020.
- [PR07] Paul A. Pearce and Jørgen Rasmussen. Solvable critical dense polymers. *J. Stat. Mech.*, 0702:P02015, 2007.
- [PRZ06] Paul A. Pearce, Jørgen Rasmussen, and Jean-Bernard Zuber. Logarithmic minimal models. *J. Stat. Mech.*, 06(11):P11017, 2006.
- [PS19] Ron Peled and Yinon Spinka. Lectures on the spin and loop  $O(n)$  models. In *Sojourns in Probability Theory and Statistical Physics I*, pages 246–320. Springer, Singapore, 2019.
- [PW19] Eveliina Peltola and Hao Wu. Global and local multiple SLEs for  $\kappa \leq 4$  and connection probabilities for level lines of GFF. *Comm. Math. Phys.*, 366(2):469–536, 2019.
- [PW23] Eveliina Peltola and Hao Wu. Crossing probabilities of multiple Ising interfaces. *Ann. Appl. Probab.*, 33(4):3169–3206, 2023.

- [PW24] Eveliina Peltola and Yilin Wang. Large deviations of multichordal  $\text{SLE}_{0+}$ , real rational functions, and zeta-regularized determinants of Laplacians. *J. Eur. Math. Soc.*, 26(2):469–535, 2024.
- [RS05] Steffen Rohde and Oded Schramm. Basic properties of SLE. *Ann. of Math.*, 161(2):883–924, 2005.
- [RS07] N. Read and H. Saleur. Associative-algebraic approach to logarithmic conformal field theories. *Nucl. Phys. B*, 777(3):316–351, 2007.
- [Rud91] Walter Rudin. *Functional Analysis*. International Series in Pure and Applied Mathematics. McGraw-Hill, 2 edition, 1991.
- [Sch00] Oded Schramm. Scaling limits of loop-erased random walks and uniform spanning trees. *Israel J. Math.*, 118(1):221–288, 2000.
- [She16] Scott Sheffield. Conformal weldings of random surfaces: SLE and the quantum gravity zipper. *Ann. Probab.*, 44(5):3474–3545, 2016.
- [Smi01] Stanislav Smirnov. Critical percolation in the plane: conformal invariance, Cardy’s formula, scaling limits. *C. R. Acad. Sci.*, 333(3):239–244, 2001. (Updated 2009, see arXiv:0909.4499).
- [Smi06] Stanislav Smirnov. Towards conformal invariance of  $2D$  lattice models. In *Proceedings of the ICM 2006, Madrid, Spain*, volume 2, pages 1421–1451. European Mathematical Society, 2006.
- [SS09] Oded Schramm and Scott Sheffield. Contour lines of the two-dimensional discrete Gaussian free field. *Acta Math.*, 202(1):21–137, 2009.
- [Str08] Daniel W. Stroock. *An introduction to partial differential equations for probabilists*, volume 112 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2008.
- [SW01] Stanislav Smirnov and Wendelin Werner. Critical exponents for two-dimensional percolation. *Math. Res. Lett.*, 8(6):729–744, 2001.
- [SW05] Oded Schramm and David B. Wilson. SLE coordinate changes. *New York J. Math.*, 11:659–669, 2005.
- [SW12] Scott Sheffield and Wendelin Werner. Conformal loop ensembles: The Markovian characterization and the loop-soup construction. *Ann. Math.*, 176(3):1827–1917, 2012.
- [SY23] Xin Sun and Pu Yu. SLE partition functions via conformal welding of random surfaces. Preprint in arXiv:2309.05177, 2023.
- [Tao09] Terence Tao. *An epsilon of room, I: Real analysis. Pages from year three of a mathematical blog*, volume 117 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2009.
- [Wer03] Wendelin Werner. Random planar curves and Schramm-Loewner evolutions. Lecture notes from the Saint-Flour Summer School (July 2002), 2003.
- [Wu20] Hao Wu. Hypergeometric SLE: conformal Markov characterization and applications. *Comm. Math. Phys.*, 374(1):433–484, 2020.

- [WW24] Yilin Wang and Hao Wu. Commutation relations for two-sided radial SLE. Preprint in `arXiv:2405.07082`, 2024.
- [Zha08] Dapeng Zhan. Duality of chordal SLE. *Invent. Math.*, 174(2):309–353, 2008.
- [Zha23] Dapeng Zhan. Existence and uniqueness of nonsimple multiple SLE. Preprint in `arXiv:2308.13886`, 2023.