

DUALITY AND THE EQUATIONS OF REES RINGS AND TANGENT ALGEBRAS

MATTHEW WEAVER

ABSTRACT. Let E be a module of projective dimension one over a Noetherian ring R and consider its Rees algebra $\mathcal{R}(E)$. We study this ring as a quotient of the symmetric algebra $\mathcal{S}(E)$ and consider the ideal \mathcal{A} defining this quotient. In the case that $\mathcal{S}(E)$ is a complete intersection ring, we employ a duality between \mathcal{A} and $\mathcal{S}(E)$ in order to study the Rees ring $\mathcal{R}(E)$ in multiple settings. In particular, when R is a complete intersection ring defined by quadrics, we consider its module of Kähler differentials $\Omega_{R/k}$ and its associated tangent algebras.

1. INTRODUCTION

The objective of this paper is to study the Rees algebra of particular ideals and modules over a Noetherian ring R . Although our treatment of the subject is purely algebraic, much of its motivation is geometric. Indeed, the Rees ring is often called the *blowup algebra* since, for I an R -ideal, the Rees algebra $\mathcal{R}(I) = R \oplus It \oplus I^2t^2 \oplus \cdots$ is the algebraic realization of the blowup of the affine scheme $\text{Spec}(R)$ along the subscheme $V(I)$. Moreover, this notion can be extended to Rees algebras of *modules* in order to treat the case of *repeated* or *successive* blowups. Indeed, for I and J ideals of R , the successive blowup along the disjoint subschemes $V(I)$ and $V(J)$ corresponds to the Rees algebra of $I \oplus J$, which is of course a module and not an ideal. As such, one is obligated to study Rees rings of modules, hence we proceed in this setting and consider the Rees algebra $\mathcal{R}(E)$ of an R -module E . We note that, as this is the more general notion, one is always able to recover the case of Rees rings of ideals.

In addition to the blowup construction, Rees algebras of modules possess many other applications to algebraic geometry. The most notable such module is the module of *Kähler differentials* $\Omega_{R/k}$ of an affine k -algebra R . The Rees ring $\mathcal{R}(\Omega_{R/k})$, and its related algebras, the symmetric algebra $\mathcal{S}(\Omega_{R/k})$ and the fiber ring $\mathcal{F}(\Omega_{R/k})$, are often called *tangent algebras*. If $X \subseteq \mathbb{P}_k^n$ is a subvariety with coordinate ring R , the fiber ring $\mathcal{F}(\Omega_{R/k})$ is the homogeneous coordinate ring of the *tangential variety* of X . Moreover, the closed fibers of the map $\text{Spec}(\mathcal{S}(\Omega_{R/k})) \rightarrow \text{Spec}(R)$ are precisely the Zariski tangent spaces to closed points of $\text{Spec}(R)$ [31]. One might also consider the *Jacobian module* \mathfrak{J} of R and its Rees algebra $\mathcal{R}(\mathfrak{J})$, which is the coordinate ring of the *conormal variety* of X .

In addition to encoding the algebraic data of blowups of affine schemes, Rees rings also serve as bihomogeneous coordinate rings of graphs of rational maps between projective varieties. As these constructions are parametric, the natural question that arises is how to find the corresponding *implicit* equations of these objects. Algebraically, this correlates to expressing the Rees ring $\mathcal{R}(E)$ as a quotient of a polynomial ring $R[y_1, \dots, y_n]$ over the base ring R . Such a description of the Rees ring provides insight into a wealth of information, and is hence a desirable form to obtain. The ideal \mathcal{J} defining this quotient ring is appropriately called the *defining ideal* of $\mathcal{R}(E)$. Unfortunately, a concrete description of the ideal \mathcal{J} and its generators, the *defining equations* of $\mathcal{R}(E)$, is difficult to attain in general. However, there has been success for Rees algebras of ideals with small codimension [2, 7, 24, 26, 27, 28, 29, 40, 41] and modules with small projective dimension [6, 7, 32, 40] in a multitude of settings.

Although it is desirable to relate the Rees ring to $R[y_1, \dots, y_n]$ and express $\mathcal{R}(E)$ as a quotient of such a ring, it is often more convenient to consider a much closer algebra mapping onto $\mathcal{R}(E)$. The natural choice is the symmetric algebra $\mathcal{S}(E)$ of the module E and the natural epimorphism $\mathcal{S}(E) \rightarrow \mathcal{R}(E)$ with kernel \mathcal{A} . In a sense, \mathcal{A} is a defining ideal of the Rees ring as well, as $\mathcal{R}(E) \cong \mathcal{S}(E)/\mathcal{A}$. Moreover, \mathcal{A} has the convenience of being simpler than its counterpart \mathcal{J} , and is often relatable to natural constructions. Moreover, one can always deduce information on \mathcal{J} from \mathcal{A} .

The theme of this paper is to apply a particular duality between \mathcal{A} and $\mathcal{S}(E)$, when $\mathcal{S}(E)$ is a complete intersection ring, in order to study the Rees ring $\mathcal{R}(E)$. We remark that this is not an entirely novel

approach, and such a duality for complete intersections was noted originally by Jouanolou [21], making such an isomorphism explicit through the use of *Morley forms*. This was later adapted to symmetric algebras of ideals by Kustin, Polini, and Ulrich [25]. We introduce Jouanolou's duality for any *bigraded* complete intersection ring, so that we may apply it to study Rees rings of modules in multiple settings.

Theorem A. *Let $R = k[x_1, \dots, x_d]$, $T = k[y_1, \dots, y_n]$, and $B = R \otimes_k T$. With the bigrading on B given by $\text{bideg } x_i = (1, 0)$ and $\text{bideg } y_i = (0, 1)$, let $\underline{f} = f_1, \dots, f_d \subseteq \mathfrak{m} = (x_1, \dots, x_d)$ be a B -regular sequence of bihomogeneous elements. Assume that $\text{bideg } f_i = (\alpha_i, \beta_i)$ and let $\mathcal{S} = B/\mathcal{I}$, where $\mathcal{I} = (\underline{f})$. There is a bihomogeneous isomorphism of B -modules*

$$H_{\mathfrak{m}}^0(\mathcal{S}) \cong \text{Hom}_T(\mathcal{S}, T)(-\delta, -\tau)$$

where $\delta = (\sum_{i=1}^d \alpha_i) - d$, $\tau = \sum_{i=1}^d \beta_i$, and $H_{\mathfrak{m}}^0(\mathcal{S})$ is the zeroth local cohomology module of \mathcal{S} with respect to \mathfrak{m} .

Under certain hypotheses, we can ensure that the symmetric algebra $\mathcal{S}(E)$ is a complete intersection ring and the $\mathcal{S}(E)$ -ideal defining $\mathcal{R}(E)$ is $\mathcal{A} = H_{\mathfrak{m}}^0(\mathcal{S}(E))$. Hence we intend to apply [Theorem A](#) to the symmetric algebra in this setting. As \mathcal{A} is bigraded, we aim to determine certain graded components using [Theorem A](#) and the resolutions of certain graded components of $\mathcal{S}(E)$. As $\mathcal{S}(E)$ is a complete intersection ring, such resolutions are readily available from the Koszul complex.

As the Koszul complex \mathcal{K}_\bullet is a bigraded B -resolution of $\mathcal{S}(E)$, each T -graded strand \mathcal{K}_i is a graded T -resolution of $\mathcal{S}(E)_i = \bigoplus_j \mathcal{S}(E)_{(i,j)}$. Hence from [Theorem A](#) we have isomorphisms of T -modules

$$\mathcal{A}_{\delta-i} \cong \text{Hom}_T(\mathcal{S}(E)_i, T(-\tau)) \cong \ker \sigma_1^*(-\tau)$$

for $0 \leq i \leq \delta$, where σ_1 is the first map of \mathcal{K}_i . Hence one needs only to understand certain dual maps in order to understand the T -modules $\mathcal{A}_{\delta-i}$. In particular, we may apply the tools developed by Kim and Mukundan in [22] to describe the kernel of this dual map when the rank of $\mathcal{S}(E)_i$ is small enough. We apply these techniques in a handful of settings, the first of which is modules which nonlinear presentation.

We remark that there is a significant amount of literature for Rees rings of ideals and modules with *linear* presentation (see e.g. [7, 24, 26, 27, 28, 29, 32, 34, 40, 41]), i.e. the entries in a presentation matrix φ are all linear forms. Additionally, there has been recent progress for ideals and modules with *almost* linear presentation [2, 6, 9], i.e. the entries of such a matrix φ are all linear, except for one column with entries of a higher degree. In this direction, our first application of [Theorem A](#) concerns Rees algebras of ideals and modules with *almost almost* linear presentation, namely when a presentation matrix φ consists of linear entries, except for two columns of quadrics. The main result of [Section 4](#) is as follows.

Theorem B. *Let $R = k[x_1, \dots, x_d]$ with $d \geq 3$ and let E be a torsion-free R -module with rank e and projective dimension one, minimally generated by $\mu(E) = d + e$ elements and presented by matrix φ . Assume that $\varphi = [\varphi' \mid \varphi'']$ where φ' has $d - 2$ columns with linear entries and φ'' consist of two columns of quadrics. Assume that $I_1(\varphi) = (x_1, \dots, x_d)$ and E satisfies G_d . The defining ideal \mathcal{A} of $\mathcal{R}(E)$ is generated as follows.*

- (i) \mathcal{A}_2 is minimally generated by one equation of bidegree $(2, d)$.
- (ii) If $\text{ht } I(\sigma_1) \geq 3$ then \mathcal{A}_1 is minimally generated by d equations of bidegree $(1, 2d - 2)$.
- (iii) If $\text{ht } I(\rho_1) \geq 2$ then \mathcal{A}_0 is minimally generated by one equation of bidegree $(0, 4d - 4)$.

Here σ_1 and ρ_1 are the first maps in the graded strands \mathcal{K}_1 and \mathcal{K}_2 , resolving $\mathcal{S}(E)_1$ and $\mathcal{S}(E)_2$ respectively.

Whereas [Theorem B](#) does not say what the equations of \mathcal{A} are, it does provide certain criteria for how many generators there are, as well as their bidegrees.

As a sequel to [Theorem B](#), we pursue a seemingly unrelated situation, namely Rees rings of ideals and modules over a *complete intersection ring* R . Measures must be taken so that [Theorem A](#) can be applied to $\mathcal{S}(E)$, as R is not a polynomial ring. However, doing so reveals how similar this setting is to that of [Theorem B](#). A similar observation was made for Rees rings of ideals and modules in *hypersurface rings* in [40], relating them to Rees rings of ideals and modules with almost linear presentation [2, 6].

Theorem C. *Let $S = k[x_1, \dots, x_{d+2}]$ with $d \geq 2$ and let $R = S/(f, g)$ where f, g is a S -regular sequence of homogeneous polynomials with $\deg f = \deg g = 2$. Let E be a torsion-free R -module with rank e and projective dimension one, minimally generated by $\mu(E) = d + e$ elements, and presented by matrix φ . Assume that*

φ consists of linear entries, E satisfies G_d , and $I_1(\varphi) = (x_1, \dots, x_{d+2})R$. The defining ideal \mathcal{A} of $\mathcal{R}(E)$ is generated as follows.

- (i) \mathcal{A}_2 is minimally generated by one equation of bidegree $(2, d)$.
- (ii) If $\text{ht } I(\sigma_1) \geq 3$ then \mathcal{A}_1 is minimally generated by $d + 2$ equations of bidegree $(1, 2d)$.
- (iii) If $\text{ht } I(\rho_1) \geq 2$ then \mathcal{A}_0 is minimally generated by one equation of bidegree $(0, 4d)$.

Here σ_1 and ρ_1 are the first maps in the graded strands \mathcal{K}_1 and \mathcal{K}_2 , resolving $\mathcal{S}(E)_1$ and $\mathcal{S}(E)_2$ respectively.

As a final application of [Theorem A](#), we consider a specific module in the setting of [Theorem C](#), namely the module of Kähler differentials $\Omega_{R/k}$, of a two-dimensional complete intersection ring R defined by quadrics. The main result of [Section 6](#) is as follows.

Theorem D. *Let $S = k[x_1, x_2, x_3, x_4]$ for k a field of characteristic zero and let f, g be a homogeneous regular sequence of quadrics such that $R = S/(f, g)$ is normal. Let $\Omega_{R/k}$ denote the module of differentials of R over k . The defining ideal \mathcal{A} of $\mathcal{R}(\Omega_{R/k})$ is generated as follows.*

- (i) \mathcal{A}_2 is minimally generated by one equation of bidegree $(2, 2)$.
- (ii) \mathcal{A}_1 is minimally generated by four equations of bidegree $(1, 4)$.
- (iii) \mathcal{A}_0 is minimally generated by one equation of bidegree $(0, n)$, for some positive integer n .

As mentioned, this is a consequence of [Theorem C](#), with many of the required conditions on the module $\Omega_{R/k}$ being satisfied by conditions on the ring R , the normality in particular.

We now briefly describe how this paper is organized. In [Section 2](#) we introduce much of the preliminary material on Rees algebras of modules required for the scope of this paper. Moreover, we recall the construction of the so-called Buchsbaum-Eisenbud multipliers associated to an acyclic complex. In [Section 3](#) we introduce the duality theorem and many of the tools used for the duration of the paper. As our first application, in [Section 4](#), we consider Rees algebras of modules with nonlinear presentation, expanding upon ideas in [\[2, 6\]](#). In [Section 5](#), we consider Rees algebras of modules over complete intersection rings, expanding upon the work in [\[40\]](#). Finally, in [Section 6](#) we apply the main result of [Section 5](#) to the defining ideal of the Rees algebra of the module of Kähler differentials of a complete intersection ring defined by quadrics. We conclude this paper by posing some open questions and directions for future work.

2. PRELIMINARIES

We now provide the preliminary material required for this paper. We begin by reviewing symmetric algebras and Rees rings of modules. We also recall certain acyclicity criteria and certain constructions associated to an acyclic complex.

2.1. Rees rings of modules. In this paper, we consider Rees algebras of ideals and modules, hence we proceed as generally as possible, opting for the latter notion. Let R denote a Noetherian ring and let E be a finitely generated R -module of rank $e \geq 1$. Writing $E = Ra_1 + \dots + Ra_n$ for a generating set of E , there is a natural homogeneous epimorphism of R -algebras

$$R[y_1, \dots, y_n] \longrightarrow \mathcal{S}(E)$$

given by sending $y_i \mapsto a_i \in [\mathcal{S}(E)]_1$, where $\mathcal{S}(E)$ denotes the symmetric algebra of E . Writing \mathcal{L} for the kernel of this map, we have an induced isomorphism

$$\mathcal{S}(E) \cong R[y_1, \dots, y_n]/\mathcal{L}.$$

Definition 2.1. The *Rees algebra* of E is the quotient $\mathcal{R}(E) = \mathcal{S}(E)/\tau_R(\mathcal{S}(E))$, where $\tau_R(\mathcal{S}(E))$ denotes the R -torsion submodule of $\mathcal{S}(E)$.

With this, the map above induces a second homogeneous epimorphism

$$R[y_1, \dots, y_n] \longrightarrow \mathcal{R}(E)$$

obtained by further factoring the R -torsion of $\mathcal{S}(E)$. Letting \mathcal{J} denote the kernel of this map, there is an isomorphism

$$\mathcal{R}(E) \cong R[y_1, \dots, y_n]/\mathcal{J}$$

as well. The ideals $\mathcal{L}, \mathcal{J} \subseteq R[y_1, \dots, y_n]$ are the *defining ideals* of $\mathcal{S}(E)$ and $\mathcal{R}(E)$, respectively.

Whereas the generators of \mathcal{J} are typically enigmatic and difficult to describe, the generators of \mathcal{L} are readily available from a presentation of the module E . Indeed, if $R^m \xrightarrow{\varphi} R^n \rightarrow E \rightarrow 0$ is such a presentation, then \mathcal{L} is generated by the linear forms ℓ_1, \dots, ℓ_m where

$$[\ell_1 \dots \ell_m] = [y_1 \dots y_n] \cdot \varphi.$$

From the prior discussion, one has the containment $\mathcal{L} \subseteq \mathcal{J}$. If $\mathcal{L} = \mathcal{J}$, or equivalently $\mathcal{S}(E) \cong \mathcal{R}(E)$, then E is said to be of *linear type*.

Typically one is most interested when E is not of linear type, and so $\mathcal{S}(E)$ and $\mathcal{R}(E)$ differ. To measure this difference, one employs the $\mathcal{S}(E)$ -ideal $\mathcal{A} = \mathcal{J}/\mathcal{L}$, as it fits into the short exact sequence

$$0 \rightarrow \mathcal{A} \rightarrow \mathcal{S}(E) \rightarrow \mathcal{R}(E) \rightarrow 0.$$

Moreover, as there is an isomorphism $\mathcal{R}(E) \cong \mathcal{S}(E)/\mathcal{A}$, one commonly refers to \mathcal{A} as a defining ideal of $\mathcal{R}(E)$ as well. The ideal \mathcal{A} tends to be simpler than \mathcal{J} and is more easily related to homological constructions, providing more insight.

Definition 2.2. For s a positive integer, the R -module E is said to satisfy G_s if $\mu(E_{\mathfrak{p}}) \leq \dim R_{\mathfrak{p}} + e - 1$ for every $\mathfrak{p} \in \text{Spec}(R)$ with $1 \leq \dim R_{\mathfrak{p}} \leq s - 1$. The module E is said to satisfy G_{∞} if it satisfies G_s for all s .

Here $\mu(E_{\mathfrak{p}})$ denotes the minimal number of generators of $E_{\mathfrak{p}}$ and, as before, e denotes the rank of E . It is often convenient to rewrite this condition in terms of Fitting ideals. Recall that for $R^m \xrightarrow{\varphi} R^n \rightarrow E \rightarrow 0$ a presentation of E as before, the i^{th} Fitting ideal is $\text{Fitt}_i(E) = I_{n-i}(\varphi)$, the ideal of $(n-i) \times (n-i)$ sized minors of φ . From [11, 20.6], it follows that E satisfies G_s if $\text{ht } \text{Fitt}_i(E) \geq i - e + 2$ for all $e \leq i \leq s + e - 2$. The condition G_s typically controls the complexity of the ideal \mathcal{A} and, in many cases, also dictates the prime ideals of R upon which \mathcal{A} is supported.

As mentioned, one is most interested when E is not of linear type, i.e. $\mathcal{L} \neq \mathcal{J}$ or rather $\mathcal{A} \neq 0$. As such, we seek a method to produce nontrivial generators, the Jacobian dual matrix being a lucrative source.

Definition 2.3. Let $R^m \xrightarrow{\varphi} R^n \rightarrow E \rightarrow 0$ be a presentation of E as above and let ℓ_1, \dots, ℓ_m denote the generators of \mathcal{L} once more. There exists an $r \times m$ matrix $\mathcal{B}(\varphi)$ consisting of linear entries in $R[y_1, \dots, y_n]$ such that

$$[y_1 \dots y_n] \cdot \varphi = [\ell_1 \dots \ell_m] = [x_1 \dots x_r] \cdot \mathcal{B}(\varphi)$$

where (x_1, \dots, x_r) is an ideal containing $I_1(\varphi)$, the ideal of entries of φ . We say that $\mathcal{B}(\varphi)$ is a *Jacobian dual* matrix of φ with respect to the sequence x_1, \dots, x_r .

As a consequence of Cramer's rule, one has that $I_r(\mathcal{B}(\varphi)) \subseteq \mathcal{J}$, hence $\mathcal{B}(\varphi)$ is a source of higher-degree equations defining $\mathcal{R}(E)$. Note that $\mathcal{B}(\varphi)$ depends on the sequence x_1, \dots, x_r and, even for a fixed sequence, $\mathcal{B}(\varphi)$ is not unique in general. However, if $R = k[x_1, \dots, x_d]$ and the entries of φ are linear, then there is a unique Jacobian dual matrix $\mathcal{B}(\varphi)$, with respect to x_1, \dots, x_d [30, p. 47]. Moreover, in this case, the entries of $\mathcal{B}(\varphi)$ belong to the subring $k[y_1, \dots, y_n]$.

Definition 2.4. If R is a local ring with maximal ideal \mathfrak{m} and residue field k , the *special fiber ring* of E is

$$\mathcal{F}(E) = \mathcal{R}(E) \otimes_R k \cong \mathcal{R}(E)/\mathfrak{m}\mathcal{R}(E).$$

Its Krull dimension is called the *analytic spread* of E and is denoted by $\ell(E) = \dim \mathcal{F}(E)$.

2.2. Acyclicity and free resolutions. One of the themes of this paper is to study certain modules through their resolutions. As such, we briefly recall necessary and sufficient criteria for a complex to be acyclic. Moreover, we also review certain constructions associated to an acyclic complex. For now, assume that R is any commutative ring and

$$C_{\bullet} : \quad 0 \longrightarrow F_n \xrightarrow{\sigma_n} \dots \xrightarrow{\sigma_2} F_1 \xrightarrow{\sigma_1} F_0 \quad (2.1)$$

is a complex of free R -modules. For each free module write $f_k = \text{rank } F_k$. Similarly, for each map write $r_k = \text{rank } \sigma_k$, where $\text{rank } \sigma_k = \max\{t \mid I_t(\sigma_k) \neq 0\}$. As before, $I_t(\sigma_k)$ denotes the ideal of $t \times t$ minors of σ_k , after choosing bases and identifying σ_k with a matrix. Whereas a matrix representation of σ_k does depend on the choice of bases, the ideal of minors $I_t(\sigma_k)$ does not. As a convention, set $I(\sigma_k) = I_{r_k}(\sigma_k)$, the ideal of rank-sized minors.

We now recall the acyclicity criterion of Buchsbaum and Eisenbud [4].

Theorem 2.5 ([4, Cor. 1]). *The complex C in (2.1) is acyclic if and only if for $k = 1, \dots, n$, one has*

- (i) $f_k = r_k + r_{k+1}$ and
- (ii) either $\text{grade } I(\sigma_k) \geq k$ or $I(\sigma_k) = R$.

Recall that any map of free R -modules $\sigma : F \rightarrow G$ induces a map of exterior powers $\bigwedge^t \sigma : \bigwedge^t F \rightarrow \bigwedge^t G$ for any t . Once bases for F and G have been chosen and a corresponding matrix representation of σ obtained, the entries of the matrix representation of $\bigwedge^t \sigma$ are precisely the signed $t \times t$ minors of φ [11, A2.3]. With this, we recall the notion of Buchsbaum-Eisenbud multipliers [5] associated to an acyclic complex.

Theorem 2.6 ([5, 3.1]). *Suppose that the conditions of Theorem 2.5 are met and the complex C in (2.1) is acyclic. For $k = 1, \dots, n$, there exists a unique R -homomorphism $a_k : R \rightarrow \bigwedge^{r_k} F_{k-1}$ such that*

- (i) $a_n = \bigwedge^{r_n} \sigma_n : R = \bigwedge^{r_n} F_n \rightarrow \bigwedge^{r_n} F_{n-1}$.
- (ii) For all $k < n$, the diagram

$$\begin{array}{ccc} \bigwedge^{r_k} F_k & \xrightarrow{\bigwedge^{r_k} \sigma_k} & \bigwedge^{r_k} F_{k-1} \\ & \searrow^{a_{k+1}^*} & \nearrow_{a_k} \\ & R & \end{array}$$

commutes, where $-^*$ denotes the functor $\text{Hom}_R(-, R)$.

- (iii) For all $k > 1$, one has $\sqrt{I(a_k)} = \sqrt{I(\sigma_k)}$.

Recall that if F is a free R -module of rank n , then $\bigwedge^n F$ is a free module of rank one. In other words, there exists an isomorphism $\omega : \bigwedge^n F \xrightarrow{\sim} R$ called an *orientation* of F . With this, it follows that the forms $\bigwedge^i F \otimes \bigwedge^{n-i} F \xrightarrow{\wedge} \bigwedge^n F \xrightarrow{\omega} R$ induce isomorphisms $\omega_i : \bigwedge^i F \rightarrow \text{Hom}_R(\bigwedge^{n-i} F, R) = (\bigwedge^{n-i} F)^*$, where \wedge is the usual multiplication in the exterior algebra.

3. BIGRADED COMPLETE INTERSECTIONS AND DUALITY

We now introduce the main tool used throughout the paper, the duality theorem and its application to symmetric algebras and the defining ideal \mathcal{A} of the Rees algebra. As noted, this is not a novel approach to the matter (see e.g. [22, 25]), however we aim to extend these techniques to previously untouched settings; see Sections 5 and 6, in particular. Our setting for much of this section is as follows.

Setting 3.1. Let $R = k[x_1, \dots, x_d]$, $T = k[y_1, \dots, y_n]$, and $B = R \otimes_k T$. Endowing B with the bigrading given by bideg $x_i = (1, 0)$ and bideg $y_i = (0, 1)$, let $\underline{f} = f_1, \dots, f_d \subseteq \mathfrak{m} = (x_1, \dots, x_d)$ be a B -regular sequence of bihomogeneous elements. Write bideg $f_i = (\alpha_i, \beta_i)$ and let $\mathcal{S} = B/\mathcal{I}$, where $\mathcal{I} = (\underline{f})$.

With the assumptions of Setting 3.1, we may now state the duality theorem and its consequences. We then describe how it may be applied to Rees algebras of modules and recall the techniques developed in [22].

3.1. Duality and complete intersections. With the general setting above established, we now introduce our main tool, the duality theorem. We note that this is only a slight reformulation of the original duality theorem of Jouanolou [21, Section 3.6], written in the bigraded setting. We present its statement and give a proof differing from Jouanolou's original arguments. Instead, we provide a proof in the style of Kustin, Polini, and Ulrich [25, 2.4], using local cohomology and its interaction with Koszul homology.

Theorem 3.2 ([21, Section 3.6], [25, 2.4]). *With the assumptions of Setting 3.1, there is a bihomogeneous isomorphism of B -modules*

$$H_{\mathfrak{m}}^0(\mathcal{S}) \cong \text{Hom}_T(\mathcal{S}, T)(-\delta, -\tau)$$

where $\delta = (\sum_{i=1}^d \alpha_i) - d$, $\tau = \sum_{i=1}^d \beta_i$, and $H_{\mathfrak{m}}^0(\mathcal{S})$ is the zeroth local cohomology module of \mathcal{S} with respect to $\mathfrak{m} = (x_1, \dots, x_d)$.

Proof. Set $m = \sum_{i=1}^d \alpha_i$ and notice that, since $f_1, \dots, f_d \subseteq \mathfrak{m}$, we have $\alpha_i \geq 1$ for all i . Hence $m \geq d$ and so δ is nonnegative. By the self-duality of the Koszul complex $\mathcal{K}_{\bullet}(\underline{f}; B)$, there is an isomorphism of complexes of bigraded B -modules

$$\text{Hom}_T(\mathcal{K}_{\bullet}(\underline{f}; B), T) \cong \mathcal{K}_{\bullet}(\underline{f}; \text{Hom}_T(B, T))[d](m, \tau) \tag{3.1}$$

where $[-]$ denotes homological shift and $(-, -)$ denotes the usual bidegree shift.

Recall that the Koszul complex $\mathcal{K}_\bullet(\underline{f}; B)$ is acyclic as \underline{f} is a B -regular sequence. As its zeroth homology is $B/\mathcal{I} = \mathcal{S}$, we have the following commutative diagram

$$\mathcal{K}_\bullet(\underline{f}; B) : \quad 0 \rightarrow K_d \rightarrow K_{d-1} \xrightarrow{\quad} K_{d-2} \rightarrow \cdots \rightarrow K_2 \xrightarrow{\quad} K_1 \xrightarrow{\quad} K_0 \rightarrow \mathcal{S} \rightarrow 0$$

where the top row is exact with the C_i its syzygies. With this, we create a series of short exact sequences

$$\begin{aligned} 0 &\longrightarrow C_1 \longrightarrow K_0 \longrightarrow \mathcal{S} \longrightarrow 0 \\ 0 &\longrightarrow C_2 \longrightarrow K_1 \longrightarrow C_1 \longrightarrow 0 \\ &\quad \quad \quad \vdots \quad \quad \quad \vdots \quad \quad \quad \vdots \\ 0 &\longrightarrow C_{d-1} \longrightarrow K_{d-2} \longrightarrow C_{d-2} \longrightarrow 0 \\ 0 &\longrightarrow K_d \longrightarrow K_{d-1} \longrightarrow C_{d-1} \longrightarrow 0. \end{aligned} \tag{3.2}$$

Notice that $\text{grad } \mathfrak{m}B = d$ and the K_i are B -free. Hence by applying the functor $H_{\mathfrak{m}}^\bullet(-)$ to the first $d-1$ sequences in (3.2) and observing the consequential vanishing, we obtain the isomorphisms

$$H_{\mathfrak{m}}^0(\mathcal{S}) \cong H_{\mathfrak{m}}^1(C_1) \cong H_{\mathfrak{m}}^2(C_2) \cong \cdots \cong H_{\mathfrak{m}}^{d-1}(C_{d-1}).$$

Now applying $H_{\mathfrak{m}}^\bullet(-)$ to the last short exact sequence in (3.2), one has

$$0 = H_{\mathfrak{m}}^{d-1}(K_{d-1}) \longrightarrow H_{\mathfrak{m}}^{d-1}(C_{d-1}) \longrightarrow H_{\mathfrak{m}}^d(K_d) \xrightarrow{\partial} H_{\mathfrak{m}}^d(K_{d-1})$$

where ∂ is precisely the d^{th} differential in $\mathcal{K}_\bullet(\underline{f}; H_{\mathfrak{m}}^d(B))$, as K_d and K_{d-1} are B -free and local cohomology commutes with direct sums. Hence

$$H_{\mathfrak{m}}^0(\mathcal{S}) \cong H_{\mathfrak{m}}^{d-1}(C_{d-1}) \cong H_d(\mathcal{K}_\bullet(\underline{f}; H_{\mathfrak{m}}^d(B))).$$

However, by Lemma 3.3 and the isomorphism in (3.1) we have

$$\begin{aligned} H_d(\mathcal{K}_\bullet(\underline{f}; H_{\mathfrak{m}}^d(B))) &\cong H_d(\mathcal{K}_\bullet(\underline{f}; \text{Hom}_T(B, T)(d, 0))) \\ &\cong H_0(\text{Hom}_T(\mathcal{K}_\bullet(\underline{f}; B), T))(-m, -\tau)(d, 0) \\ &\cong \text{Hom}_T(\mathcal{S}, T)(-m + d, -\tau) \end{aligned}$$

and the claim follows since $-m + d = -\delta$. \square

Lemma 3.3. *There is a bihomogeneous isomorphism $H_{\mathfrak{m}}^d(B) \cong \text{Hom}_T(B, T)(d, 0)$.*

Proof. Recall that $B = R \otimes_k T$ where $R = k[x_1, \dots, x_d]$ and $T = k[y_1, \dots, y_n]$. With this and noting that B is R -flat, we have

$$H_{\mathfrak{m}}^d(B) \cong H_{\mathfrak{m}}^d(R \otimes_R B) \cong H_{\mathfrak{m}}^d(R) \otimes_R B \cong H_{\mathfrak{m}}^d(R) \otimes_k T.$$

Moreover, as \mathfrak{m} is the homogeneous maximal ideal of R , by Serre duality [17, III.7.1], we have a graded isomorphism of R -modules $H_{\mathfrak{m}}^d(R) \cong \text{Hom}_k(R, k)(d)$. Hence by flatness once more, and adjusting to the bigrading of B , we have

$$H_{\mathfrak{m}}^d(R) \otimes_k T \cong \text{Hom}_k(R, k)(d) \otimes_k T \cong \text{Hom}_T(B, T)(d, 0)$$

and the claim follows. \square

Although the rings and modules here are bigraded, it is often convenient to fix one component of the bigrading to obtain singly graded modules. We introduce notation to permit this.

Notation 3.4. For any bigraded B -module $M = \bigoplus_{i,j} M_{(i,j)}$, write M_i to denote the graded T -module

$$M_i = M_{(i,*)} = \bigoplus_j M_{(i,j)}.$$

With this convention, we obtain the following consequence of [Theorem 3.2](#).

Corollary 3.5. *With the assumptions of [Setting 3.1](#), write $A = H_m^0(\mathcal{S})$.*

- (a) *There is a graded isomorphism of T -modules $A_i \cong \text{Hom}_T(\mathcal{S}_{\delta-i}, T(-\tau))$ for all i .*
- (b) *For all $i < 0$ and $i > \delta$, we have $A_i = 0$.*
- (c) *There is a graded isomorphism $A_\delta \cong T(-\tau)$.*

Proof. Assertion (a) follows immediately from [Theorem 3.2](#). Part (b) then follows from (a) noting that \mathcal{S} is nonnegatively graded as a T -module. Lastly, assertion (c) follows from (a) as $\mathcal{S}_0 \cong T$. \square

Notice that by [Corollary 3.5](#), A_δ is a cyclic T -module generated by an equation of bidegree (δ, τ) . Moreover, we can actually produce such a generator.

Proposition 3.6. *The T -module A_δ is generated as $A_\delta = \langle \det \mathcal{B} \rangle$ where \mathcal{B} is a $d \times d$ matrix \mathcal{B} consisting of bihomogeneous entries in B such that $[f_1 \dots f_d] = [x_1 \dots x_d] \cdot \mathcal{B}$.*

Proof. Writing $\mathcal{I} = (f_1, \dots, f_d)$ and $\text{bideg } f_i = (\alpha_i, \beta_i)$ as before, recall that $\alpha_i \geq 1$ for all i . Notice that the entries in column i of \mathcal{B} have bidegree $(\alpha_i - 1, \beta_i)$, hence it follows that $\det \mathcal{B}$ is a form of bidegree (δ, τ) so long as $\det \mathcal{B} \neq 0$. Moreover, the assertion will follow once $\det \mathcal{B}$ has been shown to be nonzero modulo \mathcal{I} .

We note that the matrix \mathcal{B} is not unique in general, however the ideal $\mathcal{I} + (\det \mathcal{B})$ is unique. Indeed, since \mathcal{I} and (x_1, \dots, x_d) are generated by regular sequences, it is well-known (see e.g. [\[3, 2.3.10\]](#)) that

$$\mathcal{I} : (x_1, \dots, x_d) = \mathcal{I} + (\det \mathcal{B}).$$

Notice that $\mathcal{I} : (x_1, \dots, x_d) \neq \mathcal{I}$, as A is nonzero by [Corollary 3.5\(c\)](#). Hence it follows that $\det \mathcal{B} \notin \mathcal{I}$, and so $\det \mathcal{B}$ is nonzero modulo \mathcal{I} . \square

As noted in [Section 2](#), such a transition matrix \mathcal{B} is called a Jacobian dual when f_1, \dots, f_d are the linear equations defining a symmetric algebra. This will be the case in [Section 4](#), however in [Section 5](#) we will take such a matrix \mathcal{B} to be a *modified* Jacobian dual, as introduced in [\[40\]](#).

3.2. Certain complexes. With [Corollary 3.5](#), it suffices to study the T -duals of graded components of the complete intersection \mathcal{S} , in order to understand the T -graded components of A . Moreover, information regarding the module $\text{Hom}_T(\mathcal{S}_{\delta-i}, T)$ can be obtained from a free resolution of $\mathcal{S}_{\delta-i}$. As \mathcal{S} is a complete intersection ring, it is resolved by the Koszul complex. Hence we may take its T -graded strands to resolve the graded components of \mathcal{S} .

Remark 3.7. With the assumptions of [Setting 3.1](#), recall that the Koszul complex $\mathcal{K}_\bullet(\underline{f}; B)$ is a B -resolution of $\mathcal{S} = B/(\underline{f})$. Moreover, as $\text{bideg } f_i = (\alpha_i, \beta_i)$, the bigraded Koszul complex is

$$\mathcal{K}_\bullet(\underline{f}; B) : 0 \longrightarrow B(-\sum \alpha_i, -\sum \beta_i) \longrightarrow \dots \longrightarrow \bigoplus^d B(-\alpha_i, -\beta_i) \longrightarrow B. \quad (3.3)$$

With this, we may obtain graded T -resolutions for each \mathcal{S}_t . Indeed, writing \mathcal{K}_t to denote the graded strand of [\(3.3\)](#) in degree t , the graded free resolution of \mathcal{S}_t is

$$\mathcal{K}_t : 0 \longrightarrow F_m \xrightarrow{\sigma_m} F_{m-1} \xrightarrow{\sigma_{m-1}} \dots \xrightarrow{\sigma_1} F_0 \quad (3.4)$$

for some $m \leq d$, where

$$F_i = \bigoplus_{1 \leq j_1 \leq \dots \leq j_i \leq d} T(-(\beta_{j_1} + \dots + \beta_{j_i})) \binom{t - (\alpha_{j_1} + \dots + \alpha_{j_i}) + d - 1}{d - i} \quad (3.5)$$

for $0 \leq i \leq m$, noting that $\binom{n}{k} = 0$ when $0 \leq n < k$.

With the resolution of \mathcal{S}_t in [\(3.4\)](#), notice that by [Corollary 3.5](#) we have $A_{\delta-t} \cong \ker \sigma_1^*(-\tau)$. With this, we recall the complex of Kim and Mukundan from [\[22\]](#). We present its statement slightly differently than as originally written, noting that the construction of this complex shows that it may be applied in the setting of *any* bigraded complete intersection ring, as in [Setting 3.1](#).

Theorem 3.8 ([22, Theorems 13-16]). *With the assumptions of [Setting 3.1](#) and \mathcal{K}_t the T -resolution of \mathcal{S}_t as in [Remark 3.7](#), there exists a complex of graded T -modules*

$$\bigwedge^{f_0-r_1-1} F_0 \otimes T(-s) \xrightarrow{\omega \wedge (\text{id} \otimes a_1)} F_0^* \xrightarrow{\sigma_1^*} F_1^*.$$

Here $\omega : \bigwedge^{f_0-1} F_0 \xrightarrow{\sim} F_0^*$ is the isomorphism induced by an orientation on F_0 , \wedge is multiplication within the exterior algebra, and $a_1 : T(-s) \rightarrow \bigwedge^{r_1} F_0$ is the map from [Theorem 2.6](#) with degree shift s making it a homogeneous map of free T -modules. Moreover, we have the following.

- (a) If $\text{rank } \mathcal{S}_t = 1$, then this complex is exact if and only if $\text{grade } I(a_1) \geq 2$.
- (b) Assume $\text{rank } \mathcal{S}_t = 2$ and consider the following statements:
 - (i) \mathcal{S}_t satisfies Serre's condition S_2 .
 - (ii) $\text{grade } I(\sigma_t) \geq t + 2$ for $t = 1, \dots, m$.
 - (iii) The complex is exact.
Then (i) implies (ii) which implies (iii).

In particular, if the complex in [Theorem 3.8](#) is exact, then $A_{\delta-t}$ is generated by $\binom{f_0}{f_0-r_1-1}$ elements of bidegree $(\delta-t, s+\tau)$. Note that, by [Theorem 2.6](#), it also suffices to show that $\text{ht } I(\sigma_1) \geq 2$ in [Theorem 3.8\(a\)](#), which is often more convenient.

Remark 3.9. The degree shift s for $a_1 : T(-s) \rightarrow \bigwedge^{r_1} F_0$ in [Theorem 3.8](#) may be computed by adapting [Theorem 2.6](#) to graded free modules. However, as the construction of these maps is recursive, this value must be computed iteratively. While it is possible to produce a formula for s with the assumptions of [Setting 3.1](#) using the data in (3.4) and (3.5), we forgo this option as it will be impractical for our purposes. Indeed, the complexes in the proceeding sections will be remarkably short, hence this value will be easily computed. We refer the curious reader to [22, Rem. 12] for the general procedure and a formula for s when $\beta_1 = \dots = \beta_d = 1$ in [Setting 3.1](#).

3.3. Complete intersections and symmetric algebras. We intend to use [Corollary 3.5](#) and [Theorem 3.8](#) when \mathcal{S} is the symmetric algebra $\mathcal{S}(E)$ of a module E and \mathcal{A} is the defining ideal of the Rees ring $\mathcal{R}(E)$, as a quotient of $\mathcal{S}(E)$. However, we must then ensure that $\mathcal{S}(E)$ is a complete intersection ring and that \mathcal{A} coincides with the local cohomology module $H_{\mathfrak{m}}^0(\mathcal{S}(E))$. We provide sufficient criteria for both phenomena to occur.

Proposition 3.10. *Let R be a Cohen-Macaulay local ring of dimension d with maximal ideal \mathfrak{m} and let E be a torsion-free R -module with $\text{rank } e \geq 1$. Assume that E satisfies G_d , $\mu(E) = d + e$, and E has projective dimension one.*

- (a) Writing $B = R[y_1, \dots, y_{d+e}]$ and \mathcal{L} the ideal such that $\mathcal{S}(E) \cong B/\mathcal{L}$, the ideal \mathcal{L} is generated by a B -regular sequence.
- (b) Write \mathcal{J} and \mathcal{A} for the ideals such that $\mathcal{R}(E) \cong B/\mathcal{J} \cong \mathcal{S}(E)/\mathcal{A}$. The module E is of linear type on the punctured spectrum of R , i.e. $\mathcal{A}_{\mathfrak{p}} = 0$ for all $\mathfrak{p} \in \text{Spec}(R) \setminus \{\mathfrak{m}\}$. In particular, $\mathcal{A} \cong H_{\mathfrak{m}}^0(\mathcal{S}(E))$ and $\mathcal{J} = \mathcal{L} : \mathfrak{m}^\infty$.

Proof. Since E has projective dimension 1 and $\mu(E) = d + e$, it has a minimal free resolution of the form

$$0 \longrightarrow R^d \xrightarrow{\varphi} R^{d+e} \longrightarrow E \longrightarrow 0. \quad (3.6)$$

Hence \mathcal{L} is generated by the d linear forms $\mathcal{L} = (\ell_1, \dots, \ell_d)$ where $[\ell_1 \dots, \ell_d] = [y_1 \dots y_{d+e}] \cdot \varphi$. Thus it suffices to show that $\dim \mathcal{S}(E) = d + e$ to conclude that $\text{ht } \mathcal{L} = d$, and so ℓ_1, \dots, ℓ_d is a B -regular sequence. By the Huneke-Rossi formula [20], we have

$$\dim \mathcal{S}(E) = \sup\{\mu(E_{\mathfrak{p}}) + \dim R/\mathfrak{p} \mid \mathfrak{p} \in \text{Spec}(R)\}$$

and we claim that the maximum value is $d + e$. If $\mathfrak{p} \neq \mathfrak{m}$, from the condition G_d it follows that

$$\mu(E_{\mathfrak{p}}) + \dim R/\mathfrak{p} \leq \text{ht } \mathfrak{p} + e - 1 + \dim R/\mathfrak{p} = d + e - 1.$$

Now taking $\mathfrak{p} = \mathfrak{m}$, we have $\mu(E_{\mathfrak{m}}) + \dim R/\mathfrak{m} = \mu(E) + 0 = d + e$. Hence $\dim \mathcal{S}(E) = d + e$ which shows (a).

To prove (b), it is well known that one has the containment $H_{\mathfrak{m}}^0(\mathcal{S}(E)) \subseteq \mathcal{A}$. Notice that since E satisfies G_d , each localization $E_{\mathfrak{p}}$ satisfies G_∞ for any non-maximal prime R -ideal \mathfrak{p} . Hence by [1, Prop. 3 and 4] it

follows that $E_{\mathfrak{p}}$ is of linear type for any such prime, and so $\mathcal{A}_{\mathfrak{p}} = 0$ for all $\mathfrak{p} \neq \mathfrak{m}$. Hence \mathcal{A} is annihilated by some power of \mathfrak{m} and so $\mathcal{A} \subseteq H_{\mathfrak{m}}^0(\mathcal{S}(E))$ as well. As $\mathcal{A} = \mathcal{J}/\mathcal{L}$, this also shows that \mathcal{J} is the claimed saturation. \square

We remark that, although it is stated in the local setting, [Proposition 3.10](#) also applies to a graded module E over $R = k[x_1, \dots, x_d]$ with $\mathfrak{m} = (x_1, \dots, x_d)$ the unique homogeneous maximal ideal, as in [Setting 3.1](#). We also note that, although the condition $\mu(E) = d + e$ might seem restrictive, if $\mu(E) < d + e$ with the remaining assumptions, then E satisfies G_{∞} and is hence of linear type [[1](#), Prop. 3 and 4]. Thus this is the first nontrivial setting, in regard to the number of generators $\mu(E)$.

Remark 3.11. In the setting of [Proposition 3.10](#), if $e = 1$ then E is isomorphic to a perfect ideal of grade two. Indeed, as E is torsion-free with rank $e = 1$, it is isomorphic to an R -ideal I . Since $E \cong I$ has projective dimension 1, it follows from [\(3.6\)](#) that it is presented by a $(d + 1) \times d$ matrix. Moreover, the condition G_d implies that $\text{ht } I_d(\varphi) = \text{ht Fitt}_1(I) \geq 2$ and the claim follows from the Hilbert-Burch theorem [[11](#), 20.15].

4. REES ALGEBRAS OF MODULES WITH NONLINEAR PRESENTATION

In this section, we study Rees algebras of modules with particular restrictions on their presentations. If $R = k[x_1, \dots, x_d]$ and E is a graded R -module homogeneously generated in a single degree, notice that E has a presentation matrix with entries concentrated in a single degree within each column. As these degrees dictate the bidegrees of the equations defining the symmetric algebra $\mathcal{S}(E)$, one attempts to study the Rees ring $\mathcal{R}(E)$ with certain restrictions on them. As such, we recall the notion of the *type* of a matrix from [[22](#)]. A $m \times n$ matrix is said to be of type (d_1, \dots, d_n) if it consists of entries of degree d_i in column i for $1 \leq i \leq n$.

As noted in the introduction, there has been an extensive study of Rees rings of ideals and modules with linear presentation, i.e. those with presentation matrices of type $(1, \dots, 1)$. Moreover, there has also been success for ideals and modules with *almost* linear presentation, namely those with presentation matrices of type $(1, \dots, 1, m)$ for some $m \geq 1$. To continue this study, in this section we consider modules with presentation of type $(1, \dots, 1, 2, 2)$ and their Rees rings.

Setting 4.1. Let $R = k[x_1, \dots, x_d]$ with $d \geq 3$ and let E be a torsion-free R -module with rank e and projective dimension one, minimally generated by $\mu(E) = d + e$ elements. Letting

$$0 \longrightarrow R^d \xrightarrow{\varphi} R^{d+e} \longrightarrow E \longrightarrow 0$$

denote a minimal free resolution of E , assume that φ is of type $(1, \dots, 1, 2, 2)$, i.e. $\varphi = [\varphi' \mid \varphi'']$ where φ' has $d - 2$ columns with linear entries and φ'' consists of two columns of quadrics. Further assume that $I_1(\varphi) = (x_1, \dots, x_d)$ and E satisfies the condition G_d .

As noted, one reduces to the study of Rees rings of a notable class of ideals when the rank of E is $e = 1$.

Remark 4.2. Following [Remark 3.11](#), if $e = 1$ then E is isomorphic to an R -ideal I , which is perfect of grade two. Moreover, from the type of φ and the Hilbert-Burch theorem [[11](#), 20.15], it follows that I is minimally generated by $\mu(I) = d + 1$ homogeneous forms of degree $d + 2$.

Regardless, we consider the more general situation in [Setting 4.1](#) throughout. We briefly recall the notation required to begin our treatment of the symmetric algebra $\mathcal{S}(E)$.

Notation 4.3. With the assumptions of [Setting 4.1](#), write $T = k[y_1, \dots, y_{d+e}]$ and let $B = R \otimes_k T \cong R[y_1, \dots, y_{d+e}]$ with the bigrading given by $\text{bideg } x_i = (1, 0)$ and $\text{bideg } y_i = (0, 1)$. Recall that the symmetric algebra is $\mathcal{S}(E) \cong B/\mathcal{L}$ where $[\ell_1 \dots \ell_d] = [y_1 \dots y_{d+e}] \cdot \varphi$ and $\mathcal{L} = (\ell_1, \dots, \ell_d)$. Write \mathcal{A} and \mathcal{J} to denote the ideals defining the Rees algebra $\mathcal{R}(E)$ as quotients of $\mathcal{S}(E)$ and B respectively, as in [Section 2](#).

By [Proposition 3.10](#), ℓ_1, \dots, ℓ_d is a B -regular sequence and $\mathcal{A} = H_{\mathfrak{m}}^0(\mathcal{S}(E))$ where $\mathfrak{m} = (x_1, \dots, x_d)$, hence we may apply [Corollary 3.5](#) and [Theorem 3.8](#). Moreover, with the conditions on φ , the bidegrees of $\ell_1, \dots, \ell_{d-2}$ (recall $d \geq 3$) are $(1, 1)$, while the bidegrees of ℓ_{d-1} and ℓ_d are $(2, 1)$. With this, and adopting the grading scheme in [Notation 3.4](#) throughout, we may produce T -resolutions of the graded components of $\mathcal{S}(E)$ from the bigraded Koszul complex of ℓ_1, \dots, ℓ_d following [Remark 3.7](#).

Proposition 4.4. *With R and E as in [Setting 4.1](#) and δ and τ as in [Theorem 3.2](#), we have $\delta = 2$, $\tau = d$, and $\mathcal{S}(E)_0$ is free of rank one. Moreover, $\text{rank } \mathcal{S}(E)_1 = 2$ and $\text{rank } \mathcal{S}(E)_2 = 1$.*

Proof. The first statement is clear. As for the second assertion, we construct the graded T -resolutions of $\mathcal{S}(E)_1$ and $\mathcal{S}(E)_2$ following [Remark 3.7](#). Their ranks are then computed using [Theorem 2.5](#).

(a) For the resolution of $\mathcal{S}(E)_1$, we have

$$0 \longrightarrow \bigoplus^{d-2} T(-1)^{\binom{d-1}{d-1}} \xrightarrow{\sigma} T^{\binom{d}{d-1}} \longrightarrow \mathcal{S}(E)_1 \longrightarrow 0. \quad (4.1)$$

Thus $\text{rank } \mathcal{S}(E)_1 = d - (d - 2) = 2$.

(b) For the resolution of $\mathcal{S}(E)_2$, we must consider two cases, as the projective dimension of $\mathcal{S}(E)_2$ depends on d .

(i) If $d = 3$, the graded T -resolution of $\mathcal{S}(E)_2$ is

$$0 \longrightarrow T(-1)^{\binom{3}{2}} \oplus \bigoplus^2 T(-1)^{\binom{2}{2}} \xrightarrow{\rho} T^{\binom{4}{2}} \longrightarrow \mathcal{S}(E)_2 \longrightarrow 0 \quad (4.2)$$

and so $\text{rank } \mathcal{S}(E)_2 = 6 - 3 - 2 = 1$.

(ii) If $d \geq 4$. The graded T -resolution of $\mathcal{S}(E)_2$ is

$$0 \longrightarrow \bigoplus^{\binom{d-2}{2}} T(-2)^{\binom{d-1}{d-1}} \xrightarrow{\gamma} \bigoplus^{d-2} T(-1)^{\binom{d}{d-1}} \oplus \bigoplus^2 T(-1)^{\binom{d-1}{d-1}} \xrightarrow{\rho} T^{\binom{d+1}{d-1}} \longrightarrow \mathcal{S}(E)_2 \longrightarrow 0 \quad (4.3)$$

and so $\text{rank } \mathcal{S}(E)_2 = \binom{d+1}{d-1} - d(d-2) - 2 + \binom{d-2}{2} = 1$.

Thus in either case, we have $\text{rank } \mathcal{S}(E)_2 = 1$. \square

Recall that (4.1) is the graded strand in degree 1 of the bigraded Koszul complex of ℓ_1, \dots, ℓ_d . Taking the standard monomial bases, ordered lexicographically, for the free T -modules involved, σ may be realized as the $d \times (d-2)$ matrix with linear entries of T , such that $[\ell_1, \dots, \ell_{d-2}] = [x_1 \dots x_d] \cdot \sigma$. In other words, σ is precisely the Jacobian dual of φ' , $\sigma = \mathcal{B}(\varphi')$, where φ' is the submatrix of φ as in [Setting 4.1](#).

Remark 4.5. With the map σ in (4.1), we have $\text{ht } I(\sigma) \geq 2$.

Proof. Recall that σ may be taken as the transition matrix such that $[\ell_1, \dots, \ell_{d-2}] = [x_1 \dots x_d] \cdot \sigma$. As σ consists of entries in T , it follows that $(\ell_1, \dots, \ell_{d-2})$ is the ideal defining the symmetric algebra of the T -module $\text{coker } \sigma = \mathcal{S}(E)_1$. Thus by [19, 6.6, 6.8] it suffices to show that this ring is a domain.

Notice that as σ consists of entries in T and φ' consists of entries in R , there is an isomorphism of symmetric algebras $\mathcal{S}_T(\mathcal{S}(E)_1) \cong \mathcal{S}_R(M)$, where $M = \text{coker } \varphi'$. Moreover, it can easily be seen that since E has projective dimension 1 and satisfies G_d , M does as well. However, note that $\text{rank } M = e + 2$ and $\mu(M) = d + e < d + (e + 2)$, hence M satisfies G_∞ and is thus of linear type [1, Prop. 3 and 4]. Thus $\mathcal{S}_R(M)$ is a domain and so $\mathcal{S}_T(\mathcal{S}(E)_1)$ is as well. \square

We note that $\text{ht } I(\sigma) \leq 3$ by [10, Thm. 1] and if this maximum codimension is attained, then the complex in [Theorem 3.8](#) is exact. Whereas we cannot ensure this maximum height is achieved, we remark that this is only a *sufficient* condition for exactness of this complex.

Theorem 4.6. *With E as in [Setting 4.1](#) and \mathcal{A} the ideal defining $\mathcal{R}(E)$ as a quotient of $\mathcal{S}(E)$, we have the following.*

- (a) \mathcal{A}_2 is generated as $\mathcal{A}_2 = \langle \det \mathcal{B}(\varphi) \rangle$ where $\mathcal{B}(\varphi)$ is a Jacobian dual of φ , with respect to x_1, \dots, x_d . Moreover, this is an equation of bidegree $(2, d)$.
- (b) If $\text{ht } I(\sigma) = 3$, then \mathcal{A}_1 is minimally generated by d equations of bidegree $(1, 2d - 2)$.
- (c) \mathcal{A}_0 is minimally generated by one equation. If $\text{ht } I(\rho) \geq 2$, then this generator has bidegree $(0, 4d - 4)$.

Proof. Part (a) follows from [Corollary 3.5](#) and [Proposition 3.6](#). For the first part of (c), notice that $\mathcal{A}_0 \subseteq \mathcal{S}(E)_0 = T$, and so \mathcal{A}_0 is a T -ideal. Recall that the special fiber ring of E is $\mathcal{F}(E) \cong \mathcal{R}(E)/\mathfrak{m}\mathcal{R}(E)$ and it follows that \mathcal{A}_0 defines $\mathcal{F}(E)$ as a quotient of $T = k[y_1, \dots, y_{d+e}]$, i.e. $\mathcal{F}(E) \cong T/\mathcal{A}_0$. Moreover, it is well known that E has maximal analytic spread, i.e. $\ell(E) = d + e - 1$, in this setting, which may be seen from [32, 3.10] and [34, 4.3]. As this is the dimension of $\mathcal{F}(E)$, a domain, it follows that \mathcal{A}_0 is a principal T -ideal.

Part (b) and the second assertion of (c) follow from [Theorem 3.8](#) and [Proposition 4.4](#), once the degree shift of the map a_1 has been computed for resolutions (4.1) – (4.3).

(a) For sequence (4.1), notice that $a_1 = \bigwedge^{d-2} \sigma$, following [Theorem 2.6](#). Hence

$$a_1 : \bigwedge^{d-2} (T(-1)^{d-2}) \cong T(-(d-2)) \longrightarrow \bigwedge^{d-2} (T^d)$$

and so $s = d - 2$.

(b) For the T -resolution of $S(E)_2$, we must consider the two cases from [Proposition 4.4](#).

(i) If $d = 3$, consider the T -resolution of $S(E)_2$ in (4.2). By [Theorem 2.6](#) we have $a_1 = \bigwedge^5 \rho$, hence

$$a_1 : \bigwedge^5 (T(-1)^5) \cong T(-5) \longrightarrow \bigwedge^5 (T^6)$$

and so the degree shift of a_1 is $s = 5$.

(ii) If $d \geq 4$, consider the T -resolution of $S(E)_2$ in (4.3). As this is a resolution of length two, we begin with a_2 as the construction of the maps in [Theorem 2.6](#) is recursive. Note that

$$a_2 = \bigwedge \gamma : \bigwedge^{\binom{d-2}{2}} T(-2)^{\binom{d-2}{2}} \longrightarrow \bigwedge^{\binom{d-2}{2}} T(-1)^{d(d-2)+2}$$

and so the entries in a matrix representation of a_2 have degree $\binom{d-2}{2}$. By [Theorem 2.6](#) we have the commutative diagram

$$\begin{array}{ccc} \bigwedge^r T(-1)^{d(d-2)+2} & \xrightarrow{\bigwedge^r \rho} & \bigwedge^r T^{\binom{d+1}{d-1}} \\ & \searrow^{a_2^*} & \nearrow_{a_1} \\ & T(-s) & \end{array}$$

where $r = \text{rank } \rho = d(d-2) + 2 - \binom{d-2}{2}$ by [Theorem 2.5](#). Notice that the degrees of the entries in a matrix representation of $\bigwedge^r \rho$ are all r as ρ consists of linear entries. Hence

$$s = r - \binom{d-2}{2} = d(d-2) + 2 - (d-2)(d-3) = 3d - 4.$$

Thus $s = 3d - 4$ in either case.

The claim now follows from [Theorem 3.8](#), noting that $\tau = d$ by [Proposition 4.4](#). □

Remark 4.7. We note that the shapes of the T -resolutions (4.1) – (4.3) of the graded components of $S(E)$ are independent of the rank e of the module E . Hence if $e = 1$, then one achieves the same result for the defining ideal of the Rees algebra of an ideal I as in [Remark 4.2](#).

Remark 4.8. Notice that if both $\text{ht } I(\sigma) = 3$ and $\text{ht } I(\rho) \geq 2$ in [Theorem 4.6](#), then it is understood how the defining ideals \mathcal{A} and \mathcal{J} of $\mathcal{R}(E)$ are generated. For instance, the ideal \mathcal{J} is then minimally generated as

	Bidegree	Number of generators
$\ell_1, \dots, \ell_{d-2}$	(1, 1)	$d - 2$
ℓ_{d-1}, ℓ_d	(2, 1)	2
\mathcal{A}_0	(0, $4d - 4$)	1
\mathcal{A}_1	(1, $2d - 2$)	d
$\mathcal{A}_2 = \langle \det \mathcal{B}(\varphi) \rangle$	(2, d)	1

In particular, \mathcal{J} is minimally generated by $2d + 2$ elements and \mathcal{A} is minimally generated by $d + 2$ elements.

Unfortunately, it is not necessarily the case that both conditions of [Theorem 4.6](#) are satisfied. Nevertheless, we consider the number of generators above and their bidegrees to be the *predicted* forms of \mathcal{J} and \mathcal{A} . Indeed, the conditions of [Theorem 4.6](#) are frequently satisfied, and it is likely that the behavior in [Remark 4.8](#) occurs if the entries of φ are *sufficiently general*, similar to the main result of [22]. We present an example, for Rees rings of ideals, where ideals of minors of the maps in (4.1) – (4.3) fail to have the expected codimension.

Example 4.9. Let $R = \mathbb{Q}[x_1, x_2, x_3, x_4]$, let

$$\varphi = \begin{bmatrix} x_1 & 0 & x_4^2 & x_2x_3 \\ x_2 & 0 & x_1x_2 & x_3^2 \\ 0 & 0 & x_1^2 & x_2^2 \\ 0 & x_3 & x_2^2 & x_2x_3 \\ 0 & x_4 & x_4^2 & x_2^2 \end{bmatrix}$$

be a matrix as in [Setting 4.1](#), and consider the ideal $I = I_4(\varphi)$. Computations through *Macaulay2* [14] show that $\text{ht } I = 2$, hence I is perfect of grade 2 [11, 20.15], and I satisfies the condition G_4 . Thus the assumptions of [Remark 4.2](#) and [Setting 4.1](#) are met. However, the matrix $\sigma = \mathcal{B}(\varphi')$ in (4.1) is

$$\sigma = \begin{bmatrix} y_1 & 0 \\ y_2 & 0 \\ 0 & y_4 \\ 0 & y_5 \end{bmatrix}$$

and it can be seen that $\text{ht } I(\sigma) = 2$, hence the conditions of [Theorem 4.6](#) are not met. Moreover, computations through *Macaulay2* [14] show that \mathcal{A}_1 is minimally generated by two equations of bidegree (1, 5), differing from the predicted behavior in [Remark 4.8](#). However, the map ρ in (4.3) has $\text{ht } I(\rho) = 2$, and so \mathcal{A}_0 is generated by one equation of bidegree (0, 12), following [Remark 4.7](#) and [Theorem 4.6](#).

Question 4.10. Noting that \mathcal{A}_2 is generated by the determinant of a Jacobian dual matrix, it is curious if the remaining equations of \mathcal{A} can be described by related constructions. This was shown to be the case when the presentation φ is almost linear, noted for perfect ideals of grade two ($e = 1$) in [2, 9] and more generally for modules of projective dimension one in [6].

5. REES ALGEBRAS OF MODULES OVER COMPLETE INTERSECTION RINGS

In this section, we study Rees rings of R -modules when R is a complete intersection ring. As noted in the introduction, there is much geometric motivation to study Rees rings when the ground ring R is not a polynomial ring. We note that there has been recent success in the study of Rees algebras of ideals and modules over *hypersurface rings* [40, 41]. Expanding upon this, we consider modules over complete intersection rings and their Rees algebras. Although this seems like a drastically different setting than that of the previous section, we proceed in a similar manner.

Setting 5.1. Let $S = k[x_1, \dots, x_{d+2}]$ and let $R = S/(f, g)$ where f, g is a S -regular sequence of homogeneous polynomials with $\deg f = \deg g = 2$. Let E be a torsion-free R -module with rank e and projective dimension one, minimally generated by $\mu(E) = d + e$ elements. Let

$$0 \longrightarrow R^d \xrightarrow{\varphi} R^{d+e} \longrightarrow E \longrightarrow 0$$

be a minimal free resolution of E and assume that φ consists of linear entries in R . Assume that E satisfies G_d and $I_1(\varphi) = \overline{(x_1, \dots, x_{d+2})}$, where $\overline{}$ denotes images modulo (f, g) .

As before, we recover the case of Rees algebras of perfect ideals of grade two.

Remark 5.2. If $e = 1$ in [Setting 5.1](#), then E is isomorphic to a perfect R -ideal I of grade two. Moreover, as φ consists of linear entries, by the Hilbert-Burch theorem [11, 20.15] I is minimally generated by $\mu(I) = d + 1$ homogeneous forms of degree d .

As before, we begin by considering the symmetric algebra of E . However, we first introduce notation amenable to the setting of [Corollary 3.5](#).

Notation 5.3. With the assumptions of [Setting 5.1](#), write $T = k[y_1, \dots, y_{d+e}]$ and let $B = S \otimes_k T \cong S[y_1, \dots, y_{d+e}]$ with the bigrading given by $\text{bideg } x_i = (1, 0)$ and $\text{bideg } y_i = (0, 1)$. Writing $\overline{}$ to denote images modulo (f, g) , let ψ be the $(d + e) \times d$ matrix with linear entries in S such that $\overline{\psi} = \varphi$. Let \mathcal{L} denote the B -ideal $\mathcal{L} = (\ell_1, \dots, \ell_d, f, g)$ where $[\ell_1, \dots, \ell_d] = [y_1 \dots y_{d+e}] \cdot \psi$.

Clearly the matrix ψ exists, and also it is unique as it has linear entries and $\deg f = \deg g = 2$. Notice that $\overline{\mathcal{L}} = (\overline{\ell_1}, \dots, \overline{\ell_d})$ where $[\overline{\ell_1} \dots \overline{\ell_d}] = [y_1 \dots y_{d+e}] \cdot \varphi$. Hence $\overline{\mathcal{L}}$ is the defining ideal of $\mathcal{S}(E)$, as a quotient of $R[y_1, \dots, y_{d+e}]$. By [Proposition 3.10](#), we have that $\overline{\ell_1}, \dots, \overline{\ell_d}$ is a regular sequence in this ring.

Since R is not a polynomial ring, we cannot apply [Theorem 3.2](#) with $R[y_1, \dots, y_{d+e}]$ and $\overline{\mathcal{L}}$. However, note that

$$\mathcal{S}(E) = R[y_1, \dots, y_{d+e}]/\overline{\mathcal{L}} \cong B/\mathcal{L}$$

and it follows that \mathcal{L} is generated by a B -regular sequence. With this, $\mathcal{S}(E)$ is a complete intersection ring and can be realized as a quotient of a bigraded polynomial ring, hence we use this description of $\mathcal{S}(E) \cong B/\mathcal{L}$ with [Theorem 3.2](#) and [Corollary 3.5](#).

Remark 5.4. As mentioned, we proceed with the isomorphism $\mathcal{S}(E) \cong B/\mathcal{L}$, using \mathcal{L} as the defining ideal of $\mathcal{S}(E)$, in this sense. Likewise, we may also update the ideals defining $\mathcal{R}(E)$. Differing from the convention in [Section 2](#), write \mathcal{J} to denote the kernel of the composition

$$B = S[y_1, \dots, y_{d+e}] \longrightarrow R[y_1, \dots, y_{d+e}] \longrightarrow \mathcal{R}(E)$$

where the first map quotients by (f, g) and the second is the natural map. Writing $\mathcal{A} = \mathcal{J}/\mathcal{L}$, from [Proposition 3.10](#) it follows that $\mathcal{J} = \mathcal{L} : \mathfrak{m}^\infty$ and so $\mathcal{A} = H_{\mathfrak{m}}^0(\mathcal{S}(E))$, where $\mathfrak{m} = (x_1, \dots, x_{d+2})$.

The adjustment of the defining ideals \mathcal{L} , \mathcal{A} , and \mathcal{J} is not completely novel and has also been used in [\[40, 41\]](#). Recall that the generators of $\mathcal{L} = (\ell_1, \dots, \ell_d, f, g)$ form a B -regular sequence. Moreover, ℓ_1, \dots, ℓ_d have bidegree $(1, 1)$ and f and g have bidegree $(2, 0)$. Hence one can see the similarity of $\mathcal{S}(E)$ in this setting to the symmetric algebra of [Section 4](#).

With the grading convention in [Notation 3.4](#), we may produce T -resolutions of the graded components of $\mathcal{S}(E) \cong B/\mathcal{L}$ from the bigraded Koszul complex of $\ell_1, \dots, \ell_d, f, g$ following [Remark 3.7](#).

Proposition 5.5. *Adopt the assumptions of [Setting 5.1](#) and write $\mathcal{S}(E) \cong B/\mathcal{L}$ as above. With δ and τ as in [Theorem 3.2](#), we have $\delta = 2$, $\tau = d$, and $\mathcal{S}(E)_0$ is free of rank one. Additionally $\text{rank } \mathcal{S}(E)_1 = 2$ and $\text{rank } \mathcal{S}(E)_2 = 1$.*

Proof. The first statement is clear. To verify the claim on the T -modules $\mathcal{S}(E)_1$ and $\mathcal{S}(E)_2$, we construct their graded T -resolutions following [Remark 3.7](#) and apply [Theorem 2.5](#) to compute their ranks.

(a) For the resolution of $\mathcal{S}(E)_1$, we have

$$0 \longrightarrow \bigoplus^d T(-1)^{\binom{d+1}{d+1}} \xrightarrow{\sigma} T^{\binom{d+2}{d+1}} \longrightarrow \mathcal{S}(E)_1 \longrightarrow 0 \quad (5.1)$$

and so $\text{rank } \mathcal{S}(E)_1 = (d+2) - d = 2$.

(b) For the resolution of $\mathcal{S}(E)_2$, we have

$$0 \longrightarrow \bigoplus^{\binom{d}{2}} T(-2)^{\binom{d+1}{d+1}} \xrightarrow{\gamma} \bigoplus^d T(-1)^{\binom{d+2}{d+1}} \oplus \bigoplus^2 T^{\binom{d+1}{d+1}} \xrightarrow{\rho} T^{\binom{d+3}{d+1}} \longrightarrow \mathcal{S}(E)_2 \longrightarrow 0 \quad (5.2)$$

and so $\text{rank } \mathcal{S}(E)_2 = \binom{d+3}{d+1} - d(d+2) - 2 + \binom{d}{2} = 1$. \square

Recall that [\(5.1\)](#) is the graded strand of the bigraded Koszul complex of $\ell_1, \dots, \ell_d, f, g$ in degree 1. Choosing the standard monomial bases with the lexicographic order, σ may be realized as the $(d+2) \times d$ matrix with linear entries in T such that $[\ell_1, \dots, \ell_d] = [x_1 \dots x_{d+2}] \cdot \sigma$. Hence σ may be identified with the Jacobian dual of ψ , $\sigma = \mathcal{B}(\psi)$, where ψ is the matrix in [Notation 5.3](#).

Remark 5.6. With the matrix σ as in [\(5.1\)](#), we have $\text{ht } I(\sigma) \geq 2$.

Proof. As noted, σ can be realized as the transition matrix with $[\ell_1 \dots \ell_d] = [x_1 \dots x_{d+2}] \cdot \sigma$. As σ consists of entries in T , we see that (ℓ_1, \dots, ℓ_d) is the defining ideal of the symmetric algebra of $\text{coker } \sigma = \mathcal{S}(E)_1$. Thus by [\[19, 6.6, 6.8\]](#) it is enough to show that this ring is a domain. As ψ consists of linear entries in S and σ consists of entries in T , there is an isomorphism of symmetric algebras $\mathcal{S}_S(M) \cong \mathcal{S}_T(\mathcal{S}(E)_1)$ where $M = \text{coker } \psi$. Thus it suffices to show that $\mathcal{S}(M)$ is a domain, or rather that M is an S -module of linear type.

Notice that $d = \text{rank } \varphi \leq \text{rank } \psi \leq d$, and so $\text{rank } \psi = d$. Thus the complex of S -modules

$$0 \longrightarrow S^d \xrightarrow{\psi} S^{d+e} \longrightarrow M \longrightarrow 0$$

is exact by [Theorem 2.5](#), and so M is an S -module of projective dimension one. Notice that M satisfies G_d since E satisfies this condition. Indeed, $\text{ht Fitt}_i(M) \geq \text{ht Fitt}_i(E) \geq i - e + 2$ for all $e \leq i \leq d + e - 2$, as $\bar{\psi} = \varphi$ and height can only decrease modulo (f, g) . Additionally, notice that $\text{Fitt}_{d+e-1}(M) = I_1(\psi) = (x_1, \dots, x_{d+2})$, which has height $d + 2 \geq d + 1$. Since $\mu(M) = d + e$, for any $i \geq d + e$ we see that $\text{Fitt}_i(M)$ is the unit ideal. Thus M satisfies G_∞ , and is hence of linear type by [[1](#), Prop. 3 and 4]. \square

Before we state the main result of this section, we recall the notion of a *modified* Jacobian dual matrix introduced in [[40](#)]. This matrix is an extension of the usual Jacobian dual and will inevitably be the matrix in [Proposition 3.6](#), in this setting.

Definition 5.7. With the assumptions of [Setting 5.1](#), let ψ be as in [Notation 5.3](#). We define a *modified Jacobian dual* of ψ to be a $(d + 2) \times (d + 2)$ matrix $\mathcal{B} = [\mathcal{B}(\psi) \mid \partial f \mid \partial g]$ where $\mathcal{B}(\psi)$ is the Jacobian dual of ψ , with respect to x_1, \dots, x_{d+2} , and ∂f and ∂g are columns with linear entries in S such that $f = [x_1 \dots x_{d+2}] \cdot \partial f$ and $g = [x_1 \dots x_{d+2}] \cdot \partial g$.

Note that whereas the Jacobian dual $\mathcal{B}(\psi)$ is unique, a modified Jacobian dual \mathcal{B} is not, as there are many choices for the columns ∂f and ∂g . However, there is a natural candidate for this matrix.

Remark 5.8. As the notation suggests, there is a natural choice for the columns ∂f and ∂g using differentials, if k is a field with $\text{char } k \neq 2$. Indeed, these columns may be chosen to consist of (unit multiples of) the partial derivatives of f and g , using the well-known Euler formula

$$2f = \sum_{i=1}^{d+2} \frac{\partial f}{\partial x_i} x_i, \quad 2g = \sum_{i=1}^{d+2} \frac{\partial g}{\partial x_i} x_i, \quad (5.3)$$

as f and g are homogeneous of degree 2.

Theorem 5.9. *With E as in [Setting 5.1](#) and \mathcal{A} the defining ideal of $\mathcal{R}(E)$ in the setting of [Remark 5.4](#), we have the following.*

- (a) \mathcal{A}_2 is generated as $\mathcal{A}_2 = \langle \det \mathcal{B} \rangle$ where \mathcal{B} is a modified Jacobian dual of ψ . Moreover, this is an equation of bidegree $(2, d)$.
- (b) If $\text{ht } I(\sigma) = 3$, then \mathcal{A}_1 is minimally generated by d equations of bidegree $(1, 2d)$.
- (c) \mathcal{A}_0 is minimally generated by one equation. If $\text{ht } I(\rho) \geq 2$, then this generator has bidegree $(0, 4d)$.

Proof. Part (a) follows from [Corollary 3.5](#) and [Proposition 3.6](#), noting that $[\ell_1 \dots \ell_d f g] = [x_1 \dots x_{d+2}] \cdot \mathcal{B}$. For the first assertion of (c), the same argument as in the proof of [Theorem 4.6](#) shows that E has analytic spread $\ell(E) = d + e - 1$. As this is the dimension of the special fiber ring $\mathcal{F}(E)$, it follows that \mathcal{A}_0 is a cyclic T -module.

Notice that σ is a $(d + 2) \times d$ matrix, hence $\text{ht } I(\sigma) \leq 3$ by [[10](#), Thm. 1]. Thus (b) and the second assertion of (c) follow from [Theorem 3.8](#) along with [Proposition 5.5](#), once the degree shift of the map a_1 has been computed for resolutions ([5.1](#)) and ([5.2](#)).

- (a) For the T -resolution of $S(E)_1$ in ([5.1](#)), notice that $a_1 = \bigwedge^d \sigma$, following [Theorem 2.6](#). Hence

$$a_1 : \bigwedge^d (T(-1)^d) \cong T(-d) \longrightarrow \bigwedge^d (T^{d+2})$$

and so $s = d$.

- (b) For the T -resolution of $S(E)_2$ in ([5.2](#)), we begin with a_2 as this resolution has length two and the construction of the maps in [Theorem 2.6](#) is recursive. Note that

$$a_2 = \bigwedge^{\binom{d}{2}} \gamma : \bigwedge^{\binom{d}{2}} T(-2)^{\binom{d}{2}} \longrightarrow \bigwedge^{\binom{d}{2}} (T(-1)^{d(d+2)} \oplus T^2)$$

and it can be seen that the entries in a matrix representation of a_2 have degree $\binom{d}{2}$. By [Theorem 2.6](#) we have the commutative diagram

$$\begin{array}{ccc} \bigwedge^r (T(-1)^{d(d+2)} \oplus T^2) & \xrightarrow{\bigwedge^r \rho} & \bigwedge^r T^{\binom{d+3}{d+1}} \\ & \searrow^{a_2^*} & \nearrow^{a_1} \\ & T(-s) & \end{array}$$

where $r = \text{rank } \rho = d(d+2) + 2 - \binom{d}{2}$ by [Theorem 2.5](#). Notice that the entries in a matrix representation of $\bigwedge^r \rho$ have degree $r - 2 = d(d+2) - \binom{d}{2}$ as ρ consists of linear entries and two columns with units. Hence $s = (r - 2) - \binom{d}{2} = d(d+2) - d(d-1) = 3d$.

The claim now follows from [Theorem 3.8](#), noting that $\tau = d$ by [Proposition 5.5](#). \square

Remark 5.10. Similar to the observation made in [Section 4](#), resolutions (5.1) and (5.2) do not depend on the rank of E . Thus if $e = 1$, one has the same result for the Rees algebra of an ideal I as in [Remark 5.2](#).

Remark 5.11. If the conditions of [Theorem 5.9](#) are met, then it is understood how the defining ideals \mathcal{A} and \mathcal{J} of [Remark 5.4](#) are generated. Indeed, if both $\text{ht } I(\sigma) = 3$ and $\text{ht } I(\rho) \geq 2$ in [Theorem 5.9](#), then \mathcal{J} is generated as

	Bidegree	Number of generators
ℓ_1, \dots, ℓ_d	$(1, 1)$	d
f, g	$(2, 0)$	2
\mathcal{A}_0	$(0, 4d)$	1
\mathcal{A}_1	$(1, 2d)$	$d + 2$
$\mathcal{A}_2 = \langle \det \mathcal{B} \rangle$	$(2, d)$	1

where \mathcal{B} is a modified Jacobian dual of ψ . In particular, \mathcal{J} is minimally generated by $2d + 6$ elements and \mathcal{A} is minimally generated by $d + 4$ elements.

As before, we refer to the behavior in [Remark 5.11](#) as the predicted form of \mathcal{A} and \mathcal{J} in this setting. However, we present an example to illustrate that this behavior is not always observed, even in the case of ideals. We note that this phenomenon is frequently observed, and it is suspected that the behavior in [Remark 5.11](#) always occurs if the entries of φ , and also f and g , are sufficiently general.

Example 5.12. Let $S = \mathbb{Q}[x_1, x_2, x_3, x_4]$ and let $R = S/(x_1^2, x_2^2)$. Consider the matrix φ with linear entries in R and its corresponding matrix ψ with linear entries in S ,

$$\varphi = \begin{bmatrix} \overline{x_1} & \overline{x_3} \\ \overline{x_3} & \overline{x_4} \\ \overline{x_4} & \overline{x_2} \end{bmatrix} \quad \text{and} \quad \psi = \begin{bmatrix} x_1 & x_3 \\ x_3 & x_4 \\ x_4 & x_2 \end{bmatrix}$$

and let $I = I_2(\varphi)$. Computations through *Macaulay2* [14] show that $\text{ht } I = 2$, hence I is perfect of grade 2 [11, 20.15] and satisfies G_2 . Thus the assumptions of [Remark 5.2](#) and [Setting 5.1](#) are met.

Choosing the standard monomial bases for the free modules involved, a matrix representative of the map ρ in the resolution (5.2) of $\mathcal{S}(E)_2$ is

$$\rho = \begin{bmatrix} 1 & 0 & y_1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & y_1 & 0 & 0 & y_3 & 0 & 0 & 0 \\ 0 & 0 & y_2 & 0 & y_1 & 0 & y_1 & 0 & 0 & 0 \\ 0 & 0 & y_3 & 0 & 0 & y_1 & y_2 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & y_3 & 0 & 0 \\ 0 & 0 & 0 & y_2 & 0 & 0 & 0 & y_1 & y_3 & 0 \\ 0 & 0 & 0 & y_3 & 0 & 0 & 0 & y_2 & 0 & y_3 \\ 0 & 0 & 0 & 0 & y_2 & 0 & 0 & 0 & y_1 & 0 \\ 0 & 0 & 0 & 0 & y_3 & y_2 & 0 & 0 & y_2 & y_1 \\ 0 & 0 & 0 & 0 & 0 & y_3 & 0 & 0 & 0 & y_2 \end{bmatrix}$$

and [Theorem 2.5](#) and [\(5.2\)](#) show that $\text{rank } \rho = 9$. However, $\text{ht } I(\rho) = 1$ and so the conditions of [Theorem 5.9](#) are not satisfied. Further computations show that \mathcal{A}_0 is generated by an equation of bidegree $(0, 6)$, differing from the predicted behavior in [Remark 5.11](#). However, with σ as in [\(5.1\)](#) we have $\text{ht } I(\sigma) = 3$ in this example, and so \mathcal{A}_1 is generated by four equations of bidegree $(1, 4)$ by [Remark 5.10](#) and [Theorem 5.9](#).

One possibility for why [Example 5.12](#) failed to produce the expected behavior in [Remark 5.11](#) is that the ring R is not reduced. It is curious if one obtains the predicted behavior if R is reduced or a domain. We investigate this possibility in the proceeding section, under the assumption that R is a *normal* complete intersection domain.

Question 5.13. Noting that \mathcal{A}_2 is generated by the determinant of a modified Jacobian dual \mathcal{B} , one might ask if the remaining equations of \mathcal{A} can be described by similar constructions. This was shown to be the case for Rees rings of perfect ideals of grade two and modules of projective dimension one over hypersurface rings in [\[40\]](#). This was also shown to be the case in [\[41\]](#) for Rees algebras of perfect Gorenstein ideals of grade three in hypersurface rings.

6. APPLICATIONS TO TANGENT ALGEBRAS

In this final section, we discuss applications of the results from [Section 5](#) to the Rees ring of the module of Kähler differentials $\Omega_{R/k}$ of a complete intersection ring R defined by quadrics. Whereas this situation possesses great significance, certain aspects of our setting are limited due to the relationship between $\Omega_{R/k}$ and the ring R . However, other aspects appear more natural as many conditions on $\Omega_{R/k}$ are implied by assumptions on R .

We refer the reader to [\[23\]](#) for the necessary background material on the module of differentials. We require few technical aspects of the subject for our treatment, hence much of the preliminary material is omitted. Our setting throughout is as follows.

Setting 6.1. Let $S = k[x_1, x_2, x_3, x_4]$ for k a field of characteristic zero and let $R = S/(f, g)$ where f, g is a S -regular sequence of homogeneous polynomials with $\deg f = \deg g = 2$. Assume that R is a normal ring and let $\Omega_{R/k}$ denote the module of Kähler differentials of R over k .

We note that many of the results presented here hold if k is a perfect field with $\text{char } k \neq 2$. For simplicity however, we consider the situation above.

Remark 6.2. Notice that, since R is a normal complete intersection ring, the ideal (f, g) is prime by Hartshorne's connectedness lemma [\[16\]](#). Hence R is a domain (see also [\[39, 6.15 – 6.17\]](#)).

With the properties of the ring R in [Setting 6.1](#), we note that $\Omega_{R/k}$ satisfies the assumptions of [Section 5](#). Indeed, many of the required conditions are consequences of the well-known Jacobian criterion [\[23, 7.2\]](#).

Proposition 6.3. *With the conditions of [Setting 6.1](#), $\Omega_{R/k}$ satisfies the assumptions of [Setting 5.1](#).*

Proof. Since R is a complete intersection ring, it is well known that $\Omega_{R/k}$ is an R -module of projective dimension at most one [\[13\]](#) and hence exactly one as R is not regular [\[23, 7.2\]](#). Thus there is a short exact sequence

$$0 \longrightarrow R^2 \xrightarrow{\bar{\Theta}} R^4 \longrightarrow \Omega_{R/k} \longrightarrow 0 \tag{6.1}$$

where Θ is the transpose of the Jacobian matrix of f and g and $\bar{\cdot}$ denotes images modulo (f, g) [\[23, 4.19\]](#). Since R is normal, it satisfies Serre's condition R_1 and is hence regular in codimension 1. In particular, it follows that $\text{ht Fitt}_2(\Omega_{R/k}) \geq 2$, or equivalently that $\Omega_{R/k}$ is locally free in codimension 1 [\[23, 7.2\]](#), or also equivalently that $\Omega_{R/k}$ satisfies G_2 . With this, it also follows that $\Omega_{R/k}$ is torsion-free.

For the rank and number of generators of $\Omega_{R/k}$, we refer to [\[23, ch. 7\]](#). With the conditions of [Setting 6.1](#), the rank of $\Omega_{R/k}$ agrees with the dimension of R , $\text{rank } \Omega_{R/k} = \dim R = 2$. Moreover, the minimal number of generators of $\Omega_{R/k}$ agrees with the embedding dimension of R , $\mu(\Omega_{R/k}) = \text{edim } R = 4$. With this, we see that $\mu(\Omega_{R/k}) = \dim R + \text{rank } \Omega_{R/k}$.

It remains to be shown that $I_1(\bar{\Theta}) = \overline{(x_1, \dots, x_4)}$, the homogeneous maximal ideal of R . Notice that $\bar{\Theta}$ consists of linear entries in R since f and g are quadrics. Recall that $\text{ht Fitt}_2(\Omega_{R/k}) \geq 2$ and note that this ideal is $I_2(\bar{\Theta})$, hence the containment $I_2(\bar{\Theta}) \subseteq I_1(\bar{\Theta})$ shows that $\text{ht } I_1(\bar{\Theta}) = 2$. Thus the claim will follow

once it has been shown that $I_1(\overline{\Theta})$ is a prime ideal. However, this follows as $I_1(\Theta)$ is prime, as it is generated by linear forms, and since $f, g \in I_1(\Theta)$, which can be seen from the Euler formula (5.3). \square

The ring R is assumed to be two-dimensional in Setting 6.1 in order to satisfy the generation condition in Setting 5.1. Indeed, the proof of Proposition 6.3 shows that we require $\text{edim}(R) = 2 \dim R$. However, if R is a quotient by a regular sequence of length two, this can only happen if $\dim R = 2$.

As before, we intend to study the defining ideal \mathcal{A} of the Rees ring $\mathcal{R}(\Omega_{R/k})$ using the symmetric algebra $\mathcal{S}(\Omega_{R/k})$. As in Section 5, we introduce the notation necessary for its treatment. We adopt conventions similar to those in Notation 5.3 and Remark 5.4, and restate them for clarity.

Notation 6.4. With the assumptions of Setting 6.1, write $T = k[y_1, \dots, y_4]$ and let $B = S \otimes_k T \cong S[y_1, \dots, y_4]$. As in the proof of Proposition 6.3, let Θ denote the Jacobian matrix of f and g and let \mathcal{L} denote the B -ideal $\mathcal{L} = (\ell_1, \ell_2, f, g)$, where $[\ell_1 \ell_2] = [y_1 \dots y_4] \cdot \Theta$.

As before, this allows us to update the notion of the defining ideals of $\mathcal{S}(\Omega_{R/k})$ and $\mathcal{R}(\Omega_{R/k})$. Following Remark 5.4, we have $\mathcal{S}(\Omega_{R/k}) \cong B/\mathcal{L}$ and we take \mathcal{J} to be the kernel of the composition

$$B \longrightarrow R[y_1, \dots, y_4] \longrightarrow \mathcal{R}(\Omega_{R/k})$$

so that $\mathcal{R}(\Omega_{R/k}) \cong B/\mathcal{J}$. Following Proposition 3.10 and Remark 5.4, we have $\mathcal{J} = \mathcal{L} : \mathfrak{m}^\infty$ and also $\mathcal{A} = \mathcal{J}/\mathcal{L} \cong H_{\mathfrak{m}}^0(\mathcal{S}(\Omega_{R/k}))$, where $\mathfrak{m} = (x_1, x_2, x_3, x_4)$.

Remark 6.5. Recall that the generators of \mathcal{L} form a B -regular sequence, and so $\mathcal{S}(\Omega_{R/k}) \cong B/\mathcal{L}$ is a complete intersection ring. Moreover, by Proposition 5.5 we have $\delta = \tau = 2$ and we need only analyze the T -resolutions of $\mathcal{S}(\Omega_{R/k})_1$ and $\mathcal{S}(\Omega_{R/k})_2$. From (5.1) and (5.2) we have the following.

(a) The T -resolution of $\mathcal{S}(\Omega_{R/k})_1$ is

$$0 \longrightarrow T(-1)^2 \xrightarrow{\sigma} T^4 \longrightarrow \mathcal{S}(\Omega_{R/k})_1 \longrightarrow 0. \quad (6.2)$$

(b) The T -resolution of $\mathcal{S}(\Omega_{R/k})_2$ is

$$0 \longrightarrow T(-2) \xrightarrow{\gamma} T(-1)^8 \oplus T^2 \xrightarrow{\rho} T^{10} \longrightarrow \mathcal{S}(\Omega_{R/k})_2 \longrightarrow 0. \quad (6.3)$$

As noted prior to Remark 5.6, σ may be identified with the Jacobian dual of Θ , $\sigma = \mathcal{B}(\Theta)$, with respect to x_1, \dots, x_4 . As the terminology suggests, a Jacobian dual may be seen as a Jacobian matrix. Moreover, a particularly interesting phenomenon occurs for the Jacobian dual of a Jacobian matrix of quadrics. Before we reveal this aspect, we introduce two maps allowing one to toggle between objects over the rings S and T .

Definition 6.6. With $S = k[x_1, x_2, x_3, x_4]$ and $T = k[y_1, y_2, y_3, y_4]$ as before, let Φ denote the isomorphism $\Phi : T \xrightarrow{\sim} S$ given by $\Phi(y_i) = x_i$. Moreover, let Ψ denote its inverse $\Psi : S \xrightarrow{\sim} T$ given by $\Psi(x_i) = y_i$.

In particular, one may apply the isomorphism Φ in order to treat the maps in (6.2) and (6.3) as maps between S -modules. One may then apply its inverse Ψ to transfer any discovered information back. We provide a short lemma applying this to the map σ and offer a proof by simple calculus.

Lemma 6.7. *With σ the map in (6.2), we may take $\sigma = \mathcal{B}(\Theta)$ as a matrix representation. Additionally, we have $\mathcal{B}(\Theta) = \Psi(\Theta)$ and $\Theta = \Phi(\mathcal{B}(\Theta))$.*

Proof. The first assertion has already been verified, hence we proceed with the second. Recall that, as Θ consists of linear entries in S , $\mathcal{B}(\Theta)$ is the *unique* matrix with linear entries in T such that

$$[y_1 \dots y_4] \cdot \Theta = [x_1 \dots x_4] \cdot \mathcal{B}(\Theta). \quad (6.4)$$

Hence we need only show that

$$[y_1 \dots y_4] \cdot \Theta = [x_1 \dots x_4] \cdot \Psi(\Theta) \quad (6.5)$$

as well.

Since Θ is the Jacobian matrix of f and g with respect to x_1, \dots, x_4 , it follows that $\Psi(\Theta)$ is the Jacobian matrix of $\Psi(f)$ and $\Psi(g)$, with respect to y_1, \dots, y_4 . Thus it suffices to show that

$$\sum_{t=1}^4 \frac{\partial h}{\partial x_t} y_t = \sum_{t=1}^4 \frac{\partial(\Psi(h))}{\partial y_t} x_t \quad (6.6)$$

for any homogeneous quadric $h \in S$ to conclude the equality in (6.5). However, by linearity of differentiation, it suffices to check that (6.6) holds for any monomial $h = x_i x_j$, which is easily verified. Thus $\Psi(\Theta) = \mathcal{B}(\Theta)$ and applying the inverse Φ shows that $\Theta = \Phi(\mathcal{B}(\Theta))$ as well. \square

We remark that Lemma 6.7 requires only that the entries of Θ are linear, hence it may easily be extended. In particular, the Jacobian dual of a Jacobian matrix of quadrics is essentially the same Jacobian matrix.

Example 6.8. Let $S = \mathbb{Q}[x_1, x_2, x_3, x_4]$ and let $f = x_1^2 + x_2 x_3 + x_4^2$ and $g = x_2^2 + x_1 x_4 + x_3^2$. The Jacobian matrix Θ of f and g and its Jacobian dual $\mathcal{B}(\Theta)$ are

$$\Theta = \begin{bmatrix} 2x_1 & x_4 \\ x_3 & 2x_2 \\ x_2 & 2x_3 \\ 2x_4 & x_1 \end{bmatrix} \quad \text{and} \quad \mathcal{B}(\Theta) = \begin{bmatrix} 2y_1 & y_4 \\ y_3 & 2y_2 \\ y_2 & 2y_3 \\ 2y_4 & y_1 \end{bmatrix}$$

which can be seen from the equation $[y_1 \dots y_4] \cdot \Theta = [x_1 \dots x_4] \cdot \mathcal{B}(\Theta)$. Hence $\mathcal{B}(\Theta)$ is also the Jacobian matrix of $\Psi(f) = y_1^2 + y_2 y_3 + y_4^2$ and $\Psi(g) = y_2^2 + y_1 y_4 + y_3^2$, by Lemma 6.7.

As a consequence of Lemma 6.7, properties of σ in (6.2) can be deduced from properties of Θ , heights of ideals of minors in particular. With this, we present the main result of this section.

Theorem 6.9. *With the assumptions of Setting 6.1 and \mathcal{A} as before, we have the following.*

- (a) \mathcal{A}_2 is generated as $\mathcal{A}_2 = \langle \det \mathcal{B} \rangle$ where $\mathcal{B} = [\mathcal{B}(\Theta) \mid \Theta]$. Moreover, this equation has bidegree $(2, 2)$.
- (b) \mathcal{A}_1 is minimally generated by four equations of bidegree $(1, 4)$.
- (c) \mathcal{A}_0 is minimally generated by one equation. If $\text{ht } I(\rho) \geq 2$, then this generator has bidegree $(0, 8)$.

In particular, \mathcal{A} is generated by six elements and the ideal \mathcal{J} is generated by ten elements.

Proof. By Proposition 6.3, $\Omega_{R/k}$ satisfies the assumptions of Theorem 5.9, hence assertion (c) follows immediately. By Theorem 5.9 we have that \mathcal{A}_2 is generated by the determinant of a modified Jacobian dual of Θ . Thus part (a) follows from Proposition 3.6 and Remark 5.8, noting that 2 is a unit.

For part (b), by Theorem 5.9 it suffices to show that $\text{ht } I_2(\sigma) = 3$. Recall that Φ and Ψ are isomorphisms, hence by Lemma 6.7 we have $\text{ht } I_2(\sigma) = \text{ht } I_2(\mathcal{B}(\Theta)) = \text{ht } I_2(\Theta) \leq 3$, where the inequality follows from [10, Thm. 1]. However, by [42, 2.4] we have $\text{ht } I_2(\sigma) = \text{ht } I_2(\Theta) \geq 3$, as $R = S/(f, g)$ is a complete intersection ring satisfying Serre's condition R_1 . \square

As noted, \mathcal{A}_2 is generated by the determinant of the modified Jacobian dual $\mathcal{B} = [\mathcal{B}(\Theta) \mid \Theta]$. It is particularly interesting to note that this matrix is a *total* Jacobian of the generators of $\mathcal{L} = (\ell_1, \ell_2, f, g)$, with respect to x_1, \dots, x_4 . Moreover, recall from Lemma 6.7 that $\mathcal{B}(\Theta) = \Psi(\Theta)$, hence there is a symmetry within this matrix. For instance, as a continuation of Example 6.8, we have the following.

Example 6.10. With S and f, g as in Example 6.8, let $R = S/(f, g)$ and $\Omega_{R/k}$ its module of differentials. With \mathcal{A} as above, \mathcal{A}_2 is generated by the determinant of

$$[\mathcal{B}(\Theta) \mid \Theta] = [\Psi(\Theta) \mid \Theta] = \begin{bmatrix} 2y_1 & y_4 & 2x_1 & x_4 \\ y_3 & 2y_2 & x_3 & 2x_2 \\ y_2 & 2y_3 & x_2 & 2x_3 \\ 2y_4 & y_1 & 2x_4 & x_1 \end{bmatrix}$$

following Theorem 6.9 and Lemma 6.7.

Although the number of generators of \mathcal{A} is known in Theorem 6.9, we lack certainty on the bidegree of the generator of \mathcal{A}_0 . However, we provide the following conjecture.

Conjecture 6.11. With the assumptions of Theorem 6.9 and ρ the map in (6.3), we claim that $\text{ht } I(\rho) \geq 2$. Then by Theorem 6.9, \mathcal{A}_0 is generated by an equation of bidegree $(0, 8)$.

Similar to Lemma 6.7 and the proof of Theorem 6.9, one may apply the map Φ to the map ρ in (6.3) and investigate the codimension of $I(\Phi(\rho))$ instead. As in Lemma 6.7, the entries of $\Phi(\rho)$ can be described by partial derivatives of f and g . Computations through *Macaulay2* [14] suggest that this ideal of minors of $\Phi(\rho)$ has height exactly 2, and also provides a guess as to some of the minimal primes of this ideal. Regardless, we save further investigation and optimistic confirmation of Conjecture 6.11 for a later article.

6.1. Higher dimension. As previously noted, we considered a two-dimensional ring in [Setting 6.1](#) so that $\text{edim } R = 2 \dim R$, and so $\Omega_{R/k}$ satisfies the generation condition in [Theorem 5.9](#). It is curious as to what can be said if one maintains this condition, but allows for higher dimension d .

Setting 6.12. Let $S = k[x_1, \dots, x_{2d}]$ for $d \geq 2$ and k a field of characteristic zero. Let f_1, \dots, f_d be a homogeneous regular sequence with each $\deg f_i = 2$ and let $R = S/(f_1, \dots, f_d)$. Let $\Omega_{R/k}$ denote the module of differentials of R over k , and assume that $\Omega_{R/k}$ satisfies G_d .

Repeating the argument in [Proposition 6.3](#) and [Remark 6.2](#), notice that R is a normal domain here as $\Omega_{R/k}$ satisfies G_2 . The full condition G_d must be assumed here as is not implied from normality alone, unless $d = 2$, as in [Setting 6.1](#). However, this condition will be met if R satisfies Serre's condition R_{d-1} , i.e. if R is regular in codimension at most $d - 1$ or, in other words, if R is an *isolated singularity*.

Similar arguments to the proof of [Proposition 6.3](#) show that $\Omega_{R/k}$ satisfies the assumptions of [Proposition 3.10](#), hence $\mathcal{S}(\Omega_{R/k})$ is a complete intersection ring. Moreover, we may write $\mathcal{S}(\Omega_{R/k}) \cong B/\mathcal{L}$ where $B = S[y_1, \dots, y_{2d}]$ and $\mathcal{L} = (\ell_1, \dots, \ell_d, f_1, \dots, f_d)$ where $[\ell_1 \dots \ell_d] = [y_1 \dots y_{2d}] \cdot \Theta$. Here Θ is the transposed Jacobian matrix of f_1, \dots, f_d , as before. One may apply [Theorem 3.2](#) and [Corollary 3.5](#) since $\mathcal{S}(\Omega_{R/k}) \cong B/\mathcal{L}$ is a complete intersection ring, however one cannot use [Theorem 3.8](#) to describe \mathcal{A}_i for all i , as the ranks of the T -modules $\mathcal{S}(\Omega_{R/k})_i$ become too large.

Observation 6.13. With the assumptions of [Setting 6.12](#), one finds that $\delta = \tau = d$. Moreover, The T -resolutions of $\mathcal{S}(\Omega_{R/k})_i$ can be computed using [Remark 3.7](#). From these resolutions and [Theorem 2.5](#), one observes that the rank of each T -module is $\text{rank } \mathcal{S}(\Omega_{R/k})_i = \binom{d}{i}$ for $0 \leq i \leq \delta = d$.

Certainly this was the case in [Remark 6.5](#) when $d = 2$. Moreover, this confirms that [Theorem 3.8](#) cannot be applied to study every component \mathcal{A}_i when $d \geq 3$. Regardless, since both $\mathcal{S}(\Omega_{R/k})_0$ and $\mathcal{S}(\Omega_{R/k})_d$ have rank 1, perhaps one can determine the corresponding generators of \mathcal{A}_d and \mathcal{A}_0 . As before, one has a natural candidate for the generator of \mathcal{A}_d .

Remark 6.14. With the assumptions of [Setting 6.12](#), let \mathcal{A} be the ideal defining $\mathcal{R}(\Omega_{R/k})$ as a quotient of $\mathcal{S}(\Omega_{R/k})$. The T -module \mathcal{A}_d is generated as $\mathcal{A}_d = \langle \det \mathcal{B} \rangle$ where $\mathcal{B} = [\mathcal{B}(\Theta) | \Theta]$, for $\mathcal{B}(\Theta)$ the Jacobian dual of Θ with respect to x_1, \dots, x_{2d} . Moreover, this is an equation of bidegree (d, d) .

Proof. This follows similar to the proof of part (a) of [Theorem 6.9](#). □

As noted prior to [Example 6.8](#), the statement of [Lemma 6.7](#) may be extended to this setting, and there is a similar relationship between $\mathcal{B}(\Theta)$ and Θ .

As we also have that $\text{rank } \mathcal{S}(\Omega_{R/k})_d = 1$, one may apply [Theorem 3.8](#) to the graded T -resolution of this module in order to study \mathcal{A}_0 . Similar to the proof of [Theorem 5.9](#), \mathcal{A}_0 is a cyclic T -module and [Theorem 3.8](#) can be used to determine the bidegree of its generator, so long as the ideal of minors of the syzygy matrix of $\mathcal{S}(\Omega_{R/k})_d$ has height at least two. This is the topic of [Conjecture 6.11](#), hence it may not be straightforward, however the author intends to investigate this in a subsequent article.

ACKNOWLEDGEMENTS

The author would like to thank Youngsu Kim and Vivek Mukundan for their helpful comments and suggestions on an earlier draft of this paper. The use of *Macaulay2* [[14](#)] was helpful in the preparation of this article, providing numerous examples of the results introduced here.

REFERENCES

- [1] L. Avramov, *Complete intersections and symmetric algebras*, J. Algebra **73** (1981), 248–263.
- [2] J. A. Boswell and V. Mukundan, *Rees algebras of almost linearly presented ideals*, J. Algebra **460** (2016), 102–127.
- [3] W. Bruns and J. Herzog, *Cohen-Macaulay rings*, Cambridge Studies in Advanced Mathematics **39**, Cambridge University Press, Cambridge, 1993.
- [4] D.A. Buchsbaum and D. Eisenbud, *What makes a complex exact?*, J. Algebra **25** (1973), 259–268.
- [5] D.A. Buchsbaum and D. Eisenbud, *Some structure theorems for finite free resolutions*, Advances in Math. **12** (1974), 84–139.
- [6] A. Costantini, *Cohen-Macaulay fiber cones and defining ideal of Rees algebras of modules*, Women in Commutative Algebra – Proceedings of the 2019 WICA Workshop, Association for Women in Mathematics Series, vol. 29, Springer (2022).

- [7] A. Costantini, E. F. Price, and M. Weaver, *On Rees algebras of linearly presented ideals and modules*, [arxiv:2308.16010](https://arxiv.org/abs/2308.16010). (To appear in Collect. Math.)
- [8] A. Costantini and T. Dang, *On the Cohen-Macaulay property of the Rees algebra of the modules of differentials*, Proc. Amer. Math. Soc. **150** (2022), 941–950
- [9] D. Cox, J. W. Hoffman, and H. Wang, *Syzygies and the Rees algebra*, J. Pure Appl. Algebra **212** (2008), 1787–1796.
- [10] J. A. Eagon and D. G. Northcott, *Ideals defined by matrices and a certain complex associated with them*, Proc. Roy. Soc. Ser. A **269** (1962), 188–204.
- [11] D. Eisenbud, *Commutative algebra: with a view toward algebraic geometry*, Graduate Texts in Mathematics **150**, Springer-Verlag, New York, 1995.
- [12] D. Eisenbud, C. Huneke and B. Ulrich, *What is the Rees algebra of a module?*, Proc. Amer. Math. Soc. **131** (2003), 701–708.
- [13] D. Ferrand, *Suite régulière et intersection complète*, C. R. Acad. Sci. Paris Sér. A-B **264** (1967), 247–248.
- [14] D. R. Grayson and M. E. Stillman, *Macaulay2*, a software system for research in algebraic geometry. Available at <http://www.math.uiuc.edu/Macaulay2/>
- [15] J. Harris, *Algebraic geometry*, Graduate Texts in Mathematics **133**, Springer-Verlag, New York, 1992.
- [16] R. Hartshorne, *Complete intersections and connectedness*, Amer. J. Math. **84** (1962), 497–508.
- [17] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics **52**, Springer-Verlag, New York-Heidelberg, 1977.
- [18] J. Herzog, A. Simis, and W. V. Vasconcelos, *Approximation complexes of blowing-up rings*, J. Algebra **74** (1982), 466–493.
- [19] J. Herzog, A. Simis, and W. V. Vasconcelos, *Approximation complexes of blowing-up rings. II*, J. Algebra **82** (1983), 53–83.
- [20] C. Huneke and M. Rossi, *The dimension and components of symmetric algebras*, J. Algebra **98** (1986), 200–210.
- [21] J.-P. Jouanolou, *Résultant anisotrope, compléments et application*, Electron. J. Combin. **3** (1997), research paper 2, approx. 91 pp.
- [22] Y. Kim and V. Mukundan, *Equations defining certain graphs*, Michigan Math. J. **69** (2020), 675–710.
- [23] E. Kunz, *Kähler Differentials*, Advanced Lectures in Mathematics, Friedr. Vieweg & Sohn, Braunschweig, 1986.
- [24] A. R. Kustin, C. Polini, and B. Ulrich, *The equations defining blowup algebras of height three Gorenstein ideals*, Algebra Number Theory **11** (2017), 1489–1525.
- [25] A. R. Kustin, C. Polini, and B. Ulrich, *The bi-graded structure of symmetric algebras with applications to Rees rings*, J. Algebra **469** (2017), 188–250.
- [26] S. Morey, *Equations of blowups of ideals of codimension two and three*, J. Pure Appl. Algebra **109** (1996), 197–211.
- [27] S. Morey and B. Ulrich, *Rees algebras of ideals with low codimension*, Proc. Amer. Math. Soc. **124** (1996), 3653–3661.
- [28] P. H. Lan Nguyen, *On Rees algebras of linearly presented ideals*, J. Algebra **420** (2014), 186–200.
- [29] P. H. Lan Nguyen, *On Rees algebras of linearly presented ideals in three variables*, J. Pure Appl. Algebra **221** (2017), 2180–2198.
- [30] A. Simis, B. Ulrich, and W. V. Vasconcelos, *Jacobian dual fibrations*, Amer. J. Math. **115** (1993), 47–75.
- [31] A. Simis, B. Ulrich, and W. V. Vasconcelos, *Tangent star cones*, J. Reine Angew. Math. **483** (1997), 23–59.
- [32] A. Simis, B. Ulrich, and W. V. Vasconcelos, *Rees algebras of modules*, Proc. London Math. Soc. **87** (2003), 610–646.
- [33] A. Simis, B. Ulrich, and W. V. Vasconcelos, *Tangent algebras*, Trans. Amer. Math. Soc. **364** (2012), 571–594.
- [34] B. Ulrich and W. V. Vasconcelos, *The equations of Rees algebras of ideals with linear presentation*, Math. Z. **214** (1993), 79–92.
- [35] W. V. Vasconcelos, *Ideals generated by R-sequences*, J. Algebra **6** (1967), 309–316.
- [36] W. V. Vasconcelos, *On the homology of I/I^2* , Comm. Algebra **6** (1978), 1801–1809.
- [37] W. V. Vasconcelos, *On the equations of Rees algebras*, J. Reine Angew. Math. **418** (1991), 189–218.
- [38] W. V. Vasconcelos, *Arithmetic of blowup algebras*, London Math. Soc. Lecture Note Ser. **195**, Cambridge University Press, Cambridge, 1994.
- [39] W. V. Vasconcelos, *Integral closure*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2005.
- [40] M. Weaver, *On Rees algebras of ideals and modules over hypersurface rings*, J. Algebra **636** (2023), 417–454.
- [41] M. Weaver, *The equations of Rees algebras of height three Gorenstein ideals in hypersurface rings*, J. Commut. Algebra **16** (2024), 123–149.
- [42] Y. Yoshino, *Codimension of Jacobian ideals and (R_n) conditions for complete intersections*, Hiroshima Math. J. **15** (1985), 663–667.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NOTRE DAME, 255 HURLEY BLDG, NOTRE DAME, IN 46556
 Email address: mweaver6@nd.edu