SOME GENERALIZED METRIC PROPERTIES OF *n*-SEMITOPOLOGICAL GROUPS

FUCAI LIN* AND XIXI QI

ABSTRACT. A semitopological group G is called an *n*-semitopological group, if for any $g \in G$ with $e \notin \overline{\{g\}}$ there is a neighborhood W of e such that $g \notin W^n$, where $n \in \mathbb{N}$. The class of *n*-semitopological groups $(n \geq 2)$ contains the class of paratopological groups and Hausdorff quasi-topological groups. Fix any $n \in \mathbb{N}$. Some properties of *n*-semitopological groups are studied, and some questions about *n*-semitopological groups are discussed, which contains mainly results are that (1) each Hausdorff first-countable 2-semitopological group admits a coarser Hausdorff semi-metrizable topology and has G^*_{δ} -diagonal; (2) each locally compact, Baire and σ -compact 2-semitopological groups topological groups are discussed.

1. INTRODUCTION AND PRELIMINARIES

Let G be a group, and let \mathscr{F} be a topology on G. We say that

• G is a semitopological group if the product map of $G \times G$ into G is separately continuous under the topology \mathscr{F} ;

• G is a quasitopological group if, under the topology \mathscr{F} , the space G is a semitopological group and the inverse map of G onto itself associating x^{-1} with arbitrary $x \in G$ is continuous;

• G is a paratopological group if the product map of $G \times G$ into G is jointly continuous under the topology \mathscr{F} ;

• G is a topological group if, under the topology \mathscr{F} , the space G is a paratopological group and the inverse map of G onto itself associating x^{-1} with arbitrary $x \in G$ is continuous.

The classes of semitopological groups, quasitopological groups, paratopological groups and topological groups were studied from twentieth century, see [2]. In [5], R. Ellis proved that each locally compact Hausdorff semitopological group is a topological group, which shows that each compact Hausdorff semitopological group is a topological group. Recently, the concept of almost paratopological group has been introduced by E. Reznichenko in [14], which is a generalized of paratopological groups and Hausdorff quasitopological groups. A semitopological group G is called *almost paratopological*, if for any $g \in G$ with $e \notin \overline{\{g\}}$ there is a neighborhood W of e such that $g \notin W^2$. By applying the concept of almost paratopological group, it is proved in [14] that each compact almost paratopological group is a topological group. However, there exists a

²⁰²⁰ Mathematics Subject Classification. Primary 54H11; 22A05; secondary 54A25; 54B15; 54E35.

Key words and phrases. n-semitopological group; paratopological group; quasi-topological group; locally compact; metrizable topology.

The authors are supported by the Key Program of the Natural Science Foundation of Fujian Province (No: 2020J02043), the NSFC (Nos. 11571158).

^{*}corresponding author.

FUCAI LIN* AND XIXI QI

compact T_1 quasitopological group which is not a topological group, such as, the integer group with the finite complementary topology. In this paper, we define the following concept of *n*-semitopological group $(n \in \mathbb{N})$ and ∞ -semitopological group, where each almost paratopological group is called 2-semitopological group.

Definition 1.1. Fix an $n \in \mathbb{N}$. A semitopological group G is called an *n*-semitopological group, if for any $g \in G$ with $e \notin \overline{\{g\}}$ there is a neighborhood W of e such that $g \notin W^n$. In particular, G is called an ∞ -semitopological group, if for any $n \in \mathbb{N}$ and $g \in G$ with $e \notin \overline{\{g\}}$ there is a neighborhood W of e such that $g \notin W^n$.

Remark 1.2. Clearly, each semitopological group and each almost paratopological group are just an 1-semitopological group and a 2-semitopological group respectively. In [14], E. Reznichenko proved that all paratopological groups and Hausdorff quasi-topological groups are ∞ -semitopological groups and 2-semitopological groups respectively. Obviously, there exists an ∞ -semitopological group which is neither a paratopological group nor a quasi-topological group, see the following example.

Example 1.3. Let $G = \mathbb{R}$ with the usual addition. Put $\mathscr{P} = \{[0, \frac{1}{n}) - \mathbb{Q}_+ : n \in \mathbb{N}\}$ and $\mathscr{B} = \{x+P : P \in \mathscr{P}, x \in \mathbb{R}\}, where \mathbb{Q}_+$ is the set all positive rational numbers. Let τ be a topology on G such that \mathscr{B} is a base for τ . It is easy to see that (G, τ) is a Hausdorff ∞ -semitopological group which is neither a paratopological group nor a quasi-topological group.

Remark 1.4. (1) If a T_0 -quasitopological group G is a 2-semitopological group, then G is Hausdorff.

(2) Each compact 2-semitopological group is a topological group, see [14, Theorem 6].

(3) A σ -compact regular 2-semitopological group is ccc, see [14, Corollary 4].

(4) For any T_1 -semitopological group G and $n \in \mathbb{N}$, G is an *n*-semitopological group if and only if $\bigcap_{U \in \mathcal{N}_e} U^n = \{e\}$, where \mathcal{N}_e denotes the family of all neighborhoods of the neutral element e of G.

The following question is interesting. Indeed, we give an example to show that there exists a 2-semitopological group which is not a 3-semitopological group in Section 2.

Question 1.5. For any $n \in \mathbb{N} \setminus \{1\}$, does there exists an n-semitopological group G such that G is not an (n + 1)-semitopological group?

Moreover, it is natural to pose the following question by above remark.

Question 1.6. If G is a locally compact n-semitopological group for some $n \in (\mathbb{N} \cup \{\infty\}) \setminus \{1\}$, is G a topological group?

In this paper, we give some partial answers to above question and discuss some generalized metric properties of *n*-semitopological groups, where $n \in (\mathbb{N} \cup \{\infty\}) \setminus \{1\}$. The paper is organized as follows.

In Section 2, we mainly give some topological properties of *n*-semitopological groups $(n \in \mathbb{N})$. First, we give a Hausdorff quasi-topological group G (thus a 2-semitopological group) such that G is not a 3-semitopological group. Moreover, we prove that (1) Each Hausdorff 2-semitopological group is qg-separated; (2) For any $m \in \mathbb{N} \setminus \{1\}$, a semitopological group G is a T_1 *m*-semitopological group if and only if $S_G^m = \{(x_1, \ldots, x_m) \in G^m : x_1 \cdot \ldots \cdot x_m = e\}$ is closed in G^m .

In Section 3, we mainly discuss some generalized metric properties of *n*-semitopological groups $(n \in \mathbb{N})$. We prove that (1) each Hausdorff first-countable 2-semitopological

group admits a coarser Hausdorff semi-metrizable topology and has G^*_{δ} -diagonal; (2) each locally compact, Baire and σ -compact 2-semitopological group is a topological group; (3) the condensation of a 2-semitopological group topology is given.

In Section 4, we mainly consider some cardinal invariants of *n*-semitopological groups $(n \in \mathbb{N})$. We mainly prove that (1) if G is a T_1 m-semitopological group and H a compact closed neutral subgroup of G, where $m \in \mathbb{N} \setminus \{1\}$, then G/H is an (m-1)-semitopological group; (2) if G is a regular κ -Lindelöf κ - Σ 2-semitopological group, then G is a κ -cellular space. Moreover, some interesting questions are posed.

The symbol \mathbb{N} denotes the natural numbers. The letter e denotes the neutral element of a group, and I denotes the unit interval with usual topology. Put $\mathbb{N}^* = \mathbb{N} \cup \{\infty\}$. For a semitopological group G, we denote the family of all neighborhoods of the neutral element e by \mathcal{N}_e . Readers may refer to [2, 5, 6] for notations and terminology not explicitly given here.

2. Some properties of n-semitopological groups

In this section, we mainly discuss some properties of *n*-semitopological groups, and pose some questions about *n*-semitopological groups, where $n \in \mathbb{N}^*$. First, we give a partial answer to Question 1.5.

Example 2.1. There exists a Hausdorff quasi-topological group G (thus a 2-semitopological group) such that G is not a 3-semitopological group.

Proof. We consider the strongest topology τ on the group of integers $G = \mathbb{Z}$ such that for every $z \in \mathbb{Z}$ the sequence $(z \pm n^2)_{n \in \omega}$ converges to z. We claim that G is Hausdorff. Indeed, for each $m \in \mathbb{Z}$, put $F_m = \{m \pm n^2 : n \in \omega\}$. Then each F_m is compact in τ . Let σ be determined by the countable family of compact subsets $\{F_m : m \in \mathbb{N}\}$. Obviously, we have $\sigma \subset \tau$. Since for any distinct numbers $z, z' \in Z$, the sets $\{z \pm n^2 : n \in \omega\}$ and $\{z' \pm n^2 : n \in \omega\}$ has finite intersection, it is easy to see that σ is a Hausdorff k_{ω} -topology. Therefore, it is easy to check that (G, τ) is a Hausdorff quasi-topological group. Next we claim that $1 \in \bigcap \{U + U + U : 0 \in U \in \tau\}$.

Indeed, it suffices to prove that for every even number a > 0 there exist numbers n, m, k > a such that $1 = k^2 + n^2 - m^2$. Take $k = a^2 + 1$ and observe that

$$k^{2} - 1 = (k - 1)(k + 1) = a^{2}(k + 1) = m^{2} - n^{2} = (m - n)(m + n),$$

where m, n can be found from the equation m - n = a and $m + n = a(a^2 + 2)$. Then $m = \frac{a(a^2+3)}{2} > a$ and $n = \frac{a^3+a}{2} > a$. Thus, we have $1 = k^2 + n^2 - m^2$. Therefore, (G, τ) is a 2-semitopological group, but it is not a 3-semitopological group.

Therefore, (G, τ) is a 2-semitopological group, but it is not a 3-semitopological group.

Next, we give some concepts in order to discuss some properties of n-semitopological groups.

Let (G, τ) be a semitopological group. The paratopological group reflexion $G^{pg} = (G, \tau^{pg})$ of (G, τ) we understand the group G endowed with the strongest topology $\tau^{pg} \subset \tau$ turning G into paratopological group. The quasitopological group reflexion $G^{qg} = (G, \tau^{qg})$ of (G, τ) we understand the group G endowed with the strongest topology $\tau^{qg} \subset \tau$ turning G into quasitopological group. Clearly, the following characteristic property holds: the identity map $i: G \to G^{pg}$ is continuous and for every continuous group homomorphism $h: G \to H$ from G into a paratopological group H the homomorphism $h \circ i^{-1}: G^{pg} \to H$ is continuous. The situation of quasitopological group reflexion is similar. A subset U of G is called pg-closed (pg-open) if U is closed (pg-open) in G^{qg} .

A semitopological group G is called *pg*-separated (*qg*-separated) provided its group reflexion G^{pg} (G^{qg}) is Hausdorff.

The following two propositions are obvious.

Proposition 2.2. Let G be a semitopological group, and let \mathscr{B} be a neighborhood base of e. Then the family $\{U \cup U^{-1}\}$ is a neighborhood base of e in G^{qg} .

Proposition 2.3. Let G be a semitopological group. If G^{pg} is T_1 , then G is an ∞ -semitopological group; if G^{qg} is Hausdorff, then G is a 2-semitopological group.

The following proposition shows that each Hausdorff 2-semitopological group is qg-separated.

Proposition 2.4. Let G be a Hausdorff 2-semitopological group. Then G is qg-separated.

Proof. Take any $g \neq e$. Since G is a Hausdorff 2-semitopological group, there exists an open neighborhood U of e such that $gU \cap U = \emptyset$ and $Ug \cap U = \emptyset$, $g \notin U^2$. Moreover, it follows from [14, Proposition 5 (4)] that there exists an open neighborhood $W \subset U$ of e such that $\{g, g^{-1}\} \cap (W^{-1})^2 = \emptyset$. Then $gW \cap W = \emptyset$, $Wg \cap W = \emptyset$, $g \notin W^2$ and $g \notin (W^{-1})^2$, hence $gW \cap (W \cup W^{-1}) = \emptyset$ and $gW^{-1} \cap (W \cup W^{-1}) = \emptyset$. Therefore, we have $g(W \cup W^{-1}) \cap (W \cup W^{-1}) = \emptyset$. Thus G is qg-separated by Proposition 2.2.

Let X be a space, and let $(Homeop(X), \tau_p)$ be the group of all homeomorphisms of X onto itself, with the pointwise convergence topology. Then $(Homeop(X), \tau_p)$ is a semitopological group, but it need not be a topological group, see [2, Example 1.2.12]. It is well-known that if X is a discrete space or X = I then $(Homeop(X), \tau_p)$ is a topological group, see [2, Exercises 1.2.k]. Therefore, the following question is interesting.

Question 2.5. How to given a characterization \mathcal{P} of the space X such that $(Homeop(X), \tau_p)$ is a 2-semitopological group if and only if X has the property \mathcal{P} ?

Proposition 2.6. Let X be a T_2 locally compact space and $(Homeop(X), \tau_c)$ the group of all homeomorphisms of X onto itself, with the compact-open topology. Then $(Homeop(X), \tau_c)$ is an ∞ -semitopological group.

Proof. Since X is a T_2 locally compact space, it is well known that $(Homeop(X), \tau_c)$ is a paratopological group, hence $(Homeop(X), \tau_c)$ is an ∞ -semitopological group. \Box

In particular, if X is a T_2 compact space, then $(Homeop(X), \tau_c)$ is a topological group, hence it is an ∞ -semitopological group. However, the following question is still open.

Question 2.7. How to given a characterization \mathcal{P} of the space X such that $(Homeop(X), \tau_c)$ is a 2-semitopological group if and only if X has the property \mathcal{P} ?

Let (X, τ) be a space. A subset A of X is called *regular open* if $A = int(\overline{A})$. The family of all regular open sets forms a base for a smaller topology τ_s on X, which is called the *semi-regularization* of τ . The following question is still unknown for us.

Question 2.8. Let G be an m-semitopological group for some $m \in \mathbb{N}$. Is the semiregularization G_{sr} an m-semitopological group? What if we assume the space to be ∞ semitopological group?

Next, we discuss some important properties of *m*-semitopological groups for some $m \in \mathbb{N}$.

Theorem 2.9. Let G be a semitopological group and $m \in \mathbb{N}^*$. If one of the following conditions is satisfied, then G is an m-semitopological group.

- (1) G is a paratopological group;
- (2) G is a subgroup of an m-semitopological group;
- (3) G is the product of m-semitopological groups;
- (4) there exists a continuous isomorphism of G onto a T_1 m-semitopological group.

Proof. Obviously, (1) and (2) hold.

(3) First, we consider $m \in \mathbb{N}^* \setminus \{\infty\}$. Let $\{G_\alpha : \alpha \in A\}$ be a family of *m*-semitopological groups such that $G = \prod_{\alpha \in A} G_\alpha$. Take any $g = (g_\alpha)_{\alpha \in A}$ with $e \notin \overline{\{g\}}$. It is obvious that there exists $\beta \in A$ such that $e_\beta \notin \overline{\{g_\beta\}}$, then there exists an open neighborhood U_β of e_β in G_β such that $g_\alpha \notin U_\beta^m$. Put $U = U_\beta \times \prod_{\alpha \in A \setminus \{\beta\}} G_\alpha$. Then $g \notin U^m$. The proof of the case of $m = \infty$ is similar.

(4) First, we consider $m \in \mathbb{N}^* \setminus \{\infty\}$. Suppose that $\phi : G \to H$ is a continuous isomorphism of the group G onto a T_1 *m*-semitopological group H. Take any $g \neq e$ in G. Then there exists an open neighborhood W of the neutral element in H such that $\phi(g) \notin W^m$. Put $V = \phi^{-1}(W)$. Hence $g \notin V^m$. The proof of the case of $m = \infty$ is similar. \Box

Let G be a group and any integer number $m \ge 2$. We denote

$$S_G^m = \{(x_1, \dots, x_m) \in G^m : x_1 \cdot \dots \cdot x_m = e\}, E_G^m = \bigcap_{U \in \mathcal{N}_e} \overline{(U^{-1})^{m-1}}$$

The following theorem gives some characterizations of *m*-semitopological groups for each $m \in \mathbb{N}^* \setminus \{1, \infty\}$.

Theorem 2.10. Let G be a semitopological group and $m \in \mathbb{N}^* \setminus \{1, \infty\}$. Then we have

(1)

$$\overline{\{e\}} \subset E_G^m = \bigcap_{U \in \mathcal{N}_e} (U^{-1})^m;$$

- (2) G is an m-semitopological group if and only if $E_G^m = \overline{\{e\}}$;
- (3) $\overline{S_G^m} = \mathfrak{m}^{-1}(E_G^m)$, where \mathfrak{m} is the multiplication in the group G;
- (4) the following statements are equivalent:

(i) G is a T_1 m-semitopological group; (ii) $E_G^m = \{e\};$ (iii) S_G^m is closed in G^m .

Proof. (1) From [14, Proposition 4], it follows that $\overline{\{e\}} \subset E_G^m \subset \bigcap_{U \in \mathcal{N}_e} (U^{-1})^m$. Take any $g \in G \setminus E_G^m$. Then there exists $U \in \mathcal{N}_e$ such that $gU \cap (U^{-1})^{m-1} = \emptyset$, hence $g \notin (U^{-1})^m$. Thus $E_G^m = \bigcap_{U \in \mathcal{N}_e} (U^{-1})^m$.

(2) Let G be an m-semitopological group. Then $\overline{\{e\}} \subseteq E_G^m$ by (1). Take any $g \notin G \setminus \overline{\{e\}}$. Hence $e \notin \overline{\{g^{-1}\}}$. Since G is an m-semitopological group, there exists $U \in \mathcal{N}_e$ such that $g^{-1} \notin U^m$, hence $g \notin (U^{-1})^m \supseteq E_G^m$. Therefore, $E_G^m \subseteq \overline{\{e\}}$.

Now suppose $E_G^m = \overline{\{e\}}$. Take any $g \neq e$ with $e \notin \overline{\{g\}}$. Then $g^{-1} \notin \overline{\{e\}} = E_G^m$. From (1), it follows that there exists $U \in \mathcal{N}_e$ such that $g^{-1} \notin (U^{-1})^m$. Then $g \notin U^m$.

(3) Let $(x_1, \ldots, x_m) \in G^m$. Clearly, we have

$$(x_1,\ldots,x_m)\in\overline{S_G^m}\Leftrightarrow(Ux_1\times\ldots\times Ux_m)\cap S_G^m\neq\emptyset$$

for any $U \in \mathcal{N}_e$, that is $e \in Ux_1 \dots Ux_m$ for any $U \in \mathcal{N}_e$. Hence

$$(x_1,\ldots,x_m)\in\overline{S_G^m}\Leftrightarrow e\in U^mx_1\ldots x_m$$

for any $U \in \mathcal{N}_e$, then

$$(x_1,\ldots,x_m)\in\overline{S_G^m}\Leftrightarrow x_1\ldots x_m\in (U^{-1})^m$$

for any $U \in \mathcal{N}_e$. By (1), we have $(x_1, \ldots, x_m) \in \overline{S_G^m} \Leftrightarrow x_1 \ldots x_m \in E_G^m$. (4) From (2), it follows that (i) \Leftrightarrow (ii). (ii) \Leftrightarrow (iii) since $S_G^m = \mathfrak{m}^{-1}(e)$.

By Theorem 2.10 and the definition of ∞ -semitopological group, we have the following theorem.

Theorem 2.11. Let G be a semitopological group. Then we have

- (1) G is an ∞ -semitopological group if and only if $E_G^m = \overline{\{e\}}$ for each $m \in \mathbb{N}$;
- (2) $\overline{S_G^m} = \mathfrak{m}^{-1}(E_G^m)$ for each $m \in \mathbb{N}$;
- (3) the following statements are equivalent:

(i) G is a $T_1 \infty$ -semitopological group;

- (ii) $E_G^m = \{e\}$ for each $m \in \mathbb{N}$;
- (iii) S_G^m is closed in G^m for each $m \in \mathbb{N}$.

Suppose that X and Y are spaces. We say that the mapping $f: X \to Y$ is topologypreserving if the following conditions are satisfied:

- (1) f is surjective, continuous, open and closed;
- (2) a subset U of X is open if and only if $U = f^{-1}(f(U))$ and f(U) is open.

The following proposition shows that the topology-preserving mappings can preserve and inversely preserve for the class of *m*-semitopological groups, where $m \in \mathbb{N}^*$.

Proposition 2.12. Let G and H be two semitopological groups, and let $\phi : G \to H$ be a topology-preserving homomorphism. Then G is an m-semitopological group if and only if H is an m-semitopological group, where $m \in \mathbb{N}^*$.

Proof. We divide the proof into the following two cases.

Case 1 $m \in \mathbb{N}^* \setminus \{\infty\}$.

Assume G is an m-semitopological group. Take any $h \neq e_H$ and $e_H \notin \overline{\{h\}}$ in H. Then there exists $g \in G$ such that $\phi(g) = h$. Clearly, $e_G \notin \overline{\{g\}}$ in G since $e_H \notin \overline{\{h\}}$ and ϕ is a topology-preserving mapping. Since G is an m-semitopological group, there exists an open neighborhood U of e_G such that $g \notin U^m$. We claim that $h \notin (\phi(U))^m$. Indeed, suppose $h \in (\phi(U))^m$. Then $\phi^{-1}(h) \cap \phi^{-1}((\phi(U))^m) \neq \emptyset$. Since $\phi^{-1}((\phi(U))^m) = U^m$, it follows that $U^m \cap \phi^{-1}(h) \neq \emptyset$, then $\phi^{-1}(h) \subset U^m$ since $\phi^{-1}(h)$ is antidiscrete. Hence $g \in U^m$, which is a contradiction. Therefore, $h \notin (\phi(U))^m$. Thus H is an m-semitopological group.

Assume *H* is an *m*-semitopological group. Take any $g \neq e_G$ and $e_G \notin \overline{\{g\}}$ in *G*. Since ϕ is a topology-preserving mapping, it follows from [11, Proposition 1] that $e_H \notin \overline{\{\phi(g)\}}$, hence there exists an open neighborhood *V* of e_H in *H* such that $\phi(g) \notin V^m$. Then $\phi^{-1}(\phi(g)) \cap \phi^{-1}(V^m) = \emptyset$, hence $g \notin (\phi^{-1}(V))^m$. Therefore, *G* is an *m*-semitopological group.

Case 2 $m = \infty$.

The proof is similar to Case 1.

Finally, we consider the topological direct limit of *m*-semitopological groups, $m \in \mathbb{N}^*$. First, we recall the following concept.

Definition 2.13. Given a tower

 $X_0 \subset X_1 \subset X_2 \subset \ldots \subset X_n \subset \ldots$

of spaces, the union $X = \bigcup_{n \in \mathbb{N}} X_n$ endowed with the strongest topology making each inclusion map $X_n \to X$ continuous is called the *topological direct limit* of the tower $(X_n)_{n \in \mathbb{N}}$ and is denoted by $\varinjlim X_n$.

Let $\{G_n : n \in \mathbb{N}\}$ be a tower of semitopological groups. From [17, Proposition 1.1] that $G = \varinjlim G_n$ is a semitopological group. Moreover, if each G_n is a quasitopological group, then G is a quasitopological group by [17, Proposition 1.1] again. However, there exists a tower $\{G_n : n \in \mathbb{N}\}$ of topological groups such that G is not a paratopological group, see [17, Example 1.2]. Therefore, we have the following question.

Question 2.14. Let $\{G_n : n \in \mathbb{N}\}$ be a tower of m-semitopological groups (resp., ∞ -semitopological groups), where $m \ge 2$. Is $G = \varinjlim G_n$ an m-semitopological group (resp., ∞ -semitopological group)?

The following two results are obvious.

Theorem 2.15. Let $\{H_n : n \in \mathbb{N}\}$ be a sequence of m-semitopological groups (resp., ∞ -semitopological groups), where $m \geq 2$. Then both the σ -product and Σ -product of $\prod_{i \in \mathbb{N}} H_n$ are m-semitopological groups (resp., ∞ -semitopological groups).

Corollary 2.16. Let $\{H_n : n \in \mathbb{N}\}$ be a sequence of *m*-semitopological groups (resp., ∞ -semitopological groups), where $m \ge 2$. Then $G = \varinjlim G_n$ is an *m*-semitopological groups (resp., ∞ -semitopological groups), where $G_n = \prod_{i \le n} H_i$ and each G_n is identified as a subspace of G_{n+1} for each $n \in \mathbb{N}$.

For closing this section, we give the following proposition.

Proposition 2.17. Let $\{G_n : n \in \mathbb{N}\}$ be a tower of 2-semitopological groups. If each G_n is T_1 , then $G = \lim G_n$ is T_1 .

Proof. It suffices to prove that $\{e\}$ is closed in G. Since each G_n is T_1 , it follows that $\{e\}$ is closed in each G_n . Therefore, $\{e\}$ is closed in G.

3. Generalized metric properties of n-semitopological groups

In this section, we mainly discuss some generalized metric properties of n-semitopological groups, such as, weakly first-countable, semi-metrizable, symmetrizable and etc. First, we recall a concept.

Definition 3.1. Let $\mathscr{P} = \bigcup_{x \in X} \mathscr{P}_x$ be a cover of a space X such that for each $x \in X$, (a) if $U, V \in \mathscr{P}_x$, then $W \subset U \cap V$ for some $W \in \mathscr{P}_x$; (b) the family \mathscr{P}_x is a network of x in X, i.e., $x \in \bigcap \mathscr{P}_x$, and if $x \in U$ with U open in X, then $P \subset U$ for some $P \in \mathscr{P}_x$. The family \mathscr{P} is called a *weak base* for X [1] if, for every $A \subset X$, the set A is open in X whenever for each $x \in A$ there exists $P \in \mathscr{P}_x$ such that $P \subset A$. The space X is *weakly first-countable* if \mathscr{P}_x is countable for each $x \in X$.

From [10], it follows that all weakly first-countable paratopological groups are firstcountable; moreover, there exists a Hausdorff weakly first-countable quasitopological group is not first-countable [9, Example 2.1]. Therefore, we have the following question. **Question 3.2.** Let G be an n-semitopological group (resp., ∞ -semitopological group), where $n \ge 2$. If G is weakly first-countable, when is G a first-countable space?

Let us recall that a function $d : X \times X \to [0, +\infty)$ on a set X is a symmetric if for every points x, y the following two conditions are satisfied: (1) d(x, y) = 0 if and only if x = y; (2) d(x, y) = (d(y, x)). Fro each $x \in X$ and $\varepsilon > 0$, denote by $B(x, \varepsilon) = \{y \in X : d(x, y) < \varepsilon\}$. Then

• a space X is symmetrizable if there is a symmetric d on X such that $U \subset X$ is open if and only if for each $x \in U$, there exists $\varepsilon > 0$ with $B(x, \varepsilon) \subset U$;

• a space X is *semi-metrizable* if there is a symmetric d on X such that for each $x \in X$, the family $\{B(x, \varepsilon) : \varepsilon > 0\}$ forms a neighborhood base at x;

• a space X is called a *sub-symmetrizable space* if it admits a coarser symmetrizable topology;

• a space X is called a *subsemi-metrizable space* if it admits a coarser semi-metrizable topology.

Every symmetrizable space is weakly first-countable, and a space is semi-metrizable if and only if it is first-countable and symmetrizable, see [1].

Theorem 3.3. Let (G, σ) be a T_1 weakly first-countable semitopological group. Then (G, σ) is sub-symmetrizable.

Proof. Since G is weakly first-countable, we may assume that $\mathcal{P}_e = \{P_n(e) : n \in \mathbb{N}\}$ be a countable weak base at e for G, where $P_1(x) = G$ and $P_{n+1}(x) \subset P_n(x)$ for each $n \in \mathbb{N}$. For each $x \in G$, let $\mathcal{P}_x = \{xP_n(e) : n \in \mathbb{N}\}$. Put $\mathcal{P} = \bigcup_{x \in G} \mathcal{P}_x$. Then \mathcal{P} is a countable weak base for G. For each $n \in \mathbb{N}$, put $W_n(e) = P_n(e) \cup (P_n(e))^{-1}$; then define a function $d : G \times G \to \mathbb{R}$ by setting $d(x, y) = \inf\{\frac{1}{n} : x^{-1}y \in W_n(e)\}$. We claim that d is a symmetric on G. Indeed, it is obvious that d(x, y) = d(y, x) for any $x, y \in G$. Now suppose that d(x, y) = 0 for $x, y \in G$. Then from our assumption, it follows that $x^{-1}y \in W_n(e)$ for any $n \in \mathbb{N}$, hence $x^{-1}y \in P_n(e) \cup (P_n(e))^{-1}$ for any $n \in \mathbb{N}$. Assume that $x \neq y$. Then since (G, σ) is T_1 , it follows that $e \notin \{x^{-1}y\}$ and $e \notin \{y^{-1}x\}$. Then there exists $k \in \mathbb{N}$ such that $x^{-1}y \notin P_k(e)$ and $y^{-1}x \notin P_k(e)$, hence $x^{-1}y \notin P_k(e) \cup (P_k(e))^{-1}$. This is a contradiction. Therefore, we have x = y.

Clearly, for any $n \in \mathbb{N}$ and $x \in G$, we have $xW_{n+1}(e) = B(x, \frac{1}{n})$. The topology τ which is inducted by the symmetric d on G is coarser than σ . Therefore, (G, σ) is sub-symmetrizable.

It is well known that each first-countable paratopological group is submetrizable. However, the Sorgenfrey line is a first-countable ∞ -semitopological group which is not symmetrizable. Therefore, the following question is natural.

Question 3.4. Let (G, σ) be a T_1 weakly first-countable 2-semitopological group. When is (G, σ) symmetrizable?

If we improve the conditions in Theorem 3.3, then we have the following result.

Theorem 3.5. Let (G, σ) be a Hausdorff first-countable 2-semitopological group. Then (G, σ) admits a coarser Hausdorff semi-metrizable topology and has G^*_{δ} -diagonal.

Proof. Let $\{U_n : n \in \mathbb{N}\}$ be a countable neighborhood base of e such that $U_{n+1} \subset U_n$ for each $n \in \mathbb{N}$. For any $g \in G$, put $\mathscr{B} = \{g(U_n \cup U_n^{-1}) : n \in \mathbb{N}, g \in G\}$. Let τ be the topology generated by the neighborhood system \mathscr{B} . Then it is easy to see that (G, τ) is a first-countable quasitopological group and τ is coarser than the topology of σ . We claim that (G, τ) is Hausdorff. Indeed, take any $g \neq e$. Clearly, $g^{-1} \neq e$. Since (G, σ) is a Hausdorff 2-semitopological group, there exists $n \in \mathbb{N}$ such that $U_n \cap gU_n = \emptyset$, $U_n \cap U_n g = \emptyset$, $g \notin U_n^2$ and $g^{-1} \notin U_n^2$. Then $g \notin U_n U_n^{-1}$, $g \notin U_n^{-1} U_n$, $g \notin U_n^2$ and $g^{-1} \notin U_n^2$, which implies that $g(U_n \cup U_n^{-1}) \cap (U_n \cup U_n^{-1}) = \emptyset$. Therefore, (G, τ) is Hausdorff.

By [8, Corollary 2.7], (G, τ) is Hausdorff semi-metrizable and has a G^*_{δ} -diagonal. Since τ is coarser than the topology of σ , it follows that (G, σ) has G^*_{δ} -diagonal. \Box

Next we recall some concepts, and then pose Question 3.7.

Definition 3.6. Let X be a space and $\{\mathscr{P}_n\}_n$ a sequence of collections of open subsets of X.

- (1) X is called *developable* for X if $\{st(x, \mathscr{P}_n)\}_n$ is a neighborhood base at x in X for each point $x \in X$.
- (2) X is called *Moore*, if X is regular and developable.
- (3) X is called a wM-space if for each $x \in X$ and a sequence $\{x_n\}_n$ whenever $x_n \in \operatorname{st}^2(x, \mathscr{U}_n)$ then the set $\{x_n : n \in \mathbb{N}\}$ has a cluster point in X.

In [9], C. Liu proved that each regular paratopological group G, in which each singleton is a G_{δ} -set, is metrizable if G is a wM-space, and posed that if we can replace "paratopological group" with "semitopological group". Then R. Shen in [16] gave a Moore quasitopological group which is not metrizable. Therefore, a Moore ∞ semitopological groups may not be metrizable. Hence we have the following question.

Question 3.7. Let G be an n-semitopological group (resp., ∞ -semitopological group), where $n \ge 2$. If G is a wM-space in which each singleton is a G_{δ} -set, is G metrizable?

Next we give a partial answer to Question 1.6. First, we recall some concepts. Let X be a space. Then

(1) X is said to be *locally compact* if for any point $x \in X$ there exists a compact neighborhood C of x;

(2) X is said to be σ -compact if $X = \bigcup_{n \in \mathbb{N}} K_n$, where each K_n is compact;

(3) X is said to be *Baire* if $X = \bigcup_{n \in \mathbb{N}} A_n$ then there exists $n \in \mathbb{N}$ such that the interior of $\overline{A_n}$ is nonempty.

Theorem 3.8. Each locally compact, Baire and σ -compact 2-semitopological group is a topological group.

Proof. Let (G, τ) be a locally compact, Baire and σ -compact 2-semitopological group, and let $H = \overline{\{e\}}$. Clearly, H is a normal closed antidiscrete subgroup. Since the quotient mapping $\phi : G \to G/H$ is a topology-preserving homomorphism, it follows that the the quotient group $\widehat{G} = G/H$ is a T_1 locally compact, Baire and σ -compact 2-semitopological group. By [14, Proposition 7], \widehat{G} is a topological group if and only if G is a topological group. Therefore, it suffices to prove that \widehat{G} is a topological group. Moreover, since \widehat{G} is a T_1 2-semitopological group, it follows from [14, Proposition 6.4 (a) and (b)] that $Sym(\widehat{G})$ is closed in $(\widehat{G})^2$ and $Sym(\widehat{G})$ is a Hausdorff locally compact σ -compact quasitopological group. From Ellis theorem [3, Theorem 2] that $Sym(\widehat{G})$ is a topological group. Let $\widehat{\tau}$ and $\widehat{\tau}_{Sym}$ be the topologies of \widehat{G} and $Sym(\widehat{G})$ respectively. By Ellis theorem [3, Theorem 2] again, it suffices to prove that \widehat{G} is Hausdorff.

Take any $e \neq g \in G$. Since \widehat{G} is a T_1 2-semitopological group, it follows from [14, Proposition 5(4)] that there exists $U \in \mathcal{N}(e)$ such that $g \notin \overline{U^{-1}}$, where $\mathcal{N}(e)$ is the neighborhood of e in $(G, \widehat{\tau})$. We claim that $e \in \operatorname{Int} \overline{U^{-1}}$ in $(G, \widehat{\tau})$. Indeed, since

FUCAI LIN* AND XIXI QI

 $U^{-1} \in \hat{\tau}_{Sym}$, there exists a symmetric open neighborhood V of e in $Sym(\hat{G})$ such that $V^2 \subset U^{-1}$. Since $Sym(\hat{G})$ is σ -compact, there exists a countable subset A of G such that $G = \bigcup \{aV : a \in A\}$, then there exists $a \in A$ such that $\operatorname{Int} \overline{aV} \neq \emptyset$ in $(G, \hat{\tau})$ because \hat{G} is a Baire space. Then $\operatorname{Int} \overline{V} \neq \emptyset$ in $(G, \hat{\tau})$. Take any $v \in V \cap \operatorname{Int} \overline{V}$. Hence $e \in \operatorname{Int} \overline{v^{-1}V} \subset \overline{V^2} \subset \overline{U^{-1}}$, which shows that $e \in \operatorname{Int} \overline{U^{-1}}$. Put $W = \hat{G} \setminus \overline{U^{-1}}$ and $O = \operatorname{Int} \overline{U^{-1}}$. Clearly, $W \cap O = \emptyset$, $g \in W$ and $e \in O$. Moreover, W and O are open in \hat{G} . Therefore, \hat{G} is Hausdorff.

Remark 3.9. (1) There exists a locally compact, Baire and σ -compact semitopological group G such that G is not an 2-semitopological group. Indeed, let τ be the cofinite topology on a uncountable group H. Suppose G is the Tychonoff product of H and the Euclidean space \mathbb{R} , then G is a locally compact and σ -compact semitopological group. Clearly, H is a Baire space, hence G is Baire by [5, 3.9.J(c)]. However, G is not a 2-semitopological group.

(2) There exists a Hausdorff sequentially compact ∞ -semitopological group G which is not a paratopological group, see [13, Example 3].

Clearly, a compact semitopological group may not be a Baire space, such as any cofinite topology on a countable infinite group. Therefore, we have the following question.

Question 3.10. Is each compact 2-semitopological group a Baire space?

From [14, Theorem 6], each compact 2-semitopological group is a topological group, hence each compact T_0 2-semitopological group is a Baire space. Moreover, each compact space is locally compact and σ -compact. Hence we have the following corollary by Theorem 3.8.

Corollary 3.11. Each compact T_0 2-semitopological group is a topological group.

Finally, we consider the condensation of 2-semitopological group topologies. First, we give some propositions and lemmas.

Definition 3.12. A family \mathcal{P} of subsets of a space X is called a *network* for X if for each $x \in X$ and neighborhood U of x there exists $P \in \mathcal{P}$ such that $x \in P \subset U$. The infimum of the cardinalities of all networks of X is denoted by nw(X).

The following proposition is obvious.

Proposition 3.13. Let G be a semitopological group and $nw(G) \leq \kappa$, where κ is some infinite cardinal. Then $nw(G^{qg}) \leq \kappa$.

Proposition 3.14. Let τ and σ be two topologies on group G such that (G, τ) and (G, σ) are semitopological groups with $w((G, \tau)) \leq \kappa$ and $w((G, \sigma)) \leq \kappa$, where κ is some infinite cardinal. Then $w(G, \tau \vee \sigma) \leq \kappa$.

Proof. Let \mathcal{B}_1 and \mathcal{B}_2 be bases for (G, τ) and (G, σ) respectively such that $|\mathcal{B}_1| \leq \kappa$ and $|\mathcal{B}_2| \leq \kappa$. Put $\mathcal{B} = \{U \cap V : U \in \mathcal{B}_1, V \in \mathcal{B}_2\}$. It is easily verified that \mathcal{B} is a base for $\tau \lor \sigma$ and $|\mathcal{B}| \leq \kappa$. Therefore, $w(G, \tau \lor \sigma) \leq \kappa$.

Lemma 3.15. Suppose that κ is an infinite cardinal, X is a group, τ is a Hausdorff (resp., regular, Tychonof) 2-semitopological group topology on X that has a network weight $\leq \kappa$ and τ' is a topology on X that has weight $\leq \kappa$ such that $\tau' \subset \tau$. Then one can find a topology τ^* on X with the following properties:

(i)
$$\tau' \subset \tau^* \subset \tau$$
;

(*ii*) $w(X, \tau^*) \leq \kappa$;

(iii) (X, τ^*) is a Hausdorff (resp., regular, Tychonof) 2-semitopological group.

Proof. We first prove the case of Hausdorff. By [7, Lemma 4], there exists a Hausdorff semitopological group topology σ on X such that $\tau' \subset \sigma \subset \tau$ and $w(X, \sigma) \leq \kappa$. By Proposition 2.4, X^{qg} is Hausdorff; moreover, it follows from Proposition 3.13 that X^{qg} has a network weight $\leq \kappa$. Then one can find a Hausdorff quasitopological group topology δ on X such that $\delta \subset \tau^{qg}$ and $w(X, \delta) \leq \kappa$ by [15, Theorem 1]. Clearly, (X, δ) is a 2-semitopological group by [14, Theorem 5]. Now put $\tau^* = \sigma \lor \delta$. Then $\tau^* \subset \tau$ and τ^* is a Haudorff 2-semitopological group topology on X. By Proposition 3.14, $w(X, \tau^*) \leq \kappa$. Moreover, we have $\tau' \subset \tau^* \subset \tau$.

If τ is regular (Tychonof), then it follows from the above proof and [7, Lemma 3] that there exists a topology τ^* on X which has the properties of (i) and (ii) and (X, τ^*) is a regular (Tychonof) 2-semitopological group.

Now we can prove the main theorem.

Theorem 3.16. Suppose that κ is an infinite cardinal, X is a group, τ is a Hausdorff (resp., regular, Tychonof) 2-semitopological group topology on X that has a network weight $\leq \kappa$. Then there exists a condensation $i : (X, \tau) \to (X, \tau^*)$, where τ^* is a Hausdorff (resp., regular, Tychonof) 2-semitopological group topology τ^* on X such that $\tau^* \subset \tau$ and $w(X, \tau^*) \leq \kappa$.

Proof. Since X is Hausdorff (resp., regular, Tychonof) and has a network weight $\leq \kappa$, there exists a compact Hausdorff space (X, τ_0) such that $w(X, \tau_0) \leq \kappa$. Now, it follows from Lemma 3.15 that there exists a Hausdorff (resp., regular, Tychonof) 2-semitopological group topology τ^* on X such that $\tau^* \subset \tau$ and $w(X, \tau^*) \leq \kappa$. \Box

By Theorem 3.16, we have the following corollary.

Corollary 3.17. Suppose that κ is an infinite cardinal, X is a group, τ is a Hausdorff (resp., regular, Tychonof) 2-semitopological group topology on X with a countable network. Then there exists a condensation $i : (X, \tau) \to (X, \tau^*)$, where τ^* is a Hausdorff (resp., regular, Tychonof) second-countable 2-semitopological group topology τ^* on X such that $\tau^* \subset \tau$.

However, the following question is still unknown for us.

Question 3.18. Suppose that κ is an infinite cardinal, X is a group, τ is a Hausdorff (resp., regular, Tychonof) m-semitopological group topology on X that has a network weight $\leq \kappa$, where $m \in \mathbb{N}^* \setminus \{2\}$. Can we find a Hausdorff (resp., regular, Tychonof) m-semitopological group topology τ^* on X such that $\tau^* \subset \tau$ and $w(X, \tau^*) \leq \kappa$?

4. CARDINAL INVARIANTS OF *n*-SEMITOPOLOGICAL GROUPS

In this section, we mainly consider some cardinal invariants of n-semitopological groups. Moreover, some interesting questions are posed. First, we recall some concepts.

Let κ be an ordinal. A semitopological group G is left (right) κ -narrow if for each open set U there exists a set $A \subset G$ such that $|A| \leq \kappa$ and AU = G (UA = G). Put

 $\operatorname{In}_{l}(G) = \min\{\kappa : G \text{ is left } \kappa \text{ -narrow}\}, \operatorname{In}_{r}(G) = \min\{\kappa : G \text{ is right } \kappa \text{ -narrow}\} \text{ and }$

 $ib(G) = \omega \cdot \min\{\kappa : G \text{ is left } \kappa \text{ -narrow and right } \kappa \text{ -narrow}\}.$

Moreover, we recall the following some definitions.

Character: $\chi(G) = \omega \cdot \min\{|\mathcal{B}| : \mathcal{B} \text{ is a neighborhood base at the neutral element of } G\}$. Pseudocharacter: $\psi(G) = \omega \cdot \min\{|\mathcal{U}| : \mathcal{U} \text{ is a family of open sets and } \bigcap \mathcal{U} = \{e\}\}$. Extent: $e(G) = \omega \cdot \sup\{|S| : S \text{ is a closed discrete subspace of } G\}$. Weakly Lindelöf degree: $wl(G) = \omega \cdot \min\{\kappa : \text{in each open cover } \mathcal{U} \text{ there exists a subfamily} \\ \mathcal{V} \subset \mathcal{U} \text{ with cardinality } \kappa \text{ such that } \bigcup \mathcal{V} = G \}.$

Lindelöf degree: $l(G) = \omega \cdot \min\{\kappa : \text{ in each open cover } \mathcal{U} \text{ there exists a subfamily } \mathcal{V} \subset \mathcal{U} \text{ with cardinality } \kappa \text{ such that } \bigcup \mathcal{V} = G\}.$

A semitopological group G is said to be *saturated* if, for any non-empty open set U, the interior of U^{-1} is non-empty.

The following proposition may have been proven somewhere.

Proposition 4.1. If G is a saturated semitopological group, then $In_l(G) = In_r(G)$.

Proof. Let $In_l(G) = \kappa$. Now we show that $In_r(G) \leq \kappa$. Take any open neighborhood U of e. Since G is saturated, it follows that $int(U^{-1}) \neq \emptyset$. Take any $u \in int(U^{-1})$. Then $u^{-1} \cdot int(U^{-1})$ is an open neighborhood of e, hence there exists a subset A with the cardinality of κ such that $A \cdot u^{-1} \cdot int(U^{-1}) = G$, which shows that $A \cdot u^{-1} \cdot U^{-1}$. Thus $U \cdot u \cdot A = G$ and $|u \cdot A| = |A| = \kappa$. Hence $In_r(G) \leq \kappa$. Similarly, one can prove $In_l(G) \leq In_r(G)$.

In [19, Theorem 3.2], the authors proved that $ib(G) \leq e(G)$ for each quasitopological group, and in [12] the author proved that $ib(G) \leq wl(G)$ for each saturated paratopological group. Therefore, we have the following question by applying Proposition 4.1.

Question 4.2. If G is a saturated 2-semitopological group, then is

$$ib(G) \le \max\{e(G), wl(G)\}$$
?

Moreover, we have the following question.

Question 4.3. If G is a 2-semitopological T_1 group, then does $nw(G) \leq \chi(G)l(G^2)$ hold?

Next we discuss the quotient group on m-semitopological groups. First, we give a lemma.

Lemma 4.4. Let G be a T_1 m-semitopological group and F be a compact subset with $e \notin F$, where $N \in \mathbb{N}$. Then it can find an open neighborhood U of e in G such that $e \notin FU^{m-1}$.

Proof. Since G is T_1 and $e \notin F$, we can choose, for each $x \in F$, an open neighborhood V_x of e such that $e \notin xV_x$ and $x^{-1} \notin V_x^m$. Clearly, the family $\{xV_x : x \in F\}$ covers the compact set F, hence there exists a finite set A such that $F \subset \bigcup_{a \in A} aV_a$. Now put $U = \bigcap_{a \in A} V_a$. We claim that $e \notin FU^{m-1}$. Indeed, for any $f \in F$, there is $b \in A$ such that $f \in bV_b$. Since $b^{-1} \notin V_b^m$ and $fU \subset bV_bU^{m-1} \subset bV_b^m$, it follows that $e \notin fU^{m-1}$. Thus $e \notin FU^{m-1}$.

Theorem 4.5. Let G be a T_1 m-semitopological group and H a compact closed neutral subgroup of G, where $m \in \mathbb{N} \setminus \{1\}$. Then G/H is an (m-1)-semitopological group.

Proof. Clearly, G/H is a T_1 semitopological group. Take any $g \notin H$ in G. Since G is a m-semitopological T_1 group and H a compact closed neutral subgroup of G, there exists an neighborhood U of e such that $g \notin U^m$ and $e \notin g^{-1}HU^{m-1}$ by Lemma 4.4. We claim that $\pi(g) \notin (\pi(U))^{m-1}$. Otherwise, $Hg \cap HU^{m-1} \neq \emptyset$, that is, $g \in HU^{m-1} \neq \emptyset$, which shows that $e \in g^{-1}HU^{m-1}$. This is a contradiction. Hence $\pi(g) \notin (\pi(U))^{m-1}$. Then G/H is an (m-1)-semitopological group. \Box

By Theorem 4.5, we have the following corollary.

Corollary 4.6. Let G be a $T_1 \infty$ -semitopological group and H a compact closed neutral subgroup of G. Then G/H is an ∞ -semitopological group.

The following result shows that the cardinality of some 2-semitopological groups is at most 2^{κ} .

Theorem 4.7. If G is a T_1 2-semitopological group such that $l(G^2) \leq \kappa$ and $\psi(G) \leq \kappa$, then G has cardinality at most 2^{κ} .

Proof. Since G is a T_1 2-semitopological group, it follows from [14, Proposition 6 (4)] that SymG embeds closed in G^2 , then $l(SymG) \leq \kappa$ by our assumption. Moreover, it is obvious that SymG is T_1 and $\psi(SymG) \leq \kappa$. Then SymG has cardinality at most 2^{κ} by [19, Theorem 3.5], thus G has cardinality at most 2^{κ} .

By Theorem 4.7, we have the following corollary.

Corollary 4.8. If G is a T_1 2-semitopological group such that $l(G^2) \leq \omega$ and $\psi(G) \leq \omega$, then G has cardinality at most \mathfrak{c} .

Let κ be an infinite cardinal. We say a space X is κ -cellular if for each family μ of G_{δ} -sets of X there exists a subfamily $\lambda \subset \mu$ such that $|\lambda| \leq \kappa$ and $\bigcup \mu = \bigcup \lambda$.

Finally, we discuss when a 2-semitopological group is a κ -cellular space. First, we define the class of κ - Σ -spaces and give some lemmas.

Let κ be an infinite cardinal. We say that

(1) X is κ -countably compact if each open cover of size $\leq \kappa$ has a finite subcover.

(2) X is a κ - Σ -space if there exists a family $\mathscr{P} = \bigcup_{\alpha < \kappa} \mathscr{P}_{\alpha}$ with each \mathscr{P}_{α} being locally finite and the covering of \mathscr{C} by closed κ -countably compact sets, such that if $C \in \mathscr{C}$ and $C \subset U$ is open, then $C \subset P \subset U$ for some $P \in \mathscr{P}$.

The following proposition is obvious.

Proposition 4.9. A space X is a κ - Σ -space with $e(X) \leq \kappa$ if and only if there exist a family \mathscr{P} with $|\mathscr{P}| \leq \kappa$ and the covering of \mathscr{C} by closed κ -countably compact sets, such that if $C \in \mathscr{C}$ and $C \subset U$ is open, then $C \subset P \subset U$ for some $P \in \mathscr{P}$.

Let X be a space X and κ an infinite cardinal. We define the following property:

 (\mathbf{P}_{κ}) Let $\{x_{\alpha} : \alpha < 2^{\kappa}\}$ be a subset of X and for each $\alpha < 2^{\kappa}$ let \mathscr{P}_{α} be a family of closed subsets of X with a cardinality of at most κ . Then there is $\beta < 2^{\kappa}$ such that the following conditions holds:

(*) there exists $y \in \overline{\{x_{\alpha} : \alpha < \beta\}}$ such that if $\eta < \beta$ with $x_{\beta} \in P \in \mathscr{P}_{\eta}$, then $y \in P$.

Lemma 4.10. Let X be a regular κ - Σ -space with $e(X) \leq \kappa$, where κ is an infinite cardinal. Then (P_{κ}) holds for X.

Proof. By Proposition 4.9, there exist a family \mathscr{P} with $|\mathscr{P}| \leq \kappa$ and the covering of \mathscr{C} by closed κ -countably compact sets, such that if $C \in \mathscr{C}$ and $C \subset U$ is open, then $C \subset P \subset U$ for some $P \in \mathscr{P}$. Without loss of generality, we may assume that \mathscr{P} is closed under $< \kappa$ intersections. Let $\{x_{\alpha} : \alpha < 2^{\kappa}\}$ be a subset of X and for each $\alpha < 2^{\kappa}$ let \mathscr{F}_{α} be a family of closed subsets of X with a cardinality of at most κ . For each $\mu < 2^{\kappa}$, put

$$\mathscr{F}_{\mu}^{*} = \{ \bigcap \mathscr{F} : \mathscr{F} \subset \bigcup_{\alpha < \mu} \mathscr{F}_{\alpha}, |\mathscr{F}| < \kappa \text{ and } \bigcap \mathscr{F} \neq \emptyset \}$$

$$X_{\mu} = \{ x_{\alpha} : \alpha < \mu \}.$$

By induction on $\gamma < \kappa$ we construct a family of κ ordinals $\{\beta_{\alpha} : \alpha < \kappa\}$ such that for any $0 < \alpha < \kappa$ the following two conditions are satisfied:

(i) for any $\alpha < \gamma < \kappa$, we have $\beta_{\alpha} < \beta_{\gamma}$;

(ii) if $x_{\alpha} \in P \cap F$ for $\alpha < 2^{\kappa}$, $P \in \mathscr{P}$ and $F \in \mathscr{F}^*_{\beta_{\eta}}$, then there exists $y \in \bigcap_{\gamma > \eta} \overline{X_{\beta_{\gamma}}}$ such that $y \in P \cap F$.

Indeed, let $\beta_0 = \kappa$. Assume that the family $\{\beta_\eta : \eta < \alpha\}$ has been constructed, where $\alpha < \kappa$. For $P \in \mathscr{P}$ and $F \in \bigcup_{\eta < \alpha} \mathscr{F}^*_{\beta_\eta}$, let $S(P, F) = \{\nu < 2^{\kappa} : x_{\nu} \in P \cap F\}$. If $S(P, F) \neq \emptyset$, then we put $\lambda_{P,F} = \min S(P, F)$. Now we put

$$\beta_{\alpha} = \sup\left\{\bigcup_{\eta < \alpha} \beta_{\eta}, \sup\{\lambda_{P,F} : P \in \mathscr{P}, F \in \bigcup_{\eta < \alpha} \mathscr{F}^{*}_{\beta_{\eta}}, S(P,F) \neq \emptyset\}\right\} + 1.$$

Then the family of κ ordinals $\{\beta_{\alpha} : \alpha < \kappa\}$ has been constructed. Put $\beta = \sup\{\beta_{\alpha} : \alpha < \kappa\}$. Now it suffices to check that (\star) holds in (\mathbf{P}_{κ}) definition. Clearly, there exists $C \in \mathscr{C}$ such that $x_{\beta} \in C$. We can assume that

$$\mathscr{P}_C = \{ P \in \mathscr{P} : C \subset P \} = \{ P_{\alpha, C} : \alpha < \kappa \}$$

and

$$\mathscr{F}_C = \{F \in \mathscr{F}_{\beta}^* : x_{\beta} \in F\} = \{F_{\alpha}^* : \alpha < \kappa\}$$

Take any $\nu < \kappa$. Let $P_{\nu} = \bigcap_{\alpha \in \nu} P_{\alpha,C}$ and $F_{\nu} = \bigcap_{\alpha \in \nu} F_{\alpha}^*$, and let $\lambda_{\nu} = \lambda_{P_{\nu},F_{\nu}}$ and $z_{\nu} = x_{\lambda_{\nu}}$. Clearly, $\mathscr{F}_{\beta}^* = \bigcap_{\alpha < \kappa} \mathscr{F}_{\beta_{\alpha}}^*$, hence $F_{\nu} \in \mathscr{F}_{\beta_{\alpha}}^*$ for some $\alpha < \kappa$. Since $\beta \in \lambda_{P_{\nu},F_{\nu}} \neq \emptyset$, it follows that $\lambda_{\nu} = \lambda_{P_{\nu},F_{\nu}} \leq \beta_{\alpha+1} < \beta$, hence $z_{\nu} \in X_{\beta}$.

From the definition of the families \mathscr{P} and \mathscr{C} , it follows that $\{z_v : v \in \kappa\}$ accumulates to some point $z \in C \cap \bigcap_{\nu < \kappa} F_{\nu}$. Thus, $z \in \overline{X_{\beta}}$. Assume that $F \in \mathscr{F}_{\gamma}$ for $\gamma < \beta$ and $x_{\beta} \in F$. Then $F \in \mathscr{F}_C$ and $F = F_{\alpha}^* \supset F_{\alpha}$ for some $\alpha < \kappa$. Therefore, it follows that

$$z \in \bigcap_{\nu < \kappa} F_{\nu} \subset F_{\alpha} \subset F$$

Lemma 4.11. Assume that G is a regular quasitopological group, and assume that G satisfies (P_{κ}) for some infinite cardinal κ . Then G is a κ -cellular space.

Proof. Assume that G is not a κ -cellular space. Then we can find a family $\{A_{\alpha} : \alpha < 2^{\kappa}\}$ of non-empty sets of type G_{δ} such that $A_{\gamma} \not\subset \bigcup_{\alpha < \gamma} A_{\alpha}$ for any $\gamma < 2^{\kappa}$. For each $\gamma < 2^{\kappa}$, we can pick any $g_{\gamma} \in A_{\gamma} \setminus \bigcup_{\alpha < \gamma} A_{\alpha}$, and take a sequence $(U_{\gamma,n})_{n \in \omega}$ of open sets of G such that $g_{\gamma} \in U_{\gamma,n+1} \subset \overline{U_{\gamma,n+1}} \subset U_{\gamma,n}$ for any $n \in \omega$ and $B_{\gamma} = \bigcap_{n \in \omega} U_{\gamma,n} \subset A_{\gamma}$. For each $\gamma < 2^{\kappa}$, put $\mathscr{F}_{\gamma} = \{(G \setminus U_{\beta,n})g_{\alpha}^{-1}g_{\beta} : \alpha, \beta < \gamma, n \in \omega\}$ and $\mathscr{P}_{\gamma} = \mathscr{F}_{\gamma+1}$. Then the condition (P_{κ}) is satisfied for G, it follows that there exists $\delta \in 2^{\kappa}$ and $y \in \{g_{\alpha} : \alpha < \delta\}$ such that if $\eta < \delta$, $P \in \mathscr{P}_{\eta}$ and $g_{\delta} \in P$, then $y \in P$. Therefore, $y \in P$ if $g_{\delta} \in P \in \mathscr{P}_{\delta}$. Now, for any $\eta < \delta$, put $y_{\eta} = g_{\delta}y^{-1}g_{\eta}$; we claim that $y_{\eta} \in B_{\eta}$. Suppose not, then there exists $n \in \omega$ such that $y_{\eta} \notin U_{\eta,n}$. Clearly, we have $y \in g_{\eta}U_{\eta,n+1}^{-1}y \cap g_{\eta}(G \setminus \overline{U_{\eta,n+1}})^{-1}g_{\delta}$. Since $y \in \{g_{\alpha} : \alpha < \delta\}$ and $g_{\eta}U_{\eta,n+1}^{-1}y, g_{\eta}(G \setminus \overline{U_{\eta,n+1}})^{-1}g_{\delta}$ are open, there exists $\alpha < \delta$ such that $g_{\alpha} \in g_{\eta}U_{\eta,n+1}^{-1}y \cap g_{\eta}(G \setminus \overline{U_{\eta,n+1}})^{-1}g_{\delta}$. Then $g_{\delta} \in (G \setminus U_{\eta,n+1})g_{\eta}^{-1}g_{\alpha}$ and $y \in (U_{\eta,n+1})g_{\eta}^{-1}g_{\alpha}$, which is a contradiction.

Then since $y \in \{g_{\eta} : \eta < \delta\}$, it follows that

$$g_{\delta} = g_{\delta} y^{-1} y \in \overline{\{g_{\delta} y^{-1} g_{\eta} : \eta < \delta\}} = \overline{\{y_{\eta} : \eta < \delta\}},$$

hence $g_{\delta} \in \overline{\bigcup_{\eta < \delta} B_{\eta}}$. However, it is obvious that $g_{\delta} \notin \overline{\bigcup_{\eta < \delta} B_{\eta}}$, which is a contradiction. Therefore, G is a κ -cellular space.

By Lemmas 4.10 and 4.12, we have the following lemma.

Lemma 4.12. Let X be a regular quasitopological group, which is a κ - Σ -space with $e(G) \leq \kappa$. Then G is a κ -cellular space.

Theorem 4.13. Let G be a regular 2-semitopological group and G^2 be a κ - Σ -space with $e(G) \leq \kappa$. Then G is a κ -cellular space.

Proof. From [14, Proposition 6], it follows that Sym G is a quasitopological group and embeds closed in G^2 . Then SymG is a regular a κ - Σ -space with $e(G) \leq \kappa$. By Lemma 4.12, SymG is a κ -cellular space. Since G is a continuous image of SymG, it follows that G is a κ -cellular space.

By a similar proof of the product of two Lindelöf Σ -spaces being Lindelöf Σ -space (see [18]), we have the following lemma.

Lemma 4.14. Let X be a regular κ -Lindelöf κ - Σ space. Then X^2 is a κ -Lindelöf κ - Σ space.

By Theorem 4.13 and Lemma 4.14, we have the following theorem.

Theorem 4.15. Let G be a regular κ -Lindelöf κ - Σ 2-semitopological group. Then G is a κ -cellular space.

Acknowledgements The authors wish to express our sincere thanks to professor T. Banakh who provided Example 2.1.

References

- [1] A.V. Arhangel'skii, Mappings and spaces, Russian Math. Surveys, 21(1996): 115–162.
- [2] A.V. Arhangel' skiĭ, M. Tkachenko, Topological Groups and Related Structures, Atlantis Press and World Sci., 2008.
- [3] R. Ellis, Locally compact transformation groups, Duke Math. J., 24(2)(1957): 119–125.
- [4] R. Ellis, A note on the continuity of the inverse, Proc. Amer. Math. Soc., 8(1957): 372–373.
- [5] R. Engelking, General Topology(revised and completed edition), Heldermann Verlag, Berlin, 1989.
- [6] G. Gruenhage, Generalized metric spaces, In: K. Kunen, J. E. Vaughan(Eds.), Handbook of Set-Theoretic Topology, Elsevier Science Publishers B.V., Amsterdam, 1984, 423–501.
- [7] C. Hernández, Condensations of Tychonoff universal topological algebras, Comment. Math. Univ. Car., 42(3)(2001): 529–533.
- [8] P. Li, L. Mou, On quasitopological groups, Topol. Appl., 161 (2014): 243-247.
- C. Liu, Paratopological (topological) groups with certain networks, Comment. Math. Univ. Car., 55(1)(2014): 111–119.
- [10] P. Nyikos, Metrizability and the Fréchet-Urysohn property in topological groups, Proc. Amer. Math. Soc., 83(4)(1981): 793–801.
- [11] X.X. Qi, F. Lin, Topological properties under topology-preserving mappings, J. Minnan Normal Univ. (Natural Science), 2024, 37(1): 94–98.
- [12] O.V. Ravsky, Paratopological groups I, Mat. Stud, 16(1)(2001): 37–48.
- [13] O.V. Ravsky, A note on compact-like semitopological groups, arxiv preprint arxiv: 1907.11215, 2019.
- [14] E. Reznichenko, Almost paratopological groups, Topol. Appl., 338 (2023): 108673.
- [15] D.B. Shakhmatov, Condensations of universal topological algebras preserving continuity of operations and decreasing weights, Vestnik Moskovskogo Universiteta. Seriya 1. Matematika. Mekhanika, (2)(1984): 42–45.
- [16] R.X. Shen, On generalized metrizable properties in quasitopological groups, Topol. Appl., 173 (2014): 219-226.

FUCAI LIN* AND XIXI QI

- [17] N. Tatsuuma, H. Shimomura, T. Hirai, On group topologies and unitary representations of inductive limits of topological groups and the case of the group of diffeomorphisms, J. Math. Kyoto Univ., 38(3)(1998): 551–578.
- [18] V.V. Tkachuk. Lindelöf Σ -spaces: an omnipresent class, Revista de la Real Academia de Ciencias Exactas, Fisicas y Naturales. Serie A, Matematics, 104(2)(2010): 221-224.
- [19] W. Xuan, Y. Song, Z. Tang, Quasitopological groups, covering properties and cardinal inequalities, Topol. Appl., 272 (2020): 107059.

(Fucai Lin): 1. School of mathematics and statistics, Minnan Normal University, Zhangzhou 363000, P. R. China; 2. Fujian Key Laboratory of Granular Computing and Application, Minnan Normal University, Zhangzhou 363000, P. R. China

Email address: linfucai2008@aliyun.com; linfucai@mnnu.edu.cn

(XIXI QI): 1. SCHOOL OF MATHEMATICS AND STATISTICS, MINNAN NORMAL UNIVERSITY, ZHANGZHOU 363000, P. R. CHINA

Email address: 2473148247@qq.com