

MPSDynamics.jl: Tensor network simulations for finite-temperature (non-Markovian) open quantum system dynamics

Thibaut Lacroix,¹ Briec Le Dé,² Angela Riva,³ Angus J. Dunnett,⁴ and Alex W. Chin²

¹*Institut für Theoretische Physik und IQST, Albert-Einstein-Allee 11, Universität Ulm, D-89081 Ulm, Germany*

²*Sorbonne Université, CNRS, Institut des NanoSciences de Paris, 4 place Jussieu, 75005 Paris, France*

³*LPENS, Département de physique, École normale supérieure, Centre Automatique et Systèmes (CAS), MINES ParisTech, Université PSL, Sorbonne Université, CNRS, Inria, 75005 Paris, France*

⁴*Multiverse Computing, 7 rue de la Croix Martre, 91120 Palaiseau, France*

(*Electronic mail: alex.chin@insp.jussieu.fr)

(*Electronic mail: angus.dunnett@multiversecomputing.com)

(*Electronic mail: angela.riva@inria.fr)

(*Electronic mail: lede@insp.jussieu.fr)

(*Electronic mail: thibaut.lacroix@uni-ulm.de)

(Dated: 12 June 2024)

The `MPSDynamics.jl` package provides an easy to use interface for performing open quantum systems simulations at zero and finite temperatures. The package has been developed with the aim of studying non-Markovian open system dynamics using the state-of-the-art numerically exact Thermalized-Time Evolving Density operator with Orthonormal Polynomials Algorithm (T-TEDOPA) based on environment chain mapping. The simulations rely on a tensor network representation of the quantum states as matrix product states (MPS) and tree tensor network (TTN) states. Written in the Julia programming language, `MPSDynamics.jl` is a versatile open-source package providing a choice of several variants of the Time-Dependent Variational Principle (TDVP) method for time evolution (including novel bond-adaptive one-site algorithms). The package also provides strong support for the measurement of single and multi-site observables, as well as the storing and logging of data, which makes it a useful tool for the study of many-body physics. It currently handles long-range interactions, time-dependent Hamiltonians, multiple environments, bosonic and fermionic environments, and joint system-environment observables.

I. INTRODUCTION

The description of quantum systems as interacting with an uncontrolled external environment is central in quantum physics^{1–4}, quantum chemistry^{5,6}, and quantum biology⁷. In numerous settings there is a need to go beyond the approximation of a weak coupling between the system and its environment, and beyond the Markovian approximation (i.e. memoryless environment)⁸. In these cases analytical calculations are typically not available, and one has to rely on numerical methods. However, direct many-body calculations are not tractable because of the exponential growth of the underlying Hilbert space dimension with the size of joint {System + Environment}. Moreover, methods based on reduced density techniques also become computationally expensive as the memory time and complexity (rank) of the system-environment interactions increase. In response, several numerical methods to simulate open quantum system (OQS) dynamics in this non-perturbative regime have been developed, such as the Time Evolving Matrix Product Operator (TEMPO)^{9,10} method (implemented in the OQuPy package¹¹), the Automated Compression of Environments (ACE) al-

gorithm^{12–14}, the Hierarchical Equations Of Motions (HEOM) technique^{15,16} (implemented in the QuTip package¹⁷), the Multi-Layer–Multi-Configuration Time-Dependent Hartree (ML-MCTDH) method¹⁸ (implemented in the Quantics package^{19,20}), collision models²¹, the Dissipation-Assisted Matrix Product Factorization (DAMPF) method²² and the T-TEDOPA technique^{23–28} to name a few. TEMPO, ACE, HEOM, and T-TEDOPA belong to the specific class of numerically exact methods (i.e. methods for which the numerical error can be estimated exactly in terms of the convergence parameters). To our knowledge, `MPSDynamics.jl` is currently the only maintained package available implementing the T-TEDOPA method in Julia.

In the T-TEDOPA method, the curse of dimensionality preventing a direct resolution of the Schrödinger equation for the joint {System + Environment} state is circumvented using a tensor network representation of the joint quantum state^{29–31}. The particularity of T-TEDOPA is to treat the system and its environment on an equal footing, thus leaving the internal dynamics of the environment open to inspection^{26,32,33}. The accuracy of this representation is controlled by the size of the bond dimension D of the tensor network, and the number of

environmental modes that need to be taken into account is determined by the total simulation time.

MPSDynamics.jl has already been used in several scientific publications on quantum thermodynamics³⁴, electron transfer³⁵, the calculation of linear absorption spectra^{36,37}, information backflow^{38,39}, dynamical equilibrium states and the formation of polarons³³, and nanoscale dissipation³⁹. Moreover the T-TEDOPA framework is currently used by several research groups to work on projects in quantum photo-chemistry⁴⁰, multi-dimensional non-linear optical spectra⁴¹ or exact simulations of pigment protein complexes⁴². Previous applications of the TEDOPA formalism have also been used to look at ground state properties of strongly interacting system-environment properties, where this approach has strong links to the well-established MPS implementations of DMRG and NRG^{43,44}.

We also note here that the many body T-TEDOPA approach to open quantum systems has also recently been implemented on noisy intermediate-scale quantum computers^{45–47}.

The MPSDynamics.jl package is distributed under the GPL-3.0 free software license and is publicly available on Github at <https://github.com/shareloqs/MPSDynamics>. A Zenodo record is also publicly available⁴⁸.

In Sec. II we give an overview of the package. In Sec. III we summarize the important concepts of the T-TEDOPA method. Section IV presents short and simple examples using the package. The instruction to download and install MPSDynamics.jl are provided in Sec. V. Future developments of the package are discussed in Sec. VI.

II. PACKAGE OVERVIEW

In this section we give an overview of the abilities and usage of the package. For a more detailed explanation, a documentation is provided with the package at <https://shareloqs.github.io/MPSDynamics/>, and several example scripts are available in the package (see on Github). For a detailed presentation of the underlying theory of T-TEDOPA in the formalism and context used in MPSDynamics.jl, we refer the reader to the thesis of A. Dunnett⁴⁹.

A. States and Hamiltonians

A simulation with MPSDynamics.jl requires a MPS or a TTN representing the initial wave-function and a Matrix Product Operator (MPO) (or accordingly a TTN) representing the Hamiltonian. Figure 1 shows diagrammatic representations of these objects. The package contains various functions for generating MPSs and MPOs used for simulating generic models such as the spin-boson

model, the independent boson model, XYZ Hamiltonians, spin chains or tight-binding models for instance; but no attempt is made to be comprehensive. Methods are provided to construct MPS and TTN representations of standard states such as product states, Fock states, displaced states, or single-particle electronic states for instance. For generic MPO construction, one can use the ITensors.jl package⁵⁰ and convert the resulting object into a form compatible with MPSDynamics.jl using built-in functions. One can also directly construct MPO tailored for the problem at hand. The nature of the Hamiltonian can be fairly generic, for instance long-range interactions³⁸, time-dependent Hamiltonians, bosonic and fermionic environments, multiple environments^{35,36,38,39} can be handled. The fact that TTN with an arbitrary number of branches are supported means that the coupling to several environments (more than 2) at different temperatures – or even of a different nature – can be simulated.

For finite-temperature simulations, and experimental or *ab initio* bath spectral densities (SDs), an integrated implementation of the ORTHOPOL package⁵¹ is provided in order to generate the chain parameters needed to describe the chain-mapped environment (see Sec. III)^{28,52}.

B. Time-Evolution

The time-evolution methods currently implemented belong to the family of Time-Dependent Variational Principle (TDVP)⁵³. The central point of this method, in the modern tensor-network formulation, is that instead of solving the Schrödinger equation and then truncating the MPS representation of the quantum state, one can solve the equations of motion projected into a space of restricted bond dimension^{53,54}. The major advantage of this method is that it naturally preserves the unitarity of the time evolution and conserves the energy (except in its two-site implementation). Three variants of TDVP are implemented in the MPSDynamics.jl package:

- one-site TDVP on TTN and MPS (TDVP1),
- two-site TDVP on MPS (TDVP2),
- a variant of one-site TDVP with adaptive update of bond-dimensions on MPS (DTDVP)⁵⁵.

The main advantage of the one-site TDVP (TDVP1) algorithm is that it preserves the unitarity of the MPS during the time evolution. Its main problem, conversely, is that the time evolution is constrained to happen on a manifold constituted by tensors of fixed bond dimension, a quantity closely related to the amount of entanglement in the MPS, and such a bond dimension has therefore to be fixed before the beginning of the time evolution. This strategy will necessarily be non optimal: the unavoidable growth of the bond dimensions required to describe the quantum state at later times should ideally mirror the entanglement growth induced by the time evolution. The




FIG. 1. (a) A multi-partite quantum state ψ described by the coefficients $\psi_{i_1 i_2 \dots i_N}$ can be written as a Matrix Product State (MPS). A MPS is a product of (at-most) rank-3 tensors $T_n^{\alpha_{n-1} \alpha_n}$ where $\dim(i_n) = d$ is the local Hilbert space dimension, and $\dim(\alpha_n) = D$ is the so called ‘bond dimension’. (b) Tree tensor networks (TTN) can be used to represent quantum states and Hamiltonians. TTN are especially relevant for systems interacting with more than two environments. (c) Similarly, operators (such as the Hamiltonian H) can be decomposed into a product of rank-4 tensors $W_n^{w_{n-1} w_n}$. This decomposition is called a Matrix Product Operator (MPO).

two-site implementation (TDVP2) operates in a similar way to Time-Evolving Block-Decimation (TEBD), and allows for a similar dynamical growth of the bond dimensions, and therefore better describes the entanglement in the MPS. It suffers however of other drawbacks: first of all, a truncation error is introduced (by the means of an SVD decomposition), which entails a loss of unitarity of the time-evolved MPS. Moreover, TDVP2 has worse scaling properties $\mathcal{O}(d^3)$ with the size of the local dimensions d of the MPS. This is a major issue when dealing with bosons in non-perturbative OQS problems that require large local Hilbert space dimensions d . The DTDVP algorithm combines the best features of TDVP1 and TDVP2: it preserves unitarity, it has the same scaling properties $\mathcal{O}(d^2)$ as TDVP1, and it adapts the bond dimensions to the entanglement evolution at each site and at each time-step. DTDVP does not suffer from a truncation error, but introduces only a projection error that can be controlled with a tolerance parameter.

A graphical summary of the package workflow is presented in Fig. 2. We would like to highlight that, to our knowledge, `MPSDynamics.jl` is the only package implementing all three TDVP variants for MPS and TTN. The elementary tensor operations are implemented in all cases using the `TensorOperations.jl` package⁵⁶.

C. Observables and data processing

Local one-site and two-site observables, as well as non-local two-site observables, can be efficiently computed for each time-step of the time-evolution. System and environment observables can be computed, as well as system-

and-environment non-local observables.

The data (i.e. time series of observables) is stored in the JLD format which is based on HDF5. Loading the data in Julia using the `JLD.jl` package will recover the full type information of the Julia variables that were stored. At the same time the HDF5 format is natively supported across many platforms and languages (e.g. Python, Mathematica).

III. T-TEDOPA IN A NUTSHELL

Let us consider the following Hamiltonian where an unspecified system couples linearly to a continuum of independent bosonic modes (normal modes)

$$\hat{H} = \hat{H}_S + \int_0^\infty \omega \hat{a}_\omega^\dagger \hat{a}_\omega d\omega + \hat{A}_S \int_0^\infty \sqrt{J(\omega)} (\hat{a}_\omega + \hat{a}_\omega^\dagger) d\omega \quad (1)$$

where we consider the natural units such that $\hbar = 1$. We can introduce a unitary transformation of the continuous normal modes \hat{a}_ω to an infinite discrete set of interacting modes \hat{b}_n ⁵²

$$\hat{a}_\omega = \sum_{n=0}^{\infty} U_n(\omega) \hat{b}_n = \sum_{n=0}^{\infty} \sqrt{J(\omega)} P_n(\omega) \hat{b}_n, \quad (2)$$

where $P_n(\omega)$ are orthonormal polynomials such that

$$\int_0^\infty P_n(\omega) P_m(\omega) J(\omega) d\omega = \delta_{n,m}. \quad (3)$$

Note that the orthonormality of the polynomials ensures the unitarity of the transformation defined in




FIG. 2. Package workflow diagram: the user provides a state, a Hamiltonian, observables of interest, pick a time-evolution methods, and, when the script stopped running, can analyse the dynamics of the observables.

Eq. (2). The mapping from a continuous set of modes to a (still infinite) discrete set might seem counter-intuitive, however it is a direct consequence of the separability of the underlying Hilbert space. Under this transformation, the Hamiltonian in Eq. (1) becomes

$$\hat{H} = \hat{H}_S + \sum_{n=0}^{\infty} \varepsilon_n \hat{b}_n^\dagger \hat{b}_n + t_n (\hat{b}_{n+1}^\dagger \hat{b}_n + \text{h.c.}) + c_0 \hat{A}_S (\hat{b}_0 + \hat{b}_0^\dagger). \quad (4)$$

Hence, this mapping transforms the normal bath Hamiltonian into a tight-binding Hamiltonian with on-site energies ε_n and hopping energies t_n . Another important consequence of this mapping is that now the system only interacts with the first mode $n = 0$ of the chain-mapped environment. The chain coefficients ε_n , t_n , and the coupling c_0 depend solely on the SD (see Ref.⁵²). This makes chain mapping a tool of choice for describing systems coupled to environment with highly structured SD (e.g. experimentally measured or calculated *ab initio*)^{36,41,42,57,58}. In this new representation, the Hamiltonian in Eq. (4) has naturally a 1D

chain topology. This makes its representation as a MPO and the representation of the joint {System + Environment} wave-function as a MPS suited^{29,54}. The orthogonal polynomial-based chain mapping and the subsequent representation of the joint wave-function as a MPS (and the operators as MPO) are the building blocks of the T-TEDOPA method, one of the state-of-the-art numerically exact method to simulate the dynamics of open quantum systems especially in the non-Markovian, non-perturbative regimes both at zero and finite temperatures^{23,24,28,35,38}. A peculiarity of this method is that the finite temperature case is handled with an effective zero-temperature {System + Environment} but with a temperature-dependent SD²⁸. This ‘trick’ enables the wave-function description to be kept at finite-temperature at the moderate cost of increasing the number of environmental modes, with the maximal extension being a doubling of the number of modes for very high temperatures.

IV. EXAMPLES

We briefly present a few examples of applications to showcase the package workflow and illustrate its possibilities. Further example scripts with extended comments can be found on Github. The following examples correspond to the current release (version 1.1) of the package.

In all the examples we first load the `MPSDynamics.jl` package:

```
using MPSDynamics
```

A. Pure dephasing

Let us consider the independent boson model Hamiltonian

$$\hat{H} = \frac{\hat{\sigma}_z}{2} \left(\Delta E + \int_0^{+\infty} \sqrt{J(\omega)} (\hat{a}_\omega + \hat{a}_\omega^\dagger) d\omega \right) + \int_0^{+\infty} \omega \hat{a}_\omega^\dagger \hat{a}_\omega d\omega \quad (5)$$

where ΔE is the energy gap of a two-level system (TLS), $\hat{\sigma}_z$ is the third Pauli matrix, \hat{a}_ω^\dagger creates a bosonic excitation at frequency ω . We consider the case of an Ohmic SD $J(\omega) = 2\alpha\omega_c (\omega/\omega_c)^s H(\omega_c - \omega)$ where α is a dimensionless coupling strength, s is the so called Ohmicity, $H(\omega)$ is the Heaviside step function, and ω_c is the bath cut-off frequency. The system initial state is a superposition of spin- \uparrow and spin- \downarrow states $|\psi\rangle = \frac{1}{\sqrt{2}} (|\uparrow\rangle + |\downarrow\rangle)$, and the environment is in a Gibbs state at the inverse temperature $\beta = (k_B T)^{-1}$.

We first compute the chain coefficients of the environment (truncated to N modes) that we will use to define

the chain-mapped version of the Hamiltonian in Eq. (5).

```
# chain parameters, i.e. on-site energies
↳ e_i, hopping energies t_i, and
↳ system-chain coupling c_0
cpars = chaincoeffs_finiteT(N, β; α=α, s=s)
```

At zero temperature analytical expressions of the coefficients can be calculated²⁴, and are given by the function

```
cpars = chaincoeffs_ohmic(N, α, s)
```

We use these coefficients and the local dimensionality d of the N bosonic modes to set up the MPO representation of the Hamiltonian using a built-in function

```
H = puredephasingmpo(ΔE, d, N, cpars)
```

The initial state is a product state of the system state $|\psi\rangle$ and the vacuum state for N environmental modes

```
# Initial TLS in a superposition of up and
↳ down
ψ = zeros(2)
ψ[1] = 1/sqrt(2)
ψ[2] = 1/sqrt(2)

# MPS representation of |ψ⟩|Vacuum⟩
A = productstatemps(physdims(H), state=[ψ,
↳ fill(unitcol(1,d), N)...])
```

We define the observables of interest

```
ob1 = OneSiteObservable("sx", sx, 1)
ob2 = OneSiteObservable("sy", sy, 1)
ob3 = OneSiteObservable("sz", sz, 1)
```

and we choose a bond dimension D and a time-evolution method.

```
method = :TDVP1 # time-evolution method
```

```
D = 2 # MPS bond dimension
```

We are now all set to run the simulation

```
A, dat = runsim(dt, tfinal, A, H;
name = "pure dephasing model at
↳ temperature β = $(β)",
method = method,
obs = [ob1, ob2, ob3],
convobs = [ob1],
params = @LogParams(ΔE, N, d, α,
↳ s, β),
convparams = D,
reduceddensity = true,
```

```
verbose = false,
save = true,
plot = true,
);
```

The data is stored in dictionaries

```
julia> dat["data"]
Dict{String, Any} with 4 entries:
"sx" => [1.0, 0.912818, 0.741759,
↳ 0.605797, 0.528792, 0.492497, 0.47976...
"sz" => [0.0, 0.0871825, 0.25824,
↳ 0.394201, 0.471207, 0.507503, 0.52023...
"sy" => [0.0, -0.0133489, -0.0588887,
↳ -0.0858181, -0.0759996, -0.048539...
"times" => [0.0, 0.0005, 0.001, 0.0015,
↳ 0.002, 0.0025, 0.003, 0.0035, 0.00...
```

Figure 3 (a) shows the simulation results for zero and finite temperatures compared with the analytical results for a MPS of fixed bond dimension. Taking advantage of the wave-function framework of our MPS method we can also access environmental degrees of freedom, for instance Fig. 3 (b) shows the environmental correlations $\langle \hat{a}_\omega^\dagger \hat{a}_{\omega'}^\dagger \rangle(t) - \langle \hat{a}_\omega^\dagger \rangle(t) \langle \hat{a}_{\omega'}^\dagger \rangle(t)$. Local and non-local bath observables can be easily probed within the different examples.

B. Time-dependent Hamiltonian

Time-dependent Hamiltonians can be seamlessly used in simulation. Let us consider a two-level system driven by an external field

$$\hat{H}_S(t) + \hat{H}_{\text{int}} = \frac{\omega_0}{2} \hat{\sigma}_z + (\Delta + \epsilon \sin(\omega_{\text{drive}} t)) \hat{\sigma}_x + \hat{\sigma}_x \int_0^{+\infty} \sqrt{J(\omega)} (\hat{a}_\omega + \hat{a}_\omega^\dagger) d\omega. \quad (6)$$

One just starts by defining the *time-independent* part of the Hamiltonian

```
H = spinbosonmpo(ω0, Δ, d, N, cpars) # MPO
↳ representation of the Hamiltonian
```

and creates separately list containing the *time-dependent* Hamiltonian at the different time-steps of the simulation

```
Ht = [ε*sx*sin(ωdrive*tstep) for tstep in
↳ timelist] # Time-dependent Hamiltonian
↳ term
```

Then the simulation is run specifying which site of the MPO is time dependent

```

A, dat = runsim(dt, tfinal, A, H;
  name = "Driving field on ohmic
  ↪ spin boson model",
  method = method,
  obs = [ob1],
  convobs = [ob1],
  params = @LogParams(N, d, α, Δ,
  ↪ ω0, s),
  convparams = D,
  timedep = true, # the Hamiltonian
  ↪ is time dependent
  Ndrive = 1, # the first site of
  ↪ the MPS/MPO (i.e. the system)
  ↪ is concerned
  Htime = Ht, # list of
  ↪ time-dependent terms
  verbose = false,
  save = true,
  plot = true,
  );

```

Example simulation results obtained with this script are presented in Fig. 3 (c) showing the non-trivial dynamics of $\hat{\sigma}_z$ and the growth of the bond dimension when using DTDVP.

C. Long-range system-chain couplings

If one considers an interaction Hamiltonian with space-dependent coupling coefficients such as

$$\hat{H}_{\text{int}} = \sum_r \hat{A}_r \int_{-\infty}^{+\infty} \sqrt{J(\omega_k)} \left(\hat{a}_k e^{ikr} + \hat{a}_k^\dagger e^{-ikr} \right) dk, \quad (7)$$

the chain-mapped Hamiltonian becomes long-ranged and describes a so called *correlated* environment³⁸

$$\hat{H}_{\text{int}} = \sum_{r,n} \hat{A}_r \left(\gamma_n(r) \hat{b}_n + \gamma_n(r)^* \hat{b}_n^\dagger \right), \quad (8)$$

where the new system-chain coupling coefficients are given by $\gamma_n(r) = \int_{\mathbb{R}} J(\omega_k) e^{ikr} P_n(k) dk$. This generic type of coupling is handled through a built-in method generating the environment MPO providing a list of system component positions R

```

Hchain = correlatedenvironmentmpo(R, N, d,
  ↪ chainparams=cparams,
  ↪ fnamecc="couplingcoeffs.csv", s=s, α=α)

```

The coupling coefficients $\gamma_n(r)$ can be provided by the user. If not, they will be computed and stored in order to be reused in the future.

D. Fermionic environments

In addition to bosonic environments, the methods implemented in the code also allow to deal with fermionic ones. More specifically, it is possible to implement the MPO corresponding to the single impurity Anderson model⁵⁹, where the OQS is an impurity coupled to empty/filled leads, and the fermions are considered to be spinless. In the code, the MPO is initialised as follows

```

chainparams1 = chaincoeffs_fermionic(N, β,
  ↪ 1.0; ε, J, save=false) # empty
chainparams2 = chaincoeffs_fermionic(N, β,
  ↪ 2.0; ε, J, save=false) # filled
H = tightbinding_mpo(N, εd, chainparams1,
  ↪ chainparams2)

```

Note that the chain coefficients are computed with a thermofield-like⁶⁰ transformation, which allows to perform finite temperature simulations⁶¹.

E. Non-adiabatic dynamics on continuous energy landscapes

The MPS formalism can also be used for physical chemistry problems with energy surfaces described along a coordinate⁴⁰. For instance, a proton transfer model is implemented with the description in space of a two-level system. The introduction of a reaction coordinate (RC) allows to recover the double-well viewpoint as well as the diabatic and adiabatic basis. Wave-packet dynamics and non-adiabatic transitions can easily be simulated and analyzed with the reduced density matrix. Built-in methods allow to couple environments to this extended system such as the MPO

```

H = protontransfermpo(ω0e, ω0k, x0e, x0k, Δ,
  ↪ dFockRC, d, N, cparams, cparamsRC, λreorg)

```

After the dynamics, the reduced density matrix can be expressed in the reaction coordinate dimension, resulting in a visualisation of the wave-packet in space, as shown in Fig. 3 (d).

F. Tree Tensor Networks

Using TTN, several independent environments can be included. A TTN is a generalisation of the MPS wherein each site, instead of being connected to only one other may be connected to any arbitrary number of child sites (see Fig. 1). The sites of a TTN are usually referred to as nodes, and the first node is known as the head-node. A TTN is constructed by adding branches to specified nodes. If one considers a TLS coupled with three bosonic environments, as has been done, for instance, to model the linear absorption spectrum of Methylene blue³⁶ and




FIG. 3. (a) Dynamics of the coherence in a pure dephasing model (Sec. IV A) at zero and finite temperatures for the analytical results and simulations performed with a MPS of bond dimension $D = 2$. (b) As we have access to the whole joint {System + Bath} state we can compute non-trivial bath observables such as the environmental correlations $\langle \hat{a}_\omega^\dagger \hat{a}_{\omega'}^\dagger \rangle(t) - \langle \hat{a}_\omega^\dagger \rangle(t) \langle \hat{a}_{\omega'}^\dagger \rangle(t)$. (c) Dynamics of the population of a dissipative TLS driven by an external time-dependent field (Sec. IV B). (Inset) The adaptive one-site TDVP method enables the bond dimension to be increased on the fly. (d) Reduced density matrix expressed in the reaction coordinate dimension (Sec. IV E) obtained from the MPS representing the {electronic system + RC + bath}.

the absorption and emission spectra of Proflavin in a variety of different solvents³⁷, the TTN can easily be constructed as follows

```
# Built-in TTN Hamiltonian with 3 baths
H = methylbluempo(e1, e2, δ, N1, N2, N3, d1,
  ↪ d2, d3, cpar1, cpar2, cpar3)

# Initial TTN state where the head-node (the
  ↪ system) is connected to 3 branches
  ↪ corresponding to 3 chain-mapped
  ↪ environments
A = TreeNetwork(Any[productstatemps(1, 2, D,
  ↪ state=:FullOccupy)])
addtree!(A, 1, TreeNetwork(productstatemps(N1,
  ↪ d1, D, state=:Vacuum)))
addtree!(A, 1, TreeNetwork(productstatemps(N2,
  ↪ d2, D, state=:Vacuum)))
addtree!(A, 1, TreeNetwork(productstatemps(N3,
  ↪ d3, D, state=:Vacuum)))
```

The `runsim` method can then be called as usual.

V. DOWNLOAD AND INSTALLATION

The package may be straightforwardly downloaded and installed by typing the following command into a Julia REPL

```
julia> ]
pkg> add
  ↪ https://github.com/shareloqs/MPSDynamics.git
```

VI. CONCLUSION

The open-source `MPSDynamics.jl` package is a user-friendly package for the zero-and-finite temperature simulation of the dynamics of many-body systems using

tensor-network techniques. The package is primarily designed for the study of open quantum systems in the non-Markovian, non-perturbative, regime. Nevertheless, its applicability is wider by construction (e.g. many-body physics, non-equilibrium dynamics). Implementing the state-of-the-art numerically exact T-TEDOPA method, `MPSDynamics.jl` enables to keep the full {System + Environment} information, and to put under scrutiny the environmental degrees of freedom. The package naturally handles long-range interactions, time-dependent Hamiltonians, bosonic and fermionic environments, multiple environments, and correlated initial states. Several widely used model Hamiltonians are already implemented as built-in functions (see methods section of the different MPOs in the online documentation for a complete list), and we encourage users to push their particular models of interest to make this a live repository for timely open system problems. The time-evolution methods conserve the unitarity of the dynamics and are adaptive, and will also be updated to follow new theoretical advances in time-evolution methods, particularly in the case of tree tensor networks, where long-range interactions might be tackled more effectively with entanglement renormalization methods (work in progress)²⁷. A straight forward implementation of DMRG-like methods to find interacting ground states will also allow dynamics arising from highly correlated/entangled system-environment ground states to be explored in the future, which is a major challenge for reduced density matrix approaches, due to the inherently non-Gaussian nature of the bath in such initial conditions¹. The package and its documentation are available on Github at <https://github.com/shareloqs/MPSDynamics>, and several example scripts are provided. Everyone interested is welcome to contribute to development of the package. Interested readers can either contact the authors by e-mail or open an issue on Github. `MPSDynamics.jl` is developed continuously and future releases will be devoted to its further improvement. Future features will include integration on GPUs³⁷, better parallelization, automated test units, ‘black box’ computation of continuous wave

and time-resolved optical spectra from *ab initio* modelling, and Markovian closures⁶².

ACKNOWLEDGMENTS

TL acknowledges support from the ERC Synergy grant HyperQ (Grant No. 856432) and the EU project SPINUS (Grant No. 101135699).

BLD acknowledges iSiM (Initiative Sciences et ingénierie moléculaires) from the Alliance Sorbonne Université for funding.

AR acknowledges support from the ANR grants HAMROQS and MECAFLUX (French Research Agency) and by Plan France 2030 through the project ANR-22-PETQ-0006.

AWC acknowledges ANR Project RadPolimer ANR-22-CE30-0033.

DATA AVAILABILITY STATEMENT

¹H.-P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, 2007).

²A. Rivas and S. F. Huelga, *Open Quantum Systems: An Introduction*, 1st ed., SpringerBriefs in Physics (Springer Berlin, Heidelberg, 2011).

³U. Weiss, *Quantum Dissipative Systems*, 4th ed. (World Scientific, 2012).

⁴K. Blum, *Density Matrix Theory and Applications*, 3rd ed., Springer Series on Atomic, Optical, and Plasma Physics, Vol. 64 (Springer, 2012).

⁵J. C. Tully, “Perspective: Nonadiabatic dynamics theory,” *The Journal of Chemical Physics* **137**, 22A301 (2012).

⁶F. Agostini and B. F. E. Curchod, “Different flavors of nonadiabatic molecular dynamics,” *WIREs Computational Molecular Science* **9**, e1417 (2019).

⁷M. Mohseni, Y. Omar, G. S. Engel, and M. B. Plenio, *Quantum Effects in Biology* (Cambridge University Press, 2014).

⁸L. Li, M. J. W. Hall, and H. M. Wiseman, “Concepts of quantum non-Markovianity: A hierarchy,” *Physics Reports Concepts of quantum non-Markovianity: A hierarchy*, **759**, 1–51 (2018).

⁹A. Strathearn, P. Kirton, D. Kilda, J. Keeling, and B. W. Lovett, “Efficient non-Markovian quantum dynamics using time-evolving matrix product operators,” *Nature Communications* **9**, 3322 (2018).

¹⁰A. Strathearn, *Modelling Non-Markovian Quantum Systems Using Tensor Networks*, Springer Theses (Springer International Publishing, 2020).

¹¹Tempo Collaboration, G. E. Fux, P. Fowler-Wright, D. Gribben, J. Beckles, and P. Kirton, “OQuPy: A Python 3 package to efficiently compute non-Markovian open quantum systems,” (2020), 10.5281/zenodo.7243607.

¹²M. Cygorek, M. Cosacchi, A. Vagov, V. M. Axt, B. W. Lovett, J. Keeling, and E. M. Gauger, “Simulation of open quantum systems by automated compression of arbitrary environments,” *Nature Physics*, 1–7 (2022), publisher: Nature Publishing Group.

¹³M. Cygorek, J. Keeling, B. W. Lovett, and E. M. Gauger, “Sub-linear Scaling in Non-Markovian Open Quantum Systems Simulations,” *Physical Review X* **14**, 011010 (2024).

¹⁴M. Cygorek, “mcygorek/ACE: ACE version May 2024,” (2024).

¹⁵Y. Tanimura, “Numerically “exact” approach to open quantum dynamics: The hierarchical equations of motion (HEOM),” *The Journal of Chemical Physics* **153**, 020901 (2020), publisher: American Institute of Physics.

¹⁶E. Mangaud, A. Jaouadi, A. Chin, and M. Desouter-Lecomte, “Survey of the hierarchical equations of motion in tensor-train format for non-markovian quantum dynamics,” *The European Physical Journal Special Topics* **232**, 1847–1869 (2023).

¹⁷J. R. Johansson, P. D. Nation, and F. Nori, “QuTiP 2: A Python framework for the dynamics of open quantum systems,” *Computer Physics Communications* **184**, 1234–1240 (2013).

¹⁸H.-D. Meyer, “Studying molecular quantum dynamics with the multiconfiguration time-dependent Hartree method: Multiconfiguration time-dependent Hartree,” *Wiley Interdisciplinary Reviews: Computational Molecular Science* **2**, 351–374 (2012).

¹⁹G. Worth, “Quantics: A general purpose package for quantum molecular dynamics simulations,” *Computer Physics Communications* **248**, 107040 (2020).

²⁰G. Worth, “Quantics v1,” *Mendeley Data* (2019), 10.17632/x9dpc2r5c.1.

²¹F. Ciccarello, S. Lorenzo, V. Giovannetti, and G. M. Palma, “Quantum collision models: Open system dynamics from repeated interactions,” *Physics Reports Quantum collision models: Open system dynamics from repeated interactions*, **954**, 1–70 (2022).

²²A. D. Somoza, O. Marty, J. Lim, S. F. Huelga, and M. B. Plenio, “Dissipation-Assisted Matrix Product Factorization,” *Phys. Rev. Lett.* **123**, 100502 (2019).

²³J. Prior, A. W. Chin, S. F. Huelga, and M. B. Plenio, “Efficient Simulation of Strong System-Environment Interactions,” *Physical Review Letters* **105**, 050404 (2010).

²⁴M. Woods, M. Cramer, and M. Plenio, “Simulating Bosonic Baths with Error Bars,” *Physical Review Letters* **115**, 130401 (2015).

²⁵S. Oviedo-Casado, J. Prior, A. Chin, R. Rosenbach, S. Huelga, and M. Plenio, “Phase-dependent exciton transport and energy harvesting from thermal environments,” *Physical Review A* **93**, 020102 (2016).

²⁶F. A. Schröder and A. W. Chin, “Simulating open quantum dynamics with time-dependent variational matrix product states: Towards microscopic correlation of environment dynamics and reduced system evolution,” *Physical Review B* **93**, 075105 (2016).

²⁷F. A. Schröder, D. H. Turban, A. J. Musser, N. D. Hine, and A. W. Chin, “Tensor network simulation of multi-environmental open quantum dynamics via machine learning and entanglement renormalisation,” *Nature communications* **10**, 1062 (2019).

²⁸D. Tamascelli, A. Smirne, J. Lim, S. F. Huelga, and M. B. Plenio, “Efficient Simulation of Finite-Temperature Open Quantum Systems,” *Physical Review Letters* **123**, 090402 (2019).

²⁹R. Orus, “A Practical Introduction to Tensor Networks: Matrix Product States and Projected Entangled Pair States,” *Annals of Physics* **349**, 117–158 (2014), arXiv: 1306.2164.

³⁰J. C. Bridgeman and C. T. Chubb, “Hand-waving and Interpretive Dance: An Introductory Course on Tensor Networks,” *Journal of Physics A: Mathematical and Theoretical* **50**, 223001 (2017), arXiv: 1603.03039.

³¹G. Evenbly, “Tensors.net: Everything you need to begin your exciting journey into the world of tensor networks!” *Tensors.net*

³²J. Del Pino, F. A. Schröder, A. W. Chin, J. Feist, and F. J. Garcia-Vidal, “Tensor network simulation of polaron-polaritons in organic microcavities,” *Physical Review B* **98**, 165416 (2018).

³³A. Riva, D. Tamascelli, A. J. Dunnett, and A. W. Chin, “Thermal cycle and polaron formation in structured bosonic environments,” *Physical Review B* **108**, 195138 (2023), publisher: American Physical Society.

³⁴A. J. Dunnett and A. W. Chin, “Matrix Product State Simulations of Non-Equilibrium Steady States and Transient Heat Flows in the Two-Bath Spin-Boson Model at Finite Temperatures,” *Entropy* **23**, 77 (2021), number: 1 Publisher: Multidisciplinary Digital Publishing Institute.

³⁵A. J. Dunnett and A. W. Chin, “Simulating Quantum Vibronic Dynamics at Finite Temperatures With Many Body Wave Functions at 0 K,” *Frontiers in Chemistry* **8** (2021),

- 10.3389/fchem.2020.600731, publisher: Frontiers.
- ³⁶A. J. Dunnett, D. Gowland, C. M. Isborn, A. W. Chin, and T. J. Zuehlsdorff, “Influence of non-adiabatic effects on linear absorption spectra in the condensed phase: Methylene blue,” *The Journal of Chemical Physics* **155**, 144112 (2021), arXiv:2107.06587 [physics, physics:quant-ph].
- ³⁷K. E. Hunter, Y. Mao, A. W. Chin, and T. J. Zuehlsdorff, “Environmentally driven symmetry breaking quenches dual fluorescence in proflavine,” *The Journal of Physical Chemistry Letters* **15**, 4623–4632 (2024).
- ³⁸T. Lacroix, A. Dunnett, D. Gribben, B. W. Lovett, and A. Chin, “Unveiling non-Markovian spacetime signaling in open quantum systems with long-range tensor network dynamics,” *Physical Review A* **104**, 052204 (2021).
- ³⁹T. Lacroix, B. W. Lovett, and A. W. Chin, “From Non-Markovian Dissipation to Spatiotemporal Control of Quantum Nanodevices,” *Quantum* **8**, 1305 (2024).
- ⁴⁰B. Le Dé, S. Huppert, R. Spezia, and A. W. Chin, “Extending non-perturbative simulation techniques for open-quantum systems to excited-state proton transfer and ultrafast non-adiabatic dynamics,” arXiv e-prints (2024), 10.48550/ARXIV.2405.08693.
- ⁴¹N. Lorenzoni, N. Cho, J. Lim, D. Tamascelli, S. F. Huelga, and M. B. Plenio, “Systematic Coarse Graining of Environments for the Nonperturbative Simulation of Open Quantum Systems,” *Phys. Rev. Lett.* **132**, 100403 (2024).
- ⁴²F. Caycedo-Soler, A. Mattioni, J. Lim, T. Renger, S. F. Huelga, and M. B. Plenio, “Exact simulation of pigment-protein complexes unveils vibronic renormalization of electronic parameters in ultrafast spectroscopy,” *Nat Commun* **13**, 2912 (2022).
- ⁴³A. W. Chin, J. Prior, S. F. Huelga, and M. B. Plenio, “Generalized polaron ansatz for the ground state of the sub-ohmic spin-boson model: An analytic theory of the localization transition,” *Physical Review Letters* **107**, 160601 (2011).
- ⁴⁴Z. Blunden-Codd, S. Bera, B. Bruognolo, N.-O. Linden, A. W. Chin, J. von Delft, A. Nazir, and S. Florens, “Anatomy of quantum critical wave functions in dissipative impurity problems,” *Physical Review B* **95**, 085104 (2017).
- ⁴⁵T. Nishi and K. Yamanouchi, “Simulation of a spin-boson model by iterative optimization of a parametrized quantum circuit,” *AVS Quantum Science* **6** (2024).
- ⁴⁶J. D. Guimarães, J. Lim, M. I. Vasilevskiy, S. F. Huelga, and M. B. Plenio, “Noise-assisted digital quantum simulation of open systems using partial probabilistic error cancellation,” *PRX Quantum* **4**, 040329 (2023).
- ⁴⁷J. D. Guimarães, M. I. Vasilevskiy, and L. S. Barbosa, “Digital quantum simulation of non-perturbative dynamics of open systems with orthogonal polynomials,” *Quantum* **8**, 1242 (2024).
- ⁴⁸A. J. Dunnett, T. Lacroix, A. Riva, and B. Le Dé, “shareloqs/MPSDynamics: v1.1,” (2024), 10.5281/zenodo.11400776.
- ⁴⁹A. Dunnett, *Tensor Network approaches to Open Quantum Systems at Finite Temperature with applications to Spectroscopy*, Ph.D. thesis, Sorbonne Université (2021).
- ⁵⁰M. Fishman, S. R. White, and E. M. Stoudenmire, “The ITensor Software Library for Tensor Network Calculations,” *SciPost Phys. Codebases* , 4 (2022).
- ⁵¹W. Gautschi, “Algorithm 726: ORTHPOL—a package of routines for generating orthogonal polynomials and Gauss-type quadrature rules,” *ACM Transactions on Mathematical Software* **20**, 21–62 (1994).
- ⁵²A. W. Chin, A. Rivas, S. F. Huelga, and M. B. Plenio, “Exact mapping between system-reservoir quantum models and semi-infinite discrete chains using orthogonal polynomials,” *Journal of Mathematical Physics* **51**, 092109 (2010), arXiv: 1006.4507.
- ⁵³J. Haegeman, J. I. Cirac, T. J. Osborne, I. Pižorn, H. Verschelde, and F. Verstraete, “Time-Dependent Variational Principle for Quantum Lattices,” *Physical Review Letters* **107**, 070601 (2011).
- ⁵⁴S. Paeckel, T. Köhler, A. Swoboda, S. R. Manmana, U. Schollwöck, and C. Hubig, “Time-evolution methods for matrix-product states,” *Annals of Physics* **411**, 167998 (2019).
- ⁵⁵A. J. Dunnett and A. W. Chin, “Efficient bond-adaptive approach for finite-temperature open quantum dynamics using the one-site time-dependent variational principle for matrix product states,” *Physical Review B* **104**, 214302 (2021).
- ⁵⁶L. Devos, M. V. Damme, J. Haegeman, and contributors, “TensorOperations.jl,” (2023), 10.5281/zenodo.3245496.
- ⁵⁷A. W. Chin, J. Prior, R. Rosenbach, F. Caycedo-Soler, S. F. Huelga, and M. B. Plenio, “The role of non-equilibrium vibrational structures in electronic coherence and recoherence in pigment–protein complexes,” *Nature Phys* **9**, 113–118 (2013), number: 2 Publisher: Nature Publishing Group.
- ⁵⁸A. M. Alvertis, F. A. Y. N. Schröder, and A. W. Chin, “Non-equilibrium relaxation of hot states in organic semiconductors: Impact of mode-selective excitation on charge transfer,” *J. Chem. Phys.* **151**, 084104 (2019).
- ⁵⁹P. W. Anderson, “Localized magnetic states in metals,” *Phys. Rev.* **124**, 41–53 (1961).
- ⁶⁰I. de Vega and M.-C. Bañuls, “Thermofield-based chain-mapping approach for open quantum systems,” *Phys. Rev. A* **92**, 052116 (2015).
- ⁶¹L. Kohn and G. E. Santoro, “Efficient mapping for anderson impurity problems with matrix product states,” *Phys. Rev. B* **104**, 014303 (2021).
- ⁶²A. Nüßeler, D. Tamascelli, A. Smirne, J. Lim, S. F. Huelga, and M. B. Plenio, “Fingerprint and Universal Markovian Closure of Structured Bosonic Environments,” *Phys. Rev. Lett.* **129**, 140604 (2022).