

Total Positivity of Quasi-Riordan Arrays

Tian-Xiao He¹ and Roksana Słowik²

¹Department of Mathematics, Illinois Wesleyan University,
Bloomington, IL 61702-2900, Indiana, USA

²Faculty of Applied Mathematics, Silesian University of Technology,
Kaszubska 23, 44-100 Gliwice, Poland

June 12, 2024

Abstract

In this paper the total positivity of quasi-Riordan arrays is investigated with use of the sequence characterization of quasi-Riordan arrays. Due to the correlation between quasi-Riordan arrays and Riordan arrays, this study is an in-depth discussion of the total positivity of Riordan arrays.

AMS Subject Classification: 05A15, 05A05, 15B36, 15A06, 05A19, 11B83.

Key Words and Phrases: Riordan array, Riordan group, Quasi-Riordan array, Almost-Riordan array, Quasi-Riordan group.

1 Introduction

Following Karlin [13] and Pinkus [19], an infinite matrix is called totally positive (abbreviately, TP), if its minors of all orders are nonnegative. An infinite sequence of nonnegative real numbers $(a_n)_{n \geq 0}$ is called a Pólya frequency sequence (abbreviately, PF), if its Toeplitz matrix

$$[a_{i-j}]_{i,j \geq 0} = \begin{bmatrix} a_0 & & & & \\ a_1 & a_0 & & & \\ a_2 & a_1 & a_0 & & \\ a_3 & a_2 & a_1 & a_0 & \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

is TP. We say that a finite sequence (a_0, a_1, \dots, a_n) is PF if the corresponding infinite sequence $(a_0, a_1, \dots, a_n, 0, \dots)$ is PF. Denote by \mathbb{N} the set of all nonnegative integers. A fundamental characterization for PF sequences is given by Schoenberg et al.[1, 2] (see also Karlin [13]), which states that a sequence $(a_n)_{n \geq 0}$ is PF if and only if its generating function can be written as

$$\sum_{n \geq 0} a_n t^n = C t^k e^{\gamma t} \frac{\prod_{j \geq 0} (1 + \alpha_j t)}{\prod_{j \geq 0} (1 - \beta_j t)}, \quad (1)$$

where $C > 0$, $k \in \mathbb{N}$, $\alpha_j, \beta_j, \gamma \geq 0$, and $\sum_{j \geq 0} (\alpha_j + \beta_j) < \infty$. In this case, the above generating function is called a Pólya frequency formal power series. For more preliminary materials and more relevant results see, for example, Brenti [6], Mao, Mu, and Wang [17] and Pinkus [19].

Let A and B be $m \times m$ and $n \times n$ matrices, respectively. Then we define the direct sum of A and B by

$$A \oplus B = \begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix}_{(m+n) \times (m+n)}. \quad (2)$$

Let us recall some basic definitions and introduce the notation.

First, the Riordan arrays (or Riordan matrices) are infinite, lower triangular matrices defined by the generating function of their columns. With the matrix multiplication, the set of all Riordan arrays form a group, called *the Riordan group* (see Shapiro, Getu, Woan and Woodson [22]).

More formally, let us consider the set of formal power series ring $\mathcal{F} = \mathbb{K}[[t]]$, where \mathbb{K} is the field \mathbb{R} or \mathbb{C} . The *order* of $f(t) \in \mathcal{F}$, $f(t) = \sum_{k=0}^{\infty} f_k t^k$ ($f_k \in \mathbb{K}$), is the minimum number $r \in \mathbb{N}$ such that $f_r \neq 0$, where $\mathbb{N} = \{0, 1, 2, \dots\}$ is the set of all natural numbers. We denote by \mathcal{F}_r the set of formal power series of order r . Let $g(t) \in \mathcal{F}_0$ and $f(t) \in \mathcal{F}_1$; the pair (g, f) defines the (*proper*) *Riordan array* $R = (d_{n,k})_{n,k \in \mathbb{N}} = (g, f)$ having

$$d_{n,k} = [t^n] g(t) f(t)^k \quad (3)$$

or, in other words, having $g f^k$ as the generating function of the k th column of (g, f) . Hence,

$$(g, f) := (g, g f, g f^2, g f^3, \dots), \quad (4)$$

where $g, g f, g f^2, g f^3 \dots$ are the generating functions of the 0th, 1st, 2nd, 3rd, \dots , columns of the matrix (g, f) , respectively. The *first fundamental theorem of Riordan arrays* can be represented as

$$(g(t), f(t))h(t) = g(t)(h \circ f)(t),$$

which can also be simplified to $(g, f)h = gh(f)$. Thus we immediately see that the usual row-by-column product of two Riordan arrays is also a Riordan array:

$$(g_1, f_1)(g_2, f_2) = (g_1g_2(f_1), f_2(f_1)). \quad (5)$$

The Riordan array $I = (1, t)$ is the identity matrix because its entries are $d_{n,k} = [t^n]t^k = \delta_{n,k}$.

Let $(g(t), f(t))$ be a Riordan array. Then its inverse is

$$(g(t), f(t))^{-1} = \left(\frac{1}{g(\bar{f}(t))}, \bar{f}(t) \right) \quad (6)$$

where $\bar{f}(t)$ is the compositional inverse of $f(t)$, i.e., $(f \circ \bar{f})(t) = (\bar{f} \circ f)(t) = t$. In this way, the set \mathcal{R} of all proper Riordan arrays forms a group (see [22]) called the Riordan group.

Here is a list of six important subgroups of the Riordan group (see [22, 21, 4]).

- the *Appell subgroup* $\{(g(t), t) : g \in \mathcal{F}_0\}$.
- the *Lagrange (associated) subgroup* $\{(1, f(t)) : f \in \mathcal{F}_1\}$.
- the *k-Bell subgroup* $\{(g(t), t(g(t))^k) : g \in \mathcal{F}_0\}$, where k is a fixed positive integer.
- the *hitting-time subgroup* $\{(tf'(t)/f(t), f(t)) : f \in \mathcal{F}_1\}$.
- the *derivative subgroup* $\{(f'(t), f(t)) : f \in \mathcal{F}_1\}$.
- the *checkerboard subgroup* $\{(g(t), f(t)) : g \in \mathcal{F}_0, f \in \mathcal{F}_1\}$, where g is an even function and f is an odd function.

The 1-Bell subgroup is referred to as the Bell subgroup for short, and the Appell subgroup can be considered as the 0-Bell subgroup if we allow $k = 0$ to be included in the definition of the k -Bell subgroup.

An infinite lower triangular matrix $[d_{n,k}]_{n,k \in \mathbb{N}}$ is a Riordan array if and only if a unique sequence $A = (a_0 \neq 0, a_1, a_2, \dots)$ exists such that for every $n, k \in \mathbb{N}$

$$d_{n+1,k+1} = a_0d_{n,k} + a_1d_{n,k+1} + \dots + a_nd_{n,n}. \quad (7)$$

This is equivalent to

$$f(t) = tA(f(t)) \quad \text{or} \quad t = \bar{f}(t)A(t). \quad (8)$$

Here, $A(t)$ is the generating function of the A -sequence. The first formula of (8) is also called the *second fundamental theorem of Riordan arrays*.

Moreover, there exists a unique sequence $Z = (z_0, z_1, z_2, \dots)$ such that every element in column 0 can be expressed as the linear combination

$$d_{n+1,0} = z_0d_{n,0} + z_1d_{n,1} + \dots + z_nd_{n,n}, \quad (9)$$

or equivalently,

$$g(t) = \frac{1}{1 - tZ(f(t))}, \quad (10)$$

in which and thoroughly we always assume $g(0) = g_0 = 1$, a usual hypothesis for proper Riordan arrays. From (10), we may obtain

$$Z(t) = \frac{g(\bar{f}(t)) - 1}{\bar{f}(t)g(\bar{f}(t))}.$$

A - and Z -sequence characterizations of Riordan arrays were introduced, developed, and/or studied in Merlini, Rogers, Sprugnoli, and Verri [18], Roger [20], Sprugnoli and the author [11], Jean-Louis and Nkwanta [12], Luzón, Morón, Prieto-Martinez [15, 16], and [10] as well as their references.

Secondly, we recall the definition of quasi-Riordan arrays given in [9].

Definition 1.1. [9] Let $g \in \mathcal{F}_0$ with $g(0) = 1$ and $f \in \mathcal{F}_1$. We call the following matrix a quasi-Riordan array and denote it by $[g, f]$.

$$[g, f] := (g, f, tf, t^2f, \dots), \quad (11)$$

where g, f, tf, t^2f, \dots are the generating functions of the 0th, 1st, 2nd, 3rd, \dots , columns of the matrix $[g, f]$, respectively. It is clear that $[g, f]$ can be written as

$$[g, f] = \begin{pmatrix} g(0) & 0 \\ (g - g(0))/t & (f/t, t) \end{pmatrix}, \quad (12)$$

where $(f, t) = (f, tf, t^2f, t^3f, \dots)$ and $(f/t, t)$ is an Appell Riordan array. Particularly, if $f = tg$, then the quasi-Riordan array $[g, tg] = (g, t)$, the Appell-type Riordan array.

The relationship between the Riordan arrays and quasi-Riordan arrays can be presented in the following theorem.

Theorem 1.2. [9] Let $(g, f) = (d_{n,k})_{n,k \geq 0}$ be a Riordan array, and let $([1] \oplus (g, f))$ and $[g, f]$ be defined by (2) and (11), respectively. Then (g, f) has the expression

$$(g, f) = [g, f]([1] \oplus (g, f)). \quad (13)$$

For an integer $r \geq 0$, denote the r -th truncation of a formal power series $h = \sum_{n \geq 0} h_n t^n$ by $h|_r := \sum_{n=0}^r h_n t^n$. For an integer $n \geq 0$, denote by $(g, f)_n$ the n -th order leading principle submatrix of the Riordan array $(g(t), f(t))_n$. This notation is introduced by Luzón et al in [14] and to the one used later by Luzón and her collaborators. Similarly, $[g, f]_n$ denotes the n -th order leading principle submatrix of the quasi-Riordan array $[g, f]$ defined by (11), namely

$$[g, f]_n = \begin{pmatrix} g(0) & 0 \\ ((g - g(0))/t)|_{n-1} & (f, t)_{n-1} \end{pmatrix}, \quad (14)$$

where the $n - 1$ -st truncation of $(g - g(0))/t|_{n-1}$ is

$$(g - g(0))/t|_{n-1} = \sum_{k=1}^{n-1} g_k t^{k-1},$$

and $(f, t)_{n-1} = (f, tf, t^2 f, \dots, t^{n-1} f)$. We call $[g, f]_n$ the n -th truncation of the quasi-Riordan array $[g, f]$.

Corollary 1.3. [9, 17] Let $(g, f) = (d_{n,k})_{n,k \geq 0}$ be a Riordan array with $g = \sum_{n \geq 0} g_n t^n$ and $f = \sum_{n \geq 1} f_n t^n$, and let $(g, f)_n$ be the n -th order leading principle submatrix of (g, f) . Then we have the recurrence relation of $(g, f)_n$ in the following form:

$$(g, f)_n = [g, f]_n([1] \oplus (g, f)_{n-1}). \quad (15)$$

Let $R = (g(t), f(t))$ be a Riordan array, where $g(t) = \sum_{n \geq 0} g_n t^n$ and $f(t) = \sum_{n \geq 1} f_n t^n$. If the lower triangular matrix

$$Q = \begin{bmatrix} g_0 & 0 & 0 & 0 & 0 & \cdots \\ g_1 & f_1 & 0 & 0 & 0 & \cdots \\ g_2 & f_2 & f_1 & 0 & 0 & \cdots \\ g_3 & f_3 & f_2 & f_1 & 0 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix} = (g(t), f(t), tf(t), t^2 f(t), \dots) \quad (16)$$

is totally positive (TP), then so is R [17]. This can be proved easily by using Corollary 1.3 and mathematical induction.

The question is when Q is totally positive. It is obvious that the Pólya frequency property for the formal power series g, f is not sufficient to ensure TP of Q .

Example 1.4. Let $g(t) = (1+t)^2$ and $f(t) = t/(1-t)$. Then both $g(t)$ and $f(t)$ are Pólya frequency, and (g, f) is TP. However, the minor of $[g, f]$

$$M_{0,1,2}^{1,2,3} = \det \begin{pmatrix} 2 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix} = -1 < 0.$$

Hence, $[g, f]$ is not TP although (g, f) is TP.

In this paper, we search for the conditions of TP quasi-Riordan arrays and the sequence characterization of TP quasi-Riordan arrays. From the last example, we know that the formal power series $g(t)$ plays an important role in identifying whether a quasi-Riordan array is TP. Hence, in the next section, we discuss the conditions and their equivalents for TP quasi-Riordan arrays $[g, f]$ with some special power series g . In Section 3, we study the sequence characterization of TP quasi-Riordan arrays. In particular, we present some generalizations of results given in [7]. More specifically, Theorems 3.2 and 3.4 in this section are generalizations of Theorem 2.1 and Theorem 2.3 of [7] for quasi-Riordan arrays. In Section 4 we will present some more examples with various g 's for TP and non-TP quasi-Riordan arrays $[g, f]$, respectively. From a list of examples showing in this section and the results in the previous sections, one may see a kind of independence of a TP quasi-Riordan array $[g, f]$ and the properties of that its formal power series g and f are the Pólya frequency formal power series.

2 TP Quasi-Riordan Arrays

Let us consider two special types of Riordan arrays and discuss their TP property and the TP property for the corresponding quasi-Riordan array.

If $g(t) = 1$, then from (12) we have

$$[g, f] = \begin{pmatrix} 1 & 0 \\ 0 & (f/t, t) \end{pmatrix}. \quad (17)$$

Hence,

Proposition 2.1. Let $f \in \mathcal{F}_1$. The following conditions are equivalent.

1. f is a Pólya frequency sequence.
2. The Riordan array $(f/t, t)$ is TP.
3. The Riordan array $(1, f)$ is TP.

4. The quasi-Riordan array $\left[1, \frac{f(t)}{t}\right]$ is TP.

Proof. Since $(f/t, t)$ is a Riordan array, from the definition of the TP matrices, we immediately obtain Proposition 2.1. \square

Consider the case when $g(t) = 1 + g_1t$ and $f(t) = f_1t + f_2t^2$. Then

$$[g, f] = \begin{bmatrix} 1 & 0 & & & \\ g_1 & f_1 & 0 & & \\ & f_2 & f_1 & & \\ & & f_2 & & \\ & \ddots & \ddots & \ddots & \ddots \end{bmatrix}.$$

Hence, from the proof of Proposition 2.6 of [7] we have

Corollary 2.2. *The quasi-Riordan array $[1 + g_1t, f_1t + f_2t^2]$ is TP if and only if $g_1, f_1, f_2 \geq 0$.*

Proposition 2.1 can be somewhat generalized to the following.

Proposition 2.3. *If $[g, f]$ is TP, then f is a Pólya frequency sequence.*

Clearly, the above necessary condition is absolutely not sufficient for total positivity of $[g, f]$. Primarily, if the terms of g are not nonnegative, then $[g, f]$ is definitely not TP, even if f is a Pólya frequency sequence. Moreover, it should also be emphasized again that, although the fact that both g and f are Pólya frequency sequences implies that the Riordan array (g, f) is TP [8], the analogous result for quasi-Riordan arrays does not hold as we have seen in Example 1.4.

3 Sequence Characterization of the TP of Quasi-Riordan Arrays

Let us now bring the attention to the fact that the quasi-Riordan array $[g, f]$ can be treated as almost-Riordan array $\left(g \middle| \frac{f(t)}{t}, t\right)$ [5]. From [3], we obtain the generating functions of the A -, Z -, and W -sequences of the latter one as follows:

$$A(t) = 1, \tag{18}$$

$$Z(t) = \frac{f(t) - z_0tg(t)}{f(t)} + z_0, \tag{19}$$

$$W(t) = \frac{(1 - w_0 t)g(t) - 1}{f(t)} + w_0, \tag{20}$$

where $z_0 = f_1$ and $w_0 = g_1$. For instance, for quasi-Riordan array $[1/(1 - t), t/(1 - t)]$, $A(t) = Z(t) = W(t) = 1$.

We call the following matrix the *production matrix* of the quasi-Riordan matrix $[g, f]$:

$$J = \begin{bmatrix} w_0 & z_0 & & & & \\ w_1 & z_1 & a_0 & & & \\ w_2 & z_2 & a_1 & a_0 & & \\ w_3 & z_3 & a_2 & a_1 & a_0 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix} \tag{21}$$

Proposition 3.1. *Let $[g, f]$ be a quasi-Riordan array, and let J be the matrix defined by (21). Then,*

$$[g, f]J = \overline{[g, f]}, \tag{22}$$

where $\overline{[g, f]}$ is the matrix by deleting the first row of $[g, f]$.

Proof. Equation (22) can be checked straightforwardly by using matrix multiplication and definition of quasi-Riordan array. □

Theorem 3.2. *Let J be a matrix defined by (21) and let R be the quasi-Riordan array defined by A -, Z -, and W -sequences. If J is TP, then so is R .*

Proof. It can be proved by using (22) and mathematical induction. □

Corollary 3.3. *Let $[g, f]$ and J be two matrices defined by (11) and (21), respectively. If J is TP, then so are $[g, f]$ and (g, f) .*

Proof. The corollary can be proved by using Theorems 1.2 and 3.2. □

Clearly, by (18)-(20) the matrix J shown above is of a special form. Namely

$$J = \begin{bmatrix} w_0 & z_0 & & & & \\ w_1 & z_1 & 1 & & & \\ w_2 & z_2 & 0 & 1 & & \\ w_3 & z_3 & 0 & 0 & 1 & \\ w_4 & z_4 & 0 & 0 & 0 & 1 \\ \vdots & \vdots & & & & \ddots \end{bmatrix}. \tag{23}$$

is TP.

Note that since the discriminant of the denominator $D(t)$ of the obtained g and f is equal to

$$\Delta = w_0^2 + z_1^2 - 2w_0z_1 + 4w_1z_0 = (w_0 - z_1)^2 + 4w_1z_0,$$

all the coefficients are nonnegative, so $D(t)$ has two real roots. Moreover, according to the assumption of $w_0z_1 - w_1z_0 \geq 0$, their sum $\frac{w_0+z_1}{w_0z_1-w_1z_0}$ and their product $\frac{1}{w_0z_1-w_1z_0}$ are nonnegative. Then, they must be nonnegative as well. Thus, f is of form (1), so by [1, 2, 13] this confirms that f must be Pólya frequency formal power series. If $z_1 = 0$, then by $w_0z_1 - w_1z_0 \geq 0$ we have $w_1z_0 = 0$, which implies $w_1 = 0$ due to $z_0 \neq 0$. Hence, if $z_1 = 0$, then $z_0 > 0$, $w_1 = 0$, and $w_0 > 0$, and all those make $[g, f] = \left[\frac{1}{1-w_0t}, \frac{z_0t}{1-w_0t} \right]$, where both f and g are Pólya frequency. If $z_1 > 0$, then g is not a Pólya frequency formal power series, although both $[g, f]$ and, hence, (g, f) are TP. This gives another counterexample for Item 2 in [24]. As what we have seen, if $z_1 > 0$, then g is not a Pólya frequency formal power series—its numerator has a positive root. However, as Proposition 3.2 holds, $[g(t), f(t)]$ is TP, so the coefficients of the series $g(t)$ are nonnegative. Moreover, we have already seen in Example 1.4 that, in general, fact that both series $g(t)$ and $f(t)$ are Pólya frequency is not sufficient for $[g, f]$ to be TP, so one could mistrust the necessity of this condition as well.

4 Counterexamples

Note that in Example 1.4 we have already presented a non-TP quasi-Riordan array $[g, f]$, where g and f are Pólya frequency power series. Here, in the first subsection, we give some more examples like that. In the second subsection, we give some more examples for TP of quasi-Riordan arrays $[g, f]$ with non-Pólya frequency power series g .

4.1 Quasi-Riordan arrays that are not TP

Proposition 4.1. *Let k_1, k_2 be natural numbers such that $k_2 > k_1$, $\alpha > 0$, and let $f(t) = \sum_{k \geq 1} f_k t^k$ be a Pólya frequency power series. If for some positive natural number n it holds $\alpha^{k_2-k_1} f_{k_1-n+1} > f_{k_2-n+1}$, then the quasi-Riordan array $\left[\frac{1}{1-\alpha t}, f(t) \right]$ is not TP, although both series g and f are Pólya frequency power series.*

Proof. Consider a minor of $\left[\frac{1}{1-\alpha t}, f(t)\right]$ that consists of the entries from the rows k_1, k_2 and the columns 0 and n . It is equal to

$$M_{0,n}^{k_1,k_2} = \alpha^{k_1} f_{k_2-n+1} - \alpha^{k_2} f_{k_1-n+1},$$

so by the assumption it is negative, and so $\left[\frac{1}{1-\alpha t}, f(t)\right]$ is not TP. \square

Example 4.2. *Let*

$$f(t) = \frac{t}{(1-2t)^2} = t + 4t^2 + 12t^3 + 32t^4 + 80t^5 + 192t^6 + \dots$$

Choosing $k_1 = 3, k_2 = 4, n = 1$, for $\alpha = 3$ we have

$$\alpha^{4-3} f_3 > f_4.$$

Hence,

$$\left[\frac{1}{1-3t}, \frac{t}{(1-2t)^2}\right] = \begin{bmatrix} 1 & & & & & & \\ 3 & 1 & & & & & \\ 9 & 4 & 1 & & & & \\ 27 & 12 & 4 & 1 & & & \\ 81 & 32 & 12 & 4 & 1 & & \\ \vdots & & & & & \ddots & \end{bmatrix}$$

is not TP, as

$$M_{3,4}^{0,1} = \begin{vmatrix} 27 & 12 \\ 81 & 32 \end{vmatrix} = -108 < 0,$$

although g and f are Pólya frequency power series.

In general, Proposition 4.1 implies the following.

Corollary 4.3. *For every Pólya frequency power series f , there exists a constant $\alpha > 0$ such that the quasi-Riordan array $\left[\frac{1}{1-\alpha t}, f(t)\right]$ is not TP, although both $\frac{1}{1-\alpha t}$ and $f(t)$ are Pólya frequency power series.*

Proof. Clearly, fixing arbitrarily indices $k_1, k_2, n, k_1 < k_2, n > 0$, it suffices to take

$$\alpha > \sqrt[k_2-k_1]{\frac{f_{k_2-n}}{f_{k_1-n}}}. \quad (24)$$

\square

Example 4.4. *Let*

$$f(t) = \frac{t(t+1)}{1-2t} = t + \sum_{n=2}^{\infty} (2^{n-2} + 2^{n-1}) t^n.$$

To find α appearing in Proposition 4.1 we consider the inequality

$$\alpha > \sqrt[k_2 - k_1]{\frac{f_{k_1 - n}}{f_{k_2 - n}}}.$$

Taking, for instance $k_2 = n + 2$, $k_1 = n + 1$, we get

$$\alpha > \frac{f_2}{f_1} = 3.$$

Thus, for any $\alpha > 3$ the quasi-Riordan $\left[\frac{1}{1-\alpha t}, f(t)\right]$ is not TP.

Series g , that in from Proposition 4.1 is equal to $\frac{1}{1-\alpha t}$, can be generalized to various other forms. A formula for this general form seems difficult to be found, yet let us present one more example.

Example 4.5. *Let $f(t)$ be a Pólya frequency power series and let*

$$g(t) = \frac{1}{(1-\alpha t)(1-\beta t)} = \sum_{n=0}^{\infty} \frac{\beta^{n+1} - \alpha^{n+1}}{\beta - \alpha} t^n \quad \text{with } \beta > \alpha,$$

i.e. $g(t)$ is also a Pólya frequency power series.

Looking at the minor

$$M_{0,1}^{1,2} = \begin{vmatrix} \frac{\beta^2 - \alpha^2}{\beta - \alpha} & f_0 \\ \frac{\beta^3 - \alpha^3}{\beta - \alpha} & f_1 \end{vmatrix}$$

it can be observed that it is negative if and only if

$$\frac{\alpha^2 + \alpha\beta + \beta^2}{\alpha + \beta} > \frac{f_1}{f_0} \Leftrightarrow \alpha^2 + \beta^2 + \alpha\beta - \frac{f_1}{f_0}\alpha - \frac{f_1}{f_0}\beta > 0.$$

The above inequality has infinitely many solutions. One of the examples can be the pair $\alpha = \frac{f_1}{2f_0}$, $\beta = \frac{f_1}{f_0}$. More precisely, the set of all solutions is the exterior of the ellipse

$$\frac{\left(x - \frac{\sqrt{2}f_1}{3f_0}\right)^2}{\frac{2f_1^2}{9f_0^2}} + \frac{y^2}{\frac{2f_1^2}{3f_0^2}} = 1$$

rotated by the angle $-\frac{\pi}{4}$. The example of this area for $\frac{f_1}{f_0} = 2$ is shown on Fig.1.

4.2 Quasi-Riordan arrays that are TP

Example 4.6. Let $f(t) = \frac{t}{1-2t}$, $g(t) = 1+t+t^2$. Then f is a Pólya frequency power series, g is not. Consider

$$R = [g, f] = \begin{bmatrix} 1 & & & & & \\ 1 & 1 & & & & \\ 1 & 2 & 1 & & & \\ 0 & 4 & 2 & 1 & & \\ 0 & 8 & 4 & 2 & 1 & \\ \vdots & & & & & \ddots \end{bmatrix}$$

and its minors.

Clearly, those minors that do not involve the 0th column are the minors of $[1, f(t)]$, so they are nonnegative. Moreover, only those that are either of order 1 or those that are triangular minors are positive, the other ones are equal to 0.

Consider now a minor of order n ($n \geq 2$) containing the 0th column. If it contains the 0th row, then expanding along the 0th row, we get that it is equal to some minor of $[1, f(t)]$, so it is nonnegative. Suppose that it does not contain the 0th row. Then let us expand it along the 0th column. If it contains one of the 1st row and the 2 row, expanding along the 0th column, we again get a minor of $[1, f(t)]$. Suppose that it contains both 1st and 2nd row. Again, expanding along the 0th column, we get then the difference of two minors of $[1, f(t)]$. If $n = 2$, then we deal with the minor

$$\begin{vmatrix} 1 & r_{k1} \\ 1 & r_{k2} \end{vmatrix} \in \left\{ \begin{vmatrix} 1 & 1 \\ 1 & 2 \end{vmatrix}, \begin{vmatrix} 1 & 0 \\ 1 & 1 \end{vmatrix}, \begin{vmatrix} 1 & 0 \\ 1 & 0 \end{vmatrix} \right\}$$

that is nonnegative.

If $n \geq 3$, both minors are of order greater or equal to 3, so they are equal to 0, and so is their difference. Hence, R is TP.

Example 4.6 can be extended to the following.

Corollary 4.7. Let $\alpha > 0$, and let $g_0, g_1, g_2 > 0$ be such that $g_1^2 - 4g_0g_2 < 0$. If $g_1\alpha - g_2 \geq 0$, then $R = \left[g_0 + g_1t + g_2t^2, \frac{t}{1-\alpha t} \right]$ is TP, although $g_0 + g_1t + g_2t^2$ is not a Pólya frequency power series.

Proof. Analogously as in the example, the only minors that are nonzero, are the triangular ones and the minor

$$\begin{vmatrix} r_{10} & r_{11} \\ r_{20} & r_{21} \end{vmatrix} = \begin{vmatrix} g_1 & 1 \\ g_2 & \alpha \end{vmatrix} = g_1\alpha - g_2$$

which, by assumption, is nonnegative. \square

Note that replacing $g(t) = 1 + t + t^2$ from Example 4.6 by, for instance, $g(t) = t^2 + 1$ (that does not satisfy the assumption $g_1 > 0$ of Corollary 4.7) yields the array of the form

$$\begin{bmatrix} 1 & & & & & \\ 0 & f_1 & & & & \\ 1 & f_2 & f_1 & & & \\ 0 & f_3 & f_2 & f_1 & & \\ \vdots & & & & \ddots & \end{bmatrix}$$

whose minor $M_{0,1}^{1,2}$ is equal to $-f_1 < 0$, so the above array is not TP.

5 Closing comments

In the present work we have studied total positivity of quasi-Riordan arrays. As we have mentioned, these are, in fact, some particular cases of almost-Riordan arrays. Thus, it is natural question whether the presented results can be generalized for almost-Riordan arrays. Considering the capacity of the paper, the TP of almost-Riordan arrays will be discussed in a subsequent paper. For the same reason, some other related topics, such as analogues of TP_r Riordan arrays in [7], TP_r quasi-Riordan arrays and TP_r almost-Riordan arrays, will also be discussed in a subsequent paper.

6 Data availability statement

All data generated or analyzed during this study are included in this article.

References

- [1] M. Aissen, A. Edrei, I.J. Schoenberg and A.M. Whitney, On the generating functions of totally positive sequences, *Proc. Nat. Acad. Sci. U.S.A.* 37 (1951), 303–307.
- [2] M. Aissen, I.J. Schoenberg and A.M. Whitney, On the generating functions of totally positive sequences I, *J. Analyse Math.* 2 (1952), 93–103.
- [3] Y. Alp and E. G. Kocer, Sequence characterization of almost-Riordan arrays, *Linear Algebra Appl.* 664 (2023), 1–23.
- [4] P. Barry, *Riordan Arrays: A Primer*, LOGIC Press, Kilcock, 2016.
- [5] P. Barry, On the group of almost-Riordan arrays, arXiv:1606.05077v1.

- [6] F. Brenti, Combinatorics and total positivity, *J. Combin. Theory Ser. A*, 71 (1995) 175–218.
- [7] X. Chen, H. Liang and Y. Wang, Total positivity of Riordan arrays, *European J. Combin.* 46 (2015) 68–74.
- [8] X. Chen and Y. Wang, Notes on the total positivity of Riordan arrays, *Linear Algebra Appl.* 569 (2019) 156–161.
- [9] T.-X. He, The vertical recursive relation of Riordan arrays and its matrix representation, *J. Integer Seq.* 25 (2022), no. 9, Art. 22.9.5, 22 pp.
- [10] T. -X. He, Matrix characterizations of Riordan matrices, *Linear Algebra Appl.* 465 (2015), 15-42.
- [11] T. -X. He and R. Sprugnoli. Sequence Characterization of Riordan Arrays, *Discrete Math.*, 309 (2009), 3962-3974.
- [12] C. Jean-Louis and A. Nkwanta, Some algebraic structure of the Riordan group, *Linear Algebra Appl.* 438 (2013), 2018–2035.
- [13] S. Karlin, *Total Positivity, Vol.1*, Stanford University Press, 1968.
- [14] A. Luzó n, D. Merlini, M. A. Moró n, L. F. Prieto-Martinez, and R. Sprugnoli, Some inverse limit approaches to the Riordan group, *Linear Algebra Appl.* 491 (2016), 239–262.
- [15] A. Luzó n, M. A. Moró n, and L. F. Prieto-Martinez, A formula to construct all involutions in Riordan matrix groups. *Linear Algebra Appl.* 533 (2017), 397–417.
- [16] A. Luzó n, M. A. Moró n, and L. F. Prieto-Martinez, The group generated by Riordan involutions, *Revista Matemática Complutense* <https://doi.org/10.1007/s13163-020-00382-8>, 2020.
- [17] J. Mao, L. Mu, and Y. Wang, Yet another criterion for the total positivity of Riordan arrays, *Linear Algebra Appl.* **634** (2022), 106–111.
- [18] D. Merlini, D. G. Rogers, R. Sprugnoli, and M. C. Verri, On some alternative characterizations of Riordan matrices, *Canadian J. Math.*, 49 (1997), 301–320.
- [19] A. Pinkus, *Totally Positive Matrices*, Cambridge University Press, Cambridge, 2010.
- [20] D. G. Rogers, Pascal triangles, Catalan numbers and renewal matrices, *Discrete Math.*, 22 (1978), 301–310.

- [21] L. W. Shapiro, Bijections and the Riordan group, Random generation of combinatorial objects and bijective combinatorics, *Theoret. Comput. Sci.* 307 (2003), no. 2, 403–413.
- [22] L. W. Shapiro, S. Getu, W. J. Woan and L. Woodson, The Riordan group, *Discrete Appl. Math.* 34(1991) 229–239.
- [23] L. W. Shapiro, R. Sprugnoli, P. Barry, G.-S. Cheon, T.-X. He, D. Merlini, and W. Wang, *The Riordan Group and Applications*, Springer, 2022.
- [24] R. Slowik, Some (counter)examples on totally positive Riordan arrays, *Linear Algebra Appl.* 594 (2020), 117–123.

Figure 1: The area depicting pairs (α, β) for which $\left[\frac{1}{(1-\alpha t)(1-\beta t)}, f(t)\right]$ is not TP, $\frac{f_1}{f_0} = 2$. Note that the fact that point (α, β) lies inside the ellipse does not necessarily mean that $[g, f]$ is TP.

