
CONSTRUCTIONS OF A_α -COSPECTRAL GRAPHS USING SOME CORONA OPERATIONS

Najiya V K, Chithra A V
 Department of Mathematics
 National Institute of Technology, Calicut
 Kerala, India-673601
 najiya_p190046ma@nitc.ac.in, chithra@nitc.ac.in

ABSTRACT

Let $G_1 \otimes G_2, G_1 \sqcup G_2$ and $G_1 \sqcup G_2$ denote the total corona, Q -vertex corona and Q -edge corona of two graphs G_1 and G_2 , respectively. In this paper, we compute the A_α -spectrum of $G_1 \otimes G_2, G_1 \sqcup G_2$ and $G_1 \sqcup G_2$ for regular graphs G_1 and G_2 . As an application, we construct infinitely many pairs of A_α -cospectral graphs.

1 Introduction

Throughout the manuscript, we focus only on simple undirected graphs. Let G denote a graph with vertex set $V(G) = \{u_1, u_2, \dots, u_n\}$ and edge set $E(G) = \{e_1, e_2, \dots, e_m\}$. The adjacency matrix $A(G)$ of G is a square symmetric matrix of order n whose $(i, j)^{th}$ entry is 1 if the vertices u_i and u_j are adjacent and is 0 otherwise. The degree matrix $D(G)$ is a diagonal matrix with diagonal entries corresponding to vertex degrees. The A_α -matrix [1] $A_\alpha(G)$ of G is defined as $A_\alpha(G) = \alpha D(G) + (1 - \alpha)A(G)$ for any real $\alpha \in [0, 1]$. Specifically, for $\alpha = 0, \frac{1}{2}$ and 1, $A_\alpha(G)$ becomes $A(G), \frac{1}{2}Q(G)$ and $D(G)$, respectively.

The characteristic polynomial of the matrix M of order n is defined as $\phi(M, \lambda) = |\lambda I_n - M|$, where I_n is the identity matrix of order n . The roots for this polynomial are called the M -eigenvalues. Let $\lambda_1(A_\alpha(G)) \geq \lambda_2(A_\alpha(G)) \geq \dots \geq \lambda_n(A_\alpha(G))$ represent the A_α -eigenvalues of G . The set of all $A_\alpha(G)$ eigenvalues along with their multiplicities is called the A_α -spectrum of G . For an r -regular graph G , $\lambda_i(A_\alpha(G)) = \alpha r + (1 - \alpha)\lambda_i(A(G))$, $i = 1, 2, \dots, n$. Two graphs are said to be A_α -cospectral if they have the same A_α -spectrum. Otherwise, they are non A_α -cospectral graphs. A_α -cospectral graphs are always A -cospectral graphs, but A -cospectral graphs need not always be A_α -cospectral. However, regular A -cospectral graphs are A_α -cospectral.

The adjacency energy [2] $\varepsilon(G)$ of G is defined as the sum of the absolute values of eigenvalues of G . The A_α -energy [3] $\varepsilon_\alpha(G)$ of G is defined as $\sum_{i=1}^n \left| \lambda_i(A_\alpha(G)) - \frac{2\alpha m}{n} \right|$. For a regular graph G , $\varepsilon_\alpha(G) = (1 - \alpha)\varepsilon(G)$.

The incidence matrix $R(G)$ of a graph G is the matrix of order $n \times m$, whose $(i, j)^{th}$ entry is 1 if u_i is incident to e_j and 0 otherwise. From [4], $R(G)R(G)^T = A(G) + D(G)$ and $R(G)^T R(G) = B(G) + 2I_m$, where B is the adjacency matrix of line graph of G . For an r -regular graph G , $R(G)R(G)^T = A(G) + rI_n$. Throughout this paper, I_n is the identity matrix of order n , $O_{m \times n}$ is the 0 matrix of order $m \times n$, J_n is the matrix of order n with all entries equal to one and $J_{s \times t}$ is the matrix of order $s \times t$ with all entries equal to one. Let A and B be matrices of order $a \times b$ and $c \times d$, respectively, then the Kronecker product of A and B , denoted $A \otimes B$, is the $ac \times bd$ block matrix $[a_{ij}B]$.

The corona [5] of two graphs G_1, G_2 , denoted by $G_1 \circ G_2$, is constructed by taking one copy of G_1 and n_1 copies of G_2 and joining the i th vertex of G_1 to every vertex in the i th copy of G_2 for each i . The Q graph of G , $Q(G)$, is the graph obtained from G by inserting a new vertex into every edge of G and then joining pairs of these new vertices that lie on adjacent edges of G . The total graph $T(G)$ of a graph G is the graph whose set of vertices is the union of the set of vertices and of the set of edges of G , with two vertices of $T(G)$ being adjacent if and only if the corresponding

elements of G are adjacent or incident. Let G be a graph with vertex set $V(G) = \{v_1, v_2, \dots, v_n\}$. Consider a set $U(G) = \{u_1, u_2, \dots, u_n\}$ of n vertices. Make u_i adjacent to all the vertices in the neighbourhood of v_i , for each $i = 1, 2, \dots, n$. The graph thus obtained is called the splitting graph of G , denoted by $Spl(G)$. Let G_1 and G_2 be two graphs with n_1 and n_2 vertices, respectively. The neighbourhood corona, $G_1 \star G_2$, of G_1 and G_2 is obtained by taking n_1 copies of G_2 and joining all the vertices of the i th copy of G_2 with the vertices adjacent to v_i in G_1 , for $i = 1, 2, \dots, n_1$. In this paper, we find the A_α -spectrum of a few corona products of graphs.

The rest of the paper is organised as follows. In Section 2, we present some definitions and lemmas used to prove our results. In Sections 3, 4, and 5, we compute the A_α -spectrum of the total corona, Q -vertex corona, and Q -edge corona of two graphs. Also, we construct an infinite number of pairs of A_α -cospectral graphs that are non-regular.

2 Preliminaries

Definition 2.1. [6] The total corona of G_1 and G_2 , denoted by $G_1 \otimes G_2$, is obtained by taking one copy of $T(G_1)$ and n_1 copies of G_2 , and joining the i th vertex of G_1 to every vertex in the i th copy of G_2 .

Definition 2.2. [7] The splitting corona, $G_1 \boxplus G_2$ of G_1 and G_2 is obtained from $Spl(G_1)$ and n_1 copies of G_2 by joining every vertex of the i th copy of G_2 to the i th vertex of $V(G_1)$, for $i = 1, 2, \dots, n_1$.

Definition 2.3. [7] The splitting add vertex corona, $G_1 \boxtimes G_2$ of G_1 and G_2 is obtained from $Spl(G_1)$ and n_1 copies of G_2 by joining every vertex of the i th copy of G_2 to the i th vertex of $U(G_1)$, for $i = 1, 2, \dots, n_1$.

Definition 2.4. [7] The splitting neighbourhood corona, $G_1 \boxast G_2$ of G_1 and G_2 is obtained from $Spl(G_1)$ and n_1 copies of G_2 by joining every vertex of the i th copy of G_2 to the vertices adjacent to the i th vertex in $V(G_1)$, for $i = 1, 2, \dots, n_1$.

Now, we define new graph operations based on Q -graphs as follows.

Definition 2.5. The Q -vertex corona of G_1 and G_2 , denoted by $G_1 \sqcup G_2$, is the graph obtained from $Q(G_1)$ and n_1 copies of G_2 by joining the i th vertex of $V(G_1)$ to every vertex in the i th copy of G_2 .

Definition 2.6. The Q -edge corona of G_1 and G_2 , denoted by $G_1 \sqcup G_2$, is the graph obtained from $Q(G_1)$ and m_1 copies of G_2 by joining the i th vertex of $E(G_1)$ to every vertex in the i th copy of G_2 .




Figure 1: $C_4 \sqcup K_2$




Figure 2: $C_4 \square K_2$

Lemma 2.1. [4] Let P, Q, R and S be matrices and

$$M = \begin{pmatrix} P & Q \\ R & S \end{pmatrix}.$$

If P is invertible, then $\det(M) = \det(P) \det(S - RP^{-1}Q)$.

If S is invertible, then $\det(M) = \det(S) \det(P - QS^{-1}R)$.

If P and R commute, then $\det(M) = \det(PS - QR)$.

Definition 2.7. [8] The M -coronal $\Gamma_M(\lambda)$ of a matrix M of order n is defined as the sum of the entries of the matrix $(\lambda I_n - M)^{-1}$ (if it exists), that is,

$$\Gamma_M(\lambda) = J_{n \times 1}^T (\lambda I_n - M)^{-1} J_{n \times 1}.$$

If each row sum of an $n \times n$ matrix M is constant, say a , then $\Gamma_M(\lambda) = \frac{n}{\lambda - a}$.

Lemma 2.2. [9] Let G be an r -regular graph on n vertices, then $\Gamma_{A(G)}(\lambda) = \frac{n}{\lambda - r}$.

Theorem 2.3. [10] If G is an r -regular graph on n vertices, then the adjacency spectrum of line graph of G is

$$\begin{pmatrix} -2 & \lambda_i + r - 2 \\ m - n & 1 \end{pmatrix},$$

where λ_i 's are the adjacency eigenvalues of G .

3 A_α -spectrum of total corona

In this section, we compute of A_α -spectrum of $G_1 \otimes G_2$, the total corona of the graphs G_1 and G_2 . First, we obtain an expression for the A_α -characteristic polynomial of $G_1 \otimes G_2$, where G_1 is an r_1 -regular graph and G_2 is an arbitrary graph.

Proposition 3.1. Let G_1 be an r_1 -regular graph on n_1 vertices and G_2 be an arbitrary graph on n_2 vertices. Then

$$\begin{aligned} \phi(A_\alpha(G_1 \otimes G_2), \lambda) &= (\lambda - 2(\alpha r_1 - 1 + \alpha))^{m_1 - n_1} \prod_{i=1}^{n_2} (\lambda - \alpha - \lambda_i(A_{\alpha_2}))^{n_1} \\ &\prod_{i=1}^{n_1} \left((\lambda - 2(\alpha r_1 - 1 + \alpha))(\lambda - (3r_1 + n_2)\alpha - (1 - \alpha)^2 \Gamma_{A_{\alpha_2}}(\lambda - \alpha) + r_1) \right. \\ &\quad \left. - (1 - \alpha)(\lambda_i(A_1) + r_1)(2\lambda - (5r_1 + n_2)\alpha + 3(1 - \alpha) + r_1 - (1 - \alpha)^2 \Gamma_{A_{\alpha_2}}(\lambda - \alpha)) + (1 - \alpha)^2 (\lambda_i(A_1) + r_1)^2 \right). \end{aligned}$$

Proof. With a suitable ordering of the vertices of $G_1 \otimes G_2$, we get

$$\begin{aligned}
 A_\alpha(G_1 \otimes G_2) &= \begin{pmatrix} (r_1 + n_2)\alpha I_{n_1} + A_{\alpha_1} & (1 - \alpha)R_1 & (1 - \alpha)I_{n_1} \otimes J_{1 \times n_2} \\ (1 - \alpha)R_1^T & 2\alpha r_1 I_{m_1} + (1 - \alpha)B_1 & O \\ (1 - \alpha)I_{n_1} \otimes J_{n_2 \times 1} & O & I_{n_1} \otimes (\alpha I_{n_2} + A_{\alpha_2}) \end{pmatrix}. \\
 |\lambda I - A_\alpha(G_1 \otimes G_2)| &= \begin{vmatrix} (\lambda - (r_1 + n_2)\alpha)I_{n_1} - A_{\alpha_1} & -(1 - \alpha)R_1 & -(1 - \alpha)I_{n_1} \otimes J_{1 \times n_2} \\ -(1 - \alpha)R_1^T & (\lambda - 2\alpha r_1)I_{m_1} - (1 - \alpha)B_1 & O \\ -(1 - \alpha)I_{n_1} \otimes J_{n_2 \times 1} & O & I_{n_1} \otimes ((\lambda - \alpha)I_{n_2} - A_{\alpha_2}) \end{vmatrix} \\
 &= \prod_{i=1}^{n_2} (\lambda - \alpha - \lambda_i(A_{\alpha_2}))^{n_1} \det S
 \end{aligned}$$

where

$$\det S = \begin{vmatrix} (\lambda - (r_1 + n_2)\alpha - (1 - \alpha)^2 \Gamma_{A_{\alpha_2}^T}(\lambda - \alpha))I_{n_1} - A_{\alpha_1} & -(1 - \alpha)R_1 \\ -(1 - \alpha)R_1^T & (\lambda - 2\alpha r_1)I_{m_1} - (1 - \alpha)B_1 \end{vmatrix}.$$

Now performing row transformations on rows \mathcal{R}_1 and \mathcal{R}_2 by $\mathcal{R}_2 \rightarrow R_1^T \mathcal{R}_1 + \mathcal{R}_2$ followed by $\mathcal{R}_1 \rightarrow \mathcal{R}_1 + \frac{(1 - \alpha)R_1 \mathcal{R}_2}{\lambda - 2\alpha r_1 + 2(1 - \alpha)}$ we get

$$\begin{aligned}
 \det S &= (\lambda - 2(\alpha r_1 - 1 + \alpha))^{m_1 - n_1} \prod_{i=1}^{n_1} \left((\lambda - 2(\alpha r_1 - 1 + \alpha))(\lambda - (3r_1 + n_2)\alpha - (1 - \alpha)^2 \Gamma_{A_{\alpha_2}}(\lambda - \alpha) + r_1) \right. \\
 &\quad \left. - (1 - \alpha)(\lambda_i(A_1) + r_1)(2\lambda - (5r_1 + n_2)\alpha + 3(1 - \alpha) + r_1 - (1 - \alpha)^2 \Gamma_{A_{\alpha_2}}(\lambda - \alpha)) + (1 - \alpha)^2 (\lambda_i(A_1) + r_1)^2 \right).
 \end{aligned}$$

Then the result follows. \square

Now, in the following corollary, we obtain the A_α -eigenvalues of $G_1 \otimes G_2$, where both G_1 and G_2 are regular graphs.

Corollary 3.1. Let G_i be r_i -regular graph with n_i vertices for $i = 1, 2$. Then the A_α -spectrum of $G_1 \otimes G_2$ consists of:

1. $2(\alpha r_1 - 1 + \alpha)$ repeated $m_1 - n_1$ times,
2. $\alpha + \lambda_i(A_{\alpha_2})$ repeated n_1 times for each $i = 2, 3, \dots, n_2$,
3. three roots of the equation $(\lambda - 2(\alpha r_1 - 1 + \alpha))((\lambda - \alpha - r_2)(\lambda - (3r_1 + n_2)\alpha + r_1) - (1 - \alpha)^2 n_2) - (1 - \alpha)(\lambda_i(A_1) + r_1)((\lambda - \alpha - r_2)(2\lambda - (5r_1 + n_2)\alpha + 3(1 - \alpha) + r_1) - (1 - \alpha)^2 n_2) + (1 - \alpha)^2 (\lambda - \alpha - r_2)(\lambda_i(A_1) + r_1)^2 = 0$, for $i = 1, 2, \dots, n_1$.

Finally, as an application, we construct new pairs of A_α -cospectral graphs in the following corollary.

Corollary 3.2. 1. Let G_1 and G_2 be two A -cospectral regular graphs and H be an arbitrary graph. Then the graphs $G_1 \otimes H$ and $G_2 \otimes H$ are A_α -cospectral.

2. Let H_1 and H_2 be two A_α -cospectral graphs with $\Gamma_{A_\alpha(H_1)}(\lambda) = \Gamma_{A_\alpha(H_2)}(\lambda)$ for $\alpha \in [0, 1]$. If G is a regular graph, then the graphs $G \otimes H_1$ and $G \otimes H_2$ are A_α -cospectral.

4 A_α -spectrum of corona based on splitting graph

In this section, we compute of A_α -spectrum of $G_1 \boxplus G_2, G_1 \boxtimes G_2$ and $G_1 \boxast G_2$ of the graphs G_1 and G_2 . First, we obtain an expression for the A_α -characteristic polynomial of these operations, where G_1 is an r_1 -regular graph and G_2 is an arbitrary graph.

Proposition 4.1. Let G_1 be an r_1 -regular graph on n_1 vertices and G_2 be an arbitrary graph on n_2 vertices. Then

$$\begin{aligned}
 \phi(A_\alpha(G_1 \boxplus G_2), \lambda) &= \prod_{i=1}^{n_2} (\lambda - \alpha - \lambda_i(A_{\alpha_2}))^{n_1} \\
 &\quad \prod_{i=1}^{n_1} (\lambda - \alpha r_1)(\lambda - (2r_1 + n_2)\alpha - (1 - \alpha)^2 \Gamma_{A_{\alpha_2}}(\lambda - \alpha) - (1 - \alpha)\lambda_i(A_1)) - (1 - \alpha)^2 \lambda_i(A_1).
 \end{aligned}$$

Proof. With a suitable ordering of the vertices of $G_1 \boxplus G_2$, we get

$$A_\alpha(G_1 \boxplus G_2) = \begin{pmatrix} (2r_1 + n_2)\alpha I_{n_1} + (1 - \alpha)A_1 & (1 - \alpha)A_1 & (1 - \alpha)I_{n_1} \otimes J_{1 \times n_2} \\ (1 - \alpha)A_1 & \alpha r_1 I_{n_1} & O \\ (1 - \alpha)I_{n_1} \otimes J_{n_2 \times 1} & O & I_{n_1} \otimes (\alpha I_{n_2} + A_{\alpha_2}) \end{pmatrix}.$$

$$|\lambda I - A_\alpha(G_1 \boxplus G_2)| = \begin{vmatrix} (\lambda - (2r_1 + n_2)\alpha)I_{n_1} - (1 - \alpha)A_1 & -(1 - \alpha)A_1 & -(1 - \alpha)I_{n_1} \otimes J_{1 \times n_2} \\ -(1 - \alpha)A_1 & (\lambda - \alpha r_1)I_{n_1} & O \\ -(1 - \alpha)I_{n_1} \otimes J_{n_2 \times 1} & O & I_{n_1} \otimes ((\lambda - \alpha)I_{n_2} - A_{\alpha_2}) \end{vmatrix}$$

$$= \prod_{i=1}^{n_2} (\lambda - \alpha - \lambda_i(A_{\alpha_2}))^{n_1} \det S$$

where

$$\det S = \begin{vmatrix} (\lambda - (2r_1 + n_2)\alpha)I_{n_1} - (1 - \alpha)A_1 - (1 - \alpha)^2 \Gamma_{A_{\alpha_2}}(\lambda - \alpha)I_{n_1} & -(1 - \alpha)A_1 \\ -(1 - \alpha)A_1 & (\lambda - \alpha r_1)I_{n_1} \end{vmatrix}.$$

Then by Lemma 2.1

$$= \prod_{i=1}^{n_1} (\lambda - \alpha r_1)(\lambda - (2r_1 + n_2)\alpha - (1 - \alpha)^2 \Gamma_{A_{\alpha_2}}(\lambda - \alpha) - (1 - \alpha)\lambda_i(A_1)) - (1 - \alpha)^2 \lambda_i(A_1).$$

Then the result follows. \square

Now, in the following corollary, we obtain the A_α -eigenvalues of $G_1 \boxplus G_2$, where both G_1 and G_2 are regular graphs.

Corollary 4.1. Let G_i be r_i -regular graph with n_i vertices for $i = 1, 2$. Then the A_α -spectrum of $G_1 \boxplus G_2$ consists of:

1. $\alpha + \lambda_i(A_{\alpha_2})$ repeated n_1 times for each $i = 2, 3, \dots, n_2$,
2. two roots of the equation $(\lambda - \alpha r_1)((\lambda - \alpha - r_2)(\lambda - (2r_1 + n_2)\alpha - (1 - \alpha)\lambda_i(A_1)) - (1 - \alpha)^2 n_2) - (1 - \alpha)^2 (\lambda - \alpha - r_2)\lambda_i(A_1) = 0$, for $i = 1, 2, \dots, n_1$.

Finally, as an application, we construct new pairs of A_α -cospectral graphs in the following corollary.

Corollary 4.2. 1. Let G_1 and G_2 be two A_α -cospectral regular graphs and H be an arbitrary graph. Then the graphs $G_1 \boxplus H$ and $G_2 \boxplus H$ are A_α -cospectral.

2. Let H_1 and H_2 be two A_α -cospectral graphs with $\Gamma_{A_\alpha(H_1)}(\lambda) = \Gamma_{A_\alpha(H_2)}(\lambda)$ for $\alpha \in [0, 1]$. If G is a regular graph, then the graphs $G \boxplus H_1$ and $G \boxplus H_2$ are A_α -cospectral.

Next, we compute A_α -spectrum of splitting add vertex corona of G_1 and G_2 , where G_1 is an r_1 -regular graph and G_2 is an arbitrary graph.

Proposition 4.2. Let G_1 be an r_1 -regular graph on n_1 vertices and G_2 be an arbitrary graph on n_2 vertices. Then

$$\phi(A_\alpha(G_1 \boxtimes G_2), \lambda) = \prod_{i=1}^{n_2} (\lambda - \alpha - \lambda_i(A_{\alpha_2}))^{n_1}$$

$$\prod_{i=1}^{n_1} (\lambda - 2\alpha r_1 - (1 - \alpha)\lambda_i(A_1))(\lambda - (r_1 + n_2)\alpha - (1 - \alpha)^2 \Gamma_{A_{\alpha_2}}(\lambda - \alpha)) - (1 - \alpha)^2 \lambda_i(A_1)^2.$$

Proof. With a suitable ordering of the vertices of $G_1 \boxtimes G_2$, we get

$$A_\alpha(G_1 \boxtimes G_2) = \begin{pmatrix} 2\alpha r_1 I_{n_1} + (1 - \alpha)A_1 & (1 - \alpha)A_1 & O \\ (1 - \alpha)A_1 & \alpha(r_1 + n_2)I_{n_1} & (1 - \alpha)I_{n_1} \otimes J_{1 \times n_2} \\ O & (1 - \alpha)I_{n_1} \otimes J_{n_2 \times 1} & I_{n_1} \otimes (\alpha I_{n_2} + A_{\alpha_2}) \end{pmatrix}.$$

The result follows by similar arguments as in Proposition 4.1 \square

Now, in the following corollary, we obtain the A_α -eigenvalues of $G_1 \boxtimes G_2$, where both G_1 and G_2 are regular graphs.

Corollary 4.3. Let G_i be r_i -regular graph with n_i vertices for $i = 1, 2$. Then the A_α -spectrum of $G_1 \boxtimes G_2$ consists of:

1. $\alpha + \lambda_i(A_{\alpha_2})$ repeated n_1 times for each $i = 2, 3, \dots, n_2$,
2. two roots of the equation $(\lambda - 2\alpha r_1 - (1 - \alpha)\lambda_i(A_1))((\lambda - \alpha - r_2)(\lambda - (r_1 + n_2)\alpha) - (1 - \alpha)^2 n_2) - (1 - \alpha)^2 (\lambda - \alpha - r_2)\lambda_i(A_1)^2 = 0$, for $i = 1, 2, \dots, n_1$.

Finally, as an application, we construct new pairs of A_α -cospectral graphs in the following corollary.

Corollary 4.4. 1. Let G_1 and G_2 be two A -cospectral regular graphs and H be an arbitrary graph. Then the graphs $G_1 \boxtimes H$ and $G_2 \boxtimes H$ are A_α -cospectral.

2. Let H_1 and H_2 be two A_α -cospectral graphs with $\Gamma_{A_\alpha(H_1)}(\lambda) = \Gamma_{A_\alpha(H_2)}(\lambda)$ for $\alpha \in [0, 1]$. If G is a regular graph, then the graphs $G \boxtimes H_1$ and $G \boxtimes H_2$ are A_α -cospectral.

Next, we compute A_α -spectrum of splitting neighbourhood corona of G_1 and G_2 , where G_1 is an r_1 -regular graph and G_2 is an arbitrary graph.

Proposition 4.3. Let G_1 be an r_1 -regular graph on n_1 vertices and G_2 be an arbitrary graph on n_2 vertices. Then

$$\phi(A_\alpha(G_1 * G_2), \lambda) = \prod_{i=1}^{n_2} (\lambda - \alpha r_1 - \lambda_i(A_{\alpha_2}))^{n_1} \prod_{i=1}^{n_1} (\lambda - \alpha r_1)(\lambda - (2 + n_2)r_1\alpha - (1 - \alpha)^2 \Gamma_{A_{\alpha_2}}(\lambda - \alpha r_1)\lambda_i(A_1)^2 - (1 - \alpha)\lambda_i(A_1)) - (1 - \alpha)^2 \lambda_i(A_1)^2.$$

Proof. With a suitable ordering of the vertices of $G_1 * G_2$, we get

$$A_\alpha(G_1 * G_2) = \begin{pmatrix} (n_2 + 2)r_1\alpha I_{n_1} + (1 - \alpha)A_1 & (1 - \alpha)A_1 & (1 - \alpha)J_{1 \times n_2} \otimes A_1 \\ (1 - \alpha)A_1 & \alpha r_1 I_{n_1} & O \\ (1 - \alpha)J_{n_2 \times 1} \otimes A_1 & O & (\alpha r_1 I_{n_2} + A_{\alpha_2}) \otimes I_{n_1} \end{pmatrix}.$$

The result follows by similar arguments as in Proposition 4.1 □

Now, in the following corollary, we obtain the A_α -eigenvalues of $G_1 * G_2$, where both G_1 and G_2 are regular graphs.

Corollary 4.5. Let G_i be r_i -regular graph with n_i vertices for $i = 1, 2$. Then the A_α -spectrum of $G_1 * G_2$ consists of:

1. $\alpha r_1 + \lambda_i(A_{\alpha_2})$ repeated n_1 times for each $i = 2, 3, \dots, n_2$,
2. two roots of the equation $(\lambda - \alpha r_1)((\lambda - \alpha r_1 - r_2)(\lambda - (2 + n_2)r_1\alpha - (1 - \alpha)\lambda_i(A_1)) - (1 - \alpha)^2 n_2 \lambda_i(A_1)^2) - (1 - \alpha)^2 (\lambda - \alpha r_1 - r_2)\lambda_i(A_1)^2 = 0$, for $i = 1, 2, \dots, n_1$.

Finally, as an application, we construct new pairs of A_α -cospectral graphs in the following corollary.

Corollary 4.6. 1. Let G_1 and G_2 be two A -cospectral regular graphs and H be an arbitrary graph. Then the graphs $G_1 * H$ and $G_2 * H$ are A_α -cospectral.

2. Let H_1 and H_2 be two A_α -cospectral graphs with $\Gamma_{A_\alpha(H_1)}(\lambda) = \Gamma_{A_\alpha(H_2)}(\lambda)$ for $\alpha \in [0, 1]$. If G is a regular graph, then the graphs $G * H_1$ and $G * H_2$ are A_α -cospectral.

5 A_α -spectrum of corona based on Q graph

In this section, we compute of A_α -spectrum of $G_1 \sqcup G_2$ and $G_1 \sqcup G_2$ of the graphs G_1 and G_2 . First, we obtain an expression for the A_α -characteristic polynomial of these operations, where G_1 is an r_1 -regular graph and G_2 is an arbitrary graph.

Proposition 5.1. Let G_1 be an r_1 -regular graph on n_1 vertices and G_2 be an arbitrary graph on n_2 vertices. Then

$$\phi(A_\alpha(G_1 \sqcup G_2), \lambda) = (\lambda - 2\alpha r_1 + 2(1 - \alpha))^{m_1 - n_1} \prod_{i=1}^{n_2} (\lambda - \alpha - \lambda_i(A_{\alpha_2}))^{n_1}$$

$$\prod_{i=1}^{n_1} \left((\lambda - \alpha(r_1 + n_2) - (1 - \alpha)^2 \Gamma_{A_{\alpha_2}}(\lambda - \alpha)) (\lambda - 2\alpha r_1 - (1 - \alpha)(\lambda_i(A_1) + r_1 - 2)) - (1 - \alpha)^2 (\lambda_i(A_1) + r_1) \right).$$

Proof. With a suitable ordering of the vertices of $G_1 \sqcup G_2$, we get

$$A_\alpha(G_1 \sqcup G_2) = \begin{pmatrix} \alpha(r_1 + n_2)I_{n_1} & (1 - \alpha)R_1 & (1 - \alpha)I_{n_1} \otimes J_{1 \times n_2} \\ (1 - \alpha)R_1^T & 2\alpha r_1 I_{m_1} + (1 - \alpha)B_1 & O \\ (1 - \alpha)I_{n_1} \otimes J_{n_2 \times 1} & O & I_{n_1} \otimes (\alpha I_{n_2} + A_{\alpha_2}) \end{pmatrix}.$$

$$|\lambda I - A_\alpha(G_1 \sqcup G_2)| = \begin{vmatrix} (\lambda - \alpha(r_1 + n_2))I_{n_1} & -(1 - \alpha)R_1 & -(1 - \alpha)I_{n_1} \otimes J_{1 \times n_2} \\ -(1 - \alpha)R_1^T & (\lambda - 2\alpha r_1)I_{m_1} - (1 - \alpha)B_1 & O \\ -(1 - \alpha)I_{n_1} \otimes J_{n_2 \times 1} & O & I_{n_1} \otimes ((\lambda - \alpha)I_{n_2} - A_{\alpha_2}) \end{vmatrix}$$

$$= \prod_{i=1}^{n_2} (\lambda - \alpha - \lambda_i(A_{\alpha_2}))^{n_1} \det S$$

where

$$\det S = \begin{vmatrix} (\lambda - \alpha(r_1 + n_2) - (1 - \alpha)^2 \Gamma_{A_{\alpha_2}}(\lambda - \alpha))I & -(1 - \alpha)R_1 \\ -(1 - \alpha)R_1^T & (\lambda - 2\alpha r_1)I_{m_1} - (1 - \alpha)B_1 \end{vmatrix}$$

$$= (\lambda - \alpha(r_1 + n_2) - (1 - \alpha)^2 \Gamma_{A_{\alpha_2}}(\lambda - \alpha))^{n_1}$$

$$\begin{vmatrix} (\lambda - 2\alpha r_1)I_{m_1} - (1 - \alpha)B_1 - \frac{(1 - \alpha)^2 (B_1 + 2I_{m_1})}{\lambda - \alpha(r_1 + n_2) - (1 - \alpha)^2 \Gamma_{A_{\alpha_2}}(\lambda - \alpha)} \end{vmatrix}$$

$$\det S = (\lambda - 2\alpha r_1 + 2(1 - \alpha))^{m_1 - n_1}$$

$$\prod_{i=1}^{n_1} \left((\lambda - \alpha(r_1 + n_2) - (1 - \alpha)^2 \Gamma_{A_{\alpha_2}}(\lambda - \alpha)) (\lambda - 2\alpha r_1 - (1 - \alpha)(\lambda_i(A_1) + r_1 - 2)) - (1 - \alpha)^2 (\lambda_i(A_1) + r_1) \right).$$

Then the result follows. □

Now, in the following corollary, we obtain the A_α -eigenvalues of $G_1 \sqcup G_2$, where both G_1 and G_2 are regular graphs.

Corollary 5.1. Let G_i be r_i -regular graph with n_i vertices for $i = 1, 2$. Then the A_α -spectrum of $G_1 \sqcup G_2$ consists of:

1. $2\alpha r_1 - 2(1 - \alpha)$ repeated $m_1 - n_1$ times,
2. $\alpha + \lambda_i(A_{\alpha_2})$ repeated n_1 times for each $i = 2, 3, \dots, n_2$,
3. three roots of the equation

$$((\lambda - \alpha - r_2)(\lambda - \alpha(r_1 + n_2) - (1 - \alpha)^2 n_2) - (1 - \alpha)^2 (\lambda - 2\alpha r_1 - (1 - \alpha)(\lambda_i(A_1) + r_1 - 2))) - (1 - \alpha)^2 (\lambda - \alpha - r_2)(\lambda_i(A_1) + r_1) = 0,$$

for $i = 1, 2, \dots, n_1$.

Finally, as an application, we construct new pairs of A_α -cospectral graphs in the following corollary.

Corollary 5.2. 1. Let G_1 and G_2 be two A -cospectral regular graphs and H be an arbitrary graph. Then the graphs $G_1 \sqcup H$ and $G_2 \sqcup H$ are A_α -cospectral.

2. Let H_1 and H_2 be two A_α -cospectral graphs with $\Gamma_{A_\alpha(H_1)}(\lambda) = \Gamma_{A_\alpha(H_2)}(\lambda)$ for $\alpha \in [0, 1]$. If G is a regular graph, then the graphs $G \sqcup H_1$ and $G \sqcup H_2$ are A_α -cospectral.

Next, we compute A_α -spectrum of $G_1 \boxplus G_2$, where G_1 is an r_1 -regular graph and G_2 is an arbitrary graph.

Proposition 5.2. Let G_1 be an r_1 -regular graph on n_1 vertices and G_2 be an arbitrary graph on n_2 vertices. Then

$$\begin{aligned} \phi(A_\alpha(G_1 \boxplus G_2), \lambda) &= (\lambda - \alpha r_1)^{n_1 - m_1} \prod_{i=1}^{n_2} (\lambda - \alpha - \lambda_i(A_{\alpha_2}))^{m_1} \\ &\quad ((\lambda - \alpha r_1)(\lambda - \alpha(2r_1 + n_2) - (1 - \alpha)^2 \Gamma_{A_{\alpha_2}}(\lambda - \alpha)) - 2(1 - \alpha)^2 + 2(1 - \alpha)(\lambda - \alpha r_1 + 1 - \alpha))^{m_1 - n_1} \\ &\quad \prod_{i=1}^{n_1} ((\lambda - \alpha r_1)(\lambda - \alpha(2r_1 + n_2) - (1 - \alpha)^2 \Gamma_{A_{\alpha_2}}(\lambda - \alpha)) - 2(1 - \alpha)^2 - (1 - \alpha)(\lambda - \alpha r_1 + 1 - \alpha)(\lambda_i(A_1) + r_1 - 2)). \end{aligned}$$

Proof. With a suitable ordering of the vertices of $G_1 \boxplus G_2$, we get

$$A_\alpha(G_1 \boxplus G_2) = \begin{pmatrix} \alpha r_1 I_{n_1} & (1 - \alpha) R_1 & O \\ (1 - \alpha) R_1^T & \alpha(2r_1 + n_2) I_{m_1} + (1 - \alpha) B_1 & (1 - \alpha) I_{m_1} \otimes J_{1 \times n_2} \\ O & (1 - \alpha) I_{m_1} \otimes J_{n_2 \times 1} & I_{m_1} \otimes (\alpha I_{n_2} + A_{\alpha_2}) \end{pmatrix}.$$

The proof is similar to the proof of Proposition 5.1 □

Now, in the following corollary, we obtain the A_α -eigenvalues of $G_1 \boxplus G_2$, where both G_1 and G_2 are regular graphs.

Corollary 5.3. Let G_i be r_i -regular graph with n_i vertices for $i = 1, 2$. Then the A_α -spectrum of $G_1 \boxplus G_2$ consists of:

1. αr_1 repeated $n_1 - m_1$ times,
2. $\alpha + \lambda_i(A_{\alpha_2})$ repeated m_1 times for each $i = 2, 3, \dots, n_2$,
3. three roots of the equation

$$(\lambda - \alpha r_1)((\lambda - \alpha - r_2)(\lambda - \alpha(2r_1 + n_2) - (1 - \alpha)^2 n_2) - (\lambda - \alpha - r_2)(2(1 - \alpha)^2 - 2(1 - \alpha)(\lambda - \alpha r_1 + 1 - \alpha))) = 0,$$

repeated $m_1 - n_1$,

4. three roots of the equation

$$(\lambda - \alpha r_1)((\lambda - \alpha - r_2)(\lambda - \alpha(2r_1 + n_2) - (1 - \alpha)^2 n_2) - (\lambda - \alpha - r_2)(2(1 - \alpha)^2 + (1 - \alpha)(\lambda - \alpha r_1 + 1 - \alpha)(\lambda_i(A_1) + r_1 - 2))) = 0,$$

for $i = 1, 2, \dots, n_1$.

Finally, as an application, we construct new pairs of A_α -cospectral graphs in the following corollary.

Corollary 5.4. 1. Let G_1 and G_2 be two A -cospectral regular graphs and H be an arbitrary graph. Then the graphs $G_1 \boxplus H$ and $G_2 \boxplus H$ are A_α -cospectral.

2. Let H_1 and H_2 be two A_α -cospectral graphs with $\Gamma_{A_\alpha(H_1)}(\lambda) = \Gamma_{A_\alpha(H_2)}(\lambda)$ for $\alpha \in [0, 1]$. If G is a regular graph, then the graphs $G \boxplus H_1$ and $G \boxplus H_2$ are A_α -cospectral.

References

- [1] V. Nikiforov. Merging the A - and Q -spectral theories. *Appl. Anal. Discrete Math.*, 11(1):81–107, 2017.
- [2] R. B. Bapat. *Graphs and matrices*. Universitext. Springer, London; Hindustan Book Agency, New Delhi, 2010.
- [3] S. Pirzada, Bilal A. Rather, Hilal A. Ganie, and Rezwana ul Shaban. On α -adjacency energy of graphs and Zagreb index. *AKCE Int. J. Graphs Comb.*, 18(1):39–46, 2021.
- [4] Dragoš M. Cvetković, Michael Doob, and Horst Sachs. *Spectra of graphs Theory and application*, volume 87 of *Pure and Applied Mathematics*. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, 1980.
- [5] Roberto Frucht and Frank Harary. On the corona of two graphs. *Aequationes Math.*, 4:322–325, 1970.
- [6] Xue-Qin Zhu, Gui-Xian Tian, and Shu-Yu Cui. Spectra of corona based on the total graph. *J. Math. Study*, 49(1):72–81, 2016.

- [7] Ann Susa Thomas, Sunny Joseph Kalayathankal, and Joseph Varghese Kureethara. Spectrum of corona products based on splitting graphs. *Discrete Math. Algorithms Appl.*, 15(3):Paper No. 2250087, 12, 2023.
- [8] Shu-Yu Cui and Gui-Xian Tian. The spectrum and the signless Laplacian spectrum of coronae. *Linear Algebra Appl.*, 437(7):1692–1703, 2012.
- [9] Cam McLeman and Erin McNicholas. Spectra of coronae. *Linear Algebra Appl.*, 435(5):998–1007, 2011.
- [10] S. Barik, D. Kalita, S. Pati, and G. Sahoo. Spectra of graphs resulting from various graph operations and products: a survey. *Spec. Matrices*, 6:323–342, 2018.