MEAN-FIELD MAGNETOHYDRODYNAMICS MODELS AS SCALING LIMITS OF STOCHASTIC INDUCTION EQUATIONS

FEDERICO BUTORI AND ELISEO LUONGO

ABSTRACT. We study the asymptotic properties of a stochastic model for the induction equations of the magnetic field in a three dimensional periodic domain. The turbulent velocity field driving the electromotive force on the magnetic field is modeled by a noise white in time. For this model we rigorously take a scaling limit leading to a deterministic model. While in case of isotropic turbulence this produces an additional dissipation in the limit model which influences also the decay rate of the Magnetic field in the stochastic model, the case of turbulence devoloped in a preferential direction allows us to find a dynamo effect.

1. INTRODUCTION

The evolution of the magnetic field induced by the motion of a (non-relativistic) conducting fluid with resistivity η (or conductivity $1/\eta$) is governed by the Maxwell equation and Ohm's law. Together, these laws allow to derive the so called *Induction Equation*

$$\partial_t B_t = \eta \Delta B_t + \nabla \times (u_t \times B_t), \quad \text{div } B_t = 0.$$

The two terms on the right hand side represent respectively the Ohmic dissipation of magnetic energy due to the resistivity of the media, and the magnetic field induced by the electromotive force $u_t \times B_t$, generated by the motion of the fluid. This equation is then coupled with some equation from fluid dynamics describing the evolution of the conducting fluid u_t . A correct and complete description of the system then shall include also the back reaction of B_t on u_t resulting in a system of coupled equations. This set of equations is at the core of Magneto-Hydrodynamics (MHD) and at that of the kinematic and dynamical aspects of electrically conducting fluids. In this field, an aspect that has remained obscure for many years is the mechanism with which the turbulent motion of astrophysical fluids is able to sustain large magnetic fields for timescales much longer than the dissipative one, think for example to the Sun, Earth or galaxies. Whether the fluid is in the core of a planet or on the surface of a star, it is an almost universal fact that when it is involved in rotating motion, there is the appearance of a magnetic field. The capability of these system to self excite and sustain their magnetic fields was phenomenologically explained through Dynamo Theory. However, the derivation of a precise Dynamo Theory from the induction equation above, has eluded physicists for many years. It was not until the '60s, with the work of Steenback, Krause and Radler, that the situation was substantially transformed, see the seminal paper [28] or the books [19], [23] and the references therein. The development of their two scales method, and in turn of the mean-field description of MHD allowed for a systematic treatment and understanding of the dynamo problem in turbulently moving electrically conducting media. Their approach was based on splitting the equations and the two fields in large scales $\overline{B_t}, \overline{u_t}$ and small scales components B'_t, u'_t . When the large scales are produced by some type of averaging operation, taking the average of a product is not the same as taking the product of the averages, thus the equation for the large scales produces a coupling between large and small scales:

$$\partial_t \overline{B_t} = \eta \Delta \overline{B_t} + \nabla \times (\overline{u_t} \times \overline{B_t}) + \nabla \times (\overline{u_t' \times B_t'})$$

The term $\overline{u'_t \times B'_t}$, often referred to as *Turbulent electromotive force* describes the effect of the interaction of the small scales of u and B on the large scale components. In order to close the equation for the large scales, at least in the case where the back reaction of B_t on u is neglected, Krause and Radler proposed as a first approximation to model this interaction by a mean-field effective term of the form

$$\mathcal{A}\overline{B_t} - \eta_T \nabla \times \overline{B_t}$$

Scuola Normale Superiore, Piazza dei Cavalieri, 7, 56126 Pisa, Italia

Scuola Normale Superiore, Piazza dei Cavalieri, 7, 56126 Pisa, Italia

E-mail addresses: federico.butori at sns.it, eliseo.luongo at sns.it. *Date:* June 12, 2024.

²⁰²⁰ Mathematics Subject Classification. 60H15, 76F25.

Key words and phrases. Roberts Flow, Magnetohydrodynamics, Transport Noise, Eddy Viscosity Model, Alpha Effect, Beta Effect.

where \mathcal{A} and η_T are quantities that must be derived from the statistics of u'. This two new terms are usually called *alpha* and *beta* effect terms. The mean field equation then takes the form

$$\partial_t \overline{B_t} = (\eta + \eta_T) \Delta \overline{B_t} + \nabla \times (\overline{u}_t \times \overline{B_t}) + \nabla \times (\mathcal{A}\overline{B_t})$$

From this equation we see that the role of the *beta* term $\eta_T \nabla \times \overline{B_t}$ is that of enhancing dissipation, that is, it plays against the development of a dynamo; it is instead the *alpha* term $\overline{AB_t}$ that, being of the first order, can provide the mechanism necessary to sustain the magnetic field. It is the relative magnitude of these two terms that in the end determines if the large scale magnetic field $\overline{B_t}$ is able to self-sustain or if it dissipates. Thus, the precise derivation of the quantities η_T and \mathcal{A} , is one of the main challenges of applied mean-field MHD, see [2] for a review of numerical studies on the dynamo problem.

In view of the discussion above, the aim of this work is twofold: on one hand we propose a mathematically rigorous and complete procedure to derive the mean field effective system of Krause and Radler, in the case of negligible back reaction of B_t onto u_t , from a scaling limit of a stochastic induction equation, in which the turbulent velocity field is replaced by a white noise and the product $u_t \times B_t$ is interpreted in Stratonovich form, according to a Wong-Zakai principle (*cf.* [6] and the references therein). We refer to Theorem 2.17 for the precise convergence result. On the other hand, and as a byproduct of our scheme, we are able to directly link the *alpha* and *beta* terms to the covariance of the noise, thus to the statistics of the turbulent field, so that we bypass the difficulty of having to deriving them, as they can be prescribed a priori by an appropriate choice of the noise. In this respect, the modelling choice of Gaussian noise can be seen as parallel to the second order correlation approximation (SOCA) of Krause and Radler, *cf.* [19, Chapter 4], in which the *alpha* and *beta* terms are deduced dropping correlations of order higher than two of u'. Ultimately, the Gaussian assumption, as well as the SOCA one, are too restrictive and naïve for the description of fully developed turbulence. However, we can content ourselves by dealing with this incomplete description, as doing differently would require possessing the solution of the Turbulence problem.

In order to stress the flexibility of our approach, we propose several choices of the shape of the noise that account for different turbulent regimes. In particular, we are able to explicitly identify the key role that helicity of u_t has in the appearance of the *alpha*-effect and how it is naturally linked to the size of the coefficient, in a way that is consistent with the well established models of Mean Field MHD (see for instance chapters 3 and 4 in [19]).

Our scheme is in spirit the same of our previous work [3] and falls into the category of so called Itô-Stratonovich diffusion limits initiated in [17]: the Itô-Stratonovich corrector appearing when we rewrite the term $\nabla \times (u_t \times B_t)$ from the Stratonovich form to the Itô one, corresponds precisely to the mean field term (or sum of terms) discovered in the literature under the most natural closure formulae. But in our rigorous derivation there is still the Itô term, which should disappear in the scaling limit limit. There is no obvious reason why it should disappear. Indeed, previous results in such direction where limited to consider scalar and 2D models, where stretching of vector quantities does not appear. In such case it is relatively easy to prove that the Itô term is negligible in the limit, see for example [9], [10], [4], [13], [16], [1]. In 3D, the problem was left open for a long time. Previous results in the 3D framework have drastic unphysical simplifactions as neglecting the stretching term [11], or dealing with 2D3C models [12] and their variants [3]. This work, therefore, represents the first example where the techniques of the Itô-Stratonovich diffusion limit are applied successfully in a completely 3D model. In particular it was conjectured that only a suitable geometry of the involved vector fields could remedy the excess of fluctuations in the stretching term [12]. Our approach seems to go in the opposite direction, indeed we are able to consider random vector fields u_t either in case of an isotropic structure of the covariance and in case of a preferential direction.

Although they are similar in spirit, from the physical standpoint one main differences arises between this work and [3]: in our previous work, as expected by the well known no-dynamo theorems [18, Section 3.5], since our limit equations lose any dependence from one coordinate (becomes 2D3C), the *alpha*-effect in the mean field equation cannot provide any dynamo activity for the slow-varying large scales, thus it might be classified as perturbed version of a no-dynamo theorem, displaying that a 'too simple', almost two dimensional, turbulent motion of the fluid cannot sustain a dynamo (see instead [26] for the study of the dynamo problem in 3D flows independent by one component and [25] for some others paradigmatic examples of dynamos). Here instead, even when we deduce weaker convergence results, these are enough to provide dynamo activity in the mean field equations arising from the *alpha*-effect if the covariance of the noise is sufficiently anisotropic *cf*. Corollary 2.20. On the contrary, if the breaking of the symmetry of the covariance is weak enough, the *beta*-effect is dominant, providing an enhanced decaying rate *cf*. Corollary 2.21. This includes in particular the case of isotropic mirror symmetric turbulence. In this regard, our results are in accordance with the general principle that it is really the (anisotropic) 3D features of the fluid motion that provide for dynamo activity and we confirm the famous prediction of [19, Page 12] that "Homogeneous isotropic mirror symmetric turbulence only influences the decay rate of the mean magnetic fields, which is enhanced in almost all cases of physical interest".

From the mathematical standpoint instead, the regularising effect of the thin domain is here lost, resulting in either convergence in weaker norms or in more restrictive choices of the noise coefficients. It is however important to keep in mind that, even though the choice of the noise might seems tailored to the needs of the mathematical treatise of the equation, it actually stems from physical considerations, and is tuned as to keep non-trivial and non-vanishing mean helicity of the flow throughout the limiting procedure.

The content of the paper is the following one. In section 2 we introduce rigorously our framework. In particular, we define the structure of the noise in subsection 2.3, while we state our main result and its consequences in subsection 2.4. The remainder of the paper is dedicated to proving Theorem 2.17. In section 3 we prove some estimates for B_t uniformly in the scaling limit of our noise, then we show the required convergence in section 4.

2. Functional Setting and Main Results

In subsection 2.1 we introduce the notation we follow along the paper. In subsection 2.2 we introduce some functional analysis tools we employ in order to carry on rigorously our analysis described in section 1. We describe the noise we employ in subsection 2.3 providing some insights in the meaning of the assumptions and comparisons with the existing literature. Lastly subsection 2.4 is devoted to present our main results.

2.1. Notation. In the following we denote by $\mathbb{T}^3 = \mathbb{R}^3/(2\pi\mathbb{Z})^3$ the three dimensional torus and by $\mathbb{Z}_0^3 = \mathbb{Z}^3 \setminus \{0\}$ (resp. $\mathbb{Z}_0^2 = \mathbb{Z}^2 \setminus \{0\}$) the three (resp. two) dimensional lattice made by points with integer coordinates. We introduce a partition of $\mathbb{Z}_0^3 = \Gamma_+ \cup \Gamma_-$ such that

$$\Gamma_{+} = \{k \in \mathbb{Z}_{0}^{3} : k_{3} > 0 \lor (k_{3} = 0 \land k_{2} > 0) \lor (k_{3} = k_{2} = 0 \land k_{1} > 0)\}, \quad \Gamma_{-} = -\Gamma_{+}.$$

We denote by $\{e_1, e_2, e_3\}$ the canonical basis of \mathbb{R}^3 . If $v = (v^1, v^2, v^3)^t$ and $w = (w^1, w^2, w^3)^t$ are two vector fields we will denote by $v^H = (v^1, v^2)^t$ and $\mathcal{L}_v w$ for the Lie derivative, i.e. $\mathcal{L}_v w = v \cdot \nabla w - w \cdot \nabla v$. For each $j \ge 0, l > 1$ we denote by

$$\begin{split} \zeta_{s,j}^{n} &= n^{j-3} \sum_{\substack{k \in \mathbb{Z}_{0}^{n}, \\ n \leq |k| \leq 2n}} \frac{1}{|k|^{j}}, \quad \zeta_{s,j} = 4\pi \begin{cases} \log 2 & \text{if } j = 3, \\ \frac{2^{3-j}-1}{3-j} & \text{if } j \neq 3 \end{cases}, \\ \zeta_{H,j}^{n} &= n^{j-2} \sum_{\substack{k_{H} \in \mathbb{Z}_{0}^{2} \\ n \leq |k_{H}| \leq 2n}} \frac{1}{|k_{H}|^{j}}, \quad \zeta_{H,j} = \int_{1 \leq |x| \leq 2} \frac{dx_{1}dx_{2}}{|x|^{j}} = 2\pi \begin{cases} \log 2 & \text{if } j = 2, \\ \frac{2^{2-j}-1}{2-j} & \text{if } j \neq 2 \end{cases}. \end{split}$$

In particular, by the approximation properties of Riemann sums, for each $j \ge 0$ it holds

$$\zeta_{s,j}^n = \zeta_{s,j} + O(n^{-1}), \quad \zeta_{H,j}^n = \zeta_{H,j} + O(n^{-1}).$$

Moreover, we introduce the following function of the positive octant

$$\chi: \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+ \to \mathbb{R}, \quad \chi(\alpha, \beta, \gamma) = \begin{cases} 1 & \text{if } \alpha = \beta = \gamma = 3; \\ (\gamma - 3) \wedge 1 & \text{if } \alpha = 2, \ \beta = 4, \ \gamma > 3; \\ (\alpha - 2) \wedge (\beta - 2) \wedge (\gamma - 3) & \text{if } \alpha > 2, \ \beta > 2, \ \gamma > 3; \\ 0 & \text{otherwise.} \end{cases}$$

We denote by [M, N] the quadratic covariation process, which is defined for any couple M, N of square integrable real semimartingales and we extend it by bilinearity to the analogue complex valued processes. Let a, b be two positive numbers, then we write $a \leq b$ if there exists a positive constant C such that $a \leq Cb$ and $a \leq_{\xi} b$ when we want to highlight the dependence of the constant C on a parameter ξ . If $x \in \mathbb{C}$ we denote by \overline{x} its complex conjugate. If \mathcal{Y} is a second-order tensor we denote by $\|\mathcal{Y}\|_{HS}$ its Hilbert-Schmidt norm. Lastly, if $x \in \mathbb{R}^2$ we denote by $x^{\perp} = (-x_2, x_1)^t$.

2.2. **Preliminaries.** Let us start setting some classical notation and properties of operators in the periodic setting before describing the main contributions of this work. We refer to monographs [7, 15, 22, 24, 29, 30, 32] for a complete discussion on the topics shortly recalled in this section.

2.2.1. Function Spaces in the Periodic Setting. Let $(H^{s,p}(\mathbb{T}^3), \|\cdot\|_{H^{s,p}})$, $s \in \mathbb{R}$, $p \in (1, +\infty)$ be the Bessel spaces of periodic functions. In case of p = 2, we simply write $H^s(\mathbb{T}^3)$ in place of $H^{s,2}(\mathbb{T}^3)$ and we denote by $\langle \cdot, \cdot \rangle_{H^s}$ the corresponding scalar products. In case of s = 0, we write $L^2(\mathbb{T}^3)$ instead of $H^0(\mathbb{T}^3)$ and we neglect the subscript in the notation for the norm and the inner product. With some abuse of notation, for s > 0, we denote the duality between H^{-s} and H^s by $\langle \cdot, \cdot \rangle$.

We denote by $\dot{H}^{s,p}(\mathbb{T}^3)$ (resp. $\dot{H}^s(\mathbb{T}^3)$, $\dot{L}^2(\mathbb{T}^3)$) the closed subspace of $H^{s,p}(\mathbb{T}^3)$ (resp. $H^s(\mathbb{T}^3)$, $L^2(\mathbb{T}^3)$) made by zero mean functions with norm inducted by $H^{s,p}(\mathbb{T}^3)$ (resp. $H^s(\mathbb{T}^3)$, $L^2(\mathbb{T}^3)$) and by Q the orthogonal projection of $L^2(\mathbb{T}^3)$ on $\dot{L}^2(\mathbb{T}^3)$. The restriction (resp. extension) of Q to $H^s(\mathbb{T}^3)$ (resp. $H^{-s}(\mathbb{T}^3)$) for s > 0is the orthogonal projection on $\dot{H}^s(\mathbb{T}^3)$ (resp. $\dot{H}^{-s}(\mathbb{T}^3)$). A different characterization of the fractional Sobolev spaces can be given in terms of powers of the Laplacian. Denoting by

$$\Delta: \mathsf{D}(\Delta) \subseteq \dot{L^2}(\mathbb{T}^3) \to \dot{L}^2(\mathbb{T}^3),$$

where $\mathsf{D}(\Delta) = \dot{H}^2(\mathbb{T}^3)$, it is well known that Δ is the infinitesimal generator of analytic semigroup of negative type that we denote by $e^{t\Delta} : \dot{L}^2(\mathbb{T}^3) \to \dot{L}^2(\mathbb{T}^3)$ and moreover for each s > 0, $\mathsf{D}((-\Delta)^s)$ (resp. $(\mathsf{D}((-\Delta)^s))'$) can be identified with $\dot{H}^{2s}(\mathbb{T}^3)$ (resp. $\dot{H}^{-2s}(\mathbb{T}^3)$).

Similarly, we introduce the Bessel spaces of vector fields

$$H^{s,p}(\mathbb{T}^3; \mathbb{R}^3) = \{ u = (u^1, u^2, u^3)^t : u^1, u^2, u^3 \in H^{s,p}(\mathbb{T}^3) \}$$
$$\langle u, v \rangle_{H^s} = \sum_{j=1}^3 \langle u^j, v^j \rangle_{H^s}, \quad \text{for } s \in \mathbb{R}.$$

Again, in case of s = 0 we write $L^2(\mathbb{T}^3; \mathbb{R}^3)$ instead of $H^0(\mathbb{T}^3; \mathbb{R}^3)$ and we neglect the subscript in the notation for the norm and the scalar product. Similarly we denote by $\dot{H}^s(\mathbb{T}^3; \mathbb{R}^3)$ (resp. $\dot{L}^2(\mathbb{T}^3; \mathbb{R}^3)$) the closed subspace of $H^s(\mathbb{T}^3; \mathbb{R}^3)$ (resp. $L^2(\mathbb{T}^3; \mathbb{R}^3)$) made by zero mean functions. With some abuse of notation we usually employ the same symbols that we introduced for norms and inner products of scalar functions also for the corresponding vector fields and we continue to denote the orthogonal projection between $H^s(\mathbb{T}^3; \mathbb{R}^3)$ and $\dot{H}^s(\mathbb{T}^3; \mathbb{R}^3)$, $s \in \mathbb{R}$, by Q, since it acts component-wise.

Lastly we denote by \mathbf{H}^s (resp. \mathbf{L}^2) the closed subspace of $\dot{H}^s(\mathbb{T}^3; \mathbb{R}^3)$ (resp. $\dot{L}^2(\mathbb{T}^3; \mathbb{R}^3)$) made by zero mean, divergence free vector fields with norm induces by $H^s(\mathbb{T}^3; \mathbb{R}^3)$ (resp. $L^2(\mathbb{T}^3; \mathbb{R}^3)$) and by

$$P: \dot{L}^2(\mathbb{T}^3; \mathbb{R}^3) \to \mathbf{L}^2$$

the Leray projection which, in the periodic setting, is an orthogonal projection even when acting between $\dot{H}^{s}(\mathbb{T}^{3};\mathbb{R}^{3})$ and \mathbf{H}^{s} . If $\{\frac{k}{|k|}, a_{k,1}, a_{k,2}\}$ is an orthonormal system of \mathbb{R}^{3} for each $k \in \mathbb{Z}_{0}^{3}$ then, it is well-known that the family $\{\frac{1}{(2\pi)^{3/2}}a_{k,j}e^{ik\cdot x}\}_{k\in\mathbb{Z}_{0}^{3}}$ is an orthonormal system of \mathbf{L}^{2} and is orthogonal in \mathbf{H}^{s} for each $s \in \mathbb{R}$.

We conclude this subsection introducing some classical notation when dealing with stochastic processes taking values in separable Hilbert spaces. Let Z be a separable Hilbert space, with associated norm $\|\cdot\|_Z$. We denote by $C_{\mathcal{F}}([0,T];Z)$ the space of weakly continuous adapted processes $(X_t)_{t\in[0,T]}$ with values in Z such that

$$\mathbb{E}\left[\sup_{t\in[0,T]}\|X_t\|_Z^2\right] < \infty$$

and by $L^p_{\mathcal{F}}(0,T;Z)$, $p \in [1,\infty)$, the space of progressively measurable processes $(X_t)_{t \in [0,T]}$ with values in Z such that

$$\mathbb{E}\bigg[\int_0^T \|X_t\|_Z^p \, dt\bigg] < \infty.$$

2.2.2. Operators in the Periodic Setting. In this section we are interested to the properties of differential operators of the form

$$\mu \partial_{11} + \mu \partial_{22} + \nu \partial_{33}, \ \mu \ge \nu > 0,$$

the case $\mu = \nu$ corresponding to the Laplacian. It is well known that for each $\mu \ge \nu > 0$, the operator

$$A_{\mu,\nu}: \dot{H}^2(\mathbb{T}^3) \subseteq \dot{L}^2(\mathbb{T}^3) \to \dot{L}^2(\mathbb{T}^3), \quad A_{\mu,\nu}f = \mu\partial_{11}f + \mu\partial_{22}f + \nu\partial_{33}f$$

is closed and generates an analytic semigroup of negative type which we denote by $S^{\mu,\nu}(t) : \dot{L}^2(\mathbb{T}^3) \to \dot{L}^2(\mathbb{T}^3)$. The family $\{\frac{1}{(2\pi)^{3/2}}e^{ik\cdot x}\}_{k\in\mathbb{Z}^3_0}$ is a complete orthogonal systems of $\dot{H}^s(\mathbb{T}^3)$ made by eigengunctions of $A_{\mu,\nu}$ for each $\mu \geq \nu > 0$. Moreover it is orthonormal in $\dot{L}^2(\mathbb{T}^3)$. This allows us to write the operators $A_{\mu,\nu}$, $S^{\mu,\nu}(t)$ in Fourier basis as

$$A_{\mu,\nu}f = -\frac{1}{(2\pi)^3} \sum_{k \in \mathbb{Z}_0^3} \left(\mu(k_1^2 + k_2^2) + \nu k_3^2 \right) \langle f, e^{-ik \cdot x} \rangle e^{ik \cdot x}, \ S^{\mu,\nu}(t) = \frac{1}{(2\pi)^3} \sum_{k \in \mathbb{Z}_0^3} e^{-\left(\mu(k_1^2 + k_2^2) + \nu k_3^2\right)t} \langle g, e^{-ik \cdot x} \rangle e^{ik \cdot x}$$

for $f \in \dot{H}^2(\mathbb{T}^3)$, $g \in \dot{L}^2(\mathbb{T}^3)$. The expression in Fourier basis has as immediate consequences the following facts which we recall here for the convenience of the reader

Lemma 2.1. For each $s \ge 0$, the extension (resp. restriction) of $S^{\mu,\nu}(t)$: $\dot{H}^{-s}(\mathbb{T}^3) \to \dot{H}^{-s}(\mathbb{T}^3)$ (resp. $S^{\mu,\nu}(t)$: $\dot{H}^{s}(\mathbb{T}^{3}) \rightarrow \dot{H}^{s}(\mathbb{T}^{3})$ is an analytic semigroup of negative type. Moreover, for each $\mu_{1} \geq \nu_{1} > 0, \ \mu_{2} \geq \nu_{2} > 0$ the operators $S^{\mu_1,\nu_1}(t)$ and $S^{\mu_1,\nu_1}(t)$ commute. In particular

$$S^{\mu_1,\nu_1}(t)S^{\mu_2,\nu_2}(t) = S^{\mu_2,\nu_2}(t)S^{\mu_1,\nu_1}(t) = S^{\mu_1+\mu_2,\nu_1+\nu_2}(t).$$

Lemma 2.2. Let $q \in \dot{H}^s(\mathbb{T}^3)$, $s \in \mathbb{R}$. Then:

- i) for any $\varphi \ge 0$, it holds $\|S^{\mu,\nu}(t)q\|_{\dot{H}^{s+\varphi}} \lesssim_{\varphi} (t\nu)^{-\varphi/2} \|q\|_{\dot{H}^{s}}$, the hidden constant being increasing in φ ; ii) for any $\varphi \in [0,2]$, it holds $\|(I-S^{\mu,\nu}(t))q\|_{\dot{H}^{s-\varphi}} \lesssim (t\mu)^{\varphi/2} \|q\|_{\dot{H}^{s}}$. The hidden constant being uniformly bounded for $\varphi \in [0, 2]$.

Lastly we need a closedness results on operators of this form, which has some similarities with the second item of Lemma 2.2.

Lemma 2.3. Let $q \in \dot{H}^s(\mathbb{T}^3)$, $s \in \mathbb{R}$. If $\mu_1 \ge \nu_1 > 0$ and $\mu_2 \ge \nu_2 > 0$, for any $\varphi \in [0,2]$, it holds

$$\|(S^{\mu_1,\nu_1}(t)-S^{\mu_2,\nu_2}(t))\,q\|_{\dot{H}^{s-\varphi}} \lesssim (|\mu_1-\mu_2|+|\nu_1-\nu_2|)^{\varphi/2}\,t^{\varphi/2}\|q\|_{\dot{H}^s}.$$

The hidden constant being uniformly bounded for $\varphi \in [0, 2]$.

Proof. Thanks to the Fourier expansion formula (2.1) we have

$$\begin{split} \| (S^{\mu_1,\nu_1}(t) - S^{\mu_2,\nu_2}(t)) \, q \|_{\dot{H}^{s-\varphi}}^2 &= \sum_{k \in \mathbb{Z}_0^3} \left(e^{-\left(\mu_1(k_1^2 + k_2^2) + \nu_1 k_3^2\right)t} - e^{-\left(\mu_2(k_1^2 + k_2^2) + \nu_2 k_3^2\right)t} \right)^2 \langle q, \frac{e^{-ik \cdot x}}{(2\pi)^{3/2}} \rangle^2 \frac{|k|^{2s}}{|k|^{2\varphi}} \\ &\leq \| q \|_{\dot{H}^s}^2 \sup_{k \in \mathbb{Z}_0^3} \frac{\left(e^{-\left(\mu_1(k_1^2 + k_2^2) + \nu_1 k_3^2\right)t} - e^{-\left(\mu_2(k_1^2 + k_2^2) + \nu_2 k_3^2\right)t} \right)^2}{|k|^{2\varphi}}. \end{split}$$

Therefore, the proof is complete if we are able to show that

$$\sup_{\mathbf{k}\in\mathbb{Z}_{0}^{3}}\frac{\left(e^{-\left(\mu_{1}(k_{1}^{2}+k_{2}^{2})+\nu_{1}k_{3}^{2}\right)t}-e^{-\left(\mu_{2}(k_{1}^{2}+k_{2}^{2})+\nu_{2}k_{3}^{2}\right)t}\right)^{2}}{|k|^{2\varphi}}\lesssim_{\varphi}\left(|\mu_{1}-\mu_{2}|+|\nu_{1}-\nu_{2}|\right)^{\varphi}t^{\varphi}.$$

This is indeed the case. If $\mu_1 < \mu_2$ and $\nu_1 < \nu_2$

$$\frac{\left(e^{-\left(\mu_1(k_1^2+k_2^2)+\nu_1k_3^2\right)t}-e^{-\left(\mu_2(k_1^2+k_2^2)+\nu_2k_3^2\right)t\right)^2}}{|k|^{2\varphi}} \leq \frac{\left(1-e^{-\left((\mu_2-\mu_1)(k_1^2+k_2^2)+(\nu_2-\nu_1)k_3^2\right)t\right)^2}}{|k|^{2\varphi}}$$
$$\leq t^{\varphi}(|\mu_2-\mu_1|\wedge|\nu_2-\nu_1|)^{\varphi}\frac{\left(1-e^{-\left((\mu_2-\mu_1)(k_1^2+k_2^2)+(\nu_2-\nu_1)k_3^2\right)t\right)^2}}{t^{\varphi}\left(|\mu_1-\mu_2|(k_1^2+k_2^2)+|\nu_1-\nu_2|k_3^2\right)^{\varphi}}$$
$$\lesssim t^{\varphi}(|\mu_2-\mu_1|\wedge|\nu_2-\nu_1|)^{\varphi}$$

since $\frac{1-e^{-x}}{x^{\varphi/2}}$ is bounded for x > 0, uniformly in $\varphi \in [0, 2]$. In case of $\mu_1 < \mu_2$ and $\nu_2 < \nu_1$ and either $(k_1, k_2) \neq 0$ and $k_3 \neq 0$

$$\frac{\left(e^{-\left(\mu_1(k_1^2+k_2^2)+\nu_1k_3^2\right)t}-e^{-\left(\mu_2(k_1^2+k_2^2)+\nu_2k_3^2\right)t}\right)^2}{|k|^{2\varphi}} \lesssim \frac{\left(1-e^{-\left((\mu_2-\mu_1)(k_1^2+k_2^2)\right)t}\right)^2}{(k_1^2+k_2^2)^{\varphi}} + \frac{\left(1-e^{-\left((\nu_1-\nu_2)k_3^2\right)t}\right)^2}{|k_3|^{2\varphi}} \\ \leq t^{\varphi}|\mu_2-\mu_1|^{\varphi} \frac{\left(1-e^{-(\mu_2-\mu_1)(k_1^2+k_2^2)t}\right)^2}{(t|\mu_1-\mu_2|(k_1^2+k_2^2))^{\varphi}}$$

$$+ t^{\varphi} |\nu_2 - \nu_1|^{\varphi} \frac{\left(1 - e^{-(\nu_1 - \nu_2)k_3^2 t}\right)^2}{\left(t |\nu_1 - \nu_2|k_3^2\right)^{\varphi}} \\ \lesssim t^{\varphi} \left(|\mu_2 - \mu_1|^{\varphi} + |\nu_2 - \nu_1|^{\varphi}\right).$$

The case $(k_1, k_2) = 0$ or $k_3 = 0$ easily reconducts to the one treated. This completes the proof.

With some abuse of notation we denote by

$$A_{\mu,\nu} : \mathbf{H}^2 \subseteq \mathbf{L}^2 \to \mathbf{L}^2, \quad A_{\mu,\nu}f = (A_{\mu,\nu}f^1, A_{\mu,\nu}f^2, A_{\mu,\nu}f^3)^t.$$

The Fourier expansion of (2.1), then allows us to write for $f \in \mathbf{H}^s$

$$A_{\mu,\nu}f = -\frac{1}{(2\pi)^3} \sum_{\substack{k \in \mathbb{Z}_0^3\\j \in \{1,2\}}} \left(\mu(k_1^2 + k_2^2) + \nu k_3^2 \right) \langle f, a_{k,j}e^{-ik \cdot x} \rangle e^{ik \cdot x} a_{k,j}$$

This implies that $A_{\mu,\nu}$ generates an analytic semigroup on \mathbf{L}^2 which we still denote by $S^{\mu,\nu}(t)$ and Lemma 2.1, Lemma 2.2, Lemma 2.3 continue to hold in this vector valued framework.

2.3. Description of the Stochastic System. We begin with the description of the noise. We give a general description which cover all the three regimes we are interested in: we call them the Helical noise, the perturbed 2D noise and the Isotropic noise. Choose $\rho \in [-1, 1]$, a correlation factor whose role will be that of introducing some correlation between the vertical and horizontal components of the '2D' modes of the random field. Then assume $(\Omega, \mathcal{F}, \mathcal{F}_t, \mathbb{P})$ is a filtered probability space such that $(\Omega, \mathcal{F}, \mathbb{P})$ is a complete probability space, $(\mathcal{F}_t)_{t \in [0,T]}$ is a right continuous filtration and \mathcal{F}_0 contains every \mathbb{P} null subset of Ω . Now, for each $n \in \mathbb{N}$, $\{W_t^{k,j}\}$, $k \in \mathbb{Z}_0^3$, $j \in \{1,2\}$ is a sequence of complex-valued Brownian motions adapted to \mathcal{F}_t such that $W_t^{-k,j} = (W_t^{k,j})^*$ and

$$\mathbb{E}\left[W_{1}^{k,j}, W_{1}^{l,m^{*}}\right] = \begin{cases} 2 & \text{if } k = l, \ m = j \\ 2\rho & \text{if } k = l, \ k_{3} = 0, \ m \neq j; \end{cases} \quad \begin{bmatrix}W_{\cdot}^{k,j}, W_{\cdot}^{l,m}\end{bmatrix}_{t} = \begin{cases} 2t & \text{if } k = -l, \ m = j \\ 2\rho t & \text{if } k = -l, \ k_{3} = 0, \ m \neq j. \end{cases}$$

Then, for each $k \in \mathbb{Z}_0^3$, $j \in \{1, 2\}$ we denote by $\sigma_{k,j}^n(x) = \theta_{k,j}^n a_{k,j} e^{ik \cdot x}$, where $\{\frac{k}{|k|}, a_{k,1}, a_{k,2}\}$ is an orthonormal system of \mathbb{R}^3 for $k_3 > 0$ and $a_{k,j} = a_{-k,j}$ if $k_3 < 0$, while the (complex) coefficients $\theta_{k,j}^{\varepsilon}$ will be defined later. It is well known that the family $\{a_{k,j}\frac{1}{2\pi\sqrt{2\pi}}e^{ik \cdot x}\}$, $k \in \mathbb{Z}_0^3$, $j \in \{1, 2\}$ is a complete orthogonal systems of \mathbf{H}^s made by eigenfunctions of $-\Delta$. Moreover it is orthonormal in \mathbf{L}^2 . Without losing of generality, we make a choice that will simplify some of the computations: when $k_3 = 0$ we choose $a_{k,1} = (-k_2/|k|, k_1/|k|, 0)$, $a_{k,2} = (0, 0, 1)$ if $k \in \Gamma_+$, $a_{k,j} = a_{-k,j}$ otherwise. For a physical intuition behind our construction see our previous work [3] and the discussion therein. Then we define

$$W_t^n = \sum_{\substack{k \in \mathbb{Z}_0^3 \\ j \in \{1,2\}}} \sigma_{k,j}^n W_t^{k,j}$$

with $\sigma_{k,j}^n = \theta_{k,j}^n a_{k,j} e^{ik \cdot x}$. With our choices, to ensure that W^n takes values in a space of real valued functions, the coefficients must satisfy the condition

(2.2)
$$\theta_{k,j}^{n} = \theta_{-k,j}^{n}$$

The explicit choices of the coefficients $\theta_{k,i}^n$:

(2.3)
$$\theta_{k,j}^{n} = \mathbf{1}_{\{n \le |k| \le 2n\}} \begin{cases} \frac{i}{C_{1,H}|k|^{\alpha/2}} & \text{if } k_{3} = 0, k \in \Gamma_{+}, \ j = 1\\ \frac{-i}{C_{1,H}|k|^{\alpha/2}} & \text{if } k_{3} = 0, k \in \Gamma_{-}, \ j = 1\\ \frac{1}{C_{2,H}|k|^{\beta/2}} & \text{if } k_{3} = 0, \ j = 2\\ \frac{1}{C_{V}|k|^{\gamma/2}} & \text{if } k_{3} \neq 0. \end{cases}$$

allows us to distinguish our three regimes:

Hypothesis 2.4 (Helical).

- $\alpha \ge 2$, $\beta > 2$, $\gamma > 3$, $\alpha + \beta = 6$.
- $C_{1,H}, C_{2,H}, C_V > 0 \text{ and } C_{1,H} > \sqrt{\frac{\zeta_{H,2}}{\eta}} \text{ if } \alpha = 2.$

Hypothesis 2.5 (Perturbed 2D).

- $\alpha = 2, \ \beta = 4, \ \gamma \geq 5.$
- $C_{1,H} > \sqrt{\frac{\zeta_{H,2}}{\eta}}, \ C_{2,H}, \ C_V > 0.$

Hypothesis 2.6 (Isotropic).

• $\alpha = \beta = \gamma = 3.$

•
$$C_{1,H} = C_{2,H} = C_V > \sqrt{\frac{2\zeta_{s,3}}{3\eta}}$$

Remark 2.7. The term 'isotropic' in Hypothesis 2.6 is understood with a slight abuse, indeed it refers to the fact that the scaling parameters are all equal, but not to the fact that the fields W_t^n are actually isotropic, a condition that is met only in the special case of $\rho = 0$, and if one sets all the coefficients to be real. However, since our analysis and our main result remains unchanged whether if ρ is zero or not in this regime, we included also the latter case for generality.

The Perturbed 2D regime, instead, is a sub case of the Helical regime. Its relevance comes from the fact that due to the strong suppression of true 3D modes ($\gamma \ge 5$), this model represents a small 3D perturbation of a 2D-3C flow with weak vertical components ($\beta = 4$), and akin to the works [12], [3] which previously employed similar noises, this choice is the only one that allows us to obtain a slightly stronger convergence result.

Remark 2.8. Due to our choice of the coefficients $\theta_{k,j}^n$ and the vectors $a_{k,j}$ it follows that for each k such that $k_3 = 0, n \le |k| \le 2n$

$$\theta_{k,1}^{n}a_{k,1} = \frac{i}{C_{1,H}} \frac{k^{\perp}}{|k|^{\alpha/2+1}}, \ \theta_{k,2}^{n}a_{k,2} = \frac{1}{C_{2,H}} \frac{e_3}{|k|^{\beta/2+1}}$$

The space covariance function associated to our noise is

(2.4) $Q^{n}(x,y) = \mathbb{E}\left[W_{1}^{n}(x) \otimes W_{1}^{n}(y)^{*}\right] = Q_{0}^{n}(x,y) + \overline{Q_{\rho}^{n}}(x,y) = \overline{Q^{n}}(x,y) + (Q^{n})'(x,y) + \overline{Q_{\rho}^{n}}(x,y)$ where we denoted by

$$\overline{Q^{n}}(x,y) = 2 \sum_{\substack{k \in \mathbb{Z}_{0}^{3} \\ k_{3}=0, \quad j \in \{1,2\}}} \left| \theta_{k,j}^{n} \right|^{2} a_{k,j} \otimes a_{k,j} e^{ik \cdot (x-y)}, \quad (Q^{n})'(x,y) = 2 \sum_{\substack{k \in \mathbb{Z}_{0}^{3} \\ k_{3} \neq 0, \quad j \in \{1,2\}}} \left| \theta_{k,j}^{n} \right|^{2} a_{k,j} \otimes a_{k,j} e^{ik \cdot (x-y)}, \quad (Q^{n})'(x,y) = 2 \sum_{\substack{k \in \mathbb{Z}_{0}^{3} \\ k_{3} \neq 0, \quad j \in \{1,2\}}} \left| \theta_{k,j}^{n} \right|^{2} a_{k,j} \otimes a_{k,j} e^{ik \cdot (x-y)}, \quad (Q^{n})'(x,y) = 2 \sum_{\substack{k \in \mathbb{Z}_{0}^{3} \\ k_{3} \neq 0, \quad j \in \{1,2\}}} \left| \theta_{k,j}^{n} \right|^{2} a_{k,j} \otimes a_{k,j} e^{ik \cdot (x-y)}, \quad (Q^{n})'(x,y) = 2 \sum_{\substack{k \in \mathbb{Z}_{0}^{3} \\ k_{3} \neq 0, \quad j \in \{1,2\}}} \left| \theta_{k,j}^{n} \right|^{2} a_{k,j} \otimes a_{k,j} e^{ik \cdot (x-y)},$$

Due to our choice, we have that $\overline{Q^n}(x,y)$, $(Q^n)'(x,y)$ and $\overline{Q^n_\rho}(x,y)$ are all translation invariant, therefore we simply write $Q^n(x-y)$ (resp. $Q_0^n(x-y)$, $\overline{Q^n}(x-y)$, $(Q^n)'(x-y)$, $\overline{Q^n_\rho}(x-y)$) in place of $Q^n(x,y)$ (resp. $Q_0^n(x,y)$, $\overline{Q^n}(x,y)$, $\overline{Q^n}(x,y)$, $(Q^n)'(x,y)$, $\overline{Q^n_\rho}(x,y)$). In particular $Q_0^n(x)$, $\overline{Q^n}(x)$, $(Q^n)'(x)$ are mirror symmetric, namely it holds

(2.5)
$$Q_0^n(x) = Q_0^n(-x), \ \overline{Q^n}(x) = \overline{Q^n}(-x), \ (Q^n)'(x) = (Q^n)'(-x)$$

On the contrary $\overline{Q_{\rho}^{n}}(x)$ is an odd function, in particular $\overline{Q_{\rho}^{n}}(0) = 0$. As a consequence, our noise satisfies the mirror symmetry property if and only if $\rho = 0$.

We recall here some properties of the covariance functions $\overline{Q_0^n}(x), (Q_0^n)'(x), \overline{Q_\rho^n}(x)$. Their proofs are a simple adaptation of the arguments of [3, Section 2.3] to which we refer for details.

Lemma 2.9. If either Hypothesis 2.4 or Hypothesis 2.5 hold, then

$$\overline{Q_0^n}(0) = 2\eta_T^n(e_1 \otimes e_1 + e_2 \otimes e_2) + 2\eta_R^n(e_3 \otimes e_3), \quad (Q_0^n)'(0) = 2(\eta_V^n - \eta_{R,V}^n)I + \eta_{R,V}^nI_2$$

where we set

$$\eta_T^n = \frac{\zeta_{H,\alpha}^n}{2C_{1,H}^2 n^{\alpha-2}} = O(n^{2-\alpha}), \quad \eta_R^n = \frac{\zeta_{H,\beta}^n}{C_{2,H}^2 n^{\beta-2}} = O(n^{2-\beta}), \quad \eta_V^n = \frac{2\zeta_{s,\gamma-3}^n}{3C_V^2 n^{\gamma-3}} = O(n^{3-\gamma}),$$
$$\eta_{R,V}^n = \frac{\zeta_{H,\gamma}^n}{C_V^2 n^{\gamma-2}} = O(n^{2-\gamma}).$$

In particular, if $\alpha = 2$

$$\eta_T^n = \frac{\zeta_{H,2}}{2C_{1,H}^2} + O(n^{-1}).$$

Lemma 2.10. If Hypothesis 2.6 holds, then

$$Q_0^n(0) = 2\eta_{iso}^n I$$

where we set

$$\eta_{iso}^n = \frac{2\zeta_{s,3}^n}{3C_V^2} = \frac{2\zeta_{s,3}}{3C_V^2} + O(n^{-1}).$$

Lemma 2.11. It holds

$$(2.6) \\ \left(\partial_{l}\overline{Q_{0}^{n}}\right)(0) = 2i \sum_{\substack{k \in \mathbb{Z}_{0}^{3}, \\ k_{3}=0, \ j \in \{1,2\}}} \left|\theta_{k,j}^{n}\right|^{2} k_{l} a_{k,j} \otimes a_{k,j} = 0, \quad \left(\partial_{l}Q_{0}^{n}\right)'(0) = 2i \sum_{\substack{k \in \mathbb{Z}_{0}^{3}, \\ k_{3}\neq 0, \ j \in \{1,2\}}} \left|\theta_{k,j}^{n}\right|^{2} k_{l} a_{k,j} \otimes a_{k,j} = 0.$$

0.

As a consequence, for each $l, r, s \in \{1, 2, 3\}$ and $f \in L^2(\mathbb{T}^3)$, $g \in H^1(\mathbb{T}^3)$ it holds

(2.7)
$$\sum_{\substack{k \in \mathbb{Z}_{0}^{3}, \\ k_{3}=0, \ j \in \{1,2\}}} \langle \partial_{l}g\sigma_{k,j}^{n,l}, f \partial_{r}\sigma_{-k,j}^{n,s} \rangle = \sum_{\substack{k \in \mathbb{Z}_{0}^{3}, \\ k_{3}\neq 0, \ j \in \{1,2\}}} \langle \partial_{l}g\sigma_{k,j}^{n,l}, f \partial_{r}\sigma_{-k,j}^{n,s} \rangle =$$

Lemma 2.12. We have

(2.8)
$$\nabla \overline{Q_{\rho}^{n}}(0) = \frac{2\rho}{C_{1,H}C_{2,H}} \sum_{\substack{k_{H} \in \mathbb{Z}_{0}^{2} \\ n \leq |k_{H}| \leq 2n}} \frac{1}{|k_{H}|^{2}} \left(e_{3} \otimes \frac{k_{H}^{\perp}}{|k_{H}|} \otimes \frac{k_{H}}{|k_{H}|} - \frac{k_{H}^{\perp}}{|k_{H}|} \otimes e_{3} \otimes \frac{k_{H}}{|k_{H}|} \right).$$

In particular, it holds

$$\left\|\nabla \overline{Q_{\rho}^{n}}(0)\right\|_{HS} \leq \frac{2\sqrt{2}|\rho|\zeta_{H,2}^{n}}{C_{1,H}C_{2,H}}$$

and

$$\begin{aligned} \{\mathcal{R}_{n}^{r,s}\}_{r,s\in\{1,2\}} &:= \partial_{s}\overline{Q_{\rho}^{n,3,r}}(0) = \frac{\rho\zeta_{H,2}^{N}}{C_{1,H}C_{2,H}} \begin{bmatrix} 0 & -1\\ 1 & 0 \end{bmatrix} = \frac{\rho\zeta_{H,2}}{C_{1,H}C_{2,H}} \begin{bmatrix} 0 & -1\\ 1 & 0 \end{bmatrix} + O(n^{-1}),\\ \{\mathcal{V}_{n}^{r,s}\}_{r,s\in\{1,2\}} &:= \partial_{s}\overline{Q_{\rho}^{n,r,3}}(0) = -\partial_{s}\overline{Q_{\rho}^{n,3,r}}(0). \end{aligned}$$

As discussed in [15, Chapter 3], the quantity $Q^n(0)$ is the core in order to obtain a nontrivial second order operator in our Itô-Stratonovich diffusion limit, i.e. the *beta* effect we discussed in section 1. In particular, the choice of Hypothesis 2.4 and $\alpha = 2$ or Hypothesis 2.6 allows us to obtain a non trivial *beta* effect in the scaling limit. As discussed in section 1, the *alpha* effect is instead linked to the helicity of the velocity field. The role of the helicity is usually understood in terms of distortion of the axial magnetic field lines that generates currents in the rotating direction, which in turn induce magnetic fields in the axial direction, see for example [19, Chapter 3, figure 3.2] and related comments. In our stochastic framework we denote the time average of the helicity of our random velocity field

$$\mathcal{H}^{n}(x) := \frac{1}{2T} \mathbb{E} \left[W^{n}_{T}(x) \cdot \left(\nabla \times W^{n}_{T}(x) \right) \right]$$

Exploiting the definition of our random vector field and arguing as in [3, Appendix B] we can easily see that $\mathcal{H}^n(x)$ is independent on x and it satisfies

$$\mathcal{H}^{n}(x) = -\frac{2\rho\zeta_{H,\frac{\alpha+\beta-2}{2}}^{n}}{C_{1,H}C_{2,H}n^{\frac{\alpha+\beta-6}{2}}} \longrightarrow \mathcal{H} := -\frac{4\pi\rho\log 2}{C_{1,H}C_{2,H}},$$

i.e. the choice of $\alpha + \beta = 6$ is linked to reach a finite and nontrivial *alpha* term in the limit.

2.4. Main Results. Following the ideas introduced in section 1, we fix $T \in (0, +\infty)$ and we are interested in studying the properties of the following stochastic model on \mathbb{T}^3 :

(2.9)
$$\begin{cases} dB_t^n = \eta \Delta B_t^n dt + \sum_{\substack{k \in \mathbb{Z}_0^3 \\ j \in \{1,2\}}} \mathcal{L}_{\sigma_{k,j}^n} B_t^n \circ dW_t^{k,j}, \\ \text{div} B_t^n = 0, \\ B_t^n|_{t=0} = B_0^n, \end{cases}$$

with the coefficients $\sigma_{k,j}^n$ defined in subsection 2.3. We start rewriting it in Itô form. First, let us observe that for each $k \in \mathbb{Z}_0^3$, $j \in \{1, 2\}$ we have due to the linearity of $\mathcal{L}_{\sigma_{k,j}^n}$

$$\begin{aligned} \mathcal{L}_{\sigma_{k,j}^{n}} B_{t}^{n} \circ dW_{t}^{k,j} &= \mathcal{L}_{\sigma_{k,j}^{n}} B_{t}^{n} dW_{t}^{k,j} + \frac{1}{2} \left[\mathcal{L}_{\sigma_{k,j}^{n}} B_{\cdot}^{n}, W_{\cdot}^{k,j} \right]_{t} \\ &= \mathcal{L}_{\sigma_{k,j}^{n}} B_{t}^{n} dW_{t}^{k,j} + \frac{1}{2} \mathcal{L}_{\sigma_{k,j}^{n}} \left[B_{\cdot}^{n}, W_{\cdot}^{k,j} \right]_{t} \\ &= \mathcal{L}_{\sigma_{k,j}^{n}} B_{t}^{n} dW_{t}^{k,j} + \mathcal{L}_{\sigma_{k,j}^{n}} \mathcal{L}_{\sigma_{-k,j}^{n}} B_{t}^{n} dt + \rho \mathbf{1}_{\{k_{3}=0, \ l \neq j\}} \mathcal{L}_{\sigma_{k,j}^{n}} \mathcal{L}_{\sigma_{-k,l}^{n}} B_{t}^{n} dt. \end{aligned}$$

Therefore the equation for B_t^n can be rewritten as

$$\begin{cases} dB_t^n &= \left(\eta \Delta + \sum_{\substack{k \in \mathbb{Z}_0^3 \\ j \in \{1,2\}}} \mathcal{L}_{\sigma_{k,j}^n} \mathcal{L}_{\sigma_{-k,j}^n} + \rho \sum_{\substack{k \in \mathbb{Z}_0^3 \\ k_3 = 0}} \mathcal{L}_{\sigma_{k,1}^n} \mathcal{L}_{\sigma_{-k,2}^n} + \mathcal{L}_{\sigma_{k,2}^n} \mathcal{L}_{\sigma_{-k,1}^n} \right) B_t^n dt \\ &+ \sum_{\substack{k \in \mathbb{Z}_0^3 \\ j \in \{1,2\}}} \mathcal{L}_{\sigma_{k,j}^n} B_t^n dW_t^{k,j}, \\ \operatorname{div} B_t^n &= 0, \\ B_t^n|_{t=0} &= B_0^n. \end{cases}$$

We are left to identify $\sum_{\substack{k \in \mathbb{Z}_0^3 \ j \in \{1,2\}}} \mathcal{L}_{\sigma_{k,j}^n} \mathcal{L}_{\sigma_{-k,j}^n} + \rho \sum_{\substack{k \in \mathbb{Z}_0^3 \ k_3=0}} \mathcal{L}_{\sigma_{k,1}^n} \mathcal{L}_{\sigma_{-k,2}^n} + \mathcal{L}_{\sigma_{k,2}^n} \mathcal{L}_{\sigma_{-k,1}^n}$. We sketch the proof of the following result for the convenience of the reader, in order to highlights the role of $Q_0^n(0)$ and $\nabla \overline{Q_\rho^n}(0)$ in our procedure.

Lemma 2.13. If $F : \mathbb{T}^3 \to \mathbb{R}^3$ is a smooth, zero mean divergence free vector field, then it holds

$$\sum_{\substack{k \in \mathbb{Z}_0^3\\j \in \{1,2\}}} \mathcal{L}_{\sigma_{k,j}^n} \mathcal{L}_{\sigma_{-k,j}^n} F + \rho \sum_{\substack{k \in \mathbb{Z}_0^3\\k_3 = 0}} \left(\mathcal{L}_{\sigma_{k,1}^n} \mathcal{L}_{\sigma_{-k,2}^n} + \mathcal{L}_{\sigma_{k,2}^n} \mathcal{L}_{\sigma_{-k,1}^n} \right) F = \Lambda^n F + \Lambda_\rho^n F$$

where in the formula above we denoted by Λ^n the differential operator

$$\Lambda^{n}F = \begin{cases} (\eta_{T}^{n} + \eta_{V}^{n} - \eta_{R,V}^{n}/2)(\partial_{1}^{2} + \partial_{2}^{2})F + (\eta_{R}^{n} + \eta_{V}^{n} - \eta_{R,V}^{n})\partial_{3}^{2}F & \text{if Hypothesis 2.4 holds,} \\ \eta_{iso}^{n}\Delta F & \text{if Hypothesis 2.6 holds} \end{cases}$$

and by Λ_{ρ}^{n} the differential operator

$$\begin{split} \Lambda^n_{\rho}F &= -\sum_{l \in \{1,2\}} \partial_l \overline{Q^n_{\rho}}(0) \cdot \nabla F_l = -\sum_{\substack{j \in \{1,2,3\}\\l \in \{1,2\}}} \partial_l \overline{Q^{n,\cdot,j}_{\rho}}(0) \partial_j F_l \\ &= \nabla \times (\mathcal{A}^n_{\rho}F), \end{split}$$

where

$$\mathcal{A}_{\rho}^{n} = \frac{\rho \zeta_{H,2}^{n}}{C_{1,H}C_{2,H}} \begin{bmatrix} 1 & 0 & 0\\ 0 & 1 & 0\\ 0 & 0 & 0 \end{bmatrix}.$$

In particular it holds

$$(\Lambda_{\rho}^{n}F)_{H} = \frac{\rho \zeta_{H,2}^{n}}{C_{1,H}C_{2,H}} \partial_{3}F_{H}^{\perp}, \quad (\Lambda_{\rho}^{n}F)_{3} = -\frac{\rho \zeta_{H,2}^{n}}{C_{1,H}C_{2,H}} \operatorname{div}_{H}(F_{H}^{\perp})$$

Proof. All the terms involved are Lie brackets, thus we begin with some preliminary computations. By recalling that we have $\nabla \sigma_{k,j}^n = i\theta_{k,j}^n k \otimes a_{k,j} e^{ikx}$, and also that $a_{-k,j} = a_{k,j}$ for all $k \in \mathbb{Z}_0^3$, j = 1, 2 we write

(2.10)
$$\sigma_{k,l}^{n} \cdot \nabla(\sigma_{-k,j}^{n} \cdot \nabla F) = (\sigma_{k,l}^{n} \cdot \nabla \sigma_{-k,j}^{n}) \cdot \nabla F + \sigma_{k,l}^{n} \otimes \sigma_{-k,j}^{n} : \nabla^{2} F$$
$$= -i\theta_{k,l}^{n}(a_{k,l} \cdot k)a_{-k,l} \cdot \nabla F + \sigma_{k,l}^{n} \otimes \sigma_{-k,j}^{n} : \nabla^{2} F$$
$$= \sigma_{k,l}^{n} \otimes \sigma_{-k,j}^{-n} : \nabla^{2} F$$

since $a_{k,l} \cdot k = 0$. On the other hand we have

(2.11)
$$(F \cdot \nabla \sigma_{-k,j}^n) \cdot \nabla \sigma_{k,l}^n = \theta_{-k,j}^n \theta_{k,l}^n (F \cdot k) (a_{k,j} \cdot k) a_{k,l} = 0$$

while

(2.12)
$$\sigma_{-k,j}^n \cdot \nabla \left(F \cdot \nabla \sigma_{k,l}^n \right) = i \theta_{k,l}^n \theta_{-k,j}^n e^{-ikx} a_{k,j}^r \partial_r ((F \cdot k) a_{k,l} e^{ikx})$$

$$= i\theta_{k,l}^n \theta_{-k,j}^n e^{-ikx} a_{k,j}^r ((\partial_r F \cdot k) a_{k,l} e^{ikx} + (F \cdot k) a_{k,l} k_r e^{ikx})$$

$$= i\theta_{k,l}^n \theta_{-k,j}^n a_{k,j}^r (\partial_r F \cdot k) a_{k,l}$$

and

(2.13)
$$\left(\sigma_{k,l}^n \cdot \nabla F\right) \cdot \nabla \sigma_{-k,j}^n = -i\theta_{k,l}^n \theta_{-k,j}^n a_{k,l}^r (\partial_r F \cdot k) a_{k,j} \right)$$

In particular we see that if l = j the sum of (2.12) and (2.13) is zero. We are now ready to compute the correctors. We have

$$\mathcal{L}_{\sigma_{k,j}^{n}}\mathcal{L}_{\sigma_{-k,j}^{n}}F = \sigma_{k,j}^{n} \cdot \nabla(\sigma_{-k,j}^{n} \cdot \nabla F) - \sigma_{k,j}^{n} \cdot \nabla(F \cdot \nabla \sigma_{-k,j}^{n}) - (\sigma_{-k,j}^{n} \cdot \nabla F) \cdot \nabla \sigma_{k,j}^{n} + (F \cdot \nabla \sigma_{-k,j}^{n}) \cdot \nabla \sigma_{k,j}^{n} + (F \cdot \nabla \sigma_{-k,j}^$$

By (2.11) the last term is zero, while by (2.12) and (2.13), the sum of the two middle terms is zero. Summing over $k \in \mathbb{Z}_0^3$ and j = 1, 2 we get

(2.14)
$$\sum_{\substack{k \in \mathbb{Z}_0^3\\j \in \{1,2\}}} \mathcal{L}_{\sigma_{k,j}^n} \mathcal{L}_{\sigma_{-k,j}^n} F = \sum_{\substack{k \in \mathbb{Z}_0^3\\j \in \{1,2\}}} \sigma_{k,l}^n \otimes \sigma_{-k,j}^n : \nabla^2 F$$
$$= \operatorname{div}(Q_0^n(0) \nabla F)$$
$$= \Lambda^n F$$

where we used again the fact that $\sigma_{k,j}^n \cdot \nabla \sigma_{-k,j}^n = 0$ in the second line and Lemma 2.9, Lemma 2.10 in the last equality.

We now look at the part of the corrector relative to the correlation ρ , that is

(2.15)
$$\rho \sum_{\substack{k \in \mathbb{Z}_0^n \\ k_3 = 0}} \left(\mathcal{L}_{\sigma_{k,1}^n} \mathcal{L}_{\sigma_{-k,2}^n} + \mathcal{L}_{\sigma_{k,2}^n} \mathcal{L}_{\sigma_{-k,1}^n} \right)$$

We write the general form of the two summands, for $l \neq j$, as

$$\mathcal{L}_{\sigma_{k,l}^n}\mathcal{L}_{\sigma_{-k,j}^n}F = \sigma_{k,l}^n \cdot \nabla(\sigma_{-k,j}^n \cdot \nabla F) - \sigma_{k,l}^n \cdot \nabla(F \cdot \nabla \sigma_{-k,j}^n) - (\sigma_{-k,j}^n \cdot \nabla F) \cdot \nabla \sigma_{k,l}^n + (F \cdot \nabla \sigma_{-k,j}^n) \cdot \nabla \sigma_{k,l}^n$$

Again, by (2.11) the last term is always zero. Now summing over k and $l \neq j$ we see, using first (2.10) and reasoning as in the manipulation of (2.14)

(2.16)
$$\sum_{\substack{k \in \mathbb{Z}_0^n \\ k_3 = 0}} \sigma_{k,1}^n \cdot \nabla(\sigma_{-k,2}^n \cdot \nabla F) + \sigma_{k,2}^n \cdot \nabla(\sigma_{-k,1}^n \cdot \nabla F) = \operatorname{div}(\overline{Q_{\rho}^n}(0)\nabla F) = 0.$$

We are left with the contributions of the middle terms which no longer sum to zero as in the first corrector. Recalling our choice of the coefficients (2.3), we see that if $l \neq j$ and $n \leq |k| \leq 2n$, $k_3 = 0$

$$\theta_{k,j}^n \theta_{-k,l}^n = -\theta_{k,l}^n \theta_{-k,j}^n$$

Thus, when we sum over $l \neq j$, we get from (2.12) and (2.13) that

$$(2.17) \qquad -\sigma_{k,1}^{n} \cdot \nabla (F \cdot \nabla \sigma_{-k,2}^{n}) - (\sigma_{-k,1}^{n} \cdot \nabla F) \cdot \nabla \sigma_{k,2}^{n} = 2i\theta_{k,1}\theta_{-k,2}a_{k,1}^{r}(\partial_{r}F \cdot k)a_{k,2} \\ = -\frac{2\mathrm{sgn}(k)}{|k|^{\alpha/2+\beta/2}}a_{k,1}^{r}(\partial_{r}F \cdot k)a_{k,2}$$

and similarly

$$(2.18) \qquad -(\sigma_{-k,2}^n \cdot \nabla F) \cdot \nabla \sigma_{k,1}^n - \sigma_{k,2}^n \cdot \nabla (F \cdot \nabla \sigma_{-k,1}^n) = -2i\theta_{-k,2}^n \theta_{k,1}^n a_{k,2}^r (\partial_r F \cdot k) a_{k,1} = \frac{2\mathrm{sgn}(k)}{|k|^{\alpha/2+\beta/2}} a_{k,2}^r (\partial_r F \cdot k), a_{k,1}$$

where we indicated con sgn(k) the function that takes value 1 if $k \in \Gamma^+$ and -1 if $k \in \Gamma^-$. Summing the last two expressions and then summing over $k \in \mathbb{Z}_0^3$, $k_3 = 0$ it is straightforward to see thanks to Lemma 2.12 that

$$\rho \sum_{\substack{k \in \mathbb{Z}_0^3 \\ k_3 = 0}} \left(\mathcal{L}_{\sigma_{k,1}^n} \mathcal{L}_{\sigma_{-k,2}^n} + \mathcal{L}_{\sigma_{k,2}^n} \mathcal{L}_{\sigma_{-k,1}^n} \right) = -\sum_{l \in \{1,2\}} \partial_l \overline{Q_\rho^n}(0) \cdot \nabla F_l$$
$$= \Lambda_\rho^n F.$$

According to Lemma 2.13, equation (2.9) can be rewritten in Itô form as:

(2.19)
$$\begin{cases} dB_t^n = \left(\eta\Delta + \Lambda^n + \Lambda_\rho^n\right) B_t^n dt + \sum_{\substack{k \in \mathbb{Z}_0^3 \\ j \in \{1,2\}}} \mathcal{L}_{\sigma_{k,j}^n} B_t^n dW_t^{k,j} \\ \operatorname{div} B_t^n = 0, \\ B_t^n|_{t=0} = B_0^n, \end{cases}$$

We give now the definition of weak solution for the Stochastic equation of the magnetic field (2.19).

Definition 2.14. A stochastic process

$$B^n \in C_{\mathcal{F}}([0,T];\mathbf{L}^2) \cap L^2_{\mathcal{F}}([0,T];\mathbf{H}^1)$$

is a weak solution of equation (2.19) if, for every $\phi \in \mathbf{H}^2$, we have (2.20)

$$\langle B_t^n, \phi \rangle = \langle B_0^n, \phi \rangle + \eta \int_0^t \langle B_s^n, \Delta \phi \rangle \, ds + \int_0^t \langle B_s^n, \Lambda^n \phi \rangle \, ds + \int_0^t \langle \Lambda_\rho^n B_s^n, \phi \rangle \, ds + \sum_{\substack{k \in \mathbb{Z}_0^n \\ j \in \{1,2\}}} \int_0^t \langle \mathcal{L}_{\sigma_{k,j}^n} B_s^n, \phi \rangle dW_s^{k,j} + \int_0^t \langle B_s^n, \Delta \phi \rangle \, ds + \int_0^t \langle B_s^n, \Delta \phi \rangle$$

for every $t \in [0,T]$, $\mathbb{P}-a.s$.

Remark 2.15. Due to the definition of the operator P, each vector field $\phi \in \dot{H}^2(\mathbb{T}^3; \mathbb{R}^3)$ can be decomposed in a divergence free part, i.e. $P\phi \in \mathbf{H}^2$, and a part orthogonal to \mathbf{L}^2 , i.e. $(I - P)\phi \in \dot{H}^2(\mathbb{T}^3; \mathbb{R}^3)$. Therefore for a generic $\phi \in \dot{H}^2(\mathbb{T}^3; \mathbb{R}^3)$ we get

$$\begin{split} \langle B_t^n, \phi \rangle - \langle B_0^n, \phi \rangle &= \langle B_t^n, P\phi \rangle - \langle B_0^n, P\phi \rangle \\ &= \eta \int_0^t \langle B_s^n, \Delta P\phi \rangle \, ds + \int_0^t \langle B_s^n, \Lambda^n P\phi \rangle \, ds + \int_0^t \langle \Lambda_\rho^n B_s^n, P\phi \rangle \, ds \\ &+ \sum_{\substack{k \in \mathbb{Z}_0^3 \\ j \in \{1,2\}}} \int_0^t \langle \mathcal{L}_{\sigma_{k,j}^n} B_s^n, P\phi \rangle dW_s^{k,j} \\ &= \eta \int_0^t \langle B_s^n, \Delta \phi \rangle \, ds + \int_0^t \langle B_s^n, \Lambda^n \phi \rangle \, ds + \int_0^t \langle \Lambda_\rho^n B_s^n, \phi \rangle \, ds + \sum_{\substack{k \in \mathbb{Z}_0^3 \\ j \in \{1,2\}}} \int_0^t \langle \mathcal{L}_{\sigma_{k,j}^n} B_s^n, \phi \rangle dW_s^{k,j} \end{split}$$

for every $t \in [0,T]$, $\mathbb{P} - a.s.$, due to the fact that $B^n \in C_{\mathcal{F}}([0,T]; \mathbf{L}^2) \cap L^2_{\mathcal{F}}([0,T]; \mathbf{H}^1)$ and the structure of the operators Λ^n , Λ^n_{ρ} . In particular, if we choose $\phi = (0,0,\varphi) \in \dot{H}^2(\mathbb{T}^3; \mathbb{R}^3)$ we obtain

$$\begin{split} \langle B_t^{n,3},\varphi\rangle - \langle B_0^{n,3},\varphi\rangle &= \eta \int_0^t \langle B_s^{n,3},\Delta\varphi\rangle \, ds + \int_0^t \langle B_s^{n,3},\Lambda^n\varphi\rangle \, ds + \frac{\rho\zeta_{H,2}^n}{C_{1,H}C_{2,H}} \int_0^t \langle B_s^{n,H^{\perp}},\nabla_H\varphi\rangle \, ds \\ &+ \sum_{\substack{k \in \mathbb{Z}_0^3\\j \in \{1,2\}}} \int_0^t \langle \sigma_{k,j}^n \cdot \nabla B_s^{n,3} - B^{n,3} \cdot \nabla \sigma_{k,j}^{n,3},\varphi\rangle dW_s^{k,j} \end{split}$$

for every $t \in [0, T]$, $\mathbb{P} - a.s$.

Due to the fact that equation (2.19) is linear, the existence and uniqueness of solutions in the sense of Definition 2.14 is a standard fact which follows by the abstract theory of [8, Chapters 3-5], see also [14, Section 3.1]. Indeed the following holds.

Theorem 2.16. For each $B_0^n \in \mathbf{L}^2$ there exists a unique weak solution of system (2.19) in the sense of Definition 2.14.

As discussed in section 1, we study the convergence of B_t^n in the scaling limit of the separation of scales, i.e. considering a noise which concentrates on smaller and smaller scales as described in subsection 2.3. Lastly we

introduce our limit objects. Following the idea first introduced in [17], we expect that our limit objects satisfy a PDE with an additional second order operator with intensity related to

$$\lim_{n \to +\infty} \sum_{\substack{k \in \mathbb{Z}_0^3\\ j \in \{1,2\}}} \left\| \sigma_{k,j}^n \right\|^2.$$

The quantity above is different from 0 if $\alpha = 2$ in Hypothesis 2.4, Hypothesis 2.5 or if it holds Hypothesis 2.6. Besides to the fact that the limit above may be 0 under our assumptions, the final deterministic objects keep memory of our noise thanks to the fact that $\alpha + \beta = 6$ for each *n*, which implies that our noise has a nontrivial limit helicity. Denoting by $\overline{B_t}$ the unique weak solutions of the following linear 3D PDE

(2.21)
$$\begin{cases} \partial_t \overline{B_t} &= (\eta \Delta + \Lambda_{\alpha,\beta,\gamma}) \overline{B_t} + \nabla \times (\mathcal{A}_{\rho} \overline{B_t}) \quad x \in \mathbb{T}^3, \ t \in (0,T) \\ \operatorname{div} \overline{B} &= 0 \quad x \in \mathbb{T}^3, \ t \in (0,T) \\ \overline{B_t}|_{t=0} &= \overline{B_0}, \end{cases}$$

where

$$\Lambda_{\alpha,\beta,\gamma} = \begin{cases} 0 & \text{if Hypothesis 2.4 holds and } \alpha > 2, \\ \eta_T \Delta_H & \text{if Hypothesis 2.4 holds and } \alpha = 2, \\ \eta_{iso} \Delta & \text{if Hypothesis 2.6 holds,} \end{cases} \quad \mathcal{A}_{\rho} = \frac{2\pi\rho\log 2}{C_{1,H}C_{2,H}} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix},$$

 $\eta_T = \frac{\zeta_{H,2}}{2C_{1,H}^2}, \ \eta_{iso} = \frac{2\zeta_{s,3}}{3C_V^2}.$ Our main result reads as follows:

Theorem 2.17. Let $B_0^n \in \mathbf{L}^2$, assuming Hypothesis 2.4 or Hypothesis 2.6 and $B_0^n \to \overline{B_0}$ in \mathbf{H}^{-1} , for each $\kappa \in [1,2), \ \theta \in (0,2), \ \delta \in (0,\theta)$ we have (2.22)

$$\sup_{t\in[0,T]} \mathbb{E}\left[\left\|B_t^n - \overline{B_t}\right\|_{\mathbf{H}^{-1-\theta}}^{\kappa}\right] \lesssim \begin{cases} \frac{1}{n^{\kappa\theta\chi(\alpha,\beta,\gamma)/2}} + \frac{1}{n^{\frac{\kappa(\alpha,\beta\beta,\gamma)(\theta-\delta)}{6+\delta}}} + \left\|\overline{B_0} - B_0^n\right\|_{\mathbf{H}^{-1-\theta}}^{\kappa} & \text{in case of Hypothesis 2.4,} \\ \frac{1}{n^{\frac{3\kappa(\theta-\delta)}{6+\delta}}} + \left\|\overline{B_0} - B_0^n\right\|_{\mathbf{H}^{-1-\theta}}^{\kappa} & \text{in case of Hypothesis 2.6,} \end{cases}$$

Moreover, assuming also $B^{n,3} \rightharpoonup \overline{B_0^3}$ in $\dot{L}^2(\mathbb{T}^3; \mathbb{R}^3)$, in case of Hypothesis 2.5 we have also for each $\vartheta \in (\theta, \frac{3}{2}]$

$$(2.23) \quad \sup_{t \in [0,T]} \mathbb{E}\left[\left\| B_t^{n,3} - \overline{B_t^3} \right\|_{\dot{H}^{-\vartheta}}^{\kappa} \right] \lesssim \frac{1}{n^{\frac{\kappa(\vartheta - \theta)(\theta - \delta)}{6+\delta}}} + \frac{1}{(\vartheta - \theta)^{\kappa} n^{\kappa\theta/2}} + \left\| \overline{B_0^3} - B_0^{n,3} \right\|_{\dot{H}^{-\vartheta}}^{\kappa} + \frac{\left\| \overline{B_0} - B_0^n \right\|_{\mathbf{H}^{-1-\theta}}^{\kappa}}{(\vartheta - \theta)^{\kappa}}.$$

In particular either the right hand side of (2.22) and of (2.23) converge to 0 as $n \to +\infty$.

Remark 2.18. The hidden constant depends on all the parameters of the model and blows-up as $\delta \to 0$ or if either $\eta_{iso} \nearrow \eta$ in case of Hypothesis 2.6 or $2\eta_T \nearrow \eta$ in case of Hypothesis 2.4 and $\alpha = 2$.

Remark 2.19. Due to Hypothesis 2.6 $\eta_{iso} < \eta$. In case of Hypothesis 2.4 with $\alpha = 2$ we have $\eta_T < \frac{\eta}{2}$.

Contrary to our previous work [3], here also a compactness approach inspired to [12, Section 5] works. Indeed, arguing similarly to Lemma 4.6, it is possible to study the behaviour of the time increments of B_t^n in some Sobolev spaces of negative order. Combining this control and the uniform estimates of Proposition 3.4 allows to prove the tightness of the laws of $\{B^n\}_{n\in\mathbb{N}}$ in $L^2(0,T;H^{-\delta})$ for each $\delta > 0$. Then, arguing by Prokhorov's theorem and Skorokhod's representation theorem, it is possible to find an auxiliary probability space where, up to passing to subsequences, $B^n \to \overline{B}$ in $L^2(0,T;H^{-\delta}) = -a.s$. Arguing similarly to section 4, we can identify \overline{B} as the unique weak solution of (2.21). This implies that actually the full sequence $B^n \to \overline{B}$ in $L^2(0,T;H^{-\delta})$ in probability. Similar arguments applies also to $B^{n,3}$ in case of Hypothesis 2.5. We preferred to rely on semigroup techniques and state our main result in the form of Theorem 2.17 for a twofold reason. First, this quantitative result gives explicit information on the rate of convergence which we think are of independent interest. Secondly, the convergence in $C([0,T]; L^2(\Omega; \mathbf{H}^{-1-\theta}))$ guaranteed by Theorem 2.17 gives us more information about dynamo action of the stochastic model. Indeed, as a corollary of Theorem 2.17 we can easily obtain Corollary 2.20 and Corollary 2.21.

Corollary 2.20. Let $k = \lambda e_3$, for $\lambda \in \mathbb{Z}_0$ and $B_0^n = \overline{B_0} = (\sin(k \cdot x), \cos(k \cdot x), 0)^t$. Assuming either Hypothesis 2.4 or Hypothesis 2.5, if $\lambda \rho > 0$ and $C_{2,H} < \frac{2\pi \rho \log 2}{\lambda C_{1,H} \eta}$, then for each $\theta \in (0,2)$, $\delta \in (0,\theta)$

$$\mathbb{E}\left[\left\|B_t^n\right\|^2\right] \gtrsim \frac{1}{\lambda^{2(1+\theta)}} e^{-(\lambda \mathcal{H} + 2\lambda^2 \eta)t} - \frac{1}{n^{\theta \chi(\alpha,\beta,\gamma)}} - \frac{1}{n^{\frac{2(\alpha \wedge \beta \wedge \gamma)(\theta - \delta)}{6+\delta}}}.$$

In particular, the energy of B^n is increasing in time in mean.

Proof. the unique solution of equation (2.21) with initial condition $(\sin(k \cdot x), \cos(k \cdot x), 0)^t$ is given by

$$\overline{B_t} = e^{-(\lambda \frac{\mathcal{H}}{2} + \lambda^2 \eta)t} (\sin(k \cdot x), \cos(k \cdot x), 0)^t.$$

In particular, for each $\theta \in (0, 2)$

$$\left\|\overline{B_t}\right\|_{\mathbf{H}^{-1-\theta}}^2 = \frac{8\pi^3}{\lambda^{2(1+\theta)}} e^{-(\lambda\mathcal{H}+2\lambda^2\eta)t}.$$

Under our assumptions $\lambda \frac{\mathcal{H}}{2} + \lambda^2 \eta < 0$. Therefore, since for each $\kappa \in [1, 2)$

$$\left\|\overline{B_t}\right\|_{\mathbf{H}^{-1-\theta}}^{\kappa} \le 2^{\kappa-1} \left(\left\|B_t^n\right\|_{\mathbf{H}^{-1-\theta}}^{\kappa} + \left\|B_t^n - \overline{B_t}\right\|_{\mathbf{H}^{-1-\theta}}^{\kappa} \right),$$

thanks to Theorem 2.17 and Jensen's inequality we obtain

$$\mathbb{E}\left[\left\|B_{t}^{n}\right\|^{2}\right]^{\frac{1}{2}} \geq \mathbb{E}\left[\left\|B_{t}^{n}\right\|_{\mathbf{H}^{-1-\theta}}^{\kappa}\right] \\ \geq \frac{1}{\lambda^{\kappa(1+\theta)}}e^{-\kappa(\lambda\frac{\mathcal{H}}{2}+\lambda^{2}\eta)t} - \frac{1}{n^{\kappa\theta\chi(\alpha,\beta,\gamma)/2}} - \frac{1}{n^{\frac{\kappa(\alpha\wedge\beta\wedge\gamma)(\theta-\delta)}{6+\delta}}}$$

which implies the claim.

Corollary 2.21. If Hypothesis 2.6 holds and $B_0^n = \overline{B_0} \in \mathbf{L}^2$, then for each $\kappa \in [1, 2)$, $\theta \in (0, 2)$, $\delta \in (0, \theta)$ we have

$$\mathbb{E}\left[\left\|B_t^n\right\|_{\mathbf{H}^{-1-\theta}}^{\kappa}\right] \le \frac{C}{n^{\frac{3\kappa(\theta-\delta)}{6+\delta}}} + 2^{\kappa-1} \left\|\overline{B_0}\right\|^{\kappa} e^{-\kappa \left(\eta + \frac{(8-6)\rho)|\pi \log 2}{3C_V^2}\right)t}$$

where C is a constant depending on $\overline{B_0}, C_V, \rho, \theta, \delta, \kappa$.

Proof. Thanks to simple energy estimates on (2.21) and Poincaré inequality we obtain

$$\left\|\overline{B_{t}}\right\|^{2} + 2\left(\eta + \frac{8\pi\log 2}{3C_{V}^{2}}\right)\int_{0}^{t}\left\|\nabla\overline{B_{s}}\right\|^{2}ds \leq \left\|\overline{B_{0}}\right\|^{2} + \frac{4\pi|\rho|\log 2}{C_{V}^{2}}\int_{0}^{t}\left\|\nabla\overline{B_{s}}\right\|^{2}ds.$$

Therefore, by Grönwall's inequality

$$\left\|\overline{B_t}\right\|^{\kappa} \le \left\|\overline{B_0}\right\|^{\kappa} e^{-\kappa \left(\eta + \frac{(8-6|\rho|)\pi \log 2}{3C_V^2}\right)t}.$$

As a consequence of the inequality above and Theorem 2.17 we obtain

$$\mathbb{E}\left[\left\|B_{t}^{n}\right\|_{\mathbf{H}^{-1-\theta}}^{\kappa}\right] \leq 2^{\kappa-1} \left(\mathbb{E}\left[\left\|B_{t}^{n}-\overline{B_{t}}\right\|_{\mathbf{H}^{-1-\theta}}^{\kappa}\right]+\left\|\overline{B_{t}}\right\|_{\mathbf{H}^{-1-\theta}}^{\kappa}\right)$$
$$\leq \frac{C_{B_{0},C_{V},\rho,\theta,\delta,\kappa}}{n^{\frac{3\kappa(\theta-\delta)}{6+\delta}}}+2^{\kappa-1}\left\|\overline{B_{0}}\right\|^{\kappa}e^{-\kappa\left(\eta+\frac{(8-6|\rho|)\pi\log 2}{3C_{V}^{2}}\right)t}.$$

The two corollaries above have a clear interpretation in terms of the discussion of section 1 and subsection 2.3. Indeed, Corollary 2.20 shows that if the helicity of our velocity field is large enough and the turbulent fluid is sufficiently anisotropic, the energy of our stochastic models increases in time. Therefore, a dynamo effect is active on our stochastic models in case of Hypothesis 2.4. On the contrary, in case of Hypothesis 2.6, due to Corollary 2.21, the dissipation acting on magnetic field increases due to the presence of the turbulent fluid. This is accordance to [19]. Moreover, Corollary 2.21 implies that it is not needed that the fluid is either completely isotropic or mirror symmetric in order to increase the dissipative effects acting on the magnetic field.

Remark 2.22. Contrary to Hypothesis 2.6, the matrix $Q^n(0)$ converge to a degenerate matrix as $n \to +\infty$ in case of Hypothesis 2.4. This phenomena produces a degenerate second order operator in the scaling limit. It is possible to avoid such a phenomena in case of Hypothesis 2.4 and $\alpha = 2$ considering a more general noise

$$W_t^n = \frac{1}{3} \sum_{\substack{k \in \mathbb{Z}_0^n \\ j \in \{1,2\}}} a_{k,j} e^{ik \cdot x} \left(\theta_{1,k,j}^n W_t^{1,k,j} + \theta_{2,k,j}^n W_t^{2,k,j} + \theta_{3,k,j}^n W_t^{3,k,j} \right)$$

where $W_t^{h,k,j}$ are complex Brownian motions satisfying

$$\mathbb{E}\left[W_{1}^{h,k,j}, W_{1}^{h',l,m^{*}}\right] = \begin{cases} 0 & \text{if } h \neq h' \\ 2 & \text{if } h = h', \ k = l, \ m = j \\ 2\rho_{h} & \text{if } h = h', \ k = l, \ k_{h} = 0, \ m \neq j \end{cases}$$

and

$$\theta_{h,k,j}^{n} = \mathbf{1}_{\{n \le |k| \le 2n\}} \begin{cases} \frac{i}{C_{1,H}|k|^{\alpha/2}} & \text{if } k_{h} = 0, k \in \Gamma_{+}, \ j = 1\\ \frac{-i}{C_{1,H}|k|^{\alpha/2}} & \text{if } k_{h} = 0, k \in \Gamma_{-}, \ j = 1\\ \frac{1}{C_{2,H}|k|^{\beta/2}} & \text{if } k_{h} = 0, \ j = 2\\ \frac{1}{C_{V}|k|^{\gamma/2}} & \text{if } k_{h} \neq 0. \end{cases}$$

In this case we get

$$\Lambda_{\alpha,\beta,\gamma} = \frac{2}{3}\eta_T \Delta, \quad \mathcal{A}_{\rho_1,\rho_2,\rho_3} = \frac{2\pi \log 2}{3C_{1,H}C_{2,H}} \left(\rho_1 \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} + \rho_2 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} + \rho_3 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \right)$$

and we can prove an analogous result to Theorem 2.17. Since this change would drastically complicate the notation without any change either in the proof of Theorem 2.17 and in the physical understanding of the dynamo and the dissipative properties of our stochastic systems, i.e. Corollary 2.20 and Corollary 2.21, we prefer to limit ourselves to the noise introduced in subsection 2.3. In this spirit the *helical* regime could be regarded as a stochastic perturbation of a Robert's flow [26] in which we can also allow dependence on the third space variable. This kind of flows are extensively studied in the physical and numerical literature as paradigmatic models for dynamo action (see for instance [27], [31], [5] and references therein)

3. A Priori Estimates

The goal of this section is to provide uniform bounds for the \mathbf{H}^{-1} norm of our vector fields B_t^n . In order to complete our plan it is useful to recall a well-known result about equivalence of norms for divergence free vector fields, see for example [22].

Proposition 3.1. Let $X \in \mathbf{L}^2$, $Y \in \mathbf{H}^1$. Then

(3.1)
$$||X||^2 = ||(-\Delta)^{-1/2} \nabla \times X||^2, \quad ||\nabla Y||^2 = ||\nabla \times Y||^2.$$

We start providing a control to some quantities related to the noise appearing in equation (2.19).

Lemma 3.2. Assuming Hypothesis 2.4, then

$$\sum_{\substack{k \in \mathbb{Z}_0^n \\ i \in \{1,2\}}} \|\sigma_{k,j}^n \times B_t^n\|^2 \le \left(\frac{\zeta_{H,\alpha}^n}{C_{1,H}^2 n^{2-\alpha}} + \frac{\zeta_{H,\beta}^n}{C_{2,H}^2 n^{2-\beta}} + \frac{2\zeta_{s,\gamma}^n}{C_V^2 n^{3-\gamma}}\right) \|B_t^n\|^2.$$

In case of Hypothesis 2.6, then

$$\sum_{\substack{k \in \mathbb{Z}_0^3 \\ j \in \{1,2\}}} \|\sigma_{k,j}^n \times B_t^n\|^2 = \frac{4\zeta_{s,3}^n}{3C_V^2} \|B_t^n\|^2.$$

In particular, assuming Hypothesis 2.5 we have also

$$\begin{split} &\sum_{\substack{k \in \mathbb{Z}_0^3 \\ j \in \{1,2\}}} \|\sigma_{k,j}^n \cdot \nabla B_t^{n,3}\|^2 \leq \frac{\zeta_{H,\alpha}^n \|\nabla_H B_t^{n,3}\|^2}{2C_{1,H}^2 n^{\alpha-2}} + \frac{\zeta_{H,\beta}^n \|\partial_3 B_t^{n,3}\|^2}{C_{2,H}^2 n^{\beta-2}} + \frac{2\zeta_{s,\gamma}^n \|\nabla B_t^{n,3}\|^2}{3C_V^2 n^{\gamma-3}}, \\ &\sum_{\substack{k \in \mathbb{Z}_0^3 \\ j \in \{1,2\}}} \|B_t^n \cdot \nabla \sigma_{k,j}^{n,3}\|^2 \leq \left(\frac{\zeta_{H,2}^n}{C_{2,H}^2} + \frac{\zeta_{s,\gamma-2}^n}{C_V^2 n^{\gamma-5}}\right) \|B_t^n\|^2. \end{split}$$

Proof. By Hölder's inequality we have

$$\sum_{\substack{k \in \mathbb{Z}_0^3 \\ j \in \{1,2\}}} \|\sigma_{k,j}^n \times B_t^n\|^2 \le \sum_{\substack{k \in \mathbb{Z}_0^3 \\ j \in \{1,2\}}} |\theta_{k,j}^n|^2 \|B_t^n\|^2.$$

In the first case, due to the relation above we get

$$\begin{split} \sum_{\substack{k \in \mathbb{Z}_0^3 \\ j \in \{1,2\}}} \|\sigma_{k,j}^n \times B_t^n\|^2 &\leq \left(\sum_{\substack{k \in \mathbb{Z}_0^3 \\ k_3 = 0, \quad j = 1}} |\theta_{k,j}^n|^2 + \sum_{\substack{k \in \mathbb{Z}_0^3 \\ k_3 = 0, \quad j = 2}} |\theta_{k,j}^n|^2 + \sum_{\substack{k \in \mathbb{Z}_0^3 \\ k_3 \neq 0, \quad j \in \{1,2\}}} |\theta_{k,j}^n|^2 \right) \|B_t^n\|^2 \\ &\leq \left(\sum_{\substack{k \in \mathbb{Z}_0^2 \\ n \leq |k| \leq 2n}} \frac{1}{C_{1,H}^2 |k|^\alpha} + \sum_{\substack{k \in \mathbb{Z}_0^2 \\ n \leq |k| \leq 2n}} \frac{1}{C_{2,H}^2 |k|^\beta} + 2\sum_{\substack{k \in \mathbb{Z}_0^3 \\ n \leq |k| \leq 2n}} \frac{1}{C_V^2 |k|^\gamma} \right) \|B_t^n\|^2 \\ &\leq \left(\frac{\zeta_{H,\alpha}^n}{C_{1,H}^2 n^{2-\alpha}} + \frac{\zeta_{H,\beta}^n}{C_{2,H}^2 n^{2-\beta}} + \frac{2\zeta_{s,\gamma}^n}{C_V^2 n^{3-\gamma}}\right) \|B_t^n\|^2. \end{split}$$

However, in the second case, due to the symmetries of the coefficients, we can obtain a sharp relation which is our claim. Let us denote by

$$b_t^{n,h,l} := \langle B_t^n, a_{h,l} e^{-ih \cdot x} \rangle,$$

therefore by definition

$$B_{t}^{n} = \sum_{\substack{h \in \mathbb{Z}_{0}^{3}, \\ l \in \{1,2\}}} b_{t}^{n,h,l} a_{h,l} e^{ih \cdot x},$$

$$\sigma_{k,j}^{n} \times B_{t}^{n} = \theta_{k,j}^{n} \sum_{\substack{h \in \mathbb{Z}_{0}^{3}, \\ l \in \{1,2\}}} b_{t}^{n,h,l} a_{k,j} \times a_{h,l} e^{i(h+k) \cdot x}$$

and for each k, j it holds

(3.2)

$$\begin{split} \|\sigma_{k,j}^{n} \times B_{t}^{n}\|^{2} &= |\theta_{k,j}^{n}|^{2} \sum_{\substack{h \in \mathbb{Z}_{0}^{3}, \\ l,l' \in \{1,2\}}} (a_{k,j} \times a_{h,l}) \cdot (a_{k,j} \times a_{h,l'}) b_{t}^{n,h,l} \overline{b_{t}^{n,h,l'}} \\ &= |\theta_{k,j}^{n}|^{2} \sum_{\substack{h \in \mathbb{Z}_{0}^{3}, \\ l,l' \in \{1,2\}}} (\delta_{l,l'} - a_{k,j} \otimes a_{k,j} : a_{h,l} \otimes a_{h,l'}) b_{t}^{n,h,l} \overline{b_{t}^{n,h,l'}} \\ &= |\theta_{k,j}^{n}|^{2} \|B_{t}^{n}\|^{2} - |\theta_{k,j}^{n}|^{2} \sum_{\substack{h \in \mathbb{Z}_{0}^{3}, \\ l,l' \in \{1,2\}}} a_{k,j} \otimes a_{k,j} : a_{h,l} \otimes a_{h,l'} b_{t}^{n,h,l} \overline{b_{t}^{n,h,l'}}. \end{split}$$

Therefore, in case of Hypothesis 2.6 we obtain

$$\sum_{\substack{k \in \mathbb{Z}_{0}^{3} \\ j \in \{1,2\}}} \|\sigma_{k,j}^{n} \times B_{t}^{n}\|^{2} = \frac{2\zeta_{s,3}^{n}}{C_{V}^{2}} \|B_{t}^{n}\|^{2} - \sum_{\substack{h \in \mathbb{Z}_{0}^{3}, \\ l,l' \in \{1,2\}}} \frac{2\zeta_{s,3}^{n}}{3C_{V}^{2}} I : (a_{h,l} \otimes a_{h,l'}) b_{t}^{n,h,l} \overline{b_{t}^{n,h,l'}}$$
$$= \frac{4\zeta_{s,3}^{n}}{3C_{V}^{2}} \|B_{t}^{n}\|^{2}.$$

Lastly we assume Hypothesis 2.5, in this case only few terms in the sum must be considered due to our choice of the $\sigma_{k,j}^n$'s. Then, we obtain

$$\sum_{\substack{k \in \mathbb{Z}_0^3 \\ j \in \{1,2\}}} \|\sigma_{k,j}^n \cdot \nabla B_t^{n,3}\|^2$$

$$\begin{split} &= \sum_{\substack{k \in \mathbb{Z}_{0}^{2} \\ n \leq |k| \leq 2n}} \left(\frac{\|\nabla_{H} B_{t}^{n,3}\|^{2}}{2C_{1,H}^{2} |k|^{\alpha}} + \frac{\|\partial_{3} B_{t}^{n,3}\|^{2}}{C_{2,H}^{2} |k|^{\beta}} \right) + \frac{1}{C_{V}^{2}} \sum_{\substack{k \in \mathbb{Z}_{0}^{3}, \ k_{3} \neq 0 \\ n \leq |k| \leq 2n}} \frac{1}{|k|^{\gamma}} \left\langle \left(I - \frac{k \otimes k}{|k|^{2}}\right) \nabla B_{t}^{n,3}, \nabla B_{t}^{n,3} \right\rangle \\ &\leq \frac{\zeta_{H,\alpha}^{n} \|\nabla_{H} B_{t}^{n,3}\|^{2}}{2C_{1,H}^{2} n^{\alpha-2}} + \frac{\zeta_{H,\beta}^{n} \|\partial_{3} B_{t}^{n,3}\|^{2}}{C_{2,H}^{2} n^{\beta-2}} + \frac{1}{C_{V}^{2}} \sum_{\substack{k \in \mathbb{Z}_{0}^{3} \\ n \leq |k| \leq 2n}} \frac{1}{|k|^{\gamma}} \left\langle \left(I - \frac{k \otimes k}{|k|^{2}}\right) \nabla B_{t}^{n,3}, \nabla B_{t}^{n,3} \right\rangle \\ &= \frac{\zeta_{H,\alpha}^{n} \|\nabla_{H} B_{t}^{n,3}\|^{2}}{2C_{1,H}^{2} n^{\alpha-2}} + \frac{\zeta_{H,\beta}^{n} \|\partial_{3} B_{t}^{n,3}\|^{2}}{3C_{V}^{2} n^{\gamma-3}} \end{split}$$

and by Hölder's inequality

$$\begin{split} \sum_{\substack{k \in \mathbb{Z}_0^3\\j \in \{1,2\}}} \|B_t^n \cdot \nabla \sigma_{k,j}^{n,3}\|^2 &\leq \sum_{\substack{k \in \mathbb{Z}_0^2\\n \leq |k| \leq 2n}} \frac{\|B_t^n\|^2}{C_{2,H}^2 |k|^{\beta-2}} + \sum_{\substack{|k \in \mathbb{Z}_0^3\\n \leq |k| \leq 2n}} \frac{\|B_t^n\|^2}{C_V^2 |k|^{\gamma-2}} \\ &\leq \left(\frac{\zeta_{H,2}^n}{C_{2,H}^2} + \frac{\zeta_{s,\gamma-2}^n}{C_V^2 n^{\gamma-5}}\right) \|B_t^n\|^2. \end{split}$$

Remark 3.3. Note that the simple Hölder estimate in case of Hypothesis 2.6 gives us

$$\sum_{\substack{k \in \mathbb{Z}_0^3 \\ j \in \{1,2\}}} \|\sigma_{k,j}^n \times B_t^n\|^2 \le 2 \sum_{\substack{k \in \mathbb{Z}_0^3 \\ n \le |k| \le 2n}} \frac{1}{|k|^3 C_V^2} \|B_t^n\|^2$$
$$= \frac{2\zeta_{s,3}^n}{C_V^2} \|B_t^n\|^2,$$

which is worse of the second statement of Lemma 3.2 due to the fact that we are neglecting the second term of (3.2) which produce some cancellations.

Now we are ready to prove the main result of this section.

Proposition 3.4. Assuming Hypothesis 2.4 or Hypothesis 2.6 then

(3.3)
$$\mathbb{E}\left[\sup_{t\in[0,T]} \|B_t^n\|_{\mathbf{H}^{-1}}^2\right] + \mathbb{E}\left[\int_0^T \|B_t^n\|^2\right] dt \lesssim \|B_0\|_{\mathbf{H}^{-1}}^2.$$

In particular if Hypothesis 2.5 is satisfied then also

(3.4)
$$\mathbb{E}\left[\sup_{t\in[0,T]}\left\|B_{t}^{n,3}\right\|^{2}\right] + \int_{0}^{T}\mathbb{E}\left[\left\|\nabla B_{t}^{n,3}\right\|^{2}\right]dt \lesssim \left\|B_{0}\right\|_{\mathbf{H}^{-1}}^{2} + \left\|B_{0}^{3}\right\|^{2}.$$

Proof. Since $\nabla \times (\sigma_{k,j}^n \times B_t^n) = \mathcal{L}_{\sigma_{k,j}^n} B_t^n$, the Itô formula for $\|B_t^n\|_{\mathbf{H}^{-1}}^2$ reads as

$$d\|(-\Delta)^{-1/2}B_{t}^{n}\|^{2} + 2\eta\|B_{t}^{n}\|^{2}dt = 2\left\langle(-\Delta)^{-1}\Lambda^{n}B_{t}^{n}, B_{t}^{n}\right\rangle dt + 2\left\langle(-\Delta)^{-1/2}\Lambda_{\rho}^{n}B_{t}^{n}, (-\Delta)^{-1/2}B_{t}^{n}\right\rangle dt + 2\sum_{\substack{k\in\mathbb{Z}_{0}^{3}\\ j\in\{1,2\}}}\|(-\Delta)^{-1/2}\nabla\times(\sigma_{k,j}^{n}\times B_{t}^{n})\|^{2}dt + 2\rho\sum_{\substack{k\in\mathbb{Z}_{0}^{2}\\ k\in\mathbb{Z}_{0}^{2}}}\left\langle\nabla\times(\sigma_{k,1}^{n}\times B_{t}^{n}), (-\Delta)^{-1}\nabla\times(\sigma_{-k,2}^{n}\times B_{t}^{n})\right\rangle dt + 2\rho\sum_{\substack{k\in\mathbb{Z}_{0}^{2}\\ k\in\mathbb{Z}_{0}^{2}}}\left\langle\nabla\times(\sigma_{k,2}^{n}\times B_{t}^{n}), (-\Delta)^{-1}\nabla\times(\sigma_{-k,1}^{n}\times B_{t}^{n})\right\rangle dt + dM_{t}^{n},$$
(3.5)

where

$$M^n_t = 2\sum_{\substack{k\in\mathbb{Z}^3_0\\j\in\{1,2\}}}\int_0^t \langle (-\Delta)^{-1}\nabla\times(\sigma^n_{k,j}\times B^n_s),B^n_s\rangle dW^{k,j}_s.$$

We treat first the case of Hypothesis 2.6 being easier. By the definition of Λ^n in this case we get easily (3.6) $\langle (-\Delta)^{-1}\Lambda^n B_t^n, B_t^n \rangle = -\eta_{iso}^n \|B_t^n\|^2$.

Concerning the Itô correctors we have

$$(3.7) \qquad \begin{aligned} \sum_{\substack{k \in \mathbb{Z}_0^3\\j \in \{1,2\}}} \|(-\Delta)^{-1/2} \nabla \times (\sigma_{k,j}^n \times B_t^n)\|^2 &= \sum_{\substack{k \in \mathbb{Z}_0^3\\j \in \{1,2\}}} \langle (-\Delta)^{-1} \nabla \times \nabla \times (\sigma_{k,j}^n \times B_t^n)], \overline{\sigma_{k,j}^n \times B_t^n} \rangle \\ &= \sum_{\substack{k \in \mathbb{Z}_0^3\\j \in \{1,2\}}} \langle (-\Delta)^{-1} \nabla \times \nabla \times P[Q[\sigma_{k,j}^n \times B_t^n]], \overline{\sigma_{k,j}^n \times B_t^n} \rangle \\ &= \sum_{\substack{k \in \mathbb{Z}_0^3\\j \in \{1,2\}}} \langle P[Q[\sigma_{k,j}^n \times B_t^n]], \overline{P[Q[\sigma_{k,j}^n \times B_t^n]]} \rangle \\ &\leq \sum_{\substack{k \in \mathbb{Z}_0^3\\j \in \{1,2\}}} \|\sigma_{k,j}^n \times B_t^n\|^2 = 2\eta_{iso}^n \|B_t^n\|^2, \end{aligned}$$

due to the properties of the projections P, Q, relation (3.1) and Lemma 3.2. The other terms appearing in (3.5) can be treated easily by Hölder and Young's inequalities. Indeed, again thanks to relation (3.1)

(3.8)
$$\left\langle (-\Delta)^{-1/2} \Lambda_{\rho}^{n} B_{t}^{n}, (-\Delta)^{-1/2} B_{t}^{n} \right\rangle \leq \frac{|\rho| \zeta_{H,2}^{n}}{C_{1,H} C_{2,H}} \|B_{t}^{n}\|_{\mathbf{H}^{-1}} \left\| (-\Delta)^{-1/2} \nabla \times B_{t}^{n,H} \right\| \\ \leq \frac{|\rho| \zeta_{H,2}^{n}}{C_{1,H} C_{2,H}} \|B_{t}^{n}\|_{\mathbf{H}^{-1}} \|B_{t}^{n}\| \\ \leq \frac{(\eta - \eta_{iso})}{2} \|B_{t}^{n}\|^{2} + \frac{|\rho|^{2} \left(\zeta_{H,2}^{n}\right)^{2}}{2C_{1,H}^{2} C_{2,H}^{2} (\eta - \eta_{iso})} \|B_{t}^{n}\|_{\mathbf{H}^{-1}}^{2}$$

since by assumptions $\eta > \eta_{iso}$ and

(3.9)

$$\rho \sum_{k \in \mathbb{Z}_{0}^{2}} \left\langle \nabla \times (\sigma_{k,1}^{n} \times B_{t}^{n}), (-\Delta)^{-1} \nabla \times (\sigma_{-k,2}^{n} \times B_{t}^{n}) \right\rangle \leq |\rho| \sum_{\substack{k \in \mathbb{Z}_{0}^{2} \\ n \leq |k| \leq 2n}} \frac{1}{C_{1,H}C_{2,H}|k|^{\alpha/2+\beta/2}} \|B_{t}^{n}\|^{2}$$

$$\leq |\rho| \frac{\zeta_{H,\alpha/2+\beta/2-2}^{n}}{C_{1,H}C_{2,H}n^{\alpha/2+\beta/2-2}} \|B_{t}^{n}\|^{2}$$

and analogously for the other. Therefore, integrating (3.5) between 0 and t and considering its expected value, thanks to (3.6), (3.7), (3.8), (3.9) we obtain

$$\mathbb{E}\left[\|B_{t}^{n}\|_{\mathbf{H}^{-1}}^{2}\right] + \eta \mathbb{E}\left[\int_{0}^{t}\|B_{s}^{n}\|^{2} ds\right] \leq \|B_{0}^{n}\|_{\mathbf{H}^{-1}}^{2} + \left(\eta_{iso} + O(n^{-1})\right) \mathbb{E}\left[\int_{0}^{t}\|B_{s}^{n}\|^{2} ds\right] \\ + \left(\frac{|\rho|^{2}\left(\zeta_{H,2}\right)^{2}}{C_{1,H}^{2}C_{2,H}^{2}(\eta - \eta_{iso})} + O(n^{-1})\right) \mathbb{E}\left[\int_{0}^{t}\|B_{s}^{n}\|_{\mathbf{H}^{-1}}^{2} ds\right].$$

As a consequence, if n is large enough such that $\eta_{iso} + O(n^{-1}) < \eta$ we obtain, by Grönwall's lemma

(3.10)
$$\sup_{t \in [0,T]} \mathbb{E}\left[\|B_t^n\|_{\mathbf{H}^{-1}}^2 \right] + \mathbb{E}\left[\int_0^t \|B_s^n\|^2 \, ds \right] \lesssim \|B_0^n\|_{\mathbf{H}^{-1}}^2$$

Restarting again from (3.5) and considering the expected value of the supremum in time between [0, T], thanks to (3.6), (3.7),(3.8),(3.9), for n large enough such that $\eta_{iso} + O(n^{-1}) < \eta$ we obtain

$$(3.11) \quad \mathbb{E}\left[\sup_{t\in[0,T]} \|B_t^n\|_{\mathbf{H}^{-1}}^2\right] \le \|B_0^n\|_{\mathbf{H}^{-1}}^2 + \left(\eta_{iso} + O(n^{-1})\right)T\sup_{t\in[0,T]} \mathbb{E}\left[\|B_t^n\|_{\mathbf{H}^{-1}}^2\right] + 2\mathbb{E}\left[\sup_{t\in[0,T]} |M_t^n|\right].$$

The last term can be estimated by Burkholder-Davis-Gundy inequality obtaining

$$\mathbb{E}\left[\sup_{t\in[0,T]}|M_{t}^{n}|\right] \lesssim_{|\rho|} \mathbb{E}\left[\left(\sum_{\substack{k\in\mathbb{Z}_{0}^{3}\\j\in\{1,2\}}}\int_{0}^{T}\langle(-\Delta)^{-1}\nabla\times(\sigma_{k,j}^{n}\times B_{s}^{n}),B_{s}^{n}\rangle^{2}ds\right)^{1/2}\right]$$
$$\leq \mathbb{E}\left[\sup_{t\in[0,T]}\|B_{t}^{n}\|_{\mathbf{H}^{-1}}\left(\sum_{\substack{k\in\mathbb{Z}_{0}^{3}\\j\in\{1,2\}}}|\theta_{k,j}^{n}|^{2}\int_{0}^{T}\|B_{s}^{n}\|^{2}ds\right)^{1/2}\right]$$
$$\leq \frac{1}{2}\mathbb{E}\left[\sup_{t\in[0,T]}\|B_{t}^{n}\|_{\mathbf{H}^{-1}}^{2}\right] + C_{|\rho|,\eta_{iso}}\mathbb{E}\left[\int_{0}^{T}\|B_{s}^{n}\|^{2}ds\right].$$

Combining (3.10),(3.11),(3.12) we get immediately (3.3) in case of Hypothesis 2.6. Let us now consider the case of Hypothesis 2.4. We start by noticing that

$$\Lambda^n = \left(\eta_T^n + \eta_V^n - \frac{\eta_{R,V}^n}{2}\right)\Delta + \left(\eta_R^n - \frac{\eta_{R,V}^n}{2} - \eta_T^n\right)\partial_{33}$$

As a consequence we have

$$\langle (-\Delta)^{-1}\Lambda^n B_t^n, B_t^n \rangle = -\left(\eta_T^n + \eta_V^n - \frac{\eta_{R,V}^2}{2}\right) \|B_t\|^2 + \left(\eta_T^n + \frac{\eta_{R,V}^n}{2} - \eta_R^n\right) \|(-\Delta)^{-1/2}\partial_3 B_t\|^2 \leq -\eta_V^n \|B_t^n\|^2 + \eta_R^n \|B_t^n\|^2 + \eta_{R,V}^n \|B_t^n\|^2.$$

$$(3.13)$$

While, arguing as above, due to Lemma 3.2

(3.14)
$$\sum_{\substack{k \in \mathbb{Z}_0^3 \\ j \in \{1,2\}}} \|(-\Delta)^{-1/2} \nabla \times (\sigma_{k,j}^n \times B_t^n)\|^2 \le (2\eta_T^n + \eta_R^n + 3\eta_V^n) \|B_t^n\|^2$$

Estimate (3.8) and (3.9) continue to hold, possibly replacing η_{iso} with $2\eta_T$. In conclusion, combining (3.5),(3.13), (3.14),(3.8) and (3.9) we obtain

$$\mathbb{E}\left[\|B_{t}^{n}\|_{\mathbf{H}^{-1}}^{2}\right] + \eta \mathbb{E}\left[\int_{0}^{t}\|B_{s}^{n}\|^{2} ds\right] \leq \|B_{0}^{n}\|_{\mathbf{H}^{-1}}^{2} + \left(4\eta_{T}^{n} - 2\eta_{T} + O(n^{2-\beta} + n^{3-\gamma})\right) \mathbb{E}\left[\int_{0}^{t}\|B_{s}^{n}\|^{2} ds\right] + \left(\frac{|\rho|^{2} \left(\zeta_{H,2}\right)^{2}}{C_{1,H}^{2}C_{2,H}^{2}(\eta - 2\eta_{T})} + O(n^{-1})\right) \mathbb{E}\left[\int_{0}^{t}\|B_{s}^{n}\|_{\mathbf{H}^{-1}}^{2} ds\right].$$
(3.15)

Arguing as above, by Grönwall's lemma and Burkholder-Davis-Gundy inequality we can prove (3.3) starting from (3.15). We omit the easy details. We are left to show the validity of the additional estimate in case of Hypothesis 2.5. Due to Remark 2.15 and (2.7), the Itô formula for $||B_t^{n,3}||^2$ reads

$$\begin{aligned} d\|B_{t}^{n,3}\|^{2} + 2\left(\eta + \eta_{V}^{n} - \frac{\eta_{R,V}^{n}}{2}\right)\|\nabla B_{t}^{n,3}\|^{2}dt + 2\eta_{T}^{n}\|\nabla_{H}B_{t}^{n,3}\|^{2}dt + 2\left(\eta_{R}^{n} - \frac{\eta_{RV}^{n}}{2}\right)\|\partial_{3}B_{t}^{n,3}\|^{2}dt \\ &= \frac{2\rho\zeta_{H,2}^{n}}{C_{1,H}C_{2,H}}\left\langle (B_{t}^{n,H})^{\perp}, \nabla_{H}B_{t}^{n,3}\right\rangle dt + 2\sum_{\substack{k\in\mathbb{Z}_{0}^{n}\\ j\in\{1,2\}}} \left(\|\sigma_{k,j}^{n}\cdot\nabla B_{t}^{n,3}\|^{2} + \|B_{t}^{n}\cdot\nabla\sigma_{k,j}^{n,3}\|^{2}\right)dt \\ (3.16) \qquad + 2\rho\sum_{\substack{k\in\mathbb{Z}_{0}^{n}\\ n\leq|k|\leq 2n}} \left(\left\langle \sigma_{k,1}^{n}\cdot\nabla_{H}B_{t}^{n,3}, B_{t}^{n,H}\cdot\nabla\sigma_{-k,2}^{n,3}\right\rangle + \left\langle B_{t}^{n,H}\cdot\nabla\sigma_{k,2}^{n,3}, \sigma_{-k,1}^{n}\cdot\nabla_{H}B_{t}^{n,3}\right\rangle\right)dt + dM_{t}^{n,3}, \end{aligned}$$

where

$$M_t^{n,3} = 2 \sum_{\substack{k \in \mathbb{Z}_0^3 \\ j \in \{1,2\}}} \int_0^t \left\langle \sigma_{k,j}^n \cdot \nabla B_s^3 - B_s \cdot \nabla \sigma_{k,j}^{n,3}, B_s^{n,3} \right\rangle dW_s^{k,j}$$

We can treat the different terms in the right hand side of (3.16) similarly to previous case. Thanks to Lemma 3.2 we obtain

$$\sum_{\substack{k \in \mathbb{Z}_0^n \\ j \in \{1,2\}}} \left(\|\sigma_{k,j}^n \cdot \nabla B_t^{n,3}\|^2 + \|B_t^n \cdot \nabla \sigma_{k,j}^{n,3}\|^2 \right) \le \eta_T^n \|\nabla_H B_t^{n,3}\|^2 + O(n^{-2}) \|\nabla B_t^{n,3}\|^2 + \left(\frac{\zeta_{H,2}^n}{C_{2,H}^2} + \frac{\zeta_{s,\gamma-2}^n}{C_V^2 n^{\gamma-5}}\right) \|B_t^n\|^2 + O(n^{-2}) \|\nabla B_t^{n,3}\|^2 +$$

The term associated to the correlation of our noise can be treated easily by Holder and Young's inequality

(3.18)
$$\frac{\rho \zeta_{H,2}^{n}}{C_{1,H}C_{2,H}} \left\langle (B_{t}^{n,H})^{\perp}, \nabla_{H}B_{t}^{n,3} \right\rangle \leq \frac{|\rho|\zeta_{H,2}^{n}}{C_{1,H}C_{2,H}} \|B_{t}^{n,H}\| \|\nabla_{H}B_{t}^{n,3}\| \leq \frac{\eta}{2} \|\nabla B^{n,3}\|^{2} + \frac{|\rho|^{2} \left(\zeta_{H,2}^{n}\right)^{2}}{2C_{1,H}^{2}C_{2,H}^{2} \eta} \|B_{t}^{n}\|^{2},$$

(3.19)

$$\rho \sum_{\substack{k \in \mathbb{Z}_{0}^{2} \\ n \leq |k| \leq 2n}} \left(\left\langle \sigma_{k,1}^{n} \cdot \nabla_{H} B_{t}^{n,3}, B_{t}^{n,H} \cdot \nabla \sigma_{-k,2}^{n,3} \right\rangle \right) \leq |\rho| \frac{\zeta_{H,3}^{n} \|\nabla_{H} B_{t}^{n,3}\| \|B_{t}^{n}\|}{C_{1,H} C_{2,H} n} \\ \leq \frac{\|\nabla B_{t}^{n,3}\|^{2}}{2n} + \frac{|\rho|^{2} \left(\zeta_{H,3}^{n}\right)^{2} \|B_{t}^{n}\|^{2}}{2C_{1,H}^{2} C_{2,H}^{2} n}$$

and analogously for the other. Therefore, integrating (3.16) between 0 and t and considering its expected value, thanks to (3.17), (3.18),(3.19) we obtain

$$\mathbb{E}\left[\|B_t^{n,3}\|^2\right] + \left(\eta - \frac{2}{n} + O(n^{-2})\right) \mathbb{E}\left[\int_0^t \|\nabla B_s^{n,3}\|^2 ds\right] \\ \leq \left\|B_0^{n,3}\right\|^2 + 2\left(\frac{\zeta_{H,2}^n}{C_{2,H}^2} + \frac{\zeta_{s,\gamma-2}^n}{C_V^2 n^{\gamma-5}} + \frac{|\rho|^2 \left(\zeta_{H,2}^n\right)^2}{2C_{1,H}^2 C_{2,H}^2 \eta} + O(n^{-1})\right) \mathbb{E}\left[\int_0^T \|B_s^n\|^2 ds\right]$$

Therefore, if n is large enough such that $\eta - \frac{2}{n} + O(n^{-2}) > \frac{\eta}{2}$, thanks to (3.3) we have immediately

(3.20)
$$\sup_{t \in [0,T]} \mathbb{E}\left[\|B_t^{n,3}\|^2 \right] + \mathbb{E}\left[\int_0^T \|\nabla B_s^{n,3}\|^2 ds \right] \lesssim \left\| B_0^{n,3} \right\|^2 + \left\| B_0^n \right\|_{\mathbf{H}^{-1}}^2.$$

Restarting again from (3.16) and considering the expected value of the supremum in time between [0, T], thanks to (3.17), (3.18),(3.19), we obtain for n large enough such that $\eta - \frac{2}{n} + O(n^{-2}) > \frac{\eta}{2}$

$$\mathbb{E}\left[\sup_{t\in[0,T]}\|B_{t}^{n,3}\|^{2}\right] \leq \left\|B_{0}^{n,3}\right\|^{2} + 2\left(\frac{\zeta_{H,2}^{n}}{C_{2,H}^{2}} + \frac{\zeta_{s,\gamma-2}^{n}}{C_{V}^{2}n^{\gamma-5}} + \frac{|\rho|^{2}\left(\zeta_{H,2}^{n}\right)^{2}}{2C_{1,H}^{2}C_{2,H}^{2}\eta} + O(n^{-1})\right)\mathbb{E}\left[\int_{0}^{T}\|B_{s}^{n}\|^{2}ds\right] + \mathbb{E}\left[\sup_{t\in[0,T]}|M_{t}^{n,3}|\right].$$
(3.21)
$$+\mathbb{E}\left[\sup_{t\in[0,T]}|M_{t}^{n,3}|\right].$$

The first term can be treated easily thanks to (3.3). The second one is treated via Burkholder-Davis-Gundy inequality, (2.7), Young's inequality and (3.17) obtaining

$$\mathbb{E}\left[\sup_{t\in[0,T]}|M_{t}^{n,3}|\right] \lesssim_{|\rho|} \mathbb{E}\left[\left(\sum_{\substack{k\in\mathbb{Z}_{0}^{3}\\j\in\{1,2\}}}\int_{0}^{T}\left\langle\sigma_{k,j}^{n}\cdot\nabla B_{s}^{3}-B_{s}\cdot\nabla\sigma_{k,j}^{n,3},B_{s}^{n,3}\right\rangle^{2}ds\right)^{1/2}\right]$$
$$\leq \mathbb{E}\left[\sup_{t\in[0,T]}\left\|B_{s}^{n,3}\right\|\left(\sum_{\substack{k\in\mathbb{Z}_{0}^{3}\\j\in\{1,2\}}}\int_{0}^{T}\left\|\sigma_{k,j}^{n}\cdot\nabla B_{s}^{3}\right\|+\left\|B_{s}\cdot\nabla\sigma_{k,j}^{n,3}\right\|^{2}ds\right)^{1/2}\right]$$

$$(3.22) \qquad \leq \frac{1}{2} \mathbb{E} \left[\sup_{t \in [0,T]} \|B_t^{n,3}\| \right] + C_{|\rho|,C_{1,H},C_{2,H},C_V,\gamma} \left(\mathbb{E} \left[\int_0^T \|\nabla B_s^{n,3}\|^2 ds \right] + \mathbb{E} \left[\int_0^T \|B_s^n\|^2 ds \right] \right).$$
Combining (3.21) (3.20) (3.22) and (3.3) we obtain relation (3.4)

Combining (3.21), (3.20), (3.22) and (3.3) we obtain relation (3.4).

4. Proof of Theorem 2.17

Before exploiting the convergence properties of B^n , we start with a well-posedness result for our limit object \overline{B} solution of (2.21).

Definition 4.1. We say that \overline{B} is a weak solution of equation (2.21) if

$$\overline{B} \in C(0,T;\mathbf{H}^{-1}) \cap L^2(0,T;\mathbf{L}^2)$$

and for each $\phi \in \mathbf{H}^2$, and all $t \in [0,T]$, one has

$$\begin{split} \langle \overline{B}_t, \phi \rangle - \langle \overline{B}_0, \phi \rangle &= \eta \int_0^t \langle \overline{B}_s, \Delta \phi \rangle \, ds + \int_0^t \langle \overline{B}_s, \Lambda_{\alpha, \beta, \gamma} \phi \rangle \, ds \\ &+ \int_0^t \langle \mathcal{A}_\rho \overline{B}_s, \nabla \times \phi \rangle \, ds. \end{split}$$

Since the operator $\eta \Delta + \Lambda_{\alpha,\beta,\gamma}$ is uniformly elliptic and equation (4.1) is linear, by standard theory of parabolic PDE we have existence and uniqueness of equation (4.1) in the sense Definition 4.1, see for example [20, Chapter 3], [8, Chapter 5].

Theorem 4.2. For each $B_0 \in \mathbf{H}^{-1}$, there exists a unique solution of (2.21) in the sense of Definition 4.1. Moreover if $B_0 \in \mathbf{L}^2$, then

$$\overline{B} \in C(0,T;\mathbf{L}^2) \cap L^2(0,T;\mathbf{H}^1)$$

Remark 4.3. Arguing as in Remark 2.15 we obtain easily that, given $\overline{B_0}$ in \mathbf{L}^2 , then $\overline{B_t^3}$ satisfies

$$\langle \overline{B_t^3}, \varphi \rangle - \langle \overline{B_0^3}, \varphi \rangle = \eta \int_0^t \langle \overline{B_s^3}, \Delta \varphi \rangle \, ds + \int_0^t \langle \overline{B_s^3}, \Lambda_{\alpha, \beta, \gamma} \varphi \rangle \, ds + \frac{\rho \zeta_{H, 2}}{C_{1, H} C_{2, H}} \int_0^t \langle \overline{B_s^H}^\perp, \nabla_H \varphi \rangle \, ds$$

for every $t \in [0, T]$ and $\varphi \in \dot{H}^2$.

Now we are ready to provide the proof of our main theorem. First we start with some notation. Let us denote by S^n , S the semigroups generated by $\eta \Delta + \Lambda^n$ and $\eta \Delta + \Lambda_{\alpha,\beta,\gamma}$. Since their infinitesimal generators have the form of $\mu_n(\partial_{11} + \partial_{22}) + \nu_n \partial_{33}$, $\mu_n \ge \nu_n > 0$ we are in the framework of subsubsection 2.2.2. In particular, we recall that $\mu_n, \nu_n \ge \eta$ for each $n \in \mathbb{N}$. Secondly, we denote by

$$\begin{split} Z_t^n &= \sum_{\substack{k \in \mathbb{Z}_0^n \\ j \in \{1,2\}}} \int_0^t S^n(t-s) \nabla \times (\sigma_{k,j}^n \times B_s^n) dW_s^{k,j}, \\ H_t^n &= \sum_{\substack{k \in \mathbb{Z}_0^n \\ j \in \{1,2\}}} \int_0^t S^n(t-s) \left(\sigma_{k,j}^n \cdot \nabla B_t^{n,3} - B_t^n \cdot \nabla \sigma_{k,j}^{n,3}\right) dW_s^{k,j}. \end{split}$$

The stochastic integrals above are well defined thanks to the regularity properties of B_t^n and the fact that for each n only a finite number of $\sigma_{k,j}^n \neq 0$. The first step in order to prove Theorem 2.17 is to rewrite B_t^n , $B_t^{n,3}$ in mild form. Indeed, recalling the definition of \mathcal{A}^n_{ρ} , see Lemma 2.13, the following lemma holds true.

Lemma 4.4. B_t^n and $B_t^{n,3}$ satisfy the mild formula

(4.1)
$$B_t^n = S^n(t)B_0^n + \int_0^t S^n(t-s)\nabla \times (\mathcal{A}_\rho^n B_s^n)ds + Z_t^n,$$

(4.2)
$$B_t^{n,3} = S^n(t)B_0^{n,3} - \frac{\rho\zeta_{H,2}^n}{C_{1,H}C_{2,H}} \int_0^t S^n(t-s)\operatorname{div}_H\left((B_s^{n,H})^{\perp}\right) ds + H_t^n,$$

the former seen as an equality in \mathbf{H}^{-1} and the latter in \dot{L}^2 .

Proof. Let us take $\phi = \psi_{h,l}$, $\psi_{h,l} = a_{h,l}e^{ih\cdot x}$ as test function in (2.20) obtaining

$$(4.3) \qquad \langle B_t^n, \psi_{h,l} \rangle = \langle B_0^n, \psi_{h,l} \rangle - \left(\mu_n \left(h_1^2 + h_2^2 \right) + \nu_n h_3^2 \right) \int_0^t \langle B_s^n, \psi_{h,l} \rangle \, ds + \int_0^t \langle \nabla \times (\mathcal{A}_\rho^n B_s^n), \psi_{h,l} \rangle \, ds + \sum_{\substack{k \in \mathbb{Z}_0^n \\ j \in \{1,2\}}} \int_0^t \langle \nabla \times (\sigma_{k,j}^n \times B_s^n), \psi_{h,l} \rangle \, dW_s^{k,j}$$

Therefore, applying Itô formula to $e^{t(\mu_n(h_1^2+h_2^2)+\nu_nh_3^2)}\langle B_t^n,\psi_{h,l}\rangle$ we have

$$\langle B_{t}^{n}, \psi_{h,l} \rangle = e^{-t\left(\mu_{n}\left(h_{1}^{2}+h_{2}^{2}\right)+\nu_{n}h_{3}^{2}\right)} \langle B_{0}^{n}, \psi_{h,l} \rangle + \int_{0}^{t} e^{-(t-s)\left(\mu_{n}\left(h_{1}^{2}+h_{2}^{2}\right)+\nu_{n}h_{3}^{2}\right)} \langle \nabla \times \left(\mathcal{A}_{\rho}^{n}B_{s}^{n}\right), \psi_{h,l} \rangle ds$$

$$+ \sum_{\substack{k \in \mathbb{Z}_{0}^{3} \\ j \in \{1,2\}}} \int_{0}^{t} e^{-(t-s)\left(\mu_{n}\left(h_{1}^{2}+h_{2}^{2}\right)+\nu_{n}h_{3}^{2}\right)} \langle \nabla \times \left(\sigma_{k,j}^{n} \times B_{s}^{n}\right), \psi_{h,l} \rangle dW_{s}^{k,j} \quad \mathbb{P}-a.s. \quad \forall t \in [0,T]$$

We can find $\Gamma \subseteq \Omega$ of full probability such that the above equality holds for all $t \in [0, T]$ and all $h \in \mathbb{Z}_0^3$, $l \in \{1, 2\}$. But this is exactly (4.1) written in Fourier modes. The proof of (4.2) is analogous, just replacing relation (2.20) with Remark 2.15 and choosing as test function $\phi = (0, 0, \psi_h)^t$, $\psi_h = e^{ih \cdot x}$, $h \in \mathbb{Z}_0^3$. We omit the easy details. \Box

Secondly we need to study some properties of the stochastic convolutions Z_t^n, H_t^n . Lemma 4.6 and Lemma 4.7 below are the analogous of [10, Lemma 2.5] in our framework. In particular it is important to point out that [10, Assumption 2.4] is false in our case and we have to deal also with the stochastic stretching, which was neglected in previous results. In order to reach our plan, we start with a uniform bound on our coefficients in a distributional norm.

Lemma 4.5. Assuming either Hypothesis 2.4 or Hypothesis 2.6, for each $\delta > 0$

(4.5)
$$\mathbb{E}\left[\sup_{\substack{t\in[0,T]\\j\in\{1,2\}}}\sum_{\substack{k\in\mathbb{Z}_0^3\\j\in\{1,2\}}} \|P[Q[\sigma_{k,j}^n\times B_t^n]]\|_{\mathbf{H}^{-1-\frac{\delta}{2}}}^2\right] \lesssim \|B_0^n\|_{\mathbf{H}^{-1}}^2$$

Proof. Arguing as in Lemma 3.2, if we denote by

$$b_t^{n,h,l} := \langle B_t^n, a_{h,l} e^{-ih \cdot x} \rangle,$$

then

$$\sigma_{k,j}^n \times B_t^n = \theta_{k,j}^n \sum_{\substack{h \in \mathbb{Z}_0^3, \\ l \in \{1,2\}}} b_t^{n,h,l} a_{k,j} \times a_{h,l} e^{i(h+k) \cdot x}.$$

Thanks to Proposition 3.4 we already know that

(4.6)
$$\mathbb{E}\left[\sup_{\substack{t\in[0,T]\\ l\in\{1,2\}}} \sum_{\substack{h\in\mathbb{Z}_{0}^{n},\\ l\in\{1,2\}}} \frac{|b_{t}^{n,h,l}|^{2}}{|h|^{2}}\right] \lesssim \|B_{0}^{n}\|_{\mathbf{H}^{-1}}^{2}.$$

For each $k \in \mathbb{Z}_0^3$ we have

$$\left\| P[Q[\sigma_{k,j}^n \times B_t^n]] \right\|_{\mathbf{H}^{-1-\frac{\delta}{2}}}^2 \lesssim \sum_{\substack{h \in \mathbb{Z}_0^3 \\ h \neq -k, \ l \in \{1,2\}}} \frac{|\theta_{k,j}^n|^2}{|h+k|^{2+\delta}} |b_t^{n,h,l}|^2.$$

We are left to study

$$(4.7) \qquad \mathbb{E}\left[\sup_{\substack{k\in\mathbb{Z}_{0}^{3}\\n\leq|k|\leq 2n}}\sum_{\substack{h\in\mathbb{Z}_{0}^{3}\\k\neq|k|\leq 2n}}\sum_{\substack{h\in\mathbb{Z}_{0}^{3}\\k\neq-k,\ l\in\{1,2\}}}\frac{|\theta_{k,j}^{n}b_{t}^{n,h,l}|^{2}}{|h+k|^{2+\delta}}\right] = \mathbb{E}\left[\sup_{\substack{h\in\mathbb{Z}_{0}^{3}\\l\in\{1,2\}}}\frac{|b_{t}^{n,h,l}|^{2}}{|h|^{2}}\sum_{\substack{k\in\mathbb{Z}_{0}^{3}\\k\neq-k,\ n\leq|k|\leq 2n\\j\in\{1,2\}}}\frac{|\theta_{k,j}^{n}h|^{2}}{|h+k|^{2+\delta}}\right].$$

Due to relation (4.6), it is enough to show that

$$\sum_{\substack{k \in \mathbb{Z}_0^n \\ h \neq -k, \\ n \le |k| \le 2n \\ j \in \{1,2\}}} \frac{|\theta_{k,j}^n|^2 |h|^2}{|h+k|^{2+\delta}} \lesssim 1.$$

uniformly in $h \in \mathbb{Z}_0^3$, $l \in \{1, 2\}$. By splitting the summation domain, we have

$$(4.8) \qquad \qquad \sum_{\substack{k \in \mathbb{Z}_{0}^{n} \\ h \neq -k, \\ n \leq |k| \leq 2n \\ j \in \{1,2\}}} \frac{|\theta_{k,j}^{n}|^{2}|h|^{2}}{|h+k|^{2+\delta}} \leq \sum_{\substack{k \in \mathbb{Z}_{0}^{n} \\ 1 \leq |k| \leq \frac{|h|}{2}, \\ n \leq |k| \leq 2n \\ j \in \{1,2\}}} \frac{|\theta_{k,j}^{n}|^{2}|h|^{2}}{|h+k|^{2+\delta}} + \sum_{\substack{k \in \mathbb{Z}_{0}^{n} \\ n \leq |k| \leq 2n \\ j \in \{1,2\}}} \frac{|\theta_{k,j}^{n}|^{2}|h|^{2}}{|h+k|^{2+\delta}} + \sum_{\substack{k \in \mathbb{Z}_{0}^{n} \\ n \leq |k| \leq 2n \\ j \in \{1,2\}}} \frac{|\theta_{k,j}^{n}|^{2}|h|^{2}}{|h+k|^{2+\delta}} + \sum_{\substack{k \in \mathbb{Z}_{0}^{n} \\ n \leq |k| \leq 2n \\ j \in \{1,2\}}} \frac{|\theta_{k,j}^{n}|^{2}|h|^{2}}{|h+k|^{2+\delta}} + \sum_{\substack{k \in \mathbb{Z}_{0}^{n} \\ |h+k| \geq \frac{|h|}{2}, \\ |h+k| \geq \frac{|h|}{2}, \\ |h+k| \geq \frac{|h|}{2}, \\ n \leq |k| \leq 2n \\ j \in \{1,2\}}} \frac{|\theta_{k,j}^{n}|^{2}|h|^{2}}{|h+k|^{2+\delta}} + \sum_{\substack{k \in \mathbb{Z}_{0}^{n} \\ |h+k| \geq \frac{|h|}{2}, \\ |h+k| \geq 2n \\ j \in \{1,2\}} \end{cases} |\theta_{k,j}^{n}|^{2} + \sum_{\substack{k \in \mathbb{Z}_{0}^{n} \\ n \leq |k| \leq 2n \\ j \in \{1,2\}}} \frac{|\theta_{k,j}^{n}|^{2}|h|^{2}}{|h+k|^{2+\delta}} + \sum_{\substack{k \in \mathbb{Z}_{0}^{n} \\ |h+k| \geq \frac{|h|}{2}, \\ |h+k| \geq \frac{|h|}{2}, \\ |h+k| \geq 2n \\ j \in \{1,2\}}} |\theta_{k,j}^{n}|^{2} + \sum_{\substack{k \in \mathbb{Z}_{0}^{n} \\ n \leq |k| \leq 2n \\ j \in \{1,2\}}} \frac{|\theta_{k,j}^{n}|^{2}|h|^{2}}{|h+k|^{2+\delta}} + \sum_{\substack{k \in \mathbb{Z}_{0}^{n} \\ |h+k| \geq \frac{|h|}{2}, \\ |h+k| \geq \frac{|h|}{2}, \\ |h+k| \geq 2n \\ j \in \{1,2\}}} |\theta_{k,j}^{n}|^{2} + \sum_{\substack{k \in \mathbb{Z}_{0}^{n} \\ |h+k| \geq \frac{|h|}{2}, \\ |h+k| \geq \frac$$

We start considering the case of Hypothesis 2.6 which is the easier one. In this case, due to (4.8)

$$\begin{split} \sum_{\substack{k \in \mathbb{Z}_0^3 \\ h \neq -k, \\ n \leq |k| \leq 2n \\ j \in \{1,2\}}} \frac{|\theta_{k,j}^n|^2 |h|^2}{|h+k|^{2+\delta}} &\lesssim \sum_{\substack{k \in \mathbb{Z}_0^3 \\ n \leq |k| \leq 2n}} \frac{1}{|k|^3} + \sum_{\substack{k \in \mathbb{Z}_0^3 \\ 1 \leq |h+k| \leq \frac{|h|}{2}, \\ n \leq |k| \leq 2n}} \frac{1}{|k|^{1/2}} \\ &\lesssim 1 + \left(\sum_{\substack{k \in \mathbb{Z}_0^3 \\ n \leq |k| \leq 2n}} \frac{1}{|k|^3}\right)^{1/3} \left(\sum_{\substack{k \in \mathbb{Z}_0^3 \\ 1 \leq |h+k| \leq \frac{|h|}{2}}} \frac{1}{|h+k|^{3+\frac{3}{2}\delta}}\right)^{2/3} \lesssim_{\delta} 1, \end{split}$$

where the first inequality follows from the fact that $|k| \ge |(k+h) - h| \ge \frac{|h|}{2}$ if $1 \le |h+k| \le \frac{|h|}{2}$ and the second by Hölder's inequality. This completes the proof of (4.5) in case of Hypothesis 2.6. In case of Hypothesis 2.4 we argue similarly. Due to the summability properties of the $\theta_{k,j}^n$, it is enough to consider the second term in the right hand side of (4.8), i.e.

(4.9)

n

$$\sum_{\substack{k \in \mathbb{Z}_0^3 \\ 1 \le |h+k| \le \frac{|h|}{2}, \\ n \le |k| \le 2n \\ j \in \{1,2\}}} \frac{|\theta_{k,j}^n|^2 |h|^2}{|h+k|^{2+\delta}} = \sum_{\substack{k \in \mathbb{Z}_0^2 \\ 1 \le |h+k| \le \frac{|h|}{2}, \\ n \le |k| \le 2n}} \frac{|h|^2}{|k|^\alpha |h+k|^{2+\delta}} + \sum_{\substack{k \in \mathbb{Z}_0^2 \\ 1 \le |h+k| \le \frac{|h|}{2}, \\ n \le |k| \le 2n}} \frac{|h|^2}{|k|^\beta |h+k|^{2+\delta}} + \sum_{\substack{k \in \mathbb{Z}_0^3 \\ 1 \le |h+k| \le \frac{|h|}{2}, \\ n \le |k| \le 2n}} \frac{|h|^2}{|k|^\gamma |h+k|^{2+\delta}}.$$

Since $\gamma > 3$ the third term is a simplified version of the one treated before. We are left to study the other two. It is clear that if we are able to control $\sum_{\substack{k \in \mathbb{Z}_0^n \\ 1 \le |h+k| \le \frac{|h|}{2}}} \frac{|h|^2}{|k|^2 |h+k|^{2+\delta}}$ we can control relation (4.9). Therefore we study $n \leq |k| \leq 2n$

only this case. We observe that by triangular inequality, the constraints $1 \le |h+k| \le \frac{|h|}{2}$, $n \le |k| \le 2n$ imply

$$|h| = |(h+k) + (-k)| \le |k| + \frac{|h|}{2} \le 2n + \frac{|h|}{2}.$$

As a consequence

$$\sum_{\substack{k \in \mathbb{Z}_0^2 \\ 1 \le |h+k| \le \frac{|h|^2}{2n}}} \frac{|h|^2}{|k|^2 |h+k|^{2+\delta}} \lesssim \sum_{\substack{z \in h + \mathbb{Z}_0^2 \\ 1 \le |z| \le 2n}} \frac{1}{|z|^{2+\delta}} \lesssim_{\delta} 1.$$

This concludes the proof of relation (4.5).

Lemma 4.5 was a necessary toolds in order to prove the following estimates on Z_t^n , H_t^n .

Lemma 4.6. For each $\delta > 0$

(4.10)
$$\sup_{t\in[0,T]} \mathbb{E}\left[\left\| Z_t^n \right\|_{\mathbf{H}^{-1-\delta}}^2 \right] \lesssim \left\| B_0^n \right\|_{\mathbf{H}^{-1}}^2$$

In case of Hypothesis 2.5 we have also

(4.11)
$$\sup_{t \in [0,T]} \mathbb{E}\left[\|H_t^n\|_{\dot{H}^{-\delta}}^2 \right] \lesssim \|B_0^n\|_{\mathbf{H}^{-1}}^2 + \|B_0^{n,3}\|^2.$$

Proof. By Burkholder-Davis-Gundy inequality and the regularization properties of the semigroup, see Lemma 2.2, we have

$$\begin{split} \mathbb{E}\left[\|Z_t^n\|_{\mathbf{H}^{-1-\delta}}^2\right] \lesssim_{|\rho|} \sum_{\substack{k \in \mathbb{Z}_0^n \\ j \in \{1,2\}}} \mathbb{E}\left[\int_0^t \|S^n(t-s)\nabla \times (\sigma_{k,j}^n \times B_s^n)\|_{\mathbf{H}^{-1-\delta}}^2 \, ds\right] \\ \lesssim_{\eta} \mathbb{E}\left[\int_0^t \frac{1}{(t-s)^{1-\frac{\delta}{2}}} \left\|\nabla \times (\sigma_{k,j}^n \times B_s^n)\right\|_{\mathbf{H}^{-2-\frac{\delta}{2}}}^2 \, ds\right] \\ \lesssim \mathbb{E}\left[\int_0^t \frac{1}{(t-s)^{1-\frac{\delta}{2}}} \left\|P[Q[\sigma_{k,j}^n \times B_s^n]]\right\|_{\mathbf{H}^{-1-\frac{\delta}{2}}}^2 \, ds\right] \\ \lesssim_{\delta} \|B_0^n\|_{\mathbf{H}^{-1}}^2 \end{split}$$

where the last inequality follows from (4.5) and is uniform for $t \in [0, T]$. Assuming Hypothesis 2.5, similarly we can prove (4.11). Again, by Burkholder-Davis-Gundy inequality and the regularization properties of the semigroup, Lemma 2.2, we get

$$\begin{split} \mathbb{E}\left[\|H_{t}^{n}\|_{\dot{H}^{-\delta}}^{2}\right] \lesssim_{|\rho|} \sum_{\substack{k \in \mathbb{Z}_{0}^{3} \\ j \in \{1,2\}}} \mathbb{E}\left[\int_{0}^{t} \left\|S^{n}(t-s)\left[\sigma_{k,j}^{n} \cdot \nabla B_{s}^{n,3}\right] - S^{n}(t-s)\left[B_{s}^{n} \cdot \nabla \sigma_{k,j}^{n,3}\right]\right\|_{\dot{H}^{-\delta}}^{2} ds\right] \\ \lesssim_{\eta} \sum_{\substack{k \in \mathbb{Z}_{0}^{3} \\ j \in \{1,2\}}} \mathbb{E}\left[\int_{0}^{t} \frac{1}{(t-s)^{1-\delta}} \left\|\sigma_{k,j}^{n} \cdot \nabla B_{s}^{n,3}\right\|_{\dot{H}^{-1}}^{2} ds\right] + \sum_{\substack{k \in \mathbb{Z}_{0}^{3} \\ j \in \{1,2\}}} \mathbb{E}\left[\int_{0}^{t} \left\|B_{s}^{n} \cdot \nabla \sigma_{k,j}^{n,3}\right\|^{2} ds\right] \\ \lesssim \sum_{\substack{k \in \mathbb{Z}_{0}^{3} \\ j \in \{1,2\}}} \mathbb{E}\left[\int_{0}^{t} \frac{1}{(t-s)^{1-\delta}} \left\|\sigma_{k,j}^{n} B_{s}^{n,3}\right\|^{2} ds\right] + \sum_{\substack{k \in \mathbb{Z}_{0}^{3} \\ j \in \{1,2\}}} \mathbb{E}\left[\int_{0}^{t} \left\|B_{s}^{n} \cdot \nabla \sigma_{k,j}^{n,3}\right\|^{2} ds\right]. \end{split}$$

Both the two terms can be treated easily. Indeed,

$$\sum_{\substack{k \in \mathbb{Z}_0^n \\ j \in \{1,2\}}} \mathbb{E}\left[\int_0^t \frac{1}{(t-s)^{1-\delta}} \left\|\sigma_{k,j}^n B_s^{n,3}\right\|^2 ds\right] \le \sum_{\substack{k \in \mathbb{Z}_0^n \\ j \in \{1,2\}}} |\theta_{k,j}^n|^2 \mathbb{E}\left[\int_0^t \frac{1}{(t-s)^{1-\delta}} \sup_{s \in [0,T]} \left\|B_s^{n,3}\right\|^2 ds\right] \le \delta_{\delta} \|B_0^n\|_{\mathbf{H}^{-1}}^2 + \|B_0^{n,3}\|^2.$$

23

due to Proposition 3.4. Similarly

$$\sum_{\substack{k \in \mathbb{Z}_0^3\\ j \in \{1,2\}}} \mathbb{E}\left[\int_0^t \left\| B_s^n \cdot \nabla \sigma_{k,j}^{n,3} \right\|^2 ds\right] \le \left(\sum_{\substack{k \in \mathbb{Z}_0^2\\ n \le |k| \le 2n}} \frac{1}{|k|^{\beta-2}} + \sum_{\substack{k \in \mathbb{Z}_0^3\\ n \le |k| \le 2n}} \frac{1}{|k|^{\gamma-2}}\right) \mathbb{E}\left[\int_0^T \|B_s^n\|^2 ds\right] \le \left\|B_0^n\|_{\mathbf{H}^{-1}}^2,$$

again by Proposition 3.4.

Lemma 4.7. Assuming either Hypothesis 2.4, Hypothesis 2.6 then for each $\delta > 0$

(4.12)
$$\sup_{t \in [0,T]} \mathbb{E}\left[\left\| Z_t^n \right\|_{\mathbf{H}^{-4-\frac{3}{2}\delta}}^2 \right] \lesssim \frac{\left\| B_0^n \right\|_{\mathbf{H}^{-1}}^2}{n^{\alpha \land \beta \land \gamma}}$$

In case of Hypothesis 2.5 we have also

(4.13)
$$\sup_{t \in [0,T]} \mathbb{E}\left[\left\| H_t^n \right\|_{\dot{H}^{-\frac{3}{2} - \frac{3}{2}\delta}}^2 \right] \lesssim \frac{\left\| B_0^{n,3} \right\|^2 + \left\| B_0^n \right\|_{\mathbf{H}^{-1}}^2}{n^2}$$

Proof. Let us consider for a smooth divergence free vector field ψ

$$\mathbb{E}\left[\langle Z_t^n,\psi\rangle^2\right] = \mathbb{E}\left[\left(\sum_{\substack{k\in\mathbb{Z}_0^3\\j\in\{1,2\}}}\int_0^t \langle S^n(t-s)\nabla\times(\sigma_{k,j}^n\times B_s^n),\psi\rangle dW_s^{k,j}\right)^2\right].$$

Therefore, by Burkholder-Davis-Gundy inequality, Sobolev embedding theorem and the contractivity property of our semigroup we obtain

$$\begin{split} \mathbb{E}\left[\langle Z_t^n,\psi\rangle^2\right] \lesssim_{|\rho|} \mathbb{E}\left[\sum_{\substack{k\in\mathbb{Z}_0^3\\j\in\{1,2\}}} \int_0^t \langle S^n(t-s)\nabla\times(\sigma_{k,j}^n\times B_s^n),\psi\rangle^2 ds\right] \\ &= \mathbb{E}\left[\sum_{\substack{k\in\mathbb{Z}_0^3\\j\in\{1,2\}}} |\theta_{k,j}^n|^2 \int_0^t \langle a_{k,j}e^{ik\cdot x}, B_s^n\times(\nabla\times S^n(t-s)\psi)\rangle^2 ds\right] \\ &\leq \frac{1}{n^{\alpha\wedge\beta\wedge\gamma}} \mathbb{E}\left[\sum_{\substack{k\in\mathbb{Z}_0^3\\j\in\{1,2\}}} \int_0^t \langle a_{k,j}e^{ik\cdot x}, B_s^n\times(\nabla\times S^n(t-s)\psi)\rangle^2 ds\right] \\ &\leq \frac{1}{n^{\alpha\wedge\beta\wedge\gamma}} \mathbb{E}\left[\int_0^t \|B_s^n\times(\nabla\times S^n(t-s)\psi)\|^2 ds\right] \\ &\lesssim_{\eta,\delta} \frac{1}{n^{\alpha\wedge\beta\wedge\gamma}} \|\psi\|_{H^{\frac{5}{2}+\delta}}^2 \mathbb{E}\left[\int_0^T \|B_s^n\|^2 ds\right], \end{split}$$

uniformly in $t \in [0, T]$. Therefore, choosing $\psi = a_{h,l}e^{ih \cdot x}$, $h \in Z_0^3$, $l \in \{1, 2\}$ we obtain, due to Proposition 3.4

(4.14)
$$\mathbb{E}\left[\langle Z_t^n, a_{h,l} e^{ih \cdot x} \rangle^2\right] \lesssim \frac{1}{n^{\alpha \wedge \beta \wedge \gamma}} |h|^{5+2\delta} \|B_0^n\|_{\mathbf{H}^{-1}}^2.$$

Relation (4.12) then follows from (4.14). Indeed, for each $t \in [0, T]$

$$\mathbb{E}\left[\left\|Z_t^n\right\|_{\mathbf{H}^{-4-\frac{3}{2}\delta}}^2\right] = \sum_{\substack{h \in \mathbb{Z}_0^3\\l \in \{1,2\}}} \frac{\mathbb{E}\left[\langle Z_t^n, a_{h,l}e^{ih \cdot x}\rangle^2\right]}{|h|^{8+3\delta}}$$

$$\begin{split} \lesssim \frac{1}{n^{\alpha \wedge \beta \wedge \gamma}} \sum_{h \in \mathbb{Z}_0^3} \frac{\|B_0^n\|_{\mathbf{H}^{-1}}^2}{|h|^{3+\delta}} \\ \lesssim_{\delta} \frac{\|B_0^n\|_{\mathbf{H}^{-1}}^2}{n^{\alpha \wedge \beta \wedge \gamma}}. \end{split}$$

In order to get (4.13) we argue slightly differently. By Burkholder-Davis-Gundy inequality and the regularization properties of the semigroup, see Lemma 2.2, we have

$$\begin{split} \mathbb{E}\left[\left\|H_{t}^{n}\right\|_{\mathbf{H}^{-\frac{3}{2}-\frac{3}{2}\delta}}^{2}\right] \lesssim_{|\rho|} \sum_{\substack{k \in \mathbb{Z}_{0}^{3} \\ j \in \{1,2\}}} \mathbb{E}\left[\int_{0}^{t} \left\|S^{n}(t-s)\left[\sigma_{k,j}^{n} \cdot \nabla B_{s}^{n,3}\right] - S^{n}(t-s)\left[B_{s}^{n} \cdot \nabla \sigma_{k,j}^{n,3}\right]\right\|_{\dot{H}^{-\frac{3}{2}-\frac{3}{2}\delta}}^{2} ds\right] \\ \lesssim \sum_{\substack{k \in \mathbb{Z}_{0}^{3} \\ j \in \{1,2\}}} \mathbb{E}\left[\int_{0}^{t} \frac{1}{(t-s)^{1-\delta}} \left\|\sigma_{k,j}^{n} \cdot \nabla B_{s}^{n,3}\right\|_{\dot{H}^{-\frac{5}{2}-\frac{1}{2}\delta}}^{2} ds\right] \\ + \sum_{\substack{k \in \mathbb{Z}_{0}^{3} \\ j \in \{1,2\}}} \mathbb{E}\left[\int_{0}^{t} \left\|B_{s}^{n} \cdot \nabla \sigma_{k,j}^{n,3}\right\|_{\dot{H}^{-\frac{3}{2}-\frac{3}{2}\delta}}^{2} ds\right] \\ \lesssim \sum_{\substack{k \in \mathbb{Z}_{0}^{3} \\ j \in \{1,2\}}} \mathbb{E}\left[\int_{0}^{t} \frac{1}{(t-s)^{1-\delta}} \left\|\sigma_{k,j}^{n} B_{s}^{n,3}\right\|_{\dot{H}^{-\frac{3}{2}-\frac{1}{2}\delta}}^{2} ds\right] + \sum_{\substack{k \in \mathbb{Z}_{0}^{3} \\ j \in \{1,2\}}} \mathbb{E}\left[\int_{0}^{t} \left\|B_{s}^{n} \sigma_{k,j}^{n,3}\right\|^{2} ds\right]. \end{split}$$

The second term can be analyzed easily by Proposition 3.4 obtaining

$$\sum_{\substack{k \in \mathbb{Z}_0^3\\j \in \{1,2\}}} \mathbb{E}\left[\int_0^t \left\| B_s^n \sigma_{k,j}^{n,3} \right\|^2 ds\right] \lesssim \left(\sum_{\substack{k \in \mathbb{Z}_0^2\\n \le |k| \le 2n}} \frac{1}{|k|^4} + \sum_{\substack{k \in \mathbb{Z}_0^3\\n \le |k| \le 2n}} \frac{1}{|k|^\gamma}\right) \| B_0^n \|_{\mathbf{H}^{-1}}^2$$
$$\lesssim \frac{\| B_0^n \|_{\mathbf{H}^{-1}}^2}{n^2}.$$

In order to treat the other one we need to argue in a more precise way. Indeed, by definition of the Sobolev norm on the 3D torus, we have

$$\begin{split} \sum_{\substack{k \in \mathbb{Z}_0^3\\ j \in \{1,2\}}} \mathbb{E} \left[\int_0^t \frac{1}{(t-s)^{1-\delta}} \left\| \sigma_{k,j}^n B_s^{n,3} \right\|_{\dot{H}^{-\frac{3}{2} - \frac{1}{2}\delta}}^2 ds \right] &\lesssim \frac{1}{n^2} \sum_{k \in \mathbb{Z}_0^3} \mathbb{E} \left[\int_0^t \frac{1}{(t-s)^{1-\delta}} \left\| e^{ik \cdot x} B_s^{n,3} \right\|_{\dot{H}^{-\frac{3}{2} - \frac{1}{2}\delta}}^2 ds \right] \\ &= \frac{1}{n^2} \sum_{h \in \mathbb{Z}_0^3} \frac{1}{|h|^{3+\delta}} \sum_{k \in \mathbb{Z}_0^3} \mathbb{E} \left[\int_0^t \frac{1}{(t-s)^{1-\delta}} \langle e^{i(k-h) \cdot x}, B_s^{n,3} \rangle^2 ds \right] \\ &\leq \frac{1}{n^2} \sum_{h \in \mathbb{Z}_0^3} \frac{1}{|h|^{3+\delta}} \mathbb{E} \left[\int_0^t \frac{1}{(t-s)^{1-\delta}} \sup_{t \in [0,T]} \left\| B_t^{n,3} \right\|^2 ds \right] \\ &\lesssim_{\delta} \frac{\left\| B_0^n \right\|_{\mathbf{H}^{-1}}^2 + \left\| B_0^{n,3} \right\|^2}{n^2} \end{split}$$

uniformly in $t \in [0, T]$. This concludes the proof of (4.13).

Combining Lemma 4.6 and Lemma 4.7, by interpolation we get the following result.

Corollary 4.8. Assuming either Hypothesis 2.4 or Hypothesis 2.6, for each $\theta \in (0,3]$ and $\delta \in (0,\theta]$ we have

$$\sup_{t\in[0,T]} \mathbb{E}\left[\|Z_t^n\|_{\mathbf{H}^{-1-\theta}}^2 \right] \lesssim \frac{1}{n^{\frac{(\alpha\wedge\beta\wedge\gamma)(\theta-\delta)}{3+\frac{\delta}{2}}}} \|B_0^n\|_{\mathbf{H}^{-1}}^2.$$

In case of Hypothesis 2.5 we have also for each $\vartheta \in (0, \frac{3}{2}]$ and $\delta \in (0, \vartheta]$

$$\sup_{t \in [0,T]} \mathbb{E}\left[\|H_t^n\|_{\dot{H}^{-\vartheta}}^2 \right] \lesssim \frac{\left\| B_0^{n,3} \right\|^2 + \|B_0^n\|_{\mathbf{H}^{-1}}^2}{n^{\frac{4(\vartheta-\delta)}{3}}}.$$

Moreover, we recall that by classical theory of evolution equations, see for example [21], or arguing as in Lemma 4.4, the unique weak solutions of (2.21) and its third component can be written in mild form as

$$\overline{B_t} = S(t)\overline{B_0} + \int_0^t S(t-s)\nabla \times (\mathcal{A}_{\rho}\overline{B_s})ds, \quad \overline{B_t^3} = S(t-s)\overline{B_0^3} - \frac{\rho\zeta_{H,2}}{C_{1,H}C_{2,H}} \int_0^t S(t-s)\operatorname{div}_H\left(\overline{B_s^H}^{\perp}\right)ds.$$

Now we introduce some intermediate vector fields between B_t^n and $\overline{B_t}$. Let $\widehat{B_t^n}$ the unique weak solution of the linear system

(4.15)
$$\begin{cases} \partial_t \widehat{B}^n_t &= (\eta \Delta + \Lambda^n) \widehat{B}^n_t + \nabla \times (\mathcal{A}^n_\rho \widehat{B}^n_t) \quad x \in \mathbb{T}^3, \ t \in (0,T) \\ \widehat{B}^n_t|_{t=0} &= B^n_0. \end{cases}$$

Again, by classical theory of evolution equations, see for example [20, 8], previous equation is well-posed either in \mathbf{H}^{-1} and \mathbf{L}^2 . Moreover the unique weak solution and its third component can be written in mild form as

$$\begin{split} \widehat{B_t^n} &= S^n(t-s)B_0^n + \int_0^t S^n(t-s)\nabla \times (\mathcal{A}_\rho^n \widehat{B_s^n}) ds, \\ \widehat{B_t^{3,n}} &= S^n(t-s)B_0^{n,3} - \frac{\rho \zeta_{H,2}^n}{C_{1,H}C_{2,H}} \int_0^t S^n(t-s)\operatorname{div}_H\left(\widehat{B_s^{n,H}}^{\perp}\right) ds. \end{split}$$

Now we prove that $\widehat{B_t^n}$ and $\overline{B_t}$ are close. We recall that the function χ introduced in subsection 2.1 satisfies

$$0 < \chi(\alpha, \beta, \gamma) \le 1$$

for each choice of the parameters in the range described by Hypothesis 2.4, Hypothesis 2.5 and Hypothesis 2.6. Lemma 4.9. Under the same assumptions of Theorem 2.17, for each $\theta \in (0, 2)$ we have

(4.16)
$$\sup_{t \in [0,T]} \left\| \widehat{B_t^n} - \overline{B_t} \right\|_{\dot{H}^{-1-\theta}} \lesssim_{\eta,T} \frac{1}{n^{\theta\chi(\alpha,\beta,\gamma)/2}} + \left\| \overline{B_0} - B_0^n \right\|_{\mathbf{H}^{-1-\theta}}$$

In case of Hypothesis 2.5 we have also for each $\vartheta \in (\theta, \frac{3}{2}]$

(4.17)
$$\sup_{t \in [0,T]} \left\| \widehat{B_t^{n,3}} - \overline{B_t^3} \right\|_{\dot{H}^{-\vartheta}} \lesssim_{\eta,T} \frac{1}{(\vartheta - \theta)n^{\theta/2}} + \left\| \overline{B_0^3} - B_0^{n,3} \right\|_{\dot{H}^{-\vartheta}} + \frac{\left\| \overline{B_0} - B_0^n \right\|_{\mathbf{H}^{-1-\theta}}}{\vartheta - \theta}$$

Proof. First let us observe that by assumptions in case of Hypothesis 2.4 and Hypothesis 2.6 (resp. Hypothesis 2.5)

$$B_0^n \to \overline{B_0} \text{ in } \mathbf{H}^{-1} \quad (\text{resp. } B_0^n \to \overline{B_0} \text{ in } \mathbf{H}^{-1}, \ B_0^{n,3} \to \overline{B_0^3} \text{ in } \dot{L}^2(\mathbb{T}^3))$$

we have in particular that the family $\left\{ \|B_0^n\|_{\mathbf{H}^{-1}}^2 \right\}_{n \in \mathbb{N}}$ (resp. $\left\{ \|B_0^n\|_{\mathbf{H}^{-1}}^2 \right\}_{n \in \mathbb{N}}$, $\left\{ \|B_0^{n,3}\|^2 \right\}_{n \in \mathbb{N}}$) is bounded. Moreover, $\mathcal{A}_{\rho}^n = \mathcal{A}_{\rho} + O(\frac{1}{n})$, then in particular we have

(4.18)
$$\sup_{n\in\mathbb{N}}\left\|\mathcal{A}_{\rho}^{n}\right\|_{HS} < +\infty$$

Lastly, by simple energy estimates on (4.15) we have for each $n \in \mathbb{N}$

The convergence of
$$\widehat{B_t^n}$$
 to $\overline{B_t}$ then follows by triangle inequality, Lemma 2.2, Lemma 2.3 and the uniform bounds on the initial conditions. Indeed for each $t \in [0, T]$ it holds

$$\begin{split} \left\|\widehat{B_{t}^{n}}-\overline{B_{t}}\right\|_{\dot{H}^{-1-\theta}} &\leq \left(\left\|\left(S^{n}(t)-S(t)\right)\overline{B_{0}}\right\|_{\mathbf{H}^{-1-\theta}}+\left\|S^{n}(t)\left(B_{0}^{n}-\overline{B_{0}}\right)\right\|_{\mathbf{H}^{-1-\theta}}\right) \\ &+\left\|\int_{0}^{t}\left(S^{n}(t-s)-S(t-s)\right)\nabla\times\left(\mathcal{A}_{\rho}^{n}\widehat{B_{s}^{n}}\right)ds\right\|_{\mathbf{H}^{-1-\theta}} \end{split}$$

$$+ \left\| \int_{0}^{t} S(t-s) \nabla \times \left(\left(\mathcal{A}_{\rho}^{n} - \mathcal{A}_{\rho} \right) \widehat{B_{s}^{n}} \right) ds \right\|_{\mathbf{H}^{-1-\theta}} \\ + \left\| \int_{0}^{t} S(t-s) \nabla \times \left(\mathcal{A}_{\rho} \left(\widehat{B_{s}^{n}} - \overline{B_{s}} \right) \right) ds \right\|_{\mathbf{H}^{-1-\theta}} = I_{1} + I_{2} + I_{3} + I_{4}.$$

 I_1 can be treated by Lemma 2.3 :

(4.22)

(4.21)
$$I_1 \lesssim_{\eta,T} \frac{1}{n^{\theta \chi(\alpha,\beta,\gamma)/2}} + \left\|\overline{B_0} - B_0^n\right\|_{\mathbf{H}^{-1-\theta}}.$$

Thanks to Lemma 2.3, (4.18) and (4.19) it holds

$$\begin{split} I_2 &\lesssim \frac{1}{n^{\theta\chi(\alpha,\beta,\gamma)/2}} \int_0^t |t-s|^{\theta/2} \left\| \nabla \times \left(\mathcal{A}_{\rho}^n \widehat{B_s^n} \right) ds \right\|_{\mathbf{H}^{-1}} ds \\ &\lesssim \frac{1}{n^{\theta\chi(\alpha,\beta,\gamma)/2}} \int_0^t |t-s|^{\theta/2} \left\| \widehat{B_s^n} \right\| ds \\ &\lesssim_T \frac{1}{n^{\theta\chi(\alpha,\beta,\gamma)/2}}. \end{split}$$

Due to Lemma 2.2 and (4.19) we can treat I_3 obtaining easily:

(4.23)
$$I_{3} \lesssim_{\eta,\theta} \int_{0}^{t} \left\| \nabla \times \left(\left(\mathcal{A}_{\rho}^{n} - \mathcal{A}_{\rho} \right) \widehat{B}_{s}^{n} \right) \right\|_{\mathbf{H}^{-1}} ds$$
$$\lesssim \left\| \mathcal{A}_{\rho}^{n} - \mathcal{A}_{\rho} \right\|_{HS} \int_{0}^{T} \left\| \widehat{B}_{s}^{n} \right\| ds$$
$$\lesssim_{T} \frac{1}{n}.$$

Lastly we treat I_4 by Lemma 2.2:

(4.24)
$$I_{4} \lesssim_{\eta} \int_{0}^{t} \frac{1}{(t-s)^{1/2}} \left\| \nabla \times \left(\mathcal{A}_{\rho} \left(\widehat{B_{s}^{n}} - \overline{B_{s}} \right) \right) \right\|_{\mathbf{H}^{-2-\theta}} ds$$
$$\lesssim \int_{0}^{t} \frac{1}{(t-s)^{1/2}} \left\| \widehat{B_{s}^{n}} - \overline{B_{s}} \right\|_{\mathbf{H}^{-1-\theta}} ds.$$

Combining (4.21),(4.22),(4.23),(4.24) we get (4.16) by Gronwall's lemma. We move now to $\widehat{B_t^{n,3}} - \overline{B_t^{n,3}}$. First we recall that in case of Hypothesis 2.5 $\chi(\alpha, \beta, \gamma) \equiv 1$ and we have

$$\begin{split} \left\|\widehat{B_{t}^{n,3}} - \overline{B_{t}^{3}}\right\|_{\dot{H}^{-\vartheta}} &\lesssim \left(\left\|\left(S^{n}(t) - S(t)\right)\overline{B_{0}^{3}}\right\|_{\dot{H}^{-\vartheta}} + \left\|S^{n}(t)\left(B_{0}^{n,3} - \overline{B_{0}^{3}}\right)\right\|_{\dot{H}^{-\vartheta}}\right) \\ &+ \zeta_{H,2}^{n} \left\|\int_{0}^{t} \left(S^{n}(t-s) - S(t-s)\right)\operatorname{div}_{H}\left(\widehat{B_{s}^{n,H}}^{\perp}\right) ds\right\|_{\dot{H}^{-\vartheta}} \\ &+ \left|\zeta_{H,2}^{n} - \zeta_{H,2}\right| \left\|\int_{0}^{t} S(t-s)\operatorname{div}_{H}\left(\widehat{B_{s}^{n,H}}^{\perp}\right) ds\right\|_{\dot{H}^{-\vartheta}} \\ &+ \zeta_{H,2} \left\|\int_{0}^{t} S(t-s)\operatorname{div}\left(\widehat{B_{s}^{n,H}}^{\perp} - \overline{B_{s}^{H}}^{\perp}\right) ds\right\|_{\dot{H}^{-\vartheta}} = J_{1} + J_{2} + J_{3} + J_{4}. \end{split}$$

 ${\cal J}_1$ can be treated as in the previous case obtaining

(4.25)
$$J_1 \lesssim \frac{1}{n^{\vartheta/2}} + \left\| \overline{B_0^3} - B_0^{n,3} \right\|_{\dot{H}^{-\vartheta}}.$$

Thanks to Lemma 2.1, Lemma 2.3, Lemma 2.2 and (4.19) it holds

(4.26)

$$J_{2} \lesssim \frac{1}{n^{\vartheta/2}} \int_{0}^{t} |t-s|^{\vartheta/2} \left\| e^{(t-s)\frac{\eta}{2}\Delta} \operatorname{div}_{H} \left(\widehat{B_{s}^{n,H}} \right) \right\| ds$$

$$\lesssim_{\eta} \frac{1}{n^{\vartheta/2}} \int_{0}^{t} \frac{1}{(t-s)^{1-\vartheta}} \left\| \widehat{B_{s}^{n}} \right\|_{\mathbf{H}^{-1}} ds$$

$$\lesssim_{T} \frac{1}{\vartheta n^{\vartheta/2}}.$$

Due to Lemma 2.2 we can treat J_3 obtaining:

$$J_{3} \lesssim_{\eta,\vartheta} \frac{1}{n} \int_{0}^{t} \frac{1}{(t-s)^{1-\vartheta/2}} \left\| \operatorname{div}_{H} \left(\widehat{B_{s}^{n,H}} \right) \right\|_{\dot{H}^{-2}} ds$$
$$\lesssim \frac{1}{\vartheta n}.$$

(4.27)

Lastly we treat J_4 combining Lemma 2.2 and (4.16):

(4.28)
$$J_{4} \lesssim_{\eta} \int_{0}^{t} \frac{1}{(t-s)^{1+\theta/2-\vartheta/2}} \left\|\widehat{B}_{s}^{\widehat{n}} - \overline{B}_{s}\right\|_{\dot{H}^{-1-\theta}} ds$$
$$\lesssim_{T} \frac{1}{\vartheta - \theta} \sup_{t \in [0,T]} \left\|\widehat{B}_{t}^{\widehat{n}} - \overline{B}_{t}\right\|_{\dot{H}^{-1-\theta}}$$
$$\lesssim_{\eta,T} \frac{n^{-\theta/2} + \left\|\overline{B}_{0} - B_{0}^{n}\right\|_{\mathbf{H}^{-1-\theta}}}{\vartheta - \theta}.$$

Combining (4.25), (4.26), (4.27), (4.28) we obtain relation (4.17)

Secondly we provide a quantitative result on the closeness of B_t^n (resp. $B_t^n, B_t^{n,3}$) and $\widehat{B_t^n}$ (resp. $\widehat{B_t^n}, \widehat{B_t^{n,3}}$). **Lemma 4.10.** Under the same assumptions of Theorem 2.17, for each $\kappa \in [1,2)$, $\theta \in (0,2)$, and $\delta \in (0,\theta)$ we have

(4.29)
$$\sup_{t \in [0,T]} \mathbb{E}\left[\left\| \widehat{B_t^n} - B_t^n \right\|_{\mathbf{H}^{-1-\theta}}^{\kappa} \right] \lesssim \frac{1}{n^{\frac{\kappa(\alpha \wedge \beta \wedge \gamma)(\theta - \delta)}{6+\delta}}} \left\| B_0^n \right\|_{\mathbf{H}^{-1}}^{\kappa}$$

In case of Hypothesis 2.5 we have also for each $\vartheta \in (\theta, \frac{3}{2}]$

Proof. From Lemma 4.4, we already know that B_t^n and $B_t^{n,3}$ can be rewritten in mild form as

$$B_t^n = S^n(t)B_0^n + \int_0^t S^n(t-s)\nabla \times (\mathcal{A}_{\rho}^n B_s^n)ds + Z_t^n,$$

$$B_t^{n,3} = S^n(t)B_0^{n,3} - \frac{\rho\zeta_{H,2}^n}{C_{1,H}C_{2,H}} \int_0^t S^n(t-s)\operatorname{div}_H\left((B_s^{n,H})^{\perp}\right)ds + H_t^n$$

Therefore, in case of Hypothesis 2.4 and Hypothesis 2.6 we introduce

$$d_t^n = B_t^n - \widehat{B}_t^n,$$

while in case of Hypothesis 2.5 we introduce also

$$D_t^n = B_t^{n,3} - \widehat{B_t^{n,3}}.$$

we have

$$d_t^n = \int_0^t S^n(t-s) \nabla \times (\mathcal{A}_{\rho}^n d_s^n) ds + Z_t^n, \quad D_t^n = H_t^n - \frac{\rho \zeta_{H,2}^n}{C_{1,H} C_{2,H}} \int_0^t S^n(t-s) \operatorname{div}_H \left((d_s^{n,H})^{\perp} \right) ds.$$

Due to Corollary 4.8 and Grönwall's inequality we can obtain (4.29). Indeed,

$$\mathbb{E}\left[\left\|d_{t}^{n}\right\|_{\mathbf{H}^{-1-\theta}}^{\kappa}\right] \lesssim_{\eta} \mathbb{E}\left[\left(\int_{0}^{t} \frac{1}{\sqrt{t-s}} \left\|d_{s}^{n}\right\|_{\mathbf{H}^{-1-\theta}} ds\right)^{\kappa}\right] + \mathbb{E}\left[\left\|Z_{t}^{n}\right\|_{\mathbf{H}^{-1-\theta}}^{2}\right]^{\frac{\kappa}{2}} \\ \lesssim_{T} \int_{0}^{t} \frac{1}{(t-s)^{\kappa/2}} \mathbb{E}\left[\left\|d_{s}^{n}\right\|_{\mathbf{H}^{-1-\theta}}^{\kappa}\right] ds + \mathbb{E}\left[\left\|Z_{t}^{n}\right\|_{\mathbf{H}^{-1-\theta}}^{2}\right]^{\frac{\kappa}{2}}.$$

This relation easily implies (4.29). Now we move to the analysis of D_t^n . H_t^n can be treated easily by Corollary 4.8 obtaining

(4.31)
$$\sup_{t \in [0,T]} \mathbb{E} \left[\|H_t^n\|_{\dot{H}^{-\vartheta}}^{\kappa} \right] \lesssim \frac{\left\| B_0^{n,3} \right\|^{\kappa} + \|B_0^n\|_{\mathbf{H}^{-1}}^{\kappa}}{n^{\frac{2\kappa(\vartheta-\delta)}{3}}}.$$

28

In order to treat the deterministic convolution, we first argue by interpolation obtaining thanks to Lemma 2.2

$$\begin{split} \left\| \int_{0}^{t} S^{n}(t-s) \operatorname{div}_{H} \left((d_{s}^{n,H})^{\perp} \right) ds \right\|_{\dot{H}^{-\vartheta}}^{\kappa} \\ &\leq \left\| \int_{0}^{t} S^{n}(t-s) \operatorname{div}_{H} \left((d_{s}^{n,H})^{\perp} \right) ds \right\|_{\dot{H}^{-2-\vartheta}}^{\kappa(\vartheta-\vartheta)} \left\| \int_{0}^{t} S^{n}(t-s) \operatorname{div}_{H} \left((d_{s}^{n,H})^{\perp} \right) ds \right\|_{\dot{H}^{-\vartheta}}^{\frac{\kappa(\vartheta-\vartheta)}{2}} \\ &\lesssim_{\eta} \left(\int_{0}^{t} \| d_{s}^{n} \|_{\mathbf{H}^{-1-\vartheta}} ds \right)^{\frac{\kappa(\vartheta-\vartheta)}{2}} \left(\int_{0}^{t} \frac{1}{(t-s)^{1-\vartheta/2}} \left\| \widehat{B}_{s}^{n} - B_{s}^{n} \right\|_{\mathbf{H}^{-1}} ds \right)^{\frac{\kappa(2-\vartheta+\vartheta)}{2}} \\ &\lesssim_{T} \frac{1}{\theta^{\frac{\kappa(2-\vartheta+\vartheta)}{2}}} \left(\int_{0}^{t} \| d_{s}^{n} \|_{\mathbf{H}^{-1-\vartheta}}^{\kappa} ds \right)^{\frac{\vartheta-\vartheta}{2}} \operatorname{sup}_{t\in[0,T]} \left(\| B_{t}^{n} \|_{\mathbf{H}^{-1}} + \left\| \widehat{B}_{t}^{n} \right\|_{\mathbf{H}^{-1}} \right)^{\frac{\kappa(2-\vartheta+\vartheta)}{2}}. \end{split}$$

Combining (4.32), (4.29), (3.4), (4.19) we obtain an estimate of the deterministic convolution. Indeed, by Hölder inequality, for each $t \in [0, T]$ it holds:

$$\begin{split} & \left(\frac{|\rho|\zeta_{H,2}^{n}}{C_{1,H}C_{2,H}}\right)^{\kappa} \mathbb{E}\left[\left\|\int_{0}^{t} S^{n}(t-s)\operatorname{div}_{H}\left((d_{s}^{n,H})^{\perp}\right)ds\right\|_{\dot{H}^{-\vartheta}}^{\kappa}\right] \\ & \lesssim_{\eta,T,\theta} \mathbb{E}\left[\left(\int_{0}^{t} \|d_{s}^{n}\|_{\mathbf{H}^{-1-\theta}}^{\kappa}ds\right)^{\frac{\vartheta-\theta}{2}} \operatorname{sup}_{t\in[0,T]}\left(\|B_{t}^{n}\|_{\mathbf{H}^{-1}}+\left\|\widehat{B_{t}^{n}}\right\|_{\mathbf{H}^{-1}}\right)^{\frac{\kappa(2-\vartheta+\theta)}{2}}\right] \\ & \leq \mathbb{E}\left[\int_{0}^{T} \|d_{s}^{n}\|_{\mathbf{H}^{-1-\theta}}^{\kappa}ds\right]^{\frac{\vartheta-\theta}{2}} \mathbb{E}\left[\operatorname{sup}_{t\in[0,T]}\left(\|B_{t}^{n}\|_{\mathbf{H}^{-1}}^{\kappa}+\left\|\widehat{B_{t}^{n}}\right\|_{\mathbf{H}^{-1}}^{\kappa}\right)\right]^{\frac{2-\vartheta+\theta}{2}} \\ & \lesssim_{\delta}\frac{1}{n^{\frac{\kappa(\vartheta-\theta)(\theta-\delta)}{6+\delta}}} \|B_{0}^{n}\|_{\mathbf{H}^{-1}}^{\kappa}. \end{split}$$

(4.33)

(4.32)

Combining (4.31) and (4.33) the result follows.

The proof of Theorem 2.17 then follows combining Lemma 4.9 and Lemma 4.10.

ACKNOWLEDGEMENTS. The authors thank Professor Franco Flandoli for useful discussions and valuable insight into the subject.

References

- [1] Antonio Agresti. On anomalous dissipation induced by transport noise. arXiv preprint arXiv:2405.03525, 2024.
- [2] Axel Brandenburg, Dmitry Sokoloff, and Kandaswamy Subramanian. Current status of turbulent dynamo theory: from largescale to small-scale dynamos. Space Science Reviews, 169:123–157, 2012.
- [3] Federico Butori, Franco Flandoli, and Eliseo Luongo. On the Itô-Stratonovich Diffusion Limit for the Magnetic Field in a 3D Thin Domain. arXiv preprint arXiv:2401.15701, 2024.
- [4] Giulia Carigi and Eliseo Luongo. Dissipation properties of transport noise in the two-layer quasi-geostrophic model. Journal of Mathematical Fluid Mechanics, 25(2):28, 2023.
- [5] Alice Courvoisier, Andrew D Gilbert, and Yannick Ponty. Dynamo action in flows with cat's eyes. Geophysical and Astrophysical Fluid Dynamics, 99(5):413–429, 2005.
- [6] Arnaud Debussche and Umberto Pappalettera. Second order perturbation theory of two-scale systems in fluid dynamics. arXiv preprint arXiv:2206.07775, 2022.
- [7] Klaus-Jochen Engel and Rainer Nagel. A short course on operator semigroups. Universitext. Springer, New York, 2006.
- [8] Franco Flandoli. Regularity theory and stochastic flows for parabolic SPDEs, volume 9 of Stochastics Monographs. Gordon and Breach Science Publishers, Yverdon, 1995.
- [9] Franco Flandoli, Lucio Galeati, and Dejun Luo. Scaling limit of stochastic 2D Euler equations with transport noises to the deterministic Navier–Stokes equations. Journal of Evolution Equations, 21(1):567–600, 2021.
- [10] Franco Flandoli, Lucio Galeati, and Dejun Luo. Quantitative convergence rates for scaling limit of spdes with transport noise. Journal of Differential Equations, 394:237–277, 2024.
- [11] Franco Flandoli and Dejun Luo. High mode transport noise improves vorticity blow-up control in 3D Navier-Stokes equations. Probab. Theory Related Fields, 180(1-2):309–363, 2021.
- [12] Franco Flandoli and Dejun Luo. On the Boussinesq hypothesis for a stochastic Proudman-Taylor model. to appear on SIAM J. Math. Anal., 2023.
- [13] Franco Flandoli, Dejun Luo, and Eliseo Luongo. 2D Smagorinsky-Type Large Eddy Models as Limits of Stochastic PDEs. Journal of Nonlinear Science, 34(3):54, 2024.
- [14] Franco Flandoli and Eliseo Luongo. Heat diffusion in a channel under white noise modeling of turbulence. Mathematics in Engineering, 4(4):1–21, 2022.

- [15] Franco Flandoli and Eliseo Luongo. Stochastic partial differential equations in fluid mechanics, volume 2330 of Lecture Notes in Mathematics. Springer, Singapore, 2023.
- [16] Franco Flandoli and Francesco Russo. Reduced dissipation effect in stochastic transport by Gaussian noise with regularity greater than 1/2. arXiv preprint arXiv:2305.19293, 2023.
- [17] Lucio Galeati. On the convergence of stochastic transport equations to a deterministic parabolic one. Stochastics and Partial Differential Equations: Analysis and Computations, 8(4):833–868, 2020.
- [18] Andrew D Gilbert. Dynamo theory. In Handbook of mathematical fluid dynamics, volume 2, pages 355-441. Elsevier, 2003.
- [19] Fritz Krause and K-H Rädler. Mean-field magnetohydrodynamics and dynamo theory. Elsevier, 2016.
- [20] Jacques-Louis Lions and Enrico Magenes. Non-homogeneous boundary value problems and applications. Vol. I, volume Band 181 of Die Grundlehren der mathematischen Wissenschaften. Springer-Verlag, New York-Heidelberg, 1972. Translated from the French by P. Kenneth.
- [21] Alessandra Lunardi. Analytic semigroups and optimal regularity in parabolic problems. Modern Birkhäuser Classics. Birkhäuser/Springer Basel AG, Basel, 1995. [2013 reprint of the 1995 original] [MR1329547].
- [22] Carlo Marchioro and Mario Pulvirenti. Mathematical theory of incompressible nonviscous fluids, volume 96 of Applied Mathematical Sciences. Springer-Verlag, New York, 1994.
- [23] Henry K Moffatt. Magnetic field generation in electrically conducting fluids. Cambridge University Press, Cambridge, London, New York, Melbourne, 2:5–1, 1978.
- [24] Amnon Pazy. Semigroups of linear operators and applications to partial differential equations, volume 44. Springer Science & Business Media, 2012.
- [25] Michael Richard Edward Proctor and Andrew D Gilbert. Lectures on solar and planetary dynamos, volume 2. Cambridge University Press, 1994.
- [26] Gareth Owen Roberts. Dynamo action of fluid motions with two-dimensional periodicity. Philosophical Transactions of the Royal Society of London. Series A, Mathematical and Physical Sciences, 271(1216):411-454, 1972.
- [27] Karl-Heinz R\u00e4del and Axel Brandenburg. Contributions to the theory of a two-scale homogeneous dynamo experiment. Physical review. E, Statistical, nonlinear, and soft matter physics, 67:026401, 03 2003.
- [28] Max Steenbeck, Fritz Krause, and K-H R\u00e4dler. Berechnung der mittleren Lorentz-Feldst\u00e4rke f\u00fcr ein elektrisch leitendes Medium in turbulenter, durch Coriolis-Kr\u00e4fte beeinflu\u00dfter Bewegung. Zeitschrift f\u00fcr Naturforschung A, 21(4):369-376, 1966.
- [29] Roger Temam. Navier–Stokes equations and nonlinear functional analysis. SIAM, 1995.
- [30] Roger Temam. Navier-Stokes equations: theory and numerical analysis, volume 343. American Mathematical Soc., 2001.
- [31] Andreas Tilgner and Axel Brandenburg. A growing dynamo from a saturated roberts flow dynamo. Monthly Notices of the Royal Astronomical Society, 391:1477 – 1481, 12 2008.
- [32] Hans Triebel. Interpolation Theory, Function Spaces, Differential Operators (2nd edition). Johann Ambrosius Barth Verlag, Leipzig, 1995.