

# ODD AND EVEN ELLIPTIC CURVES WITH COMPLEX MULTIPLICATION

YURI G. ZARHIN

ABSTRACT. We call an order  $O$  in a quadratic field  $K$  *odd* (resp. *even*) if its discriminant is an odd (resp. even) integer. We call an elliptic curve  $E$  over  $\mathbb{C}$  with CM *odd* (resp. *even*) if its endomorphism ring  $\text{End}(E)$  is an odd (resp. even) order in the imaginary quadratic field  $\text{End}(E) \otimes \mathbb{Q}$ .

Suppose that  $j(E) \in \mathbb{R}$  and let us consider the set  $\mathcal{J}(\mathbb{R}, E)$  of all  $j(E')$  where  $E'$  is any elliptic curve that enjoys the following properties.

- $E'$  is isogenous to  $E$ ;
- $j(E') \in \mathbb{R}$ ;
- $E'$  has the same parity as  $E$ .

We prove that the closure of  $\mathcal{J}(\mathbb{R}, E)$  in  $\mathbb{R}$  is the closed semi-infinite interval  $(-\infty, 1728]$  (resp. the whole  $\mathbb{R}$ ) if  $E$  is odd (resp. even).

This paper was inspired by a question of Jean-Louis Colliot-Thélène and Alena Pirutka about the distribution of  $j$ -invariants of certain elliptic curves of CM type.

As usual  $\mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}$  stand for the ring of integers and the fields of rational, real and complex numbers respectively. If  $n$  is a positive integer then we write  $\mathbb{Z}[1/n]$  for the subring of  $\mathbb{Q}$  generated by  $\mathbb{Z}$  and  $1/n$ . More generally, if  $F$  is a field of characteristic zero and  $A$  is a subring of  $F$  (with 1) then we write  $A[1/n]$  for the subring of  $F$  generated by  $A$  and  $1/n$ ; clearly,  $A[1/n]$  is a  $\mathbb{Z}[1/n]$ -subalgebra of  $F$ . We write  $\mathbb{Z}_{(2)}$  for the subring of  $\mathbb{Q}$  that consists of all fractions  $\frac{a}{n}$  where  $a \in \mathbb{Z}$  and  $n$  an odd integer.

The aim of this note is to introduce and study the notion of a *parity* for elliptic curves over  $\mathbb{C}$  with complex multiplication (CM) in the following way. Let  $E$  be an elliptic curve, whose endomorphism ring  $\text{End}(E)$  is an *order* in an imaginary quadratic field. We call  $E$  *odd* (resp. *even*) if the discriminant of  $\text{End}(E)$  is an *odd* (resp. *even*) integer. We call its  $j$ -invariant  $j(E) \in \mathbb{C}$  *odd* (resp. *even*) if  $E$  is *odd* (resp. *even*). It turns out that both sets of even and odd  $j$ -invariants are everywhere dense in  $\mathbb{C}$ . Surprisingly, this is not the case when we restrict ourselves to the sets of even and odd *real*  $j$ -invariants (i.e., when the corresponding curves are defined over  $\mathbb{R}$ ). Namely, while the set of real *even*  $j$ -invariants is dense in  $\mathbb{R}$ , the set of real *odd*  $j$ -invariants

---

2020 *Mathematics Subject Classification.* 11G07, 14H52.

*Key words and phrases.* Quadratic orders, discriminants, elliptic curves, complex multiplication.

The author was partially supported by Simons Foundation Collaboration grant # 585711.

lies in an open semi-infinite interval  $(-\infty, 1728]$  and everywhere dense there. Actually, we prove stronger versions of these results, dealing with isogeny classes of CM elliptic curves with the same “parity”.

The paper is organized as follows. In Section 1 we discuss various properties of quadratic orders with odd (resp. even) discriminants. Section 2 contains the statements of our main results about  $j$ -invariants of odd and even elliptic curves. In Section 3 we discuss auxiliary results about isogenies of complex elliptic curves. We prove our main results in Section 4, using properties of “odd” and “even” orders and parity preservation under isogenies of odd degrees that will be proven in Section 5 and 5 respectively. In Section 7 we “classify” (actually, list all the  $j$ -invariants of) CM elliptic curves, whose  $j$ -invariant is real and the endomorphism ring is an order with given odd discriminant.

**Acknowledgements.** I am grateful to Jean-Louis Colliot-Thélène and Alena Pirutka for an interesting and stimulating question about the distribution of  $j$ -invariants of certain elliptic curves of CM type and for their useful comments to a preliminary version of this paper, which may be viewed as an extension (or a variant) of my answer [3, Sect. 8]. I thank David Masser and Serge Vladuts for their interest in this paper and stimulating discussions.

## 1. ORDERS IN QUADRATIC FIELDS

Let us start with a discussion of *odd* and *even* orders in arbitrary quadratic fields.

Let  $K$  be a quadratic field,  $O_K$  its ring of integers, and

$$\mathrm{tr} = \mathrm{tr}_K : K \rightarrow \mathbb{Q}$$

be the corresponding trace map attached to the field extension  $K/\mathbb{Q}$ . We have

$$2\mathbb{Z} = \mathrm{tr}(\mathbb{Z}) \subset \mathrm{tr}(O_K) \subset \mathbb{Z}.$$

Let  $O$  be an order in  $K$  and let a positive integer  $\mathfrak{f} = \mathfrak{f}_O$  be the *conductor* of  $O$ . This means that

$$O = \mathbb{Z} + \mathfrak{f} \cdot O_K \subset O_K$$

and  $\mathfrak{f}$  coincides with the index  $[O_K : O]$  of the additive subgroup  $O$  in  $O_K$  [2, Ch. 2, Sect. 7, Th.1]. Since  $\mathbb{Z} \subset O$  and  $\mathrm{tr}(\mathbb{Z}) = 2\mathbb{Z}$ ,

$$\mathrm{tr}(O) = \mathbb{Z} \quad \text{or} \quad 2\mathbb{Z}. \tag{1}$$

Recall that the discriminant  $\mathrm{discr}(O) \in \mathbb{Z}$  of  $O$  is the discriminant of the symmetric bilinear form

$$O \times O \rightarrow \mathbb{Z}, \quad a, b \mapsto \mathrm{tr}(ab).$$

By definition, the discriminant  $\mathrm{discr}(K)$  of the field  $K$  is  $\mathrm{discr}(O_K)$ . It is well known [2, Ch. 2, Sect. 7, Th.1] that

$$\mathrm{discr}(O) = \mathfrak{f}_O^2 \cdot \mathrm{discr}(O_K) = \mathfrak{f}_O^2 \cdot \mathrm{discr}(K). \tag{2}$$

**Definition 1.1** (Definition-Lemma). An order  $O$  is called *odd* if it enjoys the following equivalent properties.

- (i)  $\text{tr}(O) = \mathbb{Z}$ .
- (ii)  $\text{discr}(O)$  is an odd integer.
- (iii) Both  $\text{discr}(K)$  and the conductor  $\mathfrak{f}_O$  are odd integers.
- (iv) There exists an integer  $D \equiv 1 \pmod{4}$  that is not a square and such that

$$\frac{1 + \sqrt{D}}{2} \in O.$$

- (v) There exists an integer  $D \equiv 1 \pmod{4}$  that is not a square such that

$$K = \mathbb{Q}(\sqrt{D}), \quad O = \mathbb{Z} + \mathbb{Z} \frac{1 + \sqrt{D}}{2} = \mathbb{Z} \left[ \frac{1 + \sqrt{D}}{2} \right].$$

**Definition 1.2** (Definition-Lemma). An order  $O$  in a quadratic field  $K$  is called *even* if it enjoys the following equivalent properties.

- (i)  $\text{tr}(O) = 2\mathbb{Z}$ .
- (ii)  $\text{discr}(O)$  is an even integer.
- (iii) Either  $\text{discr}(K)$  or the conductor  $\mathfrak{f}_O$  is an even integer.

Clearly, every order  $O$  is either odd or even. We prove Lemmas 1.1 and 1.2 in Section 5.

**Definition 1.3.** Let  $E$  be an elliptic curve with complex multiplication (CM) over the field  $\mathbb{C}$  of complex numbers. Then its endomorphism ring  $\text{End}(E)$  is an order in the imaginary quadratic field

$$K := \text{End}(E) \otimes \mathbb{Q} =: \text{End}^0(E).$$

We say that  $E$  is *odd* (resp. *even*) if the order  $\text{End}(E)$  is *odd* (resp. *even*).

Clearly, every elliptic curve over  $\mathbb{C}$  with CM is either odd or even. The following assertions will be proven in Section 6.

**Proposition 1.4.** *Let  $\phi : E_1 \rightarrow E_2$  be an isogeny of complex elliptic curves with CM. Suppose that  $n = \deg(\phi)$  is an odd integer. Then  $E_1$  is odd (resp. even) if and only if  $E_2$  is odd (resp. even). In other words,  $E_1$  and  $E_2$  have the same parity.*

**Proposition 1.5.** *Suppose that  $E_1$  and  $E_2$  are isogenous CM elliptic curves over  $\mathbb{C}$ . Suppose that there are elliptic curves  $E_{1,\mathbb{R}}$  and  $E_{2,\mathbb{R}}$  over the field  $\mathbb{R}$  of real numbers such that there are isomorphisms of complex elliptic curves*

$$E_1 \cong E_{1,\mathbb{R}} \times_{\mathbb{R}} \mathbb{C}, \quad E_2 = E_{2,\mathbb{R}} \times_{\mathbb{R}} \mathbb{C}.$$

*Then  $E_{1,\mathbb{R}}$  and  $E_{2,\mathbb{R}}$  are isogenous over  $\mathbb{R}$ .*

**Remark 1.6.** Let  $E$  be an elliptic curve over  $\mathbb{C}$ . It is well known [4, Ch. 3, Sect. 2, Prop. 3.7] that  $j(E) \in \mathbb{R}$  if and only if  $E$  may be defined over  $\mathbb{R}$ , i.e., there is an elliptic curve  $E_{\mathbb{R}}$  over  $\mathbb{R}$  such that  $E \cong E_{\mathbb{R}} \times_{\mathbb{R}} \mathbb{C}$ . (See also [9, Appendix A, Prop. 1.2(b)] and [7, Ch. 4, Sect. 4.1].)

- Examples 1.7.** (e) If  $E$  is an elliptic curve  $y^2 = x^3 - x$  then it is well known that the order  $\text{End}(E) = \mathbb{Z}[\sqrt{-1}]$  has even discriminant  $-4$  and therefore  $E$  is even.
- (o) If  $E$  is an elliptic curve  $y^2 = x^3 - 1$  then it is well known that the order  $\text{End}(E) = \mathbb{Z}\left[\frac{-1+\sqrt{-3}}{2}\right]$  has odd discriminant  $-3$  and therefore  $E$  is odd.

## 2. $j$ -INVARIANTS OF CM ELLIPTIC CURVES

In what follows,  $E$  is an elliptic curve over  $\mathbb{C}$ . The following assertion will be proven in Section 3.

**Proposition 2.1.** *Let us consider the subset  $J_{E,\text{is}}$  of all  $j(E')$  where  $E'$  runs through the set of elliptic curves over  $\mathbb{C}$  such that there exists an isogeny  $E \rightarrow E'$  of odd degree.*

*Then  $J_{E,\text{is}}$  is dense in  $\mathbb{C}$ .*

**Remark 2.2.** If  $E$  has CM then all  $E'$  that appear in Proposition 2.1 are elliptic curves with CM and the corresponding imaginary quadratic fields  $\text{End}^0(E')$  are isomorphic to  $\text{End}^0(E)$ .

Notice that in light of Examples 1.7, there exist an odd CM curve and an even CM curve. Now, combining Proposition 2.1 with Proposition 1.4, we obtain the following statement.

**Corollary 2.3.** *Let  $J^{\text{odd}}(\mathbb{C}) \subset \mathbb{C}$  be the set of all  $j(E^{\text{odd}})$  where  $E^{\text{odd}}$  runs through the set of all odd elliptic curves over  $\mathbb{C}$  with CM.*

*Let  $J^{\text{ev}}(\mathbb{C}) \subset \mathbb{C}$  be the set of all  $j(E^{\text{ev}})$  where  $E^{\text{ev}}$  runs through the set of all even elliptic curves over  $\mathbb{C}$  with CM.*

*Then each of two subsets  $J^{\text{odd}}(\mathbb{C})$  and  $J^{\text{ev}}(\mathbb{C})$  is dense in  $\mathbb{C}$ .*

Our main result is the following assertion.

**Theorem 2.4.** *Suppose that  $E$  is a CM curve with  $j(E) \in \mathbb{R}$ . Let us consider the set  $\mathcal{J}(\mathbb{R}, E)$  of all  $j(E')$  where  $E'$  is any elliptic curve over  $\mathbb{C}$  that enjoys the following properties.*

- $E'$  is isogenous to  $E$ ;
- $j(E') \in \mathbb{R}$ ;
- $E'$  has the same parity as  $E$ .

*Then the closure of  $\mathcal{J}(\mathbb{R}, E)$  in  $\mathbb{R}$  is the closed semi-infinite interval  $(-\infty, 1728]$  (resp. the whole  $\mathbb{R}$ ) if  $E$  is odd (resp. even).*

We prove Theorem 2.4 in Section 4. Our proof is based on auxiliary results about isogenies of complex elliptic curves with CM that will be discussed in Section 3.

**Remark 2.5.** It follows from Remark 1.6 and Proposition 1.5 that Theorem 2.4. is equivalent to the following statement.

**Theorem 2.6.** *Let  $E_{\mathbb{R}}$  be an elliptic curve over  $\mathbb{R}$  such that its complexification  $E = E_{\mathbb{R}} \times_{\mathbb{R}} \mathbb{C}$  is of CM type. Let us consider the set  $\mathcal{J}_0(\mathbb{R}, E_{\mathbb{R}})$  of all  $j(E'_{\mathbb{R}})$  where  $E'_{\mathbb{R}}$  is any elliptic curve over  $\mathbb{R}$  that enjoys the following properties.*

- $E'_{\mathbb{R}}$  is  $\mathbb{R}$ -isogenous to  $E_{\mathbb{R}}$ ;
- The complexification  $E' := E'_{\mathbb{R}} \times_{\mathbb{R}} \mathbb{C}$  of  $E'_{\mathbb{R}}$  has the same parity as  $E$ .

*Then the closure of  $\mathcal{J}_0(\mathbb{R}, E_{\mathbb{R}})$  in  $\mathbb{R}$  is the closed semi-infinite interval  $(-\infty, 1728]$  (resp. the whole  $\mathbb{R}$ ) if  $E$  is odd (resp. even).*

### 3. THE UPPER HALF-PLANE AND ELLIPTIC CURVES

**Definition 3.1.** If  $\omega_1$  and  $\omega_2$  are nonzero complex numbers such that  $\omega_1/\omega_2 \notin \mathbb{R}$  then we write  $[\omega_1, \omega_2]$  for the discrete lattice  $\mathbb{Z}\omega_1 + \mathbb{Z}\omega_2$  of rank 2 in  $\mathbb{C}$ .

Let

$$\mathfrak{H} := \{x + iy \mid x, y \in \mathbb{R}; y > 0\} \subset \mathbb{C}$$

be the upper half-plane. If  $\tau \in \mathfrak{H}$  then we write  $\mathcal{E}_{\tau}$  for the complex elliptic curve such that the complex tori  $\mathcal{E}_{\tau}(\mathbb{C})$  and  $\mathbb{C}/\Lambda_{\tau}$  are isomorphic. Here

$$\Lambda_{\tau} := [\tau, 1] = \mathbb{Z}\tau + \mathbb{Z}.$$

See [10, Ch. V, p. 408–411] for a Weierstrass equation of  $\mathcal{E}_{\tau}$ . (In the notation of [10, Ch. V], our  $\mathcal{E}_{\tau}$  is  $E_q$  with  $q = \exp(2\pi i\tau)$ .)

**Remark 3.2.** If  $\tau_1, \tau_2 \in \mathfrak{H}$  then the group  $\text{Hom}(\mathcal{E}_{\tau_1}, \mathcal{E}_{\tau_2})$  of homomorphisms from  $\mathcal{E}_{\tau_1}, \mathcal{E}_{\tau_2}$  can be canonically identified with

$$\{u \in \mathbb{C} \mid u(\Lambda_{\tau_1}) \subset \Lambda_{\tau_2}\} \subset \mathbb{C}. \quad (3)$$

Each  $u$  that satisfies (3) corresponds to the homomorphism

$$\phi_u : \mathcal{E}_{\tau_1} \rightarrow \mathcal{E}_{\tau_2}$$

of elliptic curves such that the corresponding action of  $\phi_u$  on the complex points is as follows.

$$\mathcal{E}_{\tau_1}(\mathbb{C}) = \mathbb{C}/\Lambda_{\tau_1} \rightarrow \mathbb{C}/\Lambda_{\tau_2} = \mathcal{E}_{\tau_2}(\mathbb{C}), \quad z + \Lambda_{\tau_1} \mapsto uz + \Lambda_{\tau_2}.$$

If  $u \neq 0$  then  $\phi_u$  is an isogeny, whose degree coincides with the index  $[\Lambda_{\tau_2} : u(\Lambda_{\tau_1})]$  of the subgroup  $u(\Lambda_{\tau_1})$  in  $\Lambda_{\tau_2}$  [5, p. 9]. In particular, if

$$\Lambda_{\tau_2} = u(\Lambda_{\tau_1})$$

(i.e., the index is 1),  $\phi_u$  is an isomorphism of elliptic curves  $\mathcal{E}_{\tau_1}$  and  $\mathcal{E}_{\tau_2}$ .

On the other hand, if we take  $\tau_1 = \tau_2 = \tau$  and put

$$O_{\tau} := \{u \in \mathbb{C} \mid u(\Lambda_{\tau}) \subset \Lambda_{\tau}\}, \quad (4)$$

then the map

$$O_{\tau} \rightarrow \text{End}(\mathcal{E}_{\tau}), \quad u \mapsto \phi_u \quad (5)$$

is a ring isomorphism.

**Remark 3.3.** We have

$$\mathbb{Z} \subset O_\tau, \quad \Lambda_\tau \supset O_\tau \cdot 1 = O_\tau \subset \mathbb{C},$$

i.e.,

$$\mathbb{Z} \subset O_\tau \subset \Lambda_\tau \subset \mathbb{C}. \quad (6)$$

It follows easily that if  $\Lambda_\tau$  is a *subring* of  $\mathbb{C}$  then  $O_\tau = \Lambda_\tau$ .

**Remark 3.4.** (i) It is well known [7, Ch. 4, Sect. 4.4, Prop. 4.5] that  $\mathcal{E}_\tau$  has CM if and only if  $\tau$  is an (imaginary) quadratic irrationality, i.e.,  $\mathbb{Q}(\tau) = \mathbb{Q} + \mathbb{Q}\tau$  is an imaginary quadratic field. If this is the case then  $O_\tau$  is an order in the imaginary quadratic field  $\mathbb{Q}(\tau)$ ; in particular,

$$O_\tau = \{u \in \mathbb{Q}(\tau) \mid u(\Lambda_\tau) \subset \Lambda_\tau\}. \quad (7)$$

In addition, extending isomorphism (5) by  $\mathbb{Q}$ -linearity, we get a canonical isomorphism of  $\mathbb{Q}$ -algebras

$$\mathbb{Q}(\tau) = O_\tau \otimes \mathbb{Q} = \mathbb{Q}(\tau) \rightarrow \text{End}(\mathcal{E}_\tau) \otimes \mathbb{Q} = \text{End}^0(\mathcal{E}_\tau), \quad u \otimes r \mapsto \phi_u \otimes r$$

between  $\mathbb{Q}(\tau)$  and the endomorphism algebra  $\text{End}^0(\mathcal{E}_\tau)$  of  $\mathcal{E}_\tau$ .

(ii) Suppose that  $\mathcal{E}_\tau$  has CM, i.e.,  $\mathbb{Q}(\tau)$  is an imaginary quadratic field. In light of (6),

$$\mathbb{Z} \subset O_\tau \subset \Lambda_\tau = \mathbb{Z}\tau + \mathbb{Z} \subset \mathbb{Q}(\tau). \quad (8)$$

It follows from Remark 3.3 that if  $\Lambda_\tau$  is an *order* in  $\mathbb{Q}(\tau)$  then

$$O_\tau = \Lambda_\tau.$$

(ii) Let  $\mathcal{E}_{\tau_1}$  and  $\mathcal{E}_{\tau_2}$  be two elliptic curves with complex multiplication. Then  $\mathcal{E}_{\tau_1}$  and  $\mathcal{E}_{\tau_2}$  are isogenous if and only if the corresponding imaginary quadratic fields coincide, i.e.,

$$\mathbb{Q}(\tau_1) = \mathbb{Q}(\tau_2)$$

(see [7, Ch. 4, Sect. 4.4, Prop. 4.9]).

The group  $\text{GL}_2(\mathbb{R})^+$  of two-by-two real matrices with positive determinant acts transitively on  $\mathfrak{H}$  by fractional-linear transformations. Namely, the continuous map

$$M, \tau \mapsto M(\tau) = \frac{a\tau + b}{c\tau + d} \quad \forall \tau \in \mathfrak{H}, \quad M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}_2(\mathbb{R})^+.$$

defines the transitive action of  $\text{GL}_2(\mathbb{R})^+$  on  $\mathfrak{H}$ . (Actually, even the action of the subgroup  $\text{SL}(2, \mathbb{R})$  on  $\mathfrak{H}$  is transitive.) We will mostly deal with the action of the subgroup

$$G := \text{GL}(2, \mathbb{Z}_{(2)})^+ = \{M \in \text{GL}(2, \mathbb{Z}_{(2)}) \mid \det(M) > 0\} \subset \text{GL}_2(\mathbb{R})^+.$$

**Lemma 3.5.** *Suppose that a matrix*

$$M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{GL}_2(\mathbb{Z}_{(2)})$$

*has positive determinant*

$$\det(M) = ad - bc \in \mathbb{Z}_{(2)}^*,$$

*i.e.,  $M \in G$ . Then there exists an isogeny  $E_{M(\tau)} \rightarrow \mathcal{E}_\tau$  of odd degree.*

*Proof.* By definition of  $\mathbb{Z}_{(2)}$ , there are integers  $\tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}$ , and a positive odd integer  $n$  such that

$$a = \tilde{a}/n, \quad b = \tilde{b}/n, \quad c = \tilde{c}/n, \quad d = \tilde{d}/n.$$

The conditions on  $\det(M)$  imply that  $\tilde{a}\tilde{d} - \tilde{b}\tilde{c}$  is a positive odd integer. Then

$$M(\tau) = \frac{a\tau + b}{c\tau + d} = \frac{\tilde{a}\tau + \tilde{b}}{\tilde{c}\tau + \tilde{d}} = \tilde{M}(\tau)$$

where the matrix

$$\tilde{M} = \begin{pmatrix} \tilde{a} & \tilde{b} \\ \tilde{c} & \tilde{d} \end{pmatrix}.$$

(Since

$$\tilde{a}\tilde{d} - \tilde{b}\tilde{c} = \det(\tilde{M}) = \frac{1}{n^2} \det(M) > 0,$$

we get

$$\tilde{M} \in \mathrm{GL}(2, \mathbb{Z}_{(2)})^+ \subset \mathrm{GL}_2(\mathbb{R})^+.)$$

We have

$$\Lambda_{M(\tau)} = \Lambda_{\tilde{M}(\tau)} = \left[ \frac{\tilde{a}\tau + \tilde{b}}{\tilde{c}\tau + \tilde{d}}, 1 \right].$$

This means that

$$(\tilde{c}\tau + \tilde{d})\Lambda_{M(\tau)} = [\tilde{a}\tau + \tilde{b}, \tilde{c}\tau + \tilde{d}]$$

is a subgroup in  $[\tau, 1] = \Lambda_\tau$  of odd index  $\tilde{a}\tilde{d} - \tilde{b}\tilde{c}$ . This implies that if we put

$$u := \tilde{c}\tau + \tilde{d} \in \mathbb{C}, \quad k := \tilde{a}\tilde{d} - \tilde{b}\tilde{c}$$

then  $u(\Lambda_{M(\tau)})$  is a subgroup of odd index  $k$  in  $\Lambda_\tau$ . This gives us the isogeny  $\phi_u : E_{M(\tau)} \rightarrow \mathcal{E}_\tau$  of odd degree  $k$ . This ends the proof.  $\square$

*Proof of Proposition 2.1.* Let  $B \in \mathrm{GL}(2, \mathbb{R})^+$ . The weak approximation for the field  $\mathbb{Q}$  [1, Th. 1] with respect to places  $\{\infty, 2\}$  implies the existence of a sequence  $\{B_n\}$  of  $2 \times 2$  matrices with rational entries such that  $\{B_n\}$  converges to  $B$  in the real topology and to the identity matrix  $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$  in the 2-adic topology. Removing first finite terms of the sequence, we may and will assume that for all  $n$

$$B_n \in \mathrm{GL}(2, \mathbb{Z}_{(2)}) \subset \mathrm{GL}(2, \mathbb{Q}), \quad \det(B_n) > 0;$$

the latter inequality means that all  $B_n \in \mathrm{GL}(2, \mathbb{R})^+$  and therefore

$$B_n \in \mathrm{GL}(2, \mathbb{Z}_{(2)})^+ = G.$$

It follows that  $B(\tau) \in \mathfrak{H}$  is the limit of the sequence

$$B_n(\tau) \in \mathfrak{H}$$

in the complex topology. Now the transitivity of  $\mathrm{GL}(2, \mathbb{R})^+$  implies that every orbit  $G\tau = \{M(\tau) \mid M \in G\}$  is dense in  $\mathcal{H}$  in the complex topology.

Recall that the classical holomorphic (hence, continuous) modular function

$$j : \mathfrak{H} \rightarrow \mathbb{C}$$

takes on every complex value. It follows that for every  $\tau$  the set  $\{j(M(\tau)) \mid M \in G\}$  is dense in  $\mathbb{C}$ .

Let  $E$  be an elliptic curve with complex multiplication. Then there exists  $\tau \in H$  such that the elliptic curves  $E$  and  $\mathcal{E}_\tau$  are isomorphic and therefore

$$j(E) = j(\mathcal{E}_\tau) = j(\tau).$$

In light of Lemma 3.5, if  $M \in G$  then there is an isogeny  $E_{M(\tau)} \rightarrow E$  of odd degree. Since

$$j(E_{M(\tau)}) = j(M(\tau)),$$

the set  $J_{E, \mathrm{is}}$  of complex numbers contains  $\{j(M(\tau)) \mid M \in G\}$ , which is dense in  $\mathbb{C}$ . Hence,  $J_{E, \mathrm{is}}$  is also dense in  $\mathbb{C}$ , which ends the proof.  $\square$

**Example 3.6.** Let  $\tau \in \mathfrak{H}$ . Let  $n, m$  be positive *odd* integers. Then the matrix

$$M = \begin{pmatrix} m & 0 \\ 0 & n \end{pmatrix}$$

satisfies the conditions of Lemma 3.5. We have

$$M(\tau) = \frac{m}{n}\tau.$$

It follows from Lemma 3.5 that there exists an isogeny  $E_{\frac{m}{n}\tau} \rightarrow \mathcal{E}_\tau$  of *odd* degree. This implies that if  $\tau$  is an imaginary quadratic irrationality then, by Proposition 1.4 (to be proven in Section 6), the CM elliptic curves  $E_{\frac{m}{n}\tau}$  and  $\mathcal{E}_\tau$  have the same parity.

**Example 3.7.** Let  $\tau = \frac{1+iy}{2} \in \mathfrak{H}$  where  $y \in \mathbb{R}_+$ . Let  $n, m$  be positive odd integers. Let us put

$$\tau_{m,n} := \frac{1 + \mathbf{i}\frac{m}{n}y}{2} \in \mathfrak{H}.$$

Notice that  $n - m$  is even and therefore  $\frac{n-m}{2} \in \mathbb{Z}$ . Then the matrix

$$M = \begin{pmatrix} m & \frac{n-m}{2} \\ 0 & n \end{pmatrix}$$

satisfies the conditions of Lemma 3.5. We have

$$M(\tau) = m \cdot \frac{1 + \mathbf{i}y}{2n} + \frac{n-m}{2n} = \frac{n}{2n} + m \cdot \frac{\mathbf{i}y}{2n} = \frac{1 + \mathbf{i}\frac{m}{n}y}{2} = \tau_{m,n}.$$



It follows from Lemma 3.5 that there exists an isogeny  $\mathcal{E}_{\tau_{m,n}} \rightarrow \mathcal{E}_\tau$  of odd degree. By Proposition 1.4 (to be proven in Section 6), if  $\tau$  is an imaginary quadratic irrationality then the CM elliptic curves  $\mathcal{E}_{\tau_{m,n}}$  and  $\mathcal{E}_\tau$  have the same parity.

In what follows, if  $D$  is a negative real number then we write  $\sqrt{D}$  for  $\sqrt{|D|} \cdot \mathbf{i} \in \mathfrak{H}$ .

**Example 3.8.** If  $\tau$  is a point in  $\mathfrak{H}$  such that  $\mathbb{Q}(\tau)$  is an imaginary quadratic field and the lattice  $\Lambda_\tau$  is an order in  $\mathbb{Q}(\tau)$  then  $O_\tau = \Lambda_\tau$ . E.g., let  $D$  be a negative integer that is congruent to 1 modulo 4 and

$$\tau = \frac{1 + \sqrt{D}}{2} \in \mathfrak{H}.$$

Then  $\Lambda_\tau = \mathbb{Z} + \mathbb{Z} \frac{1 + \sqrt{D}}{2}$  is an order in the imaginary quadratic field  $\mathbb{Q}(\tau) = \mathbb{Q}(\sqrt{D})$ . By Remark 3.4(ii),

$$O_\tau = \Lambda_\tau = \mathbb{Z} + \mathbb{Z} \frac{1 + \sqrt{D}}{2}.$$

It follows from Lemma 1.1 that  $O_\tau$  is an *odd* order in the quadratic field  $\mathbb{Q}(\sqrt{D}) = \mathbb{Q}(\tau) \cong \text{End}^0(\mathcal{E}_\tau)$ . This means that the elliptic curve  $\mathcal{E}_\tau$  is odd.

Now let  $n, m$  be any positive odd integers. Let us consider the complex numbers

$$\tau_{m,n}(D) = \frac{1 + \mathbf{i} \frac{m}{n} \sqrt{|D|}}{2} = \frac{1 + \frac{m}{n} \sqrt{D}}{2} \in \mathfrak{H}. \quad (9)$$

It follows from Example 3.7 (applied to  $y = \sqrt{|D|}$ ) that all the elliptic curves  $\mathcal{E}_{\tau_{n,m}(D)}$  are also odd.

**Lemma 3.9** (Key Lemma). *Let  $\tau \in \mathfrak{H}$  such that  $\mathbb{Q}(\tau)$  is an imaginary quadratic field, i.e.,  $\mathcal{E}_\tau$  is an elliptic curve with CM.*

- (i) *If  $\text{Re}(\tau) = 0$  then  $\mathcal{E}_\tau$  is even.*
- (ii) *Suppose that  $\text{Re}(\tau) = 1/2$ . Let  $D$  be a negative integer such that  $\mathbb{Q}(\tau) = \mathbb{Q}(\sqrt{D})$  and  $D \equiv 1 \pmod{4}$ . Then*

$$\frac{1 + \sqrt{D}}{2} \in O_\tau \quad (10)$$

*if and only if there is a positive odd integer  $\beta$  such that  $\beta$  divides  $D$  and*

$$\tau = 1/2 + \frac{1}{2\beta} \sqrt{D} = \frac{1 + \sqrt{D}/\beta}{2} = \tau_{1,\beta}(D).$$

*If this is the case then  $\mathcal{E}_\tau$  is odd.*

- (iii) *Let  $D$  be a negative integer such that  $D \equiv 1 \pmod{4}$ . Let  $\beta$  be a positive odd integer and  $\mathbf{d} = \mathbf{d}_{\beta,D} \geq 1$  be the greatest common divisor of  $D$  and  $\beta^2$ . Let us put*

$$\tau := \tau_{1,\beta}(D) = \frac{1 + \frac{1}{\beta} \sqrt{D}}{2} = \frac{\beta + \sqrt{D}}{2\beta} \in \mathfrak{H}.$$

Then  $\mathcal{E}_\tau$  is odd,  $\mathbb{Q}(\tau) = \mathbb{Q}(\sqrt{D})$  is an imaginary quadratic field and the order  $O_\tau$  coincides with

$$\mathbb{Z} + \mathbb{Z} \left( \frac{\beta}{\mathbf{d}} \right) \left( \frac{\beta + \sqrt{D}}{2\beta} \right)$$

and has discriminant

$$\left( \frac{\beta}{\mathbf{d}} \right)^2 D = \frac{\beta^2}{\mathbf{d}} \cdot \frac{D}{\mathbf{d}}.$$

*Proof. (i).* Assume that  $\operatorname{Re}(\tau) = 0$ . Suppose that  $\mathcal{E}_\tau$  is odd, i.e., the order  $O_\tau$  is odd. It follows from Lemma 1.1 that there is a negative integer

$$D \equiv 1 \pmod{4}$$

such that

$$\frac{1 + \sqrt{D}}{2} \in O_\tau \subset K = \mathbb{Q}(\sqrt{D}) = \mathbb{Q} + \mathbb{Q}\sqrt{D}.$$

Since  $\tau \in \mathbb{Q}(\sqrt{D}) = \mathbb{Q} + \mathbb{Q} \cdot \sqrt{D}$  and  $\operatorname{Re}(\tau) = 0$ , there is a positive rational number  $r$  such that

$$\tau = r\sqrt{D}.$$

This implies that

$$\Lambda_\tau = \mathbb{Z} + \mathbb{Z} \cdot r\sqrt{D}.$$

It follows from the definition of  $O_\tau$  (see (4)) that

$$\frac{1 + \sqrt{D}}{2} (\Lambda_\tau) \subset \Lambda_\tau.$$

This implies that there are integers  $\alpha, \beta \in \mathbb{Z}$  such that

$$\frac{1 + \sqrt{D}}{2} \cdot 1 = \alpha + \beta r\sqrt{D}.$$

Taking the real parts of both sides, we get  $\alpha = 1/2$ , which contradicts the integrality of  $\alpha$ . The obtained contradiction proves that  $\mathcal{E}_\tau$  is even, which proves (i).

**(iii).** It follows from Example 3.8 that  $\mathcal{E}_\tau$  is odd.

Since  $\beta$  is odd,  $\beta^2 \equiv 1 \pmod{4}$ . This implies that  $4 | (\beta^2 - D)$ , because  $D \equiv 1 \pmod{4}$ . On the other hand,  $\mathbf{d} | (\beta^2 - D)$ , because both  $\beta^2$  and  $D$  are divisible by  $\mathbf{d}$ . Since  $\mathbf{d}$  divides  $D$ , it is also odd. Since odd  $\mathbf{d}$  and 4 are relatively prime, their product  $4\mathbf{d}$  also divides  $\beta^2 - D$ , i.e.,

$$c := \frac{\beta^2 - D}{4\mathbf{d}} \in \mathbb{Z}.$$

Let us put

$$a := \frac{\beta^2}{\mathbf{d}} \in \mathbb{Z}, \quad b := -\frac{\beta^2}{\mathbf{d}} = -a \in \mathbb{Z}.$$

I claim that the greatest common divisor (GCD) of three integers  $a, b, c$  is 1. Indeed, by the definition of  $\mathbf{d}$ , the integers  $\beta^2/\mathbf{d}$  and  $D/\mathbf{d}$  are relatively prime. This implies that  $a = \beta^2/\mathbf{d}$  and  $4c = (\beta^2 - D)/\mathbf{d}$  are relatively

prime. Hence,  $a$  and  $c$  are relatively prime, which implies that the greatest common divisor of  $a, b, c$  is 1. Notice also that  $a \geq 1$ .

Clearly,  $\bar{\tau} = (1 - \sqrt{D}/\beta)/2$  is the complex-conjugate of  $\tau$ . Then

$$\tau + \bar{\tau} = 1, \quad \tau\bar{\tau} = \frac{\beta^2 - D}{4\beta^2}.$$

It follows that

$$\tau^2 + (-1)\tau + \frac{\beta^2 - D}{4\beta^2} = 0.$$

Multiplying it by  $\beta^2/\mathbf{d}$ , we get

$$\frac{\beta^2}{\mathbf{d}}\tau^2 + \left(-\frac{\beta^2}{\mathbf{d}}\right)\tau + \frac{\beta^2 - D}{4\mathbf{d}} = 0,$$

i.e.,

$$a\tau^2 + b\tau + c = 0. \tag{11}$$

The properties of  $a, b, c$  mentioned above combined with (11) allow us to describe explicitly

$$O_\tau = \{u \in \mathbb{Q}(\tau), u[\tau, 1] \subset [\tau, 1]\}.$$

Namely, by [2, Ch. 2, Sect. 7.4, Lemma 1],  $O_\tau$  coincides with

$$[1, a\tau] = \mathbb{Z} + \mathbb{Z}(a\tau) = \mathbb{Z}[a\tau]$$

and its discriminant is  $b^2 - 4ac$ . In order to finish the proof of (iii), it remains to notice that

$$a\tau = \frac{\beta^2}{\mathbf{d}} \cdot \frac{\beta + \sqrt{D}}{2\beta} = \frac{\beta}{\mathbf{d}} \cdot \frac{\beta + \sqrt{D}}{2},$$

$$b^2 - 4ac = \left(-\frac{\beta^2}{\mathbf{d}}\right)^2 - 4\frac{\beta^2}{\mathbf{d}} \cdot \frac{\beta^2 - D}{4\mathbf{d}} = \left(\frac{\beta}{\mathbf{d}}\right)^2 D = \frac{\beta^2}{\mathbf{d}} \cdot \frac{D}{\mathbf{d}}.$$

(ii). Assume that  $\text{Re}(\tau) = 1/2$ . Recall that we are given the *negative* integer  $D \equiv 1 \pmod{4}$  such that

$$K = \mathbb{Q}(\tau) = \mathbb{Q}(\sqrt{D}) = \mathbb{Q} \cdot 1 \oplus \mathbb{Q} \cdot \sqrt{D} \subset \mathbb{C}.$$

Then there is a positive rational number  $r \in \mathbb{Q}$  such that

$$\tau = \frac{1}{2} + r\sqrt{D}.$$

It follows that

$$\Lambda_\tau = \mathbb{Z} \cdot 1 + \mathbb{Z} \cdot \tau = \mathbb{Z} + \mathbb{Z} \left(\frac{1}{2} + r\sqrt{D}\right) \subset \mathbb{C}.$$

By definition of  $O_\tau$  (see (4)),  $\frac{1+\sqrt{D}}{2} \in O_\tau$  if and only if

$$\frac{1 + \sqrt{D}}{2} (\Lambda_\tau) \subset \Lambda_\tau. \tag{12}$$

However, since  $\{1, 1/2 + r\sqrt{D}\}$  is a basis of the  $\mathbb{Q}$ -vector space  $\mathbb{Q}(\sqrt{D})$ , there are certain rational numbers  $\alpha, \beta, \gamma, \delta \in \mathbb{Q}$  such that

$$\frac{1 + \sqrt{D}}{2} \cdot 1 = \alpha + \beta \left( \frac{1}{2} + r\sqrt{D} \right), \quad \frac{1 + \sqrt{D}}{2} \left( \frac{1}{2} + r\sqrt{D} \right) = \gamma + \delta \left( \frac{1}{2} + r\sqrt{D} \right). \quad (13)$$

The inclusion (12) is equivalent to the condition

$$\alpha, \beta, \gamma, \delta \in \mathbb{Z}. \quad (14)$$

Opening the brackets and collecting terms in (13), we get

$$\frac{1 + \sqrt{D}}{2} = (\alpha + \beta/2) + \beta r\sqrt{D}; \quad \left( \frac{1}{4} + rD/2 \right) + \frac{1}{2} \left( r + \frac{1}{2} \right) \sqrt{D} = (\gamma + \delta/2) + \delta r\sqrt{D}.$$

Taking the real parts and “imaginary” parts, we get

$$1/2 = \alpha + \beta/2, \quad 1/2 = \beta r; \quad 1/4 + rD/2 = \gamma + \delta/2; \quad \frac{1}{2} (r + 1/2) = \delta r. \quad (15)$$

The second equality of (15) implies that

$$r = \frac{1}{2\beta}, \quad \tau = \frac{1}{2} + \frac{1}{2\beta}\sqrt{D}. \quad (16)$$

Since  $r$  is positive,  $\beta$  is also positive.

It follows from first equality of (15) that  $\alpha$  is an integer if and only if  $\beta$  is an odd integer. This implies that if  $\beta$  is not an odd integer then (14) does not hold. So, in the course of the proof we may and will assume that

$$\beta \in 1 + 2\mathbb{Z}, \quad \alpha \in \mathbb{Z}. \quad (17)$$

Now let us change the strategy and apply the already proven (iii), instead of investigating the integrality of  $\gamma$  and  $\delta$ . We get

$$O_\tau = \mathbb{Z} \cdot 1 + \mathbb{Z} \cdot \left( \frac{\beta^2}{\mathbf{d}} \right) \left( \frac{\beta + \sqrt{D}}{2\beta} \right) \quad (18)$$

where  $\mathbf{d} \geq 1$  is the GCD of  $D$  and  $\beta^2$ . We need to find when  $(1 + \sqrt{D})/2 \in O_\tau$ ?

Let us put

$$w_\beta := \frac{\beta + \sqrt{D}}{2} = \beta \cdot \frac{\beta + \sqrt{D}}{2\beta} \in \mathbb{Q}(\sqrt{D}) = \mathbb{Q}(\tau). \quad (19)$$

Since  $\beta$  is odd,

$$m_\beta := (\beta - 1)/2 \in \mathbb{Z}, \quad \beta = 2m_\beta + 1.$$

This implies that

$$w_\beta = \frac{2m_\beta + 1 + \sqrt{D}}{2} = m_\beta + \frac{1 + \sqrt{D}}{2} \in \frac{1 + \sqrt{D}}{2} + \mathbb{Z}.$$

Since  $\mathbb{Z} \subset O_\tau$ , we conclude that  $(1 + \sqrt{D})/2 \in O_\tau$  if and only if  $w_\beta \in O_\tau$ . So, in order to prove (ii), it suffices to check that  $w_\beta \in O_\tau$  if and only if  $\beta$  divides  $D$ . Let us do it.

Combining (19) and (18), we get

$$O_\tau = \mathbb{Z} \cdot 1 + \mathbb{Z} \cdot \left(\frac{\beta}{\mathbf{d}}\right) w_\beta. \quad (20)$$

Since 1 and  $w_\beta$  are obviously linearly independent over  $\mathbb{Q}$ , it follows from (20) that  $w_\beta \in O_\tau$  if and only if there exists  $n \in \mathbb{Z}$  such that

$$n \cdot \frac{\beta}{\mathbf{d}} = 1.$$

This means that the ratio  $\mathbf{d}/\beta$  is an integer, i.e.,  $\beta$  divides  $\mathbf{d}$ . Equivalently (in light of the very definition of  $\mathbf{d}$ ),  $\beta$  divides both  $D$  and  $\beta^2$ . Since  $\beta$  always divides  $\beta^2$ , we get that  $\beta$  divides  $\mathbf{d}$  if and only if  $\beta$  divides  $D$ . This ends the proof of (ii).  $\square$

**Example 3.10.** Fix a negative integer  $d$  that is congruent to 1 modulo 4 (e.g., we may take  $d = -3$ ). If  $k$  and  $n$  are positive odd integers then  $k^2n^2d$  is also a negative integer that is congruent to 1 modulo 4. If we put

$$D = k^2n^2d, \quad \beta = n^2$$

then  $\beta$  is a positive odd integer dividing  $D$ . It follows that if we put

$$\tau(n, k) := \frac{1}{2} + \frac{1}{2\beta}\sqrt{D} = \frac{1}{2} + \frac{k\sqrt{d}}{2n} \quad (21)$$

then the elliptic curve  $\mathcal{E}_{\tau(n, k)}$  is *odd*.

Recall that  $n$  and  $k$  could be any odd positive integers. Notice also that all CM elliptic curves  $\mathcal{E}_{\tau(n, k)}$  are isogenous to each other, because the imaginary quadratic field

$$\mathbb{Q}(\tau(n, k)) = \mathbb{Q}(\sqrt{d})$$

does not depend on  $n, k$ .

**Corollary 3.11.** *Let  $d$  be a square-free negative integer such that  $d \equiv 1 \pmod{4}$ . If  $n$  and  $m$  are positive odd integers and*

$$\tau = \frac{1}{2} + \frac{n}{m}\sqrt{d}$$

*then  $\mathcal{E}_\tau$  is even.*

*Proof.* In light of Example 3.7 applied to  $y = 2\sqrt{|d|}$ , it suffices to check the case  $n = m = 1$ , i.e., we may (and will) assume that

$$\tau = 1/2 + \sqrt{d}.$$

Clearly,  $K := \mathbb{Q}(\tau) = \mathbb{Q}(\sqrt{d})$ . Suppose that  $\mathcal{E}_\tau$  is *odd*, i.e., the order  $O_\tau$  is odd. Let  $\mathfrak{f}$  be the conductor of  $O_\tau$ . By Lemma 1.1,  $\mathfrak{f}$  is odd, hence,

$$\frac{1 - \mathfrak{f}}{2} \in \mathbb{Z}.$$

It follows from [2, Ch. 2, Sect. 7, Th.1] that

$$O_\tau = \mathbb{Z} + \mathfrak{f}\frac{1 + \sqrt{d}}{2}\mathbb{Z} \ni \frac{1 - \mathfrak{f}}{2} + \mathfrak{f}\frac{1 + \sqrt{d}}{2} = \frac{1 + \mathfrak{f}\sqrt{d}}{2}.$$

So,

$$\frac{1 + \mathfrak{f}\sqrt{d}}{2} \in O_\tau.$$

Notice that

$$\frac{1 + \mathfrak{f}\sqrt{d}}{2} = \frac{1 + \sqrt{D}}{2}$$

where  $D = \mathfrak{f}^2 d$  is a negative integer that is also congruent to 1 modulo 4, because  $\mathfrak{f}$  is odd. By Lemma 3.9, there is an *odd* positive integer  $\beta$  such that

$$\tau = 1/2 + \frac{1}{2\beta}\sqrt{D} = 1/2 + \frac{\mathfrak{f}}{2\beta}\sqrt{d}.$$

Since  $\tau = 1/2 + \sqrt{d}$ , we get  $\frac{\mathfrak{f}}{2\beta} = 1$ , i.e.,  $\mathfrak{f} = 2\beta$ . Since  $\mathfrak{f}$  is odd, we get the desired contradiction.  $\square$

#### 4. $j$ -INVARIANTS: THE REAL CASE

We will need the following results that either contained in [10, Ch. V, Sect. 2] (see also [6, Ch. 14, Sect. 4] and [5, Ch. 1, Sect. 4.3]) or follow readily from them.

- Let us consider the *disjoint* subsets  $\mathfrak{T}_1$  and  $\mathfrak{T}_2$  of  $\mathfrak{H}$  defined as follows.

$$\mathfrak{T}_1 := \{\mathfrak{it} \mid t \in \mathbb{R}, t \geq 1\}; \quad \mathfrak{T}_2 := \left\{\frac{1}{2} + \mathfrak{it} \mid t \in \mathbb{R}, t > 1/2\right\}.$$

Let us put

$$\mathfrak{T} := \mathfrak{T}_1 \cup \mathfrak{T}_2 \subset \mathfrak{H}.$$

If  $s$  is a real number then there is *precisely one*  $\tau \in \mathfrak{T}$  such that

$$j(\mathcal{E}_\tau) = j(\tau) = s.$$

In particular, the continuous function

$$F : \mathfrak{T} \rightarrow \mathbb{R}, \quad \tau \mapsto j(\tau) = j(\mathcal{E}_\tau)$$

is a bijective map [10, Ch. V, Sect. 2, p. 417].

- The real-valued function

$$f : [1/2, \infty) \rightarrow (-\infty, \infty), \quad y \mapsto \tau = 1/2 + iy \mapsto j(\tau) = j(\mathcal{E}_\tau)$$

is continuous injective, and

$$f(1/2) = 1728, \quad f(\sqrt{3}/2) = 0, \quad \lim_{y \rightarrow \infty} f(y) = -\infty$$

(see [10, Ch. V, Sect. 2, p. 414]). It follows that  $f$  is decreasing and the image of the map  $f$  lies in the closed semi-infinite interval  $(-\infty, 1728]$ . So, one may view as a decreasing function the map

$$f : [1/2, \infty) \rightarrow (-\infty, 1728], \quad y \mapsto \tau = 1/2 + iy \rightarrow j(\tau) = j(\mathcal{E}_\tau),$$

which is obviously bijective. In addition, the image of the open semi-infinite interval  $(1/2, \infty)$  under  $f$  is the open semi-infinite interval  $(-\infty, 1728)$ .

- The bijectiveness of the continuous map  $F : \mathfrak{T} \rightarrow \mathbb{R}$  implies that the set

$$\{j(\mathcal{E}_\tau) \mid \tau \in \mathfrak{T}_1\}$$

coincides with the closed semi-infinite interval  $[1728, \infty)$ .

*Proof of Theorem 2.4.* Without loss of generality we may and will assume that  $E = \mathcal{E}_\tau$  with  $\tau \in \mathfrak{T}$ .

Suppose that  $\mathcal{E}_\tau$  is odd. In light of Lemma 3.9(i),  $\operatorname{Re}(\tau) \neq 0$ . Since  $\tau \in \mathfrak{T}$ , we get  $\operatorname{Re}(\tau) = 1/2$ , i.e.,  $\tau = \frac{1+iy}{2}$  for some real positive  $y$ . In other words,

$$\tau \in \mathfrak{T}_2.$$

Let  $E'$  be any odd CM elliptic curve with real  $j(E')$  such that  $\mathcal{E}_\tau$  and  $E'$  are isogenous over  $\mathbb{C}$ . Without loss of generality we may assume that

$$E' = \mathcal{E}_{\tau'}, \quad j(E') = j(\mathcal{E}_{\tau'}) = j(\tau')$$

for a certain  $\tau' \in \mathfrak{T}$ . Applying Lemma 3.9(i) to  $\mathcal{E}_{\tau'}$  (instead of  $\mathcal{E}_\tau$ ), we conclude that  $\operatorname{Re}(\tau') \neq 0$  and therefore

$$\tau' \in \mathfrak{T}_2.$$

It follows that  $\mathcal{J}(\mathbb{R}, E)$  lies in the open interval  $(-\infty, 1728)$  and therefore the closure of  $\mathcal{J}(\mathbb{R}, E)$  in  $\mathbb{R}$  lies in the closed interval  $(-\infty, 1728]$ .

If  $n, m$  are any positive odd integers such that

$$\frac{m}{n}y > 1$$

then

$$\frac{m}{n}y/2 > 1/2,$$

and the set of all such  $\frac{m}{n}y/2$  is dense in  $(1/2, \infty)$ . If we put

$$\tau_{m,n} := \frac{1 + \mathbf{i}\frac{m}{n}y}{2} \in \mathfrak{T}$$

then the set of all  $\tau_{m,n}$  is a dense subset of  $\mathfrak{T}_2$ . By Example 3.7, all CM elliptic curves  $\mathcal{E}_{\tau_{m,n}}$  are odd and isogenous to  $\mathcal{E}_\tau$ . In addition,

$$f\left(\frac{m}{n}y/2\right) = j(\mathcal{E}_{\tau_{m,n}}) \in \mathcal{J}(\mathbb{R}, \mathcal{E}_\tau) \subset (-\infty, 1728].$$

It follows from the continuity of  $f$  that the set of all  $j(\mathcal{E}_{\tau_{m,n}})$  is dense in  $f((1/2, \infty]) = (-\infty, 1728)$ . This implies that the closure of  $\mathcal{J}(\mathbb{R}, \mathcal{E}_\tau)$  coincides with  $(-\infty, 1728]$  when  $\mathcal{E}_\tau$  is odd.

Suppose that  $\mathcal{E}_\tau$  is even. Then  $\mathbb{Q}(\tau) = \mathbb{Q}(\sqrt{d})$  where  $d$  is a square-free negative integer. If  $n, m$  are any positive integers such that

$$\frac{m}{n}\sqrt{|d|} > 1$$

then the CM elliptic curves  $E_{\frac{m}{n}\sqrt{d}}$  are isogenous to  $\mathcal{E}_\tau$ ; in addition, they all are even, thanks to Lemma 3.9(i). On the other hand, obviously, the set of

all such  $\frac{m}{n}\sqrt{d}$  lies in  $\mathfrak{T}_1$  (in particular, all  $j\left(E_{\frac{m}{n}\sqrt{d}}\right) \in \mathbb{R}$ ) and dense there. This implies that the closure of  $\mathcal{J}(\mathbb{R}, \mathcal{E}_\tau)$  contains  $F(\mathfrak{T}_1) = [1728, \infty)$ .

Suppose that  $d \not\equiv 1 \pmod{4}$ . Then the quadratic field  $\mathbb{Q}(\sqrt{d})$  has discriminant  $4d$ , which is even. It follows from Lemma 1.2 that every elliptic curve  $\mathcal{E}_{\tau'}$  with

$$\mathbb{Q}(\tau') = \mathbb{Q}(\sqrt{d}) = \mathbb{Q}(\tau)$$

is even and isogenous to  $\mathcal{E}_\tau$ . In particular, if  $m, n$  are any positive integers such that  $\frac{m}{n}\sqrt{|d|} > 1$  then

$$\tau(m, n) := 1/2 + \frac{m}{n}\sqrt{d} \in \mathfrak{T}_2, \quad \mathbb{Q}(\tau(m, n)) = \mathbb{Q}(\sqrt{d}),$$

which implies that the elliptic curve  $\mathcal{E}_{\tau(m, n)}$  is isogenous to  $\mathcal{E}_\tau$  and even. Since  $\tau(m, n) \in \mathfrak{T}_2$ , we get

$$j(\mathcal{E}_{\tau(m, n)}) \in \mathbb{R} \text{ and therefore } j(\mathcal{E}_{\tau(m, n)}) \in \mathcal{J}(\mathbb{R}, \mathcal{E}_\tau).$$

On the other hand, the set of all such  $\tau(m, n)$  is obviously dense in  $\mathfrak{T}_2$ . It follows that the closure of  $\mathcal{J}(\mathbb{R}, \mathcal{E}_\tau)$  contains  $f(\mathfrak{T}_1) = (-\infty, 1728)$ . Recall that we have already checked that this closure contains  $[1728, \infty)$ . It follows that the closure of  $\mathcal{J}(\mathbb{R}, \mathcal{E}_\tau)$  coincides with the whole  $\mathbb{R}$  if  $d \not\equiv 1 \pmod{4}$ .

Now suppose that  $d \equiv 1 \pmod{4}$ . If  $m, n$  are any positive odd integers such that  $\frac{m}{n}\sqrt{|d|} > 1/2$  then

$$\tau(m, n) := \frac{1}{2} + \frac{m}{n}\sqrt{d} \in \mathfrak{T}_2, \quad j(\mathcal{E}_{\tau(m, n)}) \in \mathbb{R},$$

and the elliptic curve  $\mathcal{E}_{\tau(m, n)}$  is isogenous to  $\mathcal{E}_\tau$ . By Corollary 3.11,  $\mathcal{E}_{\tau(m, n)}$  is even and therefore  $j(\mathcal{E}_{\tau(m, n)}) \in \mathcal{J}(\mathbb{R}, \mathcal{E}_\tau)$ . Clearly, the set of all such  $\tau(m, n)$  is dense in  $\mathfrak{T}_2$ . It follows that the closure of  $\mathcal{J}(\mathbb{R}, \mathcal{E}_\tau)$  contains  $F(\mathfrak{T}_2) = (-\infty, 1728)$ . Recall that we have already checked that this closure contains  $[1728, \infty)$ . It follows that the closure of  $\mathcal{J}(\mathbb{R}, \mathcal{E}_\tau)$  coincides with the whole  $\mathbb{R}$  if  $d \equiv 1 \pmod{4}$ . This ends the proof of Theorem 2.4.  $\square$

**Remark 4.1.** Let  $E_{\mathbb{R}}$  be an elliptic curve over  $\mathbb{R}$  and let  $E = E_{\mathbb{R}} \times_{\mathbb{R}} \mathbb{C}$  be its complexification. Then there is  $\tau \in \mathfrak{H}$  that enjoys the following properties (see [10, Ch. V, Sect. 5.2, Th. 2.3 and its proof]).

- (i)  $\operatorname{Re}(\tau) = 0$  or  $1/2$ .
- (ii)  $E$  is isomorphic to  $\mathcal{E}_\tau$ . In particular,

$$j(E_{\mathbb{R}}) = j(E) = j(\tau) \in \mathbb{R}.$$

- (iii)  $q := \exp(2\pi i\tau)$  is a nonzero real number such that  $|q| < 1$  and the real Lie group  $E_{\mathbb{R}}(\mathbb{R})$  of  $\mathbb{R}$ -points on  $E$  is isomorphic to the quotient  $\mathbb{R}^*/q^{\mathbb{Z}}$ . (Here  $q^{\mathbb{Z}}$  is the cyclic multiplicative subgroup of  $\mathbb{R}^*$  generated by  $q$ .) In particular,  $E_{\mathbb{R}}(\mathbb{R})$  is *connected* if and only if  $q < 0$ , i.e.,  $\operatorname{Re}(\tau) = 1/2$ .
- (iv) Suppose that  $E$  is an *odd* elliptic curve with CM. It follows from Lemma 3.9(i) that  $\operatorname{Re}(\tau) = 1/2$ . In light of (iii),  $E_{\mathbb{R}}(\mathbb{R})$  is connected.



5. ODD AND EVEN ORDERS

Let  $O$  be an order in a quadratic field  $K$  with conductor  $\mathfrak{f}$ . Lemmas 1.1 and 1.2 follow from the next two assertions.

**Lemma 5.1.** *The following conditions are equivalent.*

- (i)  $\text{tr}(O) = \mathbb{Z}$ .
- (ii)  $\text{tr}(O[1/n])$  contains 1 for all odd positive integers  $n$ .
- (iii)  $\text{discr}(O)$  is an odd integer.
- (iv) Both  $\text{discr}(K)$  and the conductor  $\mathfrak{f}_O$  are odd integers.
- (v) There exists  $w \in O$  such that  $\text{tr}(w)$  is an odd integer.
- (vi) There exists an integer  $D \equiv 1 \pmod{4}$  that is not a square such that

$$\frac{1 + \sqrt{D}}{2} \in O.$$

- (vii) There exists an integer  $D \equiv 1 \pmod{4}$  that is not a square such that

$$O = \mathbb{Z} + \mathbb{Z} \frac{1 + \sqrt{D}}{2} = \mathbb{Z} \left[ \frac{1 + \sqrt{D}}{2} \right].$$

**Lemma 5.2.** *The following conditions are equivalent.*

- (i)  $\text{tr}(O) = 2\mathbb{Z}$ .
- (ii)  $\text{tr}(O[1/n])$  does not contain 1 for all odd positive integers  $n$ .
- (iii)  $\text{discr}(O)$  is an even integer.
- (iv) Either  $\text{discr}(K)$  or the conductor  $\mathfrak{f}_O$  is an even integer.
- (v) There exists an integer  $D$  that is not a square and such that

$$O = \mathbb{Z} + \mathbb{Z} \sqrt{D} = \mathbb{Z} \left[ \sqrt{D} \right].$$

*Proof of Lemmas 5.1 and 5.2.*

**Step 0** Let us start with the computation of certain discriminants. Let  $D$  be an integer that is *not* a square. Then  $\mathbb{Q}(\sqrt{D})$  is a quadratic field. Obviously,

$$\text{tr}_{\mathbb{Q}(\sqrt{D})}(\sqrt{D}) = 0. \tag{22}$$

Let us consider the lattice (*full module* in the terminology of [2])  $\mathbb{Z} + \mathbb{Z} \sqrt{D}$  in  $\mathbb{Q}(\sqrt{D})$ . Let us compute its *discriminant*  $\text{disc}(\mathbb{Z} + \mathbb{Z} \sqrt{D})$  [2, Ch. II, Sect. 2.5].

In light of (22), the matrix of the trace form with respect to the basis  $\{1, \sqrt{D}\}$  of this lattice is

$$\begin{pmatrix} 2 & 0 \\ 0 & 2D \end{pmatrix},$$

whose determinant is  $4D$ . Hence, the discriminant

$$\text{disc}(\mathbb{Z} + \mathbb{Z} \sqrt{D}) = 4D \tag{23}$$

is even. On the other hand,  $\mathbb{Z} + \mathbb{Z} \frac{1+\sqrt{D}}{2}$  is a subgroup of index 2 in

$$\mathbb{Z} + \mathbb{Z} \sqrt{D} = \mathbb{Z} + \mathbb{Z} (1 + \sqrt{D}),$$

hence, the discriminant of  $\mathbb{Z} + \mathbb{Z} \frac{1+\sqrt{D}}{2}$  is given by the formula

$$\text{disc} \left( \mathbb{Z} + \mathbb{Z} \frac{1 + \sqrt{D}}{2} \right) = \frac{\text{disc}(\mathbb{Z} + \mathbb{Z} \sqrt{D})}{2^2} = \frac{4D}{4} = D. \quad (24)$$

Now let us discuss traces. Recall ((1) and (2)) that

$$\text{tr}(O) = \mathbb{Z} \text{ or } 2\mathbb{Z}, \quad \text{discr}(O) = \text{discr}(K) \cdot \mathfrak{f}^2. \quad (25)$$

**Step 1** Let  $n$  be an odd positive integer. Then

$$\text{tr}(O[1/n]) = \mathbb{Z}[1/n] \cdot \text{tr}(O) \subset \mathbb{Q}.$$

It follows that

$$\text{tr}(O[1/n]) = \mathbb{Z}[1/n] \ni 1 \text{ if } \text{tr}(O) = \mathbb{Z};$$

$$\text{tr}(O[1/n]) = 2 \cdot \mathbb{Z}[1/n] \not\ni 1 \text{ if } \text{tr}(O) = 2\mathbb{Z}.$$

Now the equivalence of (i) and (ii) follows from first equality of (25) (for both Lemmas). The equivalence of (iii) and (iv) follows from second equality of (25) (for both Lemmas).

**Step 2** Let us concentrate for a while on the proof of Lemma 5.1. The equivalence of (i) and (v) follows from first equality of (25). In addition, (vii) obviously implies (vi) while (vi) implies (v), because if we put  $w = \frac{1+\sqrt{D}}{2}$  then

$$\text{tr}(w) = \text{tr} \left( \frac{1 + \sqrt{D}}{2} \right) = \frac{1 + \sqrt{D}}{2} + \frac{1 - \sqrt{D}}{2} = 1$$

is an odd integer. In light of (24), (vii) implies that  $\text{disc}(O) = D$ , which is odd. Therefore (vii) implies (iii).

So, in order to finish the proof of Lemma 5.1, it suffices to check that

- (1o) (i) implies (vii);
- (2o) (iii) implies (i).

Now let us switch to Lemma 5.2. Clearly, (v) implies (i). On the other hand, in light of (23), (v) implies that  $\text{disc}(O) = 4D$ , which is even. Therefore (v) implies (iii).

So, in order to finish the proof of Lemma 5.2, it suffices to check that

- (1e) (i) implies (v);
- (2e) (iii) implies (i).

**Step 3** We have  $K = \mathbb{Q}(\sqrt{d})$  where  $d \neq 1$  is a square-free integer. Clearly,

$$\text{tr}(\sqrt{d}) = 0. \quad (26)$$

Suppose that  $d \not\equiv 1 \pmod{4}$ . It is known [2, Ch. 2, Sect. 7, Th.1] that

$$\text{disc}(K) = 4d, \quad O = \mathbb{Z} + \mathfrak{f}\sqrt{d} \mathbb{Z} = \mathbb{Z} + \sqrt{\mathfrak{f}^2 d} \mathbb{Z}, \quad \text{disc}(O) = \text{disc}(K)\mathfrak{f}^2 = 4d\mathfrak{f}^2.$$

In particular,  $\text{disc}(O)$  is even. On the other hand, if we put  $D := \mathfrak{f}^2 d$  then  $D$  is not a square (because  $d$  is square-free) and

$$K = \mathbb{Q}(\sqrt{d}) = \mathbb{Q}(\sqrt{D}), \quad O = \mathbb{Z} + \sqrt{D} \mathbb{Z}, \quad \text{disc}(O) = 4d\mathfrak{f}^2 = 4D.$$

It follows from (26) that  $\text{tr}(O) = 2\mathbb{Z}$ . This implies that  $O$  enjoys properties (i), (iii), (v) of Lemma 5.2. On the other hand,  $O$  does *not* enjoy any of properties (i) and (iii) of Lemma 5.1. In light of Step 2, this proves both Lemmas in the (last remaining) case when  $d \not\equiv 1 \pmod{4}$ .

Suppose that  $d \equiv 1 \pmod{4}$ . It is known [2, Ch. 2, Sect. 7, Th.1] that

$$\text{disc}(K) = d, \quad O = \mathbb{Z} + \mathfrak{f} \frac{1 + \sqrt{d}}{2} \mathbb{Z}, \quad \text{disc}(O) = \text{disc}(K)\mathfrak{f}^2 = d\mathfrak{f}^2.$$

It follows from (26) that

$$\text{tr}(O) = \mathbb{Z} \text{ if } \mathfrak{f} \text{ is odd; } \quad \text{tr}(O) = 2\mathbb{Z} \text{ if } \mathfrak{f} \text{ is even.} \quad (27)$$

If  $\mathfrak{f}$  is even then  $\text{tr}(O) = 2\mathbb{Z}$ ,  $\text{disc}(O)$  is even,  $m = \mathfrak{f}/2$  is a positive integer, and

$$\begin{aligned} O &= \mathbb{Z} + \mathfrak{f} \frac{1 + \sqrt{d}}{2} \mathbb{Z} = \mathbb{Z} + m (1 + \sqrt{d}) \mathbb{Z} = \mathbb{Z} + m\sqrt{d} \mathbb{Z} = \\ &\quad \mathbb{Z} + \sqrt{m^2 d} \mathbb{Z} = \mathbb{Z} + \sqrt{D} \mathbb{Z} \end{aligned}$$

where  $D := m^2 d$ , which is not a square, because  $d$  is square-free. We get

$$K = \mathbb{Q}(\sqrt{d}) = \mathbb{Q}(\sqrt{D}), \quad O = \mathbb{Z} + \sqrt{D} \mathbb{Z}, \quad \text{disc}(O) = \mathfrak{f}^2 d = (2m)^2 d = 4D.$$

This implies that  $O$  enjoys all the properties (i), (iii), (v) of Lemma 5.2. On the other hand,  $O$  does *not* enjoy any of properties (i) and (iii) of Lemma 5.1. In light of Step 2, this proves both Lemmas in the case when  $d \not\equiv 1 \pmod{4}$  and  $\mathfrak{f}$  is even.

Now suppose that  $\mathfrak{f}$  is *odd*. It follows from (27) that  $\text{tr}(O) = \mathbb{Z}$  and  $\text{disc}(O)$  is odd. In addition, if we put  $m := (\mathfrak{f} - 1)/2 \in \mathbb{Z}$  then  $\mathfrak{f} = 2m + 1$  and

$$\begin{aligned} O &= \mathbb{Z} + \mathfrak{f} \frac{1 + \sqrt{d}}{2} \mathbb{Z} = \mathbb{Z} + \frac{\mathfrak{f} + \mathfrak{f}\sqrt{d}}{2} \mathbb{Z} = \mathbb{Z} + \left( m + \frac{1 + \mathfrak{f}\sqrt{d}}{2} \right) \mathbb{Z} = \\ &\quad \mathbb{Z} + \frac{1 + \mathfrak{f}\sqrt{d}}{2} \mathbb{Z} = \mathbb{Z} + \frac{1 + \sqrt{\mathfrak{f}^2 d}}{2} \mathbb{Z} = \mathbb{Z} + \frac{1 + \sqrt{D}}{2} \mathbb{Z} \end{aligned}$$

where  $D := \mathfrak{f}^2 d$  is not a square, because  $d$  is square-free. Since  $\mathfrak{f}$  is odd,  $\mathfrak{f}^2 \equiv 1 \pmod{4}$ . Since  $d \equiv 1 \pmod{4}$ , the product  $D := \mathfrak{f}^2 d$  is also congruent to 1 modulo 4. We get

$$K = \mathbb{Q}(\sqrt{d}) = \mathbb{Q}(\sqrt{D}), \quad O = \mathbb{Z} + \frac{1 + \sqrt{D}}{2} \mathbb{Z}, \quad \text{disc}(O) = \mathfrak{f}^2 d = D.$$

This implies that  $O$  enjoys properties (i), (iii), (vii) of Lemma 5.1. On the other hand,  $O$  does *not* enjoy any of properties (i) and (iii) of Lemma 5.2. In light of Step 2, this proves both Lemmas in the case when  $d \equiv 1 \pmod{4}$  and  $\mathfrak{f}$  is odd. This ends the proof of both Lemmas.  $\square$

**Remark 5.3.** Let  $n$  be a positive odd integer,  $\phi : K_1 \rightarrow K_2$  an isomorphism of quadratic fields, and  $O_1$  and  $O_2$  are orders in  $K_1$  and  $K_2$  respectively. Suppose that

$$\phi(O_1[1/n]) = O_2[1/n].$$

Then

$$\mathrm{tr}_{K_1/\mathbb{Q}}(O_1[1/n]) = \mathrm{tr}_{K_2/\mathbb{Q}}(O_2[1/n]).$$

It follows from Lemmas 5.1(ii) and 5.2(ii) that  $O_1$  and  $O_2$  have the same parity, i.e., either they both are odd or both even.

**Remark 5.4.** Recall that if  $f$  is an odd integer then  $f^2 \equiv 1 \pmod{4}$ . On the other hand, if  $f$  is an even integer then  $f^2 \equiv 0 \pmod{4}$ .

Let  $O$  be an order in a quadratic field  $K$ . Now the explicit formulas for its discriminant (see Step 3 above) imply the following.

- (e)  $O$  is even if and only if  $\mathrm{disc}(O) \equiv 0 \pmod{4}$ . If this is the case then there is an integer  $D$  that is not a square and such that

$$K = \mathbb{Q}(\sqrt{D}), \quad O = \mathbb{Z} + \mathbb{Z}\sqrt{D}, \quad \mathrm{disc}(O) = 4D.$$

- (o)  $O$  is odd if and only if  $\mathrm{disc}(O) \equiv 1 \pmod{4}$ . If this is the case then there is an integer  $D \equiv 1 \pmod{4}$  that is not a square and such that

$$K = \mathbb{Q}(\sqrt{D}), \quad O = \mathbb{Z} + \mathbb{Z}\frac{1 + \sqrt{D}}{2}, \quad \mathrm{disc}(O) = D.$$

The following assertion is a natural complement to Key Lemma 3.9(ii).

**Lemma 5.5.** Let  $D$  be a negative integer such that  $D \equiv 1 \pmod{4}$ . Let  $\beta$  be a positive integer that divides  $D$ . Let us put

$$\tau = 1/2 + \frac{1}{2\beta}\sqrt{D} \in \mathfrak{K}.$$

Then

$$O_\tau = \mathbb{Z} \left[ \frac{1 + \sqrt{D}}{2} \right] = \mathbb{Z} + \mathbb{Z}\frac{1 + \sqrt{D}}{2}$$

if and only if  $\beta$  and  $D/\beta$  are relatively prime.

*Proof.* Since  $D$  is odd, its divisor  $\beta$  is also odd. In light of Key Lemma 3.9(ii),

$$\mathcal{O}_D := \mathbb{Z} + \mathbb{Z}\frac{1 + \sqrt{D}}{2} \subset O_\tau.$$

We need to prove that the equality holds if and only if  $\beta$  and  $D/\beta$  are relatively prime. By Key Lemma 3.9(iii),

$$\mathrm{disc}(O_\tau) = (\beta/\mathbf{d})^2 D$$

where  $\mathbf{d}$  is the GCD of  $\beta^2$  and  $D$ .

Notice that  $\mathcal{O}_D$  is a subgroup of finite index, say,  $f \geq 1$  in  $O_\tau$ . Clearly,  $\mathcal{O}_D = O_\tau$  if and only if  $f = 1$ . However,

$$D = \mathrm{disc}(\mathcal{O}_D) = f^2 \mathrm{disc}(O_\tau) = f^2 (\beta/\mathbf{d})^2 D,$$

i.e.,

$$f^2(\beta/\mathbf{d})^2 = 1,$$

which means that  $f = \mathbf{d}/\beta$  (recall that all  $f, \beta, \mathbf{d}$  are positive integers). It follows that  $f = 1$  if and only if  $\mathbf{d} = \beta$ .

Since  $\mathbf{d}$  is the GCD of  $\beta^2$  and  $D$ , the integers  $\beta^2/\mathbf{d}$  and  $D/\mathbf{d}$  are relatively prime. So, if  $\mathbf{d} = \beta$  then  $\beta = \beta^2/\beta$  and  $D/\beta$  are relatively prime.

Conversely, suppose that the integers  $\beta$  and  $\tilde{\beta} = D/\beta$  are relatively prime. This implies that the GCD of  $\beta^2 (= \beta \cdot \beta)$  and  $D (= \beta \cdot \tilde{\beta})$  is  $\beta$ , i.e.,  $\mathbf{d} = \beta$ . This ends the proof.  $\square$

## 6. ISOGENIES OF ELLIPTIC CURVES

*Proof of Proposition 1.4.* There is a (dual) isogeny  $\psi : E_2 \rightarrow E_1$  such that  $\psi \circ \phi : E_1 \rightarrow E_1$  is multiplication by  $n$  in  $E_1$  and  $\phi \circ \psi : E_2 \rightarrow E_2$  is multiplication by  $n$  in  $E_2$  [5, p. 7]. Then we get an isomorphism of  $\mathbb{Q}$ -algebras

$$\Phi : \text{End}^0(E_1) = \text{End}(E_1) \otimes \mathbb{Q} \rightarrow \text{End}(E_2) \otimes \mathbb{Q} = \text{End}^0(E_2), \quad u_1 \mapsto \frac{1}{n}\phi u_1 \psi,$$

whose inverse is an isomorphism of  $\mathbb{Q}$ -algebras

$$\Psi : \text{End}^0(E_2) = \text{End}(E_2) \otimes \mathbb{Q} \rightarrow \text{End}(E_1) \otimes \mathbb{Q} = \text{End}^0(E_1), \quad u_2 \mapsto \frac{1}{n}\psi u_2 \phi.$$

Clearly, both  $\Phi$  and  $\Psi$  sends 1 to 1, sums to sums and therefore are homomorphisms of  $\mathbb{Q}$ -vector spaces. On the other hand if  $u_1 \in \text{End}^0(E_1)$  then

$$\Psi(\Phi(u_1)) = \frac{1}{n}\psi\Phi(u_1)\phi = \frac{1}{n}\psi\left(\frac{1}{n}\phi u_1 \psi\right)\phi = \frac{1}{n}\left(\frac{1}{n}\psi\phi\right)u_1\left(\frac{1}{n}\psi\phi\right) = u_1.$$

Since both  $\text{End}^0(E_1)$  and  $\text{End}^0(E_2)$  are  $\mathbb{Q}$ -vector spaces of the same dimension 2,  $\Phi$  is an isomorphism of  $\mathbb{Q}$ -vector spaces and  $\Psi$  is its inverse, hence, also an isomorphism of  $\mathbb{Q}$ -vector spaces. The only thing that remains to check is that both  $\Phi$  and  $\Psi$  are compatible with multiplication. Let us check it. If  $u_1, v_1 \in \text{End}^0(E_1)$  then

$$\Phi(u_1)\Phi(v_1) = \frac{1}{n}\phi u_1 \psi \frac{1}{n}\phi v_1 \psi = \frac{1}{n}\phi u_1 \frac{1}{n}(\psi\phi)v_1 \psi = \frac{1}{n}\phi u_1 v_1 \psi = \Phi(u_1 v_1).$$

This proves that  $\Phi$  is a homomorphism of  $\mathbb{Q}$ -algebras and therefore is an isomorphism of quadratic fields  $\text{End}^0(E_1)$  and  $\text{End}^0(E_2)$ . Hence, its inverse  $\Psi$  is also a field isomorphism.

Now I claim that

$$\Phi(\text{End}(E_1)[1/n]) = \text{End}(E_2)[1/n], \quad \Psi(\text{End}(E_2)[1/n]) = \text{End}(E_1)[1/n]. \tag{28}$$

Indeed, it follows from the definition of  $\Phi$  and  $\Psi$  that

$$\Phi(\text{End}(E_1)) \subset \frac{1}{n}\text{End}(E_2) \subset \text{End}(E_2)[1/n],$$

$$\Psi(\text{End}(E_2)) \subset \frac{1}{n}\text{End}(E_1) \subset \text{End}(E_1)[1/n].$$

It follows that

$$\Phi(\text{End}(E_1)[1/n]) \subset \text{End}(E_2)[1/n], \quad \Psi(\text{End}(E_2)[1/n]) \subset \text{End}(E_1)[1/n]$$

and therefore

$$\begin{aligned} \text{End}(E_1)[1/n] &= \Psi(\Phi(\text{End}(E_1)[1/n])) \subset \Psi(\text{End}(E_2)[1/n]), \\ \text{End}(E_2)[1/n] &= \Phi(\Psi(\text{End}(E_2)[1/n])) \subset \Phi(\text{End}(E_1)[1/n]). \end{aligned}$$

We get

$$\begin{aligned} \text{End}(E_1)[1/n] &\subset \Psi(\text{End}(E_2)[1/n]) \subset \text{End}(E_1)[1/n], \\ \text{End}(E_2)[1/n] &\subset \Phi(\text{End}(E_1)[1/n]) \subset \text{End}(E_2)[1/n], \end{aligned}$$

which proves (28).

So,  $\Phi$  is an isomorphism of quadratic fields  $\text{End}^0(E_1)$  and  $\text{End}^0(E_2)$  such that

$$\Phi(\text{End}(E_1)[1/n]) = \text{End}(E_2)[1/n].$$

Now the desired result follows from Remark 5.3.  $\square$

*Proof of Proposition 1.5.* We may assume that

$$E_1 = E_{1,\mathbb{R}} \times_{\mathbb{R}} \mathbb{C}, \quad E_2 = E_{2,\mathbb{R}} \times_{\mathbb{R}} \mathbb{C}.$$

Let  $\phi : E_1 \rightarrow E_2$  be an isogeny and  $\bar{\phi} : E_1 \rightarrow E_2$  its “complex-conjugate”. If  $\bar{\phi} = \phi$  then  $\phi$  could be descended to an isogeny  $E_{1,\mathbb{R}} \rightarrow E_{2,\mathbb{R}}$  over  $\mathbb{R}$ .

Suppose that  $\bar{\phi} \neq \phi$ . Then its difference  $\psi := \bar{\phi} - \phi : E_1 \rightarrow E_2$  is a nonzero homomorphism of elliptic curves and therefore is an isogeny. Its complex-conjugate  $\bar{\psi} = \overline{\bar{\phi} - \phi}$  coincides with  $\phi - \bar{\phi} = -\psi$ . On the other hand, it is well known that *not* all endomorphisms of the CM elliptic curve  $E_1$  are defined over  $\mathbb{R}$ , i.e., there is an endomorphism  $\alpha$  of  $E_1$  such that its complex-conjugate  $\bar{\alpha}$  does *not* coincide with  $\alpha$  (see Lemma 6.1 below). This means that  $\beta = \bar{\alpha} - \alpha : E_1 \rightarrow E_1$  is a nonzero endomorphism of  $E_1$ , i.e., is an isogeny. Clearly, the complex conjugate  $\bar{\beta}$  of  $\beta$  coincides with

$$\overline{\bar{\alpha} - \alpha} = \alpha - \bar{\alpha} = -\beta.$$

So,

$$\bar{\psi} = -\psi, \quad \bar{\beta} = -\beta.$$

This implies that the composition  $\lambda := \psi \circ \beta : E_1 \rightarrow E_2$  is an isogeny, whose complex conjugate  $\bar{\lambda} = \bar{\psi} \circ \bar{\beta}$  equals

$$\bar{\psi} \circ \bar{\beta} = (-\psi) \circ (-\beta) = \psi \circ \beta = \lambda.$$

So, the nonzero homomorphism  $\lambda$  coincides with its “complex-conjugate”, i.e., is defined over  $\mathbb{R}$ . This means that there is a  $\mathbb{R}$ -homomorphism  $\lambda_{\mathbb{R}} : E_{1,\mathbb{R}} \rightarrow E_{2,\mathbb{R}}$  of elliptic curves, whose “complexification”  $E_1 \rightarrow E_2$  coincides with  $\lambda$ . Since  $\lambda \neq 0$ , we obtain that  $\lambda_{\mathbb{R}} \neq 0$  and therefore is an *isogeny*.  $\square$

The following assertion (and its proof that I am going to reproduce) is pretty well known but I was unable to find a reference.

**Lemma 6.1.** *Let  $E_{\mathbb{R}}$  be an elliptic curve over  $\mathbb{R}$ . Then the ring  $\text{End}_{\mathbb{R}}(E_{\mathbb{R}})$  of  $\mathbb{R}$ -endomorphisms of  $E_{\mathbb{R}}$  is  $\mathbb{Z}$  and the corresponding endomorphism algebra*

$$\text{End}_{\mathbb{R}}^0(E_{\mathbb{R}}) := \text{End}_{\mathbb{R}}(E_{\mathbb{R}}) \otimes \mathbb{Q}$$

is  $\mathbb{Q}$ .

*Proof.* Let us consider the complexification  $E = E_{\mathbb{R}} \times_{\mathbb{R}} \mathbb{C}$  of  $E_{\mathbb{R}}$ . We have

$$\mathbb{Z} \subset \text{End}_{\mathbb{R}}(E_{\mathbb{R}}) \subset \text{End}(E), \quad \mathbb{Q} \subset \text{End}_{\mathbb{R}}^0(E_{\mathbb{R}}) \subset \text{End}^0(E).$$

If  $\text{End}_{\mathbb{R}}(E_{\mathbb{R}}) \neq \mathbb{Z}$  then

$$\text{End}(E) \neq \mathbb{Z}, \quad \text{End}^0(E) \neq \mathbb{Q},$$

i.e.,  $E$  is an elliptic curve with CM and  $K := \text{End}^0(E)$  is an imaginary quadratic field. This implies that

$$K = \text{End}^0(E) = \text{End}_{\mathbb{R}}^0(E_{\mathbb{R}})$$

and  $\text{End}_{\mathbb{R}}(E_{\mathbb{R}})$  is an order in the quadratic field  $K$ . Let  $\Omega_{\mathbb{R}}^1(E_{\mathbb{R}})$  be the space of differentials of the first kind on  $E_{\mathbb{R}}$ , which is a  $\mathbb{R}$ -vector space of dimension 1. By functoriality,  $\text{End}_{\mathbb{R}}(E_{\mathbb{R}})$  acts on  $\Omega_{\mathbb{R}}^1(E_{\mathbb{R}})$ , which gives us the injective ring homomorphism

$$\delta : \text{End}_{\mathbb{R}}(E_{\mathbb{R}}) \hookrightarrow \text{End}_{\mathbb{R}}(\Omega_{\mathbb{R}}^1(E_{\mathbb{R}})) = \mathbb{R}$$

[8, Ch. I, Sect. 2.8]. By  $\mathbb{Q}$ -linearity,  $\delta$  extends to the homomorphism of  $\mathbb{Q}$ -algebras

$$\delta_{\mathbb{Q}} : K = \text{End}_{\mathbb{R}}^0(E_{\mathbb{R}}) = \text{End}_{\mathbb{R}}(E_{\mathbb{R}}) \otimes \mathbb{Q} \rightarrow \text{End}_{\mathbb{R}}(\Omega_{\mathbb{R}}^1(E_{\mathbb{R}})) = \mathbb{R},$$

which is also injective (since  $K$  is a field). So,  $\delta_{\mathbb{Q}} : K \rightarrow \mathbb{R}$  is a field embedding. But such an embedding does *not* exist, since  $K$  is an *imaginary* quadratic field. The obtained contradiction proves the desired result.  $\square$

## 7. ODD ELLIPTIC CURVES REVISITED

In light of Lemma 5.5, it is natural to “classify” (count the number of) positive divisors  $\beta$  of the discriminant  $D$  that are relatively prime to  $D/\beta$ . Let us start with the following definition (notation).

**Definition 7.1** (Definition-Notation). Let  $n$  be a nonzero integer. We write  $P_n$  for the (finite) set of prime divisors of  $n$ . By Main Theorem of Arithmetic,

$$n = \pm \prod_{p \in P_n} p^{e_{n,p}}$$

where  $e_{n,p}$  are certain positive integers uniquely determined by  $n$ .

Let us call a divisor  $r$  of  $n$  a *saturated* divisor if the integers  $r$  and  $n/r$  are relatively prime.

Clearly,  $r$  is a saturated divisor of  $n$  if and only if  $|n|/r$  is one. It is also clear that  $r$  is a saturated divisor of  $n$  if and only if  $-r$  is also one.

The following assertion is an easy exercise in elementary number theory that will be proven at the end of this section.

**Proposition 7.2.** (i) *Let  $S$  be a subset of  $P_n$ . Then both integers*

$$\mathbf{r}_{n,S} = \prod_{p \in S} p^{e_{n,p}}$$

*and  $-\mathbf{r}_{n,S}$  are saturated divisors of  $n$ . (As usual, if  $S$  is the empty set then  $r_{n,\emptyset} = 1$ .) Conversely, if  $c$  is a saturated divisor of  $n$  then  $-c$  is also a saturated divisor of  $n$  and*

$$c = \pm \prod_{p \in P_c} p^{e_{n,p}} = \pm \mathbf{r}_{n,P_c}.$$

- (ii) *The number of positive saturated divisors of  $n$  is the number of subsets of  $P_n$ , i.e.,  $2^{\#(P_n)}$  where  $\#(P_n)$  is the cardinality of  $P_n$ , i.e., the number of prime divisors of  $n$ .*
- (iii) *If  $S_1$  and  $S_2$  are disjoint subsets of  $P_n$  then  $\mathbf{r}_{n,S_1}$  and  $\mathbf{r}_{n,S_2}$  are relatively prime and*

$$\mathbf{r}_{n,S_1 \cup S_2} = \mathbf{r}_{n,S_1} \mathbf{r}_{n,S_2}.$$

- (iv) *Suppose that  $n$  is a negative integer that is congruent to 1 modulo 4. Then the number of positive saturated divisors  $r$  of  $n$  with  $r < \sqrt{|n|}$  equals  $2^{\#(P_n)-1}$ .*

**Theorem 7.3.** *Let  $D$  be a negative integer that is congruent to 1 modulo 4. Let  $s_D$  be the number of prime divisors of  $D$ .*

- (1) *Let  $E$  be an elliptic curve over  $\mathbb{C}$  with  $j(E) \in \mathbb{R}$ . Then the following two conditions are equivalent.*
- (i)  *$E$  has CM and its endomorphism ring  $\text{End}(E)$  is an order of discriminant  $D$ . (In particular,  $E$  is odd.)*
- (ii) *There exists a positive saturated divisor  $\beta$  of  $D$  such that  $\beta < \sqrt{|D|}$  and*

$$j(E) = j(\tau) \text{ where } \tau = \frac{1}{2} + \frac{\sqrt{D}}{2\beta} = \frac{1}{2} + \frac{\sqrt{|D|}}{2\beta} \mathbf{i} \in \mathfrak{T}_2 \subset \mathfrak{H}.$$

- (2) *Let  $\mathcal{J}_D(\mathbb{R})$  be the set of  $j$ -invariants of all  $E$  that enjoy the equivalent properties (i)-(ii). Then  $\mathcal{J}_D(\mathbb{R})$  consists of  $2^{s_D-1}$  distinct real numbers and lies in the semi-open interval  $\left[ j\left(\frac{1+\sqrt{D}}{2}\right), 1728 \right)$ .*

*Proof.* (1) Suppose that  $E$  enjoys the properties (i). Since  $j(E) \in \mathbb{R}$ , there is (precisely) one  $\tau \in \mathfrak{T}$  such that  $E$  is isomorphic to  $\mathcal{E}_\tau$  and  $j(E) = j(\tau)$ . So, we may assume that  $E = \mathcal{E}_\tau$ . In particular,  $\mathcal{E}_\tau$  is odd. It follows from Key Lemma 3.9(i) that  $\text{Re}(\tau) \neq 0$ , i.e.,

$$\tau \in \mathfrak{T}_2, \text{Re}(\tau) = \frac{1}{2}.$$



Since  $O = \text{End}(\mathcal{E}_\tau) \cong O_\tau \subset \mathbb{C}$  is an order with *odd* discriminant  $D$ , it follows from Remark 5.4 that

$$O_\tau = \mathbb{Z} \left[ \frac{1 + \sqrt{D}}{2} \right] = \mathbb{Z} + \mathbb{Z} \frac{1 + \sqrt{D}}{2} = \mathcal{O}_D \subset \mathbb{C}. \quad (29)$$

Combining Lemmas 3.9 and 5.5, we conclude that (29) holds if and only if there is a saturated divisor  $\beta$  of  $D$  (each divisor of  $D$  is obviously odd) such that

$$\tau = \frac{1}{2} + \frac{\sqrt{D}}{2\beta} = \frac{1}{2} + \frac{\sqrt{|D|}}{2\beta}.$$

Since  $\tau \in \mathfrak{T}_2$ , the imaginary part of  $\tau$  is strictly greater than  $1/2$ , i.e.,

$$\beta < \sqrt{|D|}.$$

This proves the equivalence of (i) and (ii).

(2) In order to prove the second assertion of our theorem, recall that the function

$$[1/2, \infty] \rightarrow \mathbb{R}, \quad t \mapsto j(1/2 + \mathbf{i}t)$$

is *strictly* decreasing and  $j(1/2 + \mathbf{i}/2) = 1728$ . Now the desired result follows readily from Proposition 7.2(iv).  $\square$

*Proof of Proposition 7.2.*

(i) Let  $S' = P_n \setminus S$ . Then

$$n = \pm \mathbf{r}_{n,S} \mathbf{r}_{n,S'}, \quad \frac{n}{\mathbf{r}_{n,S}} = \pm \mathbf{r}_{n,S'}.$$

If  $p$  is a prime divisor of both  $\mathbf{r}_{n,S}$  and  $\mathbf{r}_{n,S'}$  then it must belong to both sets  $S$  and  $S'$ . However,  $S$  and  $S'$  do *not* meet each other. Hence,  $\mathbf{r}_{n,S}$  and  $\mathbf{r}_{n,S'} = \pm n/\mathbf{r}_{n,S}$  have no common prime factors and therefore are relatively prime. This implies that the divisor  $\mathbf{r}_{n,S}$  (and therefore  $-\mathbf{r}_{n,S}$ ) of  $n$  is saturated.

Conversely, suppose that  $c$  is a saturated divisor of  $n$ . Then  $P_c \subset P_n$  and

$$c = \pm \prod_{p \in P_c} p^{e_{c,p}}, \quad 1 \leq e_{c,p} \leq e_{n,p} \quad \forall p \in P_c.$$

I claim that

$$e_{c,p} = e_{n,p} \quad \forall p \in P_c, \quad \text{i.e.,} \quad d = \pm \mathbf{r}_{P_c}.$$

Indeed, suppose that  $e_{c,p} < e_{n,p}$  for some  $p \in P_c$ . Then  $p$  divides both  $c$  and  $n/c$ , which contradicts the saturatedness of  $c$ . This ends the proof.

(ii) It follows from (i) that the set of positive saturated divisors of  $n$  coincides with

$$\{\mathbf{r}_{n,S} \mid S \subset P_n\}.$$

It is also clear that if  $S_1$  and  $S_2$  are distinct subsets of  $P_n$  then  $\mathbf{r}_{n,S_1} \neq \mathbf{r}_{n,S_2}$ , because their sets of prime divisors (i.e.,  $S_1$  and  $S_2$ ) do *not* coincide. So, the number of positive saturated divisors of  $n$  equals the number of subsets of  $P_n$ , which is  $2^{\#(P_n)}$ .

(iii) Since  $S_1$  (resp.  $S_2$ ) is the set of prime divisors of  $\mathbf{r}_{n,S_1}$  (resp.  $\mathbf{r}_{n,S_2}$ ) and  $S_1$  does *not* meet  $S_2$ , the integers  $\mathbf{r}_{n,S_1}$  and  $\mathbf{r}_{n,S_2}$  have no common prime factors and therefore are relatively prime. On the other hand,

$$\mathbf{r}_{n,S_1 \cup S_2} = \prod_{p \in S_1 \cup S_2} p^{e_{n,p}} = \left( \prod_{p \in S_1} p^{e_{n,p}} \right) \left( \prod_{p \in S_2} p^{e_{n,p}} \right) = \mathbf{r}_{n,S_1} \mathbf{r}_{n,S_2}.$$

(iv) Since  $n \equiv 1 \pmod{4}$ , the positive odd integer  $|n| = (-n)$  is congruent to 3 modulo 4 and therefore is *not* a square. Hence, if  $\beta$  is a positive saturated divisor of  $n$  then  $|n|/\beta$  is also a positive saturated divisor of  $n$  while  $\beta \neq |n|/\beta$ . Clearly, there is precisely one element in the pair  $\{\beta, |n|/\beta\}$  that is strictly less than  $\sqrt{|n|}$ . In light of (ii), the number of positive saturated divisors of  $n$  that satisfy this inequality is  $2^{\#(P_n)}/2 = 2^{\#(P_n)-1}$ .  $\square$

#### REFERENCES

- [1] E. Artin and G. Whaples, *Axiomatic characterization of fields by the product formula for valuations*. Bull. Amer. Math. Soc. **51:7** (1945), 469–492. Reprinted in: Exposition by Emil Artin: A selection. American Mathematical Society, Providence, RI, 2007.
- [2] Z.I. Borevich, I.R. Shafarevich, *Number Theory*. Academic Press, New York and London, 1966.
- [3] J.-L. Colliot-Thélène and A. Pirutka, *Certaines fibrations en surfaces quadriques réelles*. arXiv:2406.00463 [math.AG].
- [4] A. Knapp, *Elliptic Curves*. Princeton University Press, Princeton, NJ, 1992.
- [5] B. Gross, *Arithmetic on elliptic curves with complex multiplication*. Lecture Notes in Math. **776**, Springer-Verlag, Berlin Heidelberg New York, 1980.
- [6] D. Husemöller, *Elliptic Curves*, 2nd edition. GTM **111**, Springer-Verlag, New York, 2004.
- [7] G. Shimura, *Introduction to the arithmetic theory of automorphic functions*. Princeton University Press, Princeton, NJ, 1971.
- [8] G. Shimura, *Abelian varieties with Complex Multiplication and Modular Functions*. Princeton University Press, Princeton, NJ, 1998.
- [9] J. Silverman, *Arithmetic of elliptic curves*, 2nd edition. GTM **106**, Springer-Verlag, New York, 2009.
- [10] J. Silverman, *Advanced topics in the arithmetic of elliptic curves*. GTM **151**, Springer-Verlag, New York, 1994.

DEPARTMENT OF MATHEMATICS, PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK, PA 16802, USA

*Email address:* zarhin@math.psu.edu