# ANY TOPOLOGICAL RECURSION ON A RATIONAL SPECTRAL CURVE IS KP INTEGRABLE

### A. ALEXANDROV, B. BYCHKOV, P. DUNIN-BARKOWSKI, M. KAZARIAN, AND S. SHADRIN

ABSTRACT. We prove that for any initial data on a genus zero spectral curve the corresponding correlation differentials of topological recursion are KP integrable. As an application we prove KP integrability of partition functions associated via ELSV-type formulas to the *r*-th roots of the twisted powers of the log canonical bundles.

#### CONTENTS

| 1. Introduction                           | 1  |
|---|----|
| 1.1. Main theorem                         | 2  |
| 1.2. Background                           | 3  |
| 1.3. Applications                         | 4  |
| 1.4. Organization of the paper            | 4  |
| 1.5. Notation                             | 4  |
| 1.6. Acknowledgments                      | 4  |
| 2. Global KP symmetries                   | 5  |
| 3. Proof of Theorem 1.1                   | 7  |
| 4. Generalized setting                    | 8  |
| 5. Application: <i>r</i> -th roots        | 9  |
| Appendix A. Variation of TR differentials | 10 |
| References                                | 11 |

## 1. INTRODUCTION

Topological recursion (TR) [EO07] is an explicit recursive procedure that associates to a small input data, the so-called spectral curve data that consists of a Riemann surface, two functions on it, and a bi-differential, a system of the so-called correlation differentials  $\{\omega_n^{(g)}\}_{g\geq 0,n\geq 1}$ , which are symmetric *n*-differentials on the *n*-th Cartesian power of the underlying Riemann surface. We omit the explicit definition of TR along with its possible variations and ramifications, as it won't be directly used in the present paper and refer the reader to *op. cit.* as well as further surveys available in [EO09; Eyn14].

Topological recursion has proved to be an ubiquitous tool that interconnects combinatorics, enumerative geometry, and integrability, with further applications to knot invariants and free probability, just to name a few. For many results of the principal interest at the interface of these areas, the only available proofs do go through topological recursion, the notable examples include e.g., [BDBKS24; DBKPS23; AS23; CLPS23].

A lot of effort is put into establishing the integrability property of the correlation differentials, see [BE12; ABDBKS23; EO24]. In particular, in [ABDBKS23] we gave a definition of KP integrability for a family of differentials  $\{\omega_n^{(g)}\}$  (not necessarily coming from topological recursion), proved that it requires the curve on which these differentials are defined to be of genus 0, and used its compatibility with the so-called x - y duality in

topological recursion in order to establish KP integrability for a large class of examples, providing, in particular, the explicit way to derive the associated tau functions.

The main goal of this paper is to complete the study of KP integrability for the TR differentials as a property of a family of differentials. We prove that for any TR input data with the spectral curve of genus 0 the associated differentials are KP integrable.

Note that, while this result is the most genera that one might hope to obtain in the framework of our understanding of KP integrability as a global property of a system of differentials, it does not exhibits the interrelations between KP and TR: there are conjectures on integrability of the system of correlation differentials associated to higher genera spectral curves that involve corrections with  $\Theta$ -functions and is understood as the property of particular expansions of the differentials, see a recent survey in [EO24].

The rest of the introduction is organized as follows: in Section 1.1 we proceed to formulate the main result of this paper, and in Section 1.2 we provide details on the results in [ABDBKS23] briefly mentioned above; they form the necessary background for our main theorem.

1.1. Main theorem. Consider any spectral curve data  $(\Sigma, x, y, B)$  such that

- $\Sigma$  is of genus zero, that is,  $\Sigma = \mathbb{C}P^1$ ;
- B is the canonical Bergman kernel, that is,  $B(z_1, z_2) = \frac{dz_1 dz_2}{(z_1 z_2)^2}$ , where z is a global affine coordinate on  $\Sigma = \mathbb{C}P^1$ ;
- dx is a meromorphic differential such that all its zeros are simple; we denote these zeros of dx by  $q_1, \ldots, q_N$ ;
- y is given as a collection of formal power expansions at the points  $q_i$  such that  $dy(q_i) \neq 0$  for all i = 1, ..., N.

Note that the notation we use is the same as in the papers we referred to above for the definitions of TR, but we relax the conditions on x and y. The topological recursion procedure associates with this initial data a system of symmetric *n*-differentials  $\omega_n^{(g)}$  on  $\Sigma^n$  for all  $g \ge 0$ ,  $n \ge 1$  such that

- $\omega_1^{(0)} = y \, dx, \, \omega_2^{(0)} = B$ . Note that  $\omega_1^{(0)}$  is only defined as a formal expansion at the points  $\{q_1, \ldots, q_N\}$ .
- For 2g 2 + n > 0, the forms  $\omega_n^{(g)}$  are global meromorphic, i.e. rational; the only possible poles that they can possess (with respect to each argument) are at the zeros of dx only. Moreover, the order of poles is at most 2(3g 3 + n) + 2 and the principal parts of the poles are odd with respect to the deck transformation of the function x considered as a local degree two ramified covering.

The main result of this paper is the following theorem that establishes integrable properties of all systems of correlation differentials obtained by topological recursion on the genus zero spectral curve.

**Theorem 1.1.** The system of TR differentials for the input data as above possesses KP integrability property.

The KP integrability as a property of a system of differentials is discussed below, see Definition 1.6. This theorem admits several generalizations. One of them is already included in the description of the input data above, as we assumed y to be given as a collection of formal power expansions at the points  $p_i$  rather than a globally defined function which is the standard setup for topological recursion. But in fact we prove that one can further relax this setup and assume that dx is also defined just in the vicinity of the points  $q_i$ . *Remark* 1.2. Note that we do not provide a detailed description of tau functions generated through topological recursion, but instead focus on proving the integrability. The general family of tau function that originates from topological recursion is much more extensive than the specific family with known explicit description, say, in terms of asymptotic description of integrals, constructed in [ABDBKS23].

1.2. **Background.** Here we mostly follow [ABDBKS23] where the inverse of Theorem 1.1 is proven, see Theorem 1.9 below.

Let  $\Sigma$  be a smooth complex curve that we refer below as the *spectral curve*. Consider a system of *n*-differentials  $\omega_n^{(g)}$  on  $\Sigma^n$  defined for all  $g \ge 0$ ,  $n \ge 1$ . We do not assume that these differentials are related by any recursion. We assume that all  $\omega_n^{(g)}$ 's are symmetric and meromorphic with no poles on the diagonals for  $(g, n) \ne (0, 2)$ , and  $\omega_2^{(0)}$  is also symmetric and meromorphic but it has a second order pole on the diagonal with biresidue 1. To pass from a system of differentials to KP tau functions, we need the following definition:

**Definition 1.3.** A point  $o \in \Sigma$  is called *regular* for the system of differentials  $\{\omega_n^{(g)}\}$  if  $\omega_n^{(g)} - \delta_{(g,n),(0,2)} \frac{d\xi_1 d\xi_2}{(\xi_1 - \xi_2)^2}$  is regular at  $(o, \ldots, o) \in \Sigma^n$  for all  $(g, n) \in \mathbb{Z}_{\geq 0} \times \mathbb{Z}_{>0}, (g, n) \neq (0, 1)$ , where  $\xi$  is a local coordinate on  $\Sigma$  near o.

*Remark* 1.4. Note that the regularity condition for  $\{\omega_n^{(g)}\}\$  is independent of a choice of local coordinate. Note also that we have no condition for  $\omega_1^{(0)}$ .

For a regular point o, and an arbitrary local coordinate  $\xi$ , the forms  $\omega_n^{(g)}$  can be expanded into a series

(1) 
$$\omega_n = \sum_{\substack{g \ge 0\\2g-2+n \ge 0}} \hbar^{2g-2+n} \omega_n^{(g)} = \delta_{n,2} \frac{d\xi_1 d\xi_2}{(\xi_1 - \xi_2)^2} + \sum_{k_1, \dots, k_n=1}^{\infty} f_{k_1, \dots, k_n} \prod_{i=1}^n k_i \xi_i^{k_i - 1} d\xi_i$$

where the coefficients  $f_{k_1,\ldots,k_n}$  expand as  $\sum_{g=0}^{\infty} \hbar^{2g-2+n} f_{k_1,\ldots,k_n}^{(g)}$ . Introduce the associated potential  $F = F_{o,\xi}$  as

(2) 
$$F(p_1, p_2, \dots) \coloneqq \sum_{\substack{g \ge 0, n \ge 1\\ 2g - 2 + n \ge 0}} \hbar^{2g - 2 + n} F_n^{(g)} = \sum_{n=1}^\infty \frac{1}{n!} \sum_{k_1, \dots, k_n = 1}^\infty f_{k_1, \dots, k_n} p_{k_1} \dots p_{k_n}$$

where  $F_n^{(g)}$  are homogeneous formal series of degree n and  $p_i$ , i = 1, 2, ..., are formal variables.

This way we associate to a given system of differentials  $\omega_n^{(g)}$  a large family of potentials  $F_{o,\xi}$ : the freedom in its definition consists of a choice of a regular point o and the choice of a local coordinate  $\xi$  at o. We would like to study conditions assuring that

(3) 
$$Z_{o,\xi} \coloneqq \exp(F_{o,\xi})$$

is a tau function of KP hierarchy.

**Theorem 1.5.** [ABDBKS23, Theorem 2.3] If  $Z_{o,\xi}$  is a tau function of KP hierarchy for some choice of a regular point o and a coordinate  $\xi$  at this point, then it is a tau function of KP hierarchy for any other choice of a regular point and a local coordinate at that point.

This allows us to give the following definition, which introduces the KP integrability as an internal property of the system of differentials:

**Definition 1.6.** A system of differentials  $\{\omega_n^{(g)}\}\$  satisfying the assumptions as above is called KP-integrable, if the associated partition function  $Z_{o,\xi}$  is a KP tau function for at least some choice of a regular point o and a local coordinate  $\xi$ .

*Remark* 1.7. The choice of  $\omega_1^{(0)}$  does not affect the KP integrability property, it can be chosen in an arbitrary way. For definiteness, in all expressions which contain a summation over genus we do not include the terms with (g, n) = (0, 1).

*Remark* 1.8. A change of the local spectral parameter is known to provide a symmetry of the KP hierarchy [Shi86], given by an element of the Virasoro subgroup of  $\widehat{GL}_{\infty}$ .

**Theorem 1.9.** [ABDBKS23, Theorem 3.8] If the differentials  $\{\omega_n^{(g)}\}$  produced by TR are KP integrable, then the spectral curve  $\Sigma$  is rational.

The upshot of Theorems 1.1 and 1.9 can be then formulated as follows:

**Corollary 1.10.** A system of differentials  $\{\omega_n^{(g)}\}$  produced by topological recursion is KP integrable if and only if the spectral curve is rational.

1.3. Applications. Theorem 1.1 immediately has a variety of applications. Namely, for any input data of topological recursion with a rational spectral curve we can immediately state the KP integrability of the resulting system of differentials and the partition functions associated to their expansions. In order to illustrate this principle, we analyze the spectral curve  $x = \log z - z^r$ ,  $y = z^s$ ,  $z \in \mathbb{C}P^1$ ,  $r, s \in \mathbb{Z}_{>0}$ , associated in [LPSZ17] to the Chern classes of the twisted *r*-th roots of the tensor powers of the log canonical bundle, the so-called  $\mathfrak{Q}$ -classes, or, in other terminology, the Chiodo classes.

In some cases the KP integrability of the associated partition functions was known — for some special choices of r, s, and local expansions one obtains instances of hypergeometric KP tau functions [DBKPS23; BDBKS24] associated to the so-called orbifold r/s-spin Hurwitz numbers (the ratio r/s is required to be integer). In this paper we generalize these results to arbitrary positive integers r and s, see Corollary 5.2.

1.4. Organization of the paper. In Section 2 we develop the theory of KP symmetries for the notion of KP integrability considered as a global property of a system of differentials. In Section 3 we use the theory of KP symmetries in combination to the deformation formulas for the differentials of topological recursion in order to reduce the KP integrability of a given system of TR differentials to the know instances of KP integrability proved in [ABDBKS23], which establishes our main theorem, Theorem 1.1. Section 4 is devoted to a generalized setup where we no longer assume that function x is globally defined. Finally, in Section 5 we provide an application to the KP integrability of the  $\Omega$ -classes.

In Appendix A we provide some background on the deformation formulas for the differentials obtained by topological recursion.

1.5. Notation. Throughout the text we use the following notation:

- [n] denotes  $\{1, ..., n\}$ .
- $z_I$  denotes  $\{z_i\}_{i \in I}$  for  $I \subseteq [[n]]$ .
- $[u^d]$  denotes the operator that extracts the corresponding coefficient from the whole expression to the right of it, that is,  $[u^d] \sum_{i=-\infty}^{\infty} a_i u^i \coloneqq a_d$ .
- $\lfloor_{u \to v}$  denotes the operator of substitution (or restriction), that is,  $\lfloor_{u \to v} f(u) \coloneqq f(v)$ .

1.6. Acknowledgments. A. A. was supported by the Institute for Basic Science (IBS-R003-D1). B.B. was supported by the ISF Grant 876/20. B.B., P.D.-B., and M.K. were supported by the Russian Science Foundation (grant No. 24-11-00366). S. S. was supported by the Netherlands Organization for Scientific Research.

## 2. Global KP symmetries

For a given system of meromorphic symmetric differentials  $\omega_n^{(g)}$  and a function x such that dx is meromorphic and  $\omega_n^{(g)} - \delta_{(g,n),(0,2)} \frac{dx_1 dx_2}{(x_1 - x_2)^2}$  is regular on the diagonals we set

(4) 
$$\omega_n = \sum_{\substack{g \ge 0\\2g-2+n \ge 0}} \hbar^{2g-2+n} \omega_n^{(g)}$$

and define

(5) 
$$\omega_2^{(0),\text{sing}}(z_1, z_2) = \frac{dx(z_1) \, dx(z_2)}{(x(z_1) - x(z_2))^2}$$

(6) 
$$\mathcal{T}_{n}(z^{+}, z^{-}; z_{[n]}) = \sum_{k=1}^{\infty} \frac{1}{k!} \int_{z^{-}}^{z^{+}} \dots \int_{z^{-}}^{z^{+}} (\omega_{n+k}(\underbrace{\cdot, \dots, \cdot}_{k}, z_{[n]}) - \delta_{(n,k),(0,2)} \omega_{2}^{(0), \text{sing}}),$$

(7) 
$$\Omega_n(z^+, z^-; z_{[n]}) = \sqrt{\omega_2^{(0), \text{sing}}(z^+, z^-)} e^{\mathcal{T}_0(z^+, z^-)} \sum_{[[n]] = \bigsqcup_{\alpha} J_{\alpha}, \ J_{\alpha} \neq \emptyset} \prod_{\alpha} \mathcal{T}_{|J_{\alpha}|}(z^+, z^-; z_{J_{\alpha}}),$$

(8) 
$$\mathcal{W}_n(x, u; z_{\llbracket n \rrbracket}) = \bigsqcup_{z^{\pm} \to e^{\pm \frac{u\hbar}{2}\partial_{x_z}}} \Omega_n(z^+, z^-; z_{\llbracket n \rrbracket}).$$

Remark 2.1. The arguments  $(z^+, z^-)$  of  $\Omega_n$  are treated as a point of the Cartesian square  $\Sigma^2$  and are considered in a certain vicinity of the diagonal  $z^+ = z^-$ . The arguments (z, u) of  $\mathcal{W}_n$  provide a particular parametrization of this vicinity through the substitutions

(9) 
$$z^{+} = e^{\frac{u\hbar}{2}\partial_{x}}z, \qquad z^{-} = e^{-\frac{u\hbar}{2}\partial_{x}}z.$$

Remark 2.2. The half-differentials  $\sqrt{dx(z^{\pm})}$  entering  $\Omega_n$  are understood formally. They attain a more definite meaning after the above substitution only. Namely, we have

(10) 
$$\lfloor_{z^{\pm} \to e^{\pm \frac{u\hbar}{2}\partial_{x_{z}}}} x^{\pm} = x \pm \frac{u\hbar}{2}, \qquad \lfloor_{z^{\pm} \to e^{\pm \frac{u\hbar}{2}\partial_{x_{z}}}} \sqrt{\omega_{2}^{(0), \text{sing}}(z^{+}, z^{-})} = \frac{dx}{u\hbar}$$

A reason to introduce  $\Omega_n$  in this form is that they are actually independent of x and are uniquely determined by the differentials  $\omega_n^{(g)}$ . Indeed, a different choice of the x-function changes both the factor  $\sqrt{\omega_2^{(0),\text{sing}}(z^+, z^-)} = \frac{\sqrt{dx^+ dx^-}}{x^+ - x^-}$  and the regularization term  $\omega_2^{(0),\text{sing}}$ of the singular form  $\omega_2^{(0)}$  under the integral in the definition of  $\mathcal{T}_0$ . An easy computation shows that these two changes compensate one another, see [ABDBKS23, Equation (3.10)].

Remark 2.3. We regard  $\mathcal{W}_n$  as an (n+1)-differential on  $\Sigma^{n+1}$  depending on an additional parameter u. One can see that in it the integration can be considered as an (infinite) combination of the differentials  $\omega_{\tilde{n}}^{(\tilde{g})}$  and their derivatives. This fact follows from the equality

$$(11) \quad \left\lfloor_{\tilde{z}^{\pm} \to e^{\pm \frac{u\hbar}{2}\partial_{x_{z}}}} \int_{z^{-}}^{z^{+}} \dots \int_{z^{-}}^{z^{+}} \omega_{n+k}(\underbrace{\cdot, \dots, \cdot}_{k}, z_{\llbracket n \rrbracket}) = \prod_{i=1}^{k} \left( \left\lfloor_{\tilde{z}_{i} \to z} u\hbar \mathcal{S}(u\hbar \partial_{\tilde{x}_{i}}) \right) \frac{\omega_{n+k}(\tilde{z}_{\llbracket k \rrbracket}, z_{\llbracket n \rrbracket})}{\prod_{i=1}^{k} d\tilde{x}_{i}} \right\rfloor$$

where  $\mathcal{S}(u) = u^{-1}(e^{u/2} - e^{-u/2})$ . As a corollary, the coefficient of each monomial in  $\hbar$  and u in  $\mathcal{W}_n$  is a differential polynomial in  $\omega$ 's. In particular, let us set  $\mathcal{W}_n^{(g),r} \coloneqq [u^r \hbar^{2g-1+n}] \mathcal{W}_n$ .

Then, we have

(12) 
$$\mathcal{W}_{n}^{(g),0}(z; z_{[n]}) = \omega_{n+1}^{(g)}(z, z_{[n]}),$$
  
(13)  $\mathcal{W}_{n}^{(g),1}(z; z_{[n]}) = \frac{1}{2 dx(z)} \Big( \omega_{n+2}^{(g-1)}(z, z, z_{[n]}) + \sum_{\substack{g_{1}+g_{2}=g\\I_{1}\sqcup I_{2}=[n]\\2g_{i}-1+|I_{i}|\geq 0}} \omega_{|I_{1}|+1}^{(g_{1})}(z, z_{I_{1}}) \, \omega_{|I_{2}|+1}^{(g_{2})}(z, z_{I_{2}}) \Big).$ 

Remark 2.4. Note that in (4) we used our convention that the sum over genus excludes the (g, n) = (0, 1) term, which affects (7) and (8). In some other situations, for instance, in the study of the loop equations and/or deformations of the differentials, these terms should not be excluded. In particular, below we'll also need a version of  $\mathcal{W}_n^{(g),1}$  with the (0, 1) terms included, which denote by

(14) 
$$\widetilde{\mathcal{W}}_{n}^{(g),1}(z; z_{[n]}) = \frac{1}{2 \, dx(z)} \Big( \omega_{n+2}^{(g-1)}(z, z, z_{[n]}) + \sum_{\substack{g_1 + g_2 = g\\I_1 \sqcup I_2 = [n]}} \omega_{|I_1|+1}^{(g_1)}(z, z_{I_1}) \, \omega_{|I_2|+1}^{(g_2)}(z, z_{I_2}) \Big).$$

**Theorem 2.5.** For  $n \ge 1$  the transformation

(15) 
$$\Delta \omega_n(z_{\llbracket n \rrbracket}) = \Omega_n(z^+, z^-; z_{\llbracket n \rrbracket})$$

is an infinitesimal KP symmetry for a system of differentials  $\{\omega_n^{(g)}\}\$  for any specialization of  $z^+$  and  $z^-$ . In other words, the above formula provides a family of infinitesimal KP symmetries with  $(z^+, z^-) \in \Sigma^2$  regarded as parameters of the family.

By an infinitesimal transformation here we mean  $\omega_n(z_{[n]}) \mapsto \omega_n(z_{[n]}) + \epsilon \Delta \omega_n(z_{[n]})$  with an infinitesimal parameter  $\epsilon$ . By a specialization  $z^+$  and  $z^-$  we mean a particular choice of these points in a vicinity of the diagonal in  $\Sigma^2$ , or the coefficient of any monomial in  $z^+, z^-$  in the power expansion in the local coordinate z at some point of  $\Sigma$ , or the integral over some contour, or the residue at some point of the spectral curve.

More explicitly, the statement of Theorem 2.5 means the following. Assume that the differentials  $\omega_n^{(g)}$  depend on an additional parameter  $\tau$  and the dependence of the differentials on  $\tau$  is govern by the differential equation of the form

(16) 
$$\frac{d}{d\tau}\omega_n(z_{\llbracket n \rrbracket}) = \Omega_n(z^+, z^-; z_{\llbracket n \rrbracket})$$

for some specialization of  $z^+$ ,  $z^-$  also possibly depending on  $\tau$  and  $\hbar$ . In what follows we consider the deformations with  $z^+$ ,  $z^-$  having at most simple poles in  $\hbar$ , thus the right hand side of (15) might contain a pole in  $\hbar$  as well; this pole should be deducted as it does not affect the KP integrability and is removed on the left hand side by our conventions, cf. Remark 1.7. Assume also that the differentials posses KP integrability property for some value of  $\tau$ . Then they posses this property for all values of  $\tau$ .

**Corollary 2.6.** For any r and any specialization of  $z \in \Sigma$  the transformation

(17) 
$$\Delta \omega_n^{(g)}(z_{\llbracket n \rrbracket}) = \mathcal{W}_n^{(g),r}(z; z_{\llbracket n \rrbracket})$$

is an infinitesimal KP symmetry for a system of differentials  $\{\omega_n^{(g)}\}$ .

Proof of Theorem 2.5. First, assume that  $z^+$  and  $z^-$  are chosen in a neighborhood of some point  $o \in \Sigma$  regular for all differentials  $\omega_n^{(g)}$ . Let us choose an arbitrary local coordinate z at this point and identify a point in a neighborhood of o with the value of the coordinate z at this point. A choice of a point o and a coordinate z determines the corresponding potential  $F_{o,z} = F_{o,z}(p_1, p_2, ...)$  defined by (1) and (2).

Now we are going to apply the logic that has earlier appeared in [ABDBKS22; ABD-BKS23], namely, we perform local computation for the formal expansions using the Fock

space semi-infinite wedge formalism (we refer the reader to the *op. cit.* for the notation that we use here, see, e.g., [ABDBKS23, Section 2.2]), but the formulas that we obtain have the global meaning on the spectral curve.

The potential  $F_{o,z}$  defines, in turn, the corresponding vector  $\hat{Z}_{o,z} |0\rangle$  in the Fock space where  $\hat{Z}_{o,z} = e^{F_{o,z}(J_{-1},J_{-2},...)}$ . With this notation, the correspondence between the potential and *n*-differentials takes the form of connected vacuum expectation values

(18) 
$$\omega_n(z_{\llbracket n \rrbracket}) = \langle 0 | J(z_1) \dots J(z_n) \hat{Z}_{o,z} | 0 \rangle^{\circ} \prod_{i=1}^n \frac{dz_i}{z_i}, \qquad J(z) = \sum_{m=-\infty}^{\infty} z^m J_m.$$

The definition of  $\Omega_n$  specialized to the setting of power expansions in z variables can be represented, in turn, in the form (19)

$$\Omega_n(z^+, z^-; z_{[n]}) - \delta_{n,0} \frac{\sqrt{dz^+ dz^-}}{z^+ - z^-} = \langle 0 | J(z_1) \dots J(z_n) \mathcal{K}(z^+, z^-) \hat{Z}_{o,z} | 0 \rangle^\circ \sqrt{dz^+ dz^-} \prod_{i=1}^n \frac{dz_i}{z_i},$$

where  $\mathcal{K}$  is the operator acting in the Fock space and defined by

(20) 
$$\mathcal{K}(z_1, z_2) = \frac{e^{\sum_{k < 0} \frac{z_1^k - z_2^k}{k} J_k} e^{\sum_{k > 0} \frac{z_1^k - z_2^k}{k} J_k}}{z_1 - z_2}.$$

It follows that the infinitesimal transformation of the vector  $\hat{Z}_{o,z}|0\rangle$  in the Fock space corresponding to the transformation (16) is given by

(21) 
$$\Delta \hat{Z}_{o,z} |0\rangle = \sqrt{dz^+ dz^-} \, \mathcal{K}(z^+, z^-) \hat{Z}_{o,z} |0\rangle.$$

It is known [MJD12; Kac90; BDBKS22] that the operator  $\mathcal{K}(z_1, z_2)$  belongs to the Lie algebra  $\widehat{\mathfrak{gl}}(\infty)$  of infinitesimal KP symmetries.

We conclude that the transformations (21) and (16) preserve infinitesimally KP integrability for the potential  $F_{o,z}$  associated with the point o and the local coordinate z at this point (the factor  $\sqrt{dz^+dz^-}$  is a numerical constant that can be set arbitrarily and does not affect KP integrability). On the other hand, by Theorem 1.5, the KP integrability does not depend on a choice of the expansion point and the local coordinate. Hence, the transformation (16) provides an infinitesimal KP symmetry for the whole globally defined system of differentials  $\{\omega_n^{(g)}\}$ . Since these arguments are applied to arbitrary choice of o, we obtain that the statement of Theorem 2.5 is valid in its full generality.

## 3. Proof of Theorem 1.1

Let us study the dependence of the TR differentials on a choice of the function y. Thus we consider a family of spectral curve data such that the spectral curve  $\Sigma$ , the Bergman kernel B, and the function x are fixed (and hence, the positions of the points  $q_1, \ldots, q_N$ are also fixed) while the function y (that is, its expansion at the points  $q_i$ ) varies in a family smoothly depending on a parameter  $\tau$ . The corresponding dependence of the TR differentials on  $\tau$  in the stable range 2g - 2 + n > 0 is described by the equation [EO07]:

(22) 
$$\frac{d}{d\tau}\omega_n^{(g)}(z_{\llbracket n \rrbracket}) = \sum_{i=1}^N \operatorname{res}_{z=q_i} \left(\omega_{n+1}^{(g)}(z_{\llbracket n \rrbracket}, z) \int_{q_i}^z \frac{dy}{d\tau} \, dx\right).$$

This equation holds true for any variation of y such that  $dy(q_i) \neq 0$ ; in particular, we allow a change of the constants  $y'(q_i)$  as long as these constants are not vanishing in the family.

By Corollary 2.6 and (12), the right hand side of Eq. (22) is an infinitesimal KP symmetry. It follows that the KP integrability property is preserved in the family, and hence the system of differentials  $\{\omega_n^{(g)}\}$  is KP integrable for some choice of y if and only if it is KP integrable for any other choice of y, provided that dy is non-vanishing at  $q_1, \ldots, q_N$ .

Thus, in the setting of Theorem 1.1, it is sufficient to prove KP integrability in the case when y = z is a global affine coordinate on  $\Sigma = \mathbb{C}P^1$ . The KP integrability for this choice of y is proved in [ABDBKS23]. This completes the proof of Theorem 1.1.

## 4. Generalized setting

The statement of Theorem 1.1 holds in a more general setting. It is sufficient to assume that x is defined in a vicinity of the points  $q_1, \ldots, q_N$  such that dx has simple zeros in these points and does not extend necessarily to the whole spectral curve, and dy is holomorphic and nonvanishing at these points. Even if dx extends beyond this vicinity and has other zeros, they do not contribute to TR, by definition.

The definition of TR differentials can be extended to this relaxed setting as well. If the Bergman kernel B entering the initial data of recursion is defined locally in a neighborhood of the points  $q_j$ , then the resulting differentials are also defined locally in a neighborhood of those points. If  $\Sigma$  is compact and B extends globally as a meromorphic bidifferential with no singularities beyond the diagonal, then the TR differentials  $\omega_n^{(g)}$  are also defined globally for 2g - 2 + n > 0, and have no poles different from  $q_1, \ldots, q_N$  (with respect to each of the arguments), even though the functions x and y of the initial data are defined locally near the points  $q_j$ .

**Theorem 4.1.** If the spectral curve is rational,  $\Sigma = \mathbb{C}P^1$ , and B is the standard Bergman kernel, then the KP integrability for the TR differentials holds in the relaxed setting of this section as well.

*Proof.* Consider a variation of the initial data of recursion such that the spectral curve and the Bergman kernel are fixed while the functions x and y vary in a one-parameter family depending on a parameter  $\tau$ . We allow the positions of the points  $q_j$  to change in the family provided that they remain pairwise distinct and the assumptions on dx and dyare satisfied for all parameter values. Then, the dependence of the TR differentials on  $\tau$ in the stable range 2g - 2 + n > 0 is described by the following equation, see [Kaz] and Appendix A

(23) 
$$\frac{d}{d\tau}\omega_n^{(g)}(z_{\llbracket n \rrbracket}) = \sum_{j=1}^N \operatorname{res}_{z=q_j} \left( \mathcal{W}_n^{(g),0}(z; z_{\llbracket n \rrbracket}) \int_{d\tau}^z \frac{d}{d\tau} (y \, dx) - \widetilde{\mathcal{W}}_n^{(g),1}(z; z_{\llbracket n \rrbracket}) \frac{d}{d\tau} x(z) \right),$$

where  $\mathcal{W}_n^{(g),0}$  and  $\widetilde{\mathcal{W}}_n^{(g),1}$  are defined by (12) and (14), respectively. By Theorem 2.5 and its Corollary 2.6, the KP integrability for the TR differentials is preserved in the family (note that we can use  $\widetilde{\mathcal{W}}_n^{(g),1}$  instead of  $\mathcal{W}_n^{(g),1}$  once the singular term in  $\hbar$  is removed). Any collection of initial data can be connected by an analytic family with those satisfying the assumptions of Theorem 1.1, that is, when dx extends globally as a meromorphic differential with no zeros except  $q_1, \ldots, q_N$ . Say, it can be connected with the case when x(z) is a polynomial of degree N + 1 with no multiple critical points. Therefore, the assertion of Theorem 4.1 follows from that one of Theorem 1.1.

### 5. Application: r-th roots

Consider the moduli space of generalized *r*-spin structures  $\overline{\mathcal{M}}_{g;\vec{a}}^{r,s}$  which parametrizes *r*-th roots of the line bundle

(24) 
$$\omega_{\log}^{\otimes s} \left( -\sum_{i=1}^{n} a_i \mathbf{x}_i \right)$$

on genus g curves with n marked points  $(C, \mathbf{x}_1, \ldots, \mathbf{x}_n)$ , where  $\omega_{\log} = \omega(\sum_i \mathbf{x}_i)$ , r and s are positive integers, and  $a_1, \ldots, a_n$  are integers satisfying

(25) 
$$a_1 + a_2 + \dots + a_n \equiv (2g - 2 + n)s \pmod{r}.$$

This condition guarantees the existence of a line bundle on the underlying curve whose *r*-th tensor power is isomorphic to  $\omega_{\log}^{\otimes s}(-\sum_{i} a_{i}x_{i})$ . Let  $\pi: \mathcal{C}_{g,\vec{a}}^{r,s} \to \overline{\mathcal{M}}_{g,\vec{a}}^{r,s}$  be the universal curve, and  $\mathcal{L} \to \overline{\mathcal{C}}_{g,\vec{a}}^{r,s}$  the universal *r*-th root. The Chiodo formula expresses the Chern characters of the derived pushforward of the universal *r*-th root  $ch_{m}(R^{\bullet}\pi_{*}\mathcal{L})$  in terms of the tautological classes, see [Chi08].

We can then consider the family of Chern classes pushed forward to the moduli spaces of stable curves under the natural projection map  $\epsilon \colon \overline{\mathcal{M}}_{g,\vec{a}}^{r,s} \to \overline{\mathcal{M}}_{g,n}$ .

(26) 
$$\mathfrak{Q}_{g,n}(r,s;a_1,\ldots,a_n) \coloneqq \epsilon_* \exp\left(\sum_{m=1}^{\infty} (-1)^m (m-1)! \operatorname{ch}_m(R^{\bullet}\pi_*\mathcal{L})\right) \in R^*(\overline{\mathcal{M}}_{g,n}).$$

These classes are called the  $\mathcal{Q}$ -classes, or, alternatively, the Chiodo classes in the literature.

Consider the spectral curve data

(27) 
$$\Sigma = \mathbb{C}P^1, \quad x = \log z - z^r, \quad y = z^s, \quad B = \frac{dz_1 dz_2}{(z_1 - z_2)^2},$$

where, as usual, z is the global affine coordinate on  $\mathbb{C}P^1$ . The following theorem is established in [LPSZ17] (see also [SSZ15] for the s = 1 case):

**Theorem 5.1** ([LPSZ17]). The expansions of the correlation differentials of the spectral curve (27) at the point  $z_1 = \cdots = z_n = 0$  in the local coordinates  $X_i = X(z_i) = \exp(x(z_i))$  are equal to

(28) 
$$\omega_n^{(g)} = d_1 \otimes \cdots \otimes d_n \sum_{\mu_1, \dots, \mu_n = 1}^{\infty} \prod_{j=1}^n \left( X_j^{\mu_j} \frac{\left(\frac{\mu_j}{r}\right)^{\left\lfloor \frac{\mu_j}{r} \right\rfloor}}{\left\lfloor \frac{\mu_j}{r} \right\rfloor!} \right) \times \frac{r^{2g-2+n+\frac{(2g-2+n)s+\sum_{j=1}^n \mu_j}{r}}}{s^{2g-2+n}}$$
$$\times \int_{\overline{\mathcal{M}}_{g,n}} \frac{\mathcal{Q}_{g,n}\left(r, s; r-r\left\langle \frac{\mu_1}{r} \right\rangle, \dots, r-r\left\langle \frac{\mu_n}{r} \right\rangle\right)}{\prod_{j=1}^n \left(1 - \frac{\mu_j}{r} \psi_j\right)},$$

where  $\frac{\mu}{r} = \lfloor \frac{\mu}{r} \rfloor + \langle \frac{\mu}{r} \rangle$  is the decomposition into the integer and the fractional parts.

Note that the spectral curve data satisfies the conditions of Theorem 1.1. Hence, we obtain the following immediate corollary:

Corollary 5.2. The following partition function is a tau function of KP hierarchy:

(29) 
$$\exp\left(\sum_{\substack{g\geq 0,n\geq 1\\2g-2+n>0}}\frac{\hbar^{2g-2+n}}{n!}\sum_{\mu_{1},\dots,\mu_{n}=1}^{\infty}\prod_{j=1}^{n}\left(p_{\mu_{j}}\frac{\left(\frac{\mu_{j}}{r}\right)^{\left\lfloor\frac{\mu_{j}}{r}\right\rfloor}}{\left\lfloor\frac{\mu_{j}}{r}\right\rfloor!}\right)\times\frac{r^{2g-2+n+\frac{(2g-2+n)s+\sum_{j=1}^{n}\mu_{j}}{r}}{s^{2g-2+n}}}{s^{2g-2+n}}\right)$$
$$\times\int_{\overline{\mathcal{M}}_{g,n}}\frac{\mathscr{Q}_{g,n}\left(r,s;r-r\left\langle\frac{\mu_{1}}{r}\right\rangle,\dots,r-r\left\langle\frac{\mu_{n}}{r}\right\rangle\right)}{\prod_{j=1}^{n}\left(1-\frac{\mu_{j}}{r}\psi_{j}\right)}}$$
$$+\frac{1}{2}\sum_{\mu_{1},\mu_{2}=1}^{\infty}p_{\mu_{1}}p_{\mu_{2}}[X_{1}^{\mu_{1}}X_{2}^{\mu_{2}}]\log\left(\frac{z_{1}-z_{2}}{x_{1}-x_{2}}\right)\right).$$

Remark 5.3. In the case r = st for some positive integer t, this function up to an extra factor of  $\exp(\hbar^{-1}\sum_{\mu} p_{\mu}[X^{\mu}] \int_{0}^{z} z^{s-1}(1-rz^{r})dz)$  is a hypergeometric KP tau function studied in [DBKPS23; BDBKS24]. So, in these cases the result of Corollary 5.2 was known.

## APPENDIX A. VARIATION OF TR DIFFERENTIALS

In this Appendix, we review formulas of [EO07; DBOSS14; Kaz] describing variation of topological recursion differentials under a deformation of initial data of recursion.

Assume that the spectral curve data  $(\Sigma, x, y, B)$  of topological recursion (or, more generally, the local spectral curve data as in Section 4) varies in a family depending analytically on one additional parameter  $\tau$ . Since zeros  $q_j$  of dx are nondegenerate, this property is preserved under variation of the spectral curve data, and by implicit function theorem, the dependence of positions of the points  $q_j$  on the parameter is analytic. We assume that no zeros of dx other than  $q_1, \ldots, q_N$  appear for the deformed spectral curve, or that additional zeros are ignored and do not contribute to the recursion, by definition. Then one may expect that the variation of the differentials  $\omega_n^{(g)}$  is determined by the variation of the functions x, y, and the bidifferential B in a neighborhood of the points  $q_i$ . In fact, this is the case and we review here explicit formulas for the variation of  $\omega_n^{(g)}$ .

In order to give a meaning to the derivative of  $\omega_n^{(g)}$  over parameter  $\tau$  we need to identify spectral curves for different parameter values, at least locally near the points  $q_j$ . Let us take an arbitrary meromorphic function  $\zeta$  on the spectral curve and denote by  $\frac{d}{d\tau}|_{\zeta}$  the derivative at fixed  $\zeta$ , that is, the partial derivative over  $\tau$  computed in any local coordinate on the spectral curve such that the expression for  $\zeta$  in this local coordinate is independent of  $\tau$ .

Denote

$$(30) \quad \omega_{n,2}^{(g),\text{st}}(z,\tilde{z};z_{\llbracket n \rrbracket}) = \frac{1}{2} \bigg( \omega_{n+2}^{(g-1)}(z,\tilde{z},z_{\llbracket n \rrbracket}) + \sum_{\substack{g_1+g_2=g, \ I_1 \sqcup J_2 = \llbracket n \rrbracket} \\ \text{no} \ (0,1) \text{ or } (0,2) \text{ terms}} \omega_{|I_1|+1}^{(g_1)}(z,z_{I_1}) \, \omega_{|I_2|+1}^{(g_2)}(\tilde{z},z_{I_1}) \bigg),$$

$$(31) \quad \widetilde{\mathcal{W}}_n^{(g),1}(z;z_{\llbracket n \rrbracket}) = \frac{1}{2 \, dx(z)} \bigg( \omega_{n+2}^{(g-1)}(z,z,z_{\llbracket n \rrbracket}) + \sum_{\substack{g_1+g_2=g \\ I_1 \sqcup J_2 = \llbracket n \rrbracket}} \omega_{|I_1|+1}^{(g_1)}(z,z_{I_1}) \, \omega_{|I_2|+1}^{(g_2)}(z,z_{I_1}) \bigg)$$

$$= \frac{1}{dx(z)} \bigg( \omega_{n,2}^{(g),\text{st}}(z,z;z_{\llbracket n \rrbracket}) + \omega_{n+1}^{(g)}(z,z_{\llbracket n \rrbracket}) \, y(z) \, dx(z)$$

$$+ \sum_{i=1}^n \omega_n^{(g)}(z,z_{\llbracket n \rrbracket \setminus \{i\}}) \, B(z,z_i) \bigg).$$

#### REFERENCES

**Theorem A.1** ([Kaz]). For 2g - 2 + n > 0 the dependence of TR differentials  $\omega_n^{(g)}$  on the parameter  $\tau$  is described by the equation

$$(32) \quad \frac{d}{d\tau} \Big|_{\zeta} \omega_n^{(g)}(z_{[n]}) = \sum_{j=1}^N \operatorname{res}_{z=q_j} \left( \omega_{n+1}^{(g)}(z, z_{[n]}) \int_{\overline{d\tau}}^{z} \Big|_{\zeta} (y \, dx) - \widetilde{\mathcal{W}}_n^{(g),1}(z; z_{[n]}) \frac{d}{d\tau} \Big|_{\zeta} x(z) \right) \\ + \sum_{i,j=1}^N \operatorname{res}_{z=q_i} \operatorname{res}_{\tilde{z}=q_j} \omega_{n,2}^{(g),\text{st}}(z, \tilde{z}, z_{[n]}) \int_{\overline{d\tau}}^{z} \Big|_{\zeta} B + \sum_{i=1}^n \sum_{j=1}^N \operatorname{res}_{z=q_j} \omega_n^{(g)}(z, z_{[n]\setminus\{i\}}) \int_{\overline{d\tau}}^{z} \Big|_{\zeta} B(\cdot, z_i).$$

Remark A.2. In the cases we apply this formula in the present paper in (22) or (23), the curve is fixed so that the derivative over  $\tau$  has an invariant meaning and coincides with the derivative at fixed  $\zeta$  for any meromorphic function  $\zeta$  independent of  $\tau$ . By that reason we drop a reference to a choice of  $\zeta$  in (22) and (23). Besides, since the Bergman kernel is also fixed, the terms on the second line in (32) do not contribute in these cases.

Though an independent proof of this formula is given in [Kaz], let us explain how it is related to the deformation formulas of [EO07; DBOSS14]. The particular formulas for  $\frac{d}{d\tau}|_{\zeta}y$  being non-trivial with  $\frac{d}{d\tau}|_{\zeta}x = 0$  and  $\frac{d}{d\tau}|_{\zeta}B = 0$  as well as  $\frac{d}{d\tau}|_{\zeta}B$  being nontrivial with  $\frac{d}{d\tau}|_{\zeta}x = 0$  and  $\frac{d}{d\tau}|_{\zeta}y = 0$  just coincide with the expressions given in *op. cit.*, see [EO07, Theorem 5.1] and [DBOSS14, Proof of Theorem 3.7]. Thus, (32) holds in a special case  $\zeta = x$ , that is, if the derivative over  $\tau$  is computed at fixed x:

$$(33) \quad \frac{d}{d\tau}\Big|_{x}\omega_{n}^{(g)}(z_{[n]}) = \sum_{j=1}^{N} \mathop{\mathrm{res}}_{z=q_{j}} \omega_{n+1}^{(g)}(z, z_{[n]}) \int_{d\tau}^{z} \frac{d}{d\tau}\Big|_{x} (y \, dx) \\ + \sum_{i,j=1}^{N} \mathop{\mathrm{res}}_{z=q_{i}} \mathop{\mathrm{res}}_{\tilde{z}=q_{j}} \omega_{n,2}^{(g),\mathrm{st}}(z, \tilde{z}, z_{[n]}) \int_{d\tau}^{z} \int_{d\tau}^{z} \frac{d}{d\tau}\Big|_{x} B + \sum_{i=1}^{n} \sum_{j=1}^{N} \mathop{\mathrm{res}}_{z=q_{j}} \omega_{n}^{(g)}(z, z_{[n]}) \int_{d\tau}^{z} \frac{d}{d\tau}\Big|_{x} B(\cdot, z_{i}).$$

Passing from the derivative at fixed x to the derivative at fixed  $\zeta$  is govern by the following relations: for a given function f, a meromorphic 1-form  $\lambda$ , and an n-differential  $\omega_n$ , respectively, we have

(34) 
$$\frac{d}{du}\Big|_{\zeta}f = \frac{d}{du}\Big|_{x}f + \dot{x}\frac{df}{dx},$$

(35) 
$$\frac{d}{du}\Big|_{\zeta}\lambda = \frac{d}{du}\Big|_{x}\lambda + d\left(\dot{x}\frac{\lambda}{dx}\right),$$

(36) 
$$\frac{d}{du}\Big|_{\zeta}\omega_n = \frac{d}{du}\Big|_x\omega_n + \sum_{i=1}^n d_i\left(\dot{x}_i\frac{\omega_n}{dx_i}\right),$$

where  $\dot{x} = \frac{d}{du}|_{\zeta} x$ . The first equation is the usual chain rule for partial derivatives, and the other two follow from the first one. Substituting these expressions to (33), we arrive, after simplifications, exactly to (32), see [Kaz] for the details.

#### References

- [ABDBKS22] A. Alexandrov, B. Bychkov, P. Dunin-Barkowski, M. Kazarian, and S. Shadrin. A universal formula for the x y swap in topological recursion. 2022. arXiv: 2212.00320 [math-ph].
- [ABDBKS23] A. Alexandrov, B. Bychkov, P. Dunin-Barkowski, M. Kazarian, and S. Shadrin. KP integrability through the x y swap relation. 2023. arXiv: 2309.12176 [math-ph].

| 12        | REFERENCES   |
|-----------|--|
| [AS23]    | A. Alexandrov and S. Shadrin. "Elements of spin Hurwitz theory: closed al-<br>gebraic formulas, blobbed topological recursion, and a proof of the Giacchetto-<br>Kramer-Lewański conjecture". In: <i>Sel. Math. New Ser.</i> 29.2 (2023), #26,<br>44 pp. DOI: 10.1007/s00029-023-00834-1 |
| [BE12]    | G. Borot and B. Eynard. "Geometry of spectral curves and all order disper-<br>sive integrable system". In: SIGMA Symmetry Integrability Geom. Meth-<br>ods Appl. 8 (2012), Paper 100, 53, DOI: 10.3842/SIGMA.2012.100.   |
| [BDBKS22] | B. Bychkov, P. Dunin-Barkowski, M. Kazarian, and S. Shadrin. "Explicit closed algebraic formulas for Orlov-Scherbin <i>n</i> -point functions". In: <i>J. Éc. Polytech.</i> , <i>Math.</i> 9 (2022), pp. 1121–1158. DOI: 10.5802/jep.202.  |
| [BDBKS24] | B. Bychkov, P. Dunin-Barkowski, M. Kazarian, and S. Shadrin. "Topolog-<br>ical recursion for Kadomtsev–Petviashvili tau functions of hypergeomet-<br>ric type". In: <i>Journal of the London Mathematical Society</i> 109.6 (2024),<br>e12946. DOI: 10.1112/jlms.12946.                  |
| [CLPS23]  | G. Carlet, J. van de Leur, H. Posthuma, and S. Shadrin. "Enumeration of hypermaps and Hirota equations for extended rationally constrained KP". In: <i>Comm. Numb. Theory Phys.</i> 17.3 (2023), pp. 643–708. DOI: 10.4310/CNTP.2023.v17.n3.a3.  |
| [Chi08]   | A. Chiodo. "Towards an enumerative geometry of the moduli space of twisted curves and <i>r</i> th roots". In: <i>Compos. Math.</i> 144.6 (2008), pp. 1461–1496. DOI: 10.1112/S0010437X08003709.  |
| [DBOSS14] | P. Dunin-Barkowski, N. Orantin, S. Shadrin, and L. Spitz. "Identifica-<br>tion of the Givental formula with the spectral curve topological recur-<br>sion procedure". In: <i>Comm. Math. Phys.</i> 328.2 (2014), pp. 669–700. DOI:<br>10.1007/s00220-014-1887-2                          |
| [DBKPS23] | P. Dunin-Barkowski, R. Kramer, A. Popolitov, and S. Shadrin. "Loop<br>equations and a proof of Zvonkine's <i>qr</i> -ELSV formula". In: <i>Ann. Sci. Éc.</i><br><i>Norm. Sunér.</i> (4) 56.4 (2023) pp. 1199–1229. DOI: 10. 24033/asens. 2553  |
| [EO07]    | B. Eynard and N. Orantin. "Invariants of algebraic curves and topological expansion". In: <i>Commun. Number Theory Phys.</i> 1.2 (2007), pp. 347–452.  |
| [Eyn14]   | B. Eynard. "An overview of the topological recursion". In: <i>Proceedings</i><br>of the International Congress of Mathematicians—Seoul 2014. Vol. III.<br>Kyung Moon Sa. Seoul 2014, pp. 1063–1085   |
| [EO09]    | B. Eynard and N. Orantin. "Topological recursion in enumerative geometry and random matrices". In: J. Phys. A 42.29 (2009), pp. 293001, 117.<br>DOI: 10.1088/1751-8113/42/29/293001.   |
| [EO24]    | B. Eynard and S. Oukassi. <i>Hirota, Fay and Geometry</i> . 2024. arXiv: 2401.   |
| [Kac90]   | V. G. Kac. <i>Infinite-dimensional Lie algebras</i> . Cambridge University Press,<br>Cambridge, 1990, pp. xxii+400, DOI: 10.1017/CB09780511626234.   |
| [Kaz]     | M. Kazarian. Symplectic geometry of topological recursion. In prepara-<br>tion. Based on the lectures of the author at IHES in 2014 available at https://youtu.be/SkZtIJmauSM.   |
| [LPSZ17]  | D. Lewanski, A. Popolitov, S. Shadrin, and D. Zvonkine. "Chiodo formulas for the <i>r</i> -th roots and topological recursion". In: <i>Lett. Math. Phys.</i> 107.5 (2017), pp. 901–919. DOI: 10.1007/s11005-016-0928-5.  |
| [MJD12]   | T. Miwa, M. Jimbo, and E. Date. Solitons: differential equations, sym-<br>metries and infinite dimensional algebras. Transl. from the Japanese by  |

#### REFERENCES

|         | <i>Miles Reid.</i> English. Reprint of the 2000 hardback edition. Vol. 135. Camb. |
|---------|---|
|         | Tracts Math. Cambridge: Cambridge University Press, 2012.                         |
| [SSZ15] | S. Shadrin, L. Spitz, and D. Zvonkine. "Equivalence of ELSV and Bouchard-         |
|         | Mariño conjectures for r-spin Hurwitz numbers". In: Math. Ann. 361.3-4            |
|         | (2015), pp. 611–645. doi: 10.1007/s00208-014-1082-y.                              |
| [Shi86] | T. Shiota. "Characterization of Jacobian varieties in terms of soliton equa-      |
|         | tions". In: Invent. Math. 83.2 (1986), pp. 333–382. DOI: 10.1007/BF               |
|         | 01388967  |

A. A.: Center for Geometry and Physics, Institute for Basic Science (IBS), Pohang 37673, Korea

Email address: alex@ibs.re.kr

B. B.: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HAIFA, MOUNT CARMEL, 3498838, HAIFA, ISRAEL

 $Email \ address: bbychkov@hse.ru$ 

P. D.-B.: Faculty of Mathematics, HSE University, Usacheva 6, 119048 Moscow, Russia; HSE–Skoltech International Laboratory of Representation Theory and Mathematical Physics, Skoltech, Bolshoy Boulevard 30 bld. 1, 121205 Moscow, Russia; and NRC "Kurchatov Institute" – ITEP, 117218 Moscow, Russia

Email address: ptdunin@hse.ru

M. K.: Faculty of Mathematics, HSE University, Usacheva 6, 119048 Moscow, Russia; and Igor Krichever Center for Advanced Studies, Skoltech, Bolshoy Boulevard 30 bld. 1, 121205 Moscow, Russia

Email address: kazarian@mccme.ru

S. S.: Korteweg-de Vries Institute for Mathematics, University of Amsterdam, Postbus 94248, 1090GE Amsterdam, The Netherlands

Email address: S.Shadrin@uva.nl