

VANISHING PROPERTIES OF KLOOSTERMAN SUMS AND DYSON'S CONJECTURES

QIHANG SUN

ABSTRACT. In a previous paper [Sun24a], the author proved the exact formulae for ranks of partitions modulo each prime $p \geq 5$. In this paper, for $p = 5$ and 7 , we prove special vanishing properties of the Kloosterman sums appearing in the exact formulae. These vanishing properties imply a new proof of Dyson's rank conjectures. Specifically, we give a new proof of Ramanujan's congruences $p(5n + 4) \equiv 0 \pmod{5}$ and $p(7n + 5) \equiv 0 \pmod{7}$.

1. INTRODUCTION

Let $p(n)$ denote the integer partition function. Ramanujan obtained the famous congruence properties of $p(n)$:

$$p(5n + 4) \equiv 0 \pmod{5}, \quad p(7n + 5) \equiv 0 \pmod{7}, \quad p(11n + 6) \equiv 0 \pmod{11}. \quad (1.1)$$

In 1944, Dyson [Dys44] defined the rank of a partition and conjectured a beautiful explanation for Ramanujan's congruences. Suppose $\Lambda = \{\Lambda_1 \geq \Lambda_2 \geq \cdots \geq \Lambda_\kappa\}$ is a partition of n , i.e. $\sum_{j=1}^\kappa \Lambda_j = n$. Then the rank of Λ is defined by

$$\text{rank}(\Lambda) := \Lambda_1 - \kappa$$

Let the quantities $N(m, n)$ and $N(a, b; n)$ be defined by

$$N(m, n) := \#\{\Lambda \text{ is a partition of } n : \text{rank } \Lambda = m\} \quad (1.2)$$

and

$$N(a, b; n) := \#\{\Lambda \text{ is a partition of } n : \text{rank } \Lambda \equiv a \pmod{b}\}. \quad (1.3)$$

Let $q = \exp(2\pi iz) = e(z)$ for $z \in \mathbb{H}$ and w be a root of unity. The generating function of $N(m, n)$ is given by (see e.g. [BO06, p. 245])

$$\mathcal{R}(w; q) := 1 + \sum_{n=1}^{\infty} \sum_{m=-\infty}^{\infty} N(m, n) w^m q^n = 1 + \sum_{n=1}^{\infty} \frac{q^{n^2}}{(wq; q)_n (w^{-1}q; q)_n}, \quad (1.4)$$

where $(a; q)_n := \prod_{j=0}^{n-1} (1 - aq^j)$. Dyson made the following conjectures which were proved by Atkin and Swinnerton-Dyer in 1953.

Theorem 1.1 ([ASD54]). *For all $n \geq 0$, we have the following identities:*

$$N(1, 5; 5n + 1) = N(2, 5; 5n + 1); \quad (5-1)$$

$$N(0, 5; 5n + 2) = N(2, 5; 5n + 2); \quad (5-2)$$

$$N(0, 5; 5n + 4) = N(1, 5; 5n + 4) = N(2, 5; 5n + 4); \quad (5-4)$$

$$N(2, 7; 7n) = N(3, 7; 7n); \quad (7-0)$$

$$N(1, 7; 7n + 1) = N(2, 7; 7n + 1) = N(3, 7; 7n + 1); \quad (7-1)$$

$$N(0, 7; 7n + 2) = N(3, 7; 7n + 2); \quad (7-2)$$

$$N(0, 7; 7n + 3) = N(2, 7; 7n + 3), \quad N(1, 7; 7n + 3) = N(3, 7; 7n + 3); \quad (7-3)$$

$$N(0, 7; 7n + 4) = N(1, 7; 7n + 4) = N(3, 7; 7n + 4); \quad (7-4)$$

$$N(0, 7; 7n + 5) = N(1, 7; 7n + 5) = N(2, 7; 7n + 5) = N(3, 7; 7n + 5); \quad (7-5)$$

$$N(0, 7; 7n + 6) + N(1, 7; 7n + 6) = N(2, 7; 7n + 6) + N(3, 7; 7n + 6). \quad (7-6)$$

Remark. By $N(a, b; n) = N(-a, b; n)$, the identity (5-4) implies

$$N(0, 5; 5n + 4) = N(1, 5; 5n + 4) = \cdots = N(4, 5; 5n + 4) = \frac{1}{5}p(5n + 4)$$

hence implies the Ramanujan congruence $p(5n + 4) \equiv 0 \pmod{5}$. The identity (7-5) implies

$$N(0, 7; 7n + 5) = N(1, 7; 7n + 5) = \cdots = N(6, 7; 7n + 5) = \frac{1}{7}p(7n + 5)$$

hence implies the Ramanujan congruence $p(7n + 5) \equiv 0 \pmod{7}$.

The proof of Theorem 1.1 in [ASD54] involves identities of generating functions

$$\sum_{n=0}^{\infty} (N(a, p; pn + k) - N(b, p; pn + k)) x^{pn} \prod_{r=1}^{\infty} (1 - x^r)$$

for $p = 5, 7$ and certain choices of the integer k . See [ASD54, Theorem 4 & Theorem 5] for details. Recently, Garvan [Gar17, §6] gave a new and simplified proof of Dyson's conjectures. For

$$\mathcal{K}_{p,0}(z) = \prod_{n=1}^{\infty} (1 - q^{pn}) \sum_{n=\lceil (p^2-1)/24p \rceil}^{\infty} \left(\sum_{k=0}^{p-1} N(k, p; pn + m - \frac{p^2-1}{24}) \zeta_p^k \right) q^n,$$

with $\zeta_p := e(\frac{1}{p})$ and $\mathcal{K}_{p,m}(z)$ defined in [Gar17, Definition 6.1], Garvan showed that $\mathcal{K}_{p,0}(z)$ is a weakly holomorphic modular form of weight 1 on $\Gamma_1(p)$. By the Valence formula, Garvan proved $\mathcal{K}_{5,0}(z) = \mathcal{K}_{7,0}(z) = 0$ and hence proved the Dyson's conjectures in [Gar17, §6.3].

For integers $b > a > 0$, denote $A(\frac{a}{b}; n)$ as the Fourier coefficient of $\mathcal{R}(\zeta_b^a; q)$:

$$\mathcal{R}(\zeta_b^a; q) =: 1 + \sum_{n=1}^{\infty} A\left(\frac{a}{b}; n\right) q^n.$$

There is an important equation which explains the relation between $A(\frac{a}{b}; n)$ and $N(a, b; n)$:

$$bN(a, b; n) = p(n) + \sum_{j=1}^{b-1} \zeta_b^{-aj} A\left(\frac{j}{b}; n\right). \quad (1.5)$$

It is not hard to show that $A(\frac{j}{b}; n) \in \mathbb{R}$ and $A(\frac{j}{b}; n) = A(1 - \frac{j}{b}; n)$ for $1 \leq j \leq b-1$, because $N(a, b; n) = N(-a, b; n)$ and $\zeta_b^{-aj} + \zeta_b^{-a(b-j)} = 2 \cos(\frac{\pi aj}{b})$. Specifically, if we know the values of $A(\frac{j}{b}; n)$ for $1 \leq j \leq b-1$, then we know the value of

$$N(a_1, b; n) - N(a_2, b; n) \quad \text{for any } 0 \leq a_1, a_2 \leq b-1.$$

Another way of approaching Dyson's conjectures is therefore via the formulae for $A(\frac{j}{b}; n)$ when $b = 5, 7$. In 2009, Bringmann [Bri09, Theorem 1.1] proved the asymptotic formula for $A(\frac{j}{b}; n)$ when $b \geq 3$ is odd and $0 \leq j \leq b-1$. Bringmann used the asymptotic formula when $b = 3$ to prove the Andrews-Lewis conjecture about comparing $N(0, 3; n)$ and $N(1, 3; n)$. In a previous paper [Sun24a], for each prime $p \geq 5$ and $1 \leq \ell \leq p-1$, the author proved that Bringmann's asymptotic formula, when summing up to infinity, is the exact formula for $A(\frac{\ell}{p}; n)$ for all $n \geq 0$. For each integer A , denote $[A]$ by

$$0 \leq [A] < 7: \quad [A] \equiv A \pmod{7}.$$

When $p = 5$ or 7 , that exact formula reduces to the following corollary.

Theorem 1.2 ([Sun24a, Corollary 2.2]). *For every positive integer n , when $p = 5$ and $1 \leq \ell \leq 4$, we have*

$$A\left(\frac{\ell}{5}; n\right) = \frac{2\pi e(-\frac{1}{8}) \sin(\frac{\pi\ell}{5})}{(24n-1)^{\frac{1}{4}}} \sum_{c>0: 5|c} \frac{S_{\infty\infty}^{(\ell)}(0, n, c, \mu_5)}{c} I_{\frac{1}{2}}\left(\frac{4\pi\sqrt{24n-1}}{24c}\right); \quad (1.6)$$

when $p = 7$ and $1 \leq \ell \leq 6$, we have

$$\begin{aligned} A\left(\frac{\ell}{7}; n\right) &= \frac{2\pi e(-\frac{1}{8}) \sin(\frac{\pi\ell}{7})}{(24n-1)^{\frac{1}{4}}} \sum_{c>0: 7|c} \frac{S_{\infty\infty}^{(\ell)}(0, n, c, \mu_7)}{c} I_{\frac{1}{2}}\left(\frac{4\pi\sqrt{24n-1}}{24c}\right) \\ &+ \frac{4\pi \sin(\frac{\pi\ell}{p})}{(24n-1)^{\frac{1}{4}}} \sum_{\substack{a>0: p \nmid a, \\ [a\ell]=1 \text{ or } 6}} \frac{S_{0\infty}^{(\ell)}(0, n, a, \mu_p; 0)}{\sqrt{7}a} I_{\frac{1}{2}}\left(\frac{4\pi\sqrt{24n-1}}{24 \times 7a}\right). \end{aligned} \quad (1.7)$$

Here $S_{\infty\infty}^{(\ell)}(0, n, c, \mu_p)$ for $p = 5, 7$ and $S_{0\infty}^{(\ell)}(0, n, a, \mu_7; 0)$ are given in (2.7), (2.8) and (2.9).

Theorem 1.2 gives us a new way to directly compute $A(\frac{\ell}{p}; n)$ for $p = 5, 7$ and all ℓ . In this paper, we give a new proof of Theorem 1.1 by establishing the following vanishing properties of the Kloosterman sums $S_{\infty\infty}^{(\ell)}(0, n, c, \mu_p)$ for $p = 5, 7$ and $S_{0\infty}^{(\ell)}(0, n, a, \mu_7; 0)$. This is totally different from the methods in [ASD54] and [Gar17].

Theorem 1.3. (i) *For all integers $n \geq 0$ and $1 \leq \ell \leq p-1$ for $p = 5, 7$ (denoted by $p|c$ below), we have the following vanishing conditions for the Kloosterman sums appeared in Corollary 1.2:*

- (5-4) *If $5|c$, we have $S_{\infty\infty}^{(\ell)}(0, 5n+4, c, \mu_5) = 0$.*
- (7-5,1) *If $7|c$, $\frac{c}{7} \cdot \ell \not\equiv 1 \pmod{7}$, and $\frac{c}{7} \cdot \ell \not\equiv -1 \pmod{7}$, then $S_{\infty\infty}^{(\ell)}(0, 7n+5, c, \mu_7; 0) = 0$.*
- (7-5,2) *If $7|c$, $7 \nmid a$, $a\ell \equiv \pm 1 \pmod{7}$, and $c = 7a$, we have*

$$e(-\frac{1}{8})S_{\infty\infty}^{(\ell)}(0, 7n+5, c, \mu_7) + 2\sqrt{7}S_{0\infty}^{(\ell)}(0, 7n+5, a, \mu_7; 0) = 0.$$

(ii) Furthermore, we denote $C_p^{a,b} := \cos(\frac{a\pi}{p}) - \cos(\frac{b\pi}{p})$ and

$$S_7^{(\ell)}(n, c) := \sin(\frac{\pi\ell}{7}) \left(e(-\frac{1}{8}) S_{\infty\infty}^{(\ell)}(0, n, c, \mu_7) + \mathbf{1}_{\substack{a:=c/7 \\ [a\ell]=1,6}} \cdot 2\sqrt{7} S_{0\infty}^{(\ell)}(0, 7n+5, a, \mu_7; 0) \right)$$

for simplicity, where $\mathbf{1}_{condition}$ equals 1 if the condition meets and equals 0 otherwise. We also have the following vanishing conditions for all $c \in \mathbb{Z}$ divisible by p , where $p = 5$ or 7 is marked at the subscript of $C_p^{a,b}$:

$$C_5^{2,4} \sin(\frac{\pi}{5}) S_{\infty\infty}^{(1)}(0, 5n+1, c, \mu_5) + C_5^{4,2} \sin(\frac{2\pi}{5}) S_{\infty\infty}^{(2)}(0, 5n+1, c, \mu_5) = 0, \quad (5-1)$$

$$C_5^{0,4} \sin(\frac{\pi}{5}) S_{\infty\infty}^{(1)}(0, 5n+2, c, \mu_5) + C_5^{0,2} \sin(\frac{2\pi}{5}) S_{\infty\infty}^{(2)}(0, 5n+2, c, \mu_5) = 0, \quad (5-2)$$

$$C_7^{4,6} S_7^{(1)}(7n, c) + C_7^{6,2} S_7^{(2)}(7n, c) + C_7^{2,4} S_7^{(3)}(7n, c) = 0, \quad (7-0)$$

$$C_7^{2,4} S_7^{(1)}(7n+1, c) + C_7^{4,6} S_7^{(2)}(7n+1, c) + C_7^{6,2} S_7^{(3)}(7n+1, c) = 0, \quad (7-1,1)$$

$$C_7^{4,6} S_7^{(1)}(7n+1, c) + C_7^{6,2} S_7^{(2)}(7n+1, c) + C_7^{2,4} S_7^{(3)}(7n+1, c) = 0, \quad (7-1,2)$$

$$C_7^{0,6} S_7^{(1)}(7n+2, c) + C_7^{0,2} S_7^{(2)}(7n+2, c) + C_7^{0,4} S_7^{(3)}(7n+2, c) = 0, \quad (7-2)$$

$$C_7^{0,4} S_7^{(1)}(7n+3, c) + C_7^{0,6} S_7^{(2)}(7n+3, c) + C_7^{0,2} S_7^{(3)}(7n+3, c) = 0, \quad (7-3,1)$$

$$C_7^{2,6} S_7^{(1)}(7n+3, c) + C_7^{4,2} S_7^{(2)}(7n+3, c) + C_7^{6,4} S_7^{(3)}(7n+3, c) = 0, \quad (7-3,2)$$

$$C_7^{0,2} S_7^{(1)}(7n+4, c) + C_7^{0,4} S_7^{(2)}(7n+4, c) + C_7^{0,6} S_7^{(3)}(7n+4, c) = 0, \quad (7-4,1)$$

$$C_7^{2,6} S_7^{(1)}(7n+4, c) + C_7^{4,2} S_7^{(2)}(7n+4, c) + C_7^{6,4} S_7^{(3)}(7n+4, c) = 0, \quad (7-4,2)$$

$$\begin{aligned} & (C_7^{0,4} + C_7^{2,6}) S_7^{(1)}(7n+6, c) + (C_7^{0,6} + C_7^{4,2}) S_7^{(2)}(7n+6, c) \\ & + (C_7^{0,2} + C_7^{6,4}) S_7^{(3)}(7n+6, c) = 0, \end{aligned} \quad (7-6)$$

Using (1.5) and Theorem 1.2, we have the following corollary.

Corollary 1.4. For any pair $(p-k)$ (or $(p-k, t)$ for both $t = 1, 2$) in Theorem 1.3 with

$$p = 5, k \in \{1, 2, 4\} \quad \text{or} \quad p = 7, k \in \{0, 1, 2, 3, 4, 5, 6\},$$

we have Dyson's conjecture $(p-k)$ in Theorem 1.1.

The paper is organized as follows. In Section 2 we review about our notations of vector-valued Kloosterman sums as established in [Sun24a]. In Section 3 we provide a detailed proof of (5-4) of Theorem 1.3. In Section 4 and Section 5 we prove (7-5,1) and (7-5,2) of Theorem 1.3, respectively. Section 6 covers the proofs of (5-1), (5-2) and (7-0) of Theorem 1.3 but does not include the repetitive proofs for the remaining parts labeled (7- k).

We have omitted detailed proofs for similar cases to avoid redundancy. The complete proof, along with the Mathematica code used for generating illustrative graphs within the proofs, are available in our GitHub repository [Sun24b].

2. NOTATIONS

In this section we define some notation involving Dedekind sums and Kloosterman sums. For the origin of these notations, see [Gar17, Sun24a].

For integers d and $m \geq 1$, let $\overline{d_{\{m\}}}$ denote the inverse of $d \pmod{m}$. If there is a subscript, e.g. d_1 , then we write $\overline{d_{1\{m\}}}$ as the inverse of $d_1 \pmod{m}$.

Define

$$((x)) := \begin{cases} x - [x] - \frac{1}{2}, & \text{when } x \in \mathbb{R} \setminus \mathbb{Z}, \\ 0, & \text{when } x \in \mathbb{Z}. \end{cases}$$

For integers $c > 0$ and $(d, c) = 1$, we define the Dedekind sum as

$$s(d, c) := \sum_{r \pmod{c}} \left(\left(\frac{r}{c} \right) \right) \left(\left(\frac{dr}{c} \right) \right). \quad (2.1)$$

The Dedekind sums have the following properties [Lew95, (4.2)-(4.5)]:

$$2\theta cs(d, c) \in \mathbb{Z}, \quad \text{where } \theta = \gcd(c, 3), \quad (2.2)$$

$$12cs(d, c) \equiv d + \overline{d_{\{\theta c\}}} \pmod{\theta c}, \quad (2.3)$$

$$12cs(d, c) \equiv c + 1 - 2\left(\frac{d}{c}\right) \pmod{8}, \quad \text{if } c \text{ is odd}, \quad (2.4)$$

$$12cs(d, c) \equiv d + \left(c^2 + 3c + 1 + 2c\left(\frac{c}{d}\right)\right) \overline{d_{\{8 \times 2^\lambda\}}} \pmod{8 \times 2^\lambda}, \quad \text{if } 2^\lambda \parallel c \text{ for } \lambda \geq 1. \quad (2.5)$$

These congruences determine $12cs(d, c) \pmod{24c}$ uniquely in every case ($2|c$ or $2 \nmid c$, $3|c$ or $3 \nmid c$).

In the proof we use the following quadratic reciprocity of the Kronecker symbol (\cdot) . For any non-zero integer n , write $n = 2^\lambda n_o$ where n_o is odd. For integers m, n with $(m, n) = 1$, we have

$$\left(\frac{m}{n}\right) \left(\frac{n}{m}\right) = \pm (-1)^{(m_o-1)(n_o-1)/4}, \quad (2.6)$$

where we take $+$ if $m \geq 0$ or $n \geq 0$, and we take $-$ if $m < 0$ and $n < 0$.

Next we define the Kloosterman sums $S_{\infty\infty}^{(\ell)}(0, n, c, \mu_p)$ for $p = 5, 7$ and $S_{0\infty}^{(\ell)}(0, n, c, \mu_7; 0)$ appearing at Corollary 1.2. We follow the notations of vector-valued Kloosterman sums in [Sun24a, §4.3]. From [Sun24a, (5.19), (5.29)], when $p|c$ we have

$$S_{\infty\infty}^{(\ell)}(0, n, c, \mu_p) = e(-\frac{1}{8}) \sum_{\substack{d \pmod{c}^* \\ ad \equiv 1 \pmod{c}}} \frac{(-1)^{\ell c} e(-\frac{3ca\ell}{2p^2})}{\sin(\frac{\pi a\ell}{p})} e^{-\pi i s(d, c)} e\left(\frac{nd}{c}\right). \quad (2.7)$$

When $p = 7$, recall that $[A\ell]$ is the least non-negative residue of $A\ell \pmod{7}$. From [Sun24a, (5.31)], when $A\ell = 7T + 1$ for some integer $T \geq 0$, we have

$$S_{0\infty}^{(\ell)}(0, n, A, \mu_7; 0) = (-1)^{A\ell - [A\ell]} \sum_{\substack{B \pmod{A}^* \\ 0 < C < 7A, 7|C \\ BC \equiv -1(A)}} e\left(\frac{(\frac{3}{2}T^2 + \frac{1}{2}T)C}{A}\right) e^{-\pi i s(B, A)} e\left(\frac{nB}{A}\right). \quad (2.8)$$

When $A\ell = 7T - 1$ for some integer $T \geq 1$, we have

$$\begin{aligned} & S_{0\infty}^{(\ell)}(0, n, A, \mu_7; 0) \\ &= (-1)^{A\ell - [A\ell]} \sum_{\substack{B \pmod{A}^* \\ 0 < C < 7A, 7|C \\ BC \equiv -1(A)}} e\left(\frac{(\frac{3}{2}(T-1)^2 + \frac{5}{2}(T-1) + 1)C}{A}\right) e^{-\pi i s(B, A)} e\left(\frac{nB}{A}\right). \end{aligned} \quad (2.9)$$

If $[A\ell] \neq 1$ and $[A\ell] \neq 6$, then $S_{0\infty}^{(\ell)}(0, n, A, \mu_7; 0) := 0$.

3. PROOF OF (5-4) OF THEOREM 1.3

In this section we prove (5-4) of Theorem 1.3. We only consider $\ell = 1, 2$ because $A(\frac{\ell}{p}; n) = A(1 - \frac{\ell}{p}; n)$.

Define $c' := c/5$. For any integer r with $(r, c') = 1$, we define

$$V(r, c) := \{d \pmod{c}^* : d \equiv r \pmod{c'}\}.$$

For example, $V(1, 30) = \{d \pmod{30}^* : d \equiv 1, 7, 13, 19 \pmod{30}\}$ and $V(4, 25) = \{d \pmod{25}^* : d \equiv 4, 9, 14, 19, 24 \pmod{25}\}$. Clearly, $|V(r, c)| = 4$ if $5 \nmid c$ and $|V(r, c)| = 5$ if $25 \mid c$. Moreover, $(\mathbb{Z}/c\mathbb{Z})^*$ is the disjoint union

$$(\mathbb{Z}/c\mathbb{Z})^* = \bigcup_{r \pmod{c'}^*} V(r, c).$$

By (2.7) we have

$$e(\frac{1}{8})S_{\infty\infty}^{(\ell)}(0, 5n+4, c, \mu_5) = \sum_{\substack{d \pmod{c}^* \\ ad \equiv 1 \pmod{c}}} \frac{(-1)^{\ell c} e(-\frac{3c' a \ell^2}{10})}{\sin(\frac{\pi a \ell}{5})} e^{-\pi i s(d, c)} e\left(\frac{(5n+4)d}{c}\right). \quad (3.1)$$

We claim the following proposition.

Proposition 3.1. *For $\ell = 1, 2$, the sum on $V(r, c)$ satisfies*

$$s_{r, c} := \sum_{\substack{d \in V(r, c) \\ ad \equiv 1 \pmod{c}}} \frac{e(-\frac{3c' a \ell^2}{10})}{\sin(\frac{\pi a \ell}{5})} e^{-\pi i s(d, c)} e\left(\frac{4d}{c}\right) = 0. \quad (3.2)$$

If Proposition 3.1 is true, then

$$S_{\infty\infty}^{(\ell)}(0, 5n+4, c, \mu_5) = e(-\frac{1}{8})(-1)^{\ell c} \sum_{r \pmod{c'}^*} s_{r, c} e\left(\frac{nr}{c'}\right) (-1)^{\ell c} = 0$$

for all $n \in \mathbb{Z}$, $\ell = 1, 2$, and we have proved (5-4) of Theorem 1.3.

In the following subsections §7.1-§7.4, we prove Proposition 3.1 when $5 \nmid c$. In §7.5, we prove Proposition 3.1 when $25 \mid c$. Suppose now that $5 \mid c$. Since $|V(r, c)| = 4$, let $\beta \in \{1, 2, 3, 4\}$ such that $\beta c' \equiv 1 \pmod{5}$ and we make a special choice of $V(r, c)$ as

$$V(r, c) = \{d_1, d_2, d_3, d_4\} \quad \text{where } d_j \equiv j \pmod{5} \text{ and } d_{j+1} = d_1 + j\beta c'. \quad (3.3)$$

We also take a_j for $j \in \{1, 2, 3, 4\}$ such that $a_j \equiv j \pmod{5}$, $a_{j+1} = a_1 + j\beta c'$, and

$$a_{\overline{j}_{\{5\}}} d_j \equiv 1 \pmod{c}. \quad (3.4)$$


These choices do not affect the sum (3.2) because $s_{r, c}$ has period c in both a and d . In (3.2), we denote each summation term as

$$P(d) := \frac{e\left(-\frac{3c' a \ell^2}{10}\right)}{\sin(\frac{\pi a \ell}{5})} \cdot e\left(-\frac{12cs(d, c)}{24c}\right) \cdot e\left(\frac{4d}{c}\right) =: P_1(d) \cdot P_2(d) \cdot P_3(d) \quad (3.5)$$

where $P_1(d) := e(-\frac{3c' a \ell^2}{10})/\sin(\frac{\pi a \ell}{5})$, $P_2(d) := \exp(-\pi i s(d, c))$, and $P_3(d) := e(\frac{4d}{c})$.

Remark. We keep $24c$ in the denominator of $P_2(d)$ because the congruence properties of the Dedekind sum are of the form $12cs(d, c)$. See (2.2)-(2.5) for details.

We claim that the set of points $P(d)$ for $d \in V(r, c)$ must have the relative position illustrated in one of the following six configurations. Here $0 < d_j < c$ for simplicity but we use (3.3) in the proof.



Here we explain the styles. Each graph above has two circles with inner one of radius $\csc(\frac{2\pi}{5})$ and outer one with radius $\csc(\frac{\pi}{5})$. When $\ell = 1$, the value of $P(d_1)$ and $P(d_4)$ will be on the outer circle ($P(d_2)$ and $P(d_3)$ on the inner circle) because the term $P_1(d_j)$ has denominator $\sin(\frac{\pi a_j \ell}{5})$. When $\ell = 2$, $P(d_1)$ and $P(d_4)$ will be on the inner circle.

We describe the relative argument differences via the following notation. Let

$$\text{Arg}_j(d_u \rightarrow d_v; \ell), \quad \text{for } j \in \{1, 2, 3\}, \quad u, v \in \{1, 2, 3, 4\}, \quad \text{and } \ell \in \{1, 2\} \quad (3.6)$$

be the argument difference (as the proportion of 2π , positive when going counter-clockwise) contributed from P_j going from d_u to d_v when $\ell \in \{1, 2\}$. To be precise, if we denote $P_j(d_u) = R_{j,u} \exp(i\Theta_{j,u})$ for $R_{j,u}, \Theta_{j,u} \in \mathbb{R}$, then

$$\text{Arg}_j(d_u \rightarrow d_v; \ell) = \alpha \iff \Theta_{j,v} - \Theta_{j,u} = \alpha \cdot 2\pi + 2k\pi \text{ for some } k \in \mathbb{Z}.$$

We say that two argument differences equal: $\text{Arg}_j(d_u \rightarrow d_v; \ell) = \text{Arg}_j(d_w \rightarrow d_x; \ell)$ or say $\text{Arg}_j(d_u \rightarrow d_v; \ell) = \alpha$ if their difference is an integer.

Although the P_2 and P_3 terms are not affected by the value of ℓ in (3.5), we still use the notations $\text{Arg}_2(d_u \rightarrow d_v; \ell)$ and $\text{Arg}_e(d_u \rightarrow d_v; \ell)$ to indicate the different cases for ℓ . Moreover, we define

$$\text{Arg}(d_u \rightarrow d_v; \ell) := \sum_{j=1}^3 \text{Arg}_j(d_u \rightarrow d_v; \ell) \quad (3.7)$$

as the argument difference in total.

The following condition ensures Proposition 3.1.

Condition 3.2. *We have the following six styles for the relative position of these four points.*

- $\ell = 1$. *First graph style: the arguments going $d_1 \rightarrow d_2 \rightarrow d_3 \rightarrow d_4 \rightarrow d_1$ are $\frac{3}{10}$, $\frac{1}{10}$, $\frac{3}{10}$, and $\frac{3}{10}$, respectively. The second graph style is that all the argument differences are $\frac{1}{2}$, while the third graph style has the reversed order of rotation compared with the first one.*

$\text{Arg}(d_u \rightarrow d_v; 1) \searrow$	d_1	\rightarrow	d_2	\rightarrow	d_3	\rightarrow	d_4	\rightarrow	d_1
$c' \equiv 1 \pmod{5}$			$\frac{3}{10}$		$\frac{1}{10}$		$\frac{3}{10}$		$\frac{3}{10}$
$c' \equiv 2, 3 \pmod{5}$			$\frac{1}{2}$		$\frac{1}{2}$		$\frac{1}{2}$		$\frac{1}{2}$
$c' \equiv 4 \pmod{5}$			$-\frac{3}{10}$		$-\frac{1}{10}$		$-\frac{3}{10}$		$-\frac{3}{10}$

- $\ell = 2$. *Here are the styles for the graphs in the second row.*

$\text{Arg}(d_u \rightarrow d_v; 2) \searrow$	d_1	\rightarrow	d_2	\rightarrow	d_3	\rightarrow	d_4	\rightarrow	d_1
$c' \equiv 3 \pmod{5}$			$-\frac{2}{5}$		$-\frac{3}{10}$		$-\frac{2}{5}$		$\frac{1}{10}$
$c' \equiv 1, 4 \pmod{5}$			0		$\frac{1}{2}$		0		$\frac{1}{2}$
$c' \equiv 2 \pmod{5}$			$\frac{2}{5}$		$\frac{3}{10}$		$\frac{2}{5}$		$-\frac{1}{10}$

One can check that, whenever the four points on \mathbb{C} satisfy any of the above cases of relative argument differences and corresponding radii, their sum becomes 0. This can be explained by

$$\frac{\cos(\frac{\pi}{10})}{\sin(\frac{2\pi}{5})} = \frac{\cos(\frac{3\pi}{10})}{\sin(\frac{\pi}{5})} = 1, \quad \text{where } \frac{1}{\sin(\frac{\pi}{5})} \text{ and } \frac{1}{\sin(\frac{2\pi}{5})} \text{ are the radii.}$$

Before we divide into the cases, we first claim the following lemma:

Lemma 3.3. *For $\ell \in \{1, 2\}$, we have*

$$\text{Arg}(d_1 \rightarrow d_2; \ell) + \text{Arg}(d_4 \rightarrow d_3; \ell) = 0 \quad \text{and} \quad \text{Arg}(d_1 \rightarrow d_3; \ell) + \text{Arg}(d_4 \rightarrow d_2; \ell) = 0. \quad (3.8)$$

Granted the above reduction, to prove that each case of the argument differences are one of the cases in Condition 3.2, we only need to verify that

$$\text{Arg}(d_1 \rightarrow d_4; \ell) \quad \text{and} \quad \text{Arg}(d_1 \rightarrow d_2; \ell) \quad \text{for } \ell = 1, 2$$

satisfy Condition 3.2. We prove this by enumerating all of the cases. We can list the argument differences for Arg_1 and Arg_3 , but for Arg_2 , we require the congruence properties of Dedekind sums in (2.2)-(2.5).

Proof of Lemma 3.3. Note that

$$\text{Arg}(d_u \rightarrow d_v; \ell) = \text{Arg}(d_u \rightarrow d_w; \ell) + \text{Arg}(d_w \rightarrow d_v; \ell)$$

for all $u, v, w \in \{1, 2, 3, 4\}$. Then it suffices to prove

$$\text{Arg}(d_1 \rightarrow d_2; \ell) = \text{Arg}(d_3 \rightarrow d_4; \ell).$$

Recall our notation for d_j and a_j in (3.3). Since $a_3 - a_1 = a_4 - a_2 = 2\beta c'$, one can show $\text{Arg}_1(d_1 \rightarrow d_2; \ell) = \text{Arg}_1(d_3 \rightarrow d_4; \ell)$ by

$$\text{sgn}\left(\sin\left(\frac{\pi a_3 \ell}{5}\right) / \sin\left(\frac{\pi a_1 \ell}{5}\right)\right) = \text{sgn}\left(\sin\left(\frac{\pi a_4 \ell}{5}\right) / \sin\left(\frac{\pi a_2 \ell}{5}\right)\right) = 1.$$

It is also easy to show $\text{Arg}_3(d_1 \rightarrow d_2; \ell) = \text{Arg}_3(d_3 \rightarrow d_4; \ell)$.

For Arg_2 , we apply (2.3), (2.4) and (2.5) with the Chinese Remainder Theorem to show that

$$12cs(d_2, c) - 12cs(d_1, c) \equiv 12cs(d_4, c) - 12cs(d_3, c) \text{ for all the corresponding congruences.}$$

When $\gcd(c, 3) = 1$, we have

$$\begin{aligned} 12cs(d_2, c) - 12cs(d_1, c) &\equiv d_2 + a_3 - d_1 - a_1 \equiv 3\beta c' \pmod{c}, \\ 12cs(d_4, c) - 12cs(d_3, c) &\equiv d_4 + a_4 - d_3 - a_2 \equiv 3\beta c' \pmod{c}, \\ 12cs(d_2, c) - 12cs(d_1, c) &\equiv 12cs(d_4, c) - 12cs(d_3, c) \equiv 0 \pmod{6}. \end{aligned}$$

When $3|c$, we apply the congruence

$$\overline{(x+y)_{\{m\}}} - \overline{x_{\{m\}}} \equiv -y \overline{(x+y)_{\{m\}}} \cdot \overline{x_{\{m\}}} \pmod{m} \quad (3.9)$$

to compute

$$\begin{aligned} d_2 + \overline{d_{2\{3c\}}} - d_1 - \overline{d_{1\{3c\}}} &\equiv \beta c' (1 - \overline{d_{2\{3c\}}} \cdot \overline{d_{1\{3c\}}}) \pmod{3c}, \\ d_4 + \overline{d_{4\{3c\}}} - d_3 - \overline{d_{3\{3c\}}} &\equiv \beta c' (1 - \overline{d_{4\{3c\}}} \cdot \overline{d_{3\{3c\}}}) \pmod{3c}, \end{aligned}$$

which imply

$$12cs(d_2, c) - 12cs(d_1, c) \equiv 12cs(d_4, c) - 12cs(d_3, c) \equiv 0 \pmod{c'}.$$

by (2.3). After dividing by c' (recall that the denominator of $P_2(d)$ is $24c$), we have

$$\begin{aligned} 60s(d_2, c) - 60s(d_1, c) &\equiv \beta (1 - \overline{d_{2\{3c\}}} \cdot \overline{d_{1\{3c\}}}) \equiv \beta (1 - a_3 a_1) \pmod{15}, \\ 60s(d_4, c) - 60s(d_3, c) &\equiv \beta (1 - \overline{d_{4\{3c\}}} \cdot \overline{d_{3\{3c\}}}) \equiv \beta (1 - a_4 a_2) \pmod{15}, \end{aligned}$$

because of (3.4) and $\overline{x_{\{un\}}} \equiv \overline{x_{\{vn\}}} \pmod{n}$. Since $a_3 \equiv a_1 \pmod{3}$ and $a_4 \equiv a_2 \pmod{3}$, we have $a_3 a_1 \equiv a_4 a_2 \equiv 1 \pmod{3}$. Moreover, $a_3 a_1 \equiv a_4 a_2 \equiv 3 \pmod{5}$. Hence $a_3 a_1 \equiv a_4 a_2 \equiv 13 \pmod{15}$ and we get

$$60s(d_2, c) - 60s(d_1, c) \equiv 60s(d_4, c) - 60s(d_3, c) \equiv 3\beta \pmod{15}.$$

When c is odd, by (2.4) and $d_{j_1} \equiv d_{j_2} \pmod{c'}$, we have

$$\begin{aligned} 12cs(d_2, c) - 12cs(d_1, c) &\equiv 2\left(\frac{d_1}{c}\right) - 2\left(\frac{d_1}{c}\right) \equiv 2\left(\frac{1}{5}\right)\left(\frac{d_1}{c'}\right) - 2\left(\frac{2}{5}\right)\left(\frac{d_2}{c'}\right) \equiv 4 \pmod{8}, \\ 12cs(d_4, c) - 12cs(d_3, c) &\equiv 2\left(\frac{d_4}{c}\right) - 2\left(\frac{d_3}{c}\right) \equiv 2\left(\frac{4}{5}\right)\left(\frac{d_4}{c'}\right) - 2\left(\frac{3}{5}\right)\left(\frac{d_3}{c'}\right) \equiv 4 \pmod{8}. \end{aligned}$$

When c is even and $2^\lambda || c$ for $\lambda \geq 1$, by (2.5) we have

$$\begin{aligned} 12cs(d_2, c) - 12cs(d_1, c) &\equiv d_2 + (c^2 + 3c + 1)\overline{d_{2\{8 \times 2^\lambda\}}} + 2c\left(\frac{c}{d_2}\right)\overline{d_{2\{8 \times 2^\lambda\}}} \\ &\quad - d_1 + (c^2 + 3c + 1)\overline{d_{1\{8 \times 2^\lambda\}}} + 2c\left(\frac{c}{d_1}\right)\overline{d_{1\{8 \times 2^\lambda\}}} \\ &\equiv \beta c' (1 - \overline{d_{2\{8 \times 2^\lambda\}}} \cdot \overline{d_{1\{8 \times 2^\lambda\}}}) \\ &\quad + 2c\left(\frac{c}{d_2}\right)\overline{d_{2\{8 \times 2^\lambda\}}} - 2c\left(\frac{c}{d_1}\right)\overline{d_{1\{8 \times 2^\lambda\}}} \pmod{8 \times 2^\lambda}. \end{aligned}$$

Hence $12cs(d_2, c) - 12cs(d_1, c)$ is a multiple of c' . Dividing c' and by $x^2 \equiv 1 \pmod{8}$ for odd x we have

$$60s(d_2, c) - 60s(d_1, c) \equiv \beta (1 - (c^2 + 3c + 1)d_2 d_1) + 2\left(\frac{c}{d_2}\right)d_2 - 2\left(\frac{c}{d_1}\right)d_1 \pmod{8}.$$

Similarly, $12cs(d_4, c) - 12cs(d_3, c)$ is a multiple of c' and

$$60s(d_4, c) - 60s(d_3, c) \equiv \beta (1 - (c^2 + 3c + 1)d_4 d_3) + 2\left(\frac{c}{d_4}\right)d_4 - 2\left(\frac{c}{d_3}\right)d_3 \pmod{8}.$$

Dividing into cases for $4|c$ or $2||c$ with $c' \equiv 2$ or $6 \pmod{8}$, one can conclude

$$d_2 d_1 \equiv d_4 d_3 \pmod{8}.$$

For the remaining part, we only need to determine $(\frac{c}{d_j})d_j \equiv \pm 1 \pmod{4}$ for $j \in \{1, 2, 3, 4\}$. Since $d_3 \equiv d_1 \pmod{4}$ and $d_2 \equiv d_4 \pmod{4}$, it is not hard to show that

$$\left(\frac{c}{d_2}\right)d_2 - \left(\frac{c}{d_1}\right)d_1 \equiv \left(\frac{c}{d_4}\right)d_4 - \left(\frac{c}{d_3}\right)d_3 \pmod{4}.$$

Combining all the congruence equations in this proof, we have shown

$$\text{Arg}_2(d_1 \rightarrow d_2; \ell) = \text{Arg}_2(d_3 \rightarrow d_4; \ell) \quad \text{for } \ell \in \{1, 2\}$$

by proving

$$\frac{12cs(d_2, c) - 12cs(d_1, c)}{24c} - \frac{12cs(d_4, c) - 12cs(d_3, c)}{24c} \in \mathbb{Z}$$

in all the cases for c ($2|c$ or $2 \nmid c$, $3|c$ or $3 \nmid c$). The lemma follows. \square

Now we begin to prove that $\text{Arg}(d_1 \rightarrow d_4; \ell)$ and $\text{Arg}(d_1 \rightarrow d_2; \ell)$ both satisfy Condition 3.2 in all the cases of $5|c$.

3.1. Case $2 \nmid c'$, $3 \nmid c'$, and $5 \nmid c'$. We first treat the case when $c' \equiv 1, 7, 11, 13, 17, 19, 23, 29 \pmod{30}$. Recall our notations in (3.3) and (3.4):

$$d_4 = d_1 + 3\beta c', \quad d_2 = d_1 + \beta c', \quad a_4 = a_1 + 3\beta c', \quad a_3 = a_1 + 2\beta c', \quad \beta c' \equiv 1 \pmod{5}.$$

The argument differences $\text{Arg}_j(d_1 \rightarrow d_4; \ell)$ for $j = 1, 2, 3$ are given by the arguments of

$$\frac{e\left(-\frac{9}{10}\beta c'^2 \ell^2\right)}{\text{sgn}\left(\sin\left(\frac{\pi a_4 \ell}{5}\right)/\sin\left(\frac{\pi a_1 \ell}{5}\right)\right)}, \quad e\left(-\frac{12cs(d_4, c) - 12cs(d_1, c)}{24c}\right), \quad \text{and } e\left(\frac{2\beta}{5}\right),$$

respectively. First we have

$$\text{sgn}\left(\sin\left(\frac{\pi a_4 \ell}{5}\right)/\sin\left(\frac{\pi a_1 \ell}{5}\right)\right) = -1 \quad \text{whenever } \begin{cases} \ell = 1 & \text{and} \\ 3\beta c' \equiv 8 \pmod{10} \end{cases} \quad \text{or } \ell = 2. \quad (3.10)$$

This is easy to prove because $3\beta c' \times 2 \equiv 6 \pmod{10}$.

By (2.3), we have $\theta = 1$ and

$$-12cs(d_4, c) + 12cs(d_1, c) \equiv -d_4 - a_4 + d_1 + a_1 \equiv -6\beta c' \equiv -\beta c' \pmod{c}. \quad (3.11)$$

Moreover, we have $-12cs(d_4, c) + 12cs(d_1, c) \equiv 0 \pmod{6}$ and

$$-12cs(d_4, c) + 12cs(d_1, c) \equiv 2\left(\left(\frac{d_4}{c}\right) - \left(\frac{d_1}{c}\right)\right) \equiv 2\left(\left(\frac{d_4}{5}\right)\left(\frac{d_4}{c'}\right) - \left(\frac{d_1}{5}\right)\left(\frac{d_1}{c'}\right)\right) \equiv 0 \pmod{8}.$$

Here we have used $\left(\frac{d_j}{5}\right) = 1$ for $j = 1, 4$ and $d_j \equiv d_1 \pmod{c'}$ for all j . Then,

$$-12cs(d_4, c) + 12cs(d_1, c) \equiv 0 \pmod{24}. \quad (3.12)$$

Combining (3.11) and (3.12), since c' is odd, we can divide both the denominator and numerator in P_2 by $24c'$. We obtain

$$\text{Arg}_2(d_1 \rightarrow d_4; \ell) = \frac{\beta}{5}.$$

Now we have Table 3.1. In the row of $\text{Arg}_1(d_1 \rightarrow d_4; 1)$, we see $+\frac{1}{2}$ because the sign difference $\text{sgn}\left(\sin\left(\frac{\pi a_4 \ell}{5}\right)/\sin\left(\frac{\pi a_1 \ell}{5}\right)\right) = -1$ when $3\beta c' \equiv 8 \pmod{10}$. The $\text{Arg}_1(d_1 \rightarrow d_4; 2)$ contains the term $+\frac{1}{2}$ because $3\beta c' \times 2 \equiv 6 \pmod{10}$. The upper half of the table is for the case $\ell = 1$ and the lower half is for $\ell = 2$.

$c' \pmod{30}$	1	7	11	13	17	19	23	29
β	1	3	1	2	3	4	2	4
$3\beta c' \pmod{10}$	3	3	3	$\boxed{8}$	3	$\boxed{8}$	$\boxed{8}$	$\boxed{8}$
$-9\beta c'^2 \pmod{10}$	1	7	1	8	7	4	7	4
$\text{Arg}_1(d_1 \rightarrow d_4; 1)$	$\frac{1}{10}$	$-\frac{3}{10}$	$\frac{1}{10}$	$-\frac{2}{10} + \frac{1}{2}$	$-\frac{3}{10}$	$-\frac{6}{10} + \frac{1}{2}$	$-\frac{3}{10} + \frac{1}{2}$	$-\frac{6}{10} + \frac{1}{2}$
$\text{Arg}_2(d_1 \rightarrow d_4; 1)$	$\frac{1}{5}$	$\frac{3}{5}$	$\frac{1}{5}$	$\frac{2}{5}$	$\frac{3}{5}$	$\frac{4}{5}$	$\frac{2}{5}$	$\frac{4}{5}$
$\text{Arg}_3(d_1 \rightarrow d_4; 1)$	$\frac{2}{5}$	$\frac{1}{5}$	$\frac{2}{5}$	$\frac{4}{5}$	$\frac{1}{5}$	$\frac{3}{5}$	$\frac{4}{5}$	$\frac{3}{5}$
Total $\text{Arg}(d_1 \rightarrow d_4; 1)$	$-\frac{3}{10}$	$\frac{1}{2}$	$-\frac{3}{10}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{3}{10}$	$\frac{1}{2}$	$\frac{3}{10}$
$3\beta c' \pmod{10}$	3	3	3	8	3	8	8	8
$-18\beta c'^2 \equiv 2c' \pmod{5}$	2	4	2	1	4	3	1	3
$\text{Arg}_1(d_1 \rightarrow d_4; 2) : \frac{1}{2} + \frac{2c'}{5}$	$-\frac{1}{10}$	$\frac{3}{10}$	$-\frac{1}{10}$	$-\frac{3}{10}$	$\frac{3}{10}$	$\frac{1}{10}$	$-\frac{3}{10}$	$\frac{1}{10}$
$\text{Arg}_2(d_1 \rightarrow d_4; 2)$	$\frac{1}{5}$	$\frac{3}{5}$	$\frac{1}{5}$	$\frac{2}{5}$	$\frac{3}{5}$	$\frac{4}{5}$	$\frac{2}{5}$	$\frac{4}{5}$
$\text{Arg}_3(d_1 \rightarrow d_4; 2)$	$\frac{2}{5}$	$\frac{1}{5}$	$\frac{2}{5}$	$\frac{4}{5}$	$\frac{1}{5}$	$\frac{3}{5}$	$\frac{4}{5}$	$\frac{3}{5}$
Total $\text{Arg}(d_1 \rightarrow d_4; 2)$	$\frac{1}{2}$	$\frac{1}{10}$	$\frac{1}{2}$	$-\frac{1}{10}$	$\frac{1}{10}$	$\frac{1}{2}$	$-\frac{1}{10}$	$\frac{1}{2}$

TABLE 3.1. Table for $\text{Arg}(d_1 \rightarrow d_4; \ell)$; $2 \nmid c$, $3 \nmid c$, $5 \nmid c$.

For $\text{Arg}_j(d_1 \rightarrow d_2; \ell)$, recall $a_3 d_2 \equiv 1 \pmod{c}$. The argument differences $\text{Arg}_j(d_1 \rightarrow d_2; \ell)$ for $j = 1, 2, 3$ are given by

$$\frac{e\left(-\frac{3}{5}\beta c'^2 \ell^2\right)}{\text{sgn}\left(\sin\left(\frac{\pi a_3 \ell}{5}\right) / \sin\left(\frac{\pi a_1 \ell}{5}\right)\right)}, \quad e\left(-\frac{12cs(d_2, c) - 12cs(d_1, c)}{24c}\right), \quad e\left(\frac{4\beta}{5}\right),$$

respectively. Since $2\beta c' \ell \equiv 2\ell \pmod{10}$, we always have

$$\text{sgn}\left(\sin\left(\frac{\pi a_3}{5}\right) / \sin\left(\frac{\pi a_1}{5}\right)\right) = 1 \quad \text{and} \quad \text{sgn}\left(\sin\left(\frac{2\pi a_3}{5}\right) / \sin\left(\frac{2\pi a_1}{5}\right)\right) = -1. \quad (3.13)$$

Moreover, from (2.3) we have

$$12cs(d_2, c) - 12cs(d_1, c) \equiv d_2 + a_3 - d_1 - a_1 \equiv 3\beta c' \pmod{c}, \quad (3.14)$$

$$12cs(d_2, c) - 12cs(d_1, c) \equiv -2\left(-\left(\frac{d_2}{c'}\right) - \left(\frac{d_1}{c'}\right)\right) \equiv 4 \pmod{8},$$

$12cs(d_2, c) - 12cs(d_1, c) \equiv 0 \pmod{6}$, and

$$12cs(d_2, c) - 12cs(d_1, c) \equiv 12 \pmod{24}. \quad (3.15)$$

Combining (3.14) and (3.15), we divide by c' and determine the unique value modulo 120:

$$-(60s(d_2, c) - 60s(d_1, c)) \quad \text{congruent to} \quad -3\beta \pmod{5} \quad \text{and} \quad 12 \pmod{24}.$$

This gives the contribution of the argument difference from P_2 . Now we can make Table 3.2.

Combining Table 3.1 and Table 3.2, we see that $\text{Arg}(d_1 \rightarrow d_4; \ell)$ and $\text{Arg}(d_1 \rightarrow d_2; \ell)$ for $\ell = 1, 2$ satisfy the styles in Condition 3.2. This finishes the proof when $2 \nmid c'$, $3 \nmid c'$ and $5 \nmid c'$.

$c' \pmod{30}$	1	7	11	13	17	19	23	29
β	1	3	1	2	3	4	2	4
$-3c' \pmod{5}$	2	4	2	1	4	3	1	3
$\text{Arg}_1(d_1 \rightarrow d_2; 1)$	$\frac{2}{5}$	$\frac{4}{5}$	$\frac{2}{5}$	$\frac{1}{5}$	$\frac{4}{5}$	$\frac{3}{5}$	$\frac{1}{5}$	$\frac{3}{5}$
$\text{Arg}_2(d_1 \rightarrow d_2; 1)$	$\frac{1}{10}$	$\frac{3}{10}$	$\frac{1}{10}$	$-\frac{3}{10}$	$\frac{3}{10}$	$-\frac{1}{10}$	$-\frac{3}{10}$	$-\frac{1}{10}$
$\text{Arg}_3(d_1 \rightarrow d_2; 1)$	$\frac{4}{5}$	$\frac{2}{5}$	$\frac{4}{5}$	$\frac{3}{5}$	$\frac{2}{5}$	$\frac{1}{5}$	$\frac{3}{5}$	$\frac{1}{5}$
Total Arg($d_1 \rightarrow d_2; 1$)	$\frac{3}{10}$	$\frac{1}{2}$	$\frac{3}{10}$	$\frac{1}{2}$	$\frac{1}{2}$	$-\frac{3}{10}$	$\frac{1}{2}$	$-\frac{3}{10}$
$-3c' \times 4 \pmod{5}$	3	1	3	4	1	2	4	2
$\text{Arg}_1(d_1 \rightarrow d_2; 2) : \frac{1}{2} - \frac{12c'}{5}$	$\frac{1}{10}$	$-\frac{3}{10}$	$\frac{1}{10}$	$\frac{3}{10}$	$-\frac{3}{10}$	$-\frac{1}{10}$	$\frac{3}{10}$	$-\frac{1}{10}$
$\text{Arg}_2(d_1 \rightarrow d_2; 2)$	$\frac{1}{10}$	$\frac{3}{10}$	$\frac{1}{10}$	$-\frac{3}{10}$	$\frac{3}{10}$	$-\frac{1}{10}$	$-\frac{3}{10}$	$-\frac{1}{10}$
$\text{Arg}_3(d_1 \rightarrow d_2; 2)$	$\frac{4}{5}$	$\frac{2}{5}$	$\frac{4}{5}$	$\frac{3}{5}$	$\frac{2}{5}$	$\frac{1}{5}$	$\frac{3}{5}$	$\frac{1}{5}$
Total Arg($d_1 \rightarrow d_2; 2$)	0	$\frac{2}{5}$	0	$-\frac{2}{5}$	$\frac{2}{5}$	0	$-\frac{2}{5}$	0

TABLE 3.2. Table for Arg($d_1 \rightarrow d_2; \ell$); $2 \nmid c$, $3 \nmid c$, $5 \nmid c$.

3.2. **Case $2 \nmid c'$, $3|c'$, and $5 \nmid c'$.** These are the cases when $c' \equiv 3, 9, 21, 27 \pmod{30}$. For $\text{Arg}_1(d_1 \rightarrow d_4; \ell)$ we use (3.10). For $\text{Arg}_2(d_1 \rightarrow d_4; \ell)$, we need the congruence (3.9). By (2.3), we have

$$12cs(d_4, c) - 12cs(d_1, c) \equiv d_4 + \overline{d_{4\{3c\}}} - d_1 - \overline{d_{1\{3c\}}} \equiv 3\beta c'(1 - \overline{d_{4\{3c\}}} \cdot \overline{d_{1\{3c\}}}) \pmod{3c}. \quad (3.16)$$

By (2.4) we also have

$$12cs(d_4, c) - 12cs(d_1, c) \equiv 0 \pmod{8}. \quad (3.17)$$

Dividing the numerator and denominator of P_2 by $24c'$, we observe that

$$-\frac{5}{2}(s(d_4, c) - s(d_1, c)) \equiv -\overline{8_{\{5\}}}\beta(1 - \overline{d_{4\{3c\}}} \cdot \overline{d_{1\{3c\}}}) \equiv \beta \pmod{5} \quad (3.18)$$

because $\overline{d_{j\{3c\}}} \equiv \overline{d_{j\{5\}}} \equiv j \pmod{5}$ for $j = 1, 4$. Now we get $\text{Arg}_2(d_1 \rightarrow d_4; \ell) = \frac{\beta}{5}$. Since $\text{Arg}_3(d_1 \rightarrow d_4; \ell) = \frac{2\beta}{5}$, we have Table 3.3.

$c' \pmod{30}$	3	9	21	27
β	2	4	1	3
$3\beta c' \pmod{10}$	8	8	3	3
$-9\beta c'^2 \pmod{10}$	8	4	1	7
$\text{Arg}_1(d_1 \rightarrow d_4; 1)$	$-\frac{2}{10} + \frac{1}{2}$	$\frac{4}{10} - \frac{1}{2}$	$\frac{1}{10}$	$-\frac{3}{10}$
$(\text{Arg}_2 + \text{Arg}_3)(d_1 \rightarrow d_4; 1) : \frac{3\beta}{5}$	$\frac{1}{5}$	$\frac{2}{5}$	$\frac{3}{5}$	$\frac{4}{5}$
Total Arg($d_1 \rightarrow d_4; 1$)	$\frac{1}{2}$	$\frac{3}{10}$	$-\frac{3}{10}$	$\frac{1}{2}$
$3\beta c' \pmod{10}$	8	8	3	3
$-18\beta c'^2 \pmod{5}$	1	3	2	4
$\text{Arg}_1(d_1 \rightarrow d_4; 2)$	$-\frac{3}{10}$	$\frac{1}{10}$	$-\frac{1}{10}$	$\frac{3}{10}$
$(\text{Arg}_2 + \text{Arg}_3)(d_1 \rightarrow d_4; 2) : \frac{3\beta}{5}$	$\frac{1}{5}$	$\frac{2}{5}$	$\frac{3}{5}$	$\frac{4}{5}$
Total Arg($d_1 \rightarrow d_4; 2$)	$-\frac{1}{10}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{10}$

TABLE 3.3. Table for Arg($d_1 \rightarrow d_4; \ell$); $2 \nmid c$, $3|c$, $5 \nmid c$.

Next we investigate $\text{Arg}(d_1 \rightarrow d_2; \ell)$. For $\text{Arg}_1(d_1 \rightarrow d_2; \ell)$, we use (3.13). For $\text{Arg}_2(d_1 \rightarrow d_2; \ell)$, by (2.3) we have

$$12cs(d_2, c) - 12cs(d_1, c) \equiv d_2 + \overline{d_{2\{3c\}}} - d_1 - \overline{d_{1\{3c\}}} \equiv \beta c'(1 - \overline{d_{2\{3c\}}} \cdot \overline{d_{1\{3c\}}}) \pmod{3c}. \quad (3.19)$$

As $15|3c$, after dividing by c' we have

$$60s(d_2, c) - 60s(d_1, c) \equiv \beta(1 - \overline{d_{2\{15\}}} \cdot \overline{d_{1\{15\}}}) \equiv \beta(1 - a_3 a_1) \pmod{15}. \quad (3.20)$$

Since $a_3 \equiv a_1 \pmod{3}$, we have $a_3 a_1 \equiv 1 \pmod{3}$. We also have $a_3 a_1 \equiv 3 \pmod{5}$ by (3.4), then $a_3 a_1 \equiv 13 \pmod{15}$ and

$$-(60s(d_2, c) - 60s(d_1, c)) \equiv -3\beta \pmod{15}. \quad (3.21)$$

By (2.4) we have

$$12cs(d_4, c) - 12cs(d_1, c) \equiv 4 \pmod{8}. \quad (3.22)$$

The congruences (3.21) and (3.22) determine a unique value modulo 120.

$c' \pmod{30}$	3	9	21	27
β	2	4	1	3
$-3c' \pmod{5}$	1	3	2	4
$\text{Arg}_1(d_1 \rightarrow d_2; 1)$	$\frac{1}{5}$	$\frac{3}{5}$	$\frac{2}{5}$	$\frac{4}{5}$
$\text{Arg}_2(d_1 \rightarrow d_2; 1)$	$-\frac{3}{10}$	$-\frac{1}{10}$	$\frac{1}{10}$	$\frac{3}{10}$
$\text{Arg}_3(d_1 \rightarrow d_2; 1)$	$\frac{3}{5}$	$\frac{1}{5}$	$\frac{4}{5}$	$\frac{2}{5}$
Total $\text{Arg}(d_1 \rightarrow d_2; 1)$	$\frac{1}{2}$	$-\frac{3}{10}$	$\frac{3}{10}$	$\frac{1}{2}$
$-12c' \pmod{5}$	4	2	3	1
$\text{Arg}_1(d_1 \rightarrow d_2; 2)$	$\frac{3}{10}$	$-\frac{1}{10}$	$\frac{1}{10}$	$-\frac{3}{10}$
$\text{Arg}_2(d_1 \rightarrow d_2; 2)$	$-\frac{3}{10}$	$-\frac{1}{10}$	$\frac{1}{10}$	$\frac{3}{10}$
$\text{Arg}_3(d_1 \rightarrow d_2; 2)$	$\frac{3}{5}$	$\frac{1}{5}$	$\frac{4}{5}$	$\frac{2}{5}$
Total $\text{Arg}(d_1 \rightarrow d_2; 2)$	$-\frac{2}{5}$	0	0	$\frac{2}{5}$

TABLE 3.4. Table for $\text{Arg}(d_1 \rightarrow d_2; \ell)$; $2 \nmid c$, $3|c$, $5 \nmid c$.

Combining Table 3.3 and Table 3.4 we finish the proof in the case $2 \nmid c'$, $3|c'$ and $5 \nmid c'$.

3.3. Case $2|c'$, $3 \nmid c'$, and $5 \nmid c'$. These are the cases $c' \equiv 2, 4, 8, 14, 16, 22, 26, 28 \pmod{30}$. For $\text{Arg}_1(d_1 \rightarrow d_4; \ell)$ we still use (3.10). By (2.3), $\theta = 1$ and we still have

$$-(12cs(d_4, c) - 12cs(d_1, c)) \equiv -(d_4 + a_4 - d_1 - a_1) \equiv -6\beta c' \equiv -\beta c' \pmod{c}, \quad (3.23)$$

and $12cs(d, c) \equiv 0 \pmod{6}$. Define the integer $\lambda \geq 1$ by $2^\lambda || c$. To determine the value modulo $24c$, we need to determine it modulo $8 \times 2^\lambda$. By (2.5) we have

$$\begin{aligned} 12cs(d_4, c) - 12cs(d_1, c) &\equiv d_4 - d_1 + (c^2 + 3c + 1)(\overline{d_{4\{8 \times 2^\lambda\}}} - \overline{d_{1\{8 \times 2^\lambda\}}}) \\ &\quad + 2c \left(\overline{d_{4\{8 \times 2^\lambda\}}}\left(\frac{c}{d_4}\right) - \overline{d_{1\{8 \times 2^\lambda\}}}\left(\frac{c}{d_1}\right) \right) \pmod{8 \times 2^\lambda} \\ &\equiv 3\beta c' \left(1 - (c^2 + 3c + 1)\overline{d_{4\{8 \times 2^\lambda\}}} \cdot \overline{d_{1\{8 \times 2^\lambda\}}} \right) \\ &\quad + 2c \left(\overline{d_{4\{8 \times 2^\lambda\}}}\left(\frac{c}{d_4}\right) - \overline{d_{1\{8 \times 2^\lambda\}}}\left(\frac{c}{d_1}\right) \right) \pmod{8 \times 2^\lambda}. \end{aligned} \quad (3.24)$$

We claim that

$$12cs(d_4, c) - 12cs(d_1, c) \equiv 0 \pmod{8 \times 2^\lambda}. \quad (3.25)$$

To see this, since $2^\lambda \parallel c'$ and $c' \mid (12cs(d_4, c) - 12cs(d_1, c))$ by (3.23), we divide (3.24) by c' and obtain

$$\begin{aligned} 60(s(d_4, c) - s(d_1, c)) &\equiv 3\beta(1 - (c^2 + 3c + 1)d_4d_1) + 2\left(d_4\left(\frac{c}{d_4}\right) - d_1\left(\frac{c}{d_1}\right)\right) \\ &\equiv 3\beta c'(3\beta d_1 - 1)(c' - 1) + 2\left(d_4\left(\frac{c}{d_4}\right) - d_1\left(\frac{c}{d_1}\right)\right) \pmod{8}. \end{aligned}$$

Define $\text{val} := 3\beta c'(3\beta d_1 - 1)(c' - 1) \pmod{8}$ as the first part of the congruence above. Note that both d_1 and $c' - 1$ are odd. We have Table 3.5 for val .

$c' \pmod{5}$	1	2	3	4
β	1	3	2	4
$3\beta c'$	$3c'$	$6c'$	$9c'$	$12c'$
$3\beta d_1 - 1 \pmod{2}$	$3d_1 - 1$	$6d_1 - 1$	$9d_1 - 1$	$12d_1 - 1$
$2 \parallel c, d_1 \equiv 1 \pmod{4}$	4	4	0	0
$2 \parallel c, d_1 \equiv 3 \pmod{4}$	0	4	4	0
$4 \mid c;$	0	0	0	0

TABLE 3.5. Table of $\text{val} := 3\beta c'(3\beta d_1 - 1)(c' - 1) \pmod{8}$; $2 \mid c$, no requirement for $(c, 3)$, $5 \nmid c$.

For the second part we only need to determine $d_4\left(\frac{c}{d_4}\right) - d_1\left(\frac{c}{d_1}\right) \pmod{4}$. When $4 \mid c$, we get $\left(\frac{2^\lambda}{d_4}\right) = \left(\frac{2^\lambda}{d_1}\right) = 1$. By quadratic reciprocity,

$$\begin{aligned} d_4\left(\frac{c}{d_4}\right) - d_1\left(\frac{c}{d_1}\right) &\equiv d_1\left(\left(\frac{5}{d_4}\right)\left(\frac{c'/2^\lambda}{d_4}\right) - \left(\frac{5}{d_1}\right)\left(\frac{c'/2^\lambda}{d_1}\right)\right) \\ &\equiv d_1\left(\frac{d_1}{c'/2^\lambda}\right)\left(\left(-1\right)^{(d_4-1)(\frac{c'}{2^\lambda}-1)/4} - \left(-1\right)^{(d_1-1)(\frac{c'}{2^\lambda}-1)/4}\right) \equiv 0 \pmod{4} \end{aligned}$$

where the last equality follows since $\frac{d_4-1}{2}$ and $\frac{d_1-1}{2}$ have the same parity. This gives the last row in Table 3.5.

When $2 \parallel c$, recall that $d_4 = d_1 + 3\beta c'$, from which

$$d_4\left(\frac{c}{d_4}\right) - d_1\left(\frac{c}{d_1}\right) \equiv \left(\frac{d_1}{c'/2}\right)\left(\left(\frac{2}{d_4}\right)\left(-1\right)^{(d_4-1)(\frac{c'}{2}-1)/4}d_4 - \left(\frac{2}{d_1}\right)\left(-1\right)^{(d_1-1)(\frac{c'}{2}-1)/4}d_1\right) \pmod{4}. \quad (3.26)$$

When $c' \equiv 2 \pmod{8}$, $\frac{c'/2-1}{2}$ is even and (3.26) becomes $\left(\frac{2}{d_4}\right)d_4 - \left(\frac{2}{d_1}\right)d_1 \pmod{4}$; when $c' \equiv 6 \pmod{8}$, $\frac{c'/2-1}{2}$ is odd and (3.26) becomes $\left(\frac{2}{d_4}\right)\left(-1\right)^{\frac{d_4-1}{2}}d_4 - \left(\frac{2}{d_1}\right)\left(-1\right)^{\frac{d_1-1}{2}}d_1 \pmod{4}$. Since $c = 5c' \equiv c' \pmod{8}$, we can use $d_4 = d_1 + 3\beta c'$ to determine $d_4 \pmod{8}$ and get Table 3.6.

Combining Table 3.5 and Table 3.6, we prove (3.25). Recall (2.2) and (3.23), we divide both the denominator and numerator in P_2 by $24c'$ and get $\text{Arg}_2(d_1 \rightarrow d_4; \ell) = \frac{\beta}{5}$. Since $\text{Arg}_3(d_1 \rightarrow d_4; \ell) = \frac{2\beta}{5}$, we have Table 3.7.

Next we deal with $\text{Arg}(d_1 \rightarrow d_2; \ell)$. For $\text{Arg}_1(d_1 \rightarrow d_2; \ell)$, we still use (3.13). By (2.3),

$$-(12cs(d_2, c) - 12cs(d_1, c)) \equiv -(d_2 + a_3 - d_1 - a_1) \equiv -3\beta c' \equiv 2\beta c' \pmod{c}. \quad (3.27)$$

(3.26) \searrow	$c' \equiv 2 \pmod{8}$				$c' \equiv 6 \pmod{8}$			
$d_1 \pmod{8}$	1	3	5	7	1	3	5	7
$\beta = 1$	2	0	2	0	2	0	2	0
$\beta = 2$	2	2	2	2	2	2	2	2
$\beta = 3$	0	2	0	2	0	2	0	2
$\beta = 4$	0	0	0	0	0	0	0	0

 TABLE 3.6. Table for (3.26); $2|c$, no requirement for $(c, 3)$, $5 \nmid c$.

$c' \pmod{30}$	2	4	8	14	16	22	26	28
β	3	4	2	4	1	3	1	2
$3\beta c' \pmod{10}$	8	8	8	8	8	8	8	8
$-9\beta c'^2 \pmod{10}$	2	4	8	4	6	2	6	8
$\text{Arg}_1(d_1 \rightarrow d_4; 1)$	$-\frac{3}{10}$	$-\frac{1}{10}$	$\frac{3}{10}$	$-\frac{1}{10}$	$\frac{1}{10}$	$-\frac{3}{10}$	$\frac{1}{10}$	$\frac{3}{10}$
$(\text{Arg}_2 + \text{Arg}_3)(d_1 \rightarrow d_4; 1): \frac{3\beta}{5}$	$\frac{4}{5}$	$\frac{2}{5}$	$\frac{1}{5}$	$\frac{2}{5}$	$\frac{3}{5}$	$\frac{4}{5}$	$\frac{3}{5}$	$\frac{1}{5}$
Total $\text{Arg}(d_1 \rightarrow d_4; 1)$	$\frac{1}{2}$	$\frac{3}{10}$	$\frac{1}{2}$	$\frac{3}{10}$	$-\frac{3}{10}$	$\frac{1}{2}$	$-\frac{3}{10}$	$\frac{1}{2}$
$-18\beta c'^2 \equiv 2c' \pmod{5}$	4	3	1	3	2	4	2	1
$\text{Arg}_1(d_1 \rightarrow d_4; 2)$	$\frac{3}{10}$	$\frac{1}{10}$	$-\frac{3}{10}$	$\frac{1}{10}$	$-\frac{1}{10}$	$\frac{3}{10}$	$-\frac{1}{10}$	$-\frac{3}{10}$
$(\text{Arg}_2 + \text{Arg}_3)(d_1 \rightarrow d_4; 2): \frac{3\beta}{5}$	$\frac{4}{5}$	$\frac{2}{5}$	$\frac{1}{5}$	$\frac{2}{5}$	$\frac{3}{5}$	$\frac{4}{5}$	$\frac{3}{5}$	$\frac{1}{5}$
Total $\text{Arg}(d_1 \rightarrow d_4; 2)$	$\frac{1}{10}$	$\frac{1}{2}$	$-\frac{1}{10}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{10}$	$\frac{1}{2}$	$-\frac{1}{10}$

 TABLE 3.7. Table for $\text{Arg}(d_1 \rightarrow d_4; 2)$; $2|c$, $3 \nmid c$, $5 \nmid c$.

This congruence shows that $12cs(d_2, c) - 12cs(d_1, c)$ is divisible by c' . Denote λ by $2^\lambda || c$. We claim that

$$-(12cs(d_2, c) - 12cs(d_1, c)) \equiv 4 \times 2^\lambda \pmod{8 \times 2^\lambda}. \quad (3.28)$$

To prove (3.28), we apply (2.5) to get

$$\begin{aligned} 12cs(d_2, c) - 12cs(d_1, c) &\equiv \beta c' (1 - (c^2 + 3c + 1) \overline{d_{4\{8 \times 2^\lambda\}}} \cdot \overline{d_{1\{8 \times 2^\lambda\}}}) \\ &\quad + 2c \left(\overline{d_{2\{8 \times 2^\lambda\}}}\left(\frac{c}{d_2}\right) - \overline{d_{1\{8 \times 2^\lambda\}}}\left(\frac{c}{d_1}\right) \right) \pmod{8 \times 2^\lambda}. \end{aligned} \quad (3.29)$$

Then as in (3.24), we have

$$60s(d_2, c) - 60s(d_1, c) \equiv \beta c' (\beta d_1 - 1)(c' - 1) + 2 \left(d_2 \left(\frac{c}{d_2} \right) - d_1 \left(\frac{c}{d_1} \right) \right) \pmod{8}. \quad (3.30)$$

See Table 3.8 for the first term $\text{val} := \beta c' (\beta d_1 - 1)(c' - 1) \pmod{8}$ and note that d_1 and $c' - 1$ are both odd.

For the second term $2 \left(d_2 \left(\frac{c}{d_2} \right) - d_1 \left(\frac{c}{d_1} \right) \right) \pmod{8}$, we argue as above using the quadratic reciprocity (2.6) and omit the details. Combining (2.2), (3.27) and (3.28), we have

$$-(12cs(d_2, c) - 12cs(d_1, c)) \equiv 12 \times 2^\lambda \pmod{24 \times 2^\lambda}.$$

After dividing c' , $-60s(d_2, c) + 60s(d_1, c) \pmod{120}$ is uniquely determined by $2\beta \pmod{5}$ and $12 \pmod{24}$. Hence

$$\text{Arg}_2(d_1 \rightarrow d_2; \ell) = \frac{1, 7, 3, 9}{10}, \quad \text{for } \beta = 1, 2, 3, 4, \text{ respectively}$$

$c' \pmod{5}$	1	2	3	4
β	1	3	2	4
$\beta c'$	c'	$3c'$	$2c'$	$4c'$
$\beta d_1 - 1$	$d_1 - 1$	$3d_1 - 1$	$2d_1 - 1$	$4d_1 - 1$
$2 \parallel c, d_1 \equiv 1 \pmod{4}$	0	4	4	0
$2 \parallel c, d_1 \equiv 3 \pmod{4}$	4	0	4	0
$4 \mid c$	0	0	0	0

TABLE 3.8. Table for $\text{val.} := \beta c' (\beta d_1 - 1) (c' - 1) \pmod{8}$; $2 \mid c$, no requirement for $(c, 3)$, $5 \nmid c$.

and we get Table 3.9.

$c' \pmod{30}$	2	4	8	14	16	22	26	28
β	3	4	2	4	1	3	1	2
$2\beta c' \pmod{10}$	2	2	2	2	2	2	2	2
$-3\beta c'^2 \equiv 2c' \pmod{5}$	4	3	1	3	2	4	2	1
$\text{Arg}_1(d_1 \rightarrow d_2; 1)$	$\frac{4}{5}$	$\frac{3}{5}$	$\frac{1}{5}$	$\frac{3}{5}$	$\frac{2}{5}$	$\frac{4}{5}$	$\frac{2}{5}$	$\frac{1}{5}$
$\text{Arg}_2(d_1 \rightarrow d_2; 1)$	$\frac{3}{10}$	$-\frac{1}{10}$	$-\frac{3}{10}$	$-\frac{1}{10}$	$\frac{1}{10}$	$\frac{3}{10}$	$\frac{1}{10}$	$-\frac{3}{10}$
$\text{Arg}_3(d_1 \rightarrow d_2; 1)$	$\frac{2}{5}$	$\frac{1}{5}$	$\frac{3}{5}$	$\frac{1}{5}$	$\frac{4}{5}$	$\frac{2}{5}$	$\frac{4}{5}$	$\frac{3}{5}$
Total $\text{Arg}(d_1 \rightarrow d_2; 1)$	$\frac{1}{2}$	$-\frac{3}{10}$	$\frac{1}{2}$	$-\frac{3}{10}$	$\frac{3}{10}$	$\frac{1}{2}$	$\frac{3}{10}$	$\frac{1}{2}$
$-12\beta c'^2 \equiv 3c' \pmod{5}$	1	2	4	2	3	1	3	4
$\text{Arg}_1(d_1 \rightarrow d_2; 2)$	$-\frac{3}{10}$	$-\frac{1}{10}$	$\frac{3}{10}$	$-\frac{1}{10}$	$\frac{1}{10}$	$-\frac{3}{10}$	$\frac{1}{10}$	$\frac{3}{10}$
$\text{Arg}_2(d_1 \rightarrow d_2; 2)$	$\frac{3}{10}$	$-\frac{1}{10}$	$-\frac{3}{10}$	$-\frac{1}{10}$	$\frac{1}{10}$	$\frac{3}{10}$	$\frac{1}{10}$	$-\frac{3}{10}$
$\text{Arg}_3(d_1 \rightarrow d_2; 2)$	$\frac{2}{5}$	$\frac{1}{5}$	$\frac{3}{5}$	$\frac{1}{5}$	$\frac{4}{5}$	$\frac{2}{5}$	$\frac{4}{5}$	$\frac{3}{5}$
Total $\text{Arg}(d_1 \rightarrow d_2; 2)$	$\frac{2}{5}$	0	$-\frac{2}{5}$	0	0	$\frac{2}{5}$	0	$-\frac{2}{5}$

TABLE 3.9. Table for $\text{Arg}(d_1 \rightarrow d_2; \ell)$; $2 \mid c$, $3 \nmid c$, $5 \nmid c$.

Combining Table 3.7 and Table 3.9, we confirm that Condition 3.2 is satisfied in these cases.

3.4. Case $2 \mid c'$, $3 \mid c'$, and $5 \nmid c'$. These are the cases $c' \equiv 6, 12, 18, 24 \pmod{30}$. For $\text{Arg}_1(d_1 \rightarrow d_4; \ell)$ we use (3.10). For $\text{Arg}_2(d_1 \rightarrow d_4; \ell)$, by (2.3) we have

$$-(12cs(d_4, c) - 12cs(d_1, c)) \equiv -3\beta c' (1 - \overline{d_{4\{3c\}}} \cdot \overline{d_{1\{3c\}}}) \pmod{3c}. \quad (3.31)$$

The proof of (3.25) in the former subsection still works for $3 \mid c$. Then $-(12cs(d_4, c) - 12cs(d_1, c))$ is a multiple of $24c'$. After dividing both the denominator and numerator in P_2 and recalling $\overline{d_{j\{3c\}}} \equiv a_j \equiv j \pmod{5}$ for $j = 1, 4$, we get $\text{Arg}_2(d_1 \rightarrow d_4; \ell) = e(\frac{\beta}{5})$. This gives Table 3.10.

Then we check $\text{Arg}(d_1 \rightarrow d_2; \ell)$. For $\text{Arg}_1(d_1 \rightarrow d_2; \ell)$, we use (3.13). For $\text{Arg}_2(d_1 \rightarrow d_2; \ell)$, by (2.3) we have

$$-(12cs(d_2, c) - 12cs(d_1, c)) \equiv -\beta c' (1 - \overline{d_{2\{3c\}}} \overline{d_{1\{3c\}}}) \pmod{3c}. \quad (3.32)$$

$c' \pmod{30}$	6	12	18	24
β	1	3	2	4
$3\beta c' \pmod{10}$	8	8	8	8
$-9\beta c'^2 \pmod{10}$	6	2	8	4
$\text{Arg}_1(d_1 \rightarrow d_4; 1)$	$\frac{1}{10}$	$-\frac{3}{10}$	$\frac{3}{10}$	$-\frac{1}{10}$
$(\text{Arg}_2 + \text{Arg}_3)(d_1 \rightarrow d_4; 1) : \frac{3\beta}{5}$	$\frac{3}{5}$	$\frac{4}{5}$	$\frac{1}{5}$	$\frac{2}{5}$
Total Arg($d_1 \rightarrow d_4; 1$)	$-\frac{3}{10}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{3}{10}$
$-18\beta c'^2 \equiv 2c' \pmod{5}$	2	4	1	3
$\text{Arg}_1(d_1 \rightarrow d_4; 2)$	$-\frac{1}{10}$	$\frac{3}{10}$	$-\frac{3}{10}$	$\frac{1}{10}$
$(\text{Arg}_2 + \text{Arg}_3)(d_1 \rightarrow d_4; 2) : \frac{3\beta}{5}$	$\frac{3}{5}$	$\frac{4}{5}$	$\frac{1}{5}$	$\frac{2}{5}$
Total Arg($d_1 \rightarrow d_4; 2$)	$\frac{1}{2}$	$\frac{1}{10}$	$-\frac{1}{10}$	$\frac{1}{2}$

 TABLE 3.10. Table for $\text{Arg}(d_1 \rightarrow d_4; \ell); 2|c, 3|c, 5 \nmid c$.

Since $3|c$, $\overline{d_{2\{3c\}}} \equiv a_3 \pmod{15}$ and $\overline{d_{1\{3c\}}} \equiv a_1 \pmod{15}$. After dividing by c' we have

$$-(60s(d_2, c) - 60s(d_1, c)) \equiv -\beta(1 - a_3 a_1) \pmod{15}.$$

We have $a_3 = a_1 + 2\beta c'$ and $a_3 a_1 \equiv 13 \pmod{15}$, so

$$-(60s(d_2, c) - 60s(d_1, c)) \equiv -3\beta \pmod{15}. \quad (3.33)$$

Denote λ by $2^\lambda || c$, then (3.28) still holds since

$$-(60s(d_2, c) - 60s(d_1, c)) \equiv 4 \pmod{8}. \quad (3.34)$$

By (3.33) and (3.34), we obtain

$$\text{Arg}_2(d_1 \rightarrow d_2; \ell) = \frac{1, 7, 3, 9}{10} \quad \text{for } \beta = 1, 2, 3, 4, \text{ respectively.}$$

This gives Table 3.11.

$c' \pmod{30}$	6	12	18	24
β	1	3	2	4
$-3\beta c'^2 \equiv 2c' \pmod{5}$	2	4	1	3
$\text{Arg}_1(d_1 \rightarrow d_2; 1)$	$\frac{2}{5}$	$\frac{4}{5}$	$\frac{1}{5}$	$\frac{3}{5}$
$\text{Arg}_2(d_1 \rightarrow d_2; 1)$	$\frac{1}{10}$	$\frac{3}{10}$	$-\frac{3}{10}$	$-\frac{1}{10}$
$\text{Arg}_3(d_1 \rightarrow d_2; 1)$	$\frac{4}{5}$	$\frac{2}{5}$	$\frac{3}{5}$	$\frac{1}{5}$
Total Arg($d_1 \rightarrow d_2; 1$)	$\frac{3}{10}$	$\frac{1}{2}$	$\frac{1}{2}$	$-\frac{3}{10}$
$-12\beta c'^2 \equiv 3c' \pmod{5}$	3	1	4	2
$\text{Arg}_1(d_1 \rightarrow d_2; 2)$	$\frac{1}{10}$	$-\frac{3}{10}$	$\frac{3}{10}$	$-\frac{1}{10}$
$\text{Arg}_2(d_1 \rightarrow d_2; 2)$	$\frac{1}{10}$	$\frac{3}{10}$	$-\frac{3}{10}$	$-\frac{1}{10}$
$\text{Arg}_3(d_1 \rightarrow d_2; 2)$	$\frac{4}{5}$	$\frac{2}{5}$	$\frac{3}{5}$	$\frac{1}{5}$
Total Arg($d_1 \rightarrow d_2; 2$)	0	$\frac{2}{5}$	$-\frac{2}{5}$	0

 TABLE 3.11. Table for $\text{Arg}(d_1 \rightarrow d_2; \ell); 2|c, 3|c, 5 \nmid c$.

Comparing Table 3.10 and Table 3.11, we have proved that Condition 3.2 is satisfied in these cases.


We have finished the proof of Proposition 3.1 by proving that the four points $P(d)$ satisfy Condition 3.2 when $5 \parallel c$. The next subsection is to prove Proposition 3.1 in the case $25 \mid c$, which is different from the former ones.

3.5. Case $5 \mid c'$. We still denote $c' = c/5$ and $V(r, c) := \{d \pmod{c}^* : d \equiv r \pmod{c'}\}$ for $r \pmod{c'}^*$. Now $|V(r, c)| = 5$ and since $(d + c', c) = 1$ when $(d, c) = 1$, we can write $V(r, c) = \{d, d + c', d + 2c', d + 3c', d + 4c'\}$ for $1 \leq d < c'$ and $d \equiv r \pmod{c'}$.

We claim that Proposition 3.1 is still true:

$$\sum_{d \in V(r, c)} \frac{e\left(-\frac{3c'ad^2}{10}\right)}{\sin\left(\frac{\pi a \ell}{p}\right)} e\left(-\frac{12cs(d, c)}{24c}\right) e\left(\frac{4d}{c}\right) = 0, \quad (3.35)$$

but this time we have five summands. We prove (3.35) by showing that there are only two possible configurations for the summands:



i.e. all at the outer circle (radius $1/\sin(\frac{\pi}{5})$) or all at the inner circle (radius $1/\sin(\frac{2\pi}{5})$) and equally distributed. As in (3.5), we still denote the factors in (3.35) by P_1 , P_2 and P_3 and investigate the argument differences contributed from each term.

For any $d \in V(r, c)$, we take $a \pmod{c}$ such that $ad \equiv 1 \pmod{c}$. We denote $d_* = d + c'$ and denote a_* by $a_*d_* \equiv 1 \pmod{c}$. Then we can pick $a_* = a - c'$ when $d \equiv 1, 4 \pmod{5}$ and pick $a_* = a + c'$ when $d \equiv 2, 3 \pmod{5}$.

Note that $P_1(d) = (-1)^{ca\ell}/\sin(\frac{\pi a \ell}{5})$ has period c' , hence $\text{Arg}_1(d \rightarrow d_*; \ell) = 0$ always. In the following two cases, we prove

$$\text{Arg}(d \rightarrow d_*; \ell) = -\frac{1}{5} \quad \text{for every } d \in V(r, c) \quad (3.36)$$

when $\ell = 1$. The other case $\ell = 2$ only affects P_1 (radii for those five points) and results in the same conclusion. This proves (3.35) when $25 \mid c$.

3.5.1. c is odd. When $d \equiv 1, 4 \pmod{5}$ and $3 \nmid c$, (2.2), (2.3) and (2.4) imply that

$$12cs(d_*, c) - 12cs(d, c) \equiv 0 \pmod{24c}, \quad (3.37)$$

hence $\text{Arg}_2(d \rightarrow d_*; \ell) = 0$ always. As $\text{Arg}_3(d \rightarrow d_*; \ell) = \frac{4}{5}$ for any $d \in V(r, c)$, we have proved (3.35) in this case.

When $3 \mid c$ and $d \equiv 1, 4 \pmod{5}$, (2.3) implies

$$-(12cs(d_*, c) - 12cs(d, c)) \equiv -c'(1 - \overline{d_{*\{3c\}}} \cdot \overline{d_{\{3c\}}}) \pmod{3c}. \quad (3.38)$$

Since $15|c$, after dividing by c' we have

$$-(60s(d_*, c) - 60s(d, c)) \equiv a^2 - 1 \pmod{15}. \quad (3.39)$$

Note that $a \equiv 1, 4 \pmod{5}$ and $a^2 \equiv 1 \pmod{15}$, hence we have $-(12cs(d_*, c) - 12cs(d, c)) \equiv 0 \pmod{24c}$ and conclude (3.36).

When $d \equiv 2, 3 \pmod{5}$ and $3 \nmid c$, recall $d_* = d + c'$ and $a_1 = a + c'$ with $a + d \equiv 0 \pmod{5}$. By (2.2), (2.3) and (2.4), we have

$$-(12cs(d_*, c) - 12cs(d, c)) \equiv -2c' \pmod{c} \quad \text{and} \quad \equiv 0 \pmod{24}.$$

Then $\text{Arg}_2(d \rightarrow d_*; \ell) = \frac{2}{5}$. Since $\text{Arg}_3(d \rightarrow d_*; \ell) = \frac{4}{5}$, we have proved (3.36) in this case.

When $d \equiv 2, 3 \pmod{5}$ and $3|c$, we still get (3.39), while this time $a \equiv 3, 2 \pmod{5}$, $a^2 - 1 \equiv 3 \pmod{15}$, and hence $a^2 - 1 \equiv 48 \pmod{120}$. We have $\text{Arg}_2(d \rightarrow d_*; \ell) = \frac{2}{5}$. Since $\text{Arg}_3(d \rightarrow d_*; \ell) = \frac{4}{5}$, we have proved (3.36) in this case.

3.5.2. *c is even.* In this case, denote λ by $2^\lambda || c$. Then by (2.5) we have

$$\begin{aligned} 12cs(d_*, c) - 12cs(d, c) &\equiv c' (1 - (c^2 + 3c + 1) \overline{d_{1\{8 \times 2^\lambda\}}} \cdot \overline{d_{\{8 \times 2^\lambda\}}}) \\ &\quad + 2c \left(\left(\frac{c}{d_*} \right) \overline{d_{1\{8 \times 2^\lambda\}}} - \left(\frac{c}{d} \right) \overline{d_{\{8 \times 2^\lambda\}}} \right) \pmod{8 \times 2^\lambda}. \end{aligned}$$

Since $c'|(12cs(d_*, c) - 12cs(d, c))$ by (3.37) and (3.38), dividing the above congruence by c' we have

$$-60(s(d_*, c) - s(d, c)) \equiv -c'(d-1)(c'-1) - 2 \left(\left(\frac{c}{d_*} \right) d_* - \left(\frac{c}{d} \right) d \right) \pmod{8}. \quad (3.40)$$

For the first term,

$$-c'(d-1)(c'-1) \equiv \begin{cases} 0 \pmod{8} & \text{if } 2||c, d \equiv 1 \pmod{4}; \\ 4 \pmod{8} & \text{if } 2||c, d \equiv 3 \pmod{4}; \\ 0 \pmod{8} & \text{if } 4|c. \end{cases} \quad (3.41)$$

When λ is even, $\left(\frac{2^\lambda}{d_*}\right) = \left(\frac{2^\lambda}{d}\right) = 1$; when $\lambda \geq 3$ is odd, $\left(\frac{2}{d_*}\right) = \left(\frac{2}{d}\right)$. In either case $\frac{d_*-1}{2}$ and $\frac{d-1}{2}$ have the same parity. Hence when $4|c$, we have

$$\left(\frac{c}{d_*} \right) d_* - \left(\frac{c}{d} \right) d \equiv 0 \pmod{4}.$$

When $2||c$, we have Table 3.12 for $\text{val} := \left(\frac{c}{d_*} \right) d_* - \left(\frac{c}{d} \right) d \pmod{4}$ using quadratic reciprocity.

$d \pmod{8}$	1	3	5	7
$d_* \pmod{8}$ when $c' \equiv 2 \pmod{8}$	3	5	7	1
val.	0	2	0	2
$d_* \pmod{8}$ when $c' \equiv 6 \pmod{8}$	7	1	3	5
val.	0	2	0	2

TABLE 3.12. Table for $\text{val} := \left(\frac{c}{d_*} \right) d_* - \left(\frac{c}{d} \right) d \pmod{4}$; $2|c$, no requirement for $(3, c)$, $5|c$.

Combining (3.41) and Table 3.12, for $2^\lambda || c$ we get

$$12cs(d_*, c) - 12cs(d, c) \equiv 0 \pmod{8 \times 2^\lambda}. \quad (3.42)$$

The argument for the cases $d \equiv 1, 4 \pmod{5}$ or $d \equiv 2, 3 \pmod{5}$, or the cases $3 \nmid c$ or $3|c$, still works as the former case.

Proof of Proposition 3.1. This is proved by Condition 3.2 and (3.36). \square

This finishes the proof of (5-4) in Theorem 1.3.

4. PROOF OF (7-5,1) OF THEOREM 1.3

Recall (2.7) in the case $p = 7$:

$$e\left(\frac{1}{8}\right)S_{\infty\infty}^{(\ell)}(0, 7n + 5, c, \mu_7) = \sum_{\substack{d \pmod{c}^* \\ ad \equiv 1 \pmod{c}}} \frac{(-1)^{\ell c} e\left(-\frac{3c' a \ell}{14}\right)}{\sin\left(\frac{\pi a \ell}{7}\right)} e^{-\pi i s(d, c)} e\left(\frac{(7n + 5)d}{c}\right). \quad (4.1)$$

We only need to consider $\ell = 1, 2, 3$ because $A\left(\frac{\ell}{p}; n\right) = A\left(1 - \frac{\ell}{p}; n\right)$.

As in the previous section, we define $c' := c/7$. For an integer r with $(r, c') = 1$, we define

$$V(r, c) = \{d \pmod{c}^* : d \equiv r \pmod{c'}\}.$$

For example, $V(1, 42) = \{1, 13, 19, 25, 31, 37\}$ and $V(4, 35) = \{4, 9, 19, 24, 29, 34\}$. Then $|V(r, c)| = 6$ if $7 \nmid c'$, $|V(r, c)| = 7$ if $49|c$, and $(\mathbb{Z}/c\mathbb{Z})^*$ is the disjoint union

$$(\mathbb{Z}/c\mathbb{Z})^* = \bigcup_{r \pmod{c'}^*} V(r, c).$$

We claim the following proposition.

Proposition 4.1. *For $\ell = 1, 2, 3$, when $7|c$, $\frac{c}{7} \cdot \ell \not\equiv 1 \pmod{7}$ and $\frac{c}{7} \cdot \ell \not\equiv -1 \pmod{7}$, the sum on $d \in V(r, c)$ for all $r \pmod{c'}^*$ is zero:*

$$s_{r, c} := \sum_{d \in V(r, c)} \frac{e\left(-\frac{3c' a \ell^2}{14}\right)}{\sin\left(\frac{\pi a \ell}{7}\right)} e\left(-\frac{12cs(d, c)}{24c}\right) e\left(\frac{5d}{c}\right) = 0 \quad (4.2)$$

If Proposition 4.1 is true, then

$$S_{\infty\infty}^{(\ell)}(0, 7n + 5, c, \mu_7) = e\left(-\frac{1}{8}\right)(-1)^{\ell c} \sum_{r \pmod{c'}^*} s_{r, c} e\left(\frac{nr}{c'}\right) = 0$$

for all $n \in \mathbb{Z}$, $\ell = 1, 2, 3$ and we have proved (7-5,1) of Theorem 1.3.

As in (3.5), we label the terms in (4.2) as

$$P(d) := \frac{e\left(-\frac{3c' a \ell^2}{14}\right)}{\sin\left(\frac{\pi a \ell}{7}\right)} \cdot e\left(-\frac{12cs(d, c)}{24c}\right) \cdot e\left(\frac{5d}{c}\right) =: P_1(d) \cdot P_2(d) \cdot P_3(d). \quad (4.3)$$


We first deal with the case $7 \nmid c'$. We denote the argument differences as in (3.6), but in this case $u, v \in \{1, 2, \dots, 6\}$ and $\ell \in \{1, 2, 3\}$, where

$$d_u \equiv a_u \equiv u \pmod{7}, \quad a_{\overline{u\{7\}}} d_u \equiv 1 \pmod{c}, \quad d_{u+1} = d_u + \beta c' \quad \text{and} \quad a_{u+1} = a_u + \beta c'. \quad (4.4)$$


Note that $a_u d_u$ may not be $1 \pmod{c}$. Let $1 \leq \beta \leq 6$ such that $\beta c' \equiv 1 \pmod{7}$.

As in Condition 3.2, we have the following styles for the six summands followed by the explanation in Condition 4.2:

- $\ell = 1, 2, 3$, first style.



- $\ell = 1, 2, 3$, reversed style from the above.



Here we explain these styles. Each graph above includes three circles centered at the origin with radii $\csc(\frac{\pi}{7})$, $\csc(\frac{2\pi}{7})$ and $\csc(\frac{3\pi}{7})$, respectively. The six points in each graph above mark $P(d)$ for $d \in V(r, c)$ on these three circles. It is not hard to prove that whenever the six points satisfy the following condition on their argument differences, they sum to zero. This proves Proposition 4.1 by using the equation

$$\frac{\cos(\frac{3\pi}{7})}{\sin(\frac{\pi}{7})} - \frac{\cos(\frac{\pi}{7})}{\sin(\frac{2\pi}{7})} + \frac{\cos(\frac{2\pi}{7})}{\sin(\frac{3\pi}{7})} = 0, \quad \text{where } \frac{1}{\sin(\frac{\pi}{7})}, \frac{1}{\sin(\frac{2\pi}{7})}, \frac{1}{\sin(\frac{3\pi}{7})} \text{ are the radii.}$$

Condition 4.2. We have the following six styles for these six points when $7 \parallel c$, $\frac{c}{7} \cdot \ell \not\equiv 1 \pmod{7}$ and $\frac{c}{7} \cdot \ell \not\equiv -1 \pmod{7}$.

- $\ell = 1$: the arguments (as a proportion of 2π) going $d_1 \rightarrow d_2 \rightarrow d_3 \rightarrow d_4 \rightarrow d_5 \rightarrow d_6 \rightarrow d_1$ are $-\frac{5}{14}, -\frac{2}{7}, -\frac{1}{7}, -\frac{2}{7}, -\frac{5}{14}$, and $\frac{3}{7}$, respectively, or the reversed style.

	$d_1 \rightarrow$	$d_2 \rightarrow$	$d_3 \rightarrow$	$d_4 \rightarrow$	$d_5 \rightarrow$	$d_6 \rightarrow$	d_1
$c' \equiv 2, 4 \pmod{7}$	$-\frac{5}{14}$	$-\frac{2}{7}$	$-\frac{1}{7}$	$-\frac{2}{7}$	$-\frac{5}{14}$	$\frac{3}{7}$	
$c' \equiv 3, 5 \pmod{7}$	$\frac{5}{14}$	$\frac{2}{7}$	$\frac{1}{7}$	$\frac{2}{7}$	$\frac{5}{14}$	$-\frac{3}{7}$	

- $\ell = 2$, second graph style:

	$d_1 \rightarrow$	$d_2 \rightarrow$	$d_3 \rightarrow$	$d_4 \rightarrow$	$d_5 \rightarrow$	$d_6 \rightarrow$	d_1
$c' \equiv 5, 6 \pmod{7}$	$\frac{3}{14}$	$\frac{1}{14}$	$\frac{2}{7}$	$\frac{1}{14}$	$\frac{3}{14}$	$\frac{1}{7}$	
$c' \equiv 1, 2 \pmod{7}$	$-\frac{3}{14}$	$-\frac{1}{14}$	$-\frac{2}{7}$	$-\frac{1}{14}$	$-\frac{3}{14}$	$-\frac{1}{7}$	

- $\ell = 3$, third graph style:

	$d_1 \rightarrow$	$d_2 \rightarrow$	$d_3 \rightarrow$	$d_4 \rightarrow$	$d_5 \rightarrow$	$d_6 \rightarrow$	d_1
$c' \equiv 1, 4 \pmod{7}$	$-\frac{3}{7}$	$\frac{5}{14}$	$\frac{3}{7}$	$\frac{5}{14}$	$-\frac{3}{7}$	$-\frac{2}{7}$	
$c' \equiv 3, 6 \pmod{7}$	$\frac{3}{7}$	$-\frac{5}{14}$	$-\frac{3}{7}$	$-\frac{5}{14}$	$\frac{3}{7}$	$\frac{2}{7}$	

Proof of Proposition 4.1 when $7 \nmid c'$. This is proved by Condition 4.2. \square

Remark. Note that (7-5,1) of Theorem 1.3 is for the case $c'\ell \not\equiv \pm 1 \pmod{7}$, so Condition 4.2 does not include all the cases of $c' \pmod{7}$. We will highlight these exceptional cases among the tables in this section by a row " $c'\ell \equiv \pm 1 \pmod{7}$ ". The corresponding entry is:

$$\begin{cases} \text{blank,} & \text{if } c'\ell \not\equiv \pm 1 \pmod{7}; \\ \text{"+"}, & \text{if } c'\ell \equiv 1 \pmod{7}; \\ \text{"-"}, & \text{if } c'\ell \equiv -1 \pmod{7}. \end{cases}$$

We will explain these exceptional styles $c'\ell \equiv \pm 1 \pmod{7}$ in the next section for (7-5,2).

In the following subsections, we show $\text{Arg}(d_1 \rightarrow d_2; \ell)$, $\text{Arg}(d_2 \rightarrow d_3; \ell)$, and $\text{Arg}(d_3 \rightarrow d_4; \ell)$ in all the cases $c' \pmod{42}$. These argument differences are sufficient to check Condition 4.2 because

$$\text{Arg}(d_1 \rightarrow d_2; \ell) = \text{Arg}(d_5 \rightarrow d_6; \ell) \text{ and } \text{Arg}(d_2 \rightarrow d_3; \ell) = \text{Arg}(d_4 \rightarrow d_5; \ell),$$

where the proof is the same as the proof of Lemma 3.3. When $7 \nmid c'$, we prove that $\text{Arg}(d_1 \rightarrow d_2; \ell)$, $\text{Arg}(d_2 \rightarrow d_3; \ell)$, and $\text{Arg}(d_3 \rightarrow d_4; \ell)$ satisfy Condition 4.2 in §4.1. When $49|c$, we prove Proposition 4.1 in §4.2.

4.1. Case $7 \nmid c'$. As in the proof of (5-4) in the previous section, we should divide into cases for the divisibility of c' by 2, 3 to apply the suitable congruence in (2.2)-(2.5). However, the whole proof will be quite long.

Here we only prove the most complicated case $2|c'$ and $3 \nmid c'$ and omit the others. In this case $c' \equiv 2, 4, 8, 10, 16, 20, 22, 26, 32, 34, 38, 40 \pmod{42}$.

For Arg_1 , we have

$$\text{Arg}_1(d_1 \rightarrow d_2; \ell) = -\frac{9\beta c'^2 \ell^2}{14} \begin{cases} +0 & \text{sgn}(\sin(\frac{\pi a \ell}{7}) / \sin(\frac{\pi a_1 \ell}{7})) = 1, \\ \pm \frac{1}{2} & \text{sgn}(\sin(\frac{\pi a_4 \ell}{7}) / \sin(\frac{\pi a_1 \ell}{7})) = -1, \end{cases} \quad (4.5)$$

$$\text{Arg}_1(d_2 \rightarrow d_3; \ell) = -\frac{3\beta c'^2 \ell^2}{14} \begin{cases} +0 & \text{if } \text{sgn}(\sin(\frac{\pi a_5 \ell}{7}) / \sin(\frac{\pi a_4 \ell}{7})) = 1, \\ \pm \frac{1}{2} & \text{if } \text{sgn}(\sin(\frac{\pi a_5 \ell}{7}) / \sin(\frac{\pi a_4 \ell}{7})) = -1, \end{cases} \quad (4.6)$$

and

$$\text{Arg}_1(d_3 \rightarrow d_4; \ell) = \frac{9\beta c'^2 \ell^2}{14} \begin{cases} +0 & \text{if } \text{sgn}(\sin(\frac{\pi a_2 \ell}{7}) / \sin(\frac{\pi a_5 \ell}{7})) = 1, \\ \pm \frac{1}{2} & \text{if } \text{sgn}(\sin(\frac{\pi a_2 \ell}{7}) / \sin(\frac{\pi a_5 \ell}{7})) = -1. \end{cases} \quad (4.7)$$

We compute $\text{Arg}(d_1 \rightarrow d_2; \ell)$ starting from (4.5). For Arg_2 we need to combine (2.3) and (2.5). We have $12cs(d, c) \equiv 0 \pmod{6}$ and

$$-12cs(d_2, c) + 12cs(d_1, c) \equiv -d_2 - a_4 + d_1 + a_1 \equiv -4\beta c' \pmod{c}. \quad (4.8)$$

Then $-12cs(d_2, c) + 12cs(d_1, c)$ is a multiple of c' .

We claim that

$$-12cs(d_2, c) + 12cs(d_1, c) \equiv 0 \pmod{8 \times 2^\lambda}. \quad (4.9)$$

Denote $\lambda \geq 1$ by $2^\lambda || c$. We have

$$\begin{aligned} -12cs(d_2, c) + 12cs(d_1, c) &\equiv -d_2 - \overline{d_{2\{8 \times 2^\lambda\}}}(c^2 + 3c + 1 + 2c(\frac{c}{d_2})) \\ &\quad + d_1 + \overline{d_{1\{8 \times 2^\lambda\}}}(c^2 + 3c + 1 + 2c(\frac{c}{d_1})) \\ &\equiv -\beta c' + \beta c \overline{d_{2\{8 \times 2^\lambda\}}} \cdot \overline{d_{1\{8 \times 2^\lambda\}}}(c^2 + 3c + 1) \\ &\quad + 2c(\overline{d_{1\{8 \times 2^\lambda\}}}(c) - \overline{d_{2\{8 \times 2^\lambda\}}}(c)) \pmod{8 \times 2^\lambda}. \end{aligned}$$

After dividing c' , we get the value modulo 8 by $\overline{x_{\{8\}}} \equiv x \pmod{8}$:

$$\begin{aligned} -84s(d_2, c) + 84s(d_1, c) &\equiv -\beta + \beta d_2 d_1 (c^2 + 3c + 1) + 6(d_1(\frac{c}{d_1}) - d_2(\frac{c}{d_2})) \\ &\equiv \beta c'(1 + d_1 \beta)(c' + 1) - 2(d_1(\frac{c}{d_1}) - d_2(\frac{c}{d_2})) \pmod{8} \end{aligned}$$

For the first value $\text{val} := \beta c'(1 + d_1 \beta)(c' + 1) \pmod{8}$, we see that both $\beta(1 + d_1 \beta)$ and c' are even, hence the result is $0, 4 \pmod{8}$. Moreover, val. is the same for c' and $c' + 7$. Then we have Table 4.1.

$c' \pmod{7}$	1	2	3	4	5	6
β	1	4	5	2	3	6
$\beta c'$	c'	$4c'$	$5c'$	$2c'$	$3c'$	$6c'$
$\beta d_1 + 1$	$d_1 + 1$	$4d_1 + 1$	$5d_1 + 1$	$2d_1 + 1$	$3d_1 + 1$	$6d_1 + 1$
$2 c, d_1 \equiv 1 \pmod{4}$	4	0	4	4	0	4
$2 c, d_1 \equiv 3 \pmod{4}$	0	0	0	4	4	4
$4 c$	0	0	0	0	0	0

TABLE 4.1. Table for $\text{val} := \beta c'(\beta d_1 + 1)(c' + 1) \pmod{8}$; $2|c$, no requirement for $(c, 3)$, $7 \nmid c$.

For the other part we determine whether

$$d_1(\frac{c}{d_1}) - d_2(\frac{c}{d_2}) \equiv 0 \text{ or } 2 \pmod{4}. \quad (4.10)$$

When $4|c$, (4.10) is always $0 \pmod{4}$, which proves (4.9) by combining the last row of Table 4.1.

When $2||c$, by $(\frac{x}{y}) = (\frac{x}{y})(-1)^{\frac{x-1}{2}}$ for odd x , we have

$$d_1(\frac{c}{d_1}) - d_2(\frac{c}{d_2}) \equiv (\frac{d_1}{c/2}) \left((-1)^{\frac{d_1-1}{2} + \frac{d_1-1}{2} \cdot \frac{c'-1}{2}} (\frac{2}{d_1}) d_1 - (-1)^{\frac{d_2-1}{2} + \frac{d_2-1}{2} \cdot \frac{c'-1}{2}} (\frac{2}{d_2}) d_2 \right) \pmod{4}. \quad (4.11)$$

Since $d_2 = d_1 + \beta c'$, we divide into cases for $c' \equiv 2, 6 \pmod{8}$, $d_1 \equiv 1, 3, 5, 7 \pmod{8}$ and β from 1 to 6 to make Table 4.2. Note that $d_2 \pmod{8}$ is derived by $c' \pmod{8}$, β and $d_1 \pmod{8}$.

(4.10) \searrow	$c' \equiv 2 \pmod{8}$				$c' \equiv 6 \pmod{8}$			
$d_1 \pmod{8}$	1	3	5	7	1	3	5	7
$\beta = 1$	2	0	2	0	2	0	2	0
$\beta = 4$	0	0	0	0	0	0	0	0
$\beta = 5$	2	0	2	0	2	0	2	0
$\beta = 2$	2	2	2	2	2	2	2	2
$\beta = 3$	0	2	0	2	0	2	0	2
$\beta = 6$	2	2	2	2	2	2	2	2

TABLE 4.2. Table for (4.10); $2|c$, no requirement for $(c, 3)$, $7 \nmid c$.

Comparing Table 4.1 and Table 4.2, we have proved (4.9). Combining (4.8) and $12cs(d, c) \equiv 0 \pmod{6}$, we divide $24c'$ to compute $\text{Arg}_2(d_1 \rightarrow d_2; \ell) = \frac{\beta}{7}$. Then $(\text{Arg}_2 + \text{Arg}_3)(d_1 \rightarrow d_2; \ell) = -\frac{\beta}{7}$ and we have Table 4.3.

$c' \pmod{42}$	2	4	8	10	16	20
β	4	2	1	5	4	6
$-3\beta c'^2 \pmod{14}$	10	6	12	8	10	2
$\text{Arg}_1(d_1 \rightarrow d_2; 1) : \frac{1}{2} - \frac{9\beta c'^2}{14}$	$\frac{3}{14}$	$-\frac{1}{14}$	$\frac{5}{14}$	$\frac{1}{14}$	$\frac{3}{14}$	$-\frac{5}{14}$
$(\text{Arg}_2 + \text{Arg}_3)(d_1 \rightarrow d_2; 1)$	$-\frac{4}{7}$	$-\frac{2}{7}$	$-\frac{1}{7}$	$-\frac{5}{7}$	$-\frac{4}{7}$	$-\frac{6}{7}$
Total Arg($d_1 \rightarrow d_2; 1$)	$-\frac{5}{14}$	$-\frac{5}{14}$	$\frac{3}{14}$	$\frac{5}{14}$	$-\frac{5}{14}$	$-\frac{3}{14}$
$c'\ell \equiv \pm 1 \pmod{7}?$			+			-
$-18\beta c'^2 \equiv 3c' \pmod{7}$	6	5	3	2	6	4
$\text{Arg}_1(d_1 \rightarrow d_2; 2) : \frac{1}{2} + \frac{3c'}{7}$	$\frac{5}{14}$	$\frac{3}{14}$	$-\frac{1}{14}$	$-\frac{3}{14}$	$\frac{5}{14}$	$\frac{1}{14}$
$(\text{Arg}_2 + \text{Arg}_3)(d_1 \rightarrow d_2; 1)$	$-\frac{3}{7}$	$-\frac{2}{7}$	$-\frac{1}{7}$	$-\frac{5}{7}$	$-\frac{4}{7}$	$-\frac{6}{7}$
Total Arg($d_1 \rightarrow d_2; 2$)	$-\frac{3}{14}$	$-\frac{1}{14}$	$-\frac{3}{14}$	$\frac{1}{14}$	$-\frac{3}{14}$	$\frac{3}{14}$
$c'\ell \equiv \pm 1 \pmod{7}?$		+		-		
$-81\beta c'^2 \pmod{14}$	6	12	10	2	6	4
$\text{Arg}_1(d_1 \rightarrow d_2; 3) : -\frac{81\beta c'^2}{14}$	$\frac{3}{7}$	$\frac{6}{7}$	$\frac{5}{7}$	$\frac{1}{7}$	$\frac{3}{7}$	$\frac{2}{7}$
$(\text{Arg}_2 + \text{Arg}_3)(d_1 \rightarrow d_2; 1)$	$-\frac{4}{7}$	$-\frac{2}{7}$	$-\frac{1}{7}$	$-\frac{5}{7}$	$-\frac{4}{7}$	$-\frac{6}{7}$
Total Arg($d_1 \rightarrow d_2; 3$)	$-\frac{1}{7}$	$-\frac{3}{7}$	$-\frac{3}{7}$	$\frac{3}{7}$	$-\frac{1}{7}$	$\frac{3}{7}$
$c'\ell \equiv \pm 1 \pmod{7}?$	-				-	
$c' \pmod{42}$	22	26	32	34	38	40
β	1	3	2	6	5	3
$-9\beta c'^2 \pmod{14}$	12	4	6	2	8	4
$\text{Arg}_1(d_1 \rightarrow d_2; 1) : \frac{1}{2} - \frac{9\beta c'^2}{14}$	$\frac{5}{14}$	$-\frac{3}{14}$	$-\frac{1}{14}$	$-\frac{5}{14}$	$\frac{1}{14}$	$-\frac{3}{14}$
$(\text{Arg}_2 + \text{Arg}_3)(d_1 \rightarrow d_2; 1)$	$-\frac{1}{7}$	$-\frac{3}{7}$	$-\frac{2}{7}$	$-\frac{6}{7}$	$-\frac{5}{7}$	$-\frac{3}{7}$
Total Arg($d_1 \rightarrow d_2; 1$)	$\frac{3}{14}$	$\frac{5}{14}$	$-\frac{5}{14}$	$-\frac{3}{14}$	$\frac{5}{14}$	$\frac{5}{14}$
$c'\ell = \pm 1 \pmod{7}?$	+			-		
$-18\beta c'^2 \equiv 3c' \pmod{7}$	3	1	5	4	2	1
$\text{Arg}_1(d_1 \rightarrow d_2; 2) : \frac{1}{2} + \frac{3c'}{7}$	$-\frac{1}{14}$	$-\frac{5}{14}$	$\frac{3}{14}$	$\frac{1}{14}$	$-\frac{3}{14}$	$-\frac{5}{14}$
$(\text{Arg}_2 + \text{Arg}_3)(d_1 \rightarrow d_2; 1)$	$-\frac{1}{7}$	$-\frac{3}{7}$	$-\frac{2}{7}$	$-\frac{6}{7}$	$-\frac{5}{7}$	$-\frac{3}{7}$
Total Arg($d_1 \rightarrow d_2; 2$)	$-\frac{3}{14}$	$\frac{3}{14}$	$-\frac{1}{14}$	$\frac{3}{14}$	$\frac{1}{14}$	$\frac{3}{14}$
$c'\ell \equiv \pm 1 \pmod{7}?$			+		-	
$-81\beta c'^2 \pmod{14}$	10	8	12	4	2	8
$\text{Arg}_1(d_1 \rightarrow d_2; 3) : -\frac{81\beta c'^2}{14}$	$\frac{5}{7}$	$\frac{4}{7}$	$\frac{6}{7}$	$\frac{2}{7}$	$\frac{1}{7}$	$\frac{4}{7}$
$(\text{Arg}_2 + \text{Arg}_3)(d_1 \rightarrow d_2; 1)$	$-\frac{1}{7}$	$-\frac{3}{7}$	$-\frac{2}{7}$	$-\frac{6}{7}$	$-\frac{5}{7}$	$-\frac{3}{7}$
Total Arg($d_1 \rightarrow d_2; 3$)	$-\frac{3}{7}$	$\frac{1}{7}$	$-\frac{3}{7}$	$\frac{3}{7}$	$\frac{3}{7}$	$\frac{1}{7}$
$c'\ell \equiv \pm 1 \pmod{7}?$		+				+

TABLE 4.3. Table for $\text{Arg}(d_1 \rightarrow d_2; \ell); 2|c, 3 \nmid c, 7 \nmid c$.

Next we deal with $\text{Arg}(d_2 \rightarrow d_3; \ell)$ with $d_2 a_4 \equiv d_3 a_5 \equiv 1 \pmod{7}$. For Arg_1 we apply (4.6). For $\text{Arg}_2(d_2 \rightarrow d_3; \ell)$ we do the similar proof as $\text{Arg}_2(d_1 \rightarrow d_2; \ell)$. First we have

$$-12cs(d_3, c) + 12cs(d_2, c) \equiv -d_3 - a_5 + d_2 + a_4 \equiv -2\beta c' \pmod{c}. \quad (4.12)$$

Then $-12cs(d_3, c) + 12cs(d_2, c)$ is a multiple of c' .

We claim that

$$-12cs(d_3, c) + 12cs(d_2, c) \equiv 4 \times 2^\lambda \pmod{8 \times 2^\lambda}. \quad (4.13)$$

For $\lambda \geq 1$ such that $2^\lambda \parallel c$, we have

$$\begin{aligned} -12cs(d_3, c) + 12cs(d_2, c) &\equiv -d_3 - \overline{d_{3\{8 \times 2^\lambda\}}}(c^2 + 3c + 1 + 2c(\frac{c}{d_3})) \\ &\quad + d_2 + \overline{d_{2\{8 \times 2^\lambda\}}}(c^2 + 3c + 1 + 2c(\frac{c}{d_2})) \\ &\equiv -\beta c' + \beta c' \overline{d_{3\{8 \times 2^\lambda\}}} \cdot \overline{d_{2\{8 \times 2^\lambda\}}}(c^2 + 3c + 1) \\ &\quad + 2c(\overline{d_{2\{8 \times 2^\lambda\}}}(\frac{c}{d_2}) - \overline{d_{3\{8 \times 2^\lambda\}}}(\frac{c}{d_3})) \pmod{8 \times 2^\lambda}, \end{aligned}$$

After dividing c' , since $2^\lambda \parallel c'$ and $\overline{x_{\{8\}}} \equiv x \pmod{8}$ for odd x , we have

$$\begin{aligned} -84s(d_3, c) + 84s(d_2, c) &\equiv -\beta + \beta d_3 d_2 (c^2 + 3c + 1) + 6(d_2(\frac{c}{d_2}) - d_3(\frac{c}{d_3})) \\ &\equiv \beta c'(1 + d_2 \beta)(c' + 1) - 2(d_2(\frac{c}{d_2}) - d_3(\frac{c}{d_3})) \pmod{8} \end{aligned}$$

The proof of (4.13) is then the same as the proof of (4.9) before, noting that in the second part we have $(\frac{d_2}{7}) = 1$ while $(\frac{d_3}{7}) = -1$. This difference makes an alternation in Table 4.2 where we should change all 2 to 0 and all 0 to 2, which results in $4 \times 2^\lambda \pmod{8 \times 2^\lambda}$ rather than $0 \pmod{8 \times 2^\lambda}$ in (4.13). We omit the details.

Combining (4.12), (4.13) and $12cs(d, c) \equiv 0 \pmod{6}$ we can determine $\text{Arg}_2(d_2 \rightarrow d_3; \ell)$ with denominator 42 and numerator by $3\beta \pmod{7}$ and $3 \pmod{6}$, hence

$$\text{Arg}_2(d_2 \rightarrow d_3; \ell) = \frac{1, 3, 5, 9, 11, 13}{14} \quad \text{when } \beta = 1, 3, 5, 2, 4, 6, \text{ resp.}$$

Now we have Table 4.4.

Finally we check $\text{Arg}(d_3 \rightarrow d_4; \ell)$ with $d_3 a_5 \equiv d_4 a_2 \equiv 1 \pmod{7}$. For Arg_1 we apply (4.7). Since c' is even, we have $-3\beta c' \equiv 4 \pmod{14}$ and the sign always changes.

For $\text{Arg}_2(d_3 \rightarrow d_4; \ell)$, first we have

$$-12cs(d_4, c) + 12cs(d_3, c) \equiv -d_4 - a_2 + d_3 + a_5 \equiv 2\beta c' \pmod{c}. \quad (4.14)$$

Then $-12cs(d_4, c) + 12cs(d_3, c)$ is a multiple of c' . We claim that

$$-12cs(d_4, c) + 12cs(d_3, c) \equiv 4 \times 2^\lambda \pmod{8 \times 2^\lambda}. \quad (4.15)$$

The proof is the same as the proof for (4.13) and we omit the details. Combining (4.14), (4.15) and (2.2), we can determine $\text{Arg}_2(d_3 \rightarrow d_4; \ell)$ with denominator 42 and numerator by $4\beta \pmod{7}$ and $3 \pmod{6}$, hence

$$\text{Arg}_2(d_2 \rightarrow d_3; \ell) = \frac{1, 3, 5, 9, 11, 13}{14} \quad \text{when } \beta = 6, 4, 2, 5, 3, 1.$$

This gives Table 4.5.

Comparing Tables 4.3, 4.4 and 4.5 we see that when $2 \mid c'$, $3 \nmid c'$ and $7 \nmid c'$, Condition 4.2 holds and we have proved Proposition 4.1 in this case.

$c' \pmod{42}$	2	4	8	10	16	20
β	4	2	1	5	4	6
$-3\beta c'^2 \pmod{14}$	8	2	4	12	8	10
$\text{Arg}_1(d_2 \rightarrow d_3; 1) : \frac{1}{2} - \frac{3\beta c'^2}{14}$	$\frac{1}{14}$	$-\frac{5}{14}$	$-\frac{3}{14}$	$\frac{5}{14}$	$\frac{1}{14}$	$\frac{3}{14}$
$(\text{Arg}_2 + \text{Arg}_3)(d_2 \rightarrow d_3; 1)$	$\frac{9}{14}$	$\frac{1}{14}$	$\frac{11}{14}$	$\frac{13}{14}$	$\frac{9}{14}$	$\frac{3}{14}$
Total Arg($d_2 \rightarrow d_3; 1$)	$-\frac{2}{7}$	$-\frac{2}{7}$	$-\frac{3}{7}$	$\frac{2}{7}$	$-\frac{2}{7}$	$\frac{3}{7}$
$c'\ell \equiv \pm 1 \pmod{7}?$			+			-
$-6\beta c'^2 \equiv c' \pmod{7}$	2	4	1	3	2	6
$\text{Arg}_1(d_2 \rightarrow d_3; 2) : \frac{3c'}{7}$	$\frac{2}{7}$	$\frac{4}{7}$	$\frac{1}{7}$	$\frac{3}{7}$	$\frac{2}{7}$	$\frac{6}{7}$
$(\text{Arg}_2 + \text{Arg}_3)(d_2 \rightarrow d_3; 2)$	$\frac{9}{14}$	$\frac{1}{14}$	$\frac{11}{14}$	$\frac{13}{14}$	$\frac{9}{14}$	$\frac{3}{14}$
Total Arg($d_2 \rightarrow d_3; 2$)	$-\frac{1}{14}$	$-\frac{5}{14}$	$-\frac{1}{14}$	$\frac{5}{14}$	$-\frac{1}{14}$	$\frac{1}{14}$
$c'\ell \equiv \pm 1 \pmod{7}?$		+		-		
$-27\beta c'^2 \pmod{14}$	2	4	8	10	2	6
$\text{Arg}_1(d_2 \rightarrow d_3; 3) : -\frac{27\beta c'^2}{14}$	$\frac{1}{7}$	$\frac{2}{7}$	$\frac{4}{7}$	$\frac{5}{7}$	$\frac{1}{7}$	$\frac{3}{7}$
$(\text{Arg}_2 + \text{Arg}_3)(d_2 \rightarrow d_3; 3)$	$\frac{9}{14}$	$\frac{1}{14}$	$\frac{11}{14}$	$\frac{13}{14}$	$\frac{9}{14}$	$\frac{3}{14}$
Total Arg($d_2 \rightarrow d_3; 3$)	$-\frac{3}{14}$	$\frac{5}{14}$	$\frac{5}{14}$	$-\frac{5}{14}$	$-\frac{3}{14}$	$-\frac{5}{14}$
$c'\ell \equiv \pm 1 \pmod{7}?$	-				-	
$c' \pmod{42}$	22	26	32	34	38	40
β	1	3	2	6	5	3
$-3\beta c'^2 \pmod{14}$	4	6	2	10	12	6
$\text{Arg}_1(d_2 \rightarrow d_3; 1) : \frac{1}{2} - \frac{3\beta c'^2}{14}$	$-\frac{3}{14}$	$-\frac{1}{14}$	$-\frac{5}{14}$	$\frac{3}{14}$	$\frac{5}{14}$	$-\frac{1}{14}$
$(\text{Arg}_2 + \text{Arg}_3)(d_2 \rightarrow d_3; 1)$	$\frac{11}{14}$	$\frac{5}{14}$	$\frac{1}{14}$	$\frac{3}{14}$	$\frac{13}{14}$	$\frac{5}{14}$
Total Arg($d_2 \rightarrow d_3; 1$)	$-\frac{3}{7}$	$\frac{2}{7}$	$-\frac{2}{7}$	$\frac{3}{7}$	$\frac{2}{7}$	$\frac{2}{7}$
$c'\ell = \pm 1 \pmod{7}?$	+			-		
$-6\beta c'^2 \equiv c' \pmod{7}$	1	5	4	6	3	5
$\text{Arg}_1(d_2 \rightarrow d_3; 2) : \frac{3c'}{7}$	$\frac{1}{7}$	$\frac{5}{7}$	$\frac{4}{7}$	$\frac{6}{7}$	$\frac{3}{7}$	$\frac{5}{7}$
$(\text{Arg}_2 + \text{Arg}_3)(d_2 \rightarrow d_3; 2)$	$\frac{11}{14}$	$\frac{5}{14}$	$\frac{1}{14}$	$\frac{3}{14}$	$\frac{13}{14}$	$\frac{5}{14}$
Total Arg($d_2 \rightarrow d_3; 2$)	$-\frac{1}{14}$	$\frac{1}{14}$	$-\frac{5}{14}$	$\frac{1}{14}$	$\frac{5}{14}$	$\frac{1}{14}$
$c'\ell \equiv \pm 1 \pmod{7}?$			+		-	
$-27\beta c'^2 \pmod{14}$	8	12	4	6	10	12
$\text{Arg}_1(d_2 \rightarrow d_3; 3) : -\frac{27\beta c'^2}{14}$	$\frac{4}{7}$	$\frac{6}{7}$	$\frac{2}{7}$	$\frac{3}{7}$	$\frac{5}{7}$	$\frac{6}{7}$
$(\text{Arg}_2 + \text{Arg}_3)(d_2 \rightarrow d_3; 3)$	$\frac{11}{14}$	$\frac{5}{14}$	$\frac{1}{14}$	$\frac{3}{14}$	$\frac{13}{14}$	$\frac{5}{14}$
Total Arg($d_2 \rightarrow d_3; 3$)	$\frac{5}{14}$	$\frac{3}{14}$	$\frac{5}{14}$	$-\frac{5}{14}$	$-\frac{5}{14}$	$\frac{3}{14}$
$c'\ell \equiv \pm 1 \pmod{7}?$		+				+

 TABLE 4.4. Table for $\text{Arg}(d_2 \rightarrow d_3; \ell); 2|c, 3 \nmid c, 7 \nmid c$.

$c' \pmod{42}$	2	4	8	10	16	20
β	4	2	1	5	4	6
$9\beta c'^2 \pmod{14}$	4	8	2	6	4	12
$\text{Arg}_1(d_3 \rightarrow d_4; 1) : \frac{1}{2} + \frac{9\beta c'^2}{14}$	$-\frac{3}{14}$	$\frac{1}{14}$	$-\frac{5}{14}$	$-\frac{1}{14}$	$-\frac{3}{14}$	$\frac{5}{14}$
$(\text{Arg}_2 + \text{Arg}_3)(d_3 \rightarrow d_4; 1)$	$\frac{1}{14}$	$\frac{11}{14}$	$\frac{9}{14}$	$\frac{3}{14}$	$\frac{1}{14}$	$\frac{5}{14}$
Total Arg($d_2 \rightarrow d_3; 1$)	$-\frac{1}{7}$	$-\frac{1}{7}$	$\frac{2}{7}$	$\frac{1}{7}$	$-\frac{1}{7}$	$-\frac{2}{7}$
$c'\ell \equiv \pm 1 \pmod{7}?$			+			-
$18\beta c'^2 \equiv 4c' \pmod{7}$	1	2	4	5	1	3
$\text{Arg}_1(d_3 \rightarrow d_4; 2) : \frac{1}{2} + \frac{4c'}{7}$	$-\frac{5}{14}$	$-\frac{3}{14}$	$\frac{1}{14}$	$\frac{3}{14}$	$-\frac{5}{14}$	$-\frac{1}{14}$
$(\text{Arg}_2 + \text{Arg}_3)(d_3 \rightarrow d_4; 1)$	$\frac{1}{14}$	$\frac{11}{14}$	$\frac{9}{14}$	$\frac{3}{14}$	$\frac{1}{14}$	$\frac{5}{14}$
Total Arg($d_2 \rightarrow d_3; 2$)	$-\frac{2}{7}$	$-\frac{3}{7}$	$-\frac{2}{7}$	$\frac{3}{7}$	$-\frac{2}{7}$	$\frac{2}{7}$
$c'\ell \equiv \pm 1 \pmod{7}?$		+		-		
$81\beta c'^2 \pmod{14}$	8	2	4	12	8	10
$\text{Arg}_1(d_3 \rightarrow d_4; 3) : \frac{1}{2} + \frac{81\beta c'^2}{14}$	$\frac{1}{14}$	$-\frac{5}{14}$	$-\frac{3}{14}$	$\frac{5}{14}$	$\frac{1}{14}$	$\frac{3}{14}$
$(\text{Arg}_2 + \text{Arg}_3)(d_3 \rightarrow d_4; 1)$	$\frac{1}{14}$	$\frac{11}{14}$	$\frac{9}{14}$	$\frac{3}{14}$	$\frac{1}{14}$	$\frac{5}{14}$
Total Arg($d_3 \rightarrow d_4; 3$)	$\frac{1}{7}$	$\frac{3}{7}$	$\frac{3}{7}$	$-\frac{3}{7}$	$\frac{1}{7}$	$-\frac{3}{7}$
$c'\ell \equiv \pm 1 \pmod{7}?$	-				-	
$c' \pmod{42}$	22	26	32	34	38	40
β	1	3	2	6	5	3
$9\beta c'^2 \pmod{14}$	2	10	8	12	6	10
$\text{Arg}_1(d_3 \rightarrow d_4; 1) : \frac{1}{2} + \frac{9\beta c'^2}{14}$	$-\frac{5}{14}$	$\frac{3}{14}$	$\frac{1}{14}$	$\frac{5}{14}$	$-\frac{1}{14}$	$\frac{3}{14}$
$(\text{Arg}_2 + \text{Arg}_3)(d_3 \rightarrow d_4; 1)$	$\frac{9}{14}$	$\frac{13}{14}$	$\frac{11}{14}$	$\frac{5}{14}$	$\frac{3}{14}$	$\frac{13}{14}$
Total Arg($d_3 \rightarrow d_4; 1$)	$\frac{2}{7}$	$\frac{1}{7}$	$-\frac{1}{7}$	$-\frac{2}{7}$	$\frac{1}{7}$	$\frac{1}{7}$
$c'\ell = \pm 1 \pmod{7}?$	+			-		
$18\beta c'^2 \equiv 4c' \pmod{7}$	4	6	2	3	5	6
$\text{Arg}_1(d_3 \rightarrow d_4; 2) : \frac{1}{2} + \frac{4c'}{7}$	$\frac{1}{14}$	$\frac{5}{14}$	$-\frac{3}{14}$	$-\frac{1}{14}$	$\frac{3}{14}$	$\frac{5}{14}$
$(\text{Arg}_2 + \text{Arg}_3)(d_3 \rightarrow d_4; 2)$	$\frac{9}{14}$	$\frac{13}{14}$	$\frac{11}{14}$	$\frac{5}{14}$	$\frac{3}{14}$	$\frac{13}{14}$
Total Arg($d_2 \rightarrow d_3; 2$)	$-\frac{2}{7}$	$\frac{2}{7}$	$-\frac{3}{7}$	$\frac{2}{7}$	$\frac{3}{7}$	$\frac{2}{7}$
$c'\ell \equiv \pm 1 \pmod{7}?$			+		-	
$81\beta c'^2 \pmod{14}$	4	6	2	10	12	6
$\text{Arg}_1(d_2 \rightarrow d_3; 3) : \frac{1}{2} + \frac{81\beta c'^2}{14}$	$-\frac{3}{14}$	$-\frac{1}{14}$	$-\frac{5}{14}$	$\frac{3}{14}$	$\frac{5}{14}$	$-\frac{1}{14}$
$(\text{Arg}_2 + \text{Arg}_3)(d_3 \rightarrow d_4; 1)$	$\frac{9}{14}$	$\frac{13}{14}$	$\frac{11}{14}$	$\frac{5}{14}$	$\frac{3}{14}$	$\frac{13}{14}$
Total Arg($d_2 \rightarrow d_3; 3$)	$\frac{3}{7}$	$-\frac{1}{7}$	$\frac{3}{7}$	$-\frac{3}{7}$	$-\frac{3}{7}$	$\frac{1}{7}$
$c'\ell \equiv \pm 1 \pmod{7}?$		+				+

 TABLE 4.5. Table for $\text{Arg}(d_3 \rightarrow d_4; \ell)$; $2|c$, $3 \nmid c$, $7 \nmid c$.

4.2. **Case $7|c'$.** We still denote $c' = c/7$ and $V(r, c) = \{d \pmod{c}^* : d \equiv r \pmod{c'}\}$ for $r \pmod{c'}^*$. Now $|V(r, c)| = 7$. Since $(d + c', c) = 1$ when $(d, c) = 1$, we can write $V(r, c) = \{d, d + c', d + 2c', \dots, d + 6c'\}$ for $1 \leq d < c'$ and $d \equiv r \pmod{c'}$.

We claim that Proposition 4.1 is still true:

$$s_{r,c} := \sum_{d \in V(r,c)} \frac{e\left(-\frac{3c'al^2}{14}\right)}{\sin\left(\frac{\pi al}{7}\right)} e\left(-\frac{12cs(d,c)}{24c}\right) e\left(\frac{5d}{c}\right) = 0, \quad (4.16)$$

while this time we have seven summands. We prove (4.16) by showing that there are only three cases for the sum: all at the outer circle (radius $1/\sin(\frac{\pi}{7})$), all at the middle circle (radius $1/\sin(\frac{2\pi}{7})$), and all at the inner circle (radius $1/\sin(\frac{3\pi}{7})$). Moreover, the seven points are equally distributed. Similar as before, we still denote P_1 , P_2 and P_3 for each term in (4.16) and investigate the argument differences contributed from each term.

For any $d \in V(r, c)$ and $a \pmod{c}$ such that $ad \equiv 1 \pmod{c}$, we define $d_* = d + c'$ and a_* by $a_*d_* \equiv 1 \pmod{c}$. Specifically, we take $a_* = a - c', a - 2c', a + 3c', a + 3c', a - 2c', a - c'$, when $d \equiv 1, 2, 3, 4, 5, 6 \pmod{7}$, respectively. Note that $P_1(d) = (-1)^{cal}/\sin(\frac{\pi al}{7})$ has period c' . Hence we always have

$$\text{Arg}_1(d \rightarrow d_*; \ell) = 0 \quad \text{and} \quad \text{Arg}_3(d \rightarrow d_*; \ell) = \frac{5}{7}.$$


In the following two cases, we prove

$$\text{Arg}(d \rightarrow d_*; \ell) = \begin{cases} -\frac{2}{7} & d \equiv 1, 6 \pmod{7}; \\ \frac{3}{7} & d \equiv 2, 5 \pmod{7}; \\ -\frac{1}{7} & d \equiv 3, 4 \pmod{7}. \end{cases} \quad (4.17)$$

when $\ell = 1$. In the other cases $\ell = 2, 3$, only P_1 is affected and we still get (4.17).

Proof of Proposition 4.1 when $49|c$. It is clear that (4.17) implies (4.16). \square

One may visualize (4.17) in the following graphs:



4.2.1. *c is odd.* First we suppose $3 \nmid c$. When $d \equiv 1, 6 \pmod{7}$, by (2.2) we have $-12cs(d_*, c) + 12cs(d, c) \equiv 0 \pmod{6}$,

$$-12cs(d_*, c) + 12cs(d, c) \equiv -d_* - a_* + d + a \equiv 0 \pmod{c}, \quad (4.18)$$

and by (2.4) we have

$$-12cs(d_*, c) + 12cs(d, c) \equiv 2\left(\frac{d_*}{c}\right) - 2\left(\frac{d}{c}\right) \equiv 0 \pmod{8}$$

because $\left(\frac{d+c'}{c}\right) = \left(\frac{d+c'}{7}\right)\left(\frac{d+c'}{c'}\right) = \left(\frac{d}{7}\right)\left(\frac{d}{c'}\right) = \left(\frac{d}{c}\right)$ always. Then $-12cs(d_*, c) + 12cs(d, c) \equiv 0 \pmod{24c}$ and $\text{Arg}_2(d \rightarrow d_*; \ell) = 0$. Since $\text{Arg}_3(d \rightarrow d_*; \ell) = \frac{5}{7}$, we have proved (4.17).

When $d \equiv 2, 5 \pmod{7}$, only (4.18) is affected and becomes

$$-12cs(d_*, c) + 12cs(d, c) \equiv -d_* - a_* + d + a \equiv c' \pmod{c}. \quad (4.19)$$

After dividing $24c'$ we get $\text{Arg}_2(d \rightarrow d_*; \ell) = \frac{5}{7}$. We have proved (4.17) in this case.

When $d \equiv 3, 4 \pmod{7}$, (4.18) becomes

$$-12cs(d_*, c) + 12cs(d, c) \equiv -d_* - a_* + d + a \equiv -4c' \pmod{c}. \quad (4.20)$$

We get $\text{Arg}_2(d \rightarrow d_*; \ell) = \frac{1}{7}$ and (4.17).

Then we investigate the case $3|c'$. The following congruence

$$-12cs(d_*, c) + 12cs(d, c) \equiv 2\left(\frac{d_*}{c}\right) - 2\left(\frac{d}{c}\right) \equiv 0 \pmod{8}$$

still holds and we compute

$$-12cs(d_*, c) + 12cs(d, c) \equiv -d_* - \overline{d_{1\{3c\}}} + d + \overline{d_{\{3c\}}} \equiv -c' + c'\overline{d_{1\{3c\}}} \cdot \overline{d_{\{3c\}}} \pmod{3c},$$

so

$$-84s(d_*, c) + 84s(d, c) \equiv -1 + a_*a \pmod{21}.$$

Since $a_*a \equiv 1 \pmod{3}$ and $a_* \equiv a \pmod{7}$, we have

$$-84s(d_*, c) + 84s(d, c) \equiv \begin{cases} 0 \pmod{21} & \text{if } d \equiv 1, 6 \pmod{7}, \\ 15 \pmod{21} & \text{if } d \equiv 2, 5 \pmod{7}, \\ 3 \pmod{21} & \text{if } d \equiv 3, 4 \pmod{7}. \end{cases} \quad (4.21)$$

Then $-28s(d_*, c) + 28s(d, c) \equiv 0, 5, 1 \pmod{7}$ and $\text{Arg}_2(d \rightarrow d_*; \ell) = \frac{0, 5, 1}{7}$, respectively. We have proved (4.17) when c is odd.

4.2.2. *c is even.* The first case is $3 \nmid c'$. Congruences (4.18), (4.19) and (4.20) are still valid here. By (2.5), we define $\lambda \geq 1$ by $2^\lambda || c'$ and claim that

$$-12cs(d_*, c) + 12cs(d, c) \equiv 0 \pmod{8 \times 2^\lambda} \quad (4.22)$$

To compute this, we have

$$\begin{aligned} -12cs(d_*, c) + 12cs(d, c) &\equiv -d_* - \overline{d_{1\{8 \times 2^\lambda\}}}(c^2 + 3c + 1) - \overline{d_{1\{8 \times 2^\lambda\}}} \cdot 2c\left(\frac{c}{d_*}\right) \\ &\quad + d + \overline{d_{\{8 \times 2^\lambda\}}}(c^2 + 3c + 1) + \overline{d_{\{8 \times 2^\lambda\}}} \cdot 2c\left(\frac{c}{d}\right) \pmod{8 \times 2^\lambda} \\ &\equiv -c' + c'(c^2 + 3c + 1)\overline{d_{1\{8 \times 2^\lambda\}}} \cdot \overline{d_{\{8 \times 2^\lambda\}}} \\ &\quad - \overline{d_{1\{8 \times 2^\lambda\}}} \cdot 2c\left(\frac{c}{d_*}\right) + \overline{d_{\{8 \times 2^\lambda\}}} \cdot 2c\left(\frac{c}{d}\right) \pmod{8 \times 2^\lambda}. \end{aligned}$$

After dividing c' we have

$$\begin{aligned} -84s(d_*, c) + 84s(d, c) &\equiv -1 + d_*d(c^2 + 3c + 1) + 2\left(\frac{c}{d_*}\right)d_* - 2\left(\frac{c}{d}\right)d \pmod{8} \\ &\equiv c'(c' + 1)(d + 1) + 2\left(\frac{c}{d_*}\right)d_* - 2\left(\frac{c}{d}\right)d \pmod{8}. \end{aligned}$$

We also get

$$c'(c' + 1)(d + 1) \equiv \begin{cases} 4 \pmod{8} & \text{if } 2||c, d \equiv 1 \pmod{4}, \\ 0 \pmod{8} & \text{if } 2||c, d \equiv 3 \pmod{4}, \\ 0 \pmod{8} & \text{if } 4|c. \end{cases} \quad (4.23)$$

When $4|c$, it is not hard to show $\left(\frac{c}{d_*}\right)d_* - \left(\frac{c}{d}\right)d \equiv 0 \pmod{4}$ and we have proved (4.22) in this case.

When $2||c$, we have Table 4.6 for $\text{val} := \left(\frac{c}{d_*}\right)d_* - \left(\frac{c}{d}\right)d \pmod{4}$ using quadratic reciprocity. Combining Table 4.6 and (4.23) we obtain (4.22).

$d \pmod{8}$	1	3	5	7
$d_* \pmod{8}$ when $c' \equiv 2 \pmod{8}$	3	5	7	1
val	2	0	2	0
$d_* \pmod{8}$ when $c' \equiv 6 \pmod{8}$	7	1	3	5
val	2	0	2	0

TABLE 4.6. Table for $\text{val} := (\frac{c}{d_*})d_* - (\frac{c}{d})d \pmod{4}$; $2|c$, no requirement for $(3, c)$, $7|c$.

Combining (4.22) with (4.18), (4.19) and (4.20), we have proved (4.17) when $2|c$ and $3 \nmid c$. When $3|c$, we use (4.21) instead of (4.18), (4.19) and (4.20). This finishes the proof of (4.17). We have proved Proposition 4.1, which implies (7-5,1) of Theorem 1.3.

5. PROOF OF (7-5,2) OF THEOREM 1.3

For all $1 \leq \ell \leq 6$, $n \geq 0$, $7|c$ and $7 \nmid A$, if $A\ell \equiv \pm 1 \pmod{7}$ and $c = 7A$, (7-5,2) becomes

$$e(\frac{1}{8})S_{\infty\infty}^{(\ell)}(0, 7n+5, c, \mu_7) + 2i\sqrt{7}S_{0\infty}^{(\ell)}(0, 7n+5, A, \mu_7; 0) = 0. \quad (5.1)$$

We still denote $c' = c/7 = A$ and $V(r, c) := \{d \pmod{c}^* : d \equiv r \pmod{c}\}$ for $(r, c') = 1$.

Recall (2.7) for $p = 7$. By $\ell c \equiv \ell A \pmod{2}$ we have

$$e(\frac{1}{8})S_{\infty\infty}^{(\ell)}(0, 7n+5, c, \mu_7) = \sum_{d \pmod{c}^*} \frac{(-1)^{\ell A} \exp\left(-\frac{3\pi i c' a \ell^2}{7}\right)}{\sin\left(\frac{\pi a \ell}{7}\right)} e^{-\pi i s(d, c)} e\left(\frac{(7n+5)d}{c}\right). \quad (5.2)$$

Recall that $[A\ell]$ is defined as the least non-negative residue of $A\ell \pmod{7}$. By (2.8), when $[A\ell] = 1$, we denote T by $A\ell = 7T + 1$ and

$$\begin{aligned} & 2i\sqrt{7}S_{0\infty}^{(\ell)}(0, 7n+5, A, \mu_7; 0) \\ &= 2i\sqrt{7}(-1)^{A\ell - [A\ell]} \sum_{B \pmod{A}^*} e\left(\frac{(\frac{3}{2}T^2 + \frac{1}{2}T)C}{A}\right) e^{-\pi i s(B, A)} e\left(\frac{(7n+5)B}{A}\right). \end{aligned} \quad (5.3)$$

By (2.9), when $[A\ell] = 6$, we denote T by $A\ell = 7T - 1$ and have

$$\begin{aligned} & 2i\sqrt{7}S_{0\infty}^{(\ell)}(0, 7n+5, A, \mu_7; 0) \\ &= 2i\sqrt{7}(-1)^{A\ell - [A\ell]} \sum_{B \pmod{A}^*} e\left(\frac{(\frac{3}{2}(T-1)^2 + \frac{5}{2}(T-1) + 1)C}{A}\right) e^{-\pi i s(B, A)} e\left(\frac{(7n+5)B}{A}\right); \end{aligned} \quad (5.4)$$

For $(r, c') = 1$ and any $d \in V(r, c)$, we define $P(d)$ as

$$P(d) := \frac{(-1)^{[A\ell]} e\left(-\frac{3c' a \ell^2}{14}\right)}{\sin\left(\frac{\pi a \ell}{7}\right)} e^{-\pi i s(d, c)} e\left(\frac{(7n+5)d}{c}\right) =: P_1(d) \cdot P_2(d) \cdot P_3(d). \quad (5.5)$$

When $A\ell = 7T + 1$, we denote $Q_1(B) = i$, $Q_3(B) = e(\frac{(7n+5)B}{A})$,

$$Q_2(B) := e\left(\frac{(\frac{3}{2}T^2 + \frac{1}{2}T)C}{A}\right) e^{-\pi i s(B, A)} \quad \text{and} \quad Q(B) =: 2\sqrt{7} \cdot Q_1(B)Q_2(B)Q_3(B); \quad (5.6)$$

when $A\ell = 7T - 1$, we only change the definition of $Q_2(B)$ to

$$Q_2(B) := e \left(\frac{\left(\frac{3}{2}(T-1)^2 + \frac{5}{2}(T-1) + 1\right)C}{A} \right) e^{-\pi i s(B,A)} \quad (5.7)$$

and still denote $Q(B) = 2\sqrt{7} \cdot Q_1(B)Q_2(B)Q_3(B)$.

We divide the cases according to $c'\ell \equiv \pm 1 \pmod{7}$, ℓ , and the divisibility of A by 2, 3. For each $r \pmod{A}^*$, recall that $d_1 \in V(r, c)$ is the unique $d_1 \pmod{c}^*$ such that $d_1 \equiv 1 \pmod{7}$. We compare the argument difference from $Q(B)$ to $P(d_1)$, where we choose

$$B = \begin{cases} -d_1 T, & A\ell = 7T + 1, \\ d_1 T, & A\ell = 7T - 1, \end{cases} \quad \text{and} \quad C = -7\overline{d_1\{A\}}. \quad (5.8)$$

We denote $\text{Arg}(Q_j \rightarrow P_j; \ell)$ in the following way: suppose $P_j(d_1) = Re^{i\Theta}$ and $Q_j(B) = R_B e^{i\Theta_B}$, then

$$\text{Arg}(Q_j \rightarrow P_j; \ell) = \alpha \quad \text{if and only if} \quad \Theta - \Theta_B = \alpha \cdot 2\pi + 2k\pi \quad \text{for } k \in \mathbb{Z}.$$

We also denote $\text{Arg}(Q \rightarrow P; \ell) = \sum_{j=1}^3 \text{Arg}(Q_j \rightarrow P_j; \ell)$. Note that if $\text{Arg}(Q_j \rightarrow P_j; \ell) = \alpha$, then $\text{Arg}(Q_j \rightarrow P_j; \ell) = \alpha + k$ for all $k \in \mathbb{Z}$.


With the notation above, we claim that the argument differences satisfy the following proposition.


Proposition 5.1. *For $c = 7c' = 7A$, any $r \pmod{c'}^*$, $d_1 \in V(r, c)$ and B chosen by (5.8), we have*

$$A\ell = 7T + 1 : \quad \text{Arg}(Q \rightarrow P; \ell) = -\frac{3}{7}, -\frac{5}{14}, \frac{3}{14} \quad \text{for } \ell = 1, 2, 3, \text{ respectively}; \quad (5.9)$$

$$A\ell = 7T - 1 : \quad \text{Arg}(Q \rightarrow P; \ell) = \frac{3}{7}, \frac{5}{14}, -\frac{3}{14} \quad \text{for } \ell = 1, 2, 3, \text{ respectively}. \quad (5.10)$$

To visualize the argument differences, here are a few examples:





The red circles in the figures are centered at the origin with radii $\csc(\frac{\pi}{7})$, $\csc(\frac{2\pi}{7})$, and $\csc(\frac{3\pi}{7})$, respectively, from the outside to the inside. The point labeled by B represents $\frac{Q(B)}{2}$.

For the styles of the six points $P(d_j)$ for $d_j \in V(r, c)$, we have the following condition. This *has already been proved* by the tables in the former section, corresponding to the rows marked with “ $c'\ell \equiv \pm 1 \pmod{7}$?” whose entries are + or $-$.

Condition 5.2. *When $c'\ell \equiv \pm 1 \pmod{7}$, we have the following six styles for these six points $P(d)$ for $d \in V(r, c)$.*

- $\ell = 1$. *When $c'\ell \equiv 1 \pmod{7}$, the arguments going $d_1 \rightarrow d_2 \rightarrow d_3 \rightarrow d_4 \rightarrow d_5 \rightarrow d_6 \rightarrow d_1$ are $\frac{3}{14}, -\frac{3}{7}, \frac{2}{7}, -\frac{3}{7}, \frac{3}{14}, \frac{1}{7}$, respectively. When $c'\ell \equiv -1 \pmod{7}$, the direction is reversed, as shown in the second line.*

	$d_1 \rightarrow$	$d_2 \rightarrow$	$d_3 \rightarrow$	$d_4 \rightarrow$	$d_5 \rightarrow$	$d_6 \rightarrow$	d_1
$c' \equiv 1 \pmod{7}$	$\frac{3}{14}$	$-\frac{3}{7}$	$\frac{2}{7}$	$-\frac{3}{7}$	$\frac{3}{14}$	$\frac{1}{7}$	
$c' \equiv 6 \pmod{7}$	$-\frac{3}{14}$	$\frac{3}{7}$	$-\frac{2}{7}$	$\frac{3}{7}$	$-\frac{3}{14}$	$-\frac{1}{7}$	

- $\ell = 2$. *The first line is for $c'\ell \equiv 1 \pmod{7}$ and the second line is for $c'\ell \equiv -1 \pmod{7}$.*

	$d_1 \rightarrow$	$d_2 \rightarrow$	$d_3 \rightarrow$	$d_4 \rightarrow$	$d_5 \rightarrow$	$d_6 \rightarrow$	d_1
$c' \equiv 4 \pmod{7}$	$-\frac{1}{14}$	$-\frac{5}{14}$	$-\frac{3}{7}$	$-\frac{5}{14}$	$-\frac{1}{14}$	$\frac{2}{7}$	
$c' \equiv 3 \pmod{7}$	$\frac{1}{14}$	$\frac{5}{14}$	$\frac{3}{7}$	$\frac{5}{14}$	$\frac{1}{14}$	$-\frac{2}{7}$	

- $\ell = 3$. *The first line is for $c'\ell \equiv 1 \pmod{7}$ and the second line is for $c'\ell \equiv -1 \pmod{7}$.*

	$d_1 \rightarrow$	$d_2 \rightarrow$	$d_3 \rightarrow$	$d_4 \rightarrow$	$d_5 \rightarrow$	$d_6 \rightarrow$	d_1
$c' \equiv 5 \pmod{7}$	$\frac{1}{7}$	$\frac{3}{14}$	$-\frac{1}{7}$	$\frac{3}{14}$	$\frac{1}{7}$	$\frac{3}{7}$	
$c' \equiv 2 \pmod{7}$	$-\frac{1}{7}$	$-\frac{3}{14}$	$\frac{1}{7}$	$-\frac{3}{14}$	$-\frac{1}{7}$	$-\frac{3}{7}$	

If the six points $P(d)$ for $d \in V(r, c)$ satisfy Condition 5.2, and $\text{Arg}(Q \rightarrow P; \ell)$ satisfies (5.9) and (5.10) in the corresponding cases, then we have

$$s_{r,c} := \sum_{d \in V(r,c)} P(d) + Q(B) = 0. \quad (5.11)$$

Note that B is chosen from $d_1 \in V(r, c)$ and A , hence from r and c . One way to prove (5.11) is by using the identity

$$\frac{\cos(\frac{\pi}{7})}{\sin(\frac{\pi}{7})} + \frac{\cos(\frac{2\pi}{7})}{\sin(\frac{2\pi}{7})} - \frac{\cos(\frac{3\pi}{7})}{\sin(\frac{3\pi}{7})} = \sqrt{7}, \quad \text{where } \frac{1}{\sin(\frac{j\pi}{7})} \text{ for } j = 1, 2, 3 \text{ are the radii.}$$

Proof of (7-5,2) of Theorem 1.3. This is implied by (5.1), which is proved by (5.11), (5.2), (5.3), (5.4), $7B \equiv r \pmod{A}$, and

$$\begin{aligned} & e\left(\frac{1}{8}\right)S_{\infty\infty}^{(\ell)}(0, 7n+5, c, \mu_7) + 2i\sqrt{7}S_{0\infty}^{(\ell)}(0, 7n+5, A, \mu_7; 0) \\ &= (-1)^{A\ell-[A\ell]} \sum_{r \pmod{A}^*} s_{r,c} e\left(\frac{nr}{A}\right) = 0. \end{aligned}$$

□

Subsections §5.1-§5.4 are devoted to prove (5.9), i.e. the cases $A\ell = c'\ell \equiv 1 \pmod{7}$. We will not repeat the proof for (5.10) but just list a few key intermediate steps at the end of this section.

5.1. Case $c'\ell \equiv 1 \pmod{7}$, $2 \nmid A$, and $3 \nmid A$. Recall $d_1 \equiv 1 \pmod{7}$ and $d_1 \equiv r \pmod{c'}$. Recall that we define $1 \leq \beta \leq 6$ as $\beta c' \equiv 1 \pmod{7}$ and here $\beta = \ell$. Note that $d_1 - \beta A \equiv 7B \pmod{7A}$:

$$7B = d_1(1 - A\ell) \equiv d_1 + (7 - d_1)\ell A \pmod{7A}, \text{ so } 7B \equiv \begin{cases} 0 \pmod{7}, \\ r \pmod{A}. \end{cases}$$

On the other hand, $d_1 - \beta c' \equiv r \pmod{A}$ and $d_1 - \beta c' \equiv 0 \pmod{7}$. The argument difference between P_3 and Q_3 is easy to compute:

$$7 \operatorname{Arg}(Q_3 \rightarrow P_3; \ell) \equiv 5d_1\ell \equiv 5\ell \pmod{7} \quad (5.12)$$

which does not depend on n .

Recall $\overline{d_{1\{7A\}}} \equiv a_1 \pmod{7A}$ and $\overline{B_{\{A\}}} \equiv \overline{7d_{1\{A\}}} \pmod{A}$. We have

$$\begin{aligned} -84A(s(d_1, 7A) - s(B, A)) &\equiv -d_1 - a_1 + d_1(1 - \beta A) + 49\overline{d_{1\{A\}}} \\ &\equiv -d_1\beta A - a_1 + 49\overline{d_{1\{A\}}} \pmod{7A}. \end{aligned}$$

Hence

$$-84A(s(d_1, 7A) - s(B, A)) \equiv \begin{cases} -2 \pmod{7} \\ 48\overline{d_{1\{A\}}} \pmod{A} \end{cases} \quad (5.13)$$

We also have

$$\begin{aligned} -84A(s(d_1, 7A) - s(B, A)) &\equiv -7A - 1 + 2\left(\frac{d_1}{7A}\right) + 7(A+1) - 14\left(\frac{B}{A}\right) \\ &\equiv 6 + 2\left(\frac{d_1}{A}\right) + 2\left(\frac{d_1}{A}\right)\left(\frac{7}{A}\right) \pmod{8}, \end{aligned}$$

where the last step is because $(A, 7) = 1$, $\left(\frac{d_1}{7}\right) = \left(\frac{1}{7}\right) = 1$ and $7B \equiv d_1 \pmod{A}$. By A is odd and $A\ell \equiv 1 \pmod{7}$, we have $\left(\frac{7}{A}\right) = \left(\frac{\ell}{7}\right)(-1)^{\frac{A-1}{2}}$. Combining $6|12cs(d_1, c)$ and $6|12As(B, A)$ we conclude

$$-84A(s(d_1, 7A) - s(B, A)) \equiv \begin{cases} 18 \pmod{24}, & \text{if } A \equiv 1 \pmod{4} (\ell = 1, 2) \\ & \text{or if } A \equiv 3 \pmod{4} (\ell = 3) \\ 6 \pmod{24}, & \text{if } A \equiv 3 \pmod{4} (\ell = 1, 2) \\ & \text{or if } A \equiv 1 \pmod{4} (\ell = 3). \end{cases} \quad (5.14)$$

Next we check the part of Q_2 other than $e^{-\pi is(B,A)}$. Since A is odd and T is even, we have

$$\begin{aligned} \left(\frac{3}{2}T^2 + \frac{1}{2}T\right) C &\equiv \frac{T}{2}(3T+1)(-7\overline{d_{1\{A\}}}) \\ &\equiv \frac{T}{2}(3-3A\ell-7)\overline{d_{1\{A\}}} \\ &\equiv -2T\overline{d_{1\{A\}}} \pmod{A}. \end{aligned}$$

Then the part of Q_2 other than $e^{-\pi is(d,c)}$ is

$$e\left(\frac{24 \cdot 2\overline{d_{1\{A\}}}(-7T)}{24 \cdot 7A}\right) = e\left(\frac{48\overline{d_{1\{A\}}}(1 - A\ell)}{168A}\right), \text{ with numerator } \equiv \begin{cases} 0 \pmod{7}, \\ 48\overline{d_{1\{A\}}} \pmod{A}, \\ 0 \pmod{24}. \end{cases} \quad (5.15)$$

We conclude that

$$24 \cdot 7A \text{Arg}(Q_2 \rightarrow P_2; \ell) \equiv R_2 \pmod{168A} \quad (5.16)$$

where R_2 is determined by (5.13), (5.14) and (5.15): $R_2 \equiv 0 \pmod{A}$, $R_2 \equiv -2 \pmod{7}$, and $R_2 \equiv 18, 6 \pmod{24}$ depending on the cases in (5.14). Therefore, by $A\ell \equiv 1 \pmod{7}$ and $A \pmod{4}$ in (5.14) we conclude

$$\text{Arg}(Q_2 \rightarrow P_2; \ell) = \frac{23, 11, 13}{28} \text{ for } \ell = 1, 2, 3. \quad (5.17)$$

Then we compute $\text{Arg}(Q_1 \rightarrow P_1; \ell)$. When $\ell = 1$, since A is odd, $A \equiv 1 \pmod{14}$. Note that both $a_1 \equiv 1, 8 \pmod{14}$ give the same result due to the sign of $\sin(\frac{\pi a}{7})$. It is direct to get

$$\text{Arg}(Q_1 \rightarrow P_1; 1) = \frac{1}{2} - \frac{3}{14} - \frac{1}{4} = \frac{1}{28}. \quad (5.18)$$

When $\ell = 2$, we get $A \equiv 4 \pmod{7}$ and

$$\text{Arg}(Q_1 \rightarrow P_1; 2) = \frac{1}{2} - \frac{3}{7} - \frac{1}{4} = -\frac{5}{28}. \quad (5.19)$$

When $\ell = 3$, we have $A \equiv 5 \pmod{14}$ and

$$\text{Arg}(Q_1 \rightarrow P_1; 3) = \frac{1}{2} - \frac{9}{14} - \frac{1}{4} = -\frac{11}{28}. \quad (5.20)$$

Combining (5.18), (5.19), (5.20), (5.17), and (5.12) proves (5.9).

5.2. Case $c'\ell \equiv 1 \pmod{7}$, $2 \nmid A$, and $3|A$. In this case (5.12) still holds. For $\text{Arg}(Q_2 \rightarrow P_2; \ell)$, by (2.3) we have

$$-84A(s(d_1, 7A) - s(B, A)) \equiv -d_1 A\ell - \overline{d_{1\{21A\}}} + 7\overline{(-d_1 T)_{\{3A\}}} \pmod{21A}.$$

We have

$$\begin{aligned} -84A(s(d_1, 7A) - s(B, A)) &\equiv -d_1 A\ell - \overline{d_{1\{3A\}}} + 49\overline{(d_1 - d_1 A\ell)_{\{3A\}}} \\ &\equiv -d_1 A\ell + (48d_1 + d_1 A\ell)\overline{d_{1\{3A\}}(d_1 - d_1 A\ell)_{\{3A\}}} \\ &\equiv d_1 A\ell \left(\overline{d_{1\{3A\}}(d_1 - d_1 A\ell)_{\{3A\}}} - 1 \right) + 48\overline{d_{1\{A\}}} \\ &\equiv 48\overline{d_{1\{A\}}} \pmod{3A} \end{aligned} \quad (5.21)$$

where in the second congruence we use

$$\overline{(x+y)_m} - 49\overline{x_{\{m\}}} \equiv \overline{x_{\{m\}}(x+y)_{\{m\}}}(-48x - 49y) \pmod{m}$$

for $(x+y, m) = (x, m) = 1$ and in the last two congruences we use

$$m_1\overline{x_{\{m_1 m_2\}}} \equiv m_1\overline{x_{\{m_2\}}} \pmod{m_1 m_2} \quad (5.22)$$

for $(x, m_1 m_2) = 1$. We still have

$$-84A(s(d_1, 7A) - s(B, A)) \equiv -2 \pmod{7}. \quad (5.23)$$

Moreover, (5.14) and (5.15) still hold except the second congruence in (5.15) should be changed to $48\overline{d_{1\{A\}}}$ (mod $3A$).

We conclude

$$24 \cdot 7A \operatorname{Arg}(Q_2 \rightarrow P_2; \ell) \equiv R_2 \pmod{168A} \quad (5.24)$$

where R_2 is determined by (5.21), (5.23), (5.14) and (5.15): $R_2 \equiv 0 \pmod{3A}$, $R_2 \equiv -2 \pmod{7}$, and $R_2 \equiv 18, 6 \pmod{24}$ depending on the cases in (5.14). Therefore, by $Al \equiv 1 \pmod{7}$ and (5.14) we conclude

$$\operatorname{Arg}(Q_2 \rightarrow P_2; \ell) = \frac{23, 11, 13}{28} \quad \text{for } \ell = 1, 2, 3. \quad (5.25)$$

The condition $3|A$ does not affect $\operatorname{Arg}(Q_1 \rightarrow P_1; \ell)$ and $\operatorname{Arg}(Q_3 \rightarrow P_3; \ell)$. Combining (5.25) with (5.18), (5.19), (5.20), and (5.12), we have proved (5.9) in this case.

5.3. Case $c'\ell \equiv 1 \pmod{7}$, $2|A$, and $3 \nmid A$. Recall (5.12). For $\operatorname{Arg}(Q_2 \rightarrow P_2; \ell)$ we have (5.13) and need to use (2.5). Let $\lambda \geq 1$ be defined as $2^\lambda || A$. Recall $B = -d_1 T$ and $7T + 1 = Al$. We have

$$\begin{aligned} & -84A(s(d_1, 7A) - s(B, A)) \\ & \equiv -d_1 - \overline{d_{1\{8 \times 2^\lambda\}}}(49A^2 + 21A + 1) - 14\overline{d_{1\{8 \times 2^\lambda\}}A}\left(\frac{7A}{d_1}\right) \\ & \quad + d_1(1 - Al) + 49\overline{(d_1 - d_1 Al)_{\{8 \times 2^\lambda\}}}(A^2 + 3A + 1) + 14\overline{B_{\{8 \times 2^\lambda\}}A}\left(\frac{A}{B}\right) \\ & \equiv -d_1 Al + 49A^2 \cdot \overline{d_1 Al(d_1 - d_1 Al)_{\{8 \times 2^\lambda\}}d_{1\{8 \times 2^\lambda\}}} \\ & \quad + 21A(6d_1 + d_1 Al)\overline{(d_1 - d_1 Al)_{\{8 \times 2^\lambda\}}d_{1\{8 \times 2^\lambda\}}} \\ & \quad + (48d_1 + d_1 Al)\overline{(d_1 - d_1 Al)_{\{8 \times 2^\lambda\}}d_{1\{8 \times 2^\lambda\}}} \\ & \quad + 14A \left(\overline{B_{\{8 \times 2^\lambda\}}}\left(\frac{A}{B}\right) - \overline{d_{1\{8 \times 2^\lambda\}}}\left(\frac{7A}{d_1}\right) \right) \pmod{8 \times 2^\lambda}. \end{aligned}$$

Since $2^\lambda || A$ with $\lambda \geq 1$, we apply (5.22) and $x^2 \equiv 1 \pmod{8}$ for odd x to get

$$\begin{aligned} -84A(s(d_1, 7A) - s(B, A)) & \equiv 6d_1 A + d_1 A^2 \ell(1 + \ell) + 48\overline{d_{1\{A\}}} \\ & \quad + 6A \left(B\left(\frac{A}{B}\right) - d_1\left(\frac{7A}{d_1}\right) \right) \pmod{8 \times 2^\lambda}. \end{aligned}$$

By (5.23), To determine $B\left(\frac{A}{B}\right) - d_1\left(\frac{7A}{d_1}\right) \pmod{4}$, we use the quadratic reciprocity (2.6). By $B < 0$ odd and $A > 0$, we compute

$$\begin{aligned} B\left(\frac{A}{B}\right) - d_1\left(\frac{7A}{d_1}\right) & \equiv -d_1 T\left(\frac{B}{A}\right)(-1)^{\frac{A-1}{2} \cdot \frac{B-1}{2}} - d_1\left(\frac{d_1}{7A}\right)(-1)^{\frac{7 \cdot \frac{A-1}{2} - 1}{2} \cdot \frac{d_1-1}{2}} \\ & \equiv -d_1 T\left(\frac{d_1 - d_1 Al}{A}\right)\left(\frac{7}{A}\right)(-1)^{\frac{A-1}{2} \cdot \frac{B-1}{2}} - d_1\left(\frac{d_1}{A}\right)(-1)^{\frac{7 \cdot \frac{A-1}{2} - 1}{2} \cdot \frac{d_1-1}{2}} \pmod{4} \end{aligned} \quad (5.26)$$

Here are the cases:

- (1) If $4|A$, then we have $T \equiv 1 \pmod{4}$, $B \equiv -d_1 \pmod{4}$. Moreover, $\left(\frac{d_1 - d_1 Al}{A}\right) = \left(\frac{d_1}{A}\right)$ always (note that A is even and we have to consider $\left(\frac{d_1}{2}\right)$). Now (5.26) simplifies to $\left(\frac{\ell}{7}\right)d_1 + 1 \pmod{4}$. In this case $d_1 A^2 \ell(1 + \ell) \equiv 0 \pmod{8 \times 2^\lambda}$ and we conclude

$$-84A(s(d_1, 7A) - s(B, A)) \equiv \begin{cases} 2A + 48\overline{d_{1\{A\}}} \pmod{8 \times 2^\lambda}, & \ell = 1, 2; \\ 6A + 48\overline{d_{1\{A\}}} \pmod{8 \times 2^\lambda}, & \ell = 3. \end{cases} \quad (5.27)$$

- (2) If $2 \parallel A$ and $\ell = 1$, then $T \equiv 3 \pmod{4}$, $B \equiv d_1 \pmod{4}$ and the above (5.26) simplifies to $d_1 - 1 \pmod{4}$. Then as $A(12d_1 - 6 + 2d_1A) \equiv 2A \pmod{8 \times 2^\lambda}$, we conclude the same as the first line of (5.27).
- (3) If $2 \parallel A$ and $\ell = 2$, then $T \equiv 1 \pmod{4}$, $B \equiv -d_1 \pmod{4}$, and $(\frac{d_1 - d_1 A \ell}{A}) = -(\frac{d_1}{A})$. Now (5.26) gives $d_1 - 1 \pmod{4}$ and we again get the first line of (5.27).
- (4) If $2 \parallel A$ and $\ell = 3$, then $T \equiv 3 \pmod{4}$, $B \equiv d_1 \pmod{4}$, and $(\frac{A}{7}) = (\frac{3A}{7})(\frac{3}{7}) = -1$. Here (5.26) results in $d_1 - 1 \pmod{4}$ again. Note that $d_1 A^2 \ell (1 + \ell) \equiv 0 \pmod{8 \times 2^\lambda}$ and we get the second line of (5.27).

Next we check the part of Q_2 other than $e^{-\pi i s(d,c)}$. In this case A is even, so $3T + 1$ is even and we have

$$\left(\frac{3}{2}T^2 + \frac{1}{2}T\right) C \equiv \frac{3T+1}{2} \cdot T(-7\overline{d_{1\{A\}}}) \equiv \frac{3T+1}{2} \overline{d_{1\{A\}}} \pmod{A}.$$

When written with denominator $24 \cdot 7A$, we have

$$e\left(\frac{\left(\frac{3}{2}T^2 + \frac{1}{2}T\right)C}{A}\right) = e\left(\frac{36A\ell\overline{d_{1\{A\}}} + 48\overline{d_{1\{A\}}}}{24 \cdot 7A}\right)$$

whose numerator is

$$36A\ell\overline{d_{1\{A\}}} + 48\overline{d_{1\{A\}}} \equiv \begin{cases} 0 \pmod{7}, \\ 48\overline{d_{1\{A\}}} \pmod{3A}, \\ 4A + 48\overline{d_{1\{A\}}} \pmod{8 \times 2^\lambda}, & \ell = 1, 3, \\ 48\overline{d_{1\{A\}}} \pmod{8 \times 2^\lambda}, & \ell = 2. \end{cases} \quad (5.28)$$

Combining the above computation with (5.13), (5.27) and (2.2), we get

$$\text{Arg}(Q_2 \rightarrow P_2; \ell) = \frac{9, 11, 27}{28} \quad \text{for } \ell = 1, 2, 3. \quad (5.29)$$

Then we compute $\text{Arg}(Q_1 \rightarrow P_1; \ell)$. When $\ell = 1$, since A is even, $\frac{A}{2} \equiv 4 \pmod{7}$. Note that $a_1 \equiv 1 \pmod{14}$ because a_1 is odd. It is direct to get (remember $Q_1 = i$)

$$\text{Arg}(Q_1 \rightarrow P_1; 1) = \frac{1}{2} - \frac{5}{7} - \frac{1}{4} = -\frac{13}{28}. \quad (5.30)$$

When $\ell = 2$, we get $\frac{A}{2} \equiv 2 \pmod{7}$ and

$$\text{Arg}(Q_1 \rightarrow P_1; 2) = \frac{1}{2} - \frac{3}{7} - \frac{1}{4} = -\frac{5}{28}. \quad (5.31)$$

When $\ell = 3$, we have $\frac{A}{2} \equiv 6 \pmod{14}$ and

$$\text{Arg}(Q_1 \rightarrow P_1; 3) = \frac{1}{2} - \frac{1}{7} - \frac{1}{4} = \frac{3}{28}. \quad (5.32)$$

Combining (5.30), (5.31), (5.32), (5.29), and (5.12), we get

$$\text{Arg}(Q \rightarrow P; \ell) = -\frac{3}{7}, -\frac{5}{14}, \frac{3}{14} \quad \text{for } \ell = 1, 2, 3. \quad (5.33)$$

This proves (5.9).

5.4. **Case $c'\ell \equiv 1 \pmod{7}$, $2|A$, and $3|A$.** Comparing to the former case, the only difference in getting $\text{Arg}(Q_2 \rightarrow P_2; \ell)$ in (5.29) is that we should using (5.21) instead of (5.13). The result (5.29) still holds in this case. The condition $3|A$ or $3 \nmid A$ does not affect the computation for $\text{Arg}(Q_1 \rightarrow P_1; \ell)$ and $\text{Arg}(Q_3 \rightarrow P_3; \ell)$, hence we still have (5.9):

$$\text{Arg}(Q \rightarrow P; \ell) = -\frac{3}{7}, -\frac{5}{14}, \frac{3}{14} \quad \text{for } \ell = 1, 2, 3. \quad (5.34)$$

Now we have finished the proof in all the cases for A when $A\ell \equiv 1 \pmod{7}$ and proved (5.9) in Proposition 5.1. For the other case $A\ell \equiv -1 \pmod{7}$, we will not repeat the same process but just list the key argument differences below. For every $r \pmod{c'}^*$, we compare $P(d_1)$ (5.5) given $d_1 \in V(r, c)$ and $Q(B)$ (5.7) given

$$T := \frac{A\ell + 1}{7} > 0, \quad B = d_1 T \quad \text{and} \quad C = -7\overline{d_{1\{A\}}}.$$

Now $7B = d_1 + d_1 A\ell$. We shall get Table 5.1.

Case $2 \nmid A$:	$\ell = 1$	$\ell = 2$	$\ell = 3$
$\text{Arg}(Q_1 \rightarrow P_1; \ell)$	$-\frac{1}{28}$	$\frac{5}{28}$	$\frac{11}{28}$
$\text{Arg}(Q_2 \rightarrow P_2; \ell)$	$\frac{5}{28}$	$-\frac{11}{28}$	$-\frac{13}{28}$
$\text{Arg}(Q_3 \rightarrow P_3; \ell)$	$\frac{2}{7}$	$-\frac{3}{7}$	$-\frac{1}{7}$
$\text{Arg}(Q \rightarrow P; \ell)$	$\frac{3}{7}$	$\frac{5}{14}$	$-\frac{3}{14}$
Case $2 A$:	$\ell = 1$	$\ell = 2$	$\ell = 3$
$\text{Arg}(Q_1 \rightarrow P_1; \ell)$	$\frac{13}{28}$	$\frac{5}{28}$	$-\frac{3}{28}$
$\text{Arg}(Q_2 \rightarrow P_2; \ell)$	$-\frac{9}{28}$	$-\frac{11}{28}$	$\frac{1}{28}$
$\text{Arg}(Q_3 \rightarrow P_3; \ell)$	$\frac{2}{7}$	$-\frac{3}{7}$	$-\frac{1}{7}$
$\text{Arg}(Q \rightarrow P; \ell)$	$\frac{3}{7}$	$\frac{5}{14}$	$-\frac{3}{14}$

TABLE 5.1. Table for the case $A\ell \equiv -1 \pmod{7}$

We have finished the proof of (7-5,2) of Theorem 1.3.

6. PART (II) OF THEOREM 1.3

For prime $p = 5, 7$ and integers a, b , recall the notation

$$C_p^{a,b} = \cos\left(\frac{a\pi}{p}\right) - \cos\left(\frac{b\pi}{p}\right).$$

6.1. **(5-1) and (5-2) of Theorem 1.3.** We still denote $c' = c/5$ and first deal with the case $25|c$. For $(r, c) = 1$, recall (3.36) with $V(r, c)$, d and d_* in that subsection. By (3.36), we have $\text{Arg}(d \rightarrow d_*; \ell) = -\frac{1}{5}$ for the $5n + 4$ case. Since we have the $5n + 1$ case here, we obtain

$$\text{Arg}(d \rightarrow d_*; \ell) = -\frac{1}{5} - \frac{3c'}{c} = -\frac{4}{5}.$$

Hence we get $S_{\infty\infty}^{(\ell)}(0, 5n + 1, c, \mu_5) = 0$ for $\ell \in \{1, 2\}$, $25|c$, and every $n \geq 0$. This proves (5-1) when $25|c$.

Similarly, in the $5n + 2$ case, we obtain

$$\text{Arg}(d \rightarrow d_*; \ell) = -\frac{1}{5} - \frac{2c'}{c} = -\frac{3}{5}.$$

We still get $S_{\infty}^{(\ell)}(0, 5n + 2, c, \mu_5) = 0$ for $\ell \in \{1, 2\}$, $25|c$, and every $n \geq 0$, which proves (5-2) when $25|c$.

Now we focus on the case $5||c$. For $(r, c) = 1$, recall the notation of $V(r, c)$, d_j and a_j in (3.3). By (3.5) and Condition 3.2, we find that the argument differences of $P(d)$ for $d \in V(r, c)$ only depends on $c' \pmod{5}$. Moreover, in (3.1), if we change $5n + 4$ to $5n + 1$ or to $5n + 2$, then only the argument of $P_3(d)$ is affected. Recall that we define $\beta \in \{1, 2, 3, 4\}$ by $\beta c' \equiv 1 \pmod{5}$.

6.1.1. *The $5n + 1$ case.* As Proposition 3.1, we denote

$$s_{r,c}^{(\ell)} = \sum_{d \in V(r,c)} P_1(d)P_2(d)P_3(d) := \sum_{d \in V(r,c)} \frac{e\left(-\frac{3c'al^2}{10}\right)}{\sin\left(\frac{\pi al}{5}\right)} e^{-\pi is(d,c)} e\left(\frac{d}{c}\right). \quad (6.1)$$

Here $P_3(d) = e\left(\frac{d}{c}\right)$ instead of $e\left(\frac{4d}{c}\right)$ in the $5n + 4$ case Proposition 3.1. To prove (5-1) of Theorem 1.3, it suffices to show

$$C_5^{2,4} \sin\left(\frac{\pi}{5}\right) s_{r,c}^{(1)} + C_5^{4,2} \sin\left(\frac{2\pi}{5}\right) s_{r,c}^{(2)} = 0. \quad (6.2)$$

When we compute $\text{Arg}_3(d_j \rightarrow d_{j+1}; \ell)$ for $j = 1, 2, 3$, previously it was $e\left(\frac{4\beta}{5}\right)$ and now it should be $e\left(\frac{\beta}{5}\right)$. Therefore, we need to subtract $\frac{3\beta}{5}$ from the argument differences in Condition 3.2 in each case.

Since we need both $\ell = 1$ and $\ell = 2$ appears at the same time, we write our new condition in the following way. It is important to note that the way we compute $\text{Arg}(d_4 \rightarrow d_1; \ell)$ is by

$$\sum_{j=1}^3 \text{Arg}(d_j \rightarrow d_{j+1}; \ell) + \text{Arg}(d_4 \rightarrow d_1; \ell) = 0$$

but not by subtracting $\frac{3\beta}{5}$.

Condition 6.1. *For the $5n + 1$ case, we have the following styles of argument differences:*

- $c' \equiv 1 \pmod{5}$, $\beta = 1$;

$c' \equiv 1 \pmod{5}$	$d_1 \rightarrow d_2$	$d_2 \rightarrow d_3$	$d_3 \rightarrow d_4$	$d_4 \rightarrow d_1$
$\text{Arg}(d_u \rightarrow d_v; 1)$	$-\frac{3}{10}$	$\frac{1}{2}$	$-\frac{3}{10}$	$\frac{1}{10}$
$\text{Arg}(d_u \rightarrow d_v; 2)$	$\frac{2}{5}$	$-\frac{1}{10}$	$\frac{2}{5}$	$\frac{3}{10}$

- $c' \equiv 2 \pmod{5}$, $\beta = 3$;

$c' \equiv 2 \pmod{5}$	$d_1 \rightarrow d_2$	$d_2 \rightarrow d_3$	$d_3 \rightarrow d_4$	$d_4 \rightarrow d_1$
$\text{Arg}(d_u \rightarrow d_v; 1)$	$-\frac{3}{10}$	$-\frac{3}{10}$	$-\frac{3}{10}$	$-\frac{1}{10}$
$\text{Arg}(d_u \rightarrow d_v; 2)$	$-\frac{2}{5}$	$\frac{1}{2}$	$-\frac{2}{5}$	$\frac{3}{10}$


- $c' \equiv 3 \pmod{5}$, $\beta = 2$;

$c' \equiv 3 \pmod{5}$	$d_1 \rightarrow d_2$	$d_2 \rightarrow d_3$	$d_3 \rightarrow d_4$	$d_4 \rightarrow d_1$
$\text{Arg}(d_u \rightarrow d_v; 1)$	$\frac{3}{10}$	$\frac{3}{10}$	$\frac{3}{10}$	$\frac{1}{10}$
$\text{Arg}(d_u \rightarrow d_v; 2)$	$\frac{2}{5}$	$\frac{1}{2}$	$\frac{2}{5}$	$-\frac{3}{10}$

$c' \equiv 4 \pmod{5}$	$d_1 \rightarrow d_2$	$d_2 \rightarrow d_3$	$d_3 \rightarrow d_4$	$d_4 \rightarrow d_1$
$\text{Arg}(d_u \rightarrow d_v; 1)$	$\frac{3}{10}$	$\frac{1}{2}$	$\frac{3}{10}$	$-\frac{1}{10}$
$\text{Arg}(d_u \rightarrow d_v; 2)$	$-\frac{2}{5}$	$\frac{1}{10}$	$-\frac{2}{5}$	$-\frac{3}{10}$

- $c' \equiv 4 \pmod{5}$, $\beta = 4$.

The condition above corresponding to the following styles of $P(d)$ for $d \in V(r, c)$:



For every two graphs close to each other in a row which satisfy the argument differences in the corresponding cases in Condition 6.1, it proves (6.2) due to the following equations:

$$C_5^{2,4} \cos\left(\frac{\pi}{10}\right) + C_5^{4,2} \sin\left(\frac{2\pi}{5}\right) \left(\frac{\cos\left(\frac{\pi}{10}\right)}{\sin\left(\frac{\pi}{5}\right)} - \frac{\cos\left(\frac{3\pi}{10}\right)}{\sin\left(\frac{2\pi}{5}\right)} \right) = 0, \quad \text{for } c' \equiv 1, 4 \pmod{5};$$

$$C_5^{2,4} \sin\left(\frac{\pi}{5}\right) \left(\frac{\cos\left(\frac{\pi}{10}\right)}{\sin\left(\frac{\pi}{5}\right)} - \frac{\cos\left(\frac{3\pi}{10}\right)}{\sin\left(\frac{2\pi}{5}\right)} \right) + C_5^{4,2} \cos\left(\frac{3\pi}{10}\right) = 0, \quad \text{for } c' \equiv 2, 3 \pmod{5}.$$

This proves (5-1) of Theorem 1.3.

6.1.2. *The $5n + 2$ case.* As (6.1), we denote $P_3(d) = e\left(\frac{2d}{c}\right)$ instead of $e\left(\frac{4d}{c}\right)$ in the $5n + 4$ case Proposition 3.1 and instead of $e\left(\frac{d}{c}\right)$ in the $5n + 1$ case (6.1). To prove (5-2) of Theorem 1.3, it suffices to show

$$C_5^{0,4} \sin\left(\frac{\pi}{5}\right) s_{r,c}^{(1)} + C_5^{0,2} \sin\left(\frac{2\pi}{5}\right) s_{r,c}^{(2)} = 0. \quad (6.3)$$

When we compute $\text{Arg}_3(d_j \rightarrow d_{j+1}; \ell)$ for $j = 1, 2, 3$, in (6.1) it was $e\left(\frac{\beta}{5}\right)$ and now it should be $e\left(\frac{2\beta}{5}\right)$. Therefore, we need to add $\frac{\beta}{5}$ to the argument differences in Condition 6.1 in each case to get the following condition.

Condition 6.2. *For the $5n + 2$ case, we have the following styles of argument differences:*

- $c' \equiv 1 \pmod{5}$, $\beta = 1$;
- $c' \equiv 2 \pmod{5}$, $\beta = 3$;
- $c' \equiv 3 \pmod{5}$, $\beta = 2$;
- $c' \equiv 4 \pmod{5}$, $\beta = 4$.


$c' \equiv 1 \pmod{5}$	$d_1 \rightarrow d_2 \rightarrow d_3 \rightarrow d_4 \rightarrow d_1$
$\text{Arg}(d_u \rightarrow d_v; 1)$	$-\frac{1}{10} \quad -\frac{3}{10} \quad -\frac{1}{10} \quad \frac{1}{2}$
$\text{Arg}(d_u \rightarrow d_v; 2)$	$-\frac{2}{5} \quad \frac{1}{10} \quad -\frac{2}{5} \quad -\frac{3}{10}$

$c' \equiv 2 \pmod{5}$	$d_1 \rightarrow d_2 \rightarrow d_3 \rightarrow d_4 \rightarrow d_1$
$\text{Arg}(d_u \rightarrow d_v; 1)$	$\frac{3}{10} \quad \frac{3}{10} \quad \frac{3}{10} \quad \frac{1}{10}$
$\text{Arg}(d_u \rightarrow d_v; 2)$	$\frac{1}{5} \quad \frac{1}{10} \quad \frac{1}{5} \quad \frac{1}{2}$

$c' \equiv 3 \pmod{5}$	$d_1 \rightarrow d_2 \rightarrow d_3 \rightarrow d_4 \rightarrow d_1$
$\text{Arg}(d_u \rightarrow d_v; 1)$	$-\frac{3}{10} \quad -\frac{3}{10} \quad -\frac{3}{10} \quad -\frac{1}{10}$
$\text{Arg}(d_u \rightarrow d_v; 2)$	$-\frac{1}{5} \quad -\frac{1}{10} \quad -\frac{1}{5} \quad \frac{1}{2}$

$c' \equiv 4 \pmod{5}$	$d_1 \rightarrow d_2 \rightarrow d_3 \rightarrow d_4 \rightarrow d_1$
$\text{Arg}(d_u \rightarrow d_v; 1)$	$\frac{1}{10} \quad \frac{3}{10} \quad \frac{1}{10} \quad \frac{1}{2}$
$\text{Arg}(d_u \rightarrow d_v; 2)$	$\frac{2}{5} \quad -\frac{1}{10} \quad \frac{2}{5} \quad \frac{3}{10}$

The condition above corresponding to the following styles of $P(d)$ for $d \in V(r, c)$.



For every two graphs with same c which satisfy the argument differences in the corresponding cases in Condition 6.2, it proves (6.3) due to the following equations:

$$C_5^{0,4} \sin\left(\frac{\pi}{5}\right) \cdot \frac{\cos\left(\frac{3\pi}{10}\right)}{\sin\left(\frac{2\pi}{5}\right)} + C_5^{0,2} \sin\left(\frac{2\pi}{5}\right) \left(\frac{\cos\left(\frac{3\pi}{10}\right)}{\sin\left(\frac{2\pi}{5}\right)} - \frac{\cos\left(\frac{\pi}{10}\right)}{\sin\left(\frac{\pi}{5}\right)} \right) = 0, \quad \text{for } c' \equiv 1, 4 \pmod{5};$$

$$C_5^{0,4} \sin\left(\frac{\pi}{5}\right) \left(\frac{\cos\left(\frac{\pi}{10}\right)}{\sin\left(\frac{\pi}{5}\right)} - \frac{\cos\left(\frac{3\pi}{10}\right)}{\sin\left(\frac{2\pi}{5}\right)} \right) - C_5^{0,2} \sin\left(\frac{2\pi}{5}\right) \cdot \frac{\cos\left(\frac{\pi}{10}\right)}{\sin\left(\frac{\pi}{5}\right)} = 0, \quad \text{for } c' \equiv 2, 3 \pmod{5}.$$

This proves (5-2) of Theorem 1.3.

6.2. Restate the condition for (7-5) of Theorem 1.3. We still denote $c = 7A = 7c'$. When $49|c$, recall the notation in §4.2 and we have (4.17) for any $d \in V(r, c)$ and $d_* = d + c'$:

$$\text{Arg}(d \rightarrow d_*; \ell) = \begin{cases} -\frac{2}{7} & d \equiv 1, 6 \pmod{7}; \\ \frac{3}{7} & d \equiv 2, 5 \pmod{7}; \\ -\frac{1}{7} & d \equiv 3, 4 \pmod{7}. \end{cases} \quad (6.4)$$

When $7||c'$, denote $A = c' = c/7$ and recall the notation of $V(r, c)$ before (4.1) and d_j and a_j in (4.4). We combine Condition 4.2, Condition 5.2, (5.9) and (5.10) and get the following condition:

Condition 6.3. For the $7n + 5$ case, we have the following conditions on $\text{Arg}(Q \rightarrow P; \ell)$ when $A\ell \equiv \pm 1 \pmod{7}$ with tables for $\text{Arg}_j(d_u \rightarrow d_v; \ell)$.

- $c' \equiv 1 \pmod{7}$. $A \cdot 1 = 7T + 1$, $\text{Arg}(Q \rightarrow P; 1) = -\frac{3}{7}$;

$c' \equiv 1 \pmod{7}$	d_1	\rightarrow	d_2	\rightarrow	d_3	\rightarrow	d_4	\rightarrow	d_5	\rightarrow	d_6	\rightarrow	d_1
$\text{Arg}(d_u \rightarrow d_v; 1)$			$\frac{3}{14}$		$-\frac{3}{7}$		$\frac{2}{7}$		$-\frac{3}{7}$		$\frac{3}{14}$		$\frac{1}{7}$
$\text{Arg}(d_u \rightarrow d_v; 2)$			$-\frac{3}{14}$		$-\frac{1}{14}$		$-\frac{2}{7}$		$-\frac{1}{14}$		$-\frac{3}{14}$		$-\frac{1}{7}$
$\text{Arg}(d_u \rightarrow d_v; 3)$			$-\frac{3}{7}$		$\frac{5}{14}$		$\frac{3}{7}$		$\frac{5}{14}$		$-\frac{3}{7}$		$-\frac{2}{7}$

- $c' \equiv 2 \pmod{7}$. $A \cdot 3 = 7T - 1$, $\text{Arg}(Q \rightarrow P; 3) = -\frac{3}{14}$;

$c' \equiv 2 \pmod{7}$	d_1	\rightarrow	d_2	\rightarrow	d_3	\rightarrow	d_4	\rightarrow	d_5	\rightarrow	d_6	\rightarrow	d_1
$\text{Arg}(d_u \rightarrow d_v; 1)$			$-\frac{5}{14}$		$-\frac{2}{7}$		$-\frac{1}{7}$		$-\frac{2}{7}$		$-\frac{5}{14}$		$\frac{3}{7}$
$\text{Arg}(d_u \rightarrow d_v; 2)$			$-\frac{3}{14}$		$-\frac{1}{14}$		$-\frac{2}{7}$		$-\frac{1}{14}$		$-\frac{3}{14}$		$-\frac{1}{7}$
$\text{Arg}(d_u \rightarrow d_v; 3)$			$-\frac{1}{7}$		$-\frac{3}{14}$		$\frac{1}{7}$		$-\frac{3}{14}$		$-\frac{1}{7}$		$-\frac{3}{7}$

- $c' \equiv 3 \pmod{7}$. $A \cdot 2 = 7T - 1$, $\text{Arg}(Q \rightarrow P; 2) = \frac{5}{14}$;

$c' \equiv 3 \pmod{7}$	d_1	\rightarrow	d_2	\rightarrow	d_3	\rightarrow	d_4	\rightarrow	d_5	\rightarrow	d_6	\rightarrow	d_1
$\text{Arg}(d_u \rightarrow d_v; 1)$			$\frac{5}{14}$		$\frac{2}{7}$		$\frac{1}{7}$		$\frac{2}{7}$		$\frac{5}{14}$		$-\frac{3}{7}$
$\text{Arg}(d_u \rightarrow d_v; 2)$			$\frac{1}{14}$		$\frac{5}{14}$		$\frac{3}{7}$		$\frac{5}{14}$		$\frac{1}{14}$		$-\frac{2}{7}$
$\text{Arg}(d_u \rightarrow d_v; 3)$			$\frac{3}{7}$		$-\frac{5}{14}$		$-\frac{3}{7}$		$-\frac{5}{14}$		$\frac{3}{7}$		$\frac{2}{7}$

- $c' \equiv 4 \pmod{7}$. $A \cdot 2 = 7T + 1$, $\text{Arg}(Q \rightarrow P; 2) = -\frac{5}{14}$;

$c' \equiv 4 \pmod{7}$	d_1	\rightarrow	d_2	\rightarrow	d_3	\rightarrow	d_4	\rightarrow	d_5	\rightarrow	d_6	\rightarrow	d_1
$\text{Arg}(d_u \rightarrow d_v; 1)$			$-\frac{5}{14}$		$-\frac{2}{7}$		$-\frac{1}{7}$		$-\frac{2}{7}$		$-\frac{5}{14}$		$-\frac{3}{7}$
$\text{Arg}(d_u \rightarrow d_v; 2)$			$-\frac{1}{14}$		$-\frac{5}{14}$		$-\frac{3}{7}$		$-\frac{5}{14}$		$-\frac{1}{14}$		$\frac{2}{7}$
$\text{Arg}(d_u \rightarrow d_v; 3)$			$-\frac{3}{7}$		$\frac{5}{14}$		$\frac{3}{7}$		$\frac{5}{14}$		$-\frac{3}{7}$		$-\frac{2}{7}$

- $c' \equiv 5 \pmod{7}$. $A \cdot 3 = 7T + 1$, $\text{Arg}(Q \rightarrow P; 3) = \frac{3}{14}$;
- $c' \equiv 6 \pmod{7}$. $A \cdot 1 = 7T - 1$, $\text{Arg}(Q \rightarrow P; 1) = \frac{3}{7}$.

We only prove (7-0) of Theorem 1.3 and omit the proof of the other (7- k) cases because the proofs are essentially the same.

$c' \equiv 5 \pmod{7}$	$d_1 \rightarrow d_2 \rightarrow d_3 \rightarrow d_4 \rightarrow d_5 \rightarrow d_6 \rightarrow d_1$
$\text{Arg}(d_u \rightarrow d_v; 1)$	$\frac{5}{14} \quad \frac{2}{7} \quad \frac{1}{7} \quad \frac{2}{7} \quad \frac{5}{14} \quad -\frac{3}{7}$
$\text{Arg}(d_u \rightarrow d_v; 2)$	$\frac{3}{14} \quad \frac{1}{14} \quad \frac{2}{7} \quad \frac{1}{14} \quad \frac{3}{14} \quad \frac{1}{7}$
$\text{Arg}(d_u \rightarrow d_v; 3)$	$\frac{1}{7} \quad \frac{3}{14} \quad -\frac{1}{7} \quad \frac{3}{14} \quad \frac{1}{7} \quad \frac{3}{7}$

$c' \equiv 6 \pmod{7}$	$d_1 \rightarrow d_2 \rightarrow d_3 \rightarrow d_4 \rightarrow d_5 \rightarrow d_6 \rightarrow d_1$
$\text{Arg}(d_u \rightarrow d_v; 1)$	$-\frac{3}{14} \quad \frac{3}{7} \quad -\frac{2}{7} \quad \frac{3}{7} \quad -\frac{3}{14} \quad -\frac{1}{7}$
$\text{Arg}(d_u \rightarrow d_v; 2)$	$\frac{3}{14} \quad \frac{1}{14} \quad \frac{2}{7} \quad \frac{1}{14} \quad \frac{3}{14} \quad \frac{1}{7}$
$\text{Arg}(d_u \rightarrow d_v; 3)$	$\frac{3}{7} \quad -\frac{5}{14} \quad -\frac{3}{7} \quad -\frac{5}{14} \quad \frac{3}{7} \quad \frac{2}{7}$

6.3. (7-0) of Theorem 1.3. As (4.1), we still denote $V(r, c) = \{d \pmod{c}^* : d \equiv r \pmod{c}\}$, $d_j \in V(r, c)$ by $d_j \equiv j \pmod{7}$. Recall the Kloosterman sums defined at (2.7), (2.8) and (2.9).

For $A := c' = c/7$, when $A\ell = 7T + 1$ for some integer $T \geq 0$, as (5.6) we define

$$Q(B) := 2\sqrt{7}Q_1(B)Q_2(B)Q_3(B) \quad (6.5)$$

with $Q_1(B) := (-1)^{[A\ell]}i$ and

$$Q_2(B) := e\left(\frac{(\frac{3}{2}T^2 + \frac{1}{2}T)C}{A}\right), \quad \text{and} \quad Q_3(B) = e\left(\frac{0 \cdot B}{A}\right) = 1 \quad (6.6)$$

and let $B = -d_1T$ with $C = -7\overline{d_1\{A\}}$. When $A\ell = 7T - 1$ for some $T \geq 0$, we still define $Q(B)$ as above while we take

$$Q_2(B) := e\left(\frac{(\frac{3}{2}(T-1)^2 + \frac{5}{2}T + 1)C}{A}\right), \quad \text{and} \quad B = d_1T. \quad (6.7)$$

instead. Note that when A is fixed, ℓ is also fixed, i.e. there is only one corresponding $Q(B)$ for every fixed c .

We define the sum on $V(r, c)$ as

$$s_{r,c}^{(\ell)} := \sin\left(\frac{\pi\ell}{7}\right) \sum_{d \in V(r,c)} P_1(d)P_2(d)P_3(d) + \sin\left(\frac{\pi\ell}{7}\right) \mathbf{1}_{A:=c/7} \mathbf{1}_{[A\ell]=1,6} Q(B), \quad \text{where} \quad (6.8)$$

$$P_1(d) := \frac{(-1)^{\ell c} e\left(-\frac{3c'ad^2}{14}\right)}{\sin\left(\frac{\pi a\ell}{7}\right)}, \quad P_2(d) := e\left(-\frac{12cs(d,c)}{24c}\right), \quad P_3(d) := e\left(\frac{0 \cdot d}{c}\right) = 1.$$

Here $\mathbf{1}_{\text{condition}}$ equals 1 if the condition meets and equals 0 otherwise.

To prove (7-0) of Theorem 1.3, it suffices to show

$$C_7^{4,6} s_{r,c}^{(1)} + C_7^{6,2} s_{r,c}^{(2)} + C_7^{2,4} s_{r,c}^{(3)} = 0. \quad (6.9)$$

First we deal with the case $49|c$ and there is no $Q(B)$. We need to subtract $\frac{5}{7}$ from (6.4) and get

$$\text{Arg}(d \rightarrow d_*; \ell) = \begin{cases} 0 & d \equiv 1, 6 \pmod{7}; \\ -\frac{2}{7} & d \equiv 2, 5 \pmod{7}; \\ \frac{1}{7} & d \equiv 3, 4 \pmod{7}. \end{cases}$$

When $r \equiv d \equiv 2, 3, 4, 5 \pmod{7}$, we get equi-distribution and (6.9) follows. When $r \equiv d \equiv 1, 6 \pmod{7}$, note that $P_1(d) = (-1)^{(a+1)c\ell} / \sin(\frac{\pi a\ell}{7})$ for $ad \equiv 1 \pmod{c}$ has the same sgn $P_1(d)$ for $\ell = 1, 2, 3$. Hence every summand for $d \in V(r, c)$ in (6.9) has the same argument and we get (6.9) by

$$C_7^{4,6} \frac{\sin(\frac{\pi}{7})}{\sin(\frac{\pi}{7})} + C_7^{6,2} \frac{\sin(\frac{2\pi}{7})}{\sin(\frac{2\pi}{7})} + C_7^{2,4} \frac{\sin(\frac{3\pi}{7})}{\sin(\frac{3\pi}{7})} = 0.$$

Next we check the condition for $7|c$. Comparing with Condition 6.3, since we have different $P_3(d)$ and $Q_3(B)$ in this case, we need to subtract $\frac{5\beta}{7}$ in $\text{Arg}(d_j \rightarrow d_{j+1}; \ell)$, $1 \leq j \leq 5$ from Condition 6.3. We also need to add $\mp \frac{5\ell}{7}$ to $\text{Arg}(Q \rightarrow P; \ell)$ when $Al \equiv \pm 1 \pmod{7}$. It is important to note that we compute $\text{Arg}(d_6 \rightarrow d_1; \ell)$ by

$$\sum_{j=1}^5 \text{Arg}(d_j \rightarrow d_{j+1}; \ell) + \text{Arg}(d_6 \rightarrow d_1; \ell) = 0$$

instead of adding $\mp \frac{5\ell}{7}$.

Condition 6.4. For the $7n$ case, we have the following conditions on $\text{Arg}(Q \rightarrow P; \ell)$ when $Al \equiv \pm 1 \pmod{7}$ with tables for $\text{Arg}_j(d_u \rightarrow d_v; \ell)$.

- $c' \equiv 1 \pmod{7}$, $\beta = 1$. $A \cdot 1 = 7T + 1$, $\text{Arg}(Q \rightarrow P; 1) = -\frac{1}{7}$;

$c' \equiv 1 \pmod{7}$	$d_1 \rightarrow d_2$	$d_2 \rightarrow d_3$	$d_3 \rightarrow d_4$	$d_4 \rightarrow d_5$	$d_5 \rightarrow d_6$	$d_6 \rightarrow d_1$
$\text{Arg}(d_u \rightarrow d_v; 1)$	$\frac{1}{2}$	$-\frac{1}{7}$	$-\frac{3}{7}$	$-\frac{1}{7}$	$\frac{1}{2}$	$-\frac{2}{7}$
$\text{Arg}(d_u \rightarrow d_v; 2)$	$\frac{1}{14}$	$\frac{3}{14}$	0	$\frac{3}{14}$	$\frac{1}{14}$	$\frac{3}{7}$
$\text{Arg}(d_u \rightarrow d_v; 3)$	$-\frac{1}{7}$	$-\frac{5}{14}$	$-\frac{2}{7}$	$-\frac{5}{14}$	$-\frac{1}{7}$	$\frac{2}{7}$

- $c' \equiv 2 \pmod{7}$, $\beta = 4$. $A \cdot 3 = 7T - 1$, $\text{Arg}(Q \rightarrow P; 3) = -\frac{1}{14}$;

$c' \equiv 2 \pmod{7}$	$d_1 \rightarrow d_2$	$d_2 \rightarrow d_3$	$d_3 \rightarrow d_4$	$d_4 \rightarrow d_5$	$d_5 \rightarrow d_6$	$d_6 \rightarrow d_1$
$\text{Arg}(d_u \rightarrow d_v; 1)$	$-\frac{3}{14}$	$-\frac{1}{7}$	0	$-\frac{1}{7}$	$-\frac{3}{14}$	$-\frac{2}{7}$
$\text{Arg}(d_u \rightarrow d_v; 2)$	$-\frac{1}{14}$	$\frac{1}{14}$	$-\frac{1}{7}$	$\frac{1}{14}$	$-\frac{1}{14}$	$\frac{1}{7}$
$\text{Arg}(d_u \rightarrow d_v; 3)$	0	$-\frac{1}{14}$	$\frac{2}{7}$	$-\frac{1}{14}$	0	$-\frac{1}{7}$

- $c' \equiv 3 \pmod{7}$. $A \cdot 2 = 7T - 1$, $\text{Arg}(Q \rightarrow P; 2) = -\frac{3}{14}$;

$c' \equiv 3 \pmod{7}$	$d_1 \rightarrow d_2$	$d_2 \rightarrow d_3$	$d_3 \rightarrow d_4$	$d_4 \rightarrow d_5$	$d_5 \rightarrow d_6$	$d_6 \rightarrow d_1$
$\text{Arg}(d_u \rightarrow d_v; 1)$	$-\frac{3}{14}$	$-\frac{2}{7}$	$-\frac{3}{7}$	$-\frac{2}{7}$	$-\frac{3}{14}$	$\frac{3}{7}$
$\text{Arg}(d_u \rightarrow d_v; 2)$	$\frac{1}{2}$	$-\frac{3}{14}$	$-\frac{1}{7}$	$-\frac{3}{14}$	$\frac{1}{2}$	$-\frac{3}{7}$
$\text{Arg}(d_u \rightarrow d_v; 3)$	$-\frac{1}{7}$	$\frac{1}{14}$	0	$\frac{1}{14}$	$-\frac{1}{7}$	$\frac{1}{7}$


- $c' \equiv 4 \pmod{7}$, $\beta = 2$. $A \cdot 2 = 7T + 1$, $\text{Arg}(Q \rightarrow P; 2) = \frac{3}{14}$;
- $c' \equiv 5 \pmod{7}$, $\beta = 3$. $A \cdot 3 = 7T + 1$, $\text{Arg}(Q \rightarrow P; 3) = \frac{1}{14}$;
- $c' \equiv 6 \pmod{7}$, $\beta = 6$. $A \cdot 1 = 7T - 1$, $\text{Arg}(Q \rightarrow P; 1) = \frac{1}{7}$.


$c' \equiv 4 \pmod{7}$	$d_1 \rightarrow d_2$	$d_2 \rightarrow d_3$	$d_3 \rightarrow d_4$	$d_4 \rightarrow d_5$	$d_5 \rightarrow d_6$	$d_6 \rightarrow d_1$
$\text{Arg}(d_u \rightarrow d_v; 1)$	$\frac{3}{14}$	$\frac{2}{7}$	$\frac{3}{7}$	$\frac{2}{7}$	$\frac{3}{14}$	$-\frac{3}{7}$
$\text{Arg}(d_u \rightarrow d_v; 2)$	$\frac{1}{2}$	$\frac{3}{14}$	$\frac{1}{7}$	$\frac{3}{14}$	$\frac{1}{2}$	$\frac{3}{7}$
$\text{Arg}(d_u \rightarrow d_v; 3)$	$\frac{1}{7}$	$-\frac{1}{14}$	0	$-\frac{1}{14}$	$\frac{1}{7}$	$-\frac{1}{7}$

$c' \equiv 5 \pmod{7}$	$d_1 \rightarrow d_2$	$d_2 \rightarrow d_3$	$d_3 \rightarrow d_4$	$d_4 \rightarrow d_5$	$d_5 \rightarrow d_6$	$d_6 \rightarrow d_1$
$\text{Arg}(d_u \rightarrow d_v; 1)$	$\frac{3}{14}$	$\frac{1}{7}$	0	$\frac{1}{7}$	$\frac{3}{14}$	$\frac{2}{7}$
$\text{Arg}(d_u \rightarrow d_v; 2)$	$\frac{1}{14}$	$-\frac{1}{14}$	$\frac{1}{7}$	$-\frac{1}{14}$	$\frac{1}{14}$	$-\frac{1}{7}$
$\text{Arg}(d_u \rightarrow d_v; 3)$	0	$\frac{1}{14}$	$-\frac{2}{7}$	$\frac{1}{14}$	0	$\frac{1}{7}$

$c' \equiv 6 \pmod{7}$	$d_1 \rightarrow d_2$	$d_2 \rightarrow d_3$	$d_3 \rightarrow d_4$	$d_4 \rightarrow d_5$	$d_5 \rightarrow d_6$	$d_6 \rightarrow d_1$
$\text{Arg}(d_u \rightarrow d_v; 1)$	$\frac{1}{2}$	$\frac{1}{7}$	$\frac{3}{7}$	$\frac{1}{7}$	$\frac{1}{2}$	$\frac{2}{7}$
$\text{Arg}(d_u \rightarrow d_v; 2)$	$-\frac{1}{14}$	$-\frac{3}{14}$	0	$-\frac{3}{14}$	$-\frac{1}{14}$	$-\frac{3}{7}$
$\text{Arg}(d_u \rightarrow d_v; 3)$	$\frac{1}{7}$	$\frac{5}{14}$	$\frac{2}{7}$	$\frac{5}{14}$	$\frac{1}{7}$	$-\frac{2}{7}$

Note that the condition for $c' \pmod{7}$ is the same as the reversed condition for $-c' \pmod{7}$. Hence we only need show the corresponding graphs for $c' \equiv 1, 2, 3 \pmod{7}$, and also for the other $7n + k$ cases in the remaining subsections. In each of the following graphs, if d_u and d_v are not shown, then $P(d_u) = P(d_v)$ are both the remaining non-labeled point.





Visualizing by the above graphs, (6.9) is proved by the following equations:

$$\begin{aligned}
 & C_7^{4,6} \sin\left(\frac{\pi}{7}\right) \left(\frac{\cos\left(\frac{2\pi}{7}\right)}{\sin\left(\frac{\pi}{7}\right)} + \frac{\cos\left(\frac{3\pi}{7}\right)}{\sin\left(\frac{2\pi}{7}\right)} - \frac{\cos\left(\frac{2\pi}{7}\right)}{\sin\left(\frac{3\pi}{7}\right)} \right) + C_7^{6,2} \sin\left(\frac{2\pi}{7}\right) \left(\frac{\cos\left(\frac{3\pi}{7}\right)}{\sin\left(\frac{\pi}{7}\right)} - \frac{\cos\left(\frac{3\pi}{7}\right)}{\sin\left(\frac{2\pi}{7}\right)} + \frac{1}{\sin\left(\frac{3\pi}{7}\right)} \right) \\
 & + C_7^{2,4} \sin\left(\frac{3\pi}{7}\right) \left(-\frac{\cos\left(\frac{2\pi}{7}\right)}{\sin\left(\frac{\pi}{7}\right)} + \frac{1}{\sin\left(\frac{2\pi}{7}\right)} + \frac{\cos\left(\frac{2\pi}{7}\right)}{\sin\left(\frac{3\pi}{7}\right)} \right) = -C_7^{4,6} \sin\left(\frac{\pi}{7}\right) \sqrt{7}, \\
 & C_7^{4,6} \sin\left(\frac{\pi}{7}\right) \left(-\frac{\cos\left(\frac{2\pi}{7}\right)}{\sin\left(\frac{\pi}{7}\right)} + \frac{1}{\sin\left(\frac{2\pi}{7}\right)} + \frac{\cos\left(\frac{2\pi}{7}\right)}{\sin\left(\frac{3\pi}{7}\right)} \right) + C_7^{6,2} \sin\left(\frac{2\pi}{7}\right) \left(\frac{1}{\sin\left(\frac{\pi}{7}\right)} + \frac{\cos\left(\frac{\pi}{7}\right)}{\sin\left(\frac{2\pi}{7}\right)} + \frac{\cos\left(\frac{\pi}{7}\right)}{\sin\left(\frac{3\pi}{7}\right)} \right) \\
 & + C_7^{2,4} \sin\left(\frac{3\pi}{7}\right) \left(\frac{\cos\left(\frac{2\pi}{7}\right)}{\sin\left(\frac{\pi}{7}\right)} + \frac{\cos\left(\frac{\pi}{7}\right)}{\sin\left(\frac{2\pi}{7}\right)} + \frac{\cos\left(\frac{\pi}{7}\right)}{\sin\left(\frac{3\pi}{7}\right)} \right) = -C_7^{2,4} \sin\left(\frac{3\pi}{7}\right) \sqrt{7}, \\
 & C_7^{4,6} \sin\left(\frac{\pi}{7}\right) \left(\frac{\cos\left(\frac{3\pi}{7}\right)}{\sin\left(\frac{\pi}{7}\right)} - \frac{\cos\left(\frac{3\pi}{7}\right)}{\sin\left(\frac{2\pi}{7}\right)} + \frac{1}{\sin\left(\frac{3\pi}{7}\right)} \right) + C_7^{6,2} \sin\left(\frac{2\pi}{7}\right) \left(-\frac{\cos\left(\frac{3\pi}{7}\right)}{\sin\left(\frac{\pi}{7}\right)} + \frac{\cos\left(\frac{3\pi}{7}\right)}{\sin\left(\frac{2\pi}{7}\right)} + \frac{\cos\left(\frac{\pi}{7}\right)}{\sin\left(\frac{3\pi}{7}\right)} \right) \\
 & + C_7^{2,4} \sin\left(\frac{3\pi}{7}\right) \left(\frac{1}{\sin\left(\frac{\pi}{7}\right)} + \frac{\cos\left(\frac{\pi}{7}\right)}{\sin\left(\frac{2\pi}{7}\right)} + \frac{\cos\left(\frac{\pi}{7}\right)}{\sin\left(\frac{3\pi}{7}\right)} \right) = -C_7^{6,2} \sin\left(\frac{2\pi}{7}\right) \sqrt{7}.
 \end{aligned}$$

This proves (7-0) of Theorem 1.3. We omit the similar proofs of (7- k) for $k \in \{1, 2, 3, 4, 6\}$.

ACKNOWLEDGEMENT

The author thanks Professor Scott Ahlgren for his careful reading in a previous version of this paper and for his plenty of insightful suggestions. The author also thanks Nick Andersen and Alexander Dunn for helpful comments.

REFERENCES

[ASD54] A. O. L. Atkin and H. P. F. Swinnerton-Dyer. Some properties of partitions. *Proc. London Math. Soc.* (3), 4:84–106, 1954.

[BO06] Kathrin Bringmann and Ken Ono. The $f(q)$ mock theta function conjecture and partition ranks. *Invent. Math.*, 165(2):243–266, 2006.

[Bri09] Kathrin Bringmann. Asymptotics for rank partition functions. *Trans. Amer. Math. Soc.*, 361:3483–3500, 2009.

[Dys44] Freeman Dyson. *Some guesses in the theory of partitions*, volume 8. Eureka (Cambridge), 1944.

[Gar17] Francis Garvan. Transformation properties for Dyson’s rank function. *Trans. Amer. Math. Soc.*, 371:1, 02 2017.

[Lew95] Richard Lewis. The components of modular forms. *J. Lond. Math. Soc.*, 52(2):245–254, 1995.

[Sun24a] Qihang Sun. Exact formulae for ranks of partitions, 2024. arXiv:2406.06294 [math.NT].

[Sun24b] Qihang Sun. GitHub repository: Vanishing Kloosterman sums, 2024. https://github.com/qihangsun1998/Vanishing_Kloosterman_Sums.