

Spread Complexity of High Energy Neutrino Propagation over Astrophysical Distances

Khushboo Dixit,¹ S. Shajidul Haque,^{2,3,4} and Soebur Razzaque^{1,5,4}

¹*Centre for Astro-Particle Physics (CAPP) and Department of Physics,
University of Johannesburg, PO Box 524, Auckland Park 2006, South Africa*

²*High Energy Physics, Cosmology & Astrophysics Theory Group and The Laboratory for Quantum Gravity & Strings,
Department of Mathematics & Applied Mathematics,
University of Cape Town, Cape Town, South Africa*

³*Department of Mathematics and Natural Sciences, Brac University, Dhaka, Bangladesh*

⁴*National Institute for Theoretical and Computational Sciences (NITheCS), Private Bag X1, Matieland, South Africa*

⁵*Department of Physics, The George Washington University, Washington, DC 20052, USA*

(Dated: June 12, 2024)

Spread complexity measures the minimized spread of quantum states over all choices of basis. It generalizes Krylov operator complexity to quantum states under continuous Hamiltonian evolution. In this paper, we study spread complexity in the context of high-energy astrophysical neutrinos and propose a new flavor ratio based on complexity. Our findings indicate that our proposal might favor an initial ratio of fluxes as $\phi_{\nu_e}^0 : \phi_{\nu_\mu}^0 : \phi_{\nu_\tau}^0 = 1 : 0 : 0$ over a more generally expected ratio of $1 : 2 : 0$, when the IceCube neutrino observatory achieves its projected sensitivity to discriminate between flavors. Additionally, complexity-based definitions of flavor ratios exhibit a slight but nonzero sensitivity to the neutrino mass ordering, which traditional flavor ratios cannot capture.

I. Introduction

Neutrinos have long been a subject of profound interest within the field of particle physics. The discovery of neutrino oscillations, which signifies the massive nature of neutrinos, has significantly impacted the scientific community, prompting an intensified quest for physics beyond the Standard Model (BSM) [1–8]. Presently, numerous long-baseline experiments, both established and planned, are dedicated to scrutinizing oscillation parameters with heightened precision and resolution [9–12]. Additionally, some experiments such as the IceCube neutrino observatory, situated at the South Pole, are actively engaged in the pursuit of high-energy astrophysical neutrinos. This pursuit is motivated by the potential for these neutrinos and their flavor composition to unveil crucial insights into their production mechanisms within celestial entities [13–26].

In addition, various analyses have demonstrated that neutrinos exhibit intriguing properties such as entanglement and several other nonlocal correlations [27–36]. These characteristics underscore their significant potential for executing diverse tasks pertinent to Quantum Information Processing (QIP). One of the important aspects related to quantum information and computation theory is known as quantum complexity, which was proposed as a useful probe [37] for the interior of a black hole by using the anti-de Sitter/conformal field theory (AdS/CFT) duality. Holographic complexity, proposed to be equivalent to the "volume" or "action," elucidates the physics behind the horizon of an eternal AdS black hole [37–41]. However, in recent days, complexity has proven to be extremely useful for understanding various aspects of quantum systems such as quantum chaos [42–47], quantum phase transitions [48], quantum decoherence [49, 50], and more. It has also been applied to investigate cosmological perturbation models and the evolution of the universe [51–53].

Complexity can be described in two ways when dealing with quantum mechanical systems. First, according to Nielsen's definition, complexity is the minimal distance in the space of the unitaries between the identity and the time evolution operator $U(t)$ of the system [54–58]. The second approach defines complexity as minimizing the spread of the state over a specified basis during the time evolution [59, 60]. The complexity of the spread of states has recently been studied in the context of neutrino oscillations in accelerator neutrino-based long baseline experiments [61]. This study not only indicated that complexity could provide more information on the CP-violating phase compared to the oscillation probability, but it also provided a perspective on the preferred value of the CP-violating phase angle. Motivated by this, in this paper, we examine the complexity of astrophysical neutrinos by considering three different initial flavor ratios based on their possible production mechanisms. IceCube Neutrino Observatory, currently operating in the Antarctica has been collecting data on astrophysical neutrinos and has potential to measure their flavor ratios [24], especially after the recent detection of astrophysical tau neutrinos [26]. Our findings suggest that the modified flavor ratios, in terms of complexities, support the idea that neutron-decay (with an initial flux ratio of $\phi_{\nu_e}^0 : \phi_{\nu_\mu}^0 : \phi_{\nu_\tau}^0 = 1 : 0 : 0$) might be favored over π -decay (with an initial flux ratio of $1 : 2 : 0$) as the mechanism responsible for the production of neutrinos in celestial bodies. We delve into the details of neutrino oscillations and flavor ratios in the context of astrophysical neutrinos in sec. II A, followed by defining spread complexity in sec. II B.

We then provide the analytical expressions of flavor ratios modified in terms of complexities in sec. II C. Finally, we present our results and discussions in sec. III and IV, respectively.

II. Neutrino Oscillations and Quantum Complexity

In this section, we discuss neutrino oscillations in the context of astrophysical neutrinos and recall the complexity embedded in a state evolution. This is followed by a discussion on the complexities for each initial neutrino flavor modified in the context of astrophysical neutrinos. Additionally, we propose the redefined flavor ratios of neutrinos in terms of complexities.

A. Astrophysical neutrino oscillations and flavor ratios

It is now well known that neutrinos can change their flavor while traveling to a distant place. Depending on their energy as well as their sensitivity to the mass squared difference associated to their mass eigenstates one can calculate the oscillation length for such flavor transitions. Neutrino oscillations occur as the flavor states of neutrino are not the mass eigenstates, but they mix via a unitary matrix to form these mass eigenstates. In the case of three flavor oscillations, the form of this unitary matrix can be given as

$$U = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & 0 & s_{13}e^{-i\delta} \\ 0 & 1 & 0 \\ -s_{13}e^{i\delta} & 0 & c_{13} \end{pmatrix} \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

where, c_{ij} and s_{ij} are $\cos \theta_{ij}$ and $\sin \theta_{ij}$ with mixing angles θ_{ij} ($i, j = 1, 2, 3$). The probability of survival or oscillation after the time evolution of the neutrino with a specific flavor can be given by

$$P_{\alpha\beta} = \delta_{\alpha\beta} - 4 \sum_{j>k} \text{Re}\{U_{\alpha j}^* U_{\beta j} U_{\alpha k} U_{\beta k}^*\} \sin^2 \left(\frac{\Delta m_{ij}^2 L}{4E} \right) - 2 \sum_{j>k} \text{Im}\{U_{\alpha j}^* U_{\beta j} U_{\alpha k} U_{\beta k}^*\} \sin \left(\frac{\Delta m_{ij}^2 L}{2E} \right)$$

In the case of astrophysical neutrinos where neutrinos travel through the distances of Mpc or larger order, the sine and cosine terms carrying the phase factor get averaged out and the probabilities take the following form

$$P_{\alpha\beta} = \delta_{\alpha\beta} - 2 \sum_{j>k} \text{Re}\{U_{\alpha j}^* U_{\beta j} U_{\alpha k} U_{\beta k}^*\}$$

The quantity of interest in the experiments searching for high energy astrophysical neutrinos is the so-called *flavor ratio*. The fraction of each neutrino flavor observed in astrophysical neutrino experiments can be defined using the flavor oscillation probabilities and the initial flavor ratios. The initial ν_τ flavor is supposed to be zero as the neutrino production mechanism in these high energy astrophysical objects like blazars or AGNs are considered to be either neutron beta decay, $n \rightarrow p + e^- + \bar{\nu}_e$, which induces initial flavor ratio of $\phi_e^0 : \phi_\mu^0 : \phi_\tau^0 = 1 : 0 : 0$, or pion decay, $\pi^+ \rightarrow e^+ + \nu_e + \bar{\nu}_\mu + \nu_\mu$, with initial flavor ratio of $1 : 2 : 0$. In the case of μ -damped π -decay, when the μ loses substantial energy before decaying [62], the initial ratio of $0 : 1 : 0$ is also possible. The fractions of each neutrino flavor arriving on the Earth can be expressed as follows

$$\begin{aligned} X_e &\equiv \frac{\phi_{\nu_e}}{\phi_{\nu_e} + \phi_{\nu_\mu} + \phi_{\nu_\tau}} = \frac{P_{ee}\phi_{\nu_e}^0 + P_{\mu e}\phi_{\nu_\mu}^0}{\phi_{\nu_e}^0 + \phi_{\nu_\mu}^0 + \phi_{\nu_\tau}^0} \\ X_\mu &\equiv \frac{\phi_{\nu_\mu}}{\phi_{\nu_e} + \phi_{\nu_\mu} + \phi_{\nu_\tau}} = \frac{P_{e\mu}\phi_{\nu_e}^0 + P_{\mu\mu}\phi_{\nu_\mu}^0}{\phi_{\nu_e}^0 + \phi_{\nu_\mu}^0 + \phi_{\nu_\tau}^0} \\ X_\tau &\equiv \frac{\phi_{\nu_\tau}}{\phi_{\nu_e} + \phi_{\nu_\mu} + \phi_{\nu_\tau}} = \frac{P_{e\tau}\phi_{\nu_e}^0 + P_{\mu\tau}\phi_{\nu_\mu}^0}{\phi_{\nu_e}^0 + \phi_{\nu_\mu}^0 + \phi_{\nu_\tau}^0} \end{aligned} \quad (1)$$

B. Complexity of spread of states

In this section, we recall the formulation of the complexity of the spread of states as introduced in [59, 60]. The evolution of a state $|\psi(0)\rangle$ is governed by the Schrodinger equation as follows

$$i\partial_t |\psi(t)\rangle = H |\psi(t)\rangle,$$

where, H is the time independent Hamiltonian of the system and the solution to this equation gives us the time evolved state

$$\begin{aligned} |\psi(t)\rangle &= e^{-iHt} |\psi(0)\rangle, \\ &= \sum_{n=0}^{\infty} \frac{(-iHt)^n}{n!} |\psi(0)\rangle, \\ &= \sum_{n=0}^{\infty} \frac{(-it)^n}{n!} |\psi_n\rangle, \end{aligned}$$

where, $|\psi_n\rangle = H^n |\psi\rangle$. We then apply the Gram-Schmidt procedure to obtain an ordered orthonormal (Krylov) basis using the set of infinite states $\{|\psi_n\rangle\}$. The number of states spanning such a basis can be equal to or less than the dynamics of the system, however, it has been seen that in case of neutrinos, the number of basis states is exactly equal to the dimension of the system [61]. We expect more complex spread of a state if the dynamics involved is chaotic. To formulate the degree of complexity a cost function can be defined as

$$\chi = \sum_n |\langle K_n |\psi(t)\rangle|^2 \quad (2)$$

It has been shown in [59] that the above cost function achieves its minimum for the Krylov basis, hence, this minimized cost function defined in Eq. (2) can be treated as the complexity embedded in the evolution of the state of the system.

C. Spread complexity in flavor oscillations of astrophysical neutrinos

Spread complexity introduced in Eq. (2) has been formulated for the general case of three flavor neutrino oscillations in [61]. The complexity for neutrino starting as ν_e flavor at the initial point can be expressed below

$$\begin{aligned} \chi_e &= P_{e\mu}(t) [N_{1e}^2 |a_1|^2 + 2N_{2e}^2 |b_1|^2] + P_{e\tau}(t) [N_{1e}^2 |a_2|^2 + 2N_{2e}^2 |b_2|^2] \\ &\quad + 2\Re [N_{1e}^2 a_1^* a_2 A_{e\mu}(t) A_{e\tau}(t)^*] + 4\Re [N_{2e}^2 b_1^* b_2 A_{e\mu}(t) A_{e\tau}(t)^*]. \end{aligned} \quad (3)$$

Here, $P_{e\mu}$ and $P_{e\tau}$ are the oscillation probabilities for $\nu_e \rightarrow \nu_\mu$ and $\nu_e \rightarrow \nu_\tau$ transitions, respectively and $A_{e\mu}$ and $A_{e\tau}$ are the corresponding oscillation amplitudes. In the case of astrophysical neutrinos where the travel distance is significantly large (of the order of Mpc or larger), the sin and cos terms carrying the phase factor depending on the mass squared differences $m_j^2 - m_i^2$ ($j > i$ and $i, j = 1, 2, 3$) corresponding to the mass eigenstates driving flavor oscillations get averaged out. Therefore, these probabilities and amplitudes take the following form

$$\begin{aligned} P_{e\mu} &= -2 (\Re(U_{e2}^* U_{\mu 2} U_{e1} U_{\mu 1}^*) + \Re(U_{e3}^* U_{\mu 3} U_{e1} U_{\mu 1}^*) + \Re(U_{e3}^* U_{\mu 3} U_{e2} U_{\mu 2}^*)) \\ &= \frac{1}{8} \sin \theta_{12} \cos \theta_{12} \cos^2 \theta_{13} (4 \cos \delta \cos 2\theta_{12} \sin \theta_{13} \sin 2\theta_{23} \\ &\quad + \sin 2\theta_{12} (2 \cos 2\theta_{13} \sin^2 \theta_{23} + 3 \cos 2\theta_{23} + 1)), \\ P_{e\tau} &= -2 (\Re(U_{e2}^* U_{\tau 2} U_{e1} U_{\tau 1}^*) + \Re(U_{e3}^* U_{\tau 3} U_{e1} U_{\tau 1}^*) + \Re(U_{e3}^* U_{\tau 3} U_{e2} U_{\tau 2}^*)) \\ &= -\cos^2 \theta_{13} \left(\frac{1}{4} \sin \theta_{13} (\cos \delta (4 \sin 2\theta_{12} + \sin 4\theta_{12}) \sin 2\theta_{23} \right. \\ &\quad \left. + 8 \sin^2 \theta_{12} \sin \theta_{13} \cos^2 \theta_{23}) - 2 \cos^4 \theta_{12} \sin^2 \theta_{13} \cos^2 \theta_{23} \right. \\ &\quad \left. - 2 \sin^2 \theta_{12} \cos^2 \theta_{12} \sin^2 \theta_{23} \right), \\ A_{e\mu} A_{e\tau}^* &= |U_{e1}|^2 U_{\mu 1}^* U_{\tau 1} + |U_{e2}|^2 U_{\mu 2}^* U_{\tau 2} + |U_{e3}|^2 U_{\mu 3}^* U_{\tau 3} \\ &= \frac{1}{16} \cos^2 \theta_{13} (\sin 2\theta_{23} (-\cos 4\theta_{12} + 7) \cos 2\theta_{13} + 3 \cos 4\theta_{12} + 5) \\ &\quad + 4 \sin 4\theta_{12} \sin \theta_{13} (\cos \delta \cos 2\theta_{23} + i \sin \delta). \end{aligned} \quad (4)$$

The effect of averaging out the sine and cosine terms will be visible in probabilities $P_{e\mu}$ & $P_{e\tau}$ and the cross term $A_{e\mu}A_{e\tau}^*$ only. Other terms such as $|a_1|^2$, $|a_2|^2$, $|b_1|^2$, $|b_2|^2$, $a_1^*a_2$ and $b_1^*b_2$ are given in the appendix.

One important point should also be noted that since the oscillation terms carrying the phase factor in terms of Δm^2 get averaged out, the discrimination between the normal ordering (NO) and inverted ordering (IO) of neutrino mass eigenstates is impossible through probabilities. However, complexities have slight but non-vanishing discrimination in the effects of NO and IO as can be seen in the expressions of $|a_1|^2$, $|a_2|^2$, $|b_1|^2$, $|b_2|^2$ and so on in Eq. (7). Similarly, the complexities associated with the evolution of neutrino states can be obtained in case of initial ν_μ and ν_τ flavors as well.




FIG. 1: Complexities χ_α (normal ordering (NO) and inverted ordering (IO) of neutrino masses in blue and red curves, respectively) and $1 - P_{\alpha\alpha}$ (black curve) plotted with respect to the CP -violating phase δ . Here $\alpha = e, \mu, \tau$ depends on the initial neutrino flavor. All other parameters are set to their best-fit values provided in a recent analysis [63].

D. Redefining Flavor Ratios Using Complexity

In this section, we will propose a new flavor ratio based on complexity. We can consider the initial flavor ratio to be $\phi_{\nu_e}^0 : \phi_{\nu_\mu}^0 : \phi_{\nu_\tau}^0 = (x : 1 - x : 0)\phi_{\nu_e}^0$, where $x = 0, 1, 1/3$ which denotes flavor ratios $\phi_{\nu_e}^0 : \phi_{\nu_\mu}^0 : \phi_{\nu_\tau}^0 = (0 : 1 : 0)$, $(1 : 0 : 0)$, $(1 : 2 : 0)$, respectively. Here, $\phi_{\nu_e} + \phi_{\nu_\mu} + \phi_{\nu_\tau} = \phi_{\nu_e}^0 + \phi_{\nu_\mu}^0 + \phi_{\nu_\tau}^0$ as a consequence of probability conservation. Hence, fractions of distinct flavors defined in Eq. (1) can be revised as

$$\begin{aligned} X_e &= xP_{ee} + (1 - x)P_{\mu e} \\ X_\mu &= xP_{\mu e} + (1 - x)P_{\mu\mu} \\ X_\tau &= (2x - 1)P_{e\tau} + (1 - x)(1 - P_{\tau\tau}). \end{aligned} \quad (5)$$

Then we introduce here the analogous equation to the Eq. (5) in terms of complexities as

$$\begin{aligned} X'_e &= x(1 - \chi_e) + (1 - x)P_{\mu e}. \\ X'_\mu &= xP_{\mu e} + (1 - x)(1 - \chi_\mu) \\ X'_\tau &= (2x - 1)P_{e\tau} + (1 - x)\chi_\tau. \end{aligned} \quad (6)$$

Here, we have replaced $1 - P_{\alpha\alpha}$ with corresponding χ_α , for flavor $\alpha = e, \mu, \tau$. We take inspiration from the findings in [61] and Fig. 1 to make this replacement, noting that χ_α and $1 - P_{\alpha\alpha}$ share similar characteristics. The flavor ratios outlined in Eq. (6) represent a combination of complexities and oscillation probabilities. Despite this, they can provide valuable insights, as complexities are known to encapsulate a greater amount of pertinent information about the system.

III. Results

In this section, we initially present our results concerning flavor ratios using traditional probability-based definitions, followed by a modification of these definitions to incorporate complexities for each neutrino flavor. In [61], a significant characteristic of complexities has been identified, indicating that the complexities χ_α , ($\alpha = e, \mu, \tau$) emulate the traits of their corresponding total oscillation probabilities, i.e., $1 - P_{\alpha\alpha}$, although χ_α provides more information. A similar observation is evident in the context of astrophysical neutrinos, as illustrated in Fig. 1. It becomes apparent that both complexities χ_e (depicted as blue(solid) and red(dashed) curves for normal and inverted mass ordering, respectively) and the total oscillation probability $1 - P_{ee}$ (denoted by the brown(dot-dashed) curve) are not impacted by the CP-phase δ . However, χ_μ , χ_τ , as well as $1 - P_{\mu\mu}$ and $1 - P_{\tau\tau}$, exhibit variations in relation to the δ phase. Furthermore, these quantities display symmetric characteristics around $\delta = 0$, signifying their CP-even nature, where CP-even denotes $f(\delta) = f(-\delta)$, with f encompassing χ_α and $1 - P_{\alpha\alpha}$. Notably, complexities χ_μ and χ_τ demonstrate enhanced sensitivity to the CP-phase. Additionally, Fig. 1 highlights a subtle sensitivity of complexities χ_α to the normal and inverted mass ordering, a feature absent in the case of $1 - P_{\alpha\alpha}$.




FIG. 2: The flavor ratios X_α (X'_α) are shown on a ternary plot representation. Dark-gray and light-gray regions represent the 68% and 95% CL of Asimov sensitivity to measure flavor composition using the current GlobalFit sample [63], while the dark-pink and light-pink regions show the same sensitivity with the inclusion of 12 years of HESE sample as well [64]. Blue and red colors depict the cases of relative fractions of X'_e , X'_μ and X'_τ corresponding to the normal ordering (NO) and inverted ordering (IO), respectively. The purple points represent the fraction of traditionally defined flavor ratios, i.e., X_e , X_μ and X_τ . Solid lines (purple, blue and red) represents the flavor ratios with varying x in the range $[0,1]$. The cases of initial flavor ratios $(1:0:0)$, $(0:1:0)$ and $(1:2:0)$ corresponding to $x = 1, 0, 1/3$, i.e., respectively are shown in different styles of points as mentioned in the legend.

In Fig. 2, the ternary plot shows the representation of flavor ratios. The traditional flavor ratios X_α are depicted in purple, while their analogous quantities X'_α are shown in blue (for NO) and red (for IO). The dark-gray and light-gray regions represent the 68% and 95% C.L. of Asimov sensitivity for measuring flavor composition using the current GlobalFit sample [64]. The dark-pink and light-pink regions show the same sensitivity when the 12 years of HESE sample is included, with 68% and 95% C.L., respectively [64].




FIG. 3: X'_e , X'_μ and X'_τ defined in Eq. (6) are plotted with respect to x . The solid and dashed curves represent X'_α with NO and X'_α with IO, while the associated initial neutrino flavor ν_e , ν_μ and ν_τ are represented by blue, orange and black color, respectively. Dot-dashed curves represent traditional ratios X_α defined in Eq. (5), respectively.

It is interesting that in contrast with the traditional flavor ratios, the complexity-based flavor ratios favor the initial ratio of neutrino flavors to be $1 : 0 : 0$. In general, the most anticipated initial neutrino flavor ratio is $1 : 2 : 0$, expected to be observed as a $1 : 1 : 1$ ratio at Earth. However, there is insufficient evidence supporting this assumption. The findings in reference [20] indicate that the best fit to IceCube data at the time is achieved with a $1 : 0 : 0$ ratio at Earth, which does not align with any of these proposed initial flavor ratios, namely $(1 : 0 : 0)$, $(0 : 1 : 0)$, $(1 : 2 : 0)$, or any other arbitrary choice as shown in Fig. 3 where we vary the initial fraction via the parameter x between $[0, 1]$.

Apart from the above observations, the following points are important to be mentioned:

- Firstly, the analogous flavor ratios X'_α defined in terms of complexities χ_e , χ_μ and χ_τ can have more spread with respect to the variation in CP -phase δ both in the case of NO and IO in comparison to the traditional quantities X_α as the complexities in Fig. (1) are shown to have more variations with respect to δ than the total oscillation probabilities. That means these newly defined flavor ratios can provide better resolution for the CP -violating phase value. Therefore, complexity-based flavor ratios can also offer the possibility to constrain or exclude a significant range of the δ -phase, in principle.
- Secondly, in Fig. (2), the newly defined flavor ratios in terms of complexities also show small but non-zero sensitivity to the neutrino mass ordering (slight difference between the blue and red lines). Specifically, the initial flavor ratio $\phi_{\nu_e}^0 : \phi_{\nu_\mu}^0 : \phi_{\nu_\tau}^0 = 1 : 0 : 0$ favors this condition the most. This sensitivity is absent in the case of traditional flavor ratios (purple). The point to be noticed here is that complexity-based ratios also favor the case of $1 : 0 : 0$ over $1 : 2 : 0$ as initial flavor ratios of the astrophysical neutrinos, while compared with the sensitivity contours shown in the plot.

The above-mentioned advantages of the redefined flavor ratios, X'_α in Eq. (6) in terms of the complexities over the traditional ones, X_α , in Eq. (5) can be attributed to the cross terms present in the complexities defined in Eq. (3) along with the terms proportional to the oscillation probabilities. We also provide individual fractions of neutrino flavors varying with respect to arbitrary values of x (the parameter defining the initial neutrino flavor ratio) in Fig. (3), which may correspond to non-traditional neutrino production mechanism.

IV. Discussion

In this work, we have attempted to redefine the neutrino flavor ratios arriving on the Earth in terms of the complexities of neutrino propagation due to the motivation that these complexities and the total oscillation probabilities of neutrino flavor states depict equivalent features, whereas the former convey more information. Comparing these complexity-based flavor ratios with the traditional ones, we find that the former favors the initial flavor ratio of $1 : 0 : 0$ in place of $1 : 2 : 0$ might be indicating the production mechanism of neutrinos to be $n - decay$ in astrophysical objects. Furthermore, these quantities are found to be sensitive to the neutrino mass ordering, unlike the case of traditional flavor ratios. Also, notably the complexity-based flavor ratios manifest a greater degree of variability in the CP -violating phase. These features might be tested in the future with observational data from the neutrino telescopes.

Acknowledgments

This work was partially supported by grants from the National Research Foundation (NRF), South Africa, through the National Institute of Theoretical and Computational Sciences (NITheCS) and from the University of Johannesburg Research Council.

$$\begin{aligned}
|a_1|^2 &= (\Delta m_{21}^2)^2 |U_{e2}|^2 |U_{\mu 2}|^2 + (\Delta m_{31}^2)^2 |U_{e3}|^2 |U_{\mu 3}|^2 + 2(\Delta m_{21}^2)(\Delta m_{31}^2) \Re(U_{e2}^* U_{\mu 2} U_{e3} U_{\mu 3}^*) \\
&= \cos^2 \theta_{13} \left((\Delta m_{21}^2)^2 \sin^2 \theta_{12} (\sin^2 \delta \sin^2 \theta_{12} \sin^2 \theta_{13} \sin^2 \theta_{23} \right. \\
&\quad \left. + (\cos \theta_{12} \cos \theta_{23} - \cos \delta \sin \theta_{12} \sin \theta_{13} \sin \theta_{23})^2 \right) + \Delta m_{31}^2 \sin \theta_{13} \sin \theta_{23} \\
&\quad \left(\Delta m_{21}^2 \cos \delta \sin 2\theta_{12} \cos \theta_{23} + \sin \theta_{13} \sin \theta_{23} (\Delta m_{31}^2 - 2\Delta m_{21}^2 \sin^2 \theta_{12}) \right) \\
|a_2|^2 &= (\Delta m_{21}^2)^2 |U_{e2}|^2 |U_{\tau 2}|^2 + (\Delta m_{31}^2)^2 |U_{e3}|^2 |U_{\tau 3}|^2 + 2(\Delta m_{21}^2)(\Delta m_{31}^2) \Re(U_{e2}^* U_{\tau 2} U_{e3} U_{\tau 3}^*) \\
&= \cos^2 \theta_{13} \left(-\frac{1}{4} \Delta m_{21}^2 \cos \delta \sin 2\theta_{12} \sin \theta_{13} \sin 2\theta_{23} (\Delta m_{21}^2 \cos 2\theta_{12} - \Delta m_{21}^2 + 2\Delta m_{31}^2) \right. \\
&\quad \left. + (\Delta m_{21}^2)^2 \sin^2 \theta_{12} \cos^2 \theta_{12} \sin^2 \theta_{23} \right. \\
&\quad \left. + \frac{1}{4} \sin^2 \theta_{13} \cos^2 \theta_{23} (\Delta m_{21}^2 (-\cos 2\theta_{12}) + \Delta m_{21}^2 - 2\Delta m_{31}^2)^2 \right) \\
|b_1|^2 &= (\Delta m_{21}^2)^2 (\Delta m_{21}^2 - 2E_\nu A_e)^2 |U_{e2}|^2 |U_{\mu 2}|^2 + (\Delta m_{31}^2)^2 (\Delta m_{31}^2 - 2E_\nu A_e)^2 |U_{e3}|^2 |U_{\mu 3}|^2 \\
&\quad + 2(\Delta m_{21}^2)(\Delta m_{31}^2)(\Delta m_{21}^2 - 2E_\nu A_e)(\Delta m_{31}^2 - 2E_\nu A_e) \Re(U_{e2}^* U_{\mu 2} U_{e3} U_{\mu 3}^*) \\
|b_2|^2 &= (\Delta m_{21}^2)^2 (\Delta m_{21}^2 - 2E_\nu A_e)^2 |U_{e2}|^2 |U_{\tau 2}|^2 + (\Delta m_{31}^2)^2 (\Delta m_{31}^2 - 2E_\nu A_e)^2 |U_{e3}|^2 |U_{\tau 3}|^2 \\
&\quad + 2(\Delta m_{21}^2)(\Delta m_{31}^2)(\Delta m_{21}^2 - 2E_\nu A_e)(\Delta m_{31}^2 - 2E_\nu A_e) \Re(U_{e2}^* U_{\tau 2} U_{e3} U_{\tau 3}^*) \\
a_1^* a_2 &= (\Delta m_{21}^2)^2 |U_{e2}|^2 U_{\mu 2}^* U_{\tau 2} + (\Delta m_{31}^2)^2 |U_{e3}|^2 U_{\mu 3}^* U_{\tau 3} \\
&\quad + (\Delta m_{21}^2)(\Delta m_{31}^2) [U_{e2} U_{\mu 2}^* U_{e3}^* U_{\tau 3} + U_{e3} U_{\mu 3}^* U_{e2}^* U_{\tau 2}] \\
&= \frac{1}{8} \cos^2 \theta_{13} \left(-8e^{-i\delta} \Delta m_{21}^2 \sin \theta_{12} \cos \theta_{12} \sin \theta_{13} (-\sin^2 \theta_{23} + e^{2i\delta} \cos^2 \theta_{23}) \right. \\
&\quad \left(\Delta m_{21}^2 \sin^2 \theta_{12} - \Delta m_{31}^2 \right) - (\Delta m_{21}^2)^2 \sin^2 2\theta_{12} \sin 2\theta_{23} \\
&\quad \left. + \sin^2 \theta_{13} \sin 2\theta_{23} (2\Delta m_{21}^2 \sin^2 \theta_{12} - 2\Delta m_{31}^2)^2 \right), \\
b_1^* b_2 &= (\Delta m_{21}^2)^2 (\Delta m_{21}^2 - 2E_\nu A_e)^2 |U_{e2}|^2 U_{\mu 2}^* U_{\tau 2} + (\Delta m_{31}^2)^2 (\Delta m_{31}^2 - 2E_\nu A_e)^2 |U_{e3}|^2 U_{\mu 3}^* U_{\tau 3} \\
&\quad + (\Delta m_{21}^2)(\Delta m_{31}^2)(\Delta m_{21}^2 - 2E_\nu A_e)(\Delta m_{31}^2 - 2E_\nu A_e) [U_{e2} U_{\mu 2}^* U_{e3}^* U_{\tau 3} + U_{e3} U_{\mu 3}^* U_{e2}^* U_{\tau 2}] \\
&= [e^{-i\delta} (\Delta m_{21}^2)^2 (\Delta m_{31}^2)^2 (\Delta m_{21}^2 - \Delta m_{31}^2)^2 \sin^2 \theta_{12} \cos^2 \theta_{12} \sin^2 \theta_{13} \cos^2 \theta_{13} (\Delta m_{21}^2 \sin \theta_{12} \\
&\quad \cos \theta_{12} \sin \theta_{23} - e^{i\delta} \sin \theta_{13} \cos \theta_{23} (\Delta m_{31}^2 - \Delta m_{21}^2 \sin^2 \theta_{12})) \\
&\quad (\sin \theta_{13} \sin \theta_{23} (\Delta m_{31}^2 - \Delta m_{21}^2 \sin^2 \theta_{12}) + e^{i\delta} \Delta m_{21}^2 \sin \theta_{12} \cos \theta_{12} \cos \theta_{23})] \\
&\quad \times [(-\Delta m_{21}^2)^2 \sin^4 \theta_{12} \cos^2 \theta_{13} + \Delta m_{21}^2 \sin^2 \theta_{12} \\
&\quad (\Delta m_{21}^2 + \Delta m_{31}^2 \cos 2\theta_{13} - \Delta m_{31}^2) + (\Delta m_{31}^2)^2 \sin^2 \theta_{13}]^2. \tag{7}
\end{aligned}$$

The normalization constants can be spelled as

$$N_{1e}^2 = [(\Delta m_{21}^2)^2 |U_{e2}|^2 (1 - |U_{e2}|^2) + (\Delta m_{31}^2)^2 |U_{e3}|^2 (1 - |U_{e3}|^2) - 2(\Delta m_{21}^2)(\Delta m_{31}^2) |U_{e2}|^2 |U_{e3}|^2]^{-1} \tag{8}$$

$$N_{2e}^2 = [(\Delta m_{21}^2)^2 (\Delta m_{21}^2 - 2E_\nu A_e)^2 |U_{e2}|^2 (1 - |U_{e2}|^2) + (\Delta m_{31}^2)^2 (\Delta m_{31}^2 - 2E_\nu A_e)^2 |U_{e3}|^2 (1 - |U_{e3}|^2) - 2(\Delta m_{21}^2)(\Delta m_{31}^2)(\Delta m_{21}^2 - 2E_\nu A_e)(\Delta m_{31}^2 - 2E_\nu A_e) |U_{e2}|^2 |U_{e3}|^2]^{-1} \tag{9}$$

where,

$$\begin{aligned}
A_e &= \frac{1}{2E} [(\Delta m_{21}^2)^3 |U_{e2}|^2 (1 - |U_{e2}|^2) + (\Delta m_{31}^2)^3 |U_{e3}|^2 (1 - |U_{e3}|^2) - \Delta m_{21}^2 \Delta m_{31}^2 |U_{e2}|^2 |U_{e3}|^2 \\
&\quad (\Delta m_{21}^2 + \Delta m_{31}^2)] [(\Delta m_{21}^2)^3 |U_{e2}|^2 (1 - |U_{e2}|^2) + (\Delta m_{31}^2)^3 |U_{e3}|^2 (1 - |U_{e3}|^2) \\
&\quad - 2(\Delta m_{21}^2)(\Delta m_{31}^2) |U_{e2}|^2 |U_{e3}|^2]^{-1}
\end{aligned}$$

-
- [1] K. S. Hirata et al., Real time, directional measurement of B-8 solar neutrinos in the Kamiokande-II detector. *Phys. Rev. D*, 44:2241, 1991. [Erratum: *Phys.Rev.D* 45, 2170 (1992)].
- [2] B. T. Cleveland, Timothy Daily, Raymond Davis, Jr., James R. Distel, Kenneth Lande, C. K. Lee, Paul S. Wildenhain, and Jack Ullman. Measurement of the solar electron neutrino flux with the Homestake chlorine detector. *Astrophys. J.*, 496:505–526, 1998.
- [3] Q. R. Ahmad et al., Direct evidence for neutrino flavor transformation from neutral current interactions in the Sudbury Neutrino Observatory. *Phys. Rev. Lett.*, 89:011301, 2002.
- [4] Y. Fukuda et al. Atmospheric muon-neutrino / electron-neutrino ratio in the multiGeV energy range. *Phys. Lett. B*, 335:237–245, 1994.
- [5] Y. Fukuda et al. Evidence for oscillation of atmospheric neutrinos. *Phys. Rev. Lett.*, 81:1562–1567, 1998.
- [6] Y. Abe et al. Indication for the disappearance of reactor electron antineutrinos in the Double Chooz experiment. *Phys.Rev.Lett.*, 108:131801, 2012.
- [7] D. Adey et al. Measurement of the Electron Antineutrino Oscillation with 1958 Days of Operation at Daya Bay. *Phys. Rev. Lett.*, 121(24):241805, 2018.
- [8] G. Bak et al. Measurement of Reactor Antineutrino Oscillation Amplitude and Frequency at RENO. *Phys. Rev. Lett.*, 121(20):201801, 2018.
- [9] B. Abi et al. [DUNE], *Eur. Phys. J. C* **80** (2020) no.10, 978 [arXiv:2006.16043 [hep-ex]].
- [10] J. Zhang [JUNO], *J. Phys. Conf. Ser.* **2156** (2021) no.1, 012110 [arXiv:2111.10112 [physics.ins-det]].
- [11] M. A. Acero *et al.* [NOvA], *Phys. Rev. D* **106** (2022) no.3, 032004 [arXiv:2108.08219 [hep-ex]].
- [12] K. Abe *et al.* [T2K], *Eur. Phys. J. C* **83** (2023) no.9, 782 [arXiv:2303.03222 [hep-ex]].
- [13] G. Barenboim and C. Quigg, *Phys. Rev. D* **67**, 073024 (2003) [arXiv:hep-ph/0301220 [hep-ph]].
- [14] Z. Z. Xing and S. Zhou, *Phys. Rev. D* **74**, 013010 (2006) [arXiv:astro-ph/0603781 [astro-ph]].
- [15] P. Lipari, M. Lusignoli and D. Meloni, *Phys. Rev. D* **75**, 123005 (2007) [arXiv:0704.0718 [astro-ph]].
- [16] S. Pakvasa, W. Rodejohann and T. J. Weiler, *JHEP* **02**, 005 (2008) [arXiv:0711.4517 [hep-ph]].
- [17] A. Esmaili and Y. Farzan, *Nucl. Phys. B* **821**, 197-214 (2009) [arXiv:0905.0259 [hep-ph]].
- [18] K. C. Lai, G. L. Lin and T. C. Liu, *Phys. Rev. D* **80**, 103005 (2009) [arXiv:0905.4003 [hep-ph]].
- [19] S. Choubey and W. Rodejohann, *Phys. Rev. D* **80**, 113006 (2009) [arXiv:0909.1219 [hep-ph]].
- [20] O. Mena, S. Palomares-Ruiz and A. C. Vincent, Flavor Composition of the High-Energy Neutrino Events in IceCube, *Phys. Rev. Lett.* **113** (2014), 091103 [arXiv:1404.0017 [astro-ph.HE]].
- [21] A. Palladino, G. Pagliaroli, F. L. Villante and F. Vissani, *Phys. Rev. Lett.* **114**, no.17, 171101 (2015) [arXiv:1502.02923 [astro-ph.HE]].
- [22] M. G. Aartsen *et al.* [IceCube], *Phys. Rev. Lett.* **114**, no.17, 171102 (2015) [arXiv:1502.03376 [astro-ph.HE]].
- [23] M. Bustamante, J. F. Beacom and W. Winter, *Phys. Rev. Lett.* **115**, no.16, 161302 (2015) [arXiv:1506.02645 [astro-ph.HE]].
- [24] M. G. Aartsen *et al.* [IceCube], *Astrophys. J.* **846** (2017) no.2, 136 doi:10.3847/1538-4357/aa8508 [arXiv:1705.02383 [astro-ph.HE]].
- [25] N. Song, S. W. Li, C. A. Argüelles, M. Bustamante and A. C. Vincent, *JCAP* **04**, 054 (2021) [arXiv:2012.12893 [hep-ph]].
- [26] R. Abbasi *et al.* [IceCube], *Eur. Phys. J. C* **82**, no.11, 1031 (2022) [arXiv:2011.03561 [hep-ex]].
- [27] M. Blasone, F. Dell’Anno, S. De Siena, M. Di Mauro, and F. Illuminati. Multipartite entangled states in particle mixing. *Phys. Rev. D*, 77:096002, 2008.
- [28] M. Blasone, F. Dell’Anno, S. De Siena, and F. Illuminati, Entanglement in neutrino oscillations. *EPL*, 85:50002, 2009.
- [29] D. Gangopadhyay, D. Home, and A. Sinha Roy. Probing the Leggett-Garg Inequality for Oscillating Neutral Kaons and Neutrinos. *Phys. Rev. A*, 88(2):022115, 2013.
- [30] A. K. Alok, S. Banerjee, and S. Uma Sankar. Quantum correlations in terms of neutrino oscillation probabilities. *Nucl. Phys. B*, 909:65–72, 2016.
- [31] S. Banerjee, A. K. Alok, R. Srikanth, and B. C. Hiesmayr. A quantum information theoretic analysis of three flavor neutrino oscillations. *Eur. Phys. J. C*, 75(10):487, 2015.
- [32] J. A. Formaggio, D. I. Kaiser, M. M. Murskyj, and T. E. Weiss. Violation of the Leggett-Garg Inequality in Neutrino Oscillations. *Phys. Rev. Lett.*, 117(5):050402, 2016.
- [33] Q. Fu and X. Chen. Testing violation of the Leggett–Garg-type inequality in neutrino oscillations of the Daya Bay experiment. *Eur. Phys. J. C*, 77(11):775, 2017.
- [34] X. K. Song, Y. Huang, J. Ling and M. H. Yung, *Phys. Rev. A* **98** (2018) no.5, 050302 [arXiv:1806.00715 [hep-ph]].
- [35] M. M. Etefaghi, Z. S. Tabatabaei Lotfi, and R. Ramezani Arani. Quantum correlations in neutrino oscillation: Coherence and entanglement. *EPL*, 132(3):31002, 2020.
- [36] S. Shafaq and P. Mehta, *J. Phys. G* **48** (2021) no.8, 085002 [arXiv:2009.12328 [hep-ph]].
- [37] L. Susskind, *Fortsch. Phys.* **64** (2016), 49-71 [arXiv:1411.0690 [hep-th]].
- [38] L. Susskind, *Fortsch. Phys.* **64** (2016), 24-43 [arXiv:1403.5695 [hep-th]].
- [39] D. Stanford and L. Susskind, *Phys. Rev. D* **90** (2014) no.12, 126007 [arXiv:1406.2678 [hep-th]].
- [40] A. R. Brown, D. A. Roberts, L. Susskind, B. Swingle and Y. Zhao, *Phys. Rev. Lett.* **116** (2016) no.19, 191301 [arXiv:1509.07876 [hep-th]].
- [41] A. R. Brown, D. A. Roberts, L. Susskind, B. Swingle and Y. Zhao, *Phys. Rev. D* **93** (2016) no.8, 086006 [arXiv:1512.04993 [hep-th]].

- [42] T. Ali, A. Bhattacharyya, S. S. Haque, E. H. Kim, N. Moynihan and J. Murugan, *Phys. Rev. D* **101** (2020) no.2, 026021 [arXiv:1905.13534 [hep-th]].
- [43] A. Bhattacharyya, W. Chemissany, S. Shajidul Haque and B. Yan, *Eur. Phys. J. C* **82** (2022) no.1, 87 [arXiv:1909.01894 [hep-th]].
- [44] A. Bhattacharyya, S. S. Haque and E. H. Kim, *JHEP* **10** (2021), 028 [arXiv:2011.04705 [hep-th]].
- [45] V. Balasubramanian, M. Decross, A. Kar and O. Parrikar, *JHEP* **01** (2020), 134 [arXiv:1905.05765 [hep-th]].
- [46] A. Bhattacharyya, W. Chemissany, S. S. Haque, J. Murugan and B. Yan, *SciPost Phys. Core* **4** (2021), 002 [arXiv:2007.01232 [hep-th]].
- [47] V. Balasubramanian, M. DeCross, A. Kar, Y. Li and O. Parrikar, *JHEP* **07** (2021), 011 [arXiv:2101.02209 [hep-th]].
- [48] T. Ali, A. Bhattacharyya, S. Shajidul Haque, E. H. Kim and N. Moynihan, *Phys. Lett. B* **811** (2020), 135919 [arXiv:1811.05985 [hep-th]].
- [49] A. Bhattacharyya, T. Hanif, S. S. Haque and A. Paul, *Phys. Rev. D* **107** (2023) no.10, 106007 [arXiv:2210.09268 [hep-th]].
- [50] A. Bhattacharyya, T. Hanif, S. S. Haque and M. K. Rahman, *Phys. Rev. D* **105** (2022) no.4, 046011 [arXiv:2112.03955 [hep-th]].
- [51] A. Bhattacharyya, S. Das, S. Shajidul Haque and B. Underwood, *Phys. Rev. D* **101** (2020) no.10, 106020 [arXiv:2001.08664 [hep-th]].
- [52] A. Bhattacharyya, S. Das, S. S. Haque and B. Underwood, *Phys. Rev. Res.* **2** (2020) no.3, 033273 [arXiv:2005.10854 [hep-th]].
- [53] S. S. Haque, C. Jana and B. Underwood, *Phys. Rev. D* **106** (2022) no.6, 063510 [arXiv:2110.08356 [hep-th]].
- [54] M. A. Nielsen, M. R. Dowling, M. Gu and A. C. Doherty, *Science* **311** (2006) no.5764, 1133-1135 [arXiv:quant-ph/0603161 [quant-ph]].
- [55] M. A. Nielsen, *Quant. Inf. Comput.* **6** (2006) no.3, 213-262 [arXiv:quant-ph/0502070 [quant-ph]].
- [56] M. R. Dowling and M. A. Nielsen, *Quant. Inf. Comput.* **8** (2008) no.10, 0861-0899 [arXiv:quant-ph/0701004 [quant-ph]].
- [57] R. Jefferson and R. C. Myers, *JHEP* **10** (2017), 107 [arXiv:1707.08570 [hep-th]].
- [58] T. Ali, A. Bhattacharyya, S. Shajidul Haque, E. H. Kim and N. Moynihan, *JHEP* **04** (2019), 087 [arXiv:1810.02734 [hep-th]].
- [59] V. Balasubramanian, P. Caputa, J. M. Magan and Q. Wu, Quantum chaos and the complexity of spread of states, *Phys. Rev. D* **106** (2022) no.4, 046007 [arXiv:2202.06957 [hep-th]].
- [60] P. Caputa and S. Liu, Quantum complexity and topological phases of matter, *Phys. Rev. B* **106** (2022) no.19, 195125 [arXiv:2205.05688 [hep-th]].
- [61] K. Dixit, S. S. Haque and S. Razzaque, Quantum spread complexity in neutrino oscillations, *Eur. Phys. J. C* **84** (2024) no.3, 260 [arXiv:2305.17025 [hep-ph]].
- [62] J. P. Rachen and P. Meszaros, *Phys. Rev. D* **58**, 123005 (1998) [arXiv:astro-ph/9802280 [astro-ph]].
- [63] I. Esteban, M. C. Gonzalez-Garcia, M. Maltoni, T. Schwetz and A. Zhou, The fate of hints: updated global analysis of three-flavor neutrino oscillations, *JHEP* **09** (2020), 178 [arXiv:2007.14792 [hep-ph]].
- [64] R. Abbasi *et al.* [IceCube], *PoS ICRC2023* (2023), 1122 [arXiv:2308.15213 [astro-ph.HE]].