UNIQUENESS ON AVERAGE OF LARGE ISOPERIMETRIC SETS IN NONCOMPACT MANIFOLDS WITH NONNEGATIVE RICCI CURVATURE

GIOACCHINO ANTONELLI, MARCO POZZETTA, AND DANIELE SEMOLA

ABSTRACT. Let (M^n, g) be a complete Riemannian manifold which is not isometric to \mathbb{R}^n , has nonnegative Ricci curvature, Euclidean volume growth, and quadratic Riemann curvature decay. We prove that there is a set $\mathcal{G} \subset (0, \infty)$ with density 1 at infinity such that for every $V \in \mathcal{G}$ there is a unique isoperimetric set of volume V in M, and its boundary is strictly volume preserving stable.

The latter result cannot be improved to uniqueness or strict stability for every large volume. Indeed, we construct a complete Riemannian surface that satisfies the previous assumptions, and with the following property: there are arbitrarily large and diverging intervals $I_n \subset (0, \infty)$ such that isoperimetric sets with volumes $V \in I_n$ exist, but they are neither unique nor they have strictly volume preserving stable boundaries.

The proof relies on a set of new ideas, as the present setting goes beyond the range of applicability of the methods based on the implicit function theorem, and no symmetry is assumed.

Contents

1. Introduction	1
Strict stability for many volumes	3
Uniqueness for many volumes	4
Conjectures and Open Questions	5
Acknowledgments	5
2. The example: Proof of Theorem 1.2	6
3. Strict stability for many volumes	8
4. Uniqueness for many volumes	15
4.1. Proof of Theorem 4.1 given Proposition 4.3 and Proposition 4.4	16
4.2. Estimates for isoperimetric graphs	21
4.3. Proof of Proposition 4.3	25
4.4. Proof of Proposition 4.4	26
Appendix A. Quantitative regularity of almost minimizers in almost Euclidean manife	olds 31
Appendix B. Uniform bounds for the reach	35
References	36

1. INTRODUCTION

In this paper we study the isoperimetric structure for large volumes of complete Riemannian manifolds (M^n, g) with nonnegative Ricci curvature, Euclidean volume growth, and quadratic Riemann curvature decay. More precisely, we shall assume that Ric ≥ 0 everywhere on M^n and that for some (and hence for every) base point $p \in M$ it holds

(1.1)
$$\operatorname{AVR}(M) := \lim_{r \to \infty} \frac{\operatorname{vol}(B_r(p))}{\omega_n r^n} > 0,$$

where ω_n is the volume of the unit ball in \mathbb{R}^n , and

(1.2)
$$|\operatorname{Riem}(x)| \le \frac{C}{\mathrm{d}^2(x,p)} \quad \text{for all } x \in M ,$$

for some constant C > 0. This class of manifolds has been studied thoroughly in the literature and includes several examples of interest. Under these assumptions, it is known (see for instance [45, 5]) that isoperimetric sets with volume V exist for each V > 0.

It is well-known that Riemannian manifolds with nonnegative Ricci curvature and Euclidean volume growth have metric cones as blow-downs, namely any pointed sequence $(M, r_i^{-2}g, o)$ for $o \in M$ and diverging r_i admits a subsequence converging in pointed measured Gromov– Hausdorff sense to an *n*-dimensional metric measure cone (C, d_C, \mathcal{H}^n) , see [19]. Assuming in addition (1.2), then all the blow-downs of (M^n, g) are metric cones $(C(N), dr^2 + r^2g_N)$ over some compact Riemannian manifold (N, g_N) , where g_N is a $C^{1,\alpha}$ Riemannian metric for each $\alpha < 1$, and $\operatorname{Ric}_N \geq n-2$ (possibly in a weak sense). Although the blow-down of (M^n, g) is not necessarily unique under these assumptions, see [52] and the more recent [27], the manifold (M^n, g) is $C^{1,\alpha}$ asymptotic to the family of its blow-down cones, for each $\alpha < 1$. Moreover, for each of these asymptotic cones the isoperimetric set with volume V is unique, it has strictly volume preserving stable boundary, and it coincides with a ball centered at the vertex for each volume V > 0, see [47], unless the blow-down is isometric to \mathbb{R}^n . However, if a blow-down is isometric to \mathbb{R}^n , then (M^n, g) is necessarily isometric to \mathbb{R}^n , see [26].

Our main result shows that uniqueness and strict volume preserving stability of the isoperimetric sets hold for *most* sufficiently large volumes also for the complete manifold (M^n, g) .

Theorem 1.1. Let (M^n, g) be a smooth complete Riemannian manifold with Ric ≥ 0 , Euclidean volume growth (1.1), and quadratic Riemann curvature decay (1.2). Assume that (M^n, g) is not isometric to \mathbb{R}^n . Then there exists a measurable set $\mathcal{G} \subset (0, \infty)$ such that

(1.3)
$$\lim_{r \to \infty} \frac{\mathcal{L}^1(\mathcal{G} \cap (r, 2r))}{r} = 1,$$

and for each $V \in \mathcal{G}$ there exists a unique isoperimetric set $E_V \subset M^n$ with $\operatorname{vol}(E_V) = V$. Moreover, the boundary of E_V is a smooth strictly volume preserving stable constant mean curvature hypersurface.

Theorem 1.1 is sharp, in the sense that neither the uniqueness nor the strict stability statement can be obtained for *all* sufficiently large volumes in this generality. Indeed, we construct examples of complete surfaces with nonnegative Gaussian curvature, quadratic volume growth, and quadratic curvature decay such that uniqueness and strict stability fail for isoperimetric sets of arbitrarily large volumes.

Theorem 1.2. There exists a complete smooth Riemannian surface (M^2, g) with nonnegative Gaussian curvature, quadratic volume growth (1.1), and quadratic curvature decay (1.2), such that the following holds. There are disjoint intervals $L_n \subset (0, \infty)$ with $\inf L_n \to \infty$ and $|L_n| \to \infty$ as $n \to \infty$, such that for every $n \in \mathbb{N}$, and for every $V \in L_n$, there are isoperimetric sets with volume V in M which are neither strictly volume preserving stable, nor unique.

The construction of Theorem 1.2 can be easily generalized to obtain analogous examples with Ric ≥ 0 satisfying (1.1) and (1.2) in any dimension $n \geq 3$.

The question of existence, uniqueness, and characterization of isoperimetric sets for large volumes, and more in general of constant mean curvature hypersurfaces on complete Riemannian manifolds under various types of curvature constraints and asymptotic conditions at infinity has received a great deal of attention in the last thirty years, see for instance [37, 55, 13, 14, 17, 16, 29, 28, 22, 21, 23, 33], and the references therein. We mention in particular [24] where the authors obtain the uniqueness of isoperimetric regions for each sufficiently large volume in complete Riemannian manifolds (M^n, g) that are $C^{2,\alpha}$ -asymptotic to a fixed cone $C(N^{n-1})$ whose link (N^{n-1}, g_N) satisfies

(1.4)
$$\operatorname{Ric}_N \ge n-2, \quad \operatorname{vol}(N) < \operatorname{vol}(\mathbb{S}^{n-1}).$$

Note that the conditions in (1.4) are satisfied for each cross-section of any blow-down of a manifold (M^n, g) as in the assumptions of Theorem 1.1. On the other hand, the assumptions of Theorem 1.1 are not sufficient to guarantee neither the uniqueness of the blow-down, nor the $C^{2,\alpha}$ convergence.

The mild curvature decay condition (1.2) makes the techniques of [24], which are based on the application of the implicit function theorem, and of all the other references mentioned above unsuitable for the present setting. Moreover, to the best of our knowledge, the *uniqueness on average* captured in Theorem 1.1 had not been studied before.

In the rest of the introduction we outline the proof of Theorem 1.1 and highlight the main novelties of our approach.

Strict stability for many volumes. To set the notation, we recall that the stability (or Jacobi) operator of a (smooth) two-sided constant mean curvature hypersurface $\Sigma^{n-1} \subset M^n$ takes the form

(1.5)
$$C^{\infty}(\Sigma) \ni f \mapsto -\Delta_{\Sigma} f - (\operatorname{Ric}(\nu, \nu) + |\operatorname{II}|^2) f \in C^{\infty}(\Sigma)$$

and it is associated to the quadratic form

(1.6)
$$C^{\infty}(\Sigma) \ni f \mapsto I(f, f) := \int_{\Sigma} \left[|\nabla_{\Sigma} f|^2 - (\operatorname{Ric}(\nu, \nu) + |\operatorname{II}|^2) f^2 \right] \operatorname{d} \operatorname{vol}_{\Sigma}$$

In (1.5) and (1.6), ν and II denote a unit normal to Σ and its second fundamental form respectively. If Σ is the boundary of an isoperimetric region, then it is a (volume preserving) stable constant mean curvature hypersurface, i.e., the stability operator in (1.5) is nonnegative definite on the subspace of functions with 0 average on Σ .¹ Equivalently, the quadratic form in (1.6) is nonnegative whenever $\int_{\Sigma} f = 0$.² We say that such constant mean curvature hypersurface Σ is strictly (volume preserving) stable provided there exists $\lambda > 0$ such that

(1.7)
$$I(f,f) \ge \lambda \int_{\Sigma} f^2 \,\mathrm{d}\,\mathrm{vol}_{\Sigma}, \quad \text{for all } f \in C^{\infty}(\Sigma) \text{ such that } \int_{\Sigma} f \,\mathrm{d}\,\mathrm{vol}_{\Sigma} = 0.$$

Under the assumptions of Theorem 1.1, for each volume V > 0 it is natural to consider the rescaled pointed manifold

(1.8)
$$M_V := (M^n, r_V^{-1} \mathbf{d}_g, p),$$

where we set

(1.9)
$$r_V := (\operatorname{AVR}(M) \,\omega_n)^{-\frac{1}{n}} \, V^{\frac{1}{n}} \,,$$

with ω_n the volume of the unit ball in \mathbb{R}^n . For any sequence $V_i \to \infty$, up to the extraction of a subsequence that we do not relabel, the pointed manifolds M_{V_i} converge to one of the blow-downs C(N) of (M^n, g) in the pointed Gromov-Hausdorff sense and in the $C^{1,\alpha}$ sense locally away from the singular vertex, for each $\alpha < 1$, see for instance [19]. Denoting by E_i a sequence of isoperimetric sets in the rescaled manifold M_{V_i} with volume equal to $\operatorname{AVR}(M)\omega_n$ in the rescaled metric, a by now standard argument shows that the isoperimetric boundaries ∂E_i converge in the Hausdorff sense to the boundary of $B_1(o)$, the ball of radius 1 centered at the vertex of the limit cone C(N), along this converging sequence. See Lemma 3.2 below for the precise statement and [7] for a proof in a more general setting.

Note that on a cone the Ricci curvature in the radial direction vanishes, and balls centered at the vertex are totally umbilical. Hence, it follows from (1.4) and the Lichnerowicz and Obata theorems that $\partial B_1(o)$ is strictly stable, unless the cone is isometric to \mathbb{R}^n . Ideally, we would

¹In order to avoid confusion we stress that in general this is not an invariant subspace under the action of the Jacobi operator.

²We note that a priori there might be a non empty singular set of ambient codimension at least 8 in Σ . However, we shall see that in the setting of Theorem 1.1 isoperimetric regions of sufficiently large volumes are always smooth and hence we will neglect the possible presence of singularities for the sake of this discussion.

like to prove that the stability operators (under volume preserving variations) of $\partial E_i \subset M_{V_i}$ converge in some sense to the stability operator of the limit isoperimetric region $B_1(o) \subset C(N)$ in the blow-down, or at least that their first eigenvalues are lower semicontinuous with respect to *i*. The lower semicontinuity would be clearly enough to prove the strict stability of ∂E_i for any sufficiently large $i \in \mathbb{N}$. However, Theorem 1.2 shows that this is too optimistic in the present generality. We shall see that the main issue is related to the Ricci curvature term of the stability operator, which should be viewed as an error term and does not converge to 0 in general. The key for us will be that this Ricci curvature term goes to 0 on average with respect to the volume, see Proposition 3.4 for a precise statement. Establishing this convergence requires a new method with respect to the above mentioned list of references.

Uniqueness for many volumes. The proof of the uniqueness part of Theorem 1.1 is based on the broad principle that strictly stable implies locally uniquely minimizing (possibly under a volume constraint). This principle was originally put forward for minimal hypersurfaces in [60], and later extended to constant mean curvature hypersurfaces in [32] and [48]. However, the implementation in the present setting requires several new insights. Most importantly, the principle cannot be implemented without restricting the set of volumes, as the examples constructed in Theorem 1.2 illustrate. Indeed, the blow-down argument discussed above shows that, for V large enough, if there exist two different isoperimetric regions $E_V, E'_V \subset M$ with volume V then they must be close to each other in a scale-invariant sense and this scaleinvariant distance converges to 0 as $V \to \infty$. If both E_V and E'_V have strictly stable boundaries, then by [32, 48] they have neighbourhoods $\partial E_V \subset U$ and $\partial E'_V \subset U'$ where they are uniquely isoperimetric. The difficulty stems from the fact that the size of these neighbourhoods might not be scale invariantly bounded away from 0. Hence it is not clear whether $\partial E'_V \subset U$ or $\partial E_V \subset U'$. We shall see that it is possible to estimate effectively the size of these neighbourhoods, up to further restricting the set of *good* volumes. This estimate requires a new method with respect to those present in the literature, due to the weak convergence of (M^n, g) to its tangent cones under rescaling.

There will be three main steps in the proof of the uniqueness. The first step will be to show that if there are two distinct isoperimetric regions E_V, E'_V with the same sufficiently large volume V and the same mean curvature of the boundary, then $\partial E'_V$ can be written as a normal graph over ∂E_V with effective scale-invariant bounds on the graphing function. See Proposition 4.3 for the precise statement. This step requires some effective versions of classical estimates in Geometric Measure Theory.

The second step of the proof will amount to get an effective control on the Ricci curvature of (M^n, g) in the direction perpendicular to the boundary of an isoperimetric set, for each isoperimetric set of volume V and for most large volumes V. The spirit is that of a maximal function estimate, see Proposition 4.4. The novelty is that the estimate will be obtained first at the level of the isoperimetric profile of M and then turned into a curvature estimate on the manifold M.

In the last step we will effectively Taylor expand the perimeter functional near ∂E_V by integrating the Laplacian of the signed distance function $d_{E_V}^s$ in the region between E_V and E'_V . We note that effective Taylor expansions of the area functional have been employed with a different aim in the recent [38]. The approach of this paper seems to be novel, even though the relationship between the Laplacian of the distance function, the mean curvature and the stability operator has been used extensively in the last decades, see for instance [43, 44, 57]. Setting

(1.10)
$$u := w - \frac{1}{P(E_V)} \int_{\partial E_V} w \, \mathrm{d}P_{E_V} \,,$$

with w denoting the graphing function obtained in the first step described above, the main term of this Taylor expansion will be the stability operator $I_{\partial E_V}(u, u)$, up to some well-controlled lower order terms and a potentially not small error term involving the Ricci curvature of M. The Taylor expansion holds for every volume. At this stage, for most large volumes we will be

The Taylor expansion holds for every volume. At this stage, for most large volumes we will be able to exploit the result of the second step, together with the strict stability of ∂E_V , to show that also the potentially bad Ricci curvature term is negligible and reach a contradiction if we assume that the isoperimetric set is non-unique.

Conjectures and Open Questions. We end this introduction by mentioning a few open questions that arose during the present work and might be worthwhile of future investigation.

In the case of surfaces, i.e., for n = 2, a slight variant of the argument discussed in Section 3 allows to prove that most isoperimetric sets have strictly volume preserving stable boundary without assuming that the Gaussian curvature decays quadratically. In this regard we raise the following:

Conjecture 1.3. If n = 2, Theorem 1.1 holds true without assuming that the Gaussian curvature decays quadratically.

We do not venture in any conjecture in higher dimensions, however we record the following:

Open Question 1.4. Let (M^3, g) be a complete Riemannian manifold with nonnegative Ricci curvature and Euclidean volume growth such that no blow-down splits a line isometrically. Understand whether there exists a measurable set $\mathcal{G} \subset (0, \infty)$ such that

(1.11)
$$\lim_{r \to \infty} \frac{\mathcal{L}^1(\mathcal{G} \cap (r, 2r))}{r} = 1,$$

and, for each $V \in \mathcal{G}$, and every isoperimetric set $E_V \subset M^n$ with $\operatorname{vol}(E_V) = V$, the boundary of E_V is a smooth strictly volume preserving stable constant mean curvature hypersurface.

The examples that we construct in the proof of Theorem 1.2 have the drawback that, although for many large volumes the isoperimetric set with that volume is not unique, all the isoperimetric sets that we can find are isometric to each other.

Open Question 1.5. Understand whether there exists a complete Riemannian manifold (M^n, g) with nonnegative Ricci curvature, Euclidean volume growth, quadratic curvature decay, and such that there exists a sequence $V_i \to \infty$ with the following property. For every $i \in \mathbb{N}$ there exist two distinct isoperimetric sets $E_i, E'_i \subset M$ with volume V_i such that ∂E_i and $\partial E'_i$ are not isometric.

By a well-known argument, a positive answer to Open Question 1.5 would follow from a positive answer to the following:

Open Question 1.6. Understand whether there exists a complete Riemannian manifold (M^n, g) with nonnegative Ricci curvature, Euclidean volume growth, quadratic curvature decay, and such that there exists a sequence $V_i \to \infty$ with the following property. For every $i \in \mathbb{N}$ the isoperimetric profile of M is not differentiable at the point V_i .

We address the reader to [50] for some related constructions in the compact case.

Acknowledgments. This paper owes a lot to the participation of Elia Bruè to several conversations on the subject. Indeed, the authors believe that he might be considered a coauthor as well.

The authors are grateful to Otis Chodosh for a stimulating email exchange that partly motivated the present paper, to Anton Petrunin for providing a proof of Proposition B.1, and to Guido De Philippis for useful suggestions concerning Appendix A, and for having pointed out the reference [1]. This project started during the Thematic Program on Nonsmooth Riemannian and Lorentzian Geometry organized at the Fields Institute in Fall 2022. The authors gratefully acknowledge the warm hospitality and the stimulating atmosphere.

G.A. acknowledges the hospitality of the IAS during his visit in December 2021 where he discussed topics related to this work with Elia Bruè. G.A. also acknowledges the financial support of the Courant Institute and the AMS-Simons Travel Grant.

M.P. is a member of INdAM - GNAMPA and he is partially supported by the PRIN Project 2022E9CF89 - PNRR Italia Domani, funded by EU Program NextGenerationEU.

D.S. was supported by the FIM ETH Zürich through a Hermann Weyl Instructorship and by the Fields Institute through a Marsden Fellowship. He acknowledges SISSA for the stimulating atmosphere and the support during a visit in Fall 2023.

2. The example: Proof of Theorem 1.2

This section is dedicated to the proof of Theorem 1.2, that we restate below for the ease of readers.

Theorem 2.1. There exists a complete smooth Riemannian surface (M^2, g) with nonnegative Gaussian curvature, quadratic volume growth (1.1), and quadratic curvature decay (1.2), such that the following holds. There are disjoint intervals $L_n \subset (0, \infty)$ with $\inf L_n \to \infty$ and $|L_n| \to \infty$ as $n \to \infty$, such that for every $n \in \mathbb{N}$, and for every $V \in L_n$, there are isoperimetric sets with volume V in M which are neither strictly volume preserving stable, nor unique.

The examples will be warped products over S^1 . We will show that it is possible to choose the warping function in such a way that balls centered at the origin of the warped product do not have volume preserving stable boundaries and all the other requirements are met. In particular, balls centered at the origin cannot be isoperimetric sets for their enclosed volume. The fact that isoperimetric sets are not unique and not strictly volume preserving stable will follow immediately by the existence of an S^1 -symmetry. We will need the following elementary lemma, whose proof is postponed at the end of the section.

Lemma 2.2. Let $(a_n), (b_n)$ be two sequences of positive real numbers such that:

i) a_n is diverging, $a_{n+1} > a_n + b_n$, $a_n \ge b_n$;

ii) there holds

$$\sum_{n=0}^{\infty} \frac{b_n}{a_n} \le \frac{1}{16} \,.$$

Denote $I_n := [a_n, a_n + b_n]$ and let $J_n \subset I_n$. Then there exists a smooth concave function $\varphi : [0, \infty) \to \mathbb{R}$, with $\varphi(r) = r$ in a neighborhood of 0, such that for every n there holds

$$(2.1) \quad |\varphi''(r)| \le \frac{2}{a_n} \quad \forall r \in I_n, \quad |\varphi''(r)| \ge \frac{1}{a_n} \quad \forall r \in J_n, \quad \varphi''(r) \equiv 0 \quad \forall r \in (0,\infty) \setminus \bigcup_{n=0}^{\infty} I_n,$$

(2.2)
$$\frac{\varphi(r)\varphi''(r) - (\varphi'(r))^2 + 1}{\varphi^2(r)} < 0 \quad \forall r \in J_n \,,$$

(2.3)
$$-\frac{\varphi''(r)}{\varphi(r)} \le \frac{16}{r^2} \quad \forall r \in (0,\infty) \,.$$

Proof of Theorem 2.1. Let $(a_n), (b_n)$ be two sequences of positive real numbers satisfying the assumptions of Lemma 2.2. We can further assume that (b_n) diverges. Let I_n, J_n, φ be given by Lemma 2.2. Note that, in particular, $\inf I_n = a_n \to \infty$, and $|I_n| = b_n \to \infty$.

We consider the manifold M given by the warped product $[0, \infty) \times S^1$ endowed with the metric $g := dr^2 + \varphi(r)^2 g_{S^1}$. The fact that φ is concave and (2.3) holds imply that Sect ≥ 0 and Sect $= O(r^{-2})$. Applying for example [6, Corollary 5.9] (see also the earlier [56] and the more

recent [9, Theorem 1.4]) we get that isoperimetric sets exist for any volume on M. Moreover, since we also have $\varphi(r) \ge 7r/8$ (see (2.4)), we get that M has Euclidean volume growth (1.1).

By (2.2), we see that all the balls $B_r(0)$ with $r \in J_n$ have unstable boundaries with respect to volume-preserving variations. Hence, in particular, they are not isoperimetric. Indeed, since $\partial B_r(0)$ is isometric to a circle of length $2\pi\varphi(r)$, there exists $f \in C^{\infty}(\partial B_r(0))$ with $\int_{\partial B_r(0)} f = 0$ and $f \neq 0$ such that

$$\int_{\partial B_r(0)} |\nabla f|^2 = \frac{1}{\varphi^2(r)} \int_{\partial B_r(0)} f^2$$

where the gradient is understood with respect to the submanifold $\partial B_r(0)$. Recalling that $II_{\partial B_r(0)} = \frac{\varphi'(r)}{\varphi(r)}(g - dr^2)$ on the tangent space along $\partial B_r(0)$, we find

$$\int_{\partial B_r(0)} |\nabla f|^2 - \left(\left| \text{II}_{\partial B_r(0)} \right|^2 + \text{Ric}(\nu, \nu) \right) f^2 = \int_{\partial B_r(0)} \left(\frac{1 - (\varphi'(r))^2 + \varphi(r)\varphi''(r)}{\varphi^2(r)} \right) f^2 < 0,$$

due to (2.2).

Let $L_n := \{|B_r(0)| : r \in J_n\}$, where we denote by $|B_r(o)|$ the Riemannian volume measure of $B_r(o)$. Isoperimetric sets with volume $V \in L_n$ always exist as we argued above. However they cannot be balls centered at o because $B_r(0)$ is not volume-preserving stable for $r \in J_n$. Since M is rotationally symmetric, rotating isoperimetric sets with volume $V \in L_n$ shows that isoperimetric sets of such volumes are not (even locally) unique, and that they are not strictly volume preserving stable.

Finally, notice that $\inf L_n \to \infty$. Moreover, since (b_n) diverges, we can choose $J_n = [a'_n, b'_n]$ for suitable a'_n, b'_n such that $b'_n - a'_n \to \infty$. Thus it is readily checked that $|L_n| = 2\pi \int_{a'_n}^{b'_n} \varphi(r) \, dr \to \infty$ as well, because $\varphi(r) \ge 7r/8$ (see (2.4)).

Proof of Lemma 2.2. We choose $\varphi(r) = r$ in a right neighborhood of zero. We now explicitly construct φ'' in such a way $\varphi''(r) \leq 0$ everywhere and such that it is nonzero only on the intervals I_n . On every I_n we take φ'' equal to a negative smooth function such that $|\varphi''| = -\varphi'' \leq \frac{2}{a_n}$ on I_n and $|\varphi''| = -\varphi'' \geq \frac{1}{a_n}$ on J_n . In particular

$$\int_{I_n} \varphi'' \ge -\frac{2b_n}{a_n}$$

In this way, for every $r \in (0, \infty)$, we have

$$\varphi'(r) - \varphi'(0) = \int_0^r \varphi'' \ge -\sum_{n=0}^\infty \frac{2b_n}{a_n},$$

hence, using $\varphi(0) = 0$, and $\varphi'(0) = 1$,

(2.4)
$$\varphi'(r) \ge \frac{7}{8}, \qquad \varphi(r) \ge \frac{7}{8}r$$

Let us now verify (2.3). The inequality is trivially satisfied outside I_n . For $r \in I_n$, since in particular $a_n \leq r \leq a_n + b_n$, and $a_n \geq b_n$, we have

$$-\frac{\varphi''(r)}{\varphi(r)} \le \frac{16}{7a_n r} \le \frac{64}{7(2a_n)^2} \le \frac{64}{7(a_n + b_n)^2} \le \frac{64}{7r^2} \le \frac{16}{r^2}$$

Let us now verify (2.2). For $r \in J_n$ we have

$$-\varphi''(r)\varphi(r) \ge \frac{7r}{8a_n} \ge \frac{7}{8}$$

and thus

$$\varphi(r)\varphi''(r) - (\varphi'(r))^2 + 1 \le -\frac{7}{8} - \left(\frac{7}{8}\right)^2 + 1 < 0$$

which implies (2.2).

3. Strict stability for many volumes

The goal of this section is to prove the strict stability part of Theorem 1.1. The statement will be important later when it comes to proving uniqueness for most large volumes.

We state below the precise result for the ease of readers.

Theorem 3.1. Let (M^n, g) be a smooth complete Riemannian manifold with Ric ≥ 0 , Euclidean volume growth (1.1), and quadratic Riemann curvature decay (1.2). Assume that (M^n, g) is not isometric to \mathbb{R}^n . Then there exist $\varepsilon_M > 0$ and a measurable set $\mathcal{G} \subset (0, \infty)$ such that

(3.1)
$$\lim_{r \to \infty} \frac{\mathcal{L}^1(\mathcal{G} \cap (r, 2r))}{r} = 1,$$

and for each $V \in \mathcal{G}$ the first eigenvalue of the Jacobi operator for volume preserving variations on the boundary of every isoperimetric set $E_V \subset M$ of volume V is at least $\varepsilon_M V^{-2/n}$. In particular, ∂E_V is a (smooth) strictly volume preserving stable constant mean curvature hypersurface.

There are three intermediate steps for the proof of Theorem 3.1, corresponding to the following statements:

Lemma 3.2. Let (M^n, g) be a smooth complete Riemannian manifold with $\text{Ric} \geq 0$, Euclidean volume growth (1.1), and quadratic Riemann curvature decay (1.2), and assume that M is not isometric to \mathbb{R}^n . Let $p \in M$ be a fixed base point. For every $\varepsilon > 0$ there exists $V_0 = V_0(\varepsilon) > 0$ such that for every $V > V_0$ and for every isoperimetric set $E_V \subset M$ of volume V, setting

(3.2)
$$r_V := (\operatorname{AVR}(M)\,\omega_n)^{-\frac{1}{n}}\,V^{\frac{1}{n}},$$

the following hold:

(3.3)
$$d_{\rm H}(B_{r_V}(p), E_V) \le \varepsilon r_V, \quad d_{\rm H}(\partial B_{r_V}(p), \partial E_V) \le \varepsilon r_V,$$

where d_H denotes Gromov-Hausdorff distance.

Proposition 3.3. Let (M^n, g) be a smooth complete Riemannian manifold with Ric ≥ 0 , Euclidean volume growth (1.1), and quadratic Riemann curvature decay (1.2). Then, for every $\varepsilon > 0$ there exists $V_0 = V_0(\varepsilon) > 0$ such that for every $V > V_0$ any isoperimetric set $E_V \subset M$ with volume V is smooth and satisfies

(3.4)
$$\sup_{x \in \partial E_V} \left| V^{\frac{2}{n}} \left| II_{\partial E_V} \right|^2 (x) - (n-1)(\omega_n AVR(M))^{\frac{2}{n}} \right| \le \varepsilon ,$$

where $\Pi_{\partial E_V}$ is the second fundamental form of ∂E_V .

Proposition 3.4. Let (M^n, g) be a complete smooth Riemannian manifold with $\operatorname{Ric} \geq 0$, and Euclidean volume growth (1.1). For every $\varepsilon > 0$ there exists a measurable set $\mathcal{G}_{\varepsilon} \subset (0, \infty)$ such that

(3.5)
$$\lim_{r \to \infty} \frac{\mathcal{L}^1(\mathcal{G}_{\varepsilon} \cap (r, 2r))}{r} = 1,$$

and for every volume $V \in \mathcal{G}_{\varepsilon}$ and every isoperimetric set $E_V \subset M$ with volume V (if there exists one), it holds

(3.6)
$$V^{\frac{2}{n}} \oint_{\partial E_V} \operatorname{Ric}(\nu_{\partial E_V}, \nu_{\partial E_V}) \le \varepsilon,$$

where ν denotes the choice of a unit normal.

Note that Proposition 3.4 does not require the quadratic curvature decay (1.2), and it makes sense as soon as isoperimetric sets exist for sufficiently large volumes. However, if we assume that the curvature decays quadratically, the L^1 estimate for the Ricci curvature in the normal direction can be immediately improved to an L^p estimate for every $1 \leq p < \infty$, up to a multiplicative constant independent of the volume V.

We start by proving Lemma 3.2 below, as this will provide us with some relevant terminology for the proof of Theorem 3.1.

Proof of Lemma 3.2. We claim that under the present assumptions (M, g) does not split any line, and no blow-down of (M, g) splits any line. Given the claim, the statement follows from [8, Theorem 1.2], and [7, Theorem 4.23]. The rest of the proof is aimed at establishing the claim.

It is sufficient to prove that no blow-down of (M, g) splits any line, as if (M, g) splits a line then every blow-down must also split a line. Since (M, g) has Euclidean volume growth, every blow-down is a metric cone C(Z) over an $\mathsf{RCD}(n-2, n-1)$ space $(Z, \mathsf{d}_Z, \mathcal{H}^{n-1})$. The first part of the statement is due to [19], see also the previous [10] for the same conclusion under the quadratic curvature decay assumption. The second part is due to [39]. By the quadratic curvature decay, (Z, d_Z) is a $C^{1,\alpha}$ Riemannian manifold for every $\alpha < 1$. If C(Z) splits a line then Z is a spherical suspension over an $\mathsf{RCD}(n-3, n-2)$ space $(Z', \mathsf{d}_{Z'}, \mathcal{H}^{n-2})$. Since (Z, d_Z) is a $C^{1,\alpha}$ Riemannian manifold, all points $z \in Z$ are regular, i.e., the tangent cone at each point is isometric to \mathbb{R}^{n-1} . In particular, the suspension points are regular and hence $(Z', \mathsf{d}_{Z'}, \mathcal{H}^{n-2})$ must be isometric to the standard sphere \mathbb{S}^{n-2} endowed with a metric with constant sectional curvature 1. Therefore Z is isometric to the standard sphere \mathbb{S}^{n-1} endowed with a standard metric with constant sectional curvature 1, and hence C(Z) is isometric to \mathbb{R}^n . By volume convergence and volume rigidity, see [26], (M, g) is also isometric to \mathbb{R}^n , a contradiction.

Before giving detailed proofs of Proposition 3.3 and Proposition 3.4, we discuss how to complete the proof of Theorem 3.1 by taking them for granted.

Proof of Theorem 3.1. The proof will be divided in three main steps. In Step 1 we argue that the stability operators of the unit spheres have a uniform spectral gap among all blow-downs of (M, g). In Step 2 we rely on Lemma 3.2 and Proposition 3.3 to show that for any sequence of volumes $V_i \to \infty$ and any sequence of isoperimetric sets E_i such that $vol(E_i) = V_i$ the boundaries ∂E_i with the induced Riemannian metrics from the ambient M converge in the $C^{1,\alpha}$ sense to a cross-section of a blow-down of M, up to rescaling. In Step 3 we show that, roughly speaking, the first eigenvalue of the stability operator is lower semicontinuous along any such converging sequence provided that the Ricci curvature term is sufficiently small. The proof will be completed with the help of Proposition 3.4, whose aim is exactly to control that Ricci curvature term for many volumes.

Step 1. We claim that there exists $\varepsilon_M > 0$ such that for every cross-section (N, g_N) of a blow-down C(N) of (M, g) the first non-zero eigenvalue of the Laplacian $\lambda_1(N)$ satisfies $\lambda_1(N) \ge n - 1 + 2\varepsilon_M$.

As we argued in the proof of Lemma 3.2 above, any cross-section of a blow-down of (M, g)is a $C^{1,\alpha}$ Riemannian manifold (N, g) with Ricci curvature bounded from below by n - 2 in the sense of RCD spaces or, equivalently given the $C^{1,\alpha}$ regularity, in the sense of distributions. Moreover, no such cross-section is isometric to the standard sphere \mathbb{S}^{n-1} with constant sectional curvature equal to 1, otherwise C(N) and hence (M, g) would be isometric to \mathbb{R}^n , by [26]. By the Lichnerowicz estimate for the spectral gap of the Laplacian, see for instance [30, Theorem 4.22] for the setting of RCD spaces, the first non-zero eigenvalue of any such cross-section satisfies $\lambda_1(N) \geq n-1$. By Obata's rigidity theorem [40], the inequality is strict for each such N. Indeed, if not, N would be a spherical suspension and we already argued that this is not possible under the present assumptions. The conclusion follows from a standard compactness argument based on the stability of the first eigenvalue of the Laplacian under (measured) Gromov-Hausdorff convergence with lower Ricci curvature bounds and on the compactness of the set of possible cross-sections of blow-downs of (M, g) with respect to the same topology (see, e.g., [3, Theorem 1.9.4]).

Step 2. Let $V_i \to \infty$ be a sequence of volumes and let $E_i \subset M$ be isoperimetric sets with $\operatorname{vol}(E_i) = V_i$. Recall the expression for r_V that was introduced in (3.2). We claim that with the metrics induced by the rescaled ambient spaces $(M^n, r_{V_i}^{-1}d)$ the boundaries ∂E_i converge in the Gromov-Hausdorff sense to a cross-section of a blow-down of M, up to the extraction of a subsequence. Moreover, the induced metrics on ∂E_i have uniform two-sided sectional curvature bounds. Hence they converge in $C^{1,\alpha}$ for every $\alpha < 1$. The rest of this step will be aimed at establishing the Gromov-Hausdorff convergence with the induced metrics.

We first note that the uniform two-sided bounds for the sectional curvatures of the induced Riemannian metrics on ∂E_i (after rescaling) follow from the quadratic curvature decay of (M, g), the second estimate in (3.3), the convergence of the second fundamental forms from Proposition 3.3 and the Gauss equations.

Up to the extraction of a subsequence that we do not relabel, $(M^n, r_{V_i}^{-1}d, p)$ converge in the pointed Gromov-Hausdorff sense to a blow-down cone C(N) of (M, g). Moreover, the convergence is $C^{1,\alpha}$ for every $\alpha < 1$ away from the vertex of the cone. We claim that ∂E_i with the induced Riemannian metrics converge to the cross-section (N, g_N) in the Gromov-Hausdorff sense. We note that the (n-1)-dimensional measures of ∂E_i , i.e., their perimeters, computed with respect to the rescaled metrics as above, converge to $\mathcal{H}^{n-1}(N)$ by [7, Theorem 4.23]. Moreover, from the forthcoming Lemma 3.6 we infer that the boundaries ∂E_i with the induced metrics have uniformly bounded diameters (after rescaling) and injectivity radii uniformly bounded away from 0. Thanks to the discussion above they also have uniform twosided bounds on the sectional curvature. In particular, up to the extraction of a subsequence that we do not relabel, by Cheeger's precompactness theorem they converge in the Gromov-Hausdorff sense to a Riemannian manifold (N', g') where g' is a $C^{1,\alpha}$ Riemannian metric for every $\alpha < 1$, and the convergence is $C^{1,\beta}$ for every $\beta < 1$. We claim that (N', g') is isometric to (N, g_N) . Thanks to the Hausdorff convergence of ∂E_i to N from Lemma 3.2 we can find a 1-Lipschitz surjective map $F : (N', g') \to (N, g_N)$. Moreover, we already argued above that

(3.7)
$$\mathcal{H}^{n-1}(N') = \lim_{i \to \infty} \mathcal{H}^{n-1}(\partial E_i) = \mathcal{H}^{n-1}(N) \,.$$

Above, it is understood that $\mathcal{H}^{n-1}(\partial E_i)$ is computed with respect to the rescaled metric on M. The well-known Lipschitz volume rigidity in Riemannian geometry (see e.g. [18, Lemma 9.1] and references therein) guarantees that F is an isometry. Therefore ∂E_i with the induced length distances Gromov-Hausdorff converge to N thus completing the proof of the claim.

Step 3. We claim that there exist $\varepsilon > 0$ and $V_0 > 0$ such that if $V > V_0$ and $V \in \mathcal{G}_{\varepsilon}$, where the set $\mathcal{G}_{\varepsilon}$ was introduced in the statement of Proposition 3.4, then the first eigenvalue (with respect to volume preserving variations) of the Jacobi operator on the boundary of each isoperimetric set $E_V \subset M$ with $\operatorname{vol}(E_V) = V$ is at least $\varepsilon_M V^{-2/n}$, where ε_M has been introduced in Step 1 above. In particular, E_V has strictly volume preserving stable boundary. By Proposition 3.4 the proof of the claim will complete the proof of Theorem 3.1.

We argue by contradiction, relying on the convergence and stability of Sobolev functions defined along Gromov-Hausdorff converging sequences of manifolds, see [31, 36, 3] for the relevant background. If the claim does not hold then there exists a sequence $V_i \to \infty$ and isoperimetric sets $E_i \subset M$ with $\operatorname{vol}(E_i) = V_i$ such that

(3.8)
$$\lim_{i \to \infty} V_i^{\frac{2}{n}} \oint_{\partial E_i} \operatorname{Ric}(\nu_{\partial E_i}, \nu_{\partial E_i}) = 0,$$

and there are smooth functions $f_i : \partial E_i \to \mathbb{R}$ with f_i not identically vanishing such that

(3.9)
$$\int_{\partial E_i} f_i \mathrm{d}P_{\partial E_i} = 0$$

and

(3.10)
$$\int_{\partial E_i} |\nabla_{\partial E_i} f_i|^2 \mathrm{d}P_{\partial E_i} \leq \int_{\partial E_i} \left[\mathrm{Ric}(\nu_{\partial E_i}, \nu_{\partial E_i}) + |\mathrm{II}_{\partial E_i}|^2 + \frac{3}{2} \varepsilon_M \right] f_i^2 \mathrm{d}P_{\partial E_i} \,,$$

for each $i \in \mathbb{N}$. Note that both (3.8), (3.9), and (3.10) are scale-invariant. We let $\Sigma_i := \partial E_i$ endowed with the metric g_i induced by the embedding into $(M^n, r_{V_i}^{-1}d)$ and the induced Riemannian (n-1)-dimensional volume vol_i. By Step 2, $(\Sigma_i, g_i, \text{vol}_i)$ converge in $C^{1,\alpha}$ for every $\alpha < 1$ to (N, g_N, vol_N) , up to the extraction of a subsequence that we do not relabel. Here N denotes the cross-section of a blow-down C(N) of (M, g). The conditions (3.9) and (3.10) can be rephrased by saying that there exists a sequence of smooth functions $\overline{f_i}: \Sigma_i \to \mathbb{R}$ such that

(3.11)
$$\int_{\Sigma_i} \bar{f}_i \mathrm{d} \operatorname{vol}_i = 0, \qquad \int_{\Sigma_i} \bar{f}_i^2 \mathrm{d} \operatorname{vol}_i = 1,$$

and

(3.12)
$$\int_{\Sigma_i} |\nabla \bar{f}_i|^2 \mathrm{d} \operatorname{vol}_i \le \int_{\Sigma_i} h_i \bar{f}_i^2 \mathrm{d} \operatorname{vol}_i$$

where we set

(3.13)
$$h_i := \operatorname{Ric}(\nu_{\Sigma_i}, \nu_{\Sigma_i}) + |\operatorname{II}_{\Sigma_i}|^2 + \frac{3}{2}\varepsilon_M$$

for each $i \in \mathbb{N}$.

By Proposition 3.3

 $(3.14) \qquad |\mathrm{II}_{\Sigma_i}|^2 \to (n-1),$

uniformly as $i \to \infty$. Hence, by (3.8) and the quadratic curvature decay (1.2), $h_i \to (n-1) + \frac{3}{2} \varepsilon_M$ in L^q as $i \to \infty$ for every $1 \le q < \infty$. By the quadratic curvature decay assumptions again and (3.14), the functions h_i are uniformly bounded with respect to i. Hence, by (3.11) and (3.12), the functions \bar{f}_i have uniformly bounded $H^{1,2}$ norms. By Step 2, a (p, 2)-Sobolev inequality with uniform constants independent of $i \in \mathbb{N}$ holds along the sequence (Σ_i, g_i) , for every p > 1, see also Lemma 3.6 below. Hence for any p > 1, up to the extraction of a subsequence that we do not relabel, the functions \bar{f}_i converge in L^p to a function $\bar{f}: N \to \mathbb{R}$. In particular, if we choose p > 2, then by (3.11) we get that

(3.15)
$$\int_{N} \bar{f} \mathrm{d} \operatorname{vol}_{N} = 0, \quad \int_{N} \bar{f}^{2} \mathrm{d} \operatorname{vol}_{N} = 1$$

Moreover, the combination of the L^q convergence of $h_i \to (n-1)$ with the L^p convergence of $\bar{f}_i \to \bar{f}$ yields that

(3.16)
$$\int_{\Sigma_i} h_i \bar{f}_i^2 \mathrm{d} \operatorname{vol}_i \to (n-1) \int_N \bar{f}^2 \mathrm{d} \operatorname{vol}_N = n-1 \quad \text{as } i \to \infty.$$

Taking into account the lower semicontinuity of the Dirichlet energy under L^2 convergence we get from (3.12) that

(3.17)
$$\int_{N} |\nabla \bar{f}|^2 \mathrm{d} \operatorname{vol}_N \le n - 1 + \frac{3}{2} \varepsilon_M$$

This is a contradiction to the uniform spectral gap estimate that was obtained in Step 1. \Box

Remark 3.5. Regarding the argument in Step 1 above, we note that in the present setting, where two-sided uniform sectional curvature bounds are in force, the set of cross-sections is compact in the $C^{1,\alpha}$ topology and it is much more elementary to check that the spectrum of the Laplacian is stable in this case, see [53, Chapter 10] for the relevant background.

Proposition 3.3 is a consequence of the effective regularity theory for (almost) minimizers of the perimeter under two-sided curvature bounds that is discussed in Appendix A and Lemma 3.2.

Proof of Proposition 3.3. Arguing by contradiction, it is enough to prove the following statement. Let $V_i \to \infty$, and let \tilde{E}_i be isoperimetric sets with volume V_i . Then, up to subsequences,

(3.18)
$$\lim_{i \to \infty} \sup_{x \in \partial \widetilde{E}_i} \left| V_i^{\frac{2}{n}} \left| \mathrm{II}_{\partial \widetilde{E}_i} \right|^2 (x) - (n-1) (\omega_n \mathrm{AVR}(M))^{\frac{2}{n}} \right| = 0.$$

Denote $r_i := (\operatorname{AVR}(M)\omega_n)^{-\frac{1}{n}}V_i^{\frac{1}{n}}$, and fix $0 < \alpha' < \alpha < 1$. Let E_i denote the set \widetilde{E}_i considered in the rescaled manifold $M_i := (M^n, r_i^{-1}d_g, p)$. Up to subsequences, let (C(N), o) be the asymptotic cone that is the limit of M_i in the pointed Gromov-Hausdorff sense, and in the $C^{1,\alpha}$ -sense outside the tip o. Let g^C be the limit metric metric on C(N), and let g_i denote the rescaled metric on M_i . We aim at proving the following assertion.

Claim. There are $\rho, \overline{C} > 0$ such that for every $p \in \partial B_1^{C(N)}(o)$ there exists a chart $\varphi : B_\rho(0^n) \to C(N)$ centered at p such that for every $\partial E_i \ni p_i \to p$ the following holds up to subsequences in i. There exist open sets $B_{\rho/2}(p) \subset \varphi(B_\rho(0^n)) \subset U_i \subset C(N)$, and $B_{\rho/2}^{M_i}(p_i) \subset V_i \subset M_i$, and diffeomorphisms $F_i : U_i \to V_i$ with $F_i(p) = p_i$ for which:

(1) It holds

(3.19)
$$\|\varphi^* g^C - (F_i \circ \varphi)^* g_i\|_{C^{1,\alpha}(B_{\rho/2}(0^n))} \to 0.$$

(2) There exist $u, u_i : B_{\rho/2}(0^{n-1}) \to \mathbb{R}$ such that

(3.20)
$$\varphi^{-1}(\partial B_1^{C(N)}(o)) \cap B_{\rho/2}(0^n) = \{(x, u(x)) : x \in B_{\rho/2}(0^{n-1})\} \cap B_{\rho/2}(0^n),$$

(3.21)
$$(F_i \circ \varphi)^{-1}(\partial E_i) \cap B_{\rho/2}(0^n) = \{(x, u_i(x)) : x \in B_{\rho/2}(0^{n-1})\} \cap B_{\rho/2}(0^n),$$

(3.22)
$$\|u_i\|_{C^{2,\alpha}(B_{\rho/4}(0^{n-1}))} \le \overline{C} \quad \text{for every } i \in \mathbb{N} ,$$

and

(3.23)
$$u_i \to u \quad \text{in } C^{2,\alpha'}(B_{\rho/8}(0^{n-1}))$$

In particular,

(3.24)
$$\lim_{i \to \infty} \sup_{x \in \partial E_i} ||\mathrm{II}_{\partial E_i}|(x) - (n-1)| = 0.$$

The proof the present proposition will be completed after the proof of the Claim. Indeed, (3.24) is equivalent to (3.18) after scaling.

In order to prove the **Claim** we rely on the results in Appendix A. Item (1) comes from the $C^{1,\alpha}$ -convergence of M_i to C(N) outside the tip o. Moreover, (3.21), and (3.22) in Item (2) come from (the proof of) Theorem A.6, and Allard's result Theorem A.5, see in particular (A.21). Recall that Theorem A.6 can be applied in the setting of the present Proposition 3.3 due to Remark A.7. Finally (3.22) in item (2) implies precompactness in $C^{2,\alpha'}(B_{\rho/8}(0^{n-1}))$, so that $u_i \to \tilde{u}$ up to subsequences in $C^{2,\alpha'}(B_{\rho/8}(0^{n-1}))$. Since $\partial E_i \to \partial B_1^{C(N)}(o)$ in the Hausdorff sense, we infer that $\tilde{u} = u$, and thus (3.23) is proved. Finally, (3.24) is a direct consequence of (3.19), and (3.23).

The next Lemma is a consequence of the results in Appendix A, and it was already used in the proof of Theorem 3.1 above.

Lemma 3.6. Let (M^n, g) be a smooth complete Riemannian manifold with Ric ≥ 0 , Euclidean volume growth (1.1), and quadratic Riemann curvature decay (1.2). There exists $V_0, C > 0$ such that the following holds. Let E_V be an isoperimetric of volume $V > V_0$, and let $\Sigma := \partial E_V$. Let g_{Σ} be the Riemannian metric induced by g on Σ . Then

(3.25)
$$\operatorname{diam}(\Sigma, g_{\Sigma}) \le CV^{1/n}, \qquad \operatorname{inj}(\Sigma, g_{\Sigma}) \ge CV^{1/n}.$$

Proof. It is enough to prove the following **Claim**, and then argue by contradiction.

Claim. Let $V_i \to \infty$, and let $\widetilde{E}_i \subset M$ be an isoperimetric set with volume V_i . Denote $r_i := (\operatorname{AVR}(M)\omega_n)^{-\frac{1}{n}}V_i^{\frac{1}{n}}$. Let E_i denote the set \widetilde{E}_i considered in the rescaled manifold $M_i := (M^n, d_i := r_i^{-1}d_g, p)$. Set $\Sigma_i := \partial E_i$, and denote by g_i , d_i the metric and distance on M_i , and by g_{Σ_i} the induced metric on Σ_i . Then

$$\limsup_{i \to \infty} \operatorname{diam}(\Sigma_i, g_{\Sigma_i}) < \infty, \qquad \liminf_{i \to \infty} \operatorname{inj}(\Sigma_i, g_{\Sigma_i}) > 0$$

By (3.19), (3.21), and (3.23) in the proof of Proposition 3.3, the intrinsic and the restricted distances on Σ_i are locally equivalent. I.e., there exist constants $\rho, C > 0$ such that, for all *i* large enough (up to subsequences) we have

(3.26)
$$\forall p \in \Sigma_i, \quad \forall q, r \in B^{M_i}_\rho(p) \cap \Sigma_i, \quad |\mathbf{d}_{\Sigma_i}(q, r) - \mathbf{d}_i(q, r)| \le C \mathbf{d}_i(q, r) \,.$$

Thus, joining (3.26) with the density estimates in [7, Corollary 4.15], we get that, possibly taking a smaller ρ , there exists a constant $\xi > 0$ such that, for all *i* large enough,

(3.27)
$$\forall p \in \Sigma_i, \operatorname{vol}_{\Sigma_i}(B_{\rho}^{\Sigma_i}(p)) \ge \xi > 0$$

Above $B_{\rho}^{\Sigma_i}$ denotes the ball in the metric g_{Σ_i} . By (3.24), the quadratic Riemann curvature decay (1.2), and the Gauss equation, there exists a constant K > 0 such that, for *i* large enough,

 $(3.28) \qquad |\operatorname{Sec}_{\Sigma_i}| \le K.$

By (3.27), (3.28), and [53, Lemma 11.4.9] we obtain that

(3.29)
$$\liminf_{i \in \mathcal{I}} \operatorname{inj}(\Sigma_i, g_{\Sigma_i}) > 0.$$

Finally, observe $|\Sigma_i| \to |\partial B_1^{C(N)}(o)|$ by [7, Theorem 4.23], where $B_1^{C(N)}(o)$ is the unit ball centered at the tip of an asymptotic cone M_i is converging to, up to subsequences. Hence $|\Sigma_i|$ is uniformly bounded above. Joining the latter information with (3.29), and (3.28), a standard covering argument using Bishop–Gromov comparison theorem gives

$$\limsup_{i\to\infty} \operatorname{diam}(\Sigma_i, g_{\Sigma_i}) < \infty \,.$$

Hence the **Claim** is proved, and the proof is concluded.

As already mentioned, controlling the Ricci curvature term of the stability operator is the most delicate part of the argument for proving strict stability. One of the reasons is that, contrary to the other terms of the stability operator, the Ricci curvature term does not pass to the limit under blow-downs in general, and a careful volume selection argument is needed.

We note that a quadratic *lower* curvature decay in (1.2), combined with (1.1) is sufficient to guarantee that the Ricci curvature of (M^n, g) weakly converges to 0 in the almost radial directions when we consider any blow-down, see [41] for the precise statement. However, even under the stronger two-sided quadratic curvature decay assumption, this weak convergence does not localize to (all) boundaries of large isoperimetric sets.

To illustrate the idea of the proof of Proposition 3.4 it is helpful to introduce the notion of isoperimetric profile of (M^n, g) . We define the isoperimetric profile $I_M = I : [0, \infty) \to [0, \infty)$ by

(3.30)
$$I(v) := \inf\{P(E) : E \subset M, \operatorname{vol}(E) = v\}.$$

and let $J_M = J := I^{\frac{n}{n-1}}$ be the normalized isoperimetric profile. In [7] it was shown that if (M^n, g) is complete and satisfies Ric ≥ 0 , then J is a concave function, see also the earlier [45] where the same result was obtained under some additional assumptions. Moreover, if AVR(M) > 0, then

$$(3.31) J \sim J_{C(N)}$$

for large volumes, where $J_{C(N)}$ denotes the normalized isoperimetric profile of one of the blowdowns of (M^n, g) . Note that $J_{C(N)}$ is an affine function whose slope depends only on AVR(M)(equivalently, on vol(N), which is independent of the blow-down), see [47, 7].

For the proof of Proposition 3.4 we will reverse the usual argument for proving the concavity of J via the second variation formula, see [11, 12, 7]. The main idea is that J is concave and asymptotically affine. Hence its second derivative converges to 0 with a rate in an integral L^1 -sense as $V \to \infty$. In particular, it converges to 0 with the same rate on sets of volumes of scale invariantly larger and larger measure. For those volumes, we shall be able to control the Ricci curvature term of the stability operator.

The proof of the forthcoming Proposition 4.4 will require an effective version of the idea discussed above. We prefer to prove the weaker Proposition 3.4 separately in order to introduced the first main idea of the proof in a simplified case.

Proof of Proposition 3.4. By [7, Theorem 1.1], letting $J : (0, \infty) \to (0, \infty)$ be defined as $J := I^{\frac{n}{n-1}}$ as above, J is a concave function. By [5, Corollary 3.6] the right-derivative J'_+ (which is well-defined at every point by concavity) has a finite limit as $v \to \infty$. Denoting by J'' the second derivative, which is well-defined \mathcal{L}^1 -a.e., we get from elementary convex analysis that

(3.32)
$$\int_{r}^{2r} J''(v) \mathrm{d}v \to 0, \quad \text{as } r \to \infty.$$

For every $\delta > 0$ we let $\mathcal{F}_{\delta} \subset (0, \infty)$ be the set of points $v \in (0, \infty)$ such that J is twice differentiable at v and $0 \geq J''(v)v \geq -\delta$. Taking into account (3.32) and the nonpositivity of J'' it is elementary to infer that

(3.33)
$$\lim_{r \to \infty} \frac{\mathcal{L}^1(\mathcal{F}_\delta \cap (r, 2r))}{r} = 1,$$

for every $\delta > 0$.

Fix $\delta > 0$ and let us consider any volume $V \in \mathcal{F}_{\delta}$. Let $E \subset M$ be any isoperimetric region in M with $\operatorname{vol}(E) = V$, if there exist any. Let E_t be the *t*-enlargement of E for all t in a small neighbourhood of 0, i.e., $E_t := \{x \in M : d(x, E) < t\}$ if t > 0 and $E_t := \{x \in M : d(x, M \setminus E) > -t\}$ for $t \leq 0$. A standard argument in Geometric Measure Theory based on the first variation formula for the perimeter shows that it is possible to find a continuous function τ defined in a neighbourhood $(V - \mu, V + \mu) \ni V$ with $\tau(V) = 0$ and such that $\operatorname{vol}(E_{\tau(v)}) = v$ for each $v \in (V - \mu, V + \mu)$. We note that, by the very definition of the isoperimetric profile, it holds

(3.34)
$$\tilde{I}_V(v) := P(E_{\tau(v)}) \ge I(v) ,$$

for every $v \in (V - \mu, V + \mu)$. Moreover, setting $\tilde{J}_V := \tilde{I}_V^{\frac{n}{n-1}}$, we have

(3.35)
$$\frac{\mathrm{d}^2}{\mathrm{d}v^2}|_{v=V}\tilde{J}_V = \frac{n}{n-1}P(E)^{\frac{2-n}{n-1}} \left[\frac{H^2}{n-1} - \oint_{\partial E} \left(|\mathrm{II}_{\partial E}|^2 + \mathrm{Ric}(\nu_{\partial E}, \nu_{\partial E})\right)\right]$$

(3.36)
$$\leq -\frac{n}{n-1}P(E)^{\frac{2-n}{n-1}} \oint_{\partial E} \operatorname{Ric}(\nu_{\partial E}, \nu_{\partial E}).$$

Above, in order to pass from (3.35) to (3.36) we applied the Cauchy-Schwartz inequality to the (n-1) eigenvalues of $II_{\partial E}$. By (3.34) it readily follows that $\tilde{J}_V \geq J$ on $(V - \mu, V + \mu)$ and $\tilde{J}_V(V) = J(V)$. Hence, since $V \in \mathcal{F}_{\delta}$, we get

(3.37)
$$\frac{n}{n-1}P(E)^{\frac{2-n}{n-1}} \oint_{\partial E} \operatorname{Ric}(\nu_{\partial E}, \nu_{\partial E}) \leq \frac{\delta}{V}$$

If ∂E is smooth, the second derivative of J_V above makes sense in the classical sense. Otherwise, the present argument can be adapted to infer (3.37) even when E has singular boundary, by using the fact that the codimension of the singular part of ∂E is at least 8, and standard

arguments in Geometric Measure Theory, compare with [12]. We note that there exists a constant C > 1 depending only on the asymptotic volume ratio of M such that

(3.38)
$$\frac{1}{C}V^{\frac{n-1}{n}} \le P(E) \le CV^{\frac{n-1}{n}},$$

for any isoperimetric set $E \subset M$ with $\operatorname{vol}(E) = V$. Therefore, the statement follows from (3.37) up to choosing $\delta < \delta(\varepsilon)$ sufficiently small and setting $\mathcal{G}_{\varepsilon} := \mathcal{F}_{\delta(\varepsilon)}$.

4. Uniqueness for many volumes

The goal of this section is to prove the uniqueness part of Theorem 1.1. We state below the precise result for the ease of readers.

Theorem 4.1. Let (M^n, g) be a smooth complete Riemannian manifold with Ric ≥ 0 , Euclidean volume growth (1.1), and quadratic Riemann curvature decay (1.2). Assume that (M^n, g) is not isometric to \mathbb{R}^n . Then there exists a measurable set $\mathcal{G} \subset (0, \infty)$ such that

(4.1)
$$\lim_{r \to \infty} \frac{\mathcal{L}^1(\mathcal{G} \cap (r, 2r))}{r} = 1$$

and for any $V \in \mathcal{G}$ the isoperimetric set $E_V \subset M$ of volume V is unique.

The key idea will be to make effective the broad principle according to which strictly stable implies locally uniquely minimizing. As mentioned in the introduction, the proof of Theorem 4.1 is divided in a few steps. A first technical tool is the fact that the reach of isoperimetric boundaries of large volumes is uniformly bounded below in a scale-invariant form. We recall that for a bounded open subset $\Omega \subset M$ with smooth boundary in a Riemannian manifold (M,g), the reach of $\partial\Omega$ is the supremum of all positive numbers r such that for every point x_0 in $\{x \in M : d(x, \partial\Omega) < r\}$ there exists a unique $y_0 \in \partial\Omega$ such that $d(x_0, y_0) = d(x_0, \partial\Omega)$. In particular the signed distance function from Ω is smooth in the open tubular neighborhood of $\partial\Omega$ having size equal to the reach of $\partial\Omega$.

Lemma 4.2. Let (M^n, g) be a smooth complete Riemannian manifold with $\operatorname{Ric} \geq 0$, Euclidean volume growth (1.1), and quadratic Riemann curvature decay (1.2). Then there exists $\overline{V}, \overline{\rho} > 0$ such that for every $V \geq \overline{V}$ the reach of the boundary of every isoperimetric set $E_V \subset M$ of volume $\operatorname{vol}(E_V) = V$ is bounded from below by $\overline{\rho}V^{\frac{1}{n}}$.

Proof. After scaling, for a constant C depending on M, we can prove the claim for isoperimetric sets E having volume $CAVR(M)\omega_n$ that are $(\Lambda, 1)$ -minimizers (Definition A.2) for some fixed $\Lambda > 0$ in manifolds with a uniform bound on |Riem|, and such that (A.7), and (A.8) hold for a suitable choice of local chart at boundary points of E (see Remark A.7, Proposition A.4, and Theorem A.1). In particular Theorem A.5 can be applied at boundary points of E and (A.10) implies that (B.1) is satisfied for some ε_0 independent of the specific isoperimetric set. Taking into account also Proposition 3.3, the conclusion follows by Proposition B.1.

The first main step for the proof of Theorem 4.1 amounts to prove that for every sufficiently large volume away from a countable set, any two isoperimetric sets with that volume can be written one as a normal graph over the boundary of the other, with suitable scale-invariant bounds on the graphing function. The set of *bad* volumes consists of the set of non-differentiability points of the isoperimetric profile I of (M, g). The precise statement is as follows.

Proposition 4.3. Let (M^n, g) be a smooth complete Riemannian manifold with Ric ≥ 0 , Euclidean volume growth (1.1), and quadratic Riemann curvature decay (1.2). Then there exists a constant C > 0 such that the following holds.

For any $\varepsilon > 0$ there exists $V_0 = V_0(\varepsilon) > 0$ such that for every $V > V_0$ away from a countable set the following holds. If $E_V, E'_V \subset M$ are two distinct isoperimetric regions with

 $\operatorname{vol}(E_V) = \operatorname{vol}(E'_V) = V$, then $\partial E'_V$ is the normal graph over ∂E_V of a function $w : \partial E_V \to \mathbb{R}$ satisfying:

(4.2)
$$\|w\|_{\infty} \le \varepsilon V^{\frac{1}{n}}, \quad \|\nabla w\|_{\infty} \le \varepsilon,$$

(4.3)
$$\|w\|_{\infty} \le \mathcal{C}V^{-\frac{n-1}{2n}} \|w\|_{L^2(\partial E_V)},$$

(4.4)
$$\|\nabla w\|_{L^2(\partial E_V)} \le \mathcal{C}V^{-\frac{1}{n}} \|w\|_{L^2(\partial E_V)}$$

In the second step of the proof, we are going to effectively control the average of the Ricci curvature in *normal* directions on tubular neighbourhoods of any size for isoperimetric sets of most volumes. The precise statement follows:

Proposition 4.4. Let (M^n, g) be a smooth complete Riemannian manifold with Ric ≥ 0 , Euclidean volume growth (1.1), and quadratic Riemann curvature decay (1.2). Then for every $\varepsilon > 0$ there exists a set $\mathcal{G}_{\varepsilon} \subset (0, \infty)$ such that

(4.5)
$$\lim_{r \to \infty} \frac{\mathcal{L}^1(\mathcal{G}_{\varepsilon} \cap (r, 2r))}{r} = 1,$$

and for every volume $V \in \mathcal{G}_{\varepsilon}$ and every isoperimetric set $E_V \subset M$ with volume V it holds

(4.6)
$$\sup_{0 < r < C_0 V^{\frac{1}{n}}} V^{\frac{2}{n}} \oint_{B_r(\partial E_V)} \operatorname{Ric}(\nabla d_{\partial E_V}, \nabla d_{\partial E_V}) \mathrm{d} \operatorname{vol} \le \varepsilon$$

In order to complete the proof of Theorem 4.1 we are going to effectively Taylor expand the area functional to compare the perimeter of two distinct isoperimetric sets E_V and E'_V with the same volume V. The identity $Per(E_V) = Per(E'_V)$ will result in a contradiction provided that the volume V is suitably chosen. In the next section we discuss the details by taking for granted Proposition 4.3 and Proposition 4.4, whose proofs are deferred to a subsequent section.

4.1. Proof of Theorem 4.1 given Proposition 4.3 and Proposition 4.4. Let $\varepsilon > 0$ and let $\mathcal{G}_{\varepsilon}$ be given by Proposition 4.4. Without loss of generality we can assume that the conclusion of Theorem 3.1 holds for all volumes $V \in \mathcal{G}_{\varepsilon}$. We claim that if ε is small enough, then the theorem follows taking $\mathcal{G} = \mathcal{G}_{\varepsilon} \cap (v_0, \infty)$ for some $v_0 > 0$.

We will denote by $\mathcal{C} > 0$ any constant depending on M only, independent of ε , that may change from line to line. Up to removing a countable (and hence negligible) set from $\mathcal{G}_{\varepsilon}$, we can assume that the isoperimetric profile is differentiable at any volume in $\mathcal{G}_{\varepsilon}$ and that Proposition 4.3 can be applied at any volume in $\mathcal{G}_{\varepsilon} \cap (V_0(\varepsilon), \infty)$. The strategy will be to argue by contradiction. Assume that there exists a sequence $V_i \in \mathcal{G}_{\varepsilon}$ such that $V_i \nearrow \infty$ and such that there exist two distinct isoperimetric sets E_{V_i}, E'_{V_i} of volume V_i . There will be three main steps. In **Step 1** we are going to rely on Proposition 4.3 to infer that $\partial E'_{V_i}$ is a graph over ∂E_V , and to use the assumption that $\operatorname{vol}(E_{V_i}) = \operatorname{vol}(E'_{V_i}) = V_i$ to obtain an effective estimate on the average of the graphing function. In **Step 2** we are going to effectively Taylor expand the perimeter in a neighbourhood of ∂E_{V_i} in terms of the stability operator and an error term depending on the ambient Ricci curvature. In **Step 3** we will conclude with the help of Proposition 4.4, whose aim is exactly to control the potentially large Ricci curvature term.

Step 1. Denote $r_i := (\operatorname{AVR}(M)\omega_n)^{-\frac{1}{n}}V_i^{\frac{1}{n}}$, and $M_{V_i} := (M^n, r_i^{-1}d_g, p)$. Let g_i be the rescaled metric on M_{V_i} . After rescaling and up to extracting a subsequence, we can assume that M_{V_i} converges in Gromov-Hausdorff sense to an asymptotic cone C(N) with $C^{1,\alpha}$ -cross section (N, g_N) and the sets E_i, E'_i corresponding to E_{V_i}, E'_{V_i} , respectively, converge to the unit ball of C(N) centered at the tip in L^1 -strong. For large i, by Proposition 4.3 there exists a sequence of functions $w_i \in C^{\infty}(\partial E_i)$ that parametrize $\partial E'_i$ over ∂E_i as a normal graph, i.e.,

$$\partial E'_i = \{ \exp_x(w_i(x)\nu_{E_i}(x)) : x \in \partial E_i \} .$$

We claim that for any $i \in \mathbb{N}$ sufficiently large it holds

(4.7)
$$\left| \int_{\partial E_i} w_i - \frac{1}{2} H_{\partial E_i} \int_{\partial E_i} w_i^2 \right| \le \varepsilon \, \mathcal{C} \int_{\partial E_i} w_i^2 \, .$$

The effect of (4.7) is to tell that the graphing functions w_i have almost zero average on ∂E_i , in a very quantitative way. This will be key later when it comes to reaching a contradiction to the strict volume preserving stability of ∂E_i .

From Proposition B.1, we know that the reach of ∂E_i is bounded below by a constant $\overline{\rho} > 0$ independent of *i*. Hence, for any large *i*, we can fix an open neighborhood U_i of ∂E_i such that $\partial E'_i \subset \{x : d(x, \partial E_i) < \overline{\rho}/2\} \subseteq U_i$, and such that the metric g_i on U_i can be written in Fermi coordinates with respect to ∂E_i . Namely,

(4.8)
$$(U_i, g_i) \simeq \left(\left(-\overline{\rho}, \overline{\rho} \right) \times \partial E_i, \mathrm{d}z^2 + g_z \right),$$

where g_z is the pull-back on ∂E_i via exponential map of the induced metric on the parallel hypersurface $\{\exp_x(z\nu_{E_i}(x)) : x \in \partial E_i\}$, for $z \in (-\overline{\rho}, \overline{\rho})$. On the manifold $((-\overline{\rho}, \overline{\rho}) \times \partial E_i, dz^2 + g_z)$ a normal graph over ∂E_i defined by a function $u \in C^{\infty}(\partial E_i)$ with $|u| < \overline{\rho}$ is a set of the form $S = \{(u(x), x) : x \in \partial E_i\}$, and the upward unit normal of S can be written as the vector field

(4.9)
$$\frac{\partial_z - \nabla^u u}{\left(1 + |\nabla^u u|_u^2\right)^{\frac{1}{2}}},$$

where ∇^u and $|\cdot|_u$ denote gradient and norm with respect to the metric g_u on ∂E_i given by $(g_u)_p := (g_{u(p)})_p$ at any $p \in \partial E_i$. Also, integration with respect to the volume form on the graph S of u can be written in terms of the volume form with respect to g_u as

(4.10)
$$\int f(z,x) \,\mathrm{d}\,\mathrm{vol}_S = \int f(u(x),x) \left(1 + |\nabla^u u|_u^2\right)^{\frac{1}{2}} \,\mathrm{d}\,\mathrm{vol}_{g_u}\,,$$

for any continuous function f on S.

Recall that $\partial_z \sqrt{\det g_z} = -\sqrt{\det g_z} \Delta d^s$ in any local chart on ∂E_i and $H_{\partial E_i}(x) = \Delta d^s(0, x)$, where d^s is the signed distance function from E_i (d^s is defined to be positive outside E_i , and negative inside E_i). Then for any $x \in \partial E_i$ working in a local chart on ∂E_i such that $\sqrt{\det g_0}(x) = 1$, we can expand

(4.11)

$$\sqrt{\det g_z}(z,x) = 1 - \int_0^z \sqrt{\det g_z}(s,x) \,\Delta \mathrm{d}^s(s,x) \,\mathrm{d}s$$

$$= 1 - H_{\partial E_i} z - \int_0^z \int_0^s |\nabla^2 \mathrm{d}^s|^2(t,x) + \operatorname{Ric}_{(t,x)}(\partial_z,\partial_z) \,\mathrm{d}t \,\mathrm{d}s + \int_0^z \int_0^s \left(\sqrt{\det g_z}(s,x) - 1\right) \,\Delta \mathrm{d}^s(s,x) \,\mathrm{d}s \,.$$

Note that $|II_{\partial E_i}|$, $H_{\partial E_i}$ are uniformly bounded with respect to *i* by Proposition 3.3. Exploiting the uniform scale invariant Riemann curvature bounds and the Riccati equation for the evolution of $\Delta d^s(s, x)$ and $\nabla^2 d^s(s, x)$ with respect to *s* (see also the proof of Lemma 4.6 below), it follows that $|\nabla^2 d^s|^2(t, x) + \operatorname{Ric}_{(z,x)}(\partial_z, \partial_z) + |\Delta d^s(t, x)| \leq C$ for any $x \in \partial E_i$ and any $|t| < \overline{\rho}$. Hence $|\sqrt{\det g_z(z, x) - 1}| \leq C|z|$ and

(4.12)
$$\left|\sqrt{\det g_z}(z,x) - 1 + H_{\partial E_i}z\right| \le \mathcal{C}z^2$$

Thus

(4.13)
$$\left| \frac{\mathrm{d} \operatorname{vol}_{g_z}}{\mathrm{d} \operatorname{vol}_{g_0}} - (1 - H_{\partial E_i} z) \right| \le \mathcal{C} z^2 \,,$$

for any $z \in (-\overline{\rho}, \overline{\rho})$, where $\frac{\mathrm{d} \operatorname{vol}_{g_z}}{\mathrm{d} \operatorname{vol}_{g_0}}$ is the Radon–Nikodým derivative of vol_{g_z} with respect to vol_{g_0} .

We proceed with the proof of the claimed (4.7). Let *i* be fixed and let $\{E_t\}_{t \in [0,1]}$ be the one-parameter family of sets having boundary

(4.14)
$$\left\{ \exp_x(tw_i(x)\nu_{E_i}(x)) : x \in \partial E_i \right\},$$

such that $E_0 = E_i$, $E_1 = E'_i$. Denote $V(t) := \operatorname{vol}_i(E_t)$. Exploiting (4.9) and (4.10) with $u = tw_i$ we can compute

(4.15)
$$V'(t) = \int_{\partial E_i} g_i \left(-\frac{\partial_z - t \nabla^{tw_i} w_i}{\left(1 + t^2 |\nabla^{tw_i} w_i|_{tw_i}^2\right)^{\frac{1}{2}}}, w_i \partial_z \right) \left(1 + t^2 |\nabla^{tw_i} w_i|_{tw_i}^2\right)^{\frac{1}{2}} \mathrm{d} \operatorname{vol}_{g_{tw_i}} \\ = -\int_{\partial E_i} w_i \operatorname{d} \operatorname{vol}_{g_{tw_i}},$$

where g_i here equals $dz^2 + g_z$ and denotes the metric tensor on M_{V_i} in the Fermi coordinates chart. Hence

(4.16)

$$0 = |E_i| - |E'_i| = -\int_0^1 V'(t) \, \mathrm{d}t = \int_0^1 \int_{\partial E_i} w_i \, \mathrm{d}\operatorname{vol}_{g_{tw_i}} \, \mathrm{d}t$$

$$= \int_0^1 \int_{\partial E_i} w_i \left(\frac{\mathrm{d}\operatorname{vol}_{g_{tw_i}}}{\mathrm{d}\operatorname{vol}_{g_0}} - (1 - H_{\partial E_i} t w_i) + (1 - H_{\partial E_i} t w_i) \right) \, \mathrm{d}\operatorname{vol}_{g_0} \, \mathrm{d}t$$

$$= \int_{\partial E_i} w_i \, \mathrm{d}\operatorname{vol}_{g_0} - \frac{1}{2} H_{\partial E_i} \int_{\partial E_i} w_i^2 \, \mathrm{d}\operatorname{vol}_{g_0} + \int_0^1 \int_{\partial E_i} w_i \left(\frac{\mathrm{d}\operatorname{vol}_{g_{tw_i}}}{\mathrm{d}\operatorname{vol}_{g_0}} - (1 - H_{\partial E_i} t w_i) \right) \, \mathrm{d}\operatorname{vol}_{g_0} \, \mathrm{d}t \, .$$

Therefore

and (4.7) follows.

Step 2. The goal of this second step is to prove that for every sufficiently large $i \in \mathbb{N}$ it holds

$$(4.18) \qquad \frac{1}{2} \int_{\partial E_i} |\nabla w_i|^2 - |\mathrm{II}_{\partial E_i}|^2 w_i^2 \,\mathrm{d}P_{\partial E_i} \le \varepsilon \,\mathcal{C} \int_{\partial E_i} w_i^2 \,\mathrm{d}P_{\partial E_i} + \int_{\partial E_i} \int_0^{w_i(x)} \int_0^z \operatorname{Ric}_{(s,x)}(\partial_z, \partial_z) \,\mathrm{d}s \,\mathrm{d}z \,\mathrm{d}\operatorname{vol}_{g_0} dz \,\mathrm{d}z \,\mathrm{d}$$

The estimate will be obtained by applying the Gauss–Green theorem to the gradient of the signed distance function from ∂E_i in the region bounded between ∂E_i and $\partial E'_i$, carefully expanding all the terms and exploiting the assumption that $P(E_i) = P(E'_i)$.

We start from the Gauss-Green identity

(4.20)
$$\int_{E'_i \Delta E_i} \Delta d_{\partial E_i} = \int_{\partial E'_i} g_i (\nabla d^s, -\nu_{E'_i}) \, dP_{\partial E'_i} - P(E_i) \,,$$

where $\nu_{E'_i}$ denotes the inner normal of E'_i and d^s is the signed distance from E_i .

Exploiting Fermi coordinates around ∂E_i again, and recalling (4.9) and (4.2), for ε small enough we can compute

$$\begin{split} &\int_{\partial E'_{i}} g_{i}(\nabla d^{s}, -\nu_{E'_{i}}) \, \mathrm{d}P_{\partial E'_{i}} = \int_{\partial E'_{i}} \frac{1}{(1+|\nabla^{w_{i}}w_{i}|^{2}_{w_{i}}(x))^{1/2}} \, \mathrm{d}P_{\partial E'_{i}}(w_{i}(x), x) \\ &\leq \int_{\partial E'_{i}} 1 - \frac{1}{2} |\nabla^{w_{i}}w_{i}|^{2}_{w_{i}}(x) + 2 |\nabla^{w_{i}}w_{i}|^{4}_{w_{i}}(x) \, \mathrm{d}P_{\partial E'_{i}}(w_{i}(x), x) \\ \stackrel{(4.10)}{=} P(E'_{i}) \\ &\quad + \int_{\partial E_{i}} \left(-\frac{1}{2} |\nabla^{w_{i}}w_{i}|^{2}_{w_{i}}(x) + 2 |\nabla^{w_{i}}w_{i}|^{4}_{w_{i}}(x) \right) \left(1 + |\nabla^{w_{i}}w_{i}|^{2}_{w_{i}}(x) \right)^{\frac{1}{2}} \frac{\mathrm{d}\operatorname{vol}_{g_{w_{i}}}}{\mathrm{d}\operatorname{vol}_{g_{0}}} \, \mathrm{d}\operatorname{vol}_{g_{0}}(x) \, . \end{split}$$

Expanding $|\cdot|_z$ with respect to z, it can be checked that $(1 - \mathcal{C}|w_i|)|\nabla w_i|^2 \leq |\nabla^{w_i}w_i|_{w_i}^2 \leq (1 + \mathcal{C}|w_i|)|\nabla w_i|^2$, where $|\nabla w_i|^2$ is understood to be computed with respect to g_0 (see the proof of Lemma 4.6, and in particular the second of (4.39)). Hence, (4.2), (4.4), and (4.13) imply

(4.21)

$$\int_{\partial E'_{i}} g_{i}(\nabla d^{s}, -\nu_{E'_{i}}) dP_{\partial E'_{i}} \leq P(E'_{i}) - \frac{1}{2} \int_{\partial E_{i}} |\nabla w_{i}|^{2} dP_{\partial E_{i}} + \varepsilon \mathcal{C} \int_{\partial E_{i}} |\nabla w_{i}|^{2} dP_{\partial E_{i}} + \varepsilon \mathcal{C} \int_{\partial E_{i}} |\nabla w_{i}|^{4} dP_{\partial E_{i}} \\ \leq P(E'_{i}) - \frac{1}{2} \int_{\partial E_{i}} |\nabla w_{i}|^{2} dP_{\partial E_{i}} + \varepsilon \mathcal{C} \int_{\partial E_{i}} w_{i}^{2} dP_{\partial E_{i}}.$$

Since $P(E_i) = P(E'_i)$, (4.20) is equivalent to

(4.22)
$$\int_{E_i'\Delta E_i} \Delta d_{\partial E_i} \leq -\frac{1}{2} \int_{\partial E_i} |\nabla w_i|^2 \, \mathrm{d}P_{\partial E_i} + \varepsilon \, \mathcal{C} \int_{\partial E_i} w_i^2 \, \mathrm{d}P_{\partial E_i} \, .$$

By the coarea formula we can also compute

$$(4.23)$$

$$\int_{E'_{i}\Delta E_{i}} \Delta d_{\partial E_{i}} = \int_{-\overline{\rho}}^{\rho} \int_{\{x\in\partial E_{i}:(z,x)\in E'_{i}\Delta E_{i}\}} \Delta d_{\partial E_{i}}(z,x) \, d\operatorname{vol}_{g_{z}} \, dz$$

$$= \int_{-\overline{\rho}}^{\overline{\rho}} \int_{\{w_{i}>0\}} \chi_{\{w_{i}>z\}}(x) \Delta d_{\partial E_{i}}(z,x) \frac{\operatorname{d}\operatorname{vol}_{g_{z}}}{\operatorname{d}\operatorname{vol}_{g_{0}}} \, d\operatorname{vol}_{g_{0}} \, dz +$$

$$+ \int_{-\overline{\rho}}^{\overline{\rho}} \int_{\{w_{i}<0\}} \chi_{\{w_{i}>z\}}(x) \Delta d_{\partial E_{i}}(z,x) \frac{\operatorname{d}\operatorname{vol}_{g_{z}}}{\operatorname{d}\operatorname{vol}_{g_{0}}} \, d\operatorname{vol}_{g_{0}} \, dz$$

$$= \int_{\{w_{i}>0\}} \int_{-\overline{\rho}}^{\overline{\rho}} \chi_{(0,w_{i}(x))}(z) \Delta d_{\partial E_{i}}(z,x) \frac{\operatorname{d}\operatorname{vol}_{g_{z}}}{\operatorname{d}\operatorname{vol}_{g_{0}}} \, dz \, \operatorname{dvol}_{g_{0}} \, +$$

$$+ \int_{\{w_{i}<0\}} \int_{-\overline{\rho}}^{\overline{\rho}} \chi_{(w_{i}(x),0)}(z) \Delta d_{\partial E_{i}}(z,x) \frac{\operatorname{dvol}_{g_{z}}}{\operatorname{dvol}_{g_{0}}} \, dz \, \operatorname{dvol}_{g_{0}} \, dz \, d\operatorname{vol}_{g_{0}} \,$$

Recall the evolution equation

(4.24)
$$\Delta \mathrm{d}^{s}(z,x) = H_{\partial E_{i}}(x) + \int_{0}^{z} |\nabla^{2}\mathrm{d}^{s}|^{2}(s,x) + \operatorname{Ric}_{(s,x)}(\partial_{z},\partial_{z}) \,\mathrm{d}s.$$

Since

(4.25)
$$\left| |\nabla^2 \mathrm{d}^s|^2(s,x) - |\nabla^2 \mathrm{d}^s|^2(0,x) \right| \le \mathcal{C}|s|$$

thanks to the scale invariant Riemann curvature bounds, we thus find

$$\begin{split} &\int_{E_i'\Delta E_i} \Delta d_{\partial E_i} = \\ &= -\int_{\partial E_i} \int_0^{w_i(x)} \left(H_{\partial E_i} + \int_0^z |\nabla^2 d^s|^2(s,x) + \operatorname{Ric}_{(s,x)}(\partial_z,\partial_z) \, \mathrm{d}s \right) (1 - H_{\partial E_i}z) \, \mathrm{d}z \, \mathrm{d}\operatorname{vol}_{g_0} + \\ &\quad -\int_{\partial E_i} \int_0^{w_i(x)} \Delta d^s(z,x) \left(\frac{\mathrm{d}\operatorname{vol}_{g_z}}{\mathrm{d}\operatorname{vol}_{g_0}} - (1 - H_{\partial E_i}(x)z) \right) \, \mathrm{d}z \, \mathrm{d}\operatorname{vol}_{g_0} \end{split}$$

$$\end{split}$$

$$\begin{split} &\stackrel{(4.13)}{\geq} -\int_{\partial E_i} H_{\partial E_i}w_i - H_{\partial E_i}^2 \frac{w_i^2}{2} + |\operatorname{II}_{\partial E_i}|^2 \frac{w_i^2}{2} \, \mathrm{d}\operatorname{vol}_{g_0} + \\ &\quad -\int_{\partial E_i} \int_0^{w_i(x)} (1 - H_{\partial E_i}z) \int_0^z \operatorname{Ric}_{(s,x)}(\partial_z,\partial_z) \, \mathrm{d}s \, \mathrm{d}z \, \mathrm{d}\operatorname{vol}_{g_0} + \\ &\quad -\varepsilon \, \mathcal{C} \int_{\partial E_i} w_i^2 \, \mathrm{d}\operatorname{vol}_{g_0} \end{split}$$

$$\end{split}$$

$$\end{split}$$

$$\begin{split} \stackrel{(4.7)}{\geq} -\int_{\partial E_i} \frac{1}{2} |\operatorname{II}_{\partial E_i}|^2 w_i^2 \, \mathrm{d}\operatorname{vol}_{g_0} - \int_{\partial E_i} \int_0^{w_i(x)} \int_0^z \operatorname{Ric}_{(s,x)}(\partial_z,\partial_z) \, \mathrm{d}s \, \mathrm{d}z \, \mathrm{d}\operatorname{vol}_{g_0} + \\ &\quad -\varepsilon \, \mathcal{C} \int_{\partial E_i} w_i^2 \, \mathrm{d}\operatorname{vol}_{g_0} - \int_{\partial E_i} \int_0^{w_i(x)} \int_0^z \operatorname{Ric}_{(s,x)}(\partial_z,\partial_z) \, \mathrm{d}s \, \mathrm{d}z \, \mathrm{d}\operatorname{vol}_{g_0} + \\ &\quad -\varepsilon \, \mathcal{C} \int_{\partial E_i} w_i^2 \, \mathrm{d}\operatorname{vol}_{g_0} - \int_{\partial E_i} \int_0^{w_i(x)} \int_0^z \operatorname{Ric}_{(s,x)}(\partial_z,\partial_z) \, \mathrm{d}s \, \mathrm{d}z \, \mathrm{d}\operatorname{vol}_{g_0} + \\ &\quad -\varepsilon \, \mathcal{C} \int_{\partial E_i} w_i^2 \, \mathrm{d}\operatorname{vol}_{g_0} \, . \end{split}$$

Recalling (4.22) we get the claimed (4.18).

Step 3. In this last step we are going to reach a contradiction to the strict volume preserving stability of ∂E_i by estimating the Ricci curvature term in (4.19) with the help of Proposition 4.4.

We start by noting that

$$\int_{\partial E_i} \int_0^{w_i(x)} \int_0^z \operatorname{Ric}_{(s,x)}(\partial_z, \partial_z) \, \mathrm{d}s \, \mathrm{d}z \, \mathrm{d}\operatorname{vol}_{g_0} \le \|w_i\|_{\infty} \int_{-\|w_i\|_{\infty}}^{\|w_i\|_{\infty}} \int_{\partial E_i} \operatorname{Ric}_{(s,x)}(\partial_z, \partial_z) \, \mathrm{d}s \, \mathrm{d}\operatorname{vol}_{g_0} = \int_{-\|w_i\|_{\infty}}^{\|w_i\|_{\infty}} \int_{\partial E_i} \operatorname{Ric}_{(s,x)}(\partial_z, \partial_z) \, \mathrm{d}s \, \mathrm{d}\operatorname{vol}_{g_0} = \int_{-\|w_i\|_{\infty}}^{\|w_i\|_{\infty}} \int_{\partial E_i} \operatorname{Ric}_{(s,x)}(\partial_z, \partial_z) \, \mathrm{d}s \, \mathrm{d}\operatorname{vol}_{g_0} = \int_{-\|w_i\|_{\infty}}^{\|w_i\|_{\infty}} \int_{\partial E_i} \operatorname{Ric}_{(s,x)}(\partial_z, \partial_z) \, \mathrm{d}s \, \mathrm{d}\operatorname{vol}_{g_0} = \int_{-\|w_i\|_{\infty}}^{\|w_i\|_{\infty}} \int_{\partial E_i} \operatorname{Ric}_{(s,x)}(\partial_z, \partial_z) \, \mathrm{d}s \, \mathrm{d}\operatorname{vol}_{g_0} = \int_{-\|w_i\|_{\infty}}^{\|w_i\|_{\infty}} \int_{\partial E_i} \operatorname{Ric}_{(s,x)}(\partial_z, \partial_z) \, \mathrm{d}s \, \mathrm{d}\operatorname{vol}_{g_0} = \int_{-\|w_i\|_{\infty}}^{\|w_i\|_{\infty}} \int_{\partial E_i} \operatorname{Ric}_{(s,x)}(\partial_z, \partial_z) \, \mathrm{d}s \, \mathrm{d}\operatorname{vol}_{g_0} = \int_{-\|w_i\|_{\infty}}^{\|w_i\|_{\infty}} \int_{\partial E_i} \operatorname{Ric}_{(s,x)}(\partial_z, \partial_z) \, \mathrm{d}s \, \mathrm{d}\operatorname{vol}_{g_0} = \int_{-\|w_i\|_{\infty}}^{\|w_i\|_{\infty}} \int_{\partial E_i} \operatorname{Ric}_{(s,x)}(\partial_z, \partial_z) \, \mathrm{d}s \, \mathrm{d}\operatorname{vol}_{g_0} = \int_{-\|w_i\|_{\infty}}^{\|w_i\|_{\infty}} \int_{\partial E_i} \operatorname{Ric}_{(s,x)}(\partial_z, \partial_z) \, \mathrm{d}s \, \mathrm{d}\operatorname{vol}_{g_0} = \int_{-\|w_i\|_{\infty}}^{\|w_i\|_{\infty}} \int_{\partial E_i} \operatorname{Ric}_{(s,x)}(\partial_z, \partial_z) \, \mathrm{d}s \, \mathrm{d}\operatorname{vol}_{g_0} = \int_{-\|w_i\|_{\infty}}^{\|w_i\|_{\infty}} \int_{\partial E_i} \operatorname{Ric}_{(s,x)}(\partial_z, \partial_z) \, \mathrm{d}s \, \mathrm{d}\operatorname{vol}_{g_0} = \int_{-\|w_i\|_{\infty}}^{\|w_i\|_{\infty}} \int_{-\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}} \int_{-\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}^{\|w_i\|_{\infty}}$$

Rewriting the integral in (4.6) exploiting Fermi coordinates around ∂E_i and using (4.13) as we did above, it is readily checked that Proposition 4.4 implies

(4.26)
$$\int_{-\|w_i\|_{\infty}}^{\|w_i\|_{\infty}} \int_{\partial E_i} \operatorname{Ric}_{(s,x)}(\partial_z, \partial_z) \,\mathrm{d}s \,\mathrm{d}\operatorname{vol}_{g_0} \le \varepsilon \,\mathcal{C} \|w_i\|_{\infty} + \mathcal{C} \|w_i\|_{\infty}^2$$

Hence

(4.27)
$$\int_{\partial E_i} \int_0^{w_i(x)} \int_0^z \operatorname{Ric}_{(s,x)}(\partial_z, \partial_z) \,\mathrm{d}s \,\mathrm{d}z \,\mathrm{d}\operatorname{vol}_{g_0} \le \varepsilon \,\mathcal{C} \|w_i\|_{\infty}^2 \stackrel{(4.3)}{\le} \varepsilon \,\mathcal{C} \int_{\partial E_i} w_i^2 \,\mathrm{d}P_{\partial E_i}$$

Combining (4.18) and (4.27) we end up with

(4.28)
$$\int_{\partial E_i} |\nabla w_i|^2 - |\mathrm{II}_{\partial E_i}|^2 w_i^2 \,\mathrm{d}P_{\partial E_i} \le \varepsilon \,\mathcal{C} \int_{\partial E_i} w_i^2 \,\mathrm{d}P_{\partial E_i}$$

Let $u_i := w_i - \int_{\partial E_i} w_i \, \mathrm{d}P_{\partial E_i}$. Notice that

$$\int_{\partial E_i} w_i^2 = \int_{\partial E_i} \left(u_i^2 + \left(\oint_{\partial E_i} w_i \right)^2 + 2u_i \oint_{\partial E_i} w_i \right) \le 2 \int_{\partial E_i} u_i^2 \, \mathrm{d}P_{\partial E_i} + \frac{3}{2} \frac{1}{P(E_i)} \left(\int_{\partial E_i} w_i \right)^2 \right)^2$$

$$\stackrel{(4.7)}{\le} 2 \int_{\partial E_i} u_i^2 \, \mathrm{d}P_{\partial E_i} + \varepsilon \, \mathcal{C} \int_{\partial E_i} w_i^2 \, \mathrm{d}P_{\partial E_i} \, .$$

Analogously,

$$\int_{\partial E_{i}} |\mathrm{II}_{\partial E_{i}}|^{2} w_{i}^{2} = \int_{\partial E_{i}} |\mathrm{II}_{\partial E_{i}}|^{2} u_{i}^{2} + 2 \oint_{\partial E_{i}} w_{i} \int_{\partial E_{i}} |\mathrm{II}_{\partial E_{i}}|^{2} u_{i} + \left(\oint_{\partial E_{i}} w_{i} \right)^{2} \int_{\partial E_{i}} |\mathrm{II}_{\partial E_{i}}|^{2} \stackrel{(4.7)}{\leq} \int_{\partial E_{i}} |\mathrm{II}_{\partial E_{i}}|^{2} u_{i}^{2} \,\mathrm{d}P_{\partial E_{i}} + \varepsilon \,\mathcal{C} \int_{\partial E_{i}} u_{i}^{2} \,\mathrm{d}P_{\partial E_{i}} \,.$$

Therefore (4.28) can be rewritten as

(4.29)
$$\int_{\partial E_i} |\nabla u_i|^2 - |\mathrm{II}_{\partial E_i}|^2 u_i^2 \,\mathrm{d}P_{\partial E_i} \le \varepsilon \,\mathcal{C} \int_{\partial E_i} u_i^2 \,\mathrm{d}P_{\partial E_i} \,.$$

Observing that $\int_{\partial E_i} u_i = 0$, the strict stability stability estimate for the Jacobi operator from Theorem 3.1 implies

(4.30)
$$\varepsilon \, \mathcal{C} \int_{\partial E_i} u_i^2 \, \mathrm{d}P_{\partial E_i} \ge \varepsilon_M \int_{\partial E_i} u_i^2 \, \mathrm{d}P_{\partial E_i}$$

for some $\varepsilon_M > 0$ depending on M. Hence, if ε was taken sufficiently small, this implies that w_i is constant. Since $w_i \neq 0$ as E_i and E'_i are distinct, this implies that $\operatorname{vol}_i(E_i) \neq \operatorname{vol}_i(E'_i)$, where the volume is computed in the metric of M_{V_i} . This results into a contradiction, thus completing the proof of Theorem 4.1.

4.2. Estimates for isoperimetric graphs. The goal of this section is to establish several technical estimates that will be useful later for the proof of Proposition 4.3.

We shall consider a smooth *n*-dimensional Riemannian manifold (M, g) and we will assume that:

- (1) There exist two isoperimetric sets E_V, E'_V having equal volume V > 0, having smooth boundary, and such that $H_{\partial E_V} = H_{\partial E'_V}$. Moreover we assume $H_{\partial E_V} \ge \vartheta$, for some $\vartheta > 0$.
- (2) $\Sigma' := \partial E'_V$ can be written as the normal graph over $\Sigma := \partial E_V$ of a function $w : \Sigma \to \mathbb{R}$. Namely,

(4.31)
$$\Sigma' = \{ \exp_p(w(p)\nu_{\Sigma}(p)) : p \in \Sigma \},\$$

where ν_{Σ} is the unit inner normal of E_V .

(3) $\Sigma' \subset U$ where U is an open neighborhood of Σ where the metric g can be written in Fermi coordinates with respect to Σ . Namely, there is a > 0 such that

(4.32)
$$(U,g) \simeq ((-a,a) \times \Sigma, dz^2 + g_z)$$

where g_z is the pull-back on Σ via exponential map of the induced metric on the parallel hypersurface

(4.33)
$$\Sigma_z := \{ \exp_p(z\nu_{\Sigma}(p)) : p \in \Sigma \},\$$

for $z \in (-a, a)$. Let $\Psi_z : \Sigma \to \Sigma_z$ be the map $\Psi_z(p) := \exp_p(z\nu_{\Sigma}(p))$.

- (4) There exists K > 1 such that $\|II_{\Sigma}\|_{\infty} + \sup_{p \in U} |\operatorname{Riem}(p)| \leq K$ for every $p \in U$.
- (5) There exists $\varepsilon_0 := \varepsilon_0(K, n, \vartheta, a)$ such that, if $C_1(K, a), C_2(n, K, a), C_3(n, K, a)$ denote the constants appearing in Lemma 4.6, (4.44), and (4.46) below, respectively, then

(4.34)
$$\frac{1}{2} < (1 - C_1 \varepsilon_0)^{n/2} \le (1 + C_1 \varepsilon_0)^{n/2} < 2,$$

(4.35)
$$\max\left\{\varepsilon_0, \varepsilon_0^2(1+C_1\varepsilon_0), C_1\varepsilon_0, C_2\varepsilon_0\right\} < \min\left\{\frac{1}{10}, \frac{\vartheta}{C_3}\right\},$$

and $||w||_{\infty} + ||\nabla w||_{\infty} < \varepsilon_0.$

We will denote by g_w the metric on Σ such that $(g_w)_p(v, v') := g_{w(p)}(v, v')$, for every $v, v' \in T_p\Sigma$. We denote with $\nabla^w := \nabla^{g_w}$ the gradient with respect to the metric g_w on Σ , and with $|\cdot|_w$ the norm in the metric g_w . In this notation g_0 is the metric induced by g on Σ . We stress that ∇w denotes the gradient with respect to the metric g_0 , and $|\cdot|$ is the norm with respect to the metric g_0 .

We will denote $\dot{g}_z := \partial_z g_z$. Notice that \dot{g}_z is a (0, 2)-tensor on Σ_z , and $\dot{g}_z = 2 \Pi_{\Sigma_z}$. We can identify \dot{g}_z with a (0, 2)-tensor on Σ via the pull-back with the map Ψ_z defined in Item 3, namely, for every $p \in \Sigma$, and every $v, v' \in T_p \Sigma$, $(\dot{g}_z)_p(v, v') = 2 \Pi_{\Sigma_z}(\mathrm{d}(\Psi_z)(v), \mathrm{d}(\Psi_z)(v'))$. We also denote $(\dot{g}_w)_p := \dot{g}_{w(p)}$.

A standard computation exploiting Jacobi fields, Item 3 and Item 4 implies that there exists a constant $\mathcal{C} := \mathcal{C}(K, a)$ such that for every $p \in \Sigma$, every $v \in T_p\Sigma$ with $(g_0)_p(v, v) = 1$, and every $z \in (-a, a)$, it holds

$$(4.36) g_z(v,v) \le \mathcal{C}$$

We record the expression for the mean curvature of Σ' in terms of w, borrowed from [49]. It will be key for the subsequent estimates.

Lemma 4.5 ([49, Proposition 4.1]). Assume Item 2, and Item 3 above hold. The mean curvature of Σ' is

(4.37)
$$H_{\Sigma'} = -\operatorname{div}_{g_w} \left(\frac{\nabla^w w}{(1+|\nabla^w w|_w^2)^{\frac{1}{2}}} \right) + \frac{1}{2} \left(1+|\nabla^w w|_w^2 \right)^{\frac{1}{2}} \operatorname{tr}_{g_w} \dot{g}_w - \frac{1}{2} \frac{\dot{g}_w (\nabla^w w, \nabla^w w)}{(1+|\nabla^w w|_w^2)^{\frac{1}{2}}}$$

In the next lemma we compare induced metrics and volume forms at different heights with respect to Σ with the background metric g_0 , with the help of the second fundamental form and Riemann curvature bounds.

Lemma 4.6. Assume that Item 2, Item 3, and Item 4 above hold. There exists $C_1 := C_1(K, a)$ such that the following holds. For every $p \in \Sigma$, and every $v \in T_p\Sigma$ with $g_0(v, v) = 1$,

(4.38)
$$\begin{aligned} |(\dot{g}_w)_p(v,v) - (\dot{g}_0)_p(v,v)| &\leq C_1 |w(p)|, \\ |(g_w)_p(v,v) - (g_0)_p(v,v)| &\leq C_1 |w(p)|. \end{aligned}$$

Moreover,

(4.39)
$$(1 - C_1 \|w\|_{\infty})^{n/2} \operatorname{vol}_{g_0} \leq \operatorname{vol}_{g_w} \leq (1 + C_1 \|w\|_{\infty})^{n/2} \operatorname{vol}_{g_0}, \\ |\nabla w|^2 (1 - C_1 |w|) \leq |\nabla^w w|_w^2 \leq |\nabla w|^2 (1 + C_1 |w|).$$

Proof. Recall that the evolution equation of the second fundamental form II_{Σ_z} is

(4.40)
$$\partial_z II_{\Sigma_z} + II_{\Sigma_z}^2 + \operatorname{Riem}(\partial_z, \cdot, \partial_z, \cdot) = 0.$$

Hence, by using Item 4 and ODE comparison, there is $\mathcal{K} := \mathcal{K}(K, a)$ such that $\|II_{\Sigma_z}\|_{\infty} \leq \mathcal{K}$ for every $z \in (-a, a)$.

Notice that $\dot{g}_z = \partial_z g_z = 2 \Pi_{\Sigma_z}$. Via the pull back with the map Ψ_z defined in Item 3 we are identifying \dot{g}_z with a (0,2)-tensor on Σ as explained above. Hence, for every $z \in (-a, a)$, since |Sect| and $\|\Pi_{\Sigma_z}\|_{\infty}$ are uniformly bounded above by K, and since (4.36), and (4.40) are in force, we have

(4.41)
$$|\dot{g}_z - \dot{g}_0| \le \int_0^{|z|} |\partial_z \dot{g}_z| = 2 \int_0^{|z|} |\partial_z \mathrm{II}_{\Sigma_z}| \le C_1(a, K) |z|$$

Here (4.41) has to be understood evaluated at every point $p \in \Sigma$, and at every $v \in T_p\Sigma$ with $(g_0)_p(v,v) = 1$. Hence the first estimate of (4.38) holds. Similarly, we get the second inequality in (4.38) by using $\partial_z g_z = 2 \Pi_{\Sigma_z}$, the fact that $\|\Pi_{\Sigma_z}\|_{\infty}$ is uniformly bounded above by \mathcal{K} , and (4.36). Finally, the two inequalities in (4.39) are a direct consequence of the second estimate in (4.38), possibly enlarging C_1 .

Our next goal is to exploit the constant mean curvature condition for Σ' to obtain $L^{\infty}-L^2$ and gradient estimates for the graphing function w, via elliptic regularity theory. The precise statement follows.

Lemma 4.7. Assume Item 1, Item 2, Item 3, Item 4, and Item 5 above hold. Then there exists a constant $C := C(n, K, a, V, \operatorname{diam}(\Sigma)) > 0$ such that the following hold.

(4.42)
$$||w||_{\infty} \leq C ||w||_{L^2(\Sigma)},$$

(4.43)
$$\|\nabla w\|_{L^2(\Sigma)} \le C \|w\|_{L^2(\Sigma)}$$

Before we can prove Lemma 4.7 we need one last technical estimate.

Lemma 4.8. Assume that Item 1, Item 2, Item 3, Item 4, and Item 5 above hold. Then there exists a constant $C_2 := C_2(n, K, a) > 0$ such that

(4.44)
$$\left| \operatorname{div}_{g_w} \left(\frac{\nabla^w w}{(1 + |\nabla^w w|_w^2)^{\frac{1}{2}}} \right) \right| \le C_2(|w| + |\nabla^w w|_w^2)$$

Proof. Let us first prove that

(4.45)
$$-\operatorname{div}_{g_w}\left(\frac{\nabla^w w}{(1+|\nabla^w w|_w^2)^{\frac{1}{2}}}\right) \le C_2(|w|+|\nabla^w w|_w^2).$$

By (4.35) in Item 5, and the second estimate in (4.39), we have $|\nabla^w w|_w \leq 1$, and $|w| \leq 1$. Taking into account (4.38), there is a constant $C_3 := C_3(n, K, a) > 0$ such that

(4.46)
$$\operatorname{tr}_{g_w} \dot{g}_w \ge \operatorname{tr}_{g_0} \dot{g}_0 - 2C_3 |w| \,.$$

Moreover, taking into account that $H_{\Sigma} \geq \vartheta > 0$, and that $C_3 \varepsilon_0 < \vartheta$ by (4.35), we get $C_3 ||w||_{\infty} < H_{\Sigma}$. Hence, again recalling $|\nabla^w w|_w \leq 1$, we have

(4.47)

$$\frac{1}{2} \left(1 + |\nabla^{w}w|_{w}^{2} \right)^{\frac{1}{2}} \operatorname{tr}_{g_{w}} \dot{g}_{w} \stackrel{(4.46)}{\geq} \left(1 + |\nabla^{w}w|_{w}^{2} \right)^{\frac{1}{2}} \left(\frac{1}{2} \operatorname{tr}_{g_{0}} \dot{g}_{0} - C_{3}|w| \right) \\
= \left(1 + |\nabla^{w}w|_{w}^{2} \right)^{\frac{1}{2}} \left(H_{\Sigma} - C_{3}|w| \right) \\
\geq \left(1 + \frac{1}{4} |\nabla^{w}w|_{w}^{2} \right) \left(H_{\Sigma} - C_{3}|w| \right) \geq H_{\Sigma} - \frac{5}{4} C_{3}|w| .$$

In addition, using the second estimate in (4.38), the fact that $\|\dot{g}_z\|_{\infty} = 2\|\Pi_{\Sigma_z}\|_{\infty}$ is uniformly bounded from above by K by Item 4, and (4.36), we get, for some $C_4 := C_4(K, a) > 0$,

(4.48)
$$\frac{1}{2} \frac{\dot{g}_w(\nabla^w w, \nabla^w w)}{(1 + |\nabla^w w|_w^2)^{\frac{1}{2}}} \le C_4 |\nabla^w w|_w^2.$$

Using that $H_{\Sigma'} = H_{\Sigma}$, and combining (4.47), (4.48), and (4.37), we get the sought (4.45) with $C_2 := \max\{C_4, 2C_3\}$. Analogously, bounding from above $\operatorname{tr}_{g_w} \dot{g}_w$, one can also get

$$\operatorname{div}_{g_w}\left(\frac{\nabla^w w}{(1+|\nabla^w w|_w^2)^{\frac{1}{2}}}\right) \le C_2(|w|+|\nabla^w w|_w^2),$$

and thus the proof is concluded.

Proof of Lemma 4.7. We claim that there exists C(n, K, a) > 0 such that for every $p \ge 1$ it holds

(4.49)
$$p \int_{\Sigma} w_{+}^{p-1} |\nabla w_{+}|^{2} \mathrm{d} \operatorname{vol}_{g_{0}} \leq C \int_{\Sigma} \left(w_{+}^{\frac{p+1}{2}} \right)^{2} \mathrm{d} \operatorname{vol}_{g_{0}}$$

where w_+ denotes the positive part of w. Recall that $|\nabla^w w|_w \leq 1$ since (4.35) and the second estimate in (4.39) hold. By exploiting the latter information, and integrating (4.44) against w_+^p , we get

(4.50)
$$p \int_{\Sigma} w_{+}^{p-1} |\nabla^{w} w_{+}|_{w}^{2} \mathrm{d} \operatorname{vol}_{g_{w}} \leq \sqrt{2} C_{2} \int_{\Sigma} \left(w_{+}^{p+1} + w_{+} |\nabla^{w} w_{+}|_{w}^{2} w_{+}^{p-1} \right) \mathrm{d} \operatorname{vol}_{g_{w}} \,.$$

By (4.35) we get $\sqrt{2}C_2w_+ \leq \sqrt{2}C_2\varepsilon_0 \leq p/2$, so we can absorb the term in the right hand side of (4.50) to obtain

(4.51)
$$p \int_{\Sigma} w_{+}^{p-1} |\nabla^{w} w_{+}|_{w}^{2} \mathrm{d} \operatorname{vol}_{g_{w}} \leq 2\sqrt{2}C_{2} \int_{\Sigma} w_{+}^{p+1} \mathrm{d} \operatorname{vol}_{g_{w}} .$$

By (4.34) and the first of (4.39), we get that

(4.52)
$$\frac{1}{2}\operatorname{vol}_{g_w} \le \operatorname{vol}_{g_0} \le 2\operatorname{vol}_{g_w}$$

Moreover by (4.35) and the second estimate of (4.39), we get

(4.53)
$$\frac{1}{2}|\nabla w_{+}|^{2} \le |\nabla^{w}w_{+}|_{w}^{2}$$

By joining (4.52) and (4.53) with (4.51), we finally get the sought (4.49).

We now aim at obtaining (4.42) by exploiting (4.49), and a standard Moser iteration argument. By using the Gauss equations, we get $|\text{Sect}_{\Sigma}| \leq C(n, K)$, since $||\text{II}_{\Sigma}||_{\infty} + |\text{Sect}| \leq K$ by Item 4. Hence there exists a constant $C(n, K, V, \text{diam}(\Sigma))$ such that, calling $2^* := (2n)/(n-2)$,

(4.54)
$$C\|f - \overline{f}\|_{L^{2^*}(\Sigma, \operatorname{vol}_{g_0})} \le \|\nabla f\|_{L^2(\Sigma, \operatorname{vol}_{g_0})}, \quad \forall f \in \operatorname{Lip}(\Sigma).$$

From now on we abbreviate $\|\cdot\|_s := \|\cdot\|_{L^s(\Sigma, \operatorname{vol}_{g_0})}$ for any $s \ge 1$. Let $q := p + 1 \ge 2$. In the following computations the constants C, \tilde{C} , only depending on $n, K, a, V, \operatorname{diam}(\Sigma)$, might change from line to line. We have

$$(4.55) \frac{q}{\sqrt{q-1}} \|w_{+}\|_{q}^{q/2} \stackrel{(4.49)}{\geq} C \|\nabla(w_{+}^{q/2})\|_{2} \stackrel{(4.54)}{\geq} C^{2} \left\|w_{+}^{q/2} - \overline{w_{+}^{q/2}}\right\|_{2^{*}} \geq C^{2} \left\|w_{+}^{q/2}\right\|_{2^{*}} - C^{2}\tilde{C} \left\|w_{+}^{q/2}\right\|_{2^{*}} \\ = C^{2} \|w_{+}\|_{nq/(n-2)}^{q/2} - C^{2}\tilde{C} \left\|w_{+}^{q/2}\right\|_{1} \geq C^{2} \|w_{+}\|_{nq/(n-2)}^{q/2} - C^{2}\tilde{C}^{2} \|w_{+}^{q/2}\|_{2} \\ = C^{2} \|w_{+}\|_{nq/(n-2)}^{q/2} - C^{2}\tilde{C} \|w_{+}\|_{q}^{q/2}.$$

Set $q^* := nq/(n-2)$. From (4.55) we get

(4.56)
$$\|w_+\|_{q^*} \le \left(C\frac{q}{\sqrt{q-1}} + C\right)^{2/q} \|w_+\|_q$$

Recalling that $p \ge 1$ was arbitrary, we can take $p = p_k$ such that $q = q_k = 2\left(\frac{n}{n-2}\right)^k$ for $k \ge 0$. Hence applying (4.56) with $q = q_k$, and letting $k \to \infty$, the usual Moser iteration scheme gives

$$(4.57) ||w_+||_{\infty} \le C||w_+||_2$$

Reasoning analogously with w_{-} in place of w_{+} , we get (4.57) for w_{-} as well. Hence (4.42) is proved.

We now aim at proving (4.43). Again, we can integrate (4.44) against w. Arguing verbatim as we did to obtain (4.49) (with p = 1) we conclude

(4.58)
$$\int_{\Sigma} |\nabla w|^2 \mathrm{d} \operatorname{vol}_{g_0} \le C \int_{\Sigma} w^2 \mathrm{d} \operatorname{vol}_{g_0},$$

which is the sought (4.43).

4.3. **Proof of Proposition 4.3.** Let \mathcal{V} be the set of volumes $V \in (0, \infty)$ such that the isoperimetric profile I_M is differentiable at V. Recalling that I_M is concave [7], then $(0, \infty) \setminus \mathcal{V}$ is a countable set. In particular, \mathcal{V} has full measure in $(0, \infty)$. We divide the proof in two steps.

Step 1. Let C be the constant in (1.2). We claim that there exist $\mathcal{C} := \mathcal{C}(M) > 0$, and $\varepsilon_0 := \varepsilon_0(C, \text{AVR}, n) > 0$ such that the following holds. For every $\varepsilon < \varepsilon_0$, every sequence $\mathcal{V} \ni V_i \to \infty$, and every isoperimetric regions $E_{V_i}, E'_{V_i} \subset M$ with $\text{vol}(E_{V_i}) = \text{vol}(E'_{V_i}) = V_i$, there is $i_0 \in \mathbb{N}$ such that for every $i \ge i_0$, $\tilde{\Sigma}'_i := \partial E'_{V_i}$ is the normal graph over $\tilde{\Sigma}_i := \partial E_{V_i}$ of a function $w_i : \tilde{\Sigma}_i \to \mathbb{R}$ satisfying

(4.59) $\|w_i\|_{\infty} \le \varepsilon V_i^{\frac{1}{n}}, \quad \|\nabla w_i\|_{\infty} \le \varepsilon,$

(4.60)
$$\|w_i\|_{\infty} \le \mathcal{C}V_i^{-\frac{n-1}{2n}} \|w_i\|_{L^2(\tilde{\Sigma}_i)}$$

(4.61)
$$\|\nabla w_i\|_{L^2(\tilde{\Sigma}_i)} \le \mathcal{C}V_i^{-\frac{1}{n}} \|w_i\|_{L^2(\tilde{\Sigma}_i)}.$$

Let $p \in M$ be a fixed point. Let ε_0 be small enough to be determined, and let $\varepsilon < \varepsilon_0$. Let us set $r_{V_i} := (\operatorname{AVR}(M) \omega_n)^{-\frac{1}{n}} V_i^{\frac{1}{n}}$, and $M_i := (M^n, r_{V_i}^{-1} \mathrm{d}_g, p)$. Let $\Sigma_i, \Sigma'_i \subset M_i$ denote the sets $\tilde{\Sigma}_i, \tilde{\Sigma}'_i \subset M$ in the rescaled manifold M_i . By the second estimate in (3.3) we get that, for *i* large enough,

(4.62)
$$d_{\mathrm{H}}^{M_{i}}(\Sigma_{i},\Sigma_{i}') \leq d_{\mathrm{H}}^{M_{i}}(\partial B_{1}^{M_{i}}(p),\Sigma_{i}) + d_{\mathrm{H}}^{M_{i}}(\partial B_{1}^{M_{i}}(p),\Sigma_{i}') \leq \varepsilon, \\ B_{1}^{M_{i}}(\Sigma_{i}) \cup B_{1}^{M_{i}}(\Sigma_{i}') \subset B_{3}^{M_{i}}(p).$$

By (3.4), (1.2), and the second line of (4.62), we get that there is a constant $C_1 := C_1(C, AVR)$ only depending on the constant C appearing in (1.2) and AVR such that, for *i* large enough,

(4.63)
$$\sup_{x \in \Sigma_i} |\mathrm{II}(x)| + \sup_{x \in B_1^{M_i}(\Sigma_i)} |\mathrm{Riem}(x)| \le C_1$$

Hence, if ε_0 is chosen small enough with respect to C_1 , we can use the first line in (4.62), and the results in Appendix A (in particular, we are using (3.20), (3.21), and (3.23) as stated in the proof of Proposition 3.3 to Σ_i, Σ'_i) to get the existence of a function $\tilde{w}_i : \Sigma_i \to \mathbb{R}$ such that Σ'_i is the normal graph over Σ_i of \tilde{w}_i , and

(4.64)
$$\|\tilde{w}_i\|_{\infty,\Sigma_i} + \|\nabla^{\Sigma_i}\tilde{w}_i\|_{\infty,\Sigma_i} \le \varepsilon.$$

Also notice that, since $V_i \in \mathcal{V}$, it holds $H_{\Sigma_i} = H_{\Sigma'_i} \geq C_2$, where $C_2 := C_2(n, \text{AVR})$ and the lower bound comes from [8, Equation (3.18), Corollary 3.5]. Thanks to the bounds (3.20), (3.21), and (3.23) as stated in the proof of Proposition 3.3, (B.1) holds. Hence Proposition B.1 applies. Thus Item 1, Item 2, Item 3, and Item 4 in Section 4.2 are satisfied. Hence, choosing $\varepsilon_0 :=$ $\varepsilon_0(n, C, \text{AVR})$ small enough, we get that Item 5 is also in force, due to (4.64). By Lemma 3.6, there is a constant C_3 depending on the manifold M such that for *i* large enough diam $\Sigma_i \leq C_3$, where Σ_i is endowed with the metric induced by the ambient M_i . Hence Lemma 4.7 applies and there exists a constant $\mathcal{C} := \mathcal{C}(n, C, \text{AVR}, C_3)$ such that

(4.65)
$$\|\tilde{w}_i\|_{\infty,\Sigma_i} \leq \mathcal{C} \|\tilde{w}_i\|_{L^2(\Sigma_i)},$$

(4.66)
$$\|\nabla^{\Sigma_i} \tilde{w}_i\|_{L^2(\Sigma_i)} \le \mathcal{C} \|\tilde{w}_i\|_{L^2(\Sigma_i)}.$$

Scaling back we get (4.59), (4.60), and (4.61) are satisfied. This completes the proof of the claim.

Step 2. Let \mathcal{V} be as at the beginning of the proof, and let $\varepsilon_0, \mathcal{C}$ be as in the **Step 1** above. We claim that for every $\varepsilon < \varepsilon_0$ there exists $V_0 := V_0(\varepsilon)$ such that for every $\mathcal{V} \ni V > V_0$ the conclusion of Proposition 4.3 holds with the constant C. This will be enough to conclude the proof.

Suppose the latter is not true. Hence there is $\varepsilon < \varepsilon_0$ such that there is $\mathcal{V} \ni V_i \to \infty$ such that the conclusion of Proposition 4.3 with the constant \mathcal{C} is false for every volume V_i . This contradicts the claim at the beginning of **Step 1**, thus completing the proof.

4.4. **Proof of Proposition 4.4.** For the proof of Proposition 4.4 we will need to make effective some of the ideas that were introduced for the proof of Proposition 3.4 above. More precisely, we are going to exploit the asymptotics of the isoperimetric profile of M to estimate the Ricci curvature in the direction normal to the isoperimetric boundaries. With respect to the proof of (3.6) several additional error terms arise, that need to be carefully controlled.

We will denote by $\mathcal{C} > 0$ a constant depending on M only that may change from line to line. Consider a volume V > 0 such that the isoperimetric profile I of M is differentiable at V. Let $E \subset M$ be any isoperimetric set of volume V. For $t \in \mathbb{R}$, we denote by E_t the *t*-enlargement of E, namely $E_t := \{x \in M : d_E^s < t\}$, where d_E^s denotes the signed distance from E. By Lemma 4.2 there exists $\overline{\rho} > 0$ independent of E, V such that the reach of ∂E is bounded below by $\overline{\rho}V^{\frac{1}{n}}$, for any V large enough. Hence, for $t \in (-\overline{\rho}V^{\frac{1}{n}}, \overline{\rho}V^{\frac{1}{n}})$ we can compute the perimeter $P(E_t)$ of *t*-enlargements as

$$P^{\frac{n}{n-1}}(E_t) = P^{\frac{n}{n-1}}(E) + \frac{n}{n-1}P^{\frac{1}{n-1}}(E)t \int_{\partial E} H_{\partial E} + \int_0^t \int_0^s \frac{\mathrm{d}^2}{\mathrm{d}r^2} P^{\frac{n}{n-1}}(E_r) \,\mathrm{d}r \,\mathrm{d}s$$

$$= P^{\frac{n}{n-1}}(E) + \frac{n P^{\frac{n}{n-1}}(E) H_{\partial E}}{n-1} t + \frac{n}{n-1} \int_0^t \int_0^s \frac{P^{\frac{2-n}{n-1}}(E_r)}{n-1} \left(\int_{\partial E_r} H_{\partial E_r}\right)^2 \,\mathrm{d}r \,\mathrm{d}s + \frac{n}{n-1} \int_0^t \int_0^s P^{\frac{1}{n-1}}(E_r) \int_{\partial E_r} H_{\partial E_r}^2 - |\mathrm{II}_{\partial E_r}|^2 - \mathrm{Ric}(\nabla \mathrm{d}_E^s, \nabla \mathrm{d}_E^s) \,\mathrm{d}r \,\mathrm{d}s \,.$$

(4.67)

Note that the last term appearing in (4.67) is exactly the term we would like to estimate.

Let $J(\cdot) := I^{\frac{n}{n-1}}(\cdot)$, where *I* is the isoperimetric profile of *M*. Since *E* is isoperimetric, denoting for brevity by $|\cdot| :=$ vol the Riemannian volume on *M*, for $t \in (0, \overline{\rho}V^{\frac{1}{n}})$ we can estimate

$$J(|E_t|) + J(|E_{-t}|) - 2J(V) \le P^{\frac{n}{n-1}}(E_t) + P^{\frac{n}{n-1}}(E_{-t}) - 2P^{\frac{n}{n-1}}(E)$$

$$= \frac{n}{n-1} \int_0^t \int_{-s}^s \frac{P^{\frac{2-n}{n-1}}(E_r)}{n-1} \left(\int_{\partial E_r} H_{\partial E_r} \right)^2$$

$$+ P^{\frac{1}{n-1}}(E_r) \int_{\partial E_r} H^2_{\partial E_r} - |\mathrm{II}_{\partial E_r}|^2 - \mathrm{Ric}(\nabla \mathrm{d}^s_E, \nabla \mathrm{d}^s_E) \,\mathrm{d}r \,\mathrm{d}s$$

$$\le \frac{n}{n-1} \int_0^t \int_{-s}^s P^{\frac{1}{n-1}}(E_r) \left(\int_{\partial E_r} H^2_{\partial E_r} \right) - P^{\frac{1}{n-1}}(E_r) \int_{\partial E_r} \mathrm{Ric}(\nabla \mathrm{d}^s_E, \nabla \mathrm{d}^s_E) \,\mathrm{d}r \,\mathrm{d}s \,.$$

By the isoperimetric inequality on manifolds with nonnegative Ricci and Euclidean volume growth, see e.g. [15, Corollary 1.3], and taking into account [7, Corollary 3.13, Corollary 4.19] and [5, Corollary 3.6], for any V large enough we can further estimate

$$(4.69) \quad CP^{\frac{n}{n-1}}(E_r) \ge |E_r| \ge |E_{-|r|}| \ge V - P(E) \int_0^{|r|} \left(1 - \frac{H_{\partial E}}{n-1}z\right)_+^{n-1} \mathrm{d}z \ge V - cV^{\frac{n-1}{n}}\overline{\rho}V^{\frac{1}{n}}$$

for some C, c depending on n, $\operatorname{AVR}(M)$, for any $r \in (-\overline{\rho}V^{\frac{1}{n}}, \overline{\rho}V^{\frac{1}{n}})$. Hence, up to possibly decreasing $\overline{\rho}$, (4.69) implies that $P^{\frac{1}{n-1}}(E_r) \geq \mathcal{C}V^{\frac{1}{n}}$, for any $r \in (-\overline{\rho}V^{\frac{1}{n}}, \overline{\rho}V^{\frac{1}{n}})$ and any V large enough. Observe also that the previous argument implies that

(4.70)
$$\frac{3}{4}V < |E_r| < \frac{5}{4}V,$$

(

(4.68)

for any $r \in (-\overline{\rho}V^{\frac{1}{n}}, \overline{\rho}V^{\frac{1}{n}})$ and any V large enough, up to decreasing $\overline{\rho}$.

Moreover, computing integrals in Fermi coordinates as in the first part of the proof of Theorem 4.1 (see (4.10)), it follows from Proposition 4.3 that

(4.71)
$$\int_{\partial E_r} \operatorname{Ric}(\nabla \mathrm{d}_E^s, \nabla \mathrm{d}_E^s) \ge \mathcal{C} \int_{\partial E} \operatorname{ric}(r, x) \, \mathrm{d}P_{\partial E}(x)$$

for any $r \in (-\overline{\rho}V^{\frac{1}{n}}, \overline{\rho}V^{\frac{1}{n}})$, where $\operatorname{ric}(r, x) := \operatorname{Ric}(\nabla d_E^s, \nabla d_E^s) \circ (\exp_x(r\nu_E^{\operatorname{ext}}(x)))$ for any $x \in \partial E$, where $\nu_E^{\operatorname{ext}}$ is the unit outer normal along ∂E .

Combining (4.69) and (4.71), for $t \in (0, \overline{\rho}V^{\frac{1}{n}})$ we get

$$\int_{0}^{t} \int_{-s}^{s} P^{\frac{1}{n-1}}(E_{r}) \int_{\partial E_{r}} \operatorname{Ric}(\nabla d_{E}^{s}, \nabla d_{E}^{s}) \, \mathrm{d}r \, \mathrm{d}s \geq \mathcal{C}V^{\frac{1}{n}} \int_{0}^{t} \int_{-s}^{s} \int_{\partial E} \operatorname{ric}(r, x) \, \mathrm{d}P_{\partial E}(x) \, \mathrm{d}r \, \mathrm{d}s$$
$$\geq \mathcal{C}V^{\frac{1}{n}} \int_{-t}^{t} \int_{|r|}^{t} \int_{\partial E} \operatorname{ric}(r, x) \, \mathrm{d}P_{\partial E}(x) \, \mathrm{d}s \, \mathrm{d}r = \mathcal{C}V^{\frac{1}{n}} \int_{-t}^{t} (t - |r|) \int_{\partial E} \operatorname{ric}(r, x) \, \mathrm{d}P_{\partial E}(x) \, \mathrm{d}r$$
$$\geq \mathcal{C}V^{\frac{1}{n}} \int_{-\frac{t}{2}}^{\frac{t}{2}} (t - |r|) \int_{\partial E} \operatorname{ric}(r, x) \, \mathrm{d}P_{\partial E}(x) \, \mathrm{d}r \geq \mathcal{C}V^{\frac{1}{n}} \frac{t}{2} \int_{-\frac{t}{2}}^{\frac{t}{2}} \int_{\partial E} \operatorname{ric}(r, x) \, \mathrm{d}P_{\partial E}(x) \, \mathrm{d}r \, .$$

Combining with (4.68) we find

(4.72)
$$t V^{\frac{1}{n}} \int_{-\frac{t}{2}}^{\frac{t}{2}} \int_{\partial E} \operatorname{ric}(r, x) \, \mathrm{d}P_{\partial E}(x) \, \mathrm{d}r \leq \mathcal{C} \left(\frac{n}{n-1} \int_{0}^{t} \int_{-s}^{s} P^{\frac{1}{n-1}}(E_{r}) \left(\int_{\partial E_{r}} H_{\partial E_{r}}^{2} \right) \, \mathrm{d}r \, \mathrm{d}s + \left(J(|E_{t}|) + J(|E_{-t}|) - 2J(V) \right) \right).$$

We now aim at estimating the difference $J(|E_t|) + J(|E_{-t}|) - 2J(V)$. This is the goal of the following lemma, whose proof is postponed to the end of this section.

Lemma 4.9. In the same setting and with the same notation as above it holds

$$(4.73) - (J(|E_t|) + J(|E_{-t}|) - 2J(V)) \leq C\eta P(E)\varepsilon t^2 V^{-\frac{1}{n}} - \int_0^t \int_{-s}^s \frac{n}{n-1} I^{\frac{1}{n-1}}(|E_r|) I'_+(|E_r|) \left(\int_{\partial E_r} H_{\partial E_r}\right) dr ds.$$

Combining (4.72) and (4.73) we deduce

$$t V^{\frac{1}{n}} \int_{-\frac{t}{2}}^{\frac{t}{2}} \int_{\partial E} \operatorname{ric}(r, x) \, \mathrm{d}P_{\partial E}(x) \, \mathrm{d}r \leq \mathcal{C} \left(\mathcal{C}\eta P(E) \varepsilon t^2 V^{-\frac{1}{n}} + \frac{n}{n-1} \int_0^t \int_{-s}^s P^{\frac{1}{n-1}}(E_r) \left(\int_{\partial E_r} H^2_{\partial E_r} \right) - I^{\frac{1}{n-1}}(|E_r|) I'(|E_r|) \left(\int_{\partial E_r} H_{\partial E_r} \right) \, \mathrm{d}r \, \mathrm{d}s \right),$$

$$(4.74) \qquad + \frac{n}{n-1} \int_0^t \int_{-s}^s P^{\frac{1}{n-1}}(E_r) \left(\int_{\partial E_r} H^2_{\partial E_r} \right) - I^{\frac{1}{n-1}}(|E_r|) I'(|E_r|) \left(\int_{\partial E_r} H_{\partial E_r} \right) \, \mathrm{d}r \, \mathrm{d}s \right),$$

for $t \in (0, \overline{\rho}V^{\frac{1}{n}})$.

The proof of Proposition 4.4 will be completed with the help of the following:

Lemma 4.10. In the same setting and with the same notation as above, up to possibly choosing a larger i_{ε} , it holds

$$(4.75) \sup_{r \in (-\overline{\rho}V^{\frac{1}{n}}/2, \overline{\rho}V^{\frac{1}{n}}/2)} P^{\frac{1}{n-1}}(E_r) \left(\int_{\partial E_r} H^2_{\partial E_r} \right) - I^{\frac{1}{n-1}}(|E_r|) I'_+(|E_r|) \left(\int_{\partial E_r} H_{\partial E_r} \right) \le \varepsilon \eta V^{\frac{n-2}{n}}.$$

Given Lemma 4.10, whose proof is postponed to the end of this section, we can combine (4.74) and (4.75) and obtain the following. For $t \in (0, \overline{\rho}V^{\frac{1}{n}}/2)$, up to choosing a larger i_{ε} it

holds

(4.76)
$$t V^{\frac{1}{n}} \int_{-\frac{t}{2}}^{\frac{t}{2}} \int_{\partial E} \operatorname{ric}(r, x) \, \mathrm{d}P_{\partial E}(x) \, \mathrm{d}r \le \mathcal{C} \left(\eta P(E) \varepsilon t^2 V^{-\frac{1}{n}} + \varepsilon \eta V^{\frac{n-2}{n}} t^2 \right) \\ < \mathcal{C}\eta P(E) \varepsilon t^2 V^{-\frac{1}{n}} \, .$$

Exploiting Fermi coordinates again, arguing similarly as in (4.71), we conclude that

(4.77)
$$V^{\frac{2}{n}} \int_{B_r(\partial E)} \operatorname{Ric}(\nabla \mathrm{d}_E^s, \nabla \mathrm{d}_E^s) \leq \mathcal{C}\eta\varepsilon \leq \varepsilon,$$

for any $r \in (0, C_0 V^{\frac{1}{n}})$ for some $C_0 > 0$. Note that in (4.77) above we estimated $\mathcal{C}\eta \leq 1$ for a choice of η depending only on the last constant \mathcal{C} appeared, hence depending only on M. This completes the proof of Proposition 4.4, modulo the proofs of Lemma 4.9 and Lemma 4.10 whose details are discussed below.

Proof of Lemma 4.9. Denote by $D^2 J$ the (negative) finite measure given by the distributional second derivative of J. For W > 4 we consider the maximal function M_W of $\chi_{(W/4,+\infty)}|D^2 J|$, namely

(4.78)
$$M_W(v) := \sup_{h>0} \frac{1}{2h} |D^2 J| ((v-h, v+h) \cap (W/4, \infty))$$

Denoting by $|A| := \mathcal{L}^1(A)$ for any measurable set $A \subset \mathbb{R}$, the classical Hardy–Littlewood inequality (whose proof works also for measures, cf. [34, Theorem 3.5.6]) implies

(4.79)
$$\left| \left\{ v \in \mathbb{R} : M_W(v) > W^{-1} \left(|D^2 J| (W/4, +\infty) \right)^{\frac{1}{2}} \right\} \right| \frac{(|D^2 J| (W/4, +\infty))^{\frac{1}{2}}}{W} \\ := |X_W| \frac{(|D^2 J| (W/4, +\infty))^{\frac{1}{2}}}{W} \le 5 |D^2 J| (W/4, +\infty) .$$

Notice that if $v \notin X_W$, then

(4.80)
$$\sup_{h>0} \frac{1}{2h} |D^2 J|((v-h,v+h) \cap (W/4,\infty)) \le W^{-1} \left(|D^2 J|(W/4,+\infty) \right)^{\frac{1}{2}}.$$

Define recursively $W_1 = 5$, $W_{i+1} = 2W_i$, and let $\widetilde{X}_i = X_{W_i} \cap (W_i, W_{i+1})$.

Let now $\varepsilon > 0$ be arbitrary. Let also $\eta \in (0,1)$ be a constant that will be chosen later depending on M only. There exists $i_{\varepsilon} \ge 1$ such that $(|D^2J|(W_i/4, +\infty))^{\frac{1}{2}} \le \varepsilon \eta$ for any $i \ge i_{\varepsilon}$. We claim that the set $\mathcal{G}_{\varepsilon}$ can be chosen as

(4.81)
$$\mathcal{G}_{\varepsilon} := \{V > W_{i_{\varepsilon}} : \exists I'(V)\} \setminus \bigcup_{i \ge i_{\varepsilon}} \widetilde{X}_{i}.$$

Indeed, we first note that for any $z > W_{i_{\varepsilon}}$ large, writing $W_{i_z} \leq z < W_{i_z+1}$ for some i_z , it holds

$$\begin{aligned} \left| (z,2z) \cap \bigcup_{i \ge i_{\varepsilon}} \widetilde{X}_{i} \right| &\leq \left| \widetilde{X}_{i_{z}} \right| + \left| \widetilde{X}_{i_{z}+1} \right| \\ &\leq 5W_{i_{z}} \left(|D^{2}J|(W_{i_{z}}/4,+\infty))^{\frac{1}{2}} + 5W_{i_{z}+1} \left(|D^{2}J|(W_{i_{z}+1}/4,+\infty))^{\frac{1}{2}} \right) \\ &\leq 10W_{i_{z}+1} \left(|D^{2}J|(W_{i_{z}}/4,+\infty))^{\frac{1}{2}} \\ &\leq 20z \left(|D^{2}J|(z/8,+\infty) \right)^{\frac{1}{2}} . \end{aligned}$$

Therefore $\lim_{z\to+\infty} \mathcal{L}^1\left((z,2z) \cap (\bigcup_{i\geq i_{\varepsilon}} \widetilde{X}_i)\right)/z = 0$ and (4.5) follows.

Hence let $V \in \mathcal{G}_{\varepsilon}$ and let $E \subset M$ be any isoperimetric set of volume V, as considered above. Let i_0 be such that $W_{i_0} < V \leq W_{i_0+1}$. Recalling (4.70), let $h_r := \max\{V - |E_{-r}|, |E_r| - V\} < V/4$ for $r \in (0, \overline{\rho}V^{\frac{1}{n}})$. Since $V - h_r \geq \frac{3}{4}V > W_{i_0}/2$, (4.80) implies

$$|D^{2}J|(|E_{-r}|, |E_{r}|) \leq |D^{2}J|(V - h_{r}, V + h_{r}) = |D^{2}J|((V - h_{r}, V + h_{r}) \cap (W_{i_{0}}/4, \infty))$$

$$\stackrel{(4.80)}{\leq} \frac{2h_{r}}{W_{i_{0}}} \varepsilon \eta \leq \frac{4h_{r}}{V} \varepsilon \eta.$$

Moreover, arguing similarly as in (4.69), one estimates $h_r \leq CP(E)r$ (see [7, Corollary 3.13]). Hence

(4.82)
$$|D^2 J|(|E_{-r}|, |E_r|) \le \mathcal{C}\eta P(E) \frac{\varepsilon}{V} r,$$

for $r \in (0, \overline{\rho}V^{\frac{1}{n}})$.

Let us now conclude the proof with a mollification argument. Let $\rho_{\tau} : \mathbb{R} \to \mathbb{R}$, for $\tau \in (0, 1)$ be a family of symmetric mollifiers with support in $(-\tau, \tau)$. Let $J_{\tau} := J * \rho_{\tau} : (1, \infty) \to \mathbb{R}$. Hence J_{τ} is concave, smooth and $J''_{\tau} = D^2 J * \rho_{\tau}$. Taylor expanding $J_{\tau}(|E_t|)$ with respect to tas done in (4.67) for $P^{\frac{n}{n-1}}(E_t)$, using [7, Proposition 3.11] and [5, Corollary 3.6], up to taking a larger i_{ε} , for $t \in (0, \overline{\rho}V^{\frac{1}{n}})$ we can estimate

$$-(J_{\tau}(|E_{t}|) + J_{\tau}(|E_{-t}|) - 2J_{\tau}(V)) = \int_{0}^{t} \int_{-s}^{s} |J_{\tau}''|(|E_{r}|)P(E_{r})^{2} - J_{\tau}'(|E_{r}|) \left(\int_{\partial E_{r}} H_{\partial E_{r}}\right) dr ds$$

$$\leq \left(1 + \frac{H_{\partial E}}{n-1}\overline{\rho}V^{\frac{1}{n}}\right)P(E) \int_{0}^{t} \int_{-|E_{s}|}^{|E_{s}|} |J_{\tau}''|(v) dv ds - \int_{0}^{t} \int_{-s}^{s} J_{\tau}'(|E_{r}|) \left(\int_{\partial E_{r}} H_{\partial E_{r}}\right) dr ds$$

$$\leq \mathcal{C}P(E) \int_{0}^{t} \int_{-|E_{s}|}^{|E_{s}|} |J_{\tau}''|(v) dv ds - \int_{0}^{t} \int_{-s}^{s} J_{\tau}'(|E_{r}|) \left(\int_{\partial E_{r}} H_{\partial E_{r}}\right) dr ds.$$

Letting $\tau \to 0$ we get

$$(4.83) \qquad (4.83) \qquad (4.82) + J(|E_{-t}|) - 2J(V)) \stackrel{(4.82)}{\leq} C\eta P(E)^2 \frac{\varepsilon}{V} t^2 - \int_0^t \int_{-s}^s J'_+(|E_r|) \left(\int_{\partial E_r} H_{\partial E_r} \right) dr ds \leq C\eta P(E)\varepsilon t^2 V^{-\frac{1}{n}} - \int_0^t \int_{-s}^s \frac{n}{n-1} I^{\frac{1}{n-1}}(|E_r|) I'_+(|E_r|) \left(\int_{\partial E_r} H_{\partial E_r} \right) dr ds.$$

Proof of Lemma 4.10. Note that (4.75) is a scale invariant estimate. Hence to prove it we can argue by contradiction and assume that there exist sequences $V_i \nearrow \infty$, $r_i \in (-\overline{\rho}V_i^{\frac{1}{n}}/2, \overline{\rho}V_i^{\frac{1}{n}}/2)$, and isoperimetric sets E_i of volume V_i such that

(4.84)
$$P^{\frac{1}{n-1}}(E_{r_i})\left(\int_{\partial E_{r_i}} H^2_{\partial E_{r_i}}\right) - I^{\frac{1}{n-1}}(|E_{r_i}|)I'_+(|E_{r_i}|)\left(\int_{\partial E_{r_i}} H_{\partial E_{r_i}}\right) > \varepsilon \eta V_i^{\frac{n-2}{n}}$$

where E_{r_i} is the r_i -enlargement of E_i in M. In the sequence of rescaled manifolds M_{V_i} this means that there exists a sequence of isoperimetric sets $\widetilde{E}_i \subset M_{V_i}$ such that $|\widetilde{E}_i| = \text{AVR}(M)\omega_n$, and a sequence $\widetilde{r}_i \in (-\widetilde{\rho}, \widetilde{\rho})$ such that

$$(4.85) P_{i}^{\frac{1}{n-1}}(\widetilde{E}_{\widetilde{r}_{i}})\left(\int_{\partial \widetilde{E}_{\widetilde{r}_{i}}}H_{\partial \widetilde{E}_{\widetilde{r}_{i}}}^{2}\right) - I_{i}^{\frac{1}{n-1}}(|\widetilde{E}_{\widetilde{r}_{i}}|)(I_{i})_{+}'(|\widetilde{E}_{\widetilde{r}_{i}}|)\left(\int_{\partial \widetilde{E}_{\widetilde{r}_{i}}}H_{\partial \widetilde{E}_{\widetilde{r}_{i}}}\right) > \mathcal{C}\varepsilon\eta,$$

where perimeters, volumes, distances and isoperimetric profile are computed on M_{V_i} .

The key for the proof will be the stability of the signed distance functions from isoperimetric sets and their Laplacians under Gromov–Hausdorff convergence.

Up to the extraction of a subsequence that we do not relabel, M_{V_i} converges to an asymptotic cone X, with isoperimetric profile I_X , and $\tilde{r}_i \to r_\infty$. Moreover $I_i \to I_X$ locally uniformly on $(0,\infty)$. Since the right derivative $(I_i^{\frac{n}{n-1}})'_+$ converges to the constant $n^{\frac{n}{n-1}}(\theta\omega_n)^{\frac{1}{n-1}}$ almost everywhere and $I_i^{\frac{n}{n-1}}$ is concave, where $\theta := \text{AVR}(M)$, then also $(I_i^{\frac{n}{n-1}})'_+ \to n^{\frac{n}{n-1}}(\theta\omega_n)^{\frac{1}{n-1}}$ locally uniformly on $(0,\infty)$. Hence $I_i^{\frac{1}{n-1}}(I_i)'_+ \to (n\theta\omega_n)^{\frac{1}{n-1}}(n-1)$ locally uniformly on $(0,\infty)$.

Denoting $H_i := H_{\partial \tilde{E}_i}$, we know from [7, Theorem 4.23] that $\tilde{E}_{\tilde{r}_i}$ converges to the ball $B_{(1+r_{\infty})}^X$ in X centered at the tip, and $H_i \to n-1$. Moreover by [7, Proposition 3.11] we get

$$\begin{aligned}
n\theta\omega_n(1+r_\infty)^{n-1} &= P_X(B_{1+r_\infty}^X) \le \liminf_i P_i(E_{\widetilde{r}_i}) \\
\le \liminf_i \left(1 + \frac{H_i}{n-1}r_\infty\right)^{n-1} P_i(\widetilde{E}_i) \\
&= n\theta\omega_n(1+r_\infty)^{n-1}.
\end{aligned}$$

Finally, by [7, Theorem 1.2],

(4.87)
$$\Delta f_i - \frac{H_i}{1 + \frac{H_i}{n-1}f_i} \ge 0$$
 on $\{0 \ge f_i > -2\tilde{\rho}\}, \quad \frac{H_i}{1 + \frac{H_i}{n-1}f_i} - \Delta f_i \ge 0$ on $\{0 \le f_i < 2\tilde{\rho}\},$

where f_i is the signed distance function from \widetilde{E}_i . Moreover, the Riccati equation for the evolution of Δf_i along geodesics perpendicular to $\partial \widetilde{E}_i$, together with Proposition 3.3, implies that Δf_i is uniformly bounded on $\{-2\widetilde{\rho} < f_i < 2\widetilde{\rho}\}$ (see also the proof of Lemma 4.6). By [3, Proposition 1.3.1], it follows that, up to subsequences, Δf_i converges weakly in L^2 in the sense of [3] along the sequence of spaces $(X_{i,\widehat{\rho}} := \{-\widehat{\rho} \leq f_i \leq \widehat{\rho}\}, d_{M_{V_i}}) \to (\overline{B}_{1+\widehat{\rho}}^X \setminus B_{1-\widehat{\rho}}^X, d_X)$ for any $\widehat{\rho} \in (0, 2\widetilde{\rho})$. By the divergence theorem, it follows that the L^2 -weak limit of Δf_i coincides with the Laplacian from the tip o on X, i.e., the function $\frac{n-1}{d_X}$.

Letting $\chi_i = \operatorname{sgn}(f_i)$, then $\chi_i \to \chi$ in L^2 -strong where $\chi = 1$ on $X \setminus B_1^X$ and $\chi = -1$ on B_1^X . It follows from (4.87) that

(4.88)
$$\int_{X_{i,\widehat{\rho}}} \left| \frac{H_i}{1 + \frac{H_i}{n-1} f_i} - \Delta f_i \right| = \int_{X_{i,\widehat{\rho}}} \chi_i \left(\frac{H_i}{1 + \frac{H_i}{n-1} f_i} - \Delta f_i \right)$$
$$\xrightarrow[i \to \infty]{} \int_{\overline{B}_{1+\widehat{\rho}}^X \setminus B_{1-\widehat{\rho}}^X} \chi \left(\frac{n-1}{d_o^X} - \frac{n-1}{d_o^X} \right) = 0$$

where the convergence follows from [3, Proposition 1.3.3(d)]. Hence [3, Proposition 1.3.3(e)] implies that Δf_i converges strongly in L^2 .

Using Fermi coordinates as in the proof of Theorem 4.1 and Fubini's theorem, it follows that the restriction of $\frac{H_i}{1+\frac{H_i}{n-1}f_i} - \Delta f_i$ on almost every geodesic from $\partial \widetilde{E}_i$ tends to zero strongly in L^1 along the geodesic. Since Δf_i are (uniformly) Lipschitz along geodesics from $\partial \widetilde{E}_i$ by the Riccati equation, then $\frac{H_i}{1+\frac{H_i}{n-1}f_i} - \Delta f_i$ tends to zero strongly in L^2 (actually, uniformly) along almost every geodesic from $\partial \widetilde{E}_i$. In particular, by using the Dominated Convergence Theorem,

$$(4.89) \quad \int_{\partial \widetilde{E}_{\widetilde{r}_i}} H^2_{\partial \widetilde{E}_{\widetilde{r}_i}} \to (n-1)^2 n \theta \omega_n (1+r_\infty)^{n-3} \,, \qquad \int_{\partial \widetilde{E}_{\widetilde{r}_i}} H_{\partial \widetilde{E}_{\widetilde{r}_i}} \to (n-1) n \theta \omega_n (1+r_\infty)^{n-2} \,.$$

Inserting (4.86), (4.89) and the convergence properties for the profile I_i collected above into (4.85) and taking the limit $i \to \infty$, we get a contradiction, thus completing the proof of (4.75).

(

Appendix A. Quantitative regularity of almost minimizers in almost Euclidean manifolds

The goal of this appendix is to discuss a series of effective regularity estimates for boundaries of isoperimetric sets in Riemannian manifolds satisfying suitable curvature bounds. These estimates play an important role in the context of the present paper.

We note that similar effective regularity estimates for perimeter minimizing hypersurfaces and stable minimal or constant mean curvature hypersurfaces have been considered in the previous literature under different sets of assumptions. We address the reader for instance to [58, 20, 25], without the aim of being exhaustive in this list.

Let us recall the following theorem, which comes directly from results of Anderson (see, e.g., [4, Theorem 3.2]), and Cheeger–Colding [19].

Theorem A.1. For every $n \ge 2$, $\delta > 0$, C > 0, $\alpha \in [0,1)$ there exists $\varepsilon_0 := \varepsilon_0(n, \delta, C, \alpha) < 1$ such that the following holds.

Let (M^n, g, p) be a pointed n-dimensional Riemannian manifold such that

(A.1)
$$-\varepsilon_0 \leq \operatorname{Ric} \leq C \quad on \ B_2(p), \qquad \operatorname{d}_{\operatorname{GH}}(B_2(p), B_2(0^n)) \leq \varepsilon_0.$$

Then there exists a (harmonic) chart around p such that

(A.2)
$$||g_{ij} - \delta_{ij}||_{C^{1,\alpha}(B_1(0^n))} \le \delta$$

We shall also need an ε -regularity result for quasi-minimizers of the perimeter whose boundary has constant mean curvature in manifolds with controlled Ricci curvature. This will imply an ε -regularity result for isoperimetric sets in manifolds with nonnegative Ricci curvature and Euclidean volume growth, see Remark A.7. Let us first recall the definition of (Λ, r) -minimizer.

Definition A.2. Let (M, g) be a Riemannian manifold of dimension n, possibly non complete. Let $\Omega \subset M$ be open, and let $E \subset \Omega$ be a Borel set. Fix $\Lambda, r_0 > 0$. We say that E is a (Λ, r_0) -minimizer in (Ω, g) if

(A.3)
$$P(E, B_r(x)) \le P(F, B_r(x)) + \Lambda \operatorname{vol}(E\Delta F),$$

whenever $E\Delta F \Subset B_r(x) \Subset \Omega$ and $r \le r_0$.

Remark A.3. Let E be a (Λ, r_0) -minimizer in (Ω, g) for some open set Ω as in Definition A.2. Then $P(E, \cdot) = \mathcal{H}^{n-1} \sqcup (\partial E \cap \Omega)$, and ∂E has generalized mean curvature bounded by Λ . More precisely there exists a vector field $H \in L^1_{loc}(\Omega, P(E, \cdot))$ such that

(A.4)
$$\int \operatorname{div}_{\mathsf{T}} X \, \mathrm{d}P(E, \cdot) = -\int g(X, H) \, \mathrm{d}P(E, \cdot) \,,$$

for any vector field $X \in C_c^{\infty}(\Omega)$, and $|H| \leq \Lambda P(E, \cdot)$ -almost everywhere, see, e.g., [51, Lemma A.2] for a proof in the Euclidean space (see also the proof of [2, Theorema 4.7.4]). Indeed, the first assertion follows by standard density estimates (see, e.g., the proof of [42, Theorem 21.11]), while the existence of the generalized mean curvature vector follows by differentiation of the Λ -minimality condition.

The following ε -regularity theorem has essentially appeared in [46, Theorem 6.8]. The version we state here can be proved arguing verbatim.

Proposition A.4. Let $n \in \mathbb{N}$ with $n \geq 2$, $v_0 > 0$, $\Lambda > 0$, and $\varepsilon > 0$ be fixed. Then there exists $\delta > 0$ for which the following holds.

Let (M^n, g, p) be a pointed n-dimensional complete Riemannian manifold with $\operatorname{Ric} \geq -\delta$ on $B_2(p)$, and $|B_2(p)| \geq v_0$. Assume E is a $(\Lambda, 1)$ -minimizer in $B_2(p)$, and

(A.5)
$$\max\left\{ d_{\mathrm{GH}}\left(\partial E \cap B_{2}(p), B_{2}(0^{n-1})\right), d_{\mathrm{GH}}\left(B_{2}(p), B_{2}(0^{n})\right), \left|\frac{|E \cap B_{2}(p)|}{\omega_{n}2^{n}} - \frac{1}{2}\right| \right\} < \delta.$$

Then, for all $q \in B_{1/2}(p) \cap \partial E$, and r < 1 it holds

(A.6)
$$d_{GH} \left(\partial E \cap B_r(q), B_r(0^{n-1}) \right) < \varepsilon r ,$$
$$\left| \frac{P(E, B_r(q))}{\omega_{n-1}r^{n-1}} - 1 \right| < \varepsilon ,$$
$$\left| \frac{|E \cap B_r(q)|}{\omega_n r^n} - \frac{1}{2} \right| < \varepsilon .$$

We are now ready to show an Allard-type result for $(\Lambda, 1)$ -minimizers in manifolds that are C^1 -close to the Euclidean space, and to deduce an ε -regularity result for quasi-minimizer with constant mean curvature boundary in manifolds with controlled Ricci curvature, see Theorem A.6.

Theorem A.5 ($C^{1,\gamma}$ -regularity of almost flat Λ -minimizers on almost Euclidean balls). Fix $n \geq 2, \gamma \in (0,1), \Lambda > 0$. Then there exist $\overline{\varepsilon}, \overline{\sigma} \in (0,1)$, and $\overline{C} > 0$ such that the following holds. Let $B_2(0^n)$ be endowed with a smooth Riemannian metric g such that

(A.7)
$$\|g_{ij} - \delta_{ij}\|_{C^1(B_2(0^n))} \leq \overline{\varepsilon} \,.$$

Let $E \subset B_2(0^n)$ be a $(\Lambda, 1)$ -minimizer in $(B_2(0^n), g)$. Suppose that $0^n \in \partial E$ and that

(A.8)
$$\left|\frac{P(E, B_r^g(0^n))}{\omega_{n-1}r^{n-1}} - 1\right| \le \overline{\varepsilon},$$

for any $r \in (0, 1)$. Then, up to rotation, there exists a Lipschitz function $u : (B_{\overline{\sigma}}(0^{n-1}), g_{eu}) \to \mathbb{R}$ such that

(A.9)
$$|u|/\overline{\sigma} + |\nabla u| \le \overline{C}, \quad \nabla u(0) = 0, \quad |\nabla u(x) - \nabla u(y)| \le \overline{C}|x - y|^{\gamma},$$

(A.10)
$$\partial E \cap B_{\overline{\sigma}}(0^n) = \{(x, u(x)) : x \in B_{\overline{\sigma}}(0^{n-1})\} \cap B_{\overline{\sigma}}(0^n)$$

Proof. We first observe that there exist $\overline{\varepsilon}_1 \in (0, 1)$ and $r_{\Lambda} = r_{\Lambda}(\Lambda, n) \in (0, 1)$ such that if g is a metric on $B_2(0^n)$ with $\|g_{ij} - \delta_{ij}\|_{C^1(B_2(0^n))} \leq \overline{\varepsilon}_1$, then any $(\Lambda, 1)$ -minimizer G in $(B_2(0^n), g)$ satisfies density estimates of the form

(A.11)
$$v_n \le \frac{\operatorname{vol}(G \cap B^g_\rho(x))}{\omega_n \rho^n} \le 1 - v_n, \qquad c_n^{-1} \le \frac{P(G, B^g_\rho(x))}{\omega_{n-1} \rho^{n-1}} \le c_n$$

for some $v_n \in (0, 1)$ and $c_n > 1$ independent of G, g, for any $x \in \partial G \cap B_2(0^n)$ and any $\rho \leq r_\Lambda$ with $B_\rho(x) \Subset B_2(0^n)$ (see, e.g., the proof of [42, Theorem 21.11]).

The proof of the claim will eventually follow from an application of Allard's regularity theorem for varifolds having bounded first variation with respect to a parametric integrand, see [1, The Regularity Theorem p. 27]. We sketch here the main steps leading to the application of the previous result.

It readily follows from the area formula that the perimeter measure of a set $G \subset B_2(0^n)$ with respect to a metric g on $B_2(0^n)$ can be written as $P(G, \cdot) = (\langle \operatorname{Cof}_x(n_G), n_G \rangle)^{\frac{1}{2}} P_{eu}(G, \cdot)$, where $\langle \cdot, \cdot \rangle$ denotes Euclidean scalar product, $P_{eu}(G, \cdot) = \mathcal{H}_{eu}^{n-1} \sqcup \partial^* G$ is the Euclidean perimeter measure, n_g is the unit Euclidean inner normal to G, and Cof_x is the cofactor matrix of the matrix (g_{ij}) at the point x. Hence, if $\overline{\varepsilon}_1 \in (0, 1)$ is small enough, the perimeter functional with respect to a metric g with $\|g_{ij} - \delta_{ij}\|_{C^1(B_2(0^n))} \leq \overline{\varepsilon}_1$ is an admissible functional corresponding to the parametric integrand $\Psi(x, u) = (\langle \operatorname{Cof}_x(u), u \rangle)^{1/2}$, in the notation of [1, 3 p. 18]. Here by admissible we mean that all the constants $M_{i,j}, N_{i,j}$ appearing in [1] corresponding to Ψ are bounded in terms of the dimension n and of $\overline{\varepsilon}_1$ only.

Take A = 0 and $\varepsilon = \lambda = \frac{1}{2}$ in [1, p. 2.2] and recall (A.11). Note that Remark A.3 implies that the first variation with respect to the integrand Ψ is bounded by Λ in the sense of [1, 3.6(4)]. Taking into account [1, Remark p. 28], [1, The Regularity Theorem p. 27] yields a parameter

 $\delta_8 \in (0,1)$ (in the notation of [1]) and constants $\overline{C} = \overline{C}(n,\gamma), c(\Lambda)$ such that if $G \subset B_2(0^n)$ is a $(\Lambda, 1)$ -minimizer for a metric g with $\|g_{ij} - \delta_{ij}\|_{C^1(B_2(0^n))} \leq \overline{\varepsilon}_1$ and

(A.12)
$$\frac{1}{2} < \frac{P(G, B_{\rho/2}(0^n))}{\omega_{n-1}(\rho/2)^{n-1}} < \frac{3}{2}, \qquad |\mathbf{q}x| < \delta_8 \rho \quad \forall x \in \partial G \cap B_{\rho}(0^n),$$

for some $\rho < c(\Lambda) \leq 1$, then $\partial G \cap B_{\overline{\sigma}_2\rho}(0^n)$ is parametrized by the graph of a $C^{1,\gamma}$ function u with bounds as in the statement, for suitable $\overline{\sigma}_2 = \overline{\sigma}_2(n,\gamma) \in (0,\frac{1}{2})$. Above $\mathbf{q} : \mathbb{R}^n \to \mathbb{R}$ is the projection $\mathbf{q}(x_1,\ldots,x_n) := x_n$. We are therefore left to verify (A.12) for a set E satisfying, for suitable choices of $\overline{\varepsilon}, \overline{\sigma}$.

Let p > 1 be such that $\gamma = 1 - (n - 1)/p$ and let $\delta_0, \overline{\sigma}_1 \in (0, 1)$ be the two parameters, depending on n, p only, in the assumptions in the classical Euclidean Allard regularity theorem [59, Theorem 5.2 p. 121]. Here $\overline{\sigma}_1$ denotes the constant called γ in [59, Theorem 5.2 p. 121]. Define $\delta := \frac{1}{2} \min\{\delta_0, (\delta_8 \overline{\sigma}_1/(5C'))^{2n+2}, \Lambda c_n^{\frac{1}{p}} \omega_{n-1}^{\frac{1}{p}}\}$ where $C' = C'(n, \gamma)$ is the constant called Cin [59, Theorem 5.2 p. 121], and $r_{\delta} := \delta/(\Lambda c_n^{\frac{1}{p}} \omega_{n-1}^{\frac{1}{p}}) < 1$. We finally define

$$\overline{\sigma} := \frac{1}{6} \min \left\{ \overline{\sigma}_1 \overline{\sigma}_2 r_\delta, c(\Lambda) \overline{\sigma}_2 \right\}.$$

We want to show that there exist $\overline{\varepsilon} \leq \overline{\varepsilon}_1$ such that if g, E are as in the assumptions, then E satisfies (A.12) with $\rho = \overline{\sigma}/\overline{\sigma}_2$, leading to the proof.

Suppose by contradiction that there exist $\overline{\varepsilon}_i \leq \overline{\varepsilon}_1$ with $\overline{\varepsilon}_i \searrow 0$, and a sequence of metrics g_i and a sequence of $(\Lambda, 1)$ -minimizers E_i as in the assumptions, but $(\Lambda.12)$ with $G = E_i$ and with $\rho = \overline{\sigma}/\overline{\sigma}_2$ does not hold. By (A.11), up to subsequence E_i converges to a nonempty Euclidean $(\Lambda, 1)$ -minimizer F. The fact that F is a Euclidean $(\Lambda, 1)$ -minimizer follows by easily adapting the proof of [42, Theorem 21.14]. Moreover, by convergence of the perimeter measures as in [42, Theorem 21.14] and applying Remark A.3 on the set F, we get

(A.13)
$$\frac{P_{\text{eu}}(F, B_r(0^n))}{\omega_{n-1}r^{n-1}} = 1, \qquad \left(\int_{B_r(0^n)} |H_{\partial F}|^p \, \mathrm{d}P_{\text{eu}}(F, \cdot)\right)^{\frac{1}{p}} r^{1-\frac{n-1}{p}} \stackrel{(\text{A.11})}{\leq} \Lambda c_n^{\frac{1}{p}} \omega_{n-1}^{\frac{1}{p}} r \leq \delta,$$

for any $r \leq r_{\delta}$. Hence the classical Euclidean Allard regularity theorem [59, Theorem 5.2 p. 121] can be applied to F, implying that, up to rotation, $\partial F \cap B_{\overline{\sigma}_1 r}$ is parametrized as the graph of a $C^{1,\gamma}$ function $f: B_{\overline{\sigma}_1 r}(0^{n-1}) \to \mathbb{R}$ with $|f|/r + |\nabla f| \leq C' \delta^{1/(2n+2)}$ and $[\nabla f]_{0,\gamma} \leq C' \delta^{1/(2n+2)} r^{\gamma}$, for any $r \leq r_{\delta}$. Since $3\overline{\sigma}/(\overline{\sigma}_1 \overline{\sigma}_2) < r_{\delta}$, we can choose $r = 3\overline{\sigma}/(\overline{\sigma}_1 \overline{\sigma}_2)$, so that there holds

(A.14)
$$|\mathbf{q}x| \le C' \delta^{1/(2n+2)} r < \frac{3}{5} \delta_8 \frac{\overline{\sigma}}{\overline{\sigma}_2} \qquad \forall x \in \partial F \cap B_{3\overline{\sigma}/\overline{\sigma}_2}(0^n)$$

Since $\partial E_i \to \partial F$ in Kuratowski sense thanks to the uniform density estimates (see, e.g., the proof of [42, Theorem 21.14 (i), (ii)]), the first equalities in (A.13) and (A.14) imply that (A.12) hold on E_i for *i* large enough with $\rho = \overline{\sigma}/\overline{\sigma}_2$, thus yielding a contradiction.

Theorem A.6. Let $n \in \mathbb{N}$ with $n \geq 2$, $C, \Lambda, v_0 > 0$, $H \in \mathbb{R}$, and $\alpha \in (0, 1)$. Then there exist $\varepsilon < 1, \sigma \in (0, 1), B > 0$ such that the following holds.

Let (M^n, g, p) be a pointed n-dimensional complete Riemannian manifold with $-\varepsilon \leq \text{Ric} \leq C$ on $B_2(p)$, and $|B_2(p)| \geq v_0$. Suppose $E \subset M$ is a set such that

E is a $(\Lambda, 1)$ -minimizer in $B_2(p)$, ∂E has a.e. generalized mean curvature $H_{\partial E} \equiv H$, and

(A.15)
$$\max\left\{ d_{\rm GH}(B_2(p), B_2(0^n)), d_{\rm GH}\left(\partial E \cap B_2(p), B_2(0^{n-1})\right), \left|\frac{|E \cap B_2(p)|}{\omega_n 2^n} - \frac{1}{2}\right| \right\} \le \varepsilon.$$

Then, denoting with $\partial_r E$ the regular part of ∂E , it holds $\partial E \cap B_{\sigma}(p) \subset \partial_r E$, and

(A.16)
$$\||\mathrm{II}|\|_{C^{0,\alpha}(\partial E \cap B_{\sigma}(p),\mathrm{d}|_{\partial E})} \coloneqq \sup_{x \in \partial E \cap B_{\sigma}(p)} |\mathrm{II}|(x) + \sup_{x,y \in \partial E \cap B_{\sigma}(p)} \frac{||\mathrm{II}|(x) - |\mathrm{II}|(y)|}{\mathrm{d}(x,y)^{\alpha}} \le B ,$$

where II denotes the second fundamental form of ∂E . In particular, $\partial E \cap B_{\sigma}(p)$ is smooth.

Moreover, if also $|\text{Sect}| \leq K$ on $B_2(p)$, then the (intrinsic) sectional curvature of $\partial E \cap B_{\sigma}(p)$ is bounded from below by a constant $B' = B'(B, K) \in \mathbb{R}$.

Proof. Let $\bar{\varepsilon}, \bar{\sigma}, \bar{C} > 0$ be the constants depending on $\Lambda, \gamma := \alpha, n$ for which Theorem A.5 holds. Let $\varepsilon_0 := \varepsilon_0(n, \bar{\varepsilon}, C, \alpha)$ be the constant for which Theorem A.1 holds. Thus, if $\varepsilon < \varepsilon_0$ in the statement of Theorem A.6, there exists a chart centered at p such that

(A.17)
$$||g_{ij} - \delta_{ij}||_{C^{1,\alpha}(B_1(0^n))} \le \bar{\varepsilon}$$

Finally let $\delta := \delta(\bar{\varepsilon}, n, v_0, \Lambda)$ be the constant appearing in Proposition A.4. We claim that in the statement of Theorem A.6 we can take $\varepsilon := \min\{\delta, \varepsilon_0\}$.

Indeed, by how we chose the constants, and subsequently applying Theorem A.1, Proposition A.4, and Theorem A.5 we get from (A.10) that $\partial E \cap B_{\bar{\sigma}/2}(p)$ is a graph of a $C^{1,\alpha}$ -function u defined on some ball $B_r(0^{n-1}) \subset B_r(0^n)$, with $\bar{\sigma}/4 < r < \bar{\sigma}$, contained in the domain of the chart given by Theorem A.1. Thus $\partial E \cap B_{\sigma}(p) \subset \partial_r E$ with $\sigma := \bar{\sigma}/2$, where $\partial_r E$ denotes the regular part of ∂E .

In order to obtain the Hölder bound on the second fundamental form we exploit a standard PDE argument based on the fact that E has smooth boundary and it has constant mean curvature. Let $u: B_r(0^{n-1}) \to \mathbb{R}$, with $\bar{\sigma}/4 < r < \bar{\sigma}$, be the function defined above such that the graph (x, u(x)) in the chart given by Theorem A.1 is (a subset of) ∂E . Let us define

(A.18)
$$h_{ij} := g_{ij} + \partial_i u \cdot g_{jn} + \partial_j u \cdot g_{in} + \partial_i u \cdot \partial_j u \cdot g_{nn}, \qquad \forall i, j = 1, \dots, n-1$$

Moreover, let $\{e_1, \ldots, e_n\}$ be the canonical basis in \mathbb{R}^n , and let $v_i := (e_i, \partial_i u) \in \mathbb{R}^n$, for every $i = 1, \ldots, n-1$. Denote $\{\hat{v}_i\}_{i=1,\ldots,n-1}$ the orthonormal vectors obtained by applying Gram-Schmidt reduction to $\{v_i\}_{i=1,\ldots,n-1}$. Denote $e_n^{\perp} := e_n - \sum_{i=1}^{n-1} g(e_n, \hat{v}_i)\hat{v}_i$.

A routine computation shows that the mean curvature of the graph of u in the chart is

(A.19)
$$H = g(e_n^{\perp}, e_n^{\perp})^{1/2} \cdot h^{ij} \cdot \partial_{ij} u + Q(g, \partial g, \partial u),$$

where $Q(g, \partial g, \partial u)$ is a rational expression involving $g, \partial g, \partial u$.

Hence, if $||g_{ij} - \delta_{ij}||_{C^0} + ||u||_{C^1} \leq \varepsilon(n)$, where $\varepsilon(n)$ is a universal small constant, (A.19) is an elliptic equation. By taking $\overline{\varepsilon} \leq \varepsilon(n)/2$, which we can always do without loss of generality, and by taking into account (A.7), and the second inequality in (A.9), we have the following. There exists $N := N(n, \alpha, \Lambda) > 1$ large enough such that

(A.20)
$$\|g_{ij} - \delta_{ij}\|_{C^0(B_{r/N}(0^n))} + \|u\|_{C^1(B_{r/N}(0^{n-1}))} \le \varepsilon(n) \, .$$

Hence (A.19) is an elliptic equation on $B_{r/N}(0^{n-1})$. We can thus apply Schauder estimates to get

(A.21)
$$\|u\|_{C^{2,\alpha}(B_{r/(2N)}(0^{n-1}))} \le C\left(r, N, \alpha, |H|, \|g\|_{C^{1,\alpha}(B_{r/N}(0^n))}, \|u\|_{C^{1,\alpha}(B_{r/N}(0^{n-1}))}\right).$$

Finally, writing II in the chart φ , we get (A.16) up to shrinking $\sigma := \bar{\sigma}/(16N)$, and where *B* only depends on $\bar{\varepsilon}$, *r*, *N*, α , \bar{C} , and an upper bound on |H|, and thus ultimately only on α , *n*, Λ , *H*.

The last assertion about the sectional curvature with respect to the induced metric being uniformly bounded readily comes from the Gauss equations, taking into account the uniform bounds for the second fundamental form and the bounds for the sectional curvature of the ambient Riemannian manifold. $\hfill \Box$

Remark A.7 (Applying Theorem A.6 to isoperimetric sets). Theorem A.6 has, as a direct consequence, an ε -regularity result for isoperimetric sets with large volumes in complete manifolds with Ric ≥ 0 , AVR > 0, and quadratic curvature decay (1.2), which is relevant to our aims.

Let (M^n, g) be a smooth Riemannian manifold with Ric ≥ 0 , and with AVR $\geq v_0 > 0$. Choosing $V_1 = \frac{1}{2}$, $V_2 = 2$, $V_3 = 3$, K = 0 in [7, Corollary 4.15], and by a scaling argument, we get that there is $\mathcal{C} := \mathcal{C}(v_0, n) \geq 1$, such that

(A.22) Any isoperimetric set $E \subset M$ with $|E| \ge 1$ is a $(\mathcal{C}|E|^{-1/n}, \mathcal{C}^{-1}|E|^{1/n})$ -minimizer, and moreover

(A.23)
$$\operatorname{Per}(E, B_r(p)) \le (1 + \mathcal{C}|E|^{-1/n}r)\operatorname{Per}(F, B_r(p)),$$

for every $p \in M$, $r < C^{-1}|E|^{1/n}$, and F such that $E\Delta F \subset B_r(p)$.

Now, let $V_1 \ge C^n$ be fixed, and define $\Lambda := CV_1^{-1/n}$. By (A.22), every isoperimetric set E with $|E| \ge V_1$ in M is a $(C|E|^{-1/n}, C^{-1}|E|^{1/n})$ -minimizer, and thus in particular a $(\Lambda, 1)$ -minimizer. Moreover, recall that the boundary of a smooth isoperimetric set has constant mean curvature, and if $|E| \ge V_1$ in M, then $|H_{\partial E}| \le C(n, V_1, v_0)$, see [8, Corollary 3.5].

Taking into account also Lemma 3.2, and the regularity of cross-sections of manifolds with Ric ≥ 0 , AVR > 0 and quadratic curvature decay (1.2), we infer that Theorem A.6 can be applied in this setting, as soon as E is an isoperimetric set with $|E| \geq V_1 \geq C(v_0, n)^n$. In this case the constants ε, σ, B will obviously depend on n, C, V_1, v_0 .

APPENDIX B. UNIFORM BOUNDS FOR THE REACH

The goal of this appendix is to prove that the reach of a smooth hypersurface Σ^{n-1} embedded in a smooth Riemannian manifold M^n can be bounded from below in terms of the second fundamental form of the hypersurface and the norm of the sectional curvature of the ambient manifold, under some minor additional technical assumptions. The statement is probably known to experts, and it follows for instance from a slight adaptation of the arguments in [35]. We provide a detailed proof as we are not able to locate a precise reference in the literature. The argument is due to Petrunin, see [54].

Proposition B.1. Let *E* be a compact subset of an *n*-dimensional smooth Riemannian manifold M^n , and assume that $\partial E := \Sigma^{n-1}$ is a codimension one hypersurface embedded in M^n . Let ν_{Σ} be the unit inner normal. Assume there is $\varepsilon_0 > 0$ such that

(B.1)
$$\exp_p(t\nu_{\Sigma}(p)) \in \begin{cases} E & \text{for all } t \in (0, \varepsilon_0), \\ M \setminus E & \text{for all } t \in (-\varepsilon_0, 0) \end{cases}$$

Let C > 0 be such that

(B.2)
$$\sup_{x \in \Sigma} |II(x)| \le C, \quad \sup_{x \in B_1(\Sigma)} |Riem(x)| \le C,$$

where we denote $B_r(\Sigma) := \{x \in M : d(x, \Sigma) < r\}$ for each r > 0. Then there exists a constant $r_0 = r_0(C, n, \varepsilon_0) > 0$ such that

(B.3)
$$F: (-r_0, r_0) \times \Sigma \to M, \qquad F(t, p) := \exp_p(t\nu_{\Sigma}(p))$$

is a diffeomorphism with its image.

Proof. Let us take C a constant chosen small enough with respect to C, and n. We claim that the choice $r_0 := \min\{\frac{\varepsilon_0}{4}, C\}$ works. First, C can be chosen small enough so that, since $r_0 < C$, by using (B.2) and classical estimates on Jacobi fields, one gets that

$$\mathrm{d}F_{(t,p)}(v) \neq 0$$
 for every $(t,p) \in (-r_0,r_0) \times \Sigma$, and every $0 \neq v \in T_{(t,p)}((-r_0,r_0) \times \Sigma)$.

Thus for every $(t, p) \in (-r_0, r_0) \times \Sigma$ we have that $dF_{(t,p)}$ is a linear isomorphism, and then F is a local diffeomorphism around every point $(t, p) \in (-r_0, r_0) \times \Sigma$. In order to prove (B.3) it is left to prove that F is injective.

Suppose by contradiction F is not injective, and consider

(B.4)
$$\ell := \inf\{0 < \vartheta \le r_0 : F : (-\vartheta, \vartheta) \times \Sigma \to M \text{ is not injective}\} > 0.$$

Hence there are $p, q \in \Sigma$ such that setting $\gamma_p(t) := \exp_p(t\nu_{\Sigma}(p))$, and $\gamma_q(t) := \exp_q(t\nu_{\Sigma}(q))$ we have

$$x = \gamma_p(\ell) = \gamma_q(\ell) \,.$$

We claim that $\gamma'_p(\ell) = -\gamma'_q(\ell)$. Indeed, if not, by taking a geodesic starting from x and forming acute angles with γ_p, γ_q at ℓ , one would get that there is a point z such that $d(z, p) < \ell$, and $d(z,q) < \ell$. Thus, by possibly taking z closer to x, and by exploiting the minimality of ℓ , zcan be chosen so that $z = \exp_{p'}(\ell_p \nu_{\Sigma}(p')) = \exp_{q'}(\ell_q \nu_{\Sigma}(q'))$ for $p', q' \in \Sigma$, with $p' \neq q'$, and $\ell_p, \ell_q < \ell$. This last property would contradict the minimality of ℓ in (B.4).

Hence, since $\gamma'_p(\ell) = -\gamma'_q(\ell)$, we get that $q = \exp_p(2\ell\nu_{\Sigma}(p))$. Thus, by the second property in (B.1) applied at q, we have that

$$\exp_p((2\ell + \varepsilon_0/3)\nu_{\Sigma}(p)) \in M \setminus E$$

which is a contradiction with the first of (B.1) applied at p since $2\ell + \varepsilon_0/3 \le 2r_0 + \varepsilon_0/3 < \varepsilon_0$. \Box

References

- William K. Allard. "An integrality theorem and a regularity theorem for surfaces whose first variation with respect to a parametric elliptic integrand is controlled". In: *Geometric measure theory and the calculus of variations (Arcata, Calif., 1984)*. Vol. 44. Proc. Sympos. Pure Math. Amer. Math. Soc., Providence, RI, 1986, pp. 1–28. ISBN: 0-8218-1470-2. DOI: 10.1090/pspum/044/840267.
- [2] Luigi Ambrosio. Corso introduttivo alla teoria geometrica della misura ed alle superfici minime. Appunti dei Corsi Tenuti da Docenti della Scuola. [Notes of Courses Given by Teachers at the School]. Scuola Normale Superiore, Pisa, 1997, pp. ii+144.
- [3] Luigi Ambrosio and Shouhei Honda. "New stability results for sequences of metric measure spaces with uniform Ricci bounds from below". English. In: *Measure theory in non-smooth spaces*. Warsaw: De Gruyter Open, 2017, pp. 1–51. ISBN: 978-3-11-055082-5; 978-3-11-055083-2.
- [4] Michael T. Anderson. "Convergence and rigidity of manifolds under Ricci curvature bounds". In: *Invent. Math.* 102.2 (1990), pp. 429–445. ISSN: 0020-9910,1432-1297. DOI: 10.1007/BF01233434.
- [5] Gioacchino Antonelli, Elia Bruè, Mattia Fogagnolo, and Marco Pozzetta. "On the existence of isoperimetric regions in manifolds with nonnegative Ricci curvature and Euclidean volume growth". English. In: *Calc. Var. Partial Differ. Equ.* 61.2 (2022). Id/No 77, p. 40. ISSN: 0944-2669. DOI: 10.1007/s00526-022-02193-9.
- [6] Gioacchino Antonelli, Mattia Fogagnolo, and Marco Pozzetta. "The isoperimetric problem on Riemannian manifolds via Gromov-Hausdorff asymptotic analysis". In: Commun. Contemp. Math. 26.1 (2024), Paper No. 2250068, 58. ISSN: 0219-1997,1793-6683. DOI: 10.1142/S0219199722500687.
- [7] Gioacchino Antonelli, Enrico Pasqualetto, Marco Pozzetta, and Daniele Semola. "Sharp isoperimetric comparison on non-collapsed spaces with lower Ricci bounds". To appear: Ann. Sci. École Norm. Sup. 2023. arXiv: 2201.04916.
- [8] Gioacchino Antonelli, Enrico Pasqualetto, Marco Pozzetta, and Daniele Semola. "Asymptotic isoperimetry on non collapsed spaces with lower Ricci bounds". In: *Math. Ann.* 389.2 (2024), pp. 1677–1730. ISSN: 0025-5831,1432-1807. DOI: 10.1007/s00208-023-02674-y.
- [9] Gioacchino Antonelli and Marco Pozzetta. Isoperimetric problem and structure at infinity on Alexandrov spaces with nonnegative curvature. 2023. arXiv: 2302.10091.
- [10] Shigetoshi Bando, Atsushi Kasue, and Hiraku Nakajima. "On a construction of coordinates at infinity on manifolds with fast curvature decay and maximal volume growth". In: *Invent. Math.* 97.2 (1989), pp. 313–349. ISSN: 0020-9910,1432-1297. DOI: 10.1007/BF01389045.

REFERENCES

- [11] Christophe Bavard and Pierre Pansu. "Sur le volume minimal de \mathbb{R}^2 . (On the minimal volume of \mathbb{R}^2)". French. In: Ann. Sci. Éc. Norm. Supér. (4) 19.4 (1986), pp. 479–490. ISSN: 0012-9593. DOI: 10.24033/asens.1514.
- [12] Vincent Bayle. "A differential inequality for the isoperimetric profile". English. In: Int. Math. Res. Not. 2004.7 (2004), pp. 311–342. ISSN: 1073-7928. DOI: 10.1155/S1073792804130079.
- [13] Hubert Lewis Bray. The Penrose inequality in general relativity and volume comparison theorems involving scalar curvature. Thesis (Ph.D.)–Stanford University. ProQuest LLC, Ann Arbor, MI, 1997, p. 103. ISBN: 978-0591-60594-5.
- Simon Brendle. "Constant mean curvature surfaces in warped product manifolds". English. In: Publ. Math., Inst. Hautes Étud. Sci. 117 (2013), pp. 247–269. ISSN: 0073-8301.
 DOI: 10.1007/s10240-012-0047-5.
- [15] Simon Brendle. "Sobolev Inequalities in Manifolds with Nonnegative Curvature". In: Communications on Pure and Applied Mathematics 76.9 (2023), pp. 2192-2218. DOI: https://doi.org/10.1002/cpa.22070.eprint: https://onlinelibrary.wiley.com/doi/pdf/10.
- [16] Simon Brendle and Michael Eichmair. "Isoperimetric and Weingarten surfaces in the Schwarzschild manifold". English. In: J. Differ. Geom. 94.3 (2013), pp. 387–407. ISSN: 0022-040X. DOI: 10.4310/jdg/1370979333.
- Simon Brendle and Michael Eichmair. "Large outlying stable constant mean curvature spheres in initial data sets". English. In: *Invent. Math.* 197.3 (2014), pp. 663–682. ISSN: 0020-9910. DOI: 10.1007/s00222-013-0494-8.
- [18] Dmitri Burago and Sergei Ivanov. "Boundary rigidity and filling volume minimality of metrics close to a flat one". In: Ann. of Math. (2) 171.2 (2010), pp. 1183–1211. ISSN: 0003-486X,1939-8980. DOI: 10.4007/annals.2010.171.1183.
- [19] Jeff Cheeger and Tobias H. Colding. "Lower bounds on Ricci curvature and the almost rigidity of warped products". In: Ann. of Math. (2) 144.1 (1996), pp. 189–237. ISSN: 0003-486X,1939-8980. DOI: 10.2307/2118589.
- [20] Jeff Cheeger and Aaron Naber. "Quantitative stratification and the regularity of harmonic maps and minimal currents". English. In: Commun. Pure Appl. Math. 66.6 (2013), pp. 965–990. ISSN: 0010-3640. DOI: 10.1002/cpa.21446.
- [21] Otis Chodosh and Michael Eichmair. "On far-outlying constant mean curvature spheres in asymptotically flat Riemannian 3-manifolds". English. In: J. Reine Angew. Math. 767 (2020), pp. 161–191. ISSN: 0075-4102. DOI: 10.1515/crelle-2019-0034.
- [22] Otis Chodosh and Michael Eichmair. "Global uniqueness of large stable CMC spheres in asymptotically flat Riemannian 3-manifolds". English. In: *Duke Math. J.* 171.1 (2022), pp. 1–31. ISSN: 0012-7094. DOI: 10.1215/00127094-2021-0043.
- [23] Otis Chodosh, Michael Eichmair, Yuguang Shi, and Jintian Zhu. "Characterization of large isoperimetric regions in asymptotically hyperbolic initial data". In: Comm. Math. Phys. 368.2 (2019), pp. 777–798. ISSN: 0010-3616. DOI: 10.1007/s00220-019-03354-2.
- [24] Otis Chodosh, Michael Eichmair, and Alexander Volkmann. "Isoperimetric structure of asymptotically conical manifolds". In: J. Differential Geom. 105.1 (2017), pp. 1–19. ISSN: 0022-040X,1945-743X. DOI: 10.4310/jdg/1483655857.
- [25] Otis Chodosh, Chao Li, and Douglas Stryker. Complete Stable Minimal Hypersurfaces in Positively Curved 4-manifolds. To appear: J. Eur. Math. Soc. arXiv: 2202.07708.
- [26] Tobias H. Colding. "Ricci curvature and volume convergence". In: Ann. of Math. (2) 145.3 (1997), pp. 477–501. ISSN: 0003-486X,1939-8980. DOI: 10.2307/2951841.
- [27] Tobias H. Colding and Aaron Naber. "Characterization of tangent cones of noncollapsed limits with lower Ricci bounds and applications". In: *Geom. Funct. Anal.* 23.1 (2013), pp. 134–148. ISSN: 1016-443X,1420-8970. DOI: 10.1007/s00039-012-0202-7.
- [28] Michael Eichmair and Jan Metzger. "Large isoperimetric surfaces in initial data sets".
 English. In: J. Differ. Geom. 94.1 (2013), pp. 159–186. ISSN: 0022-040X. DOI: 10.4310/jdg/136188906

REFERENCES

- [29] Michael Eichmair and Jan Metzger. "Unique isoperimetric foliations of asymptotically flat manifolds in all dimensions". English. In: *Invent. Math.* 194.3 (2013), pp. 591–630. ISSN: 0020-9910. DOI: 10.1007/s00222-013-0452-5.
- [30] Matthias Erbar, Kazumasa Kuwada, and Karl-Theodor Sturm. "On the equivalence of the entropic curvature-dimension condition and Bochner's inequality on metric measure spaces". In: *Invent. Math.* 201.3 (2015), pp. 993–1071. ISSN: 0020-9910,1432-1297. DOI: 10.1007/s00222-014-0563-7.
- [31] Nicola Gigli, Andrea Mondino, and Giuseppe Savaré. "Convergence of pointed non-compact metric measure spaces and stability of Ricci curvature bounds and heat flows". English. In: Proc. Lond. Math. Soc. (3) 111.5 (2015), pp. 1071–1129. ISSN: 0024-6115. DOI: 10.1112/plms/pdv047.
- [32] Karsten Grosse-Brauckmann. "Stable constant mean curvature surfaces minimize area".
 English. In: Pac. J. Math. 175.2 (1996), pp. 527–534. ISSN: 1945-5844. DOI: 10.2140/pjm.1996.175.55
- [33] Pengfei Guan, Junfang Li, and Mu-Tao Wang. "A volume preserving flow and the isoperimetric problem in warped product spaces". English. In: *Trans. Am. Math. Soc.* 372.4 (2019), pp. 2777–2798. ISSN: 0002-9947. DOI: 10.1090/tran/7661.
- [34] Juha Heinonen, Pekka Koskela, Nageswari Shanmugalingam, and Jeremy T. Tyson. Sobolev spaces on metric measure spaces. Vol. 27. New Mathematical Monographs. An approach based on upper gradients. Cambridge University Press, Cambridge, 2015, pp. xii+434. ISBN: 978-1-107-09234-1. DOI: 10.1017/CB09781316135914.
- [35] Ernst Heintze and Hermann Karcher. "A general comparison theorem with applications to volume estimates for submanifolds". English. In: Ann. Sci. Éc. Norm. Supér. (4) 11.4 (1978), pp. 451–470. ISSN: 0012-9593. DOI: 10.24033/asens.1354.
- Shouhei Honda. Elliptic PDEs on compact Ricci limit spaces and applications. English.
 Vol. 1211. Mem. Am. Math. Soc. Providence, RI: American Mathematical Society (AMS),
 2018. ISBN: 978-1-4704-2854-9; 978-1-4704-4417-4. DOI: 10.1090/memo/1211.
- [37] Gerhard Huisken and Shing-Tung Yau. "Definition of center of mass for isolated physical systems and unique foliations by stable spheres with constant mean curvature". English. In: *Invent. Math.* 124.1-3 (1996), pp. 281–311. ISSN: 0020-9910. DOI: 10.1007/s002220050054.
- [38] Daniel Ketover, Fernando C. Marques, and André Neves. "The catenoid estimate and its geometric applications". In: J. Differential Geom. 115.1 (2020), pp. 1–26. ISSN: 0022-040X. DOI: 10.4310/jdg/1586224840.
- [39] Christian Ketterer. "Cones over metric measure spaces and the maximal diameter theorem". In: J. Math. Pures Appl. (9) 103.5 (2015), pp. 1228–1275. ISSN: 0021-7824,1776-3371. DOI: 10.1016/j.matpur.2014.10.011.
- [40] Christian Ketterer. "Obata's rigidity theorem for metric measure spaces". In: Anal. Geom. Metr. Spaces 3.1 (2015), pp. 278–295. ISSN: 2299-3274. DOI: 10.1515/agms-2015-0016.
- [41] Nina Lebedeva and Anton Petrunin. Curvature tensor of smoothable Alexandrov spaces. 2022. arXiv: 2202.13420.
- [42] Francesco Maggi. Sets of finite perimeter and geometric variational problems. Vol. 135. Cambridge Studies in Advanced Mathematics. An introduction to geometric measure theory. Cambridge University Press, Cambridge, 2012, pp. xx+454. ISBN: 978-1-107-02103-7. DOI: 10.1017/CB09781139108133.
- [43] William H. Meeks III, Joaquín Pérez, and Antonio Ros. "Stable constant mean curvature surfaces". In: *Handbook of geometric analysis. No. 1*. Vol. 7. Adv. Lect. Math. (ALM). Int. Press, Somerville, MA, 2008, pp. 301–380. ISBN: 978-1-57146-130-8.
- [44] William H. Meeks III, Joaquín Pérez, and Antonio Ros. "Limit leaves of an H lamination are stable". In: J. Differential Geom. 84.1 (2010), pp. 179–189. ISSN: 0022-040X,1945-743X.

REFERENCES

- [45] Andrea Mondino and Stefano Nardulli. "Existence of isoperimetric regions in non-compact Riemannian manifolds under Ricci or scalar curvature conditions". In: Comm. Anal. Geom. 24.1 (2016), pp. 115–138. ISSN: 1019-8385,1944-9992. DOI: 10.4310/CAG.2016.v24.n1.a5.
- [46] Andrea Mondino and Daniele Semola. "Weak Laplacian bounds and minimal boundaries on non smooth spaces with Ricci curvature lower bounds". To appear: Memoirs of the AMS. 2021. arXiv: 2107.12344.
- [47] Frank Morgan and Manuel Ritoré. "Isoperimetric regions in cones". In: Trans. Amer. Math. Soc. 354.6 (2002), pp. 2327–2339. DOI: 10.1090/S0002-9947-02-02983-5.
- [48] Frank Morgan and Antonio Ros. "Stable constant-mean-curvature hypersurfaces are area minimizing in small L¹ neighborhoods". English. In: Interfaces Free Bound. 12.2 (2010), pp. 151–155. ISSN: 1463-9963. DOI: 10.4171/IFB/230.
- [49] Frank Pacard and Taoniu Sun. "Doubling construction for CMC hypersurfaces in Riemannian manifolds". Preprint https://perso.pages.math.cnrs.fr/users/frank.pacard/Public
- [50] Pierre Pansu. "Sur la régularité du profil isopérimétrique des surfaces riemanniennes compactes". In: Ann. Inst. Fourier (Grenoble) 48.1 (1998), pp. 247–264. ISSN: 0373-0956,1777-5310. DOI: 10.5802/aif.1617.
- [51] Giulio Pascale and Marco Pozzetta. *Quantitative isoperimetric inequalities for classical capillarity problems.* 2024. arXiv: 2402.04675.
- [52] Grigorij Perelman. "A complete Riemannian manifold of positive Ricci curvature with Euclidean volume growth and nonunique asymptotic cone". In: Comparison geometry (Berkeley, CA, 1993–94). Vol. 30. Math. Sci. Res. Inst. Publ. Cambridge Univ. Press, Cambridge, 1997, pp. 165–166. ISBN: 0-521-59222-4.
- [53] Peter Petersen. *Riemannian geometry*. Second. Vol. 171. Graduate Texts in Mathematics. Springer, New York, 2006, pp. xvi+401. ISBN: 978-0387-29246-5; 0-387-29246-2.
- [54] Anton Petrunin. Mathoverflow post at https://mathoverflow.net/questions/425982/existence-
- [55] Jie Qing and Gang Tian. "On the uniqueness of the foliation of spheres of constant mean curvature in asymptotically flat 3-manifolds". English. In: J. Am. Math. Soc. 20.4 (2007), pp. 1091–1110. ISSN: 0894-0347. DOI: 10.1090/S0894-0347-07-00560-7.
- [56] Manuel Ritoré. "The isoperimetric problem in complete surfaces of nonnegative curvature." English. In: J. Geom. Anal. 11.3 (2001), pp. 509–517. ISSN: 1050-6926. DOI: 10.1007/BF02922017.
- [57] Antonio Ros and Harold Rosenberg. "Properly embedded surfaces with constant mean curvature". In: Amer. J. Math. 132.6 (2010), pp. 1429–1443. ISSN: 0002-9327,1080-6377.
- [58] Harold Rosenberg, Rabah Souam, and Eric Toubiana. "General curvature estimates for stable H-surfaces in 3-manifolds applications". English. In: J. Differ. Geom. 84.3 (2010), pp. 623–648. ISSN: 0022-040X. DOI: 10.4310/jdg/1279114303.
- [59] Leon Simon. Lectures on geometric measure theory. Vol. 3. Proceedings of the Centre for Mathematical Analysis, Australian National University. Revised version - Tsinghua Lectures 2014. Australian National University, Centre for Mathematical Analysis, Canberra, 1983, pp. vii+272. ISBN: 0-86784-429-9.
- [60] Brian White. "A strong minimax property of nondegenerate minimal submanifolds". English. In: J. Reine Angew. Math. 457 (1994), pp. 203–218. ISSN: 0075-4102. DOI: 10.1515/crll.1994.45

GIOACCHINO ANTONELLI

Courant Institute Of Mathematical Sciences (NYU), 251 Mercer Street, 10012, New York, USA

Email address: ga2434@nyu.edu

Marco Pozzetta

DIPARTIMENTO DI MATEMATICA E APPLICAZIONI "RENATO CACCIOPPOLI", UNIVERSITÁ DEGLI STUDI DI NAPOLI FEDERICO II, VIA CINTIA - MONTE SANT'ANGELO, 80126 NAPOLI, ITALY Email address: marco.pozzetta@unina.it

DANIELE SEMOLA

ETH Zürich, FIM-Forschungsinstitut für Mathematik, Rämistrasse 101, 8092, Zürich, Switzer-Land

Email address: daniele.semola.math@gmail.com